ANATOMY AND PHYSIOLOGY (BOUNDLESS)



Boundless (now LumenLearning)

LibreTexts™

Book: Anatomy and Physiology (Boundless)

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/19/2026

TABLE OF CONTENTS

Licensing

1: Introduction to Anatomy and Physiology

1.1: Overview of Anatomy and Physiology

1.1A: Defining Anatomy

1.1B: Defining Physiology

1.2: Life

1.2A: Life Functions

1.2B: Survival Needs

1.2C: Levels of Organization

1.3: Homeostasis

1.3A: Homeostatic Control

1.3B: Disease as Homeostatic Imbalance

1.4: Mapping the Body

1.4A: Anatomical Position

1.4B: Directional Terms

1.4C: Regional Terms and Axes

1.4D: Body Planes and Sections

1.4E: Body Cavities

1.4F: Abdominopelvic Regions

Index

2: The Chemical Building Blocks of Life

2.1: Atomic Structure

2.1A: Overview of Atomic Structure

2.1B: Atomic Number and Mass Number

2.2: Chemical Bonds

2.2A: Covalent Bonds and Other Bonds and Interactions

2.2B: Ions and Ionic Bonds

2.2C: The Chemical Basis for Life

2.3: Inorganic Compounds

2.3A: pH, Buffers, Acids, and Bases

2.3B: Water's Cohesive and Adhesive Properties

2.3C: Water's High Heat Capacity

2.3D: Water's States: Gas, Liquid, and Solid

2.3E: Water's Solvent Properties

2.4: Organic Compounds

2.4A: Organic Molecules and Functional Groups
2.4B: Types of Biological Macromolecules
2.4C: Dehydration Synthesis

3: Organization at the Cellular Level

3.1: Introduction to the Study of Cells
3.1A: Cells as the Basic Unit of Life

3.2: Cell Membranes and the Fluid Mosaic Model
3.2A: Components of Plasma Membranes
3.2B: Fluid Mosaic Model
3.2C: Membrane Fluidity

3.3: Transport Across Membranes
3.3A: Diffusion
3.3B: Osmosis
3.3C: Tonicity
3.3D: Facilitated transport
3.3E: The Role of Passive Transport
3.3F: Primary Active Transport
3.3G: Electrochemical Gradient
3.3H: Secondary Active Transport
3.3I: Endocytosis

3.4: Cell Signaling
3.4A: Types of Receptors

3.5: The Nucleus and Ribosomes
3.5A: The Nucleus and Ribosomes
3.5B: Characteristics of Eukaryotic Cells

3.6: Organelles
3.6A: The Plasma Membrane and the Cytoplasm
3.6B: The Endoplasmic Reticulum
3.6C: The Golgi Apparatus
3.6D: Lysosomes
3.6E: Peroxisomes
3.6F: Mitochondria

3.7: The Cytoskeleton
3.7A: Intermediate Filaments and Microtubules
3.7B: Microfilaments

3.8: External Cellular Components
3.8A: Extracellular Matrix of Animal Cells
3.8B: Intercellular Junctions

4: Organization at the Tissue Level

4.1: Introduction to Tissues

4.1A: Tissues in Levels of Organization

4.2: Epithelial Tissue

4.2A: Characteristics of Epithelial Tissue

4.2B: Types of Epithelial Tissue

4.3: Connective Tissue

4.3A: Characteristics of Connective Tissue

4.3B: Types of Connective Tissue

4.4: Nervous Tissue

4.4A: Characteristics of Nervous Tissue

4.4B: Types of Nervous Tissue

4.5: Muscular Tissue

4.5A: Characteristics of Muscle Tissue

4.6: Tissue Membranes

4.6A: Epithelial Membranes

4.6B: Synovial Membranes

5: Integumentary System

5.1: The Skin

5.1A: What is Skin?

5.1B: Structure of the Skin: Epidermis

5.1C: Structure of the Skin: Dermis

5.1D: Skin Color

5.2: Cell Junctions

5.2A: Tight Junctions

5.2B: Adherens Junctions

5.3C: Gap Junctions

5.3: Accessory Structures of the Skin

5.3A: Sweat (Sudoriferous) Glands

5.3B: Sebaceous (Oil) Glands

5.3C: Nails

5.3D: Hair

5.4: Functions of the Integumentary System

5.4A: Protection

5.4B: Thermoregulation

5.4C: Cutaneous Sensation

5.4D: Metabolic Functions

5.4E: Blood Supply to the Epidermis

5.4F: Excretion and Absorption

5.5: Wound Healing

5.5A: Steps of Tissue Repair

5.5B: Epidermal Wound Healing

5.5C: Deep Wound Healing

5.6: Overview of Cancer

5.6A: Overview of Cancer

5.6B: Cell Signaling and Cell Death

6: Skeletal System

6.1: Overview of the Skeletal System

6.1A: Overview of the Musculoskeletal System

6.1B: The Axial Skeleton

6.1C: The Appendicular Skeleton

6.2: Cartilage

6.2A: Structure, Type, and Location of Cartilage

6.2B: Cartilage Growth

6.3: Introduction to Bone

6.3A: Gross Anatomy

6.3B: Supply of Blood and Nerves to Bone

6.3C: Microscopic Anatomy of Bone

6.3D: Chemical Composition of Bone

6.4: Bone Formation

6.4A: Embryonic and Fetal Bone Formation

6.4B: Postnatal Bone Growth

6.4C: Bone Remodeling

6.4D: Bone Repair

6.5: Bone and Calcium

6.5A: Bone and Calcium Homeostasis

7: Skeletal System - Parts of the Skeleton

7.1: The Skull

7.1A: General Features and Functions of the Skull

7.1B: Cranial Bones

7.1C: Facial Bones

7.1D: Orbits

7.1E: Foramina

7.1F: Fontanelles

7.1F: Sutures

7.1G: Paranasal Sinuses

7.2: The Hyoid Bone
7.2A: The Hyoid Bone

7.3: The Vertebral Column
7.3A: General Characteristics of the Spine
7.3B: Parts of a Vertebra
7.3C: Regional Vertebral Characteristics

7.4: The Thorax
7.4A: Thoracic Cage: Sternum
7.4B: Thoracic Cage: Ribs

7.5: The Shoulder
7.5A: Clavicle
7.5B: Scapula

7.6: The Upper Limb
7.6A: Humerus (The Upper Arm)
7.6B: Ulna and Radius (The Forearm)
7.6C: Carpals, Metacarpals, and Phalanges (The Hand)

7.7: The Hip
7.7A: Ilium
7.7B: Ischium
7.7C: Pubis
7.7D: False and True Pelves
7.7E: Comparison of Female and Male Pelves
7.7F: Pelvic Structure and Childbearing

7.8: The Lower Limb
7.8A: Femur (The Thigh)
7.8B: Patella (The Knee)
7.8C: Tibia and Fibula (The Leg)
7.8D: Tarsals, Metatarsals, and Phalanges (The Foot)
7.9E: Arches of the Feet

8: Joints

8.1: Classification of Joints
8.1A: Structural Classification of Joints
8.1B: Functional Classification of Joints

8.2: Fibrous Joints
8.2A: Fibrous Joints
8.2B: Sutures
8.2C: Syndesmoses
8.2D: Gomphoses

8.3: Cartilaginous Joints
8.3A: Cartilaginous Joints: Synchodroses
8.3B: Cartilaginous Joints: Symphyses

8.4: Synovial Joints
8.4A: Structure of Synovial Joints
8.4B: Nerve and Blood Supply
8.4C: Bursae and Tendon Sheaths
8.4D: Stability and Range of Motion at Synovial Joints
8.4E: Synovial Joint Movements
8.4E: Types of Synovial Joints

9: Muscular System

9.10: Muscles of the Lower Limb
9.10A: Muscles that Cause Movement at the Hip Joint
9.10B: Muscles that Cause Movement at the Knee Joint
9.10C: Muscles that Cause Movement at the Foot

9.1: Introduction to the Nervous System
9.1A: Structure and Function of the Muscular System
9.1B: Slow-Twitch and Fast-Twitch Muscle Fibers
9.1C: Sliding Filament Model of Contraction
9.1D: ATP and Muscle Contraction
9.1E: Control of Muscle Tension

9.2: Smooth Muscle
9.2A: Skeletal Muscle Fibers

9.3: Control of Muscle Tension
9.3A: Force of Muscle Contraction
9.3B: Velocity and Duration of Muscle Contraction
9.3C: Motor Units
9.3D: Muscle Tone
9.3E: Types of Muscle Contractions: Isotonic and Isometric

9.4: Muscle Metabolism
9.4A: Muscle Metabolism
9.4B: Muscle Fatigue

9.5: Exercise and Skeletal Muscle Tissue
9.5A: Impacts of Exercise on Muscles

9.6: Overview of the Muscular System
9.6A: Interactions of Skeletal Muscles
9.6B: How Skeletal Muscles Are Named
9.6C: How Skeletal Muscles Produce Movements

9.6D: Muscle Attachment Sites
9.6E: Arrangement of Fascicles
9.6F: Lever Systems

9.7: Head and Neck Muscles
9.7A: Facial Muscles
9.7B: Chewing Muscles
9.7C: Neck Muscles

9.8: Trunk Muscles
9.8A: Anterior Muscles
9.8B: Posterior Muscles

9.9: Muscles of the Upper Limb
9.9A: Muscles of the Humerus that Act on the Forearm
9.9B: Muscles of the Wrist and Hand
9.9C: Muscles of the Shoulder
9.9D: Muscles that Cause Movement at the Ankle

10: Overview of the Nervous System

10.1: Introduction to the Nervous System
10.1A: Organization of the Nervous System
10.1B: Functions of the Nervous System
10.1C: Subdivisions of the Nervous System

10.2: Neuroglia
10.2A: Neuroglia of the Central Nervous System
10.2B: Neuroglia of the Peripheral Nervous System

10.3: Neurons
10.3A: Structural Diversity of Neurons
10.3B: Classification of Neurons

10.4: Collections of Nervous Tissue
10.4A: Clusters of Neuronal Cell Bodies
10.4B: Axon Bundles
10.4C: Gray and White Matter

10.5: Neurophysiology
10.5A: Principles of Electricity
10.5B: Ion Channels
10.5D: Membrane Potentials as Signals
10.5D: Resting Membrane Potentials
10.5E: The Action Potential and Propagation
10.5F: The Synapse
10.5G: Postsynaptic Potentials and Their Integration at the Synapse

10.5H: Synaptic Transmission
10.5I: Types of Neurotransmitters by Function
10.5J: Ionotropic and Metabotropic Receptors
10.5K: Serial and Parallel Processing

Index

11: Central Nervous System

11.1: The Spinal Cord

11.1A: Overview of the Spinal Cord
11.1B: The Spine
11.1C: Spinal Cord Grey Matter and Spinal Roots
11.1D: Spinal Cord White Matter

11.2: The Brain

11.2A: The Brain

11.3: Protection of the Brain

11.3A: Blood-Brain Barrier
11.3B: Cerebrospinal Fluid and Its Circulation
11.3C: Ventricles

11.4: The Brain Stem

11.4A: Functions of the Brain Stem
11.4B: Medulla Oblongata
11.4C: Pons
11.4D: Midbrain
11.4E: Reticular Formation

11.5: The Cerebellum

11.5A: Parts of the Cerebellum
11.5B: Functions of the Cerebellum

11.6:The Diencephalon

11.6A: Functions of the Diencephalon
11.6B: Thalamus
11.6C: Hypothalamus
11.6D: Epithalamus and Pineal Gland
11.6E: Circumventricular Organs

11.7: The Cerebrum

11.7A: Overview of the Cerebrum
11.7B: Cerebral Lobes
11.7C: White Matter of the Cerebrum
11.7D: Basal Ganglia
11.7E: Limbic System

11.8: Functional Systems of the Cerebral Cortex
11.8A: Sensory Areas
11.8B: Motor Areas
11.8C: Association Areas
11.8D: Hemispheric Lateralization

Index

12: Peripheral Nervous System

12.10: Reflexes
12.10A: Components of a Reflex Arc
12.10C: Spinal Reflexes

12.11: Pain
12.11A: Pain Sensation
12.11B: Localization of Pain

12.1: Sensation
12.1A: Overview of Sensation
12.1B: Sensation to Perception
12.1C: Sensory Modalities

12.2: Sensory Receptors
12.2A: Classification of Receptors by Stimulus
12.2B: Classification of Receptors by Location

12.3: The Somatosensory System
12.3A: General Organization of the Somatosensory System
12.3B: Tactile Sensation
12.3C: Proprioceptive Sensations
12.3D: Somatic Sensory Pathways
12.3E: Mapping the Primary Somatosensory Area
12.3F: Somatic Sensory Pathways to the Cerebellum

12.4: Nerves
12.4A: Structure of a Nerve
12.4B: Classification of Nerves

12.5: Cranial Nerves
12.5A: Brief Overview of Cranial Nerves
12.5B: Olfactory (I) Nerve
12.5C: Optic (II) Nerve
12.5D: Oculomotor (III) Nerve
12.5E: Trochlear (IV) Nerve
12.5F: Trigeminal (V) Nerve
12.5G: Abducens (VI) Nerve
12.5H: Facial (VII) Nerve

12.5I: Vestibulocochlear (VIII) Nerve
12.5J: Glossopharyngeal (IX) Nerve
12.5K: Vagus (X) Nerve
12.5L: Accessory (XI) Nerve
12.5M: Hypoglossal (XII) Nerve

12.6: Spinal Nerves
12.6A: Overview of the Spinal Nerves
12.6B: Branches of Spinal Nerves
12.6C: Plexuses
12.6D: Intercostal Nerves
12.6E: Dermatomes
12.6F: Function and Physiology of the Spinal Nerves

12.7: Distribution of Spinal Nerves
12.7A: Cervical Plexus
12.7B: Brachial Plexus
12.7C: Lumbar Plexus
12.7D: Sacral and Coccygeal Plexuses
12.7E: Sensory and Motor Tracts

12.8: Motor Activity
12.8A: Peripheral Motor Endings
12.8B: Overview of Motor Integration

12.9: Motor Pathways
12.9A: Organization of Motor Neuron Pathways
12.9B: The Role of the Basal Ganglia in Movement
12.9C: Modulation of Movement by the Cerebellum
12.9D: Functions of the Cerebellum in Integrating Movements

13: Special Senses

13.1: The Vertebrate Visual System
13.1A: Anatomy of the Eye
13.1B: Transduction of Light
13.1C: Visual Processing

13.2: Taste and Olfaction
13.2A: Tastes and Odors
13.2B: Reception and Transduction

13.3: Hearing and Equilibrium
13.3A: Sound
13.3B: The Vestibular System
13.3C: Reception of Sound
13.3D: Transduction of Sound

14: Autonomic Nervous System

14.1: Introduction to the Autonomic Nervous System

14.1A: Comparing the Somatic and Autonomic Nervous Systems

14.1B: Divisions of the Autonomic Nervous System

14.2: Structure of the Autonomic Nervous System

14.2A: Preganglionic Neurons

14.2B: Autonomic Ganglia

14.2C: Postganglionic Neurons

14.2D: Autonomic Plexuses

14.2E: Parasympathetic (Craniosacral) Division

14.2F: Autonomic Reflexes

14.2G: Sympathetic Nervous System

14.3: Functions of the Autonomic Nervous System

14.3A: Sympathetic Responses

14.3B: Parasympathetic Responses

14.3C: Autonomic Interactions

14.3D: Control of Autonomic Nervous System Function

14.4: Neurotransmitters and Receptors

14.4A: Cholinergic Neurons and Receptors

14.4B: Adrenergic Neurons and Receptors

14.4C: Agonists, Antagonists, and Drugs

15: Endocrine System

15.10: The Adrenal (Suprarenal) Glands

15.10A: Overview of the Adrenal Glands

15.10B: Adrenal Cortex

15.10C: Adrenal Medulla

15.11: The Pancreas

15.11A: Overview of Pancreatic Islets

15.11B: Types of Cells in the Pancreas

15.11C: Insulin Secretion and Regulation of Glucagon

15.12: Other Endocrine Tissues and Organs

15.13: Stress

15.1: Overview of the Endocrine System

15.1A: Overview of the Endocrine System

15.1B: Comparing the Nervous and Endocrine Systems

15.2: Hormones

15.2A: Mechanisms of Hormone Action

15.2B: Hormone Receptors

15.2C: Chemistry of Hormones

15.2D: Transport of Hormones

15.3: Mechanisms of Hormone Action

15.3A: Direct Gene Activation and the Second-Messenger System

15.3B: Target Cell Specificity

15.3C: Onset, Duration, and Half-Life of Hormone Activity

15.4: Hormone Interactions

15.4A: Interactions of Hormones at Target Cells

15.5: Hormone Secretion Control

15.5A: Control of Hormone Secretion

15.6: Endocrine Glands

15.6A: Overview of Endocrine Glands

15.7: The Pituitary Gland

15.7A: Overview of the Pituitary Gland

15.7B: Control of the Pituitary Gland by the Hypothalamus

15.7C: The Anterior Pituitary

15.7D: The Posterior Pituitary

15.8: The Thyroid Gland

15.8A: Overview of the Thyroid Gland

15.8B: Formation, Storage, and Release of Thyroid Hormones

15.8C: Action of Thyroid Hormones

15.8D: Control of Thyroid Hormone Release

15.9: The Parathyroid Glands

15.9A: Overview of the Parathyroid Glands

15.9B: Parathyroid Hormone

16: Cardiovascular System - Blood

16.1: Overview of Blood

16.1A: Components of Blood

16.1B: Physical Characteristics and Volume

16.1C: Functions of Blood

16.1D: Blood Plasma

16.2: Red Blood Cells

16.2A: RBC Anatomy

16.2B: RBC Physiology

16.2C: RBC Life Cycle

16.3: White Blood Cells

16.3A: Types of WBCs

16.3B: WBC Function

16.3C: WBC Formation

16.4: Platelets
16.4A: Platelets
16.4B: Platelet Formation

16.5: Hemostasis
16.5A: Overview of Hemostasis
16.5B: Vascular Spasm
16.5C: Platelet Plug Formation
16.5D: Coagulation
16.5E: Role of Vitamin K
16.5F: Clot Retraction and Repair
16.5G: Fibrinolysis

16.6: Transfusions
16.6A: Transfusions of Whole Blood
16.6B: Plasma and Blood Volume Expanders
16.6C: Blood Groups and Blood Types
16.6D: Typing and Cross-Matching for Transfusions

17: Cardiovascular System: The Heart

17.1: The Heart
17.1A: Anatomy of the Heart
17.1B: Pericardium
17.1C: Layers of the Heart Walls
17.1D: Chambers of the Heart
17.1E: Great Vessels of the Heart
17.1F: Myocardial Thickness and Function
17.1G: Fibrous Skeleton of the Heart

17.2: Circulation and Heart Valves
17.2A: Heart Circulation
17.2B: Operation of Atrioventricular Valves
17.2C: Operation of Semilunar Valves
17.2D: Systemic and Pulmonary Circulation

17.3: Cardiac Muscle Tissue
17.3A: Mechanism and Contraction Events of Cardiac Muscle Fibers
17.3B: Microscopic Anatomy
17.3C: Energy Requirements

17.4: Physiology of the Heart
17.4A: Electrical Events
17.4B: Electrocardiogram and Correlation of ECG Waves with Systole
17.4C: Heart Sounds
17.4D: Cardiac Cycle

17.4E: Cardiac Output

17.5: Exercise and the Heart

17.5A: Effects of Exercise on the Heart

Section 6:

18: Cardiovascular System: Blood Vessels

18.10: Circulatory Routes

18.10A: The Aorta and Its Branches

18.10B: Ascending Aorta

18.10C: Arch of the Aorta

18.10D: Thoracic Aorta

18.10E: Abdominal Aorta

18.10F: Arteries of the Pelvis and Lower Limbs

18.1: Blood Vessel Structure and Function

18.1A: Blood Vessel Structure

18.1B: Blood Vessel Function

18.2: Arteries

18.2A: Artery Function

18.2B: Elastic Arteries

18.2C: Muscular Arteries

18.2D: Anastomoses

18.2E: Arterioles

18.2F: Capillaries

18.3: The Venous System

18.3A: Venules

18.3B: Veins

18.4: Physiology of Circulation

18.4A: Introduction to Blood Flow, Pressure, and Resistance

18.4B: Distribution of Blood

18.5: Systemic Blood Pressure

18.5A: Introduction to Blood Pressure

18.5B: Arterial Blood Pressure

18.5C: Venous Blood Pressure

18.6: Control of Blood Pressure

18.6A: Role of the Cardiovascular Center

18.6B: Short-Term Neural Control

18.6C: Checking Circulation

18.6C: Short-Term Chemical Control

18.6D: Long-Term Renal Regulation

18.6E: Pulse
18.6F: Measuring Blood Pressure
18.6G: Extremes in Blood Pressure

18.7: Blood Flow Through the Body
18.7A: Velocity of Blood Flow
18.7B: Local Regulation of Blood Flow
18.7C: Blood Flow in Skeletal Muscle
18.7D: Blood Flow in the Brain
18.7E: Blood Flow in the Skin
18.7F: Blood Flow in the Lungs
18.7G: Blood Flow in the Heart
18.7H: Hepatic Portal Circulation

18.8: Capillary Exchange
18.8A: Capillary Dynamics
18.8B: Transcytosis
18.8C: Bulk Flow: Filtration and Reabsorption

18.9: Circulatory Shock
18.9A: Types of Shock
18.9B: Homeostatic Responses to Shock
18.9C: Signs and Symptoms of Shock

19: Lymphatic System

19.1: Lymphatic System Structure and Function
19.1A: Structure of the Lymphatic System
19.1B: Functions of the Lymphatic System

19.2: Lymphatic Vessels
19.2A: Lymphatic Vessel Structure
19.2B: Distribution of Lymphatic Vessels
19.2C: Lymph Transport
19.2D: Lymphatic Capillaries
19.2E: Lymph Trunks and Ducts

19.3: Lymph Cells and Tissues
19.3A: Lymphoid Cells
19.3B: Lymphoid Tissue
19.3C: Lymph Nodes

19.4: Other Lymphoid Organs
19.4A: Thymus
19.4B: Spleen
19.4C: Tonsils

20: Immune System

20.1: Innate Immunity

20.1A: Skin and Mucosae (Surface Barriers)

20.1B: Phagocytes

20.1C: Natural Killer Cells

20.1D: Inflammation

20.1E: Antimicrobial Proteins

20.1F: Fever

20.2: Stress and Immunity

20.2A: Stress and Immunity

20.3: Adaptive Immunity

20.3A: Overview of Adaptive Immunity

20.3B: Types of Adaptive Immunity

20.3C: Maturation of T Cells

20.3D: Lymphocytes

20.3E: Antigen-Presenting Cells

20.4: Cytokines

20.4A: Types of Cytokines Participating in Immune Response

20.5: Antigens

20.5A: Antigens and Antigen Receptors

20.5B: Complete Antigens and Haptens

20.5C: Antigenic Determinants and Processing Pathways

20.6: Humoral Immune Response

20.6A: Clonal Selection and B-Cell Differentiation

20.6B: Structure and Function of Antibodies

20.6C: Role of the Complement System in Immunity

20.6D: Immunological Memory

20.6E: Major Histocompatibility Complex Antigens (Self-Antigens)

20.7: Cell-Mediated Immune Response

20.7A: Clonal Selection and T-Cell Differentiation

20.7B: Specific T-Cell Roles

20.7C: Active and Passive Humoral Immunity

21: Respiratory System

21.10: Respiration Control

21.10A: Neural Mechanisms (Respiratory Center)

21.10B: Neural Mechanisms (Cortex)

21.10C: Chemoreceptor Regulation of Breathing

21.10D: Proprioceptor Regulation of Breathing

21.11: Respiratory Adjustments
21.11A: Adjustments During Exercise
21.11B: Adjustments at High Altitude

21.1: Overview of the Respiratory System
21.1A: The Reason for Breathing
21.1B: Functional Anatomy of the Respiratory System

21.2: Conducting Zone
21.2A: Nose and Paranasal Sinuses
21.2B: Pharynx
21.2C: Larynx
21.2D: Structures Used in Voice Production
21.2E: Trachea

21.3: Respiratory Zone
21.3A: Bronchi and Subdivisions
21.3B: Alveoli

21.4: The Lungs and Pleurae
21.4A: Lungs
21.4B: Lobes, Fissures, and Lobules
21.4C: Blood Supply to the Lungs

21.5: Mechanics of Breathing
21.5A: Pressure Changes During Pulmonary Ventilation
21.5B: Inspiration
21.5C: Expiration
21.5D: Breathing Patterns

21.6: Factors Affecting Pulmonary Ventilation
21.6A: Factors Affecting Pulmonary Ventilation: Surface Tension of Alveolar Fluid
21.6B: Factors Affecting Pulmonary Ventilation: Compliance of the Lungs
21.6C: Factors Affecting Pulmonary Ventilation: Airway Resistance

21.7: Nonrespiratory Lung Functions
21.7A: Lung Capacity and Volume
21.7B: Nonrespiratory Air Movements

21.8: Gas Laws
21.8A: Dalton's Law of Partial Pressure
21.8B: Henry's Law

21.9: Gas Exchange
21.1A: External Respiration
21.9B: Internal Respiration
21.9C: Oxygen Transport
21.9D: Carbon Dioxide Transport

22: Digestive System

22.1: Overview of the Digestive System

22.1A: Anatomy of the Digestive System

22.1B: Processes and Functions of the Digestive System

22.1C: Organs of the Digestive System

22.2: Nervous System of the Digestive System

22.2A: Enteric Nervous System

22.2B: Gastrointestinal Reflex Pathways

22.3: The Peritoneum

22.3: The Peritoneum

22.4: The Alimentary Canal

22.4A: Mouth

22.4B: Pharynx

22.4C: Esophagus

22.5: Layers of the Alimentary Canal

22.5A: Mucosa

22.5B: Submucosa

22.5C: Muscularis

22.5D: Serosa

22.7: The Liver

22.7A: The Liver

22.7B: Anatomy of the Liver

22.7C: Histology of the Liver

22.7D: Blood Supply to the Liver

22.7E: Liver Function

22.7F: Bile

22.8: The Gallbladder

22.8A: The Gallbladder

22.8B: Anatomy of the Gallbladder

22.9: The Pancreas

22.9A: Pancreas

22.9B: Anatomy of the Pancreas

22.9C: Histology of the Pancreas

22.9D: Pancreatic Juice

22.10: The Small Intestine

22.10A: Anatomy of the Small Intestine

22.10B: Histology of the Small Intestine

22.10C: Digestive Processes of the Small Intestine

22.11: The Large Intestine

22.11A: Anatomy of the Large Intestine
22.11B: Histology of the Large Intestine
22.11C: Bacterial Flora
22.11D: Digestive Processes of the Large Intestine
22.11E: Absorption and Feces Formation in the Large Intestine
22.11F: Defecation Reflex

22.12: Chemical Digestion
22.12A: Mechanisms of Chemical Digestion
22.12B: Chemical Digestion of Carbohydrates, Proteins, Lipids, and Nucleic Acids

22.13: Absorption
22.13A: Absorption in the Small Intestine
22.13B: Absorption of Monosaccharides, Amino Acids, Dipeptides, Tripeptides, Lipids, Electrolytes, Vitamins, and Water

22.14: Phases of Digestion
22.14A: Cephalic Phase
22.14B: Gastric Phase

22.6: The Stomach
22.6A: Gross Anatomy of the Stomach
22.6B: Microscopic Anatomy of the Stomach
22.6C: Digestive Properties of the Stomach

Index

23: Nutrition and Metabolism

23.10: Diet, Digestion, and Energy Storage Regulation
23.10A: Hormonal Responses to Food
23.10B: Neural Responses to Food
23.10C: Food Energy and ATP

23.1: Nutrition
23.1A: Digestive Systems
23.1B: Carbohydrates: Sources, Uses in the Body, and Dietary Requirements
23.1C: Lipids: Sources, Uses in the Body, and Dietary Requirements
23.1D: Proteins: Sources, Uses in the Body, and Dietary Requirements
23.1E: Food Requirements and Essential Nutrients

23.2: Carbohydrates
23.2A: Carbohydrate Molecules
23.2B: Importance of Carbohydrates

23.3: Lipids
23.3A: Lipid Molecules
23.3B: Phospholipids
23.3C: Steroids

23.4: Proteins
23.4A: Types and Functions of Proteins
23.4B: Amino Acids
23.4C: Protein Structure

23.5: Metabolic Reactions
23.5A: Breakdown of Pyruvate
23.5B: Citric Acid Cycle
23.5C: Importance of Glycolysis
23.5D: Electron Transport Chain
23.5E: ATP Yield
23.5F: Control of Catabolic Pathways
23.5G: Transforming Chemical Energy

23.6: Metabolism of Nutrients
23.6A: Connecting Other Sugars to Glucose Metabolism
23.6B: Connecting Proteins to Glucose Metabolism

23.7: Metabolic Body States
23.7A: Catabolic-Anabolic Steady State
23.7B: Absorptive State
23.7C: Postabsorptive State

23.8: Metabolism and the Liver
23.8A: Cholesterol Metabolism
23.8B: Regulation of Blood Cholesterol Levels

23.9: Metabolic Rate and Thermoregulation
23.9A: Homeostatic Process
23.9B: Homeostasis: Thermoregulation
23.9C: Heat Conservation and Dissipation

24: Urinary System

24.1: Overview of the Urinary System
24.1A: Overview of the Urinary System

24.2: The Kidneys
24.2A: Location and External Anatomy of the Kidneys
24.2B: Internal Anatomy of the Kidneys
24.2C: Supply of Blood and Nerves to the Kidneys
24.2D: Nephron, Parts, and Histology

24.3: Physiology of the Kidneys
24.3A: Overview of Urine Formation
24.3B: Glomerular Filtration
24.3C: Regulation of Glomerular Filtration Rate

24.3D: Tubular Reabsorption
24.3E: Tubular Secretion

24.4: Urine
24.4A: Physical Characteristics of Urine
24.4B: Chemical Composition of Urine
24.4C: Regulation of Urine Concentration and Volume
24.4D: Urinalysis
24.4E: Renal Clearance

24.5: Urine Transport, Storage, and Elimination
24.5A: Overview of Urine Transport, Storage, and Elimination
24.5B: Ureters
24.5C: Urinary Bladder
24.5D: Urethra
24.5E: Micturition and the Micturition Reflex
24.5F: Waste Management in Other Body Systems

25: Body Fluids and Acid-Base Balance

25.1: Water Balance
25.1A: Regulation of Water Intake
25.1B: Regulation of Water Output
25.1C: Nitrogenous Waste in Terrestrial Animals: The Urea Cycle
25.1D: Water Balance Disorders

25.2: Body Fluids
25.2A: Water Content in the Body
25.2B: Fluid Compartments
25.2C: Body Fluid Composition
25.2D: Movement of Fluid Among Compartments

25.3: Electrolyte Balance
25.3A: Sodium, Electrolytes, and Fluid Balance
25.3B: Sodium Balance Regulation
25.3C: Potassium Balance Regulation
25.3D: Calcium and Phosphate Balance Regulation
25.3E: Anion Regulation

25.4: Acid-Base Balance
25.4A: pH, Buffers, Acids, and Bases
25.4B: Chemical Buffer Systems
25.4C: Regulation of H+ by the Lungs
25.4D: The Role of the Kidneys in Acid-Base Balance

26: The Reproductive System

26.1: Overview of the Reproductive System

26.1A: Overview of the Male and Female Reproductive Systems

26.2: Meiosis

26.2A: Introduction to Meiosis

26.2B: Comparing Meiosis and Mitosis

26.3: The Male Reproductive System

26.3A: Anatomy of the Male Reproductive System

26.3B: Scrotum

26.3C: Testes

26.3D: Penis

26.3E: Testes Ducts

26.3F: Accessory Sex Glands

26.4: Physiology of the Male Reproductive System

26.4A: Male Sexual Response

26.4B: Sperm

26.4C: Spermatogenesis

26.4D: Semen

26.4E: Hormonal Regulation of the Male Reproductive System

26.5: The Female Reproductive System

26.5A: Anatomy of the Female Reproductive System

26.5B: Ovaries

26.5C: Uterus

26.5D: Female Duct System

26.5E: Vagina

26.5F: Vulva

26.5G: Perineum

26.5H: Mammary Glands

26.6: Physiology of the Female Reproductive System

26.6A: Oogenesis

26.6B: Ovarian Cycle

26.6C: Uterine (Menstrual) Cycle

26.6D: Hormonal Regulation of the Female Reproductive Cycle

26.6E: Extrauterine Effects of Estrogens and Progesterone

26.6F: Female Sexual Response

27: Human Development and Pregnancy

27.10: Changes to the Mother's Body During Pregnancy

27.10A: Anatomical Changes

27.10B: Metabolic Changes

27.10C: Physiological Changes
27.10D: Exercise and Pregnancy

27.1: Overview of Pregnancy and Human Development
27.1A: Introduction to Pregnancy and Human Development

27.2: First Week of Development
27.2A: Fertilization
27.2B: Cleavage of the Zygote
27.2C: Blastocyst Formation
27.2D: Implantation

27.3: Second Week of Development
27.3A: Trophoblast Development
27.3B: Bilaminar Embryonic Disc Development
27.3C: Amnion Development
27.3D: Yolk Sac Development
27.3E: Sinusoid Development
27.3F: Development of the Extraembryonic Coelom

27.4: Third Week of Development
27.4A: Gastrulation
27.4B: Neurulation
27.4C: Somite Development
27.4D: Development of the Cardiovascular System
27.4E: Chorionic Villi and Placental Development

27.5: Fourth Week of Development
27.5A: Fourth Week of Development

27.6: Fifth Through Eighth Weeks of Development
27.6A: Fifth Through Eighth Weeks of Development

27.7: Birth (Parturition)
27.7A: Onset of Labor
27.7B: Stages of Labor
27.7C: Adjustments of the Infant at Birth

27.8: Lactation
27.8A: Physiology of Lactation

27.9: The Fetal Period
27.9A: Fetal Development
27.9B: Teratogens
27.9C: Prenatal Diagnostic Tests

28: Reproduction, Chromosomes, and Meiosis
28.10: Exceptions to Mendelian Inheritance

28.10A: Mendel's Laws of Heredity

28.1: The Link Between Genotype and Phenotype

28.1A: The Process and Purpose of Gene Expression Regulation

28.1B: The Relationship Between Genes and Proteins

28.1C: Chromosomal Theory of Inheritance

28.2: Inherited Variation

28.2A: Advantages and Disadvantages of Sexual Reproduction

28.3: Chromosomal Behavior in Meiosis and Fertilization

28.3A: The Structure and Sequence of DNA

28.4: DNA

28.4A: DNA and RNA

28.4B: The DNA Double Helix

28.4C: DNA Packaging

28.4D: Types of RNA

28.5: Cell Division

28.5A: The Role of the Cell Cycle

28.5B: Genomic DNA and Chromosomes

28.5C: Eukaryotic Chromosomal Structure and Compaction

28.6: Sources of Genetic Variation

28.6A: Genetic Variation

28.6B: Mendel's Law of Segregation

28.6C: Mendel's Law of Independent Assortment

28.7: Complex Inheritance Patterns

28.3C: Mendel's Model System

28.7A: The Punnett Square Approach for a Monohybrid Cross

28.7B: Phenotypes and Genotypes

28.8: Patterns of Inheritance

28.8A: Lethal Inheritance Patterns

28.8B: Sex-Linked Traits

28.9: Genetic Disequilibrium

28.9A: Genetic Linkage and Distances

28.9B: Chromosomal Structural Rearrangements

30: APPENDIX B: Development and Aging of the Organ Systems

30.10: Development of the Heart

30.11: Development of Blood Vessels

30.12: Development of Lymphatic Tissues

30.13: Development of the Immune System

30.14: Development of the Respiratory System

30.15: Development of the Digestive System
30.16: Development of Metabolism
30.17: Development of Bodily Fluids
30.18: Development of the Reproductive System
30.1: Integumentary System Development
30.1A: Development of the Integumentary System
30.1B: Effects of Aging on the Integumentary System
30.2: Bone Development
30.2A: Stages of Bone Development
30.2B: Exercise and Bone Tissue
30.2C: Bone Tissue and the Effects of Aging
30.3: Development of the Skeleton
30.3A: Development of the Skeleton
30.3B: Aging and the Skeleton
30.4: Joint Development
30.4A: Development of Joints
30.4B: Joints and Aging
30.4C: Arthroplasty
30.5: Muscle Development
30.5A: Muscle Development
30.6: Development of the Nervous System
30.6A: Development of the Central Nervous System
30.6B: Development of the Peripheral Nervous System
30.6C: Regeneration of Nerve Fibers
30.6D: Aging and the Nervous System
30.7: Development of the Special Senses
30.47C: Development of Hearing and Balance
30.7A: Taste and Smell at Birth and in Old Age
30.7B: Development of Vision
30.8: Development of the Endocrine System
30.9: Development of Blood
29: APPENDIX A: Diseases, Injuries, and Disorders of the Organ Systems
29.10: Circulatory Diseases and Disorders
29.10A: Angiogenesis and Disease
29.10B: Varicose Veins
29.10C: Syncope
29.10D: Hypertension
29.11: Diseases and Disorders of the Immune System

29.11A: Immunodeficiencies
29.11B: Autoimmune Diseases
29.11C: Infectious Mononucleosis
29.11D: Psoriasis

29.12: Respiratory Diseases and Disorders
29.12A: Asthma
29.12B: Respiratory Distress Syndrome
29.12C: Cystic Fibrosis
29.12D: Hypoxia

29.13: Diseases and Disorders of the Digestive System
29.13A: Peritonitis
29.13B: Mumps
29.13C: Lactose Intolerance
29.13D: Peptic Ulcer Disease

29.14: Diseases and Disorders of the Metabolism
29.14A: Phenylketonuria (PKU)
29.14B: Galactosemia and Glycogen Storage Disease

29.15: Diseases and Disorders of the Urinary System
29.15A: Renal Disease and Failure
29.15B: Nephroptosis (Floating Kidney)
29.15C: Polycystic Kidney Disease

29.16: Body Fluid-Related Diseases and Disorders
29.16A: Acidosis
29.16B: Alkalosis
29.16C: Edema

29.17: Diseases and Disorders of the Reproductive System
29.17A: Sexually Transmitted Diseases (STDs)
29.17B: Erectile Dysfunction and the Blue Pill
29.17C: Cervical Cancer

29.18: Pregnancy-Related Disorders
29.18A: Ectopic Pregnancy
29.18B: Infertility

29.1: Diseases, Injuries, and Disorders of the Integumentary System
29.1A: Burns
29.1B: Allergies
29.1C: Acne

29.2: Diseases and Disorders of the Skeletal System
29.2A: Osteoporosis

29.2B: Paget's Disease

29.3: Injuries and Disorders of the Skeleton

29.3A: Spina Bifida

29.3B: Fractures

29.4: Diseases, Disorders, and Injuries of the Joints

29.4A: Sprain and Strain

29.4B: Rheumatism and Arthritis

29.4C: Common Joint Injuries

29.5: Diseases, Disorders, and Injuries of the Muscular System

29.5A: Muscular Dystrophy

29.5B: Exercise-Induced Muscle Damage

29.5C: Carpal Tunnel Syndrome

29.6: Injuries, Diseases, and Disorders of the Central Nervous System

29.6A: Traumatic Brain Injury

29.6B: Cerebrovascular Accident

29.6C: Transient Ischemic Attacks

29.6D: Alzheimer's Disease

29.6E: Brain Tumors

29.6F: Amyotrophic Lateral Sclerosis

29.7: Diseases and Disorders of the Endocrine System

29.7A: Pancreatic Islet Disorders: Diabetes and Hyperinsulinism

29.7B: Posttraumatic Stress Disorder (PTSD)

29.8: Blood Diseases and Disorders

29.8A: Hemostasis Disorders

29.8B: Anemia

29.8C: Sickle-Cell Disease

29.8D: Leukemia

29.9: Diseases and Disorders of the Heart

29.9A: Congenital Heart Defects

29.9B: Myocarditis and Endocarditis

29.9C: Myocardial Ischemia and Infarction

29.9D: Arrhythmia

Index

Glossary

Detailed Licensing

This page titled Anatomy and Physiology (Boundless) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning) via source content that was edited to the style and standards of the LibreTexts platform.

Licensing

A detailed breakdown of this resource's licensing can be found in ***Back Matter/Detailed Licensing***.

This page titled Licensing is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1: Introduction to Anatomy and Physiology

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/19/2026

10: Overview of the Nervous System

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/19/2026

11: Central Nervous System

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/19/2026

22: Digestive System

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/19/2026

1: INTRODUCTION TO ANATOMY AND PHYSIOLOGY

1.1: Overview of Anatomy and Physiology

1.1A: Defining Anatomy

1.1B: Defining Physiology

1.2: Life

1.2A: Life Functions

1.2B: Survival Needs

1.2C: Levels of Organization

1.3: Homeostasis

1.3A: Homeostatic Control

1.3B: Disease as Homeostatic Imbalance

1.4: Mapping the Body

1.4A: Anatomical Position

1.4B: Directional Terms

1.4C: Regional Terms and Axes

1.4D: Body Planes and Sections

1.4E: Body Cavities

1.4F: Abdominopelvic Regions

Index

This page titled 1: Introduction to Anatomy and Physiology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.1: OVERVIEW OF ANATOMY AND PHYSIOLOGY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

1.1A: Defining Anatomy

1.1B: Defining Physiology

This page titled 1.1: Overview of Anatomy and Physiology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.1A: Defining Anatomy

Human anatomy is the study of the structure of the human body.

Learning Objectives

- Define the two major subtypes of anatomy

Key Points

- Gross anatomy is subdivided into surface anatomy (the external body), regional anatomy (specific regions of the body), and systemic anatomy (specific organ systems).
- Microscopic anatomy is subdivided into cytology (the study of cells) and histology (the study of tissues).
- Anatomy is closely related to physiology (study of function), biochemistry (chemical processes of living things), comparative anatomy (similarities and differences between species), and embryology (development of embryos).
- Knowledge of anatomy is needed to understand human biology and medicine.

Key Terms

- **anatomy**: The study of the body structure of animals.
- **surface Anatomy**: The study of anatomical landmarks that can be identified by observing the surface of the body. Sometimes called superficial anatomy.
- **microscopic anatomy**: The study of minute anatomical structures on a microscopic scale, including cells (cytology) and tissues (histology).
- **Gross (or macroscopic) anatomy**: The study of anatomical features visible to the naked eye, such as internal organs and external features.
- **embryology**: The science of the development of an embryo from the fertilization of the ovum to the fetal stage.
- **dissection**: The process of disassembling an organism to determine its internal structure and understand the functions and relationships of its components.

Defining Anatomy

Anatomy describes the structure and location of the different components of an organism to provide a framework for understanding. Human anatomy studies the way that every part of a human, from molecules to bones, interacts to form a functional whole.



Rembrant's "The Anatomy Lesson of Dr. Nicolaes Tulp": Human anatomy is the study of the structure of the human body, from the microscopic to the macroscopic.

There are two major types of anatomy. Gross (macroscopic) anatomy is the study of anatomical structures that can be seen by the naked eye, such as the external and internal bodily organs. Microscopic anatomy is the study of tiny anatomical structures such as tissues and cells.

Gross Anatomy

Gross anatomy can be further subdivided into three different fields:

- Surface anatomy (or superficial anatomy) is the study of external anatomical features without dissection.
- Regional anatomy focuses on specific external and internal regions of the body (such as the head or chest) and how different systems work together in that region.
- Systemic anatomy focuses on the anatomy of different organ systems, such as the respiratory or nervous system.

Regional anatomy is widely used in modern teaching because it is easier to apply to a clinical setting than systemic anatomy. The major anatomy textbook, Gray's Anatomy, has recently been reorganized from a systems format to a regional format to reflect this preference. Surface anatomy is also widely used to gauge the position and structure of deeper organs, tissues, and systems.

Microscopic Anatomy

Within microscopic anatomy, two topics of study are of great importance:

- Cytology, the study of the structure and function of cells
- Histology, the study of the organization and details of biological tissues

The human body has many layers of organization. Biological systems consist of organs that consist of tissues, and tissue in turn is made up of cells and connective tissue. Microscopic anatomy allows one to focus on these tissues and cells.

History of Anatomy

The history of anatomy has been an evolving understanding of organs and structures in the body. Beginning in Ancient Greece and developing through the Middle Ages and the Renaissance, methods of studying anatomy have advanced dramatically. This field has moved from examination of animals and cadavers through invasive dissection to the technologically complex techniques developed in the 20th century, such as non-invasive imaging and radiology.

Generally, medical and biology students learn about the human body from anatomical models, skeletons, textbooks, diagrams, photographs, lectures, and tutorials. Medical and dental students also learn through the dissection and inspection of cadavers. A thorough working knowledge of anatomy is required for all medical professionals, especially surgeons and doctors working in diagnostic specialties such as radiology.

Anatomy and Physiology

Physiology is the study of how the components of the body function, and biochemistry is the study of the chemistry of living structures. Together with anatomy, these are the three primary disciplines within the field of human biology. Anatomy provides information about structure, location, and organization of different parts of the body that is needed to truly understand physiology. Together, anatomy and physiology explain the structure and function of the different components of the human body to describe what it is and how it works.

This page titled 1.1A: Defining Anatomy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.1B: Defining Physiology

Human physiology is the study of the mechanical, physical, and biochemical processes that support the body's function.

Learning Objectives

- Describe physiology in the context of living systems

Key Points

- The study of physiology occurs at many levels, including organisms, organ systems, organs, cells, and biomolecules.
- Physiology investigates the processes and mechanisms that allow a living thing to survive, grow, and develop.
- Homeostasis is the maintenance of the overall stability of physiological processes of the body. It is monitored by the brain and nervous system and regulated by the physiology and activity of individual organs.
- While anatomy deals with the structure of the parts of an organism, physiology studies the way that the parts function together.

Key Terms


- **homeostasis**: The ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium, such as the ability of warm-blooded animals to maintain a constant internal temperature despite changes in outdoor temperature.
- **Living Systems**: Open, self-organizing living things that interact with their environments. These systems are maintained by flows of information, energy, and matter.
- **physiology**: A branch of biology that deals with the functions and activities of living matter and the physical and chemical processes and mechanisms involved.

EXAMPLES

The body's ability to digest and absorb nutrients from food.

Physiology is the science of the normal function of living systems. Physiology studies the processes and mechanisms that allow an organism to survive, grow, and develop. Physiological processes are the ways in which organ systems, organs, tissues, cells, and biomolecules work together to accomplish the complex goal of sustaining life.

Physiological mechanisms are the smaller physical and chemical events that make up a larger physiological process. Human physiology studies the functions of humans, their organs and cells, and how all of these functions combine to make life, growth, and development possible.



Leonardo da Vinci's "The Vitruvian Man": Vitruvian Man: A drawing created by Leonardo da Vinci. The drawing is based on the correlations of ideal human proportions with geometry described[4] by the ancient Roman architect Vitruvius in Book III of his treatise De Architectura.

Homeostasis

Homeostatis is the maintenance of the body's overall inner resistance to change. This resistance stabilizes the body by regulating the internal environment, even as the external environment changes. A stable internal environment is needed for normal physiological function and survival of a living system.

Maintaining a stable internal environment requires constant monitoring, mostly by the brain and nervous system. The brain, more specifically the hypothalamus, receives information from the body and responds appropriately through the release of chemical messengers such as neurotransmitters, catecholamines, and hormones. These chemical messengers signal individual organs to change their functions in order to maintain homeostasis for the whole body.

For instance, if blood oxygen levels are too low, the brain signals the muscles controlling the lungs to breathe faster to increase oxygen intake. The brain also signals the heart to beat faster so other organs and tissues receive the oxygen they need. When oxygen levels return to normal, the brain signals the lungs and heart to return to their normal rates of function, a process called feedback.

Body Systems


Traditionally, the academic discipline of physiology views the body as a collection of interacting systems, each with its own combination of functions and purposes. Each system contributes to the homeostasis of other systems and of the entire organism. No system works in isolation, and the well-being of the person depends upon the well-being of the interactions between body systems. The traditional divisions by system are somewhat arbitrary. Many organs participate in more than one system (such as the heart and kidney), and systems might be organized by function, by embryological origin, or by other categorizations.

For instance, the neuroendocrine system is the complex interactions of the neurological and endocrinological systems. Together, the neuroendocrine system regulates many physiological processes, including those that maintain homeostasis. Furthermore, many aspects of physiology are not easily categorized by traditional definitions of organ systems because they are composed of interactions between organs in multiple organ systems.

The study of how physiology is altered in disease is pathophysiology. Pathophysiology focuses on how physiological processes fail to maintain normal function, resulting in the manifestation of disease symptoms.

Anatomy and Physiology

Human anatomy deals with how all parts of the human body interact to form a functional whole. The study of anatomy is separate from the study of physiology, although the two are often taught together. While anatomy studies the structure of the parts of an organism, physiology is concerned with the way those parts function together. For example, an anatomist may study the types of tissues found in different parts of the heart, while a physiologist may study how the heart regulates blood flow to supply oxygen to other organs in the body. While anatomy and physiology study different aspects of human biology, together they provide a more complete picture of what the human body is and how it works.



The human heart and lungs: Anatomy and physiology are complimentary disciplines as the structure of body systems often influences system function.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human anatomy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Anatomy/What is anatomy?. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_...is_anatomy%3F**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Anatomy/Terminology and Organization. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_A...d_Organization**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anatomy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- surface Anatomy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/surface%20Anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dissection. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/dissection**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- embryology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/embryology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- microscopic anatomy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/microscopic%20anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- The Anatomy Lesson. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Th...omy_Lesson.jpg**. **License**:

Public Domain: No Known Copyright
- Human Anatomy/What is anatomy?. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_..._is_anatomy%3F**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Living Systems. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Living%20Systems**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homeostasis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homeostasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- physiology. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- The Anatomy Lesson. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Th...omy_Lesson.jpg**. **License**: ***Public Domain: No Known Copyright***
- Heart-and-lungs. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Heart-and-lungs.jpg**. **License**: ***Public Domain: No Known Copyright***
- Da Vinci Vitruve Luc Viatour. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Da...uc_Viatour.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 1.1B: Defining Physiology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.2: LIFE



- Topic hierarchy

1.2A: Life Functions

1.2B: Survival Needs

1.2C: Levels of Organization

This page titled 1.2: Life is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.2A: Life Functions

All life forms require certain core elements as well as physical and chemical factors from the ecosystem for biochemical functioning.

Learning Objectives

- Categorize living organisms and non-living things

Key Points

- Living organisms are systems made from cells and are capable of life functions such as reproduction, metabolism, response to stimuli, and homeostasis.
- Nonliving things are either living things that have died or inanimate objects that were never alive.
- The nonliving (abiotic) components of an ecosystem include energy, oxygen, water, nutrients, and temperature.
- Living organisms have ranges of ecosystem conditions where they can perform all physiological functions of life. Organisms cannot perform these functions as well or at all outside of these conditions, but some organisms have developed ways to adapt to environmental changes. Migration and hibernation are two examples of these adaptations.
- Organisms called extremophiles can assume forms enabling them to withstand freezing, dehydration, starvation, high levels of radiation, and other physical or chemical challenges. Extremophiles can survive exposure to such conditions for weeks, months, years, or even centuries.
- All life forms require certain core chemical elements for biochemical structure and function. These include carbon, hydrogen, nitrogen, oxygen, phosphorus, and sulfur—the elemental macronutrients for all organisms—often represented by the acronym CHNOPS.

Key Terms

- **extremophile**: A microorganism that can survive in extreme environmental conditions.
- **organism**: A contiguous living system that can metabolize, grow, respond to stimuli, reproduce, and adapt to its environment.
- **life**: A status given to any entity with the properties of reproduction and metabolism, including animals, plants, fungi, bacteria, and archaea (single-celled microorganisms).

Living Organisms

An organism is a living system capable of growth, reproduction,metabolism, response to stimuli, adaptation and homeostasis. Cells are the basic unit of life that can be found in every living organism. Organisms can be unicellular or multicellular and include plants, animals, bacteria, archaea, and fungi. Multicellular organisms are typically made up of organ systems, organs, tissues, and cells. More complex organisms are capable of communication.

The world is made up of both living (biotic) and nonliving things. Nonliving things are living things that have died or inanimate (abiotic) objects that were never alive to begin with. Living and nonliving things can be distinguished by function. Nonliving things are not capable of growth, reproduction, response to stimuli, adaptation, or homeostasis.

It is unknown whether viruses are actually alive. They are capable of reproduction (within a host) and adaptation, but they are not capable of homeostasis or metabolic functions. Because viruses exhibit some but not all characteristics of organisms, they cannot be fully considered living things.

Ecosystems Are Important for Life

The abiotic components of an ecosystem are the physical and chemical factors necessary for life—energy (sunlight or chemical energy), water, temperature, oxygen, and nutrients. In most ecosystems, the environmental conditions vary throughout the day or

over the course of seasons. Organisms who live in an ecosystem must be able to survive its range of conditions, called the “range of tolerance.” Outside the range of tolerance are the “zones of physiological stress,” where survival and reproduction are possible but not optimal. Beyond these zones are the “zones of intolerance,” where life for that organism is impossible. Many organisms can adapt to intolerance in their environment through seasonal migration, hibernation, or other adaptations.

Many microorganisms have evolved adaptations that allow them to survive in zones that are intolerant for most other organisms, such as extreme heat or cold, dehydration, starvation, or high levels of radiation. These microorganisms are called extremophiles, and they thrive outside the ranges where life is commonly found. They excel at exploiting uncontested sources of energy. While all organisms are composed of nearly identical types of molecules, evolution has enabled extremophile microbes to cope with this wide range of physical and chemical conditions.



Rainforest Trees: Plant growth in the Hoh Rainforest

Chemical Requirements for Life

All life forms require certain core chemical elements for biochemical functioning. These include carbon, hydrogen, nitrogen, oxygen, phosphorus, and sulfur—the elemental macronutrients for all organisms. Together these make up carbohydrates, nucleic acids, proteins, and lipids, the four categories of molecules that make up the structure and function of organic life. The most widespread and important of these elements is carbon, which holds molecules together through multiple, stable covalent bonds. Organic molecules generally contain carbon, and inorganic molecules generally do not contain carbon (with the key exception of carbon dioxide). Organic molecules and their properties form the basic structure of cells and allow physiological processes to occur.

This page titled 1.2A: Life Functions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.2B: Survival Needs

To sustain human life, certain physiological needs include air, water, food, shelter, sanitation, touch, sleep and personal space.

Learning Objectives

- Identify the eight minimal physiological requirements for survival

Key Points

- While the development of tools, plumbing systems and food preservation have enabled modern man to live life with many creature comforts, the truth remains that the requirements for human survival are quite basic.
- Physiological needs include air, water, food, shelter, sanitation, touch, sleep, and personal space.
- As humans have evolved to interact in community settings, both hunting and gathering in groups, touch—as in a caring caress—is often considered a basic human survival need. In fact, empirical evidence has shown touch to be essential to the early growth and development of healthy humans.

Key Terms

- **survival**: The fact or act of surviving; continued existence or life.

From the times of our first primate ancestors, innovation has progressed at a rapid clip. While the development of tools, plumbing systems and food preservation have enabled modern man to live life with many creature comforts, the truth remains that the requirements for human survival are quite basic. There are eight minimal physiological requirements for survival. These are: air, water, food, shelter, sanitation, sleep, space, and touch.

Survival Needs

1. Air: Consisting of oxygen within a particular range of pressure, concentration, and purity is vital to survival.
2. Water: Access to a safe, clean and adequate water supply is necessary for human survival.
3. Food: Our food must be adequate in calorie and nutritional standards in order to sustain life.
4. Shelter: Since exposure to cold and heat can lead to hypothermia or hyperthermia, shelter which offers protection from extremes of heat, cold, intense sun, and prolonged precipitation is a human survival need.
5. Sanitation: Proper means for the removal of human waste helps protect from deadly toxins and pathogens and is critical in promoting human survival.
6. Sleep: Seven to eight hours of uninterrupted sleep each night is optimal for human survival.
7. Space: As humans, we require personal space. In addition to the requirement for shelter, or suitable indoor living space, humans need outdoor space, to avoid overcrowding and chaos.
8. Touch: As humans have evolved to interact in community settings, both hunting and gathering in groups, touch—as in a caress—is often considered a basic human survival need. In fact, empirical evidence has shown that touch is essential for the early growth and development of healthy humans.



A Child Sleeping: Adequate sleep is necessary to sustain life.

This page titled 1.2B: Survival Needs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.2C: Levels of Organization

Living organisms are made up of four levels of organization: cells, tissues, organs, and organ systems.

Learning Objectives

- Order the levels of organization for living organisms

Key Points

- Cells are the most basic unit of life at the smallest level of organization.
- Cells can be prokaryotic (without nucleus) or eukaroyotic (with nucleus).
- The four categories of tissues are connective, muscles, epithelial, and nervous tissues.
- Organs are made of different types of tissues and perform complex functions. They can be hollow or solid.
- Organ systems are groups of organs that perform similar functions or perform functions together.
- Many physiological functions are carried out by multiple organ systems working in tandem.

Key Terms

- **cell**: The smallest unit of life capable of independent reproduction. Generally contains nucleic acid, cytoplasm, a cell membrane, and many other proteins and structures.
- **organ**: A structure made of different tissues that work together to perform physiological functions.
- **organ system**: A group of organs and tissues that work together to perform specific functions.
- **Tissues**: A group of similar cells with the same origin that work together to perform the same function.

EXAMPLES

Using the circulatory system as an example, a cell in this system is a red blood cell, the heart's cardiac muscle is a tissue, an organ is the heart itself, and the organ system is the circulatory system.

An organism is made up of four levels of organization: cells, tissues, organs, and organ systems. These levels reduce complex anatomical structures into groups; this organization makes the components easier to understand.

Level 1: Cells

The first and most basic level of organization is the cellular level. A cell is the basic unit of life and the smallest unit capable of reproduction. While cells vary greatly in their structure and function based on the type of organism, all cells have a few things in common. Cells are made up of organic molecules, contain nucleic acids (such as DNA and RNA), are filled with fluid called cytoplasm, and have a membrane made out of lipids. Cells also contain many structures within the cytoplasm called organelles, which perform various cellular functions.

Cells may be prokaryotic (without a nucleus) in bacteria and archaea (single-celled organisms), or eukaryotic (with nucleus-enclosing DNA) in plants, animals, protists, and fungi. In humans, most cells combine to form tissues, but some cells are found independent of solid tissues and have their own functions. A red blood cell found circulating in the bloodstream carrying oxygen throughout the human body is an example of an independent cell.

Level 2: Tissues

Tissues are a group of similar cells of the same origin that carry out a specific function together. Humans have four different types of basic tissues. Connective tissues such as bone tissue are made up of fibrous cells and give shape and structure to organs.


Muscle tissue is made up of cells that can contract together and allow animals to move. Epithelial tissues make up the outer layers of organs, such as the skin or the outer layer of the stomach. Nervous tissue is made of specialized cells that transmit information through electrochemical impulses, such as the tissue of nerves, the spinal cord, and the brain.

Level 3: Organs

An organ is a structure made up of different tissues that perform specific bodily functions. Most organs contain tissues such as parenchyma (used to perform the organ functions), stroma (connective tissue specific to organs) and epithelial. Organs may be solid or hollow, and vary considerably in size and complexity. The heart, lungs, and brain are all examples of organs.

Level 4: Organ Systems

An organ system is a collection of organs that that work together to perform a similar function. There are eleven different organ systems in the human body, each with its own specific functions. One example is digestive system, which is made up of many organs that work together to digest and absorb nutrients from food. While most organ systems control a few specific physiological processes, some processes are more complex and require multiple organ systems to work together. For example, blood pressure is controlled by a combination of the renal system (kidneys), the circulatory system, and the nervous system.



Levels of Organization in Animals: An organism contains organ systems made up of organs that consist of tissues, which are in turn made up of cells.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- What Matters/Physiological Needs. **Provided by**: Wikiversity. **Located at**: **en.wikiversity.org/wiki/What_...ological_Needs**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- life. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/life**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ition/organism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extremophile. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/extremophile**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Hoh rain forest trees. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ho...rest_trees.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Harry Harlow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Harry_H...infant_monkeys**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- What Matters/Physiological Needs. **Provided by**: Wikiversity. **Located at**: **en.wikiversity.org/wiki/What_...ological_Needs**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- survival. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/survival**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hoh rain forest trees. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ho...rest_trees.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- What Matters/Physiological Needs. **Provided by**: Wikiversity. **Located at**: **en.wikiversity.org/wiki/What_Matters/Physiological_Needs**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- levels of organization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Biological_organisation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Organ. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Organ_(anatomy)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tissues. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tissue_(biology)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cell. **Provided by**: Wikitionary. **Located at**: **en.wiktionary.org/wiki/cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hoh rain forest trees. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Hoh_rain_forest_trees.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- What Matters/Physiological Needs. **Provided by**: Wikiversity. **Located at**: **en.wikiversity.org/wiki/What_Matters/Physiological_Needs**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Levels of Organization in Animals. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...vels_mouse.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 1.2C: Levels of Organization is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.3: HOMEOSTASIS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

1.3A: Homeostatic Control

1.3B: Disease as Homeostatic Imbalance

This page titled 1.3: Homeostasis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.3A: Homeostatic Control

Homeostasis is maintained by the body's responses to adverse stimuli, ensuring maintenance of an optimal physiological environment.

Learning Objectives

- Model the feedback process of homeostasis

Key Points

- Homeostatic control mechanisms have at least three interdependent components: a receptor, integrating center, and effector.
- The receptor senses environmental stimuli, sending the information to the integrating center.
- The integrating center, generally a region of the brain called the hypothalamus, signals an effector (e.g. muscles or an organ) to respond to the stimuli.
- Positive feedback enhances or accelerates output created by an activated stimulus. Platelet aggregation and accumulation in response to injury is an example of positive feedback.
- Negative feedback brings a system back to its level of normal functioning. Adjustments of blood pressure, metabolism, and body temperature are all negative feedback.

Key Terms

- **homeostasis**: The ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium, such as the ability of warm-blooded animals to maintain a constant body temperature.
- **negative feedback**: A feedback loop in which the output of a system reduces the activity that causes that output.
- **positive feedback**: A feedback loop in which the output of a system is increased by the mechanism's own influence on the system that creates that output.

EXAMPLES

When an individual doesn't have enough to eat, the body adjusts by slowing down metabolism so that he or she expends fewer calories. This adaptation conserves the limited energy available from the inadequate diet.

Concept of Homeostasis

Homeostasis regulates an organism 's internal environment and maintains a stable, constant condition of properties like temperature and pH. Homeostasis can be influenced by either internal or external conditions and is maintained by many different mechanisms. All homeostatic control mechanisms have at least three interdependent components for the variable being regulated:

- A sensor or receptor detects changes in the internal or external environment. An example is peripheral chemoreceptors, which detect changes in blood pH.
- The integrating center or control center receives information from the sensors and initiates the response to maintain homeostasis. The most important example is the hypothalamus, a region of the brain that controls everything from body temperature to heart rate, blood pressure, satiety (fullness), and circadian rhythms (sleep and wake cycles).
- An effector is any organ or tissue that receives information from the integrating center and acts to bring about the changes needed to maintain homeostasis. One example is the kidney, which retains water if blood pressure is too low.

The sensors, integrating center, and effectors are the basic components of every homeostatic response. Positive and negative feedback are more complicated mechanisms that enable these three basic components to maintain homeostasis for more


complex physiological processes.

Positive Feedback

Positive feedback is a mechanism in which an output is enhanced in order to maintain homeostasis. Positive feedback mechanisms are designed to accelerate or enhance the output created by a stimulus that has already been activated. Positive feedback mechanisms are designed to push levels out of normal ranges. To achieve this, a series of events initiates a cascading process that builds to increase the effect of the stimulus. This process can be beneficial but is rarely used because it may become uncontrollable. A positive feedback example is blood platelet accumulation and aggregation, which in turn causes blood clotting in response to an injury of the blood vessels.

Negative Feedback


Negative feedback mechanisms reduce output or activity to return an organ or system to its normal range of functioning. Regulation of blood pressure is an example of negative feedback. Blood vessels have sensors called baroreceptors that detect if blood pressure is too high or too low and send a signal to the hypothalamus. The hypothalamus then sends a message to the heart, blood vessels, and kidneys, which act as effectors in blood pressure regulation. If blood pressure is too high, the heart rate decreases as the blood vessels increase in diameter (vasodilation), while the kidneys retain less water. These changes would cause the blood pressure to return to its normal range. The process reverses when blood pressure decreases, causing blood vessels to constrict and the kidney to increase water retention.



Negative Feedback Loop: The hypothalamus secretes corticotropin-releasing hormone (CRH), which directs the anterior pituitary gland to secrete adrenocorticotropic hormone (ACTH). In turn, ACTH directs the adrenal cortex to secrete glucocorticoids, such as cortisol. Glucocorticoids not only perform their respective functions throughout the body but also prevent further stimulating secretions of both the hypothalamus and the pituitary gland

Temperature control is another negative feedback mechanism. Nerve cells relay information about body temperature to the hypothalamus. The hypothalamus then signals several effectors to return the body temperature to 37 degrees Celsius (the set point). The effectors may signal the sweat glands to cool the skin and stimulate vasodilation so the body can give off more heat.

If body temperature is below the set point, muscles shiver to generate heat and the constriction of the blood vessels helps the body retain heat. This example is very complex because the hypothalamus can change the body's temperature set point, such as raising it during a fever to help fight an infection. Both internal and external events can induce negative feedback mechanisms.



Homeostatic Control: This image illustrates the feedback mechanisms of homeostatic controls.

This page titled 1.3A: Homeostatic Control is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.3B: Disease as Homeostatic Imbalance

If positive and negative feedback loops are affected or altered, homeostatic imbalance and resultant complications can occur.

Learning Objectives

- Analyze disease as a result of homeostatic imbalance

Key Points

- Many diseases are a result of homeostatic imbalance, an inability of the body to restore a functional, stable internal environment.
- Aging is a source of homeostatic imbalance as the control mechanisms of the feedback loops lose their efficiency, which can cause heart failure.
- Diseases that result from a homeostatic imbalance include heart failure and diabetes, but many more examples exist.
- Diabetes occurs when the control mechanism for insulin becomes imbalanced, either because there is a deficiency of insulin or because cells have become resistant to insulin.
- Homeostasis is the ability of a system to regulate its internal environment through maintaining a stable, relatively constant set of properties such as temperature and pH.

Key Terms

- **homeostasis**: The ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium, such as the ability of warm-blooded animals to maintain a constant body temperature.
- **diabetes**: A group of metabolic diseases in which a person or animal has high blood sugar due to an inability to produce, metabolize, or respond to insulin.
- **blood sugar regulation**: Carbohydrate and fat metabolism are regulated by insulin, a hormone produced by the pancreas.

What Is Disease?

Disease is any failure of normal physiological function that leads to negative symptoms. While disease is often a result of infection or injury, most diseases involve the disruption of normal homeostasis. Anything that prevents positive or negative feedback from working correctly could lead to disease if the mechanisms of disruption become strong enough.

Aging is a general example of disease as a result of homeostatic imbalance. As an organism ages, weakening of feedback loops gradually results in an unstable internal environment. This lack of homeostasis increases the risk for illness and is responsible for the physical changes associated with aging. Heart failure is the result of negative feedback mechanisms that become overwhelmed, allowing destructive positive feedback mechanisms to compensate for the failed feedback mechanisms. This leads to high blood pressure and enlargement of the heart, which eventually becomes too stiff to pump blood effectively, resulting in heart failure. Severe heart failure can be fatal.

Diabetes: A Disease of Failed Homeostasis


Diabetes, a metabolic disorder caused by excess blood glucose levels, is a key example of disease caused by failed homeostasis. In ideal circumstances, homeostatic control mechanisms should prevent this imbalance from occurring. However, in some people, the mechanisms do not work efficiently enough or the amount of blood glucose is too great to be effectively managed. In these cases, medical intervention is necessary to restore homeostasis and prevent permanent organ damage.

Blood Sugar Regulation

The human body maintains constant levels of glucose throughout the day, even after fasting. During long periods of fasting, glucose levels are reduced only very slightly. Insulin transports glucose to the body's cells for use in cellular metabolic function. The cells convert excess glucose to an insoluble substance called glycogen to prevent it from interfering with cellular metabolism. Because this ultimately lowers blood glucose levels, insulin is secreted to prevent hyperglycemia (high blood sugar levels). Another hormone called glucagon performs the opposite function of insulin, causing cells to convert glycogen to glucose and stimulating new glucose production (gluconeogenesis) to raise blood sugar levels. Negative feedback between insulin and glucagon levels controls blood sugar homeostasis.

Causes of Homeostatic Disruption

People with type 1 diabetes do not produce insulin due to auto-immune destruction of the insulin producing cells, while people with type 2 diabetes have chronic high blood glucose levels that cause insulin resistance. With diabetes, blood glucose is increased by normal glucagon activity, but the lack of or resistance to insulin means that blood sugar levels are unable to return to normal. This causes metabolic changes that result in diabetes symptoms like weakened blood vessels and frequent urination. Diabetes is normally treated with insulin injections, which replaces the missing negative feedback of normal insulin secretions.



Homeostasis of Glucose Metabolism: This image illustrates glucose metabolism over the course of a day. Homeostasis may become imbalanced if the pancreas is overly stressed, making it unable to balance glucose metabolism. This can lead to diabetes.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Homeostasis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Homeost...rol_mechanisms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homeostasis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homeostasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- positive feedback. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/positive_feedback**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- negative feedback. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/negative_feedback**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ACTH Negative Feedback. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...e_Feedback.svg**. **License**: ***CC BY: Attribution***
- Temperature Regulation. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...Regulation.jpg**. **License**: ***Public Domain: No Known Copyright***
- Wikipedia:Text of Creative Commons Attribution-ShareAlike 3.0 Unported License. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wikiped...ported_License**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homeostasis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homeostasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diabetes. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diabetes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- blood sugar regulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/blood%2...r%20regulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ACTH Negative Feedback. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...e_Feedback.svg**. **License**: ***CC BY: Attribution***
- Temperature Regulation. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...Regulation.jpg**. **License**: ***Public Domain: No Known Copyright***
- Glucose-day-english. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ay-english.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 1.3B: Disease as Homeostatic Imbalance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4: MAPPING THE BODY

1.4A: Anatomical Position
1.4B: Directional Terms
1.4C: Regional Terms and Axes
1.4D: Body Planes and Sections
1.4E: Body Cavities
1.4F: Abdominopelvic Regions

This page titled 1.4: Mapping the Body is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4A: Anatomical Position

When an organism is in its standard anatomical position, positional descriptive terms are used to indicate regions and features.

Learning Objectives

- Describe the standard position in human anatomy

Key Points


- In standard anatomical position, the limbs are placed similarly to the supine position imposed on cadavers during autopsy.
- The anatomical position of the skull is the Frankfurt plane. In this position, the lower margins of the orbitals (eye sockets), the lower margin of the orbits, and the upper margins of the ear canals (poria) lie in the same horizontal plane.
- Because animals can change orientation with respect to their environments and appendages can change position with respect to the body, positional descriptive terms refer to the organism only in its standard anatomical position to prevent confusion.

Key Terms

- **appendage**: A limb of the body.
- **supine**: Lying on its back, reclined.
- **anatomical position**: The standard position in which the body is standing with feet together, arms to the side, and head, eyes, and palms facing forward.

The Need for Standardization

Standard anatomical position is the body orientation used when describing an organism's anatomy. Standardization is necessary to avoid confusion since most organisms can take on many different positions that may change the relative placement of organs. All descriptions refer to the organism in its standard anatomical position, even when the organism's appendages are in another position. Thus, the standard anatomical position provides a "gold standard" when comparing the anatomy of different members of the same species.




Relative location in the anatomical position: Many terms are used to describe relative location on the body. Cranial refers to features closer to the head, while caudal refers to features closer to the feet. The front of the body is referred to as anterior or ventral, while the back is referred to as posterior or dorsal. Proximal and distal describe relative position on the limbs. Proximal refers to a feature that is closer to the torso, while distal refers to a feature that is closer to the fingers/toes. Medial and lateral refer to position relative to the midline, which is a vertical line drawn through the center of the forehead, down through the belly button to the floor. Medial indicates a feature is closer to this line, while lateral indicates features further from this line.

Standard Anatomical Position in Humans

The standard anatomical position is agreed upon by the international medical community. In this position, a person is standing upright with the lower limbs together or slightly apart, feet flat on the floor and facing forward, upper limbs at the sides with the palms facing forward and thumbs pointing away from the body, and head and eyes directed straight ahead. In addition, the arms are usually placed slightly apart from the body so that the hands do not touch the sides. The positions of the limbs, particularly the arms, have important implications for directional terms in those appendages.

The basis for the standard anatomical position in humans comes from the supine position used for examining human cadavers during autopsies. Dissection of cadavers was one of the primary ways humans learned about anatomy throughout history, which has tremendously influenced the ways by which anatomical knowledge has developed into the scientific field of today.



Standard anatomical position: The regions of the body in standard anatomical position, in which the body is erect.

In humans, the standard anatomical position of the skull is called the Frankfurt plane. In this position, the orbitales (eye sockets), lower margins of the orbits, and the poria (ear canal upper margins) all lie in the same horizontal plane. This orientation represents the position of the skull if the subject were standing upright and looking straight ahead.

It is important to note that all anatomical descriptions are based on the standard anatomical position unless otherwise stated.

This page titled 1.4A: Anatomical Position is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4B: Directional Terms

Positional terms give precise descriptions of anatomical relationships and allow for consistency when referencing anatomical positions.

Learning Objectives

- Identify the anatomical terms that define the human body in space

Key Points

- Descriptions of directional terms include: a) superior (head) and inferior (caudal), b) anterior and posterior, c) lateral and medial, d) deep and superficial, e) proximal and distal, and f) dorsal and ventral.
- Directional terms provide comparison of anatomical position by comparing the locations of different structures in the body.

Key Terms

- **directional terms**: Directional terms are words used to describe the location of an anatomical structure by comparing its position to other structures within the body or within the orientation of the body itself.

Navigating Anatomy with Language

Directional terms provide precise descriptions of a structure's location. They allow a description of anatomical position by comparing location relative to other structures or within the rest of the body. Standard anatomical terms for direction include:

- *Superior and inferior (cranial and caudal)* are used when referring to parts of the body which are toward an end of the body. Superior structures are toward the head (cranial) while inferior (caudal) structures are toward the feet. Examples include the superior and inferior vena cava, which carry deoxygenated blood away from the head (superior) and from the lower body (inferior) to the heart.
- *Anterior and posterior* are sometimes used in place of superior and inferior, respectively. These words are used more often for animal anatomy and rarely and only with very specific meaning in human anatomy. Anterior refers to the side of the structure facing up in the standard anatomical position while posterior refers to the bottom side. For example, the pituitary gland has an anterior and posterior side, each of which secretes different types of hormones.
- *Dorsal and ventral* are sometimes used in place of anterior and posterior, respectively. Dorsal means the back side or upper side, while ventral means the frontal or lower side. These are mostly used with animal anatomy, but can be used in human anatomy as long as they are describing the side of an appendage. One example is the dorsal fin in fish, found on the upper side of the fish's body.
- *Lateral* is used to describe anything closer to the sides of the body (toward the arms, in the standard anatomical position), while*medial* is used to describe anything toward the middle of the body. In general, many structures of the human body are bilateral and symmetrical with the middle of the body, such as the lungs or the arms.
- *Deep* refers to structures closer to the interior center of the body. For example, bones in an appendage are located deeper than the muscles. *Superficial* is used to describe structures that are closer to the exterior surface of the body. For example, the outer layers of skin are superficial to deeper layers of skin.
- *Proximal and Distal* describe one point relative to another. Proximal refers to a point closer to the reference point while distal refers to a point farther away. When describing appendages, the proximal end of the appendage connects the appendage to the body, while the distal end is away from the body.




Diagram of anatomical terms: Directional axes in a tetrapod.

This page titled 1.4B: Directional Terms is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4C: Regional Terms and Axes

Regional directional terms include anterior and posterior, dorsal and ventral, and lateral and medial.

Learning Objectives

- Describe how axes give direction, detail, and location when describing a region of the body

Key Points

- Regional terms describe the different parts of the body by the structures and functions of a specific region. The most basic regional terms are the axial and appendicular regions.
- Axes use directional terms to describe the location and orientation of a specific region.
- The directional term lateral is used to describe structures divided by a left-to-right axis.

Key Terms

- **ventral**: On the front side of the human body or the corresponding surface of an animal, usually the lower surface.
- **posterior**: Nearer the caudal end of the body in quadrupeds or the dorsal end in bipeds.
- **axis**: A line between two points that is used to give direction to an anatomical region.

Regional Terms in Anatomy

Regional terms describe anatomy by dividing the parts of the body into different regions that contain structures that are involved in similar functions. Two primary terms are used to describe the main regions of the body:

- *The Axial Region* makes up the main axis of the human body and includes the head, neck, chest, and trunk.
- *The Appendicular Region* makes up the parts of the human body that connect to the axial region. This includes the limbs and appendages.

These are the two basic categories of regional terms; however, many other terms are used to describe smaller regions within the axial and appendicular regions. For example, the brachial region consists of the arm as a part of the appendicular region, while the abdominal region consists of the abdomen as a smaller part of the axial region.

The abdominal region is subdivided into even smaller regions based on different functions of groups of organs and tissues in that region. If a person is experiencing pain in one part of the abdominal region, then the smaller regional divisions can help determine the organs involved in the problem to better treat symptoms.

Axes Describe Relative Positions


Another method for describing region An axis uses a straight line between two parts of the body to describe a region of the body with linear direction. For example, blood can be said to flow
in a proximal or distal direction through a region marked by that axis. The X, Y, and Z axes of the Cartesian coordinate system are used describe the specific location of an axis in standard anatomical position.

Many types of axes can give regional direction. Any pair of corresponding directional terms can be combined to form an axis (such as proximal-distal for an appendage).

- *The Dorsoventral axis (DV axis)* is formed by the connection of the dorsal and ventral points of a region. The region between the belly (ventral) and back (dorsal) is often described by a DV axis.
- *The Anterioposterior axis (AP axis)* is the axis formed by the connection of the anterior (top) and posterior (bottom) ends of

a region. The AP axis of a region is by definition perpendicular to the DV axis and vice-versa.

- *The Left-to-right axis* is the axis connecting the left and right hand sides of a region. It is used to describe the lateral sides of a region, which in humans are often symmetrical around the center of the body. It is perpendicular to both the DV and AP axes.



Different Directional AP Axes in Three Body Segments of a Horse: Axis (A) (in red) shows the AP axis of the tail, (B) shows the AP axis of the neck, and (C) shows the AP axis of the head.

Axes give more clarity and detail for describing the location of an anatomical region. They are commonly used in both zoology and human anatomy, and can be paired with body planes to give even more detail to anatomical direction, region, and location.

This page titled 1.4C: Regional Terms and Axes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4D: Body Planes and Sections

There are three basic reference planes used in anatomy: the sagittal plane, the coronal plane, and the transverse plane.

Learning Objectives

- Identify the three basic anatomical reference planes

Key Points

- A coronal or frontal plane divides the body into dorsal and ventral (back and front, or posterior and anterior) portions.
- A transverse plane, also known as an axial plane or cross-section, divides the body into cranial and caudal (head and tail) portions.
- A sagittal plane divides the body into sinister and dexter (left and right) portions.
- Body planes have several uses within the anatomy field, including in medical imaging, descriptions of body motion, and embryology.

Key Terms


- **coronal plane**: Any vertical plane that divides the body into anterior and posterior (belly and back) sections.
- **transverse plane**: Any plane that divides the body into superior and inferior parts, roughly perpendicular to the spine.
- **sagittal plane**: Any imaginary plane parallel to the median plane.

What Are Body Planes?

Body planes are hypothetical geometric planes used to divide the body into sections. They are commonly used in both human and zoological anatomy to describe the location or direction of bodily structures. Reference planes are the standard planes used in anatomical terminology and include:

- *The sagittal plane (lateral or Y-Z plane)* divides the body into sinister and dexter (left and right) sides. The midsagittal (median) plane is in the midline through the center of the body, and all other sagittal planes are parallel to it.
- *The coronal plane (frontal or Y-X plane)* divides the body into dorsal and ventral (back and front) portions. It also separates the anterior and posterior portions.
- *The transverse plane (axial or X-Z plane)* divides the body into superior and inferior (head and tail) portions. It is typically a horizontal plane through the center of the body and is parallel to the ground.

While these are the major reference planes of the body, other planes are commonly used in relation to these three. A longitudinal plane is any plane perpendicular to the transverse plane, while parasaggital planes are parallel to the saggital plane.
The coronal plane, the sagittal plane, and the parasaggital planes are examples of longitudinal
planes.



Anatomical Planes in a Human: There are three basic planes in zoological anatomy: sagittal, coronal, and transverse. A human in the anatomical position, can be described using a coordinate system with the Z-axis going from front to back, the X-axis going from left to right, and the Y-axis going from up to down.

Applications of Body Planes

Medical imaging techniques such as sonography, CT scans, MRI scans, or PET scans are one of the primary applications of body planes. By imaging a patient in standard anatomical position, a radiologist can build an X-Y-Z axis around the patient to apply body planes to the images. The planes can then be used to identify and locate the positions of the patient's internal organs. Individual organs can also be divided by planes to help identify smaller structures within that organ.

Body planes are used to describe anatomical motion in the X-Y-Z coordinate system that the body moves through. An anatomist could model a limb's range of motion by measuring which planes the limb can move through and how far it is able to travel.

Anatomical change during embryological development is also described and measured with body planes. For example, during human embryonic development the coronal plane is horizontal, but becomes vertical as the embryo develops into a fetus. In comparative embryology, body planes provide a basis for comparing the ways in which different types of organisms develop anatomically within the womb.

This page titled 1.4D: Body Planes and Sections is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4E: Body Cavities

Vertebrates have fluid-filled spaces called body cavities that contain the organs.

Learning Objectives

- Describe the major cavities of the human body


Key Points

- The dorsal cavity contains the primary organs of the nervous system, including the brain and spinal cord.
- The diaphragm is a sheet of muscle that separates the thoracic cavity from the abdominal cavity.
- Special membrane tissues surround the body cavities, such as the meninges of the dorsal cavity and the mesothelium of the ventral cavity.
- The mesothelium consists of the pleura of the lungs, the pericardium of the heart, and the peritoneum of the abdominopelvic cavity.

Key Terms

- **abdominoplevic cavity**: The ventral body chamber that contains the abdominal cavity (primarily digestive system) and the pelvic cavity (primarily reproductive system).
- **dorsal cavity**: The cavity in the back of the body that contains the cranial and vertebral cavities, which house the brain and spinal cord respectively.
- **Thoracic Cavity**: The ventral body chamber that contains the pericardial cavity (the heart) and the pleural cavity (the lungs).

By the broadest definition, a body cavity is any fluid-filled space in a multicellular organism. However, the term usually refers to the space where internal organs develop, located between the skin and the outer lining of the gut cavity."The human body cavity," normally refers to the ventral body cavity because it is by far the largest one in volume. Blood vessels are not considered cavities but may be held within cavities. Most cavities provide room for the organs to adjust to changes in the organism's position. They usually contains protective membranes and sometimes bones that protect the organs.



Anatomical terminology for body cavities: Humans have multiple body cavities, including the cranial cavity, the vertebral cavity, the thoracic cavity (containing the pericardial cavity and the pleural cavity), the abdominal cavity, and the pelvic cavity. In

mammals, the diaphragm separates the thoracic cavity from the abdominal cavity.

Dorsal

The dorsal cavity is a continuous cavity located on the dorsal side of the body. It houses the organs of the upper central nervous system, including the brain and the spinal cord. The meninges is a multi-layered membrane within the dorsal cavity that envelops and protects the brain and spinal cord.

Cranial

The cranial cavity is the anterior portion of the dorsal cavity consisting of the space inside the skull. This cavity contains the brain, the meninges of the brain, and cerebrospinal fluid.

Vertebral

The vertebral cavity is the posterior portion of the dorsal cavity and contains the structures within the vertebral column. These include the spinal cord, the meninges of the spinal cord, and the fluid-filled spaces between them. This is the most narrow of all body cavities, sometimes described as threadlike.

Ventral

The ventral cavity, the interior space in the front of the body, contains many different organ systems. The organs within the ventral cavity are also called viscera. The ventral cavity has anterior and posterior portions divided by the diaphragm, a sheet of skeletal muscle found beneath the lungs.

Thoracic

The thoracic cavity is the anterior ventral body cavity found within the rib cage in the torso. It houses the primary organs of the cardiovascular and respiratory systems, such as the heart and lungs, but also includes organs from other systems, such as the esophagus and the thymus gland. The thoracic cavity is lined by two types of mesothelium, a type of membrane tissue that lines the ventral cavity: the pleura lining of the lungs, and the pericadium lining of the heart.

Abdominopelvic

The abdominoplevic cavity is the posterior ventral body cavity found beneath the thoracic cavity and diaphragm. It is generally divided into the abdominal and pelvic cavities. The abdominal cavity is not contained within bone and houses many organs of the digestive and renal systems, as well as some organs of the endocrine system, such as the adrenal glands. The pelvic cavity is contained within the pelvis and houses the bladder and reproductive system. The abdominopelvic cavity is lined by a type of mesothelium called the peritoneum.

This page titled 1.4E: Body Cavities is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

1.4F: Abdominopelvic Regions

The abdomen is subdivided into four quadrants and nine areas.

Learning Objectives

- Distinguish among the abdominopelvic regions of the body


Key Points

- The abdominopelvic cavity can be subdivided into four quadrants and nine areas.
  The quadrants are labeled by location: the right upper, right lower, left upper, and left lower quadrants.
- The nine regions are smaller than the four abdominopelvic quadrants and include the right hypochondriac, right lumbar, right illiac, epigastric, umbilical, hypogastric (or pubic), left hypochondriac, left lumbar, and left illiac divisions.
- The perineum is sometimes considered to be the tenth division.
- The purpose of the abdominal divisions is to describe regional anatomy in the abdomen, and to help clinicians determine which organ and tissues are involved in a disease based on which regions experience pain.

Key Terms

- **left upper quadrant**: The left upper quadrant is the location of the left portion of the liver, the larger portion of the stomach, the pancreas, left kidney, spleen, portions of the transverse and descending colon, and parts of the small intestine.
- **Right upper quadrant**: The right upper quadrant contains the right portion of the liver, gallbladder, right kidney, a small portion of the stomach, portions of the ascending and transverse colon, and parts of the small intestine.
- **left lower quadrant**: The left lower quadrant houses the majority of the small intestine, some of the large intestine, the left female reproductive organs, and the left ureter.
- **nine divisions**: An alternate system for dividing the abdominopelvic cavity into regions.
- **right lower quadrant**: In the right lower quadrant sits the cecum, appendix, part of the small intestines, the right female reproductive organs, and the right ureter.

Anatomists and medical personnel divide the abdominopelvic cavity into smaller regions to facilitate study and discussion. These divisions are often used to categorize the individual abdominal organs by their location and function and are used by clinicians to help diagnose the source of abdominal pain and determine appropriate treatment. The most common divisions for the abdominopelvic region are the four quadrants and nine regions.



Abdominal nine divisions (a) and quadrant regions (b): The abdomen is subdivided into four quadrants and nine areas.

Abdominal Four Quadrants

The abdominopelvic region can be divided into four quadrants. These quadrants are defined by the intersection of the sagittal plane with the umbilical plane (the transverse plane through the navel). Clinicians use these regions to determine the organs and tissues that may be causing pain or discomfort in that region.

Right Upper Quadrant

The right upper quadrant contains the right portion of the liver, the gallbladder, right kidney, a small portion of the stomach, the duodenum, the head of the pancreas, portions of the ascending and transverse colon, and parts of small intestine. Pain in this region is associated with infection and inflammation in the gallbladder and liver or peptic ulcers in the stomach.

Left Upper Quadrant

The left upper quadrant is the location of the left portion of the liver, part of the stomach, the pancreas, left kidney, spleen, portions of the transverse and descending colon, and parts of the small intestine. Pain in this region is associated with malrotation of the intestine and colon.

Right Lower Quadrant

In the right lower quadrant sits the cecum, appendix, part of the small intestines, the right half of the female reproductive system, and the right ureter. Pain in this region is most commonly associated with appendicitis.

Left Lower Quadrant

The left lower quadrant houses the majority of the small intestine, some of the large intestine, the left half of the female reproductive system, and the left ureter. Pain in this region is generally associated with colitis (inflammation of the large intestine) as well as pelvic inflammatory disease and ovarian cysts in females.

Abdominal Nine Divisions

The nine divisions of the abdominopelvic region are smaller than the four quadrants, allowing for a more detailed discussion.

These divisions are marked by two parasagittal and two transverse planes centered around the navel. Most organs are part of multiple regions, including the gallbladder, duodenum, stomach, kidneys, spleen, small intestine and colon. The perineum (the area beneath the hypogastric region at the bottom of the pelvic cavity) is sometimes considered to be a tenth division in this system.

Right Hypochondriac

The right hypochondriac region contains the right portion of the liver, the gallbladder, the right kidney, and parts of the small intestine.

Left Hypochondriac

The left hypochondriac region contains part of the spleen, the left kidney, part of the stomach, the pancreas, and parts of the colon.

Epigastric

The epigastric (above stomach) region contains the majority of the stomach, part of the liver, part of the pancreas, part of the duodenum, part of the spleen, and the adrenal glands. This region pushes out when the diaphragm contracts during breathing.

Right Lumbar

The right lumbar region consists of the gallbladder, the right kidney, part of the liver, and the ascending colon.

Left Lumbar

The left lumbar region consists of the descending colon, the left kidney, and part of the spleen.

Umbilical

The umbilical region contains the umbilicus (navel), and many parts of the small intestine, such as part of the duodenum, the jejunum, and the illeum. It also contains the transverse colon (the section between the ascending and descending colons) and the bottom portions of both the left and right kidney.

Right Iliac

The right iliac region contains the appendix, cecum, and the right iliac fossa. It is also commonly referred to as the right inguinal region. Pain in this area is generally associated with appendicitis.

Left Iliac

The left illiac region contains part of the descending colon, the sigmoid colon, and the left illiac fossa. It is also commonly called the left inguinal region.

Hypogastric

The hypogastric region (below the stomach) contains the organs around the pubic bone. These include bladder, part of the sigmoid colon, the anus, and many organs of the reproductive system, such as the uterus and ovaries in females and the prostate in males.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Anatomical terms of location. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anatomi...man.29_anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tonye Ogele, Anatomical Terms. January 16, 2014. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/contents/1a4a3bd1-f36...7fcfe1bac6ef@1**. **License**: ***CC BY: Attribution***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...mical-position**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- supine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/supine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/List_of...Human_Body.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg**. **License**: ***CC BY: Attribution***
- **Provided by**: Wikipedia. **Located at**: **Wikipedia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tomical-planes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- directional terms. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/directional%20terms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/List_of...Human_Body.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg**. **License**: ***CC BY: Attribution***
- Horse Axes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Horse_Axes.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Terms for anatomical location. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Terms_f...ectional_terms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomical terms of location. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anatomi...ms_of_location**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventral. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventral**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- posterior. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/posterior**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/List_of...Human_Body.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg**. **License**: ***CC BY: Attribution***
- Horse Axes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Horse_Axes.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Horse Axes 2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Horse_Axes_2.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- coronal plane. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/coronal_plane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Terms for anatomical location. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Terms_for_anatomical_location**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...ansverse-plane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...sagittal-plane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/List_of...Human_Body.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg**. **License**: ***CC BY: Attribution***
- Horse Axes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Horse_Axes.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Horse Axes 2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Horse_Axes_2.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Human anatomy planes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Hu...omy_planes.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Body cavity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and Physiology of Animals/Body Organisation. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy...3Body_Cavities. **License**: *CC BY-SA: Attribution-ShareAlike*
- coelom. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/coelom. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physics/de...horacic-cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- abdominal cavity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/abdominal_cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/List_of...Human_Body.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg. **License**: *CC BY: Attribution*
- Horse Axes. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Horse_Axes.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Horse Axes 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Horse_Axes_2.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human anatomy planes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Hu...omy_planes.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Body cavities. **Provided by**: Wikimedia. **Located at**: en.Wikipedia.org/wiki/Anatomi...y_cavities.jpg. **License**: *CC BY: Attribution*
- Human anatomical terms. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_a...men_and_pelvis. **License**: *CC BY-SA: Attribution-ShareAlike*
- right lower quadrant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/right%20lower%20quadrant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...upper-quadrant. **License**: *CC BY-SA: Attribution-ShareAlike*
- left lower quadrant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/left%20lower%20quadrant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Right upper quadrant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Right%20upper%20quadrant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Regions of the body. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/List_of...Human_Body.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomical position. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/Anatomi...onal_Terms.jpg. **License**: *CC BY: Attribution*
- Horse Axes. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Horse_Axes.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Horse Axes 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Horse_Axes_2.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human anatomy planes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Hu...omy_planes.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Body cavities. **Provided by**: Wikimedia. **Located at**: en.Wikipedia.org/wiki/Anatomi...y_cavities.jpg. **License**: *CC BY: Attribution*
- Abdominal Quadrant Regions. **Provided by**: Wikimedia. **Located at**: en.Wikipedia.org/wiki/Anatomi...nt_Regions.jpg. **License**: *CC BY: Attribution*

This page titled 1.4F: Abdominopelvic Regions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2: THE CHEMICAL BUILDING BLOCKS OF LIFE

Of the 92 natural elements, 25 are essential for life. There are six main elements that are the fundamental building blocks of life. They are, in order of least to most common: sulfur, phosphorous, oxygen, nitrogen, carbon, and hydrogen.

2.1: Atomic Structure

2.1A: Overview of Atomic Structure

2.1B: Atomic Number and Mass Number

2.2: Chemical Bonds

2.2A: Covalent Bonds and Other Bonds and Interactions

2.2B: Ions and Ionic Bonds

2.2C: The Chemical Basis for Life

2.3: Inorganic Compounds

2.3A: pH, Buffers, Acids, and Bases

2.3B: Water's Cohesive and Adhesive Properties

2.3C: Water's High Heat Capacity

2.3D: Water's States: Gas, Liquid, and Solid

2.3E: Water's Solvent Properties

2.4: Organic Compounds

2.4A: Organic Molecules and Functional Groups

2.4B: Types of Biological Macromolecules

2.4C: Dehydration Synthesis

This page titled 2: The Chemical Building Blocks of Life is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.1: ATOMIC STRUCTURE



- Topic hierarchy

2.1A: Overview of Atomic Structure

2.1B: Atomic Number and Mass Number

This page titled 2.1: Atomic Structure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.1A: Overview of Atomic Structure

Atoms are made up of particles called protons, neutrons, and electrons, which are responsible for the mass and charge of atoms.

Learning Objectives

- Discuss the electronic and structural properties of an atom

Key Points

- An atom is composed of two regions: the nucleus, which is in the center of the atom and contains protons and neutrons, and the outer region of the atom, which holds its electrons in orbit around the nucleus.
- Protons and neutrons have approximately the same mass, about 1.67×10^{-24} grams, which scientists define as one atomic mass unit (amu) or one Dalton.
- Each electron has a negative charge (–1) equal to the positive charge of a proton (+1).
- Neutrons are uncharged particles found within the nucleus.

Key Terms


- **atom**: The smallest possible amount of matter which still retains its identity as a chemical element, consisting of a nucleus surrounded by electrons.
- **proton**: Positively charged subatomic particle forming part of the nucleus of an atom and determining the atomic number of an element. It weighs 1 amu.
- **neutron**: A subatomic particle forming part of the nucleus of an atom. It has no charge. It is equal in mass to a proton or it weighs 1 amu.

An atom is the smallest unit of matter that retains all of the chemical properties of an element. Atoms combine to form molecules, which then interact to form solids, gases, or liquids. For example, water is composed of hydrogen and oxygen atoms that have combined to form water molecules. Many biological processes are devoted to breaking down molecules into their component atoms so they can be reassembled into a more useful molecule.

Atomic Particles

Atoms consist of three basic particles: protons, electrons, and neutrons. The nucleus (center) of the atom contains the protons (positively charged) and the neutrons (no charge). The outermost regions of the atom are called electron shells and contain the electrons (negatively charged). Atoms have different properties based on the arrangement and number of their basic particles.

The hydrogen atom (H) contains only one proton, one electron, and no neutrons. This can be determined using the atomic number and the mass number of the element (see the concept on atomic numbers and mass numbers).



Structure of an atom: Elements, such as helium, depicted here, are made up of atoms. Atoms are made up of protons and neutrons located within the nucleus, with electrons in orbitals surrounding the nucleus.

Atomic Mass

Protons and neutrons have approximately the same mass, about 1.67×10^{-24} grams. Scientists define this amount of mass as one atomic mass unit (amu) or one Dalton. Although similar in mass, protons are positively charged, while neutrons have no charge. Therefore, the number of neutrons in an atom contributes significantly to its mass, but not to its charge.

Electrons are much smaller in mass than protons, weighing only 9.11×10^{-28} grams, or about 1/1800 of an atomic mass unit. Therefore, they do not contribute much to an element's overall atomic mass. When considering atomic mass, it is customary to ignore the mass of any electrons and calculate the atom's mass based on the number of protons and neutrons alone.

Electrons contribute greatly to the atom's charge, as each electron has a negative charge equal to the positive charge of a proton. Scientists define these charges as "+1" and "–1." In an uncharged, neutral atom, the number of electrons orbiting the nucleus is equal to the number of protons inside the nucleus. In these atoms, the positive and negative charges cancel each other out, leading to an atom with no net charge.

Protons, Neutrons, and Electrons			
	Charge	Mass (amu)	Location
Proton	+1	1	nucleus
Neutron	0	1	nucleus
Electron	–1	0	orbitals

Both protons and neutrons have a mass of 1 amu and are found in the nucleus. However, protons have a charge of +1, and neutrons are uncharged. Electrons have a mass of approximately 0 amu, orbit the nucleus, and have a charge of–1.

Exploring Electron Properties: Compare the behavior of electrons to that of other charged particles to discover properties of electrons such as charge and mass.

Volume of Atoms

Accounting for the sizes of protons, neutrons, and electrons, most of the volume of an atom—greater than 99 percent—is, in fact, empty space. Despite all this empty space, solid objects do not just pass through one another. The electrons that surround all

atoms are negatively charged and cause atoms to repel one another, preventing atoms from occupying the same space. These intermolecular forces prevent you from falling through an object like your chair.

Interactive: Build an Atom: Build an atom out of protons, neutrons, and electrons, and see how the element, charge, and mass change. Then play a game to test your ideas!

This page titled 2.1A: Overview of Atomic Structure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.1B: Atomic Number and Mass Number

The atomic number is the number of protons in an element, while the mass number is the number of protons plus the number of neutrons.

Learning Objectives

- Determine the relationship between the mass number of an atom, its atomic number, its atomic mass, and its number of subatomic particles

Key Points

- Neutral atoms of each element contain an equal number of protons and electrons.
- The number of protons determines an element's atomic number and is used to distinguish one element from another.
- The number of neutrons is variable, resulting in isotopes, which are different forms of the same atom that vary only in the number of neutrons they possess.
- Together, the number of protons and the number of neutrons determine an element's mass number.
- Since an element's isotopes have slightly different mass numbers, the atomic mass is calculated by obtaining the mean of the mass numbers for its isotopes.

Key Terms


- **mass number**: The sum of the number of protons and the number of neutrons in an atom.
- **atomic number**: The number of protons in an atom.
- **atomic mass**: The average mass of an atom, taking into account all its naturally occurring isotopes.

Atomic Number

Neutral atoms of an element contain an equal number of protons and electrons. The number of protons determines an element's atomic number (Z) and distinguishes one element from another. For example, carbon's atomic number (Z) is 6 because it has 6 protons. The number of neutrons can vary to produce isotopes, which are atoms of the same element that have different numbers of neutrons. The number of electrons can also be different in atoms of the same element, thus producing ions (charged atoms). For instance, iron, Fe, can exist in its neutral state, or in the +2 and +3 ionic states.

Mass Number

An element's mass number (A) is the sum of the number of protons and the number of neutrons. The small contribution of mass from electrons is disregarded in calculating the mass number. This approximation of mass can be used to easily calculate how many neutrons an element has by simply subtracting the number of protons from the mass number. Protons and neutrons both weigh about one atomic mass unit or amu. Isotopes of the same element will have the same atomic number but different mass numbers.



Atomic number, chemical symbol, and mass number: Carbon has an atomic number of six, and two stable isotopes with mass numbers of twelve and thirteen, respectively. Its average atomic mass is 12.11.

Scientists determine the atomic mass by calculating the mean of the mass numbers for its naturally-occurring isotopes. Often, the resulting number contains a decimal. For example, the atomic mass of chlorine (Cl) is 35.45 amu because chlorine is composed of several isotopes, some (the majority) with an atomic mass of 35 amu (17 protons and 18 neutrons) and some with an atomic mass of 37 amu (17 protons and 20 neutrons).

Given an atomic number (Z) and mass number (A), you can find the number of protons, neutrons, and electrons in a neutral atom. For example, a lithium atom (Z=3, A=7 amu) contains three protons (found from Z), three electrons (as the number of protons is equal to the number of electrons in an atom), and four neutrons (7 – 3 = 4).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Sunil Kumar Singh, Fundamental Force Types. October 27, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m14044/latest/**. **License**: ***CC BY: Attribution***
- neutron. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neutron**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- electron. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/electron**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- atom. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/atom**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- proton. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/proton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 30, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...e_02_01_01.jpg**. **License**: ***CC BY: Attribution***
- atomic number. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/atomic_number**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- A-level Chemistry/OCR/Atoms, Bonds and Groups/Atoms and Reactions/Atoms. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/A-level...eactions/Atoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...atomic-mass--2**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 30, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...e_02_01_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 27, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***

This page titled 2.1B: Atomic Number and Mass Number is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.2: CHEMICAL BONDS



- Topic hierarchy

2.2A: Covalent Bonds and Other Bonds and Interactions

2.2B: Ions and Ionic Bonds

2.2C: The Chemical Basis for Life

This page titled 2.2: Chemical Bonds is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.2A: Covalent Bonds and Other Bonds and Interactions

Covalent bonds result from a sharing of electrons between two atoms and hold most biomolecules together.

Learning Objectives

- Compare the relative strength of different types of bonding interactions

Key Points

- A polar covalent bond arises when two atoms of different electronegativity share two electrons unequally.
- A non-polar covalent bond is one in which the electrons are shared equally between two atoms.
- Hydrogen bonds and Van Der Waals are responsible for the folding of proteins, the binding of ligands to proteins, and many other processes between molecules.

Key Terms

- **hydrogen bond**: A weak bond in which a hydrogen atom in one molecule is attracted to an electronegative atom (usually nitrogen or oxygen) in the same or different molecule.
- **covalent bond**: A type of chemical bond where two atoms are connected to each other by the sharing of two or more electrons.
- **dipole**: Any object (such as a magnet, polar molecule or antenna), that is oppositely charged at two points (or poles).

EXAMPLES

Adenosine triphosphate, or ATP, is the most-commonly used cofactor in all of biology. Its biosynthesis involves breaking the triple bond of molecular nitrogen, or N_2, followed by the formation of several carbon-nitrogen single and double bonds.

The octet rule can be satisfied by the sharing of electrons between atoms to form covalent bonds. These bonds are stronger and much more common than are ionic bonds in the molecules of living organisms. Covalent bonds are commonly found in carbon-based organic molecules, such as DNA and proteins. Covalent bonds are also found in inorganic molecules such as H_2O, CO_2, and O_2. One, two, or three pairs of electrons may be shared between two atoms, making single, double, and triple bonds, respectively. The more covalent bonds between two atoms, the stronger their connection. Thus, triple bonds are the strongest.


The strength of different levels of covalent bonding is one of the main reasons living organisms have a difficult time in acquiring nitrogen for use in constructing nitrogenous molecules, even though molecular nitrogen, N_2, is the most abundant gas in the atmosphere. Molecular nitrogen consists of two nitrogen atoms triple bonded to each other. The resulting strong triple bond makes it difficult for living systems to break apart this nitrogen in order to use it as constituents of biomolecules, such as proteins, DNA, and RNA.

The formation of water molecules is an example of covalent bonding. The hydrogen and oxygen atoms that combine to form water molecules are bound together by covalent bonds. The electron from the hydrogen splits its time between the incomplete outer shell of the hydrogen atom and the incomplete outer shell of the oxygen atom. In return, the oxygen atom shares one of its electrons with the hydrogen atom, creating a two-electron single covalent bond. To completely fill the outer shell of oxygen, which has six electrons in its outer shell, two electrons (one from each hydrogen atom) are needed. Each hydrogen atom needs only a single electron to fill its outer shell, hence the well-known formula H_2O. The electrons that are shared between the two elements fill the outer shell of each, making both elements more stable.

Polar Covalent Bonds

There are two types of covalent bonds: polar and nonpolar. In a polar covalent bond, the electrons are unequally shared by the

atoms because they are more attracted to one nucleus than the other. The relative attraction of an atom to an electron is known as its electronegativity: atoms that are more attracted to an electron are considered to be more electronegative. Because of the unequal distribution of electrons between the atoms of different elements, a slightly positive (δ+) or slightly negative (δ-) charge develops. This partial charge is known as a dipole; this is an important property of water and accounts for many of its characteristics. The dipole in water occurs because oxygen has a higher electronegativity than hydrogen, which means that the shared electrons spend more time in the vicinity of the oxygen nucleus than they do near the nucleus of the hydrogen atoms.



Polar and Nonpolar Covalent Bonds: Whether a molecule is polar or nonpolar depends both on bond type and molecular shape. Both water and carbon dioxide have polar covalent bonds, but carbon dioxide is linear, so the partial charges on the molecule cancel each other out.


Nonpolar Covalent Bonds

Nonpolar covalent bonds form between two atoms of the same element or between different elements that share electrons equally. For example, molecular oxygen (O_2) is nonpolar because the electrons will be equally distributed between the two oxygen atoms. The four bonds of methane are also considered to be nonpolar because the electronegativies of carbon and hydrogen are nearly identical.

Hydrogen Bonds and Van Der Waals Interactions

Not all bonds are ionic or covalent; weaker bonds can also form between molecules. Two types of weak bonds that frequently occur are hydrogen bonds and van der Waals interactions. Without these two types of bonds, life as we know it would not exist.

Hydrogen bonds provide many of the critical, life-sustaining properties of water and also stabilize the structures of proteins and DNA, the building block of cells. When polar covalent bonds containing hydrogen are formed, the hydrogen atom in that bond has a slightly positive charge (δ+) because the shared electrons are pulled more strongly toward the other element and away from the hydrogen atom. Because the hydrogen has a slightly positive charge, it's attracted to neighboring negative charges. The weak interaction between the δ+ charge of a hydrogen atom from one molecule and the δ- charge of a more electronegative atom is called a hydrogen bond. Individual hydrogen bonds are weak and easily broken; however, they occur in very large numbers in water and in organic polymers, and the additive force can be very strong. For example, hydrogen bonds are responsible for zipping together the DNA double helix.



Adenosine Triphosphate, ATP: Adenosine Triphosphate, or ATP, is the most commonly used cofactor in nature. Its biosynthesis involves the fixation of nitrogen to provide feedstocks that eventually produce the carbon-nitrogen bonds it contains.

Like hydrogen bonds, van der Waals interactions are weak interactions between molecules. Van der Waals attractions can occur between any two or more molecules and are dependent on slight fluctuations of the electron densities, which can lead to slight temporary dipoles around a molecule. For these attractions to happen, the molecules need to be very close to one another. These bonds, along with hydrogen bonds, help form the three-dimensional structures of the proteins in our cells that are required for their proper function.

Interactions between different types of molecules: In this interactive, you can explore how different types of molecules interact with each other based on their bonds.

This page titled 2.2A: Covalent Bonds and Other Bonds and Interactions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.2B: Ions and Ionic Bonds

Ionic bonds are attractions between oppositely charged atoms or groups of atoms where electrons are donated and accepted.

Learning Objectives

- Predict whether a given element will more likely form a cation or an anion

Key Points

- Ions form from elements when they gain or lose an electron causing the number of protons to be unequal to the number of electrons, resulting in a net charge.
- If there are more electrons than protons (from an element gaining one or more electrons), the ion is negatively charged and called an anion.
- If there are more protons than electrons (via loss of electrons), the ion is positively charged and is called a cation.
- Ionic bonds result from the interaction between a positively charged cation and a negatively charged anion.


Key Terms

- **ion**: An atom, or group of atoms, bearing an electrical charge, such as the sodium and chlorine atoms in a salt solution.
- **ionic bond**: A strong chemical bond caused by the electrostatic attraction between two oppositely charged ions.

Ions and Ionic Bonds

Some atoms are more stable when they gain or lose an electron (or possibly two) and form ions. This results in a full outermost electron shell and makes them energetically more stable. Now, because the number of electrons does not equal the number of protons, each ion has a net charge. Cations are positive ions that are formed by losing electrons (as the number of protons is now greater than the number of electrons). Negative ions are formed by gaining electrons and are called anions (wherein there are more electrons than protons in a molecule). Anions are designated by their elemental name being altered to end in “-ide.” For example, the anion of chlorine is called chloride, and the anion of sulfur is called sulfide.

This movement of electrons from one element to another is referred to as electron transfer. As illustrated, sodium (Na) only has one electron in its outer electron shell. It takes less energy for sodium to donate that one electron than it does to accept seven more electrons to fill the outer shell. When sodium loses an electron, it will have 11 protons, 11 neutrons, and only 10 electrons. This leaves it with an overall charge of +1 since there are now more protons than electrons. It is now referred to as a sodium ion. Chlorine (Cl) in its lowest energy state (called the ground state) has seven electrons in its outer shell. Again, it is more energy efficient for chlorine to gain one electron than to lose seven. Therefore, it tends to gain an electron to create an ion with 17 protons, 17 neutrons, and 18 electrons. This gives it a net charge of -1 since there are now more electrons than protons. It is now referred to as a chloride ion. In this example, sodium will donate its one electron to empty its shell, and chlorine will accept that electron to fill its shell. Both ions now satisfy the octet rule and have complete outer shells. These transactions can normally only take place simultaneously; in order for a sodium atom to lose an electron, it must be in the presence of a suitable recipient like a chlorine atom.



Electron Transfer Between Na and Cl: In the formation of an ionic compound, metals lose electrons and nonmetals gain electrons to achieve an octet. In this example, sodium loses one electron to empty its shell and chlorine accepts that electron to fill its shell.

Ionic bonds are formed between ions with opposite charges. For instance, positively charged sodium ions and negatively charged chloride ions bond together to form sodium chloride, or table salt, a crystalline molecule with zero net charge. The attractive force holding the two atoms together is called the electromagnetic force and is responsible for the attraction between oppositely charged ions.

Certain salts are referred to in physiology as electrolytes (including sodium, potassium, and calcium). Electrolytes are ions necessary for nerve impulse conduction, muscle contractions, and water balance. Many sports drinks and dietary supplements provide these ions to replace those lost from the body via sweating during exercise.

This page titled 2.2B: Ions and Ionic Bonds is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.2C: The Chemical Basis for Life

Carbon is the most important element to living things because it can form many different kinds of bonds and form essential compounds.

Learning Objectives

- Explain the properties of carbon that allow it to serve as a building block for biomolecules

Key Points

- All living things contain carbon in some form.
- Carbon is the primary component of macromolecules, including proteins, lipids, nucleic acids, and carbohydrates.
- Carbon's molecular structure allows it to bond in many different ways and with many different elements.
- The carbon cycle shows how carbon moves through the living and non-living parts of the environment.

Key Terms

- **octet rule**: A rule stating that atoms lose, gain, or share electrons in order to have a full valence shell of 8 electrons (has some exceptions).
- **carbon cycle**: the physical cycle of carbon through the earth's biosphere, geosphere, hydrosphere, and atmosphere; includes such processes as photosynthesis, decomposition, respiration and carbonification
- **macromolecule**: a very large molecule, especially used in reference to large biological polymers (e.g., nucleic acids and proteins)

Carbon is the fourth most abundant element in the universe and is the building block of life on earth. On earth, carbon circulates through the land, ocean, and atmosphere, creating what is known as the Carbon Cycle. This global carbon cycle can be divided further into two separate cycles: the geological carbon cycles takes place over millions of years, whereas the biological or physical carbon cycle takes place from days to thousands of years. In a nonliving environment, carbon can exist as carbon dioxide (CO_2), carbonate rocks, coal, petroleum, natural gas, and dead organic matter. Plants and algae convert carbon dioxide to organic matter through the process of photosynthesis, the energy of light.

Carbon is Important to Life




Carbon is present in all life: All living things contain carbon in some form, and carbon is the primary component of macromolecules, including proteins, lipids, nucleic acids, and carbohydrates. Carbon exists in many forms in this leaf, including in the cellulose to form the leaf's structure and in chlorophyll, the pigment which makes the leaf green.

In its metabolism of food and respiration, an animal consumes glucose ($C_6H_{12}O_6$), which combines with oxygen (O_2) to produce carbon dioxide (CO_2), water (H_2O), and energy, which is given off as heat. The animal has no need for the carbon dioxide and releases it into the atmosphere. A plant, on the other hand, uses the opposite reaction of an animal through photosynthesis. It intakes carbon dioxide, water, and energy from sunlight to make its own glucose and oxygen gas. The glucose is used for chemical energy, which the plant metabolizes in a similar way to an animal. The plant then emits the remaining oxygen into the environment.

Cells are made of many complex molecules called macromolecules, which include proteins, nucleic acids (RNA and DNA), carbohydrates, and lipids. The macromolecules are a subset of organic molecules (any carbon-containing liquid, solid, or gas) that are especially important for life. The fundamental component for all of these macromolecules is carbon. The carbon atom has unique properties that allow it to form covalent bonds to as many as four different atoms, making this versatile element ideal to serve as the basic structural component, or "backbone," of the macromolecules.

Structure of Carbon

Individual carbon atoms have an incomplete outermost electron shell. With an atomic number of 6 (six electrons and six protons), the first two electrons fill the inner shell, leaving four in the second shell. Therefore, carbon atoms can form four covalent bonds with other atoms to satisfy the octet rule. The methane molecule provides an example: it has the chemical formula CH_4. Each of its four hydrogen atoms forms a single covalent bond with the carbon atom by sharing a pair of electrons. This results in a filled outermost shell.



Structure of Methane: Methane has a tetrahedral geometry, with each of the four hydrogen atoms spaced 109.5° apart.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Biochemistry/Metabolism and energy. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Biochem...%23Ionic_bonds**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dipole. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/dipole**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- covalent bond. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/covalent_bond**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hydrogen bond. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hydrogen_bond**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...e_02_01_11.jpg**. **License**: ***CC BY: Attribution***
- ATP structure. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi..._structure.svg**. **License**: ***Public Domain: No Known Copyright***
- ionic bond. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ionic_bond**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Biochemistry/Metabolism and energy. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Biochem...%23Ionic_bonds**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...e_02_01_11.jpg**. **License**: ***CC BY: Attribution***
- ATP structure. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi..._structure.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest...e_02_01_10.jpg**. **License**: ***CC BY: Attribution***
- octet rule. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/octet_rule**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44393/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- The Carbon Cycle. **Provided by**: climate-jigsaw Wikispace. **Located at**: **http://climate-jigsaw.wikispaces.com/The+Carbon+Cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- macromolecule. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/macromolecule**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carbon cycle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/carbon_cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest/Figure_02_01_11.jpg**. **License**: ***CC BY: Attribution***
- ATP structure. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:ATP_structure.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Atoms, Isotopes, Ions, and Molecules: The Building Blocks. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44390/latest/Figure_02_01_10.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44393/latest/Figure_02_03_01f.jpg**. **License**: ***CC BY: Attribution***
- A leaf with laminar structure andu00a0pinnateu00a0venation. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Leaf%23mediaviewer/File:Leaf_1_web.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 2.2C: The Chemical Basis for Life is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3: INORGANIC COMPOUNDS



- Topic hierarchy

2.3A: pH, Buffers, Acids, and Bases

2.3B: Water's Cohesive and Adhesive Properties

2.3C: Water's High Heat Capacity

2.3D: Water's States: Gas, Liquid, and Solid

2.3E: Water's Solvent Properties

This page titled 2.3: Inorganic Compounds is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3A: pH, Buffers, Acids, and Bases

Acids dissociate into H^+ and lower pH, while bases dissociate into OH^- and raise pH; buffers can absorb these excess ions to maintain pH.

Learning Objectives

- Explain the composition of buffer solutions and how they maintain a steady pH

Key Points

- A basic solution will have a pH above 7.0, while an acidic solution will have a pH below 7.0.
- Buffers are solutions that contain a weak acid and its a conjugate base; as such, they can absorb excess H^+ ions or OH^-ions, thereby maintaining an overall steady pH in the solution.
- pH is equal to the negative logarithm of the concentration of H^+ ions in solution: $pH = -\log[H^+]$.

Key Terms

- **alkaline**: having a pH greater than 7; basic
- **acidic**: having a pH less than 7
- **buffer**: a solution composed of a weak acid and its conjugate base that can be used to stabilize the pH of a solution

Self-Ionization of Water

Hydrogen ions are spontaneously generated in pure water by the dissociation (ionization) of a small percentage of water molecules into equal numbers of hydrogen (H^+) ions and hydroxide (OH^-) ions. The hydroxide ions remain in solution because of their hydrogen bonds with other water molecules; the hydrogen ions, consisting of naked protons, are immediately attracted to un-ionized water molecules and form hydronium ions (H_3O^+). By convention, scientists refer to hydrogen ions and their concentration as if they were free in this state in liquid water.

2H2O⇌H3O++OH−2H2O⇌H3O++OH−


The concentration of hydrogen ions dissociating from pure water is 1×10^{-7} moles H^+ ions per liter of water. The pH is calculated as the negative of the base 10 logarithm of this concentration:

$pH = -\log[H^+]$

The negative log of 1×10^{-7} is equal to 7.0, which is also known as neutral pH. Human cells and blood each maintain near-neutral pH.

pH Scale

The pH of a solution indicates its acidity or basicity (alkalinity). The pH scale is an inverse logarithm that ranges from 0 to 14: anything below 7.0 (ranging from 0.0 to 6.9) is acidic, and anything above 7.0 (from 7.1 to 14.0) is basic (or alkaline). Extremes in pH in either direction from 7.0 are usually considered inhospitable to life. The pH in cells (6.8) and the blood (7.4) are both very close to neutral, whereas the environment in the stomach is highly acidic, with a pH of 1 to 2.



The pH scale: The pH scale measures the concentration of hydrogen ions (H^+) in a solution.

Non-neutral pH readings result from dissolving acids or bases in water. Using the negative logarithm to generate positive integers, high concentrations of hydrogen ions yield a low pH, and low concentrations a high pH.

An acid is a substance that increases the concentration of hydrogen ions (H^+) in a solution, usually by dissociating one of its hydrogen atoms. A base provides either hydroxide ions (OH^-) or other negatively-charged ions that react with hydrogen ions in solution, thereby reducing the concentration of H^+ and raising the pH.


Strong Acids and Strong Bases

The stronger the acid, the more readily it donates H^+. For example, hydrochloric acid (HCl) is highly acidic and completely dissociates into hydrogen and chloride ions, whereas the acids in tomato juice or vinegar do not completely dissociate and are considered weak acids; conversely, strong bases readily donate OH^- and/or react with hydrogen ions. Sodium hydroxide (NaOH) and many household cleaners are highly basic and give up OH^- rapidly when placed in water; the OH^- ions react with H^+ in solution, creating new water molecules and lowering the amount of free H^+ in the system, thereby raising the overall pH. An example of a weak basic solution is seawater, which has a pH near 8.0, close enough to neutral that well-adapted marine organisms thrive in this alkaline environment.

Buffers

How can organisms whose bodies require a near-neutral pH ingest acidic and basic substances (a human drinking orange juice, for example) and survive? Buffers are the key. Buffers usually consist of a weak acid and its conjugate base; this enables them to readily absorb excess H^+ or OH^-, keeping the system's pH within a narrow range.

Maintaining a constant blood pH is critical to a person's well-being. The buffer that maintains the pH of human blood involves carbonic acid (H_2CO_3), bicarbonate ion (HCO_3^-), and carbon dioxide (CO_2). When bicarbonate ions combine with free hydrogen ions and become carbonic acid, hydrogen ions are removed, moderating pH changes. Similarly, excess carbonic acid can be converted into carbon dioxide gas and exhaled through the lungs; this prevents too many free hydrogen ions from building up in the blood and dangerously reducing its pH; likewise, if too much OH^- is introduced into the system, carbonic acid will combine with it to create bicarbonate, lowering the pH. Without this buffer system, the body's pH would fluctuate enough to jeopardize survival.



Buffers in the body: This diagram shows the body's buffering of blood pH levels: the blue arrows show the process of raising pH as more CO2 is made; the purple arrows indicate the reverse process, lowering pH as more bicarbonate is created.

Antacids, which combat excess stomach acid, are another example of buffers. Many over-the-counter medications work similarly to blood buffers, often with at least one ion (usually carbonate) capable of absorbing hydrogen and moderating pH, bringing relief to those that suffer "heartburn" from stomach acid after eating.

This page titled 2.3A: pH, Buffers, Acids, and Bases is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3B: Water's Cohesive and Adhesive Properties

Cohesion allows substances to withstand rupture when placed under stress while adhesion is the attraction between water and other molecules.

Learning Objectives

- Describe the cohesive and adhesive properties of water.


Key Points

- Cohesion holds hydrogen bonds together to create surface tension on water.
- Since water is attracted to other molecules, adhesive forces pull the water toward other molecules.
- Water is transported in plants through both cohesive and adhesive forces; these forces pull water and the dissolved minerals from the roots to the leaves and other parts of the plant.

Key Terms

- **adhesion**: The ability of a substance to stick to an unlike substance; attraction between unlike molecules
- **cohesion**: Various intermolecular forces that hold solids and liquids together; attraction between like molecules


Have you ever filled a glass of water to the very top and then slowly added a few more drops? Before it overflows, the water forms a dome-like shape above the rim of the glass. This water can stay above the glass because of the property of cohesion. In cohesion, water molecules are attracted to each other (because of hydrogen bonding), keeping the molecules together at the liquid-gas (water-air) interface, although there is no more room in the glass.



Surface Tension: The weight of the needle is pulling the surface downward; at the same time, the surface tension is pulling it up, suspending it on the surface of the water and keeping it from sinking. Notice the indentation in the water around the needle.


Cohesion allows for the development of surface tension, the capacity of a substance to withstand being ruptured when placed under tension or stress. This is also why water forms droplets when placed on a dry surface rather than being flattened out by gravity. When a small scrap of paper is placed onto the droplet of water, the paper floats on top of the water droplet even though paper is denser (the mass per unit volume) than the water. Cohesion and surface tension keep the hydrogen bonds of water molecules intact and support the item floating on the top. It's even possible to "float" a needle on top of a glass of water if it is placed gently without breaking the surface tension.

These cohesive forces are related to water's property of adhesion, or the attraction between water molecules and other molecules. This attraction is sometimes stronger than water's cohesive forces, especially when the water is exposed to charged surfaces such as those found on the inside of thin glass tubes known as capillary tubes. Adhesion is observed when water "climbs" up the tube placed in a glass of water: notice that the water appears to be higher on the sides of the tube than in the middle. This is because the water molecules are attracted to the charged glass walls of the capillary more than they are to each other and therefore adhere to it. This type of adhesion is called capillary action.



Adhesion: Capillary action in a glass tube is caused by the adhesive forces exerted by the internal surface of the glass exceeding the cohesive forces between the water molecules themselves.

Why are cohesive and adhesive forces important for life? Cohesive and adhesive forces are important for the transport of water from the roots to the leaves in plants. These forces create a "pull" on the water column. This pull results from the tendency of water molecules being evaporated on the surface of the plant to stay connected to water molecules below them, and so they are pulled along. Plants use this natural phenomenon to help transport water from their roots to their leaves. Without these properties of water, plants would be unable to receive the water and the dissolved minerals they require. In another example, insects such as the water strider use the surface tension of water to stay afloat on the surface layer of water and even mate there.



Cohesion & Adhesion: Water's cohesive and adhesive properties allow this water strider (Gerris sp.) to stay afloat.

This page titled 2.3B: Water's Cohesive and Adhesive Properties is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3C: Water's High Heat Capacity

Water is able to absorb a high amount of heat before increasing in temperature, allowing humans to maintain body temperature.

Learning Objectives

- Explain the biological significance of water's high specific heat

Key Points

- Water has the highest heat capacity of all liquids.
- Oceans cool slower than the land due to the high heat capacity of water.
- To change the temperature of 1 gram of water by 1 degree Celsius, it takes 1.00 calorie.

Key Terms

- **heat capacity**: The capability of a substance to absorb heat energy
- **specific heat**: the amount of heat, in calories, needed to raise the temperature of 1 gram of water by 1 degree Celsius

EXAMPLES

The high heat capacity of water has many uses. Commercial nuclear reactors release large amounts of thermal energy (heat) during radioactive decay of fission products. The heat is quickly transferred to a pool of water to cool the reactor. The water then remains hot for a long time due to its high heat capacity.

The capability for a molecule to absorb heat energy is called heat capacity, which can be calculated by the equation shown in the figure. Water's high heat capacity is a property caused by hydrogen bonding among water molecules. When heat is absorbed, hydrogen bonds are broken and water molecules can move freely. When the temperature of water decreases, the hydrogen bonds are formed and release a considerable amount of energy. Water has the highest specific heat capacity of any liquid. Specific heat is defined as the amount of heat one gram of a substance must absorb or lose to change its temperature by one degree Celsius. For water, this amount is one calorie, or 4.184 Joules. As a result, it takes water a long time to heat and a long time to cool. In fact, the specific heat capacity of water is about five times more than that of sand. This explains why the land cools faster than the sea.

$C=Q\Delta TC=Q\Delta T$

The resistance to sudden temperature changes makes water an excellent habitat, allowing organisms to survive without experiencing wide temperature fluctuation. Furthermore, because many organisms are mainly composed of water, the property of high heat capacity allows highly regulated internal body temperatures. For example, the temperature of your body does not drastically drop to the same temperature as the outside temperature while you are skiing or playing in the snow. Due to its high heat capacity, water is used by warm blooded animals to more evenly disperse heat in their bodies; it acts in a similar manner to a car's cooling system, transporting heat from warm places to cool places, causing the body to maintain a more even temperature.

This page titled 2.3C: Water's High Heat Capacity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3D: Water's States: Gas, Liquid, and Solid

The orientation of hydrogen bonds as water changes states dictates the properties of water in its gaseous, liquid, and solid forms.

Learning Objectives

- Explain the biological significance of ice's ability to float on water

Key Points

- As water is boiled, kinetic energy causes the hydrogen bonds to break completely and allows water molecules to escape into the air as gas (steam or water vapor).
- When water freezes, water molecules form a crystalline structure maintained by hydrogen bonding.
- Solid water, or ice, is less dense than liquid water.
- Ice is less dense than water because the orientation of hydrogen bonds causes molecules to push farther apart, which lowers the density.
- For other liquids, solidification when the temperature drops includes the lowering of kinetic energy, which allows molecules to pack more tightly and makes the solid denser than its liquid form.
- Because ice is less dense than water, it is able to float at the surface of water.

Key Terms


- **density**: A measure of the amount of matter contained by a given volume.

The formation of hydrogen bonds is an important quality of liquid water that is crucial to life as we know it. As water molecules make hydrogen bonds with each other, water takes on some unique chemical characteristics compared to other liquids, and since living things have a high water content, understanding these chemical features is key to understanding life. In liquid water, hydrogen bonds are constantly formed and broken as the water molecules slide past each other. The breaking of these bonds is caused by the motion (kinetic energy) of the water molecules due to the heat contained in the system. When the heat is raised as water is boiled, the higher kinetic energy of the water molecules causes the hydrogen bonds to break completely and allows water molecules to escape into the air as gas (steam or water vapor). On the other hand, when the temperature of water is reduced and water freezes, the water molecules form a crystalline structure maintained by hydrogen bonding (there is not enough energy to break the hydrogen bonds). This makes ice less dense than liquid water, a phenomenon not seen in the solidification of other liquids.

Phases of matter: See what happens to intermolecular bonds during phase changes in this interactive.

Water's lower density in its solid form is due to the way hydrogen bonds are oriented as it freezes: the water molecules are pushed farther apart compared to liquid water. With most other liquids, solidification when the temperature drops includes the lowering of kinetic energy between molecules, allowing them to pack even more tightly than in liquid form and giving the solid a greater density than the liquid.

The low density of ice, an anomaly, causes it to float at the surface of liquid water, such as an iceberg or the ice cubes in a glass of water. In lakes and ponds, ice forms on the surface of the water creating an insulating barrier that protects the animals and plant life in the pond from freezing. Without this layer of insulating ice, plants and animals living in the pond would freeze in the solid block of ice and could not survive. The detrimental effect of freezing on living organisms is caused by the expansion of ice relative to liquid water. The ice crystals that form upon freezing rupture the delicate membranes essential for the function of living cells, irreversibly damaging them. Cells can only survive freezing if the water in them is temporarily replaced by another liquid like glycerol.



Ice Density: Hydrogen bonding makes ice less dense than liquid water. The (a) lattice structure of ice makes it less dense than the freely flowing molecules of liquid water, enabling it to (b) float on water.

This page titled 2.3D: Water's States: Gas, Liquid, and Solid is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.3E: Water's Solvent Properties

Water's polarity makes it an excellent solvent for other polar molecules and ions.

Learning Objectives

- Explain why some molecules do not dissolve in water.

Key Points

- Water dissociates salts by separating the cations and anions and forming new interactions between the water and ions.
- Water dissolves many biomolecules, because they are polar and therefore hydrophilic.

Key Terms


- **dissociation**: The process by which a compound or complex body breaks up into simpler constituents such as atoms or ions, usually reversibly.
- **hydration shell**: The term given to a solvation shell (a structure composed of a chemical that acts as a solvent and surrounds a solute species) with a water solvent; also referred to as a hydration sphere.

EXAMPLES

- Sugar, sodium chloride, and hydrophilic proteins are all substances that dissolve in water.
- Oils, fats, and certain organic solvents do not dissolve in water because they are hydrophobic.

Water, which not only dissolves many compounds but also dissolves more substances than any other liquid, is considered the universal solvent. A polar molecule with partially-positive and negative charges, it readily dissolves ions and polar molecules. Water is therefore referred to as a solvent: a substance capable of dissolving other polar molecules and ionic compounds. The charges associated with these molecules form hydrogen bonds with water, surrounding the particle with water molecules. This is referred to as a sphere of hydration, or a hydration shell, and serves to keep the particles separated or dispersed in the water.

When ionic compounds are added to water, individual ions interact with the polar regions of the water molecules during the dissociation process, disrupting their ionic bonds. Dissociation occurs when atoms or groups of atoms break off from molecules and form ions. Consider table salt (NaCl, or sodium chloride): when NaCl crystals are added to water, the molecules of NaCl dissociate into Na^+ and Cl^- ions, and spheres of hydration form around the ions. The positively-charged sodium ion is surrounded by the partially-negative charge of the water molecule's oxygen; the negatively-charged chloride ion is surrounded by the partially-positive charge of the hydrogen in the water molecule.



Dissociation of NaCl in water: When table salt (NaCl) is mixed in water, spheres of hydration form around the ions.

Since many biomolecules are either polar or charged, water readily dissolves these hydrophilic compounds. Water is a poor solvent, however, for hydrophobic molecules such as lipids. Nonpolar molecules experience hydrophobic interactions in water: the water changes its hydrogen bonding patterns around the hydrophobic molecules to produce a cage-like structure called a clathrate. This change in the hydrogen-bonding pattern of the water solvent causes the system's overall entropy to greatly decrease, as the molecules become more ordered than in liquid water. Thermodynamically, such a large decrease in entropy is not spontaneous, and the hydrophobic molecule will not dissolve.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- acidic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acidic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- buffer. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/buffer**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alkaline. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/alkaline**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- cohesion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cohesion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adhesion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adhesion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest/Figure_02_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/**

latest/Figure_02_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_04.jpg. **License**: *CC BY: Attribution*
- heat capacity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/heat_capacity. **License**: *CC BY-SA: Attribution-ShareAlike*
- specific heat. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/specific_heat. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- Structural Biochemistry/Water. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structural_Biochemistry/Water%23High_Heat_Capacity. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_08.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- density. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/density. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_08.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- Structural Biochemistry/Unique Properties/Versatility as a Solvent. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structural_Biochemistry/Unique_Properties/Versatility_as_a_Solvent. **License**: *CC BY-SA: Attribution-ShareAlike*
- Structural Biochemistry/Water. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structural_Biochemistry/Water%23Universal_Solvent. **License**: *CC BY-SA: Attribution-ShareAlike*
- hydration shell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/hydration%20shell. **License**: *CC BY-SA: Attribution-ShareAlike*
- dissociation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/dissociation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clathrate compound. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clathrate_compound. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_08.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_06.jpg. **License**: *CC BY: Attribution*

- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_03.jpg. **License**: *CC BY: Attribution*

This page titled 2.3E: Water's Solvent Properties is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.4: ORGANIC COMPOUNDS

The basis of life is carbon. Carbon's importance comes mainly from the enormous variety of structures that it can form due to its unusual four valence electrons. Most important of these structures is the carbon chain, which forms the "backbone" of fatty acids and carbohydrates, among other organic molecules. Other elements do share properties similar to carbon, in this regard. However, they are not as prevalent on earth as carbon.

2.4A: Organic Molecules and Functional Groups

2.4B: Types of Biological Macromolecules

2.4C: Dehydration Synthesis

This page titled 2.4: Organic Compounds is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.4A: Organic Molecules and Functional Groups

Functional groups are groups of molecules attached to organic molecules and give them specific identities or functions.

Learning Objectives

- Describe the importance of functional groups to organic molecules

Key Points

- Functional groups are collections of atoms that attach the carbon skeleton of an organic molecule and confer specific properties.
- Each type of organic molecule has its own specific type of functional group.
- Functional groups in biological molecules play an important role in the formation of molecules like DNA, proteins, carbohydrates, and lipids.
- Functional groups include: hydroxyl, methyl, carbonyl, carboxyl, amino, phosphate, and sulfhydryl.

Key Terms

- **hydrophobic**: lacking an affinity for water; unable to absorb, or be wetted by water
- **hydrophilic**: having an affinity for water; able to absorb, or be wetted by water

Location of Functional Groups






Functional groups are groups of atoms that occur within organic molecules and confer specific chemical properties to those molecules. When functional groups are shown, the organic molecule is sometimes denoted as “R.” Functional groups are found along the “carbon backbone” of macromolecules which is formed by chains and/or rings of carbon atoms with the occasional substitution of an element such as nitrogen or oxygen. Molecules with other elements in their carbon backbone are substituted hydrocarbons. Each of the four types of macromolecules—proteins, lipids, carbohydrates, and nucleic acids—has its own characteristic set of functional groups that contributes greatly to its differing chemical properties and its function in living organisms.

Properties of Functional Groups

A functional group can participate in specific chemical reactions. Some of the important functional groups in biological molecules include: hydroxyl, methyl, carbonyl, carboxyl, amino, phosphate, and sulfhydryl groups. These groups play an important role in the formation of molecules like DNA, proteins, carbohydrates, and lipids.

Classifying Functional Groups


Functional groups are usually classified as hydrophobic or hydrophilic depending on their charge or polarity. An example of a hydrophobic group is the non-polar methane molecule. Among the hydrophilic functional groups is the carboxyl group found in amino acids, some amino acid side chains, and the fatty acid heads that form triglycerides and phospholipids. This carboxyl group ionizes to release hydrogen ions (H^+) from the COOH group resulting in the negatively charged COO^- group; this contributes to the hydrophilic nature of whatever molecule it is found on. Other functional groups, such as the carbonyl group, have a partially negatively charged oxygen atom that may form hydrogen bonds with water molecules, again making the molecule more hydrophilic. The functional groups shown here are found in many different biological molecules, where “R” is the organic molecule.

Important Functional Groups in Biology		
Functional Group	**Structure**	**Properties**
Hydroxyl		Polar
Methyl		Nonpolar
Carbonyl		Polar
Carboxyl		Charged, ionized to release H^+. Since carboxyl groups can release H^+ ions into a solution, they are considered acidic.
Amino		Charged, accepts H^+ to form NH_3^+. Since amino groups can remove H^+ from solution, they are considered basic.
Phosphate		Charged, ionizes to release H^+. Since phosphate groups can release H^+ ions into solution, they are considered acidic.
Sulfhydryl		Polar

Hydrogen Bonds between Functional Groups

Hydrogen bonds between functional groups (within the same molecule or between different molecules) are important to the function of many macromolecules and help them to fold properly and maintain the appropriate shape needed to function correctly. Hydrogen bonds are also involved in various recognition processes, such as DNA complementary base pairing and the binding of

an enzyme to its substrate.



Hydrogen bonds in DNA: Hydrogen bonds connect two strands of DNA together to create the double-helix structure.

This page titled 2.4A: Organic Molecules and Functional Groups is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

2.4B: Types of Biological Macromolecules

Biological macromolecules, the large molecules necessary for life, include carbohydrates, lipids, nucleic acids, and proteins.

Learning Objectives

- Identify the four major classes of biological macromolecules

Key Points

- Biological macromolecules are important cellular components and perform a wide array of functions necessary for the survival and growth of living organisms.
- The four major classes of biological macromolecules are carbohydrates, lipids, proteins, and nucleic acids.

Key Terms


- **polymer**: A relatively large molecule consisting of a chain or network of many identical or similar monomers chemically bonded to each other.
- **monomer**: A relatively small molecule that can form covalent bonds with other molecules of this type to form a polymer.

Nutrients are the molecules that living organisms require for survival and growth but that animals and plants cannot synthesize themselves. Animals obtain nutrients by consuming food, while plants pull nutrients from soil.



Sources of biological macromolecules: Foods such as bread, fruit, and cheese are rich sources of biological macromolecules.

Many critical nutrients are biological macromolecules. The term “macromolecule” was first coined in the 1920s by Nobel laureate Hermann Staudinger. Staudinger was the first to propose that many large biological molecules are built by covalently linking smaller biological molecules together.



Living organisms are made up of chemical building blocks: All organisms are composed of a variety of these biological macromolecules.


Monomers and Polymers

Biological macromolecules play a critical role in cell structure and function. Most (but not all) biological macromolecules are polymers, which are any molecules constructed by linking together many smaller molecules, called monomers. Typically all the monomers in a polymer tend to be the same, or at least very similar to each other, linked over and over again to build up the larger macromolecule. These simple monomers can be linked in many different combinations to produce complex biological polymers, just as a few types of Lego blocks can build anything from a house to a car.

image

Monomers and polymers: Many small monomer subunits combine to form this carbohydrate polymer.

Examples of these monomers and polymers can be found in the sugar you might put in your coffee or tea. Regular table sugar is the disaccharide sucrose (a polymer), which is composed of the monosaccharides fructose and glucose (which are monomers). If we were to string many carbohydrate monomers together we could make a polysaccharide like starch. The prefixes “mono-” (one),”di-” (two),and “poly-” (many) will tell you how many of the monomers have been joined together in a molecule.



The molecule sucrose (common table sugar): The carbohydrate monosaccharides (fructose and glucose) are joined to make the disaccharide sucrose.

Biological macromolecules all contain carbon in ring or chain form, which means they are classified as organic molecules. They usually also contain hydrogen and oxygen, as well as nitrogen and additional minor elements.

Four Classes of Biological Macromolecules

There are four major classes of biological macromolecules:

1. carbohydrates
2. lipids
3. proteins
4. nucleic acids

Each of these types of macromolecules performs a wide array of important functions within the cell; a cell cannot perform its role within the body without many different types of these crucial molecules. In combination, these biological macromolecules make up the majority of a cell's dry mass. (Water molecules make up the majority of a cell's total mass.) All the molecules both inside and outside of cells are situated in a water-based (i.e., aqueous) environment, and all the reactions of biological systems are occurring in that same environment.

Interactive: Monomers and Polymers: Carbohydrates, proteins, and nucleic acids are built from small molecular units that are connected to each other by strong covalent bonds. The small molecular units are called monomers (mono means one, or single), and they are linked together into long chains called polymers (poly means many, or multiple). Each different type of macromolecule, except lipids, is built from a different set of monomers that resemble each other in composition and size. Lipids are not polymers, because they are not built from monomers (units with similar composition).

This page titled 2.4B: Types of Biological Macromolecules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


2.4C: Dehydration Synthesis

Learning Objectives

- Explain dehydration (or condensation) reactions

Most macromolecules are made from single subunits, or building blocks, called monomers. The monomers combine with each other via covalent bonds to form larger molecules known as polymers. In doing so, monomers release water molecules as byproducts. This type of reaction is known as dehydration synthesis, which means “to put together while losing water.” It is also considered to be a condensation reaction since two molecules are condensed into one larger molecule with the loss of a smaller molecule (the water.)

In a dehydration synthesis reaction between two un-ionized monomers, such as monosaccharide sugars, the hydrogen of one monomer combines with the hydroxyl group of another monomer, releasing a molecule of water in the process. The removal of a hydrogen from one monomer and the removal of a hydroxyl group from the other monomer allows the monomers to share electrons and form a covalent bond. Thus, the monomers that are joined together are being *dehydrated* to allow for *synthesis* of a larger molecule.



A dehydration synthesis reaction involving un-ionized moners..: In the dehydration synthesis reaction between two molecules of glucose, a hydroxyl group from the first glucose is combined with a hydrogen from the second glucose, creating a covalent bond that links the two monomeric sugars (monosaccharides) together to form the dissacharide maltose. In the process, a water molecule is formed.

When the monomers are ionized, such as is the case with amino acids in an aqueous environment like cytoplasm, two hydrogens from the positively-charged end of one monomer are combined with an oxygen from the negatively-charged end of another monomer, again forming water, which is released as a side-product, and again joining the two monomers with a covalent bond.

image

A dehydration synthesis reaction involving ionized monomers.: In the dehydration synthesis reaction between two amino acids, with are ionized in aqueous environments like the cell, an oxygen from the first amino acid is combined with two hydrogens from the second amino acid, creating a covalent bond that links the two monomers together to form a dipeptide. In the process a water molecule is formed.

As additional monomers join via multiple dehydration synthesis reactions, the chain of repeating monomers begins to form a polymer. Different types of monomers can combine in many configurations, giving rise to a diverse group of macromolecules. Three of the four major classes of biological macromolecules (complex carbohydrates, nucleic acids, and proteins), are composed of monomers that join together via dehydration synthesis reactions. Complex carbohydrates are formed from monosaccharides, nucleic acids are formed from mononucleotides, and proteins are formed from amino acids.

There is great diversity in the manner by which monomers can combine to form polymers. For example, glucose monomers are the constituents of starch, glycogen, and cellulose. These three are polysaccharides, classified as carbohydrates, that have formed as a result of multiple dehydration synthesis reactions between glucose monomers. However, the manner by which glucose monomers join together, specifically locations of the covalent bonds between connected monomers and the orientation

(stereochemistry) of the covalent bonds, results in these three different polysaccharides with varying properties and functions. In nucleic acids and proteins, the location and stereochemistry of the covalent linkages connecting the monomers do not vary from molecule to molecule, but instead the multiple kinds of monomers (five different monomers in nucleic acids, A, G, C, T, and U mononucleotides; 21 different amino acids monomers in proteins) are combined in a huge variety of sequences. Each protein or nucleic acid with a different sequence is a different molecule with different properties.

Key Points

- During dehydration synthesis, either the hydrogen of one monomer combines with the hydroxyl group of another monomer releasing a molecule of water, or two hydrogens from one monomer combine with one oxygen from the other monomer releasing a molecule of water.
- The monomers that are joined via dehydration synthesis reactions share electrons and form covalent bonds with each other.
- As additional monomers join via multiple dehydration synthesis reactions, this chain of repeating monomers begins to form a polymer.
- Complex carbohydrates, nucleic acids, and proteins are all examples of polymers that are formed by dehydration synthesis.
- Monomers like glucose can join together in different ways and produce a variety of polymers. Monomers like mononucleotides and amino acids join together in different sequences to produce a variety of polymers.

Key Terms

- **covalent bond**: A type of chemical bond where two atoms are connected to each other by the sharing of two or more electrons.
- **monomer**: A relatively small molecule which can be covalently bonded to other monomers to form a polymer.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- hydrophilic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hydrophilic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hydrophobic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hydrophobic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Functional Group. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Functional_group. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_08.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_07.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44395/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- OpenStax College, Synthesis of Biological Macromolecules. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44397/latest/. **License**: ***CC BY: Attribution***
- An Introduction to Molecular Biology/Macromolecules and Cells. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/An_Intr...ules_and_Cells. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Free High School Science Texts Project, Organic Macromolecules: Biological Macromolecules. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m39433/latest/. **License**: ***CC BY: Attribution***
- monomer. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/monomer. **License**: ***CC BY-SA: Attribution-ShareAlike***
- polymer. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/polymer. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_08.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_07.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44395/latest...e_03_00_01.jpg. **License**: ***CC BY: Attribution***
- Sucrose-inkscape.svg.png. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/F...e-inkscape.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Building_blocks_of_life.png. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped...ks_of_life.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44397/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- monomer. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/monomer. **License**: ***CC BY-SA: Attribution-ShareAlike***
- covalent bond. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/covalent_bond. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_08.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Carbon. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44393/latest...e_02_03_07.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44395/latest...e_03_00_01.jpg. **License**: ***CC BY: Attribution***
- Sucrose-inkscape.svg.png. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/F...e-inkscape.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Building_blocks_of_life.png. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped...ks_of_life.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Synthesis of Biological Macromolecules. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44397/latest...e_03_01_01.jpg. **License**: ***CC BY: Attribution***

This page titled 2.4C: Dehydration Synthesis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3: ORGANIZATION AT THE CELLULAR LEVEL

3.1: Introduction to the Study of Cells

3.1A: Cells as the Basic Unit of Life

3.2: Cell Membranes and the Fluid Mosaic Model

3.2A: Components of Plasma Membranes

3.2B: Fluid Mosaic Model

3.2C: Membrane Fluidity

3.3: Transport Across Membranes

3.3A: Diffusion

3.3B: Osmosis

3.3C: Tonicity

3.3D: Facilitated transport

3.3E: The Role of Passive Transport

3.3F: Primary Active Transport

3.3G: Electrochemical Gradient

3.3H: Secondary Active Transport

3.3I: Endocytosis

3.4: Cell Signaling

3.4A: Types of Receptors

3.5: The Nucleus and Ribosomes

3.5A: The Nucleus and Ribosomes

3.5B: Characteristics of Eukaryotic Cells

3.6: Organelles

3.6A: The Plasma Membrane and the Cytoplasm

3.6B: The Endoplasmic Reticulum

3.6C: The Golgi Apparatus

3.6D: Lysosomes

3.6E: Peroxisomes

3.6F: Mitochondria

3.7: The Cytoskeleton

3.7A: Intermediate Filaments and Microtubules

3.7B: Microfilaments

3.8: External Cellular Components

3.8A: Extracellular Matrix of Animal Cells

3.8B: Intercellular Junctions

This page titled 3: Organization at the Cellular Level is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.1: INTRODUCTION TO THE STUDY OF CELLS



- Topic hierarchy

3.1A: Cells as the Basic Unit of Life

This page titled 3.1: Introduction to the Study of Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.1A: Cells as the Basic Unit of Life

A cell is the smallest unit of a living thing and is the basic building block of all organisms.

Learning Objectives

- State the general characteristics of a cell

Key Points

- A living thing can be composed of either one cell or many cells.
- There are two broad categories of cells: prokaryotic and eukaryotic cells.
- Cells can be highly specialized with specific functions and characteristics.

Key Terms

- **prokaryotic**: Small cells in the domains Bacteria and Archaea that do not contain a membrane-bound nucleus or other membrane-bound organelles.
- **eukaryotic**: Having complex cells in which the genetic material is contained within membrane-bound nuclei.
- **cell**: The basic unit of a living organism, consisting of a quantity of protoplasm surrounded by a cell membrane, which is able to synthesize proteins and replicate itself.


Close your eyes and picture a brick wall. What is the basic building block of that wall? A single brick, of course. Like a brick wall, your body is composed of basic building blocks, and the building blocks of your body are cells.

Cells as Building Blocks

A cell is the smallest unit of a living thing. A living thing, whether made of one cell (like bacteria) or many cells (like a human), is called an organism. Thus, cells are the basic building blocks of all organisms. Several cells of one kind that interconnect with each other and perform a shared function form tissues; several tissues combine to form an organ (your stomach, heart, or brain); and several organs make up an organ system (such as the digestive system, circulatory system, or nervous system). Several systems that function together form an organism (like a human being). There are many types of cells all grouped into one of two broad categories: prokaryotic and eukaryotic. For example, both animal and plant cells are classified as eukaryotic cells, whereas bacterial cells are classified as prokaryotic.

Types of Specialized Cells

Your body has many kinds of cells, each specialized for a specific purpose. Just as a home is made from a variety of building materials, the human body is constructed from many cell types. For example, epithelial cells protect the surface of the body and cover the organs and body cavities within. Bone cells help to support and protect the body. Cells of the immune system fight invading bacteria. Additionally, blood and blood cells carry nutrients and oxygen throughout the body while removing carbon dioxide. Each of these cell types plays a vital role during the growth, development, and day-to-day maintenance of the body. In spite of their enormous variety, however, cells from all organisms—even ones as diverse as bacteria, onion, and human—share certain fundamental characteristics.



Various Cell Types: (a) Nasal sinus cells (viewed with a light microscope), (b) onion cells (viewed with a light microscope), and (c) Vibrio tasmaniensis bacterial cells (seen through a scanning electron microscope) are from very different organisms, yet all share certain characteristics of basic cell structure.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44404/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Robert Bear and David Rintoul, Introduction to Cells. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m47170/latest/**. **License**: ***CC BY: Attribution***
- eukaryotic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/eukaryotic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cell. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- prokaryotic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/prokaryotic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44404/latest...e_04_00_00.jpg**. **License**: ***CC BY: Attribution***

This page titled 3.1A: Cells as the Basic Unit of Life is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.2: CELL MEMBRANES AND THE FLUID MOSAIC MODEL

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

3.2A: Components of Plasma Membranes

3.2B: Fluid Mosaic Model

3.2C: Membrane Fluidity

This page titled 3.2: Cell Membranes and the Fluid Mosaic Model is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.2A: Components of Plasma Membranes

The plasma membrane protects the cell from its external environment, mediates cellular transport, and transmits cellular signals.

Learning Objectives

- Describe the function and components of the plasma membrane

Key Points

- The principal components of the plasma membrane are lipids (phospholipids and cholesterol), proteins, and carbohydrates.
- The plasma membrane protects intracellular components from the extracellular environment.
- The plasma membrane mediates cellular processes by regulating the materials that enter and exit the cell.
- The plasma membrane carries markers that allow cells to recognize one another and can transmit signals to other cells via receptors.

Key Terms

- **plasma membrane**: The semipermeable barrier that surrounds the cytoplasm of a cell.
- **receptor**: A protein on a cell wall that binds with specific molecules so that they can be absorbed into the cell.


Structure of Plasma Membranes

The plasma membrane (also known as the cell membrane or cytoplasmic membrane) is a biological membrane that separates the interior of a cell from its outside environment.

The primary function of the plasma membrane is to protect the cell from its surroundings. Composed of a phospholipid bilayer with embedded proteins, the plasma membrane is selectively permeable to ions and organic molecules and regulates the movement of substances in and out of cells. Plasma membranes must be very flexible in order to allow certain cells, such as red blood cells and white blood cells, to change shape as they pass through narrow capillaries.

The plasma membrane also plays a role in anchoring the cytoskeleton to provide shape to the cell, and in attaching to the extracellular matrix and other cells to help group cells together to form tissues. The membrane also maintains the cell potential.

In short, if the cell is represented by a castle, the plasma membrane is the wall that provides structure for the buildings inside the wall, regulates which people leave and enter the castle, and conveys messages to and from neighboring castles. Just as a hole in the wall can be a disaster for the castle, a rupture in the plasma membrane causes the cell to lyse and die.



The plasma membrane: The plasma membrane is composed of phospholipids and proteins that provide a barrier between the external environment and the cell, regulate the transportation of molecules across the membrane, and communicate with other cells via protein receptors.

The Plasma Membrane and Cellular Transport

The movement of a substance across the selectively permeable plasma membrane can be either “passive”—i.e., occurring without the input of cellular energy —or “active”—i.e., its transport requires the cell to expend energy.

The cell employs a number of transport mechanisms that involve biological membranes:

1. Passive osmosis and diffusion: transports gases (such as O_2 and CO_2) and other small molecules and ions
2. Transmembrane protein channels and transporters: transports small organic molecules such as sugars or amino acids
3. Endocytosis: transports large molecules (or even whole cells) by engulfing them
4. Exocytosis: removes or secretes substances such as hormones or enzymes

The Plasma Membrane and Cellular Signaling

Among the most sophisticated functions of the plasma membrane is its ability to transmit signals via complex proteins. These proteins can be receptors, which work as receivers of extracellular inputs and as activators of intracellular processes, or markers, which allow cells to recognize each other.

Membrane receptors provide extracellular attachment sites for effectors like hormones and growth factors, which then trigger intracellular responses. Some viruses, such as Human Immunodeficiency Virus (HIV), can hijack these receptors to gain entry into the cells, causing infections.

Membrane markers allow cells to recognize one another, which is vital for cellular signaling processes that influence tissue and organ formation during early development. This marking function also plays a later role in the “self”-versus-“non-self” distinction of the immune response. Marker proteins on human red blood cells, for example, determine blood type (A, B, AB, or O).

This page titled 3.2A: Components of Plasma Membranes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.2B: Fluid Mosaic Model

The fluid mosaic model describes the plasma membrane structure as a mosaic of phospholipids, cholesterol, proteins, and carbohydrates.

Learning Objectives

- Describe the fluid mosaic model of cell membranes

Key Points

- The main fabric of the membrane is composed of amphiphilic or dual-loving, phospholipid molecules.
- Integral proteins, the second major component of plasma membranes, are integrated completely into the membrane structure with their hydrophobic membrane-spanning regions interacting with the hydrophobic region of the phospholipid bilayer.
- Carbohydrates, the third major component of plasma membranes, are always found on the exterior surface of cells where they are bound either to proteins (forming glycoproteins) or to lipids (forming glycolipids).

Key Terms


- **amphiphilic**: Having one surface consisting of hydrophilic amino acids and the opposite surface consisting of hydrophobic (or lipophilic) ones.
- **hydrophilic**: Having an affinity for water; able to absorb, or be wetted by water,"water-loving."
- **hydrophobic**: Lacking an affinity for water; unable to absorb, or be wetted by water,"water-fearing."

The fluid mosaic model was first proposed by S.J. Singer and Garth L. Nicolson in 1972 to explain the structure of the plasma membrane. The model has evolved somewhat over time, but it still best accounts for the structure and functions of the plasma membrane as we now understand them. The fluid mosaic model describes the structure of the plasma membrane as a mosaic of components —including phospholipids, cholesterol, proteins, and carbohydrates—that gives the membrane a fluid character. Plasma membranes range from 5 to 10 nm in thickness. For comparison, human red blood cells, visible via light microscopy, are approximately 8 µm wide, or approximately 1,000 times wider than a plasma membrane. The proportions of proteins, lipids, and carbohydrates in the plasma membrane vary with cell type. For example, myelin contains 18% protein and 76% lipid. The mitochondrial inner membrane contains 76% protein and 24% lipid.

Components of the Plasma Membrane


Component	Location
Phospholipid	Main fabric of the membrane
Cholesterol	Attached between phospholipids and between the two phospholipid layers
Integral proteins (for example, integrins)	Embedded within the phospholipid layer(s). May or may not penetrate through both layers.
Peripheral proteins	On the inner or outer surface of the phospholipid bilayer; not embedded within the phospholipids
Carbohydrates (components of glycoproteins and glycolipids)	Generally attached to outside of membrane layer

The principal components of a plasma membrane are lipids (phospholipids and cholesterol), proteins, and carbohydrates attached to some of the lipids and some of the proteins.




The fluid mosaic model of the plasma membrane: The fluid mosaic model of the plasma membrane describes the plasma membrane as a fluid combination of phospholipids, cholesterol, and proteins. Carbohydrates attached to lipids (glycolipids) and to proteins (glycoproteins) extend from the outward-facing surface of the membrane.

The main fabric of the membrane is composed of amphiphilic or dual-loving, phospholipid molecules. The hydrophilic or water-loving areas of these molecules are in contact with the aqueous fluid both inside and outside the cell. Hydrophobic, or water-hating molecules, tend to be non- polar. A phospholipid molecule consists of a three-carbon glycerol backbone with two fatty acid molecules attached to carbons 1 and 2, and a phosphate-containing group attached to the third carbon. This arrangement gives the overall molecule an area described as its head (the phosphate-containing group), which has a polar character or negative charge, and an area called the tail (the fatty acids), which has no charge. They interact with other non-polar molecules in chemical reactions, but generally do not interact with polar molecules. When placed in water, hydrophobic molecules tend to form a ball or cluster. The hydrophilic regions of the phospholipids tend to form hydrogen bonds with water and other polar molecules on both the exterior and interior of the cell. Thus, the membrane surfaces that face the interior and exterior of the cell are hydrophilic. In contrast, the middle of the cell membrane is hydrophobic and will not interact with water. Therefore, phospholipids form an excellent lipid bilayer cell membrane that separates fluid within the cell from the fluid outside of the cell.




Phospholipid aggregation: In an aqueous solution, phospholipids tend to arrange themselves with their polar heads facing outward and their hydrophobic tails facing inward.



The structure of a phospholipid molecule: This phospholipid molecule is composed of a hydrophilic head and two hydrophobic tails. The hydrophilic head group consists of a phosphate-containing group attached to a glycerol molecule. The hydrophobic tails, each containing either a saturated or an unsaturated fatty acid, are long hydrocarbon chains.

Proteins make up the second major component of plasma membranes. Integral proteins (some specialized types are called integrins) are, as their name suggests, integrated completely into the membrane structure, and their hydrophobic membrane-spanning regions interact with the hydrophobic region of the the phospholipid bilayer. Single-pass integral membrane proteins usually have a hydrophobic transmembrane segment that consists of 20–25 amino acids. Some span only part of the membrane—associating with a single layer—while others stretch from one side of the membrane to the other, and are exposed on either side. Some complex proteins are composed of up to 12 segments of a single protein, which are extensively folded and embedded in the membrane. This type of protein has a hydrophilic region or regions, and one or several mildly hydrophobic regions. This arrangement of regions of the protein tends to orient the protein alongside the phospholipids, with the hydrophobic region of the protein adjacent to the tails of the phospholipids and the hydrophilic region or regions of the protein protruding from the membrane and in contact with the cytosol or extracellular fluid.



Structure of integral membrane proteins: Integral membrane proteins may have one or more alpha-helices that span the membrane (examples 1 and 2), or they may have beta-sheets that span the membrane (example 3).

Carbohydrates are the third major component of plasma membranes. They are always found on the exterior surface of cells and

are bound either to proteins (forming glycoproteins) or to lipids (forming glycolipids). These carbohydrate chains may consist of 2–60 monosaccharide units and can be either straight or branched. Along with peripheral proteins, carbohydrates form specialized sites on the cell surface that allow cells to recognize each other. This recognition function is very important to cells, as it allows the immune system to differentiate between body cells (called “self”) and foreign cells or tissues (called “non-self”). Similar types of glycoproteins and glycolipids are found on the surfaces of viruses and may change frequently, preventing immune cells from recognizing and attacking them. These carbohydrates on the exterior surface of the cell—the carbohydrate components of both glycoproteins and glycolipids—are collectively referred to as the glycocalyx (meaning “sugar coating”). The glycocalyx is highly hydrophilic and attracts large amounts of water to the surface of the cell. This aids in the interaction of the cell with its watery environment and in the cell’s ability to obtain substances dissolved in the water.

This page titled 3.2B: Fluid Mosaic Model is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.2C: Membrane Fluidity

The mosaic nature of the membrane, its phospholipid chemistry, and the presence of cholesterol contribute to membrane fluidity.

Learning Objectives

- Explain the function of membrane fluidity in the structure of cells

Key Points


- The membrane is fluid but also fairly rigid and can burst if penetrated or if a cell takes in too much water.
- The mosaic nature of the plasma membrane allows a very fine needle to easily penetrate it without causing it to burst and allows it to self-seal when the needle is extracted.
- If saturated fatty acids are compressed by decreasing temperatures, they press in on each other, making a dense and fairly rigid membrane.
- If unsaturated fatty acids are compressed, the “kinks” in their tails push adjacent phospholipid molecules away, which helps maintain fluidity in the membrane.
- The ratio of saturated and unsaturated fatty acids determines the fluidity in the membrane at cold temperatures.
- Cholesterol functions as a buffer, preventing lower temperatures from inhibiting fluidity and preventing higher temperatures from increasing fluidity.

Key Terms

- **phospholipid**: Any lipid consisting of a diglyceride combined with a phosphate group and a simple organic molecule such as choline or ethanolamine; they are important constituents of biological membranes
- **fluidity**: A measure of the extent to which something is fluid. The reciprocal of its viscosity.

Membrane Fluidity

There are multiple factors that lead to membrane fluidity. First, the mosaic characteristic of the membrane helps the plasma membrane remain fluid. The integral proteins and lipids exist in the membrane as separate but loosely-attached molecules. The membrane is not like a balloon that can expand and contract; rather, it is fairly rigid and can burst if penetrated or if a cell takes in too much water. However, because of its mosaic nature, a very fine needle can easily penetrate a plasma membrane without causing it to burst; the membrane will flow and self-seal when the needle is extracted.



Membrane Fluidity: The plasma membrane is a fluid combination of phospholipids, cholesterol, and proteins. Carbohydrates attached to lipids (glycolipids) and to proteins (glycoproteins) extend from the outward-facing surface of the membrane.

The second factor that leads to fluidity is the nature of the phospholipids themselves. In their saturated form, the fatty acids in phospholipid tails are saturated with bound hydrogen atoms; there are no double bonds between adjacent carbon atoms. This results in tails that are relatively straight. In contrast, unsaturated fatty acids do not contain a maximal number of hydrogen atoms, although they do contain some double bonds between adjacent carbon atoms; a double bond results in a bend of approximately 30 degrees in the string of carbons. Thus, if saturated fatty acids, with their straight tails, are compressed by decreasing temperatures, they press in on each other, making a dense and fairly rigid membrane. If unsaturated fatty acids are compressed, the “kinks” in their tails elbow adjacent phospholipid molecules away, maintaining some space between the phospholipid molecules. This “elbow room” helps to maintain fluidity in the membrane at temperatures at which membranes with saturated fatty acid tails in their phospholipids would “freeze” or solidify. The relative fluidity of the membrane is particularly important in a cold environment. A cold environment tends to compress membranes composed largely of saturated fatty acids, making them less fluid and more susceptible to rupturing. Many organisms (fish are one example) are capable of adapting to cold environments by changing the proportion of unsaturated fatty acids in their membranes in response to the lowering of the temperature.

In animals, the third factor that keeps the membrane fluid is cholesterol. It lies alongside the phospholipids in the membrane and tends to dampen the effects of temperature on the membrane. Thus, cholesterol functions as a buffer, preventing lower temperatures from inhibiting fluidity and preventing higher temperatures from increasing fluidity too much. Cholesterol extends in both directions the range of temperature in which the membrane is appropriately fluid and, consequently, functional. Cholesterol also serves other functions, such as organizing clusters of transmembrane proteins into lipid rafts.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Components and Structure. October 28, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/#tab-ch05_01_01. **License**: ***CC BY: Attribution***
- OpenStax College, Components and Structure. October 28, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/#tab-ch05_01_01. **License**: ***CC BY: Attribution***
- OpenStax College, Components and Structure. October 28, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/#tab-ch05_01_01. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44415/

latest...ol11448/latest. **License**: *CC BY: Attribution*
- receptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- plasma membrane. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/plasma_membrane. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illustration of au00a0Eukaryoticu00a0cell membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cell_me..._diagram_4.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...ol11448/latest. **License**: *CC BY: Attribution*
- hydrophobic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hydrophobic. **License**: *CC BY-SA: Attribution-ShareAlike*
- amphiphilic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amphiphilic. **License**: *CC BY-SA: Attribution-ShareAlike*
- hydrophilic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hydrophilic. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illustration of au00a0Eukaryoticu00a0cell membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cell_me..._diagram_4.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...e_05_01_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 29, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...e_05_01_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...e_05_01_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/Figure_05_01_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...ol11448/latest. **License**: *CC BY: Attribution*
- fluidity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fluidity. **License**: *CC BY-SA: Attribution-ShareAlike*
- phospholipid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/phospholipid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illustration of au00a0Eukaryoticu00a0cell membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cell_me..._diagram_4.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...e_05_01_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 29, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest...e_05_01_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/Figure_05_01_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/Figure_05_01_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Components and Structure. October 28, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44416/latest/#fig-ch05_01_03. **License**: *CC BY: Attribution*

This page titled 3.2C: Membrane Fluidity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3: TRANSPORT ACROSS MEMBRANES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

3.3A: Diffusion

3.3B: Osmosis

3.3C: Tonicity

3.3D: Facilitated transport

3.3E: The Role of Passive Transport

3.3F: Primary Active Transport

3.3G: Electrochemical Gradient

3.3H: Secondary Active Transport

3.3I: Endocytosis

This page titled 3.3: Transport Across Membranes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3A: Diffusion

Diffusion is a process of passive transport in which molecules move from an area of higher concentration to one of lower concentration.

Learning Objectives

- Describe diffusion and the factors that affect how materials move across the cell membrane.

Key Points

- Substances diffuse according to their concentration gradient; within a system, different substances in the medium will each diffuse at different rates according to their individual gradients.
- After a substance has diffused completely through a space, removing its concentration gradient, molecules will still move around in the space, but there will be no net movement of the number of molecules from one area to another, a state known as dynamic equilibrium.
- Several factors affect the rate of diffusion of a solute including the mass of the solute, the temperature of the environment, the solvent density, and the distance traveled.


Key Terms

- **diffusion**: The passive movement of a solute across a permeable membrane
- **concentration gradient**: A concentration gradient is present when a membrane separates two different concentrations of molecules.

EXAMPLES

When someone is cooking food in a kitchen, the smell begins to waft through the house, and eventually everyone can tell what's for dinner! This is due to the diffusion of odor molecules through the air, from an area of high concentration (the kitchen) to areas of low concentration (your upstairs bedroom).

Diffusion is a passive process of transport. A single substance tends to move from an area of high concentration to an area of low concentration until the concentration is equal across a space. You are familiar with diffusion of substances through the air. For example, think about someone opening a bottle of ammonia in a room filled with people. The ammonia gas is at its highest concentration in the bottle; its lowest concentration is at the edges of the room. The ammonia vapor will diffuse, or spread away, from the bottle; gradually, more and more people will smell the ammonia as it spreads. Materials move within the cell 's cytosol by diffusion, and certain materials move through the plasma membrane by diffusion. Diffusion expends no energy. On the contrary, concentration gradients are a form of potential energy, dissipated as the gradient is eliminated.



Diffusion: Diffusion through a permeable membrane moves a substance from an area of high concentration (extracellular fluid, in this case) down its concentration gradient (into the cytoplasm).

Each separate substance in a medium, such as the extracellular fluid, has its own concentration gradient independent of the concentration gradients of other materials. In addition, each substance will diffuse according to that gradient. Within a system, there will be different rates of diffusion of the different substances in the medium.

Factors That Affect Diffusion

Molecules move constantly in a random manner at a rate that depends on their mass, their environment, and the amount of thermal energy they possess, which in turn is a function of temperature. This movement accounts for the diffusion of molecules through whatever medium in which they are localized. A substance will tend to move into any space available to it until it is evenly distributed throughout it. After a substance has diffused completely through a space removing its concentration gradient, molecules will still move around in the space, but there will be no net movement of the number of molecules from one area to another. This lack of a concentration gradient in which there is no net movement of a substance is known as dynamic equilibrium. While diffusion will go forward in the presence of a concentration gradient of a substance, several factors affect the rate of diffusion:

- Extent of the concentration gradient: The greater the difference in concentration, the more rapid the diffusion. The closer the distribution of the material gets to equilibrium, the slower the rate of diffusion becomes.
- Mass of the molecules diffusing: Heavier molecules move more slowly; therefore, they diffuse more slowly. The reverse is true for lighter molecules.
- Temperature: Higher temperatures increase the energy and therefore the movement of the molecules, increasing the rate of diffusion. Lower temperatures decrease the energy of the molecules, thus decreasing the rate of diffusion.
- Solvent density: As the density of a solvent increases, the rate of diffusion decreases. The molecules slow down because they have a more difficult time getting through the denser medium. If the medium is less dense, diffusion increases. Because cells primarily use diffusion to move materials within the cytoplasm, any increase in the cytoplasm's density will inhibit the movement of the materials. An example of this is a person experiencing dehydration. As the body's cells lose water, the rate of diffusion decreases in the cytoplasm, and the cells' functions deteriorate. Neurons tend to be very sensitive to this effect. Dehydration frequently leads to unconsciousness and possibly coma because of the decrease in diffusion rate within the cells.
- Solubility: As discussed earlier, nonpolar or lipid-soluble materials pass through plasma membranes more easily than polar materials, allowing a faster rate of diffusion.
- Surface area and thickness of the plasma membrane: Increased surface area increases the rate of diffusion, whereas a thicker membrane reduces it.
- Distance travelled: The greater the distance that a substance must travel, the slower the rate of diffusion. This places an upper limitation on cell size. A large, spherical cell will die because nutrients or waste cannot reach or leave the center of the cell. Therefore, cells must either be small in size, as in the case of many prokaryotes, or be flattened, as with many single-

celled eukaryotes.

A variation of diffusion is the process of filtration. In filtration, material moves according to its concentration gradient through a membrane; sometimes the rate of diffusion is enhanced by pressure, causing the substances to filter more rapidly. This occurs in the kidney where blood pressure forces large amounts of water and accompanying dissolved substances, or solutes, out of the blood and into the renal tubules. The rate of diffusion in this instance is almost totally dependent on pressure. One of the effects of high blood pressure is the appearance of protein in the urine, which is “squeezed through” by the abnormally high pressure.

This page titled 3.3A: Diffusion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3B: Osmosis

Osmosis is the movement of water across a membrane from an area of low solute concentration to an area of high solute concentration.

Learning Objectives

- Describe the process of osmosis and explain how concentration gradient affects osmosis

Key Points

- Osmosis occurs according to the concentration gradient of water across the membrane, which is inversely proportional to the concentration of solutes.
- Osmosis occurs until the concentration gradient of water goes to zero or until the hydrostatic pressure of the water balances the osmotic pressure.
- Osmosis occurs when there is a concentration gradient of a solute within a solution, but the membrane does not allow diffusion of the solute.

Key Terms

- **solute**: Any substance that is dissolved in a liquid solvent to create a solution
- **osmosis**: The net movement of solvent molecules from a region of high solvent potential to a region of lower solvent potential through a partially permeable membrane
- **semipermeable membrane**: A type of biological membrane that will allow certain molecules or ions to pass through it by diffusion and occasionally by specialized facilitated diffusion

Osmosis and Semipermeable Membranes


Osmosis is the movement of water through a semipermeable membrane according to the concentration gradient of water across the membrane, which is inversely proportional to the concentration of solutes. Semipermeable membranes, also termed selectively permeable membranes or partially permeable membranes, allow certain molecules or ions to pass through by diffusion.

While diffusion transports materials across membranes and within cells, osmosis transports only water across a membrane. The semipermeable membrane limits the diffusion of solutes in the water. Not surprisingly, the aquaporin proteins that facilitate water movement play a large role in osmosis, most prominently in red blood cells and the membranes of kidney tubules.

Mechanism of Osmosis

Osmosis is a special case of diffusion. Water, like other substances, moves from an area of high concentration to one of low concentration. An obvious question is what makes water move at all? Imagine a beaker with a semipermeable membrane separating the two sides or halves. On both sides of the membrane the water level is the same, but there are different concentrations of a dissolved substance, or solute, that cannot cross the membrane (otherwise the concentrations on each side would be balanced by the solute crossing the membrane). If the volume of the solution on both sides of the membrane is the same but the concentrations of solute are different, then there are different amounts of water, the solvent, on either side of the membrane. If there is more solute in one area, then there is less water; if there is less solute in one area, then there must be more water.

To illustrate this, imagine two full glasses of water. One has a single teaspoon of sugar in it, whereas the second one contains one-quarter cup of sugar. If the total volume of the solutions in both cups is the same, which cup contains more water? Because the large amount of sugar in the second cup takes up much more space than the teaspoon of sugar in the first cup, the first cup has more water in it.



Osmosis: In osmosis, water always moves from an area of higher water concentration to one of lower concentration. In the diagram shown, the solute cannot pass through the selectively permeable membrane, but the water can.

Returning to the beaker example, recall that it has a mixture of solutes on either side of the membrane. A principle of diffusion is that the molecules move around and will spread evenly throughout the medium if they can. However, only the material capable of passing through the membrane will diffuse through it. In this example, the solute cannot diffuse through the membrane, but the water can. Water has a concentration gradient in this system. Thus, water will diffuse down its concentration gradient, crossing the membrane to the side where it is less concentrated. This diffusion of water through the membrane—osmosis—will continue until the concentration gradient of water goes to zero or until the hydrostatic pressure of the water balances the osmotic pressure. In the beaker example, this means that the level of fluid in the side with a higher solute concentration will go up.

This page titled 3.3B: Osmosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3C: Tonicity

Tonicity, which is directly related to the osmolarity of a solution, affects osmosis by determining the direction of water flow.

LEARNING OBJECTIVES

Define tonicity and describe its relevance to osmosis

KEY TAKEAWAYS

Key Points

- Osmolarity describes the total solute concentration of a solution; solutions with a low solute concentration have a low osmolarity, while those with a high osmolarity have a high solute concentration.
- Water moves from the side of the membrane with lower osmolarity (and more water) to the side with higher osmolarity (and less water).
- In a hypotonic solution, the extracellular fluid has a lower osmolarity than the fluid inside the cell; water enters the cell.
- In a hypertonic solution, the extracellular fluid has a higher osmolarity than the fluid inside the cell; water leaves the cell.
- In an isotonic solution, the extracellular fluid has the same osmolarity as the cell; there will be no net movement of water into or out of the cell.

Key Terms

- **osmolarity**: The osmotic concentration of a solution, normally expressed as osmoles of solute per litre of solution.
- **hypotonic**: Having a lower osmotic pressure than another; a cell in this environment causes water to enter the cell, causing it to swell.
- **hypertonic**: having a greater osmotic pressure than another
- **isotonic**: having the same osmotic pressure

EXAMPLES


Tonicity is the reason why salt water fish cannot live in fresh water and vice versa. A salt water fish's cells have evolved to have a very high solute concentration to match the high osmolarity of the salt water they live in. If you place a salt water fish in fresh water, which has a low osmolarity, water in the environment will flow into the cells of the fish, eventually causing them to burst and killing the fish.

Tonicity describes how an extracellular solution can change the volume of a cell by affecting osmosis. A solution's tonicity often directly correlates with the osmolarity of the solution. Osmolarity describes the total solute concentration of the solution. A solution with low osmolarity has a greater number of water molecules relative to the number of solute particles; a solution with high osmolarity has fewer water molecules with respect to solute particles. In a situation in which solutions of two different osmolarities are separated by a membrane permeable to water, though not to the solute, water will move from the side of the membrane with lower osmolarity (and more water) to the side with higher osmolarity (and less water). This effect makes sense if you remember that the solute cannot move across the membrane, and thus the only component in the system that can move—the water—moves along its own concentration gradient. An important distinction that concerns living systems is that osmolarity measures the number of particles (which may be molecules) in a solution. Therefore, a solution that is cloudy with cells may have a lower osmolarity than a solution that is clear if the second solution contains more dissolved molecules than there are cells.

Hypotonic Solutions

Three terms—hypotonic, isotonic, and hypertonic—are used to relate the osmolarity of a cell to the osmolarity of the extracellular fluid that contains the cells. In a hypotonic situation, the extracellular fluid has lower osmolarity than the fluid inside the cell, and water enters the cell. (In living systems, the point of reference is always the cytoplasm, so the prefix hypo- means that the

extracellular fluid has a lower concentration of solutes, or a lower osmolarity, than the cell cytoplasm.) It also means that the extracellular fluid has a higher concentration of water in the solution than does the cell. In this situation, water will follow its concentration gradient and enter the cell, causing the cell to expand.



Changes in Cell Shape Due to Dissolved Solutes: Osmotic pressure changes the shape of red blood cells in hypertonic, isotonic, and hypotonic solutions.

Hypertonic Solutions

As for a hypertonic solution, the prefix hyper- refers to the extracellular fluid having a higher osmolarity than the cell's cytoplasm; therefore, the fluid contains less water than the cell does. Because the cell has a relatively higher concentration of water, water will leave the cell, and the cell will shrink.

Isotonic Solutions

In an isotonic solution, the extracellular fluid has the same osmolarity as the cell. If the osmolarity of the cell matches that of the extracellular fluid, there will be no net movement of water into or out of the cell, although water will still move in and out.

Blood cells and plant cells in hypertonic, isotonic, and hypotonic solutions take on characteristic appearances. Cells in an isotonic solution retain their shape. Cells in a hypotonic solution swell as water enters the cell, and may burst if the concentration gradient is large enough between the inside and outside of the cell. Cells in a hypertonic solution shrink as water exits the cell, becoming shriveled.

This page titled 3.3C: Tonicity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3D: Facilitated transport

Facilitated diffusion is a process by which molecules are transported across the plasma membrane with the help of membrane proteins.

Learning Objectives

- Explain why and how passive transport occurs

Key Points

- A concentration gradient exists that would allow ions and polar molecules to diffuse into the cell, but these materials are repelled by the hydrophobic parts of the cell membrane.
- Facilitated diffusion uses integral membrane proteins to move polar or charged substances across the hydrophobic regions of the membrane.
- Channel proteins can aid in the facilitated diffusion of substances by forming a hydrophilic passage through the plasma membrane through which polar and charged substances can pass.
- Channel proteins can be open at all times, constantly allowing a particular substance into or out of the cell, depending on the concentration gradient; or they can be gated and can only be opened by a particular biological signal.
- Carrier proteins aid in facilitated diffusion by binding a particular substance, then altering their shape to bring that substance into or out of the cell.

Key Terms

- **facilitated diffusion**: The spontaneous passage of molecules or ions across a biological membrane passing through specific transmembrane integral proteins.
- **membrane protein**: Proteins that are attached to, or associated with the membrane of a cell or an organelle.


EXAMPLES

Channel-mediated facilitated diffusion functions much like a bridge over a river that must raise and lower in order to allow boats to pass. When the bridge is lowered, boats cannot pass through to the other side of the river. Similarly, a gated channel protein often remains closed, not allowing substances into the cell until it receives a signal (like the binding of an ion) to open. When this signal is received, the bridge (gate) opens, allowing the boats (substance) to pass through the bridge and into the other side of the river (cell).

Facilitated transport is a type of passive transport. Unlike simple diffusion where materials pass through a membrane without the help of proteins, in facilitated transport, also called facilitated diffusion, materials diffuse across the plasma membrane with the help of membrane proteins. A concentration gradient exists that would allow these materials to diffuse into the cell without expending cellular energy. However, these materials are ions or polar molecules that are repelled by the hydrophobic parts of the cell membrane. Facilitated transport proteins shield these materials from the repulsive force of the membrane, allowing them to diffuse into the cell.

The material being transported is first attached to protein or glycoprotein receptors on the exterior surface of the plasma membrane. This allows the material that is needed by the cell to be removed from the extracellular fluid. The substances are then passed to specific integral proteins that facilitate their passage. Some of these integral proteins are collections of beta-pleated sheets that form a channel through the phospholipid bilayer. Others are carrier proteins which bind with the substance and aid its diffusion through the membrane.

Channels




Channel Proteins in Facilitated Transport: Facilitated transport moves substances down their concentration gradients. They may cross the plasma membrane with the aid of channel proteins.

The integral proteins involved in facilitated transport are collectively referred to as transport proteins; they function as either channels for the material or carriers. In both cases, they are transmembrane proteins. Channels are specific for the substance that is being transported. Channel proteins have hydrophilic domains exposed to the intracellular and extracellular fluids; they additionally have a hydrophilic channel through their core that provides a hydrated opening through the membrane layers. Passage through the channel allows polar compounds to avoid the nonpolar central layer of the plasma membrane that would otherwise slow or prevent their entry into the cell. Aquaporins are channel proteins that allow water to pass through the membrane at a very high rate.

Channel proteins are either open at all times or they are “gated,” which controls the opening of the channel. The attachment of a particular ion to the channel protein may control the opening or other mechanisms or substances may be involved. In some tissues, sodium and chloride ions pass freely through open channels, whereas in other tissues, a gate must be opened to allow passage. An example of this occurs in the kidney, where both forms of channels are found in different parts of the renal tubules. Cells involved in the transmission of electrical impulses, such as nerve and muscle cells, have gated channels for sodium, potassium, and calcium in their membranes. Opening and closing of these channels changes the relative concentrations on opposing sides of the membrane of these ions, resulting in the facilitation of electrical transmission along membranes (in the case of nerve cells) or in muscle contraction (in the case of muscle cells).

Carrier Proteins

Another type of protein embedded in the plasma membrane is a carrier protein. This protein binds a substance and, in doing so, triggers a change of its own shape, moving the bound molecule from the outside of the cell to its interior; depending on the gradient, the material may move in the opposite direction. Carrier proteins are typically specific for a single substance. This adds to the overall selectivity of the plasma membrane. The exact mechanism for the change of shape is poorly understood. Proteins can change shape when their hydrogen bonds are affected, but this may not fully explain this mechanism. Each carrier protein is specific to one substance, and there are a finite number of these proteins in any membrane. This can cause problems in transporting enough of the material for the cell to function properly.



Carrier Proteins: Some substances are able to move down their concentration gradient across the plasma membrane with the aid of carrier proteins. Carrier proteins change shape as they move molecules across the membrane.

An example of this process occurs in the kidney. Glucose, water, salts, ions, and amino acids needed by the body are filtered in one part of the kidney. This filtrate, which includes glucose, is then reabsorbed in another part of the kidney. Because there are only a finite number of carrier proteins for glucose, if more glucose is present than the proteins can handle, the excess is not transported; it is excreted from the body in the urine. In a diabetic individual, this is described as “spilling glucose into the urine.” A different group of carrier proteins called glucose transport proteins, or GLUTs, are involved in transporting glucose and other hexose sugars through plasma membranes within the body.

Channel and carrier proteins transport material at different rates. Channel proteins transport much more quickly than do carrier proteins. Channel proteins facilitate diffusion at a rate of tens of millions of molecules per second, whereas carrier proteins work at a rate of a thousand to a million molecules per second.

This page titled 3.3D: Facilitated transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3E: The Role of Passive Transport

Passive transport, such as diffusion and osmosis, moves materials of small molecular weight across membranes.

Learning Objectives

- Indicate the manner in which various materials cross the cell membrane

Key Points

- Plasma membranes are selectively permeable; if they were to lose this selectivity, the cell would no longer be able to sustain itself.
- In passive transport, substances simply move from an area of higher concentration to an area of lower concentration, which does not require the input of energy.
- Concentration gradient, size of the particles that are diffusing, and temperature of the system affect the rate of diffusion.
- Some materials diffuse readily through the membrane, but others require specialized proteins, such as channels and transporters, to carry them into or out of the cell.


Key Terms

- **concentration gradient**: A concentration gradient is present when a membrane separates two different concentrations of molecules.
- **passive transport**: A movement of biochemicals and other atomic or molecular substances across membranes that does not require an input of chemical energy.
- **permeable**: Of or relating to substance, substrate, membrane or material that absorbs or allows the passage of fluids.

Introduction: Passive Transport

Plasma membranes must allow or prevent certain substances from entering or leaving a cell. In other words, plasma membranes are selectively permeable; they allow some substances to pass through, but not others. If they were to lose this selectivity, the cell would no longer be able to sustain itself, and it would be destroyed. Some cells require larger amounts of specific substances than other cells; they must have a way of obtaining these materials from extracellular fluids. This may happen passively, as certain materials move back and forth, or the cell may have special mechanisms that facilitate transport. Some materials are so important to a cell that it spends some of its energy (hydrolyzing adenosine triphosphate (ATP)) to obtain these materials. Red blood cells use some of their energy to do this. All cells spend the majority of their energy to maintain an imbalance of sodium and potassium ions between the interior and exterior of the cell.

The most direct forms of membrane transport are passive. Passive transport is a naturally-occurring phenomenon and does not require the cell to exert any of its energy to accomplish the movement. In passive transport, substances move from an area of higher concentration to an area of lower concentration. A physical space in which there is a range of concentrations of a single substance is said to have a concentration gradient.



Passive Transport: Diffusion is a type of passive transport. Diffusion through a permeable membrane moves a substance from an area of high concentration (extracellular fluid, in this case) down its concentration gradient (into the cytoplasm).

The passive forms of transport, diffusion and osmosis, move materials of small molecular weight across membranes. Substances diffuse from areas of high concentration to areas of lower concentration; this process continues until the substance is evenly distributed in a system. In solutions containing more than one substance, each type of molecule diffuses according to its own concentration gradient, independent of the diffusion of other substances. Many factors can affect the rate of diffusion, including, but not limited to, concentration gradient, size of the particles that are diffusing, and temperature of the system.

In living systems, diffusion of substances in and out of cells is mediated by the plasma membrane. Some materials diffuse readily through the membrane, but others are hindered; their passage is made possible by specialized proteins, such as channels and transporters. The chemistry of living things occurs in aqueous solutions; balancing the concentrations of those solutions is an ongoing problem. In living systems, diffusion of some substances would be slow or difficult without membrane proteins that facilitate transport.

This page titled 3.3E: The Role of Passive Transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3F: Primary Active Transport

The sodium-potassium pump maintains the electrochemical gradient of living cells by moving sodium in and potassium out of the cell.

Learning Objectives

- Describe how a cell moves sodium and potassium out of and into the cell against its electrochemical gradient

Key Points


- The sodium-potassium pump moves K+ into the cell while moving Na+ at a ratio of three Na+ for every two K+ ions.
- When the sodium-potassium- ATPase enzyme points into the cell, it has a high affinity for sodium ions and binds three of them, hydrolyzing ATP and changing shape.
- As the enzyme changes shape, it reorients itself towards the outside of the cell, and the three sodium ions are released.
- The enzyme's new shape allows two potassium to bind and the phosphate group to detach, and the carrier protein repositions itself towards the interior of the cell.
- The enzyme changes shape again, releasing the potassium ions into the cell.
- After potassium is released into the cell, the enzyme binds three sodium ions, which starts the process over again.

Key Terms

- **electrogenic pump**: An ion pump that generates a net charge flow as a result of its activity.
- **Na+-K+ ATPase**: An enzyme located in the plasma membrane of all animal cells that pumps sodium out of cells while pumping potassium into cells.

Primary Active Transport

The primary active transport that functions with the active transport of sodium and potassium allows secondary active transport to occur. The secondary transport method is still considered active because it depends on the use of energy as does primary transport.



Active Transport of Sodium and Potassium: Primary active transport moves ions across a membrane, creating an electrochemical gradient (electrogenic transport).

One of the most important pumps in animals cells is the sodium-potassium pump (Na^+-K^+ ATPase), which maintains the electrochemical gradient (and the correct concentrations of Na^+ and K^+) in living cells. The sodium-potassium pump moves two K^+ into the cell while moving three Na^+ out of the cell. The Na+-K+ ATPase exists in two forms, depending on its orientation to the interior or exterior of the cell and its affinity for either sodium or potassium ions. The process consists of the following six steps:

- With the enzyme oriented towards the interior of the cell, the carrier has a high affinity for sodium ions. Three sodium ions bind to the protein.
- ATP is hydrolyzed by the protein carrier, and a low-energy phosphate group attaches to it.
- As a result, the carrier changes shape and re-orients itself towards the exterior of the membrane. The protein's affinity for sodium decreases, and the three sodium ions leave the carrier.
- The shape change increases the carrier's affinity for potassium ions, and two such ions attach to the protein. Subsequently, the low-energy phosphate group detaches from the carrier.
- With the phosphate group removed and potassium ions attached, the carrier protein repositions itself towards the interior of the cell.
- The carrier protein, in its new configuration, has a decreased affinity for potassium, and the two ions are released into the cytoplasm. The protein now has a higher affinity for sodium ions, and the process starts again.

Several things have happened as a result of this process. At this point, there are more sodium ions outside of the cell than inside and more potassium ions inside than out. For every three ions of sodium that move out, two ions of potassium move in. This results in the interior being slightly more negative relative to the exterior. This difference in charge is important in creating the conditions necessary for the secondary process. The sodium-potassium pump is, therefore, an electrogenic pump (a pump that creates a charge imbalance), creating an electrical imbalance across the membrane and contributing to the membrane potential.

This page titled 3.3F: Primary Active Transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3G: Electrochemical Gradient

To move substances against the membrane's electrochemical gradient, the cell utilizes active transport, which requires energy from ATP.

Learning Objectives

- Define an electrochemical gradient and describe how a cell moves substances against this gradient


Key Points

- The electrical and concentration gradients of a membrane tend to drive sodium into and potassium out of the cell, and active transport works against these gradients.
- To move substances against a concentration or electrochemical gradient, the cell must utilize energy in the form of ATP during active transport.
- Primary active transport, which is directly dependent on ATP, moves ions across a membrane and creates a difference in charge across that membrane.
- Secondary active transport, created by primary active transport, is the transport of a solute in the direction of its electrochemical gradient and does not directly require ATP.
- Carrier proteins such as uniporters, symporters, and antiporters perform primary active transport and facilitate the movement of solutes across the cell's membrane.

Key Terms

- **adenosine triphosphate**: a multifunctional nucleoside triphosphate used in cells as a coenzyme, often called the "molecular unit of energy currency" in intracellular energy transfer
- **active transport**: movement of a substance across a cell membrane against its concentration gradient (from low to high concentration) facilitated by ATP conversion
- **electrochemical gradient**: The difference in charge and chemical concentration across a membrane.

Electrochemical Gradients



Electrochemical Gradient: Electrochemical gradients arise from the combined effects of concentration gradients and electrical gradients.

Simple concentration gradients are differential concentrations of a substance across a space or a membrane, but in living systems, gradients are more complex. Because ions move into and out of cells and because cells contain proteins that do not move across the membrane and are mostly negatively charged, there is also an electrical gradient, a difference of charge, across the plasma membrane. The interior of living cells is electrically negative with respect to the extracellular fluid in which they are bathed. At the same time, cells have higher concentrations of potassium (K^+) and lower concentrations of sodium (Na^+) than does the extracellular fluid. In a living cell, the concentration gradient of Na^+ tends to drive it into the cell, and the electrical gradient of Na^+ (a positive ion) also tends to drive it inward to the negatively-charged interior. The situation is more complex, however, for other elements such as potassium. The electrical gradient of K^+, a positive ion, also tends to drive it into the cell, but the concentration gradient of K^+ tends to drive K^+ out of the cell. The combined gradient of concentration and electrical charge that affects an ion is called its electrochemical gradient.


Moving Against a Gradient

To move substances against a concentration or electrochemical gradient, the cell must use energy. This energy is harvested from adenosine triphosphate (ATP) generated through the cell's metabolism. Active transport mechanisms, collectively called pumps, work against electrochemical gradients. Small substances constantly pass through plasma membranes. Active transport maintains concentrations of ions and other substances needed by living cells in the face of these passive movements. Much of a cell's supply of metabolic energy may be spent maintaining these processes. For example, most of a red blood cell's metabolic energy is used to maintain the imbalance between exterior and interior sodium and potassium levels required by the cell. Because active transport mechanisms depend on a cell's metabolism for energy, they are sensitive to many metabolic poisons that interfere with the supply of ATP.

Two mechanisms exist for the transport of small-molecular weight material and small molecules. Primary active transport moves ions across a membrane and creates a difference in charge across that membrane, which is directly dependent on ATP. Secondary active transport describes the movement of material that is due to the electrochemical gradient established by primary active transport that does not directly require ATP.

Carrier Proteins for Active Transport

An important membrane adaption for active transport is the presence of specific carrier proteins or pumps to facilitate movement. There are three types of these proteins or transporters: uniporters, symporters, and antiporters. A uniporter carries one specific ion or molecule. A symporter carries two different ions or molecules, both in the same direction. An antiporter also carries two different ions or molecules, but in different directions. All of these transporters can also transport small, uncharged organic molecules like glucose. These three types of carrier proteins are also found in facilitated diffusion, but they do not require ATP to work in that process. Some examples of pumps for active transport are Na^+-K^+ ATPase, which carries sodium and potassium ions, and H^+-K^+ ATPase, which carries hydrogen and potassium ions. Both of these are antiporter carrier proteins. Two other carrier protein pumps are Ca^{2+} ATPase and H^+ATPase, which carry only calcium and only hydrogen ions, respectively.



Uniporters, Symporters, and Antiporters: A uniporter carries one molecule or ion. A symporter carries two different molecules or ions, both in the same direction. An antiporter also carries two different molecules or ions, but in different directions.

This page titled 3.3G: Electrochemical Gradient is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3H: Secondary Active Transport

In secondary active transport, a molecule is moved down its electrochemical gradient as another is moved up its concentration gradient.

Learning Objectives

- Differentiate between primary and secondary active transport

Key Points

- While secondary active transport consumes ATP to generate the gradient down which a molecule is moved, the energy is not directly used to move the molecule across the membrane.
- Both antiporters and symporters are used in secondary active transport.
- Secondary active transport brings sodium ions into the cell, and as sodium ion concentrations build outside the plasma membrane, an electrochemical gradient is created.
- If a channel protein is open via primary active transport, the ions will be pulled through the membrane along with other substances that can attach themselves to the transport protein through the membrane.
- Secondary active transport is used to store high-energy hydrogen ions in the mitochondria of plant and animal cells for the production of ATP.
- The potential energy in the hydrogen ions is translated into kinetic energy as the ions surge through the channel protein ATP synthase, and that energy is used to convert ADP into ATP.


Key Terms

- **secondary active transport**: A method of transport in which the electrochemical potential difference created by pumping ions out of the cell is used to transport molecules across a membrane.

Secondary Active Transport (Co-transport)

Unlike in primary active transport, in secondary active transport, ATP is not directly coupled to the molecule of interest. Instead, another molecule is moved up its concentration gradient, which generates an electrochemical gradient. The molecule of interest is then transported down the electrochemical gradient. While this process still consumes ATP to generate that gradient, the energy is not directly used to move the molecule across the membrane, hence it is known as secondary active transport. Both antiporters and symporters are used in secondary active transport. Co-transporters can be classified as symporters and antiporters depending on whether the substances move in the same or opposite directions across the cell membrane.

Secondary active transport brings sodium ions, and possibly other compounds, into the cell. As sodium ion concentrations build outside the plasma membrane because of the action of the primary active transport process, an electrochemical gradient is created. If a channel protein exists and is open, the sodium ions will be pulled through the membrane. This movement is used to transport other substances that can attach themselves to the transport protein through the membrane. Many amino acids, as well as glucose, enter a cell this way. This secondary process is also used to store high-energy hydrogen ions in the mitochondria of plant and animal cells for the production of ATP. The potential energy that accumulates in the stored hydrogen ions is translated into kinetic energy as the ions surge through the channel protein ATP synthase, and that energy is used to convert ADP into ATP.



Secondary Active Transport: An electrochemical gradient, created by primary active transport, can move other substances against their concentration gradients, a process called co-transport or secondary active transport.

This page titled 3.3H: Secondary Active Transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.3I: Endocytosis

Endocytosis takes up particles into the cell by invaginating the cell membrane, resulting in the release of the material inside of the cell.

Learning Objectives

- Describe endocytosis, including phagocytosis, pinocytosis, and receptor-mediated endocytosis.

Key Points

- Endocytosis consists of phagocytosis, pinocytosis, and receptor -mediated endocytosis.
- Endocytosis takes particles into the cell that are too large to passively cross the cell membrane.
- Phagocytosis is the taking in of large food particles, while pinocytosis takes in liquid particles.
- Receptor-mediated endocytosis uses special receptor proteins to help carry large particles across the cell membrane.


Key Terms

- **endosome**: An endocytic vacuole through which molecules internalized during endocytosis pass en route to lysosomes
- **neutrophil**: A cell, especially a white blood cell that consumes foreign invaders in the blood.

Endocytosis

Endocytosis is a type of active transport that moves particles, such as large molecules, parts of cells, and even whole cells, into a cell. There are different variations of endocytosis, but all share a common characteristic: the plasma membrane of the cell invaginates, forming a pocket around the target particle. The pocket pinches off, resulting in the particle being contained in a newly-created intracellular vesicle formed from the plasma membrane.

Phagocytosis




Phagocytosis: In phagocytosis, the cell membrane surrounds the particle and engulfs it.

Phagocytosis (the condition of “cell eating”) is the process by which large particles, such as cells or relatively large particles, are taken in by a cell. For example, when microorganisms invade the human body, a type of white blood cell called a neutrophil will remove the invaders through this process, surrounding and engulfing the microorganism, which is then destroyed by the neutrophil.

In preparation for phagocytosis, a portion of the inward-facing surface of the plasma membrane becomes coated with a protein called clathrin, which stabilizes this section of the membrane. The coated portion of the membrane then extends from the body of the cell and surrounds the particle, eventually enclosing it. Once the vesicle containing the particle is enclosed within the cell, the clathrin disengages from the membrane and the vesicle merges with a lysosome for the breakdown of the material in the newly-formed compartment (endosome). When accessible nutrients from the degradation of the vesicular contents have been extracted, the newly-formed endosome merges with the plasma membrane and releases its contents into the extracellular fluid. The endosomal membrane again becomes part of the plasma membrane.

Pinocytosis




Pinocytosis: In pinocytosis, the cell membrane invaginates, surrounds a small volume of fluid, and pinches off.

A variation of endocytosis is called pinocytosis. This literally means “cell drinking” and was named at a time when the assumption was that the cell was purposefully taking in extracellular fluid. In reality, this is a process that takes in molecules, including water, which the cell needs from the extracellular fluid. Pinocytosis results in a much smaller vesicle than does phagocytosis, and the vesicle does not need to merge with a lysosome.

Potocytosis, a variant of pinocytosis, is a process that uses a coating protein, called caveolin, on the cytoplasmic side of the plasma membrane, which performs a similar function to clathrin. The cavities in the plasma membrane that form the vacuoles have membrane receptors and lipid rafts in addition to caveolin. The vacuoles or vesicles formed in caveolae (singular caveola) are smaller than those in pinocytosis. Potocytosis is used to bring small molecules into the cell and to transport these molecules through the cell for their release on the other side of the cell, a process called transcytosis.

Receptor-mediated Endocytosis



Receptor-Mediated Endocytosis: In receptor-mediated endocytosis, uptake of substances by the cell is targeted to a single type of substance that binds to the receptor on the external surface of the cell membrane.

A targeted variation of endocytosis, known as receptor-mediated endocytosis, employs receptor proteins in the plasma membrane that have a specific binding affinity for certain substances. In receptor-mediated endocytosis, as in phagocytosis, clathrin is attached to the cytoplasmic side of the plasma membrane. If uptake of a compound is dependent on receptor-mediated endocytosis and the process is ineffective, the material will not be removed from the tissue fluids or blood. Instead, it will stay in those fluids and increase in concentration. Some human diseases are caused by the failure of receptor-mediated endocytosis. For example, the form of cholesterol termed low-density lipoprotein or LDL (also referred to as “bad” cholesterol) is removed from the blood by receptor-mediated endocytosis. In the human genetic disease familial hypercholesterolemia, the LDL receptors are defective or missing entirely. People with this condition have life-threatening levels of cholesterol in their blood, because their cells cannot clear LDL particles from their blood.

Although receptor-mediated endocytosis is designed to bring specific substances that are normally found in the extracellular fluid into the cell, other substances may gain entry into the cell at the same site. Flu viruses, diphtheria, and cholera toxin all have sites that cross-react with normal receptor-binding sites and gain entry into cells.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- concentration gradient. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/concentration%20gradient**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...tion/diffusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest...e_05_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest...ol11448/latest**. **License**: ***CC BY: Attribution***

- osmosis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osmosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- solute. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/solute. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- hypotonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypotonic. **License**: *CC BY-SA: Attribution-ShareAlike*
- hypertonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypertonic. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...ol11448/latest. **License**: *CC BY: Attribution*
- isotonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/isotonic. **License**: *CC BY-SA: Attribution-ShareAlike*
- osmolarity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osmolarity. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44417/latest...e_05_02_07.png. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...ol11448/latest. **License**: *CC BY: Attribution*
- membrane protein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/membrane%20protein. **License**: *CC BY-SA: Attribution-ShareAlike*
- facilitated diffusion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/facilitated%20diffusion. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44417/latest...e_05_02_07.png. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...ol11448/latest. **License**: *CC BY: Attribution*
- passive transport. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/passive%20transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- permeable. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/permeable. **License**: *CC BY-SA: Attribution-ShareAlike*
- concentration gradient. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/concentration%20gradient. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44417/latest...e_05_02_07.png. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/

m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...ctrogenic-pump. **License**: *CC BY-SA: Attribution-ShareAlike*
- Na -K ATPase. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Na%20-K%20%20ATPase. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44417/latest...e_05_02_07.png. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest...e_05_03_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- active transport. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/active%20transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- electrochemical gradient. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/electro...cal%20gradient. **License**: *CC BY-SA: Attribution-ShareAlike*
- adenosine triphosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adenosine%20triphosphate. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest...e_05_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44417/latest...e_05_02_07.png. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest/Figure_05_03_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest...e_05_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest...e_05_03_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44418/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- Structural Biochemistry/Membrane Proteins. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structural_Biochemistry/Membrane_Proteins%23Secondary_Active_Transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- secondary active transport. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/secondary%20active%20transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44417/latest/Figure_05_02_06.jpg. **License**: *CC BY: Attribution*

- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **cnx.org/content/m44417/latest...e_05_02_07.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_04.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_05.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest...e_05_03_04.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44419/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- neutrophil. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neutrophil**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endosome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endosome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_06.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **cnx.org/content/m44417/latest...e_05_02_07.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_04.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_05.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Passive Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44417/latest/Figure_05_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Active Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44418/latest/Figure_05_03_04.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Bulk Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44419/latest...e_05_04_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Bulk Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44419/latest...e_05_04_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Bulk Transport. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44419/latest...e_05_04_03.jpg**. **License**: ***CC BY: Attribution***

This page titled 3.3I: Endocytosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.4: CELL SIGNALING



- Topic hierarchy

3.4A: Types of Receptors

This page titled 3.4: Cell Signaling is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.4A: Types of Receptors

Receptors, either intracellular or cell-surface, bind to specific ligands, which activate numerous cellular processes.

Learning Objectives

- Compare internal receptors with cell-surface receptors

Key Points

- Intracellular receptors are located in the cytoplasm of the cell and are activated by hydrophobic ligand molecules that can pass through the plasma membrane.
- Cell-surface receptors bind to an external ligand molecule and convert an extracellular signal into an intracellular signal.
- Three general categories of cell-surface receptors include: ion -channel, G- protein, and enzyme -linked protein receptors.
- Ion channel -linked receptors bind a ligand and open a channel through the membrane that allows specific ions to pass through.
- G-protein-linked receptors bind a ligand and activate a membrane protein called a G-protein, which then interacts with either an ion channel or an enzyme in the membrane.
- Enzyme-linked receptors are cell-surface receptors with intracellular domains that are associated with an enzyme.

Key Terms

- **integral protein**: a protein molecule (or assembly of proteins) that is permanently attached to the biological membrane
- **transcription**: the synthesis of RNA under the direction of DNA


EXAMPLES

Bacteria that are pathogenic to humans can release poisons that interrupt specific G-protein-linked receptor function, leading to illness. In cholera, the water-borne bacterium *Vibrio cholerae* produces a toxin, choleragen, that binds to cells lining the small intestine. The toxin then enters these intestinal cells, where it modifies a G-protein that controls the opening of a chloride channel and causes it to remain continuously active, resulting in large losses of fluids from the body that can lead to potentially fatal dehydration.

Receptors are protein molecules in the target cell or on its surface that bind ligands. There are two types of receptors: internal receptors and cell-surface receptors.

Internal receptors

Internal receptors, also known as intracellular or cytoplasmic receptors, are found in the cytoplasm of the cell and respond to hydrophobic ligand molecules that are able to travel across the plasma membrane. Once inside the cell, many of these molecules bind to proteins that act as regulators of mRNA synthesis to mediate gene expression. Gene expression is the cellular process of transforming the information in a cell's DNA into a sequence of amino acids that ultimately forms a protein. When the ligand binds to the internal receptor, a conformational change exposes a DNA-binding site on the protein. The ligand-receptor complex moves into the nucleus, binds to specific regulatory regions of the chromosomal DNA, and promotes the initiation of transcription. Internal receptors can directly influence gene expression without having to pass the signal on to other receptors or messengers.



Intracellular Receptors: Hydrophobic signaling molecules typically diffuse across the plasma membrane and interact with intracellular receptors in the cytoplasm. Many intracellular receptors are transcription factors that interact with DNA in the nucleus and regulate gene expression.

Cell-Surface Receptors


Cell-surface receptors, also known as transmembrane receptors, are cell surface, membrane-anchored, or integral proteins that bind to external ligand molecules. This type of receptor spans the plasma membrane and performs signal transduction, converting an extracellular signal into an intracellular signal. Ligands that interact with cell-surface receptors do not have to enter the cell that they affect. Cell-surface receptors are also called cell-specific proteins or markers because they are specific to individual cell types.

Each cell-surface receptor has three main components: an external ligand-binding domain (extracellular domain), a hydrophobic membrane-spanning region, and an intracellular domain inside the cell. The size and extent of each of these domains vary widely, depending on the type of receptor.

Cell-surface receptors are involved in most of the signaling in multicellular organisms. There are three general categories of cell-surface receptors: ion channel-linked receptors, G-protein-linked receptors, and enzyme-linked receptors.

Ion Channel-Linked Receptors

Ion channel-linked receptors bind a ligand and open a channel through the membrane that allows specific ions to pass through. To form a channel, this type of cell-surface receptor has an extensive membrane-spanning region. In order to interact with the phospholipid fatty acid tails that form the center of the plasma membrane, many of the amino acids in the membrane-spanning region are hydrophobic in nature. Conversely, the amino acids that line the inside of the channel are hydrophilic to allow for the passage of water or ions. When a ligand binds to the extracellular region of the channel, there is a conformational change in the protein's structure that allows ions such as sodium, calcium, magnesium, and hydrogen to pass through.




Gated-Ion Channels: Gated ion channels form a pore through the plasma membrane that opens when the signaling molecule binds. The open pore then allows ions to flow into or out of the cell.

G-Protein Linked Receptors

G-protein-linked receptors bind a ligand and activate a membrane protein called a G-protein. The activated G-protein then interacts with either an ion channel or an enzyme in the membrane. All G-protein-linked receptors have seven transmembrane domains, but each receptor has its own specific extracellular domain and G-protein-binding site.

Cell signaling using G-protein-linked receptors occurs as a cyclic series of events. Before the ligand binds, the inactive G-protein can bind to a newly-revealed site on the receptor specific for its binding. Once the G-protein binds to the receptor, the resultant shape change activates the G-protein, which releases GDP and picks up GTP. The subunits of the G-protein then split into the α subunit and the β subunit. One or both of these G-protein fragments may be able to activate other proteins as a result. Later, the GTP on the active α subunit of the G-protein is hydrolyzed to GDP and the β subunit is deactivated. The subunits reassociate to form the inactive G-protein, and the cycle starts over.



G-proteins: Heterotrimeric G proteins have three subunits: α, β, and γ. When a signaling molecule binds to a G-protein-coupled receptor in the plasma membrane, a GDP molecule associated with the α subunit is exchanged for GTP. The β and γ subunits

dissociate from the α subunit, and a cellular response is triggered either by the α subunit or the dissociated β pair. Hydrolysis of GTP to GDP terminates the signal.

Enzyme-Linked Receptors

Enzyme-linked receptors are cell-surface receptors with intracellular domains that are associated with an enzyme. In some cases, the intracellular domain of the receptor itself is an enzyme or the enzyme-linked receptor has an intracellular domain that interacts directly with an enzyme. The enzyme-linked receptors normally have large extracellular and intracellular domains, but the membrane-spanning region consists of a single alpha-helical region of the peptide strand. When a ligand binds to the extracellular domain, a signal is transferred through the membrane and activates the enzyme, which sets off a chain of events within the cell that eventually leads to a response. An example of this type of enzyme-linked receptor is the tyrosine kinase receptor. The tyrosine kinase receptor transfers phosphate groups to tyrosine molecules. Signaling molecules bind to the extracellular domain of two nearby tyrosine kinase receptors, which then dimerize. Phosphates are then added to tyrosine residues on the intracellular domain of the receptors and can then transmit the signal to the next messenger within the cytoplasm.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44451/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- transcription. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/transcription**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- integral protein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/integral%20protein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Signaling Molecules and Cellular Receptors. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44451/latest...e_09_01_05.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Signaling Molecules and Cellular Receptors. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44451/latest...e_09_01_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Signaling Molecules and Cellular Receptors. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44451/latest...e_09_01_04.jpg**. **License**: ***CC BY: Attribution***

This page titled 3.4A: Types of Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.5: THE NUCLEUS AND RIBOSOMES



- Topic hierarchy

3.5A: The Nucleus and Ribosomes

3.5B: Characteristics of Eukaryotic Cells

This page titled 3.5: The Nucleus and Ribosomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.5A: The Nucleus and Ribosomes

Found within eukaryotic cells, the nucleus contains the genetic material that determines the entire structure and function of that cell.

Learning Objectives

- Explain the purpose of the nucleus in eukaryotic cells

Key Points

- The nucleus contains the cell ‘s DNA and directs the synthesis of ribosomes and proteins.
- Found within the nucleoplasm, the nucleolus is a condensed region of chromatin where ribosome synthesis occurs.
- Chromatin consists of DNA wrapped around histone proteins and is stored within the nucleoplasm.
- Ribosomes are large complexes of protein and ribonucleic acid (RNA) responsible for protein synthesis when DNA from the nucleus is transcribed.


Key Terms

- **histone**: any of various simple water-soluble proteins that are rich in the basic amino acids lysine and arginine and are complexed with DNA in the nucleosomes of eukaryotic chromatin
- **nucleolus**: a conspicuous, rounded, non-membrane bound body within the nucleus of a cell
- **chromatin**: a complex of DNA, RNA, and proteins within the cell nucleus out of which chromosomes condense during cell division


The Nucleus

One of the main differences between prokaryotic and eukaryotic cells is the nucleus. As previously discussed, prokaryotic cells lack an organized nucleus while eukaryotic cells contain membrane-bound nuclei (and organelles) that house the cell’s DNA and direct the synthesis of ribosomes and proteins.

The nucleus stores chromatin (DNA plus proteins) in a gel-like substance called the nucleoplasm. To understand chromatin, it is helpful to first consider chromosomes. Chromatin describes the material that makes up chromosomes, which are structures within the nucleus that are made up of DNA, the hereditary material. You may remember that in prokaryotes, DNA is organized into a single circular chromosome. In eukaryotes, chromosomes are linear structures. Every eukaryotic species has a specific number of chromosomes in the nuclei of its body’s cells. For example, in humans, the chromosome number is 46, while in fruit flies, it is eight. Chromosomes are only visible and distinguishable from one another when the cell is getting ready to divide. In order to organize the large amount of DNA within the nucleus, proteins called histones are attached to chromosomes; the DNA is wrapped around these histones to form a structure resembling beads on a string. These protein-chromosome complexes are called chromatin.




DNA is highly organized: This image shows various levels of the organization of chromatin (DNA and protein). Along the chromatin threads, unwound protein-chromosome complexes, we find DNA wrapped around a set of histone proteins.



The nucleus stores the hereditary material of the cell: The nucleus is the control center of the cell. The nucleus of living cells contains the genetic material that determines the entire structure and function of that cell.

The nucleoplasm is also where we find the nucleolus. The nucleolus is a condensed region of chromatin where ribosome synthesis occurs. Ribosomes, large complexes of protein and ribonucleic acid (RNA), are the cellular organelles responsible for protein synthesis. They receive their “orders” for protein synthesis from the nucleus where the DNA is transcribed into messenger RNA (mRNA). This mRNA travels to the ribosomes, which translate the code provided by the sequence of the nitrogenous bases in the mRNA into a specific order of amino acids in a protein.



Ribosomes are responsible for protein synthesis: Ribosomes are made up of a large subunit (top) and a small subunit (bottom). During protein synthesis, ribosomes assemble amino acids into proteins.

Lastly, the boundary of the nucleus is called the nuclear envelope. It consists of two phospholipid bilayers: an outer membrane and an inner membrane. The nuclear membrane is continuous with the endoplasmic reticulum, while nuclear pores allow substances to enter and exit the nucleus.

This page titled 3.5A: The Nucleus and Ribosomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.5B: Characteristics of Eukaryotic Cells

A eukaryotic cell has a true membrane-bound nucleus and has other membranous organelles that allow for compartmentalization of functions.

Learning Objectives

- Describe the structure of eukaryotic cells

Key Points

- Eukaryotic cells are larger than prokaryotic cells and have a “true” nucleus, membrane-bound organelles, and rod-shaped chromosomes.
- The nucleus houses the cell’s DNA and directs the synthesis of proteins and ribosomes.
- Mitochondria are responsible for ATP production; the endoplasmic reticulum modifies proteins and synthesizes lipids; and the golgi apparatus is where the sorting of lipids and proteins takes place.
- Peroxisomes carry out oxidation reactions that break down fatty acids and amino acids and detoxify poisons; vesicles and vacuoles function in storage and transport.
- Animal cells have a centrosome and lysosomes while plant cells do not.
- Plant cells have a cell wall, a large central vacuole, chloroplasts, and other specialized plastids, whereas animal cells do not.

Key Terms

- **eukaryotic**: Having complex cells in which the genetic material is organized into membrane-bound nuclei.
- **organelle**: A specialized structure found inside cells that carries out a specific life process (e.g. ribosomes, vacuoles).
- **photosynthesis**: the process by which plants and other photoautotrophs generate carbohydrates and oxygen from carbon dioxide, water, and light energy in chloroplasts

Eukaryotic Cell Structure

Like a prokaryotic cell, a eukaryotic cell has a plasma membrane, cytoplasm, and ribosomes. However, unlike prokaryotic cells, eukaryotic cells have:


1. a membrane-bound nucleus
2. numerous membrane-bound organelles (including the endoplasmic reticulum, Golgi apparatus, chloroplasts, and mitochondria)
3. several rod-shaped chromosomes

Because a eukaryotic cell’s nucleus is surrounded by a membrane, it is often said to have a “true nucleus.” Organelles (meaning “little organ”) have specialized cellular roles, just as the organs of your body have specialized roles. They allow different functions to be compartmentalized in different areas of the cell.

The Nucleus & Its Structures

Typically, the nucleus is the most prominent organelle in a cell. Eukaryotic cells have a true nucleus, which means the cell’s DNA is surrounded by a membrane. Therefore, the nucleus houses the cell’s DNA and directs the synthesis of proteins and ribosomes, the cellular organelles responsible for protein synthesis. The nuclear envelope is a double-membrane structure that constitutes the outermost portion of the nucleus. Both the inner and outer membranes of the nuclear envelope are phospholipid bilayers. The nuclear envelope is punctuated with pores that control the passage of ions, molecules, and RNA between the nucleoplasm and cytoplasm. The nucleoplasm is the semi-solid fluid inside the nucleus where we find the chromatin and the nucleolus. Furthermore, chromosomes are structures within the nucleus that are made up of DNA, the genetic material. In prokaryotes, DNA

is organized into a single circular chromosome. In eukaryotes, chromosomes are linear structures.




Eukaryotic Nucleus: The nucleus stores chromatin (DNA plus proteins) in a gel-like substance called the nucleoplasm.The nucleolus is a condensed region of chromatin where ribosome synthesis occurs.The boundary of the nucleus is called the nuclear envelope.It consists of two phospholipid bilayers: an outer membrane and an inner membrane.The nuclear membrane is continuous with the endoplasmic reticulum.Nuclear pores allow substances to enter and exit the nucleus.

Other Membrane-Bound Organelles

Mitochondria are oval-shaped, double membrane organelles that have their own ribosomes and DNA. These organelles are often called the “energy factories” of a cell because they are responsible for making adenosine triphosphate (ATP), the cell’s main energy-carrying molecule, by conducting cellular respiration. The endoplasmic reticulum modifies proteins and synthesizes lipids, while the golgi apparatus is where the sorting, tagging, packaging, and distribution of lipids and proteins takes place. Peroxisomes are small, round organelles enclosed by single membranes; they carry out oxidation reactions that break down fatty acids and amino acids. Peroxisomes also detoxify many poisons that may enter the body. Vesicles and vacuoles are membrane-bound sacs that function in storage and transport. Other than the fact that vacuoles are somewhat larger than vesicles, there is a very subtle distinction between them: the membranes of vesicles can fuse with either the plasma membrane or other membrane systems within the cell. All of these organelles are found in each and every eukaryotic cell.


Animal Cells Versus Plant Cells

While all eukaryotic cells contain the aforementioned organelles and structures, there are some striking differences between animal and plant cells. Animal cells have a centrosome and lysosomes, whereas plant cells do not. The centrosome is a microtubule-organizing center found near the nuclei of animal cells while lysosomes take care of the cell’s digestive process.



Animal Cells: Despite their fundamental similarities, there are some striking differences between animal and plant cells.Animal cells have centrioles, centrosomes, and lysosomes, whereas plant cells do not.

In addition, plant cells have a cell wall, a large central vacuole, chloroplasts, and other specialized plastids, whereas animal cells do not. The cell wall protects the cell, provides structural support, and gives shape to the cell while the central vacuole plays a key role in regulating the cell's concentration of water in changing environmental conditions. Chloroplasts are the organelles that carry out photosynthesis.



Plant Cells: Plant cells have a cell wall, chloroplasts, plasmodesmata, and plastids used for storage, and a large central vacuole, whereas animal cells do not.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 21, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- histone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/histone. **License**: *CC BY-SA: Attribution-ShareAlike*
- chromatin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chromatin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...n/nucleolus--2. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, The Nucleus and DNA Replication. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46073/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Cells. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m45432/latest/. **License**: *CC BY: Attribution*
- organelle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/organelle. **License**: *CC BY-SA: Attribution-ShareAlike*
- photosynthesis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/photosynthesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- eukaryotic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/eukaryotic. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, The Nucleus and DNA Replication. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46073/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Cells. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m45432/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Cells. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m45432/latest/. **License**: *CC BY: Attribution*

This page titled 3.5B: Characteristics of Eukaryotic Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6: ORGANELLES



- Topic hierarchy

3.6A: The Plasma Membrane and the Cytoplasm

3.6B: The Endoplasmic Reticulum

3.6C: The Golgi Apparatus

3.6D: Lysosomes

3.6E: Peroxisomes

3.6F: Mitochondria

This page titled 3.6: Organelles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6A: The Plasma Membrane and the Cytoplasm

The plasma membrane is made up of a phospholipid bilayer that regulates the concentration of substances that can permeate a cell.

Learning Objectives

- Explain the structure and purpose of the plasma membrane of a cell

Key Points


- All eukaryotic cells have a surrounding plasma membrane, which is also known as the cell membrane.
- The plasma membrane is made up by a phospholipid bilayer with embedded proteins that separates the internal contents of the cell from its surrounding environment.
- Only relatively small, non- polar materials can easily move through the lipid bilayer of the plasma membrane.
- Passive transport is the movement of substances across the membrane that does not require the use of energy while active transport is the movement of substances across the membrane using energy.
- Osmosis is the diffusion of water through a semi- permeable membrane down its concentration gradient; this occurs when there is an imbalance of solutes outside of a cell compared to the inside the cell.

Key Terms

- **phospholipid**: Any lipid consisting of a diglyceride combined with a phosphate group and a simple organic molecule such as choline or ethanolamine; they are important constituents of biological membranes
- **hypertonic**: having a greater osmotic pressure than another
- **hypotonic**: Having a lower osmotic pressure than another; a cell in this environment causes water to enter the cell, causing it to swell.


The Plasma Membrane

Despite differences in structure and function, all living cells in multicellular organisms have a surrounding plasma membrane (also known as the cell membrane). As the outer layer of your skin separates your body from its environment, the plasma membrane separates the inner contents of a cell from its exterior environment. The plasma membrane can be described as a phospholipid bilayer with embedded proteins that controls the passage of organic molecules, ions, water, and oxygen into and out of the cell. Wastes (such as carbon dioxide and ammonia) also leave the cell by passing through the membrane.



Eukaryotic Plasma Membrane: The eukaryotic plasma membrane is a phospholipid bilayer with proteins and cholesterol embedded in it.

The cell membrane is an extremely pliable structure composed primarily of two adjacent sheets of phospholipids. Cholesterol, also present, contributes to the fluidity of the membrane. A single phospholipid molecule consists of a polar phosphate "head," which is hydrophilic, and a non-polar lipid "tail," which is hydrophobic. Unsaturated fatty acids result in kinks in the hydrophobic tails. The phospholipid bilayer consists of two phospholipids arranged tail to tail. The hydrophobic tails associate with one another, forming the interior of the membrane. The polar heads contact the fluid inside and outside of the cell.



Phospholipid Bilayer: The phospholipid bilayer consists of two adjacent sheets of phospholipids, arranged tail to tail. The hydrophobic tails associate with one another, forming the interior of the membrane. The polar heads contact the fluid inside and outside of the cell.


The plasma membrane's main function is to regulate the concentration of substances inside the cell. These substances include ions such as Ca^{++}, Na^{+}, K^{+}, and Cl^{-}; nutrients including sugars, fatty acids, and amino acids; and waste products, particularly carbon dioxide (CO_2), which must leave the cell.

The membrane's lipid bilayer structure provides the cell with access control through permeability. The phospholipids are tightly packed together, while the membrane has a hydrophobic interior. This structure causes the membrane to be selectively permeable. A membrane that has selective permeability allows only substances meeting certain criteria to pass through it

unaided. In the case of the plasma membrane, only relatively small, non-polar materials can move through the lipid bilayer (remember, the lipid tails of the membrane are nonpolar). Some examples of these materials are other lipids, oxygen and carbon dioxide gases, and alcohol. However, water-soluble materials—such as glucose, amino acids, and electrolytes—need some assistance to cross the membrane because they are repelled by the hydrophobic tails of the phospholipid bilayer.

Transport Across the Membrane

All substances that move through the membrane do so by one of two general methods, which are categorized based on whether or not energy is required. Passive (non-energy requiring) transport is the movement of substances across the membrane without the expenditure of cellular energy. During this type of transport, materials move by simple diffusion or by facilitated diffusion through the membrane, down their concentration gradient. Water passes through the membrane in a diffusion process called osmosis. Osmosis is the diffusion of water through a semi-permeable membrane down its concentration gradient. It occurs when there is an imbalance of solutes outside of a cell versus inside the cell. The solution that has the higher concentration of solutes is said to be hypertonic and the solution that has the lower concentration of solutes is said to be hypotonic. Water molecules will diffuse out of the hypotonic solution and into the hypertonic solution (unless acted upon by hydrostatic forces).



Osmosis: Osmosis is the diffusion of water through a semipermeable membrane down its concentration gradient. If a membrane is permeable to water, though not to a solute, water will equalize its own concentration by diffusing to the side of lower water concentration (and thus the side of higher solute concentration). In the beaker on the left, the solution on the right side of the membrane is hypertonic.

In contrast to passive transport, active (energy-requiring) transport is the movement of substances across the membrane using energy from adenosine triphosphate (ATP). The energy is expended to assist material movement across the membrane in a direction against their concentration gradient. Active transport may take place with the help of protein pumps or through the use of vesicles. Another form of this type of transport is endocytosis, where a cell envelopes extracellular materials using its cell membrane. The opposite process is known as exocytosis. This is where a cell exports material using vesicular transport.

Cytoplasm

The cell's plasma membrane also helps contain the cell's cytoplasm, which provides a gel-like environment for the cell's organelles. The cytoplasm is the location for most cellular processes, including metabolism, protein folding, and internal transportation.

This page titled 3.6A: The Plasma Membrane and the Cytoplasm is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6B: The Endoplasmic Reticulum

The endoplasmic reticulum is an organelle that is responsible for the synthesis of lipids and the modification of proteins.

Learning Objectives

- Describe the structure of the endoplasmic reticulum and its role in synthesis and metabolism

Key Points

- If the endoplasmic reticulum (ER) has ribosomes attached to it, it is called rough ER; if it does not, then it is called smooth ER.
- The proteins made by the rough endoplasmic reticulum are for use outside of the cell.
- Functions of the smooth endoplasmic reticulum include synthesis of carbohydrates, lipids, and steroid hormones; detoxification of medications and poisons; and storage of calcium ions.


Key Terms

- **lumen**: The cavity or channel within a tube or tubular organ.
- **reticulum**: A network

The Endoplasmic Reticulum

The endoplasmic reticulum (ER) is a series of interconnected membranous sacs and tubules that collectively modifies proteins and synthesizes lipids. However, these two functions are performed in separate areas of the ER: the rough ER and the smooth ER. The hollow portion of the ER tubules is called the lumen or cisternal space. The membrane of the ER, which is a phospholipid bilayer embedded with proteins, is continuous with the nuclear envelope.

Rough ER



Rough Endoplasmic Reticulum: This transmission electron micrograph shows the rough endoplasmic reticulum and other organelles in a pancreatic cell.

The rough endoplasmic reticulum (RER) is so named because the ribosomes attached to its cytoplasmic surface give it a studded appearance when viewed through an electron microscope. Ribosomes transfer their newly synthesized proteins into the lumen of the RER where they undergo structural modifications, such as folding or the acquisition of side chains. These modified proteins will be incorporated into cellular membranes—the membrane of the ER or those of other organelles —or secreted from the

cell (such as protein hormones, enzymes). The RER also makes phospholipids for cellular membranes. If the phospholipids or modified proteins are not destined to stay in the RER, they will reach their destinations via transport vesicles that bud from the RER's membrane. Since the RER is engaged in modifying proteins (such as enzymes, for example) that will be secreted from the cell, the RER is abundant in cells that secrete proteins. This is the case with cells of the liver, for example.

Smooth ER

The smooth endoplasmic reticulum (SER) is continuous with the RER but has few or no ribosomes on its cytoplasmic surface. Functions of the SER include synthesis of carbohydrates, lipids, and steroid hormones; detoxification of medications and poisons; and storage of calcium ions. In muscle cells, a specialized SER called the sarcoplasmic reticulum is responsible for storage of the calcium ions that are needed to trigger the coordinated contractions of the muscle cells.

This page titled 3.6B: The Endoplasmic Reticulum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6C: The Golgi Apparatus

The Golgi apparatus sorts and packages materials before they leave the cell to ensure they arrive at the proper destination.

Learning Objectives

- Describe the structure of the Golgi apparatus and its role in protein modification and secretion

Key Points


- The Golgi apparatus is a series of flattened sacs that sort and package cellular materials.
- The Golgi apparatus has a cis face on the ER side and a trans face opposite of the ER.
- The *trans* face secretes the materials into vesicles, which then fuse with the cell membrane for release from the cell.

Key Terms

- **vesicle**: A membrane-bound compartment found in a cell.

The Golgi Apparatus

We have already mentioned that vesicles can bud from the ER and transport their contents elsewhere, but where do the vesicles go? Before reaching their final destination, the lipids or proteins within the transport vesicles still need to be sorted, packaged, and tagged so that they wind up in the right place. Sorting, tagging, packaging, and distribution of lipids and proteins takes place in the Golgi apparatus (also called the Golgi body), a series of flattened membranes.



The Golgi apparatus sorts and packages cellular products: The Golgi apparatus in this white blood cell is visible as a stack of semicircular, flattened rings in the lower portion of the image. Several vesicles can be seen near the Golgi apparatus.

The receiving side of the Golgi apparatus is called the cis face. The opposite side is called the trans face. The transport vesicles that formed from the ER travel to the cis face, fuse with it, and empty their contents into the lumen of the Golgi apparatus. As the proteins and lipids travel through the Golgi, they undergo further modifications that allow them to be sorted. The most frequent modification is the addition of short chains of sugar molecules. These newly-modified proteins and lipids are then tagged with phosphate groups or other small molecules so that they can be routed to their proper destinations.

Finally, the modified and tagged proteins are packaged into secretory vesicles that bud from the trans face of the Golgi. While some of these vesicles deposit their contents into other parts of the cell where they will be used, other secretory vesicles fuse with the plasma membrane and release their contents outside the cell.

In another example of form following function, cells that engage in a great deal of secretory activity (such as cells of the salivary glands that secrete digestive enzymes or cells of the immune system that secrete antibodies) have an abundance of Golgi. In plant cells, the Golgi apparatus has the additional role of synthesizing polysaccharides, some of which are incorporated into the cell wall and some of which are used in other parts of the cell.

This page titled 3.6C: The Golgi Apparatus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6D: Lysosomes

Lysosomes are organelles that digest macromolecules, repair cell membranes, and respond to foreign substances entering the cell.

Learning Objectives

- Describe how lysosomes function as the cell's waste disposal system

Key Points


- Lysosomes breakdown/digest macromolecules (carbohydrates, lipids, proteins, and nucleic acids), repair cell membranes, and respond against foreign substances such as bacteria, viruses and other antigens.
- Lysosomes contain enzymes that break down the macromolecules and foreign invaders.
- Lysosomes are composed of lipids and proteins, with a single membrane covering the internal enzymes to prevent the lysosome from digesting the cell itself.
- Lysosomes are found in all animal cells, but are rarely found within plant cells due to the tough cell wall surrounding a plant cell that keeps out foreign substances.

Key Terms

- **enzyme**: a globular protein that catalyses a biological chemical reaction
- **lysosome**: An organelle found in all types of animal cells which contains a large range of digestive enzymes capable of splitting most biological macromolecules.

A lysosome has three main functions: the breakdown/digestion of macromolecules (carbohydrates, lipids, proteins, and nucleic acids), cell membrane repairs, and responses against foreign substances such as bacteria, viruses and other antigens. When food is eaten or absorbed by the cell, the lysosome releases its enzymes to break down complex molecules including sugars and proteins into usable energy needed by the cell to survive. If no food is provided, the lysosome's enzymes digest other organelles within the cell in order to obtain the necessary nutrients.

In addition to their role as the digestive component and organelle-recycling facility of animal cells, lysosomes are considered to be parts of the endomembrane system. Lysosomes also use their hydrolytic enzymes to destroy pathogens (disease-causing organisms) that might enter the cell. A good example of this occurs in a group of white blood cells called macrophages, which are part of your body's immune system. In a process known as phagocytosis or endocytosis, a section of the plasma membrane of the macrophage invaginates (folds in) and engulfs a pathogen. The invaginated section, with the pathogen inside, then pinches itself off from the plasma membrane and becomes a vesicle. The vesicle fuses with a lysosome. The lysosome's hydrolytic enzymes then destroy the pathogen.



Lysosomes digest foreign substances that might harm the cell: A macrophage has engulfed (phagocytized) a potentially pathogenic bacterium and then fuses with a lysosomes within the cell to destroy the pathogen. Other organelles are present in the cell but for simplicity are not shown.

A lysosome is composed of lipids, which make up the membrane, and proteins, which make up the enzymes within the membrane. Usually, lysosomes are between 0.1 to 1.2μm, but the size varies based on the cell type. The general structure of a lysosome consists of a collection of enzymes surrounded by a single-layer membrane. The membrane is a crucial aspect of its structure because without it the enzymes within the lysosome that are used to breakdown foreign substances would leak out and digest the entire cell, causing it to die.

Lysosomes are found in nearly every animal-like eukaryotic cell. They are so common in animal cells because, when animal cells take in or absorb food, they need the enzymes found in lysosomes in order to digest and use the food for energy. On the other hand, lysosomes are not commonly-found in plant cells. Lysosomes are not needed in plant cells because they have cell walls that are tough enough to keep the large/foreign substances that lysosomes would usually digest out of the cell.

This page titled 3.6D: Lysosomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6E: Peroxisomes

Peroxisomes neutralize harmful toxins and carry out lipid metabolism and oxidation reactions that break down fatty acids and amino acids.

Learning Objectives

- Name the various functions that peroxisomes perform inside the cell

Key Points


- Lipid metabolism and chemical detoxification are important functions of peroxisomes.
- Peroxisomes are responsible for oxidation reactions that break down fatty acids and amino acids.
- Peroxisomes oversee reactions that neutralize free radicals, which cause cellular damage and cell death.
- Peroxisomes chemically neutralize poisons through a process that produces large amounts of toxic H_2O_2, which is then converted into water and oxygen.
- The liver is the organ primarily responsible for detoxifying the blood before it travels throughout the body; as a result, liver cells contain large amounts of peroxisomes.

Key Terms

- **enzyme**: a globular protein that catalyses a biological chemical reaction
- **free radical**: Any molecule, ion or atom that has one or more unpaired electrons; they are generally highly reactive and often only occur as transient species.

Peroxisomes

A type of organelle found in both animal cells and plant cells, a peroxisome is a membrane-bound cellular organelle that contains mostly enzymes. Peroxisomes perform important functions, including lipid metabolism and chemical detoxification. They also carry out oxidation reactions that break down fatty acids and amino acids.



Peroxisomes: Peroxisomes are membrane-bound organelles that contain an abundance of enzymes for detoxifying harmful

substances and lipid metabolism.

In contrast to the digestive enzymes found in lysosomes, the enzymes within peroxisomes serve to transfer hydrogen atoms from various molecules to oxygen, producing hydrogen peroxide (H_2O_2). In this way, peroxisomes neutralize poisons, such as alcohol, that enter the body. In order to appreciate the importance of peroxisomes, it is necessary to understand the concept of reactive oxygen species.

Reactive oxygen species (ROS), such as peroxides and free radicals, are the highly-reactive products of many normal cellular processes, including the mitochondrial reactions that produce ATP and oxygen metabolism. Examples of ROS include the hydroxyl radical OH, H_2O_2, and superoxide (O_{-2}). Some ROS are important for certain cellular functions, such as cell signaling processes and immune responses against foreign substances. Many ROS, however, are harmful to the body. Free radicals are reactive because they contain free unpaired electrons; they can easily oxidize other molecules throughout the cell, causing cellular damage and even cell death. Free radicals are thought to play a role in many destructive processes in the body, from cancer to coronary artery disease.

Peroxisomes oversee reactions that neutralize free radicals. They produce large amounts of the toxic H_2O_2 in the process, but contain enzymes that convert H_2O_2 into water and oxygen. These by-products are then safely released into the cytoplasm. Like miniature sewage treatment plants, peroxisomes neutralize harmful toxins so that they do not cause damage in the cells. The liver is the organ primarily responsible for detoxifying the blood before it travels throughout the body; liver cells contain an exceptionally high number of peroxisomes.

This page titled 3.6E: Peroxisomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.6F: Mitochondria

Mitochondria are organelles that are responsible for making adenosine triphosphate (ATP), the cell's main energy-carrying molecule.

Learning Objectives

- Explain the role of the mitochondria.

Key Points

- Mitochondria contain their own ribosomes and DNA; combined with their double membrane, these features suggest that they might have once been free-living prokaryotes that were engulfed by a larger cell.
- Mitochondria have an important role in cellular respiration through the production of ATP, using chemical energy found in glucose and other nutrients.
- Mitochondria are also responsible for generating clusters of iron and sulfur, which are important cofactors of many enzymes.


Key Terms

- **alpha-proteobacteria**: A taxonomic class within the phylum Proteobacteria — the phototropic proteobacteria.
- **adenosine triphosphate**: a multifunctional nucleoside triphosphate used in cells as a coenzyme, often called the "molecular unit of energy currency" in intracellular energy transfer
- **cofactor**: an inorganic molecule that is necessary for an enzyme to function

One of the major features distinguishing prokaryotes from eukaryotes is the presence of mitochondria. Mitochondria are double-membraned organelles that contain their own ribosomes and DNA. Each membrane is a phospholipid bilayer embedded with proteins. Eukaryotic cells may contain anywhere from one to several thousand mitochondria, depending on the cell's level of energy consumption. Each mitochondrion measures 1 to 10 micrometers (or greater) in length and exists in the cell as an organelle that can be ovoid to worm-shaped to intricately branched.

Mitochondria Structure

Most mitochondria are surrounded by two membranes, which would result when one membrane-bound organism was engulfed into a vacuole by another membrane-bound organism. The mitochondrial inner membrane is extensive and involves substantial infoldings called cristae that resemble the textured, outer surface of alpha-proteobacteria. The matrix and inner membrane are rich with the enzymes necessary for aerobic respiration.



Mitochondrial structure: This electron micrograph shows a mitochondrion as viewed with a transmission electron microscope. This organelle has an outer membrane and an inner membrane. The inner membrane contains folds, called cristae, which increase its surface area. The space between the two membranes is called the intermembrane space, and the space inside the inner membrane is called the mitochondrial matrix. ATP synthesis takes place on the inner membrane.

Mitochondria have their own (usually) circular DNA chromosome that is stabilized by attachments to the inner membrane and carries genes similar to genes expressed by alpha-proteobacteria. Mitochondria also have special ribosomes and transfer RNAs that resemble these components in prokaryotes. These features all support the hypothesis that mitochondria were once free-living prokaryotes.

Mitochondria Function

Mitochondria are often called the “powerhouses” or “energy factories” of a cell because they are responsible for making adenosine triphosphate (ATP), the cell’s main energy-carrying molecule. ATP represents the short-term stored energy of the cell. Cellular respiration is the process of making ATP using the chemical energy found in glucose and other nutrients. In mitochondria, this process uses oxygen and produces carbon dioxide as a waste product. In fact, the carbon dioxide that you exhale with every breath comes from the cellular reactions that produce carbon dioxide as a by-product.

It is important to point out that muscle cells have a very high concentration of mitochondria that produce ATP. Your muscle cells need a lot of energy to keep your body moving. When your cells don’t get enough oxygen, they do not make a lot of ATP. Instead, the small amount of ATP they make in the absence of oxygen is accompanied by the production of lactic acid.

In addition to the aerobic generation of ATP, mitochondria have several other metabolic functions. One of these functions is to generate clusters of iron and sulfur that are important cofactors of many enzymes. Such functions are often associated with the reduced mitochondrion-derived organelles of anaerobic eukaryotes.

Origins of Mitochondria

There are two hypotheses about the origin of mitochondria: endosymbiotic and autogenous, but the most accredited theory at present is endosymbiosis. The endosymbiotic hypothesis suggests mitochondria were originally prokaryotic cells, capable of implementing oxidative mechanisms. These prokaryotic cells may have been engulfed by a eukaryote and became endosymbionts living inside the eukaryote.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Laura Martin, Discovering the Structure of the Plasma Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/col10470/latest/. **License**: ***CC BY: Attribution***
- phospholipid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/phospholipid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypertonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypertonic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypotonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypotonic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- reticulum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/reticulum. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lumen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lumen. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest..._04_02_new.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- vesicle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vesicle. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest..._04_02_new.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_03.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Lysosome. **Provided by**: epiehonorsbiology Wikispace. **Located at**: http://epiehonorsbiology.wikispaces.com/Lysosome. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lysosome. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lysosome. **License**: ***CC BY-SA: Attribution-ShareAlike***
- enzyme. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/enzyme. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: ***CC BY: Attribution***

- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest..._04_02_new.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 21, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, The Cytoplasm and Cellular Organelles. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46023/latest/. **License**: *CC BY: Attribution*
- enzyme. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/enzyme. **License**: *CC BY-SA: Attribution-ShareAlike*
- free radical. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/free_radical. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest..._04_02_new.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cytoplasm and Cellular Organelles. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46023/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest/?collection=col11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Origins. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44614/latest/. **License**: *CC BY: Attribution*
- cofactor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cofactor. **License**: *CC BY-SA: Attribution-ShareAlike*
- adenosine triphosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adenosine%20triphosphate. **License**: *CC BY-SA: Attribution-ShareAlike*
- alpha-proteobacteria. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/alpha-proteobacteria. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mitochondrion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mitochondrion. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest...e_04_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest..._04_02_new.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Endomembrane System and Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44435/latest...e_04_04_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cytoplasm and Cellular Organelles. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46023/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Eukaryotic Cells. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44407/latest/Figure_04_03_07.jpg. **License**: *CC BY: Attribution*

This page titled 3.6F: Mitochondria is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.7: THE CYTOSKELETON



- Topic hierarchy

3.7A: Intermediate Filaments and Microtubules

3.7B: Microfilaments

This page titled 3.7: The Cytoskeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.7A: Intermediate Filaments and Microtubules

Microtubules are part of the cell's cytoskeleton, helping the cell resist compression, move vesicles, and separate chromosomes at mitosis.

Learning Objectives

- Describe the roles of microtubules as part of the cell's cytoskeleton

Key Points


- Microtubules help the cell resist compression, provide a track along which vesicles can move throughout the cell, and are the components of cilia and flagella.
- Cilia and flagella are hair-like structures that assist with locomotion in some cells, as well as line various structures to trap particles.
- The structures of cilia and flagella are a "9+2 array," meaning that a ring of nine microtubules is surrounded by two more microtubules.
- Microtubules attach to replicated chromosomes during cell division and pull them apart to opposite ends of the pole, allowing the cell to divide with a complete set of chromosomes in each daughter cell.

Key Terms

- **microtubule**: Small tubes made of protein and found in cells; part of the cytoskeleton
- **flagellum**: a flagellum is a lash-like appendage that protrudes from the cell body of certain prokaryotic and eukaryotic cells
- **cytoskeleton**: A cellular structure like a skeleton, contained within the cytoplasm.

Microtubules


As their name implies, microtubules are small hollow tubes. Microtubules, along with microfilaments and intermediate filaments, come under the class of organelles known as the cytoskeleton. The cytoskeleton is the framework of the cell which forms the structural supporting component. Microtubules are the largest element of the cytoskeleton. The walls of the microtubule are made of polymerized dimers of α-tubulin and β-tubulin, two globular proteins. With a diameter of about 25 nm, microtubules are the widest components of the cytoskeleton. They help the cell resist compression, provide a track along which vesicles move through the cell, and pull replicated chromosomes to opposite ends of a dividing cell. Like microfilaments, microtubules can dissolve and reform quickly.



Micrtubule Structure: Microtubules are hollow, with walls consisting of 13 polymerized dimers of α-tubulin and β-tubulin (right image). The left image shows the molecular structure of the tube.

Microtubules are also the structural elements of flagella, cilia, and centrioles (the latter are the two perpendicular bodies of the centrosome). In animal cells, the centrosome is the microtubule-organizing center. In eukaryotic cells, flagella and cilia are quite different structurally from their counterparts in prokaryotes.

Intermediate Filaments




Stained Keratin Intermediate filaments: Keratin cytoskeletal intermediate filaments are concentrated around the edge of the cells and merge into the surface membrane. This network of intermediate filaments from cell to cell holds together tissues like skin.

Intermediate filaments (IFs) are cytoskeletal components found in animal cells. They are composed of a family of related proteins sharing common structural and sequence features. Intermediate filaments have an average diameter of 10 nanometers, which is between that of 7 nm actin (microfilaments), and that of 25 nm microtubules, although they were initially designated 'intermediate' because their average diameter is between those of narrower microfilaments (actin) and wider myosin filaments found in muscle cells. Intermediate filaments contribute to cellular structural elements and are often crucial in holding together tissues like skin.

Flagella and Cilia

Flagella (singular = flagellum) are long, hair-like structures that extend from the plasma membrane and are used to move an entire cell (for example, sperm, *Euglena*). When present, the cell has just one flagellum or a few flagella. When cilia (singular = cilium) are present, however, many of them extend along the entire surface of the plasma membrane. They are short, hair-like structures that are used to move entire cells (such as paramecia) or substances along the outer surface of the cell (for example, the cilia of cells lining the Fallopian tubes that move the ovum toward the uterus, or cilia lining the cells of the respiratory tract that trap particulate matter and move it toward your nostrils).

Despite their differences in length and number, flagella and cilia share a common structural arrangement of microtubules called a "9 + 2 array." This is an appropriate name because a single flagellum or cilium is made of a ring of nine microtubule doublets surrounding a single microtubule doublet in the center.



Microtubules are the structural component of flagella: This transmission electron micrograph of two flagella shows the 9 + 2 array of microtubules: nine microtubule doublets surround a single microtubule doublet.

This page titled 3.7A: Intermediate Filaments and Microtubules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.7B: Microfilaments

Microfilaments, which are the thinnest part of the cytoskeleton, are used to give shape to the cell and support all of its internal parts.

Learning Objectives

- Describe the structure and function of microfilaments

Key Points


- Microfilaments assist with cell movement and are made of a protein called actin.
- Actin works with another protein called myosin to produce muscle movements, cell division, and cytoplasmic streaming.
- Microfilaments keep organelles in place within the cell.

Key Terms

- **actin**: A globular structural protein that polymerizes in a helical fashion to form an actin filament (or microfilament).
- **filamentous**: Having the form of threads or filaments
- **myosin**: a large family of motor proteins found in eukaryotic tissues, allowing mobility in muscles

Microfilaments

If all the organelles were removed from a cell, the plasma membrane and the cytoplasm would not be the only components left. Within the cytoplasm there would still be ions and organic molecules, plus a network of protein fibers that help maintain the shape of the cell, secure some organelles in specific positions, allow cytoplasm and vesicles to move within the cell, and enable unicellular organisms to move independently. This network of protein fibers is known as the cytoskeleton. There are three types of fibers within the cytoskeleton: microfilaments, intermediate filaments, and microtubules. Of the three types of protein fibers in the cytoskeleton, microfilaments are the narrowest. They function in cellular movement, have a diameter of about 7 nm, and are made of two intertwined strands of a globular protein called actin. For this reason, microfilaments are also known as actin filaments.



Microfilaments are the thinnest component of the cytoskeleton.: Microfilaments are made of two intertwined strands of actin.

Actin is powered by ATP to assemble its filamentous form, which serves as a track for the movement of a motor protein called myosin. This enables actin to engage in cellular events requiring motion such as cell division in animal cells and cytoplasmic streaming, which is the circular movement of the cell cytoplasm in plant cells. Actin and myosin are plentiful in muscle cells. When your actin and myosin filaments slide past each other, your muscles contract.

Microfilaments also provide some rigidity and shape to the cell. They can depolymerize (disassemble) and reform quickly, thus enabling a cell to change its shape and move. White blood cells (your body's infection-fighting cells) make good use of this ability. They can move to the site of an infection and engulf the pathogen.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 21, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Cell Biology/Cytoskeleton/Microtubules. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Cell_Bi...n/Microtubules**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intermediate Filaments. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intermediate_filament**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cytoskeleton. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cytoskeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- microtubule. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/microtubule**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- flagellum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/flagellum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithelial-cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ep...lial-cells.jpg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, The Cytoskeleton. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest...04_05_04ab.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, The Cytoskeleton. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest...e_04_05_05.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax, Biology. May 18, 2016. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.9**. **License**: ***CC BY: Attribution***
- actin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/actin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myosin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myosin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- filamentous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/filamentous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithelial-cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Epithelial-cells.jpg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, The Cytoskeleton. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest/Figure_04_05_04ab.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, The Cytoskeleton. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest/Figure_04_05_05.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, The Cytoskeleton. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44412/latest/Figure_04_05_02.jpg**. **License**: ***CC BY: Attribution***

This page titled 3.7B: Microfilaments is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.8: EXTERNAL CELLULAR COMPONENTS



- Topic hierarchy

3.8A: Extracellular Matrix of Animal Cells

3.8B: Intercellular Junctions

This page titled 3.8: External Cellular Components is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.8A: Extracellular Matrix of Animal Cells

The extracellular matrix of animal cells holds cells together to form a tissue and allow tissues to communicate with each other.

Learning Objectives

- Explain the role of the extracellular matrix in animal cells

Key Points


- The extracellular matrix of animal cells is made up of proteins and carbohydrates.
- Cell communication within tissue and tissue formation are main functions of the extracellular matrix of animal cells.
- Tissue communication is kick-started when a molecule within the matrix binds a receptor; the end results are conformational changes that induce chemical signals that ultimately change activities within the cell.

Key Terms

- **collagen**: Any of more than 28 types of glycoprotein that forms elongated fibers, usually found in the extracellular matrix of connective tissue.
- **proteoglycan**: Any of many glycoproteins that have heteropolysaccharide side chains
- **extracellular matrix**: All the connective tissues and fibres that are not part of a cell, but rather provide support.

Extracellular Matrix of Animal Cells

Most animal cells release materials into the extracellular space. The primary components of these materials are proteins. Collagen is the most abundant of the proteins. Its fibers are interwoven with carbohydrate-containing protein molecules called proteoglycans. Collectively, these materials are called the extracellular matrix. Not only does the extracellular matrix hold the cells together to form a tissue, but it also allows the cells within the tissue to communicate with each other.



The Extracellular Matrix: The extracellular matrix consists of a network of proteins and carbohydrates.

How does this cell communication occur? Cells have protein receptors on the extracellular surfaces of their plasma membranes.

When a molecule within the matrix binds to the receptor, it changes the molecular structure of the receptor. The receptor, in turn, changes the conformation of the microfilaments positioned just inside the plasma membrane. These conformational changes induce chemical signals inside the cell that reach the nucleus and turn “on” or “off” the transcription of specific sections of DNA. This affects the production of associated proteins, thus changing the activities within the cell.

An example of the role of the extracellular matrix in cell communication can be seen in blood clotting. When the cells lining a blood vessel are damaged, they display a protein receptor called tissue factor. When a tissue factor binds with another factor in the extracellular matrix, it causes platelets to adhere to the wall of the damaged blood vessel and stimulates the adjacent smooth muscle cells in the blood vessel to contract (thus constricting the blood vessel). Subsequently, a series of steps are initiated which then prompt the platelets to produce clotting factors.

This page titled 3.8A: Extracellular Matrix of Animal Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

3.8B: Intercellular Junctions

Intercellular junctions provide plant and animal cells with the ability to communicate through direct contact.

Learning Objectives

- Describe the purpose of intercellular junctions in the structure of cells

Key Points

- Plasmodesmata are intercellular junctions between plant cells that enable the transportation of materials between cells.
- A tight junction is a watertight seal between two adjacent animal cells, which prevents materials from leaking out of cells.
- Desmosomes connect adjacent cells when cadherins in the plasma membrane connect to intermediate filaments.
- Similar to plasmodesmata, gap junctions are channels between adjacent cells that allow for the transport of ions, nutrients, and other substances.

Key Terms

- **plasmodesma**: A microscopic channel traversing the cell walls of plant cells and some algal cells, enabling transport and communication between them.
- **connexon**: An assembly of six connexins forming a bridge called a gap junction between the cytoplasms of two adjacent cells.
- **occludin**: Together with the claudin group of proteins, it is the main component of the tight junctions.


Intercellular Junctions

The extracellular matrix allows cellular communication within tissues through conformational changes that induce chemical signals, which ultimately transform activities within the cell. However, cells are also capable of communicating with each other via direct contact through intercellular junctions.

There are some differences in the ways that plant and animal cells communicate directly. Plasmodesmata are junctions between plant cells, whereas animal cell contacts are carried out through tight junctions, gap junctions, and desmosomes.

Junctions in Plant Cells


In general, long stretches of the plasma membranes of neighboring plant cells cannot touch one another because they are separated by the cell wall that surrounds each cell. How then can a plant transfer water and other soil nutrients from its roots, through its stems, and to its leaves? This transport primarily uses the vascular tissues (xylem and phloem); however, there are also structural modifications called plasmodesmata (singular: plasmodesma) that facilitate direct communication in plant cells. Plasmodesmata are numerous channels that pass between cell walls of adjacent plant cells and connect their cytoplasm; thereby, enabling materials to be transported from cell to cell, and thus throughout the plant.



Plasmodesmata: A plasmodesma is a channel between the cell walls of two adjacent plant cells. Plasmodesmata allow materials to pass from the cytoplasm of one plant cell to the cytoplasm of an adjacent cell.


Junctions in Animal Cells

Communication between animal cells can be carried out through three types of junctions. The first, a tight junction, is a watertight seal between two adjacent animal cells. The cells are held tightly against each other by proteins (predominantly two proteins called claudins and occludins). This tight adherence prevents materials from leaking between the cells. These junctions are typically found in epithelial tissues that line internal organs and cavities and comprise most of the skin. For example, the tight junctions of the epithelial cells lining your urinary bladder prevent urine from leaking out into the extracellular space.



Tight Junctions: Tight junctions form watertight connections between adjacent animal cells. Proteins create tight junction adherence.

Also found only in animal cells are desmosomes, the second type of intercellular junctions in these cell types. Desmosomes act like spot welds between adjacent epithelial cells, connecting them. Short proteins called cadherins in the plasma membrane connect to intermediate filaments to create desmosomes. The cadherins join two adjacent cells together and maintain the cells in a sheet-like formation in organs and tissues that stretch, such as the skin, heart, and muscles.



Desmosomes: A desmosome forms a very strong spot weld between cells. It is created by the linkage of cadherins and intermediate filaments.

Lastly, similar to plasmodesmata in plant cells, gap junctions are the third type of direct junction found within animal cells. These junctions are channels between adjacent cells that allow for the transport of ions, nutrients, and other substances that enable cells to communicate. Structurally, however, gap junctions and plasmodesmata differ. Gap junctions develop when a set of six proteins (called connexins) in the plasma membrane arrange themselves in an elongated doughnut-like configuration called a connexon. When the pores ("doughnut holes") of connexons in adjacent animal cells align, a channel between the two cells forms. Gap junctions are particularly important in cardiac muscle. The electrical signal for the muscle to contract is passed efficiently through gap junctions, which allows the heart muscle cells to contract in tandem.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- extracellular matrix. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/extracellular_matrix. **License**: ***CC BY-SA: Attribution-ShareAlike***
- proteoglycan. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/proteoglycan. **License**: ***CC BY-SA: Attribution-ShareAlike***
- collagen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/collagen. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Connections between Cells and Cellular Activities. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...e_04_06_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- plasmodesma. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/plasmodesma. **License**: ***CC BY-SA: Attribution-ShareAlike***
- occludin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/occludin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- connexon. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/connexon. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- OpenStax College, Connections between Cells and Cellular Activities. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...e_04_06_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Connections between Cells and Cellular Activities. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...e_04_06_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Connections between Cells and Cellular Activities. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...e_04_06_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Connections between Cells and Cellular Activities. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44413/latest...e_04_06_04.jpg. **License**: *CC BY: Attribution*

This page titled 3.8B: Intercellular Junctions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4: ORGANIZATION AT THE TISSUE LEVEL

- Topic hierarchy

4.1: Introduction to Tissues

4.1A: Tissues in Levels of Organization

4.2: Epithelial Tissue

4.2A: Characteristics of Epithelial Tissue

4.2B: Types of Epithelial Tissue

4.3: Connective Tissue

4.3A: Characteristics of Connective Tissue

4.3B: Types of Connective Tissue

4.4: Nervous Tissue

4.4A: Characteristics of Nervous Tissue

4.4B: Types of Nervous Tissue

4.5: Muscular Tissue

4.5A: Characteristics of Muscle Tissue

4.6: Tissue Membranes

4.6A: Epithelial Membranes

4.6B: Synovial Membranes

This page titled 4: Organization at the Tissue Level is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.1: INTRODUCTION TO TISSUES



- Topic hierarchy

4.1A: Tissues in Levels of Organization

This page titled 4.1: Introduction to Tissues is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.1A: Tissues in Levels of Organization

The human body is organized at several levels of scale that can each be examined.

Learning Objectives

- Characterize where tissues fall in levels of organization

Key Points

- The human body has many levels of structural organization: atoms, cells, tissues, organs, and organ system.
- The simplest level is the chemical level, which includes tiny building blocks such as atoms.
- Cells are the smallest functional units of life.
- Tissues are groups of similar cells that have a common function.
- An organ is a structure that is composed of at least two or more tissue types and performs a specific set of functions for the body.
- Many organs working together to accomplish a common purpose is called an organ system.

Tissues in the Human Body

The human body has many levels of structural organization. The simplest level is the chemical level, which includes tiny building blocks such as atoms. Cells are the smallest functional units of life. The simplest living creatures are single cell creatures, but in complex life forms, such as human beings, cells also exist in the tissue level.

Tissues are groups of similar cells that have a common function. The four basic tissue types are epithelial, muscle, connective, and nervous tissue. Each tissue type has a characteristic role in the body:


1. Epithelium covers the body surface and lines body cavities.
2. Muscle provides movement.
3. Connective tissue supports and protects body organs.
4. Nervous tissue provides a means of rapid internal communication by transmitting electrical impulses.

Organs: Made of Tissues

An organ is a structure that is composed of at least two or more tissue types and performs a specific set of functions for the body. The liver, stomach, brain, and blood are all different organs and perform different functions. Each organ is a specialized functional center responsible for a specific function of the body.

At the organ level, complex functions become possible because of the specialized activities of various tissues. Most organs contain more than one tissue type. For example, the stomach consists of smooth muscle tissue for churning movement while it is innervated, but it is also supplied by blood, which is a connective tissue.

The next level is the organ system level. Many organs working together to accomplish a common purpose create an organ system. For example, the heart and the blood vessels of the cardiovascular system circulate blood and transport oxygen and nutrients to all the body cells.



Levels of Organization: Molecules form cells. Cells form tissues, and tissues form organs. Organs that fulfill related functions are called organ systems. An organism is made up of interconnected organ systems.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Anatomy & Physiology. **Provided by**: OpenStax. **Located at**: http://cnx.org/content/col11496/latest/. **License**: ***CC BY: Attribution***

This page titled 4.1A: Tissues in Levels of Organization is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.2: EPITHELIAL TISSUE



- Topic hierarchy

4.2A: Characteristics of Epithelial Tissue

4.2B: Types of Epithelial Tissue

This page titled 4.2: Epithelial Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.2A: Characteristics of Epithelial Tissue

The human body consists of four types of tissue: epithelial, connective, muscular, and nervous. Epithelial tissue covers the body, lines all cavities, and composes the glands.

Learning Objectives

- Describe the primary functions and characteristics of epithelial tissue

Key Points

- Epithelial tissue is composed of cells laid together in sheets with the cells tightly connected to one another. Epithelial layers are avascular, but innervated.
- Epithelial cells have two surfaces that differ in both structure and function.
- Glands, such as exocrine and endocrine, are composed of epithelial tissue and classified based on how their secretions are released.

Key Terms

- **epithelium**: A membranous tissue composed of one or more layers of cells that form the covering of most internal and external surfaces of the body and its organs.
- **avascular**: Lacking blood vessels.
- **vascular**: Containing blood vessels.

Functions of the Epithelium

Epithelia tissue forms boundaries between different environments, and nearly all substances must pass through the epithelium. In its role as an interface tissue, epithelium accomplishes many functions, including:

1. Protection for the underlying tissues from radiation, desiccation, toxins, and physical trauma.
2. Absorption of substances in the digestive tract lining with distinct modifications.
3. Regulation and excretion of chemicals between the underlying tissues and the body cavity.
4. The secretion of hormones into the blood vascular system. The secretion of sweat, mucus, enzymes, and other products that are delivered by ducts come from the glandular epithelium.
5. The detection of sensation.

Characteristics of Epithelial Layers




Epithelial tissue is composed of cells laid out in sheets with strong cell-to-cell attachments. These protein connections hold the cells together to form a tightly connected layer that is avascular but innervated in nature.



The epithelial cells are nourished by substances diffusing from blood vessels in the underlying connective tissue. One side of the epithelial cell is oriented towards the surface of the tissue, body cavity, or external environment and the other surface is joined to a basement membrane. The basement layer is non-cellular in nature and helps to cement the epithelial tissue to the underlying structures.



Types of Epithelial Tissue


Epithelial tissues are identified by both the number of layers and the shape of the cells in the upper layers. There are eight basic types of epithelium: six of them are identified based on both the number of cells and their shape; two of them are named by the

type of cell (squamous) found in them. Epithelial tissue is classified based on the number of cells, the shape of those cells, and the types of those cells.

Epithelial Tissue Cells		
Cells	**Locations**	**Function**
Simple squamous epithelium 	Air sacs of the lungs and the lining of the heart, blood vessels and lymphatic vessels	Allows materials to pass through by diffusion and filtration, and secretes lubricating substances
Simple cuboidal epithelium 	In ducts and secretory portions of small glands and in kidney tubules	Secretes and absorbs
Simple columnar epithelium 	Ciliated tissues including the bronchi, uterine tubes, and uterus; smooth (nonciliated tissues) are in the digestive tract bladder	Absorbs; it also secretes mucous and enzymes.

Epithelial Tissue Cells		
Cells	**Locations**	**Function**
Pseudostratified columnar epithelium 	Ciliated tissue lines the trachea and much of the upper respiratory tract	Secrete mucous; ciliated tissue moves mucous
Stratified squamous epithelium 	Lines the esophagus, mouth, and vagina	Protects against abrasion

Epithelial Tissue Cells		
Cells	**Locations**	**Function**
Stratified cuboidal epithelium 	Sweat glands, salivary glands, and mammary glands	Protective tissue
Stratified columnar epithelium 	The male urethra and the ducts of some glands.	Secretes and protects

Epithelial Tissue Cells		
Cells	**Locations**	**Function**
Transitional epithelium 	Lines the bladder, urethra and ureters	Allows the urinary organs to expand and stretch

This page titled 4.2A: Characteristics of Epithelial Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.2B: Types of Epithelial Tissue

Epithelial tissue is classified by cell shape and the number of cell layers.

Learning Objectives

- Classify epithelial tissue by cell shape and layers


Key Points

- There are three principal cell shapes associated with epithelial cells: squamous epithelium, cuboidal epithelium, and columnar epithelium.
- There are three ways of describing the layering of epithelium: simple, stratified, and pseudostratified.
- Pseudostratified epithelium possesses fine hair-like extensions called cilia and unicellular glands called goblet cells that secrete mucus. This epithelium is described as ciliated pseudostratified epithelium.
- Stratified epithelium differs from simple epithelium in that it is multilayered. It is therefore found where body linings have to withstand mechanical or chemical insult.
- In keratinized epithelia, the most apical layers (exterior) of cells are dead and and contain a tough, resistant protein called keratin. An example of this is found in mammalian skin that makes the epithelium waterproof.
- Transitional epithelia are found in tissues such as the urinary bladder where there is a change in the shape of the cell due to stretching.

Key Terms

- **simple columnar**: A columnar epithelium that is uni-layered.
- **pseudostratified epithelium**: A type of epithelium that, though comprising only a single layer of cells, has its cell nuclei positioned in a manner suggestive of stratified epithelia.
- **squamous**: Flattened and scale-like.
- **cuboidal**: Resembling a cube.
- **Keratinized**: To produce or become like keratin.
- **columnar**: Having the shape of a column.

Most epithelial tissue is described with two names. The first name describes the number of cell layers present and the second describes the shape of the cells. For example, simple squamous epithelial tissue describes a single layer of cells that are flat and scale-like in shape.



Epithelial Tissue: There are three principal classifications associated with epithelial cells. Squamous epithelium has cells that are wider than they are tall. Cuboidal epithelium has cells whose height and width are approximately the same. Columnar

epithelium has cells taller than they are wide.

Simple Epithelia

Simple epithelium consists of a single layer of cells. They are typically where absorption, secretion and filtration occur. The thinness of the epithelial barrier facilitates these processes.

Simple epithelial tissues are generally classified by the shape of their cells. The four major classes of simple epithelium are: 1) simple squamous; 2) simple cuboidal; 3) simple columnar; and 4) pseudostratified.

Simple Squamous

Simple squamous epithelium cells are flat in shape and arranged in a single layer. This single layer is thin enough to form a membrane that compounds can move through via passive diffusion. This epithelial type is found in the walls of capillaries, linings of the pericardium, and the linings of the alveoli of the lungs.

Simple Cuboidal

Simple cuboidal epithelium consists of a single layer cells that are as tall as they are wide. The important functions of the simple cuboidal epithelium are secretion and absorption. This epithelial type is found in the small collecting ducts of the kidneys, pancreas, and salivary glands.

Simple Columnar

Simple columnar epithelium is a single row of tall, closely packed cells, aligned in a row. These cells are found in areas with high secretory function (such as the wall of the stomach), or absorptive areas (as in small intestine). They possess cellular extensions (e.g., microvilli in the small intestine, or the cilia found almost exclusively in the female reproductive tract).

Pseudostratified

These are simple columnar epithelial cells whose nuclei appear at different heights, giving the misleading (hence pseudo) impression that the epithelium is stratified when the cells are viewed in cross section.

Pseudostratified epithelium can also possess fine hair-like extensions of their apical (luminal) membrane called cilia. In this case, the epithelium is described as ciliated pseudostratified epithelium. Ciliated epithelium is found in the airways (nose, bronchi), but is also found in the uterus and fallopian tubes of females, where the cilia propel the ovum to the uterus.

Stratified Epithelium

Stratified epithelium differs from simple epithelium by being multilayered. It is therefore found where body linings have to withstand mechanical or chemical insults.

Stratified epithelia are more durable and protection is one their major functions. Since stratified epithelium consists of two or more layers, the basal cells divide and push towards the apex, and in the process flatten the apical cells.

Stratified epithelia can be columnar, cuboidal, or squamous type. However, it can also have the following specializations:

Keratinized Epithelia

In keratinized epithelia, the most apical layers (exterior) of cells are dead and lose their nucleus and cytoplasm. They contain a tough, resistant protein called keratin. This specialization makes the epithelium waterproof, and it is abundant in mammalian skin. The lining of the esophagus is an example of a non-keratinized or moist stratified epithelium.

Transitional Epithelia

Transitional epithelia are found in tissues that stretch and it can appear to be stratified cuboidal when the tissue is not stretched, or stratified squamous when the organ is distended and the tissue stretches. It is sometimes called the urothelium since it is almost exclusively found in the bladder, ureters, and urethra.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Epithelial tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithelial_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithelium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithelial tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithelial_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epithelium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- avascular. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/avascular**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vascular. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vascular**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ummary showing different epithelial cells/tissues and their characteristics.. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Epithel...sue_CellsN.jpg**. **License**: ***CC BY: Attribution***
- Epithelial tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithelial_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cuboidal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cuboidal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Keratinized. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Keratinized**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pseudostratified epithelium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pseudos...d%20epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- columnar. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/columnar**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- simple columnar. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/simple%20columnar**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- squamous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/squamous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithelium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ummary showing different epithelial cells/tissues and their characteristics.. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Epithel...sue_CellsN.jpg**. **License**: ***CC BY: Attribution***
- Illu%252520epithelium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_epithelium.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 4.2B: Types of Epithelial Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.3: CONNECTIVE TISSUE



- Topic hierarchy

4.3A: Characteristics of Connective Tissue

4.3B: Types of Connective Tissue

This page titled 4.3: Connective Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.3A: Characteristics of Connective Tissue

Connective tissue is incredibly diverse and contributes to energy storage, the protection of organs, and the body's structural integrity.

Learning Objectives

- Describe the main characteristics and functions of connective tissue

Key Points

- Connective tissue is the most abundant and widely distributed of the primary tissues.
- Connective tissue has three main components: cells, fibers, and ground substance. Together the ground substance and fibers make up the extracellular matrix.
- Connective tissue is classified into two subtypes: soft and specialized connective tissue.
- Major functions of connective tissue include: 1) binding and supporting, 2) protecting, 3) insulating, 4) storing reserve fuel, and 5) transporting substances within the body.
- Connective tissues can have various levels of vascularity. Cartilage is avascular, while dense connective tissue is poorly vascularized. Others, such as bone, are richly supplied with blood vessels.

Key Terms

- **extracellular matrix**: Cells of the connective tissue are suspended in a non-cellular matrix that provides structural and biochemical support to the surrounding cells.
- **fibroblast**: A type of cell found in connective tissue that synthesizes the extracellular matrix and collagen.
- **connective tissue**: A type of tissue found in animals whose main function is to bind, support, and anchor the body.


Connective tissue (CT) is a one of the four main classes of tissues. Although it is the most abundant and widely distributed of the primary tissues, the amount of connective tissue in a particular organ varies. Like to the timber framing of a house, the connective tissue provides structure and support throughout the body.

Structure of Connective Tissue

Connective tissue has three main components:

1. Ground substance
2. Fibers
3. Cells

Together the ground substance and fibers make up the extracellular matrix. The composition of these three elements vary tremendously from one organ to the other. This offers great diversity in the types of connective tissue.




Structural elements of connective tissue: Connective tissues consist of three parts: cells suspended in a ground substance or matrix; and most have fibers running through it.

Ground substance is a clear, colorless, viscous fluid that fills the space between the cells and fibers. It is composed of proteoglycans and cell adhesion proteins that allow the connective tissue to act as glue for the cells to attach to the matrix. The ground substance functions as a molecular sieve for substances to travel between blood capillaries and cells.

Connective tissue fibers provide support. Three types of fibers are found in connective tissue:

1. Collagen
2. Elastic fibers
3. Reticular fibers

Collagen Fibers



Collagen: Collagen fibers are the strongest and most abundant of all the connective tissue fibers.

Collagen fibers are fibrous proteins and are secreted into the extracellular space and they provide high tensile strength to the matrix.

Elastic Fibers

Elastic fibers are long, thin fibers that form branching network in the extracellular matrix. They help the connective tissue to

stretch and recoil.

Reticular Fibers

Reticular fibers are short, fine collagenous fibers that can branch extensively to form a delicate network.

Function of Connective Tissue

The major functions of connective tissue include:

1. Binding and supporting.
2. Protecting.
3. Insulating.
4. Storing reserve fuel.
5. Transporting substances within the body.

This page titled 4.3A: Characteristics of Connective Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.3B: Types of Connective Tissue

Connective tissues encompass a diverse array of tissue types that are involved in binding and supporting body structure and tissues.

Learning Objectives

- Describe the diverse types of connective tissue

Key Points

- The lymphatic system is a part of the circulatory system, comprising a network of conduits called lymphatic vessels that carry a clear fluid called lymph unidirectionally towards the heart.
- Blood is considered a specialized form of connective tissue. In vertebrates, it is composed of blood cells suspended in a liquid called blood plasma.
- The primary tissue of bone, osseous tissue, is a relatively hard and lightweight composite material, formed mostly of calcium phosphate in the chemical arrangement termed calcium hydroxylapatite.
- Adipose tissue or body fat is loose connective tissue composed of adipocytes.
- Cartilage is a flexible connective tissue found in many areas in the bodies of humans and other animals, including the joints between bones, the rib cage, the ear, the nose, the elbow, the knee, the ankle, the bronchial tubes, and the intervertebral discs.
- In humans, adipose tissue is located beneath the skin (subcutaneous fat), around internal organs (visceral fat), in bone marrow (yellow bone marrow), and in breast tissue.

Key Terms

- **cartilage**: A type of dense, non-vascular connective tissue, usually found at the end of joints, the rib cage, the ear, the nose, in the throat, and between intervertebral disks.
- **adipose tissue**: Connective tissue that stores fat and cushions and insulates the body.
- **blood**: A vital liquid flowing in the bodies of many types of animals that usually conveys nutrients and oxygen. In vertebrates, it is colored red by hemoglobin, is conveyed by arteries and veins, is pumped by the heart, and is usually generated in bone marrow.

Connective tissue is divided into four main categories:

1. Connective proper
2. Cartilage
3. Bone
4. Blood


Connective tissue proper has two subclasses: loose and dense. Loose connective tissue is divided into 1) areolar, 2) adipose, 3) reticular. Dense connective tissue is divided into 1) dense regular, 2) dense irregular, 3) elastic.

Areolar Connective Tissue

These tissues are widely distributed and serve as a universal packing material between other tissues. The functions of areolar connective tissue include the support and binding of other tissues.

It also helps in defending against infection. When a body region is inflamed, the areolar tissue in the area soaks up the excess fluid as a sponge and the affected area swells and becomes puffy, a condition called edema.

Adipose Tissue or Body Fat



Adipose tissue: Yellow adipose tissue in paraffin section with lipids washed out.

This is loose connective tissue composed of adipocytes. It is technically composed of roughly only 80% fat. Its main role is to store energy in the form of lipids, although it also cushions and insulates the body.

The two types of adipose tissue are white adipose tissue (WAT) and brown adipose tissue (BAT). Adipose tissue is found in specific locations, referred to as adipose depots.

Reticular Connective Tissue

This tissue resembles areolar connective tissue, but the only fibers in its matrix are the reticular fibers, which form a delicate network. The reticular tissue is limited to certain sites in the body, such as internal frameworks that can support lymph nodes, spleen, and bone marrow.

Dense Regular Connective Tissue

This consists of closely packed bundles of collagen fibers running in the same direction. These collagen fibers are slightly wavy and can stretch a little bit.

With the tensile strength of collagen, this tissue forms tendons, aponeurosis and ligaments. This tissue forms the fascia, which is a fibrous membrane that wraps around the muscles, blood vessels, and nerves.

Dense Irregular Tissue

This has the same structural elements as dense regular tissue, but the bundles of collagen fibers are much thicker and arranged irregularly. This tissue is found in areas where tension is exerted from many different directions. It is part of the skin dermis area and in the joint capsules of the limbs.

Elastic Connective Tissue

The main fibers that form this tissue are elastic in nature. These fibers allow the tissues to recoil after stretching. This is especially seen in the arterial blood vessels and walls of the bronchial tubes.

Cartilage

This is a flexible connective tissue found in many areas in the bodies of humans and other animals, including the joints between bones, the rib cage, the ear, the nose, the elbow, the knee, the ankle, the bronchial tubes, and the intervertebral discs.

Cartilage is composed of specialized cells called chondroblasts and, unlike other connective tissues, cartilage does not contain blood vessels. Cartilage is classified in three types: 1) elastic cartilage, 2) hyaline cartilage, and 3) fibrocartilage, which differ in the relative amounts of these three main components.

Elastic Cartilage

This is similar to hyaline cartilage but is more elastic in nature. Its function is to maintain the shape of the structure while allowing flexibility. It is found in the external ear (known as an auricle) and in the epiglottis.

Hyaline Cartilage

This is is the most abundant of all cartilage in the body. Its matrix appears transparent or glassy when viewed under a microscope. It provides strong support while providing pads for shock absorption. It is a major part of the embryonic skeleton, the costal cartilages of the ribs, and the cartilage of the nose, trachea, and larynx.

Fibrocartilage

This is a blend of hyaline cartilage and dense regular connective tissue. Because it is compressible and resists tension well, fibrocartilage is found where strong support and the ability to withstand heavy pressure are required. It is found in the intervertebral discs of the bony vertebrae and knee meniscus.

Bone tissue is also called the osseous tissue. The osseous tissue is relatively hard and lightweight in nature. It is mostly formed of calcium phosphate in the chemical arrangement termed calcium hydroxyapatite, which gives bones their rigidity. It has relatively high compressive strength, but poor tensile strength, and very low shear stress strength.

The hard outer layer of bones is composed of compact bone tissue, so-called due to its minimal gaps and spaces. Its porosity is 5–30%. This tissue gives bones their smooth, white, and solid appearance, and accounts for 80% of the total bone mass of an adult skeleton.

Filling the interior of the bone is the trabecular bone tissue (an open cell porous network also called cancellous or spongy bone), which is composed of a network of rod and plate-like elements that make the overall organ lighter and allow room for blood vessels and marrow.

Blood

This is considered a specialized form of connective tissue. Blood is a bodily fluid in animals that delivers necessary substances, such as nutrients and oxygen, to the cells and transports metabolic waste products away from those same cells.

It is an atypical connective tissue since it does not bind, connect, or network with any body cells. It is made up of blood cells and is surrounded by a nonliving fluid called plasma.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- extracellular matrix. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/extracellular_matrix. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connective tissue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Connect...sue%23See_also. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connective tissue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Connective_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and Physiology of Animals/Body Organisation. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy...ective_Tissues. **License**: *CC BY-SA: Attribution-ShareAlike*
- fibroblast. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fibroblast. **License**: *CC BY-SA: Attribution-ShareAlike*
- connective tissue. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/connective_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals loose connective tissue. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_loose_connective_tissue.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Fibers of Collagen Type I - TEM. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Fibers_of_Collagen_Type_I_-_TEM.jpg. **License**: *Public Domain: No Known Copyright*
- Blood. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphatic_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Adipose tissue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adipose_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- General Biology/Tissues and Systems/Connective Tissue. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/General_Biology/Tissues_and_Systems/Connective_Tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- blood. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/blood. **License**: *CC BY-SA: Attribution-ShareAlike*
- cartilage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- adipose tissue. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/adipose_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals loose connective tissue. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_loose_connective_tissue.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Fibers of Collagen Type I - TEM. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Fibers_of_Collagen_Type_I_-_TEM.jpg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/2/27/Yellow_adipose_tissue_in_paraffin_section_-_lipids_washed_out.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 4.3B: Types of Connective Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.4: NERVOUS TISSUE



- Topic hierarchy

4.4A: Characteristics of Nervous Tissue

4.4B: Types of Nervous Tissue

This page titled 4.4: Nervous Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.4A: Characteristics of Nervous Tissue

Learning Objectives

- Describe the characteristics of nervous tissue


Nervous tissue is one of four major classes of tissues. It is specialized tissue found in the central nervous system and the peripheral nervous system. It consists of neurons and supporting cells called neuroglia. The nervous system is responsible for the control of the body and the communication among its parts. Nervous tissue contains two categories of cells—neurons and neuroglia.

Neurons

Neurons are highly specialized nerve cells that generate and conduct nerve impulses. A typical neuron consists of dendrites, the cell body, and an axon.

Dendrites

Dendrites are responsible for responding to stimuli; they receive incoming signals towards the cell body. The axons are responsible for transmitting impulses over long distances from cell body. The cell body is like a factory for the neuron. It produces all the proteins and contains specialized organelles such as nucleus, granules and Nissl bodies.



Neuron: This image illustrates the parts of a neuron. The dendrites receive incoming signals while axons propagate signals away from the neuron cell body. The myelin sheath surrounds and insulates the axon.

Dendrite

The axon is surrounded by a whitish, fatty layer called the myelin sheath. Outside the myelin sheath there is a cellular layer called the neurilemma.

Schwann Cells

In the peripheral nervous system, Schwann cells are neuroglia cells that support neuronal function by increasing the speed of impulse propagation. The Schwann cells are underlain by the medullary sheath. The medullary sheath is interrupted at intervals by the nodes of Ranvier.




Illustration of the Schwann Cells and the Myelin Sheath: Transmission electron micrograph of a myelinated axon. The myelin layer (concentric) surrounds the axon of a neuron, showing Schwann cells.

Key Points

- Nervous tissue is one of four major classes of tissues and makes up the central nervous system and the peripheral nervous system.
- Integration and communication are the two major functions of nervous tissue.
- Nervous tissue contains two categories of cells — neurons and neuroglia.
- Neurons are highly specialized nerve cells that generate and conduct nerve impulses.
- Neuroglia are supporting cells that provide physical sport, remove debris, and provide electrical insulation.

Key Terms

- **myelin**: A substance produced by cells of the neuroglia that increases the speed of impulses along the axon of the neuronal fiber.
- **nervous tissue**: The principal constituent of the central and peripheral nervous system, comprised neurons and neuroglia cells.
- **brain**: The control center of the central nervous system, located in the skull.

This page titled 4.4A: Characteristics of Nervous Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.4B: Types of Nervous Tissue

Learning Objectives

- Describe the main cells that comprise nervous tissue

Nervous tissue, one of the four main tissue types, is composed of neurons and supporting cells called neuroglia. Neuroglia are also called “glial cells.”

Neuroglia

There are six types of neuroglia—four in the central nervous system and two in the PNS. These glial cells are involved in many specialized functions apart from support of the neurons. Neuroglia in the CNS include astrocytes, microglial cells, ependymal cells and oligodendrocytes. In the PNS, satellite cells and Schwann cells are the two kinds of neuroglia.

Astrocytes

Astrocytes are shaped like a star and are the most abundant glial cell in the CNS. They have many radiating processes which help in clinging to the neurons and capillaries. They support and brace the neurons and anchor them to the nutrient supply lines. They also help in the guiding the migration of young neurons. Astrocytes control the chemical environment around the neurons.

Microglial Cells

Microglial cells are small and ovoid un shape with thorny processes. They are found in the CNS. When invading microorganism or dead neurons are present, the microglial cells can transform into a phagocytic macrophage and help in cleaning the neuronal debris.

Ependymal Cells

Ependymal cells are ciliated and line the central cavities of the brain and spinal cord where they form a fairly permeable barrier between the cerebrospinal fluid that fills these cavities and the tissue cells of the CNS.

Oligodendrocytes

Oligodendrocytes line up along the nerves and produce an insulating cover called myelin sheath. They are found in the CNS.

Satellite Cells

Satellite cells surround neuron cell bodies in the peripheral nervous system (PNS). They are analogous to the astrocytes in the CNS.

Schwann Cells

Schwann cells surround all nerve fibers in the peripheral nervous system and form myelin sheaths around the nerve fibers. They are found in the PNS. Their function is similar to oligodendrocytes.

Neurons

Neurons consist of cell body and one or more slender processes. The neuronal cell body consists of a nucleus and rough endoplasmic reticulum or Nissl Bodies. The cell body is the major biosynthetic center of a neuron and contains the usual organelles for the synthesis of proteins and other chemicals. Arm like processes extend from the cell body to all neurons.

The two types of neuron processes are called dendrites and axons. Dendrites are motor neurons that are short and have a large surface area for receiving signals from other neurons. Dendrites convey incoming messages towards the cell body and are therefore called the receptive input region.

The axon arises from the cone shaped portion of the cell body called the axon hillock. Functionally, the axon is the conducting region of the neuron and is responsible for generating and transmitting impulses typically away from the cell body. A single axon routes the nerve impulse from the cell body to another neuron or an effector organ. The axon can have many terminal branches, so each time the nerve fires, it can stimulate more than one cell.

Key Points

- Nervous tissue is composed of neurons and supporting cells called neuroglia, or " glial cells."
- There are six types of neuroglia. Four are found in the central nervous system, while two are found in the peripheral nervous system.
- The four types of neuroglia found in the central nervous system are astrocytes, microglial cells, ependymal cells, and oligodendrocytes.
- The two types of neuroglia found in the peripheral nervous system are satellite cells and Schwann cells.
- Neurons are the other the other type of cell that comprise nervous tissue. Neurons have cell bodies, dendrites, and axons.

Key Terms

- **neuron**: The main cell type in nervous tissue.
- **neuroglia**: Supporting cells in nervous tissue.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Nervous tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- brain. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/brain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myelin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/myelin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nervous tissue. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nervous_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Neuron.svg**. **License**: ***Public Domain: No Known Copyright***
- Myelinated neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:My...ted_neuron.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Neuron.svg**. **License**: ***Public Domain: No Known Copyright***
- Myelinated neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:My...ted_neuron.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 4.4B: Types of Nervous Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.5: MUSCULAR TISSUE



- Topic hierarchy

4.5A: Characteristics of Muscle Tissue

This page titled 4.5: Muscular Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.5A: Characteristics of Muscle Tissue

The three types of muscle tissue are skeletal, smooth, and cardiac.

Learning Objectives

- Describe the types of muscle tissue

Key Points

- Muscle tissue can be divided functionally, is it under voluntary or involuntary control; and morphologically, striated or non-striated.
- By applying these classifications three muscle types can be described; skeletal, cardiac and smooth.
- Skeletal muscle is voluntary and striated, cardiac muscle is involuntary and straited and smooth muscle is involuntary and non-striated.

Key Terms

- **skeletal muscle**: The voluntary muscle of vertebrates, which is striated and anchored by tendons to bone, is used to effect skeletal movement such as locomotion.
- **involuntary**: A muscle movement not under conscious control e.g. the beating of the heart.
- **striated**: The striped appearance of certain muscle types in which myofibrils are aligned to produce a constant directional tension.
- **voluntary**: A muscle movement under conscious control e.g. deciding to move the forearm.
- **smooth muscle**: Involuntary muscle that is found within the intestines, throat, uterus, and blood vessel walls.
- **cardiac muscle**: The striated and involuntary muscle of the vertebrate heart.

Muscle Tissue

Muscle is a soft tissue that is highly specialized for the production of tension which results in the generation of force. Muscle cells, or myocytes, contain myofibrils comprised of actin and myosin myofilaments which slide past each other producing tension that changes the shape of the myocyte. Numerous myocytes make up muscle tissue and the controlled production of tension in these cells can generate significant force.

Types of Muscle Tissue

Muscle tissue can be classified functionally, voluntary or involuntary and morphologically striated or non-striated. Voluntary refers to whether the muscle is under conscious control, striation refers to the presence of visible banding within myocytes which occurs due to organization of myofibrils to produce a constant direction of tension.

By applying the above classifications it is possible to describe three forms of muscle tissue which perform the wide range of functions described.

Skeletal Muscle

Skeletal muscle mainly attaches to the skeletal system via tendons to maintain posture and control movement for example contraction of the biceps muscle, attached to the scapula and radius, will raise the forearm. Some skeletal muscle can attach directly to other muscles or the skin, as seen in the face where numerous muscles control facial expression.

Skeletal muscle is under voluntary control, although this can be subconscious for example when maintaining posture or balance.

Morphologically skeletal myocytes are elongated and tubular and appear striated with multiple peripheral nuclei.

Cardiac Muscle Tissue


Cardiac muscle tissue is found only in the heart where cardiac contractions pump blood throughout the body and maintain blood pressure.

As with skeletal muscle cardiac muscle is striated, however it is not consciously controlled and so is involuntary. Cardiac muscle can be further differentiated from skeletal muscle by the presence of intercalated discs which control the synchronized contraction of cardiac tissues. Cardiac myocytes are shorter than skeletal equivalents and contain only one or two centrally located nuclei.

Smooth Muscle Tissue

Smooth muscle tissue is found associated with numerous other organs and tissue systems such as the digestive system or respiratory system. It plays an important role in the regulation of flow in such tissues for example aiding the movement of food through the digestive system via peristalsis.

Smooth muscle is non-striated, although it contains the same myofilaments they are just organized differently, and involuntary. Smooth muscle myocytes are spindle shaped with a single centrally located nucleus.



Muscle Types: Cardiac and skeletal muscle are both striated in appearance, while smooth muscle is not. Both cardiac and smooth muscle are involuntary while skeletal muscle is voluntary.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- skeletal muscle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/skeletal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle%23Types_of_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- smooth muscle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/smooth_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiac muscle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cardiac_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu muscle tissues. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...le_tissues.jpg**. **License**:

This page titled 4.5A: Characteristics of Muscle Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.6: TISSUE MEMBRANES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

4.6A: Epithelial Membranes

4.6B: Synovial Membranes

This page titled 4.6: Tissue Membranes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.6A: Epithelial Membranes

The mucous membranes are linings of mostly endodermal origin, covered in epithelium, which are involved in absorption and secretion.

Learning Objectives

- Describe the function of mucous membranes

Key Points

- The mucous membranes are linings of ectodermal origin. It consists of an epithelium layer and an underlying lamina propria of loose connective tissue.
- The mucus membranes are involved in absorption and secretion.
- Most mucosal membranes contain stratified squamous or simple columnar epithelial tissue types.
- Submucosal exocrine glands secrete mucus to facilitate the movement of particles along the body's various tubes, such as the throat and the intestines.

Key Terms


- **mucous membrane**: Linings of cavities that are exposed to the external environment and to internal organs.

The mucous membranes are linings of ectodermal origin. It consists of an epithelium layer and an underlying lamina propria of loose connective tissue. These mucus membranes are involved in absorption and secretion. They line cavities that are exposed to the external environment and internal organs. These membranes exist in the hollow organs of the digestive, respiratory, and urogenital tracts.

The term "mucous membrane" refers to where they are found in the body; not every mucous membrane secretes mucus. Secreted mucous traps the pathogens in the body, preventing any further progression of microbes.

Most mucous membranes contain stratified squamous or simple columnar epithelial tissue. The epithelial tissue sheet lies directly over the layer of loose connective tissue called lamina propria. In some mucosa, the lamina propria rests on a deeper, third layer of smooth muscle.

The submucosa is the tissue that connects the mucosa to the muscle outside the tube. Submucosal glands consist of exocrine glands that secrete mucus. These glands excrete mucus to facilitate the movement of particles along the body's various tubes, such as the throat and intestines. The submucosal glands are a companion to unicellular goblet cells, which also produce mucus, and are found lining the same tubes.



General organization of the gastrointestinial tract: Illustration of mucosa in relation to other lining components.

This page titled 4.6A: Epithelial Membranes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

4.6B: Synovial Membranes

A synovial membrane is the soft tissue found between the articular capsule (joint capsule) and the joint cavity of synovial joints.

Learning Objectives

- Describe the role of synovial membranes in joint function

Key Points

- Connective tissue membranes do not contain an epithelial cell layer. Synovial membranes line the inner surface of the capsule of a synovial joint and contain synovial fluid within; which functions to lubricate joint movement.
- Synovial membranes often have two layers: a fibrous outer layer, or subintima, and an inner layer, or intima.
- The intimal cells are termed synoviocytes and are of two types, fibroblastic (type B synoviocytes) and macrophagic (type A synoviocytes).
- The meninges is the system of membranes that envelopes the central nervous system.
- The meninges is comprised of three layers, the dura mater, arachnoid mater and pia mater.

Key Terms

- **meninges**: The system of membranes that envelopes the central nervous system.
- **synovial membrane**: The connective tissue which lines the inner surface of the capsule of a synovial joint.
- **Connective tissue membranes**: A membrane which does not contain an epithelial cell layer.
- **synoviocyte**: The cell that forms the initima layer of a snynovial membrane.

Membranes are thin sheets of tissue found within the body which can line cover tissues or line cavities. Connective tissue membranes do not contain an epithelial cell layer and there are two forms found in the body; synovial and meninges membranes.

Synovial Membrane

The synovial membrane (or synovium) is the connective tissue which lines the inner surface of the capsule of a synovial joint and secretes synovial fluid which serves a lubricating function, allowing joint surfaces to smoothly move across each other.

The morphology of synovial membranes may vary, but it often consists of two layers. The outer layer, or subintima, is a thicker and fibrous protecting the single cell initma layer which is composed of synoviocytes.


Synoviocytes

The intimal cells are termed synoviocytes and are of two types: fibroblastic (type B) and macrophagic (type A). It is the lack of epithelial cells within the initma which defines the synovial membrane as connective rather than epithelial.

The type B synoviocytes manufacture a long-chain sugar polymer called hyaluronan, which makes the synovial fluid together with a molecule called lubricin, which lubricates the joint surfaces. The water component of synovial fluid is effectively trapped in the joint space by the hyaluronan, due to its large, highly negatively charged moeties.

The type A synoviocytes are responsible for the removal of undesirable substances from the synovial fluid.

Structure of Synovium



Synovial Membrane: A synovial joint showing the location of the synovial membrane.

The surface of synovium may be flat or may be covered with finger-like projections (villi), to allow the soft tissue to change shape as the joint surfaces move on one another.

Just beneath the intima, most synovium has a dense net of small blood vessels that provide nutrients, not only for synovium, but also for the avascular cartilage. In any one position, much of the cartilage is close enough to get nutrition directly from the synovium.

Meninges

The meninges is the system of membranes that envelopes the central nervous system. In mammals, the meninges consist of three layers: the dura mater, the arachnoid mater, and the pia mater. The primary function of the meninges and of the cerebrospinal fluid is to protect the central nervous system.


Dura Mater

The dura mater is a thick, durable membrane which lies closest to the skull. It consists of two layers, the periosteal layer which lies closest to the calvaria, and the inner meningeal layer which lies closer to the brain. It is composed of dense fibrous tissue, and its inner surface is covered by flattened cells like those present on the surfaces of the pia mater and arachnoid.

Arachnoid Mater

The middle layer of the meninges is the arachnoid mater, so named because of its spider web-like appearance. It provides a cushioning effect for the central nervous system. The arachnoid mater is a thin, transparent membrane composed of fibrous tissue and, like the pia mater, is covered by flat cells also thought to be impermeable to fluid.

Pia Mater



The Meninges: This figure displays the meninges with respect to the skull and surface of the brain.

The pia mater is the innermost layer of the meninges. It firmly adheres to the surface of the brain and spinal cord, following the brain's minor contours. As such it is a very thin, delicate membrane composed of fibrous tissue covered on its outer surface by a sheet of flat cells thought to be impermeable to fluid.

The subarachnoid space is the space that normally exists between the arachnoid and the pia mater, which is filled with cerebrospinal fluid. Normally, the dura mater is attached to the skull or to the bones of the vertebral canal in the spinal cord. The arachnoid is attached to the dura mater, while the pia mater is attached to the central nervous system tissue.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Mucous membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mucous_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Submucosal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Submucosal_glands**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epithelium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mucous membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mucous_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Synovial membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synovial_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibroblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibroblast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- macrophage. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/macrophage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synoviocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/synoviocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mucous membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mucous_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 20100052912574721Meninges-en.svg.png. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/archive/8/8e/20100529125747%21Meninges-en.svg**. **License**: ***Public Domain: No Known Copyright***

- Synovial membranes. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fc/Illu_synovial_joint.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 4.6B: Synovial Membranes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5: INTEGUMENTARY SYSTEM

5.1: The Skin

5.1A: What is Skin?

5.1B: Structure of the Skin: Epidermis

5.1C: Structure of the Skin: Dermis

5.1D: Skin Color

5.2: Cell Junctions

5.2A: Tight Junctions

5.2B: Adherens Junctions

5.3C: Gap Junctions

5.3: Accessory Structures of the Skin

5.3A: Sweat (Sudoriferous) Glands

5.3B: Sebaceous (Oil) Glands

5.3C: Nails

5.3D: Hair

5.4: Functions of the Integumentary System

5.4A: Protection

5.4B: Thermoregulation

5.4C: Cutaneous Sensation

5.4D: Metabolic Functions

5.4E: Blood Supply to the Epidermis

5.4F: Excretion and Absorption

5.5: Wound Healing

5.5A: Steps of Tissue Repair

5.5B: Epidermal Wound Healing

5.5C: Deep Wound Healing

5.6: Overview of Cancer

5.6A: Overview of Cancer

5.6B: Cell Signaling and Cell Death

This page titled 5: Integumentary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.1: THE SKIN



- Topic hierarchy

5.1A: What is Skin?

5.1B: Structure of the Skin: Epidermis

5.1C: Structure of the Skin: Dermis

5.1D: Skin Color

This page titled 5.1: The Skin is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.1A: What is Skin?

The skin is the soft outer covering of vertebrates that guards the underlying muscles, bones, ligaments, and internal organs.

LEARNING OBJECTIVE

Identify the layers of the skin

Key Takeaways

Key Points

- The outer layer of skin, the epidermis, provides waterproofing and serves as a barrier to infection.
- The middle layer of skin, the dermis, contains blood vessels, nerves, and glands that are important for our skin's function. The inner layer of the skin, the subcutis, contains fat that protects us from trauma.

Key Terms

- **epidermis**: The outermost layer of the skin.
- **subcutis**: The inner layer of skin that is also called the hypodermis or subcutaneous layer.
- **dermis**: The middle layer of the skin.
- **cutaneous membrane**: The formal name for the skin.

The Cutaneous Membrane

The cutaneous membrane is the technical term for our skin. The skin's primary role is to help protect the rest of the body's tissues and organs from physical damage such as abrasions, chemical damage such as detergents, and biological damage from microorganisms. For example, while the skin harbors many permanent and transient bacteria, these bacteria are unable to enter the body when healthy, intact skin is present.

Our skin is made of three general layers. In order from most superficial to deepest they are the epidermis, dermis, and subcutaneous tissue.

The Epidermis

The epidermis is a thin layer of skin. It is the most superficial layer of skin, the layer you see with your eyes when you look at the skin anywhere on your body. Functions of the epidermis include touch sensation and protection against microorganisms.

This skin is further divided into five, separate layers. In order from most superficial to deepest, they are the:

- Stratum Corneum
- Stratum Lucidum
- Stratum Granulosum
- Stratum Spinosum
- Stratum Basale

Stratum Corneum

This layer is composed of the many dead skin cells that you shed into the environment—as a result, these cells are found in dust throughout your home. This layer helps to repel water.

Stratum Lucidum

This layer is found only on the palms of the hands, fingertips, and the soles of the feet.

Stratum Granulosum

This is the layer where part of keratin production occurs. Keratin is a protein that is the main component of skin.

Stratum Spinosum

This layer gives the skin strength as well as flexibility.

Stratum Basale

This is where the skin's most important cells, called keratinocytes, are formed before moving up to the surface of the epidermis and being shed into the environment as dead skin cells.

This layer also contains melanocytes, the cells that are largely responsible for determining the color of our skin and protecting our skin from the harmful effects of UV radiation. These harmful effects include burns in the short term and cancer in the long run.

The Dermis

Underneath the epidermis lies the dermis. The dermis contains:


- Blood vessels that nourish the skin with oxygen and nutrients. The blood vessels also allow immune system cells to come to the skin to fight an infection. These vessels also help carry away waste products.
- Nerves that help us relay signals coming from the skin. These signals include touch, temperature, pressure, pain, and itching.
- Various glands.
- Hair follicles.
- Collagen, a protein that is responsible for giving skin strength and a bit of elasticity.

The Subcutaneous Tissue

The deepest layer of the skin is called the subcutaneous layer, the subcutis, or the hypodermis. Like the dermis, the layer contains blood vessels and nerves for much the same reasons.

Importantly, the subcutis contains a layer of fat. This layer of fat works alongside the blood vessels to maintain an appropriate body temperature. The layer of fat here acts as a cushion against physical trauma to internal organs, muscles, and bones.

Additionally, the body will turn to this fat in times of starvation to provide power to its various processes, especially brain function.



Layers of cutaneous membranes (skin): This image details features of the epidermal and dermal layers of the skin.

This page titled 5.1A: What is Skin? is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.1B: Structure of the Skin: Epidermis

LEARNING OBJECTIVE


- Describe the layers of the epidermis

Layers of the Epidermis

The epidermis is the outermost layer of our skin. It is the layer we see with our eyes. It contains no blood supply of its own—which is why you can shave your skin and not cause any bleeding despite losing many cells in the process. Assuming, that is, you don't nick your skin to deep, where the blood supply is actually found.


The epidermis is itself divided into at least four separate parts. A fifth part is present in some areas of our body. In order from the deepest layer of the epidermis to the most superficial, these layers (strata) are the:

- Stratum basale
- Stratum spinosum
- Stratum granulosum
- Stratum lucidum
- Stratum corneum



Skin overview: Skin layers, of both hairy and hairless skin.

Stratum Basale



Human skin: This image details the parts of the integumentary system.

The stratum basale, also called the stratum germinativum, is the basal (base) layer of the epidermis. It is the layer that's closest to the blood supply lying underneath the epidermis.


This layer is one of the most important layers of our skin. This is because it contains the only cells of the epidermis that can divide via the process of mitosis, which means that skin cells germinate here, hence the word germinativum.

In this layer, the most numerous cells of the epidermis, called keratinocytes, arise thanks to mitosis. Keratinocytes produce the most important protein of the epidermis.

This protein is appropriately called keratin. Keratin makes our skin tough and provides us with much-needed protection from microorganisms, physical harm, and chemical irritation.

Millions of these new cells arise in the stratum basale on a daily basis. The newly produced cells push older cells into the upper layers of the epidermis with time. As these older cells move up toward the surface, they change their shape, nuclear, and chemical composition. These changes are, in part, what give the strata their unique characteristics.

Stratum Spinosum and Granulosum



Layers of the epidermis: The epidermis is made up of 95% keratinocytes but also contains melanocytes, Langerhans cells, Merkel cells, and inflammatory cells. The stratum basale is primarily made up of basal keratinocyte cells, which can be considered the stem cells of the epidermis. They divide to form the keratinocytes of the stratum spinosum, which migrate superficially.

From the stratum basale, the keratinocytes move into the stratum spinosum, a layer so called because its cells are spiny-shaped cells. The stratum spinosum is partly responsible for the skin's strength and flexibility.

From there the keratinocytes move into the next layer, called the stratum granulosum. This layer gets its name from the fact that the cells located here contain many granules.

The keratinocytes produce a lot of keratin in this layer—they become filled with keratin. This process is known as keratinization. The keratinocytes become flatter, more brittle, and lose their nuclei in the stratum granulosum as well.

Stratum Lucidum

Once the keratinocytes leave the stratum granulosum, they die and help form the stratum lucidum. This death occurs largely as a result of the distance the keratinocytes find themselves from the rich blood supply the cells of the stratum basale lie on top off. Devoid of nutrients and oxygen, the keratinocytes die as they are pushed towards the surface of our skin.

The stratum lucidum is a layer that derives its name from the lucid (clear/transparent) appearance it gives off under a microscope. This layer is only easily found in certain hairless parts of our body, namely the palms of our hands and the soles of our feet. Meaning, the places where our skin is usually the thickest.

Stratum Corneum

From the stratum lucidum, the keratinocytes enter the next layer, called the stratum corneum (the horny layer filled with cornified cells). This the only layer of skin we see with our eyes.

The keratinocytes in this layer are called corneocytes. They are devoid of almost all of their water and they are completely devoid of a nucleus at this point. They are dead skin cells filled with the tough protein keratin. In essence, they are a protein mass more so than they are a cell.

The corneocytes serve as a hard protective layer against environmental trauma, such as abrasions, light, heat, chemicals, and microorganism. The cells of the stratum corneum are also surrounded by lipids (fats) that help repel water as well. These

corneocytes are eventually shed into the environment and become part of the dandruff in our hair or the dust around us, which dust mites readily munch on.

This entire cycle, from new keratinocyte in the straum basale to a dead cell flaked off into the air, takes between 25–45 days.

Key Points

- The epidermis provides a protective waterproof barrier that also keeps pathogens at bay and regulates body temperature.
- The main layers of the epidermis are: stratum corneum, stratum lucidium, stratum granulosm, stratum spinosum, stratum germinativum (also called stratum basale).
- Keratinocytes in the stratum basale proliferate during mitosis and the daughter cells move up the strata, changing shape and composition as they undergo multiple stages of cell differentiation.

Key Terms

- **keratinocyte**: The predominant cell type in the epidermis, the outermost layer of the skin, constituting 95% of the cells found there. Those keratinocytes found in the basal layer (stratum germinativum) of the skin are sometimes referred to as basal cells or basal keratinocytes.
- **stratum germinativum**: The basal layer—sometimes referred to as stratum basale—is the deepest of the five layers of the epidermis.
- **stratum corneum**: The most superficial layer of the epidermis from which dead skin sheds.
- **epidermis**: The outermost layer of skin.
- **stratum lucidum**: A layer of our skin that is found on the palms of our hands and the soles of our feet.

This page titled 5.1B: Structure of the Skin: Epidermis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.1C: Structure of the Skin: Dermis

The dermis consists of a papillary and a reticular layer that serve to protect and cushion the body from stress and strain.

LEARNING OBJECTIVE

Differentiate among the regions of the dermis and the hypodermis

Key Takeaways

Key Points

- The dermis is divided into a papillary region and a reticular region.
- The primary function of the dermis is to cushion the body from stress and strain, and to also provide: elasticity to the skin, a sense of touch, and heat.
- The dermis contains hair roots, sebaceous glands, sweat glands, nerves, and blood vessels.
- The hypodermis lies below the dermis and contains a protective layer of fat.

Key Terms


- **the reticular layer**: The deepest layer of the dermis.
- **hypodermis**: A subcutaneous layer of loose connective tissue containing fat cells, lying beneath the dermis.
- **the dermis**: The layer of skin underneath the epidermis.
- **the papillary layer**: The most superficial layer of the dermis.

The Dermis

Lying underneath the epidermis—the most superficial layer of our skin—is the dermis (sometimes called the corium). The dermis is a tough layer of skin. It is the layer of skin you touch when buying any leather goods.

The dermis is composed of two layers. They are the papillary layer (the upper layer) and the reticular layer (the lower layer).

The Papillary Layer



Human Skin: This image details the parts of the integumentary system.

The papillary layer provides the layer above it, the epidermis, with nutrients to produce skin cells called keratinocytes. It also helps regulate the temperature of our skin and thus the body as a whole.

Both the nutrient supply and temperature regulation occur thanks to an extensive network of blood vessels in this layer. These blood vessels also help remove cellular waste products that would otherwise kill the skin cells if they were allowed to accumulate.

The pink tint to the skin of light-skinned individuals is due to the blood vessels found here. In fact, when you blush, it is the dilation of these blood vessels that causes you to turn red. The uneven projections found in this layer, called dermal papillae, also form people's fingerprints and give this layer its name.

The Reticular Layer

The reticular layer serves to strengthen the skin and also provides our skin with elasticity. Elasticity refers to how our skin is able to spring back into shape if deformed by something like a pinch. The reticular layer also contains hair follicles, sweat glands, and sebaceous glands.

The sweat gland can either be apocrine, such as those found in the armpits and the groin area, or the eccrine glands, which are found all over the body. The former help contribute to body odor (along with the bacteria on our skin), and the latter help regulate our body temperature through the process of evaporation.

The sebaceous glands found in the dermis secrete a substance called sebum that helps to lubricate and protect our skin from drying out.

The dermis also contains:

- Nerve endings that transmit various stimuli such as pain, itch, pressure, and temperature.
- Lymphatic vessels that transport immune system cells, the cells that help destroy infectious organisms that may have found their way into our body via a scratch on the skin.
- Collagen, a protein that helps strengthen our skin, and elastin, a protein that helps keep our skin flexible.


The Hypodermis

Beneath the dermis is the deepest layer of our skin. It is alternatively termed hypodermis, subcutis, or subcutaneous tissue. It

contains many collagen cells as well as fat.

Fat, in particular, helps insulate our body from the cold and act as a cushion for our internal structures (such as muscles and organs) when something hits us. Fat can also be called upon by the body in times of great need as an energy source.

Given the alternative names for this layer, it should come as no surprise that this is the layer where subcutaneous injections are given into via a hypodermic needle.



Skin sensory receptors: Those nearest the surface of the skin include receptors that detect gentle pressure, temperature, and vibrations, as well as naked nerve endings (dendrites) that detect pain. Deeper in the dermis are naked dendrites that wind around the bases of hair follicles and detect motions of the hairs, as well as receptors such as Pacinian corpuscles that respond to strong pressure and vibrations.

This page titled 5.1C: Structure of the Skin: Dermis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.1D: Skin Color

Skin color is determined largely by the amount of melanin pigment produced by melanocytes in the skin.

LEARNING OBJECTIVE

Explain how differing degrees of pigmentation are produced

Key Takeaways

Key Points


- Skin color is mainly determined by a pigment called melanin.
- Melanin is produced by melanocytes through a process called melanogenesis.
- The difference in skin color between lightly and darkly pigmented individuals is due to their level of melanocyte activity; it is not due to the number of melanocytes in their skin.

Key Terms

- **melanin**: Any of a group of naturally occurring dark pigments responsible for the color of skin.
- **melanocyte**: A cell in the skin that produces the pigment melanin.
- **keratinocytes**: Cells that take up and store melanin.
- **eumelanin**: The type of melanin mainly responsible for brown and black skin.
- **stratum basale**: The epidermal layer where melanocytes are found.

Melanin

Skin color is largely determined by a pigment called melanin but other things are involved. Your skin is made up of three main layers, and the most superficial of these is called the epidermis. The epidermis itself is made up of several different layers.



Melanocyte: Cross-section of skin showing melanin in melanocytes

The deepest of the epidermal layers is called the stratum basale or stratum germinativum. In this layer lie important cells called

melanocytes. Their name is derived from two parts: melano-, which means black or darkness, and -cyte, which means cell.

Melanocytes are irregularly shaped cells that produce and store a pigment called melanin. The most abundant type of melanin is called eumelanin. This pigment is stored in organelles called melanosomes.

Eumelanin is responsible for the brown and black pigmentation of human skin or the lack thereof if little of it is produced. The production of melanin is called melanogenesis—genesis means formation or development.

How Skin Color is Determined

Regardless of background, every person has largely the same number of melanocytes, but the genetics of each person is what determines how much melanin is produced and how it is distributed throughout the skin. For example, light skinned individuals may have darker places like nipples and moles. Conversely, dark skinned individuals have a lighter tone to the palms of their hands.

Another critical factor, exposure to sunlight, triggers the production of melanin as well. This is what gives us a tan. The melanin produced in response to the sun's rays protects our skin and the rest of the body from the harmful effects of the sun's burn and cancer-inducing U.V. radiation.


The Role of Keratinocytes

People with darker skin have more active melanocytes compared to people with lighter skin. However, the pigment of our skin also involves the most abundant cells of our epidermis, the keratinocytes.

While melanocytes produce, store, and release melanin, keratinocytes are the largest recipients of this pigment. The transfer of melanin from melanocytes to keratinocytes occurs thanks to the long tentacles each melanocyte extends to upwards of 40 keratinocytes.

If a person is unable to produce melanin, they have a condition called albinism.

Other Skin Color Determinants



Tanned Skin: Exposure to UV radiation through tanning causes changes in the pigmentation of the skin by increasing melanin production.

Besides melanin, other factors play a role in general or local skin color. These include:

1. The amount of carotene found in the stratum corneum of the epidermis and the deepest layer of the skin, the hypodermis. Carotene is a yellow-orange pigment found in carrots. Your skin may turn this color if you eat a lot of carotene-rich foods. The skin may turn yellow due to another factor, called icterus or jaundice, which occurs with serious liver disease. In this instance, bile pigments are deposited within the skin and impart a yellow color to it.
2. The amount of oxygen-saturated hemoglobin found in the blood vessels of the middle layer of our skin, the dermis. Hemoglobin is the iron-containing protein pigment of our blood cells. A lack of oxygen saturation imparts a paler, grayer, or bluer color to the skin. Skin may also become paler as a result of anemia (a reduced number of hemoglobin and/or red blood cells), low blood pressure, or poor circulation of blood.

3. Conversely, light-skinned individuals (compared to dark-skinned ones) may have a rosy effect to their skin thanks to the relatively more oxygen-rich hemoglobin flowing through the blood vessels of their dermis. Red-colored skin may also occur as a result of blood vessels in or near the skin dilating (expanding) due to embarrassment, fever, allergy, or inflammation.
4. Finally, the skin may have red, black, blue, purple, and green bruises—all as a result of the escape of blood into surrounding tissues. As the blood (namely, the hemoglobin) disintegrates and is processed and removed by various cells, it and the bruise changes color with time.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Cutaneous. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cutaneous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- integumentary system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/integumentary_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skin Layers. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...kin_layers.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- stratum germinativum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/stratum%20germinativum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epidermis (skin). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epidermis_(skin)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- keratinocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/keratinocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Integumentary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P..._System%23Skin**. **Project**: Human Physiology. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Merkel cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Merkel%20cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human skin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vitamin D. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vitamin...on_in_the_skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Dermis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Dermis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypodermis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypodermis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- connective tissue. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/connective_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- basement membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/basement%20membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mechanoreceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mechanoreceptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Melanocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Melanocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pigmentation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pigmentation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- melanin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/melanin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- melanocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/melanocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- stratum corneum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/stratum_corneum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skin tanning. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Skin_tanning.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu skin02. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_skin02.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 5.1D: Skin Color is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.2: CELL JUNCTIONS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

5.2A: Tight Junctions

5.2B: Adherens Junctions

5.3C: Gap Junctions

This page titled 5.2: Cell Junctions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.2A: Tight Junctions

Tight junctions serve as selectively permeable seals in our body's internal and external surfaces.

LEARNING OBJECTIVE

Describe the characteristics of tight junctions

Key Takeaways

Key Points

- Tight junctions are the closely associated areas of two cells whose membranes join together to form a virtually impermeable barrier to fluid.
- Tight junctions perform vital functions—such as holding cells together—and form protective and functional barriers.
- Tight junctions are composed of a branching network of sealing strands with each strand acting independently from the others.
- The major types of proteins in junctions are the claudins and the occludins.
- Each strand is formed from a row of transmembrane proteins embedded in both plasma membranes, with extracellular domains joining one another directly.

Key Terms

- **blood-brain barrier**: A structure in the central nervous system (CNS) that keeps various substances found in the bloodstream out of the brain while allowing in the substances essential to metabolic function, e.g., oxygen.
- **Claudins**: Proteins that form the backbone of the tight junction strands.
- **cell adhesion molecule**: Molecules that help cells stick to each other and to their surroundings. The proteins located on the cell surface bind with other cells or with the extracellular matrix (ECM).
- **cytoskeleton**: A cellular structure like a skeleton, contained within the cytoplasm.
- **epithelia**: The covering of internal and external body surfaces, where tight junctions are found.
- **zonula occludens**: Another name for tight junctions.



Tight Junction: An electron micrograph showing a tight junction in rat kidney tissue. The three dark lines of density correspond to the tight junction and the light lines in between correspond to the paracellular space.

Imagine a largely waterproof zipper connecting the sides of two different jackets. That zipper is like a tight junction (TJ), also called an occluding junction. A TJ creates a small zone that occludes the extracellular space (the space between cells).


This is why tight junctions are also called zonula occludens. The word zonula comes from words that mean small zone or encircling belt, while occludens comes from the Latin word occludere, which means to close up.

Location and Function

Tight junctions are virtually (but also partly selectively) impermeable seals that encircle cells and bind them together into leakproof sheets. In other words, the plasma membranes of adjacent cells essentially fuse together tightly in order to limit the leakage of various substances between the two cells.

What can and cannot go through all depends on the substance's size, charge, as well as the location and precise composition of the tight junctions in the part of the body in question.

Tight junctions are located within our body's epithelia. Epithelia is the plural of epithelium. Epithelium is a word that refers to the covering of the body's internal and external surfaces. This includes organs (such as skin), blood vessels, and cavities.



Occludin: Model of the protein structure of the coiled-coil domain of human occludin.

Thus, these tight junctions serve various functions, depending on what epithelium is in question. In the skin, they keep us somewhat watertight and help keep allergens out of our body. In the digestive system, they help prevent the leakage of digestive enzymes into our bloodstream.

Tight junctions also serve as a structural support mechanism that help keep the epithelium together.


Composition

A tight junction—a kind of symmetrical cell junction—is composed of numerous important proteins that are either directly involved in its composition or intimately involved with connecting the tight junction to and between the cells in one way or another. These proteins include:

- Occludins, which maintain the barrier between adjacent cells.
- Claudins, which form the backbone of tight junction strands.

- Junctional adhesion molecules (JAMs) are immunoglobulin (antibody) proteins that help seal the intercellular space between two cells.
- Zonula occludens (ZO) are proteins that help link the tight junction to each cell's internal skeleton (cytoskeleton).

The occludins and claudins are the major components of tight junction strands. When fully formed, a tight junction is not one, long, continuous seal. Instead, it looks like a series of local seals joined together in a maze-like fashion.



Tight junction: Diagram of tight junction components.

This page titled 5.2A: Tight Junctions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.2B: Adherens Junctions

Adherens junctions provide strong mechanical attachments between adjacent cells through the linkage of cytoplasmic face with cytoskeleton.

LEARNING OBJECTIVE

Describe the characteristics of adherens junctions

Key Takeaways

Key Points

- Adherens junctions are involved in a number of critical functions, including providing additional structural support. For example, they hold cardiac muscle cells tightly together as the heart expands and contracts.
- Adherens junctions are built primarily from cadherins, whose extracellular segments bind to each other and whose intracellular segments bind to catenins. Catenins are connected to actin filaments.

Key Terms

- **cadherin**: Any of a class of transmembrane proteins important in maintaining tissue structure.
- **adherens junctions**: Protein complexes that occur at cell–cell junctions in epithelial tissues; they are usually more basal than tight junctions.
- **catenin**: Any of a class of proteins that have a role in cell adhesion.

Adherens junctions are also referred to as zonula adherens, intermediate junction, or as belt desmosomes. Zonula means small zone or belt-like, and adherens refers to adhesion (sticking together). As a result, the zonula adherens often runs like a belt around the entire cell in a continuous fashion, and it acts as an adhesion belt.

Location and Function

This type of cell junction is located right below tight junctions and provides a strong bond between the sides of adjacent epithelial cell membranes. While other junctions, like tight junctions, provide some support for and fusion of adjacent cells, their resistance to mechanical stress is relatively small compared to the much stronger adherens junctions.

Structure and Composition


The zonula adherens is composed of several different proteins:

- The actin microfilaments of the cytoskeleton (the internal skeleton of the cell).
- Anchor proteins, found inside each cell. These are called alpha-catenin, beta-catenin, gamma-catenin (aka plakoglobin), vinculin, and alpha-actinin. They link the actin microfilaments to the cadherins.
- Cadherins, namely E-cadherin. These are transmembrane adhesion proteins, whose main portions are located in the extracellular space.

The extracellular part of one cell's cadherin binds to the extracellular part of the adjacent cell's cadherin in the space between the two cells. Each cell's cadherin molecule also contains a tail that inserts itself inside its respective cell.

This intracellular (within the cell) tail then links up to catenin proteins to form the cadherin–catenin complex. This complex binds to vinculin and alpha-actinin; these two proteins are what link the cadherin–catenin complex to the cell's internal skeletal framework (the actin microfilaments).

The extracellular portions of the cadherin molecules of adjacent cells are bonded together by calcium ions (or another protein in some cases). This means that the functional as well as morphological integrity of the adherens junctions are calcium dependent. If you were to remove calcium from the equation, this type of cell junction would disintegrate as a result.



The structural proteins in an adherens junction: These are the principal interactions of structural proteins at a cadherin-based plasma membrane adherens junction. Actin filaments are associated with adherens junctions in addition to several other actin-binding proteins.

This page titled 5.2B: Adherens Junctions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3C: Gap Junctions

A gap junction is a specialized cell junction that directly connects the cytoplasm of two cells.

LEARNING OBJECTIVE

Describe the characteristics of gap junctions

Key Takeaways

Key Points

- Gap junctions allow various molecules and ions to pass freely between cells.
- A gap junction channel is composed of two connexons, also known as hemichannels that line up across the intercellular space.
- Most gap junction hemichannels are composed of a complex of six connexin proteins, each characterized by four transmembrane domains. Six connexin sub-units assemble to create one connexon, or hemichannel.
- Channel composition influences the function of the gap junction.
- Gap junctions allow for electrical communication between cells, and also allow the passage of small second messengers.
- Gap junctions are expressed in virtually all tissues and cells, but most notably in cell types that are involved in direct electrical communication, such as neurons and cardiac muscle.

Key Terms

- **cytoplasm**: The contents of a cell except for the nucleus. It includes cytosol, organelles, vesicles, and the cytoskeleton.
- **connexin**
- **connexon**
- **nexus**: An alternative name for a gap junction.

Gap junctions are also called communicating junctions, macula communicans, or nexuses. These are connections that allow for the direct passage of molecules between two cells.


Gap junctions consist of a number of transmembrane channels called pores that are found in a closely packed arrangement. The number of gap junctions shared between two cells can vary as well.

Structure

Each gap junction channel is made up of two half channels (hemichannels), one in each cell's membrane. These half channels join together, bridge the extracellular space in the process, and form the entire channel that spans both cell membranes.

Each of these half channels is called a connexon. Each connexon is made up of six symmetrical integral membrane protein units called connexins. This means each channel is made up of 12 circularly arranged protein units.

Function



Intercalated disk in heart muscle contains gap junctions: Intercalated disks consist of three different types of cell–cell junctions: actin filaments anchoring adherens junctions, intermediate filaments anchoring desmosomes, and gap junctions. Gap junctions are responsible for electrochemical and metabolic coupling

The molecules that may cross this channel include the likes of ions, regulatory proteins, and metabolites (products of metabolism). Examples of this includes calcium ions and cAMP (cyclic adenosine monophosphate).


Depending on the type of gap junction in question, molecules can pass evenly in both directions, or asymmetrically, so in some gap junctions the molecules will move in one direction faster than in the other direction.

The channels in a gap junction aren't always open. They fluctuate between being open and closed. The ability of the channel to open or close is made possible in part to calcium ions, which induce a reversible conformational change in the connexin molecules, which leads to the closure of a channel at its extracellular surface. The cytoplasmic end of each connexon can also be closed, if necessary.

Location

Gap junctions are found in many places throughout the body. This includes epithelia, which are the coverings of body surfaces, as well as nerves, cardiac (heart) muscle, and smooth muscle (such as that of the intestines).

Their primary role is to coordinate the activity of adjacent cells. For instance, when heart cells need to beat in unison, gap junctions allow for the transmission of electrical signals between the cells.



Gap junction: The major molecular components of the gap junction.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Cell junctions. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cell_junctions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tight junction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tight_junction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Occludin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Occludin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cell adhesion molecule. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cell%20adhesion%20molecule**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cytoskeleton. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cytoskeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Claudins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Claudins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- blood-brain barrier. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/blood-brain_barrier**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adherens junctions. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/adherens%20junctions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cadherin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cadherin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- catenin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/catenin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Junctions Between Cells. **Provided by**: Kimball's Biology Pages. **Located at**: **http://biology-pages.info/J/Junctions.html**. **License**: ***CC BY: Attribution***
- Adherens junction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adherens_junction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gap junction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gap_junction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synapse. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/synapse**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cytoplasm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cytoplasm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intercellular. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/intercellular**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gap cell junction-en. **Authored by**: LadyofHats. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/**

Fi...unction-en.svg. **License**: ***Public Domain: No Known Copyright***
- Glanzstreifen. **Authored by**: Dr. S. Girod, Anton Becker. **Located at**: **en.Wikipedia.org/wiki/File:Glanzstreifen.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 5.3C: Gap Junctions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3: ACCESSORY STRUCTURES OF THE SKIN

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

5.3A: Sweat (Sudoriferous) Glands

5.3B: Sebaceous (Oil) Glands

5.3C: Nails

5.3D: Hair

This page titled 5.3: Accessory Structures of the Skin is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3A: Sweat (Sudoriferous) Glands

Sweat glands, also known as sudoriferous glands, are distributed over most of the body surface.

LEARNING OBJECTIVE

Classify eccrine and apocrine sweat glands

Key Takeaways

Key Points

- Sweat glands are located deep within the skin and primarily regulate temperature.
- The two main types of sweat glands are eccrine sweat glands and apocrine sweat glands.
- Eccrine sweat glands are smaller sweat glands. They are coiled tubular glands that discharge their secretions directly onto the surface of the skin.
- Apocrine sweat glands are coiled tubular glands that discharge in the canals of hair follicles. The sweat produced may be acted upon by bacteria, causing a noticeable odor.

Key Terms

- **eccrine gland**: The major sweat glands of the human body, found in virtually all skin, produce a clear, odorless substance, consisting primarily of water and NaCl.
- **apocrine sweat gland**: The type of sweat gland that is least responsible for thermoregulation and most responsible for body odor.


Sweat glands, also called sudoriferous glands, are simple tubular glands found almost everywhere on our body. Each sweat gland is made up of two portions:

1. A secretory section
2. An excretory duct

The secretory portion is found in the dermis, the middle layer of the skin. Sometimes it's also found in the hypodermis, the deepest layer of our skin.

The secretory portion of a sweat gland is a twisted and coiled tube that has an opening at its very top. It is in the coiled secretory portion of the sweat gland where the sweat is actually produced. The excretory duct moves from the secretion portion, through the dermis, and into the topmost layer of the skin, the epidermis, where it opens up at the surface of our skin.

Eccrine Glands



Human Skin: Cross sectional image of skin showing a sweat gland and a sebaceous gland.

The most numerous types of sweat glands in our skin, found almost everywhere on the body, are called eccrine glands. These are the true sweat glands in the sense of helping to regulate body temperature. In other words, sweating causes the loss of body heat and thus cools us down on a hot day or when performing strenuous exercise. This is because as the water in sweat evaporates, it takes body heat with it.

Apocrine Glands

The other kind of sweat glands are known as apocrine glands. The apocrine glands are found in places like the armpits, scrotum, anus, and labia majora. They are typically larger than eccrine glands and their ducts tend to open into hair follicles instead of hairless areas of skin.

These glands, unlike the eccrine glands, serve virtually no role in the regulation of body temperature. These are also the glands largely responsible for body smells, as their excretions are converted by skin bacteria into various chemicals we associated with body odor.

Unlike eccrine glands, the exact function of apocrine glands is unknown and debated. We do know they are activated during times of stress, pain, and sexual foreplay but for what reasons is yet to be made clear.

This page titled 5.3A: Sweat (Sudoriferous) Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3B: Sebaceous (Oil) Glands

Sebaceous glands are found in most of the skin (except the palms of the hands and the soles of the feet).

LEARNING OBJECTIVE

Describe the location and function of sebaeous glands

Key Takeaways

Key Points

- Sebaceous glands are located throughout the skin except in the palms of the hands and soles of the feet.
- Sebum is an oily substance composed of fat (lipids) and the debris of dead fat-producing cells.
- Sebaceous glands are classified as holocrine glands.

Key Terms


- **sebum**: A thick oily substance, secreted by the sebaceous glands of the skin, that consists of fat and cellular debris.
- **holocrine gland**: The sebaceous gland is an example of a holocrine gland because its product of secretion (sebum) is released with remnants of dead cells.

Sebaceous glands are the oil secreting glands of your body. This is why they are also called the oil glands. They are a type of holocrine simple saccular (alveolar) gland. Their function is to secrete a substance called sebum, a mixture of fatty substances, entire sebum-producing cells, and epithelial cell debris. The sebaceous glands are located in the dermis, the middle layer of the skin, and they develop from the epithelial cells of the hair follicle itself (the external root sheath of the hair follicle).

Sebaceous gland ducts thus usually open up into the upper part of a hair follicle, called the infundibulum. The infundibulum is part of the pilosebaceous canal, the one responsible for discharging sebum and one that is composed of the infundibulum and the short duct of the sebaceous gland itself. However, some sebaceous gland ducts open directly onto our skin surface such as at the corner of the mouth and the glans penis. Regardless, the secretion of sebum out of the gland is helped along by the contraction of the arrector pili muscle.

While the sebaceous glands are present just about all over the skin, they are notably absent on the palms of the hands and the soles of the feet. The sebum being excreted by your body today began production around 8 days ago.

The Function of Sebum



Sebaceous Gland: Schematic view of a hair follicle with sebaceous gland.

The sebum produced by these glands plays numerous important roles:

1. Sebum is a lubricant and inasmuch it helps to moisturize the skin. It does so by preventing the excess evaporation of water from the skin.
2. Sebum serves to keep us healthy by keeping in check the growth of certain bacteria on our skin. That's because sebum contains chemicals that kill bacteria. This helps ensure bacteria don't invade into deeper layer of our skin.
3. It helps to condition the hair. Meaning, it ensures our hair doesn't become too dry and brittle.

Sebaceous glands are involved in numerous conditions. During puberty, various hormones cause them to produce a lot of sebum and this therefore contributes to oily skin. If a duct of a sebaceous gland is clogged with sebum, a whitehead results. If this material is allowed to dry and oxidize, it will become darker, forming a blackhead. If a sebaceous gland becomes infected, moderate and severe forms of acne are the result.

The glands lining the ear canal that produce earwax (cerumen) are called ceruminous glands. They are modified sebaceous glands.

This page titled 5.3B: Sebaceous (Oil) Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3C: Nails

Fingernails are made of keratin and they perform two major functions: protection and sensation.

LEARNING OBJECTIVE

Describe the structure of fingernails

Key Takeaways

Key Points

- The nail bed contains the blood vessels, nerves, and melanocytes or melanin-producing cells. As the nail is produced by the root, it streams down along the nail bed, which adds material to the undersurface of the nail and makes it thicker.
- The nail plate is the actual fingernail, composed of translucent keratin. The pink appearance of the nail comes from the blood vessels underneath the nail.
- The eponychium, or cuticle, is situated between the skin of the finger and the nail plate. It fuses these structures together and provides a waterproof barrier.
- Deformity or disease of the nails is referred to as onychosis. There are many diseases that can occur with the fingernails and toenails. The most common of these diseases are ingrown nails and fungal infections.
- Ingrown nails, also known as onychocryptosis, can affect either the fingers or the toes. In this condition, the nail cuts into one or both sides of the nail bed, resulting in inflammation and possibly infection.

Key Term

- **keratin**: A protein that makes up hair and nails.

Function of the Fingernail

The fingernail is an important structure made of keratin. The fingernail generally serves two purposes: it acts as a protective plate and enhances sensation of the fingertip. Nails can also help grasp small things.

The protection function of the fingernail is commonly known, but the sensation function is equally important. The fingertip has many nerve endings in it that allow it to receive volumes of information about the objects we touch. The nail acts as a counterforce to the fingertip, providing even more sensory input when an object is touched.

Nails grow from the nail bed continuously but they slow down their growth rate with age, poor nutrition, or poor circulation.

Anatomy of the Fingernail

The structure of the fingernail is divided into six specific parts:

- root
- nail bed
- nail plate
- eponychium (cuticle)
- perionychium
- hyponychium

Root and Nail Sinus

The nail sinus (sinus unguis) is where the nail root is—at the base of the nail underneath the skin. It originates from the actively growing tissue below, the matrix. The root of the fingernail is also known as the germinal matrix.

This portion of the nail is actually beneath the skin, behind the fingernail, and extends several millimeters into the finger. The fingernail root produces most of the volume of the nail and the nail bed. This portion of the nail does not have any melanocytes, or melanin-producing cells. The edge of the germinal matrix is seen as a white, crescent shaped structure called the lunula.

Nail Bed

The nail bed is a part of the nail matrix called the sterile matrix. It extends from the edge of the germinal matrix, or lunula, to the hyponychium. The nail bed contains the blood vessels, nerves, and melanocytes, or melanin-producing cells. As the nail is produced by the root, it streams down along the nail bed, which adds material to the undersurface of the nail and makes it thicker.

Nail Plate

The nail plate is the actual fingernail, composed of translucent keratin. The pink appearance of the nail comes from the blood vessels underneath the nail. The underneath surface of the nail plate has grooves along the length of the nail that help anchor it to the nail bed. The free margin or distal edge is the anterior margin of the nail plate corresponding to the abrasive or cutting edge of the nail.

Eponychium

The eponychium, or cuticle, is situated between the skin of the finger; the nail plate fuses these structures together and provides a waterproof barrier.

Perionychium

The perioncyhium is the skin that overlies the nail plate on its sides; it is also known as the paronychial edge. The perionychium is the site of hangnails, ingrown nails, and an infection of the skin called paronychia.

Hyponychium

The hyponychium is the area between the nail plate and the fingertip. It is the junction between the free edge of the nail and the skin of the fingertip, also providing a waterproof barrier.

This page titled 5.3C: Nails is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.3D: Hair

Hair growth occurs from the hair follicle.

LEARNING OBJECTIVE

Describe the characteristics of body hair

Key Takeaways

Key Points


- Hair primarily serves for protection, warmth, and sensation.
- Human hair is made of keratin.
- The structure of the hair follicle includes the papilla, matrix, root, and bulb.
- The different types of hair on the human body include lanugo, vellus hair, and terminal hair.
- Attached to the follicle is a tiny bundle of muscle fiber called the arrector pili.

Key Terms

- **hair follicle**: The structure that produces hair.
- **papilla**: A structure that provides nutrients that help our hair grow.

Hair is present on almost the entire surface of our body, excluding certain regions such as the palms of our hands, soles of our feet, and some genital areas. In certain places, hair is so small that it is virtually invisible to the naked eye and in other places it is quite obvious, like on our head and in our armpits.

Structure



Hair follicle: Cross section of a hair follicle.

A hair can be divided into two main parts lengthwise:

- The root is part of the hair enclosed by the hair follicle, which is itself a tube-like involution of the skin.
- The shaft is the part of the hair projecting from the surface of our skin. A round shaft results in straight and coarse hair. An oval hair shaft is responsible for wavy hair. A flat shaft causes curly hair.

Each hair can also be broken into three main parts. They are, from the most superficial to the deepest:

- Cuticle, the outermost portion. This layer is composed of scale-like cells that seem to overlap in a shingle-like manner. Such an arrangement helps prevent hair from matting.
- Cortex, the middle portion. This portion is made up of layers of elongated and flattened cells.
- Medulla, the inner (central) portion. This is made up of large cells with many sides and air spaces.

There are three types of hairs:

- Lanugo, found on the fetus and for a bit after birth.
- Vellus, fine body hair (peach-fuzz hair).
- Terminal hair, the coarse hair.

Growth and Composition

In the deepest portion of the each hair follicle lies the hair bulb. It is supplied with nutrients via blood by a structure called the hair papilla. The hair bulb has a growth zone called the matrix. Here, stratum basale epithelial cells divide via mitosis to form the hair.

As new cells form, they push older cells up to the surface. These older cells die and become keratinized in the process. This means that most hair is made up of protein (keratin).

The hair bulb also contains cells called melanocytes that produce various kinds of melanin pigments. Their varying combinations are what help to produce the different natural hair colors people have.


Hair grows in three stages:

- Anagen phase, the fast growing phase.
- Catagen phase, the involution phase.
- Telogen phase, the rest phase.

Purpose

Depending on the kind of hair and location, hair can have one of several purposes:

- Expression. Hair on our eyebrows may have stuck around on our body to help gauge a person's emotions or intent.
- Protection. Eyelashes help protect our eyes and nose hairs keep things out of our respiratory system. Hair on the head helps add a small cushion against bumps.
- Sensation. Hair helps us sense light touches.
- Warmth and defense. Attached to a hair follicle is a bundle of muscle fibers. They help comprise the arrector pili muscle that causes the hairs on our body to stand on their ends when we're cold or we're scared. The standing of the hair on end also dimples the skin and thus produces goosebumps. In non-human mammals this action may help add an insulating layer of air between the hair (called fur in non-human mammals) and skin, or it may be used as a way to make a scared animal look larger to its enemy. Neither function is all that relevant or useful in humans, although some warmth is provided by scalp hair.



Skin overview: Skin layers, of both hairy and hairless skin.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- eccrine gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Eccrine_sweat_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Integumentary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Integumentary_System**. **Project**: Human Physiology. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sebaceous glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sebaceous_glands**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- holocrine gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Holocrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sebum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sebum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nail (anatomy). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nail_(anatomy)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- keratin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/keratin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hair follicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hair_follicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hair. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hair**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- papilla. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/papilla**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hair fiber. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hair%20fiber**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human skin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 5.3D: Hair is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4: FUNCTIONS OF THE INTEGUMENTARY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

5.4A: Protection

5.4B: Thermoregulation

5.4C: Cutaneous Sensation

5.4D: Metabolic Functions

5.4E: Blood Supply to the Epidermis

5.4F: Excretion and Absorption

This page titled 5.4: Functions of the Integumentary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4A: Protection

The skin provides an overlaying protective barrier from the environment and pathogens while contributing to the adaptive immune system.

LEARNING OBJECTIVE

Describe the ways in which the integumentary system protects the body

Key Takeaways

Key Points


- The skin provides a protective barrier from the external environment and prevents dehydration.
- Langerhans cells in the skin also contribute to protection as they are part of the adaptive immune system.
- The integumentary system protects the body's internal living tissues and organs, protects against invasion by infectious organism, and protects the body from dehydration.

Key Terms

- **vitamin D**: An important vitamin synthesized thanks to the skin.
- **melanocytes**: Cells that help protect our body from radiological damage.
- **Langerhans cells**: Langerhans cells are dendritic cells (antigen-presenting immune cells) of the skin and mucosa that contain large granules.

The skin helps protect our body's internal structures from physical, chemical, biological, radiological, and thermal damage as well as damage from starvation and malnutrition.

Physical and Chemical Damage



Human skin: A diagram of human skin.

The skin is composed of tough skin cells as well as a tough protein called keratin that guard tissues, organs, and structures underneath the skin against physical damage from minor cuts, scratches, and abrasions. Because our skin is tough and largely waterproof, it helps protect internal structures from chemical irritants such as man-made detergents or even natural irritants like poison ivy.

Otherwise, these dangerous chemicals would seep into our sensitive internal environment. The waterproof nature of our skin also ensures that important molecules stay within our body.

Biological Damage

The skin also contains important cells called Langerhans cells. These cells help our immune system fight off infectious biological agents, like bacteria that try to get further into our body through skin that may have been compromised by physical damage.

Sebaceous glands associated with the skin secrete substances that help fight off potentially dangerous microorganisms as well. These glands also secrete substances that help keep our skin hydrated, and thus more resistant to bacterial invasion.

Radiological Damage

Our skin also contains melanocytes that produce a pigment called melanin. This protects the body from radiological damage via the sun's UV radiation (or that from tanning beds).

Other Protective Roles

Part of our skin is made up of fat. This fat serves three large purposes:

1. It helps cushion internal structures against any physical blows.
2. It acts as a food source, protecting our body from the effects of starvation.
3. It helps insulate us against cold temperatures.

Our skin is also closely associated with sweat glands that help protect us from high temperatures by cooling us off through the process of evaporation. These glands also help to excrete potentially dangerous substances, like urea, out of the body.

All sorts of sensory receptors are found within the skin as well. These help move our body parts away from potential sources of damage, like hot stoves, when they sense danger, thereby protecting our body from great harm.

Finally, the skin is also important for the synthesis of vitamin D, which is an important vitamin for the building of strong and healthy bones. Ergo, the skin protects the body from fractures if we do not otherwise get enough of this vitamin from food-based sources.

This page titled 5.4A: Protection is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4B: Thermoregulation

The integumentary system keeps body temperature within limits even when environmental temperature varies; this is called thermoregulation.

LEARNING OBJECTIVE

Explain the skin's role in thermoregulation

Key Takeaways


Key Points

- The skin's immense blood supply helps regulate temperature: dilated vessels allow for heat loss, while constricted vessels retain heat.
- The skin regulates body temperature with its blood supply.
- The skin assists in homeostasis.
- Humidity affects thermoregulation by limiting sweat evaporation and thus heat loss.

Key Terms

- **Evaporation**: What happens when water crosses the skin via sweat glands and then dissipates into the air; this process cools body temperature to within the body's tolerance range.
- **homeostasis**: The ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium; such as the ability of warm-blooded animals to maintain a constant temperature.
- **vasoconstriction**: The constriction (narrowing) of a blood vessel.
- **arrector pili**: Any of the small muscles attached to hair follicles in mammals; when the muscles contract they cause the hairs to stand on end.

The integumentary system functions in thermoregulation—the ability of an organism to keep its body temperature within certain boundaries—even when the surrounding temperature is very different. This process is one aspect of homeostasis: a dynamic state of stability between an animal's internal and external environment.



Human skin: This image details the parts of the integumentary system.

The skin assists in homeostasis (keeping different aspects of the body constant, e.g., temperature). It does this by reacting differently to hot and cold conditions so that the inner body temperature remains more or less constant.

The Skin's Role in Cooling the Body

The skin is an incredibly large organ. It is about 2 meters squared (depending on the size of the individual). Owing to its location at the barrier of the environment and our internal selves, and its relatively very large surface area, it is plays an incredibly important role in thermoregulation.

This is because in a healthy individual, when all else is held equal, their body is constantly generating heat as a result of its various metabolic and physical processes. At rest, such an individual is expected to increase their body temperature by 1 C every 5 minutes as a result of these processes. Left unregulated, this would kill a person quite quickly.

The process of skin-based thermoregulation occurs through several means. The first way involves the abundance of blood vessels found in the dermis, the middle layer of the skin. If the body must cool down, the body vasodilates these blood vessels.

Radiation

Vasodilation refers to the process of expanding (-dilation) the size of the blood vessels (vaso-). The now enlarged peripheral vessels of the skin allow for greater amounts of blood to flow near the surface of the skin. This allows for our body to release a lot of body heat through radiation. Radiation, in this case, refers to thermal radiation, which is the process of transferring heat through space via electromagnetic waves.

Convection

At the same time, if a fluid such as circulating air or water in a pool comes into contact with the skin when we are very hot, this will allow for heat loss through the process of convection. The higher the amount of our body surface exposed to this (usually) circulating air (e.g. as little clothing as possible), the higher the speed of the circulating air (e.g. it's really windy), and the smaller the distance between the skin surface and the blood vessels, the greater the loss of heat from our body via convection.

Conduction

If our skin touches a cold object (like a cold drink), then we will lose heat via the process of conduction, which is the direct heat transfer of heat from a hotter surface, to a colder surface touching that hotter surface.

Perspiration

The body also thermoregulates via the process of sweating (perspiration). Roughly speaking, sweating begins when the body temperature rises above 37 C. Sweat production can be increased or lowered as necessary.

For instance, if we must cool down, sweat production increases. As drops of sweat form on and then evaporate from our skin surface, they take body heat away with them. All else held equal, the greater the skin surface area and the higher the sweat rate, the greater the rate of cooling via sweating.

With respect to body heat loss, the processes of radiation and convection are most effective when the environmental temperature is below 20 C, while evaporative cooling accounts for the most heat loss when the environmental temperature is above 20 C, and especially when it's hotter than 35 C.

Increased humidity, however, limits the ability of our body to dissipate heat via perspiration.


Arrector Pili Muscles

The hairs on the skin lie flat and prevent heat from being trapped by the layer of still air between the hairs. This is caused by tiny

muscles under the surface of the skin, called arrector pili muscles.

When these muscles relax their attached hair follicles are not erect. These flat hairs increase the flow of air next to the skin and increase heat loss by convection. The exact extent to which this process help keep us cool is debated (read below).

The Skin's Role in Keeping Us Warm



Anatomy of the skin: The skin is the largest organ of the integumentary system, made up of multiple layers of ectodermal tissue, and guards the underlying muscles, bones, ligaments, and internal organs.

On the other hand, if the body needs to prevent the loss of excess heat, such as on a cool day, it will end up constricting the blood vessels of our skin. This process is known as vasoconstriction.

Since the blood vessels are narrower than they were before, less blood flows through the skin and thus less heat can escape into the environment via radiation, convection, and conduction. The body will also limit or stop the process of sweating to minimize any evaporative heat loss.

In addition, our body thermoregulates using our hair. The arrector pili muscles contract (piloerection) and lift the hair follicles upright. This makes the hairs stand on end, which acts as an insulating layer, trapping heat. This is also how goose bumps are caused, since humans don't have very much hair and the contracted muscles can easily be seen.

While this hair-based method of thermoregulation is effective in many mammals and birds owing to their large and thick amounts of fur and feathers (respectively), the relative effectiveness of this method of thermoregulation in humans is in question since we have little to no body hair in comparison.

Finally, while technically not a thermoregulatory mechanism, the fat associated with our skin does help insulate our body and therefore increases body temperature as a result.

This page titled 5.4B: Thermoregulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4C: Cutaneous Sensation

The somatosensory system is composed of the receptors and processing centers to produce the sensory modalities, such as touch and pain.

LEARNING OBJECTIVE

Differentiate among the types of cutaneous mechanoreceptors

Key Takeaways

Key Points

- The somatosensory is the system of nerve cells that responds to changes to the external or internal state of the body.
- Receptors are spread throughout the body, with large numbers found in the skin.
- Several distinct receptor types form the somatosensory system including thermoreceptors (heat), nociceptors (pain), and mechanoreceptors (pressure).
- There are four types of mechanoreceptors that respond to different pressure stimui and provide a wide range of mechanical sensitivity—they are the keys for fine motor control.

Key Terms

- **somatosensory system**: A diverse sensory system composed of the receptors and processing centers to produce the sensory modalities such as touch, temperature, proprioception (body position), and nociception (pain).
- **sensory receptor**: A sensory nerve ending that recognizes a stimulus in the internal or external environment of an organism.

The Somatosensory System

The somatosensory is the system of nerve cells that responds to changes to the external or internal state of the body, predominately through the sense of touch, but also by the senses of body position and movement.

Spread through all major parts of the body, it consists of sensory receptors and sensory neurons in the periphery (for example, skin, muscle, and organs), along with deeper neurons within the central nervous system.

While touch is considered one of the five traditional senses, the impression of touch is actually formed from several diverse stimuli using different receptors:

- Thermoreceptors (temperature)
- Nociceptors (pain)
- Mechanoreceptors (pressure)

Transmission of information from the receptors passes via sensory nerves through tracts in the spinal cord and into the brain. Processing primarily occurs in the primary somatosensory area in the parietal lobe of the cerebral cortex.

Thermoreceptors

Mammals have at least two types of sensors: those that detect heat and those that detect cold.

Upon deviation from the norm ,sensory receptors trigger an action potential that can provide feedback or lead to alterations in behavior in order to maintain homoeostasis. Two receptors that exhibit the ability to detect changes in temperature include Krause end bulbs (cold) and Ruffini endings (heat).

Nociceptors

A nociceptor is a sensory nerve cell that responds to damaging or potentially damaging stimuli by sending signals to the spinal cord and brain. Nociceptors can respond to excessive thermal, mechanical, or chemical stimulation and often result the generation of an involuntary motor respons—for example, pulling a hand away from a hot surface.

Mechanoreceptors

Mechanoreceptors are sensory receptors that respond to pressure and vibration. Four key types of mechanoreceptor have been described based on their response to stimulation and receptive field.

Receptors can either induce a slow response to stimulation, whereby a constant activation is initiated, or a fast response, whereby activation is only initiated at the beginning and end of stimulation. The receptive field—the region in which a receptor can sense an effect—can vary from small to large.

- The Merkel receptor is a disk-shaped receptor located near the border between the epidermis and dermis. It demonstrates a slow response and has a small receptive field; it is useful for detecting steady pressure from small objects, such as when gripping something with the hand.
- The Meissner corpuscle is a stack of flattened cells located in the dermis, near the epidermis. It demonstrates a rapid response and has a small receptive field; it is useful for detecting texture or movement of objects against the skin.
- The Ruffini cylinder is located in the dermis and has many branched fibers inside a cylindrical capsule. It demonstrates a slow response and has a large receptive field; it is good for detecting steady pressure or stretching, such as during the movement of a joint.
- The Pacinian corpuscle is a layered, onion-like capsule surrounding a nerve fiber. It is located deep in the dermis, in the subcutaneous fat. It demonstrates a fast response and has a large receptive field; it is useful for detecting large changes in the environment, such as vibrations.

Together they provide a wide range of mechanical sensitivity that enables fine motor control.

This page titled 5.4C: Cutaneous Sensation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4D: Metabolic Functions

One of the metabolic functions of the skin is the production of vitamin D3 when ultraviolet light reacts with 7-dehydrocholesterol.

LEARNING OBJECTIVE

Describe the integumentary system's role in producing vitamin D

Key Takeaways

Key Points

- Vitamin D refers to a group of fat-soluble steroids responsible for increasing intestinal absorption of calcium, iron, magnesium, phosphate, and zinc.
- Foods rich in vitamin D are relatively scarce and so the body synthesises the majority of vitamin D itself, in the skin.
- Vitamin D deficiency is associated with poor development of bones in children and a softening of bones in adults.
- Vitamin D_3 is made in the skin when 7-dehydrocholesterol reacts with ultraviolet light. Vitamin D is produced in the two innermost strata of the epidermis. Cholecalciferol (D_3) is produced photochemically in the skin from 7-dehydrocholesterol.
- Vitamin D is produced in the two innermost strata of the epidermis, the stratum basale and stratum spinosum.

Key Term

- **7-dehydrocholesterol**: 7-dehydrocholesterol is a cholesterol precursor that is converted to vitamin D3 in the skin, therefore functioning as provitamin-D3.


The integumentary system is the largest of the body's organ systems, made up of the skin and its associated appendages. The integumentary system distinguishes, separates, and protects the organism from its surroundings, but also plays a key metabolic function, as the major region for vitamin D production.

What is Vitamin D?

Vitamin D refers to a group of fat-soluble steroids responsible for increasing intestinal absorption of calcium, iron, magnesium, phosphate, and zinc. In humans, the most important compounds in this group are vitamin D_3 (also known as cholecalciferol) and vitamin D_2 (ergocalciferol). Cholecalciferol and ergocalciferol can be ingested from the diet and from supplements, however very few foods are rich in vitamin D; and so synthesis within the skin is a key source.

Vitamin D deficiency is associated with impaired bone development in children, which leads to the development of rickets and a softening of bones in adults. Deficiency in vitamin D has been termed a modern disorder associated with both a poorer diet and reduced time spent outside.

Vitamin D Synthesis




Vitamin D: The chemical structure of vitamin D.

The human skin consists of three major layers: the epidermis, dermis, and hypodermis. The epidermis forms the outermost layer, providing the initial barrier to the external environment. Beneath this, the dermis comprises two sections, the papillary and reticular layers, and contains connective tissues, vessels, glands, follicles, hair roots, sensory nerve endings, and muscular tissue. The deepest layer is the hypodermis, which is primarily made up of adipose tissue.

Vitamin D is produced in the two innermost strata of the epidermis, the stratum basale and stratum spinosum.

Vitamin D_3 is made in the skin when the 7-dehydrocholesterol reacts with ultraviolet light of UVB type at wavelengths between 280 and 315 nm, with peak synthesis occurring between 295 and 297 nm.

Depending on the intensity of UVB rays and the minutes of exposure, an equilibrium can develop in the skin, and vitamin D degrades as fast as it is generated.
Vitamin D from the diet or that is synthesized by the body is biologically inactive; activation requires enzymatic conversion in the liver and kidney.



Metabolism and pathway map for vitamin D: Vitamin D synthesis pathway

This page titled 5.4D: Metabolic Functions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4E: Blood Supply to the Epidermis

The blood vessels in the dermis provide nourishment and remove waste from its own cells and from the stratum basale of the epidermis.

LEARNING OBJECTIVE

Identify the source of the blood supply for the integumentary system

Key Takeaways


Key Points

- The epidermis contains no blood vessels, and cells in the deepest layers are nourished by diffusion from blood capillaries present in the upper layers of the dermis.
- The papillary region of the dermis is composed of loose areolar connective tissue. This is named for its fingerlike projections called papillae, that extend toward the epidermis and contain terminal networks of blood capillaries.
- The control of blood vessels within the dermis forms a key part of the body's thermoregulatory capacity.

Key Terms

- **papillary region**: The uppermost region of the dermis that is adjacent to the epidermis.
- **reticular region**: The lower region of the dermis.

The epidermis does not contain blood vessels; instead, cells in the deepest layers are nourished by diffusion from blood capillaries that are present in the upper layers of the dermis. Diffusion provides nourishment and waste removal from the cells of the dermis, as well as for the cells of the epidermis.



The dermis: The distribution of the blood vessels in the skin of the sole of the foot. Corium—labeled at upper right—is an alternate term for dermis. Blood vessels that supply the capillaries of the papillary region are seen running through the reticular layer.

The dermis is the layer of skin beneath the epidermis that consists of connective tissue and cushions the body from stress and strain. The dermis is tightly connected to the epidermis by a basement membrane.

The dermis is structurally divided into two areas: a superficial area adjacent to the epidermis, called the papillary region, and a deep, thicker area known as the reticular region.

The papillary region is composed of loose areolar connective tissue. This is named for its fingerlike projections called papillae, that extend toward the epidermis and contain terminal networks of blood capillaries.

The reticular region lies under the papillary region and is usually much thicker. It is composed of dense, irregular connective tissue. The reticular region receives its name from the dense concentration of collagenous, elastic, and reticular fibers that weave throughout it.

These protein fibers give the dermis its typical properties of strength, extensibility, and elasticity. Blood vessels that supply the capillaries of the papillary region run through the reticular region.

Control of the blood supply to the dermis forms part of the body's thermoregulatory capacity. Increasing blood flow, which makes the skin appear redder, will increase the loss of radiant heat through the skin, whereas constricting blood flow, making the skin appear paler, reduces heat loss.

This page titled 5.4E: Blood Supply to the Epidermis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.4F: Excretion and Absorption

The integumentary system functions in absorption (oxygen and some medications) and excretion (e.g., perspiration via the eccrine glands).

LEARNING OBJECTIVE

Describe the role of glands in excretion and absorption

Key Takeaways

Key Points

- Eccrine glands, the major sweat glands of the human body, produce a clear, odorless substance, consisting primarily of water and NaCl. NaCl is reabsorbed in the duct to reduce salt loss.
- Apocrine sweat glands are found only in certain locations of the body: the axillae (armpits), areola and nipples of the breast, ear canal, perianal region, and some parts of the external genitalia.
- The sebaceous glands secrete an oily/waxy matter called sebum to lubricate and waterproof the skin and hair of mammals. In humans, they are found in greatest abundance on the face and scalp, though they are distributed throughout all skin sites except the palms and soles.

A major function of the integumentary system is absorption and excretion.


Excretion

There are numerous secretory glands present in the skin which secrete a large range of distinct fluids.

Perspiration, or sweating, is the production of fluids secreted by the sweat glands in the skin of mammals. Two types of sweat glands can be found in humans: eccrine glands and apocrine glands.

Eccrine glands are the major sweat glands of the human body, found in virtually all skin. They produce a clear, odorless substance consisting primarily of water and NaCl (note that the odor from sweat is due to bacterial activity on the secretions of the apocrine glands).

NaCl is reabsorbed in the duct to reduce salt loss. Eccrine glands are active in thermoregulation and are stimulated by the sympathetic nervous system.



Sweat gland: A sectional view of the skin (magnified), with the eccrine glands highlighted.

Apocrine sweat glands are inactive until they are stimulated by hormonal changes in puberty. Apocrine sweat glands are mainly thought to function as olfactory pheromones, chemicals important in attracting a potential mate. The stimulus for the secretion of apocrine sweat glands is adrenaline, which is a hormone carried in the blood.

The sebaceous glands are microscopic glands in the skin that secrete an oily/waxy matter, called sebum, to lubricate and waterproof the skin and hair of mammals. In humans, they are found in greatest abundance on the face and scalp, though they are distributed throughout all skin sites except the palms and soles. In the eyelids, meibomian sebaceous glands secrete a special type of sebum into tears.

Absorption

Due to the absorptive capabilities of skin, the cells comprising the outermost 0.25–0.40 mm of the skin can be supplied by external oxygen rather than via the underlying capillary network. Additionally certain medications can be administered through the skin.

The most common mechanism of administration through the skin is the use of ointments or an adhesive patch, such as the nicotine patch or iontophoresis. Iontophoresis, also called electromotive drug administration, is a technique that uses a small electric charge to deliver a medicine or other chemical through the skin.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Integumentary system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Integumentary_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anatomical barrier. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/anatomical%20barrier**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Langerhans cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Langerhans%20cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adaptive immune system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adaptive_immune_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cuticle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cuticle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Skin Diagram. **Provided by**: Daniel de Souza Telles. **Located at**: **en.Wikipedia.org/wiki/File:HumanSkinDiagram.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thermoregulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thermoregulation**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- Evaporation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Evaporation. **License**: *CC BY-SA: Attribution-ShareAlike*
- arrector pili. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/arrector_pili. **License**: *CC BY-SA: Attribution-ShareAlike*
- vasoconstriction. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasoconstriction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Integumentary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/Integumentary_System. **Project**: Human Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- homeostasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/homeostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- somatosensory system. **Authored by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/somatosensory. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cutaneous mechanoreceptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mechanoreceptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- central nervous system. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/central_nervous_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- sensory receptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sensory_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- 7-dehydrocholesterol. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/7-dehydrocholesterol. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vitamin D. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vitamin...on_in_the_skin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Integumentary system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Integumentary_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- papillae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/papillae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypodermis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypodermis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dermis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dermis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human skin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_skin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Perspiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perspiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- Eccrine sweat gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Eccrine_sweat_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sebaceous gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sebaceous_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Apocrine sweat glands. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Apocrine_sweat_glands. **License**: *CC BY-SA: Attribution-ShareAlike*
- excretion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/excretion. **License**: *CC BY-SA: Attribution-ShareAlike*
- Iontophoresis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Iontophoresis. **License**: *CC BY-SA: Attribution-ShareAlike*
- areola. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/areola. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940 - sweat gland. **Provided by**: Henry Gray. **Located at**: en.Wikipedia.org/wiki/File:Gray940_-_sweat_gland.png. **Project**: Gray's Anatomy. **License**: *Public Domain: No Known Copyright*
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 5.4F: Excretion and Absorption is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.5: WOUND HEALING

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

5.5A: Steps of Tissue Repair

5.5B: Epidermal Wound Healing

5.5C: Deep Wound Healing

This page titled 5.5: Wound Healing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.5A: Steps of Tissue Repair

Wound healing is divided into four overlapping states: 1) homeostasis, 2) inflammatory, 3) proliferative, and 4) remodeling.

Learning Objectives

- Describe the overlapping phases of tissue repair

Key Points

- Wound healing is the process whereby the skin repairs itself after injury.
- Wound healing can be divided into four overlapping processes; maintenance of homoeostasis, an inflammatory response, a proliferative phase, and remodeling.
- Maintenance of homoeostasis is achieved by clotting in any damaged regions of the circulatory system.
- The inflammatory response clears the wound site of debris and prevents infection.
- During the proliferative phase new tissue and an extra cellular matrix to support tissue repair are laid down.
- During the remodelling stage, the rapidly laid down tissue is altered to more closely mimic surrounding, mature tissues.
- The wound healing process can be interrupted, leading to the formation of chronic wounds or the development of fibrosis.

Key Terms

- **fibrin**: A fibrous protein involved in blood clotting.
- **fibronectin**: An extra cellular matrix protein, often laid down as a component of the temporary granulation tissue seen in wound healing.
- **Granulation**: The temporary connective tissue laid down during wound healing to facilitate wound closure.

Wound Healing


Wound healing is the process by which the skin, or any injured organ, repairs itself after injury. The main aim of wound healing is to prevent or limit further damage, to clean and seal the wound against infection, to restore tissue strength, and, if possible, tissue function.

Wounds in the skin can either be classed as epidermal (shallow, in which the dermis remains intact) or deep (in which the dermis is damaged; this is sometimes referred to as a full thickness wound).

Phases of the Wound Healing Process

The wound healing process can be characterized by four overlapping phases:

1. An initial response to maintain homoeostasis.
2. An inflammatory response to prevent infection.
3. A proliferative phase to reconstitute the wound site.
4. A remodelling phase where tissue strength and function are restored.



Critical developments that occur during a localized inflammatory response: The phases of wound healing during an inflammatory response to infection. Limits vary within faded intervals, mainly by wound size and healing conditions. The image does not include major impairments that cause chronic wounds.

Homoeostasis

Upon wounding, the first phase of the wound response is concerned with maintaining homoeostasis within the body. Most wounds, even superficial shallow wounds, result in damage to the circulatory system. To prevent blood loss and reduce the chance of infection spreading throughout the body, circulation platelets within the blood begin to form a fibrin clot, which seals the wound site.

Additionally, vasoconstriction initially occurs around the wound site as a means of isolating the wound site. However, this is soon followed by vasodilation so the required cells are able to be recruited to the wound site. Factors are released from damaged cells, and those around the wound site initiate the inflammatory response. This phase is very quick.

Inflammatory

Immune cells, such as neutrophils and macrophages, are attracted by factors released from the wound site and begin to accumulate, travelling through the circulatory system. These cells are responsible for the removal of debris and killing of bacteria that easily colonize the wound site, and prepare the wound for the proliferative/remodelling phase.

Proliferative

The proliferative phase can itself be divided into four phases; in the case of shallow wounds the first two steps may not occur:

- Re-vascularisation: New blood vessels are formed around the wound site in order to supply the cells and nutrients required to remodel the wound.
- Granulation: Fibroblasts attracted to the wound site quickly lay down a temporary extra cellular matrix, comprised of collagen and fibronectin, upon which the epidermis can be reconstituted.
- Re-epithelialization: The exact mechanism of re-epithelialization is poorly understood. It is thought that surviving epithelial cells around the wound edge become more motile and stretch to cover the wound site. Once a continuous epidermis is formed they lose this motility and begin to divide.
- Contraction: Re-epithelization is thought to occur simultaneously with contraction, where myo-fibroblasts recruited around the wound site pull against each other to contract the size of the wound.

Remodeling

Following closure of the wound, remodeling can occur. The epidermis proliferates and returns to its normal character; fibroblasts and immune cells which were recruited to the site are degraded; and the temporary extra cellular matrix that was laid down is remodelled into a stronger, more permanent structure.

Issues with Wound Healing

The above mechanism describes a best-case scenario that results in the restoration of a fully functional dermis and epidermis.

However, as the main aim of the wound healing process in the initial stage is to prevent further damage or infection, the intial stages can lead to a less than optimum result, as evidenced by the formation of scars. The larger and more severe the wound the more likely this is to occur.

The response to wounding is also susceptible to disruptions that can lead to the formation of chronic, non-resolving wounds such as ulcers; or the development of fibrosis if the proliferative phase does not resolve.

This page titled 5.5A: Steps of Tissue Repair is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.5B: Epidermal Wound Healing

Epidermal wound healing describes the mechanism by which the skin repairs itself after injury.

Learning Objectives

- Characterize epidermal wound healing

Key Points

- Epidermal wounds are typically less severe than those affecting the dermis.
- Clotting may not occur if there is no breaching of the vasculature; however, an immune response is still generated as the wound site is susceptible to infection.
- Proliferation is not required as the dermis remains intact and is able to independently re-constitute the basement membrane required for re-epithelialization.
- Keratinocytes surrounding the wound site, and epidermal cells found in dermal appendages (such as hair follicles), are able to re-epithelialize the wound site.
- Little remodeling will occur as the original, mature extracellular matrix remains intact.
- Epidermal wounds often resolve quickly and have fewer potential issues than deeper wounds.

Epidermal wound healing refers to the repair of the epidermis in response to wounding. Epidermal only wounds are typically less severe than those affecting the dermis and so stages of the wound healing response may be missed.

As the epidermis is itself not vascularised—it is receiving blood from the dermis—a clotting and vasoconstrictive response is often not necessary. Immune cells may still be recruited to the wound site because the removal of the epidermal barrier makes the wound susceptible to infection.

Since the dermis is intact, local fibroblasts are able to contribute to the formation of a new basement membrane, upon which the epidermis sits. In very minor wounds even the basement membrane might remain intact, allowing for rapid re-epithelialization.

Keratinocytes—epidermal epithelial cells—around the wound site migrate across the wound and close it. Additionally, epidermal cells from dermal appendages, such as hair follicles, can contribute to wound closure.

Since the dermis and underlying tissue have not been damaged very little remodelling is required. As such, small wounds only in the epidermis typically heal rapidly and are often not observable (e.g., via the formation of scar tissue) within a period of months.

This page titled 5.5B: Epidermal Wound Healing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.5C: Deep Wound Healing

A deep wound involves the inner, deeper layers of the skin (dermis).

Learning Objectives

- Describe the process of deep wound healing

Key Points


- Deep wounds are more difficult to heal, as the basement membrane and/or skin appendages (hair follicles and sweat and oil glands) may be destroyed.
- The healing of deep wounds may occur only at the edges of the wound, with scar tissue covering the center of the wound site.
- Loss of physiological function is often associated with deep wounds as they do not heal correctly.
- After healing, the tissue underlying a deep wound may be weaker than the surrounding mature tissue, making repeat injury more likely.

Key Terms

- **basement membrane**: Controls the traffic of cells and molecules between the dermis and epidermis, and provides support to epidermal cells.
- **skin appendage**: Skin-associated structures that serve a particular function, including sensation, contractility, lubrication, and heat loss.

Deep wounds that damage the dermis, or even the underlying muscle and fat, are more difficult to heal than shallow, epidermal-only wounds. The wound healing processes may be extended and scar tissue is likely to form due to improper re-epithelialization.

Additionally, deep wounds are more susceptible to infection, and also to the development of systemic infection through the circulatory system, as well as dysregulation that results in chronic wounds such as ulcers.



Wound healing phases: This image illustrates the phases of wound healing. Limits vary within faded intervals, mainly by wound size and healing conditions, but the image does not include major impairments that cause chronic wounds.

The wound healing process for deep wounds is similar to that of shallow wounds. However, with the removal of the dermis and its associated skin appendages, re-epithelialization can only occur from the wound edge, with no contribution from the dermal compartment.

Therefore, proper reconstitution of the epidermis is often only seen at the edge of the wound, with fibrous scar tissue—formed from the extracellular matrix (ECM) deposited during the proliferative phase—covering the rest of the wound site.

With the formation of a scar, the original physiological properties of the tissue are lost. For example scars are less flexible than skin, and do not feature sweat glands or hair follicles.

The ECM formed during wound healing may also be weaker in deep wounds, making the site susceptible to additional later wounding. The provisional ECM laid down during the proliferative phase is rich in fibronectin and collagen III that combine to allow quicker cell movement through the wound, which is very important during wound healing.

However, the ECM of mature skin is rich in collagen I. In large, deep wounds the remodelling of a fibronectin and collagen III-rich ECM to a collagen-I rich ECM may not occur, leading to a weakening of the tissue.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epithelialization and contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/epithel...%20contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibrin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibrin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dermis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/dermis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing phases. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Wo...ing_phases.png**. **License**:
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibrin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibrin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/wound%20healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- granulation tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/granulation%20tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing phases. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Wo...ing_phases.png**. **License**:
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Wound_healing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basement membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Basement_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- skin appendage. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/skin%20appendage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- deep wound. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/deep%20wound**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- basement membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/basement%20membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wound healing phases. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Wo...ing_phases.png**. **License**:
- Wound healing phases. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Wound_healing_phases.png**. **License**:

This page titled 5.5C: Deep Wound Healing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.6: OVERVIEW OF CANCER

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

5.6A: Overview of Cancer

5.6B: Cell Signaling and Cell Death

This page titled 5.6: Overview of Cancer is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.6A: Overview of Cancer

Cancer, known medically as a malignant neoplasm, is a broad group of various diseases, all involving unregulated cell growth.

Learning Objectives


- Explain the primary characteristics of cancer

Key Points

- In cancer, cells divide and grow uncontrollably, forming malignant tumors, and invade nearby parts of the body.
- The cancer may also spread to more distant parts of the body through the lymphatic system or bloodstream; this is called metastasis.
- Cancers are primarily associated with environmental factors such as smoking or excessive alcohol consumption.
- Some cancers have a very strong genetic risk associated with their development.
- Oncogenes are genes that promote cell growth and reproduction.
- Tumor suppressor genes inhibit cell division and survival.
- Malignant transformation can occur through the formation of novel oncogenes, the inappropriate over-expression of normal oncogenes, or by the under-expression or disabling of tumor suppressor genes.
- Cancer treatment options include surgery, chemotherapy, and radiation. Alternative treatments are being explored through clinical trials.

Key Terms

- **benign**: A non-malignant form of cancer.
- **neoplasm**: A cell or group of cells that have undergone unregulated growth and form a mass of tissue.
- **malignant**: In relation to cancer, it refers to the tendency of the cancer to become worse, e.g., by spreading throughout the body.
- **metastasis**: The transference of a bodily function or disease to another part of the body, specifically the development of a secondary area of disease remote from the original site.



Chest x-ray showing lung cancer in the left lung: A tumor present in the lung is marked via the black box.

Cancer describes a broad group of diseases, all involving unregulated cell growth caused by genetic mutations. Cancers comprise the malignant (having a tendency to become worse) subset of neoplasms —a cell or group of cells that undergo unregulated growth and form a mass of tissue —often referred to as a tumor. A major hallmark of cancers is metastasis, the ability of the cancer to spread between tissues and organs within the body.

Non-malignant tumors are referred to as benign; they are typically slower growing and are often surrounded by a membrane of connective tissue that prevents metatasis. A common example of a benign tumor is a skin mole.

While typically asymptomatic, benign tumors can impact health, usually by impairing organ function through the compression of blood vessels or nerve fibers.

Hallmarks of Cancer

Due to the wide variety of cancerous diseases, six hallmarks are used to group and define cancers:

1. Unregulated cell growth and division.
2. Continuous growth and division even given contrary signals.
3. Avoidance of programmed cell death.
4. Unlimited cell division.
5. Promoting vascularisation.
6. Invasion of tissue and formation of metastases.

Signs and Symptoms

Cancer is typically asymptomatic at onset; its symptoms appear as the tumor grows and invades other tissues. Initial symptoms are typically associated with the loss of organ function at the tumor site.

For example, patients with lung cancer often present with symptoms including breathlessness and chronic cough, but can vary widely between patients. Due to this variation and the association of symptoms with other disorders, an early diagnosis of cancer is often difficult.

As the disease progresses, systemic symptoms such as weight loss, fever, and tiredness can develop. Additionally, symptoms associated with metastasis—such as enlarged lymph nodes, an enlarged liver, or an enlarged spleen—can develop.

Classification

Cancers are classified by the type of cell that the tumor cells resemble with the presumption that this shows the origin of the tumor. These include:

- Carcinoma: Cancers derived from epithelial cells. This group includes many of the most common cancers, particularly in the aged, and include nearly all those developing in the breast, prostate, lung, pancreas, and colon.
- Sarcoma: Cancers arising from connective tissue (i.e., bone, cartilage, fat, nerve), each of which develop from cells originating in the mesenchymal cells outside the bone marrow.
- Lymphoma and leukemia: These two classes of cancer arise from hematopoietic (blood-forming) cells that leave the marrow and tend to mature in the lymph nodes and blood, respectively.
- Germ cell tumor: Cancers derived from pluripotent cells, most often presenting in the testicle or the ovary (seminoma and dysgerminoma, respectively).
- Blastoma: Cancers derived from immature precursor cells or embryonic tissue. These are also most common in children.

Causes of Cancer

Cancers are primarily associated with environmental factors such as smoking, obesity, high alcohol consumption, radiation exposure, and environmental pollutants. However, it is important to note that exposure to these factors merely imparts an increased risk of the development of cancer, rather than ensuring its development.

There is some element of genetic risk in the development of cancers, with some specific subsets having a very high genetic risk

component. For example, familial mutations of the BRCA1 and BRCA2 genes are associated with an increased risk of developing breast cancer.


Genetics of Cancer

Cancer is a disease of dysregulated cell proliferation. At the genetic level, cell proliferation can either be promoted or repressed via the expression of key genes; with alteration of these genes, expression or function via mutation, as induced by the factors discussed above, result in dysregulated proliferation.

Affected genes can be divided into two broad categories.

1. Oncogenes are genes that promote cell growth and reproduction, such as the gene MYC.
2. Tumor suppressor genes are genes that inhibit cell division and survival, such as the TP53 gene.

Malignant transformation can occur through the formation of novel oncogenes, the inappropriate over-expression of normal oncogenes, or by the under-expression or disabling of tumor suppressor genes. Typically, changes in multiple genes are required to transform a normal cell into a cancer cell.



Pathway of cancer development: Cancer develops due to mutations in several key genes that control cell proliferation and survival.

Therapies

Cancer therapy varies by patient, tumor type, and location. The most common treatments include:

- Surgery: This is the primary method of treatment for most isolated, solid cancers. It is also typically an important part of making the definitive diagnosis of tumor type and character through biopsies.
- Chemotherapy: In addition to surgery, this has proven useful in a number of different cancer types. Chemotherapy refers to the administration of a toxin that targets rapidly dividing cells and promotes their cell death. It is nonspecific to tumors and it's effect on other rapidly dividing cells, such as those of the hair follicle, give rise to the side-effects associated with chemotherapy.
- Radiotherapy: Radiation therapy involves the use of ionizing radiation to specifically target a tumor within the body, leading to its death. Radiation is typically used in addition to surgery and/or chemotherapy. For certain types of cancer, such as early head and neck cancer, it may be used alone.

This page titled 5.6A: Overview of Cancer is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

5.6B: Cell Signaling and Cell Death

When a cell is damaged, unnecessary, or dangerous to an organism, a cell can initiate the mechanism for cell death known as apoptosis.

Learning Objectives

- Describe how apoptosis is initiated

Key Points

- Apoptosis allows a cell to die in a controlled manner by preventing the release of damaging molecules from inside the cell.
- Internal checkpoints to monitor a cell's health exist; if abnormalities are observed, a cell can also spontaneously initiate the process of apoptosis.
- In some cases, such as a viral infection or cancer, the cell's normal checks and balances fail.
- External signaling can also initiate apoptosis.
- Apoptosis is also essential for normal embryological development; unnecessary cells that appear during the early stages of development will eventually be eliminated through cell signaling.

Key Terms

- **apoptosis**: a process of programmed cell death
- **glycoprotein**: a protein with covalently-bonded carbohydrates

Apoptosis

When a cell is damaged, superfluous, or potentially dangerous to an organism, a cell can initiate a mechanism to trigger programmed cell death, or apoptosis. Apoptosis allows a cell to die in a controlled manner that prevents the release of potentially damaging molecules from inside the cell.

Internal Signaling

There are many internal checkpoints that monitor a cell's health; if abnormalities are observed, a cell can spontaneously initiate the process of apoptosis. However, in some cases such as a viral infection or uncontrolled cell division due to cancer, the cell's normal checks and balances fail.

External Signaling



Apoptosis: The histological section of a foot of a 15-day-old mouse embryo, visualized using light microscopy, reveals areas of tissue between the toes which apoptosis will eliminate before the mouse reaches its full gestational age at 27 days.

External signaling can also initiate apoptosis. For example, most normal animal cells have receptors that interact with the extracellular matrix, a network of glycoproteins that provides structural support for cells in an organism. The binding of cellular receptors to the extracellular matrix initiates a signaling cascade within the cell. However, if the cell moves away from the extracellular matrix, the signaling ceases, and the cell undergoes apoptosis. This system keeps cells from traveling through the body and proliferating out of control, as happens with tumor cells that metastasize.

Another example of external signaling that leads to apoptosis occurs in T-cell development. T-cells are immune cells that bind to foreign macromolecules and particles, targeting them for destruction by the immune system. Normally, T-cells do not target "self" proteins (those of their own organism), a process that can lead to autoimmune diseases. In order to develop the ability to discriminate between self and non-self, immature T-cells undergo screening to determine whether they bind to so-called self proteins. If the T-cell receptor binds to self proteins, the cell initiates apoptosis to remove the potentially dangerous cell.

Apoptosis and Embryos

Apoptosis is also essential for normal embryological development. In vertebrates, for example, early stages of development include the formation of web-like tissue between individual fingers and toes. During the course of normal development, these unnecessary cells must be eliminated, enabling fully separated fingers and toes to form. A cell signaling mechanism triggers apoptosis, which destroys the cells between the developing digits.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Alcohol and cancer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Alcohol_and_cancer. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cancer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cancer. **License**: *CC BY-SA: Attribution-ShareAlike*
- metastasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/metastasis. **License**: *CC BY-SA: Attribution-ShareAlike*

- Thorax_pa_peripheres_Bronchialcarcinom_li_OF_markiert.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3AThorax_pa_peripheres_Bronchialcarcinom_li_OF_markiert.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cancer_requires_multiple_mutations_from_NIHen.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ACancer_requires_multiple_mutations_from_NIHen.png**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44453/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- glycoprotein. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glycoprotein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- apoptosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/apoptosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thorax_pa_peripheres_Bronchialcarcinom_li_OF_markiert.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3AThorax_pa_peripheres_Bronchialcarcinom_li_OF_markiert.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cancer_requires_multiple_mutations_from_NIHen.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ACancer_requires_multiple_mutations_from_NIHen.png**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Response to the Signal. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44453/latest...e_09_03_02.jpg**. **License**: ***CC BY: Attribution***

This page titled 5.6B: Cell Signaling and Cell Death is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6: SKELETAL SYSTEM

6.1: Overview of the Skeletal System

6.1A: Overview of the Musculoskeletal System

6.1B: The Axial Skeleton

6.1C: The Appendicular Skeleton

6.2: Cartilage

6.2A: Structure, Type, and Location of Cartilage

6.2B: Cartilage Growth

6.3: Introduction to Bone

6.3A: Gross Anatomy

6.3B: Supply of Blood and Nerves to Bone

6.3C: Microscopic Anatomy of Bone

6.3D: Chemical Composition of Bone

6.4: Bone Formation

6.4A: Embryonic and Fetal Bone Formation

6.4B: Postnatal Bone Growth

6.4C: Bone Remodeling

6.4D: Bone Repair

6.5: Bone and Calcium

6.5A: Bone and Calcium Homeostasis

This page titled 6: Skeletal System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.1: OVERVIEW OF THE SKELETAL SYSTEM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

6.1A: Overview of the Musculoskeletal System

6.1B: The Axial Skeleton

6.1C: The Appendicular Skeleton

This page titled 6.1: Overview of the Skeletal System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.1A: Overview of the Musculoskeletal System

The musculoskeletal system is an organ system that enables an organism to move, support itself, and maintain stability during locomotion.

Learning Objectives

- Explain the purpose of the musculoskeletal system

Key Points

- The musculoskeletal system 's primary functions include supporting the body, allowing motion, and protecting vital organs.
- The musculoskeletal system is made up of the body's bones (the skeleton), muscles, cartilage, tendons, ligaments, joints, and other connective tissue that support and bind tissues and organs together.
- The skeleton serves as the main storage system for calcium and phosphorus.
- The skeleton also contains critical components of the hematopoietic (blood production) system and fat storage. These functions occur in red marrow and yellow marrow, respectively.
- To allow motion, different bones are connected by articulating joints. Cartilage prevents the bone ends from rubbing directly on to each other while the muscles contract to move the bones associated with the joint.

Key Terms

- **red marrow**: Red marrow or medulla ossium rubra, consists mainly of hematopoietic tissue, and gives rise to red blood cells (RBCs), platelets and most white blood cells (WBCs).
- **musculoskeletal system**: An organ system that gives animals (and humans) the ability to move, using the combined actions of the muscular and skeletal systems. It provides form, support, stability, and movement to the body.
- **hematopoeisis**: A biological process in which new blood cells are formed from hematopoietic stem cells (HSCs) within the marrow. All cellular blood components are derived from HSCs.


The musculoskeletal system (also known as the locomotor system) is an organ system that gives animals (including humans) the ability to move, using the muscular and skeletal systems. It provides form, support, stability, and movement to the body.

The musculoskeletal system is made up of the body's bones (the skeleton), muscles, cartilage, tendons, ligaments, joints, and other connective tissue that supports and binds tissues and organs together.

Its primary functions include supporting the body, allowing motion, and protecting vital organs.
The bones of the skeletal system provide stability to the body analogous to a reinforcement bar in concrete construction.

Muscles keep bones in place and also play a role in their movement. To allow motion, different bones are connected by articulating joints, and cartilage prevents the bone ends from rubbing directly onto each other.

Skeletal System



A human skeleton: Image as overview of the human skeletal system.

The skeletal portion of the system serves as the main storage system for calcium and phosphorus. The importance of this storage is to help regulate mineral balance in the bloodstream. When the fluctuation of minerals is high, these minerals are stored in bone; when it is low, minerals are withdrawn from the bone.

The skeleton also contains critical components of the hematopoietic (blood production) system. Located in long bones are two distinctions of bone marrow: yellow and red. The yellow marrow has fatty connective tissue and is found in the marrow cavity. In times of starvation, the body uses the fat in yellow marrow for energy.

The red marrow of some bones is an important site for hematopoeisis or blood cell production that replaces cells that have been destroyed by the liver. Here, all erythrocytes, platelets, and most leukocytes form in bone marrow from where they migrate to the circulation.

Muscular System

Muscles contract (shorten) to move the bone attached at the joint. Skeletal muscles are attached to bones and arranged in opposing groups around joints. Muscles are innervated—the nerves conduct electrical currents from the central nervous system that cause the muscles to contract.

Three types of muscle tissue exist in the body. These are skeletal, smooth, and cardiac muscle.


- Only skeletal and smooth muscles are considered part of the musculoskeletal system.
- Skeletal muscle is involved in body locomotion.
- Examples of smooth muscles include those found in intestinal and vessel walls.
- Cardiac and smooth muscle are characterized by involuntary movement (not under conscious control).
- Cardiac muscles are found in the heart.

Tendons, Joints, Ligaments, and Bursae

A tendon is a tough, flexible band made of fibrous connective tissue, and functions to connect muscle to bone. Joints are the bone articulations allowing movement. A ligament is a dense, white band of fibrous elastic tissue.

Ligaments connect the ends of bones together in order to form a joint. These help to limit joint dislocation and restrict improper hyperextension and hyperflexion. Also made of fibrous tissue are bursae. These provide cushions between bones and tendons

and/or muscles around a joint.



Musculoskeletal system: Image depicting the human muscular system (skeletal muscle)

This page titled 6.1A: Overview of the Musculoskeletal System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.1B: The Axial Skeleton

The axial skeleton functions to support and protect the organs of the dorsal and ventral cavities and serves as a surface for the attachment of muscles and parts of the appendicular skeleton.

Learning Objectives

- List the components of the axial skeleton


Key Points

- The axial skeleton is the part of the skeleton that consists of the bones of the head and trunk of a vertebrate animal, including humans.
- The primary divisions of the skeleton system are the head, thorax, and vertebral column.
- The human cranium supports the structures of the face and forms the brain cavity.
- The rib cage functions as protection for the vital organs of the chest such as the heart and lungs.
- The cervical vertebrae make up the junction between the vertebral column and the cranium, and the bone makes up the junction between the vertebral column and the pelvic bones.

Key Terms

- **flat bones**: Thin bones (although often curved) that serve as points of attachment for muscles and protect internal organs (examples, cranium, sternum).
- **cranial vault**: The space in the skull occupied by the brain.
- **sutures**: Fibrous joints that are only found in the cranium.

The axial skeleton is the part of the skeleton that consists of the bones of the head and trunk of a vertebrate animal, including humans.



Axial skeleton: Image depicting the human skeleton with the axial skeleton.

The word axial is from the word axis, and refers to how the bones of the axial skeleton are located along the central axis of the body.

The axial skeleton functions to support and protect the organs of the dorsal and ventral cavities. It also serves as a surface for the attachment of muscles and parts of the appendicular skeleton.

The human's axial skeleton is composed of 80 bones and is the central core of the body. The primary divisions of the skeleton system are:

- Head, including the bones of the skull (cranium), face, auditory ossicles, and hyoid bone.
- Thorax, including the rib cage and sternum.
- Vertebral column.

Bones of the Head

Skull (Cranium)

The human cranium consists of the flat bones of the cranium and includes the facial bones. The cranium protects the brain that is contained in the cranial vault. The cranium is formed from eight bones connected by sutures.

Fourteen facial bones form the lower front part of the cranium. Important facial bones include the lower jaw or mandible, the upper jaw or maxilla, the zygomatic or cheek bone, and the nasal bone.

The immature cranium has separate plates to allow the flexibility needed for a newborn to pass through the birth canal and pelvis.

These plates fuse as the skull matures (except the mandible). The human cranium supports the structures of the face and forms the brain cavity.

Ossicle

The ossicles (also called auditory ossicles) consist of three bones (malleus, incus, and stapes) that are the smallest in the body. These are located in the middle ear and serve to transmit sounds from the air to the fluid-filled labyrinth.

Hyoid Bone

The hyoid bone is a horseshoe-shaped bone situated in the anterior midline of the neck between the chin and the thyroid cartilage. It provides attachment to the muscles of the floor of the mouth, the tongue above, larynx below, and the epiglottis and pharynx behind.

Rib Cage

The rib cage is composed of 25 bones that include the 12 pairs of ribs plus the sternum. It functions as protection for the vital organs of the chest, such as the heart and lungs. The rounded ends are attached at joints to the thoracic vertebrae posteriorly and the flattened ends come together at the sternum anteriorly.

The first seven pairs of ribs attach to the sternum with costal cartilage and are known as true ribs. Thelength of each rib pair increases from number one to seven. After rib seven, the size begins to decrease. The 8th through 10th ribs have noncostal cartilage that connects them to the ribs above.

The last two ribs are called floating ribs because they do not attach to the sternum or to other ribs.

Vertebral Column

There are normally thirty-three vertebrae in the human vertebral column. The upper twenty-four articulate and are unfused, the lower nine are fused. The fused vertebrae are the five in the sacrum and four in the coccyx.

The articulating vertebrae are named according regions:

- Cervical vertebrae (seven vertebrae).
- Thoracic (twelve vertebrae).
- Lumbar (five vertebrae).

The first and second cervical vertebrae are the atlas and axis, respectively, on which the head rests. The cervical vertebrae make up the junction between the vertebral column and the cranium, and the bone makes up the junction between the vertebral column and the pelvic bones.

This page titled 6.1B: The Axial Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.1C: The Appendicular Skeleton

The appendicular skeleton includes the skeletal elements within the limbs, as well as supporting pectoral and pelvic girdles.

Learning Objectives

- List the components of the appendicular skeleton

Key Points

- The appendicular skeleton comprises 126 bones and is involved in locomotion and manipulation of objects in the environment.
- The bones of the appendicular skeleton are divided into two groups: the bones that are located within the limbs themselves, and the girdle bones that attach the limbs to the axial skeleton.
- The bones of the pectoral girdle anchor the upper limb to the thoracic cage of the axial skeleton.
- The pelvic girdle is formed by a single bone and serves as the attachment point for each lower limb.

Key Terms


- **Girdle**: A group of bones that connect the appendages to the axial skeleton.
- **phalanges**: The digital bones of the hands and feet (singular, phalanx).
- **appendages**: The parts of the body that extend from the axial trunk.

The appendicular skeleton of vertebrates, including humans, consists of the bones that support and compose the appendages (for example, the arms and legs of humans). The word appendicular is the adjective of the noun appendage.

The appendicular skeleton includes the skeletal elements within the limbs, as well as supporting the pectoral and pelvic girdles.

The appendicular skeleton comprises 126 bones and is involved in locomotion and manipulation of objects in the environment. It is unfused, allowing for greater range of motion.

Divisions of the Appendicular Skeleton



A diagram of the apendicular skeleton: Image depicting the human skeleton with the appendicular skeleton colored red.

The appendicular skeleton is divided into six major regions:

1. The pectoral girdles consist of 4 bones: The left and right clavicle (2) and the scapula (2).
2. The upper arms and forearms are made up of 6 bones: The left and right humerus (upper arm, 2), the ulna (2), and the radius (forearm, 2).
3. The hands have 54 bones: The left and right carpals (wrist, 16), metacarpals (10), proximal phalanges (10), intermediate phalanges (8), and the distal phalanges (10).
4. The pelvis has 2 bones: The left and right hip bone (2).
5. The thighs and legs have 8 bones: The left and right femur (thigh, 2), patella (knee, 2), tibia (2) and fibula (leg, 2).
6. The feet and ankles have 52 bones: The left and right tarsals (ankle, 14), metatarsals (10), proximal phalanges (10), intermediate phalanges (8), and distal phalanges (10).

Pectoral Girdle

The bones of the pectoral girdle consist of two bones (scapula and clavicle) and anchor the upper limb to the thoracic cage of the axial skeleton.

The three regions of the upper limb are: arm (humerus), forearm (ulna medially and radius laterally), and the hand.

The base of the hand contains eight bones (carpal bones), and the palm is formed by five bones (metacarpal bones). The fingers and thumb contain a total of 14 bones, called phalanges.

Pelvic Girdle

The pelvic girdle is formed by a single bone, the hip or coxal bone, and serves as the attachment point for each lower limb. Each hip bone is joined to the axial skeleton by its attachment to the sacrum of the vertebral column. The right and left hip bones attach to each other anteriorly.

The lower limb contains 30 bones and is divided into three regions, the thigh, leg, and foot. These consist of the femur, patella, tibia, fibula, tarsal bones, metatarsal bones, and phalanges.

- The femur is the single bone of the thigh.
- The patella (kneecap) articulates with the distal femur.
- The tibia is located on the medial side of the leg,
- The fibula is the thin bone of the lateral leg.

The bones of the foot are divided into three groups, the tarsal bones, metatarsal bones, and phalanges of the foot.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human musculoskeletal system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_m...keletal_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- musculoskeletal system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/musculoskeletal%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- red marrow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/red%20marrow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hematopoeisis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hematopoeisis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscular System. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/**

File:Muscular_System.jpg. **License**: ***Public Domain: No Known Copyright***
- Grey Skeleton. **Provided by**: PublicDomainPictures.net. **Located at**: **http://www.publicdomainpictures.net/...=grey-skeleton**. **License**: ***Public Domain: No Known Copyright***
- Anatomical Terms of Bones. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anatomical_terms_of_bone#Flat_bones**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vertebral Column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ossicles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ossicles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Suture (Joint). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Suture_(joint)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cranial Vault. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cranial_vault**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Axial Skeleton. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Axial_skeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscular System. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Muscular_System.jpg**. **License**: ***Public Domain: No Known Copyright***
- Grey Skeleton. **Provided by**: PublicDomainPictures.net. **Located at**: **http://www.publicdomainpictures.net/...=grey-skeleton**. **License**: ***Public Domain: No Known Copyright***
- Axial_skeleton_diagram.svg. **Provided by**: Wikipedian. **Located at**: **en.Wikipedia.org/wiki/File:Axial_skeleton_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- The Appendicular Skeleton. **Provided by**: Open Stax College. **Located at**: **http://openstaxcollege.org**. **License**: ***CC BY: Attribution***
- Appendicular Skeleton. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Appendicular_skeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscular System. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Muscular_System.jpg**. **License**: ***Public Domain: No Known Copyright***
- Grey Skeleton. **Provided by**: PublicDomainPictures.net. **Located at**: **http://www.publicdomainpictures.net/...=grey-skeleton**. **License**: ***Public Domain: No Known Copyright***
- Axial_skeleton_diagram.svg. **Provided by**: Wikipedian. **Located at**: **en.Wikipedia.org/wiki/File:Axial_skeleton_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- appendicular_skeleton_diagram.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:A...on_diagram.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 6.1C: The Appendicular Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.2: CARTILAGE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

6.2A: Structure, Type, and Location of Cartilage

6.2B: Cartilage Growth

This page titled 6.2: Cartilage is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.2A: Structure, Type, and Location of Cartilage

Cartilage is an avascular, flexible connective tissue located throughout the body that provides support and cushioning for adjacent tissues.

LEARNING OBJECTIVE

Differentiate among the types of cartilage

Key Takeaways

Key Points

- Cartilage is a flexible connective tissue that differs from bone in several ways; it is avascular and its microarchitecture is less organized than bone.
- Cartilage is not innervated and therefore relies on diffusion to obtain nutrients. This causes it to heal very slowly.
- The main cell types in cartilage are chondrocytes, the ground substance is chondroitin sulfate, and the fibrous sheath is called perichondrium.
- There are three types of cartilage: hyaline, fibrous, and elastic cartilage.
- Hyaline cartilage is the most widespread type and resembles glass. In the embryo, bone begins as hyaline cartilage and later ossifies.
- Fibrous cartilage has many collagen fibers and is found in the intervertebral discs and pubic symphysis.
- Elastic cartilage is springy, yellow, and elastic and is found in the internal support of the external ear and in the epiglottis.

Key Terms

chondroitin sulfate: An important structural component of cartilage that provides much of its resistance to compression.

connective tissue: A type of tissue found in animals whose main function is to bind other tissue systems (such as muscle to skin) or organs. It consists of the following three elements: cells, fibers, and a ground substance (or extracellular matrix).

hyaline cartilage: A type of cartilage found on many joint surfaces; it contains no nerves or blood vessels, and its structure is relatively simple.

temporal mandibular joint: A joint of the jaw that connects it to the temporal bones of the skull.

Chondrocytes: Cells that form and maintain the cartilage.

What is Cartilage?


Cartilage is a flexible connective tissue that differs from bone in several ways. For one, the primary cell types are chondrocytes as opposed to osteocytes. Chondrocytes are first chondroblast cells that produce the collagen extracellular matrix (ECM) and then get caught in the matrix. They lie in spaces called lacunae with up to eight chondrocytes located in each.

Chondrocytes rely on diffusion to obtain nutrients as, unlike bone, cartilage is avascular, meaning there are no vessels to carry blood to cartilage tissue. This lack of blood supply causes cartilage to heal very slowly compared with bone.

The base substance of cartilage is chondroitin sulfate, and the microarchitecture is substantially less organized than in bone. The cartilage fibrous sheath is called the perichondrium. The division of cells within cartilage occurs very slowly, and thus growth in cartilage is usually not based on an increase in size or mass of the cartilage itself.

Articular cartilage function is dependent on the molecular composition of its ECM, which consists mainly of proteoglycans and

collagens. The remodeling of cartilage is predominantly affected by changes and rearrangements of the collagen matrix, which responds to tensile and compressive forces experienced by the cartilage.



Cartilage types: Images of microscopic views of the different types of cartilage: elastic, hyaline, and fibrous. Elastic cartilage has the most ECM; hyaline a middle amount; and fibrous cartilage has the least amount of ECM.

Types of Cartilage

There are three major types of cartilage: hyaline cartilage, fibrocartilage, and elastic cartilage.

Hyaline Cartilage

Hyaline cartilage is the most widespread cartilage type and, in adults, it forms the articular surfaces of long bones, the rib tips, the rings of the trachea, and parts of the skull. This type of cartilage is predominately collagen (yet with few collagen fibers), and its name refers to its glassy appearance.

In the embryo, bones form first as hyaline cartilage before ossifying as development progresses. Hyaline cartilage is covered externally by a fibrous membrane, called the perichondrium, except at the articular ends of bones; it also occurs under the skin (for instance, ears and nose).

Hyaline cartilage is found on many joint surfaces. It contains no nerves or blood vessels, and its structure is relatively simple.

If a thin slice of cartilage is examined under the microscope, it will be found to consist of cells of a rounded or bluntly angular form, lying in groups of two or more in a granular or almost homogeneous matrix. These cells have generally straight outlines where they are in contact with each other, with the rest of their circumference rounded.

They consist of translucent protoplasm in which fine interlacing filaments and minute granules are sometimes present. Embedded in this are one or two round nuclei with the usual intranuclear network.

Fibrocartilage

Fibrous cartilage has lots of collagen fibers (Type I and Type II), and it tends to grade into dense tendon and ligament tissue. White fibrocartilage consists of a mixture of white fibrous tissue and cartilaginous tissue in various proportions.

It owes its flexibility and toughness to the fibrous tissue, and its elasticity to the cartilaginous tissue. It is the only type of cartilage that contains type I collagen in addition to the normal type II.

Fibrocartilage is found in the pubic symphysis, the annulus fibrosus of intervertebral discs, menisci, and the temporal mandibular

joint.

Elastic Cartilage

Elastic or yellow cartilage contains elastic fiber networks and collagen fibers. The principal protein is elastin.

Elastic cartilage is histologically similar to hyaline cartilage but contains many yellow elastic fibers lying in a solid matrix. These fibers form bundles that appear dark under a microscope. They give elastic cartilage great flexibility so it can withstand repeated bending.

Chondrocytes lie between the fibers. Elastic cartilage is found in the epiglottis (part of the larynx) and the pinnae (the external ear flaps of many mammals, including humans).

This page titled 6.2A: Structure, Type, and Location of Cartilage is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.2B: Cartilage Growth

Chondrification is the process by which cartilage is formed from condensed mesenchyme tissue.

LEARNING OBJECTIVE

Describe cartilage growth and repair

Key Takeaways

Key Points


- The division of cells within cartilage occurs very slowly.
- Articular cartilage function is dependent on the molecular composition of its extracellular matrix (ECM).
- The remodeling of cartilage is predominantly affected by changes and rearrangements of the collagen matrix in response to forces experienced by the cartilage.
- Cartilage growth mainly refers to matrix deposition, but can include both growth and remodeling of the extracellular matrix.

Key Terms

- **collagen matrix**: The most abundant protein in the human body and accounts for 90% of bone matrix protein content.
- **mesenchyme**: Undifferentiated cells of the early embryo are able to develop into the different tisue types, including bone and cartilage.
- **chondrocyte**: A cell that makes up the tissue of cartilage.
- **chondrification**: The process by which cartilage is formed from condensed mesenchyme tissue.

Formation

Chondrification (also known as chondrogenesis) is the process by which cartilage is formed from condensed mesenchyme tissue.



A chondrocyte: A chondrocyte, stained for calcium, showing its nucleus (N) and mitochondria (M).

Mesenchyme tissue differentiates into chondroblasts and begins secreting the molecules that form the extracellular matrix (ECM). Mesenchymal stem cells (MSCs) are undifferentiated, meaning they can give rise to different cell types. Under the appropriate conditions and at sites of cartilage formation, they are referred to as chondrogenic cells.

During cartilage formation, undifferentiated MSCs are highly proliferative and form dense aggregates of chondrogenic cells at the center of chondrification. These condrogenic cells then differentiate to chondroblasts, which will then synthesize the cartilage ECM.



Cartilage: Hyaline cartilage showing chondrocytes and organelles, lacunae and matrix.

The extracellular matrix consists of ground substance (proteoglycans and glycosaminoglycans) and associated fibers, such as collagen. The chondroblasts then trap themselves in lacunae, small spaces that are no longer in contact with the newly created matrix and contain extracellular fluid. The chondroblast is now a chondrocyte, which is usually inactive but can still secrete and degrade the matrix depending on the conditions.

Growth

The majority of body cartilage is synthesized from chondroblasts that are largely inactive at later developmental stages compared to earlier years (pre-pubescence). The division of cells within cartilage occurs very slowly.

Therefore, growth in cartilage is usually not based on an increase in size or mass of the cartilage itself. Remodeling of cartilage is predominantly affected by changes and rearrangements of the collagen matrix, which responds to tensile and compressive forces experienced by the cartilage. Cartilage growth thus mainly refers to matrix deposition, but can include both growth and remodeling of the ECM.

Early in fetal development, the greater part of the skeleton is cartilaginous. This temporary cartilage is gradually replaced by bone (endochondral ossification), a process that ends at puberty. In contrast, the cartilage in the joints remains permanently unossified during life.

Repair

Once damaged, cartilage has limited repair capabilities because chondrocytes are bound in lacunae and cannot migrate to damaged areas. Also, because cartilage does not have a blood supply, the deposition of new matrix is slow.

Damaged hyaline cartilage is usually replaced by fibrocartilage scar tissue. Over the last few years, surgeons and scientists have elaborated a series of cartilage repair procedures that help to postpone the need for joint replacement.

These include marrow stimulation techniques, including surgeries, stem cell injections, and grafting of cartilage into damaged areas.

However, due to the extremely slow growth of cartilage and its avascular properties, regeneration and growth of cartilage post-injury is still very slow.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Fibrocartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Fibrocartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hyaline cartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hyaline_cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- chondroitin sulfate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chondroitin_sulfate. **License**: *CC BY-SA: Attribution-ShareAlike*
- Elastic cartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Elastic_cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- connective tissue. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/connective_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connective Tissue. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/General_Biology/Tissues_and_Systems/Connective_Tissue. **Project**: General Biology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cartilage Types. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Cartilage_types.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cartilage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cartilage. **License**: *CC BY: Attribution*
- collagen matrix. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/collagen%20matrix. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chondrogenesis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chondrogenesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chondrocytes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chondrocytes. **License**: *CC BY-SA: Attribution-ShareAlike*
- chondrocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chondrocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- mesenchyme. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mesenchyme. **License**: *CC BY-SA: Attribution-ShareAlike*
- chondrification. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/chondrification. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chondrocytes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chondrocytes. **License**: *CC BY-SA: Attribution-ShareAlike*
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

PUBLIC DOMAIN CONTENT

- Hyaline Cartilage. **Authored by**: Robert M. Hunt. **Located at**: commons.wikimedia.org/wiki/File:Hypertrophic_Zone_of_Epiphyseal_Plate.jpg. **License**: *Public Domain: No Known Copyright*

This page titled 6.2B: Cartilage Growth is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.3: INTRODUCTION TO BONE

6.3A: Gross Anatomy

6.3B: Supply of Blood and Nerves to Bone

6.3C: Microscopic Anatomy of Bone

6.3D: Chemical Composition of Bone

This page titled 6.3: Introduction to Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.3A: Gross Anatomy

Learning Objectives

- Differentiate long bones from flat bones

Bone Tissue

Bones support and protect the body and its organs. They also produce various blood cells, store minerals, and provide support for mobility in conjunction with muscle. Bone is made of bone tissue, a type of dense connective tissue. Bone (osseous) tissue is the structural and supportive connective tissue of the body that forms the rigid part of the bones that make up the skeleton. Overall, the bones of the body are an organ made up of bone tissue, bone marrow, blood vessels, epithelium, and nerves.

There are two types of bone tissue: **cortical** and **cancellous bone**. Cortical bone is compact bone, while cancellous bone is trabecular and spongy bone. Cortical bone forms the extremely hard exterior while cancellous bone fills the interior. The tissues are biologically identical but differ in the arrangement of their microstructure.

Bone Cells

The following are the different types of bone cells:

- Osteoblasts-involved in the creation and mineralisation of bone
- Osteocytes and osteoclasts: These are involved in the reabsorption of bone tissue. The mineralized matrix of bone tissue has an organic component—mainly made of collagen—and an inorganic component of bone mineral made up of various salts.

Bone Types

There are different types of bone. These are:

- Long bones
- Short bones
- Flat bones
- Sesamoid bones
- Irregular bones




Figure 6.3A.1: Bone types: This image show the different bone classifications, based on shape, that are found in a human skeleton. These are flat bone, sutural bone, short bone, irregular, sesamoid bone, and long bone.

Long Bones

Long bones grow primarily by elongation of the diaphysis (the central shaft), with an epiphysis at each end of the growing bone. The ends of epiphyses are covered with hyaline cartilage (articular cartilage). At the cessation of growth, the epiphyses fuse to the diaphysis, thus obliterating the intermediate area known as the epiphyseal plate or growth plate. The long bones in the body are as follows:

- Legs: The femur, tibia, and fibula.
- Arms: The humerus, radius, and ulna.
- The clavicles or collar bones.
- Metacarpals, metarsals, phalanges.

The outside of the bone consists of a layer of connective tissue called the periosteum. The outer shell of the long bone is compact bone, below which lies a deeper layer of cancellous bone (spongy bone), as shown in the following figure. The interior part of the long bone is called the medullary cavity; the inner core of the bone cavity is composed of marrow.




Figure 6.3A.2: Long bone: A long bone is longer than it is wide. Growth occurs by a lengthening of the diaphysis. located in the center of the long bone.

Short Bones

Short bones are about as wide as they are long. These provide support with less movement. Examples of short bones include the carpal and tarsal bones of the wrist and feet. They consist of a thin layer of cortical bone with cancellous interiorly.




Figure 6.3A.3: Compact bone and spongy bone: The hard outer layer of bones is composed of compact bone tissue, so-called due to its minimal gaps and spaces. Its porosity is 5–30%. Inside the interior of the bone is the trabecular bone tissue, an open cell, porous network that is also called cancellous or spongy bone.

Flat Bones

Flat bones are broad bones that provide protection or muscle attachment. They are composed of two thin layers of compact bone surrounding a layer of cancellous (spongy) bone.

These bones are expanded into broad, flat plates, as in the cranium (skull), ilium (pelvis), sternum, rib cage, sacrum, and scapula. The flat bones are named:

- Occipital
- Parietal
- Frontal
- Nasal
- Lacrimal
- Vomer
- Scapula
- Os coxæ (hip bone)
- Sternum
- Ribs

Sesamoid Bone

Sesamoid bones are smaller bones that are fixed in tendons to protect them. An example is the patella (knee cap) located in the patellar tendon. Other examples include the small bones of the metatarsals and the pisiform bones of the carpus.

Irregular Bone

The irregular bones are named for their nonuniform shape. Examples include the bones of the vertebrae. These typically have a thin cortical layer with more cancellous bone in their tissue.

Key Points

- Long bones are those that are longer than they are wide.
- The end of the long bone is the epiphysis and the shaft is the diaphysis. When a human finishes growing these parts fuse together.
- The outside of the flat bone consists of a layer of connective tissue called the periosteum.
- The interior part of the long bone is the medullary cavity with the inner core of the bone cavity being composed of marrow.
- Flat bones have broad surfaces for protection or muscular attachment.
- Flat bones are composed of two thin layers of compact bone that surround a layer of cancellous (spongy) bone. In an adult, most red blood cells are formed in the marrow in flat bones.

Key Terms

- **endosteum**: A thin vascular membrane of connective tissue that lines the surface of the bone tissue that forms the medullary cavity of long bones.
- **medullary cavity**: The medullary cavity, also known as the marrow cavity, is the central cavity of bone shafts where red bone marrow and/or yellow bone marrow (adipose tissue) is stored.
- **diaphysis**: The central shaft of any long bone.
- **epiphyseal plate**: A hyaline cartilage plate in the metaphysis, located at each end of a long bone where growth occurs in children and adolescents.


This page titled 6.3A: Gross Anatomy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.3B: Supply of Blood and Nerves to Bone

Learning Objectives


- Describe the blood and nerve supply of bones

Blood is supplied to mature compact bone through the Haversian canal. Haversian canals are formed when individual lamellae form concentric rings around larger longitudinal canals (approx. 50 µm in diameter) within the bone tissue. Haversian canals typically run parallel to the surface and along the long axis of the bone. The canals and the surrounding lamellae (8–15) are called a Haversian system or an osteon. A Haversian canal generally contains one or two capillaries and nerve fibers. The Haversian canals also surround nerve cells throughout the bone and communicate with osteocytes in lacunae (spaces within the dense bone matrix that contain the living bone cells) through canaliculi. This unique arrangement is conducive to the storage of mineral salt deposits that give bone tissue its strength.



Haversian canal: The Haversian canals surround blood vessels and nerve cells throughout the bone.

The vascular supply of long bones depends on several points of inflow, which feed complex sinusoidal networks within the bone. These in turn drain to various channels through all surfaces of the bone except that covered by articular cartilage.



Epiphyseal plate: Image shows the location of the epiphyseal plates (or lines) and the articular surfaces of long bones.

Volkmann's canals are channels that assist with blood and nerve supply from the periosteum to the Haversian canal. One or two main diaphyseal nutrient arteries enter the shaft obliquely through one or two nutrient foramina leading to nutrient canals. Their sites of entry and angulation are almost constant and characteristically directed away from the growing epiphysis.

Except for a few with double or no foramina, 90% of long bones have a single nutrient foramen in the middle third of the shaft. The nutrient arteries divide into ascending and descending branches in the medullary cavity. These approach the epiphysis dividing into smaller rami. Near the epiphysis, they anastomose with the metaphyseal and epiphyseal arteries.

The blood supply of the immature bones is similar, but the epiphysis is a discrete vascular zone separated from the metaphysis by the growth plate. Epiphyseal and metaphyseal arteries enter on both sides of the growth cartilage, with anastamoses between them being few or absent.

Growth cartilage receives its blood supply from both sources and also from an anastamotic collar in the adjoining perichondrium. Young periosteum is more vascular, has more metaphyseal branches, and its vessels communicate more freely with those of the shaft than adult periosteum.

Key Points

- Haversian canals typically run parallel to the surface and along the long axis of the bone and generally contain one or two capillaries and nerve fibers.
- Volkmann's canals are channels that assist with blood and nerve supply from the periosteum to the Haversian canal.
- The vascular supply of long bones depends on several points of inflow.
- Except for a few with double or no foramina (places in bone where capillaries enervate), 90% of long bones have a single nutrient foramen in the middle third of the shaft.
- Young periosteum is more vascular and its vessels communicate more freely with those of the shaft compared to adult periosteum.

Key Terms

- **perichondrium**: A layer of dense irregular connective tissue that surrounds the cartilage of developing bone.
- **Volkmann's canal**: Also known as perforating holes, these are microscopic structures found in the compact bone that carry small arteries throughout the bone.
- **anastomose**: Joined or run together.
- **Haversian canal**: A hollow channel in the center of an osteon, running parallel to the length of a bone.

This page titled 6.3B: Supply of Blood and Nerves to Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.3C: Microscopic Anatomy of Bone


Learning Objectives

- Classify woven bone and lamellar bone

Bones are composed of bone matrix, which has both organic and inorganic components. Bone matrix is laid down by osteoblasts as collagen, also known as osteoid. Osteoid is hardened with inorganic salts, such as calcium and phosphate, and by the chemicals released from the osteoblasts through a process known as mineralization.


The basic microscopic unit of bone is an osteon (or Haversian system). Osteons are roughly cylindrical structures that can measure several millimeters long and around 0.2 mm in diameter.

Each osteon consists of a lamellae of compact bone tissue that surround a central canal (Haversian canal). The Haversian canal contains the bone's blood supplies. The boundary of an osteon is called the cement line. Osteons can be arranged into woven bone or lamellar bone.



Osteon: A photo taken through a microscope that shows the anatomy of compact bone with a detailed view of an osteon.

Woven Bone



Woven bone: Woven bone is characterized by the irregular organization of collagen fibers and is mechanically weak.

Woven bone is found on the growing ends of an immature skeleton or, in adults, at the site of a healing fracture. Woven bone is characterized by the irregular organization of collagen fibers and is mechanically weak, but forms quickly.

The criss-cross appearance of the fibrous matrix is why it is referred to as woven. It has a high proportion of osteocytes to hard inorganic salts that leads to its mechanical weakness.

Woven bone is replaced by lamellar bone during development. In contrast to woven bone, lamellar bone is highly organized in concentric sheets with a much lower proportion of osteocytes to surrounding tissue. The regular parallel alignment of collagen into sheets, or, lamellae, causes lamellar bone to be mechanically strong.

Lamellar Bone



Femur head showing trabecular bone: A cross-section of the head of the femur showing lamellar bone on the borders and trabecular bone in the center.

Lamellar bone makes up the compact or cortical bone in the skeleton, such as the long bones of the legs and arms. In a cross-section, the fibers of lamellar bone can be seen to run in opposite directions in alternating layers, much like in plywood, assisting in the bone's ability to resist torsion forces.

When the same lamellar bone is loosely arranged, it is referred to as trabecular bone. Trabecular bone gets its name because of the spongy pattern it displays in an x-ray. The spaces within trabecular bone are filled with active bone marrow. After a fracture, woven bone forms initially, but it is gradually replaced by lamellar bone during a process known as bony substitution.

Key Points

- Woven bone is found on the growing ends of an immature skeleton or, in adults, at the site of a healing fracture.
- Woven bone is characterized by the irregular organization of collagen fibers and is mechanically weak, but forms quickly.
- Lamellar bone is much stronger than woven bone, and is highly organized in concentric sheets with a much lower proportion of osteocytes to mineralized tissue.
- When the same lamellar bone is loosely arranged, it is referred to as trabecular bone. Trabecular bone gets its name because of the spongy pattern it displays on an x-ray.
- After a fracture, woven bone forms initially and is gradually replaced by lamellar bone during a process known as bony substitution.

Key Terms

- **osteoblast**: A mononucleate cell from which bone develops.
- **osteocytes**: A star-shaped type of bone cell that is found in the cells of mature bone.
- **lamellar bone**: A bone with a regular, parallel alignment of collagen into sheets (lamellae) that is mechanically strong.
- **woven bone**: Characterized by an irregular organization of collagen fibers, this bone is mechanically weak.

This page titled 6.3C: Microscopic Anatomy of Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.3D: Chemical Composition of Bone

Learning Objectives

- Differentiate among the acid-base disorders

EXAMPLES

Traveling to a high altitude can cause an acid-base imbalance due to reduced levels of oxygen in the atmosphere, and, therefore, in the blood. To compensate for this, the traveler begins to hyperventilate, trying to expel excess carbon dioxide and bring pH back to normal. However, if the traveler stays at high altitude, it may take several days for their pH to fully return to normal.

Acid-Base Disorders

Acid-base imbalance is an abnormality of the human body's normal balance of acids and bases that causes the plasma pH to deviate out of normal range (7.35 to 7.45). In the fetus, the normal range differs based on which umbilical vessel is sampled (umbilical vein pH is normally 7.25 to 7.45; umbilical artery pH is normally 7.18 to 7.38). Acid-base imbalances can exist in varying levels of severity, some life-threatening.

An excess of acid is called acidosis and an excess in bases is called alkalosis. The process that causes the imbalance is classified based on the etiology of the disturbance (respiratory or metabolic) and the direction of change in pH (acidosis or alkalosis).

Mixed disorders may feature an acidosis and alkalosis excess at the same time that partially counteract each other, or there can be two different conditions affecting the pH in the same direction. The phrase mixed acidosis, for example, refers to metabolic acidosis in conjunction with respiratory acidosis.

Metabolic Acidosis

In medicine, metabolic acidosis is a condition that occurs when the body produces too much acid or when the kidneys are not removing enough acid from the body. If unchecked, metabolic acidosis leads to acidemia, that is, blood pH is less than 7.35 due to increased production of hydrogen by the body, or because of the body's inability to form bicarbonate (HCO_3-) in the kidneys.

Acidosis refers to a low pH in tissue. Acidemia refers to a low pH in the blood. Symptoms may include chest pain, palpitations, headache, altered mental status such as severe anxiety due to hypoxia, decreased visual acuity, nausea, vomiting, abdominal pain, altered appetite (either loss of or increased) and weight loss (longer term), muscle weakness, and bone pains.

Rapid deep breaths increase the amount of carbon dioxide exhaled, thus lowering the serum carbon dioxide levels, resulting in some degree of compensation. Overcompensation via respiratory alkalosis to form an alkalemia does not occur.

Neurological complications include lethargy, stupor, coma, seizures. Cardiac complications include arrhythmias (ventricular tachycardia) and decreased response to epinephrine; both lead to hypotension (low blood pressure).

Metabolic Alkalosis

Metabolic alkalosis is a metabolic condition in which the pH of tissue is elevated beyond the normal range (7.35 to 7.45). This is the result of decreased hydrogen ion concentration, leading to increased bicarbonate concentration, or as a direct result of increased bicarbonate concentrations. Alkalosis refers to a high pH in tissue.

Alkalemia refers to a high pH in the blood. The causes of metabolic alkalosis can be divided into two categories, depending upon urine chloride levels. Chloride-responsive causes result from the loss of hydrogen ions via vomiting or the kidneys. Vomiting results in the loss of hydrochloric acid (hydrogen and chloride ions) with the stomach contents.

The kidneys compensate for these losses by retaining sodium in the collecting ducts at the expense of hydrogen ions (sparing sodium/potassium pumps to prevent further loss of potassium), and leads to metabolic alkalosis. The excess sodium increases extracellular volume and the loss of hydrogen ions creates a metabolic alkalosis.

Later, the kidneys respond through the aldosterone escape to excrete sodium and chloride in urine. Compensation for metabolic alkalosis occurs mainly in the lungs, which retain carbon dioxide (CO_2) through slower breathing, or hypoventilation (respiratory compensation).

CO_2 is then consumed towards the formation of the carbonic acid intermediate, thus decreasing pH. Renal compensation for metabolic alkalosis, less effective than respiratory compensation, consists of increased excretion of HCO_3– (bicarbonate), as the filtered load of HCO_3– exceeds the ability of the renal tubule to reabsorb it.

Respiratory Acidosis

Respiratory acidosis is a medical condition in which decreased ventilation (hypoventilation) causes an increase in blood carbon dioxide concentration and decreased pH (a condition generally called acidosis). Carbon dioxide is produced constantly as the body's cells respire, and this CO_2 will accumulate rapidly if the lungs do not adequately expel it through alveolar ventilation.

Acute respiratory acidosis occurs when an abrupt failure of ventilation occurs. This failure in ventilation may be caused by depression of the central respiratory center by cerebral disease or drugs, an inability to ventilate adequately due to neuromuscular disease (e.g., myasthenia gravis, amyotrophic lateral sclerosis, Guillain-Barré syndrome, muscular dystrophy), or airway obstructions related to asthma or chronic obstructive pulmonary disease (COPD) exacerbation.

Respiratory Alkalosis

Respiratory alkalosis is a medical condition in which increased respiration (hyperventilation) elevates the blood pH (a condition generally called alkalosis). There are two types of respiratory alkalosis: chronic and acute.

Acute respiratory alkalosis occurs rapidly. During acute respiratory alkalosis, the person may lose consciousness whereupon the rate of ventilation will resume to normal.

Chronic respiratory alkalosis is a more long-standing condition. Respiratory alkalosis may be produced accidentally (iatrogenically) during excessive mechanical ventilation. Other causes include: psychiatric causes, drug use, fever, and pregnancy.

Key Points

- Metabolic acidosis can produce, among other symptoms, chest pains, altered mental states, nausea, abdominal pain, and muscle weakness.
- Rapid, deep breathing during metabolic acidosis is an attempt to lower carbon dioxide levels and return pH to normal.
- Extreme acidemia can lead to coma, seizures, heart arrhythmias, and low blood pressure.
- Slowed breathing, which results in retaining more CO_2, is the primary method of reducing metabolic alkalosis.
- Chronic respiratory acidosis is a result of COPD, obesity hypoventilation syndrome, ALS, and thoracic deformities.
- Respiratory alkalosis can be caused by excessive mechanical ventilation, psychiatric problems, stroke, drug use, traveling to high altitude regions, lung disease, fever, and pregnancy, among other factors.

Key Terms

- **metabolic alkalosis**: A metabolic condition in which the pH of tissue is elevated beyond the normal range (7.35 to 7.45). This is the result of decreased hydrogen ion concentration, leading to increased bicarbonate concentration, or a direct result of increased bicarbonate concentration.
- **respiratory acidosis**: A medical condition in which decreased ventilation (hypoventilation) causes increased blood carbon dioxide concentration and decreased pH (a condition generally called acidosis).
- **metabolic acidosis**: A condition that occurs when the body produces too much acid or when the kidneys are not removing enough acid from the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Endosteum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endosteum. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone Tissue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- Flat bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Flat_bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- Long bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Long_bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- epiphyseal plate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/epiphyseal%20plate. **License**: *CC BY-SA: Attribution-ShareAlike*
- medullary cavity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/medullary%20cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- diaphysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/diaphysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Long bone diagram. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Classifcation of Bones. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blausen_0229_ClassificationofBones.png. **License**: *CC BY: Attribution*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Perichondrium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perichondrium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Orthopaedic Surgery/Blood Supply of Long Bones. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Orthopaedic_Surgery/Blood_Supply_of_Long_Bones. **License**: *CC BY-SA: Attribution-ShareAlike*
- General Anatomy/Skeletal System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/General_Anatomy/Skeletal_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- Haversian canal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Haversian_canal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Volkmann's canal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Volkmann's%20canal. **License**: *CC BY-SA: Attribution-ShareAlike*
- anastomose. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/anastomose. **License**: *CC BY-SA: Attribution-ShareAlike*
- Haversian canal. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Haversian_canal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Long bone diagram. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Classifcation of Bones. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blausen_0229_ClassificationofBones.png. **License**: *CC BY: Attribution*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Epiphyseal Plate. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Osteon. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Osteon. **License**: *CC BY-SA: Attribution-ShareAlike*
- Orthopaedic Surgery/Histology of Bone. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Orthopaedic_Surgery/Histology_of_Bone%23Microscopic_Structure. **License**: *CC BY-SA: Attribution-ShareAlike*
- osteoblast. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osteoblast. **License**: *CC BY-SA: Attribution-ShareAlike*

- woven bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/woven%20bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- lamellar bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lamellar%20bone. **License**: *CC BY-SA: Attribution-ShareAlike*
- Long bone diagram. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Classifcation of Bones. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blausen_0229_ClassificationofBones.png. **License**: *CC BY: Attribution*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Epiphyseal Plate. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Compact Bone Histology. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Compact_bone_histology_2014.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Caput femoris cortex medulla. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Caput_femoris_cortex_medulla.jpg. **License**: *Public Domain: No Known Copyright*
- Woven bone matrix. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Woven_bone_matrix.jpg. **License**: *Public Domain: No Known Copyright*
- Respiratory alkalosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Respiratory_alkalosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Metabolic acidosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Metabolic_acidosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Respiratory acidosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Respiratory_acidosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Metabolic alkalosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Metabolic_alkalosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acidu2013base imbalance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acid%E2%80%93base_imbalance. **License**: *CC BY-SA: Attribution-ShareAlike*
- respiratory acidosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/respiratory%20acidosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- metabolic acidosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/metabolic%20acidosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- metabolic alkalosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/metabolic%20alkalosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Long bone diagram. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Classifcation of Bones. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blausen_0229_ClassificationofBones.png. **License**: *CC BY: Attribution*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu compact spongy bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_compact_spongy_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Epiphyseal Plate. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg. **License**: *Public Domain: No Known Copyright*
- Compact Bone Histology. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Compact_bone_histology_2014.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Caput femoris cortex medulla. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Caput_femoris_cortex_medulla.jpg. **License**: *Public Domain: No Known Copyright*
- Woven bone matrix. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Woven_bone_matrix.jpg. **License**: *Public Domain: No Known Copyright*

This page titled 6.3D: Chemical Composition of Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.4: BONE FORMATION

6.4A: Embryonic and Fetal Bone Formation

6.4B: Postnatal Bone Growth

6.4C: Bone Remodeling

6.4D: Bone Repair

This page titled 6.4: Bone Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.4A: Embryonic and Fetal Bone Formation

During fetal development, bone tissue is created through intramembranous ossification and endochondral ossification.

Learning Objectives

- Differentiate intramembranous ossification from enchondral ossification in embryonic development


Key Points

- Intramembranous ossification occurs during fetal development and does not involve cartilage.
- Embryologic mesenchymal cells differentiate into osteogenic cells that direct bone growth from spicules to trabeculae, to woven bone, and finally to lamellar bone.
- Endochondral ossification creates fetal long bones from a cartilage template.
- Osteoblasts are involved in both intramembranous and endochondral ossification.
- When osteoblasts become trapped in the matrix they differentiate into osteocytes.
- Osteons are units or principal structures of compact bone.

Key Terms

- **osteon**: Any of the central canals, and surrounding bony layers, found in compact bone.
- **canaliculi**:
- **endochondral ossification**: A process that occurs during fetal development by which bone tissue is created using a cartilage template.
- **intramembranous ossification**: A process that occurs during fetal development to produce bone tissue without a cartilage template. The membrane that occupies the place of the future bone resembles connective tissue and ultimately forms the periosteum, or outer bone layer.

Fetal Development



Mesenchymal stem cell: Mesenchymal stem cells initiate the process of intramembranous ossification.

Embryonic/fetal development proceeds from rostral (nose and mouth area) to caudal (posterior). The skull and vertebral column are produced by intramembranous ossification. As development proceeds down the body axis, the long bones of the arms and legs are produced by endochondral ossification.


Intramembranous ossification is one of the two essential processes during fetal development of the mammalian skeletal system. It is the process by which bone tissue is created.

Unlike the other process of bone creation— endochondral ossification—intramembranous ossification does not involve cartilage. It is also an essential process during the natural healing of bone fractures and the rudimentary formation of the bones of the head.

The first step in the process is the formation of bone spicules (aggregates of bony matrix) that eventually fuse with each other and become trabeculae. The periosteum is formed and bone growth continues at the surface of trabeculae.

Much like spicules, the increasing growth of trabeculae result in interconnection, and this network is called woven bone. Eventually, woven bone is replaced by lamellar bone.

Embryonic mesenchymal cells (MSC) condense into layers of vascularized primitive connective tissue. Certain mesenchymal cells group together, usually near or around blood vessels, and differentiate into osteogenic cells that deposit bone matrix constitutively. Separate mesenchymal cells differentiate into osteoblasts, which line up along the surface of the spicule and secrete more osteoid, increasing the size of the spicule.



Osteoblasts: This figure shows osteoblasts creating rudimentary bone tissue.

When osteoblasts become trapped in the matrix that they secrete, they differentiate into osteocytes. Osteoblasts continue to line up on the surface, which increases their size. As growth continues, trabeculae become interconnected and woven bone is formed.

The primary center of ossification is the area where bone growth occurs between the periosteum and the bone.

Osteons are units or principal structures of compact bone. During the formation of bone spicules, cytoplasmic processes from osteoblasts interconnect. This becomes the canaliculi of osteons.

Since bone spicules tend to form around blood vessels, the perivascular space is greatly reduced as the bone continues to grow. When replacement with compact bone occurs, this blood vessel becomes the central canal of the osteon.

Endochondral Ossification



Cartilage: Hyaline cartilage showing chondrocytes and organelles, lacunae and matrix.

Endochondral ossification is the other essential bone creation process during fetal development of the mammalian skeletal system. Unlike intramembranous ossification, cartilage is present during endochondral ossification. It is also an essential process during the rudimentary formation of long bones, the growth of the length of long bones, and the natural healing of bone fractures.

The first site of ossification occurs in the primary center of ossification, which is in the middle of diaphysis (shaft). The perichondrium becomes the periosteum. The periosteum contains a layer of undifferentiated cells (osteoprogenitor cells) that later become osteoblasts.

The osteoblasts secrete osteoid against the shaft of the cartilage model (appositional growth). This serves as support for the new bone. Chondrocytes in the primary center of ossification begin to grow (hypertrophy). They stop secreting collagen and other proteoglycans and begin secreting alkaline phosphatase, an enzyme essential for mineral deposition. Then calcification of the matrix occurs.

This page titled 6.4A: Embryonic and Fetal Bone Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.4B: Postnatal Bone Growth


Learning Objectives

- Differentiate among the zones of development from cartilage to bone in postnatal ossification

Postnatal Ossification


Secondary ossification occurs after birth. It forms the epiphyses of long bones and the extremities of irregular and flat bones.

The diaphysis and both epiphyses of a long bone are separated by a growing zone of cartilage (the epiphyseal plate). When a child reaches skeletal maturity (18 to 25 years of age), all of the cartilage is replaced by bone, fusing the diaphysis and both epiphyses together (epiphyseal closure). This process involves replacing the hyaline cartilage, initially present at the epiphyseal region, with active osteoblasts that deposit bone structural proteins.



Epiphyseal plate: The epiphysis is the rounded end of a long bone located at its joint with adjacent bone(s). Between the epiphysis and diaphysis (the long midsection of the long bone) lies the metaphysis, including the epiphyseal plate (growth plate).

The Role of Osteoblasts



Developing femur: Pictured is part of a longitudinal section of a rabbit's developing femur, with parts including: a) Flattened cartilage cells; b) Enlarged cartilage cells; c), d) Newly formed bone; e) Osteoblasts; f) Giant cells or osteoclasts; g), h) Shrunken cartilage cells.

Osteoblasts are mononucleate cells that are responsible for bone formation. In essence, osteoblasts are specialized fibroblasts that, in addition to fibroblastic products, express bone sialoprotein and osteocalcin.

Osteoblasts produce a matrix of osteoid that is composed mainly of Type I collagen. Osteoblasts are also responsible for the mineralization of this matrix. Minerals required for mineralization and related processes include zinc, copper, and sodium.


Bone is a dynamic tissue that is constantly being reshaped by osteoblasts and osteoclasts. Osteoblasts produce bone matrix and mineral, and osteoclasts break down the tissue. The number of osteoblasts tends to decrease with age, affecting the balance of the formation and resorption in the bone tissue, and potentially leading to osteoporosis.

During postnatal bone formation, endochondral ossification initiates bone deposition by first generating a structural framework at the ends of long bones, within which the osteoblasts can synthesize a new bone matrix.

Cartilage to Bone

1. Zone of reserve cartilage: This region is farthest from the marrow cavity and consists of hyaline cartilage that does not actively transform into bone. Quiescent chondrocytes are found here.
2. Zone of cell proliferation: Closer to the marrow cavity, chondrocytes in this region multiply and arrange themselves into longitudinal columns of flattened lacunae.
3. Zone of cell hypertrophy: They stop dividing and begin to hypertrophy (enlarge). The walls of the matrix between the lacunae become very thin.
4. Zone of calcification: The region where the cartilagenous matrix begins to calcify.
   Minerals are deposited in the matrix between the columns of lacunae, but are not the permanent bone mineral deposits. This acts as a temporary support for the cartilage that would otherwise be weakened due to the breakdown of the lacunae.
5. Zone of bone deposition (ossification): The walls between the lacunae break down and the chondrocytes die. Each column then becomes a longitudinal channel that is immediately invaded by blood vessels and marrow from the marrow cavity. Osteoblasts begin depositing concentric lamellae of matrix, while osteoclasts dissolve the temporarily calcified cartilage.

The growth in the diameter of bones around the diaphysis occurs through the deposition of bone beneath the periosteum. Osteoclasts in the interior cavity continue to degrade bone until its ultimate thickness is achieved. At this point the rate of formation on the outside and degradation from the inside is constant. This process is termed appositional growth.



Endochondral ossification: The development of the primary and secondary ossification centers.

Key Points

- The epiphyseal plate is the growth zone between the diaphysis (shaft) of the long bone and the epiphysis (end) of the long bone.
- At skeletal maturity, growth ceases when the epiphyses fuse with the diaphyses, indicating that all the cartilage has been replaced with bone and epiphyseal closure has been achieved.
- Osteoblasts are mononucleate cells that are responsible for bone formation.
- Bone is a dynamic tissue that is constantly being reshaped by osteoblasts and osteoclasts.
- Osteoblasts produce a matrix of osteoid, which is composed mainly of Type I collagen.

Key Terms

- **osteocalcin**: Secreted solely by osteoblasts, it is pro-osteoblastic (bone-building), and is implicated in bone mineralization and calcium ion homeostasis.
- **sialoprotein**: A component of mineralized tissues, such as bone, that acts as a nucleus for the formation of the first apatite crystals.
- **secondary ossification**: A process that occurs after birth, and forms the epiphyses of long bones and the extremities of irregular and flat bones.
- **diaphyses**: The main or mid section (shaft) of a long bone that is made up of cortical bone.
- **epiphyseal closure**: The fusion of the epiphysis to the diaphysis.

This page titled 6.4B: Postnatal Bone Growth is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


6.4C: Bone Remodeling

Learning Objectives

- Explain the effects of growth factors on bone

Remodeling or bone turnover is the process of resorption followed by the replacement of bone with limited change in shape; this process occurs throughout a person's life. Repeated stress, such as weight-bearing exercise or bone healing, results in the bone thickening at the points of maximum stress.

It has been hypothesized that this is a result of bone's piezoelectric properties that cause bone to generate small electrical potentials under stress. Osteoblasts and osteoclasts, coupled together via paracrine cell signaling, are referred to as bone remodeling units. The purpose of remodeling is to regulate calcium homeostasis, repair micro-damaged bones (from everyday stress), and to shape and sculpture the skeleton during growth.



Bone remodeling: Bone tissue is removed by osteoclasts, and then new bone tissue is formed by osteoblasts. Both processes utilize cytokine (TGF-β, IGF) signaling.

Bone volume is determined by the rates of bone formation and bone resorption. The action of osteoblasts and osteoclasts are controlled by a number of chemical factors that either promote or inhibit the activity of the bone remodeling cells, controlling the rate at which bone is made, destroyed, or changed in shape. The cells also use paracrine signalling to control the activity of each other.

Role of Growth Factors

Recent research has suggested that certain growth factors may work to locally alter bone formation by increasing osteoblast activity. Numerous bone-derived growth factors have been isolated and classified via bone cultures. These factors include insulin-like growth factors I and II, transforming growth factor beta, fibroblast growth factor, platelet-derived growth factor, and bone morphogenetic proteins.

- Insulin-like growth factors protect cartilage cells, and are associated with the activation of osteocytes.
- The transforming growth factor beta superfamily includes bone morphogenic proteins involved in osteogenesis.
- Fibroblast growth factor activates various cells of the bone marrow including osteoclasts and osteoblasts.
- Platelet-derived growth factor has been found to enhance bone collagen degradation.

Evidence suggests that bone cells produce growth factors for extracellular storage in the bone matrix. The release of these growth factors from the bone matrix could cause the proliferation of osteoblast precursors. Essentially, bone growth factors may act as potential determinants of local bone formation.

Research has suggested that trabecular bone volume in postmenopausal osteoporosis may be determined by the relationship between the total bone forming surface and the percent of surface resorption.

Clinical Note

Osteoporosis means porous bone, which is caused by an over-reaction to osteoclastic bone resorption, and makes bones quite fragile for the elderly. Falls are dangerous for the elderly because they are more likely to break a bone. Hip fractures are especially troublesome as they result in a long recovery period during which complications that may lead to death are quite common.

Key Points

- Bone remodeling involves resorption by osteoclasts and replacement by osteoblasts. Osteoblasts and osteoclasts are referred to as bone remodeling units.
- The purpose of bone remodeling is to regulate calcium homeostasis, repair micro-damage to bones from everyday stress, and to shape the skeleton during growth.
- Bone growth factors affect the process of bone remodeling. These factors include insulin-like growth factors I and II, transforming growth factor beta, fibroblast growth factor, platelet-derived growth factor, and bone morphogenetic proteins.
- Bone volume is determined by the rates of bone formation and bone resorption. The action of osteoblasts and osteoclasts are controlled by a number of chemical factors that either promote or inhibit the activity of the bone remodeling cells.
- Postmenopausal osteoporosis is the result of imbalances in the relationship between bone resorption and replacement.

Key Terms

- **growth factors**: Naturally occurring substances that stimulate cell growth, proliferation, healing, and cellular differentiation.
- **piezoelectric**: Accumulation of an electric charge due to mechanical stress.
- **bone remodeling**: The resorption of bone by osteoclasts and replacement by osteoblasts.

This page titled 6.4C: Bone Remodeling is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.4D: Bone Repair

Learning Objectives

- Explain the process of bone repair



Bone repair: This figure depicts a communitive midshaft humeral fracture with callus formation.

Bone healing, or fracture healing, is a proliferative physiological process in which the body facilitates the repair of a bone fracture. Generally, bone fracture treatment consists of a doctor reducing (pushing) dislocated bones back into place via relocation with or without anesthetic, stabilizing their position, and then waiting for the bone's natural healing process to occur.

While immobilization and surgery may facilitate healing, a fracture ultimately heals through physiological processes. The healing process is mainly determined by the periosteum (the connective tissue membrane covering the bone).

The periosteum is one source of precursor cells that develop into the chondroblasts and osteoblasts that are essential to heal bone. The bone marrow (when present), endosteum, small blood vessels, and fibroblasts are other sources of precursor cells.

Days after a fracture, the cells of the periosteum replicate and transform. The periosteal cells proximal (closest) to the fracture gap develop into chondroblasts that form hyaline cartilage.

The periosteal cells distal to (further from) the fracture gap develop into osteoblasts that form woven bone. The fibroblasts within the granulation tissue develop into chondroblasts that also form hyaline cartilage.

These two new tissues grow in size until they unite with their counterparts from other parts of the fracture. These processes culminate in a new mass of heterogeneous tissue that is known as the fracture callus.

Eventually, the fracture gap is bridged by the hyaline cartilage and woven bone, restoring some of its original strength.



Healing fracture: This figure depicts a radiograph of a child's healing supracondylar humeral fracture that has been treated with closed reduction and pinning. This image, taken three weeks post injury, demonstrates the benign periosteal reaction of normal healing bone.

The next phase is the replacement of the hyaline cartilage and woven bone with lamellar bone. The replacement process is known as endochondral ossification with respect to the hyaline cartilage and bony substitution with respect to the woven bone.

Substitution of the woven bone with lamellar bone precedes the substitution of the hyaline cartilage with lamellar bone. The lamellar bone begins forming soon after the collagen matrix of either tissue becomes mineralized. At this point, the mineralized matrix is penetrated by channels, each containing a microvessel and numerous osteoblasts.

The osteoblasts form new lamellar bone upon the recently exposed surface of the mineralized matrix. This new lamellar bone is in the form of trabecular bone. Eventually, all of the woven bone and cartilage of the original fracture callus is replaced by trabecular bone, restoring most of the bone's original strength.

The remodeling process continues with substitution of the trabecular bone with compact bone. The trabecular bone is first resorbed by osteoclasts, creating a shallow resorption pit known as Howship's lacuna, and then osteoblasts deposit compact bone within the resorption pit.

Eventually, the fracture callus is remodeled into a new shape that closely duplicates the bone's original shape and strength. The remodeling phase takes three to five years depending on factors such as age or general condition.

When the humerus in the upper arm is fractured and properly set, bone healing can repair the bone. However, if the bone is not set or improperly set, the chondroblasts and osteoblasts will still try to heal the bone but will be unable to return the bone to full proper functioning.

Key Points

- Cells of the periosteum (connective tissue membrane covering the bone) replicate and transform into chondroblasts, which form hyaline cartilage proximal to the gap, and osteoblasts, which form woven bone distal to the gap. The components join to form a fracture callus.
- Endochondral ossification then replaces the cartilage and woven bone with lamellar bone that restores the integrity and strength of the bone.
- The remodeling process substitutes the trabecular bone with compact bone. The trabecular bone is first resorbed by osteoclasts, creating a shallow resorption pit. Then osteoblasts deposit compact bone within the resorption pit. The callus is remodeled to duplicate the original bone.

Key Terms

- **lacuna**: A small opening, a small pit or depression, a small blank space, a gap or vacancy, or a hiatus.
- **fracture callus**: A fibrocartilage or fracture callus is a temporary formation of fibroblasts and chondroblasts that form at the area of a bone fracture as the bone attempts to heal itself.
- **compact bone**: One of the two types of osseous tissue that form bones.
- **bone healing**: Bone healing, or fracture healing, is a proliferative physiological process in which the body facilitates the repair of a bone fracture.
- **endosteum**: A membranous vascular layer of cells that line the medullary cavity of a bone.
- **chondroblast**: A cell that originates from a mesenchymal stem cell and forms chondrocytes.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Bone Canaliculus. **Provided by**: Wikipecia. **Located at**: **en.Wikipedia.org/wiki/Bone_canaliculus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endochondral ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Endocho...l_ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intramembranous ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intrame...s_ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intramembranous ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intrame...20ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endochondral ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/endocho...20ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/osteon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mesenchymal stem cells. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi..._Stem_Cell.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hyaline cartilage. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...seal_Plate.jpg**. **License**: ***Public Domain: No Known Copyright***
- Osteoblasts. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ny_nidus_3.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Osteocalcin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Osteocalcin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone Sailoprotein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone_sialoprotein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endochondral ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Endocho...l_ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epiphyseal plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epiphyseal_plate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone%23Formation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone%23...l_ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Osteoblasts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Osteoblasts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epiphyseal closure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/epiphyseal%20closure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diaphyses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/diaphyses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secondary ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/secondary%20ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mesenchymal stem cells. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f4/**

Mesenchymal_Stem_Cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hyaline cartilage. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/e/ee/Hypertrophic_Zone_of_Epiphyseal_Plate.jpg**. **License**: ***Public Domain: No Known Copyright***
- Osteoblasts. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/b8/Bony_nidus_3.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Epiphysis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: ***Public Domain: No Known Copyright***
- Primary and secondary ossification. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/c/cf/Bone_growth.png**. **License**: *Public Domain: No Known Copyright*
- Gray80. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray80.png**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Growth Factor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Growth_factor**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone%23Remodeling**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone%23Remodeling**. **License**: *CC BY-SA: Attribution-ShareAlike*
- bone remodeling. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bone%20remodeling**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mesenchymal stem cells. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f4/Mesenchymal_Stem_Cell.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hyaline cartilage. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/e/ee/Hypertrophic_Zone_of_Epiphyseal_Plate.jpg**. **License**: *Public Domain: No Known Copyright*
- Osteoblasts. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/b8/Bony_nidus_3.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Epiphysis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: *Public Domain: No Known Copyright*
- Primary and secondary ossification. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/c/cf/Bone_growth.png**. **License**: *Public Domain: No Known Copyright*
- Gray80. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray80.png**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone metabolism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bonemetabolism.svg**. **License**: *CC BY: Attribution*
- Bone healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone_healing**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Compact bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Compact_bone**. **License**: *CC BY-SA: Attribution-ShareAlike*
- fracture callus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fracture%20callus**. **License**: *CC BY-SA: Attribution-ShareAlike*
- lacuna. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lacuna**. **License**: *CC BY-SA: Attribution-ShareAlike*
- bone healing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bone%20healing**. **License**: *CC BY-SA: Attribution-ShareAlike*
- endosteum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endosteum**. **License**: *CC BY-SA: Attribution-ShareAlike*
- chondroblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chondroblast**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mesenchymal stem cells. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f4/Mesenchymal_Stem_Cell.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hyaline cartilage. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/e/ee/Hypertrophic_Zone_of_Epiphyseal_Plate.jpg**. **License**: *Public Domain: No Known Copyright*
- Osteoblasts. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/b8/Bony_nidus_3.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Epiphysis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: *Public Domain: No Known Copyright*
- Primary and secondary ossification. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/c/cf/Bone_growth.png**. **License**: *Public Domain: No Known Copyright*
- Gray80. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray80.png**. **License**: *CC BY-SA: Attribution-ShareAlike*

- Bone metabolism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bonemetabolism.svg**. **License**: ***CC BY: Attribution***
- Callus formation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f5/Communitive_midshaft_humeral_fracture_callus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Healing supracondylar humeral fracture. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/7/7a/Healing_supracondylar_fracture.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 6.4D: Bone Repair is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.5: BONE AND CALCIUM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

6.5A: Bone and Calcium Homeostasis

This page titled 6.5: Bone and Calcium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

6.5A: Bone and Calcium Homeostasis

Calcium metabolism or calcium homeostasis is the mechanism by which the body maintains adequate calcium levels.

Learning Objectives

- Describe the factors involved in calcium homeostasis

Key Points


- Calcium homeostasis regulates calcium flow to and from the bones.
- Inadequate calcium levels can result in osteoporosis.
- Calcium release from bone is regulated by parathyroid hormone.
- Vitamin D is converted to calcidiol (also called calcifediol) in the liver, which is then converted to calcitriol in the kidneys, the biologically active form of vitamin D.
- Calcitriol regulates the levels of calcium and phosphorus in the blood and helps maintain a healthy skeletal system.
- Bone resorption by osteoclasts releases calcium into the bloodstream, which helps regulate calcium homeostasis.

Key Terms

- **calcitriol**: The active metabolite—1,25-dihydroxycholecalciferol—that is involved in the absorption of calcium.
- **Calcitonin**: Produced in humans by the thyroid gland, it acts to reduce blood calcium, opposing the effects of parathyroid hormone.
- **calcium homeostasis**: Calcium homeostasis is the mechanism by which the body maintains adequate calcium levels in order to prevent hypercalcemia or hypocalcemia, both of which can have important consequences for health.
- **calcidiol**: A prehormone that is produced in the liver by the hydroxylation of vitamin D3 (cholecalciferol) by the enzyme cholecalciferol 25-hydroxylase.

Calcium metabolism or calcium homeostasis is the mechanism by which the body maintains adequate calcium levels. Derangements of this mechanism lead to hypercalcemia or hypocalcemia, both of which can have important consequences for health.

Although calcium flow to and from the bone is neutral, about five mmol is turned over a day. Bone serves as an important storage point for calcium, as it contains 99% of the total body calcium. Calcium release from bone is regulated by parathyroid hormone. Calcitonin stimulates incorporation of calcium in bone.



Calcium regulation: Parathyroid hormone regulates the levels of calcium in the blood.

Low calcium intake may be a risk factor in the development of osteoporosis. With a better bone balance, the risk of osteoporosis is lowered. Supplementation with vitamin D and calcium slightly improves bone mineral density.

Vitamin D and Calcium Homeostasis

Vitamin D is converted to calcidiol in the liver. Part of the calcidiol is converted by the kidneys to calcitriol, the biologically active form of vitamin D. It circulates as a hormone in the blood, regulating the concentration of calcium and phosphate in the bloodstream and promoting the healthy growth and remodeling of bone.

Responses to Blood Calcium Changes

The process of bone resorption by the osteoclasts releases stored calcium into systemic circulation and is an important process for regulating calcium balance. As bone formation actively fixes circulating calcium in its mineral form by removing it from the bloodstream, resorption actively unfixes it, thereby increasing circulating calcium levels.

When blood calcium concentration rises, the parafollicular cells of the thyroid gland increase calcitonin secretion into the blood. At the same time, the parathyroid glands reduce parathyroid hormone secretion into the blood. The resulting high levels of calcitonin in the blood stimulate the bone to remove calcium from the blood plasma and deposit it as bone.

Removal of calcium from the bone is also inhibited. When the blood calcium level is too low, calcitonin secretion is inhibited and PTH secretion is stimulated. This results in the removal of calcium from the bone to correct blood calcium levels.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Calcitonin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Calcitonin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Calcium metabolism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Calcium...e_role_of_bone**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Vitamin D. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vitamin_D%23Bone_health. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone%23Calcium_balance. **License**: *CC BY-SA: Attribution-ShareAlike*
- calcidiol. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/calcidiol. **License**: *CC BY-SA: Attribution-ShareAlike*
- calcium homeostasis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/calcium%20homeostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- calcitriol. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/calcitriol. **License**: *CC BY-SA: Attribution-ShareAlike*
- Calcium regulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ca...regulation.png. **License**: *Public Domain: No Known Copyright*

This page titled 6.5A: Bone and Calcium Homeostasis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7: SKELETAL SYSTEM - PARTS OF THE SKELETON

7.1: The Skull

7.1A: General Features and Functions of the Skull

7.1B: Cranial Bones

7.1C: Facial Bones

7.1D: Orbits

7.1E: Foramina

7.1F: Fontanelles

7.1F: Sutures

7.1G: Paranasal Sinuses

7.2: The Hyoid Bone

7.2A: The Hyoid Bone

7.3: The Vertebral Column

7.3A: General Characteristics of the Spine

7.3B: Parts of a Vertebra

7.3C: Regional Vertebral Characteristics

7.4: The Thorax

7.4A: Thoracic Cage: Sternum

7.4B: Thoracic Cage: Ribs

7.5: The Shoulder

7.5A: Clavicle

7.5B: Scapula

7.6: The Upper Limb

7.6A: Humerus (The Upper Arm)

7.6B: Ulna and Radius (The Forearm)

7.6C: Carpals, Metacarpals, and Phalanges (The Hand)

7.7: The Hip

7.7A: Ilium

7.7B: Ischium

7.7C: Pubis

7.7D: False and True Pelves

7.7E: Comparison of Female and Male Pelves

7.7F: Pelvic Structure and Childbearing

7.8: The Lower Limb

7.8A: Femur (The Thigh)

7.8B: Patella (The Knee)

7.8C: Tibia and Fibula (The Leg)
7.8D: Tarsals, Metatarsals, and Phalanges (The Foot)
7.9E: Arches of the Feet

This page titled 7: Skeletal System - Parts of the Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1: THE SKULL



-  Topic hierarchy 

7.1A: General Features and Functions of the Skull

7.1B: Cranial Bones

7.1C: Facial Bones

7.1D: Orbits

7.1E: Foramina

7.1F: Fontanelles

7.1F: Sutures

7.1G: Paranasal Sinuses

This page titled 7.1: The Skull is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1A: General Features and Functions of the Skull

The human skull is the part of the skeleton that supports the structures of the face and forms a cavity for the brain.

Learning Objectives

- Describe the parts of the skull

Key Points

- The adult human skull consists of two regions of different embryological origins: the neurocranium and the viscerocranium.
- The neurocranium is a protective shell surrounding the brain and brain stem.
- The viscerocranium (or facial skeleton) is formed by the bones supporting the face.
- Except for the mandible, all skull bones are joined together by sutures —synarthrodial (immovable) joints.
- The skull contains air-filled cavities called sinuses. Their functions are debatable, but may be related to lessening skull weight, contributing to voice resonance, and warming and moistening inspired air.

Key Terms

- **viscerocranium**: The skeleton that supports facial structure.
- **sutures**: A fairly rigid joint between bones of the neurocranium.
- **neurocranium**: The protective vault surrounding the brain and brain stem.


The skull supports the musculature and structures of the face and forms a protective cavity for the brain. The skull is formed of several bones which, with the exception of the mandible, are joined together by sutures—synarthrodial (immovable) joints.

Composition of the Skull

The adult human skull is comprised of twenty-two bones which are divided into two parts of differing embryological origin: the neurocranium and the viscerocranium.

Neurocranium


The neurocranium forms the cranial cavity that surrounds and protects the brain and brainstem. The neurocranium is formed from the occipital bone, two temporal bones, two parietal bones, the sphenoid, ethmoid and frontal bones; they are all joined together with sutures.



Components of neurocranium: Neurocranium consists of 8 parts: frontal, sphenoid, ethmoid, occipital, 2 temporal, and 2 parietal bones.

Viscerocranium

The viscerocranium bones form the anterior and lower regions of the skull and include the mandible, which attaches through the only truly motile joint found in the skull. The facial skeleton contains the vomer, two nasal conchae, two nasal bones, two maxilla, the mandible, two palatine bones, two zygomatic bones, and two lacrimal bones.



Components of viscerocranium : The face is made of 2 nasal, 2 lacrimal, 2 palantine, 2 inferior nasal concha, 2 zygomatic, 2 maxilla, 1 mandible, and 1 vomer.

Other Features

The skull also contains the sinuses. These are air-filled cavities that contribute to lessening the weight of the skull with a minimal reduction in strength. They contribute to resonance of the voice and assist in the warming and moistening of air inhaled via the nose.

This page titled 7.1A: General Features and Functions of the Skull is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


7.1B: Cranial Bones

Learning Objectives

- Differentiate the bones of the neurocranium

The neurocranium forms the cranial cavity that surrounds and protects the brain and brainstem. The neurocranium consists of the occipital bone, two temporal bones, two parietal bones, the sphenoid, ethmoid, and frontal bones—all are joined together with sutures.

Evolutionary,it is the expansion of the neurocranium that has facilitated the expansion of the brain and its associated developments.



Neurocranium: A lateral view showing the bones that structure the neurocranium.

Occipital Bone

The occipital bone forms the base of the skull at the rear of the cranium. It articulates with the first vertebra of the spinal cord and also contains the foramen magnum, the large opening of the skill through which the spinal cord passes as it enters the vertebral column. The occipital bone borders the parietal bones through the heavily serrated lambdoidal suture, and also the temporal bones through occipitomastoid suture.

Temporal Bones

The temporal bones are situated at the base and sides of the skull, lateral to the temporal lobes of the brain. The temporal bones consist of four regions the squamous, mastoid, petrous and tympanic regions.

The squamous region is the largest and most superior region. Inferior to the squamous is the mastoid region, and fused between the squamous and mastoid regions is the petrous region. Finally, the small and inferior tympanic region lies anteriorly to the mastoid.

There are two processes that originate from the temporal bone:

1. The zygomatic process that projects from the lower squamous region and articulates with the zygomatic bone of the cheek.
2. The styloid process projects downwards from the interior of the temporal bone and provides attachment for several muscles associated with the tongue.

The temporal bones have four borders:

1. The occipitomastoid suture separates the occipital bone and mastoid portion of temporal bone.
2. The squamosal suture separates the parietal bone and squama portion of temporal bone.
3. The sphenosquamosal suture separates the sphenoid bone and squama portion of temporal bone.
4. The zygomaticotemporal suture separates the zygomatic bone and zygomatic process of temporal bone.

Parietal Bones

The two large parietal bones are connected and make up part of the roof and sides of the human skull. The two bones articulate to form the sagittal suture. In the front, the parietal bones form the coronal suture with the frontal bone, and in the rear, the lambdoid suture is formed by the occipital bone. Finally, the squamosal suture separates the parietal and temporal bones.

Sphenoid Bone

The sphenoid bone is situated in the middle of the skull towards the front and forms the rear of the orbit. It has been described as resembling a butterfly due to its wing-like processes. The sphenoid bone is divided into several parts: the body of the bone, two greater wings, two lesser wings, and the pterygoid processes.

The sphenoid bone is one of the most complex in the body due to its interactions with numerous facial bones, ligaments, and muscles. The body that forms the middle of the sphenoid bone articulates with the ethmoid and occipital bone and forms a key part of the nasal cavity; it also contains the sphenoidal sinuses.

The greater wings form the floor of the middle cranial fossa that houses the frontal lobes and pituitary gland, and also the posterior wall of the orbit. The lesser wings project laterally and form the floor of the anterior cranial fossa and the superior orbital fissure through which several key optical nerves pass.

Ethmoid Bone

The ethmoid bone is a small bone in the skull that separates the nasal cavity from the brain. It is lightweight due to its spongy, air-filled construction and is located at the roof of the nose and between the two orbits.

The ethmoid bone forms the medial wall of the orbit, the roof of the nasal cavity, and due to its central location it articulates with numerous bones of the viscerocranium. Inside the neurocranium it articulates with the frontal and sphenoid bones.

Frontal Bone

The frontal bone forms the front of the skull and is divided into three parts:

1. Squamous: This part is large and flat and forms the main region of the forehead.
2. Orbital: This part lies inferiorly and forms the superior border of the orbit.
3. Nasal: this part is smaller and articulates with the nasal bones and maxilla to contribute to the roof of the nose.

The frontal bone borders two other neurocranial bones—the parietal bones through the coronal sutures and the sphenoid bone through the sphenofrontal suture. It also articulates with the zygomatic and nasal bones and the maxilla.

Key Points

- The eight bones of the neurocranium form major portions of the skull and protect the brain.
- The neurocranium consists of two temporal bones situated to the base and side of the skull, and two parietal bones that make up the roof of the skull.

- A single occipital bone forms the base of the skull, and the frontal bone forms the forehead.
- The sphenoid and ethmoid bones located to the front of the skull form parts of the orbital sockets and nasal cavity; they also support and protect key organs found in the skull.

Key Terms

- **neurocranium**: The part of the skull that encloses and protects the brain and brain stem.

This page titled 7.1B: Cranial Bones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1C: Facial Bones

The viscerocranium (face) includes these bones: vomer, 2 inferior nasal conchae, 2 nasals, maxilla, mandible, palatine, 2 zygomatics, and 2 lacrimals.

Learning Objectives

- List the facial bones of the viscerocranium

Key Points

- The several bones of the viscerocranium are joined by sutures to each other and the neurocranium, except for the mandible, which articulates with the temporal bones.
- The hyoid bone, ethmoid bone, and sphenoid bones are sometimes included in the viscerocranium.

Key Terms

- **viscerocranium**: The facial skeleton that is formed by the bones of the anterior and lower skull, which are derived from branchial arches.

The Viserocranium

The viscerocranium or facial bones supports the soft tissue of the face. The viscerocranium consists of 14 individual bones that fuse together. However, the hyoid bone, ethmoid bone, and sphenoid bones are sometimes included in the viscerocranium.

Zygomatic Bones

The two zygomatic bones form the cheeks and contribute to the orbits. They articulate with the frontal, temporal, maxilla, and sphenoid bones.

Lacrimal Bones

The two lacrimal bones form the medial wall of the orbit and articulate with the frontal, ethmoid, maxilla, and inferior nasal conchae. The lacrimal bones are the two smallest bones located in the face.

Nasal Bones

The two slender nasal bones located in the midline of the face fuse to form the bridge of the noise and also articulate with the frontal, ethmoid and maxilla bones. The inferior nasal conchae are located within the nasal cavity. They are spongy and curled in shape; their primary function is to increase the surface area of the nasal cavity, which also increases the amount of air that contacts the mucous membranes and cilia of the nose, thus filtering, warming, and humidifying the air before it enters the lungs. At the base of the nasal cavity is the small vomer bone which forms the nasal septum.

Maxilla Bones


The maxilla bones fuse in the midline and form the upper jaw. They provide the bed for the upper teeth, the floor of the nose, and the base of the orbits. The maxilla articulates with the zygomatic, nasal, lacrimal, and palatine bones.

Palatine Bones

The palatine bones fuse in the midline to form the palatine, located at the back of the nasal cavity that forms the roof of the mouth and the floor of the orbit.

The Mandible

Finally, the mandible forms the lower jaw of the skull. The joint between the mandible and the temporal bones of the neurocranium, known as the temporomandibular joint, forms the only non-sutured joint in the skull.



Facial bones: There are fourteen facial bones. Some, like the lacrimal and nasal bones, are paired. Others, like the mandible and vomer, are singular.

This page titled 7.1C: Facial Bones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1D: Orbits

Learning Objectives

- Locate the orbits in the skull

The orbit, or eye socket, is the cavity located in the skull in which the eye and its associated appendages are housed. The orbits are conical, sometimes described as four-sided pyramidal, cavities that open in the midline of the face and point backwards. To the rear of the orbit, the optic foramen opens into the optical canal through which the optic nerve and ophthalmic artery pass.

The primary functions of the orbit include protection of its delicate contents and, through muscle attachment and a smooth coating fascia, to also promote the smooth, delicate movements of the eye.

Structure of the Orbit

The orbital cavity is formed from seven bones. The frontal bone forms the superior border of the orbital rim and also the superior wall (roof) of the orbital surface.

The zygomatic bone forms the lateral (and half of the basal) border of the orbital rim, and also the lateral wall of the orbital surface—this is the thickest region of the orbit as it is most exposed to external trauma.

Completing the basal and medial border of the orbital rim is the maxillary bone, which also forms the inferior wall (floor) of the orbital surface.

The lacrimal and ethmoid bones contribute to the medial wall of the orbit and also to the medial wall of the orbital canal. The small palatine bone contributes to the floor of the orbit.

Finally, the sphenoid bone forms the posterior wall of the orbit and also contributes to the formation of the optic canal.

Key Points

- The orbits are conical or four-sided pyramidal cavities, which open into the midline of the face and point backwards.
- To the rear of the orbit the optical foramen opens into the optic canal, which transmits the optic nerve and opthalmic artery.
- The orbit protects the eye from mechanical injury and provides access for the optic nerve to the brain.

Key Terms

- **optic canal**: The canal that transmits the optic nerve and ophthalmic artery into the orbital cavity.
- **orbit**: The bony cavity containing the eyeball; the eye socket.

This page titled 7.1D: Orbits is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1E: Foramina

The human skull has numerous holes known as foramina through which cranial nerves, arteries, veins, and other structures pass.

Learning Objectives


- Describe the purpose of foramina in the skull

Key Points

- A foramen (plural: foramina) is an opening inside the body that allows key structures to connect one part of the body to another.
- The skull bones that contain foramina include the frontal, ethmoid, sphenoid, maxilla, palatine, temporal, and occipital.
- There are 21 foramina in the human skull.

Key Terms

- **foramina**: The openings inside the body that typically allow muscles, nerves, arteries, veins, or other structures to connect one part of the body to another.



Base of the skull (upper surface): This image details the foramina of the skull.

In anatomy, a foramen is any opening. Foramina inside the body of humans and other animals typically allow muscles, nerves, arteries, veins, or other structures to connect one part of the body with another.

The human skull has numerous foramina through which cranial nerves, arteries, veins, and other structures pass. The skull bones that contain foramina include the frontal, ethmoid, sphenoid, maxilla, palatine, temporal, and occipital lobes.

Key foramina in the skull include:

- Supraorbital foramen: Located in the frontal bone, it allows passage of the supraorbital vein, artery, and nerve into the orbit.
- Optic foramen: Located in the sphenoid, it allows the passage of the ophthalmic artery and nerve from the optic canal into the orbit.

- Foramen magnum: Located in the occipital bone, it allows the passage of the spinal and vertebral arteries and the spinal cord to pass from the skull into the vertebral column.
- Foramina of cribriform plate: Located in the ethmoid bone, it allows the passage of the olfactory nerve.
- Foramen rotundum: Located in the sphenoid bone, it allows passage of the maxillary nerve.

This page titled 7.1E: Foramina is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1F: Fontanelles

A fontanelle is an anatomical feature on an infant's skull that allows its plates to be flexible to pass through the birth canal.

Learning Objectives


- Identify the evolutionary purpose of skull fontanelles

Key Points

- Fontanelles are soft spots on a baby's head that, during birth, enable the bony plates of the skull to flex and allow the child's head to pass through the birth canal.
- At birth, the skull features a small posterior fontanelle (an open area covered by a tough membrane) where the two parietal bones adjoin the occipital bone (at the lambda); it usually closes in the first two to three months of life through intramembranous ossification.
- The much larger, diamond-shaped anterior fontanelle—where the two frontal and two parietal bones join—generally remains open until the child is about two years of age.
- Two smaller fontanelles are located on each side of the head. The more anterior one is the sphenoidal (between the sphenoid, parietal, temporal, and frontal bones), while the more posterior one is the mastoid (between the temporal, occipital, and parietal bones).

Key Terms

- **fontanelle**: A fontanelle is a soft membraneous spot on the head of a baby due to incomplete fusion of the cranial bones.



Superior view of infant skull: This image shows the location of the anterior (frontal) and posterior fontanelles.


Fontanelles are soft spots on a baby's head that, during birth, enable the bony plates of the skull to flex and allow the child's head to pass through the birth canal. The ossification of the bones of the skull causes the fontanelles to close over a period of 18 to 24 months; they eventually form the sutures of the neurocranium.

The cranium of a newborn consists of five main bones: two frontal bones, two parietal bones, and one occipital bone. These are joined by fibrous sutures that allow movement that facilitates childbirth and brain growth.

At birth, the skull features a small posterior fontanelle (an open area covered by a tough membrane) where the two parietal bones adjoin the occipital bone (at the lambda). This fontanelle usually closes during the first two to three months of an infant's life. This is called intramembranous ossification. The mesenchymal connective tissue turns into bone tissue.

The much larger, diamond-shaped anterior fontanelle—where the two frontal and two parietal bones join—generally remains open until a child is about two years old. The anterior fontanelle is useful clinically, as examination of an infant includes palpating the anterior fontanelle.

Two smaller fontanelles are located on each side of the head. The more anterior one is the sphenoidal (between the sphenoid, parietal, temporal, and frontal bones), while the more posterior one is the mastoid (between the temporal, occipital, and parietal bones).



Lateral view of infant skull: This image show the location of the sphenoidal and mastoid fontanelles.

The fontanelle may pulsate. Although the precise cause of this is not known, it is perfectly normal and seems to echo the heartbeat, perhaps via the arterial pulse within the brain vasculature, or in the meninges. This pulsating action is how the soft spot got its name: fontanelle means little fountain.

Parents may worry that their infant may be more prone to injury at the fontanelles. In fact, although they may colloquially be called soft spots, the membrane covering the fontanelles is extremely tough and difficult to penetrate.

The fontanelles allow the infant brain to be imaged using ultrasonography. Once they are closed, most of the brain is inaccessible to ultrasound imaging because the bony skull presents an acoustic barrier.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human skull. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_skull**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human skull. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_skull**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- skull. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/skull**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/**

2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png. License: *Public Domain: No Known Copyright*
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png. License: *Public Domain: No Known Copyright*
- Neurocranium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neurocranium. License: *CC BY-SA: Attribution-ShareAlike*
- Human skull. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_skull. License: *CC BY-SA: Attribution-ShareAlike*
- neurocranium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurocranium. License: *CC BY-SA: Attribution-ShareAlike*
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png. License: *Public Domain: No Known Copyright*
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png. License: *Public Domain: No Known Copyright*
- Neurocranium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg. License: *Public Domain: No Known Copyright*
- Facial skeleton. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_skeleton. License: *CC BY-SA: Attribution-ShareAlike*
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png. License: *Public Domain: No Known Copyright*
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png. License: *Public Domain: No Known Copyright*
- Neurocranium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg. License: *Public Domain: No Known Copyright*
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Facial_skeleton_-_en.svg. License: *Public Domain: No Known Copyright*
- Orbit (anatomy). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Orbit_(anatomy). License: *CC BY-SA: Attribution-ShareAlike*
- orbit. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/orbit. License: *CC BY-SA: Attribution-ShareAlike*
- optic canal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/optic%20canal. License: *CC BY-SA: Attribution-ShareAlike*
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png. License: *Public Domain: No Known Copyright*
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png. License: *Public Domain: No Known Copyright*
- Neurocranium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg. License: *Public Domain: No Known Copyright*
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Facial_skeleton_-_en.svg. License: *Public Domain: No Known Copyright*
- Foramina of the skull. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Foramina_of_the_skull. License: *CC BY-SA: Attribution-ShareAlike*
- Foramen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Foramen. License: *CC BY-SA: Attribution-ShareAlike*
- Foramina of the skull. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Foramina_of_the_skull. License: *CC BY-SA: Attribution-ShareAlike*
- Foramen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Foramen. License: *CC BY-SA: Attribution-ShareAlike*
- foramina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/foramina. License: *CC BY-SA: Attribution-ShareAlike*
- foramen ovale. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/foramen_ovale. License: *CC BY-SA: Attribution-ShareAlike*
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png. License: *Public Domain: No Known Copyright*
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png. License: *Public Domain: No Known*

Copyright
- Neurocranium. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg**. **License**: ***Public Domain: No Known Copyright***
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Facial_skeleton_-_en.svg**. **License**: ***Public Domain: No Known Copyright***
- The location of foramina in the skull. . **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...81/Gray193.png**. **License**: ***Public Domain: No Known Copyright***
- Skull suture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skull_suture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skull suture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skull_suture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cranium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cranium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- suture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/suture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Neurocranium. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg**. **License**: ***Public Domain: No Known Copyright***
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Facial_skeleton_-_en.svg**. **License**: ***Public Domain: No Known Copyright***
- The location of foramina in the skull. . **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/81/Gray193.png**. **License**: ***Public Domain: No Known Copyright***
- Lateral view of a skull showing sutures.. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...uras_right.svg**. **License**: ***Public Domain: No Known Copyright***
- Paranasal sinuses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Paranasal_sinuses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- paranasal sinuses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/paranasal%20sinuses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Neurocranium. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg**. **License**: ***Public Domain: No Known Copyright***
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Facial_skeleton_-_en.svg**. **License**: ***Public Domain: No Known Copyright***
- The location of foramina in the skull. . **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/81/Gray193.png**. **License**: ***Public Domain: No Known Copyright***
- Lateral view of a skull showing sutures.. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3e/Human_skull_side_suturas_right.svg**. **License**: ***Public Domain: No Known Copyright***
- Paranasal sinuses. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi.../Nnh_front.svg**. **License**: ***Public Domain: No Known Copyright***
- Fontanelle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fontanelle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fontanelle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fontanelle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Components of Viscerocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/2/25/Facial_skeleton_-_en.svg/2000px-Facial_skeleton_-_en.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Components of the Neurocranium. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/thumb/3/39/Cranial_bones_en_v2.svg/2000px-Cranial_bones_en_v2.svg.png**. **License**: ***Public Domain: No Known Copyright***
- Neurocranium. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/8b/Cranial_bones_en.svg**. **License**: ***Public Domain: No Known Copyright***
- facilabones.png. **Provided by**: Wikimedia Commons. **Located at**: **https://commons.wikimedia.org/wiki/**

File:Facial_skeleton_-_en.svg. **License**: *Public Domain: No Known Copyright*
- The location of foramina in the skull. . **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/8/81/Gray193.png. **License**: *Public Domain: No Known Copyright*
- Lateral view of a skull showing sutures.. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/3e/Human_skull_side_suturas_right.svg. **License**: *Public Domain: No Known Copyright*
- Paranasal sinuses. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fb/Nnh_front.svg. **License**: *Public Domain: No Known Copyright*
- Top view of fontanelles. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...b3/Gray197.png. **License**: *Public Domain: No Known Copyright*
- Infant skull fontanelles. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...a0/Gray198.png. **License**: *Public Domain: No Known Copyright*

This page titled 7.1F: Fontanelles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1F: Sutures

A suture is a type of fibrous joint (or synarthrosis) that only occurs in the skull (or cranium).

Learning Objectives

- List the sutures of the skull

Key Points

- It is normal for many of the bones of the skull to be unfused at birth. This allows a tiny amount of movement at the sutures, which contributes to the compliance and elasticity of the skull.
- Sutures become fused as individuals age; thus, examining sutures can provide an estimate of age postmortem.
- There are 17 named sutures on the human skull.


Key Terms

- **suture**: A fairly rigid joint between two or more hard elements, such as the bony plates of the skull.

A suture is a type of fibrous joint (or synarthrosis) that only occurs in the skull. The bones are bound together by Sharpey's fibers, a matrix of connective tissue which provide a firm joint.

A small amount of movement is permitted through these sutures that contributes to the compliance and elasticity of the skull. The joint between the mandible and the cranium, known as the temporomandibular joint, forms the only non-sutured joint in the skull. Most sutures are named for the bones that they articulate.

At birth, many of the bones of the skull remain unfused to the soft spots described as fontanelle. The bones fuse relatively rapidly through a process known as craniosynotosis, although the relative positions of the bones can continue to change through life. In old age the cranial sutures may ossify completely, reducing the amount of elasticity present in the skull. As such, the degree of ossification can be a useful tool in determining age postmortem.



Lateral view of a skull showing sutures: The dotted red lines indicate the location of skull sutures.

Sutures primarily visible from the side include:

- Coronal suture: Located between the frontal and parietal bones.
- Lambdoid suture: Located between the parietal, temporal and occipital bones.
- Occipitomastoid suture.
- Parietomastoid suture.
- Sphenofrontal suture.
- Sphenoparietal suture.
- Sphenosquamosal suture.
- Sphenozygomatic suture.
- Squamosal suture: Located between the parietal and the temporal bone.
- Zygomaticotemporal suture.
- Zygomaticofrontal suture.

Sutures primarily visible from front or above include:

- Frontal suture or metopic suture: Located between the two frontal bones, prior to the fusion of the two into a single bone.
- Sagittal suture: Located along the midline, between the parietal bones.

Sutures primarily visible from below or inside include:

- Frontoethmoidal suture.
- Petrosquamous suture.
- Sphenoethmoidal suture.
- Sphenopetrosal suture.

This page titled 7.1F: Sutures is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.1G: Paranasal Sinuses

The paranasal sinuses (four, paired, air-filled spaces) surround the nasal cavity, and are located above and between the eyes, and behind the ethmoids.

Learning Objectives

- Describe the structure and function of the paranasal sinuses


Key Points

- The sinuses are named for the facial bones that they are located behind.
- There are four sinuses; the maxillary, frontal, ethmoid, and sphenoid.
- Paranasal sinuses form developmentally through excavation of bone by air-filled sacs from the nasal cavity.
- This process begins prenatally and continues through the course of an individual's lifetime.
- The biological role of the sinuses is debated, but a number of possible functions have been proposed: decreasing weight of the face; increasing resonance of voice; buffer against blows; insulating sensitive structures from rapid temperature fluctuations; and humidifying and heating of inhaled air.

Key Terms

- **paranasal sinuses**: A group of four, paired, air-filled spaces that surround the nasal cavity (maxillary sinuses), above the eyes (frontal sinuses), between the eyes (ethmoid sinuses), and behind the eyes (sphenoid sinuses).

Structure of the Paranasal Sinuses



Skull Sinuses: This image shows the position of the sinuses in the human skull.

Paranasal sinuses are a group of four, paired, air-filled spaces that surround the nasal cavity (maxillary sinuses), above the eyes (frontal sinuses), between the eyes (ethmoid sinuses), and behind the eyes (sphenoid sinuses). The sinuses are named for the facial bones that they are located behind.

- The maxillary sinuses (also called the maxillary antrechea, the largest of the paranasal sinuses) are located under the orbits in the maxillary bones.

- The frontal sinuses are superior to the orbits and are in the frontal bone.
- The ethmoid sinuses are formed from several discrete air cells within the ethmoid bone between the nose and the orbits.
- The sphenoid sinuses are in the sphenoid bone at the center of the skull base under the pituitary gland.
- The paranasal sinuses are lined with respiratory epithelium.

The paranasal sinuses form developmentally through excavation of bone by air-filled sacs (pneumatic diverticula) from the nasal cavity. This process begins prenatally and continues through the course of an individual's lifetime.

Function of the Paranasal Sinuses

The biological role of the sinuses is debated, but a number of possible functions have been proposed. These include:

- Decreasing the relative weight of the front of the skull, and especially the bones of the face.
- Increasing resonance of the voice.
- Providing a buffer against blows to the face.
- Insulating sensitive structures like dental roots and eyes from rapid temperature fluctuations in the nasal cavity.
- Humidifying and heating of inhaled air because of slow air turnover in this region.
- Regulation of intranasal and serum gas pressures.
- Immunological defense.

This page titled 7.1G: Paranasal Sinuses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.2: THE HYOID BONE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

7.2A: The Hyoid Bone

This page titled 7.2: The Hyoid Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.2A: The Hyoid Bone

The hyoid, a horseshoe-shaped bone situated in the anterior of the neck, between chin and thyroid, aids tongue movement and swallowing.

Learning Objectives

- Describe the hyoid bone


Key Points

- The hyoid bone is a horseshoe shaped bone found in the neck.
- It functions to protect the essophagus and facilitates a wide range of movements involved in speaking and swallowing.
- The hyoid bone is only distantly articulated to other bones by muscles or ligaments.
- The hyoid consists of a central part called the body, and two pairs of cornua: the greater cornu and the lesser cornu.
- Ossification begins in the greater cornua toward the end of fetal life, in the hyoid body shortly afterward, and in the lesser cornua during the first or second year after birth.


Key Terms

- **hyoid bone**: A bone in the neck which protects the esophagus and facillitates a wide range of muscle movements required for speaking and swallowing.

The hyoid bone is a horseshoe shaped bone found in the neck. Located anteriorly between the mandible and the thyroid cartilage, the hyoid bone protects the esophagus and also facilitates the wide range of muscle activity required for speaking and swallowing. It is visible upon extension of the neck.



Position of the hyoid bone in the neck: This image shows where the hyoid bone is located.



Hyoid Bone: Anterior view of hyoid bone with sections labeled.

The hyoid bone consists of a central body and two pairs of cornua, or horns, termed greater and lesser cornua. The greater horns project backwards from the body and provide a platform for key muscles and ligaments to attach to including the stylohyoid and throhyoid muscles.

The lesser horns are two smaller eminences which project superiorly to the join between the greater cornua and body and are attached to the body by fibrous tissue. As with the greater cornua the lesser cornua provide a platform for muscle and ligament attachment specifically for the stylohyoid ligament.

The hyoid ossifies towards the end of fetal development, commencing in the greater cornua before completing in the body shortly after birth. The greater cornua and body are initially connected by fibrous material although this ossifies towards middle age.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Hyoid bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hyoid_bone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pharynx. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pharynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- larynx. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/larynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hyoid bone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hyoid_bone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...1f/Gray186.png**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...4/Gray1194.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 7.2A: The Hyoid Bone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.3: THE VERTEBRAL COLUMN

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

7.3A: General Characteristics of the Spine

7.3B: Parts of a Vertebra

7.3C: Regional Vertebral Characteristics

This page titled 7.3: The Vertebral Column is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.3A: General Characteristics of the Spine

The spine is made of vertebrae that link together to protect the spinal cord.

Learning Objectives


- Describe the structure of the vertebral column

Key Points

- The main functions of the vertebral bones are for structure (posture) and protection of the spinal cord.
- The spine is split into five regions: the cervical, thoracic, lumbar, sacral, and coccyx.
- The vertebrae of the sacrum and coccyx are fused, but those of the cervical, thoracic, and lumber regions are free to articulate.
- Viewed laterally, the vertebral column presents several curves that correspond to the column's different regions—the cervical, thoracic, lumbar, and sacral.

Key Terms

- **vertebral column**: The series of vertebrae that protect the spinal cord; the spinal column.
- **vertebrae**: The bones that make up the spinal column.



Human vertebral column: The vertebral column has 33 bones. Each color represents a section of the column.

The vertebral column (also known as the backbone or spine), is a tall, thin organ located dorsally that extends from the base of the spine to the pelvis. It protects the spinal cord and provides a key attachment point for numerous muscle groups.

There are 33 vertebrae in the human spine that are split into four regions that correspond to the curvature of the spine; the cervical, thoracic, lumbar, sacrum, and coccyx. The vertebrae of the sacrum and coccyx are fused, but those of the cervical, thoracic and lumbar regions are separated by intervertebral discs.

Vertebrae are given an alphanumeric descriptor, with the initial letter derived from the region they are located in followed by a digit; the digit increases moving down the region. For example, the most superior cervical vertebra is termed C1 and the most

inferior C7, which is then followed by the T1 vertebrae of the thoracic region.

Viewed laterally the vertebral column presents several curves that correspond to the different regions of the column. These are called the cervical, thoracic, lumbar, and pelvic regions.

- The cervical curve covers the region between vertebrae C1 and T2, it is the least marked of all the spinal curves.
- The thoracic curve covers the region between vertebrae T2 and T12.
- The lumbar curve covers the region between vertebrae T12 and L5 and is more marked in the females than in males due to differences in pelvic structure.
- The sacral curve begins at the sacrovertebral articulation, and ends at the point of the coccyx.

The thoracic and sacral curves are termed primary curves because they alone are present during fetal life. The cervical and lumbar curves are secondary curves that are developed after birth; the former when the child is able to maintain an upright posture, the latter when the child begins to walk.

This page titled 7.3A: General Characteristics of the Spine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.3B: Parts of a Vertebra

A vertebra consists of two parts: an anterior segment, or the vertebral body; and a posterior part, or the vertebral (neural) arch.

Learning Objectives

- Identify the parts of a vertebra

Key Points

- The vertebral processes can give the structure rigidity, help it articulate with ribs, or serve as muscle attachment points.
- When the vertebrae are articulated with each other, their bodies form a strong pillar for the support of the head and trunk, and the vertebral foramina constitute a canal for the protection of the spinal cord.
- Two transverse processes and one spinous process are posterior to the vertebral body.
- Two superior and two inferior articular processes articulate with the adjoining vertebrae. They allow for a small degree of movement in the spine, but greatly strengthen it.In between every pair of vertebrae are two apertures, the intervertebral foramina, one on either side, for the transmission of the spinal nerves and vessels.

Key Terms

- **process**: An outgrowth of tissue or cell.
- **vertebral arch**: The posterior part of a vertebra through which the spinal cord passes.

When articulated together the vertebrae form a strong yet flexible structure that encloses the vertebral foramen, or opening, where the spinal cord sits. It also provides a base for numerous muscle attachments and articulations with other bones.

The strength and flexibility of this structure is generated by the structure of the individual vertebrae. Comprised of bone and cartilage, the configuration of a vertebra varies based on its location within the body, although there are common features associated with those of the upper region.


A typical vertebra of the upper region of the spine consists of two regions:

1. The anterior vertebral body which is the point of articulation between the vertebrae.
2. The posterior vertebral or neural arch that encloses the spinal cord.

Located between each pair of vertebrae are two laterally located openings:

1. The intervertebral foramina that facilitate access to the spinal cord for nerves and vessels.
2. The intervertebral discs that act as ligaments between the vertebral bodies.

The vertebral arch is formed from two, short, thick processes called pediments that extend posteriorly from the lateral sides of the vertebral body, before joining together at the midline with the laminae.



Oblique view of cervical vertebra: The parts of a human vertebra.

Vertebral Processes

There are seven processes that project from a typical vertebra.

Four articular processes originate from the joint between the pedicles and laminae, two point superiorly and two point inferiorly. They interact with the zyhapophysis, a socket for the articular processes, of the adjacent vertebrae to make the spine more stable and to facilitate a small degree of articulation.

A single spinous process projects backwards and downwards from the center of the vertebral arch and it serves as a major attachment point for muscles and ligaments of the back.

The two transverse processes project laterally from the join between the pedicle and lamina and also serve as an attachment point for muscles and ligaments of the back. The transverse processes articulate with the ribs in conjunction with the vertebral body.

This page titled 7.3B: Parts of a Vertebra is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.3C: Regional Vertebral Characteristics

The vertebrae of the spinal column are divided into five regions: cervical, thoracic, lumbar, sacral, and coccyx.

Learning Objectives

- Identify the different types of vertebrae

Key Points


- The upper three regions of the vertebral column are articulating vertebrae grouped under the names cervical, thoracic and lumbar, according to the regions they occupy.
- Vertebrae are given an alphanumeric descriptor, with the initial letter derived from the region they are located in followed by a digit that increases moving down the region. For example, the most superior cervical vertebrae is termed C1 and the most inferior C7, which is then followed by the T1 vertebrae of the thoracic region.
- The two lower most regions of vertebrae are the sacrum and the coccyx.

The vertebrae comprising the spinal column can be divided into five regions, based on the five varying curvatures of the spine. The upper three regions of the spinal column are termed the cervical, thoracic, and lumbar; they contain individually jointed vertebrae. The two lower regions—the sacrum and coccyx, or tailbone—are formed from fused vertebrae.

Upper Vertebrae


Vertebrae are given an alphanumeric descriptor, with the initial letter derived from the region they are located in followed by a digit that increases moving down the region. For example, the most superior cervical vertebrae is termed C1 and the most inferior C7, which is then followed by the T1 vertebrae of the thoracic region.

The cervical region of the spine is the most superior and contains seven small vertebrae. The main function of the cervical region is to facilitate attachment of the skull to the spine, protect the spinal cord over the exposed neck and shoulder region, and support the body.




Cervical vertebra, lateral view: The lateral view of a typical cervical vertebra.

The twelve thoracic vertebrae are located inferiorly to the cervical region. They are larger than the cervical vertebrae and increase in size moving inferiorly to the lumbar region.



Thoracic vertebra: Image of a typical thoracic vertebra.

The five lumbar vertebrae are the largest vertebral bones and increase in size when moving inferiorly. The lumbar vertebrae play a key role in supporting the body and facilitating locomotion.



Lumbar vertebra: Image of typical lumbar vertebra.


Lower Vertebrae

During childhood the five vertebrae of the sacral region are distinct. In adulthood the five bones fuse to form the sacrum, although it is still often divided into regions termed S1–S5 based on the formation of the original individual bones. The sacrum functions to support the body and protect organs of the pelvis and lower back.

The final region of the spine is the coccyx, or tailbone. As with the sacrum, the coccyx is formed from several vertebrae that have fused together.

As it's alternative name suggests, the coccyx forms the basis of a tail that has been lost in humans, although it is incorrect to think of it as a vestigial structure since is a key attachment point for many muscles and ligaments and plays a key role in supporting

the body while sitting.



Coccyx: Lateral view of coccyx shown beneath the sacrum.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- vertebral column. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vertebrae. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vertebrae**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray 111 - Vertebral column-coloured. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:G...n-coloured.png**. **License**: ***Public Domain: No Known Copyright***
- Human vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- process. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/process**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vertebra. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vertebra**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vertebral arch. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vertebral_arch**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray 111 - Vertebral column-coloured. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray_111_-_Vertebral_column-coloured.png**. **License**: ***Public Domain: No Known Copyright***
- ACDF oblique annotated english. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ACDF_oblique_annotated_english.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Vertebral column. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vertebral_column**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vertebra. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vertebra**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray 111 - Vertebral column-coloured. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray_111_-_Vertebral_column-coloured.png**. **License**: ***Public Domain: No Known Copyright***
- ACDF oblique annotated english. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ACDF_oblique_annotated_english.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical vertebra. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/5/52/Gray85.png**. **License**: ***Public Domain: No Known Copyright***
- Coccyx . **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/5/5a/Gray97.png**. **License**: ***Public Domain: No Known Copyright***
- Lumbar vertebra. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/62/Gray94.png**. **License**: ***Public Domain: No Known Copyright***
- Thoracic vertebra. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/30/Gray82.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 7.3C: Regional Vertebral Characteristics is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.4: THE THORAX



- Topic hierarchy

7.4A: Thoracic Cage: Sternum

7.4B: Thoracic Cage: Ribs

This page titled 7.4: The Thorax is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


7.4A: Thoracic Cage: Sternum

Learning Objectives

- Describe the structure and function of the sternum

The sternum, or breastbone, is a long, flat, bony plate that forms the most anterior section of the ribcage. The sternum is highly vascular in nature and covered with a thin layer of compact bone providing a degree of flexibility.

Function of the Sternum



Sternum: This image shows the parts of the sternum.

Together with the anterior ribs, the sternum helps to protect the heart and lungs from damage, as well as facilitates the expansion and contraction of the thoracic cavity during respiration. During development the sternum is comprised of four individual sections called sternebrae, which fuse to form the sternum in adulthood.

Structure of the Sternum

The sternum is divided into three regions:

1. The manubrium.
2. The gladiolus.
3. The xiphoid process.

The manubrium is the most superior region of the sternum and articulates with the clavicles or collarbones and the first pair of ribs. The manubrium is the thickest portion of the sternum as it carries the greatest physical load.

Located below the manubrium, the gladiolus is the longest portion of the sternum and articulates with the ribs, either directly or indirectly, through the costal cartilage

The thin, pointed xiphoid process forms the most inferior region of the sternum to which the costal cartilage and cartilage of the celiac, or solar, plexus attaches.

Key Points

- The sternum, or breastbone, is a long, flat, bony plate that forms the most anterior section of the rib cage.
- The primary function of the sternum is the protection of the heart, lungs, and blood vessels from physical damage.
- The sternum is made of the manubrium, the gladiolus, and the xiphoid process.

Key Terms

- **xiphoid process**: The most inferior region of the thorax.
- **gladiolus**: The body of the thorax.
- **manubrium**: The most superior region of the thorax.

This page titled 7.4A: Thoracic Cage: Sternum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


7.4B: Thoracic Cage: Ribs

Learning Objectives

- Describe the ribs of the thoracic cage

Ribs are long, curved bones that form the rib cage surrounding the thorax. The thoracic cage can expand and contract to facilitate breathing in association with the diaphragm; it also protects the lungs, heart, and other organs of the thoracic cavity.

Structure of a Rib



Ribs: Lateral view of rib cage showing its connections to vertebrae. The first thoracic vertebra has a rib attached to it that curves across to the sternum.

A rib consists of a head, neck, and shaft. The head of the rib is the most posterior region of the rib and articulates with the vertebral column. The flattened neck region provides an attachment point for numerous muscles located within the back. Finally, the shaft forms the majority of the length of the rib as it curves around the thoracic cavity forming the rib cage.

Organization of the Ribs

Humans have 24 ribs split into pairs that are named numerically passing inferiorly from the neck. The first five ribs are termed true ribs because they attach directly to the sternum through the costal cartilage.

The next five ribs are termed false ribs because they attach to the sternum indirectly through the costal cartilage.

Finally, the last two ribs are termed floating ribs because they do not attach to the sternum at all.

Rib 1 is the shortest and most curved of the ribs. Descending from rib 1, the ribs increase in size up until rib 7 before decreasing in length again.

Key Points

- Ribs are long, curved bones that form the rib cage surrounding the thorax.
- Articulating from the spine, the majority attach to the sternum either directly or indirectly.
- Humans have 24 ribs split into pairs. Pairs 1–5 attach directly to the sternum through the costal cartilage, 6–10 attach indirectly though the costal cartilage, and 11–12 are termed floating ribs and do not attach.

Key Terms

- **rib**: A long, curved bone that forms the rib cage surrounding the thorax.
- **costal**: Pertaining to a rib.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human sternum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_sternum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- comminuted. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/comminuted**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sternum. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/23/Gray116.png**. **License**: ***Public Domain: No Known Copyright***
- Rib. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rib**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tetrapod. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tetrapod**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- costal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/costal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sternum. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/23/Gray116.png**. **License**: ***Public Domain: No Known Copyright***
- Rib. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/4/4b/Gray114.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 7.4B: Thoracic Cage: Ribs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.5: THE SHOULDER

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

7.5A: Clavicle

7.5B: Scapula

This page titled 7.5: The Shoulder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.5A: Clavicle

The clavicle or collar bone is a long, curved bone on the upper portion of the shoulder that connects with the scapula and the sternum.

Learning Objectives

- Describe the characteristics of the clavicle


Key Points

- The clavicle is a long, doubly curved bone that connects the arm to the body, located directly above the first rib. It acts as a strut to keep the scapula in place so the arm can hang freely.
- The clavicle is an attachment point for several muscles.
- Structurally, the clavicle can be divided into three parts: medial end, lateral end, and shaft.
- There are sex differences in clavicle shape—female clavicles are shorter and thinner than male clavicles.

Key Terms

- **acromion**: The outermost point of the shoulder blade.

The clavicle, or collarbone, is a slender s-shaped bone that extends between the sternum and the scapula and is located directly above the first rib. It functions to attach the upper arm to the trunk and provides support to allow free movement around the shoulder.



Left Clavicle: The left clavicle, viewed from above. Muscle attachment sites (pectoralis major, subclavius muscle, deltoid, and sterno-hyoid) are highlighted.

Medially the clavicle is quadrangular in shape and articulates with the manubrium of the sternum forming the sternoclavicular joint. Laterally, the clavicle is flattened and attaches to the acromion process of the scapula forming the acromioclavicular joint.

The shaft of the clavicle acts as the origin and attachment point for numerous muscles and ligaments. At the medial end of the shaft the pectoralis major originates from the anterior surface, the posterior surface gives origin to the sternohyoid muscle and the superior surface the sternocleidomastoid muscle.

The costoclavicular ligament attaches to the inferio≠r surface. Laterally the deltoid muscle originates from the anterior surface and the trapezius muscle attaches to the posterior surface at the trapezoid line. Adjacent to this is the conoid tubercle which is an attachment point for the conoid ligament.

The clavicle in males is typically thicker and longer than a female's clavicle to account for the larger muscle mass operating through it.

This page titled 7.5A: Clavicle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.5B: Scapula

The scapula, or shoulder bone, is a flat, triangular bone that connects to the humerus and the clavicle.

Learning Objectives

- Describe the shape and function of the scapula in humans

Key Points


- The scapula articulates with the humerus and the clavicle.
- The scapula is flat and triangular.
- The scapula articulates with the humerus at the glenoid fossa and the clavicle at the acromion process.
- The scapula provides attachment sites for many muscles including the pectoralis minor, coracobrachialis, serratus anterior, triceps brachii, biceps brachii, and the subscapularis.
- The scapula has two main surfaces: the costal (front facing) surface and the dorsal (rear facing) surface.

Key Terms

- **acromion**: The outermost point of the shoulder blade.
- **glenoid**: A shallow depression in a bone, especially in the scapula.


The scapula, or shoulder blade, is a flat, triangular bone located to the posterior of the shoulder. The scapula articulates with the clavicle through the acromion process, a large projection located superiorly on the scapula forming the acromioclavicular joint. The scapula also articulates with the humerus of the upper arm to form the shoulder joint, or glenohumeral joint, at the glenoid cavity.

Due to its flat nature, the scapula presents two surfaces and three borders; the front-facing costal surface and the rear-facing dorsal surface, as well as the superior, lateral, and medial borders.



Costal surface: Costal surface of the left scapula. The subscapular fossia for subscapularis, serratus, pector minor regions are highlighted.

The serratus anterior originates from the costal surface, which also provides an attachment for the subscapularis muscle. The dorsal surface gives origin to the supraspinatus and infraspinatus muscles, and inferiorly to the teres minor and major. It is divided by a ridge-like structure called the spine of the scapula, from which the deltoid and trapezius muscles originate.



Dorsal surface: Dorsal surface of the left scapula.

The lateral border is the thickest border of the scapula and extends downwards from the glenoid cavity. Immediately below the glenoid cavity is the infraglenoid tuberosity, which is the origin for the long head of the triceps brachii.

Immediately above the glenoid cavity is the supraglenoid tubercle and its associated hook-like coracoid process, from which the long and short heads of the biceps brachii originate.

Four muscles attach to the medial border of the scapula. To the anterior side the serratis anterior attaches, whilst posteriorly the levator scapulae and rhomboids minor and major attach.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Clavicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Clavicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Clavicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Clavicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- acromion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acromion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- manubrium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/manubrium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray201. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray201.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scapula. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scapula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scapula. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scapula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glenoid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glenoid**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- fossa. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fossa**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- acromion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acromion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray201. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray201.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray203. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray203.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray202. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray202.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 7.5B: Scapula is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.6: THE UPPER LIMB

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

7.6A: Humerus (The Upper Arm)

7.6B: Ulna and Radius (The Forearm)

7.6C: Carpals, Metacarpals, and Phalanges (The Hand)

This page titled 7.6: The Upper Limb is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.6A: Humerus (The Upper Arm)

The bone forming the upper arm is the humerus.

Learning Objectives

- Describe the anatomy of the humerus


Key Points

- The humerus articulates with the scapula at the shoulder and with the radius and ulna at the elbow.
- The humerus is typically described as having a proximal region, shaft, and distal region.
- Both the anatomical and surgical neck of the humerus are highly susceptible to fracture.

Key Terms

- **humerus**: The bone of the upper arm.
- **surgical neck**: A constriction below the tubercles of the greater tubercle and lesser tubercle.
- **capitulum**: At the distal head of the humerus, it articulates with the radius of the forearm.
- **trochlea**: At the distal head of the humerus, it articulates with the ulna of the forearm.
- **anatomical neck**: A constriction adjacent to the humeral head that increases the range of movement possible at the shoulder joint.

The humerus is a bone in the upper arm. It runs from the shoulder to the elbow. Proximally it articulates with the scapula to form the shoulder joint, or glenohumeral joint. Distally, the humerus articulates with the radius and ulna to form the elbow joint.



Humerus: The humerus attaches proximally to the scapula (shoulderblade) at the humeral head and distally with the radius and ulna (lower-arm bones) at the trochlea and capitulum, respectively.

The proximal portion of the humerus can be divided into three parts.

1. The rounded humeral head projects medially and articulates with the glenoid cavity of the scapula.
2. Immediately adjacent to the head is the narrower anatomical neck, which allows for a wider range of movements of the head

within the shoulder joint.
3. Finally, the greater and lesser tubercles are found at the most superior end of the main shaft of the humerus.

The four rotator cuff muscles attach to these tubercles, strengthening and maintaining the shoulder joint. Between the two tubercles lies a deep grove called the intertubercular sulcus, through which the tendon of the long head of the biceps brachii runs.

Below this proximal region lies the shaft, which is separated from the proximal region by the surgical neck, so termed as this in an area of frequent fracture. A major feature of the shaft is the deltoid tuberosity located laterally to which the deltoid muscle attaches.

The deltoid, corocobrachialis, brachialis, and brachioradialis muscles attach to the anterior surface, with the triceps brachii attaching to the posterior.

Distally, the humerus flattens to articulate with the ulna and radius at the elbow joint. The medially located trochlea articulates with the ulna. Located laterally to this is the capitulum that articulates with the radius.

Several muscles of the forearm responsible for extension at the wrist attach to the humerus immediately above the capitulum and trochlea.

This page titled 7.6A: Humerus (The Upper Arm) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.6B: Ulna and Radius (The Forearm)

The forearm contains two bones, the radius and the ulna.

Learning Objectives


- Discuss the radius and ulna

Key Points

- The radius and the ulna are long, slightly curved bones that lie parallel from the elbow, where they articulate with the humerus, to the wrist, where they articulate with the carpals.
- The radius is located laterally, near the thumb, and the ulna medially, near the little finger. The radius and the ulna have a styloid process at the distal end; they are also attachment sites for many muscles.
- The radius is smaller than the ulna.

Key Terms

- **radius**: One of two forearm bones, it is located laterally to the ulna.
- **ulna**: One of two forearm bones, it is located medially to the radius.



The forearm: The radius and ulna are the bones of the forearm.

The forearm contains two bones—the radius and the ulna—that extend in parallel from the elbow, where they articulate with the humerus to the wrist, where they articulate with the carpals. The space between the two bones is spanned by the interosseous membrane.

The Ulna

Anatomically, the ulna is located medially to the radius, placing it near the little finger. The ulna is slightly larger than the radius.

Proximally, there are five key regions of the ulna:

1. The olecranon is a projection of bone that extends proximally from the ulna.
2. The triceps brachii muscle attaches to the ulna superiorly.
3. The cornoid process, together with the olecranon, forms the trochlear notch where it articulates with the trochlea of the humerus.
4. Laterally to the trochlear notch lies the radial notch, which articulates with the head of the radius to form the proximal radioulnar joint.
5. Immediately distal to the coronoid process is the tuberosity of ulna, to which the brachialis muscle attaches.

The shaft of the ulna is triangular and numerus muscles involved in pronation and flexion of the forearm attach to its surface.

Distally, the ulna is much smaller and terminates with a rounded head that articulates with the ulnar notch of the radius to form the distal radioulnar joint. The styloid process of the ulna extends distally and is the site of attachment for ligaments found in the wrist.

The Radius

Anatomically, the radius is located laterally to the ulna placing it near the thumb. The radius is slightly smaller than the ulna and pivots around the ulna to produce movement at the proximal and distal radioulnar joints.

Proximally, the radius terminates with a disk-shaped head that articulates with the capitulum of the humerus and the radial notch of the ulna. Immediately below the head lies the radial tuberosity to which the biceps brachii attaches. As with the ulna, the shaft of the radius is triangular in shape and numerous muscles, including the protonator teres, attach to it.

Distally the radius expands, medially the ulnar notch articulates with the head of the ulnar. Immediately adjacent to the ulnar notch, the radius articulates with the scaphoid and lunate carpal bones to form part of the wrist.

This page titled 7.6B: Ulna and Radius (The Forearm) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.6C: Carpals, Metacarpals, and Phalanges (The Hand)

Each hand consists of 27 bones, divided between the wrist bones (carpals), the palm bones (metacarpals), and the finger bones (phalanges).

Learning Objectives

- Describe the types of bones in the hand

Key Points

- There are eight carpal bones in each wrist.
- There are five metacarpal bones in each hand.
- There are proximal, intermediate, and distal phalanges in each digit except for the thumb, which lacks an intermediate phalange.

Key Terms

- **metacarpal**: Any of the bones of the palm.
- **carpal**: Any of the eight bones of the wrist.
- **phalange**: One of the bones of the digits.



Carpals of the left hand: There are eight carpal bones in each wrist: scaphoid, lunate, triquetral, pisiform, trapezium, trapezoid, capitate, and hamate.


The hand contains 27 bones. Each one belongs to one of three regions: the carpals, (wrist), the metacarpals, (the palm), and the phalanges (the digits).

Carpals

The eight, irregularly shaped carpals are the most proximal bones of the hand. The carpals are often split into two rows, the proximal row containing the scaphoid, lunate, triquetrum, and pisiform, moving lateral to medial.

The scaphoid and lunate articulate with the radius, and the lunate and triquetrum articulate with the articular disk of the wrist. The pisiform carpal is a sesamoid bone, located within a tendon and is not involved in movement at the wrist.

The distal row contains the trapezium, trapezoid, capitate, and hamate, moving lateral to medial. The trapezium articulates with the scaphoid proximally and the first, thumb, and second metacarpal distally. The trapezoid articulates with the scaphoid proximally and the second metacarpal distally.



Metacarpal bones of the left hand: The metacarpals connect the carpal bones of the wrist with the phalanges (finger bones).

The capitate articulates with the scaphoid and lunate proximally and the third and fourth metacarpal. Finally, the hamate articulates with the lunate and triquetral proximally and the fourth and fifth, little finger, metacarpals distally.


Metacarpals

The hand contains five metacarpal bones that articulate proximally with the carpals and distally with the proximal phalanges. They are numbered moving lateral to medial, and start with the thumb, which is metacarpal I, and end with metacarpal V, the little finger.

Each metacarpal consists of a base, shaft, and head, with the concave lateral and medial borders of the shaft allowing attachment of the interossei muscles.

Phalanges

The digits are named in a similar fashion to the metacarpals, moving lateral to medial, and starting at the thumb. With the exception of the thumb, each digit contains a proximal, intermediate, and distal phalange; the thumb lacks an intermediate phalange. The length of the phalanges decreases distally.



Human hand bones: Fingers are made up of proximal, intermediate, and distal phalanges. The thumb lacks an intermediate phalange.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Humerus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Humerus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- humerus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/humerus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- surgical neck. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/surgical%20neck**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- HumerusFront.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...merusFront.png**. **License**: ***Public Domain: No Known Copyright***
- Radius (bone). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Radius_(bone)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ulna. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ulna**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- forearm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/forearm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- radius. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/radius**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ulna. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ulna**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- HumerusFront.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...merusFront.png**. **License**: ***Public Domain: No Known Copyright***
- Gray214.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray214.png**. **License**: ***Public Domain: No Known Copyright***
- phalange. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/phalange**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carpal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/carpal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- metacarpal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/metacarpal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- HumerusFront.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...merusFront.png**. **License**: ***Public Domain: No Known Copyright***
- Gray214.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray214.png**. **License**: ***Public***

Domain: No Known Copyright

- Scheme_human_hand_bones-en.svg. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...d_bones-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Metacarpal_bones_(left_hand)_01_palmar_view_with_label.png. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...with_label.png**. **License**: ***Public Domain: No Known Copyright***
- Carpus_(left_hand)_-_animation01a.gif. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...imation01a.gif**. **License**: ***Public Domain: No Known Copyright***

This page titled 7.6C: Carpals, Metacarpals, and Phalanges (The Hand) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7: THE HIP

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

7.7A: Ilium

7.7B: Ischium

7.7C: Pubis

7.7D: False and True Pelves

7.7E: Comparison of Female and Male Pelves

7.7F: Pelvic Structure and Childbearing

This page titled 7.7: The Hip is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7A: Ilium

The ilium is the uppermost and largest bone of the pelvis.

Learning Objectives

- Describe the ilium

Key Points


- The ilium articulates with the ischium, sacrum, and pubis.
- The ilium is divisible into the body and the ala, or wing.
- The body and ala are separated by the arcuate line on the posterior surface and the margin of the acetabulum on the inferior surface.
- The iliac crest is on the upper margin of the ala.
- The external surface of the ilium is partly articular (relating to a joint) and partly non-articular.
- Bi-iliac width is an anatomical term referring to the widest measure of the pelvis between the outer edges of the upper iliac bones.

Key Terms

- **acetabulum**: A concave structure formed from three bones in the pelvis that articulates with the femoral head to form the hip joint.
- **ilium**: The upper and widest of the three bones that make up each side of the hipbone and pelvis.
- **pelvis**: The large compound bone structure at the base of the spine that supports the legs. It consists of hip bone, sacrum, and coccyx.

The ilium is the uppermost and largest bone of the pelvis. It articulates with the sacrum, ischium, and pubis.

The human ilium is divisible into the body and the ala, or wing. These two parts are separated on the superior surface by a curved line known as the arcuate line, and on the inferior surface by the margin of the acetabulum.



Ilium: The ilium is the uppermost bone of the pelvis and extends laterally.

The body contributes to the formation of the acetabulum, a concave structure where the head of the femur articulates to form the hip joint. The internal surface of the body forms part of the wall of the lesser pelvis and gives origin to some fibers of the obturator internus muscle.

The Ala

The ala or wing, is the large expanded portion that extends laterally. It has an external and internal surface, a crest (illiac crest), and an anterior and posterior border.

The external surface is smooth and divided by three lines: the posterior, anterior and inferior gluteal lines. The gluteus maximus originates from the external surface adjacent to the posterior gluteal line, the gluteus medius from adjacent to the anterior line, and the gluteus minimus from the inferior line.

The internal surface of the ala is smooth and concave and forms the iliac fossa. The sacrospinalis and multifidus muscles of the back originate in the posterior.

In humans, bi-iliac width is an anatomical term referring to the widest measure of the pelvis between the outer edges of the upper iliac bones. The bi-iliac width measure is helpful in obstetrics because a pelvis that is significantly too small or too large can have obstetrical complications. It is also used by anthropologists to estimate body mass.

This page titled 7.7A: Ilium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7B: Ischium

The ischium forms the lower and posterior portion of the hip bones of the pelvis.

Learning Objectives

- Describe the ischium


Key Points

- The ischium articulates with the ilium and the pubis.
- The ischium is divided into the body, superior ramus of the ischium, and inferior ramus of the ischium.
- The ischial tuberosity, which supports weight when sitting, is located on the ischium.

Key Terms

- **ischium**: The most inferior of the three bones that make up each side of the pelvis.

The ischium forms the lower and back part of the hip bone. Situated below the ilium and behind the pubis, the superior portion of this bone forms approximately one third of the acetabulum, which articulates with the femoral head to form the hip joint.



Lateral view of ischium: Left hip joint, opened by removing the floor of the acetabulum from within the pelvis. The ischium is labeled at the bottom left of the ilium.

The ischium is divisible into three portions; the body, and the superior and inferior rami.


The body contains a prominent spine that is the origin for the gemellus superior muscle. Two indentations run parallel to the spine—superiorly,the greater sciatic notch and, inferiorly, the lesser sciatic notch, through which key nervous and vascular vessels pass.

The superior ramus of the ischium extends inferiorly and posteriorly from the body. It is the partial origin for the obturator internus and obturator externus muscles.

Posteriorly the ramus forms a large swelling termed the tuberosity of the ischium, or ischial tuberosity, which supports weight while sitting and is the origin for the gemellus inferior and adductor magnus muscles.

Dorsally, the ramus contributes to the obturator foramen, a large opening in the pelvis through which key nervous and vascular vessels pass.

The inferior ramus of the ischium is thin and flattened and ascends from the superior ramus of the ischium to join the inferior ramus of the pubis. It is the partial origin for the gracillis and adductor magnus muscles.



Ischium: The ischium is located below the ilium and behind the pubis.

This page titled 7.7B: Ischium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7C: Pubis

The pubis is the lowest and most anterior portion of the hip bones of the pelvis.

Learning Objectives

- Describe the pubic bone

Key Points

- The most anterior portion of the pubis, the pubic symphysis, is where the two hip bones of the pelvis are fused together.
- The pubis has a body, a superior ramus, and an inferior ramus.
- The body of the pubis contributes to the lunate surface and acetabular fossa in the acetabulum.

Key Terms


- **pubic symphysis**: A cartilaginous joint between the two bones of the pubis.

The pubic bone forms the anterior region of the pelvis and contributes to the acetabulum, which articulates with the femoral head to form the hip joint.

The pubic bone is covered by a layer of fat that is covered by the mons pubis. The two pubic bones joint anteriorly through the pubic symphysis, a cartilaginous joint.

The pubic bone articulates with the ilium and the ischium on each hip. The internal surface forms the wall of the lesser pelvis and is the point of origin for a portion of the obturator internus muscle. The pubic bone is divisible into a body, a superior ramus, and an inferior ramus.

The Body of the Pubic Bone



Body of pubic bone: (1) sacrum, (2) ilium, (3) ischium, (4) pubis: 4a-body; 4b-superior ramus; 4c-inferior ramus, (5) pubic symphisis, (6) acetabulum, (7) obturator foramen, (8) coccyx, (red dotted line) linea terminalis.

The body of the pubis is a wide, strong, medial, and flat portion of the pubic bone that unites with the pubic symphisis.

The rough superior edge of the corpus, known as the pubic crest, ends laterally in the pubic tubercle. This tubercle, found roughly 3 cm from the pubic symphysis, is a distinctive feature on the lower part of the abdominal wall and is useful when attempting to localize the superficial inguinal ring and the femoral canal of the inguinal canal.

Its internal surface enters into the formation of the wall of the lesser pelvis and gives origin to a portion of the obturator internus muscle.

Superior Pubic Ramus

The superior pubic ramus is one third of the pubic bone. It forms a portion of the obturator foramen and extends from the body to the median plane where it articulates with its counterpart from the opposite side. It is described in two portions, which are a medial flattened part and a narrow lateral prismoid portion.

Inferior Pubic Ramus

The inferior pubic ramus is a thin and flat bone that makes up one third of the pubis. It passes laterally and downward from the medial end of the superior ramus, and becomes narrower as it descends and joins with the inferior ramus of the ischium below the obturator foramen.

This page titled 7.7C: Pubis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7D: False and True Pelves

The false (greater) pelvis is larger and superior to the true (lesser) pelvis where the pelvic inlet is located.

Learning Objectives

- Describe the differences between false and true pelves

Key Points

- Some believe that the false pelvis is actually part of the abdominal cavity and therefore that the true pelvis is the only true portion of the pelvis.
- The true pelvis contains the pelvic inlet and is a short, curved canal, deeper on its posterior than on its anterior wall.
- The true pelvis contains the pelvic colon, rectum, bladder, and some of the reproductive organs.
- The false pelvis supports the intestines (specifically, the ileum and sigmoid colon) and transmits part of their weight to the anterior wall of the abdomen.

Key Terms

- **true pelvis**: Bounded in front and below by the pubic symphysis and the superior rami of the pubis; above and behind, by the sacrum and coccyx; and laterally, by a broad, smooth, quadrangular area of bone, corresponding to the inner surfaces of the body and superior ramus of the ischium, and the part of the ilium below the arcuate line.
- **false pelvis**: Bounded on either side by the ilium; in front it is incomplete, presenting a wide interval between the anterior borders of the ilia; behind is a deep notch on either side between the ilium and the base of the sacrum.

There is some disagreement as to what constitutes the pelvis. Depending on what is included in the description these groupings are often termed true (lesser) or false (greater) pelves.

The True Pelvis

The true (or lesser) pelvis is bounded in front and below by the pubic symphysis and the superior rami of the pubis; above and behind, by the sacrum and coccyx; and laterally, by a broad, smooth, quadrangular area of bone, corresponding to the inner surfaces of the body and superior ramus of the ischium, and the part of the ilium below the arcuate line.

This cavity is a short, curved canal, deeper on its posterior than on its anterior wall, and contains the pelvic inlet. Some consider this region to be the entirety of the pelvic cavity. Others define the pelvic cavity as the larger space including the false greater pelvis, just above the pelvic inlet.



Greater and lesser pelvis: The greater pelvis (yellow) is larger and superior to the lesser pelvis (red) where the pelvic inlet is located.

The true pelvis contains the pelvic colon, rectum, bladder, and some of the reproductive organs. The rectum is at the back, in the curve of the sacrum and coccyx; the bladder is in front, behind the pubic symphysis.

In the female, the uterus and vagina occupy the interval between these viscera. The pelvic splanchnic nerves arising at S2–S4 are in the lesser pelvis.

The False Pelvis

The false (or greater) pelvis is bounded on either side by the ilium. In front it is incomplete, presenting a wide interval between the anterior borders of the ilia; behind is a deep notch on either side between the ilium and the base of the sacrum.

Some consider this region to be part of the pelvic cavity, while others consider it part of the abdominal cavity (hence the name false pelvis). Others compromise by referring to the area as the abdominopelvic cavity.

The false pelvis supports the intestines (specifically, the ileum and sigmoid colon), and transmits part of their weight to the anterior wall of the abdomen.

This page titled 7.7D: False and True Pelves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7E: Comparison of Female and Male Pelves

The female pelvis has evolved to its maximum width for childbirth and the male pelvis has been optimized for bipedal locomotion.

Learning Objectives

- Apply the diagnostic criteria to determine whether a pelvis is male or female

Key Points

- The pelvis is one of the most useful skeletal elements for differentiating between males and females.
- Female pelves are larger and wider than male pelves and have a rounder pelvic inlet.
- Male iliac crests are higher than females, causing their false pelves to look taller and narrower.
- The male sacrum is longer, narrower, straighter, and has a pronounced sacral promontory relative to the female sacrum.
- The angle between the inferior pubic rami is acute (70 degrees) in men, but obtuse (90–100 degrees) in women. Accordingly, the angle is called the subpubic angle in men and the pubic arch in women.

Key Terms

- **pelvis**: The large compound bone structure at the base of the spine that supports the legs. It consists of hip bone, sacrum, and coccyx.

Different Sex, Different Pelvis

Like the skull, the pelvis is highly useful for determining a skeleton's biological sex. A wide pelvis is beneficial for child birth, however a narrow pelvis is beneficial for locomotion when walking upright. These conflicting demands are often termed the obstetrical dilemma.

The female pelvis has evolved to its maximum width for childbirth—a wider pelvis would make women unable to walk. In contrast, human male pelves are not constrained by the need to give birth and therefore are optimized for bipedal locomotion.

Diagnostic Criteria

There are several diagnostic criteria for differentiating a male from a female pelvis:

1. The female pelvis is larger and broader than the male pelvis, which is taller (owing to a higher iliac crest), narrower, and more compact.
2. The distance between the ischium bones is small in males. This causes the sides of the male pelvis to converge from the inlet to the outlet, whereas the sides of the female pelvis are wider apart. This results in the female inlet being large and oval in shape, while the male inlet is more heart shaped.
3. The angle between the inferior pubic rami is acute (70 degrees) in men, but obtuse (90–100 degrees) in women. Accordingly, the angle is called the subpubic angle in men and pubic arch in women.
4. The greater sciatic notch is wider in females.
5. The ischial spines and tuberosities are heavier and project farther into the pelvic cavity in males.
6. The male sacrum is long, narrow, straighter, and has a pronounced sacral promontory. The female sacrum is shorter, wider, more curved posteriorly, and has a less pronounced promontory.
7. The acetabula are wider apart and face more medially in females than in males. This change in the angle of the femoral head gives the female gait its characteristic swinging of hips.

This page titled 7.7E: Comparison of Female and Male Pelves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.7F: Pelvic Structure and Childbearing

The human pelvis has evolved to be narrow enough for efficient upright locomotion, while still being wide enough to facilitate childbirth.

Learning Objectives

- Describe the evolution of the human pelvis

Key Points

- The human pelvis is narrower and smaller than that of our closest living relatives, the apes. This narrower pelvis enables efficient bipedal locomotion.
- Since the pelvis is vital for both efficient locomotion and childbirth, natural selection has been forced to strike a compromise between a wide pelvis to facilitate birthing large-brained infants and having a narrow pelvis to increase locomotive efficiency.
- The human head and brain is also larger in proportion to the rest of the body in comparison to apes. The female pelvis has evolved to be as wide as possible, to make childbirth easier, without becoming so wide as to make bipedal locomotion too inefficient.
- The male pelvis is narrower, since they do not have to give birth. Men therefore have greater locomotive efficiency.

Key Terms

- **bipedal locomotion**: Walking upright; a form of terrestrial locomotion where an organism moves by means of its two rear limbs, or legs.
- **pubic symphysis**: A cartilaginous joint between the two bones of the pubis.


Human sex differences in pelvic shape have emerged over the course of human evolution. The human pelvis is narrower and smaller than that of our closest living relatives, the apes. These changes in the pelvis enable bipedal locomotion, or upright walking.

This narrowing of the pelvis has also affected the way humans give birth, as a narrow pelvis makes it more difficult for an infant to move through the birth canal. This problem is compounded, because as human pelvises became smaller, the heads of infants became larger to accommodate increased brain size.


Since the pelvis is vital for both efficient locomotion and childbirth, natural selection has been forced to strike a compromise between a wide pelvis to facilitate birthing large-brained infants and having a narrow pelvis to increase locomotive efficiency. This compromise has been referred to as the obstetrical dilemma. Thus, the female pelvis has evolved to be as wide as possible, to make childbirth easier, without becoming so wide as to make bipedal locomotion too inefficient.

Additionally, the female pubic symphysis, which is the cartilaginous joint connecting the left and right side of the pelvis, is remodeled by hormones released during pregnancy, allowing it to stretch during childbirth.

Male pelves are not constrained by the issue of childbirth, and thus are narrower and more optimal for bipedal locomotion. Wider hips in females cause an increased valgus angle, which is the angle between the femur and lower leg. This increases the risk of torsional knee injuries.



The Male Pelvis: The male pelvis is narrower than that of the female, as can be seen by the less than 90 degree angle of the pubic arch.



The Female Pelvis: The female pelvis is wider than that of the male, as can be seen by the greater than 90 degree angle of the pubic arch.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Ilium (bone). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ilium_(bone)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pelvis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pelvis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ilium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ilium**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...is_diagram.png. **License**: ***Public Domain: No Known Copyright***
- Ischium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ischium. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ischium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ischium. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f2/Pelvis_diagram.png. **License**: ***Public Domain: No Known Copyright***
- Pelvis diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pelvis_diagram.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray341. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray341.png. **License**: ***Public Domain: No Known Copyright***
- pubic symphysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pubic_symphysis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Body of pubic bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_of_pubic_bone. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Inferior pubic ramus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Inferior_pubic_ramus. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Superior pubic ramus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Superior_pubic_ramus. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pubis (bone). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pubis_(bone). **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f2/Pelvis_diagram.png. **License**: ***Public Domain: No Known Copyright***
- Pelvis diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pelvis_diagram.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray341. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray341.png. **License**: ***Public Domain: No Known Copyright***
- Body of pubic bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_of_pubic_bone. **License**: ***Public Domain: No Known Copyright***
- Pelvic cavity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pelvic_cavity. **License**: ***CC BY-SA: Attribution-ShareAlike***
- false pelvis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/false%20pelvis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- true pelvis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/true%20pelvis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f2/Pelvis_diagram.png. **License**: ***Public Domain: No Known Copyright***
- Pelvis diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pelvis_diagram.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray341. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray341.png. **License**: ***Public Domain: No Known Copyright***
- Body of pubic bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_of_pubic_bone. **License**: ***Public Domain: No Known Copyright***
- Pelvis. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi.../b9/Pelvis.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human pelvis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_p...ual_dimorphism. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pelvis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pelvis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f2/Pelvis_diagram.png. **License**: ***Public Domain: No Known Copyright***
- Pelvis diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pelvis_diagram.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray341. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray341.png. **License**: ***Public Domain: No Known Copyright***
- Body of pubic bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_of_pubic_bone. **License**: ***Public Domain: No Known Copyright***
- Pelvis. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/b/b9/Pelvis.jpg. **License**: ***CC***

BY-SA: Attribution-ShareAlike
- pubic symphysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pubic_symphysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pubic symphysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pubic_symphysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human pelvis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_pelvis%23Sexual_dimorphism. **License**: *CC BY-SA: Attribution-ShareAlike*
- bipedal locomotion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/bipedal%20locomotion. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ilium. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f2/Pelvis_diagram.png. **License**: *Public Domain: No Known Copyright*
- Pelvis diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pelvis_diagram.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray341. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray341.png. **License**: *Public Domain: No Known Copyright*
- Body of pubic bone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Body_of_pubic_bone. **License**: *Public Domain: No Known Copyright*
- Pelvis. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/b/b9/Pelvis.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray241. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray241.png. **License**: *Public Domain: No Known Copyright*
- Gray242. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray242.png. **License**: *Public Domain: No Known Copyright*

This page titled 7.7F: Pelvic Structure and Childbearing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.8A: Femur (The Thigh)

The femur—the bone of the upper leg—is the longest bone in the human body and one of the strongest.

Learning Objectives

- Describe the femur

Key Points


- The femur is the longest bone in the human skeleton.
- It functions in supporting the weight of the body and allowing motion of the leg.
- The femur articulates proximally with the acetabulum of the pelvis forming the hip joint, and distally with the tibia and patella to form the knee joint.

Key Terms

- **condyle**: A smooth prominence on a bone where it forms a joint with another bone.

The femur or thigh bone is found in the upper leg and is the longest bone in the body. The femur articulates proximally with the acetabulum of the pelvis to form the hip joint, and distally with the tibia and patella to form the knee joint.

Proximal



Femur: The anterior surface of the femur with parts labeled

Proximally, the femur exhibits four key regions. The femoral head projects medially and superiorly and articulates with the acetabulum of the pelvis to form the hip joint. Immediately lateral to the head is the neck that connects the head with the shaft. It is narrower than the head to permit a greater range of movement at the hip joint.

Located superiorly on the main shaft, lateral to the joining of the neck, the greater trochanter is a projection to which the abductor and lateral rotator muscles of the leg attach.

Also located on the main shaft, but inferiorly to the neck joint, is the lesser trochanter. A much smaller projection than the greater trochanter, the psoas major and iliacus muscles attach here.

Shaft

The shaft descends in a slightly medial direction that is designed to bring the knees closer to the body's center of gravity, increasing stability. Due to the widening of the female pelvis this angle is greater in women and can lead to increased knee instability.

Two key features of the shaft are the proximal gluteal tuberosity to which the gluteus maximus attaches, and the distal adductor tubercle to which the adductor magnus attaches.

Distal

Distally, the femur exhibits five key regions. Two rounded regions, termed the medial and lateral condyles, articulate with the tibia at the most anterior projection of the patella.

Between the two condyles lies the intercondylar fossa, a depression in which key knee ligaments attach; this significantly strengthens the knee joint and protects it against torsional damage.

Finally, the two epicondyles, the medial and lateral, lie immediately proximal to the condyles; they are also regions where key internal knee ligaments attach.

This page titled 7.8A: Femur (The Thigh) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.8B: Patella (The Knee)

The patella (knee cap) is the bone between the fibula and femur.

Learning Objectives

- Identify the purpose of the patella

Key Points

- The primary functions of the patella are to enhance leg extension and protect the joint of the knee.
- The patella is a bone embedded within a tendon. This means it is a sesamoid bone.

Key Terms


- **patella**: A sesamoid bone found in the knee, commonly known as the knee cap.
- **sesamoid**: A bone embedded within a tendon.

The patella or knee cap is the bone between the fibula and femur. Each leg has a patella to protect its knee joint. The patella serves two functions:

1. To protect the knee from physical trauma.
2. To enhance the leverage that the quadriceps tendon can exert on the femur, thereby increasing muscle efficiency.

The apex of the patella faces inferiorly and connects to the tibia tuberosity through the patella ligament that attaches to the anterior surface. The base of the patella faces superiorly and is the attachment point for the quadriceps tendon.

The posterior surface of the patella contains the medial and lateral facets that articulate with the condyles of the femur. The lower posterior region of the patella has vascular canaliculi, small channels within the bone, which form the infrapatellar fat pad.



Knee Joint: This image shows the position of the patella relative to the articulation of the femur and the tibia.

This page titled 7.8B: Patella (The Knee) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.8C: Tibia and Fibula (The Leg)

The tibia and the smaller fibula bones comprise the lower leg and articulate at the knee and ankle.

Learning Objectives

- Describe the fibula and tibia

Key Points


- The tibia is more commonly called the shinbone. It is located between the ankle and the patella.
- The fibula is a long, thin bone also located between the ankle and the patella. It runs parallel to the tibia.
- Like the femur, the tibia bears much of the body's weight and plays an essential role in movement and locomotion. The fibula, along with the tibia and the tarsals, forms the ankle.

Key Terms

- **tibia**: The inner and usually the larger of the two bones of the lower leg.
- **fibula**: The smaller of the two bones in the lower leg, the calf bone.

The tibia and fibula are the two bones of the lower leg. The tibia is located medially to the fibula and is much larger. Both are bound together with the interosseous membrane.

The Tibia



The leg: Tibia and fibula in anatomical position with parts labeled.

The tibia, or shin bone, spans the lower leg, articulating proximally with the femur and patella at the knee joint, and distally with the tarsal bones, to form the ankle joint. It is the major weight-bearing bone of the lower leg.

Proximally, there are five key features of the tibia:

1. It widens and forms two condyles —the lateral and medial—that articulate with the condyles of the femur.
2. Between the two condyles is the intercondylar fossa, a small grove, into which two intercondylar tubercles sit. Numerous internal ligaments of the knee joint attach to these tubercles and strengthen it significantly.
3. On the anterior surface of the proximal region and inferiorly to the condyles is the tibial tuberosity to which the patella ligament attaches.
4. The shaft of the tibia is triangular and the soleus muscle, which gives the calf its characteristic shape, originates on the posterior surface.
5. Distally, the tibia also widens to aid with weight bearing and it displays two key features. The medial malleolus is a bony projection that articulates with the tarsal bones to form the ankle joint. Laterally, there is the fibular notch that articulates with the fibula.

The Fibula

The fibula also spans the lower leg, although proximally it does not articulate with the femur or patella. It serves more as an attachment point for muscles rather than a weight-bearing bone.

Proximally, the fibula head articulates with the lateral condyle of the tibia, and the biceps femoris attaches to the fibula head. As with the tibia, the shaft of the fibula is triangular and numerus muscles are involved in the extension and flexion of the foot. These muscles originate from the fibula's surface and include the extensor digitorum longus, soleus, and flexor hallucis longus, among others.

Distally, the fibula forms the lateral malleolus, which is more prominent than the medial malleolus of the tibia. It also articulates with the tarsal bones to form the ankle joint.

This page titled 7.8C: Tibia and Fibula (The Leg) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.8D: Tarsals, Metatarsals, and Phalanges (The Foot)

The human ankle and foot bones include tarsals (ankle), metatarsals (middle bones), and phalanges (toes).

Learning Objectives

- Describe the different types of bones in the foot


Key Points

- The human foot contains 26 bones.
- The foot can be subdivided into the tarsals, metatarsals, and phalanges.

Key Terms

- **phalange**: A bone within the digit.
- **metatarsal**: A bone from the center of the foot that articulates with the tarsals and phalanges.
- **tarsal**: A bone forming part of the ankle or heel.

The foot contains 26 bones that are divided into three regions: the tarsals (or ankle and heel), the metatarsals (forming the sole of the foot), and the phalanges (forming the digits). While sharing a similar underlying structure with the hand, the foot is visibly and structurally different to account for its greater load-bearing and locomotive duties, and reduced fine movements.



The Foot: The foot contains the proximal tarsals that form the ankle and heel; intermediate metatarsals; and the distal phalanges that form the toes.

Tarsals

The tarsal bones of the foot are organized into three rows: proximal, intermediate, and distal. The proximal row contains the talus, which is the most superior of the tarsals and articulates with the tibia and fibula to form the ankle joint. The talus is responsible for transmitting forces from the tibia to the heel and acts as an attachment point for numerous ligaments that strengthen the ankle joint.

The calcaneus is the thickest tarsal and forms the heel of the foot. It articulates with the talus superiorly and anteriorly with the cuboid of the distal group. Posteriorly the calcaneal tuberosity is the attachment point for the Achilles tendon.

The intermediate group contains only the navicular bone, which articulates with all of the tarsals—with the exception of the calcaneus. The navicular bone plays a key role in maintaining the medial longitudinal arch of the foot.

There are four distal tarsals: the lateral cuboid and the three cuneiforms, located medially. The distal tarsals articulate with the metatarsals and also maintain the transverse arch of the foot.

Metatarsals

The foot contains five metatarsals that are numbered I–V, moving medial to lateral, big toe to little toe. Each metatarsal consists of a head, shaft, and base.

The proximal base articulates with the cuboid bones, and distally with the proximal phalanges, and each metatarsal also articulates laterally with adjacent metatarsals. The interossei of the foot originate from the shafts of the metatarsals.

Phalanges

The digits are named in a similar fashion to the metatarsals, medial to lateral from the big toe. With the exception of the big toe, each digit contains a proximal, intermediate, and distal phalange; the big toe lacks an intermediate phalange. The length of the phalanges decreases distally.

This page titled 7.8D: Tarsals, Metatarsals, and Phalanges (The Foot) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.8: THE LOWER LIMB

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

7.8A: Femur (The Thigh)

7.8B: Patella (The Knee)

7.8C: Tibia and Fibula (The Leg)

7.8D: Tarsals, Metatarsals, and Phalanges (The Foot)

7.9E: Arches of the Feet

This page titled 7.8: The Lower Limb is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

7.9E: Arches of the Feet

The arches of the foot are formed by the tarsal and metatarsal bones; they dissipate impact forces and store energy for the subsequent step.

Learning Objectives

- Differentiate among the arches of the foot


Key Points

- The arches of the foot are formed by the tarsal and metatarsal bones and strengthened by ligaments and tendons. They allow the foot to support the weight of the body in the erect posture with the least weight.
- The slight mobility of the arches when weight is applied to and removed from the foot makes walking and running more economical in terms of energy.
- The longitudinal arch of the foot can be broken down into several smaller arches. The main arches are the antero- posterior arches, which may, for descriptive purposes, be regarded as divisible into two types—a medial and a lateral.
- The two longitudinal arches serve as pillars for the transverse arch that runs obliquely across the tarsometatarsal joints.

Key Terms

- **arches of the foot**: The area of the foot formed by the tarsal and metatarsal bones and strengthened by ligaments and tendons. They allow the foot to support the weight of the body in the erect posture with the least weight.

Arches of the Foot



Arches of the Foot: Skeleton of foot. Lateral aspect.

The arches of the foot are formed by the tarsal and metatarsal bones. Strengthened by ligaments and tendons, the elastic properties of arches allow the foot to act as a spring, dissipating impact forces and storing energy to be transfered into the subsequent step improving locomotion.


The two longitudinal arches and a transverse arch are maintained by the interlocking shapes of the foot bones, strong ligaments, and pulling muscles during activity. The slight mobility of these arches when weight is applied to and removed from the foot makes walking and running more economical in terms of energy.

Excessive strain on the tendons and ligaments of the feet can result in fallen arches or flat feet.

Longitudinal Arches

The longitudinal arch of the foot can be broken down into several smaller arches. The main arches are the antero-posterior arches, which may, for descriptive purposes, be regarded as divisible into two types—a medial and a lateral.

Medial Arch



Arches of Foot: Skeleton of foot. Medial aspect.

As can be examined in a footprint, the medial longitudinal arch curves above the ground. It is made by the calcaneus, the talus, the navicular, the three cuneiforms, and the first, second, and third metatarsals.

Its summit is at the superior articular surface of the talus. Its two extremities or piers, on which it rests in standing, are the tuberosity on the plantar surface of the calcaneus posteriorly, and the heads of the first, second, and third metatarsal bones anteriorly.

The chief characteristic of this arch is its elasticity, due to its height and to the number of small joints between its component parts. Its weakest part (i.e., the part most liable to yield from too much pressure) is the joint between the talus and navicular, but this portion is braced by the plantar calcaneonavicular ligament, also called the spring ligament, which is elastic and is thus able to quickly restore the arch to its original condition when the disturbing force is removed.

The ligament is strengthened medially by blending with the deltoid ligament of the ankle joint, and is supported inferiorly by the tendon of the tibialis posterior, which is spread out in a fan-shaped insertion and prevents undue tension of the ligament or such an amount of stretching as would permanently elongate it.

The arch is further supported by the plantar aponeurosis, by the small muscles in the sole of the foot, by the tendons of the peroneus longus and the tibialis anterior and posterior, and by the ligaments of all the articulations involved.

Lateral Arch

In contrast, the lateral longitudinal arch is very low. It is composed of the calcaneus, the cuboid, and the fourth and fifth metatarsals.

Its summit is at the talocalcaneal articulation, and its chief joint is the calcaneocuboid, which possesses a special mechanism for locking and allows only a limited movement. The most marked features of this arch are its solidity and its slight elevation.

Two strong ligaments—the long plantar and the plantar calcaneocuboid—the extensor tendons, and the short muscles of the little toe preserve its integrity.

Fundamental Longitudinal Arch

While these medial and lateral arches may be readily demonstrated as the component antero-posterior arches of the foot, the fundamental longitudinal arch is contributed to by both, and consists of the calcaneus, cuboid, third cuneiform, and third metatarsal: all the other bones of the foot may be removed without destroying this arch.

Transversal Arches

In addition to the longitudinal arches, the foot presents a series of transverse arches. The arches are complete at the posterior part of the metatarsus and the anterior part of the tarsus, but in the middle of the tarsus they present more of the characteristics of concavities.

These are directed downward and medially, so that when the medial borders of the feet are placed in apposition, a complete tarsal dome is formed. The transverse arches are strengthened by the interosseous, plantar, and dorsal ligaments; by the short muscles of the first and fifth toes (especially the transverse head of the adductor hallucis), and by the peroneus longus, whose tendon stretches between the piers of the arches.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Femur. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Femur**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lower extremity of femur. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lower_extremity_of_femur**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Femur**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- condyle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/condyle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2b/Gray244.png**. **License**: ***Public Domain: No Known Copyright***
- Patella. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Patella**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Patella. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Patella**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- exostoses. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/exostoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- patella. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/patella**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2b/Gray244.png**. **License**: ***Public Domain: No Known Copyright***
- Patella. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ee_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Fibula. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tibia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tibia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibula. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tibia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tibia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Prismatoid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Prismatoid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibula. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tibia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tibia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2b/Gray244.png**. **License**: ***Public Domain: No Known Copyright***
- Patella. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ee_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Tibia and fibula. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...45/Gray258.png**. **License**: ***Public Domain: No Known Copyright***
- Foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Foot%23Human_foot**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Foot%23Skeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phalange. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/phalange**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- metatarsus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/metatarsus**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- tarsus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tarsus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2b/Gray244.png**. **License**: ***Public Domain: No Known Copyright***
- Patella. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ee_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Tibia and fibula. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/4/45/Gray258.png**. **License**: ***Public Domain: No Known Copyright***
- Foot. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...52/Gray291.png**. **License**: ***Public Domain: No Known Copyright***
- Arches of the foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arches_of_the_foot**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Foot%23Arches**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Arches of the foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arches_of_the_foot**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- arches of the foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/arches%20of%20the%20foot**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Femur. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2b/Gray244.png**. **License**: ***Public Domain: No Known Copyright***
- Patella. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ee_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Tibia and fibula. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/4/45/Gray258.png**. **License**: ***Public Domain: No Known Copyright***
- Foot. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/5/52/Gray291.png**. **License**: ***Public Domain: No Known Copyright***
- Arches of foot. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/5/52/Gray291.png**. **License**: ***Public Domain: No Known Copyright***
- Arches of a foot. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/81/Gray290.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 7.9E: Arches of the Feet is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8: JOINTS

- Topic hierarchy

8.1: Classification of Joints

8.1A: Structural Classification of Joints

8.1B: Functional Classification of Joints

8.2: Fibrous Joints

8.2A: Fibrous Joints

8.2B: Sutures

8.2C: Syndesmoses

8.2D: Gomphoses

8.3: Cartilaginous Joints

8.3A: Cartilaginous Joints: Synchodroses

8.3B: Cartilaginous Joints: Symphyses

8.4: Synovial Joints

8.4A: Structure of Synovial Joints

8.4B: Nerve and Blood Supply

8.4C: Bursae and Tendon Sheaths

8.4D: Stability and Range of Motion at Synovial Joints

8.4E: Synovial Joint Movements

8.4E: Types of Synovial Joints

This page titled 8: Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.1: CLASSIFICATION OF JOINTS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

8.1A: Structural Classification of Joints

8.1B: Functional Classification of Joints

This page titled 8.1: Classification of Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.1A: Structural Classification of Joints

There are three structural classifications of joints: fibrous, cartilaginous, and synovial.

Learning Objectives

- Describe the three structural categories of joints

Key Points

- The type and characteristics of a given joint determine the degree and type of movement.
- Structural classification categorizes joints based on the type of tissue involved in their formations.
- There are three structural classifications of joints: fibrous, cartilaginous, and synovial.
- Of the three types of fibrous joints, syndesmoses are the most movable.
- Cartilaginous joints allow more movement than fibrous joints but less than synovial joints.
- Synovial joints (diarthroses) are the most movable joints of the body and contain synovial fluid.

Key Terms

- **periosteum**: A membrane that covers the outer surface of all bones.
- **manubrium**: The broad upper part of the sternum.
- **synovial fluid**: A viscous fluid found in the cavities of synovial joints that reduces friction between the articular cartilage during movement.

A joint, also known as an articulation or articular surface, is a connection that occurs between bones in the skeletal system. Joints provide the means for movement. The type and characteristics of a given joint determines its degree and type of movement. Joints can be classified based on structure and function.

Structural classification of joints categorizes them based on the type of tissue involved in formation. There are three structural classifications of joints: fibrous, cartilaginous, and synovial.


Fibrous Joints

Fibrous joints are connected by dense, tough connective tissue that is rich in collagen fibers. These fixed or immovable joints are typically interlocked with irregular edges. There are three types of fibrous joints.

Sutures are the types of joint found in the cranium (skull). The bones are connected by Sharpey's fibres. The nature of cranial sutures allows for some movement in the fetus. However, they become mostly immovable as the individual ages, although very slight movement allows some necessary cranial elasticity. These rigid joints are referred to as synarthrodial.

Syndesmoses are found between long bones of the body, such as the radio-ulnar and tibio-fibular joints. These moveable fibrous joints are also termed amphiarthrodial. They have a lesser range of movement than synovial joints.


Gomphosis is a type of joint found at the articulation between teeth and the sockets of the maxilla or mandible (dental-alveolar joint). The fibrous tissue that connects the tooth and socket is called the periodontal ligament.




Fibrous joints: Image demonstrating the three types of fibrous joints. (a) Sutures (b) Syndesmosis (c) Gomphosis.

Cartilaginous Joints

Cartilaginous joints are connected by fibrocartilage or hyaline cartilage. They allow more movement than fibrous joints but less than that of synovial joints. These types of joints are further subdivided into primary (synchondroses) and secondary (symphyses) cartilaginous joints. The epiphyseal (growth) plates are examples of synchondroses. Symphyses are found between the manubrium and sternum (manubriosternal joint), intervertebral discs, and the pubic symphysis.



Cartilaginous Joints: Image demonstrates a synchondrosis joint with epiphyseal plate (temporary hyaline cartilage joint) indicated (a) and a symphysis joint (b).



Synovial Joint: This diagram of a synovial joint delineates the articular cartilage, articular capsule, bone, synovial membrane, and joint cavity containing synovial fluid.

Synovial Joints

This is the most common and movable joint type in the body. These joints (also called diarthroses) have a synovial cavity. Their bones are connected by dense irregular connective tissue that forms an articular capsule surrounding the bones' articulating surfaces.

A synovial joint connects bones with a fibrous joint capsule that is continuous with the bones' periosteum. This joint capsule constitutes the outer boundary of a synovial cavity and surrounds the bones' articulating surfaces.

Synovial cavities are filled with synovial fluid. The knees and elbows are examples of synovial joints.

This page titled 8.1A: Structural Classification of Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.1B: Functional Classification of Joints

Functional classification of joints is based on the type and degree of movement permitted.

Learning Objectives

- Describe the three functional categories of joints


Key Points

- Synarthrosis joints are immobile or have limited mobility and include fibrous joints.
- Amphiarthrosis joints allow a small amount of mobility and include cartilaginous joints.
- Diarthrosis joints are the freely movable synovial joints.
- Synovial joints can also be classified as nonaxial, monoaxial, biaxial, and multiaxial.
- The various movements permitted by synovial joints are abduction, adduction, extension, flexion, and rotation.

Key Terms

- **fibrous joints**: Fixed or immobile joints that are connected by dense, tough connective tissue that is rich in collagen fibers.
- **cartilaginous joints**: Joints connected by fibrocartilage or hyaline cartilage. They allow more movement than fibrous joints but less than synovial joints.
- **gomphosis joints**: Joints of very limited mobility. These are found at the articulation between teeth and the sockets of maxilla or mandible (dental-alveolar joint).

Joints or articulations (connections between bones) can be classified in a number of ways. The primary classifications are structural and functional. Functional classification is based on the type and degree of movement permitted.



Types of Synovial Joints.jpg: Image of a skeleton and skematics of the different classes of synovial joints.

Three Categories of Functional Joints

- Synarthrosis: These types of joints are immobile or allow limited mobility. This category includes fibrous joints such as suture joints (found in the cranium) and gomphosis joints (found between teeth and sockets of the maxilla and mandible).
- Amphiarthrosis: These joints allow a small amount of mobility. Most joints in this category include cartilaginous joints such as those found between vertebrae and the pubic symphysis.
- Diarthrosis: These are the freely-movable synovial joints. Synovial joints are further classified based on the different types of movement they provide, including:
  - Plane joint
  - Ball and socket joint
  - Hinge joint
  - Pivot joint
  - Condyloid joint
  - Saddle joint

Movement of Synovial Joints

Joints can also be classified by the number of axes of movement they permit:

- Nonaxial (gliding): Found between the proximal ends of the ulna and radius.
- Monoaxial (uniaxial): Movement occurs in one plane. An example is the elbow joint.
- Biaxial: Movement can occur in two planes. An example is the wrist.
- Multiaxial: Includes the ball and socket joints. An example is the hip joint.

The movements possible with synovial joints are:

- Abduction: movement away from the body's midline
- Adduction: movement toward the body's midline
- Extension: straightening limbs at a joint
- Flexion: bending the limbs at a joint
- Rotation: a circular movement around a fixed point

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Joint. **License**: *CC BY-SA: Attribution-ShareAlike*
- Synovial Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synovial_joint. **License**: *CC BY: Attribution*
- Cartilaginous Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cartilaginous_joint. **License**: *CC BY-SA: Attribution-ShareAlike*
- Syndesmosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Syndesmosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Suture (joint). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Suture_(joint). **License**: *CC BY-SA: Attribution-ShareAlike*
- Gomphosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gomphosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Synovial Joints. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:907_Synovial_Joints.jpg. **License**: *CC BY: Attribution*
- Fibrous Joints. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:904_Fibrous_Joints.jpg. **License**: *CC BY: Attribution*
- Cartiliginous Joints. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:906_Cartiliginous_Joints.jpg. **License**: *CC BY: Attribution*
- Synarthrosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synarthrosis. **License**: *CC BY-SA: Attribution-*

ShareAlike

- Synovial Joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synovial_joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amphiarthrosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Amphiarthrosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Synovial Joints. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:907_Synovial_Joints.jpg**. **License**: ***CC BY: Attribution***
- Fibrous Joints. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:904_Fibrous_Joints.jpg**. **License**: ***CC BY: Attribution***
- Cartiliginous Joints. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:906_Cartiliginous_Joints.jpg**. **License**: ***CC BY: Attribution***
- Types of Synovial Joints. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:909_Types_of_Synovial_Joints.jpg**. **License**: ***CC BY: Attribution***

This page titled 8.1B: Functional Classification of Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.2: FIBROUS JOINTS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

8.2A: Fibrous Joints

8.2B: Sutures

8.2C: Syndesmoses

8.2D: Gomphoses

This page titled 8.2: Fibrous Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.2A: Fibrous Joints

Fibrous joints are also called fixed or immovable joints because they do not move.

Learning Objectives

- Describe fibrous joints


Key Points

- A joint is the location at which two or more bones make contact.
- Joints are classified based on structural and functional properties.
- Fibrous joints, such as sutures, syndesmoses, and gomphoses, have no joint cavity.
- Fibrous joints are connected by dense connective tissue consisting mainly of collagen.
- Fibrous joints are called “fixed” or “immovable” joints because they do not move.

Key Terms

- **syndesmoses**: Slightly movable articulations where the contiguous bony surfaces are united by an interosseous ligament, as in the inferior tibiofibular articulation.
- **suture**: In anatomy, a suture is a fairly rigid joint between two or more hard elements such as the bony plates of the skull.
- **gomphoses**: A joint that binds the teeth to bony sockets (dental alveoli) in the maxillary bone and mandible.

A joint is the location at which two or more bones make contact. They are constructed to allow movement (except for skull bones), provide mechanical support, and are classified structurally and functionally. Structural classification is determined by how the bones connect to each other, while functional classification is determined by the degree of movement between the articulating bones. In practice, there is significant overlap between the two types of classifications.



Fibrous Joints: The adult skull is normally made up of 22 bones. Except for the mandible, all are joined together by sutures, semi-rigid articulations formed by bony ossification. The presence of Sharpey’s fibers permit a little flexibility.

Fibrous joints are joined by dense irregular connective tissue that is rich in collagen fibers.

Characteristics of Fibrous Joints

Fibrous joints are connected by dense connective tissue consisting mainly of collagen. These joints are also called fixed or immovable joints because they do not move. Fibrous joints have no joint cavity and are connected via fibrous connective tissue. The skull bones are connected by fibrous joints called sutures. The skull bones of a fetus are unfused so that they can move over each other slightly to compress skull size during birth. After birth, the bones slowly begin to fuse to become fixed, making the skull bones immovable in order to protect the brain from impact.

Syndesmoses of long bones and gomphoses of teeth are also types of fibrous joints. The movement of the root within a gomphosis has a threefold effect. It lessens some of the impact between the upper and lower teeth in biting. It also pumps blood and lymph from the periodontal membrane into the dental veins and lymph channels and stimulates sensory nerve terminals in the membrane to send signals to the brain centers that control the muscles of mastication.

This page titled 8.2A: Fibrous Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.2B: Sutures

A suture is a type of fibrous joint (synarthrosis) bound by Sharpey's fibers that only occurs in the skull (cranium).

Learning Objectives

- Describe a suture joint

Key Points


- A suture 's fibrous connective tissue helps protect the brain and form the face by strongly uniting the adjacent skull bones.
- Sutures form a tight union that prevents most movement between the bones. Most sutures are named for the bones they articulate.
- Skull sutures visible from the side (norma lateralis) include the frontal, parietal, temporal, occipital, sphenoid, and zygomatic bones, while skull sutures visible from the front (norma frontalis) and above (norma verticalis) include those related to the frontal and parietal bones.
- Skull sutures visible from below (norma basalis) include the frontal, ethmoid, and sphenoid bones.

Key Terms

- **fontanelle**: An anatomical feature of the infant human skull comprising the soft membranous gaps.
- **Sharpey's fibres**: A matrix of connective tissue consisting of bundles of strong collagenous fibers connecting periosteum to bone.
- **suture**: A type of fibrous joint which only occurs in the skull (cranium).

A suture is a type of fibrous joint which only occurs in the cranium, where it holds bony plates together. Sutures are bound together by a matrix of connective tissues called Sharpey's fibers, which grow from each bone into the adjoining one. A tiny amount of movement is permitted at sutures, which contributes to the compliance and elasticity of the skull. These joints are synarthroses (immovable joints).

Cranial Sutures



Cranial Sutures: Lateral view of skull showing the location of some of the cranial sutures.

Most sutures are named for the bones they articulate, but some have special names of their own. Sutures primarily visible from the side of the skull (norma lateralis) include:

- Coronal suture: between the frontal and parietal bones
- Lambdoid suture: between the parietal, temporal, and occipital bones
- Occipitomastoid suture
- Parietomastoid suture
- Sphenofrontal suture
- Sphenoparietal suture
- Sphenosquamosal suture
- Sphenozygomatic suture
- Squamosal suture: between the parietal and the temporal bone
- Zygomaticotemporal suture
- Zygomaticofrontal suture

Sutures primarily visible from front of the skull (norma frontalis) or above the skull (norma verticalis) include:


- Frontal suture / Metopic suture: between the two frontal bones, prior to the fusion of the two into a single bone
- Sagittal suture: along the midline, between parietal bones.

Sutures primarily visible from below the skull (norma basalis) or inside the skull include:

- Frontoethmoidal suture
- Petrosquamous suture
- Sphenoethmoidal suture
- Sphenopetrosal suture

The fibrous connective tissue found at a suture (to bind or sew) strongly unites the adjacent skull bones and thus helps to protect the brain and form the face. In adults, the skull bones are closely opposed and fibrous connective tissue fills the narrow gap between the bones. The suture is frequently convoluted, forming a tight union that prevents most movement between the bones.

Fontanelles



Frontal suture top view: Drawing of human baby skull seen from the top. Cranial sutures are depicted with the frontal suture highlighted in blue.

It is normal for many of the bones of the skull to remain unfused at birth. The fusion of the skull's bones at birth is known as craniosynostosis. The joint between the mandible and the cranium, the temporomandibular joint, forms the only non-sutured joint in the skull. In newborns and infants, the areas of connective tissue between the bones are much wider, especially in those areas on the top and sides of the skull that will become the sagittal, coronal, squamous, and lambdoid sutures.

These broad areas of connective tissue are called fontanelles. During birth, the fontanelles provide flexibility to the skull, allowing the bones to push closer together or to overlap slightly, thus aiding movement of the infant's head through the birth canal. After birth, these expanded regions of connective tissue allow for rapid growth of the skull and enlargement of the brain. The fontanelles greatly decrease in width during the first year after birth as the skull bones enlarge. When the connective tissue between the adjacent bones is reduced to a narrow layer, these fibrous joints are now called sutures.

Synostosis

At some sutures, the connective tissue will ossify and be converted into bone, causing the adjacent bones to fuse to each other. This fusion between bones is called a synostosis (joined by bone). Examples of synostosis fusions between cranial bones are found both early and late in life. At the time of birth, the frontal and maxillary bones consist of right and left halves joined together by sutures, which disappear by the eighth year as the halves fuse together to form a single bone. Late in life, the sagittal, coronal, and lambdoid sutures of the skull will begin to ossify and fuse, causing the suture line to gradually disappear.

This page titled 8.2B: Sutures is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.2C: Syndesmoses

Syndesmoses are slightly movable joints formed where an interosseous ligament joins two bones.

Learning Objectives

- Describe syndesmoses

Key Points

- A syndesmosis, a subcategory of fibrous joints, is a slightly movable (amphiarthrodial) articulation where the contiguous bony surfaces are united by an interosseous ligament, such as the tibiofibular articulation.
- Tears in this joint are generally repaired with a syndesmotic screw.
- Due to the lack of flexibility in these joint structures, ligament injuries in syndesmosis joints are common, particularly at the wrist and ankle.

Key Terms

- **symphysis**: The cartilaginous material that adjoins and facilitates the junction of such bones, with or without synovia.
- **prime mover**: A muscle that acts directly to bring about a desired movement.
- **diastasis**: A separation between two parts of a bone, without fracture.

A syndesmosis is a type of articulation or joint in which two adjacent bones are joined by an interosseous membrane.


Interosseous Membrane

The interosseous membrane is a type of connective tissue found between certain bones, such as those in syndesmosis joints. The membrane is important in creating compartments to separate different structures, distributing the impact of forces and separating the joints. For example, the long bones of the lower arm and the leg both have attached interosseous membranes. In the leg, the interosseous membrane extends between the tibia and the fibula, running along the crests of the bones. The muscles in the leg are separated into sections in the front and back with this membrane. The strength of the membrane allows absorption and distribution of impacts to either bone.

The interosseous membrane in the lower arm extends between the radius and the ulna. It is involved in the elbow joint and helps to stabilize the lower arm bones for strength, durability, and flexibility. Like other joint tissue, it is designed to be able to deform and flex rather than shred or fracture on impact, allowing the joint to absorb considerable stress before damage occurs.

Syndesmoses and Amphiarthrosis Joints

Along with symphysis joints, syndesmoses are classified as amphiarthrosis joints in that they allow slight movement. Joints of this kind are found at several points in the human body, including the intermediate radioulnar joint where the radius and ulna meet above the wrist, in the spine between the spinous processes of various adjacent vertebra, and above the ankle joint where the tibia and fibula converge.



Fibrous Joints: Image of fibrous joints with the tibiofibular syndesmosis demonstration in figure (b).

Located directly above the ankle joint, which is a synovial hinge joint, the ankle syndesmosis is held together by four ligaments. The anterior inferior tibiofibular ligament crosses in front of the tibia and fibula bones. The posterior inferior tibiofibular ligament and the transverse ligament connect the two bones from behind, and the interosseous ligament runs between the contiguous bony surfaces of the two bones.

Due to the limited flexibility in these joint structures, ligament injuries in syndesmosis joints are common, particularly at the wrist and ankle. When the wrist or ankle joint is bent beyond its normal range of motion, a sprain or even a tear in these ligaments can occur. Mild syndesmosis injury may involve the sprain of a single ligament.

More severe injuries can involve damage to multiple ligaments at once or even the separating of the bones at the joint (known as diastasis). Players of rough sports such as football or rugby have an increased risk of fracturing their fibulas and tearing the interosseous ligament between it and the tibia. When that happens, the surgeon temporarily replaces the ligament with a syndesmotic screw.

This page titled 8.2C: Syndesmoses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.2D: Gomphoses

A gomphosis is a fibrous joint that binds the teeth to bony sockets in the bones of the maxilla mandible.

Learning Objectives

- Describe a gomphosis joint


Key Points

- The gomphosis is the only joint type in which a bone does not join another bone, because teeth are not technically bone.
- The motion of a gomphosis is minimal, though considerable movement can be achieved with pressure over time, which is why using braces can realign teeth.
- One disorder that can affect the gomphosis is scurvy, which is a disease of connective tissue.

Key Terms

- **scurvy**: A disease resulting from a lack of vitamin C.
- **synarthrosis**: A type of joint in which two bones are connected rigidly by fibrous tissue.
- **gomphoses**: A joint that binds the teeth to bony sockets (dental alveoli) in the maxillary bone and mandible.

A gomphosis is a joint that anchors a tooth to its socket. Gomphoses line the upper and lower jaw in each tooth socket and are also known as peg and socket joints. These joints have a very limited range of mobility so the teeth are held firmly in place. However, as illustrated with braces, it is possible to move them incrementally over time. Each tooth has bony protrusions or pegs that latch into the socket with the assistance of the gomphosis. Disorders of the mouth sometimes involve these joints.



Gomphoses: This image illustrates the gomphoses joints of teeth within the jaw.

This particular joint is an example of a synarthrosis, a joint with limited to no movement. Several other joints of this type can be found in the body, including the connections between the plates of the skull. The gomphosis is made up of fibrous tissue, a collection of tough ligaments that attach to the socket and base of the tooth. As people age and lose their initial set of baby teeth, the new teeth develop gomphoses to anchor them in the jaw.

One disorder that can affect the gomphosis is scurvy, a disease of connective tissue. Connective tissues such as the ligaments around the teeth start to dissolve. Patients with untreated scurvy develop loose teeth that may eventually fall out because the joints are too unstable. Periodontal infection and inflammation can also damage the joint, causing pain and erosion in the soft tissue. Chronic dental problems may loosen the ligaments and lead to tooth loss or instability.

Patients with braces and retainers take advantage of the limited range of movement offered by the gomphosis to pull teeth into new positions. This may be necessary for a variety of reasons. The goal is to align the teeth evenly to create a strong, healthy bite. Braces are adjusted incrementally over time to pull and push the teeth into place. Between each adjustment, the teeth and jaw have time to recover.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Fibrous joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrous_joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- suture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/suture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gomphoses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/gomphoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- syndesmoses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/syndesmoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skull bones diagram. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...ed_(bones).svg**. **License**: ***Public Domain: No Known Copyright***
- Fontanelle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fontanelle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Suture (anatomical). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Suture_(anatomical)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sharpey's fibers. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sharpey's_fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Suture (joint). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Suture_(joint)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tonye Ogele, Joints. January 16, 2014. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m46182/latest/**. **License**: ***CC BY: Attribution***
- OpenStax College, Fibrous Joints. January 16, 2014. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/contents/edcb337f-44b...104fdb0c53d1@3**. **License**: ***CC BY: Attribution***
- suture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/suture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sharpey's fibres. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sharpey's%20fibres**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skull bones diagram. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/1/1a/Human_skull_front_simplified_(bones).svg**. **License**: ***Public Domain: No Known Copyright***
- Cranial Sutures. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2c/Gray188.png**. **License**: ***Public Domain: No Known Copyright***
- Frontal suture top view.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Frontal_suture_top_view.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrous joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrous_joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Syndesmoses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Syndesmoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/prime-mover**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- symphysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/symphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diastasis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diastasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Skull bones diagram. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/1/1a/Human_skull_front_simplified_(bones).svg**. **License**: ***Public Domain: No Known Copyright***
- Cranial Sutures. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/2c/Gray188.png**. **License**: ***Public Domain: No Known Copyright***
- Frontal suture top view.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Frontal_suture_top_view.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrous Joints. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrous_joint#/media/File:904_Fibrous_Joints.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gomphoses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gomphoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synarthrosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/synarthrosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gomphoses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/gomphoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skull bones diagram. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/1/1a/Human_skull_front_simplified_(bones).svg**. **License**: ***Public Domain: No Known Copyright***
- Cranial Sutures. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/2c/Gray188.png**. **License**: ***Public Domain: No Known Copyright***
- Frontal suture top view.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Frontal_suture_top_view.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrous Joints. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrous_joint#/media/File:904_Fibrous_Joints.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gompnoses. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/6c/Gray1003.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 8.2D: Gomphoses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.3: CARTILAGINOUS JOINTS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

8.3A: Cartilaginous Joints: Synchodroses

8.3B: Cartilaginous Joints: Symphyses

This page titled 8.3: Cartilaginous Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.3A: Cartilaginous Joints: Synchodroses

Cartilaginous joints connected by hyaline cartilage are termed synchondroses.

Learning Objectives

- Describe synchondroses


Key Points

- The first sternocostal joint where the first rib meets the sternum is a synchondrosis.
- The epiphyseal growth plate is a temporary cartilaginous joint formed as the cartilage is converted to bone during growth and development.
- Cartilaginous joints are connected entirely by cartilage and allow more movement between bones than a fibrous joint, but less than the highly mobile synovial joint.

Key Terms

- **synchondrosis**: A slightly moveable articulation between bones joined by hyaline cartilage.
- **apoptosis**: A type of "cell suicide" called programmed cell death that occurs in multicellular organisms.
- **epiphyseal plate**: The epiphyseal plate is a hyaline cartilage plate where growth occurs in children and adolescents, located in the metaphysis at each end of a long bone.

Cartilaginous joints are connected entirely by cartilage (fibrocartilage or hyaline). Cartilaginous joints allow more movement between bones than a fibrous joint but less than the highly mobile synovial joint. The joint between the manubrium and the sternum is an example of a cartilaginous joint. This type of joint also forms the growth regions of immature long bones and the intervertebral discs of the spinal column.



Synchondroses: Section through occipitosphenoid synchondrosis of an infant, including the cartilage, perichrondrium, and periosteum.

Where the connecting medium is hyaline cartilage, a cartilaginous joint is termed a synchondrosis or primary cartilaginous joint. A synchondrosis joint is the first sternocostal joint (where the first rib meets the sternum). In this example, the rib articulates with the sternum via the costal cartilage. The rest of the sternocostal joints are synovial plane joints.

A temporary form of joint called an epiphyseal (growth) plate, is one where the cartilage is converted into bone before adult life. Such joints are found between the epiphyses and diaphyses of long bones, between the occipital and the sphenoid bones, and during the early years of life, between the petrous portion of the temporal and the jugular process of the occipital bone. The epiphyseal plate is a hyaline cartilage plate in the metaphysis at each end of a long bone.

The epiphyseal plate is found in children and adolescents. In puberty, increasing levels of estrogen, in both females and males, leads to increased apoptosis of chondrocytes in the epiphyseal plate. Depletion of chondrocytes due to apoptosis leads to less ossification, and growth slows down and later stops when the cartilage has been completely replaced by bone. This process leaves only a thin epiphyseal scar that later disappears. In adults who have stopped growing, the plate is replaced by an epiphyseal line.

Those with achondroplasia (a form of dwarfism) have premature closure of the epiphyseal growth plates, which results in shorter than average arms and legs.

This page titled 8.3A: Cartilaginous Joints: Synchodroses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.3B: Cartilaginous Joints: Symphyses

A symphysis is a secondary cartilaginous joint that is permanent and slightly movable.

Learning Objectives

- Differentiate among the types of symphyses between two bones

Key Points

- Symphyses include the pubic symphysis and the intervertebral disc between two vertebrae, among others.
- The pubic symphysis or symphysis pubis is the midline cartilaginous joint uniting the superior rami of the left and right pubic bones. It widens slightly whenever the legs are stretched far apart and can become dislocated.
- Intervertebral discs lie between adjacent vertebrae in the spine. Each disc forms a cartilaginous joint to allow slight movement of the vertebrae and acts as a ligament to hold the vertebrae together.


Key Terms

- **annulus fibrosus**: Fibrous ring of intervertebral disk.
- **nucleus pulposus**: Inner gel-like center of the vertebral disc.
- **sciatica**: Pain that travels down the leg from the lower back region.
- **symphysis**: The cartilaginous material that adjoins and facilitates the junction of such bones, with or without synovia.
- **intervertebral disc**: A cartilaginous joint that allows slight movement of the vertebrae by lying between adjacent vertebrae in the spine. It also acts as a ligament to hold the vertebrae together.

A symphysis, a type of secondary cartilaginous joint, is a fibrocartilaginous fusion between two bones. It is an amphiarthrosis (slightly movable) joint, and an area where two parts or structures grow together. Unlike synchondroses, symphyses are permanent. The more prominent symphyses are the pubic symphysis; the symphyses between the bones of the skull, most notably the mandible (symphysis menti); sacrococcygeal symphysis; the intervertebral disc between two vertebrae; and in the sternum, between the manubrium and body, and between the body and xiphoid process.

Pubic Symphysis

The pubic symphysis or symphysis pubis is the midline cartilaginous joint (secondary cartilaginous) uniting the superior rami of the left and right pubic bones. It is a nonsynovial amphiarthrodial joint connected by fibrocartilage, and may contain a fluid-filled cavity. The ends of both pubic bones are covered by a thin layer of hyaline cartilage attached to the fibrocartilage.



Symphyses: Diagrammatic section of a symphysis including the ligament, disc of fibrocartilage, and articular cartilage.

The pubic symphysis is located anterior to the urinary bladder and superior to the external genitalia, above the vulva in females and above the penis in males. The suspensory ligament of the penis attaches to the pubic symphysis. In females, the pubic

symphysis is intimately close to the clitoris. In normal adults, it can be moved roughly two mm and with one degree of rotation. Mobility of this joint increases for women at the time of childbirth.

During birth, the pubic symphysis of relaxes to slightly widen the birth canal. This movement is minimal, but along with the compression of the unfused fetal skull generally allows an infant to be born vaginally.

The pubic symphysis widens slightly whenever the legs are stretched far apart. In sports in which this movement is frequent, the risk of a pubic symphysis blockage is high. This injury occurs when the bones at the symphysis do not realign correctly after completion of the movement and get jammed in a dislocated position. The resulting pain can be quite severe, especially if further strain is put upon the affected joint. In most cases, the joint can only be successfully reduced into its normal position by a trained medical professional.

Pubic symphyses have importance in the field of forensic anthropology, as they can be used to estimate the age of adult skeletons. Throughout life, the surfaces become worn at a more or less predictable rate. By examining the wear of the pubic symphysis, it is possible to estimate the age of the person at death.

Mandible

The external surface of the mandible is marked in the median line by a faint ridge, indicating the symphysis menti, mandibular symphysis, or line of junction. This line delineates the two pieces of bone that compose the mandible during the first years of life.

Intervertebral Discs

Intervertebral discs (or intervertebral fibrocartilage) lie between adjacent vertebrae in the spine. Each disc forms a cartilaginous joint to allow slight movement of the vertebrae and acts as a ligament to hold the vertebrae together. The discs consist of an outer annulus fibrosus that surrounds the inner nucleus pulposus. The annulus fibrosus and the nucleus pulposus distribute pressure evenly across the disc.

The nucleus pulposus contains loose fibers suspended in a mucoprotein gel with the consistency of jelly. The nucleus of the disc acts as a shock absorber, absorbing the impact of the body's daily activities and keeping the two vertebrae separated. The disc can be likened to a jelly doughnut with the annulus fibrosis as the dough and the nucleus pulposis as the jelly. If one presses down on the front of the doughnut, the jelly moves posteriorly. When one develops a prolapsed disc, the jelly (the nucleus pulposus) is forced out of the doughnut (the disc) and may put pressure on the nerve located near the disc, potentially causing symptoms of sciatica.




Diagram of Invertebral Disc: The lateral and superior view of an invertebral disc, including the vertebral body, intervertebral foramen, anulus fibrosis, and nucleus pulposus.

Aging causes disc degeneration, in which the nucleus pulposus begins to dehydrate and the concentration of proteoglycans in the matrix decreases, limiting the ability of the disc to absorb shock. This general shrinking of disc size is partially responsible for the common decrease in height as humans age.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Apoptosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Apoptosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cartilaginous joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cartilaginous_joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Synchondroses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synchondroses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epiphyseal plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epiphyseal_plate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epiphyseal plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/epiphyseal%20plate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synchondrosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/synchondrosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/7/76/Gray297.png**. **License**: ***Public Domain: No Known Copyright***
- intervertebral disc. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/intervertebral_disc**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Invertebral disc. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Invertebral_disc**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pubic symphysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pubic_symphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Symphyses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Symphyses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- symphysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/symphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/7/76/Gray297.png**. **License**: ***Public Domain: No Known Copyright***
- intervertebral disc. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:716_Intervertebral_Disk.jpg**. **License**: ***CC BY: Attribution***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/b1/Gray298.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 8.3B: Cartilaginous Joints: Symphyses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4: SYNOVIAL JOINTS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

8.4A: Structure of Synovial Joints

8.4B: Nerve and Blood Supply

8.4C: Bursae and Tendon Sheaths

8.4D: Stability and Range of Motion at Synovial Joints

8.4E: Synovial Joint Movements

8.4E: Types of Synovial Joints

This page titled 8.4: Synovial Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4A: Structure of Synovial Joints

A synovial joint or diarthrosis occurs at articulating bones to allow movement. It is distinguished by a surrounding synovial capsule.

Learning Objectives

- Identify the structures of the synovial joint that allow it to move freely

Key Points

- The bones of a synovial joint are surrounded by a synovial capsule, which secretes synovial fluid to lubricate and nourish the joint while acting as a shock absorber.
- The ends of the joint bones are covered with smooth, glass-like hyaline cartilage which reduces friction during movement.
- A synovial joint contains a synovial cavity and dense, irregular connective tissue that forms the articular capsule normally associated with accessory ligaments.


Key Terms

- **articulation**: A joint or the collection of joints at which something is articulated, or hinged, for bending.
- **synovial membrane**: A thin membrane of joints comprised of smooth connective tissue and that secretes synovial fluid.
- **synovial fluid**: A viscous, non-Newtonian fluid found in the cavities of synovial joints. With its yolk-like consistency, its principal role is to reduce friction between the articular cartilage of synovial joints during movement.
- **articular cartilage**: A tough, elastic, fibrous connective tissue found in various parts of the body such as the joints, outer ear, and larynx. A major constituent of the embryonic and young vertebrate skeleton, converted largely to bone with maturation.
- **diarthrosis**: A joint that can move freely in various planes.

A synovial joint, also known as a diarthrosis, is the most common and most movable type of joint in a mammal's body. Diarthroses are freely movable articulations. In these joints, the contiguous bony surfaces are covered with articular cartilage and connected by ligaments lined by synovial membrane. The joint may be divided, completely or incompletely, by an articular disk or meniscus, the periphery of which is continuous with the fibrous capsule while its free surfaces are covered by synovial membrane.

The articular capsule is fibrous and continuous with the periosteum of articulating bones, surrounding the diarthrosis and uniting the articulating bones. The articular capsule also consists of two layers: (1) the outer fibrous membrane that may contain ligaments and (2) the inner synovial membrane that secretes the lubricating, shock-absorbing, and joint-nourishing synovial fluid. The bones of a synovial joint are covered by a layer of hyaline cartilage that lines the epiphyses of joint ends of bone with a smooth, slippery surface that does not bind them together. This articular cartilage functions to absorb shock and reduce friction during movement.

Synovial Membrane and Components



Synovial Joint: An illustration of the structure of a synovial joint.

A synovial membrane (or synovium) is the soft tissue found between the articular capsule (joint capsule) and the joint cavity of synovial joints. Synovial fluid is the clear, viscid, lubricating fluid secreted by synovial membranes. The morphology of synovial membranes may vary, but it often consists of two layers. The outer layer, or subintima, can be fibrous, fatty, or loosely areolar. The inner layer, or intima, consists of a sheet of cells thinner than a piece of paper.

Where the underlying subintima is loose, the intima sits on a pliable membrane called the synovial membrane. This membrane, together with the cells of the intima, acts like an inner tube, sealing the synovial fluid from the surrounding tissue and effectively stopping the joints from being squeezed dry when subjected to impact (such as when running). As with most other joints, synovial joints achieve movement at the point of contact of the articulating bones. The main structural differences between synovial and fibrous joints are the existence of capsules surrounding the articulating surfaces of a synovial joint and the presence of lubricating synovial fluid within those capsules (synovial cavities).

Synoviocytes

The intimal cells are termed synoviocytes and can be either fibroblastic (type B synoviocytes) and macrophagic (type A synoviocytes). Both types have differences from similar cells in other tissues. The type B synoviocytes manufacture a long-chain sugar polymer called hyaluronan, which combines with a molecule called lubricin to give the synovial fluid a stringy, egg-white consistency. The water component of synovial fluid is effectively trapped in the joint space by the hyaluronan due to its large, highly negatively charged moieties. The macrophages are responsible for the removal of undesirable substances from the synovial fluid.

Structure of Synovium

The surface of a synovium may be flat or covered with finger-like projections (villi) to allow the soft tissue to change shape as the joint surfaces move on one another. Just beneath the intima, most synovia have a dense net of small blood vessels that provide nutrients for the synovia and the avascular cartilage.

In any one position, much of the cartilage is close enough to get nutrition directly from the synovium. Some areas of cartilage have to obtain nutrients indirectly and may do so either from diffusion through cartilage or by the stirring of synovial fluid.

Synovial Bursa

The synovial bursa is a small, fluid-filled sac lined by synovial membrane containing synovial fluid. It provides a cushion between bones and tendons and/or muscles around a joint.

This page titled 8.4A: Structure of Synovial Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4B: Nerve and Blood Supply

Synovial joints are highly innervated but vascularized indirectly by nearby tissues.

Learning Objectives


- Identify the nerve and blood supply of synovial joints

Key Points

- Although the articular capsule is innervated with the nerves necessary for movement, it lacks blood vessels because the arteries wrap around the joint in an anastomosis, bypassing direct capillary contact with the capsule.
- The articular and epiphyseal branches given off by the neighboring arteries form a periarticular arterial plexus.
- Exchange of gases (oxygen and carbon dioxide) and nutrients is achieved, albeit slowly, via diffusion or more efficiently during exercise via convection.

Key Terms

- **convection**: The movement of groups of molecules within fluids such as liquids or gases.
- **osteomyelitis**: An infection of the bone and bone marrow characterized by inflammation.
- **anastomosis**: A cross-connection between two blood vessels.



Elbow Joint: Diagram of the anastomosis around the elbow joint.

The blood supply of a synovial joint comes from the arteries sharing in anastomosis around the joint. The articular and epiphyseal branches of the neighboring arteries form a periarticular arterial plexus. The articular capsule is highly innervated but avascular (lacking blood and lymph vessels), and receives nutrition from the surrounding blood supply via either the slow process of diffusion or convection, a far more efficient process.

Numerous vessels from this plexus pierce the fibrous capsule and form a rich vascular plexus in the deeper part of the synovial membrane. The blood vessels of the synovial membrane terminate around the articular margins in a fringe of looped anastomoses termed the circulus vasculosus (circulus articularis vasculosus). It supplies the capsule, synovial membrane, and the epiphyses. After epiphyseal fusion in the growth of long bones, communication between the circulosus vasculosus and the end arteries of

the metaphysis is established. This minimizes the chances of osteomyelitis in the metaphysis.

The synovial cartilage in the capsule acts somewhat like a sponge. A sponge will absorb fluid, but it will release little of that fluid unless it is squeezed. Exercising the joint, in effect, squeezes the synovial "sponge," allowing gas exchange to occur and nutrients to flow into the cartilage. Flexing and extending the joint alternately squeezes the sponge and releases it to reabsorb more fluid.

This page titled 8.4B: Nerve and Blood Supply is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4C: Bursae and Tendon Sheaths

Joints are cushioned by small fluid-filled sacs called bursae and stabilized by tough bands of fibrous connective tissue called tendons.

Learning Objectives

- List the components of a joint


Key Points

- Synovial joints are made up of five classes of tissues. These include bone, cartilage, synovium, synovial fluid, and tensile tissues composed of tendons and ligaments.
- Tendons are tough bands of fibrous connective tissue that connect muscles to bones.
- Bursae are sacs filled with synovial fluid that provide cushioning around a joint between the bones and the muscles and tendons crossing the joint.

Key Terms

- **retinacula**: A band around tendons that holds them in place for stabilization.
- **tendon**: A tough band of inelastic fibrous tissue that connects a muscle with its bony attachment.
- **synovial fluid**: A viscous, non-Newtonian fluid found in the cavities of synovial joints. With its yolk-like consistency, its principal role is to reduce friction between the articular cartilage of synovial joints during movement.
- **connective tissue**: A type of tissue found in animals that functions in binding other tissue systems (such as muscle to skin) or organs. It consists of the cells, fibers, and a ground substance or extracellular matrix.

Synovial joints are made up of five classes of tissues: bone, cartilage, synovium, synovial fluid, and tensile tissues composed of tendons and ligaments. The synovial lining in the bursae and tendon sheaths, similar to that within joints, is a slippery, non-adherent surface allowing movement between planes of tissue. Synovial tendon sheaths line tendons only where they pass through narrow passages or retinacula, as in the palm, at the wrist, and around the ankle. Elsewhere, the tendon lies in a bed of loose fibrous tissue.




Tendons: Illustration of the location of tendons in the hand

A tendon or sinew is a tough band of fibrous connective tissue that usually connects muscle to bone and is capable of withstanding tension. Tendons are similar to ligaments and fasciae as they are all made of collagen, but a ligament joins one bone to another and fasciae connect muscles to other muscles. Tendons connect muscle to bone and move the bones or structures to

which they are attached.

A bursa (plural bursae) is a small, fluid-filled sac lined by synovial membrane with an inner capillary layer of fluid (synovial fluid) with the consistency of raw egg white. It provides a cushion between bones and tendons or muscles around a joint. This helps to reduce friction between the bones and allows for free movement.

Bursae occur at sites of shearing in subcutaneous tissue or between deeper tissues such as muscle groups and fascia. Many bursae develop during growth, but new or adventitious bursae can occur at sites of occupational friction. Bursae are found around most major joints of the body, such as the shoulder and the knee. For example, to protect the knee and reduce friction from the various muscles, tendons, and ligaments that attach to and cross the knee joint, knees are cushioned by 14 different bursae: five in front, four laterally, and five medially.



Knee joint: Diagram of the knee joint.

This page titled 8.4C: Bursae and Tendon Sheaths is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4D: Stability and Range of Motion at Synovial Joints

Tendons provide stability at joints.

Learning Objectives

- Explain the roles of tendons in movement and flexibility

Key Points


- Although tendons have long been considered just a way to attach muscles to bones, research has shown that their springy properties also allow them to provide stability during locomotion with no active work.
- The elasticity of tendons enables them to release stored energy during walking, allowing the muscles to generate greater force without changing length.
- Many factors influence joint stability and range of motion.

Key Terms

- **pronation**: The action of rotating the forearm so that the palm of the hand is turned down or back.
- **supination**: The action of rotating the forearm so that the palm of the hand is turned up or forward
- **eversion**: The condition of being turned outward.
- **plantarflexion**: The movement that increases the approximate 90 degree angle between the front part of the foot and the shin.
- **dorsiflexion**: The movement which decreases the angle between the dorsum (superior surface) of the foot and the leg, so that the toes are brought closer to the shin.

A tendon is a mechanism by which muscles connect to bone and that transmits force. However, over the past two decades, research has also characterized the elastic properties of tendons and their ability to function as springs. This characteristic allows tendons to passively modulate forces during locomotion, thus providing additional stability with no active work. It also allows tendons to store and recover energy with high efficiency.

Effect of Tendon Elasticity



Achilles Tendon: The Achilles tendon, also called the calcaneus, provides stability and limits the range of motion at the ankle joint. It's depicted in this diagram in relation to the tendo calcaneus.

During a human stride, the Achilles (calcaneal) tendon stretches as the ankle joint undergoes dorsiflexion. During the last portion of the stride, as the foot undergoes plantar flexion (pointing the toes downward), the stored elastic energy is released. Because the tendon stretches, the muscle is able to function with less or even no change in length, allowing it to generate greater force.

Joint Stability

Certain joints exhibit special movements including elevation, depression, protraction, retraction, inversion, eversion, dorsiflexion, plantar flexion, supination, pronation, and opposition. A number of factors influence joint stability. These include:

- Shape of articular surfaces (how close they fit)
- Strength and tension of capsule and ligaments (dependent on position)
- Arrangement and tension of muscles
- Contact with soft parts such as adipose tissue
- Hormones
- Disuse, causing decrease in synovial fluid, flexibility of ligaments and tendons, and muscle atrophy
- Gravity and atmospheric pressure.

Typically, the more stable the joint is, the less is its range of motion and vice versa. Aging is another factor that influences motion due to decreased fluid, thinning of cartilage, shortening of ligaments, and loss of flexibility.

This page titled 8.4D: Stability and Range of Motion at Synovial Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4E: Synovial Joint Movements

Synovial joints allow an individual to achieve a wide range of movements.

Learning Objectives

- Identify the different types of synovial joints

Key Points


- Synovial joints achieve movement at the point of contact of the articulating bones.
- Synovial joints allow bones to slide past each other or to rotate around each other. This produces movements called abduction (away), adduction (towards), extension (open), flexion (close), and rotation.
- There are six types of synovial joints. Some are relatively immobile but more stable than mobile joints.

Key Terms

- **synovial joint**: Also known as a diarthrosis, the most common and most movable type of joint in the body of a mammal.
- **abduction**: The movement that separates a limb or other part from the axis, or middle line, of the body.
- **flexion**: The act of bending a joint. The counteraction of extension.
- **adduction**: The action by which the parts of the body are drawn toward its axis.

A synovial joint, also known as a diarthrosis, is the most common and most movable type of joint in the body of a mammal. Synovial joints achieve movement at the point of contact of the articulating bones. Structural and functional differences distinguish synovial joints from cartilaginous joints (synchondroses and symphyses) and fibrous joints (sutures, gomphoses, and syndesmoses). The main structural differences between synovial and fibrous joints are the existence of capsules surrounding the articulating surfaces of a synovial joint and the presence of lubricating synovial fluid within those capsules (synovial cavities).


Several movements may be performed by synovial joints. Abduction is the movement away from the midline of the body. Adduction is the movement toward the middle line of the body. Extension is the straightening of limbs (increase in angle) at a joint. Flexion is bending the limbs (reduction of angle) at a joint. Rotation is a circular movement around a fixed point.



Body Movements I: Image demonstrating the various joint movements.

There are six types of synovial joints. Some are relatively immobile but more stable than mobile joints. Others have multiple degrees of freedom, but at the expense of greater risk of injury. The six types of joints include:

- Gliding joints: only allow sliding movement
- Hinge joints: allow flexion and extension in one plane
- Pivot joints: allow bone rotation about another bone
- Condyloid joints: perform flexion, extension, abduction, and adduction movements
- Saddle joints: permit the same movement as condyloid joints and combine with them to form compound joints
- Ball and socket joints: allow all movements except gliding



Six Types of Synovial Joints: Image demonstrating the six different types of synovial joints.

This page titled 8.4E: Synovial Joint Movements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

8.4E: Types of Synovial Joints

There are six different types of synovial joint based on their shapes, each allowing a different kind of movement.

Learning Objectives

- Describe the different types of synovial joints

Key Points

- Plane joints are flat and have slipping and gliding properties.
- Hinge joints are formed between the cylindrical end of a bone and the trough-shaped surface of another bone, allowing flexion and extension in one plane.
- Pivot joints are formed between the rounded end of the bone and a sleeve or ring of a bone, allowing up and down and side-to-side movement.
- Condyloid joints occur where an egg-shaped surface of a bone fits into a concave space in another bone, allowing flexion, extension, abduction, and adduction movements (circumduction). The saddle joint resembles a saddle and permits the same movements as the condyloid joints.
- Ball-and-socket joints occur where one bone ends in a spherical head and the other has a round socket, allowing all movements except gliding.
- Saddle joints
  are biaxial, and movements are the same as those for condyloid
  joints; however, no axial
  rotation is possible.

Key Terms

- **ball-and-socket joint**: A joint in which the ball-shaped surface of one rounded bone fits into the cup-like depression of another bone.
- **acromioclavicular joint**: A joint at the top of the shoulder that is the junction between the acromion (a bony process on the scapula) and the clavicle.
- **circumduction**: A conical movement of a body part consisting of a combination of flexion, extension, adduction, and abduction.
- **condyle**: A smooth prominence on a bone where it forms a joint with another bone.
- **Synovial joint**: The most common and most movable type of joint in the body of a mammal.


There are six basic types of synovial joints. Anatomical joints may consist of a combination of two or more joint types. Some synovial joints are relatively immobile but stable. Others have multiple degrees of freedom, but at the expense of greater risk of injury. The types of the synovial joints are based on their shapes and can be classified as plane, hinge, pivot, condyloid, saddle, and ball-and-socket. The following descriptions are in ascending order of mobility:

- The articulating surfaces of the plane joint are usually flat to allow slipping and gliding properties. Examples include the carpals of the wrist and the acromioclavicular joint.
- A hinge joint (ginglymus) is formed when the cylindrical end of a bone fits into a trough-shaped surface of another bone, like that of an elbow joint (between the humerus and the ulna). These joints act as a hinge, allowing flexion and extension in just one plane.
- In a pivot joint, the rounded end of the bone fits into a sleeve or ring of bone. The atlanto-axial joint, proximal radioulnar joint, and distal radioulnar joint are examples of pivot joints.
- The condyloid joint occurs where an egg-shaped surface of a bone fits into a concavity in another bone. Examples include the wrist joint (radiocarpal joint) and the temporomandibular joint. Some classifications make a distinction between condyloid and ellipsoid joints, but both allow flexion, extension, abduction, adduction, and circumduction movements.
- The surface of a saddle joint has both convex and concave areas which resemble a saddle and permit the same movements


as the condyloid joints. The carpometacarpal or trapeziometacarpal joint of the thumb (between the metacarpal and carpal, the trapezium) and the sternoclavicular joint are examples of saddle joints.

- A ball-and-socket joint occurs where one bone ends in a spherical head and the other bone has a round socket. This joint creates the ball-and-socket movement found in such places as the hip and shoulder (glenohumeral). This type of joint allows for all movements except gliding.


The knee joint is an example of a compound joint/modified hinge joint where different types of joints combine. In this example, the condyles of the femur join with condyles of tibia and the saddle joint, where the lower end of the femur joins with the patella.



Plane Joint: The left shoulder and acromioclavicular joints, and the proper ligaments of the scapula.



Ball and Socket Joint: Hip joint: the ball of the femur head fits in the socket of the acetabulum of the pelvis.



Saddle Joint: Sternoclavicular articulation. Anterior view.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- synovial membrane. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/synovial_membrane. **License**: ***CC BY-SA: Attribution-ShareAlike***
- III. Syndesmology. **Provided by**: Saylor. **Located at**: http://www.saylor.org/site/wp-conten...BIO302-ch3.pdf. **License**: ***Public Domain: No Known Copyright***
- Synovial joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synovial_joint. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diarthrosis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/diarthrosis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- articulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/articulation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- articular cartilage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/articular_cartilage. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synovial fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/synovial_fluid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Joint.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Synovial joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synovial_joint. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anastomosis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/anastomosis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteomyelitis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osteomyelitis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Joint.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray526. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray526.png. License: ***Public Domain: No Known Copyright***
- Synovial joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synovial_joint. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Bursae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bursae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Tendons. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Tendons. **License**: *CC BY-SA: Attribution-ShareAlike*
- tendon. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tendon. **License**: *CC BY-SA: Attribution-ShareAlike*
- connective tissue. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/connective_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- synovial fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/synovial_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Joint.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray526. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray526.png. **License**: *Public Domain: No Known Copyright*
- Bursa.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:908_Bursa.jpg. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ndon_(PSF).svg. **License**: *Public Domain: No Known Copyright*
- Tendons. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Tendons. **License**: *CC BY-SA: Attribution-ShareAlike*
- pronation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pronation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Plantar flexion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Plantar_flexion. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsiflexion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsiflexion. **License**: *CC BY-SA: Attribution-ShareAlike*
- eversion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/eversion. **License**: *CC BY-SA: Attribution-ShareAlike*
- supination. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/supination. **License**: *CC BY-SA: Attribution-ShareAlike*
- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Joint.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray526. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray526.png. **License**: *Public Domain: No Known Copyright*
- Bursa.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:908_Bursa.jpg. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a1/Tendon_(PSF).svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...les-tendon.jpg. **License**: *Public Domain: No Known Copyright*
- Synovial joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synovial_joint%23Movements_possible. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...tion/abduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...tion/adduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- flexion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/flexion. **License**: *CC BY-SA: Attribution-ShareAlike*
- Joint. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Joint.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray526. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray526.png. **License**: *Public Domain: No Known Copyright*
- Bursa.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:908_Bursa.jpg. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a1/Tendon_(PSF).svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/3c/Achilles-tendon.jpg. **License**: *Public Domain: No Known Copyright*
- Body Movements. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Body_Movements_I.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Synovial_Joints.jpg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/

File:909_Types_of_Synovial_Joints.jpg. **License**: *CC BY: Attribution*
- Synovial joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synovial_joint%23Types**. **License**: *CC BY-SA: Attribution-ShareAlike*
- General Anatomy/Skeletal System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/General...ynovial_joints**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circumduction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/circumduction**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acromioclavicular Joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acromioclavicular_joint**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Synovial joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synovial%20joint**. **License**: *CC BY-SA: Attribution-ShareAlike*
- condyle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/condyle**. **License**: *CC BY-SA: Attribution-ShareAlike*
- ball-and-socket joint. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ball-and-socket_joint**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Joint.png**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray526. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray526.png**. **License**: *Public Domain: No Known Copyright*
- Bursa.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:908_Bursa.jpg**. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/a/a1/Tendon_(PSF).svg**. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3c/Achilles-tendon.jpg**. **License**: *Public Domain: No Known Copyright*
- Body Movements. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Body_Movements_I.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Synovial_Joints.jpg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:909_Types_of_Synovial_Joints.jpg**. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/9f/Hip.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sternoclavical Articulation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3f/Gray325.png**. **License**: *Public Domain: No Known Copyright*
- Plane Joint. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3b/Gray326.png**. **License**: *Public Domain: No Known Copyright*

This page titled 8.4E: Types of Synovial Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9: MUSCULAR SYSTEM

- Topic hierarchy

9.10: Muscles of the Lower Limb

9.10A: Muscles that Cause Movement at the Hip Joint

9.10B: Muscles that Cause Movement at the Knee Joint

9.10C: Muscles that Cause Movement at the Foot

9.1: Introduction to the Nervous System

9.1A: Structure and Function of the Muscular System

9.1B: Slow-Twitch and Fast-Twitch Muscle Fibers

9.1C: Sliding Filament Model of Contraction

9.1D: ATP and Muscle Contraction

9.1E: Control of Muscle Tension

9.2: Smooth Muscle

9.2A: Skeletal Muscle Fibers

9.3: Control of Muscle Tension

9.3A: Force of Muscle Contraction

9.3B: Velocity and Duration of Muscle Contraction

9.3C: Motor Units

9.3D: Muscle Tone

9.3E: Types of Muscle Contractions: Isotonic and Isometric

9.4: Muscle Metabolism

9.4A: Muscle Metabolism

9.4B: Muscle Fatigue

9.5: Exercise and Skeletal Muscle Tissue

9.5A: Impacts of Exercise on Muscles

9.6: Overview of the Muscular System

9.6A: Interactions of Skeletal Muscles

9.6B: How Skeletal Muscles Are Named

9.6C: How Skeletal Muscles Produce Movements

9.6D: Muscle Attachment Sites

9.6E: Arrangement of Fascicles

9.6F: Lever Systems

9.7: Head and Neck Muscles

9.7A: Facial Muscles

9.7B: Chewing Muscles
9.7C: Neck Muscles

9.8: Trunk Muscles
9.8A: Anterior Muscles
9.8B: Posterior Muscles

9.9: Muscles of the Upper Limb
9.9A: Muscles of the Humerus that Act on the Forearm
9.9B: Muscles of the Wrist and Hand
9.9C: Muscles of the Shoulder
9.9D: Muscles that Cause Movement at the Ankle

This page titled 9: Muscular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.10: MUSCLES OF THE LOWER LIMB

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

9.10A: Muscles that Cause Movement at the Hip Joint

9.10B: Muscles that Cause Movement at the Knee Joint

9.10C: Muscles that Cause Movement at the Foot

This page titled 9.10: Muscles of the Lower Limb is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.10A: Muscles that Cause Movement at the Hip Joint

The four main groups of hip muscles are gluteal, adductor, iliopsoas, and lateral rotator, defined by the type of movement they mediate.

Learning Objectives

- Differentiate among the muscles involved in moving the hip joint

Key Points


- The gluteus maximus extends the hip, while the gluteus medius and minimus are involved in hip rotation and abduction (moving hip out from the midline).
- The adductor group (adductor brevis, longus, and magnus along with petineus and gracilis) moves the femur towards the midline from an abducted position.
- The iliopsoas group of muscles (iliacus and psoas major) is responsible for hip flexion.
- The lateral rotator group of muscles (externus and internus obturators, the piriformis, the superior and inferior gemelli, and the quadratus femoris) turns the anterior surface of the femur outward. This motion is aided by the gluteus maximus and the adductor magnus.

Key Terms

- **adductor group**: The adductor brevis, adductor longus, adductor magnus, pectineus, and gracilis.
- **lateral rotator group**: The externus and internus obturators, the piriformis, the superior and inferior gemelli, and the quadratus femoris.
- **gluteal group**: The gluteus maximus, gluteus medius, gluteus minimus, and tensor fasciae latae.
- **iliopsoas group**: The iliacus and psoas major.

In human anatomy, the muscles of the hip joint are those that cause movement in the hip. Hip joint muscles are divided into four groups according to their orientation and function. Movement at the hip is similar to that of the shoulder joint, but due to increased weight-bearing requirements the range of potential movements is reduced.

Gluteal Group



Key muscles of the hip: The gluteus maximus can be seen at the top, cut away to expose the underlying muscles.

Muscles in the gluteal group are superficially located and act mainly to abduct and extend the thigh at the hip.


- *Gluteus Maximus*: The gluteus maximus is the largest of the gluteal muscles and gives structure to the buttocks.
  - Attachments: Originates from the posterior of the pelvis and coccyx (tailbone) and attaches to the femur.
  - Actions: Extends of the thigh and assists with rotation. Is only used when the generation of force is required (e.g. when climbing).
- *Gluteus Medius*: The fan-shaped gluteus medius muscle lies between the gluteus maximus and minimus and performs a similar function to the gluteus minimus.
  - Attachments: Originates from the posterior of the pelvis and attaches to the femur.
  - Actions: Abducts and medially rotates the thigh and fixes the pelvis during walking.
- *Gluteus Minimus*: The gluteus minimus is the deepest and smallest of the superficial gluteal muscles and performs a similar function to the gluteus medius.
  - Attachments: Originates from the pelvis and attaches to the femur.
  - Actions: Abducts and medially rotates the thigh and fixes the pelvis during walking.

Lateral Rotator Group

The muscles of the lateral rotator group are deeply located and as the name suggests, act to laterally rotate the thigh at the hip. All of the lateral rotator group muscles originate from the pelvis and attach to the femur.

- *Piriformis*: The piriformis is the most superior of the lateral rotator group muscles.
  - Actions: Lateral rotation and abduction of the thigh at the hip.
- *Obturator Internus*: The obturator internus lines the internal wall of the pelvis.
  - Actions: Lateral rotation and abduction of the thigh at the hip.
- *Gemelli*: The gemelli are two (superior and inferior) narrow and triangular muscles, separated by the obturator internus tendon.
  - Actions: Lateral rotation and abduction of the thigh at the hip.
- *Quadratus Femoris*: The quadratus femoris is a flat, square-shaped muscle (actually composed of four distinct muscles). It is the most inferior of the lateral rotator group muscles, located below the gemelli and obturator internus.
  - Actions: Lateral rotation of the thigh at the hip, plays a major role in extension of the lower leg at the knee as well.

Adductor Group



(a) Adductor Group Muscles and (b) Key muscles associated with movement at the hip: The deep-lying adductor group muscles originate from the pubis and attach to the length of the femur. The iliacus and psoas major comprise the iliopsoas group and can be seen at the pelvis and lower spine.

The five muscles of the adductor group are responsible for the adduction of the thigh, although several have additional functions.

- *Adductor Longus*: The adductor longus is a large, flat muscle covering the adductor magnus and adductor brevis.
  - Attachments: Originates from the pubis and broadly attaches to the femur.
  - Actions: Adduction and medial rotation of the thigh.
- *Adductor Magnus*: The adductor magnus is the largest and most posterior of the adductor group muscles.
  - Attachments: Originates from the pubis and attaches to the femur.
  - Actions: Adducts, flexes, and extends the thigh.
- *Adductor Brevis*: The adductor brevis is a short muscle lying underneath the adductor longus.
  - Attachments: Originates from the pubis and attaches to the femur.
  - Actions: Adduction of the thigh.
- *Obturator Externus*: This is one of the smaller muscles of the medial thigh, and it is located most superiorly.
  - Attachments: Originates from the pubis and attaches to the femur.
  - Actions: Laterally rotates the thigh.
- *Gracilis*: The gracilis is the most superficial and medial of the adductor group muscles. Crossing both the hip and knee joints, it can induce movement at both the hip and knee.
  - Attachments: Originates from the pubis and attaches to the tibia.
  - Actions: Adduction of the thigh at the hip, and flexing of the thigh at the knee.

Other Muscles

There are several other muscles that induce movement around the hip joint.

- *Psoas Major*: The psoas major is located deep in the back near the midline immediately adjacent to the spine. The iliacus and psoas major comprise the iliopsoas group.
  - Attachments: Originates from the base of the spine, combining with the iliacus to attach to the femur.
  - Actions: Flexing of the thigh at the hip joint.
- *Iliacus*: The iliacus muscle is a large, fan-shaped muscle which lines the interior of the pelvis. The iliacus and psoas major comprise the iliopsoas group.
  - Attachments: Originates from the pelvis and the base of the spine, combining with the psoas major to attach to the femur.

    - Actions: Flexing of the thigh at the hip joint.
- *Sartorius*: The sartorius is a long thin muscle in the thigh, the longest muscle in the body.
    - Attachments: Originates from the pelvis and attaches to the tibia.
    - Actions: Flexing, abducting and rotation of the thigh at the hip joint.
- *Pectineus*: The pectineus muscle is a large flat muscle found in the thigh.
    - Attachments: Originates from the pelvis and attaches to the femur.
    - Actions: Adduction and flexing at the thigh at the hip joint.
- *Biceps Femoris*: A similar muscle to the biceps brachii in the upper arm, also double-headed. Two synergistic muscles are associated with the biceps femoris, the semitendinosus, and the semimembranosus.
    - Attachments: Originates from the pelvis and femur and attaches to the fibula.
    - Actions: Extends and laterally rotates at the hip. The main action is flexing of the lower leg at the knee.

KEY MOVEMENTS

- **Extension** (bringing thigh behind body) Produced by the gluteus maximus, adductor magnus, and biceps femoris. *Flexion* (extending thigh to front of body): Produced by the gracilis, psoas major, iliacus, and pectineus.
- **Abduction** (moving thigh laterally away from pelvis): Produced by the gluteus medius and minimus, obturator externus, gemelli, and sartorius.
- **Adduction** (returning thigh to midline): Produced by the adductor group of muscles.
- **Rotation** (rotation of the thigh around the hip joint): Produced by the lateral rotator group of muscles and the biceps femoris, sartorius, and gluteus medius and minimus.

This page titled 9.10A: Muscles that Cause Movement at the Hip Joint is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.10B: Muscles that Cause Movement at the Knee Joint

Three sets of muscles (popliteus, quadriceps and hamstrings) allow for movement, balance, and stability at the knee joint.

Learning Objectives

- Differentiate among the muscles that enable the knee joint to move

Key Points

- At full extension, the tibia and femur “lock” into position, providing stability in the leg and improving load-bearing capacity. The popliteus muscle at the back of the leg unlocks the knee by rotating the femur on the tibia, allowing flexion of the joint.
- The quadriceps femoris muscle group (rectus femoris, vastus lateralis, vastus medius, and vastus intermedius) crosses the knee via the patella and acts to extend the leg.
- The hamstring group muscles (semitendinosus, semimembranosus, and biceps femoris) flex the knee and extend the hip.

Key Terms

- **hamstring group**: A group of three muscles found in the posterior region of the thigh, responsible for flexing of the lower leg at the knee.
- **quadriceps femoris**: A group of four muscles found in the anterior region of the thigh, responsible for extension of the lower leg at the knee.
- **popliteus**: A muscles located behind the knee which “unlocks” the fully extended knee joint allowing for flexion.

The knee joint allows for movement of the lower leg relative to the thigh across the knee joint. The knee joint is in fact comprised of two joints: the tibiofemoral joint between the femur and tibia, which is the weight-bearing knee joint, and the patellofemoral joint, which joins the patella (kneecap) with the femur.


The tibiofemoral joint is relatively weak and easily damaged, so it relies on muscles and ligaments to ensure stability. When the knee is fully extended the femur rotates slightly on the tibia to lock the joint into place, allowing for efficient load bearing.

The patella is the attachment point for the quadriceps femoris muscle and is the attached by a ligament to the tibia. This increases the leverage afforded to the quadriceps femoris muscle, thus increasing its efficiency when extending the lower leg. The patella additionally protects the knee joint from damage.

The patellofemoral has two key functions: increasing leverage of the quadriceps tendon to improve muscle stability and protecting the knee joint from damage.

Muscles that generate movement across the knee are mainly located in the thigh and can be split into anterior and posterior compartments. The popliteus muscle, located in the lower leg, is responsible for “unlocking” the knee joint after extension.

Anterior Muscles of the Thigh



(a) Posterior muscles of the thigh and (b) posterior region of the lower leg: The biceps femoris and synergistic semitendinosus and the semimembranosus muscles are responsible for flexing of the lower leg at the knee. Posterior view of muscles of the lower leg, the popliteus can be seen at the top located behind the knee.

There are four muscles in anterior region of the thigh. The pectineus and iliopsoas muscles are responsible for movement at the hip and are discussed elsewhere.

- *Sartorius*: The sartorius, a thin muscle in the thigh, the is the body's longest muscle.
  - Attachments: Originates from the pelvis and attaches to the tibia.
  - Actions: Flexing of the lower leg at the knee joint.
- *Quadriceps Femoris*: The quadriceps femoris is actually composed of four muscles that comprise the front of the thigh: three deep-lying vastus muscles (lateralis, intermedius, and medialis) and the rectus femoris which covers them. All four muscles are the key extensors of the lower leg at the knee joint and also stabilize and protect the patella.
  - Attachments: The vastus lateralis, intermedius, and medialis originate from the femur and attach to the patella. The rectus femoris originates from the pelvis and attaches to the patella.
  - Actions: Extends the lower leg at the knee joint and stabilizes the patella. The rectus femoris additionally facilitates rotation at the hip.

Posterior Muscles of the Thigh

There are three muscles in the posterior compartment of the thigh: the biceps femoris and two synergistic muscles (the semitendinosus and semimembranosus). These muscles are sometimes termed the hamstring group. The posterior region of the thigh displays similarity with the anterior region of the upper arm in both structure and function.

- *Biceps Femoris*: A similar muscle to the biceps brachii in the upper arm and also double-headed. Two synergistic muscles are associated with the biceps femoris, the semitendinosus and the semimembranosus.
  - Attachments: Originates from the pelvis and femur and attaches to the fibula.
  - Actions: Extends and laterally rotates at the hip, main action is flexing of the lower leg at the knee.

Other Muscles

- *Popliteus*: The popliteus is located behind the knee joint and acts to "unlock" the knee by rotating the femur on the tibia allowing for the lower leg to be flexed.
  - Attachments: Originates from the posterior of the tibia and attaches to the femur.

- Actions: Laterally rotates the femur on the tibia “unlocking” the knee joint so that flexion can occur.

KEY MOVEMENTS

- **Extension:** Produced by the sartorius and quadriceps femoris group of muscles.
- **Flexion:** Produced by the biceps femoris, semitendinosus, and semimembranosus muscles. The popliteus muscle facilitates this movement by unlocking the fully extended knee joint.
- **Rotation:** The knee joint allows for slight rotation when flexed, which is produced by the biceps femoris, semitendinosus, semimembranosus, gracilis, and sartorius.

This page titled 9.10B: Muscles that Cause Movement at the Knee Joint is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.10C: Muscles that Cause Movement at the Foot

Movement of the foot and toes requires the action of many muscles.

Learning Objectives

- Differentiate among the muscles that cause feet to move

Key Points

- The hallux or large toe is extended by the extensor hallucis brevis on the top of the foot.
- The flexor hallucis brevis and abductor hallucis flex and abduct the big toe. The adductor hallucis adducts the big toe.
- The remaining toes are flexed by the flexor digitorum longus, lumbricals, flexor digitorum brevis, and quadratus plantae.
- The little toe is also controlled by the flexor digiti minimi and abductor digiti minimi.
- The toes (other than the big toe) are extended by the extensor digitorum brevis.
- The dorsal and plantar interossei are muscles between the metatarsals that help maintain the foot's arch. The also aid in flexion and extension.

Key Terms

- **plantar**: The sole of the foot.
- **dorsal**: The top surface of foot.

The foot plays a fundamental role in maintaining upright walking posture and balance, so it must provide a strong base while allowing for small, accurate movements. The foot can be split into dorsal and plantar compartments, the latter of which contains significantly more muscles.

Dorsal Compartment


Although many extrinsic muscles attach within the dorsal compartment, there are only three intrinsic muscles which act on the foot and two extrinsic muscles which act on the foot rather than the ankle.

- *Extensor Digitorum Longus*: The extensor digitorum longus is a deep-lying extrinsic muscle that lies runs down the length of the tibia.
  - Attachments: Originates from the tibia and transitions into a tendon, passes into the foot, splits into four, and attaches to the toes.
  - Actions: Extension of the toes.
- *Extensor Digitorum Brevis*: The extensor digitorum brevis muscle is a deep-lying intrinsic muscle lying beneath the tendon of the extensor digitorum longus.
  - Attachments: Originates from the heel and attaches to the toes.
  - Actions: Extension of the toes.
- *Extensor Hallucis Longus*: The extensor hallucis longus is a deep-lying extrinsic muscle lying beneath the extensor digitorum longus.
  - Attachments: Originates from the fibula and attaches to the big toe.
  - Actions: Extension of the big toe.
- *Extensor Hallucis Brevis*: The extensor hallucis brevis muscle is an intrinsic muscle of the foot lying between the extensor digitorum longus and lateral to extensor hallucis longus.
  - Attachments: Originates from the heel and attaches to the big toe.
  - Actions: Extension of the big toe.
- *Dorsal Interossei*: There are four dorsal interossei located between the metatarsals. Each arises from two metatarsals.
  - Attachments: Originates from the sides of metatarsals one to five (big toe to little toe). The first muscle attaches to the medial side of the phalanx of the second toe. The second to fourth interossei attach to the lateral sides of the phalanxes

of toes two to four.
- Actions: Abduct and flexes digits two to four.

Plantar Compartment

The muscles of the plantar compartment play a key role in stabilizing the arch of the foot and controlling the digits, providing the necessary strength and fine movements to maintain balance and posture and promote walking.



Plantar View of Foot: The muscles, tendons, and ligaments of the sole of the foot with flexor digitorum brevis shown in red.

- *Abductor Hallucis*: The abductor hallucis muscle is located on the medial side of the sole.
  - Attachments: Originates from the heel and attaches to the base of the big toe.
  - Actions: Abducts and flexes the big toe.
- *Flexor Digitorum Brevis*: The flexor digitorum brevis muscle is located laterally to the abductor hallucis and lies in the middle of the sole of the foot.
  - Attachments: Originates from the heel and attaches to all the toes excluding the little toe.
  - Actions: Flexes the toes at the proximal interphalangeal (between the phalanges preoxmales and phalanges mediae) joints.
- *Abductor Digiti Minimi*: The abductor digiti minimi muscle is located on the lateral side of the foot.
  - Attachments: Originates from the heel and attaches to the little toe.
  - Actions: Abducts and flexes the little toe.
- *Quadratus Plantae*: The quadratus plantae muscle is located superior to the flexor digitorum longus tendons.
  - Attachments: Originates from the heel and attaches to the tendons of flexor digitorum longus.
  - Actions: Assists flexor digitorum longus in flexing the lateral four toes.
- *Lumbricals*: The lumbricals are four small skeletal muscles, accessory to the tendons of the flexor digitorum longus and numbered from the medial side of the foot.
  - Attachments: Originates from the tendons of flexor digitorum longus. Attaches to the toes.
  - Actions: Flexes at the metatarsophalangeal (between the metatarsals and phalanges proximale) joints, while extending the interphalangeal joints (between the phalanges proximale and phalanges mediae).
- *Flexor Hallucis Brevis*: The flexor hallucis brevis muscle is located on the medial side of the foot.
  - Attachments: Originates from the plantar surfaces of the cuboid and lateral cuneiform bones in the sole of the foot and attaches to the big toe.
  - Actions: Flexes the big toe.
- *Adductor Hallucis*: The adductor hallucis muscle is located laterally to the flexor hallucis brevis.
  - Attachments: The adductor hallucis originates from the bases of the first four metatarsals, and also from the plantar ligaments before attaching to the big toe.
  - Actions: Adduct the big toe and maintains the arch of the foot.
- *Plantar Interossei*: The three plantar interossei lie beneath rather than between the metatarsal bones. Each is connected with

one metatarsal bone.
    - Attachments: Originates from the medial side of metatarsals three to five. Attaches to the medial sides of the phalanges of digits three to five.
    - Actions: Adducts and flexes digits three to five.
- *Flexor Digiti Minimi Brevis*: Similar to the interossei in structure, the flexor digiti minimi brevis muscle is located on the lateral side of the foot underneath the metatarsal of the little toe.
    - Attachments: Originates from the base of the fifth metatarsal and attaches to the base of phalanx of the little toe
    - Actions: Flexes the little toe.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Muscles of the hip. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscles_of_the_hip**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology.../gluteal-group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...liopsoas-group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...-rotator-group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...adductor-group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- image433.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus433.html**. **License**: ***Public Domain: No Known Copyright***
- image430.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus430.html**. **License**: ***Public Domain: No Known Copyright***
- Hamstrings. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hamstrings**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Popliteus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Popliteus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Patellar reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Patellar_reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Quadriceps femoris muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Quadric...femoris_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- quadriceps. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/quadriceps**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- popliteus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/popliteus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- patellar reflex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/patellar_reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- image433.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus433.html**. **License**: ***Public Domain: No Known Copyright***
- image430.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus430.html**. **License**: ***Public Domain: No Known Copyright***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...c1/Gray439.png**. **License**: ***Public Domain: No Known Copyright***
- Ankle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ankle%23Ligaments**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Fibularis longus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibularis_longus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tibialis anterior muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tibialis_anterior_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ltoid-ligament**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- inversion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/inversion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ialis-anterior**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...bular-ligament**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- image433.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus433.html**. **License**: ***Public Domain: No Known Copyright***
- image430.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus430.html**. **License**: ***Public Domain: No Known Copyright***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...c1/Gray439.png**. **License**: ***Public Domain: No Known Copyright***
- image438.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus438.html**. **License**: ***Public Domain: No Known Copyright***
- Tibialis anterior 2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ti...anterior_2.png**. **License**: ***Public Domain: No Known Copyright***
- Extensor hallucis brevis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extenso..._brevis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lumbrical muscle (foot). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lumbrical_muscle_(foot)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Abductor digiti minimi muscle (foot). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Abducto..._muscle_(foot)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor digitorum brevis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_..._brevis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Dorsal interossei of the foot. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Dorsal_...ei_of_the_foot**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor digiti minimi brevis muscle (foot). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_..._muscle_(foot)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Plantar interossei. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Plantar_interossei**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Abductor hallucis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Abductor_hallucis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adductor hallucis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adductor_hallucis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Quadratus plantae muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Quadratus_plantae_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor hallucis brevis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_..._brevis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hallux. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hallux**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plantar. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plantar**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dorsal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/dorsal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- image433.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus433.html**. **License**: ***Public Domain: No Known Copyright***
- image430.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus430.html**.

License: ***Public Domain: No Known Copyright***

- image434.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus434.html**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...c1/Gray439.png**. **License**: ***Public Domain: No Known Copyright***
- image438.gif. **Provided by**: Anatomy of the Human Body. **Located at**: **http://www.bartleby.com/107/illus438.html**. **License**: ***Public Domain: No Known Copyright***
- Tibialis anterior 2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Tibialis_anterior_2.png**. **License**: ***Public Domain: No Known Copyright***
- Flexor Digitorum Brevis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/69/Flexor_digitorum_brevis.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 9.10C: Muscles that Cause Movement at the Foot is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1: INTRODUCTION TO THE NERVOUS SYSTEM



- Topic hierarchy

9.1A: Structure and Function of the Muscular System

9.1B: Slow-Twitch and Fast-Twitch Muscle Fibers

9.1C: Sliding Filament Model of Contraction

9.1D: ATP and Muscle Contraction

9.1E: Control of Muscle Tension

This page titled 9.1: Introduction to the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1A: Structure and Function of the Muscular System

The muscular system controls numerous functions, which is possible with the significant differentiation of muscle tissue morphology and ability.

Learning Objectives

- Describe the three types of muscle tissue

Key Points


- The muscular system is responsible for functions such as maintenance of posture, locomotion, and control of various circulatory systems.
- Muscle tissue can be divided functionally (voluntarily or involuntarily controlled) and morphologically (striated or non-striated).
- These classifications describe three distinct muscle types: skeletal, cardiac and smooth. Skeletal muscle is voluntary and striated, cardiac muscle is involuntary and striated, and smooth muscle is involuntary and non-striated.

Key Terms

- **myofibril**: A fiber made up of several myofilaments that facilitates the generation of tension in a myocyte.
- **myofilament**: A filament composed of either multiple myosin or actin proteins that slide over each other to generate tension.
- **myosin**: A motor protein which forms myofilaments that interact with actin filaments to generate tension.
- **actin**: A protein which forms myofilaments that interact with myosin filaments to generate tension.
- **striated**: The striped appearance of certain muscle types in which myofibrils are aligned to produce a constant directional tension.
- **voluntary**: A muscle movement under conscious control (e.g. deciding to move the forearm).
- **involuntary**: A muscle movement not under conscious control (e.g. the beating of the heart).
- **myocyte**: A muscle cell.

The Musculoskeletal System

The muscular system is made up of muscle tissue and is responsible for functions such as maintenance of posture, locomotion and control of various circulatory systems. This includes the beating of the heart and the movement of food through the digestive system. The muscular system is closely associated with the skeletal system in facilitating movement. Both voluntary and involuntary muscular system functions are controlled by the nervous system.



The muscular system: Skeletal muscle of the muscular system is closely associated with the skeletal system and acts to maintain posture and control voluntary movement.

Muscle is a highly-specialized soft tissue that produces tension which results in the generation of force. Muscle cells, or myocytes, contain myofibrils comprised of actin and myosin myofilaments which slide past each other producing tension that changes the shape of the myocyte. Numerous myocytes make up muscle tissue and the controlled production of tension in these cells can generate significant force.

Muscle tissue can be classified functionally as voluntary or involuntary and morphologically as striated or non-striated. Voluntary refers to whether the muscle is under conscious control, while striation refers to the presence of visible banding within myocytes caused by the organization of myofibrils to produce constant tension.

Types of Muscle

The above classifications describe three forms of muscle tissue that perform a wide range of diverse functions.

Skeletal Muscle

Skeletal muscle mainly attaches to the skeletal system via tendons to maintain posture and control movement. For example, contraction of the biceps muscle, attached to the scapula and radius, will raise the forearm. Some skeletal muscle can attach directly to other muscles or to the skin, as seen in the face where numerous muscles control facial expression.

Skeletal muscle is under voluntary control, although this can be subconscious when maintaining posture or balance. Morphologically skeletal myocytes are elongated and tubular and appear striated with multiple peripheral nuclei.

Cardiac Muscle Tissue


Cardiac muscle tissue is found only in the heart, where cardiac contractions pump blood throughout the body and maintain blood pressure.

As with skeletal muscle, cardiac muscle is striated; however it is not consciously controlled and so is classified as involuntary. Cardiac muscle can be further differentiated from skeletal muscle by the presence of intercalated discs that control the synchronized contraction of cardiac tissues. Cardiac myocytes are shorter than skeletal equivalents and contain only one or two centrally located nuclei.

Smooth Muscle Tissue

Smooth muscle tissue is associated with numerous organs and tissue systems, such as the digestive system and respiratory system. It plays an important role in the regulation of flow in such systems, such as aiding the movement of food through the digestive system via peristalsis.

Smooth muscle is non-striated and involuntary. Smooth muscle myocytes are spindle shaped with a single centrally located nucleus.



Skeletal muscle | Smooth muscle | Cardiac muscle

Types of muscle: The body contains three types of muscle tissue: skeletal muscle, smooth muscle, and cardiac muscle, visualized here using light microscopy. Visible striations in skeletal and cardiac muscle are visible, differentiating them from the more randomised appearance of smooth muscle.

This page titled 9.1A: Structure and Function of the Muscular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1B: Slow-Twitch and Fast-Twitch Muscle Fibers

Skeletal muscle contains different fibers which allow for both rapid short-term contractions and slower, repeatable long-term contractions.

Learning Objectives

- Describe the different types of skeletal muscle fibers and their respective functions

Key Points

- Slow-twitch fibers rely on aerobic respiration to fuel muscle contractions and are ideal for long term endurance.
- Fast-twitch fibers rely on anaerobic respiration to fuel muscle contractions and are ideal for quick contractions of short duration.

Key Terms

- **aerobic**: A combination of glycolysis and the Krebs cycle, an efficient but slow way of producing ATP.
- **anaerobic**: Glycolysis alone, an inefficient but quick way of producing ATP with pyruvate converted to lactate.
- **glycolysis**: The breakdown of glucose (or other carbohydrates) by enzymes, generating ATP and pyruvate.
- **slow-twitch**: Type I fibers characterized as muscles with long contraction duration, associated with endurance.
- **Krebs cycle**: A sequence of reactions which converts pyruvate into carbon dioxide and water, generating further adenosine triphosphate (ATP).
- **fast-twitch**: Type II fibers which are characterized by fast muscle contractions of short duration.

Skeletal muscle fibers can be further subdivided into slow and fast-twitch subtypes depending on their metabolism and corresponding action. Most muscles are made up of combinations of these fibers, although the relative number substantially varies.

Slow Twitch (Type 1)

Slow-twitch fibers are designed for endurance activities that require long-term, repeated contractions, like maintaining posture or running a long distance. The ATP required for slow-twitch fiber contraction is generated through aerobic respiration (glycolysis and Krebs cycle), whereby 30 molecules of ATP are produced from each glucose molecule in the presence of oxygen. The reaction is slower than anaerobic respiration and thus not suited to rapid movements, but much more efficient, which is why slow-twitch muscles do not tire quickly. However, this reaction requires the delivery of large amounts of oxygen to the muscle, which can rapidly become rate-limiting if the respiratory and circulatory systems cannot keep up.

Due to their large oxygen requirements, slow-twitch fibers are associated with large numbers of blood vessels, mitochondria, and high concentrations of myoglobin, an oxygen-binding protein found in the blood that gives muscles their reddish color. One muscle with many slow-twitch fibers is the soleus muscle in the leg (~80% slow-twitch), which plays a key role in standing.

Fast Twitch (Type II)

Fast-twitch fibers are good for rapid movements like jumping or sprinting that require fast muscle contractions of short duration. Unlike slow-twitch fibers, fast twitch-fibers rely on anaerobic respiration (glycolysis alone) to produce two molecules of ATP per molecule of glucose. While much less efficient than aerobic respiration, it is ideal for rapid bursts of movement since it is not rate limited by need for oxygen. Lactate (lactic acid), a byproduct of anaerobic respiration, accumulates in the muscle tissue reducing the pH (making it more acidic, and producing the stinging feeling in muscles when exercising). This inhibits further anaerobic respiration. While this may seem counter-intuitive, it is a feedback cycle in place to protect the muscles from over-exertion and resultant damage.

As fast-twitch fibers generally do not require oxygenation, they contain fewer blood vessels and mitochondria than slow-twitch fibers and less myoglobin, resulting in a paler color. Muscles controlling eye movements contain high numbers of fast-twitch fibers (~85% fast-twitch).

Determination and Alteration of Muscle Type

While there is evidence that each person has a unique proportion of fast-twitch versus slow-twitch muscles determined by genetics, more research is required. Regardless, repeated exercise that prioritizes one type of muscle fiber use over the other can lead to improvements in an individual's ability to perform that activity through alterations in the number and composition of fibers associated with improvements in the respiratory and circulatory systems.

This page titled 9.1B: Slow-Twitch and Fast-Twitch Muscle Fibers is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1C: Sliding Filament Model of Contraction

In the sliding filament model, the thick and thin filaments pass each other, shortening the sarcomere.

Learning Objectives

- Describe the sliding filament model of muscle contraction

Key Points

- The sarcomere is the region in which sliding filament contraction occurs.
- During contraction, myosin myofilaments ratchet over actin myofilaments contracting the sarcomere.
- Within the sarcomere, key regions known as the I and H band compress and expand to facilitate this movement.
- The myofilaments themselves do not expand or contract.

Key Terms

- **I-band**: The area adjacent to the Z-line, where actin myofilaments are not superimposed by myosin myofilaments.
- **A-band**: The length of a myosin myofilament within a sarcomere.
- **M-line**: The line at the center of a sarcomere to which myosin myofilaments bind.
- **Z-line**: Neighbouring, parallel lines that define a sarcomere.
- **H-band**: The area adjacent to the M-line, where myosin myofilaments are not superimposed by actin myofilaments.

Movement often requires the contraction of a skeletal muscle, as can be observed when the bicep muscle in the arm contracts, drawing the forearm up towards the trunk. The sliding filament model describes the process used by muscles to contract. It is a cycle of repetitive events that causes actin and myosin myofilaments to slide over each other, contracting the sarcomere and generating tension in the muscle.

Sarcomere Structure

To understand the sliding filament model requires an understanding of sarcomere structure. A sarcomere is defined as the segment between two neighboring, parallel Z-lines. Z lines are composed of a mixture of actin myofilaments and molecules of the highly elastic protein titin crosslinked by alpha-actinin. Actin myofilaments attach directly to the Z-lines, whereas myosin myofilaments attach via titin molecules.

Surrounding the Z-line is the I-band, the region where actin myofilaments are not superimposed by myosin myofilaments. The I-band is spanned by the titin molecule connecting the Z-line with a myosin filament.


The region between two neighboring, parallel I-bands is known as the A-band and contains the entire length of single myosin myofilaments. Within the A-band is a region known as the H-band, which is the region not superimposed by actin myofilaments. Within the H-band is the M-line, which is composed of myosin myofilaments and titin molecules crosslinked by myomesin.

Titin molecules connect the Z-line with the M-line and provide a scaffold for myosin myofilaments. Their elasticity provides the underpinning of muscle contraction. Titin molecules are thought to play a key role as a molecular ruler maintaining parallel alignment within the sarcomere. Another protein, nebulin, is thought to perform a similar role for actin myofilaments.

Model of Contraction

The molecular mechanism whereby myosin and acting myofilaments slide over each other is termed the cross-bridge cycle. During muscle contraction, the heads of myosin myofilaments quickly bind and release in a ratcheting fashion, pulling themselves along the actin myofilament.

At the level of the sliding filament model, expansion and contraction only occurs within the I and H-bands. The myofilaments themselves do not contract or expand and so the A-band remains constant.



The sarcomere and the sliding filament model of contraction: During contraction myosin ratchets along actin myofilaments compressing the I and H bands. During stretching this tension is release and the I and H bands expand. The A-band remains constant throughout as the length of the myosin myofilaments does not change.

The amount of force and movement generated generated by an individual sarcomere is small. However, when multiplied by the number of sarcomeres in a myofibril, myofibrils in a myocyte and myocytes in a muscle, the amount of force and movement generated is significant.

This page titled 9.1C: Sliding Filament Model of Contraction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1D: ATP and Muscle Contraction

ATP is critical for muscle contractions because it breaks the myosin-actin cross-bridge, freeing the myosin for the next contraction.

Learning Objectives

- Discuss how energy is consumed during movement


Key Points

- ATP prepares myosin for binding with actin by moving it to a higher- energy state and a “cocked” position.
- Once the myosin forms a cross-bridge with actin, the Pi disassociates and the myosin undergoes the power stroke, reaching a lower energy state when the sarcomere shortens.
- ATP must bind to myosin to break the cross-bridge and enable the myosin to rebind to actin at the next muscle contraction.

Key Terms

- **M-line**: the disc in the middle of the sarcomere, inside the H-zone
- **troponin**: a complex of three regulatory proteins that is integral to muscle contraction in skeletal and cardiac muscle, or any member of this complex
- **ATPase**: a class of enzymes that catalyze the decomposition of ATP into ADP and a free phosphate ion, releasing energy that is often harnessed to drive other chemical reactions

ATP and Muscle Contraction



Cross-bridge muscle contraction cycle: The cross-bridge muscle contraction cycle, which is triggered by Ca2+ binding to the actin active site, is shown. With each contraction cycle, actin moves relative to myosin.

Muscles contract in a repeated pattern of binding and releasing between the two thin and thick strands of the sarcomere. ATP is critical to prepare myosin for binding and to “recharge” the myosin.

The Cross-Bridge Muscle Contraction Cycle

ATP first binds to myosin, moving it to a high-energy state. The ATP is hydrolyzed into ADP and inorganic phosphate (P_i) by the enzyme ATPase. The energy released during ATP hydrolysis changes the angle of the myosin head into a “cocked” position, ready to bind to actin if the sites are available. ADP and Pi remain attached; myosin is in its high energy configuration.

The muscle contraction cycle is triggered by calcium ions binding to the protein complex troponin, exposing the active-binding sites on the actin. As soon as the actin-binding sites are uncovered, the high-energy myosin head bridges the gap, forming a cross-bridge. Once myosin binds to the actin, the P_i is released, and the myosin undergoes a conformational change to a lower energy state. As myosin expends the energy, it moves through the “power stroke,” pulling the actin filament toward the M-line. When the actin is pulled approximately 10 nm toward the M-line, the sarcomere shortens and the muscle contracts. At the end of the power stroke, the myosin is in a low-energy position.

After the power stroke, ADP is released, but the cross-bridge formed is still in place. ATP then binds to myosin, moving the myosin to its high-energy state, releasing the myosin head from the actin active site. ATP can then attach to myosin, which allows the cross-bridge cycle to start again; further muscle contraction can occur. Therefore, without ATP, muscles would remain in their contracted state, rather than their relaxed state.

This page titled 9.1D: ATP and Muscle Contraction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.1E: Control of Muscle Tension

Muscle tension is influenced by the number of cross-bridges that can be formed.

Learning Objectives

- Describe the factors that control muscle tension

Key Points


- The more cross-bridges that are formed, the more tension in the muscle.
- The amount of tension produced depends on the cross-sectional area of the muscle fiber and the frequency of neural stimulation.
- Maximal tension occurs when thick and thin filaments overlap to the greatest degree within a sarcomere; less tension is produced when the sarcomere is stretched.
- If more motor neurons are stimulated, more myofibers contract, and there is greater tension in the muscle.

Key Terms

- **tension**: condition of being held in a state between two or more forces, which are acting in opposition to each other

Control of Muscle Tension

Neural control initiates the formation of actin – myosin cross-bridges, leading to the sarcomere shortening involved in muscle contraction. These contractions extend from the muscle fiber through connective tissue to pull on bones, causing skeletal movement. The pull exerted by a muscle is called tension. The amount of force created by this tension can vary, which enables the same muscles to move very light objects and very heavy objects. In individual muscle fibers, the amount of tension produced depends primarily on the amount of cross-bridges formed, which is influenced by the cross-sectional area of the muscle fiber and the frequency of neural stimulation.



Muscle tension: Muscle tension is produced when the maximum amount of cross-bridges are formed, either within a muscle with a large diameter or when the maximum number of muscle fibers are stimulated. Muscle tone is residual muscle tension

that resists passive stretching during the resting phase.

Cross-bridges and Tension

The number of cross-bridges formed between actin and myosin determine the amount of tension that a muscle fiber can produce. Cross-bridges can only form where thick and thin filaments overlap, allowing myosin to bind to actin. If more cross-bridges are formed, more myosin will pull on actin and more tension will be produced.

Maximal tension occurs when thick and thin filaments overlap to the greatest degree within a sarcomere. If a sarcomere at rest is stretched past an ideal resting length, thick and thin filaments do not overlap to the greatest degree so fewer cross-bridges can form. This results in fewer myosin heads pulling on actin and less muscle tension. As a sarcomere shortens, the zone of overlap reduces as the thin filaments reach the H zone, which is composed of myosin tails. Because myosin heads form cross-bridges, actin will not bind to myosin in this zone, reducing the tension produced by the myofiber. If the sarcomere is shortened even more, thin filaments begin to overlap with each other, reducing cross-bridge formation even further, and producing even less tension. Conversely, if the sarcomere is stretched to the point at which thick and thin filaments do not overlap at all, no cross-bridges are formed and no tension is produced. This amount of stretching does not usually occur because accessory proteins, internal sensory nerves, and connective tissue oppose extreme stretching.

The primary variable determining force production is the number of myofibers (long muscle cells) within the muscle that receive an action potential from the neuron that controls that fiber. When using the biceps to pick up a pencil, for example, the motor cortex of the brain only signals a few neurons of the biceps so only a few myofibers respond. In vertebrates, each myofiber responds fully if stimulated. On the other hand, when picking up a piano, the motor cortex signals all of the neurons in the biceps so that every myofiber participates. This is close to the maximum force the muscle can produce. As mentioned above, increasing the frequency of action potentials (the number of signals per second) can increase the force a bit more because the tropomyosin is flooded with calcium.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- intercalated disc. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intercalated%20disc**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Animal locomotion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Animal_locomotion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- actin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/actin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myosin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myosin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Bougle_whole2_retouched.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ABougle_whole2_retouched.png**. **License**: ***Public Domain: No Known Copyright***
- Structural Biochemistry/Cell Signaling Pathways/Muscular System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structural_Biochemistry/Cell_Signaling_Pathways/Muscular_System%23Oxidative_and_Glycolytic_Fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycolytic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glycolytic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...on/fast-twitch**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...on/slow-twitch**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Bougle_whole2_retouched.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ABougle_whole2_retouched.png**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Sliding filament model. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sliding_filament_model**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/definition/i-band**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...n/cross-bridge**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...1-25e30f147dbb**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- Bougle_whole2_retouched.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ABougle_whole2_retouched.png**. **License**: ***Public Domain: No Known Copyright***
- Sarcomere.png. **Provided by**: Wikipedia - Adapted From. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- M-line. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/M-line**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- troponin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/troponin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ATPase. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/ATPase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- Bougle_whole2_retouched.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ABougle_whole2_retouched.png**. **License**: ***Public Domain: No Known Copyright***
- Sarcomere.png. **Provided by**: Wikipedia - Adapted From. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Muscle Contraction and Locomotion. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest...e_38_04_05.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- tension. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. December 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- Bougle_whole2_retouched.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3ABougle_whole2_retouched.png**. **License**: ***Public Domain: No Known Copyright***
- Sarcomere.png. **Provided by**: Wikipedia - Adapted From. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Muscle Contraction and Locomotion. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/Figure_38_04_05.png**. **License**: ***CC BY: Attribution***
- Skeletal muscle. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Skeletal_muscle.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 9.1E: Control of Muscle Tension is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.2: SMOOTH MUSCLE



- Topic hierarchy

9.2A: Skeletal Muscle Fibers

This page titled 9.2: Smooth Muscle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.2A: Skeletal Muscle Fibers

Skeletal muscles are composed of striated subunits called sarcomeres, which are composed of the myofilaments actin and myosin.

Learning Objectives

- Outline the structure of a skeletal muscle fiber

Key Points

- Muscles are composed of long bundles of myocytes or muscle fibers.
- Myocytes contain thousands of myofibrils.
- Each myofibril is composed of numerous sarcomeres, the functional contracile region of a striated muscle. Sarcomeres are composed of myofilaments of myosin and actin, which interact using the sliding filament model and cross-bridge cycle to contract.

Key Terms

- **sarcoplasm**: The cytoplasm of a myocyte.
- **sarcoplasmic reticulum**: The equivalent of the smooth endoplasmic reticulum in a myocyte.
- **sarcolemma**: The cell membrane of a myocyte.
- **sarcomere**: The functional contractile unit of the myofibril of a striated muscle.

Skeletal Muscle Fiber Structure

Myocytes, sometimes called muscle fibers, form the bulk of muscle tissue. They are bound together by perimysium, a sheath of connective tissue, into bundles called fascicles, which are in turn bundled together to form muscle tissue. Myocytes contain numerous specialized cellular structures which facilitate their contraction and therefore that of the muscle as a whole.

The highly specialized structure of myocytes has led to the creation of terminology which differentiates them from generic animal cells.


- Generic cell > Myocyte
- Cytoplasm > Sarcoplasm
- Cell membrane > Sarcolemma
- Smooth endoplasmic reticulum > Sarcoplasmic reticulum

Myocyte Structure

Myocytes can be incredibly large, with diameters of up to 100 micrometers and lengths of up to 30 centimeters. The sarcoplasm is rich with glycogen and myoglobin, which store the glucose and oxygen required for energy generation, and is almost completely filled with myofibrils, the long fibers composed of myofilaments that facilitate muscle contraction.

The sarcolemma of myocytes contains numerous invaginations (pits) called transverse tubules which are usually perpendicular to the length of the myocyte. Transverse tubules play an important role in supplying the myocyte with Ca^{+} ions, which are key for muscle contraction.

Each myocyte contains multiple nuclei due to their derivation from multiple myoblasts, progenitor cells that give rise to myocytes. These myoblasts asre located to the periphery of the myocyte and flattened so as not to impact myocyte contraction.




Myocyte: Skeletal muscle cell: A skeletal muscle cell is surrounded by a plasma membrane called the sarcolemma with a cytoplasm called the sarcoplasm. A muscle fiber is composed of many myofibrils, packaged into orderly units.

Myofibril Structure

Each myocyte can contain many thousands of myofibrils. Myofibrils run parallel to the myocyte and typically run for its entire length, attaching to the sarcolemma at either end. Each myofibril is surrounded by the sarcoplasmic reticulum, which is closely associated with the transverse tubules. The sarcoplasmic reticulum acts as a sink of Ca^{+} ions, which are released upon signalling from the transverse tubules.

Sarcomeres



Sarcomere: The sarcomere is the functional contractile region of the myocyte, and defines the region of interaction between a set of thick and thin filaments.

Myofibrils are composed of long myofilaments of actin, myosin, and other associated proteins. These proteins are organized into regions termed sarcomeres, the functional contractile region of the myocyte. Within the sarcomere actin and myosin, myofilaments are interlaced with each other and slide over each other via the sliding filament model of contraction. The regular organization of these sarcomeres gives skeletal and cardiac muscle their distinctive striated appearance.

Myofilaments (Thick and Thin Filaments)

Myofibrils are composed of smaller structures called myofilaments. There are two main types of myofilaments: thick filaments and thin filaments. Thick filaments are composed primarily of myosin proteins, the tails of which bind together leaving the heads exposed to the interlaced thin filaments. Thin filaments are composed of actin, tropomyosin, and troponin. The molecular model of contraction which describes the interaction between actin and myosin myofilaments is called the cross-bridge cycle.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Myofibril. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myofibril**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skeletal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeletal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- Skeletal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeletal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myofibril. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myofibril**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sarcomere. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sarcomere**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Muscle Contraction and Locomotion. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/Figure_38_04_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Muscle Contraction and Locomotion. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44788/latest/Figure_38_04_03.jpg**. **License**: ***CC BY: Attribution***

This page titled 9.2A: Skeletal Muscle Fibers is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3: CONTROL OF MUSCLE TENSION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

9.3A: Force of Muscle Contraction

9.3B: Velocity and Duration of Muscle Contraction

9.3C: Motor Units

9.3D: Muscle Tone

9.3E: Types of Muscle Contractions: Isotonic and Isometric

This page titled 9.3: Control of Muscle Tension is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3A: Force of Muscle Contraction

The force a muscle generates is dependent on its length and shortening velocity.

Learning Objectives

- Differentiate between force-length and force-velocity of muscle contraction

Key Points

- The force -length relationship indicates that muscles generate the greatest force when at their resting (ideal) length, and the least amount of force when shortened or stretched relative to the resting length.
- The force-velocity relationship demonstrates that power produced is controlled by the velocity and force of muscle contraction, with an optimum power output at one third of maximum velocity.

Key Terms


- **force**: Any influence that causes an object to undergo a certain change concerning its movement, direction, or geometrical construction.
- **resting length**: Often the ideal length of a muscle and the length at which it can create the greatest active force.
- **power**: A measure of force x velocity, a measurable output for muscle contraction
- **Force-Velocity Relationship**: The relationship between the speed and force of muscle contraction, outputted as power.
- **Force-Length Relationship**: The relationship between sarcomere length and force produced in the muscle, modulated by actin and myosin myofilament overlap.

Muscle Force Generation

The force a muscle generates is dependent on the length of the muscle and its shortening velocity. These two fundamental properties limit many key biomechanical properties, including running speed, strength, and jumping distance.

Force-Length Relationship

Due to the presence of titin, muscles are innately elastic. Skeletal muscles are attached to bones via tendons that maintain the muscle under a constant level of stretch called the resting length. If this attachment was removed, for example if the bicep was detached from the scapula or radius, the muscle would shorten in length.




The Ideal Length of a Sarcomere: Sarcomeres produce maximal tension when thick and thin filaments overlap between about 80 percent to 120 percent, approximately 1.6 to 2.6 micrometers.

Muscles exist in this state to optimize the force produced during contraction, which is modulated by the interlaced myofilaments of the sarcomere. When a sarcomere contracts, myosin heads attach to actin to form cross-bridges. Then, the thin filaments slide over the thick filaments as the heads pull the actin. This results in sarcomere shortening, creating the tension of the muscle contraction. If a sarcomere is stretched too far, there will be insufficient overlap of the myofilaments and the less force will be produced. If the muscle is over-contracted, the potential for further contraction is reduced, which in turn reduces the amount of force produced.

Simply put, the tension generated in skeletal muscle is a function of the magnitude of overlap between actin and myosin myofilaments.

In mammals, there is a strong overlap between the optimum and actual resting length of sarcomeres.

Force-Velocity Relationship



Force-Velocity Relationship: As velocity increases force and therefore power produced is reduced. Although force increases

due to stretching with no velocity, zero power is produced. Maximum power is generated at one-third of maximum shortening velocity.

The force-velocity relationship in muscle relates the speed at which a muscle changes length with the force of this contraction and the resultant power output (force x velocity = power). The force generated by a muscle depends on the number of actin and myosin cross-bridges formed; a larger number of cross-bridges results in a larger amount of force. However, cross-bridge formation is not immediate, so if myofilaments slide over each other at a faster rate the ability to form cross bridges and resultant force are both reduced.

At maximum velocity no cross-bridges can form, so no force is generated, resulting in the production of zero power (right edge of graph). The reverse is true for stretching of muscle. Although the force of the muscle is increased, there is no velocity of contraction and zero power is generated (left edge of graph). Maximum power is generated at approximately one-third of maximum shortening velocity.

This page titled 9.3A: Force of Muscle Contraction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3B: Velocity and Duration of Muscle Contraction

The shortening velocity affects the amount of force generated by a muscle.

Learning Objectives

- Explain the interaction of velocity and duration in muscle contraction

Key Points

- Twitch contractions, which are short in duration, do not reach peak force.
- Tetanic contractions, which are long in duration, reach peak force and plateau.
- The shortening velocity is the speed at which a muscle changes length during a contraction.
- The force of a muscle contraction declines with increasing velocity.


Key Terms

- **Force-Velocity Relationship**: The relationship between the speed and force of muscle contraction, outputted as power.
- **summation.**: The occurrence of another twitch contraction before complete relaxation of the prior twitch has occurred.
- **tetanic**: A longer contraction of a muscle which reaches peak force.
- **twitch**: A short contraction of a muscle which does not reach peak force.

Muscle Contraction Velocity

Skeletal muscle contractions can be broadly separated into twitch and tetanic contractions. In a twitch contraction, a short burst of stimulation causes the muscle to contract, but the duration is so brief that the muscle begins relaxing before reaching peak force. If another contraction occurs before complete relaxation of a muscle twitch, then the next twitch will simply sum onto the previous twitch, a phenomenon called summation. If the stimulation is long enough, the muscle reaches peak force and plateaus at this level, resulting in a tetanic contraction.

Force-Velocity Relationship



Force-Velocity Relationship: As velocity increases force and power produced is reduced. Although force increases due to stretching with no velocity, zero power is produced. Maximum power is generated at one-third of maximum shortening velocity.

The force-velocity relationship in muscle relates the speed at which a muscle changes length to the force of this contraction and the resultant power output (force x velocity = power). The force generated by a muscle depends on the number of actin and myosin cross-bridges formed; a larger number of cross-bridges results in a larger amount of force. However, cross-bridge formation is not immediate and if myofilaments slide over each other at a faster rate, their ability to form cross-bridges and subsequent force are both reduced.

At a maximum velocity no cross-bridges can form so no force is generated, resulting in the production of zero power (right edge of graph). The reverse is true for stretching of muscle; although the force of the muscle is increased, there is no velocity of contraction and zero power is generated (left edge of graph). Maximum power is generated at approximately one-third of maximum shortening velocity.

Twitch contractions are short in duration. Though they have high velocity, they begin resting before reaching peak force. Tetanic contractions, which are long in duration, reach peak force and plateau.

This page titled 9.3B: Velocity and Duration of Muscle Contraction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3C: Motor Units

The motor unit is the functional unit of muscle contraction and includes the motor nerve fiber and the muscle fibers it innervates.

Learning Objectives

- Describe a motor unit and its function

Key Points

- A motor unit, the functional unit of muscle contraction, is a single motor nerve and the associated muscle fibers that are innervated upon stimulation from the nerve.
- A collection of motor units is referred to as a motor pool.
- The number of muscle fibers within each unit can vary within a particular muscle, which impacts precision and force generation.
- Differential activation of single or multiple motor units within a motor pool can therefore control precision and force of movement.

Key Terms

- **Motor Pool**: A collection of motor units.
- **Motor Unit**: A grouping of a motor neuron and the muscle fibers innervated by it.

Motor Units

A motor unit consists of the motor neuron and the grouping of muscle fibers innervated by the neuron. The number of muscle fibers within a motor unit varies, and is a function of the muscle's ability for accurate and refined motion. Precision is inversely proportional to the size of the motor unit. Thus, small motor units can exercise greater precision of movement compared to larger motor units. For instance, thigh muscles, responsible for large powerful movements, can have a thousand fibers in each unit, while eye muscles, requiring small precise movements, might only have ten. Groups of motor units are innervated to coordinate contraction of a whole muscle and generate appropriate movement; all of the motor units within a muscle are considered a motor pool.

Motor Pools and Force

There are often multiple sizes of motor unit within a motor pool as a means of modulating the precision and force produced by a single muscle. For example, a small motor unit in the biceps can be activated for small precise movements, while a larger motor unit can be activated to facilitate more forceful actions.

These multiple motor units of different sizes within a motor pool allow for very fine control of force either spatially or temporally.

- Spatial: The recruitment of more or larger motor units to increase force.
- Temporal: The repeated stimulation of a motor unit so that the force of contraction generated is summed.

This page titled 9.3C: Motor Units is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3D: Muscle Tone

Muscle tone is a measure of a muscle's resistance to stretching while in a passive resting state.

Learning Objectives

- Describe the factors involved in muscle tone

Key Points


- Muscle tone is the maintenance of partial contraction of a muscle, important for generating reflexes, maintaining posture and balance, and controlling proper function of other organ systems.
- Tone is controlled by the sensory muscle spindle, which measures muscle stretch.
- Tone is not limited to skeletal muscles, but is also a property of cardiac and smooth muscles.

Key Terms

- **muscle spindle**: A sensory unit associated with muscle tissue that is responsible for maintaining muscle tone.
- **muscle tone**: The continuous and passive partial contraction of the muscles, which helps maintain posture.
- **golgi tendon organ**: A sensory unit associated with a tendon that is responsible for preventing damage to the associated muscle.

Even when at rest, muscle fibers are at least partially contracted, possessing a small degree of tension which is termed muscle tone or tonus. Muscle tone is controlled by neuronal impulses and influenced by receptors found in the muscle and tendons.

This influence leads to the generation of reflexes in the spinal cord, such as the immediately obvious knee jerk reaction but also including key functions such as the posture maintenance and proper digestive system function..



Sliding Filament Model of Contraction: Muscle fibers in relaxed and contracted positions. Muscle tone ensures that even when at rest the muscle is at least partially contracted.

Control of Muscle Tone

The main regulator of muscle tone is the muscle spindle, a small sensory unit that is closely associated with and lies parallel to a muscle. Connecting to the endomysium of a muscle fiber, muscle spindles are composed of nuclear bag fibers and nuclear chain fibers. Both are similar to muscle fibers in that they contain actin and myosin myofilaments that allow them to stretch with the muscle. However, unlike skeletal muscle fibers where the nuclei are spread out and located at the periphery of the cell, in nuclear bag and nuclear chain fibers the nuclei are located in a central region which is enlarged in nuclear bag fibers.

Both cells of the muscle spindle contain sensory neurons. When stretched, muscle spindles become activated, triggering impulses to the spinal cord that can generate an immediate reflex. Spindles can also trigger impulses to the cerebral cortex providing information about the degree of stretch within the muscle.

To maintain tone, spindles also operate a feedback loop by directly triggering motor neurons linked to their associated muscles. If tone decreases and the muscle stretches the spindle, an impulse results in a muscle contraction. With this contraction, the spindle is no longer stretched.

A similar system is found in the tendons attaching muscle to bone. Distinct stretch receptors called golgi tendon organs assess the level of stretch within the tendon. The sensitivity of the golgi tendon organ is significantly less than that of the spindle, so it is thought they exist to prevent damage rather than control muscle tone.

Smooth and Cardiac Muscles

Smooth and cardiac muscles do not have specialized muscle spindles. Tone is maintained through autonomous feedback from the muscle fibers, neurons, and associated tissues.

This page titled 9.3D: Muscle Tone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.3E: Types of Muscle Contractions: Isotonic and Isometric

Muscle contractions are defined by changes in the length of the muscle during contraction.

Learning Objectives

- Differentiate among the types of muscle contractions

Key Points

- Isotonic contractions generate force by changing the length of the muscle and can be concentric contractions or eccentric contractions.
- A concentric contraction causes muscles to shorten, thereby generating force.
- Eccentric contractions cause muscles to elongate in response to a greater opposing force.
- Isometric contractions generate force without changing the length of the muscle.

Key Terms

- **Isometric**: A muscular contraction in which the length of the muscle does not change.
- **isotonic**: A muscular contraction in which the length of the muscle changes.
- **eccentric**: An isotonic contraction where the muscle lengthens.
- **concentric**: An isotonic contraction where the muscle shortens.

A muscle fiber generates tension through actin and myosin cross-bridge cycling. While under tension, the muscle may lengthen, shorten, or remain the same. Although the term contraction implies shortening, when referring to the muscular system, it means the generation of tension within a muscle fiber. Several types of muscle contractions occur and are defined by the changes in the length of the muscle during contraction.

Isotonic Contractions


Isotonic contractions maintain constant tension in the muscle as the muscle changes length. Isotonic muscle contractions can be either concentric or eccentric.

Concentric Contractions

A concentric contraction is a type of muscle contraction in which the muscles shorten while generating force, overcoming resistance. For example, when lifting a heavy weight, a concentric contraction of the biceps would cause the arm to bend at the elbow, lifting the weight towards the shoulder. Cross-bridge cycling occurs, shortening the sarcomere, muscle fiber, and muscle.

Eccentric Contractions

An eccentric contraction results in the elongation of a muscle while the muscle is still generating force; in effect, resistance is greater than force generated. Eccentric contractions can be both voluntary and involuntary. For example, a voluntary eccentric contraction would be the controlled lowering of the heavy weight raised during the above concentric contraction. An involuntary eccentric contraction may occur when a weight is too great for a muscle to bear and so it is slowly lowered while under tension. Cross-bridge cycling occurs even though the sarcomere, muscle fiber, and muscle are lengthening, controlling the extension of the muscle.



Types of Muscle Contraction: An isotonic concentric contraction results in the muscle shortening, an isotonic eccentric contraction results in the muscle lengthening. During an isometric contraction the muscle is under tension but neither shortens nor lengthens.

Isometric Contractions

In contrast to isotonic contractions, isometric contractions generate force without changing the length of the muscle, common in the muscles of the hand and forearm responsible for grip. Using the above example, the muscle contraction required to grip but not move a heavy object prior to lifting would be isometric. Isometric contractions are frequently used to maintain posture.

Isometric contractions are sometimes described as yielding or overcoming.

Yielding

A yielding contraction occurs when a muscle contraction is opposed by resistance. For example, when holding a heavy weight steady, neither raising nor lowering it.

Overcoming

An overcoming contraction occurs when a muscle contraction is opposed by an immovable object, such as the contraction generated in the muscles when pushing against a wall.

In both instances, cross-bridge cycling is maintaining tension in the muscle; the sarcomere, muscle fibers, and muscle are not changing length.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: *CC BY: Attribution*

- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction%23Force-length_and_force-velocity_relationships**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle contraction. **Provided by**: Saylor. **Located at**: **http://www.saylor.org/site/wp-content/uploads/2010/11/Smooth-Muscle-Contraction.pdf**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lengthtension. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lengthtension.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/resting-length**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/shortening-velocity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- force. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/force**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: ***CC BY: Attribution***
- Muscle_Force_Velocity_relationship.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction%23Classification_of_voluntary_muscular_contractions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction%23Force-length_and_force-velocity_relationships**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tetanic contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tetanic%20contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/shortening-velocity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: ***CC BY: Attribution***
- Muscle_Force_Velocity_relationship.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Force-Velocity Relationship. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Motor unit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_unit**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor unit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_unit**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor unit recruitment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_unit_recruitment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- motor nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/motor%20nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- innervation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/innervation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: ***CC BY: Attribution***
- Muscle_Force_Velocity_relationship.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Force-Velocity Relationship. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Muscle tone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_tone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle tone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_tone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/tonus-of-residual-muscle-tension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- muscle tone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/muscle_tone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: ***CC BY: Attribution***
- Muscle_Force_Velocity_relationship.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***

- Force-Velocity Relationship. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Isometric contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Isometric_contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction%23Contractions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction%23Contractions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- isotonic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/isotonic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- isometric. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/isometric**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- eccentric. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/eccentric**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- concentric. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/concentric**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Anatomy & Physiology. January 20, 2014. **Provided by**: OpenStax CNX. **Located at**: **openstaxcollege.org/files/tex...ol11496-op.pdf**. **License**: ***CC BY: Attribution***
- Muscle_Force_Velocity_relationship.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Force-Velocity Relationship. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muscle_Force_Velocity_relationship.png**. **License**: ***Public Domain: No Known Copyright***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 1015_Types_of_Contraction_new.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:1015_Types_of_Contraction_new.jpg**. **License**: ***CC BY: Attribution***

This page titled 9.3E: Types of Muscle Contractions: Isotonic and Isometric is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.4: MUSCLE METABOLISM



- Topic hierarchy

9.4A: Muscle Metabolism

9.4B: Muscle Fatigue

This page titled 9.4: Muscle Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.4A: Muscle Metabolism

Muscle contraction occurs via metabolism of adenosine triphosphate (ATP) derived primarily from the simple sugar glucose.

Learning Objectives

- Explain the process involved in muscle metabolism during aerobic exercise

Key Points

- ATP is required for muscle contraction. Four sources of this substance are available to muscle fibers: free ATP, phosphocreatine, glycolysis and cellular respiration.
- A small amount of free ATP is available in the muscle for immediate use.
- Phosphocreatine provides phosphates to ADP molecules, producing high-energy ATP molecules. It is present in low levels in the muscle.
- Glycolysis converts glucose to pyruvate, water and NADH, producing two molecules of ATP. Excess pyruvate is converted to lactic acid which causes muscle fatigue.
- Cellular respiration produces further molecules of ATP from pyruvate in the mitochondria. It is also required to resynthesize glycogen from lactic acid and restore stores of phosphocreatine and ATP in the muscle.

Key Terms

- **phosphocreatine**: A phosphorylated creatine molecule that serves as a rapidly mobilizable reserve of high-energy phosphates in skeletal muscle.
- **ATP**: The molecule containing high-energy bonds used to transfer energy between systems within a cell.

Muscle contractions are fueled by adenosine triphosphate (ATP), an energy-storing molecule. Four potential sources of ATP power muscle contractions.

Free ATP

Low levels of ATP exist within the muscle fibers and can immediately provide energy for contraction. However, the pool is very small and after a few muscle twitches will be exhausted.

Phosphocreatine

Phosphocreatine, also known as creatine phosphate, can rapidly donate a phosphate group to ADP to form ATP and creatine under anaerobic conditions. Enough phosphocreatine is present in the muscle to provide ATP for up to 15 seconds of contraction.

The reaction of phosphocreatine + ADP to ATP + creatine is reversible. During periods of rest, the store of phosphocreatine is regenerated from ATP.

Glycolysis

Glycolysis is the metabolic reaction which produces two molecules of ATP through the conversion of glucose into pyruvate, water, and NADH in the absence of oxygen.

The glucose for glycolysis can be provided by the blood supply, but is more often converted from glycogen in the muscle fibers. If glycogen stores in the muscle fibers are expended, glucose can be created from fats and proteins. However, this conversion is not as efficient.

Pyruvate is continually processed into lactic acid. With pyruvate accumulation, the amount of lactic acid produced is also increased. This lactic acid accumulation in the muscle tissue reduces the pH, making it more acidic and producing the stinging feeling in muscles when exercising. This inhibits further anaerobic respiration, inducing fatigue.

Glycolysis alone can provide energy to the muscle for approximately 30 seconds, although this interval can be increased with muscle conditioning.

Cellular Respiration

While the pyruvate generated through glycolysis can accumulate to form lactic acid, it can also be used to generate further molecules of ATP. Mitochondria in the muscle fibers can convert pyruvate into ATP in the presence of oxygen via the Krebs Cycle, generating an additional 30 molecules of ATP.

Cellular respiration is not as rapid as the above mechanisms; however, it is required for exercise periods longer than 30 seconds. Cellular respiration is limited by oxygen availability, so lactic acid can still build up if pyruvate in the Krebs Cycle is insufficient.

Cellular respiration plays a key role in returning the muscles to normal after exercise, converting the excess pyruvate into ATP and regenerating the stores of ATP, phosphocreatine, and glycogen in the muscle that are required for more rapid contractions.

This page titled 9.4A: Muscle Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.4B: Muscle Fatigue

Muscle fatigue occurs following a period of sustained activity.

Learning Objectives

- Describe the factors involved in metabolic muscle fatigue

Key Points

- Muscle fatigue refers to the decline in muscle force generated over time.
- Several factors contribute to muscle fatigue, the most important being lactic acid accumulation.
- With sufficient exercise the onset of muscle fatigue can be delayed.

Key Terms

- **Lactic Acid**: A byproduct of anaerobic respiration which strongly contributes to muscle fatigue.

Muscle fatigue refers to the decline in muscle force generated over sustained periods of activity or due to pathological issues. Muscle fatigue has a number of possible causes including impaired blood flow, ion imbalance within the muscle, nervous fatigue, loss of desire to continue, and most importantly, the accumulation of lactic acid in the muscle.

Lactic Acid Accumulation

Long-term muscle use requires the delivery of oxygen and glucose to the muscle fiber to allow aerobic respiration to occur, producing the ATP required for muscle contraction. If the respiratory or circulatory system cannot keep up with demand, then energy will be generated by the much less efficient anaerobic respiration.

In aerobic respiration, pyruvate produced by glycolysis is converted into additional ATP molecules in the mitochondria via the Krebs Cycle. With insufficient oxygen, pyruvate cannot enter the Krebs cycle and instead accumulates in the muscle fiber. Pyruvate is continually processed into lactic acid. With pyruvate accumulation, lactic acid production is also increased. This lactic acid accumulation in the muscle tissue reduces the pH, making it more acidic and producing the stinging feeling in muscles when exercising. This further inhibits anaerobic respiration, inducing fatigue.

Lactic acid can be converted back to pyruvate in well-oxygenated muscle cells; however, during exercise the focus in on maintaining muscle activity. Lactic acid is transported to the liver where it can be stored prior to conversion to glucose in the presence of oxygen via the Cori Cycle. The amount of oxygen required to restore the lactic acid balance is often referred to as the oxygen debt.

Ion Imbalance

Contraction of a muscle requires Ca^{+} ions to interact with troponin, exposing the actin binding site to the myosin head. With extensive exercise, the osmotically active molecules outside of the muscle are lost through sweating. This loss changes the osmotic gradient, making it more difficult for the required Ca^{+} ions to be delivered to the muscle fiber. In extreme cases, this can lead to painful, extended maintenance of muscle contraction or cramp.

Nervous Fatigue and Loss of Desire

Nerves are responsible for controlling the contraction of muscles, determining the number, sequence, and force of muscular contractions. Most movements require a force far below what a muscle could potentially generate, and barring disease nervous

fatigue is seldom an issue. However, loss of desire to exercise in the face of increasing muscle soreness, respiration, and heart rate can have a powerful negative impact on muscle activity.

Metabolic Fatigue

Depletion of required substrates such as ATP or glycogen within a muscle result in fatigue as the muscle is not able to generate energy to power contractions. Accumulation of metabolites from these reactions other than lactic acid, such as Mg^{2+} ions or reactive oxygen species, can also induce fatigue by interfering with the release of Ca^{+} ions from the sarcoplasmic reticulum or through reduction in the sensitivity of troponin to Ca^{+}.

Exercise and Aging

With sufficient training, the metabolic capacity of a muscle can change, delaying the onset of muscle fatigue. Muscle specified for high-intensity anaerobic exercise will synthesise more glycolytic enzymes, whereas muscle for long endurance aerobic exercise will develop more capillaries and mitochondria. Additionally, with exercise, improvements to the circulatory and respiratory systems can facilitate better delivery of oxygen and glucose to the muscle.

With aging, levels of ATP, CTP, and myoglobin begin to decline, reducing the muscle's ability to function. Muscle fibers shrink or are lost and surrounding connective tissue hardens, making muscle contraction slower and more difficult. Exercise throughout life can help reduce the impact of aging by maintaining a healthy oxygen supply to the muscle.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- A computational model of skeletal muscle metabolism linking cellular adaptations induced by altered loading states to metabolic responses during exercise. **Provided by**: BioMedical Engineering OnLine. **Located at**: http://www.biomedical-engineering-on...content/6/1/14. **License**: ***CC BY: Attribution***
- Muscles. **Provided by**: Kimball's Biology Pages. **Located at**: http://biology-pages.info/M/Muscles.html. **License**: ***CC BY: Attribution***
- ATP. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ATP. **License**: ***CC BY-SA: Attribution-ShareAlike***
- creatine phosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/creatine%20phosphate. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactic acid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lactic%20acid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Muscular System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...%23Lactic_Acid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle fatigue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscle_fatigue. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle fatigue. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscle_fatigue. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactic acid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lactic%20acid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycolysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glycolysis. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 9.4B: Muscle Fatigue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.5: EXERCISE AND SKELETAL MUSCLE TISSUE



- Topic hierarchy

9.5A: Impacts of Exercise on Muscles

This page titled 9.5: Exercise and Skeletal Muscle Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.5A: Impacts of Exercise on Muscles

Sustained, repeated overload of a muscle group leads to hypertrophy and strengthening of those muscles.

Learning Objectives

- Describe the effects of exercise on muscles

Key Points

- In the short term, exercise can lead to muscle fatigue and soreness.
- In the long term, exercise can lead to muscle hypertrophy and other physiological changes.
- These physiological changes alter the muscle to make repeat exercise easier.

Key Terms

- **muscle hypertrophy**: Increase in muscle mass due to exercise, particularly weight training; a noticeable long-term effect of exercise.
- **micro-trauma**: The damage to muscle fibers that occurs during excessive eccentric contraction.

Exercise involves a series of sustained muscle contractions of either long or short duration depending on the nature of the physical activity.

Short-Term Effects

In the short term muscle can become fatigued and sore for reasons like impaired blood flow, ion imbalance within the muscle, nervous fatigue, loss of desire to continue exercising, and most importantly, the accumulation of lactic acid in the muscle.

Muscle soreness, once thought to be due to lactic acid accumulation, has more recently been attributed to small tearing, or micro-trauma, of the muscles fibers caused by eccentric contraction. With repeated cycles of eccentric contraction this soreness will be reduced.

Long-term Effects

Muscle hypertrophy, or the increase in muscle mass due to exercise, particularly weight training, is a noticeable long-term effect of exercise. Exercise of specific muscles can often result in hypertrophy in the opposite muscles as well, a phenomenon known as *cross education*.

Increases in muscle mass are not the only long-term effect of exercise. With sufficient training the metabolic capacity of a muscle can change, delaying the onset of muscle fatigue. Muscle specified for high intensity anaerobic exercise will synthesize more glycolytic enzymes, whereas muscle for long endurance aerobic exercise will develop more capillaries and mitochondria. Additionally, with exercise improvements to the circulatory and respiratory systems can facilitate better delivery of oxygen and glucose to the muscle.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle%23Exercise**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle hypertrophy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_...tein_synthesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...le-hypertrophy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactic acid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lactic%20acid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...-muscle-fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 9.5A: Impacts of Exercise on Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6: OVERVIEW OF THE MUSCULAR SYSTEM



- Topic hierarchy

9.6A: Interactions of Skeletal Muscles

9.6B: How Skeletal Muscles Are Named

9.6C: How Skeletal Muscles Produce Movements

9.6D: Muscle Attachment Sites

9.6E: Arrangement of Fascicles

9.6F: Lever Systems

This page titled 9.6: Overview of the Muscular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6A: Interactions of Skeletal Muscles

Skeletal muscles interact to produce movements by way of anatomical positioning and the coordinated summation of innervation signals.

Learning Objectives

- Explain the summation interactions of skeletal muscles and how they affect movement

Key Points

- Muscle contractions can be termed twitch, summation or tetanus.
- A twitch contraction is the
  period of contraction and relaxation of a muscle after a single stimulation.
- Summation is the
  occurrence of additional twitch contractions before the previous twitch has completely relaxed.
- Summation can be achieved by increasing the frequency of stimulation, or by recruiting additional muscle fibers within a muscle.
- Tetanus
  occurs when the frequency of muscle contraction is such that the maximal force is tension is generated without any relaxation of the muscle.


Key Terms

- **tetanus**: When the frequency of muscle contraction is such that the maximal force is tension is generated without any relaxation of the muscle.
- **summation**: The occurrence of additional twitch contractions before the previous twitch has completely relaxed.
- **twitch**: The period of contraction and relaxation of a muscle after a single stimulation.

Skeletal muscle contractions can be grouped based on the length and frequency of contraction.

Twitch

When stimulated by a single action potential a muscle contracts and then relaxes. The time between the stimulus and the initiation of contraction is termed the latent period, which is followed by the contraction period. At peak contraction the muscle relaxes and returns to its resting position. Taken all together these three periods are termed a twitch.



Muscle Twitch Contraction: The time between stimulation and contraction is termed the latent period. After contraction the muscle relaxes back to a resting level of tension. Together these three periods form a single muscle twitch,

Summation

If an additional action potential were to stimulate a muscle contraction before a previous muscle twitch had completely relaxed then it would sum onto this previous twitch increasing the total amount of tension produced in the muscle. This addition is termed summation. Within a muscle summation can occur across motor units to recruit more muscle fibers, and also within motor units by increasing the frequency of contraction.

Multiple fiber summation


When a weak signal is sent by the central nervous system to contract a muscle, the smaller motor units, being more excitable than the larger ones, are stimulated first. As the strength of the signal increases, more (and larger) motor units are excited. The largest motor units have as much as 50 times the contractile strength as the smaller ones; thus, as more and larger motor units are activated, the force of muscle contraction becomes progressively stronger. A concept known as the size principle allows for a gradation of muscle force during weak contraction to occur in small steps, which become progressively larger as greater amounts of force are required.

Frequency summation

For skeletal muscles, the force exerted by the muscle can be controlled by varying the frequency at which action potentials are sent to muscle fibers. Action potentials do not arrive at muscles synchronously, and, during a contraction, only a certain percentage of the fibers in the muscle will be contracting at any given time. In a typical circumstance, when a human is exerting as much muscular force as they are consciously able, roughly one-third of the fibers in that muscle will be contracting at once. This relatively low level of contraction is a protective mechanism to prevent damage to the muscle tissue and attaching tendons and structures.

Tetanus

If the frequency of action potentials generated increases to such a point that muscle tension has reached its peak and plateaued and no relaxation is observed then the muscle contraction is described as a tetanus.



Summation and Tetanus Contractions: Repeated twitch contractions, where the previous twitch has not relaxed completely are called a summation. If the frequency of these contractions increases to the point where maximum tension is generated and no relaxation is observed then the contraction is termed a tetanus.

This page titled 9.6A: Interactions of Skeletal Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6B: How Skeletal Muscles Are Named

The anatomical arrangement of skeletal muscle fascicles can be described as parallel, convergent, pennate, or sphincter.

Learning Objectives

- Differentiate among parallel, pennate, convergent, and sphincter muscle types

Key Points

- Parallel muscles are the most abundant and typical, with fascicles arranged parallel to one another.
- Convergent muscles are similar to parallel muscles in attachment, although fascicles do not run parallel to one another, producing a broader muscle.
- In a pennate muscle the tendon runs through the length of the muscle, with the fascicles attaching at an angle.
- Sphincter muscles are characterized by a circular arrangement of fascicles around an opening. With contraction, the opening becomes smaller.

Key Terms

- **Parallel**: A muscle with a common point of attachment, with fascicles running parallel to each other.
- **Circular**: A ring like band of muscle that surrounds a bodily opening, constricting and relaxing to control flow.
- **Pennate**: A feather shaped muscle with fascicles that attach obliquely (at an angle) to a central tendon.
- **Convergent**: A muscle with a common point of attachment, although individual fascicles do not necessarily run parallel to each other.

Skeletal muscle can be categorised into four groups based on its anatomical arrangement.

Parallel

Parallel muscles are characterized by fascicles that run parallel to one another, and contraction of these muscle groups acts as an extension of the contraction of a single muscle fiber. Most skeletal muscles in the body are parallel muscles; although they can be seen in a variety of shapes such as flat bands, spindle shaped, and some can have large protrusions in their middle known as the belly of the muscle.

Parallel muscles can be divided into fusiform and non-fusiform types based on their shape. Fusiform muscles are more spindle shaped (their diameter at the center is greater than at either end), whereas, non-fusiform muscles are more rectangular with a constant diameter.

The biceps brachii is an example of a fusiform parallel muscle, and is responsible for flexing the forearm.

Convergent

Convergent muscles have a common point of attachment, from which the muscle fascicles extend outward, not necessarily in a specific spatial pattern, allowing the muscle to cover a broad surface. These muscles do not tend to exert as much force on their tendons. Muscle fibers can often exert opposing effects during contraction, such as not pulling in the same direction depending on the location of the muscle fiber. Covering a broad surface these fibers allow for more versatile types of movement. Because the fascicles pull on the tendons at an angle, they do not move the tendon as far as their parallel muscle counterparts. Despite this they generate greater tension because they possess a greater amount of muscle fibers than similarly sized parallel muscles.

The pectoralis major found in the chest is an example of a convergent muscle, and is responsible for flexing the upper arm.

Pennate

In Pennate muscles, the tendon runs through the length of the muscle. Fascicles pull on the tendon at an angle, thus not moving as far at the parallel muscles during a contraction. However, these muscles tend to have relatively more muscle fibers than similarly sized parallel muscles, and thus carry more tension.


If all the fascicles of a pennate muscle are on the same side of the tendon, the pennate muscle is called unipennate. If the fascicles lie to either side of the tendon the muscle is called bipennate. If the central tendon branches within a pennate muscle, the muscle is called multipennate.

The rectus femoris found in the thigh, and responsible for its flexion, is an example of a bipennate muscle.

Circular

The fibers of the circular or sphincter muscles are arranged concentrically around an opening or recess. As the muscle contracts, the opening it circumvents gets smaller. For this reason, these muscles are often found at the entrances and exits of external and internal passageways. Skeletal circular muscles are different from smooth muscle equivalents due to their structure and because they are under voluntary control

The orbicularis oris which controls the opening of the mouth is an example of a circular muscle.



Types of muscle in the body: The four types of muscle; parallel (fusiform and non-fusiform), circular, convergent and pennate (uni, bi and multi).

This page titled 9.6B: How Skeletal Muscles Are Named is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6C: How Skeletal Muscles Produce Movements

Muscles are arranged in groupings of agonist, antagonist, and synergists that produce and modulate movement.

Learning Objectives

- Differentiate between agonist and antagonist muscles

Key Points

- Agonist muscles shorten with contraction to produce a movement.
- Following contraction, the antagonist muscle paired to the agonist muscle returns the limb to the previous position.
- Synergist muscles act around a movable joint to produce motion similar to or in concert with agonist muscles, allowing for a range of possible movements.

Key Terms

- **antagonist**: This type of muscle acts as opposing muscle to agonists, usually contracting as a means of returning the limb to its original resting position.
- **agonist**: These muscles are typically associates with the movement itself, and are sometimes referred to as prime movers. They contract while another muscle relaxes.
- **synergist**: This type of muscle acts around a movable joint to produce motion similar to or in concert with agonist muscles.

Muscles exist in groupings that work to produce movements by muscle contraction. Muscles are classified according to their actions during contractions as agonists, antagonists, or synergists.

For muscle pairings referred to as antagonistic pairs, one muscle is designated as the extensor muscle, which contracts to open the joint, and the flexor muscle, which acts opposite to the extensor muscle. These pairs exist in places in the body in which the body cannot return the limb back to its original position through simple lack of contraction. Typical muscle pairings include the biceps brachii and triceps brachii, which act to flex or extend the forearm.

Agonist Muscles

Agonist muscles are those we typically associate with movement itself, and are thus sometimes referred to as prime movers. Agonist muscles produce the primary movement or series of movements through their own contractions. To generate a movement, agonist muscles must physically be arranged so that they cross a joint by way of the tendon. Contraction will move limbs associated with that joint. In this sense, the bone acts as a lever with the attached muscle fiber's contraction, driving movement.

During flexing of the forearm the biceps brachii is the agonist muscle, pulling the forearm up towards the shoulder.

Antagonist Muscles


The majority of muscles are grouped in pairs, with an antagonist to each agonist muscle. Exceptions include those muscles such as sphincter muscles that act to contract in a way that is opposite to the resting state of the muscle. Antagonist muscles act as opposing muscles to agonists, usually contracting as a means of returning the limb to its original, resting position.

During flexing of the forearm, the triceps brachii is the antagonist muscle, resisting the movement of the forearm up towards the shoulder.

Synergist Muscles

Synergist muscles act around a moveable joint to produce motion similar to or in concert with agonist muscles. They often act to reduce excessive force generated by the agonist muscle and are referred to as neutralizers. Synergists are useful because they fix certain joints to allow a range of contractions, in contrast with the sheer power of an agonist contraction that limits the range of possible movements.

During flexing of the forearm, the brachioradialis and brachialis act as synergist muscles, aiding the biceps brachii in pulling the forearm up towards the shoulder. The muscles of the rotator cuff are also synergists in that they fix the shoulder joint allowing the bicepps brachii to exert a greater force.



Flexing of the forearm by the biceps brachii: The biceps brachii is the agonist, or primer mover, responsible for flexing the forearm. The triceps brachii (not shown) acts as the antagonist. The brachioradialis and brachialis are synergist muscles, and the rotator cuff (not shown) fixes the shoulder joint allowing the biceps brachii to exert greater force.

This page titled 9.6C: How Skeletal Muscles Produce Movements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6D: Muscle Attachment Sites

Tendons are composed of connective tissue that attaches muscle to bone.

Learning Objectives

- Describe the function of tendons

Key Points


- Tendons have elasticity, which allows them to withstand tension and act as springs.
- Tendons mainly consists of closely-packed collagen fibers running parallel to the force generated by the muscle to which they are attached.
- Tendons also contain elastin fibers to improve the elastic properties and proteoglycans, which maintain tendon organisation during extension and compression.
- Aponeuroses are large flat sheets of connective tissue similar to tendons. They are responsible for binding muscle to bone and to the fascia of other muscles.

Key Terms

- **tendon**: A tough band of fibrous tissue that usually connects a muscle with a bone.
- **aponeuroses**: A tough flat sheet of fibrous tissue that connects muscle with bones or with the fascia of other muscles.

Most skeletal muscle attaches to bone in order to produce movement. However, some skeletal muscle attaches directly to other muscles, fascia, or tissues such as the skin.

Tendons



Achilles Tendon: The Achilles tendon provides stability and limits the range of motion at the ankle joint. It is the thickest and strongest tendon in the body. Tendons are a common tissue that connect muscle to bone.

A tendon is a cord-like, fibrous connective tissue that connects muscle to bone and is capable of withstanding tension. At either

end of the tendon, its fibers intertwine with the fascia of a muscle or the periosteum (a dense fibrous covering of a bone), allowing force to be dissipated across the bone or muscle.

Tendons mainly consists of closely-packed collagen fibers running parallel to the force generated by the muscle to which they are attached. Intertwined with the collagen fibers are elastin molecules, which improve the tendons' elasticity, and various proteoglycans, proteins to which many carbohydrate molecules are attached. These proteins play a key role in maintaining the organization of the tendon, especially during compression and extension.

Tendons were once thought to play only a passive connective role. However, research into their elastic properties has demonstrated that they can also act as springs. The elasticity of tendons allows them to passively store energy for later release. The most widely-researched example is the Achilles tendon which stores and releases elastic energy during walking, improving efficiency and reducing muscle load.

Aponeuroses

Not all muscle attaches via tendons. Aponeuroses are large, sheet-like layers of connective tissue with a similar composition to tendons. Aponeuroses can also attach to bone, as in the scalp aponeuroses, and to the fascia of other muscles or tissues, such as the anterior abdominal aponeuroses. Their large form and shape provides structure and distributes tension across a wider area or large number of muscle groups.

Other Attachments

Muscles can also attach directly to other tissues, which is most evident in the face. The skeletal muscles involved in controlling expression attach directly onto the fascia of the skin.

This page titled 9.6D: Muscle Attachment Sites is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6E: Arrangement of Fascicles

Skeletal muscles are grouped into fascicles, which are bunches of muscle fibers surrounded by a perimysium.

Learning Objectives

- Outline the construction of a muscle fascicle


Key Points

- Skeletal muscle is surrounded by a thick connective sheet termed the fascia
- Underneath this is another layer of connective tissue called the epimysium, which extends inwards as the perimysium into the muscle, splitting fibers into bundles termed fascicles.
- Each fascicle is surrounded by another layer of connective tissue termed the endomysium.
- This structure separates and protects the muscle and also acts to spread force throughout the muscle, preventing damage.

Key Terms

- **perimysium**: The continuation of the epimysium into the muscle, splitting fibers into fascicles.
- **epimysium**: A sheet of connective tissue lying below the fascia, also surrounding a muscle.
- **fascia**: A sheet of thick connective tissue which surrounds a muscle.
- **endomysium**: A sheet of connective tissue which wraps each fascicle.
- **fascicle**: A group of muscle of fibers surrounded by the perimysium.

Muscle Fascia



Muscle Structure: Skeletal muscle is surrounded by a thick outer layer of connective tissue termed the fascia. Within this is a layer termed the epimysium which splits inwards into the muscle as the perimysium dividing muscle fibers into groups termed fascicle. Each fascicle is surrounded by another layer of connective tissue termed the endomysium.

Skeletal muscle tissue is composed of numerous muscle fibers which are separated from adjacent muscles and other tissues by a layer of dense, elastic connective tissue termed the fascia. This fascia can project beyond the end of the muscle and attach

to bones, other muscles, and other tissues. Key muscle groups and the associated vascular and nervous systems can also be separated from other tissue, such as in the upper arm. These groupings are called fascial compartments.

This fascia is interlinked with a serious of fascia found throughout the body, including the superficial fascia which is the lowermost layer of the skin and the visceral fascia which surrounds internal organs. The fascia surrounding a muscle or muscle group does not contain many blood vessels, but is rich with sensory receptors.

Muscle fascia is predominately composed of cross-linked collagen and elastin fibers oriented parallel to the direction of muscle force, making them able to resist high-tension forces while remaining somewhat elastic.

Fascicles

Beneath the fascia in skeletal muscle is another layer of connective tissue termed the epimysium which is closely associated with the fascia. It extends inwards and becomes the perimysium, then into the muscle separating muscle fibers into small bundles termed fascicles. Fascicles can be arranged in a variety of anatomical positions within a muscle, producing different movements.

Each individual fiber within a fascicle is surrounded by a thin connective layer termed the endomysium, which helps maintain close association between the muscle fiber and associated vascular and nervous systems.

The organization of connective tissue throughout and around a muscle provides strength and flexibility while distributing the force evenly. It also maintains the close association of the vascular and nervous system with the muscle, which is required to deliver necessary metabolites and nerve impulses.

Cardiac and Smooth Muscle Tissue

Whilst both cardiac and smooth muscles are also wrapped in connective tissue, they are not differentiated in the same way as skeletal muscles.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- skeletal muscle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/skeletal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skeletal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeletal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skeletal muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeleta..._muscle_fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Muscular System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...e_Contractions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- summation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/summation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1...ch_Myogram.jpg**. **License**: ***CC BY: Attribution***
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg**. **License**: ***CC BY: Attribution***
- Skeletal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeleta...3Muscle_fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skeletal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeleta..._muscle_fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- sphincter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sphincter. **License**: *CC BY-SA: Attribution-ShareAlike*
- convergent. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/convergent. **License**: *CC BY-SA: Attribution-ShareAlike*
- pennate. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pennate. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1012_Muscle_Twitch_Myogram.jpg. **License**: *CC BY: Attribution*
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg. **License**: *CC BY: Attribution*
- Fascicle_Muscle_Shapes.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Fascicle_Muscle_Shapes.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...nition/agonist. **License**: *CC BY-SA: Attribution-ShareAlike*
- The human body.. **Provided by**: educacionfisicamaruxamallo Wikispace. **Located at**: http://educacionfisicamaruxamallo.wikispaces.com/The+human+body.#The+human+body.-The+skeletal+system.-The+joints.-Joints+and+movements.. **License**: *CC BY-SA: Attribution-ShareAlike*
- Antagonist (muscle). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Antagonist_(muscle). **License**: *CC BY-SA: Attribution-ShareAlike*
- Agonist (muscle). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Agonist_(muscle). **License**: *CC BY-SA: Attribution-ShareAlike*
- Synergist. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synergist. **License**: *CC BY-SA: Attribution-ShareAlike*
- antagonist. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antagonist. **License**: *CC BY-SA: Attribution-ShareAlike*
- synergist. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/synergist. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1012_Muscle_Twitch_Myogram.jpg. **License**: *CC BY: Attribution*
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg. **License**: *CC BY: Attribution*
- Fascicle_Muscle_Shapes.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Fascicle_Muscle_Shapes.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Biceps_Muscle_CNX.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Biceps_Muscle_CNX.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Tendon. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Tendon. **License**: *CC BY-SA: Attribution-ShareAlike*
- tendon. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tendon. **License**: *CC BY-SA: Attribution-ShareAlike*
- connective tissue. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/connective_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1012_Muscle_Twitch_Myogram.jpg. **License**: *CC BY: Attribution*
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg. **License**: *CC BY: Attribution*
- Fascicle_Muscle_Shapes.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Fascicle_Muscle_Shapes.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Biceps_Muscle_CNX.jpg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Biceps_Muscle_CNX.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/3c/Achilles-tendon.jpg. **License**: *Public Domain: No Known Copyright*
- Muscle fascicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscle_fascicle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Perimysium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perimysium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Endomysium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endomysium. **License**: *CC BY-SA: Attribution-ShareAlike*
- fascicle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fascicle. **License**: *CC BY-SA: Attribution-ShareAlike*
- endomysium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/endomysium. **License**: *CC BY-SA:*

Attribution-ShareAlike
- perimysium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/perimysium**. License: *CC BY-SA: Attribution-ShareAlike*
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1012_Muscle_Twitch_Myogram.jpg**. License: *CC BY: Attribution*
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg**. License: *CC BY: Attribution*
- Fascicle_Muscle_Shapes.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Fascicle_Muscle_Shapes.jpg**. License: *CC BY-SA: Attribution-ShareAlike*
- Biceps_Muscle_CNX.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Biceps_Muscle_CNX.jpg**. License: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3c/Achilles-tendon.jpg**. License: *Public Domain: No Known Copyright*
- Muscle Structure. **Provided by**: CNX. **Located at**: **http://cnx.org/contents/14fb4ad7-39a1-4eee-ab6e-3ef2482e3e22@6.27.**. License: *CC BY: Attribution*
- Anatomical terms of motion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anatomical_terms_of_motion**. License: *CC BY-SA: Attribution-ShareAlike*
- Muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle%23Strength**. License: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/definition/abduction**. License: *CC BY-SA: Attribution-ShareAlike*
- extensor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/extensor**. License: *CC BY-SA: Attribution-ShareAlike*
- flexor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/flexor**. License: *CC BY-SA: Attribution-ShareAlike*
- rotation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/rotation**. License: *CC BY-SA: Attribution-ShareAlike*
- 1012_Muscle_Twitch_Myogram.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1012_Muscle_Twitch_Myogram.jpg**. License: *CC BY: Attribution*
- 1013_Summation_Tetanus.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1013_Summation_Tetanus.jpg**. License: *CC BY: Attribution*
- Fascicle_Muscle_Shapes.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Fascicle_Muscle_Shapes.jpg**. License: *CC BY-SA: Attribution-ShareAlike*
- Biceps_Muscle_CNX.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Biceps_Muscle_CNX.jpg**. License: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3c/Achilles-tendon.jpg**. License: *Public Domain: No Known Copyright*
- Muscle Structure. **Provided by**: CNX. **Located at**: **http://cnx.org/contents/14fb4ad7-39a1-4eee-ab6e-3ef2482e3e22@6.27.**. License: *CC BY: Attribution*
- Body_Movements_I.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Body_Movements_I.jpg**. License: *CC BY-SA: Attribution-ShareAlike*

This page titled 9.6E: Arrangement of Fascicles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.6F: Lever Systems

Arrangement of muscles allows them to move relative to one another, while the insertion joint acts as the pivot point for a lever system.

Learning Objectives

- Differentiate between flexors and extensors and between adductors and abductors

Key Points

- A simple lever system can be used to describe skeletal muscle action. Levers are composed of a pivot, a fixed rod, and a load to which a force is applied.
- The relative positions of the pivot, load, and force determine the type of lever and subsequent action.
- Flexors and extensors work to adjust the angle between two body parts. Flexion decreases the angle and extension increases the angle.
- Abduction and adduction are movements relative to the midline of the body. Abduction is the movement away from this midline while adduction is movement towards the midline.
- Internal rotation brings the distal portions of the limbs inward towards the midline. External rotation is the opposite movement, pointing the distal portion of limbs away from the midline.
- The elevation of a limb or body part moves it in a superior direction, while depression moves body parts in an inferior direction.

Key Terms

- **abduction**: Movement away from the midline of the body.
- **rotation**: The act of turning around a centre or an axis.
- **first class lever**: Force and load to either side of the pivot.
- **second-class lever**: Load between force and pivot.
- **third-class lever**: Force between pivot and load.
- **origin**: The relatively immovable point of attachment for a muscle.
- **insertion**: The movable point of attachment for a muscle.
- **flexion**: Reduces the angle between two body parts.
- **extension**: Increases the angle between two body parts.
- **adduction**: Movement towards the midline of the body.

Skeletal muscle is usually attached to a relatively immovable part of the body at one end and more mobile region on the other end of a joint. The attachment at the immovable end is referred to as the origin and at the moveable end, the insertion. Upon contraction, the insertion is pulled towards the origin. This movement can be described using a simple lever system. Muscles can have multiple origins and insertions which modulate the type of movement they produce.

Levers

A lever is composed of three parts: a fixed rod which is attached to a fulcrum (pivot) and a load. Depending on the relative position of the three components, levers can move heavy loads or move loads further or faster when a force is applied.

If a load is close to a pivot and the force is applied far from the pivot, then the lever is said to operate at mechanical advantage. A large but relatively small force can move a heavy object. The classic example of such a lever is a car jack. With big movements of the lever, the heavy car is lifted in small increments off the ground.

If a load is far from a pivot and a force is applied near to the pivot, then the lever is said to operate at a mechanical disadvantage. A large force is required to move a relatively small load, but the speed and distance at which that load can be moved is greatly

increased. An example of this is a spade combined with forceful movements of the muscles in the arm, which results in large movements of the spade head.

In muscles, the joints are the pivots and the bones are the fixed rods. The load is the weight of the bone, associated tissues, and other objects being moved, and the force is applied by the muscle at its insertion point.

Classes of Levers

Levers can also vary based on the relative position of the load, pivot and point of force application. Classes of levers include:

First-class Lever

In a first class lever, the load and force sit on either side of the pivot like a seesaw. First-class levers are relatively uncommon in the body, but one example is the triceps brachii muscle of the upper arm which acts to extend the forearm. The force is applied at its point of insertion on the ulna in the forearm, the elbow is the pivot, and the load is the humerus in the upper arm. Thinking of the relative distance between the points of attachment, the triceps brachii can be said to act at a mechanical disadvantage.

Second-class Lever

In a second-class lever, the force is applied at one end with the pivot at the other and the load in between. Second-class levers are also relatively uncommon in the body. One example is raising yourself up on your toes. The pivot is based at the front of the foot, the load is the weight of the body, and the force is applied through the Achilles tendon in the heel. All second-class levers in the body act at a mechanical advantage since the force is always applied closer to the load than to the pivot.

Third-class Lever

In a third-class lever the force is applied between the load and the pivot. The majority of muscles in the body are third-class levers and all act at a mechanical disadvantage as the force is applied closer to the pivot than the load. An example of a third-class lever in the body is the biceps brachii, which flexes the forearm towards the shoulder. Originating at the scapula, the pivot is the elbow, with the force applied immediately after the elbow at the point of insertion on the radius of the forearm. The load is the forearm and any objects a person carries.

Types of Movement

Working together using the lever principles discussed above, skeletal muscles can induce a wide range of movements.

Flexors and Extensors

Flexors and extensors adjust and change the angle between two body parts. Flexion decreases the angle and extension increases the angle. For example, extension of the arm opens the angle of the elbow joint, while flexion allows for the bending of the arm. Flexion can also move inwards towards the body or forward, such as with hips or shoulders. Extension in this case moves the limbs towards the posterior side of the body lever.

Abduction and Adduction

Abduction and adduction are movements relative to the midline of the body. Abduction is the movement away from the midline of the body and adduction is movement towards this line. For example, moving arms or legs laterally away from the body is abduction, and bringing the limbs back to the midline is adduction.


Internal Rotation

Internal or medial rotation is specific to the shoulder or hip and brings the distal portions of the limbs inwards towards the midline. Internal rotation can also move the humerus and femur inward. External or lateral rotation is the opposite movement,

pointing the distal portion of limbs as well as the humerus and femur away from the mid-line.

Depression and Elevation

A limb or a body part can be moved upward (or in a superior direction) through elevation. For example, the trapezius elevates the apex of the shoulder upward. Depression is the opposite of elevation, or moving body parts in an inferior direction.



Types of body movements: Muscle positioning around a joint determines the type of movement that is produced.

This page titled 9.6F: Lever Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.7: HEAD AND NECK MUSCLES



- Topic hierarchy

9.7A: Facial Muscles

9.7B: Chewing Muscles

9.7C: Neck Muscles

This page titled 9.7: Head and Neck Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.7A: Facial Muscles

The human face is composed of multiple muscles that control the fine movements that produce facial expressions.

Learning Objectives

- Characterize the muscles involved in creating facial expressions

Key Points


- The human face is composed of numerous muscles that control fine movement to produce facial expressions.
- Unlike other muscles, these muscles originate on bone or fascia of the face and attach directly to the skin, allowing it to be manipulated.

Key Terms

- **depressor labii inferioris**: An analogous muscle that lowers the bottom lipEndFragment
- **Buccinator**: This muscle is located between the upper and lower jaws in the cheek, deep to the other muscles of the face.
- **zygomatic**: This muscle controls the cheeks to create smiles and frowns.
- **Procerus**: The most superior of all facial muscles.
- **depressor anguli oris**: This muscle is opposite to the levator anguli oris and pulls the corners of the mouth downward, producing a frown.
- **levator labii superiori**: Abroad muscle responsible for elevation of the upper lip.
- **Orbicularis Oris**: Muscle fibers that enclose the opening to the oral cavity.
- **Nasalis**: The largest of the nasal muscles. It is split into two sections: alar and transverse.
- **risorius**: This muscle pulls the mouth back mimicking a smile, but does not affect the skin around the smile.
- **Corrugator Supercilii**: A small muscle located superiorly to the orbicularis oculi.
- **Orbicularis Oculi**: A thin muscle that surrounds the eye socket.
- **mentalis**: This muscle pushes the lower lip uppers and wrinkles the chin.
- **levator labii superioris alaeque nasi**: The muscle of the upper lip. It acts to lift the upper lift and dilates nostril, producing a snarling expression.

The human face is composed of numerous muscles that control fine movement to produce facial expressions. Unlike other muscles, these muscles originate on the bone or fascia of the face and attach directly onto the skin, allowing it to be manipulated. The facial muscles can be split into three groups: orbital, nasal and oral.

Orbital Group



Schematic of head and neck muscles.: Locations of facial muscles noted.


- *Orbicularis Oculi*: A thin muscle that surrounds the eye socket.
  - Attachments: Originates from the skull around the eye socket and attaches to ligaments found in the eyelid.
  - Actions: Closes the eyelid.
- *Corrugator Supercilii*: A small muscle located superiorly to the orbicularis oculi.
  - Attachments: Originates from the bone below the eyebrow and attaches to the skin directly above.
  - Actions: Draws the eyebrows together.

Nasal Group

The nasal group of muscles are associated with movements of the nose and surrounding skin.

- *Nasalis*: The largest of the nasal muscles. It is split into two sections: alar and transverse.
  - Attachments: Originates from the upper jaw. The alar section attaches to the cartilage of the nose and the transverse section to an aponeurosis covering the bridge of the nose.
  - Actions: The transverse section closes the nostrils and the alar part opens them.
- *Procerus*: The most superior of all facial muscles.
  - Attachments: Originates from the nasal bone attaching to the skin of the forehead.
  - Actions: Pulls the eyebrows down.

Oral Group



Location of the buccinator muscle: Highlighted in orange, the buccinator is associated with the cheeks directly lateral to the mouth

Muscles of the oral group play key roles in respiration, communication, eating, and drinking. The lips in particular are controlled by numerous small muscles.

- *Orbicularis Oris*: Muscle fibers that enclose the opening to the oral cavity.
  - Attachments: Originates from the upper jaw and muscles of the cheek and attaches to the lips.
  - Action: Purses the lips.
- *Buccinator*: This muscle is located between the upper and lower jaws in the cheek, deep to the other muscles of the face.
  - Attachments: It originates from upper and lower jaw and attaches to the lips and orbicularis oris.
  - Actions: The buccinator pulls the cheek inwards.

Other Muscles of the Mouth

- The *levator labii superioris alaeque nasi* is the muscle of the upper lip. It acts to lift the upper lift and dilates nostril, producing a snarling expression.
- The *levator anguli oris* (caninus) inserts at the corners of the mouth at an angle, and is associated with other muscles including the zygomaticus, triangularis, and orbicularis oris. When innervated, this muscle contracts to lift the corners of the mouth, producing part of the expression of a smile.
- The *depressor anguli oris* (triangularis) is also associated with the corners of the mouth. Located opposite to the levator anguli oris, it pulls the corners of the mouth downward, producing a frown.
- The *levator labii superioris* is a broad muscle responsible for elevation of the upper lip. This muscle originates at the side of the nose and has several insertion points on either side of the nose, extending down to the lip, and inserting at both lateral and frontal portions of the upper lip.
- The *depressor labii inferioris* is an analogous muscle that lowers the bottom lip.
- The *zygomatic* muscle, associated with the cheeks, is divided into two parts: the major and the minor. The zygomaticus major draws the mouth upward and outward to generate a smile. The zygomaticus minor inserts into the outer part of the upper lip, not the angle of the mouth as with the major. When innervated, contraction of this muscle draws the lip backward, upward, and outward and is associated with facial expressions conveying sadness.
- The *mentalis*, associated with the tip of the chin, is a paired muscle. Sometimes referred to as the pouting muscle, contraction of the mentalis causes the lower lip to be pushed upwards and wrinkles the chin.
- The *risorius* muscle is lateral to the orbicularis oris and inserts into the angle of the mouth. When innervated, the risorius pulls the mouth back mimicking a smile, but does not affect the skin around the smile As a result, this facial expression is often interpreted as insincere.

This page titled 9.7A: Facial Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.7B: Chewing Muscles

Mastication, or chewing, involves the adduction and lateral motions of the jaw bone. It is controlled by four muscles of the face.

Learning Objectives

- Differentiate between the actions of the masseter and the temporalis muscles in chewing


Key Points

- The masseter elevates the jaw, closing the mouth.
- The temporalis elevates and retracts the jaw.
- The lateral pterygoid is the only muscle of mastication that actively opens the jaw. Unilateral action of a lateral pterygoid produces lateral movement in the jaw, usually performed in concert with the medial pterygoids.
- The medial pterygoid elevates and closes the jaw, contributes to protrusion of the mandible, and assists in mastication.

Key Terms

- **medial pterygoid**: A muscle of mastication with two heads. It lies inferiorly to the medial pterygoid.
- **temporalis**: A broad muscle that fans out to cover much of the temporal bone on the side of the skull.
- **lateral pterygoid**: A muscle of mastication with two heads. It lies superiorly to the medial pterygoid.
- **masseter**: The large muscle which raises the lower jaw, and assists in mastication.


Mastication, or the act of chewing, involves adduction and lateral motion of the jaw bone. It is controlled by four bilateral muscles in the face. The lower jaw, or mandible, connects to the temporal bone of the skull via the temporomandibular joint, which allows movement in all planes.



Location of the masseter muscle: The masseter muscle spans the opening of the mouth.

- *Masseter*: The most powerful muscle of mastication. It is quadrangular in shape and split into two regions, deep and superficial. It covers the other muscles of mastication.
  - Attachments: The superficial region originates from the skull below the eye socket, while the deep part originates from the skull above the jaw. Both parts attach to the jaw.

- Actions: Elevates the jaw.
- *Temporalis*: A broad muscle that fans out to cover much of the temporal bone on the side of the skull.
  - Attachments: The temporalis muscle has a wide, fan-shaped origin on the side of the skull and condenses into a tendon which attaches to the jaw.
  - Actions: Elevates and retracts the jaw.
- *Lateral Pterygoid*: The lateral pterygoid muscle has a triangular shape with two head, superior and inferior. It is the major protractor (opener) of the jaw.
  - Attachments: Both heads originate adjacent to the eye socket and converge into a tendon which attaches to the jaw.
  - Actions: Together, the lateral pterygoids protract the jaw, working independently to produce lateral movement.
- *Medial Pterygoid*: The medial pterygoid muscle has a quadrangular shape with two heads, deep and superficial. It is located inferior to the lateral pterygoid.
  - Attachments: The superficial head originates from the front of the skull and the deep head attaches to the skull adjacent to the eye socket. Both heads attach to the jaw.
  - Actions: Elevates the jaw and assists in the production of lateral movement.



Location of the temporalis muscle and the lateral pterygoid: (a) Highlighted in orange, the temporalis muscle is a broad muscle extending from zygomatic bone. (b) Arrows indicate the location of the lateral pterygoid, highlighted with the medial pterygoid in orange.

KEY MOVEMENTS

- **Elevation of the Jaw:** Produced by the masseter, temporalis and medial pterygoid.
- **Depression of the Jaw:** Produced by the lateral pterygoid, assisted by the digastric, mylohyoid and geniohyoid muscles found in the neck.
- **Protraction of the Jaw:** Produced by the lateral pterygoid.
- **Retraction of the Jaw:** Produced by the temporalis.
- **Lateral Movement of the Jaw:** Produced by the lateral and medial pterygoid.

This page titled 9.7B: Chewing Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.7C: Neck Muscles

Cervical muscles are those associated with the front of the neck; vertebral muscles are associated with the vertebral column.

Learning Objectives

- Outline the neck muscles and their movements

Key Points

- Numerous muscles contribute to the processes of speaking and swallowing.
- These muscles can be divided into suprahyoid and ingrahyoid groups based on their locations relative to the hyoid bone.
- The hyoid bone, located beneath the mandible, acts as a key attachment point for muscles involved in speaking and swallowing.
- Numerous muscles contribute to both the stabilization and fine movements of the head and neck.

Key Terms


- **suprahyoid muscles**: A group of muscles located above the hyoid bone, responsible for its elevation which widens the esophagus.
- **hyoid bone**: A -shaped bone which sits below the mandible
and in front of the esophagus, facilitating the wide range of movements associated with speaking and swallowing.
- **infrahyoid muscles**: A group of muscles located below the hyoid bone, responsible for its depression which narrows the esophagus.

Muscles of the neck play important roles in mastication (chewing), swallowing, speaking and supporting and moving the head. All muscles found in the neck are paired, meaning they exist to both the left and right side of the spine. The muscles involved in chewing have been discussed previously so only those involved in swallowing and support will be discussed below.

Muscles Involved in Swallowing and Speaking

Located to the anterior of the neck, these muscles are split into two based on their location relative to the hyoid bone. The U-shaped hyoid bone sits below the mandible and in front of the esophagus, providing a level of protection and facilitating the wide range of muscle activity required for speaking and swallowing.

Suprahyoid Muscles



Suprahyoid and infrahyoid muscles of the neck: Suprahyoid and infrahyoid muscle groups are named based on their location relative to the hyoid bone. The hyoid bone sits below the mandible and in front of the esophagus, providing a level of protection but also facilitating the wide range of muscle activity required for speaking and swallowing.

The four suprahyoid muscles found above the hyoid bone act in concert to elevate the hyoid bone, assisting with swallowing by widening the esophagus.


- *Stylohyoid*: The most superior of the suprahyoid muscles, the stylohyoid originates from the skull and attaches to the hyoid bone.
- *Digastric*: The digastric muscle is split into two parts that are connected by a tendon attached to the hyoid bone. The anterior section originates from the mandible and the posterior section from the skull.
- *Mylohyoid*: The mylohyoid is a broad flat muscle which forms the floor of the oral cavity. It originates from the mandible and attaches to the hyoid bone.
- *Geniohyoid*: The deepest of the suprahyoid muscles, the geniohyoid muscle originates from the mandible and attaches to the hyoid bone.

Infrahyoid Muscles

The four infrahyoid muscles found below the hyoid bone act in concert to depress the hyoid bone during swallowing and speaking, compressing the esophagus.

- *Sternohyoid*: A superficial muscle which originates from the sternum and attaches onto the hyoid bone.
- *Omohyoid*: Located laterally to the sternohyoid, the omohyoid muscle is split in two parts attached by a tendon. The inferior region originates from the scapula, joins the superior region, and attaches to the hyoid bone.
- *Sternothyroid*: Sitting deeper than the sternohyoid, the sternothyroid originates from the sternum and attaches to the thyroid cartilage associated with the hyoid bone.
- *Thyrohyoid*: A short continuation of the sternothyroid muscle, the thyrohyoid originates from the thyroid cartilage and attaches to the hyoid bone.

Muscles of the Back and Neck



Muscles of the back and neck: Muscles of the back and neck play an important role in maintaining posture and the movement of the head and neck.

The muscles of the back and neck are responsible for maintaining posture and facilitating movement of the head and neck. They are divided into three layers.

Superficial Layer

Two muscles in the superficial layer are responsible for rotation of the head.

- *Splenius Capitis*: A thick rectangular muscle, the most superior of the neck muscles.
  - Attachments: Originates from the upper spine and attaches to the skull.
  - Actions: Rotates and extend the head and neck.
- *Splenius Cervicis*: A small triangular-shaped muscle located immediately below the splenius capitis.
  - Attachments: Originates from the spine and attaches several vertebrae higher.
  - Actions: Rotate and extend the head and neck.

Intermediate Layer

Three columnar muscles in the intermediate layer are responsible for flexion and extension of the neck as well as posture maintenance. All three originate from a common tendon associated with the pelvis and can be divided into thoracic, cervicis, and capitis regions.

- *Iliocostalis*: The most laterally located of the three intermediate muscles.
  - Attachments: Originates from the common tendon and attaches to the ribs and lower neck.
  - Actions: Extends and controls abduction and adduction of the spine and neck.
- *Longissimus*: Located between the iliocostalis and spinalis muscles, this is the largest of the intermediate layer muscles.
  - Attachments: Originates from the common tendon and attaches to the lower ribs, the spine, and the skull.
  - Actions: Extends and controls abduction and adduction of the spine and neck.
- *Spinalis*: The most medially located and smallest of the three intermediate layer muscles.
  - Attachments: Originates from the common tendon and attaches to the upper spine and skull.
  - Actions: Extends and flexes to control abduction and adduction of the spine and neck.

Deep Layer

Two muscles in the deep layer are responsible for maintenance of posture and rotation of the neck.

- *Semispinalis*: The semispinalis is the most superficial of the deep muscles.
  - Attachments: A broad origin on the upper regions of the spine, with each origin attaching several vertebrae higher or to the skull.
  - Actions: Extends and rotates the head and maintains posture.
- *Multifidus*: The multifidus is located underneath the semispinalis muscle and is key in maintaining posture.
  - Attachments: A broad origin up the length of the spine, with each origin attaching several vertebrae higher.
  - Actions: Maintains posture through the spine.

Other Muscles That Act on the Neck

Several other muscles act on the head and neck. Below are three with a larger impact.

- *Trapezius*: The trapezius is the most superficial muscle of the back and forms a broad flat triangle.
  - Attachments: The trapezius originates from the skull and spine of the upper back and neck. It attaches to the clavicle and scapula.
  - Actions: The superior region supports the arm and elevates and rotates the scapula. It controls adduction, abduction and rotation of the head, the intermediate region retracts the scapula, and the inferior region rotates and depresses the scapula.
- *Sternocleidomastoid*: A thick rectangular muscle that is responsible for many movements within the neck.
  - Attachments: Dual-headed, the sternocleidomastoid originates from the clavicle and the sternum and attaches to the mandible.
  - Actions: Abduction, adduction, extension, flexion, and rotation of the neck depending on intra and inter-muscle contractions.
- *Platysma*: A broad sheet of muscle arising from the fascia covering the pectorals.
  - Attachments: Originates from the fascia covering the pectorals and attaches to various locations within the mandible and dermis of the face and neck.
  - Actions: Depresses the mandible and angles the lip and mouth, wrinkling the skin upon the neck flexing.

KEY MOVEMENTS

- **Extension** (tilting head backwards): Produced by the semispinalis, splenus capitis, longissimus, trapezius (superior fibers), and sternocleidomastoid (posterior fibers).
- **Flexion** (tilting head forwards): Produced by the sternocleidomastoid (anterior fibers).
- **Abduction** (tilting head towards shoulder): Produced by the sternocleidomastoid, longissimus, splenius capitis, semispinalis, and trapezius (superior fibers)
- **Adduction** (returning head to midline): Produced by the sternocleidomastoid, longissimus, splenius capitis, semispinalis, and trapezius (superior fibers)
- **Rotation** (rotation head to left or right): Produced by the sternocleidomastoid, longissimus, splenius capitis, semispinalis, and trapezius (superior fibers)

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Depressor supercilii muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Depress...ercilii_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Levator anguli oris. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Levator_anguli_oris**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Nasalis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nasalis_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Levator palpebrae superioris muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Levator...erioris_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Levator labii superioris alaeque nasi muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Levator...ue_nasi_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Occipitofrontalis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Occipitofrontalis_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dilator naris muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dilator_naris_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Superior tarsal muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Superior_tarsal_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Depressor septi nasi muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Depress...ti_nasi_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Levator labii superioris. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Levator_labii_superioris. **License**: *CC BY-SA: Attribution-ShareAlike*
- Buccinator muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Buccinator_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Procerus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Procerus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Zygomaticus major muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Zygomaticus_major_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Zygomaticus minor muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Zygomaticus_minor_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Risorius. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Risorius. **License**: *CC BY-SA: Attribution-ShareAlike*
- Depressor anguli oris muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Depressor_anguli_oris_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mentalis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mentalis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Depressor labii inferioris muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Depressor_labii_inferioris_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Orbicularis oris muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Orbicularis_oris_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...bicularis-oris. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...ontalis-muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- buccinator. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/buccinator. **License**: *CC BY-SA: Attribution-ShareAlike*
- Buccinator. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Buccinator.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray378. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray378.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Temporalis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Temporalis_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lateral pterygoid muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lateral_pterygoid_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Table of muscles of the human body. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Table_of_muscles_of_the_human_body. **License**: *CC BY-SA: Attribution-ShareAlike*
- Medial pterygoid muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Medial_pterygoid_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Masseter muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Masseter_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- masseter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/masseter. **License**: *CC BY-SA: Attribution-ShareAlike*
- lateral pterygoid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lateral%20pterygoid. **License**: *CC BY-SA: Attribution-ShareAlike*

- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/temporalis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Buccinator. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Buccinator.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray378. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray378.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1024. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1024.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Musculuspterygoideuslateralis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Musculuspterygoideuslateralis.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray382. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray382.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Suprahyoid muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Suprahyoid_muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Omohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Omohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platysma muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platysma_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sternocleidomastoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sternocleidomastoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scalene muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scalene_muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sternothyroid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sternothyroid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Longus capitis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Longus_capitis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Stylohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Stylohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scalenus medius. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scalenus_medius**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rectus capitis anterior muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rectus_capitis_anterior_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scalenus anterior. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scalenus_anterior**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Digastric muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digastric_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Longus colli muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Longus_colli_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyrohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thyrohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Geniohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Geniohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sternohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sternohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mylohyoid muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mylohyoid_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scalenus posterior. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scalenus_posterior**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rectus capitis lateralis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rectus_capitis_lateralis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Infrahyoid muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Infrahyoid_muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- suprahyoid muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/suprahyoid%20muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...n/longus-colli**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- platysma muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/platysma%20muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Buccinator. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Buccinator.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray378. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray378.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1024. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1024.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Musculuspterygoideuslateralis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Musculuspterygoideuslateralis.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray382. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray382.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- image389.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus389.html. **License**: *Public Domain: No Known Copyright*
- image386.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus386.html. **License**: *Public Domain: No Known Copyright*

This page titled 9.7C: Neck Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.8: TRUNK MUSCLES



- Topic hierarchy

9.8A: Anterior Muscles

9.8B: Posterior Muscles

This page titled 9.8: Trunk Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.8A: Anterior Muscles

The anterior muscles of the torso (trunk) are those on the front of the body, including the muscles of the chest, abdomen, and pelvis.

Learning Objectives

- Outline the anterior muscles of the torso

Key Points

- The intercostal muscles form the chest wall and function in respiration.
- The diaphragm is a sheet-like muscle that extends underneath the rib cage and aids in respiration by physically moving the lungs.
- The obliques are abdominal muscles that assist during bending and twisting of the torso.
- The rectus abdominis are the muscles often referred to as the “six-pack abs” and are involved in numerous aspects of trunk stabilization and bending.

Key Terms

- **diaphragm**: The key muscle in the control of respiration.
- **abdominal wall**: A layer of muscle and fascia which protects and encloses the abdominal cavity, allowing for its compression as well as torso movement.
- **linea alba**: A tough, fibrous line running down the midline of the abdomen, formed from the aponeuroses of the abdominal muscles.
- **intercostal**: Muscles forming the chest wall, which aid in respiration.

The anterior muscles of the trunk (torso) are associated with the front of the body, include chest and abdominal muscles. Chest muscles function in respiration while abdominal muscles function in torso movement and in maintenance of balance and posture.


Muscles of the Chest

Pectoral Muscles

Pectoral muscles lie in the chest and exert force through the shoulder to move the upper arm.

- *Pectoralis Major*: The pectoralis major is a fan-shaped muscle covering the chest and comprised of clavicular and sternocostal regions.
  - Attachments: The clavicular region originates from the clavicle and the sternocostal region originates from the sternum and the fascia of the oblique muscles of the abdomen. Both attach to the humerus.
  - Actions: Adducts and rotates the upper arm.
- *Pectoralis Minor*: The smaller pectoralis minor muscle lies beneath the pectoralis major.
  - Attachments: The pectoralis minor originates from the third to fifth ribs and attaches to the scapula.
  - Actions: Supports and depresses the scapula.
- *Serratus Anterior*: The serratus anterior is located in the lateral wall of the chest.
  - Attachments: The muscle is formed of several strips originating from the second to eighth ribs, each of which attaches to the scapula.
  - Actions: Supports the scapula, allowing for elevation of the upper arm.

Intercostal Muscles



Intercostal muscles of the anterior trunk: Deep muscles of the chest and front of the arm, with the boundaries of the axilla. The intercostals are muscles between the ribs that form the chest cavity wall.

Lying below the pectoral muscles, the intercostal muscles form the chest wall and play a key role in respiration. All intercostal muscles originate on the lower border of a rib and attach to the upper border of the rib below.

- *External Intercostals*: The external intercostals are the most superficial of the intercostal muscles. They are continuous with the external oblique muscle of the abdomen.
  - Actions: Elevate the ribs.
- *Internal Intercostals*: Lying below the external intercostals, the internal intercostals are continuous with the internal oblique muscle of the abdomen.
  - Actions: Elevate or depress the ribs.
- *Innermost Intercostals*: The deepest lying of the intercostals, these muscles are similar in structure to the internal intercostals.
  - Actions: Elevate or depress the ribs.

Other Muscles

Diaphragm: The diaphragm is a large, flat, sheet-like muscle that extends horizontally underneath the rib cage.


Functionally, the diaphragm separates the thoracic cavity, containing the lungs and heart and enclosed by the rib cage from the abdominal cavity, which contains the digestive organs. The diaphragm's position allows it to aid in respiration. When it contracts, it physically moves the lungs and deforms the volume of the thoracic cavity.

- Attachments: The diaphragm has several points of origin along the sternum, the lower ribs, and lower vertebrae. The muscle fibers combine into a central tendon, which ascends and attaches to the surface of the pericardium.
- Actions: Contracts, flattening and increasing the volume of the thoracic cavity. Relaxes and returns to original shape, reducing the volume of the thoracic cavity.

Muscles of the Abdomen

The skeletal muscles of the abdomen form part of the abdominal wall, which holds and protects the gastrointestinal system. Five muscles form the abdominal wall, divided into vertical and flat groups. The flat muscles act to flex, laterally flex, and rotate the trunk. The fibers run in different directions and cross each other, strengthening the abdominal wall. The vertical muscles aid in compressing the abdominal cavity, stabilizing the pelvis, and depressing the ribs when a person is walking. Toward the midline,

the muscles form aponeuroses, which merge into the linea alba.



Location of the external obliques: Highlighted in orange, the external obliques lie inferior to the pectoral muscles

- *External Oblique*: The external oblique is the largest and most superficial of the flat muscles.
  - Attachments: Originates from the lower ribs and attaches to the pelvis, forming an aponeurosis toward the midline and linea alba.
- *Internal Oblique*: Lying deep to the external oblique, the internal oblique is smaller and thinner. Its fibers run perpendicular to the external oblique, improving the strength of the abdominal wall.
  - Attachments: Originates from the pelvis and thoracolumbar fascia, running through the back. Attaches to the lower ribs and forms an aponeurosis toward the midline and linea alba.
- *Transversus Abdominis*: The deepest of the flat muscles, the transversus abdominis consists of transversely-running fibers.
  - Attachments: Originates from the lower ribs, thoracolumbar fascia, and pelvis, forming an aponeurosis toward the midline and linea alba.
- *Rectus Abdominis*: A long vertical muscle that covers the abdomen, lying below the flat muscles. It is split through the midline by the linea alba formed from the aponeuroses of the abdominal muscles and separated by horizontal tendinous intersections which give rise to the six pack.
  - Attachments: Originates from the pubis and attaches to the lower edge of the rib cage and sternum.
- *Pyramidalis*: Lying superficial to the rectus abdomini,s the pyramidalis is a small, triangular vertical muscle.
  - Attachments: Originates from the pubis and attaches to the linea alba.

This page titled 9.8A: Anterior Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.8B: Posterior Muscles

Muscles of the posterior portion of the trunk include muscles of the back, suboccipital region, and perineum region.

Learning Objectives


- Outline the posterior muscles of the torso

Key Points

- The back is characterized by numerous muscle groups which allow movement of the shoulder, head, and neck, as well as aid in respiration and maintain posture and balance.
- The superficial muscles of the back are responsible for movement of the shoulder.
- The intermediate muscles of the back assist in the movement of the rib cage during respiration.
- The intrinsic back muscles facilitate movement of the head and neck and are fundamental in maintaining posture and balance.

The posterior or back muscles perform a wide range of functions, including movement of the shoulder, head, and neck and assisting in respiration, posture, and balance. Posterior muscles are split into three groups depending on their physiological location.

Superficial Posterior Muscles



Location of the latissimus dorsi muscle: Highlighted in orange, the latissimus dorsi is a muscle of the posterior torso.

The superficial posterior muscles are associated with movement of the shoulder. As the name suggests, they are the most superficially located of the muscles covering the intermediate and intrinsic layers.

- *Trapezius*: The trapezius is the most superficial muscle of the back and forms a broad flat triangle.
  - Attachments: The trapezius originates from the skull and spine of the upper back and neck. It attaches to the clavicle and scapula.
  - Actions: The superior region supports the arm and elevates and rotates the scapula, the intermediate region retracts the scapula, and the inferior region rotates and depresses the scapula.

- *Latissimus Dorsi*: The latissimus dorsi originates from the lower back and covers a wide area.
  - Attachments: The latissimus dorsi originates from the lower spine and ribs and the upper pelvis and fascia of the deep trunk muscles. The muscle converges into a tendon attaching to the humerus.
  - Actions: Extends, adducts, and medially rotates the upper arm.
- *Levator Scapulae*: A small, strap-like muscle that joins the neck to the scapula.
  - Attachments: Originates from the side of the spine in the neck and attaches to the scapula.
  - Actions: Elevates the scapula.
- *Rhomboid Major*: Sits inferiorly to the levator scapulae.
  - Attachments: Originates from the spine in the upper back and attaches to the scapula inferior to the levator scapulae attachment.
  - Actions: Retracts and rotates the scapula.
- *Rhomboid Minor*: Sits between the levator scapulae and rhomboid major, with which it is paired in action and function, this retracts and rotates the scapula.

Intermediate Posterior Muscles

The intermediate muscles of the posterior contribute to movements of the ribcage during respiration.

Serratus Posterior Superior – The serratus posterior superior is a thin, rectangular-shaped muscle lying below the rhomboid muscles.

- Attachments: Originates from the lower spine and attaches to ribs 2 through 5.
- Actions: Elevates ribs 2 through 5.


Serratus Posterior Inferior: The serratus posterior inferior is a broad muscle lying beneath the latissimus dorsi.

- Attachments: Originates from the spine and attaches to ribs 9 through 12.
- Actions: Depresses ribs 9 through 12.

Intrinsic Posterior Muscles

The intrinsic muscles of the posterior are responsible for maintaining posture and facilitating movement of the head and neck. They are divided into three layers.

Superficial Layer



Location of the splenius muscle.: The splenius capitis is highlighted in orange, with the splenius cervicis directly below

Two muscles in the superficial layer are responsible for rotation of the head.

- *Splenius Capitis*: This thick rectangular muscle is the most superior of the next muscles.
  - Attachments: Originates from the upper spine and attaches to the skull.
  - Actions: Rotates and extends the head and neck.
- *Splenius Cervicis*: A small triangular-shaped muscle located immediately below the splenius capitis.
  - Attachments: Originates from the spine and attaches several vertebrae higher.
  - Actions: Rotates and extends the head and neck.

Intermediate Layer


Three columnar muscles in the intermediate layer are responsible for flexing and extending the neck as well as maintaining posture. All three originate from a common tendon associated with the pelvis.

- *Iliocostalis*: The most laterally located of the three intermediate muscles.
  - Attachments: Originates from the common tendon and attaches to the ribs and lower neck.
  - Actions: Extends and controls abduction and adduction of the spine and neck.
- *Longissimus*: Located between the iliocostalis and spinalis muscles, it is the largest of the intermediate layer muscles.
  - Attachments: Originates from the common tendon and attaches to the lower ribs, spine, and skull.
  - Actions: Extends and controls abduction and adduction of the spine and neck.
- *Spinalis*: The most medially-located and smallest of the three intermediate layer muscles.
  - Attachments: Originates from the common tendon and attaches to the upper spine and skull.
  - Actions: Extends, flexes, and controls abduction and adduction of the spine and neck.

Deep Layer

Two muscles in the deep layer are responsible for maintenance of posture and rotation of the neck.

- *Semispinalis*: The semispinalis is the most superficial of the deep muscles.
  - Attachments: A broad origin on the upper regions of the spine, with each origin attaching several vertebrae higher or to the skull.
  - Actions: Extends and rotates the head and maintains posture.
- *Multifidus*: The multifidus is located underneath the semispinalis muscle, and is key in maintaining posture.
  - Attachments: A broad origin up the length of the spine, with each origin attaching several vertebrae higher.
  - Actions: Maintains posture through the spine.



Location of the multifidus muscle: Highlighted in orange, the multifidus muscle is a muscle of the posterior trunk and lies interior to a majority of muscles

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Innermost intercostal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Innermost_intercostal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pelvic floor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pelvic_floor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pyramidalis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pyramidalis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Transversus abdominis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Transversus_abdominis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Levatores costarum muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Levatores_costarum_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- External intercostal muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/External_intercostal_muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Transversus thoracis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Transversus_thoracis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Abdominal internal oblique muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Abdominal_internal_oblique_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intercostal muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intercostal_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Subcostales muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Subcostales_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Abdominal external oblique muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Abdominal_external_oblique_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Internal intercostal muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Internal_intercostal_muscles**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thoracic diaphragm. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thoracic_diaphragm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Iliococcygeus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Iliococcygeus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Puborectalis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Puborectalis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coccygeus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coccygeus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rectus abdominis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rectus_abdominis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Quadratus lumborum muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Quadratus_lumborum_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pelvic diaphragm. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pelvic_diaphragm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cremaster muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cremaster_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pubococcygeus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pubococcygeus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Levator ani. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Levator_ani**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- rectus abdominis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/rectus%20abdominis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- levatores costarum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/levatores%20costarum**. **License**: ***CC BY-***

SA: Attribution-ShareAlike
- Gray392. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray392.png. **License**: CC BY-SA: Attribution-ShareAlike
- Gray411. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray411.png. **License**: CC BY-SA: Attribution-ShareAlike
- Superficial perineal pouch. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Superfi...perineal_pouch. **License**: CC BY-SA: Attribution-ShareAlike
- Rectus capitis posterior minor muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rectus_...r_minor_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Transversus perinei superficialis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Transve...icialis_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Serratus posterior inferior muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serratu...nferior_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Spinalis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinalis. **License**: CC BY-SA: Attribution-ShareAlike
- Splenius cervicis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenius_cervicis_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Splenius capitis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenius_capitis_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Erector spinae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Erector_spinae. **License**: CC BY-SA: Attribution-ShareAlike
- Iliocostalis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Iliocostalis. **License**: CC BY-SA: Attribution-ShareAlike
- Longissimus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Longissimus. **License**: CC BY-SA: Attribution-ShareAlike
- Serratus posterior superior muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serratus_posterior_superior_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Transversospinales muscles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Transversospinales_muscles. **License**: CC BY-SA: Attribution-ShareAlike
- Sphincter ani internus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sphincter_ani_internus_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Transversus perinei profundus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Transversus_perinei_profundus_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Semispinalis dorsi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Semispinalis_dorsi. **License**: CC BY-SA: Attribution-ShareAlike
- Rectus capitis posterior major muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rectus_capitis_posterior_major_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Rotatores muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rotatores_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Deep perineal pouch. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Deep_perineal_pouch. **License**: CC BY-SA: Attribution-ShareAlike
- Semispinalis capitis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Semispinalis_capitis. **License**: CC BY-SA: Attribution-ShareAlike
- Multifidus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Multifidus_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Latissimus dorsi muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Latissimus_dorsi_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Bulbospongiosus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bulbospongiosus_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Obliquus capitis superior muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Obliquus_capitis_superior_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Intertransversarii muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intertransversarii_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Obliquus capitis inferior muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Obliquus_capitis_inferior_muscle. **License**: CC BY-SA: Attribution-ShareAlike
- Sphincter urethrae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sphincter_urethrae. **License**: CC BY-SA: Attribution-ShareAlike
- Sphincter ani externus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/

Sphincter_ani_externus_muscle. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Semispinalis cervicis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Semispinalis_cervicis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ischiocavernosus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ischiocavernosus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Interspinales muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Interspinales_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- compressor urethrae. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/compressor%20urethrae**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- multifidus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/multifidus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/internal-urethral-sphincter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray392. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray392.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray411. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray411.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Multifidi. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Multifidi.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Musculus splenius capitis marked. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Musculus_splenius_capitis_marked.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Latissimus dorsi. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Latissimus_dorsi_.PNG**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 9.8B: Posterior Muscles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.9: MUSCLES OF THE UPPER LIMB

- Topic hierarchy

9.9A: Muscles of the Humerus that Act on the Forearm

9.9B: Muscles of the Wrist and Hand

9.9C: Muscles of the Shoulder

9.9D: Muscles that Cause Movement at the Ankle

This page titled 9.9: Muscles of the Upper Limb is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.9A: Muscles of the Humerus that Act on the Forearm

Humerus that act on the forearm are primarily involved in flexion and extension.

Learning Objectives

- Diagram the movements of the humerus muscles that act on the forearm

Key Points

- Muscles of both the upper arm and forearm control movement of the forearm.
- The biceps brachii flex the forearm and work with the supinator of the forearm to rotate it so the palm faces upward.
- The triceps brachii extend the forearm.
- The pronator teres and quadratus control pronation, or rotation of the forearm so that the palm faces downward.

Key Terms


- **Pronator Teres**: A muscle of the anterior compartment of the forearm that controls pronation.
- **Supinator**: A muscle of the posterior compartment of the forearm that controls supination.
- **Pronator Quadraturs**: A muscle of the anterior compartment of the forearm that controls pronation.
- **Brachioradialis**: A muscle of the posterior compartment of the forearm that flexes the forearm.
- **Biceps Brachii**: A muscle of the anterior compartment of the upper arm that flexes the forearm.
- **Triceps Brachii**: A muscle of the posterior compartment of the upper arm that extends the forearm.

The humerus is a long bone in the arm that runs from the shoulder to the elbow. Anatomically, it interacts with the scapula to form the shoulder joint and the radius and ulna of the lower arm to form the elbow joint. Forearm rotation is controlled by two joints: the proximal radioulnar joint which exists immediately below the elbow, and the distal radioulnar joint located immediately before the wrist.

Upper Arm

There are four muscles in the upper arm split into an anterior and posterior compartment.

Anterior Compartment



Superficial muscles of the chest and upper arm: The biceps brachii is located in the anterior compartment of the upper arm and flexes and supinates the forearm at the elbow.

Three muscles are located in the anterior compartment of the upper arm.


- *Biceps Brachii*: The biceps brachii is a two-headed muscle. Although the majority of the muscle mass is located anteriorly to the humerus, it has no attachment to the bone itself.
  - Attachments: Both heads originate from the scapula and attach via the bicipital aponeurosis to the fascia of the forearm.
  - Action: Supination of the forearm. It also flexes the arm at the elbow and at the shoulder.
- *Coracobrachialis*: The coracobrachialis lies within the two heads of the biceps brachii.
  - Attachments: Originates from the scapula and attaches to the humerus.
  - Action: Flexing of the arm at the shoulder, and weak adduction.
- *Brachialis*: The brachialis muscle lies within the distal region of the biceps brachii.
  - Attachments: Originates from the humerus and attaches to the ulna.
  - Action: Flexing of the arm at the elbow.

Posterior Compartment

The posterior compartment of the upper arm contains only one muscle.

- *Triceps Brachii*: The triceps brachii is a three-headed muscle.
  - Attachments: The long head originates from the scapula, the lateral head from the proximal region of the humerus, and the medial head from the distal region of the humerus. All three converge into one tendon which attaches to the ulna.
  - Action: Extension of the arm at the elbow.

Forearm



Superficial muscles of the posterior forearm: The anconeus, located in the superficial region of the posterior forearm compartment, moves the ulna during pronation and extends the forearm at the elbow.

As with the upper arm, the forearm is split into anterior and posterior compartments. Only those responsible for movement of the forearm are discussed below; the muscles responsible for movement of the hand and wrist are discussed in a later section.

Anterior

The anterior compartment of the forearm is split into superficial, intermediate, and deep regions.

- *Pronator Teres*: A rectangular muscle located in the superficial region of the anterior compartment.
  - Attachments: The pronator teres has two origins, one on the proximal end of the humerus and one of the distal end of the ulna. It attaches to the mid region of the radius.
  - Action: Pronates the forearm.
- *Pronator Quadratus*: A square shaped muscle located adjacent to the wrist in the deep region of the anterior compartment.
  - Attachments: Originates from the ulna and attaches to the radius.
  - Action: Pronates the forearm.

Posterior

The posterior compartment of the forearm is split into superficial and deep regions.

- *Anconeus*: The anconeus is located in the superficial region of the forearm posterior compartment and is blended with the triceps brachii.
  - Attachments: Originates from the humerus and attaches to the ulna.
  - Action: Moves the ulna during pronation and extends the forearm at the elbow.
- *Brachioradialis*: The brachioradialis is located in the superficial region of the forearm posterior compartment,
  - Attachments: Originates from the humerus and attaches to the distal end of the radius.
  - Action: Flexes the forearm at the elbow.
- *Supinator*: The supinator is located in the deep region of the forearm posterior compartment.
  - Attachments: The supinator has two heads: one originating from the humerus, the other from the ulna. Together they attach to the radius.
  - Action: Supinates the forearm.

KEY MOVEMENTS

- **Extension** (forearm away from upper arm): Produced by the triceps brachii and anconeus of the forearm.
- **Flexion** (forearm towards upper arm): Produced by the brachialis, biceps brachii, and brachioradialis of the forearm.
- **Pronation** (rotation of the forearm so the palm faces downwards): Produced by the pronator quadratus and pronator teres of the forearm.
- **Supination** (rotation of the forearm so the palm faces upwards): Produced by the supinator of the forearm and biceps brachii.

This page titled 9.9A: Muscles of the Humerus that Act on the Forearm is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.9B: Muscles of the Wrist and Hand

Muscles in the forearm move the wrists, and hand movement is caused by both forearm and hand muscles.

Learning Objectives

- Outline the movements of the wrist and hand muscles


Key Points

- Movements of the wrist include abduction, adduction, extension, and flexion.
- Movements of the fingers and thumb include abduction, adduction, extension, and flexion.
- Rotation of the thumb and little finger allows for opposition.
- Muscles of the forearm that act on the wrist and hand are referred to as extrinsic muscles, or external to the hand.
- Muscles within the wrist and hand are referred to as intrinsic muscles.

Key Terms

- **Palmaris Longus**: A long muscle originating near the elbow and passing through into the wrist, attaching to the base of the hand.
- **Flexor Digitorum Superficialis**: A key muscle controlling wrist and finger flex.
- **Flexor Carpi Ulnaris**: A long muscle originating near the elbow and passing through into the wrist, attaching to one of the carpal bones in the wrist.
- **Flexor Carpi Radialis**: A long muscle originating near the elbow and passing through into the wrist, attaching to the base of the digits (fingers).
- **Flexor Digitorum Profundus**: A long muscle originating near the elbow and passing through into the wrist, flexing the wrist and the most distant regions of the fingers.
- **Pronator Teres**: A rectangular muscle that pronates the forearm.
- **Flexor Pollicis Longus**: A long, deep muscle responsible for flexing the thumb.
- **Pronator quadratus**: A square-shaped muscle located adjacent to the wrist.

Muscles associated with the wrist include those of the forearm and hand that move the wrist and digits. The wrist and hand exhibit a remarkable range of movement, key for grasping and interaction with objects. These muscles can generate highly variable force, from the strong grip required when lifting a heavy object to the delicate movements required to write.



Muscles and tendons of the forearm and hand: The extrinsic muscles of the forearm are responsible for movement of the wrist and fingers. Often providing the more forceful movements required.

Muscles of the forearm that act on the wrist and hand are referred to as extrinsic muscles, or external to the hand. Those located within the hand are referred to as intrinsic.

Muscles of the Forearm

As with the upper arm, the forearm is split into anterior and posterior compartment. Each contains many more muscles than described below due to the requirement for more complex movements in the wrist and hand.

Anterior

The anterior compartment of the forearm is split into superficial, intermediate, and deep layers. These muscles are generally responsible for flexing of the wrist and fingers and pronation of the forearm.

Superficial Layer

Three muscles are located in the superficial layer of the anterior compartment of the forearm.

- *Flexor Carpi Ulnaris*: A long muscle originating near the elbow and passing through into the wrist.
  - Attachments: Originates from the humerus and ulna and attaches to one of the carpal bones in the wrist.
  - Actions: Flexion and adduction at the wrist.
- *Palmaris Longus*: A long muscle originating near the elbow and passing through into the wrist.
  - Attachments: Originates from the humerus and attaches to the base of the hand.
  - Actions: Flexion at the wrist.
- *Flexor Carpi Radialis*: A long muscle originating near the elbow and passing through into the wrist.
  - Attachments: Originates from the humerus and attaches to the base of the digits.
  - Actions: Flexion and abduction at the wrist.
- *Pronator Teres*: A rectangular muscle.
  - Attachments: The pronator teres has two origins, one on the proximal end of the humerus and one of the distal end of the ulna. It attaches to the mid region of the radius.
  - Actions: Pronates the forearm.

Intermediate Layer

There is just one muscle in the intermediate layer of the anterior compartment of the forearm.

- *Flexor Digitorum Superficialis*: Lying below the superficial region, the flexor digitorum superficialis is a key muscle controlling wrist and finger flex.
  - Attachments: Originates from the humerus and the radius, splitting into four tendons at the wrist which travel through the carpal tunnel and attach to the fingers.
  - Actions: Flexes fingers and wrist.

Deep Layer

There are three muscles in the deep layer of the anterior compartment of the forearm.

- *Flexor Digitorum Profundus*: A long muscle originating near the elbow and passing through into the wrist, lying adjacent to the flexor pollicis longus.
  - Attachments: Originates from the ulna, splitting into four tendons at the wrist which travel through the carpal tunnel and attach distally to the fingers.
  - Actions: Flexes the wrist and the most distal regions of the fingers.
- *Flexor Pollicis Longus*: A long muscle originating near the elbow and passing through into the wrist, lying adjacent to the flexor digitorum profundus.
  - Attachments: Originates from the radius and attaches to the base of the thumb.
  - Actions: Flexes the thumb.
- *Pronator quadratus*: A square-shaped muscle located adjacent to the wrist.
  - Attachments: Originates from the ulna and attaches to the radius.
  - Actions: Pronates the forearm.

Posterior

The posterior compartment of the forearm is split into superficial and deep regions. The muscles are generally responsible for extension of the wrist and fingers.

Superficial Layer

The superficial layer of the posterior forearm contains seven muscles.

- *Aconeus*: The aconeus is located in the superficial region of the forearm posterior compartment and is blended with the triceps brachii.
  - Attachments: Originates from the humerus and attaches to the ulna.
  - Actions: Moves the ulna during pronation and extends the forearm at the elbow.
- *Brachioradialis*: The brachioradialis is located in the superficial region of the forearm posterior compartment.
  - Attachments: Originates from the humerus and attaches to the distal end of the radius.
  - Actions: Flexes the forearm at the elbow.
- *Extensor Carpi Radialis Longus and Brevis*: A pair of muscles located on the side of the forearm, allowing them to control extension and abduction of the wrist.
  - Attachments: Both originate from the humerus and attach to the base of the hand.
  - Actions: Extend and abduct the wrist.
- *Extensor Digitorum*: The extensor digitorum is the main extensor of the fingers.
  - Attachments: Originates from the humerus, splitting into four tendons at the wrist which travel through the carpal tunnel and attach to the digits.
  - Actions: Extends fingers.
- *Extensor Digiti Minimi*: Originates from the extensor digitorum. In some people, these muscles cannot be individually defined.
  - Attachments: Originates from the humerus and attaches to the little finger.
  - Actions: Extends the little finger, and contributes to extension at the wrist.
- *Extensor Carpi Ulnaris*: Located on the other side of the forearm to the extensor carpi radialis longus and brevis, it performs a similar role.

- Attachments: Originates from the humerus and attaches to the base of the hand.
- Actions: Extension and adduction of wrist.

Deep Layer


There are four muscles in the deep layer of the posterior compartment of the forearm.

- *Supinator*: The supinator is located in the deep region of the forearm posterior compartment.
  - Attachments: The supinator has two heads: one originates from the humerus, the other from the ulna. Together they attach to the radius.
  - Actions: Supinates the forearm.
- *Abductor Pollicis Longus*: The abductor pollicis longus is situated immediately distal to the supinator muscle.
  - Attachments: Originates from the radius and ulna attaching to the base of the thumb.
  - Actions: Abducts the thumb.
- *Extensor Pollicis Brevis*: The extensor pollicis brevis is located below the abductor pollicis longus.
  - Attachments: Originates from radius and attaches to the base of the thumb.
  - Actions: Extends the thumb.
- *Extensor Indicis Proprius*: This muscle allows the index finger to be independent of the other fingers during extension.
  - Attachments: Originates from the ulna and attaches to the index finger.
  - Actions: Extends the index finger.

Muscles of the Hand

The extrinsic muscles of the hand are responsible for the larger scale, stronger movements of the wrist and hand. The intrinsic muscles produce finer, more controlled movements and play important roles in maintaining grip.

Thenar Muscles



Muscles of the hand

The thenar muscles are three short muscles located at the base of the thumb and responsible for its fine movement.

- *Opponens Pollicis*: The opponens pollicis is the largest and deepest-lying of the thenar muscles.
  - Attachments: Originates from the wrist and attaches to the thumb.
  - Actions: Rotates the thumb towards the palm, producing opposition and improving grip.
- *Abductor Pollicis Brevis* : Located anteriorly to the opponens pollicis and proximal to the flexor pollicis brevis.
  - Attachments: Originates from the wrist and attaches to the thumb.

    - Actions: Abducts the thumb.
- *Flexor Pollicis Brevis*: The smallest and most distal of the thenar muscles.
    - Attachments: Originates from the wrist and attaches to the thumb.
    - Actions: Flexes the thumb.

Hypothenar Muscles

The hypothenar muscles are located at the base of the little finger. Their naming, function, and organization are similar to those of the thenar muscles.

- *Opponens Digiti Minimi*: The opponens digit minimi is the deepest-lying of the hypothenar muscles.
    - Attachments: Originates from the wrist and attaches to the little finger.
    - Actions: Rotates little finger towards the palm, producing opposition and improving grip.
- *Abductor Digiti Minimi*: The most superficial of the hypothenar muscles.
    - Attachments: Originates from the wrist and attaches to the little finger.
    - Actions: Abducts the little finger.
- *Flexor Digiti Minimi Brevis*: Located laterally to the digiti minimi.
    - Attachments: Originates from the wrist and attaches to the little finger.
    - Actions: Flexes little finger.

Lumbricals

These are four lumbricals in the hand, each associated with an individual finger.

- Attachments: Originates from a tendon of attached to the flexor digitorum profundus of the forearm, each attaching to an individual finger
- Actions: Flexes and extends the fingers.

Interossei

The interossei muscles are located between the fingers; they can be split into two groups.

- *Dorsal Interossei*: Located superficially on the dorsal side of the hand, there are four dorsal interossei muscles.
    - Attachments: Originates from the base of the finger, each attaching after the first finger joint.
    - Actions: Abducts the fingers.
- *Palmar Interossei*: Located on the anterior side of the hand, there are three palmar interossei, with the index finger controlled by the extensor indicis proprius.
    - Attachments: Originates from the base of the finger, each attaching after the first finger joint.
    - Actions: Adducts the fingers.

Other Muscles

One other muscle in the hand is not easily grouped with the above categories.

- *Palmaris Brevis*: The palmaris brevis is a small superficial muscle found in the palm.
    - Attachments: Originates from the fascia of the palm and attaches to the dermis.
    - Actions: Wrinkles the skin and deepens the curvature of the palm improving grip.

This page titled 9.9B: Muscles of the Wrist and Hand is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.9C: Muscles of the Shoulder

Muscles of the shoulder include those that attach to the bones of the shoulder to move and stabilize the joint.

Learning Objectives

- Outline the movements of the muscles of the shoulder

Key Points

- The shoulder exhibits a wide range of movement, which makes it susceptible to dislocation and injury.
- The trapezius muscles rotate the scapulae upward.
- The rhomboid major and the rhomboid minor press the scapula against the thoracic wall, retracting the scapula towards the spine.
- The deltoid is a complex muscle that forms the rounded edge of the shoulder and participates in many articulations of the shoulder joint.
- The rotator cuff are the muscles that stabilize movement of the shoulder.
- The pectoralis minor and pectoralis major are large muscles of the chest that participate in many movements, including flexion of the humerus.

Key Terms

- **pectoralis major**: A large, fan-shaped muscle of the chest.
- **rotator cuff**: A set of four smaller muscles in the shoulder responsible for rotating the humerus (upper arm bone).
- **trapezius**: A large vertebrate skeletal muscle divided into an ascending, descending, and transverse portion, attaching the neck and central spine to the outer extremity of the scapula. It functions in scapular elevation, adduction, and depression.
- **deltoid**: The deltoid muscle, a triangular muscle on the human shoulder.

The shoulder or glenohumeral joint is a ball and socket joint formed between the humerus and scapula. Due to the shallowness of the socket and relatively loose connections, the shoulder joint allows for a wide range of motion; however, this wide range makes the joint unstable and thus more prone to dislocation and injury than other joints.


Two other joints make up the shoulder; the acromioclavicular joint of the clavicle and scapula, which allows the arm to be raised above the head, and the sternoclavicular joint of the clavicle and sternum, which plays an important role in facilitating movement of the upper arm and connecting it to the rest of the skeleton.

Muscles that act on the shoulder can be classified as extrinsic, intrinsic, pectoral, or upper arm. Upper arm muscles will be discussed in a later section since they primarily promote forearm movement.

Extrinsic Shoulder Muscles

Extrinsic muscles of the shoulder originate from the trunk and attach to the bones of the shoulder. They can be further subdivided into superficial and deep layers.

Superficial



Location of the trapezius muscle: Highlighted in orange, the trapezius is a large, broad muscle of the back that acts on the shoulder.

As suggested by the name, superficial muscles lie on the surface. There are two superficial extrinsic muscles.


- *Trapezius*: The trapezius is the most superficial muscle of the back and forms a broad flat triangle.
  - Attachments: The trapezius originates from the skull and spine of the upper back and neck. It attaches to the clavicle and scapula.
  - Actions: The superior region supports the arm and elevates and rotates the scapula, the intermediate region retracts the scapula, and the inferior region rotates and depresses the scapula.
- *Latissimus Dorsi*: The latissimus dorsi originates from the lower back and covers a wide area.
  - Attachments: The latissimus dorsi originates from the lower spine and ribs and the upper pelvis and fascia of the deep trunk muscles. The muscle converges into a tendon attaching to the humerus.
  - Actions: Extends, adducts, and medially rotates the upper arm.

Deep

Three deep muscles lie below the superficial muscles of the shoulder.

- *Levator Scapulae*: A small, strap-like muscle that joins the neck to the scapula.
  - Attachments: Originates from the side of the spine in the neck and attaches to the scapula.
  - Actions: Elevates the scapula.
- *Rhomboid Major*: Sits inferiorly to the levator scapulae.
  - Attachments: Originates from the spine in the upper back and attaches to the scapula in an inferior position to the levator scapulae attachment.
  - Actions: Retracts and rotates the scapula.
- *Rhomboid Minor*: Sits between the levator scapulae and rhomboid major, with which it is paired in action and function. It retracts and rotates the scapula.

Intrinsic




Location of the deltoid muscles: Highlighted in orange, the deltoids cover the rounding of the shoulder joint.

Intrinsic muscles originate from the scapula or clavicle and attach to the humerus. There are six intrinsic muscles, four of which form the rotator cuff.

- *Deltoid*: The deltoid muscle is a triangular muscle which covers the shoulder. The action of the muscle is complex, with the components acting in opposing and separate ways during the course of a contraction.
  - Attachments: The deltoid muscle originates from the scapula and clavicle and attaches to the lateral surface of the humerus.
  - Actions: The anterior region assists the pectoralis major during transverse flexion of the shoulder and acts weakly in strict transverse flexion. The lateral region assists in shoulder flexion when the shoulder is rotating, although it also assists the transverse abduction of the shoulder. The posterior region is the hyperextensor of the shoulder, contributing to transverse
    extension.
- *Teres Major*: The teres major is a thick flattened muscle connecting the lower scapula with the humerus.
  - Attachments: Originates from the posterior of the scapula and attaches to the humerus.
  - Actions: Adducts the shoulder and assists in rotation of the arm.

Rotator Cuff

The rotator cuff is a group of four muscles that pull the ball of the humerus into the shallow socket of the scapula, adding required stability. The rotator cuff complex is composed of the supraspinatus, infraspinatus, subscapularis, and teres minor all of which originate from the scapula and connect to the humerus. The supraspinatus is involved in abduction of the arm in association with the deltoid, while the other muscles facilitate rotation of the arm.



Muscles of the rotator cuff: Muscles of the rotator cuff and presented with the triceps brachii.

Pectoral

Pectoral muscles lie in the chest and exert force through the shoulder to move the upper arm. Three pectoral muscles interact with the shoulder.

- *Pectoralis Major*: The pectoralis major is a large, fan-shaped muscle covering the chest. It is comprised of clavicular and sternocostal regions.
  - Attachments: The clavicular region originates from the clavicle and the sternocostal region originates from the sternum and the fascia of the oblique muscles of the abdomen. Both attach to the humerus.
  - Actions: Adducts and rotates the upper arm.
- *Pectoralis Minor*: The pectoralis minor muscle is smaller and lies beneath the pectoralis major.
  - Attachments: The pectoralis minor originates from the third to fifth ribs and attaches to the scapula.
  - Actions: Supports and depresses the scapula.
- *Serratus Anterior*: The serratus anterior is located in the lateral wall of the chest.
  - Attachments: The muscle is formed of several strips originating from the second to eight ribs, each of which attaches to the scapula.
  - Actions: Supports the scapula allowing for elevation of the upper arm.

KEY MOVEMENTS

- **Extension** (upper limb backwards behind back): Produced by the posterior deltoid, latissimus dorsi, and teres major.
- **Flexion** (upper limb forwards past chest): Produced by the biceps brachii (both heads), pectoralis major, anterior deltoid, and corocobrachialis.
- **Abduction** (upper limb away from trunk, spreading arms wide): Produced by the supraspinatus and deltoid. Past 90 degrees, the scapula needs to be rotated by the trapezius and serratus anterior to achieve abduction.
- **Adduction** (upper limb towards trunk, bringing arms down to side): Produced by contraction of pectoralis major, latissimus dorsi, and teres major.
- **Medial Rotation** (rotation of arm inwards to cover abdomen): Produced by contraction of subscapularis, pectoralis major, latissimus dorsi, teres major, and anterior deltoid.
- **Lateral Rotation** (rotation of arm outwards away from the abdomen): Produced by contraction of the infraspinatus and teres minor.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Biceps brachii. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Biceps_brachii**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Humerus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Humerus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Humeroulnar joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Humeroulnar_joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brachialis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brachialis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor digitorum superficialis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_digitorum_superficialis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Supinator. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Supinator**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Triceps brachii muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Triceps_brachii_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anconeus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anconeus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brachioradialis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brachioradialis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor pollicis longus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_pollicis_longus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anconeus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anconeus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pronator teres. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pronator_teres**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brachioradialis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brachioradialis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Extensor carpi radialis longus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extensor_carpi_radialis_longus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor carpi radialis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_carpi_radialis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor digitorum profundus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_digitorum_profundus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Palmaris longus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Palmaris_longus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor carpi ulnaris. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_carpi_ulnaris**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- flexion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/flexion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extension. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/extension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray410.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray410.png**. **License**: ***Public Domain: No Known Copyright***
- Gray418. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray418.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Extensor carpi ulnaris muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extensor_carpi_ulnaris_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor carpi ulnaris muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_carpi_ulnaris_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor carpi radialis muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_carpi_radialis_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Flexor digitorum profundus muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Flexor_digitorum_profundus_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Extensor digiti minimi muscle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extensor_digiti_minimi_muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Extensor indicis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Extensor_indicis_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Flexor pollicis longus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Flexor_pollicis_longus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Extensor pollicis longus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Extensor_pollicis_longus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Extensor carpi radialis longus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Extensor_carpi_radialis_longus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Extensor digitorum muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Extensor_digitorum_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Flexor digitorum superficialis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Flexor_digitorum_superficialis_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Wrist. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wrist%23Movements_and_muscles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Abductor pollicis longus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Abductor_pollicis_longus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- flexion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/flexion. **License**: *CC BY-SA: Attribution-ShareAlike*
- extension. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/extension. **License**: *CC BY-SA: Attribution-ShareAlike*
- extrinsic muscles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/extrinsic%20muscles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray410.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Gray410.png. **License**: *Public Domain: No Known Copyright*
- Gray418. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray418.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- image427.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus427.html. **License**: *Public Domain: No Known Copyright*
- image415.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus415.html. **License**: *Public Domain: No Known Copyright*
- Rhomboid minor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rhomboid_minor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Deltoid muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Deltoid_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pectoralis major. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pectoralis_major. **License**: *CC BY-SA: Attribution-ShareAlike*
- Supraspinatus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Supraspinatus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Shoulder. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Shoulder%23Muscles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Infraspinatus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Infraspinatus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Subclavius. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Subclavius. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serratus anterior. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serratus_anterior. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sternocleidomastoid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sternocleidomastoid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pectoralis minor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pectoralis_minor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Levator scapulae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Levator_scapulae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Rotator cuff. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rotator...g_rotator_cuff. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trapezius. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trapezius. **License**: *CC BY-SA: Attribution-ShareAlike*
- Teres minor muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Teres_minor_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Subscapularis muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Subscapularis_muscle. **License**: *CC*

BY-SA: Attribution-ShareAlike
- Rhomboid major. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Rhomboid_major. **License**: *CC BY-SA: Attribution-ShareAlike*
- trapezius. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/trapezius. **License**: *CC BY-SA: Attribution-ShareAlike*
- deltoid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/deltoid. **License**: *CC BY-SA: Attribution-ShareAlike*
- rotator cuff. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/rotator_cuff. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray410.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Gray410.png. **License**: *Public Domain: No Known Copyright*
- Gray418. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray418.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- image427.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus427.html. **License**: *Public Domain: No Known Copyright*
- image415.gif. **Provided by**: Anatomy of the Human Body. **Located at**: http://www.bartleby.com/107/illus415.html. **License**: *Public Domain: No Known Copyright*
- Trapezius Gray409. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Tr...us_Gray409.PNG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray412. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray412.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Deltoideus posterior. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:De..._posterior.PNG. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 9.9C: Muscles of the Shoulder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

9.9D: Muscles that Cause Movement at the Ankle

Muscles of the leg insert into ankle and foot bones to facilitate ankle movement.

Learning Objectives

- Describe the muscles that cause the ankle to move

Key Points

- The ankle consists of two joints which permit dorsiflexion, plantarflexion, inversion, and eversion of the foot.
- Strong ligaments hold the ankle joint in place, although it is susceptible to damage.
- Muscles controlling movement at the ankle are found in the leg and can be split into anterior, posterior, and lateral compartments.

Key Terms


- **plantarflexion**: Movement of the foot downwards away from the lower leg.
- **eversion**: Tilting of the foot so the sole faces away from the midline.
- **inversion**: Tilting of the foot so the sole faces into the midline.
- **dorsiflexion**: Movement of the foot upwards towards the lower leg.

Movement at the ankle is controlled by two joints. The ankle or talocrural joint is formed from the tibia and fibula of the lower leg and talus of the foot. Functionally, it acts as a hinge, allowing dorsiflexion (pulling the foot upwards towards the lower leg) and plantarflexion (pulling the foot downwards away from the lower leg). Eversion (tilting of the sole of the foot away from the midline) and inversion (tilting of the sole of the foot inwards towards the midline) is controlled by the subtalar joint formed between the talus and calcaneus bones of the foot.

The ankle joint is held in place by numerous strong ligaments that can be easily damaged when excessive force is placed on the ankle, particularly during strenuous inversion and eversion. Movement at the ankle is key for maintenance of posture and balance, but is most important in locomotion. Variation in muscle activation can control the movement of the ankle joint, allowing the foot to generate graduated force.

Muscles that generate movement at the ankle are generally found in the lower leg and can be split into three categories.

Anterior Compartment



(a) Anterior Compartment of the Leg and (b) Posterior Compartment of the leg: Anterior view of leg showing the muscles and tendons involved in ankle movement. : Posterior view of leg showing muscles and tendons involved in ankle movement.

Three muscles in the anterior compartment of the leg act to dorsiflex and invert the foot at the ankle joint.

- *Tibialis Anterior*: The tibialis anterior muscle is located alongside the lateral surface of the tibia and is the strongest dorsiflexor of the foot.
  - Attachments: Originates from the lateral surface of the tibia and attaches to the base of the big toe.
  - Actions: Dorsiflexion and inversion of the foot.
- *Extensor Digitorum Longus*: The extensor digitorum longus is a deep-lying extrinsic muscle that runs the length of the tibia.
  - Attachments: Originates from the tibia and transitions into a tendon, passes into the foot, splits into four, and attaches to the toes.
  - Actions: Extension of the toes and dorsiflexion of the foot.
- *Extensor Hallucis Longus*: The extensor hallucis longus is a deep lying extrinsic muscle beneath the extensor digitorum longus.
  - Attachments: Originates from the fibula and attaches to the big toe.
  - Actions: Extension of the big toe, and dorsiflexion of the foot.

Posterior Compartment

Several muscles are located in the posterior compartment of the leg, typically grouped into superficial and basal groups. The majority of these muscles work to plantarflex the foot at the ankle.

Superficial Muscles

The superficial muscles give rise to the characteristic shape of the lower leg.

- *Gastrocnemius*: The gastrocnemius, a two-headed muscle, is the most superficial of the muscles in the posterior compartment.
  - Attachments: Both heads originate from the femur. The fibers converge to form the calcaneal tendon which attaches to the heel.
  - Actions: Plantarflexes the foot, can also flex the lower leg at the knee but is not key in this movement.
- *Plantaris*: The plantaris is a small muscle lying between the gastrocnemius and soleus. It is absent in 10% of people.
  - Attachments: Originates from the femur and attaches to the heel via the calcaneal tendon.
  - Actions: Plantarflexes the foot, can also flex the lower leg at the knee but is not key in this movement.

- *Soleus*: The soleus is a large flat muscle which is the deepest lying of the superficial muscles.
  - Attachments: Originates from the tibia and fibula and attaches to the heel via the calcaneal tendon.
  - Actions: Plantarflexes the foot.

Deep Muscles

- *Tibialis Posterior*: The tibialis posterior is the deepest lying of the muscles in the posterior compartment.
  - Attachments: Originates from the tibia and fibula and attaches to the plantar surfaces of the toes.
  - Actions: Inverts and plantarflexes the foot, maintains the arch of the foot.

Lateral Compartment

Two muscles found in the lateral compartment function to control eversion of the foot. Physiologically, there is a preference for the foot to invert, so these muscles also prevent excessive inversion.

- *Fibularis Longus*: The fibularis longus is the longer and more superficial of the two muscles.
  - Attachments: Originates from the fibula and tibia. The fibers converge into a tendon which passes under the foot and attaches to the medial side of the foot.
  - Actions: Eversion and plantarflexion of the foot.
- *Fibularis Brevis*: The fibularis brevis muscles is the deeper and shorter of the two muscles.
  - Attachments: Originates from the lateral surface of the fibula and attaches to the little toe.
  - Actions: Eversion of the foot.

KEY MOVEMENTS

- **Eversion of the Foot** (tilting of the sole of the foot away from the midline): Performed by the fibularis brevis and fibularis longus.
- **Inversion of the Foot** (tilting of the sole of the foot inwards towards the midline): Performed by the tibialis posterior and tibialis anterior.
- **Dorsiflexion of the Foot** (pulling the foot upwards towards the leg): Performed by the tibialis anterior, extensor hallucis longus and extensor digitorum longus.
- **Plantarflexion of the Foot** (pulling the foot downwards away from the lower leg): Performed by the gastrocnemius, plantaris, soleus and fibularis longus.

This page titled 9.9D: Muscles that Cause Movement at the Ankle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10: OVERVIEW OF THE NERVOUS SYSTEM



- Topic hierarchy

10.1: Introduction to the Nervous System

10.1A: Organization of the Nervous System

10.1B: Functions of the Nervous System

10.1C: Subdivisions of the Nervous System

10.2: Neuroglia

10.2A: Neuroglia of the Central Nervous System

10.2B: Neuroglia of the Peripheral Nervous System

10.3: Neurons

10.3A: Structural Diversity of Neurons

10.3B: Classification of Neurons

10.4: Collections of Nervous Tissue

10.4A: Clusters of Neuronal Cell Bodies

10.4B: Axon Bundles

10.4C: Gray and White Matter

10.5: Neurophysiology

10.5A: Principles of Electricity

10.5B: Ion Channels

10.5D: Membrane Potentials as Signals

10.5D: Resting Membrane Potentials

10.5E: The Action Potential and Propagation

10.5F: The Synapse

10.5G: Postsynaptic Potentials and Their Integration at the Synapse

10.5H: Synaptic Transmission

10.5I: Types of Neurotransmitters by Function

10.5J: Ionotropic and Metabotropic Receptors

10.5K: Serial and Parallel Processing

Index

This page titled 10: Overview of the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.1: INTRODUCTION TO THE NERVOUS SYSTEM



- Topic hierarchy

10.1A: Organization of the Nervous System

10.1B: Functions of the Nervous System

10.1C: Subdivisions of the Nervous System

This page titled 10.1: Introduction to the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.1A: Organization of the Nervous System

The nervous system is a network of cells called neurons that coordinate actions and transmit signals between different parts of the body.

Learning Objectives

- Describe the organization of the nervous system

Key Points

- Neurons (specialized cells of the nervous system) send signals along thin fibers called axons and communicate with other cells by releasing chemicals called neurotransmitters at cell-cell junctions called synapses.
- Glial cells are non-neuronal cells that provide support and nutrition in the nervous system.
- In humans, the nervous system consists of both the central and peripheral nervous systems.
- The human central nervous system contains the brain, spinal cord, and retina.
- The peripheral nervous system consists of sensory neurons, clusters of neurons called ganglia, and nerves connecting them to each other and to the central nervous system.

Key Terms

- **sensory receptor**: A nerve ending that recognizes stimulus in the internal or external environment of an organism.
- **peripheral nervous system**: This system consists of the nerves and ganglia outside of the brain and spinal cord.
- **glia**: Non-neuronal cells that maintain homeostasis, form myelin, and provide support and protection for neurons in the brain and other parts of the nervous system.


EXAMPLES

A nervous system allows us to react to the changing environment around us.

The nervous system is an organ system that coordinates voluntary and involuntary actions and responses by transmitting signals between different parts of our bodies.

Neurons


Central to the functioning of the nervous system is an extensive network of specialized cells called neurons. Neurons feature many thin projecting fibers called axons, which penetrate deep into tissues. They are able to communicate with other cells by chemical or electrical means at synapses. Neuronal function is supported by neuroglia, specialized cells which provide nutrition, mechanical support, and protection.



Major elements in neuron-to-neuron communication: Electrical impulses travel along the axon of a neuron. When this signal reaches a synapse, it provokes release of neurotransmitter molecules, which bind to receptor molecules located in the the target cell.

Divisions of the Nervous System

In most animals, including humans, the nervous system consists of two parts: central and peripheral. The central nervous system (CNS) is composed of the brain, spinal cord, and cerebellum. The peripheral nervous system (PNS) consists of sensory neurons, motor neurons, and neurons that communicate either between subdivisions of the PNS or connect the PNS to the CNS



The Human Nervous System: The major organs and nerves of the human nervous system. The CNS is comprised of the brain, cerebellum and spinal cord. Remaining neurons, and associated cells, distributed throughout the body form the PNS.

The nervous system has three broad functions: sensory input, information processing, and motor output. In the PNS, sensory receptor neurons respond to physical stimuli in our environment, like touch or temperature, and send signals that inform the CNS of the state of the body and the external environment. This sensory information is then processed by the CNS, predominantly by the brain.

After information is processed, motor neurons return signals to the muscles and glands of the PNS, which responds with motor output. Central neurons, which in humans greatly outnumber the sensory and motor neurons, make all of their input and output connections with other neurons. The connections of these neurons form neural circuits that are responsible for our perceptions of the world and determine our behavior. Along with neurons, the nervous system relies on the function of other specialized cells called glial cells, or glia, that provide structural and metabolic support to the nervous system.

This page titled 10.1A: Organization of the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.1B: Functions of the Nervous System

The primary function of the nervous system is to coordinate and control the various body functions.

Learning Objectives

- Describe the functions of the nervous system

Key Points

- The nervous system is a highly integrated system. The nervous system has three overlapping functions based on sensory input, integration, and motor output.
- At a more integrative level, the primary function of the nervous system is to control and communicate information throughout the body.

Key Terms

- **hormone**: A molecule released by a cell or a gland in one part of the body that sends out messages affecting cells in other parts of the organism.
- **nervous system**: The organ system that coordinates the activities of muscles, monitors organs, constructs and processes data received from the senses, and initiates actions.

The nervous system has three overlapping functions based on the sensory input, integration, and motor output. The nervous system is a highly integrated system.

Sensory Input

Sensory input comes from the many sensory receptors that monitor changes occurring both inside and outside the body. The total sum of the information gathered by these receptors is called sensory input. The nervous system processes and interprets sensory input and decides what actions should be taken. The nervous system activates effector organs such as muscles and glands to cause a response called motor output.


Integration

At a more integrative level, the primary function of the nervous system is to control and communicate information throughout the body. It does this by extracting information from the environment using sensory receptors. This sensory input is sent to the central nervous system, which determines an appropriate response.

Motor Response

Once the response is activated, the nervous system sends signals via motor output to muscles or glands to initiate the response.

In humans, the sophistication of the nervous system allows for language, abstract representation of concepts, transmission of culture, and many other features of society that would not otherwise exist.



Major elements in neuron-to-neuron communication: Electrical impulses travel along the axon of a neuron. When this signal reaches a synapse, it provokes release of neurotransmitter molecules, which bind to receptor molecules located in the the target cell.

This page titled 10.1B: Functions of the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.1C: Subdivisions of the Nervous System

The CNS includes the brain and spinal cord, while the PNS is a network of nerves linking the body to the brain and spinal cord.

Learning Objectives

- Describe the subdivisions of the nervous system

Key Points


- The nervous system is often divided into components called gray matter and white matter. Gray matter contains a relatively high proportion of neuron cell bodies and white matter is composed mainly of axons.
- The peripheral nervous system is subdivided into nerves, the autonomic system, and the somatic system. The autonomic nervous system is further subdivided into the parasympathetic and sympathetic nervous systems.
- The enteric nervous system is an independent subsystem of the peripheral nervous system.
- The central nervous system includes the brain and spinal cord and has various centers that integrate of all the information in the body. These can be subdivided into lower centers that carry out essential body functions and higher centers that control more sophisticated information processing.

Key Terms

- **gray matter**: A major component of the central nervous system, consisting of neuronal cell bodies, neuropil (dendrites and unmyelinated axons), glial cells (astroglia and oligodendrocytes), and capillaries.
- **central nervous system**: In vertebrates, the part of the nervous system comprising the brain, brainstem, and spinal cord.
- **white matter**: A region of the central nervous system containing myelinated nerve fibers and no dendrites.
- **peripheral nervous system**: This system consists of the nerves and ganglia outside of the brain and spinal cord.

The nervous system is comprised of two major subdivisions, the central nervous system (CNS) and the peripheral nervous system (PNS).

Central Nervous System



The Central Nervous System: The central nervous system (2) is a combination of the brain (1) and the spinal cord (3).

The CNS includes the brain and spinal cord along with various centers that integrate all the sensory and motor information in the body. These centers can be broadly subdivided into lower centers, including the spinal cord and brain stem, that carry out essential body and organ-control functions and higher centers within the brain that control more sophisticated information processing, including our thoughts and perceptions. Further subdivisions of the brain will be discussed in a later section.

Gray Matter and White Matter

The nervous system is often divided into components called gray matter and white matter. Gray matter, which is gray in preserved tissue but pink or light brown in living tissue, contains a relatively high proportion of neuron cell bodies. Conversely, white matter is composed mainly of axons and is named because of the color of the fatty insulation called myelin that coats many axons. White matter includes all of the nerves of the PNS and much of the interior of the brain and spinal cord. Gray matter is found in clusters of neurons in the brain and spinal cord and in cortical layers that line their surfaces.

By convention, a cluster of neuron cell bodies in the gray matter of the brain or spinal cord is called a nucleus, whereas a cluster of neuron cell bodies in the periphery is called a ganglion. However, there are a few notable exceptions to this rule, including a part of the brain called the basal ganglia, which will be discussed later.

Peripheral Nervous System

The PNS is a vast network of nerves consisting of bundles of axons that link the body to the brain and the spinal cord. Sensory nerves of the PNS contain sensory receptors that detect changes in the internal and external environment. This information is sent to the CNS via afferent sensory nerves. Following information processing in the CNS, signals are relayed back to the PNS by way of efferent peripheral nerves.

Autonomic and Somatic Nervous Systems


The PNS is further subdivided into the autonomic nervous system (ANS) and the somatic nervous system. The autonomic system has involuntary control of internal organs, blood vessels, and smooth and cardiac muscles. The somatic system has voluntary control of our movements via skeletal muscle.

As mentioned, the autonomic nervous system acts as a control system and most functions occur without conscious thought. The ANS affects heart rate, digestion, respiratory rate, salivation, perspiration, pupil diameter, urination, and sexual arousal. While most of its actions are involuntary, some, such as breathing, work in tandem with the conscious mind. The ANS is classically divided into two subsystems: the parasympathetic nervous system (PSNS) and sympathetic nervous system (SNS).

Parasympathetic and Sympathetic Nervous Systems

Broadly, the parasympathetic system is responsible for stimulation of “rest-and-digest” activities that occur when the body is at rest, including sexual arousal, salivation, lacrimation (tears), urination, digestion, and defecation. The sympathetic nervous syste is responsible for stimulating activities associated with the “fight-or-flight” response: mobilizing the systems of the body for escape or attacking sources of danger. In truth, the functions of both the parasympathetic and sympathetic nervous systems are not so straightforward, but this division is a useful rule of thumb.

The enteric nervous system (ENS) controls the gastrointestinal system and is sometimes considered part of the autonomic nervous system. However, it is sometimes considered an independent system because it can operate independently of the brain and the spinal cord.



The Nervous System of a Vertebrate: The brain and the spinal cord are the central nervous system (CNS) (shown in yellow). The left-right pair of cranial nerves, spinal nerves, and ganglia make up the peripheral nervous system (shown in dark gold).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- sensory receptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sensory_receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Nervous_System%23Overview_of_the_entire_nervous_systemhttp://en.Wikipedia.org/wiki/Nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/glia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peripheral nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/periphe...rvous%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous...tem%23Function**. **License**: ***Public Domain: No Known Copyright***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Neural_circuits_and_systems**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nervous system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Function**. **License**: ***Public Domain: No Known Copyright***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Nervous_System%23Overview_of_the_entire_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Function**. **License**:

Public Domain: No Known Copyright
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Enteric nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Enteric_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parasympathetic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Anatomy_in_vertebrates**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Nervous_System%23Overview_of_the_entire_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gray matter. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/gray%20matter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peripheral nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/peripheral%20nervous%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- white matter. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/white_matter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- central nervous system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/central_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Function**. **License**: ***Public Domain: No Known Copyright***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Nervous_System%23Overview_of_the_entire_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Function**. **License**: ***Public Domain: No Known Copyright***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Nervous_System%23Overview_of_the_entire_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- TE-Nervous system diagram. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...em_diagram.svg**. **License**: ***CC BY: Attribution***

This page titled 10.1C: Subdivisions of the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.2: NEUROGLIA



- Topic hierarchy

10.2A: Neuroglia of the Central Nervous System

10.2B: Neuroglia of the Peripheral Nervous System

This page titled 10.2: Neuroglia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.2A: Neuroglia of the Central Nervous System

Glia, named from the Greek word for “glue,” support and scaffold neurons while performing other unique functions.

Learning Objectives

- Identify the types of neuroglia in central nervous system

Key Points

- Neuroglia in the CNS include astrocytes, microglial cells, ependymal cells and oligodendrocytes.
- Neuroglia in the PNS include Schwann cells and satellite cells.Astrocytes support and brace the neurons and anchor them to their nutrient supply lines. They also play an important role in making exchanges between capillaries and neurons.
- Microglial cells can transform into a special type of macrophage that can clear up the neuronal debris, while monitoring the health of the neuron.
- Ependymal cells are another glial subtype that line the ventricles of the CNS to help circulate the CSF.
- Oligodendrocytes are cells that wrap their process tightly around the fibers producing an insulating covering called myelin sheath.
- Schwann cells are similar in function to oligodendrocytes and microglial cells.
- Satellite cells perform a similar function to astrocytes,

Key Terms


- **myelin**: A white, fatty material, composed of lipids and lipoproteins, that surrounds the axons of nerves.
- **glia**: Non-neuronal cells that maintain homeostasis, form myelin, and provide support and protection for neurons in the brain, and for neurons in other parts of the nervous system such as in the autonomic nervous system.
- **astrocyte**: a neuroglial cell, in the shape of a star, in the brain

Neurogila or glial cells, are non-neuronal cells that maintain homeostasis, form myelin, and provide support and protection for neurons in the central (CNS) and peripheral nervous systems (PNS). It was long believed that neuroglia did not play any role in neuro-transmission, however recent advances have demonstrated that neuroglia play a key role in synapse formation and maintenance.

Neuroglia of the CNS

Neuroglia in the CNS include astrocytes, microglial cells, ependymal cells, and oligodendrocytes. In the human brain, it is estimated that the total number of glia roughly equals the number of neurons, although the proportions vary in different brain areas.

- Astrocytes are delicate, star-shaped branching glial cells. Their numerous radiating processes cling to neurons and their synaptic endings. These astrocytes cover nearly all the capillaries in the CNS. They support and brace the neurons and anchor them to their nutrient supply lines. They also play an important role in making exchanges between capillaries and neurons. Astrocytes also regulate the external chemical environment of neurons by removing excess ions and recycling neurotransmitters released during synaptic transmission.
- Microglial cells are small and have thorny processes that can touch the neighboring neurons. Microglial cells can transform into a special type of macrophage that can clear up the neuronal debris. They are also able to monitor the health of neurons by detecting injuries to the neuron.
- Ependymal cells are another glial subtype that line the ventricles of the CNS, forming a permeable barrier between the cerebrospinal fluid (CSF) and underlying cells, and also aid in the circulation of CSF through cilial beat.
- Oligodendrocytes are cells that have fewer processes compared to astrocytes. They line up along the nerve fibers in the CNS and wrap their process tightly around the fibers producing the insulating myelin sheath.




Oligodendrocyte: Oligodendrocytes form the electrical insulation around the axons of CNS nerve cells.

Neuroglia of the PNS

Neuroglia in the PNS include Schwann cells and satellite cells.

- Schwann cells are similar in function to oligodendrocytes and microglial cells, providing myelination to axons in the PNS. They also have phagocytotic activity and clear cellular debris that allows for regrowth of PNS neurons.
- Satellite cells are similar in function to astrocytes small cells that surround neurons in sensory, sympathetic, and parasympathetic ganglia, helping to regulate the external chemical environment. They are highly sensitive to injury and inflammation, and appear to contribute to pathological states, such as chronic pain.



Neuroglia: Types of neuroglia found in the CNS and PNS.

This page titled 10.2A: Neuroglia of the Central Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.2B: Neuroglia of the Peripheral Nervous System

The two kinds of glia cells in the PNS, schwann cells and satellite cells; each have unique functions.

Learning Objectives

- Identify the neuroglia of the peripheral nervous system

Key Points


- There are two kinds of neuroglia in the peripheral nervous system (PNS): Schwann cells and satellite cells.
- Schwann cells provide myelination to peripheral neurons. Functionally, the schwann cells are similar to oligodendrocytes of the central nervous system (CNS).
- Satellite cells play an important role in modulating the PNS following injury and inflammation. These resemble the astrocytes of the CNS and assist in regulating the external chemical environment.

Key Terms

- **glia**: Non-neuronal cells that maintain homeostasis, form myelin, and provide support and protection for neurons in the brain and in other parts of the nervous system.
- **Schwann cells**: The principal glia of the PNS, these cells are involved in the conduction of nervous impulses along axons, nerve development and regeneration, trophic support for neurons, production of the nerve extracellular matrix, modulation of neuromuscular synaptic activity, and presentation of antigens to T-lymphocytes.
- **Satellite glial cells**: These cells line the exterior surface of neurons in the PNS and neuron cell bodies within ganglia.

The PNS has two kinds of neuroglia: schwann cells and satellite cells. Schwann cells provide myelination to peripheral neurons. They also perform phagocytic functions and clear cellular debris, allowing for the regrowth of PNS neurons. Functionally, the schwann cells are similar to oligodendrocytes of the CNS.

Satellite cells are small glia that surround neurons' sensory ganglia in the ANS. These resemble the astrocytes of the CNS and assist in regulating the external chemical environment. PNS satellite glia are very sensitive to injury and may exacerbate pathological pain.



Schwann Cell: A Schwann cell in culture

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 10.2B: Neuroglia of the Peripheral Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.3: NEURONS



- Topic hierarchy

10.3A: Structural Diversity of Neurons

10.3B: Classification of Neurons

This page titled 10.3: Neurons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.3A: Structural Diversity of Neurons

A number of anatomically neuron types have evolved to participate in different organismal functions.

Learning Objectives

- Describe the structural diversity of neurons

Key Points


- Neurons are electrically excitable cells that are the structural unit of the nervous system.
- A typical neuron consists of a cell body and neuronal processes such as dendrites and axon.
- Neurons can generally be anatomically characterized as unipolar, bipolar, or multipolar.
- A number of anatomically distinct neuron types, such as sensory, motor, and interneurons, have evolved to participate in different organismal functions.

Key Terms

- **Neurons**: Electrically excitable cells that are the structural unit of the nervous system.
- **dendrites**: Short, tapering extensions that convey incoming messages toward the body of the neuron.
- **Axons**: The conducting region of the neuron.


Neurons are electrically excitable cells that are the structural unit of the nervous system. A typical neuron consists of a cell body and neuronal processes such as dendrites and axon. The dendrites are short, tapering extensions that are the receptive regions and help in conveying incoming messages towards the cell body. Axons arise from a cone-shaped area of the cell body called axon hillock. These extensions are the conducting region of the neuron. Nerve impulses are generated in the axon and transmitted away from the cell body towards the synapse. The cell body is the major biosynthetic center of the neuron. It contains neurotransmitters and other organelles needed to synthesize proteins and chemicals. The cell body is the focal point for the outgrowth of neuronal process during embryonic development.

A number of anatomically distinct types of neurons have evolved to participate in different organismal functions. For example, sensory neurons respond to touch, sound, light, and other sensory inputs. Motor neurons receive signals from the brain and spinal cord to initiate muscle contractions and affect glands. Interneurons act as relays between neurons in close proximity to one another.



Neurotransmission at a Chemical Synapse: A signal propagating down an axon to the cell body and dendrites of the next cell

Neurons can generally be grouped according to the number of processes extending from their cell bodies. Three major neuron groups make up this classification: multipolar, bipolar, and unipolar. Unipolar neurons have a single short process that emerges from the cell body and divides T-like into proximal and distal branches. Bipolar neurons have two processes, an axon and a dendrite, that extend from opposite ends of the soma. Multipolar neurons, the most common type, have one axon and two or more dendrites.



types of neurons: 1: Unipolar neuron, 2: Bipolar neuron, 3: Multipolar neuron, 4: Pseudounipolar neuron

This page titled 10.3A: Structural Diversity of Neurons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.3B: Classification of Neurons

Neurons can be classified by direction of travel, neurotransmitter utilized, or electrophysiological properties.

Learning Objectives

- Describe the various methods used to classify neurons

Key Points


- Neurons can be classified by the direction of the action potential or route by which information travels. Afferent neurons convey information from tissues and organs to the brain and efferent signals transmit information from the brain to effector cells in the body.
- Neurons can have excitatory, inhibitory, or modulatory effects on target neurons depending on the neurotransmitter they release.
- Afferent neurons convey information from tissues and organs into the central nervous system.
- Interneurons connect neurons within specific regions of the central nervous system.
- Efferent neurons carry information away from a brain region.

Key Terms

- **efferent**: Efferent neurons transmit signals from the central nervous system to the effector cells (e.g. motor neurons).
- **afferent**: Afferent neurons convey information from tissues and organs into the central nervous system (e.g. sensory neurons).
- **tonic or regular spiking**: Neurons that are typically constantly (or tonically) active are called tonic or regular spiking.

Direction of Nerve Impulse


The functional classification of neurons is based on the direction the action potential (i.e. information) travel relative to the central nervous system. Afferent neurons convey information from tissues and organs into the central nervous system (e.g. sensory neurons). Efferent neurons transmit signals from the central nervous system (CNS) to the effector cells (e.g. motor neurons). Afferent and efferent also refer generally to neurons that bring information to or send information from a brain region. Interneurons connect neurons within specific regions of the central nervous system.



The organization of the nervous system: Gross organization of the nervous system, with the peripheral nervous system, the spinal, and the cortical levels.

Neurotransmitter Type

Neurons are also classified by their effect on target neurons. A neuron releases a neurotransmitter that binds to chemical receptors on the target neuron. The combination of neurotransmitter and receptor properties results in an excitatory, inhibitory, or modulatory change to the target neuron. For example, the two most common neurotransmitters in the brain (released by 90% of neurons), glutamate and GABA, have opposing actions. Glutamate acts on several different types of receptors with largely excitatory effects. GABA acts on several different classes of receptors to exert inhibitory effects. Other types of neurons include excitatory motor neurons in the spinal cord that release acetylcholine and inhibitory spinal neurons that release glycine.



Major elements in neuron-to-neuron communication: Electrical impulses travel along the axon of a neuron. When this signal reaches a synapse, it provokes release of neurotransmitter molecules, which bind to receptor molecules located in the the target cell.

Firing Properties

A third, less common method of neuron classification is according to their electrophysiological characteristics. Neurons that are typically constantly (or tonically) active are called tonic or regular spiking. Neurons that are intermittently active are called phasic or bursting. Neurons with high activity rates are classified as fast spiking.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neuron%23Classes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neuron**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neuron. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neuron**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neurons uni bi multi pseudouni. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:N..._pseudouni.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neuron%23Classes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neuron%23Classes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...egular-spiking**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ition/efferent**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- afferent. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/afferent**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neurons uni bi multi pseudouni. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:N..._pseudouni.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neuron%23Classes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous...tem%23Function**. **License**:

This page titled 10.3B: Classification of Neurons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.4: COLLECTIONS OF NERVOUS TISSUE



- Topic hierarchy

10.4A: Clusters of Neuronal Cell Bodies

10.4B: Axon Bundles

10.4C: Gray and White Matter

This page titled 10.4: Collections of Nervous Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.4A: Clusters of Neuronal Cell Bodies

Clusters of cell bodies in the central nervous system are called nuclei, while the cell bodies lining the nerves in the peripheral nervous system are called ganglia.

Learning Objectives

- Identify the types of clusters of neuronal cell bodies

Key Points

- A cluster of neurons is called a nucleus if found in the central nervous system; it is called a ganglion if found in the peripheral nervous system (PNS).
- Ganglia are the intermediate structures between the central and peripheral nervous systems.
- Satellite glial cells (SGC) line the exterior surface of neurons in the PNS and surround neuron cell bodies within ganglia.


Key Terms

- **ganglion**: A cluster of interconnecting nerve cells outside the brain.
- **nucleus**: A cluster of neuronal bodies where synapsing occurs.

In neuroanatomy, a nucleus is a brain structure consisting of a relatively compact cluster of neurons. It is one of the two most common forms of nerve cell organization along with layered structures such as the cerebral cortex or cerebellar cortex. In anatomical sections, a nucleus shows up as a region of gray matter often bordered by white matter. The vertebrate brain contains hundreds of distinguishable nuclei varying widely in shape and size. A nucleus may itself have a complex internal structure, with multiple types of neurons arranged in clumps (subnuclei) or layers.

In addition, the term nucleus can refer to a distinct group of neurons that spread over an extended area. For example, the reticular nucleus of the thalamus is a thin layer of inhibitory neurons that surround the thalamus.

In the peripheral nervous system, a cluster of neurons is called a ganglion. One exception is the basal ganglia, located not in the periphery but rather in the forebrain. Ganglia are composed mainly of neuron cell bodies (somata) and dendritic structures. They are the intermediary connections between the peripheral and central nervous systems.



Innervation of the Autonomic Nervous System: Satellite glial cells are expressed throughout the sympathetic and parasympathetic ganglia in their respective nervous system divisions.

Satellite glial cells line the exterior surface of neurons in the PNS. Satellite glial cells (SGCs) also surround neuron cell bodies within ganglia. They are of a similar embryological origin to Schwann cells of the PNS, as both are derived from the neural crest of the embryo during development. SGCs have a variety of roles, including control over the microenvironment of sympathetic ganglia. They are thought to have a similar role to astrocytes in the central nervous system (CNS). They supply nutrients to the surrounding neurons and also have some structural function. Satellite cells also act as protective, cushioning cells. Additionally, they express a variety of receptors that allow for a range of interactions with neuroactive chemicals.

This page titled 10.4A: Clusters of Neuronal Cell Bodies is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.4B: Axon Bundles

A bundle of axons is called a nerve in the peripheral nervous system and a tract in the central nervous system.

Learning Objectives

- Describe bundles of axons in the central and peripheral nervous systems

Key Points

- In the peripheral nervous system a bundle of axons is called a nerve.
- In the central nervous system a bundle of axons is called a tract.
- Each axon is surrounded by a delicate endoneurium layer.
- The course connective tissue layer called perineurium, binds the fibers into bundles called fascicles.
- A tough fibrous sheath called epineurium encloses all the fascicles to form the nerve.

Key Terms

- **endoneurial fluid**: A protein liquid surrounding individual nerve axons.
- **fascicle**: A bundle of axons.

Axons

Neurons feature many long, slender projections termed axons, along which electrochemical nerve impulses are transmitted. In the central nervous system (CNS) bundles of these axons are called tracts, whereas in the peripheral nervous system (PNS) they are called nerves.

Each nerve is covered externally by a dense sheath of connective tissue, the epineurium. Underlying this layer of flat cells, the perineurium, forms a complete sleeve around a bundle of axons called fascicles. Surrounding each axon is the endoneurium. The endoneurium consists of an inner sleeve of material called the glycocalyx and an outer delicate meshwork of collagen fibers. Within the endoneurium, the individual nerve axons are surrounded by a protein liquid called endoneurial fluid. The endoneurium has properties analogous to the blood-brain barrier, in that it prevents certain molecules from crossing from the blood into the endoneurial fluid.

Axon length and diameter can vary greatly from between 1 m to 1 mm in length and 1 µm to 20 µm in diameter. The longest axons in the human body are those of the sciatic nerve, which run from the base of the spinal cord to the big toe of each foot. Axons in the central nervous system typically show complex trees with many branch points allowing for the simultaneous transmission of messages to a large number of target neurons.

Myelin

Axons are described as either un-myelinated or myelinated. Myelin is a layer of a fatty insulating substance, which is formed by two types of glial cells: Schwann cells en-sheathing peripheral neurons and oligodendrocytes insulating those of the central nervous system. Myelination enables an especially rapid mode of electrical impulse propagation called saltatory conduction. De-myelination of axons causes the multitude of neurological symptoms found in the disease multiple sclerosis.

Localization

Nerves in the PNS are typically divided into cranial and spinal nerves. There are twelve pairs of cranial nerves and thirty one pair of spinal nerves. Cranial nerves innervate parts of the head and connect directly to the brain (especially to the brainstem). They

are typically assigned Roman numerals from 1 to 12, although cranial nerve zero is sometimes included. In addition, cranial nerves have descriptive names. Spinal nerves innervate much of the body, and connect through the spinal column to the spinal cord. They are given letter-number designations according to the vertebra through which they connect to the spinal column.

This page titled 10.4B: Axon Bundles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.4C: Gray and White Matter

The central nervous system consists of a central cavity surrounded by gray matter made of neuronal cell bodies and white matter made of myelinated axons.

Learning Objectives

- Categorize gray versus white matter

Key Points

- Gray matter is a major component of the central nervous system, consisting of neuronal cell bodies and other cells such as glia and dendrites.
- The white matter is composed of bundles of myelinated axons (few cell bodies) that connect various grey matter regions.
- Myelin is a thin layer, around the axons of white matter neurons and provides the white coloration.

Key Terms


- **gray matter**: A major component of the CNS consisting of neuronal cell bodies.

The central nervous system (CNS) is comprised of white and gray matter. In the spinal cord and cerebrum gray matter is surrounded by white matter. However, in the cerebellum and cerebral hemispheres this is reversed with the grey matter surrounding underlying white matter.

Gray Matter

Gray matter is a major component of the CNS, it refers to un-myelinated neurons and other cells of the central nervous system such as glial cells and dendrites. It is present in the brain, brainstem and cerebellum, and present throughout the spinal cord.

Gray matter contains most of the brain's neuronal cell bodies. The grey matter includes regions of the brain involved in muscle control, and sensory perception such as seeing and hearing, memory, emotions, speech, decision making, and self-control.




Grey and White Matter: Micrograph showing grey matter, with the characteristic neuronal cell bodies (right of image – darker pink), and white matter with its characteristic fine mesh work-like appearance (left of image – lighter pink).

White Matter

A second major component of the CNS is white matter. It is composed mainly of bundles of myelinated axons, with very few neuronal bodies. White matter connects the various grey matter regions of the nervous system to each other and carries nerve impulses between neurons.

Myelin, the lipid that forms a thin layer, known as the myelin sheath, around the axons providing electrical insulation is white in color, giving rise to the name white matter.



White Matter on an MRI: This MRI highlights the location of white matter in the brain.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Satellite glial cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Satellite_glial_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ganglion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ganglion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nucleus (neuroanatomy). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nucleus_(neuroanatomy)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ganglion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ganglion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nucleus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***
- Nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Neural tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neural_tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray839. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray839.png. **License**: *Public Domain: No Known Copyright*
- White matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/White_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- Grey matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Grey_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- White matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/White_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- grey matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/grey%20matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- white matter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/white_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray839. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray839.png. **License**: *Public Domain: No Known Copyright*
- Grey matter and white matter - very high mag. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Grey_matter_and_white_matter_-_very_high_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- 3DSlicer-KubickiJPR2007-fig6. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:3DSlicer-KubickiJPR2007-fig6.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.4C: Gray and White Matter is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5: NEUROPHYSIOLOGY

- Topic hierarchy

10.5A: Principles of Electricity

10.5B: Ion Channels

10.5D: Membrane Potentials as Signals

10.5D: Resting Membrane Potentials

10.5E: The Action Potential and Propagation

10.5F: The Synapse

10.5G: Postsynaptic Potentials and Their Integration at the Synapse

10.5H: Synaptic Transmission

10.5I: Types of Neurotransmitters by Function

10.5J: Ionotropic and Metabotropic Receptors

10.5K: Serial and Parallel Processing

This page titled 10.5: Neurophysiology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5A: Principles of Electricity

When a neuron is stimulated, an electrical impulse is generated and conducted along the length of its axon. This process, called action potential, underlies many nervous system functions.

Learning Objectives

- Identify principles of electricity that are important in neurophysiology

Key Points

- Voltage and current are two important factors to consider in the study of neurons.
- The flow of electrical charge from one point to another is called current.
- Across the cellular membranes, potential difference is established between the outside of a cell and the inside of the cell which can affect the flow of current across the cell membrane.


Key Terms

- **Voltage**: The measure of potential energy generated by separated charge.
- **current**: The flow of electrical charge from one point to another.
- **resistance**: Hindrance to the flow of charge.

Voltage and current are two important factors to consider in the study of neurons. Voltage is the measure of potential energy generated by separated charge. It is measured in volts or millivolts. The greater the difference in charge between two points, the higher the voltage. Voltage is equal to the work that would have to be done per unit charge to move the charge between two points against a static electric field. Voltage may represent either a source of energy (electromotive force) or lost or stored energy (potential drop).

The flow of electrical charge from one point to another is called current. The amount of charge that moves between two points depends on two factors: voltage and resistance. Resistance is the hindrance to the flow of charge. Some substances with high resistance are insulators, like the myelin sheath.

In the body, electrical currents reflect the flow of ions across cell membranes. Since there is a slight difference in the number of positive and negative ions on the two sides of the cellular plasma membrane, there is a potential difference across the membranes. Diffusion arises from the statistical tendency of particles to redistribute from regions where they are highly concentrated to regions where the concentration is low. The difference in potential established between the outside of a cell and the inside of the cell can affect the flow of current across the cell membrane.



Cellular Membrane: Differences in concentration of ions on opposite sides of a cellular membrane lead to a voltage called the membrane potential. Many ions have a concentration gradient across the membrane, including potassium (K+), which is at a high inside and a low concentration outside the membrane. Sodium (Na+) and chloride (Cl–) ions are at high concentrations in the extracellular region and low concentrations in the intracellular regions. These concentration gradients provide the potential energy to drive the formation of the membrane potential.

In electrically active tissue, the potential difference between any two points can be measured by inserting an electrode at each point and connecting both electrodes to to a specialized voltmeter.

This page titled 10.5A: Principles of Electricity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5B: Ion Channels

Ion channels are membrane proteins that allow ions to travel into or out of a cell.

Learning Objectives

- Distinguish between types of ion channels

Key Points

- Ion channels can be voltage-sensitive, ligand-gated, or mechanically-gated in nature.
- Ligand-gated ion channels open when a chemical ligand such as a neurotransmitter binds to the protein.
- Voltage channels open and close in response to changes in membrane potential.
- Mechanically-gated channels open in response to physical deformation of the receptor, as in sensory receptors of touch and pressure.


Key Terms

- **leakage channel**: The simplest type of ion channel, with more or less constant permeability.
- **ligand-gated channel**: A group of transmembrane ion channels that open or close in response to the binding of a chemical messenger (ligand) such as a neurotransmitter.
- **ion pump**: Critical membrane proteins that carry out active transport by using cellular energy (ATP) to "pump" the ions against their concentration gradient.

EXAMPLES

Voltage-gated sodium ion channels contribute to the "spike" of a neuron's action potential.

Properties of Ion Channels



Ion channel: A schematic representation of an ion channel. The cell membrane is depicted by #6 and the ion channel pore by

Plasma membrane is studded with a variety of membrane proteins that act as ion channels. Each channel only allows certain types of ions to pass across the membrane. Most channels are specific (selective) for one ion. The channel pore is typically so small that ions must pass through it in single file.

A channel may have several different states (corresponding to different conformations of the protein), but each such state is either open or closed. In general, closed states correspond either to a contraction of the pore—making it impassable to the ion—or to a separate part of the protein, stoppering the pore. When a channel is open, ions permeate through the channel pore down the transmembrane concentration gradient for that particular ion. Rate of ionic flow through the channel is determined by the maximum channel conductance and electrochemical driving force for that ion.

Types of Ion Channels

Ion channels can be classified by how they respond to their environment. For example, ion channels can be voltage-sensitive in that they open and close in response to the voltage across the membrane. Ligand-gated channels form another important class; these ion channels open and close in response to the binding of a ligand molecule such as a neurotransmitter. Other ion channels open and close with mechanical forces. Still others, such as those of sensory neurons, open and close in response to other stimuli, such as light, temperature, or pressure. The most common types of ion channels are described below.


Leakage Channels

Leakage channels are the simplest type of ion channel, in that their permeability is more or less constant. The types of leakage channels with the greatest significance in neurons are potassium and chloride channels. Although they are the simplest in theory, most conduct better in one direction than the other (they are rectifiers) and some are capable of being shut off by ligands even though they do not require ligands in order to operate.

Gated Channels

There are three main types of gated channels: chemically-gated or ligand-gated channels, voltage-gated channels, and mechanically-gated channels.


Ligand-gated ion channels are channels whose permeability is greatly increased when some type of chemical ligand binds to the protein structure. A large subset function as neurotransmitter receptors—they occur at postsynaptic sites, and the chemical ligand that gates them is released by the presynaptic axon terminal. Ligand-gated channels can be activated by ligands that appear in the extracellular area or by interactions on the intracellular side.



A Ligand-gated Ion Channel: Ligand-gated ion channels (LGICs) are one type of ionotropic receptor or channel-linked receptor. They are a group of transmembrane ion channels that open or close in response to the binding of a chemical messenger (ligand), such as a neurotransmitter.

Voltage-gated ion channels, also known as voltage-dependent ion channels, are channels whose permeability is influenced by the membrane potential. They form another very large group, with each member having a particular ion selectivity and a particular voltage dependence. Many are also time-dependent—in other words, they do not respond immediately to a voltage change, but only after a delay. Voltage-gated channels are essential for the generation and propagation of action potentials.

Ion pumps are not ion channels, but are critical membrane proteins that carry out active transport by using cellular energy (ATP) to "pump" the ions against their concentration gradient. Such ion pumps take in ions from one side of the membrane (decreasing its concentration there) and release them on the other side (increasing its concentration there).



Ion pump example: Example of primary active transport, where energy from hydrolysis of ATP is directly coupled to the movement of a specific substance across a membrane independent of any other species.

This page titled 10.5B: Ion Channels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5D: Membrane Potentials as Signals

The membrane potential allows a cell to function as a battery, providing electrical power to activities within the cell and between cells.

Learning Objectives

- Describe how membrane potentials are used as signals

Key Points

- A membrane potential is the difference in electrical potential between the interior and the exterior of a biological cell.
- In electrically excitable cells, changes in membrane potential are used for transmitting signals within the cell.
- The opening and closing of ion channels can induce changes from the resting potential.
- Depolarization is when the interior voltage becomes more positive and hyperpolarization is when it becomes more negative.

Key Terms

- **action potential**: A short-term change in the electrical potential that travels along a cell such as a nerve or muscle fiber.
- **depolarization**: A change in a cell's membrane potential, making it more positive or less negative, that may result in generation of an action potential.
- **graded potentials**: These vary in size and arise from the summation of the individual actions of ligand-gated ion channel proteins, and decrease over time and space.

EXAMPLES

In neurons, a sufficiently large depolarization can evoke an action potential in which the membrane potential changes rapidly.

Membrane Potentials

Membrane potential (also transmembrane potential or membrane voltage) is the difference in electrical potential between the interior and the exterior of a biological cell. All animal cells are surrounded by a plasma membrane composed of a lipid bilayer embedded with various protein types. The membrane serves as both an insulator and a semi-permeable diffusion barrier to the movement of ions. Ions are moved across the cell membrane either through active (using energy) or passive (not using energy) transport.


For example, ion transporter/pump proteins actively push ions across the membrane to establish concentration gradients across the membrane and ion channels allow ions to move across the membrane down those concentration gradients, a process known as facilitated diffusion.

Virtually all eukaryotic cells (including cells from animals, plants, and fungi) maintain a nonzero transmembrane potential, usually with a negative voltage in the cell interior compared to the cell exterior.

The Membrane Potential as a Signal

The membrane potential has two basic functions. First, it allows a cell to function as a battery, providing power to operate a variety of "molecular devices" embedded in the membrane. Second, in electrically excitable cells such as neurons and muscle cells, it is used for transmitting signals between different parts of a cell. Signals are generated by opening or closing of ion channels at one point in the membrane, producing a local change in the membrane potential that causes electric current to flow rapidly to other points in the membrane.

In non-excitable cells, and in excitable cells in their baseline states, the membrane potential is held at a relatively stable value called the resting potential. For neurons, typical values of the resting potential range from –70 to –80 millivolts; that is, the interior of a cell has a negative baseline voltage of a bit less than one tenth of a volt. The opening and closing of ion channels can induce a departure from the resting potential. This is called a depolarization if the interior voltage becomes more positive (say from –70 mV to –60 mV), or a hyperpolarization if the interior voltage becomes more negative (say from –70 mV to –80 mV). The changes in membrane potential can be small or larger (graded potentials) depending on how many ion channels are activated and what type they are. In excitable cells, a sufficiently large depolarization can evoke an action potential in which the membrane potential changes rapidly and significantly for a short time (on the order of 1 to 100 milliseconds), often reversing its polarity. Action potentials are generated by the activation of certain voltage-gated ion channels.



Action potential: A. Schematic and B. actual action potential recordings. The action potential is a clear example of how changes in membrane potential can act as a signal.

This page titled 10.5D: Membrane Potentials as Signals is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5D: Resting Membrane Potentials

Learning Objectives

- Describe the resting membrane potential of cells

The potential difference in a resting neuron is called the resting membrane potential. This causes the membrane to be polarized. The value of the resting membrane potential varies from –40mV to –90mV in a different types of neurons.

The resting membrane potential exists only across the membrane. Most of the time, the difference in ionic composition of the intracellular and extracellular fluids and difference in ion permeability generates the resting membrane potential difference.

The interactions that generate the resting potential are modeled by the Goldman equation. It is based on the charges of the ions in question, as well as the difference between their inside and outside concentrations and the relative permeability of the plasma membrane to each ion where:

$$Em = RTFln(PK[K+]out + PNa[NA+]out + PCl[Cl-]inPK[K+]in + PNa[Na+]in + PCl[Cl-]out) \quad (10.5\text{D}.1)$$

Goldman equation: R is the universal gas constant, equal to 8.314 joules·K–1·mol^{-1} T is the absolute temperature, measured in kelvins (= K = degrees Celsius + 273.15) F is the Faraday constant, equal to 96,485 coulombs·mol–1 or J·V–1·mol^{-1}

The three ions that appear in this equation are potassium (K^+), sodium (Na^+), and chloride (Cl^-). The Goldman formula essentially expresses the membrane potential as an average of the reversal potentials for the individual ion types, weighted by permeability. In most animal cells, the permeability to potassium is much higher in the resting state than the permeability to sodium. Consequently, the resting potential is usually close to the potassium reversal potential.

Key Points

- The potential difference in a resting neuron is called the resting membrane potential.
- The value of the resting membrane potential varies from -40mV to -90mV in a different types of neurons.
- Most of the time, the difference in ionic composition of the intracellular and extracellular fluids and difference in ion permeability generates the resting membrane potential difference.
- The Goldman formula essentially expresses the membrane potential as an average of the reversal potentials for the individual ion types weighted by permeability

Key Terms

- **resting membrane potential**: The potential difference in a resting neuron that causes its membrane to be polarized.
- **Goldman equation**: Models the interactions that generate resting membrane potential.

This page titled 10.5D: Resting Membrane Potentials is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5E: The Action Potential and Propagation

Neurons typically send signals over long distances by generating and propagating action potentials over excitable axonal membrane.

Learning Objectives

- Describe the stages of an action potential and its propogation

Key Points

- Action potential is a brief reversal of membrane potential in which the membrane potential changes from -70mV to +30mV
- The action potential has three main stages: depolarization, repolarization, and hyperpolarization.
- Depolarization is caused when positively charged sodium ions rush into a neuron with the opening of voltage-gated sodium channels.
- Repolarization is caused by the closing of sodium ion channels and the opening of potassium ion channels.
- Hyperpolarization occurs due to an excess of open potassium channels and potassium efflux from the cell.


Key Terms

- **action potential.**: A brief reversal of membrane potential.
- **repolarization**: Also called the falling phase,
  caused by the slow closing of sodium channels and the opening of voltage-gated potassium channels.
- **absolute refractory period**: The period from the opening of the sodium channels until the sodium channels begin to reset.
- **hyperpolarization**: A phase where some potassium channels remain open and sodium channels reset.
- **depolarization**: Also called the rising phase, when positively charged sodium ions (Na+) suddenly rush through open voltage-gated sodium channels into a neuron.

Action Potential

Action potential is a brief reversal of membrane potential where the membrane potential changes from -70mV to +30mV. When the membrane potential of the axon hillock of a neuron reaches threshold, a rapid change in membrane potential occurs in the form of an action potential.

This moving change in membrane potential has three phases. First is depolarization, followed by repolarization and a short period of hyperpolarization. These three events happen over just a few milliseconds.



Action potential: A. Schematic and B. actual action potential recordings. The action potential is a clear example of how changes in membrane potential can act as a signal.

- The depolarization, also called the rising phase, is caused when positively charged sodium ions (Na+) suddenly rush through open voltage-gated sodium channels into a neuron. As additional sodium rushes in, the membrane potential actually reverses its polarity. During this change of polarity the membrane actually develops a positive value for a moment (+40 millivolts).
- The repolarization or falling phase is caused by the slow closing of sodium channels and the opening of voltage-gated potassium channels. As a result, the membrane permeability to sodium declines to resting levels. As the sodium ion entry declines, the slow voltage-gated potassium channels open and potassium ions rush out of the cell. This expulsion acts to restore the localized negative membrane potential of the cell.
- Hyperpolarization is a phase where some potassium channels remain open and sodium channels reset. A period of increased potassium permeability results in excessive potassium efflux before the potassium channels close. This results in hyperpolarization as seen in a slight dip following the spike.

The propagation of action potential is independent of stimulus strength but dependent on refractory periods. The period from the opening of the sodium channels until the sodium channels begin to reset is called the absolute refractory period. During this period, the neuron cannot respond to another stimulus, no matter how strong.

This page titled 10.5E: The Action Potential and Propagation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5F: The Synapse

A synapse is a structural junction that mediates information transfer from one neuron to the next or from one neuron to an effector cell as in muscle or gland.

Learning Objectives

- Describe the role of a synapse in nervous tissue

Key Points

- In the nervous system, a synapse is a structure that permits a neuron to pass an electrical or chemical signal to another cell.
- The neuron conducting impulses towards the synapse is called pre-synaptic neuron. The neuron transmitting the electrical impulse away from the synapse is called post-synaptic neuron, if the post-synaptic cell is not neuronal it is sometimes referred to as an effector cell.
- There are two varieties of synapses: electrical and chemical synapse based on the type of transmission.
- Chemical synapses rely on the secretion of neurotransmitters across a synaptic cleft between cells. Receptors on the post-synaptic cell bind these neurotransmitters and induce a signalling response.
- Electrical synapses form when the plasma membranes of two neurons are fused, and punctured by gap junction proteins, allowing for the diffusion of ions across the plasma cell membrane.
- Chemical synapses are slower than electrical synapses, but allow for gain, or amplification of signal.

Key Terms


- **axon**: A long, slender projection of a nerve cell, or neuron, that typically conducts electrical impulses away from the neuron's cell body.
- **synapse**: A structure that permits a neuron to pass an electrical or chemical signal to another cell.

In the nervous system, a synapse is a structure that permits the axon of a neuron to pass an electrical or chemical signal the axon of another neuron or to another cell type.

The neuron conducting impulses towards the synapse is called pre-synaptic neuron. The neuron transmitting the electrical impulse away from the synapse is called post-synaptic neuron, if the post-synaptic cell is not neuronal it is sometimes referred to as an effector cell.

Synapses can be classified by the type of cellular structures serving as the pre- and post-synaptic components. The vast majority of synapses in the mammalian nervous system are axo-axonal (axon synapsing with another axon) or axo-dendritic synapses (axon synapsing upon a dendrite). However, a variety of other arrangements exist.

Chemical and Electrical Synapses




Chemical Synapse: A: Pre-synaptic neuron B: Post-synaptic-neuron. Neurotransmitters are stored in synaptic vesicles within the pre-synaptic neuron (2). Upon receipt of an action-potential the neurotransmitter is secreted into the synaptic cleft (3 & 4) and binds to the receptors on the post-synaptic neuron (5).

There are two varieties of synapses: electrical and chemical synapse.

In a chemical synapse, the plasma membrane of the pre-synaptic neuron is closely associated with the plasma membrane of the post-synaptic cell, with the gap between termed the synaptic cleft. The synapse is stabilized by the expression of synaptic adhesion molecules projecting from both the pre- and post-synaptic cells maintaining the close association.

Upon arrival of an action potential at the pre-synaptic axon neurotransmitters are released into the synaptic cleft via the action of voltage-gated calcium channels. This neurotransmitter binds to receptors located in the plasma membrane of the post-synaptic cell which can elicit an electrical response or the activation of a secondary messenger pathway. Because of the complexity of receptor signal transduction, chemical synapses can have complex effects on the post-synaptic cell, and are able to induce effects such as gain, or amplification, whereby the strength of the signal is increased in the post-synaptic cell.

In an electrical synapse, the pre-synaptic and post-synaptic cell membranes are fused and connected by special channels called gap junctions that are capable of passing electrical current. These gap junctions contain connexion proteins which allow ions and small molecules to flow directly from one neuron to the next. The neurons are electrically coupled and transmission across these synapses is very rapid, allowing for faster signal processing than chemical synapses. However, due to their nature electical synapses cannot induce gain of signal strength.



Electrical Synapse: The membranes of pre and post-synaptic cells are fused and punctured by gap-junctions. When open they allow the rapid diffusion of ions across the plasma membranes allowing for rapid, continuous signal processing across the synapse.

This page titled 10.5F: The Synapse is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5G: Postsynaptic Potentials and Their Integration at the Synapse

Postsynaptic potentials are excitatory or inhibitory changes in the graded membrane potential in the postsynaptic terminal of a chemical synapse.

Learning Objectives

- Describe the role of postsynaptic potentials

Key Points

- Postsynaptic potentials are graded changes in the membrane potential of a postsynaptic synapse.
- Postsynaptic potentials are graded potentials and should not be confused with action potentials, although their function is to initiate or inhibit action potentials.
- Excitatory postsynaptic potentials (EPSP) bring the neuron's potential closer to its firing threshold.
- Inhibitory postsynaptic potentials (IPSP) change the charge across the membrane to be further from the firing threshold.
- Postsynaptic potentials are subject to spatial and temporal summation.

Key Terms

- **Postsynaptic potentials**: Changes in the membrane potential of the postsynaptic terminal of a chemical synapse.
- **inhibitory postsynaptic potential**: This occurs when the opening of the ion channels results in a net gain of negative charge, the potential moves further from zero and is referred to as hyperpolarization.
- **excitatory postsynaptic potential**: A net depolarization that brings the neuron's potential closer to its firing threshold.


Postsynaptic potentials are changes in the membrane potential of the postsynaptic terminal of a chemical synapse. Postsynaptic potentials are graded potentials and should not be confused with action potentials, although their function is to initiate or inhibit action potentials.

Many postsynaptic membrane receptors at chemical synapses are specialized to open ion channels. This converts a chemical signal into an electrical signal. Chemical synapses are either excitatory or inhibitory depending on how they affect the membrane potential of the postsynaptic neuron. The neurotransmitters bind to receptors on the postsynaptic terminal resulting in an opening of ion channels. At excitatory synapses, neurotransmitter binding depolarizes the postsynaptic membrane. Unlike the action potential in axonal membranes, chemically-gated ion channels open on postsynaptic membranes. Sodium and potassium diffuse simultaneously but in opposite directions.

Since the electrochemical gradient of sodium is steeper than that of potassium, a net depolarization occurs. If enough neurotransmitter binds, depolarization of the postsynaptic membrane can reach 0mV, which is higher than threshold of -30-50mV. This is an excitatory postsynaptic potential (EPSP) as it brings the neuron's potential closer to its firing threshold (about -50mV).

Neurotransmitter binding at inhibitory synapses reduces a postsynaptic neuron's ability to generate an action potential. Most inhibitory neurotransmitters hyperbolize the postsynaptic membrane by making it more permeable to potassium or chloride. When the opening of the ion channels results in a net gain of negative charge, the potential moves further from zero and is referred to as hyperpolarization. This is an inhibitory postsynaptic potential (IPSP).

EPSPs and IPSPs are transient changes in the membrane potential. A single EPSP at one synapse is generally far too small to trigger an action potential in the postsynaptic neuron. Postsynaptic potentials are subject to spatial and temporal summation.



Temporal summation: This figure depicts the mechanism of temporal summation in which multiple action potentials in the presynaptic cell cause a threshold depolarization in the postsynaptic cell.

This page titled 10.5G: Postsynaptic Potentials and Their Integration at the Synapse is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5H: Synaptic Transmission

Synaptic transmission is a chemical event which is involved in the transmission of the impulse via release, diffusion, receptor binding of neurotransmitter molecules and unidirectional communication between neurons.

Learning Objectives


- Describe the process of synaptic transmission

Key Points

- In a chemical synapse, the pre and post synaptic membranes are separated by a synaptic cleft, a fluid filled space.
- The chemical event is involved in the transmission of the impulse via release, diffusion, receptor binding of neurotransmitter molecules and unidirectional communication between neurons.
- The neurotransmitter termination can occur in three ways – reuptake, enzymatic degradation in the cleft and diffusion.

In a chemical synapse, the pre and post synaptic membranes are separated by a synaptic cleft, a fluid filled space. The chemical event is involved in the transmission of the impulse via release, diffusion, receptor binding of neurotransmitter molecules and unidirectional communication between neurons.

Chemical Synapse




Synaptic vesicles inside a neuron: This pseudocolored image taken with a scanning electron microscope shows an axon terminal that was broken open to reveal synaptic vesicles (blue and orange) inside the neuron.

Neurotransmission at a chemical synapse begins with the arrival of an action potential at the presynaptic axon terminal. When an action potential reaches the axon terminal, it depolarizes the membrane and opens voltage-gated Na^{+}channels. Na^{+} ions enter the cell, further depolarizing the presynaptic membrane. This depolarization causes voltage-gated Ca^{2+} channels to open. Calcium ions entering the cell initiate a signaling cascade. A calcium sensing protein binds calcium and interacts with SNARE proteins. These SNARE proteins are involved in the membrane fusion. The synaptic vesicles fuse with the presynaptic axon terminal membrane and empty their contents by exocytosis into the synaptic cleft. Calcium is quickly removed from the terminal.

Fusion of a vesicle with the presynaptic membrane causes neurotransmitters to be released into the synaptic cleft. The

neurotransmitter diffuses across the synaptic cleft, binding to receptor proteins on the postsynaptic membrane.



Communication at a chemical synapse: Communication at chemical synapses requires release of neurotransmitters. When the presynaptic membrane is depolarized, voltage-gated Ca2+ channels open and allow Ca2+ to enter the cell. The calcium entry causes synaptic vesicles to fuse with the membrane and release neurotransmitter molecules into the synaptic cleft. The neurotransmitter diffuses across the synaptic cleft and binds to ligand-gated ion channels in the postsynaptic membrane, resulting in a localized depolarization or hyperpolarization of the postsynaptic neuron.

The binding of a specific neurotransmitter causes particular ion channels, in this case ligand-gated channels, on the postsynaptic membrane to open. The binding of a neurotransmitter to its receptor is reversible. As long as it is bound to a post synaptic receptor, a neurotransmitter continues to affect membrane potential. The effects of the neurotransmitter generally lasts few milliseconds before being terminated. The neurotransmitter termination can occur in three ways. First, reuptake by astrocytes or presynaptic terminal where the neurotransmitter is stored or destroyed by enzymes. Second, degradation by enzymes in the synaptic cleft such as acetylcholinesterase. Third, diffusion of the neurotransmitter as it moves away from the synapse.

This page titled 10.5H: Synaptic Transmission is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5I: Types of Neurotransmitters by Function

Neurotransmitters are endogenous chemicals that transmit signals from a neuron to a target cell across a synapse.

Learning Objectives

- Describe types of neurotransmitters by their function

Key Points


- Neurotransmitters are endogenous chemicals that transmit signals from a neuron to a target cell across a synapse.
- Release of neurotransmitters usually follows arrival of an action potential at the synapse, but may also follow a graded electrical potential.
- The major types of neurotransmitters include acetylcholine, biogenic amines, and amino acids.
- Biogenic amines include the catecholamines such as dopamine, norepinephrine (NE), and epinephrine, as well as indolamines such as serotonin and histamine.

Key Terms

- **neurotransmitter**: An endogenous chemical that transmits signals from a neuron to a target cell across a synapse.
- **acetylcholine**: This neurotransmitter acts on the neuromuscular juncture and is synthesized from acetic acid and choline.
- **Biogenic amines**: Neurotransmitters distributed in the brain, where they play a role in emotional behavior and help in regulating the biological clock.
- **glutamate**: An amino acid that promotes excitatory effects by increasing the probability that the target cell will fire an action potential.

Neurotransmitters are endogenous chemicals that transmit signals from a neuron to a target cell across a synapse. Although some neurons produce and release only one kind of neurotransmitter, most make two or more and may release one or more at any given time. The coexistence of more than one neurotransmitter in the synapse makes it possible for the cell to exert several influences at the same time.

Neurotransmitter Structure



Major elements in neuron-to-neuron communication: Chemical synapses are specialized junctions through which neurons signal to each other and to non-neuronal cells such as those in muscles or glands.

Neurotransmitters are packaged into synaptic vesicles clustered beneath the membrane in the axon terminal on the presynaptic side of a synapse. They are released into and diffuse across the synaptic cleft, where they bind to specific receptors in the membrane on the postsynaptic side of the synapse. Release of neurotransmitters usually follows arrival of an action potential at the synapse, but may also follow a graded electrical potential. Low-level baseline release also occurs without electrical stimulation.

Neurotransmitters are synthesized from plentiful and simple precursors such as amino acids, which are readily available from the diet and require only a small number of biosynthetic steps to convert.

Classification of Neurotransmitters

Neurotransmitters fall into several chemical classes based on the molecular structure. The major types of neurotransmitters include acetylcholine, biogenic amines, and amino acids. The neurotransmitters can also be classified based on function (excitatory or inhibitory) and action (direct or neuromodulatory).

Acetylcholine

Acetylcholine, which acts on the neuromuscular junction, was the first neurotransmitter identified. It is synthesized from acetic acid and choline. Once released, acetylcholine binds to post-synaptic receptors and is degraded by acetylcholinesterase. Acetylcholine-releasing neurons are also found in the central nervous system (CNS).

Biogenic Amines

Biogenic amines include the catecholamines, such as dopamine, norepinephrine (NE), and epinephrine, as well as indolamines such as serotonin and histamine. Dopamine and NE are synthesized from amino acid tyrosine. Serotonin is synthesized from tryptophan. Histamine is synthesized from amino acid
histidine.

Biogenic amines are distributed in the brain, where they play a role in emotional behavior and help in regulating the biological clock. Additionally, some motor neurons of the ANS release catecholamines like NE. NE, dopamine, and histamine can be excitatory or inhibitory depending on the receptor type. Addictive drugs such as cocaine and amphetamine exert their effects primarily on the dopamine system, while addictive opiates and functional analogs of opioid peptides which regulate dopamine

levels.

Amino Acids

Glutamate and gamma-aminobutyric acid (GABA) are amino acid-based neurotransmitters. The most prevalent transmitter in the human brain is glutamate, which promotes excitatory effects by increasing the probability that the target cell will fire an action potential. The next most prevalent is GABA, which is inhibitory at more than 90% of the synapses that do not use glutamate.

Neuropeptides such as Substance P and endorphins are strings of amino acids that are important in the mediation of pain signals. Enkephalin activity increases dramatically in pregnant women in labor. Endorphins are released in the so called “runner’s high.” There is also evidence that neuropeptides such as gut-brain peptides are produced by non-neural tissues in the gastrointestinal tract.

This page titled 10.5I: Types of Neurotransmitters by Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5J: Ionotropic and Metabotropic Receptors

Although both ionotropic and metabotropic receptors are activated by neurotransmitters, ionotropic receptors are channel-linked while metabotropic receptors initiate a cascade of molecules via G-proteins.

Learning Objectives

- Distinguish between ionotropic and metabotropic receptors

Key Points

- Two types of membrane bound receptors (ionotropic and metabotropic) are activated with the binding of neurotransmitters.
- Ionotropic receptors such as nicotinic acetylcholine are a group of transmembrane ion channels that open or close in response to the binding of a chemical messenger.
- Metabotropic receptor are a type of G protein -coupled receptor
- When a metabotropic receptor is activated, a series of intracellular events are triggered that can also result in ion channels opening but must involve a range of second messenger chemicals.

Key Terms

- **nicotinic acetylcholine receptor**: A pentamer of protein subunits with two binding sites for acetylcholine which, when bound, alter the receptor's configuration and cause an internal pore to open.
- **Ionotropic receptors**: A group of transmembrane ion channels that open or close in response to the binding of a chemical messenger (ligand) such as a neurotransmitter.
- **metabotropic receptors**: A subtype of membrane receptors that do not form an ion channel pore but use signal transduction mechanisms, often G proteins, to activate a series of intracellular events using second messenger chemicals.

Two types of membrane-bound receptors are activated with the binding of neurotransmitters: ligand-gated ion channels (LGICs) inotropic receptors and metabotropic G- protein coupled receptors.

Ionotropic Receptors

Ionotropic receptors are a group of transmembrane ion channels that open or close in response to the binding of a chemical messenger (ligand) such as a neurotransmitter.The binding site of endogenous ligands on LGICs protein complexes are normally located on a different portion of the protein (an allosteric binding site) than the location of the ion conduction pore.The ion channel is regulated by a ligand and is usually very selective to one or more ions such as Na^+, K^+, Ca^{2+}, or Cl^-.

The prototypic ligand-gated ion channel is the nicotinic acetylcholine receptor. It consists of a pentamer of protein subunits with two binding sites for acetylcholine which, when bound, alter the receptor's configuration and cause an internal pore to open. This pore allows Na^+ ions to flow down their electrochemical gradient into the cell. With a sufficient number of channels opening at once, the inward flow of positive charges carried by Na^+ ions depolarizes the postsynaptic membrane enough to initiate an action potential.

Metabotropic Receptors

Metabotropic receptors are a subtype of membrane receptors that do not form an ion channel pore but use signal transduction mechanisms, often G proteins, to activate a series of intracellular events using second messenger chemicals. Examples of metabotropic receptors include glutamate receptors, muscarinic acetylcholine receptors, $GABA_B$ receptors, most serotonin receptors, and receptors for norepinephrine, epinephrine, histamine, dopamine, neuropeptides, and endocannabinoids.

The metabotropic G protein-coupled receptors have seven hydrophobic transmembrane domains. When the neurotransmitter binds to the receptor, there is an activation via the G-protein that later activates the secondary messengers. Metabotropic receptors on the presynaptic membrane can inhibit or more rarely facilitate neurotransmitter release from the presynaptic neuron.

Since opening channels by metabotropic receptors involves activating a number of molecules in the intracellular mechanism, these receptors take longer to open than the inotropic receptors. They have a much longer effect than ionotropic receptors, which open quickly but only remain open for a few milliseconds. While ionotropic channels have an effect only in the immediate region of the receptor, the effects of metabotropic receptors can be more widespread throughout the cell.

This page titled 10.5J: Ionotropic and Metabotropic Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

10.5K: Serial and Parallel Processing

Serial memory processing compares a memory to a target stimulus, while parallel processing carries out multiple operations simultaneously.

Learning Objectives

- Distinguish between serial and parallel processing by nervous tissue

Key Points

- Serial memory processing can be either self-terminating or exhaustive.
- Self-terminating serial memory processing implies that comparisons stop abruptly as soon as the target is found, and then the response is generated.
- Exhaustive serial memory processing implies that comparisons continue until the entire set is compared and then a response is generated.
- Parallel processing is the ability of the brain to simultaneously process incoming stimuli of differing quality.
- Parallel processing becomes most important in vision, as the brain divides what it sees into four components: color, motion, shape, and depth.
- Serial memory processing uses internal representations of the memory set in order to compare them to a target stimulus or item.

Key Terms

- **Serial memory processing**: The act of attending to and processing one item at a time.
- **self-terminating**: Self-terminating implies that comparisons stop abruptly as soon as the target is found, and then the response is generated.
- **parallel processing**: Parallel processing is the ability to carry out multiple operations or tasks simultaneously.


Serial memory processing is the act of attending to and processing one item at a time, while parallel memory processing is the act of attending to and processing all items simultaneously. In short-term memory tasks, participants are given a set of items (e.g. letters, digits) one at a time and then, after varying periods of delay, are asked for recall of the items. Participants could also be asked whether a specific target item was present in their original set. The serial order of items and the relationships between them can have varying effects on speed and accuracy of recall.

Serial Memory Processing

Serial memory processing compares internal representations of the memory set to a target stimulus or item is being presented, one at a time. Reaction time increases linearly with the set size, which means the more items in the memory set, the longer it will take to compare.

Serial memory processing can be either self-terminating or exhaustive.

Self-Terminating Processing



Self-Terminating Search: This line graph depicts both positive and negative self-terminating search, comparing length of list to mean reaction time recall.

Self-terminating implies that comparisons stop abruptly as soon as the target is found, and then the response is generated. Evidence for this method is found in reaction time studies. If the reaction time slope for a positive trial (where the target was present in the memory set) is about half of the slope for a negative trial (where the target was not present in the memory set), this demonstrates self-terminating processing. This is because on average, participants (on positive trials) will stop comparisons about halfway through when they find the target match, but participants (on negative trials) must compare until the end when no target match is found.

Exhaustive Processing

On the other hand, exhaustive implies that comparisons continue until the entire set is compared and then a response is generated. Evidence for this method is also found in reaction time studies. In this case, the reaction time slope is equal for both positive and negative trials, as comparisons are made to the end in both cases. Participants may process some serial memory sets using the self-terminating method and others with the exhaustive method. There is no clear distinction about which method is better to use.

Parallel Processing

Parallel processing is the ability to carry out multiple operations or tasks simultaneously. The term is used in the contexts of human cognition, particularly in the ability of the brain to simultaneously process incoming stimuli, and in parallel computing by machines.

In parallel processing, the brain simultaneously processes incoming stimuli of differing quality. This is most important in vision, as the brain divides what it sees into four components: color, motion, shape, and depth. These are individually analyzed and then compared to stored memories, which helps the brain identify what you are viewing. The brain then combines all four components into the field of view that you see and comprehend. This is a continual and seamless operation.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- potential difference. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/potential_difference**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Membrane potential. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/**

Membrane_potential%23Physical_basis. License: CC BY-SA: Attribution-ShareAlike
- Voltage. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Voltage. License: CC BY-SA: Attribution-ShareAlike
- voltage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/voltage. License: CC BY-SA: Attribution-ShareAlike
- voltmeter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/voltmeter. License: CC BY-SA: Attribution-ShareAlike
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. License: CC BY-SA: Attribution-ShareAlike
- Membrane potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Membrane_potential%23Ion_pumps. License: CC BY-SA: Attribution-ShareAlike
- ligand-gated channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/ligand-gated%20channel. License: CC BY-SA: Attribution-ShareAlike
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/ion-pump. License: CC BY-SA: Attribution-ShareAlike
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/leakage-channel. License: CC BY-SA: Attribution-ShareAlike
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. License: CC BY-SA: Attribution-ShareAlike
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. License: Public Domain: No Known Copyright
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. License: Public Domain: No Known Copyright
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:S...um_pump-en.svg. License: Public Domain: No Known Copyright
- Membrane potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Membrane_potential%23Resting_potential. License: CC BY-SA: Attribution-ShareAlike
- Goldman equation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Goldman%20equation. License: CC BY-SA: Attribution-ShareAlike
- ion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ion. License: CC BY-SA: Attribution-ShareAlike
- resting membrane potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/resting...ne%20potential. License: CC BY-SA: Attribution-ShareAlike
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. License: CC BY-SA: Attribution-ShareAlike
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. License: Public Domain: No Known Copyright
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. License: Public Domain: No Known Copyright
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:S...um_pump-en.svg. License: Public Domain: No Known Copyright
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped...48677b9305.png. License: CC BY-SA: Attribution-ShareAlike
- Membrane potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Membrane_potential. License: CC BY-SA: Attribution-ShareAlike
- graded potentials. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/graded%20potentials. License: CC BY-SA: Attribution-ShareAlike
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...depolarization. License: CC BY-SA: Attribution-ShareAlike
- action potential. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/action_potential. License: CC BY-SA: Attribution-ShareAlike
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. License: CC BY-SA: Attribution-ShareAlike
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. License: Public Domain: No Known Copyright
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. License: Public Domain: No Known Copyright
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:S...um_pump-en.svg. License: Public Domain: No Known Copyright

- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped...48677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Action_...physical_basis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...tion_potential. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...depolarization. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...ory-conduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- action potential. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/action_potential. **License**: *CC BY-SA: Attribution-ShareAlike*
- node of Ranvier. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/node_of_Ranvier. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain: No Known Copyright*
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped...48677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neurotransmitter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synapse...urotransmitter. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neurotransmitter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synapse...urotransmitter. **License**: *CC BY-SA: Attribution-ShareAlike*
- postsynaptic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/postsynaptic. **License**: *CC BY-SA: Attribution-ShareAlike*
- synapse. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/synapse. **License**: *CC BY-SA: Attribution-ShareAlike*
- action potential. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/action_potential. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain: No Known Copyright*
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F...unction-en.svg. **License**: *Public Domain: No Known Copyright*
- Synapse diag1. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/F...apse_diag1.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

- Postsynaptic potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postsynaptic_potential. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neuromodulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neuromodulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...tial-summation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...oral-summation. **License**: *CC BY-SA: Attribution-ShareAlike*
- postsynaptic potential. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/postsynaptic%20potential. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain: No Known Copyright*
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F...unction-en.svg. **License**: *Public Domain: No Known Copyright*
- Synapse diag1. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/F...apse_diag1.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Temporal summation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Te..._summation.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44748/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Synaptic transmission. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Synaptic_transmission. **License**: *CC BY-SA: Attribution-ShareAlike*
- postsynaptic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/postsynaptic. **License**: *CC BY-SA: Attribution-ShareAlike*
- presynaptic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/presynaptic. **License**: *CC BY-SA: Attribution-ShareAlike*
- synapse. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/synapse. **License**: *CC BY-SA: Attribution-ShareAlike*
- neurotransmitter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurotransmitter. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Basis_of_Membrane_Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain: No Known Copyright*
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File%3AGap_cell_junction-en.svg. **License**: *Public Domain: No Known Copyright*

- Synapse diag1. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Synapse_diag1.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Temporal summation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Te..._summation.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44748/latest...e_35_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44748/latest...e_35_02_07.jpg. **License**: *CC BY: Attribution*
- Neurotransmitter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neurotr...rotransmitters. **License**: *CC BY-SA: Attribution-ShareAlike*
- opioid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/opioid. **License**: *CC BY-SA: Attribution-ShareAlike*
- dopamine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/dopamine. **License**: *CC BY-SA: Attribution-ShareAlike*
- acetylcholine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/acetylcholine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Basis_of_Membrane_Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain: No Known Copyright*
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ac...ntial_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Action potential vert. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Action_potential_vert.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File%3AGap_cell_junction-en.svg. **License**: *Public Domain: No Known Copyright*
- Synapse diag1. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Synapse_diag1.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Temporal summation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Te..._summation.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44748/latest...e_35_02_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44748/latest...e_35_02_07.jpg. **License**: *CC BY: Attribution*
- Nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nervous...tem%23Function. **License**: *Public Domain: No Known Copyright*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...ropic-receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Metabotropic receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Metabotropic_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ionotropic. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ionotropic. **License**: *CC BY-SA: Attribution-ShareAlike*
- receptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...upled-receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- electrochemical gradient. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/electro...cal%20gradient. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Basis_of_Membrane_Potential2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- LGIC. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LGIC.png. **License**: *Public Domain: No Known Copyright*
- Ion channel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ion_channel.png. **License**: *Public Domain:*

No Known Copyright

- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Action potential vert. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Action_potential_vert.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Action potential vert. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Action_potential_vert.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3AGap_cell_junction-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Synapse diag1. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Synapse_diag1.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Temporal summation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Te..._summation.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44748/latest...e_35_02_06.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44748/latest...e_35_02_07.jpg**. **License**: ***CC BY: Attribution***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous...tem%23Function**. **License**: ***Public Domain: No Known Copyright***
- Serial memory processing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Serial_memory_processing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parallel processing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parallel_processing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...lel-processing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...lf-terminating**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basis of Membrane Potential2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Basis_of_Membrane_Potential2.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- LGIC. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:LGIC.png**. **License**: ***Public Domain: No Known Copyright***
- Ion channel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ion_channel.png**. **License**: ***Public Domain: No Known Copyright***
- Scheme sodium-potassium pump-en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Scheme_sodium-potassium_pump-en.svg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **https://upload.wikimedia.org/Wikipedia/en/math/5/9/3/5939d2ef41c086b20b25a948677b9305.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Action potential vert. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Action_potential_vert.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Action potential vert. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Action_potential_vert.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gap_cell_junction-en.svg.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File%3AGap_cell_junction-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Synapse diag1. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Synapse_diag1.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Temporal summation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Temporal_summation.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44748/latest/Figure_35_02_06.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, How Neurons Communicate. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44748/latest/Figure_35_02_07.jpg**. **License**: ***CC BY: Attribution***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Function**. **License**: ***Public Domain: No Known Copyright***
- Graph showing Self-Terminating Search. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Graph_showing_Self-Terminating_Search.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 10.5K: Serial and Parallel Processing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11: CENTRAL NERVOUS SYSTEM



-  Topic hierarchy

11.1: The Spinal Cord

11.1A: Overview of the Spinal Cord

11.1B: The Spine

11.1C: Spinal Cord Grey Matter and Spinal Roots

11.1D: Spinal Cord White Matter

11.2: The Brain

11.2A: The Brain

11.3: Protection of the Brain

11.3A: Blood-Brain Barrier

11.3B: Cerebrospinal Fluid and Its Circulation

11.3C: Ventricles

11.4: The Brain Stem

11.4A: Functions of the Brain Stem

11.4B: Medulla Oblongata

11.4C: Pons

11.4D: Midbrain

11.4E: Reticular Formation

11.5: The Cerebellum

11.5A: Parts of the Cerebellum

11.5B: Functions of the Cerebellum

11.6:The Diencephalon

11.6A: Functions of the Diencephalon

11.6B: Thalamus

11.6C: Hypothalamus

11.6D: Epithalamus and Pineal Gland

11.6E: Circumventricular Organs

11.7: The Cerebrum

11.7A: Overview of the Cerebrum

11.7B: Cerebral Lobes

11.7C: White Matter of the Cerebrum

11.7D: Basal Ganglia

11.7E: Limbic System

11.8: Functional Systems of the Cerebral Cortex

11.8A: Sensory Areas

11.8B: Motor Areas

11.8C: Association Areas

11.8D: Hemispheric Lateralization

Index

This page titled 11: Central Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.1: THE SPINAL CORD

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

11.1A: Overview of the Spinal Cord

11.1B: The Spine

11.1C: Spinal Cord Grey Matter and Spinal Roots

11.1D: Spinal Cord White Matter

This page titled 11.1: The Spinal Cord is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.1A: Overview of the Spinal Cord

The spinal cord runs along the inside of the vertebral column and serves as the signaling conduit between the brain and the periphery.

Learning Objectives

- Describe the features of the spinal cord

Key Points

- The spinal cord extends from the occipital bone of the skull until it terminates near the second lumbar vertebra.
- The spinal cord is protected by three layers of meninges: the dura mater, the arachnoid mater, and the pia mater.
- The central nervous system (CNS) is made up of the brain and spinal cord. The area between the arachnoid space and the pia mater contains cerebral spinal fluid (CSF).
- The spinal cord is divided into 31 segments that send nerve rootlets out into the body through intervertebral foramen.
- Each segment of the spinal cord is associated with a pair of ganglia called dorsal root ganglia, which are situated just outside of the spinal cord and contain cell bodies of sensory neurons. These neurons travel into the spinal cord via the dorsal roots.
- Ventral roots consist of axons from motor neurons, which bring information to the periphery from cell bodies within the CNS. Dorsal roots and ventral roots come together and exit the intervertebral foramina as they become spinal nerves.


Key Terms

- **peripheral nervous system**: The part of the nervous system that consists of the nerves and ganglia on the outside of the brain and spinal cord.
- **efferent**: The conduction of impulses outward from the brain or spinal cord.
- **afferent**: The conduction of impulses inwards to the brain or spinal cord.
- **cauda equina**: A bundle of nerve roots at the base of the spinal column.
- **spinal cord**: A thick, whitish cord of nerve tissue which is a major part of the vertebrate central nervous system. It extends from the brain stem down through the spine, with nerves branching off to various parts of the body.

EXAMPLES

- A lumbar puncture (spinal tap) is an example of a medical procedure that directly targets the spinal cord.
- The birth defect spina bifida is a failure of the vertebral arch to close, exposing the spinal cord.

The spinal cord is a long, thin, tubular bundle of nervous tissue and support cells that extends from the medulla oblongata of the brain to the level of the lumbar region. The brain and spinal cord together make up the central nervous system (CNS). The spinal cord, protected by the vertebral column, begins at the occipital bone and extends down to the space between the first and second lumbar vertebrae. The spinal cord has a varying width, ranging from 0.5 inch thick in the cervical and lumbar regions to 0.25 inch thick in the thoracic area. The length of the spinal cord is approximately 45 cm (18 in) in men and about 43 cm (17 in) long in women.



Spinal Cord and Vertebrae.png: Relationship between the spinal cord and vertebral column, delineating the cervical, thoracic, and lumbar sections.

Layers and Regions of the Spinal Cord


The spinal cord is protected by three layers of tissue called meninges and divided into three regions.

Spinal Cord Tissue Layers

The dura mater is the outermost layer of spinal cord tissue, forming a tough protective coating. The space between the dura mater and the surrounding bone of the vertebrae is called the epidural space. The epidural space is filled with adipose tissue and contains a network of blood vessels. The middle layer is called the arachnoid mater. The pia mater is the innermost protective layer and is tightly associated with the surface of the spinal cord. The space between the arachnoid and pia maters is called the subarachnoid space and is where the CSF is located. It is from this location at the level of the lumbar region that CSF fluid is obtained in a spinal tap.

Spinal Cord Regions

In cross-section, the peripheral region of the cord displays neuronal white matter tracts containing sensory and motor neurons. Internal to this peripheral region is the gray, butterfly-shaped central region made up of nerve cell bodies. This central region surrounds the central canal, which is an anatomic extension of the spaces in the brain known as the ventricles and like the ventricles, contains cerebrospinal fluid.



Spinal Cord Regions: Cross-sectional perspective of the spinal cord regions

The spinal cord is divided into cervical, thoracic, and lumbar regions. The cervical region is divided into eight levels that are related to different motor and sensory functions in the neck and the arms. The spinal nerves of the thoracic region supply the thorax and abdomen. The nerves of the lumbosacral spinal cord supply the pelvic region, legs, and feet.

Spinal Cord Nerve Branches

Thirty-one pairs of spinal nerves (sensory and motor) branch from the human spinal cord. Each spinal nerve is formed from the combination of nerve fibers from its posterior and anterior roots. The posterior root is the sensory (afferent) root that carries sensory information to the brain from other areas of the body. The anterior root is the motor (efferent) root that carries motor information to the body from the brain.

The spinal nerve emerges from the spinal column through the opening (intervertebral foramen) between adjacent vertebrae. An exception is the first spinal nerve pair (C1), which emerges between the occipital bone and the atlas (the first vertebra). The swelling found in the posterior root is the posterior (dorsal) root ganglion, which contains the cell bodies of sensory neurons. The anterior (ventral) root contains axons of motor neurons that conduct nerve impulses from the CNS to other parts of the body such as the muscles.

The cauda equina (“horse’s tail”) is the name for the collection of nerves in the vertebral column that extends beyond the cord. The nerves that compose the cauda equina supply the pelvic organs and lower limbs, including motor innervation for the hips, knees, ankles, feet, and internal and external anal sphincters. In addition, the cauda equina extends to sensory innervation of the perineum.

Primary Spinal Cord Function

The spinal cord functions primarily in the transmission of neural signals between the brain and the rest of the body, but it also contains neural circuits that can independently control numerous reflexes and central pattern generators. The three major functions of the spinal cord are the conduction of motor information traveling down the spinal cord, the conduction of sensory information in the reverse direction, and acting as the center for conducting certain reflexes. The spinal cord is the main pathway for information connecting the brain and peripheral nervous system.

This page titled 11.1A: Overview of the Spinal Cord is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.1B: The Spine

The spine encases the spinal cord for protection and support.

Learning Objectives

- Describe the vertebral column, the protective structure of the spinal cord

Key Points

- The human spine consists of 24 articulating vertebrae grouped into cervical, thoracic, and lumbar regions. Nine more vertebrae make up the sacrum and coccyx.
- Typical vertebrae consist of the anterior vertebral body and the posterior section, which contains the vertebral foramen through which the spinal cord passes.
- There are four main curves of the spine: cervical, thoracic, lumbar, and pelvic.
- Facets of the vertebrae restrict range of movement to prevent shearing of the spinal cord.
- Blood vessels and nerves exit the spinal column at intervertebral foramina.
- There are four main curves of the spine: cervical, thoracic, lumbar and pelvic.


Key Terms

- **vertebrae**: The bones that make up the spinal column.
- **laminae**: Plates of bone that form the posterior walls of each vertebra.
- **pedicle**: A segment of bone connecting the lamina to the vertebral body.
- **vertebral foramen**: Formed by the vertebral body and vertebral arch and containing the spinal cord.
- **vertebral column**: The series of vertebrae that protect the spinal cord; the spinal column.

EXAMPLES

- Kyphosis is an exaggerated concave (kyphotic) curvature of the thoracic vertebral column; it is commonly known as “humpback.”
- Lordosis is an exaggerated convex (lordotic) curvature of the lumbar region; it is commonly known as “swayback.”
- Scoliosis is an abnormal lateral curvature of the vertebral column.

Number of Vertebrae




Vertebral Column: The sections of the vertebral column.

In human anatomy, the vertebral column (backbone or spine) usually consists of 24 articulating vertebrae and nine fused vertebrae in the sacrum and the coccyx. Situated in the dorsal aspect of the torso and separated by intervertebral discs, it houses and protects the spinal cord in its spinal canal. There are normally 33 vertebrae in humans, including the five that are fused to form the sacrum, the four coccygeal bones that form the tailbone, and the others separated by intervertebral discs. The upper three regions comprise the remaining 24, and are grouped as cervical (seven vertebrae), thoracic (12 vertebrae) and lumbar (five vertebrae).

Vertebral Shape

A typical vertebra consists of the vertebral body and vertebral arch. These parts together enclose the vertebral foramen that contains the spinal cord. The vertebral arch is formed by a pair of pedicles and a pair of laminae. Two transverse processes and one spinous process are posterior to (behind) the vertebral body. The spinous process projects toward the posterior direction, while one transverse process projects to the left and the other to the right. The spinous processes of the cervical and lumbar regions can be felt through the skin. Facet joints are located above and below each vertebra. These restrict the range of movement. Between each pair of vertebrae are two small openings called intervertebral foramina through which the spinal nerves exit.



Vertebrae: Oblique view of cervical vertebrae.

Vertebral Curvature

When viewed laterally, the vertebral column presents several curves corresponding to the different regions of the column: cervical, thoracic, lumbar, and pelvic.

Cervical and Thoracic Curves

The cervical curve convexes forward and begins at the apex of the odontoid (tooth-like) process. It ends at the middle of the second thoracic vertebra. The thoracic curve convexes dorsally, begins at the middle of the second thoracic vertebra, and ends at the middle of the 12th thoracic vertebra.

Lumbar and Pelvic Curves

The lumbar curve, which is more pronounced in women than in men, begins at the middle of the last thoracic vertebra and ends at the sacrovertebral angle. It is convex anteriorly with the lower three vertebrae much more convex than the upper two. This curve is described as a lordotic curve. The pelvic curve begins at the sacrovertebral articulation and ends at the point of the coccyx; its concavity is directed downward and forward.

Primary and Secondary Curves

The thoracic and sacral curvatures are termed primary curves because they are present in the fetus and remain the same in the adult. As the child grows, lifts the head, and begins to assume an upright position, the secondary curves (cervical and lumbar) develop. The cervical curve forms when the infant is able to hold up his or her head (at three or four months) and sit upright (at nine months). The lumbar curve forms between twelve to eighteen months when the child begins to walk.

This page titled 11.1B: The Spine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.1C: Spinal Cord Grey Matter and Spinal Roots

The grey matter of the spinal cord contains neuronal cell bodies, dendrites, axons, and nerve synapses.

Learning Objectives

- Describe the grey matter and spinal roots of the spinal cord

Key Points

- Each segment of the spinal cord is associated with a pair of ganglia called dorsal root ganglia, situated just outside of the spinal cord.
- The dorsal root ganglia contain the cell bodies of sensory neurons. Axons of these sensory neurons travel into the spinal cord via the dorsal roots.
- The grey matter in the center of the cord contains interneurons and the cell bodies of motor neurons, axons, and dendrites.
- Projections of the grey matter (the “wings”) are called horns. Together, the grey horns and the grey commissure form the H-shaped grey matter.
- The dorsal root ganglia develops in the embryo from neural crest cells. The spinal ganglia can thus be regarded as grey matter of the spinal cord that was translocated to the periphery.

Key Terms


- **neural crest**: A strip of ectodermal material in the early vertebrate embryo inserted between the prospective neural plate and the epidermis.
- **grey matter**: A major component of the central nervous system consisting of neuronal cell bodies, neuropil (dendrites and unmyelinated axons), glial cells (astroglia and oligodendrocytes), and capillaries.
- **neural tube**: The embryonic precursor to the central nervous system (CNS).

EXAMPLES

- The spine acts as the conduit to relay information to and from the brain from the rest of the body.
- Damage to the grey matter (eg, the ventral gray horn) may lead to tingling and muscle weakness.

The spinal cord is the main pathway for information connecting the brain and peripheral nervous system. The spinal cord is much shorter in length than the bony spinal column. The human spinal cord extends from the foramen magnum of the occipital bone of the skull and continues to the conus medullaris near the second lumbar vertebra, terminating in a fibrous extension known as the filum terminale.

Spinal Cord Topography and Roots



Cross-Section of Spinal Cord, Mid-Thoracic Level: Note the commissure, rendered in grey.

The spinal cord is compressed dorsoventrally, giving it an elliptical shape. The cord has grooves in the dorsal and ventral sides. The posterior median sulcus is the groove in the dorsal side, and the anterior median fissure is the groove in the ventral side.

Each segment of the spinal cord is associated with a pair of ganglia called dorsal root ganglia, situated just outside of the spinal cord. These ganglia contain cell bodies of sensory neurons. Axons of these sensory neurons travel into the spinal cord via the dorsal roots.

The grey matter, in the center of the cord, is shaped like a butterfly and consists of cell bodies of interneurons and motor neurons, as well as neuroglia cells and unmyelinated axons. Projections of the grey matter (the “wings”) are called horns. Together, the grey horns and the grey commissure form the H-shaped grey matter.

Dorsal and Ventral Roots

The dorsal root ganglia lie along the vertebral column by the spine. The dorsal root ganglia develops in the embryo from neural crest cells, not the neural tube. Hence, the spinal ganglia can be regarded as grey matter of the spinal cord that became translocated to the periphery.

The axons of dorsal root ganglion neurons are known as afferents. In the peripheral nervous system, afferents refer to the axons that relay sensory information into the central nervous system. These neurons are of the pseudo-unipolar type, meaning that they have an axon with two branches that act as a single axon, often referred to as distal and proximal processes. Ventral roots consist of axons from motor neurons, which bring information to the periphery from cell bodies within the CNS. Dorsal roots and ventral roots come together and exit the intervertebral foramina as they become spinal nerves.

The nerve endings of dorsal root ganglion neurons have a variety of sensory receptors that are activated by mechanical, thermal, chemical, and noxious stimuli. In these sensory neurons, a group of ion channels thought to be responsible for somatosensory transduction has been identified. Compression of the dorsal root ganglion by a mechanical stimulus lowers the voltage threshold needed to evoke a response and causes action potentials to be fired. This firing may even persist after the removal of the stimulus.

Impulse Transmission

The dendrite receives information from another neuron’s axon at the synapse, and the axon sends information to the next neuron’s dendrites. Unlike the majority of neurons found in the CNS, an action potential in a dorsal root ganglion neuron may initiate in the distal process in the periphery, bypass the cell body, and continue to propagate along the proximal process until reaching the

synaptic terminal in the dorsal horn of the spinal cord.

The distal section of the axon may either be a bare nerve ending or encapsulated by a structure that helps relay specific information to nerve. For example, a Meissner's corpuscle or a Pacinian corpuscle may encapsulate the nerve ending, rendering the distal process sensitive to mechanical stimulation, such as stroking or vibration.

Ion Channels

Two distinct types of mechanosensitive ion channels have been found in the dorsal root ganglia, broadly classified as either high-threshold (HT) or low-threshold (LT). As their names suggest, they have different thresholds as well as different sensitivities to pressure. These are cationic channels whose activity appears to be regulated by the proper functioning of the cytoskeleton and cytoskeleton-associated proteins. The presence of these channels in the dorsal root ganglion gives reason to believe that other sensory neurons may contain them as well.

High-threshold channels have a possible role in nociception. These channels are found predominantly in smaller sensory neurons in the dorsal root ganglion cells and are activated by higher pressures, two attributes that are characteristic of nociceptors. Also, the threshold of HT channels was lowered in the presence of PGE2 (a compound that sensitizes neurons to mechanical stimuli and mechanical hyperalgesia), which further supports a role for HT channels in the transduction of mechanical stimuli into nociceptive neuronal signals.

This page titled 11.1C: Spinal Cord Grey Matter and Spinal Roots is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.1D: Spinal Cord White Matter

The white matter of the spinal cord is composed of bundles of myelinated axons.

Learning Objectives

- Describe the function and compositon of spinal cord white matter

Key Points

- White matter is one of the two components of the central nervous system and consists mostly of glial cells and myelinated axons.
- The white matter is white because of the fatty substance (myelin) that surrounds the nerve fibers. Myelin acts as an electrical insulation. It allows the messages to pass quickly from place to place.
- Cerebral and spinal white matter do not contain dendrites, which can only be found in grey matter along with neural cell bodies, and shorter axons.
- White matter modulates the distribution of action potentials, acting as a relay and coordinating communication between different brain regions.
- White matter in the spinal cord functions as the “wiring”; primarily to carry information.

Key Terms

- **myelin**: A white, fatty, material composed of lipids and lipoproteins, that surrounds the axons of nerves.
- **white matter**: A region of the central nervous system containing myelinated nerve fibers and no dendrites.
- **cerebral ventricles**: Interconnected cavities in the brain where the cerebrospinal fluid is produced.
- **glial cell**: A type of cell, in the nervous system, that provides support for the neurons.

White matter is one of the two components of the central nervous system. It consists mostly of glial cells and myelinated axons and forms the bulk of the deep parts of the brain and the superficial parts of the spinal cord. It is the tissue through which messages pass between different areas of grey matter within the nervous system.

Composition of White Matter

White matter is composed of bundles of myelinated nerve cell processes (or axons). The axons connect various grey matter areas (the locations of nerve cell bodies) of the brain to each other and carry nerve impulses between neurons. The axonal myelin acts as an insulator and increases the speed of transmission of all nerve signals. White matter does not contain dendrites, which are only found in grey matter along with neural cell bodies and shorter axons.

In a freshly cut brain, the tissue of white matter appears pinkish white to the naked eye because myelin is composed largely of lipid tissue that contains capillaries. In nonelderly adults, 1.7-3.6% of the white matter is blood. Myelin is found in almost all long nerve fibers and acts as electrical insulation. This is important because it allows the messages to pass quickly from place to place.

Spinal Cord Columns

The spinal cord white matter is subdivided into columns. The dorsal columns carry sensory information from mechanoreceptors (cells that respond to mechanical pressure or distortion). The axons of the lateral columns (corticospinal tracts) travel from the cerebral cortex to contact spinal motor neurons. The ventral columns carry sensory pain and temperature information and some motor information.

Function of White Matter

Long thought to be passive tissue, white matter actively affects how the brain learns and functions. While grey matter is primarily associated with processing and cognition, white matter modulates the distribution of action potentials, acting as a relay and coordinating communication between different brain regions. The brain in general (and especially a child's brain) can adapt to white-matter damage by finding alternative routes that bypass the damaged white-matter areas; therefore, it can maintain good connections between the various areas of grey matter. Using a computer network as an analogy, the grey matter can be thought of as the actual computers themselves, whereas the white matter represents the network cables connecting the computers together.


Axon Tracts

Within white matter, there are three different kinds of tracts or bundles of axons that connect one part of the brain to another and to the spinal cord:

1. Projection tracts extend vertically between higher and lower brain and spinal cord centers. They carry information between the cerebrum and the rest of the body. The corticospinal tracts, for example, carry motor signals from the cerebrum to the brainstem and spinal cord.
2. Commissural tracts cross from one cerebral hemisphere to the other through bridges called commissures. Commissural tracts enable the left and right sides of the cerebrum to communicate with each other.
3. Association tracts connect different regions within the same hemisphere of the brain. Among their roles, association tracts link perceptual and memory centers of the brain.

White Matter-Grey Matter Interactions

White matter forms the bulk of the deep parts of the brain and the superficial parts of the spinal cord. Aggregates of grey matter, such as the basal ganglia and brain stem nuclei, are spread within the cerebral white matter. The cerebellum is structured in a similar manner as the cerebrum, with a superficial mantle of cerebellar cortex, deep cerebellar white matter (called the "arbor vitae"), and aggregates of grey matter surrounded by deep cerebellar white matter (dentate nucleus, globose nucleus, emboliform nucleus, and fastigial nucleus). The fluid-filled cerebral ventricles (lateral ventricles, third ventricle, cerebral aqueduct, and fourth ventricle) are also located deep within the cerebral white matter.



White matter in spinal cord: The spinal cord diagram showing location of the white matter surrounding grey matter.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- spinal cord. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- cauda equina. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cauda_equina. **License**: *CC BY-SA: Attribution-ShareAlike*
- cross section.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Denticulate_ligaments. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Cord and Vertebrae.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human vertebral column. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_vertebral_column. **License**: *CC BY-SA: Attribution-ShareAlike*
- vertebrae. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vertebrae. **License**: *CC BY-SA: Attribution-ShareAlike*
- vertebral column. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vertebral_column. **License**: *CC BY-SA: Attribution-ShareAlike*
- cross section.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Denticulate_ligaments. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Cord and Vertebrae.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vertebral Column. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f8/Illu_vertebral_column.jpg. **License**: *Public Domain: No Known Copyright*
- ACDF oblique annotated english. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ACDF_oblique_annotated_english.svg. **License**: *Public Domain: No Known Copyright*
- Spinal cord. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- grey matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/grey%20matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsal root. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsal%20root. **License**: *CC BY-SA: Attribution-ShareAlike*
- neural crest. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neural_crest. **License**: *CC BY-SA: Attribution-ShareAlike*
- cross section.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Denticulate_ligaments. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Cord and Vertebrae.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vertebral Column. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f8/Illu_vertebral_column.jpg. **License**: *Public Domain: No Known Copyright*
- ACDF oblique annotated english. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ACDF_oblique_annotated_english.svg. **License**: *Public Domain: No Known Copyright*
- Gray664. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray664.png. **License**: *Public Domain: No Known Copyright*
- Ventricular System. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventricular_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Cord. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_cord. **License**: *CC BY-SA: Attribution-ShareAlike*
- White matter. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/White_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- myelin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/myelin. **License**: *CC BY-SA: Attribution-ShareAlike*
- white matter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/white_matter. **License**: *CC BY-SA: Attribution-ShareAlike*
- glial cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glial_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- cross section.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Denticulate_ligaments. **License**: *CC BY-SA: Attribution-ShareAlike*

- Spinal Cord and Vertebrae.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_cord**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vertebral Column. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f8/Illu_vertebral_column.jpg**. **License**: ***Public Domain: No Known Copyright***
- ACDF oblique annotated english. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ACDF_oblique_annotated_english.svg**. **License**: ***Public Domain: No Known Copyright***
- Gray664. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray664.png**. **License**: ***Public Domain: No Known Copyright***
- Anatomy and physiology of animals The spinal cord. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_The_spinal_cord.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 11.1D: Spinal Cord White Matter is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.2: THE BRAIN

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

11.2A: The Brain

This page titled 11.2: The Brain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.2A: The Brain

The brain is the neurological center of an organism.

Learning Objectives

- Distinguish between the cerebellum, cerebral cortex, and brain stem regions of the brain

Key Points

- The brain is protected by the thick bones of the skull, suspended in cerebrospinal fluid, and isolated from the bloodstream by the blood-brain barrier.
- The brain stem consists of the midbrain, pons, and medulla.
- Each of the two hemispheres is divided into four separate lobes: the frontal (control of specialized motor control, learning, planning, and speech); parietal (control of somatic sensory functions); occipital (control of vision); and temporal lobes (control of hearing and some speech).
- The seahorse-shaped hippocampus is responsible for memory.
- The amygdala controls mood and emotions and is the center for danger identification and self-preservation.
- The cortex is divided into three functional categories: sensory, motor, and association.

Key Terms

- **hippocampus**: A part of the brain located inside the temporal lobe, consisting mainly of gray matter. It is a component of the limbic system and plays a role in memory and emotion.
- **amygdala**: Located in the medial temporal lobe, this brain region is believed to play a key role in emotions such as fear and pleasure in both animals and humans.
- **cerebrum**: The upper part of the brain, divided into the two cerebral hemispheres. In humans, it is the largest part of the brain and the seat of motor and sensory functions, as well as higher mental functions such as consciousness, thought, reason, emotion, and memory.

EXAMPLES

Lesions of the hypothalamus interfere with several vegetative functions and some behaviors, such as sexuality, combativeness, and hunger.

The human brain is the center of the human nervous system. It has the same general structure as the brains of other mammals, but is larger than expected on the basis of body size when compared to other primates. Estimates for the number of neurons (nerve cells) in the human brain range from 80 to 120 billion. Most of the expansion comes from the cerebral cortex, especially the frontal lobes, which are associated with executive functions such as self-control, planning, reasoning, and abstract thought. Despite being protected by the thick bones of the skull, suspended in cerebrospinal fluid, and isolated from the bloodstream by the blood-brain barrier, the human brain is susceptible to many types of damage and disease. This includes degenerative disorders such as Parkinson's disease, multiple sclerosis, and Alzheimer's disease. A number of psychiatric conditions, such as schizophrenia and depression, are thought to be associated with brain dysfunction, although the nature of such brain anomalies is not well understood.

Cerebral Cortex

The cerebral hemispheres form the largest part of the human brain and are situated above most other brain structures. They are covered with a cortical layer and have a convoluted topography. The cerebral cortex is essentially a sheet of neural tissue folded in a way that allows a large surface area to fit within the confines of the skull. Anatomists call each cortical fold a sulcus and the smooth area between folds a gyrus. As a rule, the smaller the cerebrum, the less convoluted the cortex. The cortex of a rat or

mouse is almost completely smooth. The cortex of a dolphin or whale, on the other hand, is more convoluted than the cortex of a human.

Cortex Divisions




Diagram of Brain Regions: Demonstration of brain regions, including the four lobes and internal structures.

The left and right hemispheres of the cerebral cortex are nearly symmetrical. The hemispheres are connected by the corpus callosum, the largest white-matter structure in the brain. Anatomists conventionally divide each hemisphere into four lobes: the frontal (control of specialized motor control, learning, planning, and speech), parietal (control of somatic sensory functions), occipital (control of vision), and temporal lobes (control of hearing and some speech). The division into lobes does not actually arise from the structure of the cortex itself. Instead, each section is named after the skull bone that covers it. The borders between lobes are placed beneath the sutures that link the skull bones together. The only exception is the border between the frontal and parietal lobes, which is shifted backward from the corresponding suture to the central sulcus. This deep fold marks the line where the primary somatosensory cortex (main sensory receptive area for the sense of touch) and primary motor cortex (one of the principal areas of the brain involved in motor function) come together. Functionally, the cortex is commonly described as comprising three parts: sensory, motor, and association areas.

Brain Stem and Cerebellum

The cerebrum is attached to a stalk-like structure called the brain stem, which consists of the midbrain, pons, and medulla. At the rear of the brain beneath the cerebrum and behind the brainstem is the cerebellum. This structure has a horizontally-furrowed surface with an appearance that is distinct from all other brain areas. These same structures are present in other mammals, although the cerebellum is not so large relative to the rest of the brain in non-human mammals.

Hippocampus

Found deep in the temporal lobe, the seahorse-shaped hippocampus is responsible for memory. The amygdala is a small, almond-shaped structure deep inside the anteroinferior region of the temporal lobe. It plays an important role in the mediation and control of activities and feelings such as love, friendship, affection, and mood expression. The amygdala is the center for danger identification, a fundamental part of self-preservation.

Thalamus

The thalamus is perched on top of the brainstem near the center of the brain, with nerve fibers projecting out to the

cerebral cortex in all directions. Its functions includes relaying sensory and motor signals to the cerebral cortex and regulating consciousness, sleep, and alertness. It likely acts as a relay between a variety of subcortical areas and the cerebral cortex.

Hypothalamus

The hypothalamus is a small part of the brain located just below the thalamus on both sides of the third ventricle. The hypothalamus also plays a role in emotion. Specifically, the lateral parts seem to be involved with pleasure and rage, while the medial part is linked to aversion, displeasure, and a tendency toward uncontrollably loud laughing. When the physical symptoms of emotion appear, the threat they pose returns to the limbic centers via the hypothalamus, then to the prefrontal nuclei, increasing anxiety.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...System%23Brain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human brain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_brain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebrum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebrum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- amygdala. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amygdala**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hippocampus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hippocampus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain Parts. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...rain_parts.jpg**. **License**: ***CC BY: Attribution***

This page titled 11.2A: The Brain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.3: PROTECTION OF THE BRAIN

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

11.3A: Blood-Brain Barrier

11.3B: Cerebrospinal Fluid and Its Circulation

11.3C: Ventricles

This page titled 11.3: Protection of the Brain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.3A: Blood-Brain Barrier

Tight junctions present in the blood-brain barrier separate circulating blood from cerebrospinal fluid, regulating diffusion into the brain.

Learning Objectives

- Describe the function of the blood-brain barrier

Key Points

- The blood – brain barrier (BBB) endothelial cells restrict the passage of substances from the bloodstream to a greater extent than endothelial cells in capillaries elsewhere in the body.
- The BBB results from the selectivity of the tight junctions between endothelial cells in central nervous system (CNS) vessels that restrict the passage of solutes.
- Several areas of the human brain are not protected by the BBB, including the circumventricular organs.
- Tight junctions are composed of transmembrane proteins such as occludin and the claudins.
- The BBB effectively protects the brain from many common bacterial infections. However, since antibodies and antibiotics are too large to cross the BBB, infections of the brain that do occur are often difficult to treat.

Key Terms

- **astrocyte**: A star-shaped neuroglial cell.
- **claudins**: This family of proteins is the most important component of tight junctions, where they establish the paracellular barrier that controls the flow of molecules in the intercellular space between the cells of an epithelium.
- **blood-brain barrier**: A structure in the CNS that keeps substances found in the bloodstream out of the brain while allowing in substances essential to metabolic function such asoxygen.
- **occludin**: A protein forming the main component of the tight junctions, along with the claudin group of proteins.

EXAMPLES

- An exception to the bacterial exclusion are the diseases caused by spirochetes, such as *Borrelia*, which causes Lyme disease, and *Treponema pallidum*, which causes syphilis. These harmful bacteria seem to breach the BBB by physically tunneling through the blood vessel walls.
- Modalities for drug delivery through the BBB entail its disruption by osmotic means, biochemically by the use of vasoactive substances, or by localized exposure to high-intensity focused ultrasound.

The blood-brain barrier (BBB) is a separation of circulating blood from the brain extracellular fluid in the central nervous system (CNS). Bacteriologist Paul Ehrlich observed that chemical dye injected into an animal would stain all of its organs except for the brain. In a later experiment, his student Edwin Goldmann found that when dye is directly injected into the cerebrospinal fluid (CSF) of animals' brains, the brains were dyed while the rest of the organs were unaffected. This clearly demonstrated the existence of some sort of compartmentalization between the brain and the rest of the body. The concept of the BBB (then termed hematoencephalic barrier) was proposed by Lewandowsky in 1900. It was not until the introduction of the scanning electron microscope that the actual membrane could be observed and proven to exist.

Blood-Brain Barrier Structure



The Blood Brain Barrier at the Small Scale: Astrocytes send projections to completely encircle capillaries. This allows tight regulation of passage of molecules into and out of the CNS.

The BBB results from the selectivity of the tight junctions between endothelial cells in CNS vessels that restrict the passage of solutes. At the interface between blood and the brain, endothelial cells are joined by these tight junctions, which are composed of smaller subunits, frequently biochemical dimers that are transmembrane proteins such as occludin, claudins, and junctional adhesion molecule. Each of these transmembrane proteins is anchored into the endothelial cells by another protein complex. This barrier also includes a thick basement membrane and astrocyte cell projections called astrocytic feet (forming the thin barrier called the glia limitans) that surround the endothelial cells of the BBB, providing biochemical support to those cells.

Function and Importance of the Blood-Brain Barrier

The BBB endothelial cells restrict the passage of substances from the bloodstream to a greater extent than endothelial cells in capillaries elsewhere in the body. The diffusion of microscopic particles (e.g., bacteria), large molecules, and hydrophilic molecules into the CSF is restricted, while the diffusion of small hydrophobic molecules (O2, CO2, hormones) is permitted. Also, BBB cells actively transport metabolic products such as glucose across the barrier.

Non-Protected Areas of the Brain

Several areas of the human brain are not protected by the BBB. These include the circumventricular organs such as the area postrema, median eminence of the hypothalamus, pineal gland, and the posterior pituitary. The posterior pituitary and pineal gland are not covered by the BBB because they secrete hormones into circulation. The median eminence is not covered by BBB because the pituitary secretions collect in this area before release into circulation. The area postrema detects noxious substances present in the blood and is therefore not covered by the BBB.

Role of Blood-Brain Barrier in Infectious Processes

The BBB effectively protects the brain from many common bacterial infections, so brain infections are very rare. However, since antibodies and antibiotics are too large to cross the BBB, brain infections that do occur are often very serious and difficult to treat. However, the BBB becomes more permeable during inflammation. This allows some antibiotics and phagocytes to move across the BBB, but also allows bacteria and virus to cross. Diseases caused by spirochetes are exceptions to this bacterial exclusion. These include Borrelia (the cause of Lyme disease), and Treponema pallidum, which causes syphilis. These harmful bacteria seem to breach the BBB by physically tunneling through the blood vessel walls. Some toxins are made up of large molecules that cannot pass through the BBB. Neurotoxins such as botulinum toxin in food might affect peripheral nerves, but the BBB can often prevent such toxins from reaching the CNS, where they could cause serious or fatal damage.

This page titled 11.3A: Blood-Brain Barrier is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.3B: Cerebrospinal Fluid and Its Circulation

Cerebrospinal fluid is a clear fluid that acts as a cushion for the brain and maintains overall central nervous system homeostasis.

Learning Objectives

- Describe the role and circulation of cerebrospinal fluid in the nervous system

Key Points

- Cerebrospinal fluid (CSF) is a clear, colorless bodily fluid that occupies the subarachnoid space and the ventricular system around and inside the brain and spinal cord.
- CSF acts as a cushion or buffer for the cortex, providing a basic mechanical and immunological protection to the brain inside the skull and serving a vital function in cerebral autoregulation of cerebral blood flow.
- CSF serves five primary purposes: buoyancy, protection, chemical stability, waste removal, and prevention of brain ischemia.
- CSF can be tested for the diagnosis of a variety of neurological diseases through the use of a procedure called lumbar puncture.
- CSF is produced in the choroid plexus in the brain by modified ependymal cells.

Key Terms

- **glymphatic system**: Functional waste clearance pathway for the vertebrate central nervous system (CNS) that consists of a para-arterial influx route for CSF to enter the brain coupled to a clearance mechanism for the removal of interstitial fluid and extracellular solutes from the interstitial compartments of the brain and spinal cord.
- **choroid plexus**: A structure in the ventricles of the brain where CSF is produced.
- **lumbar puncture**: A diagnostic and at times therapeutic procedure performed to collect a CSF for biochemical, microbiological, and cytological analysis, or rarely to relieve increased intracranial pressure.


EXAMPLES

A 2010 study showed that analysis of CSF for three protein biomarkers can indicate the presence of Alzheimer's disease. The three biomarkers are CSF amyloid beta 1-42, total CSF tau protein, and P-Tau181P. In the study, the biomarker test showed good sensitivity, identifying 90% of persons with Alzheimer's disease, but poor specificity, as 36% of control subjects were positive for the biomarkers. The researchers suggested that the low specificity may be explained by developing but not yet symptomatic disease in controls.


Cerebrospinal fluid (CSF) is a clear, colorless bodily fluid that occupies the subarachnoid space and the ventricular system around and inside the brain and spinal cord. The CSF occupies the space between the arachnoid mater (the middle layer of the brain cover, the meninges) and the pia mater (the layer of the meninges closest to the brain). It constitutes the content of all intracerebral ventricles, cisterns, and sulci (singular sulcus), as well as the central canal of the spinal cord. It acts as a cushion or buffer for the cortex, providing a basic mechanical and immunological protection for the brain inside the skull and serving a vital function in cerebral autoregulation of cerebral blood flow.

CSF Production

Between 50 to 70% of CSF is produced in the brain by modified ependymal cells in the choroid plexus, and the remainder is formed around blood vessels and along ventricular walls. It circulates from the lateral ventricles to the foramen of Monro (interventricular foramen), third ventricle, aqueduct of Sylvius (cerebral aqueduct), fourth ventricle, foramen of Magendie (median aperture), foramen of Luschka (lateral apertures), and the subarachnoid space over the brain and the spinal cord. CSF is reabsorbed into venous sinus blood via arachnoid granulations.



Subarachnoid Cavity: Diagrammatic representation of a section across the top of the skull, showing the membranes of the brain with the subarachnoid cavity visible on the left.



The Choroid Plexus: This diagram indicates the (1) posterior medullary velum (2) choroid plexus (3) cisterna cerebellomedullaris of subarachnoid cavity (4) central canal (5) corpora quadrigemina (6) cerebral peduncle (7) anterior medullary velum (8) ependymal lining of ventricle (9) cisterna pontis of subarachnoid cavity

CSF is produced at a rate of 500 ml/day. Since the subarachnoid space around the brain and spinal cord can contain only 135 to 150 ml, large amounts are drained into the blood through arachnoid granulations in the superior sagittal sinus. Thus, the CSF turns over about 3.7 times a day. This continuous flow into the venous system dilutes the concentration of larger, lipid-insoluble molecules penetrating the brain and CSF. The CSF contains approximately 0.3% plasma proteins, or approximately 15 to 40 mg/dL, depending on the sampling site.

Functions of CSF

The functions of CSF include:

1. Buoyancy: The actual mass of the human brain is about 1400 grams; however, the net weight of the brain suspended in the

CSF is equivalent to a mass of 25 grams. The brain therefore exists in neutral buoyancy, which allows the brain to maintain its density without being impaired by its own weight.
2. Protection: CSF protects the brain tissue from injury when jolted or hit.
3. Chemical stability: CSF flows throughout the inner ventricular system in the brain and is absorbed back into the bloodstream, rinsing the metabolic waste from the central nervous system (CNS) through the blood-brain barrier. This allows for homeostatic regulation of the distribution of neuroendocrine factors, to which slight changes can cause problems or damage to the nervous system.
4. Prevention of brain ischemia: Decreasing the amount of CSF in the limited space inside the skull decreases total intracranial pressure and facilitates blood perfusion.
5. Clearing waste: CSF has been shown by the research group of Maiken Nedergaard to be critical in the brain's glymphatic system, the functional waste clearance pathway for the vertebrate CNS.

CSF as a Diagnostic Tool

When CSF pressure is elevated, cerebral blood flow may be constricted. When disorders of CSF flow occur, they may therefore affect not only CSF movement but also craniospinal compliance and intracranial blood flow, with subsequent neuronal and glial vulnerabilities. The venous system is also important in this equation. CSF can be tested for the diagnosis of various neurological diseases, usually with a procedure called lumbar puncture. Lumbar puncture is performed in an attempt to count the cells in the fluid and to detect the levels of protein and glucose. These parameters alone may be extremely beneficial in the diagnosis of subarachnoid hemorrhage and CNS infections such as meningitis. Moreover, a CSF culture examination may yield the microorganism that has caused the infection.

This page titled 11.3B: Cerebrospinal Fluid and Its Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.3C: Ventricles

The ventricular system is a set of hollow cavities in the brain filled with cerebrospinal fluid.

Learning Objectives

- Describe the function and locations of the ventricles in the brain

Key Points

- The ventricular system is continuous with the central canal of the spinal cord.
- The ventricles are filled with cerebrospinal fluid (CSF), which bathes and cushions the brain and spinal cord within their bony confines.
- CSF flows from the lateral ventricles via the foramina of Monro into the third ventricle.
- CSF flows from the third ventricle to the fourth ventricle via the cerebral aqueduct in the brainstem.
- CSF flows from the fourth ventricle into the central canal of the spinal cord or into the cisterns of the subarachnoid space via three small foramina: the central foramen of Magendie and the two lateral foramina of Luschka.
- The cerebral aqueduct and the foramina are very small and easily blocked, which would cause high pressure in the lateral ventricles and hydrocephalus.

Key Terms

- **cerebrospinal fluid (CSF)**: A clear, colorless bodily fluid produced in the choroid plexus of the brain that acts as a cushion or buffer for the cortex, providing a basic mechanical and immunological protection to the brain inside the skull.
- **cerebral aqueduct**: The channel in the brain which connects the third ventricle to the fourth ventricle. Also called the aqueduct of sylvius. It is surrounded by the periaqueductal gray.
- **lateral foramina of Luschka**: Also known as the lateral aperture, an opening in each lateral extremity of the fourth ventricle of the human brain that provides a conduit for cerebrospinal fluid to flow from the brain's ventricular system into the subarachnoid space.

EXAMPLES

In the late 1970s, CT scans of the ventricles revolutionized the study of mental disorders. Researchers found that on average, individuals with schizophrenia had enlarged ventricles compared to healthy subjects. This became the first evidence that mental disorders are biological in origin and led to a reinvigoration of the study of such conditions via modern scientific techniques.

The ventricular system is a set of four interconnected cavities (ventricles) in the brain and the location of CSF production. This system is continuous with the central canal of the spinal cord. The system comprises four ventricles:


- right and left lateral ventricles (the first and second ventricles)
- third ventricle
- fourth ventricle

CSF Production

The cavities of the cerebral hemispheres are called lateral ventricles or first and second ventricles. These two ventricles open into the third ventricle by a common opening called the foramen of Monro. CSF is produced by modified ependymal cells of the choroid plexus found in all components of the ventricular system except for the cerebral aqueduct and the posterior and anterior horns of the lateral ventricles. The brain and spinal cord are covered by a series of tough membranes called meninges, which protect these organs from rubbing against the bones of the skull and spine. The CSF within the skull and spine is found between the pia mater and the arachnoid and provides further cushioning. EndFra

CSF Flow Within Ventricles

CSF flows from the lateral ventricles via the foramina of Monro into the third ventricle, and then into the fourth ventricle via the cerebral aqueduct in the brainstem. From there, it passes into the central canal of the spinal cord and into the cisterns of the subarachnoid space via three small foramina: the central foramen of Magendie and the two lateral foramina of Luschka. The fluid then flows around the superior sagittal sinus to be reabsorbed via the arachnoid villi into the venous system. CSF within the spinal cord can flow all the way down to the lumbar cistern at the end of the cord around the cauda equina.



Brain Ventricles: Lateral and anterior views of the brain ventricles, including the third and fourth ventricle, lateral ventricles, interventricular foramen, cerebral aqueduct, and central canal.

Ventricular System Dysfunction and Disease

Diseases of the ventricular system include abnormal enlargement (hydrocephalus) and inflammation of the CSF spaces (meningitis, ventriculitis) caused by infection or introduction of blood following trauma or hemorrhage. The aqueduct between the third and fourth ventricles is very small, as are the foramina. This means they can be easily blocked, causing high pressure in the lateral ventricles. This is a common cause of hydrocephalus (known colloquially as “water on the brain”), an extremely serious condition due to the damage caused by the pressure as well as the nature of the block (e.g., a tumor or inflammatory swelling).

Embryonic Development of the Ventricles

The structures of the ventricular system are embryologically derived from the center of the neural tube (the neural canal). As the future brain stem aspect of the primitive neural tube develops, the neural canal expands dorsally and laterally, creating the fourth ventricle. The cerebral aqueduct is formed from the part of the neural canal that does not expand and remains the same at the level of the midbrain superior to the fourth ventricle. The fourth ventricle narrows at the obex, where the fourth ventricle narrows to become the central canal in the caudal medulla.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Circumventricular Organs. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circum...ricular_organs**. **License**:

CC BY-SA: Attribution-ShareAlike

- Bloodu2013brain barrier. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%E...3brain_barrier**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- astrocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/astrocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- claudins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/claudins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- occludin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/occludin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- blood-brain barrier. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/blood-brain_barrier**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood Brain Barriere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bl...n_Barriere.jpg**. **License**: ***CC BY: Attribution***
- lumbar puncture. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lumbar_puncture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glymphatic System. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Glymphatic_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebrospinal fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebrospinal_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Choroid plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Choroid_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebrospinal fluid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebrospinal_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood Brain Barriere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bl...n_Barriere.jpg**. **License**: ***CC BY: Attribution***
- Gray708. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray708.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray769-en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray769-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ventricular system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ventricular_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebrospinal fluid (CSF). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cerebro...0fluid%20(CSF)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lateral foramina of Luschka. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lateral...20of%20Luschka**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebral aqueduct. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebral_aqueduct**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood Brain Barriere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bl...n_Barriere.jpg**. **License**: ***CC BY: Attribution***
- Gray708. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray708.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray769-en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray769-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain ventricles. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...cles_Brain.png**. **License**: ***CC BY: Attribution***

This page titled 11.3C: Ventricles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4: THE BRAIN STEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

11.4A: Functions of the Brain Stem

11.4B: Medulla Oblongata

11.4C: Pons

11.4D: Midbrain

11.4E: Reticular Formation

This page titled 11.4: The Brain Stem is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4A: Functions of the Brain Stem

The brainstem regulates vital cardiac and respiratory functions and acts as a vehicle for sensory information.

Learning Objectives

- Describe the functions of the brainstem

Key Points

- In vertebrate anatomy, the brainstem is the posterior part of the brain adjoining, and structurally continuous with, the spinal cord.
- Though small, the brainstem is an extremely important part of the brain, as the nerve connections from the motor and sensory systems of the cortex pass through it to communicate with the peripheral nervous system.
- The brainstem also plays an important role in the regulation of cardiac and respiratory function, consciousness, and the sleep cycle.
- The brainstem consists of the medulla oblongata, pons, and midbrain.

Key Terms

- **pons**: Contains nuclei that relay signals from the forebrain to the cerebellum, along with nuclei that deal primarily with sleep, respiration, swallowing, bladder control, hearing, equilibrium, taste, eye movement, facial expressions, facial sensation, and posture.
- **midbrain**: Associated with vision, hearing, motor control, sleep and wake cycles, alertness, and temperature regulation.
- **medulla**: The lower half of the brainstem that contains the cardiac, respiratory, vomiting, and vasomotor centers and regulates autonomic, involuntary functions such as breathing, heart rate, and blood pressure.

EXAMPLES


Diseases of the brainstem can result in abnormalities in cranial nerve function, leading to visual and hearing disturbances, changes in sensation, muscle weakness, vertigo, coordination problems, swallowing and speech difficulty, and voice changes.

Location and Basic Physiology

In vertebrate anatomy, the brainstem is the most inferior portion of the brain, adjoining and structurally continuous with the brain and spinal cord. The brainstem gives rise to cranial nerves 3 through 12 and provides the main motor and sensory innervation to the face and neck via the cranial nerves. Though small, it is an extremely important part of the brain, as the nerve connections of the motor and sensory systems from the main part of the brain that communicate with the peripheral nervous system pass through the brainstem. This includes the corticospinal tract (motor), the posterior column-medial lemniscus pathway (fine touch, vibration sensation, and proprioception) and the spinothalamic tract (pain, temperature, itch, and crude touch). The brain stem also plays an important role in the regulation of cardiac and respiratory function. It regulates the central nervous system (CNS) and is pivotal in maintaining consciousness and regulating the sleep cycle.

Components of the Brainstem

The three components of the brainstem are the medulla oblongata, midbrain, and pons.



Brainstem Anatomy: Structures of the brainstem are depicted on these diagrams, including the midbrain, pons, medulla, basilar artery, and vertebral arteries.

The medulla oblongata (myelencephalon) is the lower half of the brainstem continuous with the spinal cord. Its upper part is continuous with the pons. The medulla contains the cardiac, respiratory, vomiting, and vasomotor centers regulating heart rate, breathing, and blood pressure.


The midbrain (mesencephalon) is associated with vision, hearing, motor control, sleep and wake cycles, alertness, and temperature regulation.

The pons (part of metencephalon) lies between the medulla oblongata and the midbrain. It contains tracts that carry signals from the cerebrum to the medulla and to the cerebellum. It also has tracts that carry sensory signals to the thalamus.

Brainstem Function

The brainstem has many basic functions, including regulation of heart rate, breathing, sleeping, and eating. It also plays a role in conduction. All information relayed from the body to the cerebrum and cerebellum and vice versa must traverse the brainstem. The ascending pathways from the body to the brain are the sensory pathways, including the spinothalamic tract for pain and temperature sensation and the dorsal column, fasciculus gracilis, and cuneatus for touch, proprioception, and pressure sensation. The facial sensations have similar pathways and also travel in the spinothalamic tract and the medial lemniscus.

Descending tracts are upper motor neurons destined to synapse on lower motor neurons in the ventral horn and intermediate horn of the spinal cord. In addition, upper motor neurons originate in the brain stem's vestibular, red, tectal, and reticular nuclei, which also descend and synapse in the spinal cord. The brainstem also has integrative functions, including cardiovascular system control, respiratory control, pain sensitivity control, alertness, awareness, and consciousness.



Human Brain with Cranial Nerves: Cranial nerves are nerves that emerge directly from the brain, in contrast to spinal nerves, which emerge from segments of the spinal cord. In humans, there are traditionally twelve pairs of cranial nerves. Only the first and the second pair emerge from the cerebrum; the remaining ten pairs emerge from the brainstem.

This page titled 11.4A: Functions of the Brain Stem is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4B: Medulla Oblongata

The medulla oblongata controls autonomic functions and connects the higher levels of the brain to the spinal cord.

Learning Objectives

- Describe the location and function of the medulla oblongata region of the brain stem

Key Points

- The medulla oblongata is the lower half of the brainstem. It controls autonomic functions and connects the higher levels of the brain to the spinal cord.
- The medulla oblongata is responsible for regulating several basic functions of the autonomic nervous system, including respiration, cardiac function, vasodilation, and reflexes like vomiting, coughing, sneezing, and swallowing.


Key Terms

- **tuberculum cinereum**: A raised area between the rootlets of the accessory nerve and posterolateral sulcus that overlies the spinal tract of trigeminal nerve.
- **cerebellar peduncle**: The structure that connects the medulla to the cerebellum.
- **sympathetic system**: The division of the autonomic nervous system responsible for stimulating the body's fight-or-flight response.
- **olivary body**: Either of a pair of prominent oval structures in the medulla oblongata containing the olivary nuclei. These structures are involved in cerebellar motor learning and the perception of sound.
- **parasympathetic system**: The division of the autonomic nervous system responsible for the relaxation or inhibition of various body functions.

EXAMPLES

A stroke can injure the pyramidal tract, medial lemniscus, and the hypoglossal nucleus. This causes a syndrome called medial medullary syndrome, a type of alternating hemiplegia characterized by recurrent episodes of paralysis on one side of the body.

The medulla oblongata is the lower half of the brainstem. In discussions of neurology and similar contexts where no ambiguity will result, it is often referred to as simply the medulla. The medulla contains the cardiac, respiratory, vomiting, and vasomotor centers and regulates autonomic, involuntary functions such as breathing, heart rate, and blood pressure.



The Brain Stem with Pituitary and Pineal Glands: Medulla oblongata labeled at bottom left, in relation to the pons, pituitary gland, spinal cord, pineal gland and cerebellum.

The medulla is often divided into two parts:

1. An open or superior part where the dorsal surface of the medulla is formed by the fourth ventricle.
2. A closed or inferior part where the metacoel (caudal part of fourth ventricle) lies within the medulla oblongata.

Structure of the Medulla Oblongata

The region between the anterior median and anterolateral sulci is occupied by an elevation on either side known as the pyramid of medulla oblongata. This elevation is caused by the corticospinal tract. In the lower part of the medulla, some of these fibers cross each other, thus obliterating the anterior median fissure. This is known as the decussation of the pyramids. Other fibers that originate from the anterior median fissure above the decussation of the pyramids and run laterally across the surface of the pons are known as the external arcuate fibers.

The region between the anterolateral and posterolateral sulcus in the upper part of the medulla is marked by a swelling known as the olivary body, caused by a large mass of gray matter known as the inferior olivary nucleus.

The posterior part of the medulla between the posterior median and posterolateral sulci contains tracts that enter it from the posterior funiculus of the spinal cord. These are the fasciculus gracilis, lying medially next to the midline, and the fasciculus cuneatus, lying laterally.

The fasciculi end in rounded elevations known as the gracile and cuneate tubercles. They are caused by masses of gray matter known as the nucleus gracilis and the nucleus cuneatus. Just above the tubercles, the posterior aspect of the medulla is occupied by a triangular fossa, which forms the lower part of the floor of the fourth ventricle. The fossa is bounded on either side by the inferior cerebellar peduncle, which connects the medulla to the cerebellum.

The lower part of the medulla, immediately lateral to the fasciculus cuneatus, is marked by another longitudinal elevation known as the tuberculum cinereum. It is caused by an underlying collection of gray matter known as the spinal nucleus of the trigeminal nerve. The gray matter of this nucleus is covered by a layer of nerve fibers that form the spinal tract of the trigeminal nerve.

The base of the medulla is defined by the commissural fibers, crossing over from the ipsilateral side in the spinal cord to the contralateral side in the brain stem; below this is the spinal cord.

Embryonic Development

During development, the medulla oblongata forms from the myelencephalon. The final neuroblasts from the alar plate of the neural tube produce the sensory nuclei of the medulla. The basal plate neuroblasts give rise to the motor nuclei.

Function of the Medulla Oblongata

The medulla oblongata controls autonomic functions and connects the higher levels of the brain to the spinal cord. It is also responsible for regulating several basic functions of the autonomic nervous system, including:

- **Respiration:** chemoreceptors
- **Cardiac center:** sympathetic system, parasympathetic system
- **Vasomotor center:** baroreceptors
- Reflex centers of vomiting, coughing, sneezing, and swallowing

This page titled 11.4B: Medulla Oblongata is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4C: Pons

The pons is a relay station between the forebrain and cerebellum that passes sensory information from the periphery to the thalamus.

Learning Objectives


- Describe the role and location of the pons region of the brainstem

Key Points

- The pons is a structure located on the brainstem, named after the Latin word for “bridge.”
- This white matter includes tracts that conduct signals from the cerebrum down to the cerebellum and medulla, as well as tracts that carry the sensory signals up into the thalamus.
- The pons contains nuclei that relay signals from the forebrain to the cerebellum, along with nuclei that deal primarily with sleep, respiration, swallowing, bladder control, hearing, equilibrium, taste, eye movement, facial expressions, facial sensation, and posture.
- Within the pons is the pneumotaxic center, a nucleus that regulates the change from inspiration to expiration.
- The pons also contains the sleep paralysis center of the brain and plays a role in generating dreams.
- The functions of these four nerves include sensory roles in hearing, equilibrium, taste, and in facial sensations such as touch and pain. They also have motor roles in eye movement, facial expressions, chewing, swallowing, urination, and the secretion of saliva and tears.

Key Terms

- **pons**: Contains nuclei that relay signals from the forebrain to the cerebellum, along with nuclei that regulate sleep, respiration, swallowing, bladder control, hearing, equilibrium, taste, eye movement, facial expressions, facial sensation, and posture.
- **pneumotaxic center**: A network of neurons in the rostral dorsal lateral pons that regulates the respiratory rate; also known as the pontine respiratory group (PRG).
- **Basal plate**: The region of the neural tube ventral to the sulcus limitans and containing primarily motor neurons.
- **alar plate**: Also called the alar lamina, it is a neural structure in the embryonic nervous system; the caudal part later becomes the sensory axon aspect of the spinal cord.



Pons/Brainstem: Structure of the brainstem showing the location of the pons in relation to the midbrain and medulla.

The pons is a structure located on the brainstem, named after the Latin word for “bridge.” It is above the medulla, below the midbrain, and anterior to the cerebellum. The white matter of the pons includes tracts that conduct signals from the cerebrum down to the cerebellum and medulla, and tracts that carry the sensory signals up into the thalamus.

Structure

The pons measures about 2.5 cm in length in adults. Most of it appears as a broad anterior bulge rostral to the medulla. Posteriorly, it consists mainly of two pairs of thick stalks called cerebellar peduncles. These connect the cerebellum to the pons and midbrain.

The pons contains nuclei that relay signals from the forebrain to the cerebellum, along with nuclei that regulate sleep, respiration, swallowing, bladder control, hearing, equilibrium, taste, eye movement, facial expressions, facial sensation, and posture. Within the pons is the pneumotaxic center, a nucleus that regulates the change from inspiration to expiration. The pons also contains the sleep paralysis center of the brain and also plays a role in generating dreams.

Development

During embryonic development, the metencephalon develops from the rhombencephalon and gives rise to two structures: the pons and the cerebellum. The alar plate produces sensory neuroblasts, which will give rise to the solitary nucleus and its special visceral afferent column, the cochlear and vestibular nuclei (which form the special somatic afferent fibers of the vestibulocochlear nerve), the spinal and principal trigeminal nerve nuclei (which form the general somatic afferent column of the trigeminal nerve), and the pontine nuclei, which is involved in motor activity. Basal plate neuroblasts give rise to the abducens nucleus (forms the general somatic efferent fibers), the facial and motor trigeminal nuclei (form the special visceral efferent column), and the superior salivatory nucleus, which forms the general visceral efferent fibers of the facial nerve.

Cranial Nerves of the Pons

A number of cranial nerve nuclei are present in the pons:

- The chief or pontine nucleus of the trigeminal nerve sensory nucleus (V)- mid-pons
- The motor nucleus for the trigeminal nerve (V)-mid-pons
- Abducens nucleus (VI)-lower pons
- Facial nerve nucleus (VII)-lower pons
- Vestibulocochlear nuclei (VIII)-lower pons

Functional Characteristics

The functions of the four nerves of the pons include sensory roles in hearing, equilibrium, taste, and facial sensations such as touch and pain. They also have motor roles in eye movement, facial expressions, chewing, swallowing, urination, and the secretion of saliva and tears. Central pontine myelinosis is a demyelination disease that causes difficulty with sense of balance, walking, sense of touch, swallowing, and speaking. If it is not diagnosed and treated, it can lead to death or locked-in syndrome (a condition in which a person is conscious but cannot move or communicate).

This page titled 11.4C: Pons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4D: Midbrain

The midbrain plays a major role in both wakefulness and regulation of homeostasis.

Learning Objectives

- Describe the location and functions of the midbrain

Key Points


- The midbrain or mesencephalon is a portion of the central nervous system (CNS) associated with vision, hearing, motor control, sleep and wake cycles, arousal (alertness), and temperature regulation.
- Anatomically, the midbrain comprises the tectum (or corpora quadrigemina), tegmentum, ventricular mesocoelia (or “iter”), and cerebral peduncles, as well as several nuclei and fasciculi.
- During embryonic development, the midbrain arises from the second vesicle, (mesencephalon) of the neural tube.
- The mesencephalon is considered part of the brainstem.

Key Terms

- **mesencephalon**: A part of the brain located rostral to the pons and caudal to the thalamus and the basal ganglia, composed of the tectum (dorsal portion) and the tegmentum (ventral portion).
- **substantia nigra**: Brain structure located in the midbrain that plays an important role in reward and movement.
- **tectum**: The dorsal part of the midbrain, responsible for auditory and visual reflexes.
- **tegmentum**: The ventral portion of the midbrain, a multisynaptic network of neurons involved in many unconscious homeostatic and reflexive pathways.

The midbrain or mesencephalon (from the Greek *mesos,* middle, and *enkephalos,* brain) is a portion of the central nervous system (CNS) associated with vision, hearing, motor control, sleep and wake cycles, arousal (alertness), and temperature regulation. Anatomically, it comprises the tectum (or corpora quadrigemina), tegmentum, ventricular mesocoelia (or “iter”), and the cerebral peduncles, as well as several nuclei and fasciculi. Caudally (posteriorly) the mesencephalon adjoins the pons (metencephalon), and rostrally it adjoins the diencephalon (eg., thalamus, hypothalamus). The midbrain is located below the cerebral cortex and above the hindbrain placing it near the center of the brain.

Primary Midbrain Components



Brainstem Anatomy: Brainstem anatomy showing the location of the midbrain in relation to the midbrain, pons, medulla, basilar artery, and vertebral arteries.

The tectum (Latin for “roof”) is formed by the superior and inferior colliculi and comprises the rear portion of the midbrain. The superior colliculus regulates preliminary visual processing and eye movement, while the inferior colliculus is involved in auditory processing. Collectively, the colliculi is referred to as the corpora quadrigemina.

The tegmentum is involved in many unconscious homeostatic and reflexive pathways, and is the motor center that relays inhibitory signals to the thalamus and basal nuclei to prevent unwanted body movement. It extends from the substantia nigra to the cerebral aqueduct (also called the ventricular mesocoeli). The nuclei of cranial nerves III and IV are located in the tegmentum portion of the midbrain.

The substantia nigra is closely associated with motor system pathways of the basal ganglia. The human mesencephalon is archipallian in origin, sharing its general architecture with the most ancient of vertebrates. Dopamine produced in the substantia nigra plays a role in motivation and habituation of species from humans to the most elementary animals such as insects. The midbrain is the smallest region in the brain and helps to relay information for vision and hearing.

The cerebral peduncles are located on either side of the midbrain and are its most anterior part, acting as the connectors between the rest of the midbrain and the thalamic nuclei. The cerebral peduncles assist in motor movement refinement, motor skill learning, and converting proprioceptive information into balance and posture maintenance.

Embryonic Development

During embryonic development, the midbrain arises from the second vesicle, also known as the mesencephalon, of the neural tube. Unlike the other two vesicles (the prosencephalon and rhombencephalon), the mesencephalon remains undivided for the remainder of neural development. It does not split into other brain areas while the prosencephalon, for example, divides into the telencephalon and the diencephalon. Throughout embryonic development, the cells within the midbrain continually multiply and compress the still-forming aqueduct of sylvius or cerebral aqueduct. Partial or total obstruction of the cerebral aqueduct during development can lead to congenital hydrocephalus.

This page titled 11.4D: Midbrain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.4E: Reticular Formation

The reticular formation assists in regulation of the sleep cycle and detecting sensory salience.

Learning Objectives

- Describe the functions of the reticular formation region of the pons

Key Points

- The reticular formation is a region in the pons involved in regulating the sleep-wake cycle and filtering incoming stimuli to discriminate irrelevant background stimuli.
- The reticular formation consists of more than 100 small neural networks with varied functions including motor control, cardiovascular control, pain modulation, sleep, and habituation.
- Bilateral damage to the reticular formation of the midbrain may lead to coma or death.
- Traditionally, the nuclei of the reticular formation are divided into three columns: the median column or the Raphe nuclei, the medial column or the magnocellular nuclei, and the lateral column or parvocellular nuclei.

Key Terms

- **magnocellular nuclei**: Nuclei within the reticular formation involved in motor coordination.
- **parvocellular nuclei**: Nuclei within the reticular formation that are involved in the regulation of expiration during breathing and other motor functions.
- **raphe nuclei**: Located in the pons of the brainstem, the principal site of the synthesis of the neurotransmitter serotonin. Serotonin plays an important role in mood regulation, particularly when stress is associated with depression and anxiety.

The reticular formation is a region in the pons involved in regulating the sleep-wake cycle and filtering incoming stimuli to discriminate irrelevant background stimuli. It is essential for governing some of the basic functions of higher organisms, and is one of the phylogenetically oldest portions of the brain.

Divisions of the Reticular Formation

Traditionally, the nuclei are divided into three columns:


1. Raphe nuclei (medium column)
2. Magnocellular red nucleus (medial zone)
3. Parvocellular reticular nucleus (lateral zone)

Sagittal division reveals more morphological distinctions. The raphe nuclei form a ridge in the middle of the reticular formation, and directly to its periphery, there is a division called the medial reticular formation. The medial reticular formation is large, has long ascending and descending fibers, and is surrounded by the lateral reticular formation. The lateral reticular formation is close to the motor nuclei of the cranial nerves and mostly mediates their function. The raphe nuclei is the place of synthesis of the neurotransmitter serotonin, which plays an important role in mood regulation.

The medial reticular formation and lateral reticular formation are two columns of neuronal nuclei with ill-defined boundaries that send projections through the medulla and into the mesencephalon (midbrain). The nuclei can be differentiated by function, cell type, and projections of efferent or afferent nerves. The magnocellular red nucleus is involved in motor coordination, and the parvocellular nucleus regulates exhalation.

The original functional differentiation was a division of caudal and rostral, based on the observation that damage to the rostral reticular formation induces a hypersomnia in the cat brain. In contrast, damage to the more caudal portion of the reticular formation produces insomnia in cats. This study led to the idea that the caudal portion inhibits the rostral portion of the reticular

formation.



Cross Section of the Pons: A cross section of the lower part of the pons showing the pontine reticular formation labeled as #9.

Functions

The reticular formation consists of more than 100 small neural networks, with varied functions including:

1. **Somatic motor control:** Some motor neurons send their axons to the reticular formation nuclei, giving rise to the reticulospinal tracts of the spinal cord. These tracts play a large role in maintaining tone, balance, and posture, especially during movement. The reticular formation also relays eye and ear signals to the cerebellum so that visual, auditory, and vestibular stimuli can be integrated in motor coordination. Other motor nuclei include gaze centers, which enable the eyes to track and fixate objects, and central pattern generators, which produce rhythmic signals to the muscles of breathing and swallowing.
2. **Cardiovascular control:** The reticular formation includes the cardiac and vasomotor centers of the medulla oblongata.
3. **Pain modulation:** The reticular formation is one means by which pain signals from the lower body reach the cerebral cortex. It is also the origin of the descending analgesic pathways. The nerve fibers in these pathways act in the spinal cord to block the transmission of some pain signals to the brain.
4. **Sleep and consciousness:** The reticular formation has projections to the thalamus and cerebral cortex that allow it to exert some control over which sensory signals reach the cerebrum and come to our conscious attention. It plays a central role in states of consciousness like alertness and sleep. Injury to the reticular formation can result in irreversible coma.
5. **Habituation:** This is a process in which the brain learns to ignore repetitive, meaningless stimuli while remaining sensitive to others. A good example of this is when a person can sleep through loud traffic in a large city, but is awakened promptly by the sound of an alarm or crying baby. Reticular formation nuclei that modulate activity of the cerebral cortex are part of the reticular activating system.

Effects of Damage

Mass lesions in the brainstem cause severe alterations in the level of consciousness (such as coma) because of their effects on the reticular formation. Lesions in the reticular formation have been found in the brains of people who have post-polio syndrome. Some imaging studies have shown abnormal activity in this area in people with chronic fatigue syndrome, indicating a high likelihood that damage to the reticular formation is responsible for the fatigue associated with these syndromes.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- medulla. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/medulla**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pons. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pons**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en-2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br...abels_en-2.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Alternating Hemiplegia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alternating_hemiplegia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medulla oblongata. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Medulla_oblongata**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tuberculum cinereum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tuberculum%20cinereum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebellar peduncle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cerebellar%20peduncle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- olivary body. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/olivary_body**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en-2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br...abels_en-2.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pituitary pineal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...eal_glands.jpg**. **License**: ***Public Domain: No Known Copyright***
- Basal Plate. **Provided by**: Wipedia. **Located at**: **en.Wikipedia.org/wiki/Basal_..._(neural_tube)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Alar Plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alar_plate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pons. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pons**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pneumotaxic center. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pneumotaxic%20center**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pons. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pons**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en-2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br...abels_en-2.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pituitary pineal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...eal_glands.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:1311 Brain Stem.jpg. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:1...Brain_Stem.jpg**. **License**: ***CC BY: Attribution***
- Cerebral peduncle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebral_peduncle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Midbrain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Midbrain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mesencephalon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mesencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tegmentum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tegmentum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tectum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tectum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en-2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/**

File:Br...abels_en-2.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pituitary pineal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pituitary_pineal_glands.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:1311 Brain Stem.jpg. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:1311_Brain_Stem.jpg**. **License**: ***CC BY: Attribution***
- File:Blausen 0114 BrainstemAnatomy.png. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:B...temAnatomy.png**. **License**: ***CC BY: Attribution***
- Reticular formation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Reticular_formation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parvocellular nuclei. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/parvocellular%20nuclei**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- magnocellular nuclei. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/magnocellular%20nuclei**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- raphe nuclei. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/raphe%20nuclei**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainstem. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brainstem**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en-2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en-2.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pituitary pineal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pituitary_pineal_glands.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:1311 Brain Stem.jpg. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:1311_Brain_Stem.jpg**. **License**: ***CC BY: Attribution***
- File:Blausen 0114 BrainstemAnatomy.png. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Blausen_0114_BrainstemAnatomy.png**. **License**: ***CC BY: Attribution***
- Lower pons horizontal KB. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lower_pons_horizontal_KB.svg**. **License**: ***CC BY: Attribution***

This page titled 11.4E: Reticular Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.5: THE CEREBELLUM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

11.5A: Parts of the Cerebellum

11.5B: Functions of the Cerebellum

This page titled 11.5: The Cerebellum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.5A: Parts of the Cerebellum

The cerebellum, which looks like a separate structure attached to the bottom of the brain, plays an important role in motor control.

Learning Objectives

- Describe the parts of the cerebellum


Key Points

- The cerebellum can be separated into three lobes: the flocculonodular lobe, anterior lobe, and posterior lobe.
- The medial zone of the anterior and posterior lobes constitutes the spinocerebellum, or paleocerebellum.
- There are about 3.6 times as many neurons in the cerebellum as in the neocortex.
- Based on surface appearance, three lobes can be distinguished in the cerebellum: the flocculonodular lobe, anterior lobe (above the primary fissure), and the posterior lobe (below the primary fissure).

Key Terms


- **granule cells**: An extremely small type of neuron that is the the smallest cell found in the brain.
- **neocortex**: The largest part of the cerebral cortex of the human brain, covering the two cerebral hemispheres.
- **Purkinje cells**: A class of GABAergic neurons located in the cerebellar cortex that are some of the largest neurons in the human brain.
  GABA is the chief inhibitory neurotransmitter in the mammalian central nervous system.
- **cerebellum**: Part of the hindbrain. In humans, it lies between the brainstem and the cerebrum. It plays an important role in sensory perception, motor output, balance, and posture.

The cerebellum has the appearance of a separate structure attached to the bottom of the brain, tucked underneath the cerebral hemispheres. The surface of the cerebellum is covered with finely spaced parallel grooves, in striking contrast to the broad irregular convolutions of the cerebral cortex.



Cerebellum: Brain section showing cerebellum position and structure

These parallel grooves conceal the fact that the cerebellum is actually a continuous thin layer of tissue (the cerebellar cortex), tightly folded in the style of an accordion. This thin layer contains several types of neurons with a highly regular arrangement, most importantly Purkinje cells and granule cells. This complex neural network gives rise to a massive signal-processing capability, but almost all of its output is directed to a set of small deep cerebellar nuclei lying in the interior of the cerebellum.




Cells of the Cerebellum: Transverse section of a cerebellar folium, showing principal cell types and connections.

The cerebellum is separated from the overlying cerebrum by a layer of leathery dura mater. Anatomists classify the cerebellum as part of the metencephalon, which also includes the pons, and all its connections with other parts of the brain travel through the pons. The metencephalon is the upper part of the rhombencephalon, or hindbrain. Like the cerebral cortex, the cerebellum is divided into two hemispheres. It also contains a narrow midline zone called the vermis. A set of large folds is, by convention, used to divide the overall structure into 10 smaller lobules. Due to its large number of tiny granule cells, the cerebellum contains more neurons than the rest of the brain put together, but comprises only 10% of total brain volume.

Based on surface appearance, three lobes can be distinguished in the cerebellum: the flocculonodular lobe, anterior lobe (above the primary fissure), and the posterior lobe (below the primary fissure). Excluding the flocculonodular lobe, which has distinct connections and functions, the cerebellum can be parsed functionally into a medial sector called the spinocerebellum and a larger

lateral sector called the cerebrocerebellum. A narrow strip of protruding tissue along the midline is called the vermis (Latin for "worm").



Divisions of the cerebellum: Schematic representation of the major anatomical subdivisions of the cerebellum. Superior view of an "unrolled" cerebellum, placing the vermis in one plane.

The smallest region, the flocculonodular lobe, is often called the vestibulocerebellum. It is the oldest part of the brain in evolutionary terms (archicerebellum) and participates mainly in balance and spatial orientation. Its primary connections are with the vestibular nuclei, although it also receives visual and other sensory input.

The medial zone of the anterior and posterior lobes constitutes the spinocerebellum, also known as the paleocerebellum. It receives proprioception input from the dorsal columns of the spinal cord (including the spinocerebellar tract) and from the trigeminal nerve, as well as from visual and auditory systems. It sends fibers to deep cerebellar nuclei that in turn project to both the cerebral cortex and the brain stem, thus providing modulation of descending motor systems.

The lateral zone, which in humans is by far the largest part, constitutes the cerebrocerebellum, also known as the neocerebellum. It receives input exclusively from the cerebral cortex (especially the parietal lobe) via the pontine nuclei (forming corticopontocerebellar pathways), and sends output mainly to the ventrolateral thalamus (in turn connected to motor areas of the premotor cortex and primary motor area of the cerebral cortex) and to the red nucleus.

This page titled 11.5A: Parts of the Cerebellum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.5B: Functions of the Cerebellum

Cerebellar function was once believed to be motor-specific, but newer findings suggest the cerebellum is also involved in higher-level brain processing.

Learning Objectives

- Describe the function of the cerebellum

Key Points

- The cerebellum is essential for making fine adjustments to motor actions.
  Cerebellar dysfunction primarily results in problems with motor control.
- Four principles are important to cerebellar processing: feedforward processing, divergence and convergence, modularity, and plasticity.
- Signal processing in the cerebellum is almost entirely feedforward. Signals move through the system from input to output with very little internal transmission.
- The cerebellum both receives input and transmits output via a limited number of cells.
- The cerebellar system is divided into thousands of independent modules with similar structure.

Key Terms

- **neural divergence**: When a neuron fires and the signal is sent to many other neurons.
- **neuroplasticity**: Changes in neural pathways and synapses due to changes in behavior, environment, neural processes, or bodily injury.
- **feedforward processing**: A property of some neural circuits where signals move unidirectionally through the system from input to output with very little recurrent internal transmission.

Examining the consequences of damage to the the cerebellum provides the strongest clues to its function. Animals and humans with cerebellar dysfunction show problems with motor control. They can still generate motor activity, but lose precision and produce erratic, uncoordinated, or incorrectly timed movements. Functional imaging studies have also shown cerebellar activation in relation to language, attention, and mental imagery. Additionally, correlation studies have shown interactions between the cerebellum and non-motor areas of the cerebral cortex.

Principles of Cerebellar Function

The comparative simplicity and regularity of the cerebellar anatomy led to an early hope that a similar simplicity of computational function could be implied. This was expressed in one of the first books on cerebellar electrophysiology, *The Cerebellum as a Neuronal Machine* by John C. Eccles, Masao Ito, and Janos Szentágothai. Although a full understanding of cerebellar function has remained elusive, at least four principles have been identified as important: feedforward processing, divergence and convergence, modularity, and plasticity.

1. **Feedforward processing:** The cerebellum differs from other parts of the brain (especially the cerebral cortex) in that the signal processing is almost entirely feedforward. This means signals move unidirectionally through the system from input to output with very little recurrent internal transmission. The small amount of recurrence that does exist consists of mutual inhibition. There are no mutually excitatory circuits. This feedforward mode of operation means that the cerebellum, in contrast to the cerebral cortex, cannot generate self-sustaining patterns of neural activity. Signals enter the circuit, are processed by each stage in sequential order, and then leave. This provides a quick, concise response to any combination of inputs.
2. **Divergence and convergence:** In the human cerebellum, information from 200 million mossy fiber inputs is expanded to 40 billion granule cells. This neural divergence is followed by parallel fiber outputs that converge onto 15 million Purkinje cells. Due to their longitudinal alignment, the approximately 1000 Purkinje cells belonging to a microzone may receive input

via neural convergence from as many as 100 million parallel fibers. The cells then focus their own output down to a group of less than 50 deep nuclear cells. Therefore, the cerebellar network only receives a modest number of inputs to process and send results via a limited number of output cells.

3. **Modularity:** The cerebellar system is functionally divided into thousands of independent modules. All modules have a similar internal structure but different inputs and outputs. A module consists of a small cluster of neurons in the inferior olivary nucleus, a set of long narrow strips of Purkinje cells in the cerebellar cortex (microzones), and a small cluster of neurons in one of the deep cerebellar nuclei. Different modules share input, but also appear to function independently. The output of one module does not seem to significantly influence the activity of other modules.
4. **Plasticity:** The synapses between parallel fibers and Purkinje cells and between mossy fibers and deep nuclear cells are both susceptible to modification of their strength. In a single cerebellar module, input from as many as a billion parallel fibers converge onto a group of less than 50 deep nuclear cells, and the influence of each parallel fiber on those nuclear cells is adjustable. This arrangement gives tremendous flexibility for fine-tuning the relationship between cerebellar inputs and outputs.

Role of the Cerebellum in Motor Learning

There is considerable evidence that the cerebellum plays an essential role in some types of motor learning, most clearly in tasks in which fine adjustments must be made to an action's performance. There has been much dispute about whether learning takes place within the cerebellum itself, or whether it merely serves to provide signals that promote learning in other brain structures.

One of the most extensively studied cerebellar learning tasks is the eyeblink conditioning paradigm. A blink response is elicited when a neutral conditioned stimulus, such as a tone or a light, is repeatedly paired with an unconditioned stimulus, such as an air puff.

After many conditioned-unconditioned stimuli (CS-US) pairings, an association is formed whereby a learned blink, or conditioned response, occurs and precedes US onset. The magnitude of learning is measured by the percentage of all paired CS-US trials that result in a CR.

Experiments showed that lesions localized either to a specific part of the interpositus nucleus (one of the deep cerebellar nuclei), or to a few specific points in the cerebellar cortex, abolished learning of a correctly timed blink response. If cerebellar outputs are pharmacologically inactivated while leaving the inputs and intracellular circuits intact, learning takes place even while the animal fails to show any response. However, if intracerebellar circuits are disrupted, no learning takes place; these facts taken together make a strong case that learning occurs inside the cerebellum and that its cells exhibit neuroplasticity.

Clinical Relevance

Motor abnormalities are the primary symptoms of cerebellar dysfunction, and the nature of those depends on the part of the cerebellum affected.

EXAMPLE SYMPTOMS OF CEREBELLAR DYSFUNCTION

- Damage to the flocculonodular lobe (vestibulocerebellum): loss of equilibrium causing an altered walking gait
- Lateral zone damage: problems with skilled voluntary and planned movements leading to errors in intended movements (eg., dysdiadochokinesia, the inability to perform rapid alternating movements).
- Damage to the midline portion: disruption of whole-body movements
- Damage to the upper part of the cerebellum: gait impairments and other problems with leg coordination (ie, ataxia).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cerebellum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebellum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Purkinje cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Purkinje%20cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- granule cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/granule%20cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebellum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebellum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray706. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray706.png**. **License**: ***Public Domain: No Known Copyright***
- cerebellum.png. **Provided by**: Wikipedia commons. **Located at**: **commons.wikimedia.org/wiki/File%3ASobo_1909_624.png**. **License**: ***Public Domain: No Known Copyright***
- CerebellumDiv. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:CerebellumDiv.png**. **License**: ***Public Domain: No Known Copyright***
- Eyeblink Conditioning. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Eyeblink_conditioning**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebellum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebellum%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neural divergence. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/neural%20divergence**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neuroplasticity. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neuroplasticity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- feedforward processing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/feedforward%20processing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray706. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray706.png**. **License**: ***Public Domain: No Known Copyright***
- cerebellum.png. **Provided by**: Wikipedia commons. **Located at**: **commons.wikimedia.org/wiki/File%3ASobo_1909_624.png**. **License**: ***Public Domain: No Known Copyright***
- CerebellumDiv. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:CerebellumDiv.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 11.5B: Functions of the Cerebellum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6:THE DIENCEPHALON

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

11.6A: Functions of the Diencephalon

11.6B: Thalamus

11.6C: Hypothalamus

11.6D: Epithalamus and Pineal Gland

11.6E: Circumventricular Organs

This page titled 11.6:The Diencephalon is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6A: Functions of the Diencephalon

Distinct parts of diencephalon perform numerous vital functions, from regulating wakefulness to controlling the autonomic nervous system.

Learning Objectives

- Describe the functions of the diencephalon region of the brain


Key Points

- The diencephalon is made up of four main components: the thalamus, the subthalamus, the hypothalamus, and the epithalamus.
- The hypothalamus is an integral part of the endocrine system, with the key function of linking the nervous system to the endocrine system via the pituitary gland.
- The thalamus is critically involved in a number of functions including relaying sensory and motor signals to the cerebral cortex and regulating consciousness, sleep, and alertness.
- The epithalamus functions as a connection between the limbic system to other parts of the brain. Some functions of its components include the secretion of melatonin by the pineal gland (involved in circadian rhythms) and regulation of motor pathways and emotions.

Key Terms

- **subthalamus**: Receives afferent connections from the substantia nigra and striatum and regulates skeletal muscle movements.
- **thalamus**: Either of two large, ovoid structures of gray matter within the forebrain that relay sensory impulses to the cerebral cortex.
- **hypothalamus**: A region of the forebrain located below the thalamus, forming the basal portion of the diencephalon, and functioning to regulate body temperature, some metabolic processes, and the autonomic nervous system.
- **epithalamus**: The dorsal posterior segment of the diencephalon, involved in the maintenance of circadian rhythms and regulation of motor pathways and emotions.
- **limbic system**: A set of brain structures located on both sides of the thalamus, right under the cerebrum. Supports a variety of functions including emotion, behavior, motivation, long-term memory, and olfaction.


The diencephalon (“interbrain”) is the region of the vertebrate neural tube that gives rise to posterior forebrain structures. In development, the forebrain develops from the prosencephalon, the most anterior vesicle of the neural tube that later forms both the diencephalon and the telencephalon. In adults, the diencephalon appears at the upper end of the brain stem, situated between the cerebrum and the brain stem. It is made up of four distinct components: the thalamus, the subthalamus, the hypothalamus, and the epithalamus.



Diencephalon: Three-dimensional view of the diencephalon

Other structures that are part of the diencephalon are:

- Anterior and posterior paraventricular nuclei
- Medial and lateral habenular nuclei
- Stria medullaris thalami
- Posterior commissure
- Pineal gland



Embryonic Brain: Subdivisions of the embryonic vertebrate brain that later differentiate into forebrain, midbrain, and hindbrain structures.

Functions of Primary Diencephalon Structures

The thalamus is a kind of switchboard of information, believed to act as a relay between a variety of subcortical areas and the cerebral cortex. In particular, every sensory system (with the exception of the olfactory system) includes a thalamic nucleus that receives sensory signals and sends them to the associated primary cortical area. The thalamus also plays an important

role in regulating states of sleep and wakefulness. Thalamic nuclei have strong reciprocal connections with the cerebral cortex, forming thalamo-cortico-thalamic circuits that are believed to be involved with consciousness. The thalamus plays a major role in regulating arousal, awareness level, and activity. Damage to the thalamus can lead to permanent coma.

The subthalamus connects to the globus pallidus, a basal nucleus of the telencephalon. It receives afferent connections from the substantia nigra and striatum and regulates skeletal muscle movements.

The hypothalamus performs numerous vital functions (e.g., regulation of certain metabolic processes), most of which relate directly or indirectly to the regulation of visceral activities by way of other brain regions and the autonomic nervous system. It synthesizes and secretes certain neurohormones, often called hypothalamic-releasing hormones, and these in turn stimulate or inhibit the secretion of pituitary hormones. The hypothalamus controls body temperature, hunger, thirst, fatigue, sleep, and circadian cycles.

The epithalamus functions as a connection between the limbic system and other parts of the brain. Some functions of its components include the secretion of melatonin by the pineal gland (involved in circadian rhythms) and regulation of motor pathways and emotions.

This page titled 11.6A: Functions of the Diencephalon is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6B: Thalamus

The thalamus is a small structure in the center of the brain that acts as a relay center for sensory and motor information.

Learning Objectives

- Describe the thalamus and its functions

Key Points

- The thalamus is the largest structure derived from the embryonic diencephalon.
- Together, the two halves of the thalamus are a prominent bulb-shaped mass, about 5.7 cm in length, located obliquely and symmetrically on each side of the third ventricle.
- The thalamus has a system of myelinated fibers that separate the different thalamic subparts. These areas are defined by distinct clusters of neurons.
- Every sensory system (with the exception of the olfactory system) has a thalamic nucleus that receives sensory signals and sends them to the associated primary cortical area.
- The thalamus's functions include relaying sensory and motor signals to the cerebral cortex and the regulation of consciousness, sleep, and alertness.

Key Terms

- **nuclear**: In neuroanatomy, a nucleus is a brain structure consisting of a relatively compact cluster of neurons.
- **medial geniculate nucleus**: A part of the auditory thalamus that is the relay between the inferior colliculus and the auditory cortex.
- **lateral geniculate nucleus**: A relay center in the thalamus for the visual pathway that receives major sensory input from the retina.


The thalamus (derived from the Greek meaning "inner chamber") is a midline symmetrical structure within the brain, situated between the cerebral cortex and midbrain. Its functions include relaying sensory and motor signals to the cerebral cortex and regulating consciousness, sleep, and alertness. The thalamus surrounds the third ventricle and is the main product of the embryonic diencephalon.

Overall Structure of the Thalamus

Each half of the thalamus is about the size and shape of a walnut: approximately 3 centimeters long, 2.5 centimeters across at the widest point, and about 2 centimeters high. Together, the two halves of the thalamus are a prominent bulb-shaped mass, about 5.7 cm in length, located obliquely (about 30°) and symmetrically on each side of the third ventricle.

The thalamus is part of a nuclear complex of composed of four parts: the hypothalamus, epithalamus, the ventral thalamus, and the dorsal thalamus. The thalamus has a system of myelinated fibers that separate the different thalamic subparts. These areas are defined by distinct clusters of neurons.

The thalamus derives its blood supply from four arteries including the polar artery (posterior communicating artery), paramedian thalamic-subthalamic arteries, inferolateral (thalamogeniculate) arteries, and posterior (medial and lateral) choroidal arteries. These are all derived from the vertebrobasilar arterial system except the polar artery. The thalamus is connected to the hippocampus via the mammillothalamic tract.



The Nuclei of the Thalamus: The graphic details the various nuclei of the thalamus.

Primary Functions of the Thalamus

The thalamus has multiple functions, serving as a sort of switchboard of information. It is generally believed to act as a relay between a variety of subcortical areas and the cerebral cortex. In particular, every sensory system (with the exception of the olfactory system) has a thalamic nucleus that receives sensory signals and sends them to the associated primary cortical area. For the visual system, for example, inputs from the retina are sent to the lateral geniculate nucleus of the thalamus, which in turn projects to the primary visual cortex in the occipital lobe. The thalamus is believed to both process and relay sensory information. Each of the primary sensory relay areas receives strong back projections from the cerebral cortex. Similarly, the medial geniculate nucleus acts as a key auditory relay between the inferior colliculus of the midbrain and the primary auditory cortex. The ventral posterior nucleus is a key somatosensory relay, which sends touch and proprioceptive information to the primary somatosensory cortex.

The thalamus also plays an important role in regulating states of sleep and wakefulness. Thalamic nuclei have strong reciprocal connections with the cerebral cortex, forming thalamo-cortico-thalamic circuits that are believed to be involved with consciousness. The thalamus plays a major role in regulating arousal, awareness level, and activity. Damage to the thalamus can lead to permanent coma. Fatal familial insomnia is a hereditary prion disease characterized by degeneration of the thalamus and leading to a gradual progression to a state of total insomnia that is eventually fatal.

This page titled 11.6B: Thalamus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6C: Hypothalamus

The hypothalamus serves as a gateway between the nervous system and endocrine system.

Learning Objectives

- Describe the functions of the hypothalamus of the brain.


Key Points

- The hypothalamus produces and secretes a wide variety of neurohormones that lead to the release or inhibition of pituitary gland
  hormone.
- The posterior pituitary is composed of tissue derived from the hypothalamus, whereas the anterior pituitary is derived from epithelial tissue.
- The hypothalamus can pass signals back and forth between the anterior and posterior pituitary by the neurohypophysis and median eminence.
- The hypothalamus can sample the blood composition at two sites, the subfornical organ and the organum vasculosum of the lamina terminalis. This is important for the uptake of circulating hormones and to determine concentration of substances in the blood.
- The outputs of the hypothalamus can be divided into neural projections and endocrine hormones. The neural projections tend to run bidirectionally.

Key Terms


- **neurohypophysis**: The posterior lobe of the pituitary gland, responsible for the release of oxytocin and antidiuretic hormone (ADH), also called vasopressin.
- **median eminence**: Part of the inferior boundary for the hypothalamus and one of the seven areas of the brain devoid of a blood-brain barrier.
- **anterior pituitary**: Also called the adenohypophysis, the glandular anterior lobe of the pituitary gland. The anterior pituitary regulates several physiological processes including stress, growth, reproduction, and lactation.

The hypothalamus (derived from the Greek for “under chamber”) is a portion of the brain that contains a number of small, distinct nuclei with various functions and less anatomically distinct areas. The hypothalamus is located below the thalamus just above the brain stem. In the terminology of neuroanatomy, it forms the ventral part of the diencephalon. All vertebrate brains contain a hypothalamus. In humans, it is roughly the size of an almond.



The location of hypothalamus and pituitary: At center, hypothalamus is located just superior to the pituitary, with which it closely interacts.

General Functions



Location of the hypothalamus: This is the location of the human hypothalamus in relation to the thalamus, pituitary gland, sella turcica, and optic chiasm.

One of the most important functions of the hypothalamus is linking the nervous system to the endocrine system via the pituitary gland (hypophysis).

The hypothalamus contains thyrotropin-releasing hormone, gonadotropin-releasing hormone, growth hormone-releasing hormone, corticotropin-releasing hormone, somatostatin, and dopamine, as well as vasopressin and oxytocin. These hormones are released into the bloodstream and target other organ systems, most notably the pituitary. The hypothalamus affects the endocrine system and governs emotional behavior such as anger and sexual activity. Most of the hypothalamic hormones generated are distributed to the pituitary via the hypophyseal portal system. The hypothalamus maintains homeostasis, including the regulation of blood pressure, heart rate, and temperature.

The hypothalamus coordinates hormonal and behavioral circadian rhythms, complex patterns of neuroendocrine outputs, complex homeostatic mechanisms, and important behaviors. It must therefore respond to many different signals, some of which are generated externally and some internally. The hypothalamus is thus richly connected with many parts of the central nervous system, including the brainstem, reticular formation and autonomic zones, and the limbic forebrain (particularly the amygdala, septum, diagonal band of Broca, olfactory bulbs, and cerebral cortex).

The hypothalamus can sample the blood composition at the subfornical organ and the organum vasculosum of the lamina terminalis. This is important for the uptake of circulating hormones and to determine concentration of substances in the blood.

Temperature Regulation

The hypothalamus functions as a type of thermostat for the body. It sets a desired body temperature and stimulates either heat production and retention to raise the blood temperature to a higher level, or sweating and vasodilation to cool the blood to a lower temperature. All fevers result from a raised setting in the hypothalamus; elevated body temperatures due to any other cause are classified as hyperthermia. Rarely, direct damage to the hypothalamus such as from a stroke causes a fever; this is sometimes called a hypothalamic fever. However, such damage more commonly causes abnormally low body temperatures.

Appetite

The extreme lateral part of the ventromedial nucleus of the hypothalamus is responsible for control of food intake. Stimulation of this area causes increased food intake. Bilateral lesion in this area causes complete cessation of food intake. Medial parts of the nucleus have a controlling effect on the lateral part. Bilateral lesion of the medial part of the ventromedial nucleus causes hyperphagia and obesity. Further lesion of the lateral part of the ventromedial nucleus in the same animal produces complete cessation of food intake.

Fear Processing

The medial zone of hypothalamus is part of a circuitry that controls motivated behaviors, such as defensive behaviors and social defeat. Analyses of Fos-labeling showed that a series of nuclei in the behavioral control column is important in regulating the expression of innate and conditioned defensive behaviors. Nuclei in the medial zone are also mobilized during an encounter with an aggressor. The defeated animal has an increase in Fos levels in sexually dimorphic structures.

Sexual Dimorphism

Several hypothalamic nuclei are sexually dimorphic, with clear differences in both structure and function between males and females. Some differences are apparent even in gross neuroanatomy, most notably is the sexually dimorphic nucleus within the preoptic area. However, most are subtle changes in the connectivity and chemical sensitivity of particular sets of neurons. In neonatal life, gonadal steroids influence the development of the neuroendocrine hypothalamus. For instance, they determine the ability of females to exhibit a normal reproductive cycle and of males and females to display appropriate reproductive behaviors in adult life.

In 2004 and 2006, two studies by Berglund, Lindström, and Savic used Positron Emission Tomography (PET) to observe how the hypothalamus responds to smelling common odors, the scent of testosterone found in male sweat, and the scent of estrogen found in female urine. These studies showed that the hypothalamus of heterosexual men and homosexual women both respond to estrogen. Also, the hypothalamus of homosexual men and heterosexual women both respond to testosterone. In all four groups, common odors were processed similarly involving only the olfactory brain.

This page titled 11.6C: Hypothalamus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6D: Epithalamus and Pineal Gland

The epithalamus connects the limbic system to other parts of the brain.

Learning Objectives

- Describe the function of the epithalamus of the brain


Key Points

- The epithalamus is a dorsal posterior segment of the diencephalon, which includes the habenula and their interconnecting fibers, the habenular commissure, the stria medullaris, and the pineal body.
- A main function of the epithalamus is the secretion of melatonin by the pineal gland.
- The epithalamus is connected with both the limbic system and the basal ganglia.

Key Terms

- **circadian rhythm**: The "internal body clock" that regulates the 24-hour cycle of biological processes in animals and plants.
- **pineal gland**: A small, pinecone-shaped endocrine gland found near the centre of the brain that produces melatonin.
- **melatonin**: A hormone related to serotonin that is secreted by the pineal gland and stimulates color change in the skin of reptiles. It is involved in the sleep/wake and reproductive cycles in mammals.

The epithalamus is a dorsal posterior segment of the diencephalon (as shown in the figure below).



The Epithalamus: A brain sectioned in the median sagittal plane. The epithalamus is labeled in red.

It includes the habenula and their interconnecting fibers (the habenular commissure), the stria medullaris, and the pineal gland. The habenular commissure is a band of nerve fibers situated in front of the pineal gland that connects the habenular nuclei on both sides of the diencephalon. The stria medullaris, also known as stria medullaris thalami, is a fiber bundle containing afferent fibers from the septal nuclei, lateral preoptic hypothalamic region, and anterior thalamic nuclei to the habenula.

Pineal Gland

The pineal gland (also called the pineal body, epiphysis cerebri, epiphysis, conarium, or the “third eye”) is the only unpaired midline brain structure. It is about the size of a grain of rice (5–8 mm) in humans. The pineal gland lies between the laterally positioned thalamic bodies and behind the habenular commissure. It is located behind the third ventricle and is bathed in cerebrospinal fluid supplied through a small pineal recess of the third ventricle.

Epithalamic Function

The epithalamus acts as a connection between the limbic system and other parts of the brain. Some functions of its components include the secretion of melatonin by the pineal gland (involved in circadian rhythms) and regulation of motor pathways and emotions. It is wired with the limbic system and basal ganglia.

This page titled 11.6D: Epithalamus and Pineal Gland is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.6E: Circumventricular Organs

Circumventricular organs are situated adjacent to the brain ventricles and sense concentrations of various compounds in the blood.

Learning Objectives

- Describe the circumventricular organs of the brain

Key Points


- Circumventricular organs have incomplete blood-brain barriers.
- Circumventricular organs secrete or are sites of action of a variety of different hormones, neurotransmitters, and cytokines. They are sometimes classified by whether they are secretory or sensory.
- The sensory organs are able to sense plasma molecules and then pass that information into other regions of the brain.
- The secretory organs are responsible for secreting hormones and glycoproteins into the peripheral vascular system using feedback from both the brain environment and external stimuli.

Key Terms

- **circumventricular organs**: So named because they are positioned at distinct sites around the margin of the ventricular system of the brain. They are among the few sites in the brain which have an incomplete blood-brain barrier and, as a result, can directly sense the concentrations of various compounds, particularly peptide hormones, in the bloodstream.
- **median eminence**: Part of the inferior boundary for the hypothalamus and one of the seven areas of the brain devoid of a blood-brain barrier.
- **subcommissural organ**: A gland in the brain and one of the circumventricular organs, consisting of ependymal cells which secrete SCO-spondin, a protein that contributes to neuron growth and maintenance.

Circumventricular organs (CVOs) are positioned at distinct sites around the margin of the ventricular system of the brain. They are among the few sites in the brain that have an incomplete blood-brain barrier. As a result, neurons located in circumventricular organs can directly sense the concentrations of various compounds, particularly peptide hormones, in the bloodstream without the need for specialized transport systems that move those compounds across the blood-brain barrier. A useful mnemonic device for remembering this aspect of their function, though not the source of the name, is that they allow factors to circumvent' the blood-brain barrier. These organs secrete or are sites of action of a variety of different hormones, neurotransmitters, and cytokines. They are sometimes classified by whether they are secretory or sensory.

CVOs allow for linkage between the central nervous system (CNS) and peripheral blood flow, and are an integral part of neuroendocrine function.



CVO: View of the third and fourth ventricles. The CVOs border these ventricles.

Sensory Circumventricular Organs

The sensory organs are able to sense plasma molecules and pass that information into other regions of the brain. Therefore, they provide direct information to the autonomic nervous system from the systemic circulation. These organs include:

- Area postrema: Site of the chemoreceptor trigger zone for vomiting, sends major and minor efferents to sections of the brain involved in the autonomic control of cardiovascular and respiratory activities.
- Subfornical organ: Active in osmoregulation, cardiovascular regulation, and energy homeostasis.
- Vascular organ of lamina terminalis: Responsible for the homeostatic conservation of osmolarity.

Secretory Circumventricular Organs

The secretory organs are responsible for secreting hormones and glycoproteins into the peripheral vascular system using feedback from both the brain environment and external stimuli. These include:

- Subcommissural organ (SCO): Secretion of the glycoprotein SCO-spondin.
- Posterior pituitary: Stores and releases oxytocin and vasopressin, also known as antidiuretic hormone, produced in the hypothalamus.
- Pineal gland: The main function is the secretion of melatonin.
- Median eminence: Allows for the transport of neurohormones between the CSF and the peripheral blood supply.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Subthalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Subthalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypothalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypothalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thalamus%23Function**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- Diencephalon. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Diencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypothalamus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypothalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epithalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/epithalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thalamus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Diencephalon small.gif. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Diencephalon_small.gif**. **License**: ***Public Domain: No Known Copyright***
- EmbryonicBrain.svg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:EmbryonicBrain.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lateral Geniculate Nucleus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lateral_geniculate_nucleus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medial Geniculate Nucleus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Medial_geniculate_nucleus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nuclear. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/nuclear**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Diencephalon small.gif. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Diencephalon_small.gif**. **License**: ***Public Domain: No Known Copyright***
- EmbryonicBrain.svg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:EmbryonicBrain.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thalmus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thalmus.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypothalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypothalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- median eminence. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/median%20eminence**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neurohypophysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neurohypophysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anterior pituitary. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/anterior%20pituitary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Diencephalon small.gif. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Diencephalon_small.gif**. **License**: ***Public Domain: No Known Copyright***
- EmbryonicBrain.svg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:EmbryonicBrain.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thalmus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thalmus.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypothalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Hypothalamus.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- LocationOfHypothalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:LocationOfHypothalamus.jpg**. **License**: ***Public Domain: No Known Copyright***
- Pineal body. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pineal_body**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Diencephalon. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Diencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Habenular commissure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Habenular_commissure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Stria medullaris. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Stria_medullaris**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epithalamus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epithalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- melatonin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/melatonin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- circadian rhythm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/circadian_rhythm**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- pineal gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pineal_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- File:Diencephalon small.gif. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Diencephalon_small.gif. **License**: *Public Domain: No Known Copyright*
- EmbryonicBrain.svg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:EmbryonicBrain.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thalmus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Thalmus.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypothalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Hypothalamus.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- LocationOfHypothalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:LocationOfHypothalamus.jpg. **License**: *Public Domain: No Known Copyright*
- Epithalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Epithalamus.png. **License**: *Public Domain: No Known Copyright*
- Circumventricular organs. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circumventricular_organs. **License**: *CC BY-SA: Attribution-ShareAlike*
- circumventricular organs. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/circumventricular%20organs. **License**: *CC BY-SA: Attribution-ShareAlike*
- median eminence. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/median%20eminence. **License**: *CC BY-SA: Attribution-ShareAlike*
- subcommissural organ. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/subcommissural%20organ. **License**: *CC BY-SA: Attribution-ShareAlike*
- File:Diencephalon small.gif. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Diencephalon_small.gif. **License**: *Public Domain: No Known Copyright*
- EmbryonicBrain.svg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:EmbryonicBrain.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thalmus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Thalmus.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypothalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Hypothalamus.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- LocationOfHypothalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:LocationOfHypothalamus.jpg. **License**: *Public Domain: No Known Copyright*
- Epithalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Epithalamus.png. **License**: *Public Domain: No Known Copyright*
- Ventricular System. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventricular_system. **License**: *CC BY: Attribution*

This page titled 11.6E: Circumventricular Organs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.7: THE CEREBRUM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

11.7A: Overview of the Cerebrum

11.7B: Cerebral Lobes

11.7C: White Matter of the Cerebrum

11.7D: Basal Ganglia

11.7E: Limbic System

This page titled 11.7: The Cerebrum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.7A: Overview of the Cerebrum

With the assistance of the cerebellum, the cerebrum controls all voluntary actions in the body.

LEARNING OBJECTIVE

Describe the function of the cerebrum

Key Takeaways

Key Points

- The cerebrum is the largest and most developed of the five major divisions of the brain.
- The brain contains two hemispheres, the left and the right, connected by a bundle of nerve fibers called the corpus callosum.
- The cerebrum directs the conscious or volitional motor functions of the body. Damage to this area of the brain can result in loss of muscular power and precision rather than total paralysis.
- The primary sensory areas of the cerebral cortex receive and process visual, auditory, somatosensory, gustatory, and olfactory information.
- Each hemisphere of the mammalian cerebral cortex can be broken down into four functionally and spatially defined lobes: frontal, parietal, temporal, and occipital.

Key Terms

- **sulci**: Any of the grooves that mark the convolutions of the surface of the brain (plural of sulcus).
- **cerebral cortex**: The cerebrum's outer layer of neural tissuecomposed of folded gray matter. The cerebral cortex plays a key role in memory, attention, perception, awareness, thought, language, and consciousness.
- **olfactory bulb**: A neural structure of the vertebrate forebrain involved in olfaction (sense of smell).
- **Broca's area**: A region in the frontal lobe of the dominant hemisphere of the human brain with functions linked to speech production.
- **Wernicke's area**: Involved in the comprehension or understanding of written and spoken language.
- **aphasia**: A combination speech and language disorder often caused by a stroke.
- **gyri**: A ridge on the cerebral cortex (plural of gyrus).




Cerebrum Animation: Location of the cerebrum (in red).

The cerebrum, which lies in front or on top of the brainstem, comprises a large portion of the brain. In humans, it is the largest and best-developed of the brain's five major divisions. The cerebrum is the newest structure in the phylogenetic sense, with mammals having the largest and most developed among all species.

The cerebrum contains the cerebral cortex (of the two cerebral hemispheres), as well as several subcortical structures, including the hippocampus, basal ganglia, and olfactory bulb. In larger mammals, the cerebral cortex is folded into many gyri and sulci, which allows it to expand in surface area without taking up much greater volume. With the assistance of the cerebellum, the cerebrum controls all voluntary actions in the body.

Cerebral Cortex




The cerebral cortex: The cerebral cortex is the outer layer depicted in dark violet. Notice the folded structure of the cortex: the "valleys" of the cortex are known as sulci.

The cortex is composed of two hemispheres, right and left, separated by a large sulcus. A thick fiber bundle, the corpus callosum, connects the two hemispheres, allowing information to be passed from one side to the other. The right hemisphere controls and processes signals from the left side of the body, while the left hemisphere controls and processes signals from the right side of the body.

The Four Brain Lobes

Each hemisphere of the mammalian cerebral cortex can be broken down into four functionally and spatially defined lobes: frontal, parietal, temporal, and occipital.

The frontal lobe is located at the front of the brain, over the eyes, and contains the olfactory bulb. The frontal lobe also contains the motor cortex, which is important for planning and implementing movement.



Cerebral Lobes: Locations of the cerebral lobes

Two of the parietal lobe's main functions are processing somatosensation (touch sensations such as pressure, pain, heat, cold) and proprioception (the sense of how parts of the body are oriented in space).

The temporal lobe is located at the base of the brain by the ears. It is primarily involved in processing and interpreting sounds. It also contains the hippocampus, which processes memory formation.

The occipital lobe is located at the back of the brain. It is primarily involved in vision: seeing, recognizing, and identifying the visual world.

Cerebrum Function

The cerebrum directs the conscious or volitional motor functions of the body. These functions originate within the primary motor cortex and other frontal lobe motor areas where actions are planned. Upper motor neurons in the primary motor cortex send their axons to the brainstem and spinal cord to synapse on the lower motor neurons, which innervate the muscles. Damage to motor areas of cortex can lead to certain types of motor neuron disease. This kind of damage results in loss of muscular power and precision rather than total paralysis.

The olfactory sensory system is unique in that neurons in the olfactory bulb send their axons directly to the olfactory cortex, rather than to the thalamus first. Damage to the olfactory bulb results in a loss of the sense of smell. The olfactory bulb also receives "top-down" information from such brain areas as the amygdala, neocortex, hippocampus, locus coeruleus, and substantia nigra. Its potential functions can be placed into four non-exclusive categories: discriminating among odors, enhancing sensitivity of odor detection, filtering out background odors, and permitting higher brain areas involved in arousal and attention to modify the detection or the discrimination of odors.

Speech and language are mainly attributed to parts of the cerebral cortex. Motor portions of language are attributed to Broca's area within the frontal lobe. Speech comprehension is attributed to Wernicke's area, at the temporal-parietal lobe junction. Damage to the Broca's area results in expressive aphasia (non-fluent aphasia) while damage to Wernicke's area results in receptive aphasia.

This page titled 11.7A: Overview of the Cerebrum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.7B: Cerebral Lobes

The cortex is divided into four main lobes: frontal, parietal, occipital, temporal.

LEARNING OBJECTIVE

Distinguish between the frontal, temporal, parietal, and occipital lobes of the cerebral cortex


Key Takeaways

Key Points

- Each lobe contributes to overall functionality of the brain and each lobe has many different roles.
- The frontal lobe is involved in conscious thought.
- The parietal lobe is important for spatial reasoning.
- The occipital lobe is required for visual processing.
- The temporal lobe contributes to language and face recognition.

Key Terms

- **frontal lobe**: The frontal lobe is an area in the brain of mammals, located at the front of each cerebral hemisphere and positioned anterior to the parietal lobe and superior and anterior to the temporal lobes. In humans, it contributes to a number of higher cognitive functions including attention, planning, and motivation.
- **temporal lobe**: A region of the cerebral cortex that is located behind the temples and beneath the Sylvian fissure on both cerebral hemispheres of the human brain. This region is involved in auditory perception, speech and vision processing, and the formation of long-term memory as it houses the hippocampus.
- **parietal lobe**: A part of the brain positioned superior to the occipital lobe and posterior to the frontal lobe that integrates sensory information from different modalities, particularly spatial sense and navigation.
- **occipital lobe**: Located at the back of the head, this is the visual processing center of the mammalian brain containing most of the anatomical region of the visual cortex.



Cerebral lobes: The four lobes (frontal, parietal, occipital, and temporal) of the human brain are depicted along with the

cerebellum.

Brain lobes were originally a purely anatomical classification, but we now know they are also associated with specific brain functions. The telencephalon (cerebrum), the largest portion of the human brain, is divided into lobes like the cerebellum. If not specified, the expression “lobes of the brain” refers to the telencephalon. There are four uncontested lobes of the telencephalon:

The Frontal Lobe

The frontal lobe is an area in the mammalian brain located at the front of each cerebral hemisphere and positioned anterior to (in front of) the parietal lobe and superior and anterior to the temporal lobes. It is separated from the parietal lobe by a space between tissues called the central sulcus and from the temporal lobe by a deep fold called the lateral (Sylvian) sulcus. The precentral gyrus, forming the posterior border of the frontal lobe, contains the primary motor cortex, which controls voluntary movements of specific body parts.

The frontal lobe contains most of the dopamine-sensitive neurons in the cerebral cortex. The dopamine system is associated with reward, attention, short-term memory tasks, planning, and motivation. Dopamine tends to limit and select sensory information that the thalamus sends to the forebrain. A report from the National Institute of Mental Health indicates that a gene variant that reduces dopamine activity in the prefrontal cortex is related to poorer performance in that region during memory tasks; this gene variant is also related to slightly increased risk for schizophrenia.

The frontal lobe is considered to contribute to our most human qualities. Damage to the frontal lobe can result in changes in personality and difficulty planning. The frontal lobes are the most uniquely human of all the brain structures.

The Parietal Lobe

The parietal lobe is a part of the brain positioned above (superior to) the occipital lobe and behind (posterior to) the frontal lobe. The parietal lobe integrates sensory information from different modalities, particularly spatial sense and navigation. For example, it comprises the somatosensory cortex and the dorsal stream of the visual system. This enables regions of the parietal cortex to map objects perceived visually into body coordinate positions.

Several portions of the parietal lobe are also important in language processing. Also, this lobe integrates information from various senses and assists in the manipulation of objects. Portions of the parietal lobe are involved with visuospatial processing.

The Occipital Lobe

The two occipital lobes are the smallest of the four paired lobes in the human cerebral cortex. Located in the rearmost portion of the skull, the occipital lobes are part of the forebrain. At the front edge of the occipital there are several lateral occipital gyri separated by lateral occipital sulci. The occipital lobe is involved in the sense of sight; lesions in this area can produce hallucinations.

The Temporal Lobe

The temporal lobe is a region of the cerebral cortex located beneath the lateral fissure on both cerebral hemispheres of the mammalian brain. The temporal lobes are involved in many functions, such as retaining visual memories, processing sensory input, comprehending language, storing new memories, feeling and expressing emotion, and deriving meaning. The temporal lobe contains the hippocampus and plays a key role in the formation of explicit long-term memory, modulated by the amygdala. It is involved in the senses of smell and sound as well as in processing of complex stimuli.

Adjacent areas in the superior, posterior, and lateral parts of the temporal lobes are involved in high-level auditory processing. The temporal lobe is involved in primary auditory perception such as hearing and holds the primary auditory cortex. The superior temporal gyrus includes an area where auditory signals from the ear first reach the cerebral cortex and are processed by the primary auditory cortex in the left temporal lobe.

The areas associated with vision in the temporal lobe interpret the meaning of visual stimuli and establish object recognition. The ventral part of the temporal cortices appear to be involved in high-level visual processing of complex stimuli such as faces

(fusiform gyrus) and scenes (parahippocampal gyrus). Anterior parts of this ventral stream for visual processing are involved in object perception and recognition.

This page titled 11.7B: Cerebral Lobes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.7C: White Matter of the Cerebrum

White matter is composed of myelinated axons and glia and connects distinct areas of the cortex.

LEARNING OBJECTIVE

Describe the white matter of the cerebrum

Key Takeaways

Key Points

- White matter modulates the distribution of action potentials, acting as a relay and coordinating communication between different brain regions.
- There are three main kinds of white matter tracts: projection, commissural, and association.
- The largest white matter structure of the brain is the corpus collosum, a form of commissural tract that connects the right and left hemispheres.

Key Terms

- **corpus collosum**: A wide, flat bundle of neural fibers beneath the cortex that connects the left and right cerebral hemispheres and facilitates interhemispheric communication.
- **grey matter**: A major component of the central nervous system, consisting of neuronal cell bodies, neuropil (dendrites and unmyelinated axons), glial cells (astroglia and oligodendrocytes), and capillaries.
- **gyri**: Ridges on the cerebral cortex, generally surrounded by one or more sulci.



A lateral cross-section of the human brain: White matter appears white in this dissected human brain, while gray matter appears darker. White matter is composed largely of myelinated axons.

White matter is one of the two components of the central nervous system (CNS). It consists mostly of glial cells and myelinated axons and forms the bulk of the deep parts of the cerebrum and the superficial parts of the spinal cord. In a freshly cut brain, the tissue of white matter appears pinkish white to the naked eye because myelin is composed largely of lipid tissue containing capillaries. The axons of white matter transmit signals from various grey matter areas (the locations of nerve cell bodies) of the cerebrum to each other and carry nerve impulses between neurons. While grey matter is primarily associated with processing and cognition, white matter modulates the distribution of action potentials, acting as a relay and coordinating communication between different brain regions.


Tracts

There are three different kinds of tracts (bundles of axons) that connect one part of the brain to another within the white matter:

- Projection tracts extend vertically between higher and lower brain areas and spinal cord centers, and carry information between the cerebrum and the rest of the body. Other projection tracts carry signals upward to the cerebral cortex. Superior to the brainstem, such tracts form a broad, dense sheet called the internal capsule between the thalamus and basal nuclei, then radiate in a diverging, fanlike array to specific areas of the cortex.
- Commissural tracts cross from one cerebral hemisphere to the other through bridges called commissures. The great majority of commissural tracts pass through the large corpus callosum. A few tracts pass through the much smaller anterior and posterior commissures. Commissural tracts enable the left and right sides of the cerebrum to communicate with each other.
- Association tracts connect different regions within the same hemisphere of the brain. Long association fibers connect different lobes of a hemisphere to each other, whereas short association fibers connect different gyri within a single lobe. Among their roles, association tracts link perceptual and memory centers of the brain.

Corpus Callosum

The corpus callosum (Latin: “tough body”), also known as the colossal commissure, is a wide, flat bundle of neural fibers beneath the cortex in the eutherian brain at the longitudinal fissure. It connects the left and right cerebral hemispheres and facilitates interhemispheric communication. It is the largest white matter structure in the brain, consisting of 200 to 250 million contralateral axonal projections.



Corpus Callosum: Location of the corpus callosum in the cerebrum.

The posterior portion of the corpus callosum is called the splenium, the anterior is called the genu (or “knee”), and the area between the two is the truncus or body of the corpus callosum. The part between the body and the splenium is often markedly thin and thus called the isthmus. The rostrum is the part of the corpus callosum that projects posteriorly and inferiorly from the anteriormost genu. The rostrum is so named for its resemblance to a bird’s beak.

Agenesis of the corpus callosum (ACC) is a rare congenital disorder in which the corpus callosum is partially or completely absent. It is usually diagnosed within the first two years of life and may manifest as a severe syndrome in infancy or childhood, as a milder condition in young adults, or as an asymptomatic incidental finding. Initial symptoms of ACC usually include seizures that may be followed by feeding problems and delays in holding the head erect, sitting, standing, and walking. Hydrocephaly may also occur.

Other possible symptoms include impairments in mental and physical development, hand-eye coordination, and visual and auditory memory. In mild cases, symptoms such as seizures, repetitive speech, or headaches may not appear for years.

This page titled 11.7C: White Matter of the Cerebrum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.7D: Basal Ganglia

The basal ganglia is important for the control of movement and forming habits, and each of its components has a complex internal anatomical and neurochemical organization.

LEARNING OBJECTIVE

Describe the functions of the basal ganglia

Key Takeaways

Key Points


- The basal ganglia are a group of nuclei of varied origin in the brains of vertebrates that act as a cohesive functional unit.
- The basal ganglia are associated with a variety of functions including voluntary motor control, procedural learning relating to routine behaviors or “habits” such as bruxism, eye movements, cognitive, and emotional functions.
- The basal ganglia are composed of the striatum, the pallidum, the substantia nigra, and the subthalamic nucleus.
- The basal ganglia play a central role in a number of neurological conditions including Parkinson’s disease and Huntington’s disease.

Key Terms

- **Huntington’s disease**: A neurodegenerative genetic disorder that affects muscle coordination and leads to cognitive decline and psychiatric problems.
- **Parkinson’s disease**: A chronic neurological disorder resulting in lack of control over movement; poor balance and coordination; and similar symptoms.
- **executive functions**: Also known as cognitive control and supervisory attentional system. Refers to a set of cognitive processes, including attentional control, inhibitory control, working memory, and cognitive flexibility, reasoning, problem solving, and planning.
- **bruxism**: Excessive grinding and clenching of the teeth.

The basal ganglia (or basal nuclei) are a group of nuclei of varied origin in the brains of vertebrates that act as a cohesive functional unit. They are situated at the base of the forebrain and are strongly connected with the cerebral cortex, thalamus, and other brain areas. The components of the basal ganglia include the striatum, pallidum, substantia nigra, and subthalamic nucleus. Each of these components has a complex internal anatomical and neurochemical organization.

Structure



The Basal Ganglia: The basal nuclei are often referred to as the basal ganglia. The main components of the basal nuclei are labeled in purple.

The main components of the basal ganglia are:

- Striatum, or neostriatum: This component consists of 3 divisions: the caudate, putamen, and ventral striatum (includes the nucleus accumbens). The striatum receives input from many brain areas, but sends output only to other components of the basal ganglia.
- Globus pallidus, or pallidum: This component is composed of the globus pallidus externa (GPe) and globus pallidus interna (GPi). The pallidum receives its most important input from the striatum (either directly or indirectly), and sends inhibitory output to a number of motor-related areas, including the part of the thalamus that projects to the motor-related areas of the cortex.
- Substantia nigra: This component consists of the substantia nigra pars compacta (SNc) and substantia nigra pars reticulata (SNr). The SNr functions similarly to the pallidum, and the SNc cells contain neuromelanin and produce dopamine (a neurotransmitter) for input to the striatum.
- Subthalamic nucleus (STN): The STN receives input mainly from the striatum and cortex, and projects to a portion of the pallidum (interna portion or GPi). It is the only portion of the ganglia that produces an excitatory neurotransmitter, glutamate. The role of the subthalamic nucleus is to stimulate the SNr-GPi complex, and it receives inhibitory input from the GPe and sends excitatory signal to the GPi.

Function

The basal ganglia are associated with a variety of functions, including voluntary motor control, procedural learning relating to routine behaviors or habits such as bruxism, eye movements, and cognitive, emotional functions. Currently, popular theories implicate the basal ganglia primarily in action selection, that is, the decision of which several possible behaviors to execute at a given time. Experimental studies show that the basal ganglia exert an inhibitory influence on a number of motor systems and that a release of this inhibition permits a motor system to become active. The behavior switching that takes place within the basal ganglia is influenced by signals from many parts of the brain, including the prefrontal cortex, which plays a key role in executive functions.

The basal ganglia play a central role in a number of neurological conditions, including several movement disorders. The most notable are Parkinson's disease, which involves degeneration of the melanin-pigmented dopamine-producing cells in the substantia nigra pars compacta (SNc), and Huntington's disease, which primarily involves damage to the striatum. Basal ganglia dysfunction is also implicated in some other disorders of behavior control such as Tourette's syndrome, ballismus (particularly hemibalismus), obsessive-compulsive disorder, and Wilson's disease (hepatolenticular degeneration). With the exception of

Wilson's disease and hemiballismus, the neuropathological mechanisms underlying diseases of ganglia such as Parkinsons' and Huntington's are not very well understood or are at best still developing theories.

The basal ganglia have a limbic sector whose components are the nucleus accumbens, ventral pallidum, and ventral tegmental area (VTA). This limbic sector is thought to play a central role in reward learning, particularly a pathway from the VTA to the nucleus accumbens that uses the neurotransmitter dopamine. A number of highly addictive drugs, including cocaine, amphetamine, and nicotine, are thought to work by increasing the efficacy of this dopamine signal.

This page titled 11.7D: Basal Ganglia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


11.7E: Limbic System

LEARNING OBJECTIVE

- Describe the functions of the limbic system

The limbic system, or paleomammalian brain, is a set of brain structures in the precortex and subcortex of the brain. It includes the hippocampus, amygdala, anterior thalamic nuclei, septum, limbic cortex, and fornix, and supports a variety of functions including emotion, behavior, motivation, long-term memory, and olfaction. The term “limbic” comes from the Latin limbus, for “border” or “edge,” because the limbic system forms the inner border of the cortex.

The limbic system consists of various structures that each support distinctive brain functions.



The Limbic System: This diagram of the limbic system delineates components of the diencephalon and cerebrum.

Hippocampus and Associated Structures

- Hippocampus: Required for the formation of long-term memories and implicated in maintenance of cognitive maps for navigation.
- Amygdala: Involved in signaling the cortex of motivationally-significant stimuli, such as those related to reward and fear, and in social functions, such as mating.
- Fornix: A white matter structure that carries signals from the hippocampus to the mammillary bodies and septal nuclei.
- Mammillary body: Important for the formation of memory.

Septal Nuclei

- These lie below the rostrum of the corpus callosum and anterior to the lamina terminalis. The septal nuclei receive reciprocal connections from the olfactory bulb, hippocampus, amygdala, hypothalamus, midbrain, habenula, cingulate gyrus, and thalamus.

Limbic Lobe

A phylogenetically old structure consisting of the following structures:

- Parahippocampal gyrus: Plays a role in the formation of spatial memory
- Cingulate gyrus: Conducts autonomic functions regulating heart rate, blood pressure, and cognitive and attentional processing
- Dentate gyrus: Thought to contribute to the formation of new memories

Additional Structures

- Entorhinal cortex: Important memory and associative components
- Piriform cortex: Processes olfactory information
- Fornicate gyrus: Region encompassing the cingulate and parahippocampal gyrus
- Nucleus accumbens: Involved in reward, pleasure, and addiction
- Orbitofrontal cortex: Involved in cognitive processing during decision-making

Limbic System Function

The limbic system operates by influencing the endocrine system and the autonomic nervous system. It is highly interconnected with the nucleus accumbens, the brain's pleasure center, which plays a role in sexual arousal and the "high" derived from certain recreational drugs. The structures of the limbic system are involved in motivation, emotion, learning, and memory.

The limbic system is also tightly connected to the prefrontal cortex. Some scientists contend that this connection is related to the pleasure obtained from solving problems. To cure severe emotional disorders, this connection was sometimes surgically severed, a procedure of psychosurgery called a prefrontal lobotomy. Patients who underwent this procedure often became passive and lacked motivation.

Key Points

- The lymbic system includes the hippocampus, amygdala, anterior thalamic nuclei, septum, limbic cortex, and fornix, which together support a variety of functions including emotion, behavior, motivation, long-term memory, and olfaction.
- The limbic system operates by influencing the endocrine system and the autonomic nervous system. It is highly interconnected with the nucleus accumbens, the brain's pleasure center, which plays a role in sexual arousal and the "high" derived from certain recreational drugs.
- The limbic system is also tightly connected to the prefrontal cortex. Some scientists contend that this connection is related to the pleasure obtained from solving problems.

Key Terms

- **nucleus accumbens**: A collection of neurons that forms the main part of the ventral striatum. It is thought to play an important role in reward, pleasure, laughter, addiction, aggression, fear, and the placebo effect.
- **hippocampus**: A part of the brain located inside the temporal lobe consisting mainly of gray matter. It is a component of the limbic system and plays a role in memory and emotion.
- **mammillary bodies**: Act as a relay for impulses coming from the amygdalae and hippocampi.
- **prefrontal cortex**: The anterior part of the frontal lobes of the brain, lying in front of the motor and premotor areas. This brain region has been implicated in planning complex cognitive behavior, personality expression, decision making, and moderating social behavior, but its primary function is the orchestration of thoughts and actions in accordance with internal goals.
- **septal nucle**: Play a role in reward and reinforcement along with the nucleus accumbens.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Wernicke's area. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wernicke%27s_area.. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Broca's area. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Broca%27s_area. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Olfactory_bulb. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Cerebral Hemisphere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebral_hemisphere**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebrum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebrum%23Structure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebral cortex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebral_cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Aphasia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Aphasia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gyri. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gyri**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sulci. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sulci**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebrum animation small. **Authored by**: BodyParts3D. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Cerebrum_animation_small.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cerebrum Lobes. **Authored by**: Jkwchui. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Cerebrum_lobes.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brainmaps macaque hippocampus. **Provided by**: Brainmaps.org. **Located at**: **en.Wikipedia.org/wiki/File:Brainmaps-macaque-hippocampus.jpg**. **License**: ***CC BY: Attribution***
- Lobes of the brain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lobes_of_the_brain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Temporal lobe. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Temporal_lobe**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Frontal lobe. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Frontal_lobe**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parietal lobe. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parietal_lobe**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Occipital lobe. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Occipital_lobe**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Corpus callosum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Corpus_callosum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- White matter. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/White_matter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- grey matter. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/grey%20matter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human brain right dissected lateral view description. **Authored by**: John A Beal. **Located at**: **commons.wikimedia.org/wiki/F...escription.JPG**. **License**: ***CC BY: Attribution***
- Corpus_callosum.jpg. **Authored by**: John A. Beal, Ph.D; Looie496. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Corpus_callosum.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parkinson. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parkinson%27s_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Huntington's disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Huntington%27s_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basal ganglia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Basal_ganglia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basal Ganglia and Related Structures. **Authored by**: John Henkel; Leevanjackson. **Provided by**: Food and Drug Administration. **Located at**: **commons.wikimedia.org/wiki/F...Structures.svg**. **License**: ***Public Domain: No Known Copyright***
- Limbic system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Limbic_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fornicate Gyrus. **Provided by**: Wikipedia. **Located at**: **en.wiktionary.org/wiki/fornicate_gyrus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- prefrontal cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/prefrontal_cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hippocampus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hippocampus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleus accumbens. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nucleus_accumbens**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blausen 0614 LimbicSystem. **Authored by**: BruceBlaus. **Provided by**: Wikimedia Commons. **Located at**:

commons.wikimedia.org/wiki/F...mbicSystem.png. **License**: ***CC BY: Attribution***
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

PUBLIC DOMAIN CONTENT

- Brainlobes. **Authored by**: King of Hearts. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/F...Brainlobes.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 11.7E: Limbic System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.8: FUNCTIONAL SYSTEMS OF THE CEREBRAL CORTEX

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

11.8A: Sensory Areas

11.8B: Motor Areas

11.8C: Association Areas

11.8D: Hemispheric Lateralization

This page titled 11.8: Functional Systems of the Cerebral Cortex is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.8A: Sensory Areas

Sensory areas of the brain receive and process sensory information, including sight, touch, taste, smell, and hearing.

Learning Objectives

- Describe the sensory areas of the cerebral cortex

Key Points

- The cortex can be divided into three functionally distinct areas: sensory, motor, and associative.
- The main sensory areas of the brain include the primary auditory cortex, primary somatosensory cortex, and primary visual cortex.
- In general, the two hemispheres receive information from the opposite side of the body. For example, the right primary somatosensory cortex receives information from the left limbs, and the right visual cortex receives information from the left eye.
- Sensory areas are often represented in a manner that makes topographical sense.

Key Terms

- **calcarine sulcus**: An anatomical landmark located at the caudal end of the medial surface of the brain.
- **primary somatosensory cortex**: The main sensory receptive area for the sense of touch.
- **primary auditory cortex**: A region of the brain that processes sound and thereby contributes to our ability to hear.
- **primary visual cortex**: Located in the posterior pole of the occipital cortex, the simplest, earliest cortical visual area. It is highly specialized for processing information about static and moving objects and is excellent in pattern recognition.

Sensory areas are the areas of the brain that receive and process sensory information. The cerebral cortex is connected to various subcortical structures such as the thalamus and the basal ganglia. Most sensory information is routed to the cerebral cortex via the thalamus. Olfactory information, however, passes through the olfactory bulb to the olfactory cortex, bypassing the thalamus. The cortex is commonly described as composed of three parts: sensory, motor, and association areas. Parts of the cortex that receive sensory inputs from the thalamus are called primary sensory areas. Each of the five senses relates to specific groups of brain cells that categorize and integrate sensory information.

The Five Sensory Modalities

The five commonly recognized sensory modalities, including sight, hearing, taste, touch, and smell, are processed as follows:

Somatosensory System

The primary somatosensory cortex, located across the central sulcus and behind the primary motor cortex, is configured to generally correspond with the arrangement of nearby motor cells related to specific body parts.

Taste

The primary gustatory area is near the face representation within the postcentral gyrus.

Olfaction

The olfactory cortex is located in the uncus, found along the ventral surface of the temporal lobe. Olfaction is the only sensory system that is not routed through the thalamus.

Vision

The visual area is located on the calcarine sulcus deep within the inside folds of the occipital lobe.

Hearing


The primary auditory cortex is located on the transverse gyri that lie on the back of the superior temporal convolution of the temporal lobes.

Organization of Sensory Maps

In general, each brain hemisphere receives information from the opposite side of the body. For example, the right primary somatosensory cortex receives information from the left limbs, and the right visual cortex receives information from the left eye. The organization of sensory maps in the cortex reflects that of the corresponding sensing organ, in what is known as a topographic map. Neighboring points in the primary visual cortex, for example, correspond to neighboring points in the retina. This topographic map is called a retinotopic map.

Similarly, there is a tonotopic map in the primary auditory cortex and a somatotopic map in the primary sensory cortex. This somatotopic map has commonly been illustrated as a deformed human representation, the somatosensory homunculus, in which the size of different body parts reflects the relative density of their innervation.

A cortical homunculus is a physical representation of the human body located within the brain. This neurological map of the anatomical divisions of the body depicts the portion of the human brain directly associated with the activity of a particular body part. Simply put, it is the view of the body from the brain's perspective. Areas with lots of sensory innervation, such as the fingertips and the lips, require more cortical area to process finer sensation.



Sensory Homunculus: Cortical Homunculus: A depiction of the human brain areas directly associated with the activity of a particular body part.

This page titled 11.8A: Sensory Areas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.8B: Motor Areas

The motor areas, arranged like a pair of headphones across both cortex hemispheres, are involved in the control of voluntary movements.

Learning Objectives

- Describe the motor areas of the cerebral cortex

Key Points


- The primary motor cortex is involved in the planning of movements.
- The posterior parietal cortex guides movements in space.
- The dorsolateral prefrontal cortex acts as a decision maker for which planned movements will actually be made.
- The basal nuclei receive input from the substantia nigra of the midbrain and motor areas of the cerebral cortex and send signals back to both of these locations.

Key Terms

- **primary motor cortex**: A brain region located in the posterior portion of the frontal lobe of humans. It plans and executes movements in association with other motor areas including the premotor cortex, supplementary motor area, posterior parietal cortex, and several subcortical brain regions.
- **cognitive flexibility**: Ability to switch between thinking about two different concepts and to think about multiple concepts simultaneously.
- **dorsolateral prefrontal cortex**: The highest cortical area responsible for motor planning, organization, and regulation. It plays an important role in the integration of sensory and mnemonic information and the regulation of intellectual function and action.
- **posterior parietal cortex**: Plays an important role in producing planned movements by receiving input from the three sensory systems that help localize the body and external objects in space.

The motor areas of the brain are located in both hemispheres of the cortex. They are positioned like a pair of headphones stretching from ear to ear. The motor areas are very closely related to the control of voluntary movements, especially fine movements performed by the hand. The right half of the motor area controls the left side of the body, and the left half of the motor area controls the right side of the body.

Motor Cortex Divisions



Motor Cortex: Topography of the human motor cortex, including the premotor cortex, SMA, primary motor cortex, primary somatosensory cortex, and posterior parietal cortex.

The motor cortex is divided into three areas:

1. Primary motor cortex: Main contributor to the generation of neural impulses that control the execution of movement.
2. Premotor cortex: Located anterior to the primary motor cortex and responsible for some aspects of motor control.
3. Supplementary motor area (SMA): Functions include internally generated planning of movement, planning of sequences of movement, and the coordination of the two sides of the body. It is located on the midline surface of the hemisphere anterior to the primary motor cortex.

Motor Cortex Functions


Motor functions are also controlled by these additional structures:

- Posterior parietal cortex: Guides planned movements, spatial reasoning, and attention.
- Dorsolateral prefrontal cortex: Important for executive functions, including working memory, cognitive flexibility, and abstract reasoning.

Buried deep in the white matter of the cerebral cortex are interconnected subcortical masses of cerebral gray matter called basal nuclei (or basal ganglia) that are involved in motor control. The basal nuclei receive input from the substantia nigra of the midbrain and motor areas of the cerebral cortex and send signals back to both of these locations.

Motor Cortex Map

The majority of neurons in the motor cortex project to the spinal cord synapse on interneuron circuitry in the spinal cord. The view that each point in the motor cortex controls a muscle or a limited set of related muscles has been debated. Various experiments examining the motor cortex map showed that each point in motor cortex influences a range of muscles and joints, indicating significant overlapping in the map.



Cortex Map: Map of the body in the human brain.

This page titled 11.8B: Motor Areas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.8C: Association Areas

Learning Objectives

- Describe the association areas of the cerebral cortex


Association areas produce a meaningful perceptual experience of the world, enable us to interact effectively, and support abstract thinking and language. The parietal, temporal, and occipital lobes, all located in the posterior part of the cortex, organize sensory information into a coherent perceptual model of our environment centered on our body image. The frontal lobe or prefrontal association complex is involved in planning actions and movement, as well as abstract thought.

Language abilities are localized in the left hemisphere in Broca's area for language expression and Wernicke's area for language reception. The association areas are organized as distributed networks, and each network connects areas distributed across widely spaced regions of the cortex. Distinct networks are positioned adjacent to one another, yielding a complex series of interwoven networks. In humans, association networks are particularly important to language function.

The processes of language expression and reception occur in areas other than just the perisylvian structures such as the prefrontal lobe, basal ganglia, cerebellum, pons, caudate nucleus, and others. The association areas integrate information from different receptors or sensory areas and relate the information to past experiences. Then the brain makes a decision and sends nerve impulses to the motor areas to elicit responses.

Methods of Brain Function Analysis

Behavioral and neuroscientific methods are used to get a better understanding of how our brain influences the way we think, feel, and act. Many different methods help us analyze the brain and give an overview of the relationship between brain and behavior. This promotes understanding of the ways in which associations are made by multiple brain regions, allowing the appropriate responses to occur in a given situation. Well-known techniques are EEG (electroencephalography), which records the brain's electrical activity, and fMRI (functional magnetic resonance imaging), which tells us more about brain functions. Other methods, such as the lesion method, are not as well-known, but still very influential in modern neuroscientific research.



Cortical Areas of the Brain: Locations of brain areas historically associated with language processing. Associated cortical regions involved in vision, touch sensation, and non-speech movement are also shown.

In the lesion method, patients with brain damage are examined to determine which brain structures were damaged and to what

extent this influences the patient's behavior. The concept of the lesion method is based on the idea of finding a correlation between a specific brain area and an occurring behavior. From experiences and research observations, it can be concluded that damage to part of the brain causes behavioral changes or interferes in performing a specific task.

For example, a patient with a lesion in the parietal-temporal-occipital association area has an agraphia, which means he is unable to write although he has no deficits in motor skills. Consequently, researchers deduce that if structure X is damaged and changes in behavior Y occur, X has a relation to Y.

Key Points

- Many areas of the brain are required to form a cohesive view of the world and permit perception.
- The prefrontal association cortex is involved in planning actions and abstract thought.
- The association areas integrate information from different receptors or sensory areas and relate the information to past experiences. Then the brain makes a decision and sends nerve impulses to the motor areas to generate responses.

Key Terms

- **Wernicke's area**: The posterior section of the superior temporal gyrus in the dominant cerebral hemisphere, one of two parts of the cerebral cortex linked with speech (the other being Broca's area).
- **prefrontal association complex**: A region of the brain located in the frontal lobe that is involved in planning actions and movement, as well as abstract thought.
- **agraphia**: An acquired neurological disorder causing a loss in the ability to communicate through writing.
- **Broca's area**: A region in the frontal lobe of the dominant hemisphere (usually the left) of the hominid brain with functions linked to speech production.


This page titled 11.8C: Association Areas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

11.8D: Hemispheric Lateralization

Learning Objectives

- Describe the impact of hemispheric lateralization on brain function

A longitudinal fissure separates the human brain into two distinct cerebral hemispheres connected by the corpus callosum. The two sides resemble each other and each hemisphere's structure is generally mirrored by the other side. Yet despite the strong anatomical similarities, the functions of each cortical hemisphere are distinct.



The hemispheres of the cerebral cortex: The human brain is divided into two hemispheres–left and right. Scientists continue to explore how some cognitive functions tend to be dominated by one side or the other; that is, how they are lateralized.

Broad generalizations are often made in popular psychology about one hemisphere having a broad label, such as "logical" for the left side or "creative" for the right. But although measurable lateral dominance occurs, most functions are present in both hemispheres. The extent of specialization by hemisphere remains under investigation. If a specific region of the brain or even an entire hemisphere is either injured or destroyed, its functions can sometimes be taken over by a neighboring region even in the opposite hemisphere, depending upon the area damaged and the patient's age. When injury interferes with pathways from one area to another, alternative (indirect) connections may develop to communicate information with detached areas, despite the inefficiencies.

While many functions are lateralized, this is only a tendency. The implementation of a specific brain function significantly varies by individual. The areas of exploration of this causal or effectual difference of a particular brain function include gross anatomy, dendritic structure, and neurotransmitter distribution. The structural and chemical variance of a particular brain function, between the two hemispheres of one brain or between the same hemisphere of two different brains, is still being studied. Short of having a hemispherectomy (removal of a cerebral hemisphere), no one is a "left-brain only" or "right-brain only" person.

Lateralization and Handedness

Brain function lateralization is evident in the phenomena of right- or left-handedness, but a person's preferred hand is not a clear indication of the location of brain function. Although 95% of right-handed people have left-hemisphere dominance for language, 18.8% of left-handed people have right-hemisphere dominance for language function. Additionally, 19.8% of left-handed people have bilateral language functions. Even within various language functions (e.g., semantics, syntax, prosody), degree and even hemisphere of dominance may differ.

Language functions such as grammar, vocabulary and literal meaning are typically lateralized to the left hemisphere, especially in right-handed individuals. While language production is left-lateralized in up to 90% of right-handed subjects, it is more bilateral or even right-lateralized in approximately 50% of left-handers. In contrast, prosodic language functions, such as intonation and accentuation, often are lateralized to the right hemisphere of the brain.

Further Lateral Distinctions

The processing of visual and auditory stimuli, spatial manipulation, facial perception, and artistic ability are represented bilaterally, but may show right-hemisphere dominance. Numerical estimation, comparison, and online calculation depend on bilateral parietal regions. Exact calculation and fact retrieval are associated with left parietal regions, perhaps due to their ties to linguistic processing. Dyscalculia is a neurological syndrome associated with damage to the left temporoparietal junction. This syndrome is associated with poor numeric manipulation, poor mental arithmetic skill, and the inability to understand or apply mathematical concepts.

Lateralization and Evolution

Specialization of the two hemispheres is general in vertebrates including fish, frogs, reptiles, birds, and mammals, with the left hemisphere specialized to categorize information and control routine behavior. The right hemisphere is responsible for responses to novel events and behavior in emergencies, including the expression of intense emotions. Feeding is an example of a routine left-hemisphere behavior, while escape from predators is an example of a right-hemisphere behavior. This suggests that the evolutionary advantage of lateralization comes from the capacity to perform separate parallel tasks in each hemisphere of the brain.

Split-Brain Phenomenon

Patients with split-brain are individuals who have undergone corpus callosotomy, a severing of a large part of the corpus callosum (usually as a treatment for severe epilepsy). The corpus callosum connects the two hemispheres of the brain and allows them to communicate. When these connections are cut, the two halves of the brain have a reduced capacity to communicate with each other.

The widespread lateralization of many vertebrate animals indicates an evolutionary advantage associated with the specialization of each hemisphere. The evolutionary advantage of lateralization comes from the capacity to perform separate parallel tasks in each hemisphere of the brain. In a 2011 study published in the journal of *Brain Behavioral Research,* lateralization of a few specific functions as opposed to overall brain lateralization was correlated with parallel tasks efficiency.

Key Points

- The corpus collosum connects the hemispheres of the brain.
- Lateralization of function between the two hemispheres does occur but after injury, other regions of cortex can often compensate.
- There is no such thing as being left-brained or right-brained.
- Functional lateralization often varies between individuals.

Key Terms

- **corpus collosum**: A wide, flat bundle of neural fibers beneath the cortex that connects the left and right cerebral hemispheres and facilitates interhemispheric communication.
- **lateralization**: Localization of a function such as speech to the right or left side of the brain.
- **hemisphere**: Either of the two halves of the cerebrum..
- **prosody**: Properties of syllables and larger units of speech that contribute to linguistic functions such as intonation, tone, stress, and rhythm.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Calcarine Sulcus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Calcarine_sulcus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cerebral cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebra...%23Connections. **License**: *CC BY-SA: Attribution-ShareAlike*
- primary visual cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/primary...isual%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- primary somatosensory cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/primary...nsory%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- primary auditory cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/primary...itory%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus.jpg. **Provided by**: Wikipedia Commoons. **Located at**: commons.wikimedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Cognitive Flexibility. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cognitive_flexibility. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cerebral cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebra...%23Connections. **License**: *CC BY-SA: Attribution-ShareAlike*
- primary motor cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/primary%20motor%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsolateral prefrontal cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsolateral%20prefrontal%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- posterior parietal cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/posterior%20parietal%20cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus.jpg. **Provided by**: Wikipedia Commoons. **Located at**: commons.wikimedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Agraphia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Agraphia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Broca's Area. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Broca%27s_area. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cerebral cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebral_cortex%23Connections. **License**: *CC BY-SA: Attribution-ShareAlike*
- Wernickes area. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wernickes_area. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cognitive Psychology and Cognitive Neuroscience/Behavioural and Neuroscience Methods. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Cognitive_Psychology_and_Cognitive_Neuroscience/Behavioural_and_Neuroscience_Methods. **License**: *CC BY-SA: Attribution-ShareAlike*
- prefrontal association complex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/prefron...tion%20complex. **License**: *CC BY-SA: Attribution-ShareAlike*
- percept. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/percept. **License**: *CC BY-SA: Attribution-ShareAlike*
- Wernicke's area. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Wernicke's_area. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus.jpg. **Provided by**: Wikipedia Commoons. **Located at**: commons.wikimedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*

- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Brain. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_brain. **License**: *CC BY-SA: Attribution-ShareAlike*
- Prosody. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Prosody_(linguistics). **License**: *CC BY-SA: Attribution-ShareAlike*
- Lateralization of brain function. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lateralization_of_brain_function. **License**: *CC BY-SA: Attribution-ShareAlike*
- corpus collosum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/corpus%20collosum. **License**: *CC BY-SA: Attribution-ShareAlike*
- hemisphere. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hemisphere. **License**: *CC BY-SA: Attribution-ShareAlike*
- lateralization. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lateralization. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus.jpg. **Provided by**: Wikipedia Commoons. **Located at**: commons.wikimedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Motor Cortex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Brain. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_brain. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cerebral lobes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Cerebral_lobes.png. **License**: *Public Domain: No Known Copyright*

This page titled 11.8D: Hemispheric Lateralization is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12: PERIPHERAL NERVOUS SYSTEM

- Topic hierarchy

12.10: Reflexes

12.10A: Components of a Reflex Arc

12.10C: Spinal Reflexes

12.11: Pain

12.11A: Pain Sensation

12.11B: Localization of Pain

12.1: Sensation

12.1A: Overview of Sensation

12.1B: Sensation to Perception

12.1C: Sensory Modalities

12.2: Sensory Receptors

12.2A: Classification of Receptors by Stimulus

12.2B: Classification of Receptors by Location

12.3: The Somatosensory System

12.3A: General Organization of the Somatosensory System

12.3B: Tactile Sensation

12.3C: Proprioceptive Sensations

12.3D: Somatic Sensory Pathways

12.3E: Mapping the Primary Somatosensory Area

12.3F: Somatic Sensory Pathways to the Cerebellum

12.4: Nerves

12.4A: Structure of a Nerve

12.4B: Classification of Nerves

12.5: Cranial Nerves

12.5A: Brief Overview of Cranial Nerves

12.5B: Olfactory (I) Nerve

12.5C: Optic (II) Nerve

12.5D: Oculomotor (III) Nerve

12.5E: Trochlear (IV) Nerve

12.5F: Trigeminal (V) Nerve

12.5G: Abducens (VI) Nerve

12.5H: Facial (VII) Nerve

12.5I: Vestibulocochlear (VIII) Nerve
12.5J: Glossopharyngeal (IX) Nerve
12.5K: Vagus (X) Nerve
12.5L: Accessory (XI) Nerve
12.5M: Hypoglossal (XII) Nerve

12.6: Spinal Nerves

12.6A: Overview of the Spinal Nerves
12.6B: Branches of Spinal Nerves
12.6C: Plexuses
12.6D: Intercostal Nerves
12.6E: Dermatomes
12.6F: Function and Physiology of the Spinal Nerves

12.7: Distribution of Spinal Nerves

12.7A: Cervical Plexus
12.7B: Brachial Plexus
12.7C: Lumbar Plexus
12.7D: Sacral and Coccygeal Plexuses
12.7E: Sensory and Motor Tracts

12.8: Motor Activity

12.8A: Peripheral Motor Endings
12.8B: Overview of Motor Integration

12.9: Motor Pathways

12.9A: Organization of Motor Neuron Pathways
12.9B: The Role of the Basal Ganglia in Movement
12.9C: Modulation of Movement by the Cerebellum
12.9D: Functions of the Cerebellum in Integrating Movements

This page titled 12: Peripheral Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.10: REFLEXES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

12.10A: Components of a Reflex Arc

12.10C: Spinal Reflexes

This page titled 12.10: Reflexes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.10A: Components of a Reflex Arc

A reflex arc defines the pathway by which a reflex travels—from the stimulus to sensory neuron to motor neuron to reflex muscle movement.

Learning Objectives

- Describe the components of a reflex arc

Key Points

- Reflexes, or reflex actions, are involuntary, almost instantaneous movements in response to a specific stimulus.
- Reflex arcs that contain only two neurons, a sensory and a motor neuron, are considered monosynaptic. Examples of monosynaptic reflex arcs in humans include the patellar reflex and the Achilles reflex.
- Most reflex arcs are polysynaptic, meaning multiple interneurons (also called relay neurons) interface between the sensory and motor neurons in the reflex pathway.

Key Terms

- **motor neuron**: A neuron located in the central nervous system that projects its axon outside the CNS and directly or indirectly control muscles.
- **sensory neuron**: These are typically classified as the neurons responsible for converting various external stimuli that come from the environment into corresponding internal stimuli.
- **reflex arc**: A neural pathway that controls an action reflex. In higher animals, most sensory neurons do not pass directly into the brain, but synapse in the spinal cord. This characteristic allows reflex actions to occur relatively quickly by activating spinal motor neurons without the delay of routing signals through the brain, although the brain will receive sensory input while the reflex action occurs. There are two types of reflex arcs: autonomic reflex arc (affecting inner organs) and somatic reflex arc (affecting muscles).


Description

A reflex action, also known as a reflex, is an involuntary and nearly instantaneous movement in response to a stimulus. When a person accidentally touches a hot object, they automatically jerk their hand away without thinking. A reflex does not require any thought input.

The path taken by the nerve impulses in a reflex is called a reflex arc. In higher animals, most sensory neurons do not pass directly into the brain, but synapse in the spinal cord. This characteristic allows reflex actions to occur relatively quickly by activating spinal motor neurons without the delay of routing signals through the brain, although the brain will receive sensory input while the reflex action occurs.

Most reflex arcs involve only three neurons. The stimulus, such as a needle stick, stimulates the pain receptors of the skin, which initiate an impulse in a sensory neuron. This travels to the spinal cord where it passes, by means of a synapse, to a connecting neuron called the relay neuron situated in the spinal cord.

The relay neuron in turn makes a synapse with one or more motor neurons that transmit the impulse to the muscles of the limb causing them to contract and pull away from the sharp object. Reflexes do not require involvement of the brain, although in some cases the brain can prevent reflex action.



Reflex arc: The path taken by the nerve impulses in a reflex is called a reflex arc. This is shown here in response to a pin in the paw of an animal, but it is equally adaptable to any situation and animal (including humans).

Types of Reflex Arcs

There are two types of reflex arcs:the autonomic reflex arc, affecting inner organs, and the somatic reflex arc, affecting muscles. When a reflex arc consists of only two neurons, one sensory neuron, and one motor neuron, it is defined as monosynaptic.

Monosynaptic refers to the presence of a single chemical synapse. In the case of peripheral muscle reflexes (patellar reflex, achilles reflex), brief stimulation to the muscle spindle results in the contraction of the agonist or effector muscle.

By contrast, in polysynaptic reflex arcs, one or more interneurons connect afferent (sensory) and efferent (motor) signals.
For example, the withdrawal reflex (nociceptive or flexor withdrawal reflex) is a spinal reflex intended to protect the body from damaging stimuli. It causes the stimulation of sensory, association, and motor neurons.

This page titled 12.10A: Components of a Reflex Arc is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.10C: Spinal Reflexes

Learning Objectives

- Distinguish between the types of spinal reflexes

Spinal reflexes include the stretch reflex, the Golgi tendon reflex, the crossed extensor reflex, and the withdrawal reflex.

Stretch Reflex

The stretch reflex (myotatic reflex) is a muscle contraction in response to stretching within the muscle. This reflex has the shortest latency of all spinal reflexes. It is a monosynaptic reflex that provides automatic regulation of skeletal muscle length.

When a muscle lengthens, the muscle spindle is stretched and its nerve activity increases. This increases alpha motor neuron activity, causing the muscle fibers to contract and thus resist the stretching. A secondary set of neurons also causes the opposing muscle to relax. The reflex functions to maintain the muscle at a constant length.

Golgi Tendon Reflex

The Golgi tendon reflex is a normal component of the reflex arc of the peripheral nervous system. The tendon reflex operates as a feedback mechanism to control muscle tension by causing muscle relaxation before muscle force becomes so great that tendons might be torn.

Although the tendon reflex is less sensitive than the stretch reflex, it can override the stretch reflex when tension is great, making you drop a very heavy weight, for example. Like the stretch reflex, the tendon reflex is ipsilateral.

The sensory receptors for this reflex are called Golgi tendon receptors, and lie within a tendon near its junction with a muscle. In contrast to muscle spindles, which are sensitive to changes in muscle length, tendon organs detect and respond to changes in muscle tension that are caused by a passive stretch or muscular contraction.

Crossed Extensor Reflex



Jendrassik maneuver: The Jendrassik maneuver is a medical maneuver wherein the patient flexes both sets of fingers into a hook-like form and interlocks those sets of fingers together (note the hands of the patient in the chair). This maneuver is used often when testing the patellar reflex, as it forces the patient to concentrate on the interlocking of the fingers and prevents conscious inhibition or influence of the reflex.

The crossed extensor reflex is a withdrawal reflex. The reflex occurs when the flexors in the withdrawing limb contract and the extensors relax, while in the other limb, the opposite occurs. An example of this is when a person steps on a nail, the leg that is stepping on the nail pulls away, while the other leg takes the weight of the whole body.

The crossed extensor reflex is contralateral, meaning the reflex occurs on the opposite side of the body from the stimulus. To produce this reflex, branches of the afferent nerve fibers cross from the stimulated side of the body to the contralateral side of the spinal cord. There, they synapse with interneurons, which in turn, excite or inhibit alpha motor neurons to the muscles of the contralateral limb.


Withdrawal Reflex

The withdrawal reflex (nociceptive or flexor withdrawal reflex) is a spinal reflex intended to protect the body from damaging stimuli. It is polysynaptic, and causes the stimulation of sensory, association, and motor neurons.

When a person touches a hot object and withdraws his hand from it without thinking about it, the heat stimulates temperature and danger receptors in the skin, triggering a sensory impulse that travels to the central nervous system. The sensory neuron then synapses with interneurons that connect to motor neurons. Some of these send motor impulses to the flexors to allow withdrawal.

Some motor neurons send inhibitory impulses to the extensors so flexion is not inhibited—this is referred to as reciprocal innervation. Although this is a reflex, there are two interesting aspects to it:

1. The body can be trained to override that reflex.
2. An unconscious body (or even drunk or drugged bodies) will not exhibit the reflex.



Golgi tendon organ: The Golgi tendon organ, responsible for the Golgi tendon reflex, is diagrammed with its typical position in a muscle (left), neuronal connections in spinal cord (middle), and expanded schematic (right). The tendon organ is a stretch receptor that signals the amount of force on the muscle and protects the muscle from excessively heavy loads by causing the muscle to relax and drop the load.

Key Points

- The stretch reflex is a monosynaptic reflex that regulates muscle length through neuronal stimulation at the muscle spindle.

The alpha motor neurons resist stretching by causing contraction, and the gamma motor neurons control the sensitivity of the reflex.
- The stretch and Golgi tendon reflexes work in tandem to control muscle length and tension. Both are examples of ipsilateral reflexes, meaning the reflex occurs on the same side of the body as the stimulus.
- The crossed extensor reflex is a contralateral reflex that allows the body to compensate on one side for a stimulus on the other. For example, when one foot steps on a nail, the crossed extensor reflex shifts the body's weight onto the other foot, protecting and withdrawing the foot on the nail.
- The withdrawal reflex and the more-specific pain withdrawal reflex involve withdrawal in response to a stimulus (or pain). When pain receptors, called nociceptors, are stimulated, reciprocal innervations stimulate the flexors to withdraw and inhibit the extensors to ensure they are unable to prevent flexion and withdrawal.

Key Terms

- **golgi tendon reflex**: A normal component of the reflex arc of the peripheral nervous system. In this reflex, a skeletal muscle contraction causes the agonist muscle to simultaneously lengthen and relax. This reflex is also called the inverse myotatic reflex because it is the inverse of the stretch reflex. Although muscle tension is increasing during the contraction, the alpha motor neurons in the spinal cord that supply the muscle are inhibited. However, antagonistic muscles are activated.
- **alpha motor neuron**: These are large, lower motor neurons of the brainstem and spinal cord. They innervate the extrafusal muscle fibers of skeletal muscle and are directly responsible for initiating their contraction. Alpha motor neurons are distinct from gamma motor neurons that innervate the intrafusal muscle fibers of muscle spindles.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and Physiology of Animals/Nervous System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy...tem%23Reflexes. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Reflex arc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Reflex_arc. **License**: ***CC BY-SA: Attribution-ShareAlike***
- reflex arc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/reflex%20arc. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sensory neuron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sensory%20neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Withdrawal Reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Withdrawal_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- motor neuron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/motor_neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and physiology of animals A reflex arc. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...reflex_arc.jpg. **License**: ***CC BY: Attribution***
- Pain withdrawal reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pain_withdrawal_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Crossed extensor reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Crossed_extensor_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Golgi tendon reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Golgi_tendon_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Withdrawal reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Withdrawal_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Stretch reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stretch_reflex. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alpha motor neuron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/alpha%20motor%20neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***

- golgi tendon reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/golgi%20tendon%20reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- List of reflexes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/List_of_reflexes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and physiology of animals A reflex arc. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_A_reflex_arc.jpg**. **License**: ***CC BY: Attribution***
- Tendon organ model. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Tendon_organ_model.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gehuchten Jendrassik maneuver. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gehuchten_Jendrassik_maneuver.JPG**. **License**: ***Public Domain: No Known Copyright***

This page titled 12.10C: Spinal Reflexes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.11: PAIN

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

12.11A: Pain Sensation

12.11B: Localization of Pain

This page titled 12.11: Pain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.11A: Pain Sensation

Pain is an unpleasant sensation caused by the activation of nociceptors by thermal, mechanical, chemical, or other stimuli.

Learning Objectives

- Describe the sensation of pain in the peripheral nervous system

Key Points

- Sleeping or silent nociceptors do not respond to these types of signals, but may respond during inflammation of the surrounding tissue.
- Nociceptors receive and send pain signals through myelinated fast Aδ fibers and nonmyelinated slow C fibers that are only activated with intense or prolonged input.
- Nociceptive pain may also be divided into visceral, deep somatic, and superficial somatic pain.
- Neuropathic pain is caused by damage to the nervous system.
- Phantom pain is pain perceived as from amputated or paralyzed limbs.
- Psychogenic pain is caused or exacerbated by mental, emotional, and behavioral factors.

Key Terms

- **visceral**: Relating to the internal organs.
- **ischemia**: A restriction in blood supply to tissues, causing a shortage of oxygen and glucose needed for cellular metabolism.
- **pain**: An ache or bodily suffering, or an instance of this; an unpleasant sensation, resulting from a derangement of functions, disease, or injury by violence; hurt.
- **nociceptor**: A sensory receptor that sends signals that cause the perception of pain in response to a potentially damaging stimulus.

Overview

Pain is an unpleasant feeling often caused by intense or damaging stimuli, such as stubbing a toe, burning a finger, putting alcohol on a cut, and bumping the funny bone. The International Association for the Study of Pain's widely used definition states: "Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage."

Pain motivates the individual to withdraw from damaging situations, to protect a damaged body part while it heals, and to avoid similar experiences in the future. Most pain resolves promptly once the painful stimulus is removed and the body has healed, but sometimes pain persists despite removal of the stimulus and apparent healing of the body. Pain sometimes arises in the absence of any detectable stimulus, damage, or disease.


Nociceptors

A nociceptor is a sensory receptor that responds to potentially damaging stimuli by sending nerve signals to the spinal cord and brain. This process, called nociception, usually causes the perception of pain. There are several types and functions of nociceptors:

- Thermal nociceptors are activated by noxious heat or cold at various temperatures.
- Mechanical nociceptors respond to excess pressure or mechanical deformation. They also respond to incisions that break the skin surface. These mechanical nociceptors frequently have polymodal characteristics. So it is possible that some of the transducers for thermal stimuli are the same for mechanical stimuli.
- Chemical nociceptors respond to a wide variety of spices commonly used in cooking. The one that sees the most response

and is very widely tested is capsaicin. Other chemical stimulants are environmental irritants like acrolein, a World War I chemical weapon and a component of cigarette smoke. Besides these external stimulants, chemical nociceptors have the capacity to detect endogenous ligands, and certain fatty acid amines that arise from changes in internal tissues.

The peripheral terminal of the mature nociceptor is where the noxious stimuli are detected and transduced into electrical energy. When the electrical energy reaches a threshold value, an action potential is induced and driven towards the central nervous system. This leads to the train of events that allows for the conscious awareness of pain.



Gray matter in the spinal cord: A delta fibers (Aδ fibers), a type of sensory fiber, are associated with the sensation of cold and pressure. Aδ fibers are thinly myelinated; therefore, they conduct signals more rapidly than unmyelinated C fibers, but more slowly than other, more thickly myelinated “A” class fibers. Aδ fibers terminate at Rexed laminae I and V (labeled I and V in the diagram). C fibers respond to thermal, mechanical, and chemical stimuli and terminate at the Rexed lamina II (labeled II in the diagram).

Nociceptors have two different types of axons.

1. The Aδ fiber axons are myelinated and can allow an action potential to travel at a rate of about 20 meters/second towards the central nervous system.
2. The other type is the more slowly conducting C fiber axons. These only conduct at speeds of around 2 meters/second. This is due to the light or nonmyelination of the axon.

Phases of Pain

Pain comes in two phases. The first phase is mediated by the fast-conducting Aδ fibers, and the second part is due to C fibers. The pain associated with the Aδ fibers can be associated to an initial extremely sharp pain.

The second phase is a more prolonged and slightly less intense feeling of pain as a result of the damage. If there is massive or prolonged input to a C fiber, there is a progressive build up in the spinal cord dorsal horn; this phenomenon is similar to tetanus in muscles but is called wind-up. If wind-up occurs, there is a probability of increased sensitivity to pain.

Although each nociceptor can have a variety of possible threshold levels, some do not respond at all to chemical, thermal, or mechanical stimuli unless injury actually has occurred. These are typically referred to as silent or sleeping nociceptors since their response comes only at the onset of inflammation to the surrounding tissue.

Types of Nociceptive Pain

Nociceptive pain can be divided into visceral, deep somatic and superficial somatic pain.

1. Visceral structures are highly sensitive to stretch, ischemia, and inflammation, but relatively insensitive to other stimuli that normally evoke pain in other structures, such as burning and cutting. Visceral pain is diffuse, difficult to locate, and often referred to a distant, usually superficial, structure. It may be accompanied by nausea and vomiting and may be described as sickening, deep, squeezing, and dull.
2. Deep somatic pain is initiated by stimulation of nociceptors in ligaments, tendons, bones, blood vessels, fasciae and muscles, and is a dull, aching, poorly localized pain. Examples include sprains and broken bones.
3. Superficial pain is initiated by the activation of nociceptors in the skin or other superficial tissue, and is sharp, well-defined, and clearly located. Examples of injuries that produce superficial somatic pain include minor wounds and minor (first degree) burns.

This page titled 12.11A: Pain Sensation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.11B: Localization of Pain

Localization of pain is determined by whether the pain is superficial somatic, visceral, or deep somatic.

Learning Objectives

- Describe the localization of pain in the peripheral nervous system

Key Points

- Nociceptive pain is caused by stimulation of peripheral nerve fibers that respond only to stimuli approaching or exceeding harmful intensity (nociceptors).
- Nociceptive pain may be classified according to the mode of noxious stimulation, the most common categories being thermal (heat or cold), mechanical (crushing, tearing, etc.), and chemical (iodine in a cut, chili powder in the eyes).
- Superficial pain is initiated by activation of nociceptors in the skin or other superficial tissue, and is sharp, well-defined, and clearly located.
- Visceral pain is diffuse and difficult to locate; deep somatic pain is dull aching and difficult to locate; and superficial somatic pain is sharp, well-defined, and easily located.
- Deep somatic pain is initiated by stimulation of nociceptors in ligaments, tendons, bones, blood vessels, fasciae and muscles, and is dull, aching, poorly localized pain.
- Referred pain is characterized by pain felt in a location away from the site of the painful stimulus.

Key Terms

- **visceral pain**: Visceral pain is diffuse, difficult to locate, and often referred to a distant, usually superficial, structure. It may be accompanied by nausea and vomiting and may be described as sickening, deep, squeezing, and dull.
- **deep somatic pain**: Pain initiated by stimulation of nociceptors in ligaments, tendons, bones, blood vessels, fasciae and muscles; it is dull, aching, poorly localized pain. Examples include sprains and broken bones.
- **superficial somatic pain**: Pain initiated by activation of nociceptors in the skin or other superficial tissue; it is sharp, well-defined, and clearly located. Examples of injuries that produce superficial somatic pain include minor wounds and minor (first degree) burns.

Pain Categories

Nociceptive pain is caused by stimulation of peripheral nerve fibers that respond only to stimuli approaching or exceeding harmful intensity (nociceptors), and may be classified according to the mode of noxious stimulation.

The most common categories are thermal (heat or cold), mechanical (crushing, tearing, etc.), and chemical (iodine in a cut, chili powder in the eyes). Nociceptive pain may also be divided into visceral, deep somatic, and superficial somatic pain.

Visceral Pain

Visceral structures are highly sensitive to stretch, ischemia, and inflammation, but relatively insensitive to other stimuli that normally evoke pain in other structures, such as burning and cutting. Visceral pain is diffuse, difficult to locate, and often referred to a distant, usually superficial, structure. It may be accompanied by nausea and vomiting and may be described as sickening, deep, squeezing, and dull.

Deep Somatic Pain

Deep somatic pain is initiated by the stimulation of nociceptors in ligaments, tendons, bones, blood vessels, fasciae, and muscles,

and is a dull, aching, poorly localized pain. Examples include sprains and broken bones. Superficial pain is initiated by activation of nociceptors in the skin or other superficial tissue, and is sharp, well-defined, and clearly located. Examples of injuries that produce superficial somatic pain include minor wounds and minor (first degree) burns.

Referred Pain


Referred pain (also reflective pain) is pain perceived at a location other than the site of the painful stimulus. An example is the case of ischemia brought on by a myocardial infarction (heart attack), where pain is often felt in the neck, shoulders, and back rather than in the chest, the site of the injury.

The International Association for the Study of Pain has not officially defined referred pain as of 2001; hence, several authors have defined the term differently. Radiation is different from referred pain.


The pain related to a myocardial infarction could either be referred pain or pain radiating from the chest. Classically, the pain associated with a myocardial infarction is located in the middle or left side of the chest, where the heart is actually located. The pain can radiate to the left side of the jaw and into the left arm.

Referred pain occurs when the pain is located away from or adjacent to the organ involved. An example would be when a person has pain only in their jaw or left arm, but not in the chest. The size of referred pain is related to the intensity and duration of ongoing/evoked pain.

Also, temporal summation is a potent mechanism for the generation of referred muscle pain. Central hyperexcitability is also important for the extent of referred pain.



Temporal summation: Temporal summation, shown in the diagram, is the transmitting of signals with increased frequency of impulse, thus increasing the strength of signals in each fiber. Temporal summation is a potent mechanism for the generation of referred muscle pain.



Brain freeze: In an ice cream headache, known colloquially as brain freeze and medically as a cold-stimulus headache, the trigeminal nerve, shown in yellow, conducts signals from dilating blood vessels in the palate to the brain that interpret the pain as coming from the forehead.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Pain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pain%23Classification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pain sensation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pain_se...on%23Mechanism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nociceptors. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nocicep..._and_functions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pain. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nociceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nociceptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ischemia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ischemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medulla spinalis - Substantia grisea - English. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Me..._-_English.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Referred pain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Referred_pain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pain%23Nociceptive**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- visceral pain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/visceral%20pain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medulla spinalis - Substantia grisea - English. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Me..._-_English.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Temporal summation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Te..._summation.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray778 Trigeminal. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gr...Trigeminal.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 12.11B: Localization of Pain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.1: SENSATION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

12.1A: Overview of Sensation

12.1B: Sensation to Perception

12.1C: Sensory Modalities

This page titled 12.1: Sensation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.1A: Overview of Sensation

Sensation refers to our ability to detect or sense the physical qualities of our environment.

Learning Objectives

- Describe what sensation means in terms of the peripheral nervous system

Key Points

- Sensation refers to our ability to detect and sense the internal and external physical qualities of our environment.
- Our senses include both exteroception (stimuli that occur outside of our body) and interoception (stimuli occurring inside of our bodies).
- Our primary senses are considered to be sight, hearing, taste, smell, and touch.
- All senses require one of four fundamental sensory capacities: chemoreception, photoreception, mechanoreception, or thermoreception.
- The peripheral nervous system (PNS) consists of sensory receptors to communicate with other parts of the body.

Key Terms

- **chemoreception**: A physiological response to chemical stimuli.
- **mechanoreception**: A physiological response to mechanical forces like pressure, touch, and vibration.
- **photoreception**: A physiological response to light, as occurs during vision in animals.
- **thermoreception**: A physiological response to relative or absolute changes in temperature.

Our senses can be broadly grouped into exteroception, for the detection of stimuli that occur outside of our body, and interoception, for stimuli occurring inside of our bodies. However, what constitutes a sense is a matter of great debate, leading to difficulties in precisely defining what it is. Traditionally, human beings are considered to have five main senses: sight, hearing, taste, smell, and touch.

The peripheral nervous system (PNS) consists of sensory receptors that extend from the central nervous system (CNS) to communicate with other parts of the body. These receptors respond to changes and stimuli in the environment. Sense organs (made up of sensory receptors and other cells) operate the senses of vision, hearing, equilibrium, smell, and taste.

Sight

Sight or vision (ophthalmoception) is the ability of the eye(s) to focus and detect images of visible light on photoreceptors in the retina that generate electrical nerve impulses for varying colors, hues, and brightness. There are two types of photoreceptors: rods and cones. Rods are very sensitive to light, but do not distinguish colors. Cones distinguish colors, but are less sensitive to dim light. The inability to see is called blindness.

Hearing

Hearing or audition (audioception) is the sense of sound perception. Mechanoreceptors in the inner ear turn vibration motion into electrical nerve pulses. The vibrations are mechanically conducted from the eardrum through a series of tiny bones to hair-like fibers in the inner ear that detect the mechanical motion of the fibers.

Sound can also be detected as vibrations conducted through the body by tactition. The inability to hear is called deafness or hearing impairment.

Taste

Taste (gustaoception) refers to the ability to detect substances such as food, certain minerals, poisons, etc. The sense of taste is often confused with the concept of flavor, which is a combination of taste and smell perception. Flavor depends on odor, texture, and temperature as well as on taste.

Humans receive tastes through sensory organs called taste buds, or gustatory calyculi, concentrated on the upper surface of the tongue. Five basic tastes exist: sweet, bitter, sour, salty, and umami. The inability to taste is called ageusia.

Smell

The olfactory system is the sensory system used for the sense of smell (olfaction). This sense is mediated by specialized sensory cells of the nasal cavity. In humans, olfaction occurs when odorant molecules bind to specific sites on the olfactory receptors in the nasal cavity. These receptors are used to detect the presence of smell. They come together at a structure (the glomerulus) that transmits signals to the olfactory bulb in the brain. The inability to smell is called anosmia.

Touch

Touch or somatosensation (tactioception, tactition, or mechanoreception), is a perception resulting from the activation of neural receptors in the skin, including hair follicles, tongue, throat, and mucosa. A variety of pressure receptors respond to variations in pressure (firm, brushing, sustained, etc.).

The touch sense of itching is caused by insect bites or allergies that involve special itch-specific neurons in the skin and spinal cord. The loss or impairment of the ability to feel anything touched is called tactile anesthesia.

Paresthesia is a sensation of tingling, pricking, or numbness of the skin that may result from nerve damage and may be permanent or temporary.

Additional Senses

Many scientists and philosophers argue that humans have additional senses including:


- Pain or nociception (physiological pain): Signals nerve and other tissue damage.
- Balance or equilibrioception: Allows the sensing of body movement, direction, and acceleration, and to attain and maintain postural equilibrium and balance.
- Body awareness or proprioception: Provides the parietal cortex of the brain with information on the relative positions of the parts of the body.
- Sense of time or chronoception: Refers to how the passage of time is perceived and experienced but is not associated with a specific sensory system. According to psychologists and neuroscientists, however, human brains have a system governing the perception of time.
- Temperature sensing or thermoception: The sensation of heat and the absence of heat (cold).

If interoceptive senses are also considered, sensation can be expanded to include stretch (as in muscles or organs like the lungs), oxygen and carbon dioxide sensing, pH sensing, and more.

While the exact definition of sensation is still controversial, most scientists agree that all senses rely on four fundamental sensory capacities:

1. Chemical detection (chemoreception).
2. Light detection (photoreception).
3. Force detection (mechanoreception).
4. Temperature detection (thermoreception).

Our nervous system has sensory systems and organs that mediate each sense and these systems rely on chemoreceptors, photoreceptors, mechanoreceptors, or thermoreceptors to detect the state of the internal or external environment.



The five senses: Photographic depiction of the five senses.

This page titled 12.1A: Overview of Sensation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.1B: Sensation to Perception

The goal of sensation is detection, while the goal of perception is to create useful information about our environment.

Learning Objectives

- Describe how sensations become perceptions


Key Points

- Sensation is a function of the low-level, biochemical and neurological mechanisms that allow the receptor cells of a sensory organ to detect an environmental stimulus.
- Perception refers to the mental processes that represent understanding of the real-world causes of sensory input.
- Neural signals are transmitted to the brain and processed; the resulting mental recreation of the distal stimulus is the percept. Perception is particularly important to our ability to understand speech. After processing the initial auditory signal, speech sounds are further processed to extract acoustic cues and phonetic information.

Key Terms

- **Perception**: The organization, identification, and interpretation of sensory information in order to construct a mental representation through the process of transduction, during which sensors in the body transform signals from the environment into encoded neural signals.
- **reverberation**: The persistence of sound after a sound is produced (such as an echo).
- **transduction**: The conversion of a stimulus from one form to another.
- **sensation**: The function of the low-level biochemical and neurological events that occur when a stimulus activates the receptor cells of a sensory organ.

Sensation and perception are two distinct stages of processing during human sensing. Sensation is a function of the low level, biochemical, and neurological mechanisms that allow the receptor cells of a sensory organ to detect an environmental stimulus.



Color optical illusion: The brain interprets the pink cube on the light gray background as being a darker shade than the pink cube on the dark gray background.


Perception refers to the mental processes that are reflected in statements like “I see a blue wall” that represent understanding of the real-world causes of sensory input. In other words, the goal of sensation is detection, while the goal of perception is to create useful information about the environment.

Stimuli from the environment (distal stimuli) are transformed into neural signals, which are then interpreted by the brain through a process called transduction. Transduction can be likened to a bridge connecting sensation to perception. This raw pattern of neural activity is called the proximal stimulus.

The neural signals are transmitted to the brain and processed. The resulting mental recreation of the distal stimulus is the percept. The sound stimulating a person’s auditory receptors is the proximal stimulus, and the brain’s interpretation of this as the ringing of a telephone is the percept.

All perception involves signals in the nervous system that result from physical stimulation of the sense organs. For example, vision involves light striking the retinas of the eyes, smell is mediated by odor molecules, and hearing involves sound waves.

However, perception is not the passive receipt of these signals, but is a process of organization, identification, and interpretation. Although the senses were traditionally viewed as passive receptors, the study of illusions and ambiguous images, has demonstrated that the brain’s perceptual systems actively influence sensory systems in an attempt to construct useful representations of our environment.



The Necker cube and Rubin vase: These are two optical illusions that illustrate how perception may differ from reality. On the left, we see a cube when in fact it is a flat image on our screen. On the right, the vase actually resembles two faces looking at each other.

Perception is particularly important to our ability to understand speech. The sound of a word can vary widely according to the words around it and the tempo of the speech, as well as the physical characteristics, accent, and mood of the speaker. Listeners manage to perceive words across this wide range of different conditions. Another variation is that reverberation can make a large difference in sound, such as hearing a word spoken from the far side of a room and the same word spoken up close. The process of perceiving speech begins at the level of the sound within the auditory signal and the process of audition. After processing the initial auditory signal, speech sounds are further processed to extract acoustic cues and phonetic information. This speech information can then be used for higher-level language processes, such as word recognition.

This page titled 12.1B: Sensation to Perception is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.1C: Sensory Modalities

A sensory modality (also called a stimulus modality) is an aspect of a stimulus or what is perceived after a stimulus.

Learning Objectives

- Describe the sensory modalities of the peripheral nervous system

Key Points

- The basic sensory modalities include: light, sound, taste, temperature, pressure, and smell.
- A broadly acceptable definition of a sense is: A system that consists of a group of sensory cell types, responding to a specific physical phenomenon, and corresponding to a particular group of regions within the brain where the signals are received and interpreted.
- Multimodal perception is the ability of the mammalian nervous system to combine different inputs of the sensory system. Nociception (physiological pain) signals nerve damage or damage to tissue. The three types of pain receptors are cutaneous (skin), somatic (joints and bones), and visceral (body organs).
- Proprioception, the kinesthetic sense, provides the parietal cortex of the brain with information on the relative positions of the parts of the body.

Key Terms

- **thermoreception**: A physiological response to relative or absolute changes in temperature.
- **modality**: Also known as stimulus modality, it is one feature of a complex stimulus; for example, temperature, pressure, sound, or taste.
- **utricle**: Stimulates hair cells of the inner ear to detect motion and orientation.
- **saccule**: A bed of sensory cells situated in the inner ear that translates head movements into neural impulses that the brain can interpret.
- **circadian**: Any biological process that displays an endogenous, entrainable oscillation of about 24 hours.
- **ultradian**: A recurrent period or cycle repeated throughout a 24-hour circadian day.
- **mechanoreception**: A physiological response to mechanical forces like pressure, touch, and vibration.
- **bipolar cell**: Specialized sensory neuron for the transmission of special senses.

Sensing

Senses are transducers from the physical world to the realm of the mind. Another broadly acceptable definition of a sense is: A system that consists of a group of sensory cell types, responding to a specific physical phenomenon, and corresponding to a particular group of regions within the brain where the signals are received and interpreted.

Disputes about the number of senses typically arise around classification of the various cell types, and their mapping to regions of the brain.

Sensory Modalities

A sensory modality (also called a stimulus modality) is an aspect of a stimulus or what is perceived after a stimulus. The term sensory modality is often used interchangeably with sense. The basic sensory modalities include: light, sound, taste, temperature, pressure, and smell.

Light Modality

The sensory modality for vision is light. To perceive a light stimulus, the eye must first refract the light so that it directly hits the retina. The transduction of light into neural activity occurs via the photoreceptors in the retina.

When a particle of light hits the photoreceptors of the eye, the photopigment of the photoreceptor undergoes a chemical change leading to a chain of chemical reactions occur. A message is sent to a neuron called the bipolar cell through the use of a nerve impulse. Finally, a message is sent to the ganglion cell and then, finally, the brain.

Sound Modality

The sensory modality for audition is sound. Sound is created through air pressure. A vibrating object compresses the surrounding molecules of air as it moves towards a given point, and expands the molecules as it moves away from the point.

The eardrum is stimulated by vibrations in the air. It collects and sends these vibrations to receptor cells. The ossicles (three tiny bones in the middle ear) pass the vibrations to the fluid-filled cochlea (a spiral, shell-shaped auditory organ of the inner ear). The vibrations move through the liquid in the cochlea where the receptive organ is able to sense it.

Taste Modality

Taste stimuli are encountered by receptor cells located in taste buds on the tongue and pharynx. Receptor cells disseminate onto different neurons and convey the message of a particular taste in a single medullar nucleus.

Taste perception is created by combining multiple sensory inputs. Different modalities help determine the perception of taste.

Temperature Modality

Temperature modality excites or elicits a symptom through cold or hot temperature. The cutaneous somatosensory system detects changes in temperature.

Thermal stimuli from a homeostatic set point excite temperature specific sensory nerves in the skin. Specific thermosensory fibers respond to warmth and to cold.

Pressure Modality

Tactile stimulation can be direct, such as through bodily contact, or indirect, such as through the use of a tool or probe. Tactual perception gives information regarding cutaneous stimuli (pressure, vibration, and temperature), kinesthetic stimuli (limb movement), and proprioceptive stimuli (position of the body).

Smell Modality


The sense of smell is called olfaction. Materials constantly shed molecules, which float into the nose or are taken in through breathing. Inside the nasal chambers is the neuroepithelium lining.

It contains the receptors responsible for detecting molecules that are small enough to smell. These receptor neurons then synapse at the olfactory cranial nerve, which sends the information to the olfactory bulbs in the brain for initial processing.

Multimodal Perception

Multimodal perception is the ability of the mammalian nervous system to combine all of the different inputs of the sensory system to result in an enhanced detection or identification of a particular stimulus.

Integration of all sensory modalities occurs when multimodal neurons receive sensory information that overlaps with different modalities. Multimodal perception comes into effect when a unimodal stimulus fails to produce a response.




Multisensory perception: This is a diagram of how multimodal perception is created by the overlapping and combining of different inputs from the sensory systems.

Additional Senses

Balance (or equilibrioception) is the sense that allows an organism to sense body movement, direction, and acceleration, and also attain and maintain postural equilibrium and balance. The organ of equilibrioception is the vestibular labyrinthine system found in both of the inner ears.

In technical terms, this organ is responsible for two senses: angular momentum and acceleration (known together as equilibrioception). The vestibular nerve conducts information from sensory receptors in three ampulla, each of which sense fluid motion in three semicircular canals caused by a three-dimensional rotation of the head.

The vestibular nerve also conducts information from the utricle and the saccule; these contain hair-like sensory receptors that bend under the weight of otoliths (small crystals of calcium carbonate) that provide the inertia needed to detect head rotation, linear acceleration, and the direction of gravitational force.



Inner ear: Inner ear anatomy showing utricle, saccule, and vestibular nerve.

Thermoception is the sense of heat or absence of heat (cold) by the skin and internal skin passages. Perceiving changes in temperature is referred to as heat flux (the rate of heat flow) in these areas.

There are specialized receptors for cold (declining temperature) and heat. The cold receptors infer wind direction, an important part in the animal's sense of smell. The heat receptors are sensitive to infrared radiation and can occur in specialized organs, for instance in pit vipers.

The thermoceptors in the skin are quite different from the homeostatic thermoceptors in the brain (hypothalamus), which provides feedback on internal body temperature.

Proprioception, the kinesthetic sense, provides the parietal cortex of the brain with information on the relative positions of the parts of the body. Neurologists test this sense by telling patients to close their eyes and touch their own nose with the tip of a finger. Assuming proper proprioceptive function, at no time will the person lose awareness of where their hand actually is, even though it is not being detected by any of the other senses. Proprioception and touch are related in subtle ways, and their impairment results in deep and surprising deficits in perception and action.

Nociception (physiological pain) signals nerve or other tissue damage. The three types of pain receptors are cutaneous (skin), somatic (joints and bones), and visceral (body organs).

It was previously believed that pain was simply the overloading of pressure receptors, but research in the first half of the 20th century showed that pain is a distinct phenomenon that intertwines with all of the other senses, including touch.

Chronoception refers to how the passage of time is perceived and experienced. Although the sense of time is not associated with a specific sensory system, psychological and neuro-scientific research indicates that human brains do have a system governing the perception of time.

It is composed of a highly distributed system involving the cerebral cortex, cerebellum and basal ganglia. One particular component, the suprachiasmatic nucleus, is responsible for the circadian (daily) rhythm, while other cell clusters appear capable of shorter-range (ultradian) timekeeping.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Senses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Senses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Senses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Definition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Senses. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Senses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Other_internal_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Culture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Senses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...hermoreception**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...chanoreception**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/photoreception**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Five senses. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Five_senses.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Perception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perception. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensation (psychology). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensation_(psychology). **License**: *CC BY-SA: Attribution-ShareAlike*
- Perception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perception. **License**: *CC BY-SA: Attribution-ShareAlike*
- Perception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perception. **License**: *CC BY-SA: Attribution-ShareAlike*
- sensation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sensation. **License**: *CC BY-SA: Attribution-ShareAlike*
- transduction. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/transduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Five senses. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Five_senses.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Color optical illusion - pink. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ion_-_pink.jpg. **License**: *Public Domain: No Known Copyright*
- Perception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perception. **License**: *Public Domain: No Known Copyright*
- Sense. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sense. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cochlea. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cochlea. **License**: *CC BY-SA: Attribution-ShareAlike*
- Utradian Rythm. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ultradian_rhythm. **License**: *CC BY-SA: Attribution-ShareAlike*
- Saccule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Saccule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circadian Rythm. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circadian_rhythm. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stimulus Modality. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stimulus_modality. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ossicles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ossicles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensory_system%23Modality. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sense. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sense%23Other_senses. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sense. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sense%23Traditional_senses. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sense. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sense. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensation (psychology). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensation_(psychology). **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/thermoreception. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/mechanoreception. **License**: *CC BY-SA: Attribution-ShareAlike*
- modality. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/modality. **License**: *CC BY-SA: Attribution-ShareAlike*
- Five senses. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Five_senses.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Color optical illusion - pink. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Color_optical_illusion_-_pink.jpg. **License**: *Public Domain: No Known Copyright*
- Perception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perception. **License**: *Public Domain: No Known Copyright*
- Saccule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Saccule. **License**: *CC BY: Attribution*
- Multisensory.jpg. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Multisensory.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 12.1C: Sensory Modalities is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.2: SENSORY RECEPTORS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

12.2A: Classification of Receptors by Stimulus

12.2B: Classification of Receptors by Location

This page titled 12.2: Sensory Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.2A: Classification of Receptors by Stimulus

Sensory receptors are primarily classified as chemoreceptors, thermoreceptors, mechanoreceptors, or photoreceptors.

Learning Objectives

- Differentiate among the types of stimuli to which receptors respond

Key Points


- Chemoreceptors detect the presence of chemicals.
- Thermoreceptors detect changes in temperature.
- Mechanoreceptors detect mechanical forces.
- Photoreceptors detect light during vision.
- More specific examples of sensory receptors are baroreceptors, propioceptors, hygroreceptors, and osmoreceptors.
- Sensory receptors perform countless functions in our bodies mediating vision, hearing, taste, touch, and more.

Key Terms

- **photoreceptor**: A specialized neuron able to detect and react to light.
- **mechanoreceptor**: Any receptor that provides an organism with information about mechanical changes in its environment such as movement, tension, and pressure.
- **baroreceptor**: A nerve ending that is sensitive to changes in blood pressure.

Sensory receptors can be classified by the type of stimulus that generates a response in the receptor. Broadly, sensory receptors respond to one of four primary stimuli:

1. Chemicals (chemoreceptors)
2. Temperature (thermoreceptors)
3. Pressure (mechanoreceptors)
4. Light (photoreceptors)



A schematic of the classes of sensory receptors: Sensory receptor cells differ in terms of morphology, location, and

stimulus.

All sensory receptors rely on one of these four capacities to detect changes in the environment, but may be tuned to detect specific characteristics of each to perform a specific sensory function. In some cases, the mechanism of action for a receptor is not clear. For example, hygroreceptors that respond to changes in humidity and osmoreceptors that respond to the osmolarity of fluids may do so via a mechanosensory mechanism or may detect a chemical characteristic of the environment.

Sensory receptors perform countless functions in our bodies. During vision, rod and cone photoreceptors respond to light intensity and color. During hearing, mechanoreceptors in hair cells of the inner ear detect vibrations conducted from the eardrum. During taste, sensory neurons in our taste buds detect chemical qualities of our foods including sweetness, bitterness, sourness, saltiness, and umami (savory taste). During smell, olfactory receptors recognize molecular features of wafting odors. During touch, mechanoreceptors in the skin and other tissues respond to variations in pressure.

Classification of Sensory Receptors

Adequate Stimulus

Adequate stimulus can be used to classify sensory receptors. A sensory receptor's adequate stimulus is the stimulus modality for which it possesses the adequate sensory transduction apparatus.

Sensory receptors with corresponding stimuli to which they respond.

Receptor	Stimulus
Apmullae of Lorenzini (primarily function as electroreceptors)	Electric fields, salinity, and temperature
Baroreceptors	Pressure in blood vessels
Chemo receptors	Chemical stimuli
Electromagnetic radiation receptors	Electromagnetic radiation
Electroreceptors	Electrofields
Hydroreceptors	Humidity
Infrared receptors	Infrared radiation
Magnetoreceptors	Magnetic fields
Mechanoreceptors	Mechanical stress or strain
Nociceptors	Damage or threat of damage to body tissues (leads to pain perception)
Osmoreceptors	Osmolarity of fluids
Photoreceptors	Visible light
Proprioceptors	Sense of position
Thermoreceptors	Temperature
Ultraviolet receptors	Ultraviolet radiation

Location

Sensory receptors can be classified by location:

- Cutaneous receptors are sensory receptors found in the dermis or epidermis.
- Muscle spindles contain mechanoreceptors that detect stretch in muscles.

Morphology

Somatic sensory receptors near the surface of the skin can usually be divided into two groups based on morphology:

1. Free nerve endings characterize the nociceptors and thermoreceptors.
2. Encapsulated receptors consist of the remaining types of cutaneous receptors. Encapsulation exists for specialized functioning.

Rate of Adaptation

A tonic receptor is a sensory receptor that adapts slowly to a stimulus, while a phasic receptor is a sensory receptor that adapts rapidly to a stimulus.

This page titled 12.2A: Classification of Receptors by Stimulus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.2B: Classification of Receptors by Location

Some sensory receptors can be classified by the physical location of the receptor.

Learning Objectives

- Differentiate among sensory receptors by location

Key Points

- Sensory receptors that share a common location often share a related function.
- Sensory receptors code four aspects of a stimulus: modality (or type), intensity, location, and duration.
- Cutaneous touch receptors and muscle spindle receptors are both mechanoreceptors, but they differ in location.

Key Terms

- **cutaneous touch receptor**: A type of sensory receptor found in the dermis or epidermis of the skin.
- **muscle spindle**: Sensory receptors within the belly of a muscle that primarily detect changes in the length of this muscle.

Types of Receptors

As we exist in the world, our bodies are tasked with receiving, integrating, and interpreting environmental inputs that provide information about our internal and external environments. Our brains commonly receive sensory stimuli from our visual, auditory, olfactory, gustatory, and somatosensory systems.

Remarkably, specialized receptors have evolved to transmit sensory inputs from each of these sensory systems. Sensory receptors code four aspects of a stimulus:


1. Modality (or type)
2. Intensity
3. Location
4. Duration

Receptors are sensitive to discrete stimuli and are often classified by both the systemic function and the location of the receptor.

Sensory receptors are found throughout our bodies, and sensory receptors that share a common location often share a common function. For example, sensory receptors in the retina are almost entirely photoreceptors. Our skin includes touch and temperature receptors, and our inner ears contain sensory mechanoreceptors designed for detecting vibrations caused by sound or used to maintain balance.

Force -sensitive mechanoreceptors provide an example of how the placement of a sensory receptor plays a role in how our brains process sensory inputs. While the cutaneous touch receptors found in the dermis and epidermis of our skin and the muscle spindles that detect stretch in skeletal muscle are both mechanoreceptors, they serve discrete functions.

In both cases, the mechanoreceptors detect physical forces that result from the movement of the local tissue, cutaneous touch receptors provide information to our brain about the external environment, while muscle spindle receptors provide information about our internal environment.



Muscle spindle: Mammalian muscle spindle showing typical position in a muscle (left), neuronal connections in spinal cord (middle), and an expanded schematic (right). The spindle is a stretch receptor with its own motor supply consisting of several intrafusal muscle fibers. The sensory endings of a primary (group Ia) afferent and a secondary (group II) afferent coil around the non-contractile central portions of the intrafusal fibers.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...itional_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...itional_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Other_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Other_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Other_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...itional_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory receptors. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory...quate_stimulus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...nternal_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%23Other_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...itional_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory receptors. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory...quate_stimulus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sense. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sense%2...itional_senses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- photoreceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/photoreceptor**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- baroreceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/baroreceptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mechanoreceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mechanoreceptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Original figure by Stacy Matthews Branch. Licensed CC BY-SA 4.0. **Provided by**: Stacy Matthews Branch. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Structure of sensory system (4 models) E. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi..._models)_E.PNG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hair cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hair_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory...s_and_function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory...%23by_location**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cutaneous touch receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cutaneo...uch%20receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- muscle spindle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/muscle%20spindle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Original figure by Stacy Matthews Branch. Licensed CC BY-SA 4.0. **Provided by**: Stacy Matthews Branch. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Structure of sensory system (4 models) E. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi..._models)_E.PNG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle spindle model. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mu...ndle_model.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 12.2B: Classification of Receptors by Location is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3: THE SOMATOSENSORY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

12.3A: General Organization of the Somatosensory System
12.3B: Tactile Sensation
12.3C: Proprioceptive Sensations
12.3D: Somatic Sensory Pathways
12.3E: Mapping the Primary Somatosensory Area
12.3F: Somatic Sensory Pathways to the Cerebellum

This page titled 12.3: The Somatosensory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3A: General Organization of the Somatosensory System

The somatosensory system is composed of the neurons that make sensing touch, temperature, and position in space possible.

Learning Objectives

- Describe how the somatosensory system is composed of neurons that make sensing touch, temperature, and position in space possible

Key Points

- The somatosensory system consists of primary, secondary, and tertiary neurons.
- Sensory receptors housed in the dorsal root ganglia project to secondary neurons of the spinal cord that decussate and project to the thalamus or cerebellum.
- Tertiary neurons project to the postcentral gyrus of the parietal lobe, forming a sensory homunculus.
- A sensory homunculus maps sub-regions of the cortical postcentral gyrus to certain parts of the body.

Key Terms

- **decussate**: Where nerve fibers obliquely cross from one lateral part of the body to the other.
- **postcentral gyrus**: A prominent structure in the parietal lobe of the human brain and an important landmark that is the location of the primary somatosensory cortex, the main sensory receptive area for the sense of touch.
- **organization**: The quality of being constituted of parts, each having a special function, act, office, or relation; to systematize.
- **thalamus**: Either of two large, ovoid structures of gray matter within the forebrain that relay sensory impulses to the cerebral cortex.

The somatosensory system is distributed throughout all major parts of our body. It is responsible for sensing touch, temperature, posture, limb position, and more. It includes both sensory receptor neurons in the periphery (eg., skin, muscle, and organs) and deeper neurons within the central nervous system.

Structure

A somatosensory pathway will typically consist of three neurons: primary, secondary, and tertiary.

1. In the periphery, the primary neuron is the sensory receptor that detects sensory stimuli like touch or temperature. The cell body of the primary neuron is housed in the dorsal root ganglion of a spinal nerve or, if sensation is in the head or neck, the ganglia of the trigeminal or cranial nerves.
2. The secondary neuron acts as a relay and is located in either the spinal cord or the brainstem. This neuron's ascending axons will cross, or decussate, to the opposite side of the spinal cord or brainstem and travel up the spinal cord to the brain, where most will terminate in either the thalamus or the cerebellum.
3. Tertiary neurons have cell bodies in the thalamus and project to the postcentral gyrus of the parietal lobe, forming a sensory homunculus in the case of touch. Regarding posture, the tertiary neuron is located in the cerebellum.


Processing

The primary somatosensory area of the human cortex is located in the postcentral gyrus of the parietal lobe. The postcentral gyrus is the location of the primary somatosensory area, the area of the cortex dedicated to the processing of touch information. At this location there is a map of sensory space referred to as a sensory homunculus.

A cortical homunculus is the brain's physical representation of the human body; it is a neurological map of the anatomical divisions of the body. The surface area of cortex dedicated to a body part correlates with the amount of somatosensory input

from that area.

For example, there is a large area of cortex devoted to sensation in the hands, while the back requires a much smaller area. Somatosensory information involved with proprioception and posture is processed in the cerebellum.



Homunculus: Image representing the cortical sensory homunculus. It shows how the anatomical portions of the body, such as the tongue, elbow, and hip, are mapped out on the homonculus. The surface area of cortex dedicated to a body part correlates with the amount of somatosensory input from that area.

Functions

The somatosensory system functions in the body's periphery, spinal cord, and the brain.

- Periphery: Sensory receptors (i.e., thermoreceptors, mechanoreceptors, etc.) detect the various stimuli.
- Spinal cord: Afferent pathways in the spinal cord serve to pass information from the periphery and the rest of the body to the brain.
- Brain: The postcentral gyrus contains Brodmann areas (BA) 3a, 3b, 1, and 2 that make up the somatosensory cortex. BA3a is involved with the sense of relative position of neighboring body parts and the amount of effort being used during movement. BA3b is responsible for distributing somatosensory information to BA1 and shape and size information to BA2.

This page titled 12.3A: General Organization of the Somatosensory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3B: Tactile Sensation

Touch is sensed by mechanoreceptive neurons that respond to pressure in various ways.

Learning Objectives

- Describe how touch is sensed by mechanoreceptive neurons responding to pressure

Key Points

- Our sense of touch, or tactile sensation, is mediated by cutaneous mechanoreceptors located in our skin.
- There are four main types of cutaneous mechanoreceptors: Pacinian corpuscles, Meissner's corpuscles, Merkel's discs, and Ruffini endings.
- Cutaneous mechanoreceptors are categorized by morphology, by the type of sensation they perceive, and by the rate of adaptation. Furthermore, each has a different receptive field.

Key Terms

- **receptive field**: The particular region of the sensory space (e.g., the body surface, space inside the ear) in which a stimulus will trigger the firing of that neuron.
- **adaptation**: A change over time in the responsiveness of the sensory system to a constant stimulus.
- **Aβ fiber**: A type of sensory nerve fiber that carries cold, pressure, and some pain signals.
- **Aδ fiber**: Carries sensory information related to muscle spindle secondary endings, touch, and kinesthesia.

A mechanoreceptor is a sensory receptor that responds to mechanical pressure or distortion. For instance, in the periodontal ligament, there are mechanoreceptors that allow the jaw to relax when biting down on hard objects; the mesencephalic nucleus is responsible for this reflex.

In the skin, there are four main types in glabrous (hairless) skin:

1. Ruffini endings.
2. Meissner's corpuscles.
3. Pacinian corpuscles.
4. Merkel's discs.

There are also mechanoreceptors in hairy skin. The hair cells in the cochlea are the most sensitive mechanoreceptors, transducing air pressure waves into nerve signals sent to the brain.

Cutaneous Mechanoreceptors

Cutaneous mechanoreceptors are located in the skin, like other cutaneous receptors. They provide the senses of touch, pressure, vibration, proprioception, and others. They are all innervated by Aβ fibers, except the mechanoreceiving free nerve endings, which are innervated by Aδ fibers.

They can be categorized by morphology, by the type of sensation they perceive, and by the rate of adaptation. Furthermore, each has a different receptive field:

- Ruffini's end organs detect tension deep in the skin.
- Meissner's corpuscles detect changes in texture (vibrations around 50 Hz) and adapt rapidly.
- Pacinian corpuscles detect rapid vibrations (about 200–300 Hz).
- Merkel's discs detect sustained touch and pressure.
- Mechanoreceiving free nerve endings detect touch, pressure, and stretching.

- Hair follicle receptors are located in hair follicles and sense the position changes of hair strands.

Ruffini Ending

The Ruffini ending (Ruffini corpuscle or bulbous corpuscle) is a class of slowly adapting mechanoreceptors thought to exist only in the glabrous dermis and subcutaneous tissue of humans. It is named after Angelo Ruffini.

This spindle-shaped receptor is sensitive to skin stretch, and contributes to the kinesthetic sense of and control of finger position and movement. It is believed to be useful for monitoring the slippage of objects along the surface of the skin, allowing the modulation of grip on an object.

Ruffini endings are located in the deep layers of the skin. They register mechanical information within joints, more specifically angle change, with a specificity of up to two degrees, as well as continuous pressure states. They also act as thermoreceptors that respond for a long time, such as holding hands with someone during a walk. In a case of a deep burn to the body, there will be no pain as these receptors will be burned off.

Meissner's Corpuscles

Meissner's corpuscles (or tactile corpuscles) are responsible for sensitivity to light touch. In particular, they have the highest sensitivity (lowest threshold) when sensing vibrations lower than 50 hertz. They are rapidly adaptive receptors.

Pacinian Corpuscles

Pacinian corpuscles (or lamellar corpuscles) are responsible for sensitivity to vibration and pressure. The vibrational role may be used to detect surface texture, e.g., rough versus smooth.

Merkel Nerve

Merkel nerve endings are mechanoreceptors found in the skin and mucosa of vertebrates that provide touch information to the brain. The information they provide are those regarding pressure and texture. Each ending consists of a Merkel cell in close apposition with an enlarged nerve terminal.

This is sometimes referred to as a Merkel cell–neurite complex, or a Merkel disk receptor. A single afferent nerve fiber branches to innervate up to 90 such endings. They are classified as slowly adapting type I mechanoreceptors.

This page titled 12.3B: Tactile Sensation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3C: Proprioceptive Sensations

Proprioception refers to the sense of knowing how one's body is positioned in three-dimensional space.

Learning Objectives

- Describe how propioception is the sense of the position of parts of our body in a three dimensional space

Key Points

- Proprioception is the sense of the position of parts of our body and force being generated during movement.
- Proprioception relies on two, primary stretch receptors: Golgi tendon organs and muscle spindles.
- Muscle spindles are sensory receptors within the belly of a muscle that primarily detect changes in the length of this muscle. They convey length information to the central nervous system via sensory neurons. This information can be processed by the brain to determine the position of body parts.
- The Golgi organ (also called Golgi tendon organ, tendon organ, neurotendinous organ, or neurotendinous spindle) is a proprioceptive sensory receptor organ that is located at the insertion of skeletal muscle fibers into the tendons of skeletal muscle.

Key Terms

- **alpha motor neuron**: Large, multipolar lower motor neurons of the brainstem and spinal cord that are directly responsible for initiating muscle contraction.
- **proprioreceptor**: A sensory receptor that responds to position and movement and that receives internal bodily stimuli.
- **posterior (dorsal) column-medial lemniscus pathway**: A sensory pathway of the central nervous system that conveys localized sensations of fine touch, vibration, two-point discrimination, and proprioception from the skin and joints.
- **Law of Righting**: A reflex rather than a law, this refers to the sudden movement of the head to level the eyes with the horizon in the event of an accidental tilting or imbalance of the body.
- **Golgi tendon organ**: A proprioceptive sensory receptor organ that is located at the insertion of skeletal muscle fibers into the tendons of skeletal muscle.
- **muscle spindle**: Sensory receptors within the belly of a muscle that primarily detect changes in the length of this muscle.
- **proprioception**: The sense of the position of parts of the body, relative to other neighboring parts of the body.


Proprioception is the sense of the relative position of neighboring parts of the body and the strength of effort being employed in movement. It is distinguished from exteroception, perception of the outside world, and interoception, perception of pain, hunger, and the movement of internal organs, etc.

The initiation of proprioception is the activation of a proprioreceptor in the periphery. The proprioceptive sense is believed to be composed of information from sensory neurons located in the inner ear (motion and orientation) and in the stretch receptors located in the muscles and the joint-supporting ligaments (stance).

Conscious proprioception is communicated by the posterior (dorsal) column–medial lemniscus pathway to the cerebrum. Unconscious proprioception is communicated primarily via the dorsal and ventral spinocerebellar tracts to the cerebellum.

An unconscious reaction is seen in the human proprioceptive reflex, or Law of Righting. In the event that the body tilts in any direction, the person will cock their head back to level the eyes against the horizon. This is seen even in infants as soon as they gain control of their neck muscles. This control comes from the cerebellum, the part of the brain that affects balance.


Muscle spindles are sensory receptors within the belly of a muscle that primarily detect changes in the length of a muscle. They convey length information to the central nervous system via sensory neurons. This information can be processed by the brain to determine the position of body parts. The responses of muscle spindles to changes in length also play an important role in regulating the contraction of muscles.



Muscle spindle: Mammalian muscle spindle showing typical position in a muscle (left), neuronal connections in spinal cord (middle), and expanded schematic (right). The spindle is a stretch receptor with its own motor supply consisting of several intrafusal muscle fibers. The sensory endings of a primary (group Ia) afferent and a secondary (group II) afferent coil around the non-contractile central portions of the intrafusal fibers.

The Golgi organ (also called Golgi tendon organ, tendon organ, neurotendinous organ or neurotendinous spindle) is a proprioceptive sensory receptor organ that is located at the insertion of skeletal muscle fibers onto the tendons of skeletal muscle. It provides the sensory component of the Golgi tendon reflex.

The Golgi organ should not be confused with the Golgi apparatus—an organelle in the eukaryotic cell —or the Golgi stain, which is a histologic stain for neuron cell bodies.



Golgi tendon organ: The Golgi tendon organ contributes to the Golgi tendon reflex and provides proprioceptive information about joint position.

The Golgi tendon reflex is a normal component of the reflex arc of the peripheral nervous system. In a Golgi tendon reflex, skeletal muscle contraction causes the agonist muscle to simultaneously lengthen and relax. This reflex is also called the inverse myotatic reflex, because it is the inverse of the stretch reflex.

Although muscle tension is increasing during the contraction, alpha motor neurons in the spinal cord that supply the muscle are inhibited. However, antagonistic muscles are activated.

This page titled 12.3C: Proprioceptive Sensations is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3D: Somatic Sensory Pathways

The somatosensory pathway is composed of three neurons located in the dorsal root ganglion, the spinal cord, and the thalamus.

Learning Objectives

- Describe the somatosensory area in the human cortex


Key Points

- A somatosensory pathway will typically have three neurons: primary, secondary, and tertiary.
- The cell bodies of the three neurons in a typical somatosensory pathway are located in the dorsal root ganglion, the spinal cord, and the thalamus.
- A major target of somatosensory pathways is the postcentral gyrus in the parietal lobe of the cerebral cortex.
- A major somatosensory pathway is the dorsal column–medial lemniscal pathway.
- The postcentral gyrus is the location of the primary somatosensory area that takes the form of a map called the sensory homunculus.

Key Terms

- **parietal lobe**: A part of the brain positioned superior to the occipital lobe and posterior to the frontal lobe that integrates sensory information from different modalities and is particularly important for determining spatial sense and navigation.
- **reticular activating system**: A set of connected nuclei in the brain responsible for regulating wakefulness and sleep-to-wake transitions.
- **postcentral gyrus**: A prominent structure in the parietal lobe of the human brain that is the location of the primary somatosensory cortex, the main sensory receptive area for the sense of touch.
- **thalamus**: Either of two large, ovoid structures of gray matter within the forebrain that relay sensory impulses to the cerebral cortex.

A somatosensory pathway will typically have three long neurons: primary, secondary, and tertiary. The first always has its cell body in the dorsal root ganglion of the spinal nerve.



Dorsal root ganglion: Sensory nerves of a dorsal root ganglion are depicted entering the spinal cord.

The second has its cell body either in the spinal cord or in the brainstem; this neuron's ascending axons will cross to the opposite side either in the spinal cord or in the brainstem. The axons of many of these neurons terminate in the thalamus, and others terminate in the reticular activating system or the cerebellum.

In the case of touch and certain types of pain, the third neuron has its cell body in the ventral posterior nucleus of the thalamus and ends in the postcentral gyrus of the parietal lobe.

In the periphery, the somatosensory system detects various stimuli by sensory receptors, such as by mechanoreceptors for tactile sensation and nociceptors for pain sensation. The sensory information (touch, pain, temperature, etc.,) is then conveyed to the central nervous system by afferent neurons, of which there are a number of different types with varying size, structure, and properties.


Generally, there is a correlation between the type of sensory modality detected and the type of afferent neuron involved. For example, slow, thin, unmyelinated neurons conduct pain whereas faster, thicker, myelinated neurons conduct casual touch.

Ascending Pathways

In the spinal cord, the somatosensory system includes ascending pathways from the body to the brain. One major target within the brain is the postcentral gyrus in the cerebral cortex. This is the target for neurons of the dorsal column–medial lemniscal pathway and the ventral spinothalamic pathway.

Note that many ascending somatosensory pathways include synapses in either the thalamus or the reticular formation before they reach the cortex. Other ascending pathways, particularly those involved with control of posture, are projected to the cerebellum, including the ventral and dorsal spinocerebellar tracts.

Another important target for afferent somatosensory neurons that enter the spinal cord are those neurons involved with local segmental reflexes.



Spinal nerve: The formation of the spinal nerve from the dorsal and ventral roots.

Parietal Love: Primary Somatosensory Area


The primary somatosensory area in the human cortex is located in the postcentral gyrus of the parietal lobe. This is the main sensory receptive area for the sense of touch.

Like other sensory areas, there is a map of sensory space called a homunculus at this location. Areas of this part of the human brain map to certain areas of the body, dependent on the amount or importance of somatosensory input from that area.

For example, there is a large area of cortex devoted to sensation in the hands, while the back has a much smaller area. Somatosensory information involved with proprioception and posture also target an entirely different part of the brain, the cerebellum.

Cortical Homunculus

This is a pictorial representation of the anatomical divisions of the primary motor cortex and the primary somatosensory cortex; namely, the portion of the human brain directly responsible for the movement and exchange of sensory and motor information of the body.




Homunculus: Image representing the cortical sensory homunculus.

Thalamus

The thalamus is a midline symmetrical structure within the brain of vertebrates including humans; it is situated between the cerebral cortex and midbrain, and surrounds the third ventricle.

Its function includes relaying sensory and motor signals to the cerebral cortex, along with the regulation of consciousness, sleep, and alertness.



Thalamic nuclei: The ventral posterolateral nucleus receives sensory information from the body.

This page titled 12.3D: Somatic Sensory Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3E: Mapping the Primary Somatosensory Area

The cortical sensory homunculus is located in the postcentral gyrus and provides a representation of the body to the brain.

Learning Objectives

- Describe how primary somatosensory areas can be mapped

Key Points

- A sensory homunculus is a pictorial representation of the primary somatosensory cortex.
- Somatotopy is the correspondence of an area of the body to a specific point in the brain.
- Wilder Penfield was a researcher and surgeon who created maps of the somatosensory cortex.

Key Terms

- **somesthetic cortex**: The primary mechanism of cortical processing for sensory information originating at body surfaces and other tissues (eg., muscles, joints).
- **postcentral gyrus**: A prominent structure in the parietal lobe of the human brain that is the location of the primary somatosensory cortex, the main sensory receptive area for the sense of touch.
- **precentral gyrus**: The precentral gyrus lies in front of the postcentral gyrus and is the site of the primary motor cortex (Brodmann area 4).

Cortical Homunculus


A cortical homunculus is a pictorial representation of the anatomical divisions of the primary motor cortex and the primary somatosensory cortex; it is the portion of the human brain directly responsible for the movement and exchange of sensory and motor information of the body.

It is a visual representation of the concept of the body within the brain—that one's hand or face exists as much as a series of nerve structures or a neuron concept as it does in a physical form. There are two types of homunculus: sensory and motor. Each one shows a representation of how much of its respective cortex innervates certain body parts.

The primary somesthetic cortex (sensory) pertains to the signals within the postcentral gyrus coming from the thalamus, and the primary motor cortex pertains to signals within the precentral gyrus coming from the premotor area of the frontal lobes.

These are then transmitted from the gyri to the brain stem and spinal cord via corresponding sensory or motor nerves. The reason for the distorted appearance of the homunculus is that the amount of cerebral tissue or cortex devoted to a given body region is proportional to how richly innervated that region is, not to its size.


The homunculus is like an upside-down sensory or motor map of the contralateral side of the body. The upper extremities such as the facial body parts and hands are closer to the lateral sulcus than lower extremities such as the leg and toes.



Homunculus: The idea of the cortical homunculus was created by Wilder Penfield and serves as a rough map of the receptive fields for regions of primary somatosensory cortex.

The resulting image is a grotesquely disfigured human with disproportionately huge hands, lips, and face in comparison to the rest of the body. Because of the fine motor skills and sense nerves found in these particular parts of the body, they are represented as being larger on the homunculus. A part of the body with fewer sensory and/or motor connections to the brain is represented to appear smaller.

Somatotopy



Postcentral gyrus: The postcentral gyrus is located in the parietal lobe of the human cortex and is the primary somatosensory region of the human brain.

This is the point-for-point correspondence of an area of the body to a specific point on the central nervous system. Typically, the area of the body corresponds to a point on the primary somatosensory cortex (postcentral gyrus).

This cortex is typically represented as a sensory homunculus which orients the specific body parts and their respective locations upon the homunculus. Areas such as the appendages, digits, and face can draw their sensory locations upon the somatosensory

cortex.

Areas that are finely controlled, such as the digits, have larger portions of the somatosensory cortex, whereas areas that are coarsely controlled, such as the trunk, have smaller portions. Areas such as the viscera do not have sensory locations on the postcentral gyrus.

Montreal Procedure

Wilder Penfield was a groundbreaking researcher and highly original surgeon. With his colleague, Herbert Jasper, he invented the Montreal procedure, in which he treated patients with severe epilepsy by destroying nerve cells in the brain where the seizures originated.

Before operating, he stimulated the brain with electrical probes while the patients were conscious on the operating table (under only local anesthesia), and observed their responses. In this way he could more accurately target the areas of the brain responsible, reducing the side-effects of the surgery.

This technique also allowed him to create maps of the sensory and motor cortices of the brain, showing their connections to the various limbs and organs of the body. These maps are still used today, practically unaltered.

Along with Herbert Jasper, he published this landmark work in 1951 as *Epilepsy and the Functional Anatomy of the Human Brain*. This work contributed a great deal to understanding the lateralization of brain function.

Penfield's maps showed considerable overlap between regions (for instance, the motor region controlling muscles in the hand sometimes also controlled muscles in the upper arm and shoulder), a feature that he put down to individual variation in brain size and localization; we now know that this is due to the fractured somatotropy of the motor cortex.

This page titled 12.3E: Mapping the Primary Somatosensory Area is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.3F: Somatic Sensory Pathways to the Cerebellum

The ventral and dorsal spinocerebellar tracts convey proprioceptive information from the body to the cerebellum.

Learning Objectives

- Describe the somatic sensory pathways to the cerebellum

Key Points

- The main somatosensory pathways that communicate with the cerebellum are the ventral (or anterior) and dorsal (or posterior) spinocerebellar tracts.
- The ventral spinocerebellar tract will cross to the opposite side of the body then cross again to end in the cerebellum (referred to as a double cross). The dorsal spinocerebellar tract does not decussate or cross sides at all through its path.
- The dorsal spinocerebellar tract (also called the posterior spinocerebellar tract, Flechsig's fasciculus, or Flechsig's tract) conveys inconscient proprioceptive information from the body to the cerebellum.

Key Terms


- **Clarke's nucleus**: A group of interneurons important in proprioception that is found in the intermediate zone of the spinal cord.
- **first order neuron**: Conducts impulses from proprioceptors and skin receptors to the spinal cord or brain stem.

A sensory system is a part of the nervous system responsible for processing sensory information. A sensory system consists of sensory receptors, neural pathways, and the parts of the brain involved in sensory perception. Commonly recognized sensory systems are those for vision, hearing, somatic sensation (touch), taste, and olfaction (smell).

In short, senses are transducers from the physical world to the realm of the mind where we interpret the information, creating our perception of the world around us.

The ventral spinocerebellar tract conveys proprioceptive information from the body to the cerebellum. It is part of the somatosensory system and runs in parallel with the dorsal spinocerebellar tract.

Both tracts involve two neurons. The ventral spinocerebellar tract will cross to the opposite side of the body then cross again to end in the cerebellum (referred to as a double cross). The dorsal spinocerebellar tract does not decussate, or cross sides, at all through its path.



The major tracts of the spinal cord: The anterior and posterior spinocerebellar tracts are the major somatosensory pathways communicating with the cerebellum.

The ventral tract (under L2/L3) gets its proprioceptive/fine touch/vibration information from a first order neuron, with its cell body in a dorsal ganglion. The axon runs via the fila radicularia (nerve rootlets) to the dorsal horn of the gray matter. There it makes a synapse with the dendrites of two neurons that send their axons bilaterally to the ventral border of the lateral funiculi (transmit the contralateral corticospinal and
spinothalamic tracts). The ventral spinocerebellar tract then enters the cerebellum via the superior cerebellar peduncle (connects the cerebellum to the midbrain).

This is in contrast with the dorsal spinocerebellar tract (C8 – L2/L3), which only has one unilateral axon that has its cell body in Clarke's nucleus (only at the level of C8 – L2/L3). The fibers of the ventral spinocerebellar tract then eventually enter the cerebellum via the superior cerebellar peduncle.

This is one of the few afferent tracts through the superior cerebellar peduncle. Axons first cross midline in the spinal cord and run in the ventral border of the lateral funiculi. These axons ascend to the pons where they join the superior cerebellar peduncle to enter the cerebellum.

Once in the deep, white matter of the cerebellum, the axons recross the midline, give off collaterals to the globose and emboliform nuclei (deep cerebellar nuclei), and terminate in the cortex of the anterior lobe and vermis of the posterior lobe.

The dorsal spinocerebellar tract (also called the posterior spinocerebellar tract, Flechsig's fasciculus, or Flechsig's tract) conveys inconscient proprioceptive information from the body to the cerebellum. It is part of the somatosensory system and runs in parallel with the ventral spinocerebellar tract.

Proprioceptive information is taken to the spinal cord via central processes of the dorsal root ganglia (where first order neurons reside). These central processes travel through the dorsal horn where they synapse with second order neurons of Clarke's nucleus.

Axon fibers from Clarke's nucleus convey this proprioceptive information in the spinal cord to the peripheral region of the posterolateral funiculus ipsilaterally until it reaches the cerebellum, where unconscious proprioceptive information is processed. This tract involves two neurons and ends up on the same side of the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Somatosensory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Somatos...stem%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- postcentral gyrus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/postcentral%20gyrus. **License**: *CC BY-SA: Attribution-ShareAlike*
- organization. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/organization. **License**: *CC BY-SA: Attribution-ShareAlike*
- decussate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/decussate. **License**: *CC BY-SA: Attribution-ShareAlike*
- thalamus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thalamus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1...Homunculus.jpg. **License**: *CC BY: Attribution*
- Neural Adaptation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neural_adaptation. **License**: *CC BY-SA: Attribution-ShareAlike*
- A Delta Fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/A_delta_fiber. **License**: *CC BY-SA: Attribution-ShareAlike*
- Receptive Field. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Receptive_field. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nervie Fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_fiber#Sensory_fibers_of_the_A_group. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mechanoreceptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mechanoreceptor%23Cutaneous. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thermoreceptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thermoreceptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ruffini ending. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ruffini_ending. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pacinian corpuscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pacinian_corpuscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mechanoreceptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mechanoreceptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Meissner's corpuscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Meissner's_corpuscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Merkel nerve ending. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Merkel_nerve_ending. **License**: *CC BY-SA: Attribution-ShareAlike*
- Merkel's disc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Merkel's%20disc. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pacinian corpuscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pacinian%20corpuscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ruffini ending. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ruffini%20ending. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Alpha Motor Neuron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Alpha_motor_neuron. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stretch receptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stretch_receptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscle spindle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscle_spindle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Proprioception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Proprioception%23Basis_of_proprioceptive_sense. **License**: *CC BY-SA: Attribution-ShareAlike*
- Proprioception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Proprioception. **License**: *CC BY-SA: Attribution-ShareAlike*
- Golgi tendon reflex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Golgi_tendon_reflex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Golgi organ. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Golgi_organ. **License**: *CC BY-SA: Attribution-ShareAlike*

- Proprioception. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Proprioception%23Impairment. **License**: *CC BY-SA: Attribution-ShareAlike*
- muscle spindle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/muscle%20spindle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Golgi tendon organ. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Golgi%20tendon%20organ. **License**: *CC BY-SA: Attribution-ShareAlike*
- proprioception. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/proprioception. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Muscle spindle model. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mu...ndle_model.jpg. **License**: *Public Domain: No Known Copyright*
- Gray938. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray938.png. **License**: *Public Domain: No Known Copyright*
- Ventral posterior nucleus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventral_posterior_nucleus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Postcentral gyrus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postcentral_gyrus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thalamus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thalamus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dorsal column-medial lemniscus pathway. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dorsal_column-medial_lemniscus_pathway. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dorsal root ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dorsal_root_ganglion. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cortical homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Somatosensory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Somatosensory_system%23General_somatosensory_pathway. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parietal lobe. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parietal_lobe. **License**: *CC BY-SA: Attribution-ShareAlike*
- parietal lobe. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/parietal%20lobe. **License**: *CC BY-SA: Attribution-ShareAlike*
- thalamus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thalamus. **License**: *CC BY-SA: Attribution-ShareAlike*
- postcentral gyrus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/postcentral%20gyrus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Muscle spindle model. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mu...ndle_model.jpg. **License**: *Public Domain: No Known Copyright*
- Gray938. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray938.png. **License**: *Public Domain: No Known Copyright*
- Gray675. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray675.png. **License**: *Public Domain: No Known Copyright*
- Thalamus-schematic. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...-schematic.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Spinal_nerve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY: Attribution*
- Sensory homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensory...nculus%23Types. **License**: *CC BY-SA: Attribution-ShareAlike*
- Postcentral gyrus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postcentral_gyrus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Somatotopic map. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Somatotopic_map. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensory_homunculus. **License**: *CC BY-SA: Attribution-ShareAlike*

- Wilder Penfield. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wilder_Penfield. **License**: *CC BY-SA: Attribution-ShareAlike*
- somatotopy. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somatotopy. **License**: *CC BY-SA: Attribution-ShareAlike*
- postcentral gyrus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/postcentral%20gyrus. **License**: *CC BY-SA: Attribution-ShareAlike*
- sensory homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sensory%20homunculus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Muscle spindle model. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mu...ndle_model.jpg. **License**: *Public Domain: No Known Copyright*
- Gray938. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray938.png. **License**: *Public Domain: No Known Copyright*
- Gray675. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray675.png. **License**: *Public Domain: No Known Copyright*
- Thalamus-schematic. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...-schematic.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Spinal_nerve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY: Attribution*
- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY: Attribution*
- Postcentral gyrus superior view. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F....png#filelinks. **License**: *Public Domain: No Known Copyright*
- Ventral spinocerebellar tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventral...rebellar_tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dorsal spinocerebellar tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dorsal_...rebellar_tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lateral Funiculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lateral_funiculus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Globose Nucleus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Globose_nucleus. **License**: *CC BY-SA: Attribution-ShareAlike*
- ventral spinocerebellar tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/ventral...bellar%20tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsal spinocerebellar tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsal%...bellar%20tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory Homunculus. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1421_Sensory_Homunculus.jpg. **License**: *CC BY: Attribution*
- Muscle spindle model. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Muscle_spindle_model.jpg. **License**: *Public Domain: No Known Copyright*
- Gray938. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray938.png. **License**: *Public Domain: No Known Copyright*
- Gray675. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray675.png. **License**: *Public Domain: No Known Copyright*
- Thalamus-schematic. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...-schematic.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Spinal_nerve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY: Attribution*
- Cortical Homunculus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cortical_homunculus. **License**: *CC BY: Attribution*
- Postcentral gyrus superior view. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Postcentral_gyrus_-_superior_view2.png#filelinks. **License**: *Public Domain: No Known Copyright*
- Spinal cord tracts - English. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Spinal_cord_tracts_-_English.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 12.3F: Somatic Sensory Pathways to the Cerebellum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.4: NERVES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

12.4A: Structure of a Nerve

12.4B: Classification of Nerves

This page titled 12.4: Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.4A: Structure of a Nerve

A nerve is the primary structure of the peripheral nervous system and is composed of bundles of axons.

Learning Objectives

- Describe the structure of nerves


Key Points

- A nerve is the primary structure of the peripheral nervous system (PNS) that encloses the axons of peripheral neurons.
- A nerve provides a structured pathway that supports neuron function.
- A nerve consists of many structures including axons, glycocalyx, endoneurial fluid, endoneurium, perineurium, and epineurium.
- The axons are bundled together into groups called fascicles, and each fascicle is wrapped in a layer of connective tissue called the perineurium.
- Magnetic resonsance neurography is a technology used to detect nerve damage.

Key Terms

- **endoneurial fluid**: A low protein liquid that is the peripheral nervous system equivalent to cerebrospinal fluid in the central nervous system.
- **perineurium**: A protective sheath covering nerve fascicles.
- **glycocalyx**: A glycoprotein-polysaccharide covering that surrounds cell membranes.
- **endoneurium**: A layer of connective tissue that surrounds axons.
- **fascicles**: A small bundle of nerve fibers enclosed by the perineurium.
- **epineurium**: The outermost layer of dense, irregular connective tissue surrounding a peripheral nerve.

Nerve Anatomy



Nerves: An illustration of the main nerves of the arm.

A nerve is an enclosed, cable-like bundle of axons (the projections of neurons) in the peripheral nervous system (PNS). A nerve provides a structured pathway that supports the electrochemical nerve impulses transmitted along each of the axons.

In the central nervous system, the analogous structures are known as tracts. Neurons are sometimes referred to as nerve cells, although this term is misleading since many neurons do not occupy nerves, and nerves also include non-neuronal support cells (glial cells) that contribute to the health of enclosed neurons.


Each nerve contains many axons that are sometimes referred to as fibers. Within a nerve, each axon is surrounded by a layer of connective tissue called the endoneurium. The axons are bundled together into groups called fascicles. Each fascicle is wrapped in a layer of connective tissue called the perineurium.

Finally, the entire nerve is wrapped in a layer of connective tissue called the epineurium. See the following illustrations of these structures.

The endoneurium consists of an inner sleeve of material called the glycocalyx and a mesh of collagen. Nerves are bundled along with blood vessels, which provide essential nutrients and energy to the enclosed, and metabolically demanding, neurons.

Within the endoneurium, individual nerve fibers are surrounded by a liquid called the endoneurial fluid. The endoneurium has properties analogous to the blood–brain barrier. It prevents certain molecules from crossing from the blood into the endoneurial fluid.

In this respect, endoneurial fluid is similar to cerebrospinal fluid in the central nervous system. During nerve irritation or injury, the amount of endoneurial fluid may increase at the site of damage. This increase in fluid can be visualized using magnetic resonance neurography to diagnose nerve damage.



(a) Anatomy of a nerve and (b) Cross-section of a nerve: The primary structures of a nerve. An illustration of a cross-section of a nerve highlighting the epineurium and perineurium. Individual axons can also be seen as tiny circles within each perineurium.

Basic Function

A nerve conveys information in the form of electrochemical impulses (known as nerve impulses or action potentials) carried by the individual neurons that make up the nerve. These impulses are extremely fast, with some myelinated neurons conducting at speeds up to 120 m/s. The impulses travel from one neuron to another by crossing a synapse, and the message is converted from electrical to chemical and then back to electrical.

Nerves can be categorized into two groups based on function:

1. Sensory nerves conduct sensory information from their receptors to the central nervous system, where the information is then processed. Thus they are synonymous with afferent nerves.
2. Motor nerves conduct signals from the central nervous system to muscles. Thus they are synonymous with efferent nerves.

Neurologists usually diagnose disorders of the nerves by a physical examination, including the testing of reflexes, walking and other directed movements, muscle weakness, proprioception, and the sense of touch. This initial exam can be followed with tests such as nerve conduction study, electromyography, or computed tomography.

This page titled 12.4A: Structure of a Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.4B: Classification of Nerves

Learning Objectives


- List the different ways that nerves can be classified

Nerve Classifications

Direction of Signal Transmission

Nerves are categorized into three, primary groups based on the direction of signal transmission within the nervous system.

1. Afferent nerves conduct signals from sensory neurons to the central nervous system, for example from mechanoreceptors in skin.
2. Efferent nerves conduct signals away from the central nervous system to target muscles and glands.
3. Mixed nerves contain both afferent and efferent axons, and thus conduct both incoming sensory information and outgoing muscle commands in the same nerve bundle.



Afferent and efferent nerve transmission: Schematic of efferent and afferent nerve transmission to and from peripheral tissue and spinal cord.

Central Nervous System Connection

Nerves can be further categorized based on where they connect to the central nervous system. Spinal nerves innervate much of the body and connect through the spinal column to the spinal cord. Spinal nerves are assigned letter-number designations according to the vertebra where they connect to the spinal column. Cranial nerves innervate parts of the head and connect directly to the brain. Cranial nerves are typically assigned Roman numerals from 0 to 12.

Diameter, Conduction Velocity, Myelination State


Peripheral nerve fibers are grouped based on the diameter, signal conduction velocity, and myelination state of the axons. These classifications apply to both sensory and motor fibers. Fibers of the A group have a large diameter, high conduction velocity, and

are myelinated.

The A group is further subdivided into four types (A-alpha, A-beta, A-delta, and A-gamma fibers) based on the information carried by the fibers and the tissues they innervate.

- A-alpha fibers are the primary receptors of the muscle spindle and golgi tendon organ.
- A-beta fibers act as secondary receptors of the muscle spindle and contribute to cutaneous mechanoreceptors.
- A-delta fibers are free nerve endings that conduct painful stimuli related to pressure and temperature.
- A-gamma fibers are typically motor neurons that control the intrinsic activation of the muscle spindle.

Fibers of the B group are myelinated with a small diameter and have a low conduction velocity. The primary role of B fibers is to transmit autonomic information. Fibers of the C group are unmyelinated, have a small diameter, and low conduction velocity. The lack of myelination in the C group is the primary cause of their slow conduction velocity.



Saltatory conduction: Demonstrates the faster propagation of an action potential in myelinated neurons than that of unmyelinated neurons.

C fiber axons are grouped together into what is known as Remak bundles. These occur when an unmyelinated Schwann cell bundles the axons close together by surrounding them. The Schwann cell keeps them from touching each other by squeezing its cytoplasm between the axons.

C fibers are considered polymodal because they can often respond to combinations of thermal, mechanical, and chemical stimuli.

A-delta and C fibers both contribute to the detection of diverse painful stimuli. Because of their higher conduction velocity, A-delta fibers are responsible for the sensation of a sharp, initial pain and respond to a weaker intensity of stimulus.

These nerve fibers are associated with acute pain and therefore constitute the afferent portion of the reflex arc that results in pulling away from noxious stimuli. An example is the retraction or your hand from a hot stove. Slowly conducting, unmyelinated C fibers, by contrast, carry slow, longer-lasting pain sensations.

Key Points

- Nerves can be categorized as afferent, efferent, and mixed based on the direction of signal transmission within the nervous system. Nerves can be further categorized as spinal nerves or cranial nerves based on where they connect to the central nervous system.
- Individual peripheral nerve fibers are classified based on the diameter, signal conduction velocity, and myelination state of the axons, as well as by the type of information transmitted and the organs they innervate.

Key Terms

- **mixed nerve**: Nerves that contain both afferent and efferent axons, and thus conduct both incoming sensory information and outgoing muscle commands in the same bundle.
- **Afferent nerve**: Carries nerve impulses from sensory receptors or sense organs toward the central nervous system.
- **Schwann cell**: The principal glia of the peripheral nervous system.
- **efferent nerve**: Nerves that conduct signals from the central nervous system along motor neurons to their target muscles and glands.
- **spinal nerve**: The term generally refers to a mixed nerve that carries motor, sensory, and autonomic signals between the spinal cord and the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Epineurium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epineurium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glycocalyx. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Glycocalyx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Perineurium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Perineurium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerve Fascicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_fascicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endoneurium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Endoneurium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerves%23Anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fascicle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fascicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- perineurium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/perineurium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endoneurial fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/endoneurial%20fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerves%23Anatomy**. **License**: ***Public Domain: No Known Copyright***
- Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerves%23Anatomy**. **License**: ***Public Domain: No Known Copyright***
- Nerve fascicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_fascicle**. **License**: ***Public Domain: No Known Copyright***
- Schwann Cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Schwann_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerve fiber. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_f...of_the_A_group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerve fiber. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_f...of_the_A_group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerve fiber. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_f...ve_fiber_types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sensory fiber types. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sensory..._types%23Motor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerves%23Anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nerve fiber. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_f...l_nerve_fibers**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- B fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/B_fiber. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sensory fiber types. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sensory...ypes%23Sensory. **License**: *CC BY-SA: Attribution-ShareAlike*
- A delta fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/A_delta_fiber. **License**: *CC BY-SA: Attribution-ShareAlike*
- Group C nerve fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Group_C...re_and_Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Group C nerve fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Group_C...ber%23Function. **License**: *CC BY-SA: Attribution-ShareAlike*
- efferent nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/efferent%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- mixed nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/mixed%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- spinal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/spinal%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerves%23Anatomy. **License**: *Public Domain: No Known Copyright*
- Nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerves%23Anatomy. **License**: *Public Domain: No Known Copyright*
- Nerve fascicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_fascicle. **License**: *Public Domain: No Known Copyright*
- Saltatory Conduction. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Saltatory_Conduction.gif. **License**: *CC BY: Attribution*
- Afferent. **Provided by**: Wikepedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Afferent_(PSF).png. **License**: *Public Domain: No Known Copyright*

This page titled 12.4B: Classification of Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5: CRANIAL NERVES

12.5A: Brief Overview of Cranial Nerves
12.5B: Olfactory (I) Nerve
12.5C: Optic (II) Nerve
12.5D: Oculomotor (III) Nerve
12.5E: Trochlear (IV) Nerve
12.5F: Trigeminal (V) Nerve
12.5G: Abducens (VI) Nerve
12.5H: Facial (VII) Nerve
12.5I: Vestibulocochlear (VIII) Nerve
12.5J: Glossopharyngeal (IX) Nerve
12.5K: Vagus (X) Nerve
12.5L: Accessory (XI) Nerve
12.5M: Hypoglossal (XII) Nerve

This page titled 12.5: Cranial Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5A: Brief Overview of Cranial Nerves

Learning Objectives

- Describe the function of the cranial nerves

Cranial nerves are the nerves that emerge directly from the brain (including the brainstem). In contrast, spinal nerves emerge from segments of the spinal cord. Cranial nerves relay information between the brain and parts of the body, primarily to and from regions of the head and neck.

Cranial Nerve Anatomy and Terminology


Spinal nerves emerge sequentially from the spinal cord with the spinal nerve closest to the head (C1) emerging in the space above the first cervical vertebra. The cranial nerves emerge from the central nervous system above this level.

Each cranial nerve is paired and is present on both sides. The numbering of the cranial nerves is based on the order in which they emerge from the brain, front to back (brainstem).

The terminal nerves, olfactory nerves (I) and optic nerves (II) emerge from the cerebrum or forebrain, and the remaining ten pairs arise from the brainstem, which is the lower part of the brain. The cranial nerves are considered components of the peripheral nervous system.

However, on a structural level, the olfactory, optic, and terminal nerves are more accurately considered part of the central nervous system.

The twelve cranial nerves are shown in the figure below followed by brief descriptions.



The cranial nerves: The locations of the cranial nerves within the brain.

- The olfactory nerve (I): This is instrumental for the sense of smell, it is one of the few nerves that are capable of regeneration.
- The optic nerve (II): This nerve carries visual information from the retina of the eye to the brain.
- The oculomotor nerve (III): This controls most of the eye's movements, the constriction of the pupil, and maintains an open eyelid.
- The trochlear nerve (IV): A motor nerve that innervates the superior oblique muscle of the eye, which controls rotational

movement.
- The trigeminal nerve (V): This is responsible for sensation and motor function in the face and mouth.
- The abducens nerve (VI): A motor nerve that innervates the lateral rectus muscle of the eye, which controls lateral movement.
- The facial nerve (VII): This controls the muscles of facial expression, and functions in the conveyance of taste sensations from the anterior two-thirds of the tongue and oral cavity.
- The vestibulocochlear nerve (VIII): This is responsible for transmitting sound and equilibrium (balance) information from the inner ear to the brain.
- The glossopharyngeal nerve (IX): This nerve receives sensory information from the tonsils, the pharynx, the middle ear, and the rest of the tongue.
- The vagus nerve (X): This is responsible for many tasks, including heart rate, gastrointestinal peristalsis, sweating, and muscle movements in the mouth, including speech and keeping the larynx open for breathing.
- The spinal accessory (XI): This nerve controls specific muscles of the shoulder and neck.
- The hypoglossal nerve (XII): This nerve controls the tongue movements of speech, food manipulation, and swallowing.

There are many mnemonic devices to remember the cranial nerves. One that may be helpful is: Old Opie Occasionally Tries Trigonometry And Feels Very Gloomy, Vague And Hypoactive.

Key Points

- The cranial nerves serve functions such as smell, sight, eye movement, and feeling in the face. The cranial nerves also control balance, hearing, and swallowing.
- The twelve cranial nerves, in order from I to XII are: olfactory nerve, optic nerve, oculomotor nerve, trochlear nerve, trigeminal nerve, abducens nerve, facial nerve, vestibulocochlear nerve, glossopharengeal nerve, vagus nerve, spinal accessory nerve, and hypoglossal nerve.
- The vagus nerve (X) has many branches and is responsible for tasks including heart rate, gastrointestinal peristalsis, sweating, and muscle movements in the mouth, including speech and keeping the larynx open for breathing.

Key Terms

- **innervate**: To supply an organ or other body part with nerves.
- **peripheral nervous system**: The nerves and ganglia outside of the brain and spinal cord.
- **peristalsis**: A radially symmetrical contraction and relaxation of muscles that propagates in a wave down a tube (such as the gastrointestinal tract) in an anterograde direction.

This page titled 12.5A: Brief Overview of Cranial Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5B: Olfactory (I) Nerve

The olfactory nerve, or cranial nerve I, is the first of 12 cranial nerves and is responsible for the sense of smell.

Learning Objectives


- Describe the olfactory nerve (cranial nerve I)

Key Points

- The olfactory nerve consists of a collection of many sensory nerve fibers that extend from the olfactory epithelium to the olfactory bulb.
- Olfactory receptors within the olfactory mucosa in the nasal cavity receive information about smells that travel to the brain through the cranial nerve that extends from the olfactory epithelium to the olfactory bulb.
- Olfactory receptor neurons continue to emerge throughout life and extend new axons to the olfactory bulb.
- The olfactory nerve is the shortest of the 12 cranial nerves and only one of two cranial nerves (the other being the optic nerve) that do not join with the brainstem.

Key Terms

- **olfactory receptors**: Expressed in the cell membranes of olfactory receptor neurons, these are responsible for the detection of odor molecules. Activated olfactory receptors are the initial player in a signal transduction cascade that ultimately produces a nerve impulse that is transmitted to the brain. The olfactory receptors form a multigene family consisting of over 900 genes in humans and 1,500 genes in mice.
- **cribriform plate**: A sieve-like structure of the ethmoid bone that supports the olfactory bulb.
- **olfactory tract**: A narrow white band made of a bundle of axons connecting the olfactory bulb to several brain regions.
- **ethmoid bone**: An unpaired bone in the skull that separates the nasal cavity from the brain.
- **olfactory mucosa**: Located in the upper region of the nasal cavity, this is made up of the olfactory epithelium and the underlying lamina propria, connective tissue containing fibroblasts, blood vessels, Bowman's glands, and bundles of fine axons from the olfactory neurons.



Olfactory bulb: Sagittal section of human head showing the olfactory bulb.

The olfactory nerve, or cranial nerve I, is the first of the 12 cranial nerves. It is instrumental in the sense of smell. The olfactory nerve is the shortest of the 12 cranial nerves and only one of two cranial nerves (the other being the optic nerve) that do not join with the brainstem.

The specialized olfactory receptor neurons of the olfactory nerve are located in the olfactory mucosa of the upper parts of the nasal cavity. The olfactory nerves consist of a collection of many sensory nerve fibers that extend from the olfactory epithelium to the olfactory bulb, passing through the many openings of the cribriform plate of the ethmoid bone.

Olfactory receptor neurons continue to emerge throughout life and extend new axons to the olfactory bulb. Olfactory-ensheathing glia wrap bundles of these axons and are thought to facilitate their passage into the central nervous system.

The sense of smell (olfaction) arises from the stimulation of olfactory (or odorant) receptors by small molecules of different spatial, chemical, and electrical properties that pass over the nasal epithelium in the nasal cavity during inhalation. These interactions are transduced into electrical activity in the olfactory bulb, which then transmits the electrical activity to other parts of the olfactory system and the rest of the central nervous system via the olfactory tract.

This page titled 12.5B: Olfactory (I) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5C: Optic (II) Nerve

The optic nerve (cranial nerve II) receives visual information from photoreceptors in the retina and transmits it to the brain.

Learning Objectives

- Describe the optic nerve (cranial nerve II)

Key Points


- The optic nerve is considered part of the central nervous system. The myelin on the optic nerve is produced by oligodendrocytes rather than Schwann cells and it is encased in the meningeal layers instead of the standard endoneurium, perineurium, and epineurium of the peripheral nervous system.
- The optic nerve travels through the optic canal, partially decussates in the optic chiasm, and terminates in the lateral geniculate nucleus where information is transmitted to the visual cortex.
- The axons responsible for reflexive eye movements terminate in the pretectal nucleus.

Key Terms

- **oligodendrocyte**: A type of neuroglia that provides support and insulation to axons in the central nervous system.
- **retina**: The thin layer of cells at the back of the eyeball where light is converted into neural signals sent to the brain.
- **optic nerve**: Either of a pair of nerves that carry visual information from the retina to the brain.
- **visual cortex**: The visual cortex of the brain is the part of the cerebral cortex responsible for processing visual information. It is located in the occipital lobe, in the back of the brain.
- **pretectal nucleus**: This mediates behavioral responses to acute changes in ambient light, such as the pupillary light reflex and the optokinetic reflex.

The optic nerve is also known as cranial nerve II. It transmits visual information from the retina to the brain.

Each human optic nerve contains between 770,000 and 1.7 million nerve fibers. The eye's blind spot is a result of the absence of photoreceptors in the area of the retina where the optic nerve leaves the eye.



Optic nerve: An illustration of the brain highlighting the optic nerve and optic tract.

The optic nerve is the second of twelve paired cranial nerves. It is considered by physiologists to be part of the central nervous system, as it is derived from an outpouching of the diencephalon during embryonic development.

As a consequence, the fibers are covered with myelin produced by oligodendrocytes, rather than Schwann cells that are found in the peripheral nervous system. The optic nerve is ensheathed in all three meningeal layers (dura, arachnoid, and pia mater) rather than the epineurium, perineurium, and endoneurium found in the peripheral nerves.

The fiber tracks of the mammalian central nervous system are incapable of regeneration. As a consequence, optic nerve damage produces irreversible blindness.

The optic nerve leaves the orbit, which is also known as an eye socket, via the optic canal, running posteromedially toward the optic chiasm, where there is a partial decussation (crossing) of fibers from the nasal visual fields of both eyes.

Most of the axons of the optic nerve terminate in the lateral geniculate nucleus (where information is relayed to the visual cortex), while other axons terminate in the pretectal nucleus and are involved in reflexive eye movements.

The optic nerve transmits all visual information including brightness perception, color perception, and contrast. It also conducts the visual impulses that are responsible for two important neurological reflexes: the light reflex and the accommodation reflex.

The light reflex refers to the constriction of both pupils that occurs when light is shone into either eye; the accommodation reflex refers to the swelling of the lens of the eye that occurs when one looks at a near object, as in reading.

This page titled 12.5C: Optic (II) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5D: Oculomotor (III) Nerve

The oculomoter nerve (cranial nerve III) controls eye movement, such as constriction of the pupil and open eyelids.

Learning Objectives

- Describe the oculomotor nerve (cranial nerve III)

Key Points

- The oculomotor nerve is the third paired cranial nerve.
- The oculomotor nerve contains two nuclei, including the Edinger-Westphal nucleus that supplies parasympathetic nerve fibers to the eye to control pupil constriction and accommodation.
- The oculomotor nerve originates at the superior colliculus and enters through the superior orbital fissure to control the levator palpebrae superioris muscles that hold the eyelids open.

Key Terms

- **levator palpebrae superioris**: A muscle that elevates the upper eyelid.
- **superior colliculus**: The primary integrating center for eye movements.
- **cavernous sinus**: A venous channel found between layers of dura mater in the brain.
- **tentorium cerebelli**: An extension of the dura mater that separates the cerebellum from the inferior portion of the occipital lobes.
- **superior orbital fissure**: The superior orbital fissure is a foramen in the skull, although strictly it is more of a cleft, lying between the lesser and greater wings of the sphenoid bone.
- **plexus**: A network or interwoven mass, especially of nerves, blood vessels, or lymphatic vessels.

The oculomotor nerve is the third paired cranial nerve. It enters the orbit via the superior orbital fissure and controls most of the eye's movements, including constriction of the pupil and maintaining an open eyelid by innervating the levator palpebrae superiors muscle.

The occulomotor nerve is derived from the basal plate of the embryonic midbrain. Cranial nerves IV and VI also participate in control of eye movement.

There are two nuclei for the oculomotor nerve:

1. The oculomotor nucleus originates at the level of the superior colliculus. The muscles it controls are the striated muscle in the levator palpebrae superioris and all extraocular muscles, except for the superior oblique muscle and the lateral rectus muscle.
2. The Edinger-Westphal nucleus supplies parasympathetic fibers to the eye via the ciliary ganglion, and controls the pupillae muscle (affecting pupil constriction) and the ciliary muscle (affecting accommodation).


Sympathetic postganglionic fibers also join the nerve from the plexus on the internal carotid artery in the wall of the cavernous sinus and are distributed through the nerve, for example, to the smooth muscle of levator palpebrae superioris.

Emergence from Brain

On emerging from the brain, the oculomotor nerve is invested with a sheath of pia mater and enclosed in a prolongation from the arachnoid mater. It passes between the superior cerebellar and posterior cerebral arteries, and then pierces the dura mater anterior and lateral to the posterior clinoid process (to give attachment to the tectorium cerebella), passing between the free and attached borders of the tentorium cerebelli.

It then runs along the lateral wall of the cavernous sinus, above the other orbital nerves, receiving in its course one or two filaments from the cavernous plexus of the sympathetic nervous system, and a communicating branch from the ophthalmic division of the trigeminal nerve.

It then divides into two branches that enter the orbit through the superior orbital fissure, between the two heads of the lateral rectus (a muscle on the lateral side of the eyeball in the orbit). Here the nerve is placed below the trochlear nerve and the frontal and lacrimal branches of the ophthalmic nerve, while the nasociliary nerve is placed between its two rami (the superior and inferior branch of oculomotor nerve).



Cranial nerves: Image of cranial nerves showing the position of the oculomotor nerve.

This page titled 12.5D: Oculomotor (III) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5E: Trochlear (IV) Nerve

The trochlear nerve (cranial nerve IV) is a motor nerve that innervates a single muscle: the superior oblique muscle of the eye.

Learning Objectives

- Describe the trochlear nerve (cranial nerve IV)


Key Points

- The trochlear nerve innervates the superior oblique muscle of the eye.
- The trochlear nerve contains the smallest number of axons of all the cranial nerves and has the greatest intracranial length.
- The two major clinical syndromes that can arise from damage to the trochlear nerve are vertical and torsional diplopia.

Key Terms

- **contralateral**: On the opposite side of the body.
- **cerebral aqueduct**: The channel in the brain that connects the third ventricle to the fourth ventricle. Also called the aqueduct of Sylvius, it is surrounded by periaqueductal gray matter.

The trochlear nerve (cranial nerve IV) is a motor nerve that innervates a single muscle: the superior oblique muscle of the eye.



The trochlear nerve: The trocheal nerve and where it innervates.

The trochlear nerve is unique among the cranial nerves in several respects.

- It is the smallest nerve in terms of the number of axons it contains and it has the greatest intracranial length.
- Other than the optic nerve (cranial nerve II), it is the only cranial nerve that decussates (crosses to the other side) before innervating its target.
- It is the only cranial nerve that exits from the dorsal aspect of the brainstem.

The nucleus of the trochlear nerve is located in the caudal mesencephalon beneath the cerebral aqueduct. It is immediately below the nucleus of the oculomotor nerve (III) in the rostral mesencephalon.

The trochlear nucleus is unique in that its axons run dorsally and cross the midline before emerging from the brainstem—so a lesion of the trochlear nucleus affects the contralateral eye. Lesions of all other cranial nuclei affect the ipsilateral side (except of course the optic nerve, cranial nerve II, which innervates both eyes).

Homologous trochlear nerves are found in all jawed vertebrates. The unique features of the trochlear nerve, including its dorsal exit from the brainstem and its contralateral innervation, are seen in the primitive brains of sharks.

The human trochlear nerve is derived from the basal plate of the embryonic midbrain.

Clinical Syndromes

There are two major clinical syndromes that can manifest through damage to the trochlear nerve:

- Vertical diplopia: Injury to the trochlear nerve causes weakness of downward eye movement with consequent vertical diplopia (double vision).
- Torsional diplopia: Weakness of intorsion results in torsional diplopia, in which two different visual fields, tilted with respect to each other, are seen at the same time. To compensate for this, patients with trochlear nerve palsies tilt their heads to the opposite side, in order to fuse the two images into a single visual field.

The clinical syndromes can originate from both peripheral and central lesions. A peripheral lesion is damage to the bundle of nerves, in contrast to a central lesion, which is damage to the trochlear nucleus.

This page titled 12.5E: Trochlear (IV) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5F: Trigeminal (V) Nerve

The trigeminal nerve is the fifth cranial nerve and it is responsible for sensation and motor function in the face and mouth.

Learning Objectives

- Describe the trigeminal nerve (cranial nerve V)


Key Points

- The sensory function of the trigeminal nerve is to provide tactile, motion, position, and pain sensations for the face and mouth; its motor function activates the muscles of the jaw, mouth, and inner ear.
- The trigeminal nerve has three major branches on each side—the opthalmic nerve, maxillary nerve, and mandibular nerve—that converge on the trigeminal ganglion.
- The trigeminal ganglion is analogous to the dorsal root ganglia of the spinal cord, which contain the cell bodies of incoming sensory fibers from the rest of the body.

Key Terms

- **trigeminal ganglion**: The trigeminal ganglion (also called the Gasserian ganglion, semilunar ganglion, or Gasser's ganglion) is a sensory ganglion of the trigeminal nerve (cranial nerve V) that occupies a cavity (Meckel's cave) in the dura mater, covering the trigeminal impression near the apex of the petrous part of the temporal bone.
- **trigeminal nerve**: The nerve responsible for sensation and motor function in the face and mouth.

The trigeminal nerve (cranial nerve V), and it contains both sensory and motor fibers. It is responsible for sensation in the face and certain motor functions such as biting, chewing, and swallowing.



Trigeminal nerve: Schematic illustration of the trigeminal nerve (labeled Sensory root above) and the structures it innervates in the face and mouth.


The sensory function of the trigeminal nerve is to provide the tactile, motion, position, and pain sensations of the face and mouth. The motor function activates the muscles of the jaw, mouth, and inner ear.

Structure

The trigeminal nerve is the largest of the cranial nerves. Its name, trigeminal, means three twins. It is derived from the fact that each nerve, one on each side of the pons, has three major branches: the ophthalmic nerve (V1 in the illustration below), the maxillary nerve (V2), and the mandibular nerve (V3).

The ophthalmic and maxillary nerves are purely sensory. The mandibular nerve has both sensory and motor functions.

The three branches converge on the trigeminal ganglion that is located within the trigeminal cave in the brain; it contains the cell bodies of incoming sensory nerve fibers. The trigeminal ganglion is analogous to the dorsal root ganglia of the spinal cord, which contain the cell bodies of incoming sensory fibers from the rest of the body.



Areas of the face innervated by the trigeminal nerve: The ophthalmic nerve branch (V1) innervates the bright red area, the maxillary nerve branch (V2) innervates the light red area, and the mandibular nerve branch (V3) innervates the yellow area.

From the trigeminal ganglion, a single large sensory root enters the brainstem at the level of the pons. Immediately adjacent to the sensory root, a smaller motor root emerges from the pons at the same level.

Motor fibers pass through the trigeminal ganglion on their way to peripheral muscles, but their cell bodies are located in the nucleus of the trigeminal nerve, deep within the pons.

Function

The sensory function of the trigeminal nerve is to provide tactile, proprioceptive, and nociceptive afferents to the face and mouth. The motor component of the mandibular division (V3) of the trigeminal nerve controls the movement of eight muscles, including the four muscles of mastication: the masseter, the temporal, and the medial and lateral pterygoids.

The other four muscles are the tensor veli palatini, the mylohyoid, the anterior belly of the digastric, and the tensor tympani. With the exception of the tensor tympani, all of these muscles are involved in biting, chewing and swallowing, and all have bilateral cortical representation.

This page titled 12.5F: Trigeminal (V) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5G: Abducens (VI) Nerve

The abducens nerve (cranial nerve VI) controls the lateral movement of the eye through innervation of the lateral rectus muscle.

Learning Objectives

- Describe the abducens nerve (cranial nerve VI)


Key Points

- The abducens nerve exits the brainstem at the junction of the pons and the medulla and runs upward to reach the eye, traveling between the dura and the skull.
- The long course of the abducens nerve between the brainstem and the eye makes it vulnerable to injury at many levels.
- In most mammals besides humans, it also innervates the musculus retractor bulbi, which can retract the eye for protection.

Key Terms

- **abducens nerve**: A nerve that controls the lateral rectus muscle in the eye.
- **clivus**: A part of the cranium at the base of the skull. It forms a gradual sloping process at the anterior-most portion of the basilar occipital bone at its junction with the sphenoid bone.
- **petrous temporal bone**: A pyramid-shaped bone that is wedged in at the base of the skull between the sphenoid and occipital bones and is part of the endocranium.
- **lateral rectus muscle**: A muscle in the orbit. It is one of six extraocular muscles that control the movements of the eye (abduction in this case) and the only muscle innervated by the abducens nerve, cranial nerve VI, functioning to bring the pupil away from the midline of the body.

The abducens nerve (cranial nerve VI) is a somatic efferent nerve that, in humans, controls the movement of a single muscle: the lateral rectus muscle of the eye that moves the eye horizontally. In most other mammals it also innervates the musculus retractor bulbi, which can retract the eye for protection. Homologous abducens nerves are found in all vertebrates except lampreys and hagfishes.



Abducens nerve: Schematic of cranial nerves showing cranial nerve VI, the abducens nerve.

The abducens nerve leaves the brainstem at the junction of the pons and the medulla, medial to the facial nerve. In order to reach the eye, it runs upward (superiorly) and then bends forward (anteriorly).

The nerve enters the subarachnoid space when it emerges from the brainstem. It runs upward between the pons and the clivus, and then pierces the dura mater to run between the dura and the skull.

At the tip of the petrous temporal bone, it makes a sharp turn forward to enter the cavernous sinus. In the cavernous sinus it runs alongside the internal carotid artery. It then enters the orbit through the superior orbital fissure and innervates the lateral rectus muscle of the eye.

The long course of the abducens nerve between the brainstem and the eye makes it vulnerable to injury at many levels. For example, fractures of the petrous temporal bone can selectively damage the nerve, as can aneurysms of the intracavernous carotid artery.

Mass lesions that push the brainstem downward can damage the nerve by stretching it between the point where it emerges from the pons and the point where it hooks over the petrous temporal bone.

This page titled 12.5G: Abducens (VI) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5H: Facial (VII) Nerve

The facial nerve (cranial nerve VII) determines facial expressions and the taste sensations of the tongue.

Learning Objectives


- Describe the facial nerve (cranial nerve VII)

Key Points

- The facial nerve (cranial nerve VII) is responsible for the muscles that determine facial expression, as well as the sensation of taste in the front of the tongue and oral cavity.
- The facial nerve's motor component begins in the facial nerve nucleus in the pons, and the sensory component begins in the nervus intermedius. The nerve then runs through the facial canal, passes through the parotid gland, and divides into five branches.
- Voluntary facial movements, such as wrinkling the brow, showing teeth, frowning, closing the eyes tightly (inability to do so is called lagophthalmos), pursing the lips, and puffing out the cheeks, all test the facial nerve.

Key Terms

- **nervus intermedius**: A part of the facial nerve (cranial nerve VII) located between the motor component of the facial nerve and the vestibulocochlear nerve (cranial nerve VIII). It contains the sensory and parasympathetic fibers of the facial nerve.
- **Bell's Palsy**: Bell's palsy is a form of facial paralysis resulting from a dysfunction of the cranial nerve VII (the facial nerve) that results in the inability to control facial muscles on the affected side.



The facial nerve: Illustration of the facial nerve and its branches.

The facial nerve is the seventh (cranial nerve VII) of the 12, paired cranial nerves. It emerges from the brainstem between the pons and the medulla and controls the muscles of facial expression.

It also functions in the conveyance of taste sensations from the anterior two-thirds of the tongue and oral cavity, and it supplies preganglionic parasympathetic fibers to several head and neck ganglia.

Location


The motor part of the facial nerve arises from the facial nerve nucleus in the pons, while the sensory part of the facial nerve arises from the nervus intermedius. The motor and sensory parts of the facial nerve enter the petrous temporal bone into the internal auditory meatus (intimately close to the inner ear), then runs a tortuous course (including two tight turns) through the facial canal, emerges from the stylomastoid foramen, and passes through the parotid gland, where it divides into five major branches.

Although it passes through the parotid gland, it does not innervate the gland (this is the responsibility of cranial nerve IX, the glossopharyngeal nerve). The facial nerve forms the geniculate ganglion prior to entering the facial canal.

The path of the facial nerve can be divided into six segments.

1. The intracranial (cisternal) segment.
2. The meatal segment (brainstem to internal auditory canal).
3. The labyrinthine segment (internal auditory canal to geniculate ganglion),
4. The tympanic segment (from geniculate ganglion to pyramidal eminence).
5. The mastoid segment (from pyramidal eminence to stylomastoid foramen).
6. The extratemporal segment (from stylomastoid foramen to post parotid branches).

Function



Bell's Palsy: A person attempting to show his teeth and raise his eyebrows with Bell's palsy on his right side (left side of the image).

Voluntary facial movements, such as wrinkling the brow, showing teeth, frowning, closing the eyes tightly (inability to do so is called lagophthalmos), pursing the lips, and puffing out the cheeks, all test the facial nerve. There should be no noticeable asymmetry.

In an upper motor neuron lesion, called central seven (central facial palsy), only the lower part of the face on the contralateral side will be affected due to the bilateral control to the upper facial muscles (frontalis and orbicularis oculi).

Lower motor neuron lesions can result in a cranial nerve VII palsy (Bell's palsy is the idiopathic form of facial nerve palsy), manifested as both upper and lower facial weakness on the same side of the lesion.

Taste can be tested on the anterior 2/3 of the tongue. This can be tested with a swab dipped in a flavored solution, or with electronic stimulation (similar to putting your tongue on a battery).

In regards to the corneal reflex, the afferent arc is mediated by the general sensory afferents of the trigeminal nerve. The efferent arc occurs via the facial nerve.

The reflex involves the consensual blinking of both eyes in response to stimulation of one eye. This is due to the facial nerve's innervation of the muscles of facial expression, namely the orbicularis oculi, responsible for blinking. Thus, the corneal reflex effectively tests the proper functioning of both cranial nerves V and VII.

This page titled 12.5H: Facial (VII) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5I: Vestibulocochlear (VIII) Nerve

The vestibulocochlear nerve (cranial nerve VIII) carries information about hearing and balance.

Learning Objectives

- Describe the vestibulocochlear nerve (cranial nerve VIII)

Key Points

- The vestibulocochlear nerve comprises the cochlear nerve that transmits hearing information, and the vestibular nerve that transmits balance information.
- The cochlear nerve travels away from the cochlea of the inner ear where it starts as the spiral ganglia.
- The vestibular nerve travels from the vestibular system of the inner ear.


Key Terms

- **cochlear nerve**: A sensory nerve that conducts information about the environment to the brain, in this case acoustic energy impinging on the tympanic membrane (sound waves reaching the ear drum). The cochlear nerve arises from within the cochlea and extends to the brainstem where its fibers make contact with the cochlear nucleus, the next stage of neural processing in the auditory system.
- **vestibulocochlear nerve**: Also known as the auditory vestibular nerve, this is the eighth of twelve cranial nerves, and it is responsible for transmitting sound and equilibrium (balance) information from the inner ear to the brain.
- **vestibular nerve**: One of the two branches of the vestibulocochlear nerve (the cochlear nerve being the other). It connects to the semicircular canals via the vestibular ganglion and receives positional information.

The vestibulocochlear nerve (also known as the auditory vestibular nerve and cranial nerve VIII) has axons that carry the modalities of hearing and equilibrium.

It consists of the cochlear nerve that carries information about hearing, and the vestibular nerve that carries information about balance.

This is the nerve along which the sensory cells (the hair cells) of the inner ear transmit information to the brain. It emerges from the pons and exits the inner skull via the internal acoustic meatus (or internal auditory meatus) in the temporal bone.



Vestibular system's semicircular canal: An illustration of the inner ear showing its semicircular canal, hair cells, ampulla, cupula, vestibular nerve, and fluid.

The vestibulocochlear nerve consists mostly of bipolar neurons and splits into two large divisions: the cochlear nerve and the vestibular nerve. The cochlear nerve travels away from the cochlea of the inner ear where it starts as the spiral ganglia.

Processes from the organ of Corti (the receptor organ for hearing) conduct afferent transmission to the spiral ganglia. It is the inner hair cells of the organ of Corti that are responsible for activating the afferent receptors in response to pressure waves reaching the basilar membrane through the transduction of sound.

The vestibular nerve travels from the vestibular system of the inner ear. The vestibular ganglion houses the cell bodies of the bipolar neurons and extends processes to five sensory organs.

Three of these are the cristae, located in the ampullae of the semicircular canals. Hair cells of the cristae activate afferent receptors in response to rotational acceleration.

The other two sensory organs supplied by the vestibular neurons are the maculae of the saccule and utricle. Hair cells of the maculae activate afferent receptors in response to linear acceleration.

The vestibulocochlear nerve has axons that carry the modalities of hearing and equilibrium. Damage to the vestibulocochlear nerve may cause hearing loss, vertigo, a false sense of motion, loss of equilibrium in dark places, nystagmus, motion sickness, and gaze-evoked tinnitus.

A benign primary intracranial tumor of vestibulocochlear nerve is called a vestibular schwannoma (also called acoustic neuroma).

This page titled 12.5I: Vestibulocochlear (VIII) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5J: Glossopharyngeal (IX) Nerve

The glossopharyngeal nerve (cranial nerve IX) serves many distinct functions, including providing sensory innervation to various head and neck structures.

Learning Objectives

- Describe the glossopharyngeal nerve (cranial nerve IX)

Key Points


- The glossopharyngeal nerve (cranial nerve IX) is responsible for swallowing and the gag reflex, along with other functions.
- The glossopharyngeal nerve receives input from the general and special sensory fibers in the back of the throat.
- The glossopharyngeal nerve has five components: branchial motor, visceral motor, visceral sensory, general sensory, and special sensory components.

Key Terms

- **otic ganglion**: A small parasympathetic ganglion located immediately below the foramen ovale in the infratemporal fossa that is associated with the glossopharyngeal nerve.
- **medulla**: The lower half of the brainstem that contains the cardiac, respiratory, vomiting, and vasomotor centers and is associated with autonomic, involuntary functions such as breathing, heart rate, and blood pressure.
- **stylopharyngeus muscle**: A muscle in the head that stretches between the temporal styloid process and the pharynx.

Structure

The glossopharyngeal nerve is the ninth of 12 pairs of cranial nerves. It exits the brainstem out from the sides of the upper medulla, just rostral (closer to the nose) to the vagus nerve.



Glossopharyngeal nerve: Image of head structures including the glossopharyngeal nerve.

The motor division of the glossopharyngeal nerve is derived from the basal plate of the embryonic medulla oblongata, while the

sensory division originates from the cranial neural crest.

Function

There are a number of functions of the glossopharyngeal nerve. It controls muscles in the oral cavity and upper throat, as well as part of the sense of taste and the production of saliva.

Along with taste, the glossopharyngeal nerve relays general sensations from the pharyngeal walls. The various functions of the glossopharyngeal nerve are that:

- It receives general sensory fibers (ventral trigeminothalamic tract) from the tonsils, the pharynx, the middle ear, and the posterior 1/3 of the tongue.
- It receives special sensory fibers (taste) from the posterior 1/3 of the tongue.
- It receives visceral sensory fibers from the carotid bodies, carotid sinus.
- It supplies parasympathetic fibers to the parotid gland via the otic ganglion.
- It supplies motor fibers to the stylopharyngeus muscle.
- It contributes to the pharyngeal plexus.

Five Functional Components

The glossopharyngeal nerve consists of five components with distinct functions:

1. Branchial motor (special visceral efferent): Supplies the stylopharyngeus muscle.
2. Visceral motor (general visceral efferent): Provides parasympathetic innervation of the parotid gland.
3. Visceral sensory (general visceral afferent): Carries visceral sensory information from the carotid sinus and body.
4. General sensory (general somatic afferent): Provides general sensory information from the skin of the external ear, internal surface of the tympanic membrane, upper pharynx, and the posterior 1/3 of the tongue.
5. Special sensory (special afferent): Provides taste sensation from the posterior 1/3 of the tongue.

This page titled 12.5J: Glossopharyngeal (IX) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5K: Vagus (X) Nerve

The vagus nerve (cranial nerve X) is responsible for parasympathetic output to the heart and visceral organs.

Learning Objectives

- Describe the vagus nerve (cranial nerve X)

Key Points


- The vagus nerve (cranial nerve X) sends information about the body's organs to the brain and carries some motor information back to the organs.
- The vagus nerve has axons that originate from or enter the dorsal nucleus of the vagus nerve, the nucleus ambiguus, the solitary nucleus in the medulla, and the spinal trigeminal nucleus.
- The vagus nerve is responsible for heart rate, gastrointestinal peristalsis, and sweating, to name a few.

Key Terms

- **nucleus ambiguus**: A region of histologically disparate cells located just dorsal (posterior) to the inferior olivary nucleus in the lateral portion of the upper (rostral) medulla. It receives upper motor neuron innervation directly via the corticobulbar tract.
- **Valsalva maneuver**: Performed by making a moderately forceful attempt to exhale against a closed airway; this is usually done by closing one's mouth and pinching one's nose shut while pressing out as if blowing up a balloon.
- **solitary nucleus**: Structures in the brainstem that carry and receive visceral sensation and taste from the facial (VII), glossopharyngeal (IX), and vagus (X) cranial nerves.
- **dorsal nucleus of the vagus nerve**: Also called the posterior motor nucleus of vagus, this is a cranial nerve nucleus for the vagus nerve in the medulla that lies under the floor of the fourth ventricle. It mostly serves parasympathetic vagal functions in the gastrointestinal tract, lungs, and other thoracic and abdominal vagal innervations.
- **spinal trigeminal nucleus**: A nucleus in the medulla that receives information about deep/crude touch, pain, and temperature from the ipsilateral face.

Vagus Nerve Anatomy

The vagus nerve, also known as the pneumogastric nerve or cranial nerve X, is the tenth of twelve paired cranial nerves. Upon leaving the medulla between the medullary pyramid and the inferior cerebellar peduncle, it extends through the jugular foramen, then passes into the carotid sheath between the internal carotid artery and the internal jugular vein below the head, to the neck, chest and abdomen, where it contributes to the innervation of the viscera.



Vagus nerve: Diagram demonstrating the course of the vagus nerve.

Besides output to the various organs in the body, the vagus nerve conveys sensory information about the state of the body's organs to the central nervous system. Eighty to 90% of the nerve fibers in the vagus nerve are afferent (sensory) nerves that communicate the state of the viscera to the brain.

The vagus nerve includes axons that emerge from or converge onto four nuclei of the medulla.

- The dorsal nucleus of vagus nerve: Sends parasympathetic output to the viscera, especially the intestines.
- The nucleus ambiguus: Sends parasympathetic output to the heart (slowing it down).
- The solitary nucleus: Receives afferent taste information and primary afferents from visceral organs.
- The spinal trigeminal nucleus: Receives information about deep/crude touch, pain, and temperature of the outer ear, the dura of the posterior cranial fossa, and the mucosa of the larynx.

Function

The vagus nerve supplies motor parasympathetic fibers to all the organs, except the suprarenal (adrenal) glands, from the neck down to the second segment of the transverse colon. The vagus also controls a few skeletal muscles, most notably:

- Cricothyroid muscle.
- Levator veli palatini muscle.
- Salpingopharyngeus muscle.
- Palatoglossus muscle.
- Palatopharyngeus muscle.
- Superior, middle, and inferior pharyngeal constrictors.
- Muscles of the larynx (speech).

This means that the vagus nerve is responsible for such varied tasks as heart rate, gastrointestinal peristalsis, sweating, and quite a few muscle movements in the mouth, including speech (via the recurrent laryngeal nerve), swallowing, and keeping the larynx open for breathing (via action of the posterior cricoarytenoid muscle, the only abductor of the vocal folds).

It also has some afferent fibers that innervate the inner (canal) portion of the outer ear, via the auricular branch (also known as Alderman's nerve) and part of the meninges. This explains why a person may cough when tickled on the ear (such as when trying to remove ear wax with a cotton swab).

Afferent vagus nerve fibers that innervate the pharynx and back of the throat are responsible for the gag reflex. In addition, 5-HT3 receptor-mediated afferent vagus stimulation in the gut due to gastroenteritis and other insults is a cause of vomiting.

Cardiovascular Influence

Parasympathetic innervation of the heart is partially controlled by the vagus nerve and is shared by the thoracic ganglia. Activation of the vagus nerve typically leads to a reduction in heart rate and/or blood pressure.

This occurs commonly in cases of viral gastroenteritis, acute cholecystitis, or in response to stimuli such as the Valsalva maneuver or pain. Excessive activation of the vagal nerve during emotional stress can also cause vasovagal syncope due to a sudden drop in cardiac output, causing cerebral hypoperfusion.

This page titled 12.5K: Vagus (X) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5L: Accessory (XI) Nerve

The accessory nerve (cranial nerve XI) controls the muscles of the shoulder and neck.

Learning Objectives

- Describe the accessory nerve (cranial nerve XI)

Key Points

- Cranial nerve XI is responsible for tilting and rotating the head, elevating the shoulders, and adducting the scapula.
- Most of the fibers of the accessory nerve originate in neurons situated in the upper spinal cord. The fibers that make up the accessory nerve enter the skull through the foramen magnum and proceed to exit the jugular foramen with cranial nerves IX and X.
- Due to its unusual course, the accessory nerve is the only nerve that enters and exits the skull.


Key Terms

- **sternocleidomastoid**: A paired muscle in the superficial layers of the side of the neck. Its actions of the muscle are rotation of the head to the opposite side and flexion of the neck.
- **trapezius**: A large vertebrate skeletal muscle divided into an ascending, descending, and transverse portion, attaching the neck and central spine to the outer extremity of the scapula; it functions in scapular elevation, adduction, and depression.
- **accessory nerve**: The 11th cranial nerve of higher vertebrates, it controls the pharynx and the muscles of the upper chest and shoulders.
- **foramen**: An opening, an orifice, or a short passage.

Anatomic Description

The accessory nerve (cranial nerve XI) controls the sternocleidomastoid and trapezius muscles of the shoulder and neck. It begins in the central nervous system (CNS) and exits the cranium through a foramen.

Unlike the other 11 cranial nerves, the accessory nerve begins outside the skull. In fact, most of the fibers of the nerve originate in neurons situated in the upper spinal cord.



The accessory nerve: Upon exiting the skull via the jugular foramen, the spinal accessory nerve pierces the sternocleidomastoid muscle before terminating on the trapezius muscle.

The fibers that make up the accessory nerve enter the skull through the foramen magnum and proceed to exit the jugular foramen with cranial nerves IX and X. Due to its unusual course, the accessory nerve is the only nerve that enters and exits the skull.

Traditional descriptions of the accessory nerve divide it into two components: a spinal component and a cranial component. However, more modern characterizations of the nerve regard the cranial component as separate and part of the vagus nerve.

Therefore, in contemporary discussions of the accessory nerve, it is common to disregard the cranial component when referencing the accessory nerve and assume reference to the spinal accessory nerve.

Accessory Nerve Function

The accessory nerve provides motor innervation from the CNS to the sternocleidomastoid and trapezius muscles of the neck. The sternocleidomastoid muscle tilts and rotates the head, while the trapezius muscle has several actions on the scapula, including shoulder elevation and adduction of the scapula.

During neurological examinations, the function of the spinal accessory nerve is often measured by testing the range of motion and strength of the aforementioned muscles. Limited range of motion or diminished muscle strength often indicates injury of the accessory nerve.

Patients with spinal accessory nerve palsy may exhibit signs of lower motor neuron disease, such as atrophy and fasciculations of both the sternocleidomastoid and trapezius muscles.

This page titled 12.5L: Accessory (XI) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.5M: Hypoglossal (XII) Nerve

The hypoglossal nerve (cranial nerve XII) controls the muscles of the tongue.

Learning Objectives

- Describe the hypoglossal nerve (cranial nerve XII)

Key Points

- It controls tongue movements of speech, food manipulation, and swallowing.
- While the hypoglossal nerve controls the tongue's involuntary activities of swallowing to clear the mouth of saliva, most of the functions it controls are voluntary, meaning that the execution of these activities requires conscious thought.
- Proper function of the hypoglossal nerve is important for executing tongue movements associated with speech. Many languages require specific uses of the nerve to create unique speech sounds, which may contribute to the difficulties some adults encounter when learning a new language.


Key Terms

- **digastric muscle**: A small muscle located under the jaw.
- **hypoglossal nerve**: The twelfth cranial nerve (XII) controls the muscles of the tongue.
- **olivary body**: Located on the anterior surface of the medulla, lateral to the pyramid.
- **medullary pyramid**: Paired white matter structures of the brainstem's medulla oblongata that contain motor fibers of the corticospinal and corticobulbar tracts.

Structure and Location

The hypoglossal nerve is the twelfth cranial nerve (XII) and innervates all extrinsic and intrinsic muscles of the tongue, except for the palatoglossus. The hypoglossal nerve emerges from the medulla oblongata in the preolivary sulcus where it separates the olive (olivary body) and the pyramid (medullary pyramid).

It goes on to traverse the hypoglossal canal and, upon emerging, it branches and merges with a branch from the anterior ramus of C1. It passes behind the vagus nerve and between the internal carotid artery and internal jugular vein which lies on the carotid sheath. After passing deep to the posterior belly of the digastric muscle it proceeds to the submandibular region to enter the tongue.



Hypoglossal nerve: Schematic image of the hypoglossal nerve and the structures it innervates.

Function

The hypoglossal nerve controls tongue movements of speech, food manipulation, and swallowing. It supplies motor fibers to all of the muscles of the tongue, with the exception of the palatoglossus muscle, which is innervated by the vagus nerve (cranial nerve X) or, according to some classifications, by fibers from the glossopharyngeal nerve (cranial nerve IX) that hitchhike within the vagus.

While the hypoglossal nerve controls the tongue's involuntary activities of swallowing to clear the mouth of saliva, most of the functions it controls are voluntary, meaning that the execution of these activities requires conscious thought.

Proper function of the hypoglossal nerve is important for executing the tongue movements associated with speech. Many languages require specific and sometimes unusual uses of the nerve to create unique speech sounds, which may contribute to the difficulties some adults encounter when learning a new language. Several corticonuclear-originating fibers supply innervation and aid in the unconscious movements required upon engaging in speech and articulation.

Progressive bulbar palsy is a neuromuscular atrophy associated with the combined lesions of the hypoglossal nucleus and the nucleus ambiguous, upon atrophy of the motor nerves of the pons and medulla. This condition causes dysfunctional tongue movements that lead to speech and chewing impairments and swallowing difficulties. Tongue muscle atrophy may also occur.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cranial Nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cranial_nerves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peristalsis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peristalsis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_...Nervous_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peripheral nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/periphe...rvous%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br..._labels_en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ethmoid Bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ethmoid_bone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Olfactory_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- olfactory mucosa. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/olfactory%20mucosa**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- olfactory receptors. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/olfactory%20receptors**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br..._labels_en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:H...ry_bulb_en.png**. **License**: ***CC BY: Attribution***
- Oligodendrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Oligodendrocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pretectal Area. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pretectal_area**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Optic nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Optic_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- retina. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/retina**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- visual cortex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/visual%20cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- optic nerve. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/optic_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br..._labels_en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:H...ry_bulb_en.png**. **License**: ***CC BY: Attribution***
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...ptic_Tract.jpg**. **License**: ***CC BY: Attribution***
- Oculomotor nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Oculomotor_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- superior orbital fissure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/superio...ital%20fissure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plexus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- levator palpebrae superioris. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/levato...rae_superioris**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br..._labels_en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:H...ry_bulb_en.png**. **License**: ***CC BY: Attribution***
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...ptic_Tract.jpg**. **License**: ***CC BY: Attribution***
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:B..._labels_en.svg**. **License**: ***CC BY: Attribution***
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Trochlear_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Trochlear_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- contralateral. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/contralateral**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebral aqueduct. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebral_aqueduct**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Br..._labels_en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:H...ry_bulb_en.png**. **License**: ***CC BY: Attribution***

- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...ptic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:B..._labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trigeminal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trigeminal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- trigeminal ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/trigeminal%20ganglion. **License**: *CC BY-SA: Attribution-ShareAlike*
- trigeminal nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/trigeminal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Br..._labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:H...ry_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...ptic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:B..._labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Tr...nnervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Clivus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clivus_(anatomy). **License**: *CC BY-SA: Attribution-ShareAlike*
- Abducent nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Abducent_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- lateral rectus muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lateral...ectus%20muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- abducens nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/abducens_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Br..._labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:H...ry_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...ptic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:B..._labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Tr...nnervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bell's Palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bell's%20Palsy. **License**: *CC BY-SA: Attribution-ShareAlike*
- nervus intermedius. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/nervus%20intermedius. **License**: *CC BY-SA: Attribution-ShareAlike*

- facial nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/facial_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Br..._labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:H...ry_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...ptic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Tr...nnervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bellspalsy.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY: Attribution*
- Vestibulocochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vestibulocochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- cochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cochlear%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- vestibulocochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vestibu...chlear%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- vestibular nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vestibular%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Br..._labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:H...ry_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:1204_Optic_Nerve_vs_Optic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Tr...nnervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bellspalsy.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY: Attribution*
- Vestibular system's semicircular canal-a cross-section. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Vestibular_system%27s_semicircular_canal-_a_cross-section.jpg. **License**: *Public Domain: No Known Copyright*
- Stylopharyngeus Muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stylopharyngeus_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Peripheral Nervous System. **Provided by**: Openstax. **Located at**: http://cnx.org/contents/Hw6GpH20@1/P...Nervous-System. **License**: *CC BY-SA: Attribution-ShareAlike*
- Otic Ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Otic_ganglion. **License**: *CC BY-SA: Attribution-ShareAlike*
- Glossopharyngeal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glossopharyngeal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*

- medulla. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/medulla. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Br..._labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Head_olfactory_nerve_-_olfactory_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:1204_Optic_Nerve_vs_Optic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trig_innervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bellspalsy.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY: Attribution*
- Vestibular system's semicircular canal-a cross-section. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Vestibular_system%27s_semicircular_canal-_a_cross-section.jpg. **License**: *Public Domain: No Known Copyright*
- Gray794. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray794.png. **License**: *Public Domain: No Known Copyright*
- Spinal Trigeminal Nucleus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_trigeminal_nucleus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Valsalva Maneuver. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Valsalva_maneuver. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vagus nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vagus_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsal nucleus of the vagus nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsal%...0vagus%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- solitary nucleus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/solitary%20nucleus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...cleus-ambiguus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Head_olfactory_nerve_-_olfactory_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:1204_Optic_Nerve_vs_Optic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trig_innervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bellspalsy.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY: Attribution*
- Vestibular system's semicircular canal-a cross-section. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Vestibular_system%27s_semicircular_canal-_a_cross-section.jpg. **License**: *Public Domain: No Known Copyright*

- Gray794. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray794.png. **License**: *Public Domain: No Known Copyright*
- Recurrent Laryngeal Nerve. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Recurrent_laryngeal_nerve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sternocleidomastoid Muscle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sternocleidomastoid_muscle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Accessory nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Accessory_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Accessory nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Accessory_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Accessory nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Accessory_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- foramen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/foramen. **License**: *CC BY-SA: Attribution-ShareAlike*
- trapezius. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/trapezius. **License**: *CC BY-SA: Attribution-ShareAlike*
- accessory nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/accessory_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Head_olfactory_nerve_-_olfactory_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:1204_Optic_Nerve_vs_Optic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Trochlear_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trig_innervation.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray778.png. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray785.png. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bellspalsy.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Facial_nerve. **License**: *CC BY: Attribution*
- Vestibular system's semicircular canal-a cross-section. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Vestibular_system%27s_semicircular_canal-_a_cross-section.jpg. **License**: *Public Domain: No Known Copyright*
- Gray794. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray794.png. **License**: *Public Domain: No Known Copyright*
- Recurrent Laryngeal Nerve. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Recurrent_laryngeal_nerve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Accessory nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Accessory_nerve. **License**: *Public Domain: No Known Copyright*
- Olivary Body. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Olivary_body. **License**: *CC BY-SA: Attribution-ShareAlike*
- Medullary Pyramids. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Medullary_pyramids_(brainstem). **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypoglossal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoglossal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- hypoglossal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/hypoglossal%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain human normal inferior view with labels en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Olfactory Bulb. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Head_olfactory_nerve_-_olfactory_bulb_en.png. **License**: *CC BY: Attribution*
- Optic Nerve vs Optic Tract. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/

File:1204_Optic_Nerve_vs_Optic_Tract.jpg. **License**: *CC BY: Attribution*
- Brain Human Normal Inferior view with labels. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Brain_human_normal_inferior_view_with_labels_en.svg**. **License**: *CC BY: Attribution*
- Trochlear nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Trochlear_nerve**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trig innervation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Trig_innervation.svg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray778 (Trigeminal Nerve). **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Gray778.png**. **License**: *Public Domain: No Known Copyright*
- Gray785 (Abducens nerve). **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Gray785.png**. **License**: *Public Domain: No Known Copyright*
- Bell's palsy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bellspalsy.JPG**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Facial nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Facial_nerve**. **License**: *CC BY: Attribution*
- Vestibular system's semicircular canal-a cross-section. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Vestibular_system%27s_semicircular_canal-_a_cross-section.jpg**. **License**: *Public Domain: No Known Copyright*
- Gray794. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray794.png**. **License**: *Public Domain: No Known Copyright*
- Recurrent Laryngeal Nerve. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Recurrent_laryngeal_nerve.svg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Accessory nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Accessory_nerve**. **License**: *Public Domain: No Known Copyright*
- Lawrence_1960_17.26. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lawrence_1960_17.26.png**. **License**: *Public Domain: No Known Copyright*

This page titled 12.5M: Hypoglossal (XII) Nerve is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6A: Overview of the Spinal Nerves

Spinal nerves, a part of the peripheral nervous system (PNS), are mixed nerves that send motor, sensory, and autonomic signals between the CNS and the body.

Learning Objectives

- Describe spinal nerves of the peripheral nervous system

Key Points

- Afferent sensory axons bring sensory information from the body to the spinal cord and brain; they travel through the dorsal roots of the spinal cord.
- Efferent motor axons bring motor information from the brain to the body; they travel through the ventral roots of the spinal cord.
- All spinal nerves—except the first pair—emerge from the spinal column through an opening between vertebrae, called an intervertebral foramen.
- The spinal nerves are typically labeled by their location in the body: thoracic, lumbar, or sacral.


Key Terms

- **ventral root**: Also called the anterior root, it is the efferent motor root of a spinal nerve.
- **autonomic**: Acting or occurring involuntarily, without conscious control.
- **dorsal root**: Also known as the posterior root, the afferent sensory root of a spinal nerve.
- **intervertebral foramen**: The foramen allows for the passage of the spinal nerve root, dorsal root ganglion, the spinal artery of the segmental artery, the communicating veins between the internal and external plexuses, recurrent meningeal (sinu-vertebral) nerves, and transforaminal ligaments.

Spinal Nerve Anatomy


The term spinal nerve generally refers to a mixed spinal nerve that carries motor, sensory, and autonomic signals between the spinal cord and the body.

Humans have 31 left–right pairs of spinal nerves, each roughly corresponding to a segment of the vertebral column: eight cervical spinal nerve pairs (C1–C8), 12 thoracic pairs (T1–T12), five lumbar pairs (L1–L5), five sacral pairs (S1–S5), and one coccygeal pair. The spinal nerves are part of the peripheral nervous system (PNS).



A spinal nerve: Spinal nerves arise from a combination of nerve fibers from the dorsal and ventral roots of the spinal cord.

Location



Intervertebral foramina: Intervertebral foramina are indicated by arrows.

Each spinal nerve is formed by the combination of nerve fibers from the dorsal and ventral roots of the spinal cord. The dorsal roots carry afferent sensory axons, while the ventral roots carry efferent motor axons.

The spinal nerve emerges from the spinal column through an opening (intervertebral foramen) between adjacent vertebrae.

This is true for all spinal nerves except for the first spinal nerve pair, which emerges between the occipital bone and the atlas (the first vertebra). Thus the cervical nerves are numbered by the vertebra below, except C8, which exists below C7 and above T1.

The thoracic, lumbar, and sacral nerves are then numbered by the vertebra above. In the case of a lumbarized S1 vertebra (i.e., L6) or a sacralized L5 vertebra, the nerves are typically still counted to L5 and the next nerve is S1.

Spinal Nerve Innervation

Outside the vertebral column, the nerve divides into branches. The dorsal ramus contains nerves that serve the dorsal portions of the trunk; it carries visceral motor, somatic motor, and somatic sensory information to and from the skin and muscles of the back (epaxial muscles).

The ventral ramus contains nerves that serve the remaining ventral parts of the trunk and the upper and lower limbs (hypaxial muscles); they carry visceral motor, somatic motor, and sensory information to and from the ventrolateral body surface, structures in the body wall, and the limbs.

The meningeal branches (recurrent meningeal or sinuvertebral nerves) branch from the spinal nerve and re-enter the intervertebral foramen to serve the ligaments, dura, blood vessels, intervertebral discs, facet joints, and periosteum of the vertebrae.

The rami communicantes contain autonomic nerves that serve visceral functions, such as carrying visceral motor and sensory information to and from the visceral organs.

Cervical Nerves

The posterior distribution of the cervical nerves includes the suboccipital nerve (C1), the greater occipital nerve (C2), and the third occipital nerve (C3). The anterior distribution includes the cervical plexus (C1–C4) and brachial plexus (C5–T1).

The muscles innervated by the cervical nerves are the sternohyoid, sternothyroid, and omohyoid muscles.

A loop of nerves called ansa cervicalis is also part of the cervical plexus.

Thoracic Nerves

Thoracic nerve branches exit the spine and go directly to the paravertebral ganglia of the autonomic nervous system, where they are involved in the functions of organs and glands in the head, neck, thorax, and abdomen.

Anterior Divisions

The intercostal nerves come from thoracic nerves T1–T11, and run between the ribs. The subcostal nerve comes from nerve T12, and runs below the twelfth rib.

Posterior Divisions

The medial branches (ramus medialis) of the posterior branches of the upper six thoracic nerves run between the semispinalis dorsi and multifidus, which they supply.

They then pierce the rhomboid and trapezius muscles, and reach the skin by the sides of the spinous processes. This branch is called the medial cutaneous ramus.

The medial branches of the lower six thoracic nerves are distributed chiefly to the multifidus and longissimus dorsi, occasionally they give off filaments to the skin near the middle line. This sensitive branch is called the posterior cutaneous ramus.

Lumbar Nerves

The lumbar nerves are divided into posterior and anterior divisions.

Posterior Divisions

The medial branches of the posterior divisions of the lumbar nerves run close to the articular processes of the vertebrae and end

in the multifidus muscle. The lateral branches supply the erector spinae muscles.

Anterior Divisions

The anterior divisions of the lumbar nerves (rami anteriores) consist of long, slender branches that accompany the lumbar arteries around the sides of the vertebral bodies, beneath the psoas major.

The first and second, and sometimes the third and fourth, lumbar nerves are each connected with the lumbar part of the sympathetic trunk by a white ramus communicans.

The nerves pass obliquely outward behind the psoas major, or between its fasciculi, distributing filaments to it and the quadratus lumborum.

The first three and the greater part of the fourth are connected by anastomotic loops and form the lumbar plexus.

The smaller part of the fourth joins with the fifth to form the lumbosacral trunk, which assists in the formation of the sacral plexus. The fourth nerve is named the furcal nerve, from the fact that it is subdivided between the two plexuses.

Sacral Nerves

There are five paired sacral nerves, half of them arising through the sacrum on the left side and the other half on the right side. Each nerve emerges in two divisions: one division through the anterior sacral foramina and the other division through the posterior sacral foramina.

The sacral nerves have both afferent and efferent fibers, thus they are responsible for part of the sensory perception and the movements of the lower extremities of the human body.

The pudendal nerve and parasympathetic fibers arise from S2, S3, and S4. They supply the descending colon and rectum, urinary bladder, and genital organs. These pathways have both afferent and efferent fibers.

Coccygeal Nerve

The coccygeal nerve is the 31st pair of spinal nerves and arises from the conus medullaris. Its anterior root helps form the coccygeal plexus.

Function

Spinal nerve motor functions are summarized in the table below.

Actions of the spinal nerves	
Level	**Motor Function**
C1–C6	Neck flexors
C1–T1	Neck extensors
C3, C4, C5	Supply diaphragm (mostly C4)
C5, C6	Move shoulder, raise arm (deltoid); flex elbow (biceps)
C6	Externally rotate (supinate) the arm
C6, C7	Extend the elbow and wrist (triceps and wrist extensors); pronate wrist
C7, C8	Flex wrist; supply small muscles of the hand
T1–T6	Intercostals and trunk above the waist
T7–L1	Abdominal muscles

Actions of the spinal nerves	
Level	**Motor Function**
L1–L4	Flex thigh
L2, L3, L4	Adduct thigh; extend leg at the knee (quadriceps femoris)
L4, L5, S1	Abduct thigh; flex leg at the knee (hamstrings); dorsiflex foot (tibialis anterior); extend toes
L5, S1, S2	Extend leg at the hip (gluteus maximus); plantar flex foot and flex toes

This page titled 12.6A: Overview of the Spinal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6B: Branches of Spinal Nerves

The spinal nerves branch into the dorsal ramus, ventral ramus, the meningeal branches, and the rami communicantes.

Learning Objectives


- Describe branches of the peripheral nervous system

Key Points

- The dorsal and ventral rami contain nerves that provide visceral motor, somatic motor, and sensory information, with the dorsal ramus feeding the dorsal trunk (skin and muscles of the back), and the ventral ramus feeding the ventral trunk and limbs through the ventrolateral surface.
- The meningeal branches supply nerve function to the vertebrae themselves, including the ligaments, dura, blood vessels, intervertebral discs, facet joints, and periosteum.
- The rami communicantes contain autonomic nerves that carry visceral motor and sensory information to and from the visceral organs.

Key Terms

- **nerve plexus**: A branching network of intersecting nerves.
- **dorsal ramus**: The posterior (or dorsal) branches (or divisions) of the spinal nerves are, as a rule, smaller than the anterior divisions. They are also referred to as the dorsal rami. They are directed backwards and—with the exceptions of those of the first cervical, the fourth and fifth sacral, and the coccygeal—divide into medial and lateral branches for the supply of the muscles and skin of the posterior part of the trunk.
- **meningeal branches**: Also known as recurrent meningeal nerves, sinuvertebral nerves, or recurrent nerves of Luschka, these are a number of small nerves that branch from the spinal nerve (or the posterior ramus) near the origin of the anterior and posterior rami. They then re-enter the intervertebral foramen, and innervate the facet joints, the annulus fibrosus of the intervertebral disk, and the ligaments and periosteum of the spinal canal, carrying pain sensation.



Course and branches of thoracic spinal nerve: This diagram depicts the course and branches of a typical thoracic spinal nerve. The posterior division (dorsal ramus) is labeled at the top right.

Outside the vertebral column, the spinal nerves divide into branches.

- The dorsal ramus: Contains nerves that serve the dorsal portions of the trunk carrying visceral motor, somatic motor, and sensory information to and from the skin and muscles of the back.
- The ventral ramus: Contains nerves that serve the remaining ventral parts of the trunk and the upper and lower limbs carrying visceral motor, somatic motor, and sensory information to and from the ventrolateral body surface, structures in the body wall, and the limbs.
- Some ventral rami merge with adjacent ventral rami to form a nerve plexus, a network of interconnecting nerves. Nerves emerging from a plexus contain fibers from various spinal nerves, which are now carried together to some target location. Major plexuses include the cervical, brachial, lumbar, and sacral plexuses.
- The meningeal branches (recurrent meningeal or sinuvertebral nerves): These branch from the spinal nerve and re-enter the intervertebral foramen to serve the ligaments, dura, blood vessels, intervertebral discs, facet joints, and periosteum of the vertebrae.
- The rami communicantes: Contain autonomic nerves that carry visceral motor and sensory information to and from the visceral organs.

This page titled 12.6B: Branches of Spinal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6C: Plexuses

A nerve plexus is a network of intersecting nerves that serve the same part of the body.

Learning Objectives

- Describe nerve plexuses in the peripheral nervous system

Key Points

- Nerve plexuses throughout the body tend to be named after the area in which the plexus occurs and the organs, limbs, and tissues it serves. Examples include the cervical, brachial, lumbar, sacral, celiac, and coccygeal plexuses.
- Auerbach's plexus, which serves the gastrointestinal tract, is named after the first person to describe this plexus, Leopold Auerbach, rather than the area of the body it serves.
- The brachial plexus serves the chest, shoulders, arms and hands and is formed by the ventral rami of C5, C8, and T1 spinal nerves, and the lower and upper halves of the C4 and T2 spinal nerves.

Key Terms

- **nerve plexus**: A network of intersecting nerves.
- **brachial plexus**: A network of nerve fibers, running from the spine, formed by the ventral rami of the lower four cervical and first thoracic nerve roots (C5–C8, T1). It proceeds through the neck, the axilla (armpit region), and into the arm. It also supplies the brachium, the antebrachium, and the hand.
- **coccygeal plexus**: A plexus of nerves near the coccyx bone.
- **Autonomic**: A division of the peripheral nervous system that influences the function of internal organs.

A nerve plexus is a network of intersecting nerves; multiple nerve plexuses exist in the body. Nerve plexuses are composed of afferent and efferent fibers that arise from the merging of the anterior rami of spinal nerves and blood vessels.

There are five spinal nerve plexuses—except in the thoracic region—as well as other forms of autonomic plexuses, many of which are a part of the enteric nervous system.

Spinal Plexuses

Cervical Plexus—Serves the Head, Neck and Shoulders

The cervical plexus is formed by the ventral rami of the upper four cervical nerves and the upper part of fifth cervical ventral ramus. The network of rami is located deep within the neck.

Brachial Plexus—Serves the Chest, Shoulders, Arms and Hands

The brachial plexus is formed by the ventral rami of C5–C8 and the T1 spinal nerves, and lower and upper halves of the C4 and T2 spinal nerves. The plexus extends toward the armpit (axilla).

Lumbar Plexus—Serves the Back, Abdomen, Groin, Thighs, Knees, and Calves

The lumbar plexus is formed by the ventral rami of L1–L5 spinal nerves with a contribution of T12 form the lumbar plexus. This plexus lies within the psoas major muscle.

Sacral Plexus—Serves the Pelvis, Buttocks, Genitals, Thighs, Calves, and Feet

The sacral plexus is formed by the ventral rami of L4-S3, with parts of the L4 and S4 spinal nerves. It is located on the posterior wall of the pelvic cavity.


Coccygeal Plexus—Serves a Small Region over the Coccyx

The coccygeal plexus serves a small region over the coccyx and originates from S4, S5, and Co1 spinal nerves. It is interconnected with the lower part of sacral plexus.


In addition, the celiac plexus serves the internal organs, and Auerbach's plexus serves the gastrointestinal tract.

Autonomic Plexuses

- Celiac plexus (solar plexus)—Serves internal organs.
- Auerbach's plexus—Serves the gastrointestinal tract.
- Meissner's plexus (submucosal plexus)—Serves the gastrointestinal tract.



Brachial plexus: Cervical (C5–C8) and thoracic (T1) nerves comprise the brachial plexus, which is a nerve plexus that provides sensory and motor function to the shoulders and upper limbs.



Lumbar plexus: The lumbar plexus is comprised of the ventral rami of the lumbar spinal nerves (L1–L5) and a contribution from thoracic nerve (T12). The posterior (green) and anterior (yellow) divisions of the lumbar plexus are shown in the diagram.

This page titled 12.6C: Plexuses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6D: Intercostal Nerves

The anterior divisions of the thoracic spinal nerves (T1–T11) are called the intercostal nerves.

Learning Objectives

- Describe the function of intercostal nerves


Key Points

- The intercostal nerves are part of the somatic nervous system. This enables them to control the contraction of muscles, as well as provide specific sensory information regarding the skin and parietal pleura.
- Intercostal nerves connect to the appropriate ganglion in the sympathetic trunk through rami communicantes and serve the thoracic pleura and the abdominal peritoneum.
- Unlike most other anterior divisions of spinal nerves, the intercostal nerves do not form a plexus.

Key Terms

- **thoracic spinal nerves**: The spinal nerves emerging from the thoracic vertebrae. Branches also exit the spine and go directly to the sympathetic chain ganglia of the autonomic nervous system where they are involved in the functions of organs and glands in the head, neck, thorax, and abdomen.
- **sympathetic trunk**: Also called the sympathetic chain or gangliated cord, these are a paired bundle of nerve fibers that run from the base of the skull to the coccyx.
- **abdominal peritoneum**: The serous membrane that forms the lining of the abdominal cavity. It covers most of the intra-abdominal organs. It is composed of a layer of mesothelium supported by a thin layer of connective tissue. The peritoneum supports the abdominal organs and serves as a conduit for their blood and lymph vessels and nerves.

The intercostal nerves are part of the somatic nervous system and arise from anterior divisions (rami anteriores, ventral divisions) of the thoracic spinal nerves T1 to T11. The intercostal nerves are distributed chiefly to the thoracic pleura and abdominal peritoneum.



Intercostal nerves: An image of the intercostal brachial nerves.

They differ from the anterior divisions of the other spinal nerves in that each pursues an independent course without plexus formation.

First Thoracic Nerve

The anterior division of the first thoracic nerve divides into two branches:

1. The larger branch leaves the thorax in front of the neck of the first rib and enters the brachial plexus.
2. The other smaller branch, the first intercostal nerve, runs along the first intercostal space and ends on the front of the chest as the first anterior cutaneous branch of the thorax.

The Upper Thoracic Nerves (2nd–6th)

These are limited in their distribution to the parietes (wall) of the thorax. The anterior divisions of the second, third, fourth, fifth, and sixth thoracic nerves, and the small branch from the first thoracic, are confined to the walls of the thorax and are named thoracic intercostal nerves.

Near the sternum, they cross in front of the internal mammary artery and transversus thoracis muscle, pierce the intercostales interni, the anterior intercostal membranes, and pectoralis major, and supply the integument of the front of the thorax and over the mamma, forming the anterior cutaneous branches of the thorax.

The branch from the second nerve unites with the anterior supraclavicular nerves of the cervical plexus.

The Lower Thoracic Nerves (7th–12th)

The seventh intercostal nerve terminates at the xyphoid process, at the lower end of the sternum.

The anterior divisions of the seventh, eighth, ninth, tenth, and eleventh thoracic intercostal nerves are continued anteriorly from the intercostal spaces into the abdominal wall; hence they are named thoraco-abdominal nerves or thoracicoabdominal intercostal nerves.

The tenth intercostal nerve terminates at the umbilicus.

The twelfth (subcostal) thoracic nerve is distributed to the abdominal wall and groin.

Unlike the nerves from the autonomic nervous system that innervate the visceral pleura of the thoracic cavity, the intercostal nerves arise from the somatic nervous system. This enables them to control the contraction of muscles, as well as provide specific sensory information regarding the skin and parietal pleura.

This explains why damage to the internal wall of the thoracic cavity can be felt as a sharp pain localized in the injured region. Damage to the visceral pleura is experienced as an unlocalized ache.

This page titled 12.6D: Intercostal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6E: Dermatomes

A dermatome is an area of skin that is supplied by a single spinal nerve, and a myotome is a group of muscles that a single spinal nerve root innervates.

Learning Objectives

- Describe dermatomes and how they relate to the peripheral nervous system


Key Points

- There are eight cervical nerves, twelve thoracic nerves, five lumbar nerves and five sacral nerves. Each of these nerves relays sensation, including pain, from a particular region of skin to the brain.
- Along the thorax and abdomen, the dermatomes are like a stack of discs, with each section supplied by a different spinal nerve. The pattern is different along the arms and the legs. The dermatomes run longitudinally along the limbs, so that each half of the limb has a different dermatome.
- Dermatomes have clinical significance, especially in the diagnosis of certain diseases. Symptoms that follow a dermatome, such as pain or a rash, may indicate a pathology that involves the related nerve root. Examples include dysfunction of the spine or a viral infection.

Key Terms

- **shingles**: Also known as herpes zoster, shingles is an acute viral inflammation of the sensory ganglia of spinal and cranial nerves associated with a vesicular eruption and neuralgic pains and caused by reactivation of the poxvirus causing chicken pox.
- **chickenpox**: A common childhood disease caused by the varicella-zoster virus.

A dermatome is an area of skin that is supplied by a single spinal nerve. There are eight cervical nerves, twelve thoracic nerves, five lumbar nerves and five sacral nerves. Each of these nerves relays sensation, including pain, from a particular region of the skin to the brain.



Dermatomes: Dermatomes are areas of skin supplied by sensory neurons that arise from a spinal nerve ganglion. Dermatomes and the associated major cutaneous nerves are shown here in a ventral view.

Along the thorax and abdomen, the dermatomes are like a stack of discs, with each section supplied by a different spinal nerve. Along the arms and the legs, the pattern is different. The dermatomes run longitudinally along the limbs, so that each half of the limb has a different dermatome.

Although the general pattern is similar in all people, the precise areas of innervation are as unique to an individual as fingerprints.

Dermatomes have clinical significance, especially in the diagnosis of certain diseases. Symptoms that follow a dermatome, such as pain or a rash, may indicate a pathology that involves the related nerve root. Examples include dysfunction of the spine or a viral infection.

Viruses that remain dormant in nerve ganglia, such as the varicella zoster virus that causes both chickenpox and shingles, often cause either pain, rash, or both in a pattern defined by a dermatome.



Shingles rash: The shingles rash appears across a dermatome. In this patient, one of the dermatomes in the arm is affected, restricting the rash to the length of the back of the arm.

Shingles is one of the only diseases that causes a rash in a dermatomal pattern, and as such, this is its defining symptom. The rash of shingles is almost always restricted to a specific dermatome, such as on the chest, leg, or arm caused by the residual varicella zoster virus infection of the nerve that supplies that area of skin. Shingles typically appears years or decades after recovery from chickenpox.

Mytome

A myotome is the group of muscles that a single spinal nerve root innervates. The myotome is the motor equivalent of a dermatome.

The myotome distributions of the upper and lower extremities are listed below:

- C1/C2: Neck flexion / extension
- C3: Neck lateral flexion
- C4: Shoulder elevation
- C5: Shoulder abduction
- C6: Elbow flexion/wrist extension
- C7: Elbow extension/wrist flexion
- C8: Finger flexion
- T1: Finger abduction
- L2: Hip flexion
- L3: Knee extension
- L4: Ankle dorsi-flexion
- L5: Great toe extension
- S1: Ankle plantar flexion/ankle eversion/hip extension
- S2: Knee flexion

- S3–S4: Anal reflex

The testing of myotomes provides the clinician with information about the level in the spine where a lesion may be present. During testing, the clinician looks for muscle weakness of a particular group of muscles. Results may indicate lesions to the spinal cord nerve root, or intervertebral disc herniation that presses on the spinal nerve roots.

This page titled 12.6E: Dermatomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6: SPINAL NERVES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

12.6A: Overview of the Spinal Nerves

12.6B: Branches of Spinal Nerves

12.6C: Plexuses

12.6D: Intercostal Nerves

12.6E: Dermatomes

12.6F: Function and Physiology of the Spinal Nerves

This page titled 12.6: Spinal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.6F: Function and Physiology of the Spinal Nerves

Spinal nerves connect the brain and spinal cord to the limbs and organs of the body.

Learning Objectives

- Describe the function and physiology of the spinal nerves

Key Points

- Spinal nerves, considered part of the peripheral nervous system, generally refer to mixed spinal nerves, which carry motor, sensory, and autonomic information between the brain and spinal cord and the rest of the body.
- The cervical spinal nerves innervate the muscles and provide sensation for the head, neck, and diaphragm, as well as the upper limbs and back.
- The lumbar, sacral, and coccygeal nerves combine to form the lumbosacral plexus.
- The spinal cord can be divided into the lateral, posterior, and medial cord, each segment of which gives rise to specific nerves and serves specific areas of the body.
- The somatic nervous system is responsible for voluntary body movements, receiving information from afferent fibers and contracting muscles with efferent fibers.
- The autonomic nervous system involves the visceral organs and regulates involuntary movements or unconscious actions.
- The sympathetic nervous system is responsible for the fight or flight reaction under stressful conditions, while the parasympathetic nervous system conserves energy after high stress situations or during rest and digesting.
- The primary neurotransmitters of the peripheral nervous system (PNS) are acetylcholine and noradrenaline, though other neurotransmitters are also present. Acetylcholine acts on two sets of receptors, muscarinic and nicotinic cholinergic receptors.


Key Terms

- **nicotinic**: Excitatory acetylcholine receptors.
- **mixed spinal nerve**: A nerve that carries motor, sensory, and autonomic signals between the spinal cord and the body.
- **peripheral nervous system**: The nerves and ganglia outside of the brain and spinal cord.
- **muscarinic**: Acetylcholine receptors
  that can be both excitatory and inhibitory.

Review of Peripheral Nervous System Structure

The peripheral nervous system (PNS) consists of the nerves and ganglia outside of the brain and spinal cord. The main function of the PNS is to connect the central nervous system (CNS) to the limbs and organs.

Unlike the CNS, the PNS is not protected by the bones of the spine and skull, or by the blood –brain barrier, leaving it exposed to toxins and mechanical injuries. The peripheral nervous system is divided into the somatic nervous system and the autonomic nervous system.



Spinal nerve: This diagram indicates the formation of a typical spinal nerve from the dorsal and ventral roots. Numbers indicate the types of nerve fibers: 1 somatic efferent, 2 somatic afferent, 3–5 sympathetic efferent, 6–7 sympathetic afferent.

The peripheral nervous system includes 12 cranial nerves and 31 pairs of spinal nerves that provide communication from the CNS to the rest of the body by nerve impulses to regulate the functions of the human body. The term spinal nerve generally refers to a mixed spinal nerve, which carries motor, sensory, and autonomic signals between the spinal cord and the body.

Spinal Nerve Correspondences

Each pair of spinal nerves roughly correspond to a segment of the vertebral column: 8 cervical spinal nerve pairs (C1–C8), 12 thoracic pairs (T1–T12), 5 lumbar pairs (L1–L5), 5 sacral pairs (S1–S5), and 1 coccygeal pair.

- The first 4 cervical spinal nerves, C1 through C4, split and recombine to produce a variety of nerves that subserve the neck and back of the head.
- The spinal nerve C1 (suboccipital nerve) provides motor innervation to muscles at the base of the skull.
- C2 and C3 form many of the nerves of the neck, and provides both sensory and motor control. These include the greater occipital nerve that provides sensation to the back of the head, the lesser occipital nerve that provides sensation to the area behind the ears, the greater auricular nerve, and the lesser auricular nerve.
- The phrenic nerve arises from nerve roots C3, C4, and C5. It innervates the diaphragm to enable breathing. If the spinal cord is transected above C3, then spontaneous breathing is not possible.
- The last four cervical spinal nerves, C5 through C8, and the first thoracic spinal nerve, T1, combine to form the brachial plexus, or plexus brachialis, a tangled array of nerves, splitting, combining and recombining to form the nerves that subserve the upper limb region and upper back. Although the brachial plexus may appear tangled, it is highly organized and predictable with little variation among people.

Lumbosacral Plexus

The anterior divisions of the lumbar, sacral, and coccygeal nerves form the lumbosacral plexus, the first lumbar nerve being frequently joined by a branch from the twelfth thoracic. For descriptive purposes, this plexus is usually divided into three parts: lumbar plexus, sacral plexus, and pudendal plexus.

Autonomic Nervous System Function (ANS)

The sympathetic division typically functions in actions that need quick responses. The parasympathetic division functions with actions that do not require immediate reaction.

The sympathetic system is often considered the fight or flight system, while the parasympathetic system is often considered the rest and digest or feed and breed system.

Some typical actions of the sympathetic and parasympathetic systems are listed below.

Sympathetic Nervous System


- Diverts blood flow away from the gastrointestinal (GI) tract and skin via vasoconstriction.
- Enhances blood flow to skeletal muscles and the lungs.
- Dilates bronchioles of the lung by circulating epinephrine to allow for greater alveolar oxygen exchange.
- Increases the heart rate and contractility of cardiac muscle for enhanced blood flow to skeletal muscles.
- Dilates pupils and relaxes the ciliary muscle to the lens for far vision.
- Provides vasodilation for the coronary vessels of the heart.
- Constricts all the intestinal sphincters and the urinary sphincter.
- Inhibits peristalsis.
- Stimulates orgasm.

Parasympathetic Nervous System

- Dilates blood vessels that lead to the GI tract to increase blood flow; this is important following food consumption due to the greater metabolic demands placed on the body by the gut.
- Constricts the bronchiolar diameter when the need for oxygen has diminished.
- Manages heart control via dedicated cardiac branches of the vagus and thoracic spinal accessory nerves.
- Constricts the pupil and contracts the ciliary muscles to facilitate accommodation for closer vision.
- Stimulates salivary gland secretion and accelerates peristalsis to mediate the digestion of food.
- PNS nerves are involved in the erection of genital tissues via the pelvic splanchnic nerves 2–4. They are also responsible for stimulating sexual arousal.

Neurotransmitters

- Acetylcholine is the preganglionic neurotransmitter for both divisions of the ANS, as well as the postganglionic neurotransmitter of parasympathetic neurons.
- Nerves that release acetylcholine are said to be cholinergic. In the parasympathetic system, ganglionic neurons use acetylcholine as a neurotransmitter to stimulate muscarinic receptors.
- At the adrenal medulla, there is no postsynaptic neuron. Instead the presynaptic neuron releases acetylcholine to act on nicotinic receptors.
- Stimulation of the adrenal medulla releases adrenaline (epinephrine) into the bloodstream, which acts on adrenoceptors, producing a widespread increase in sympathetic activity.



Autonomic nervous sytem: The functions of the parasympathetic and sympathetic nervous systems are detailed.

Somatic Nervous System Function (SoNS)

The somatic nervous system consists of afferent and efferent nerves and is associated with the voluntary control of skeletal muscle movements. The afferent nerves are responsible for relaying sensations from the body to the central nervous system (CNS), while the efferent nerves are responsible for sending out commands from the CNS to the body to stimulate muscle contraction.

Upper motor neurons release acetylcholine. Acetylcholine is released from the axon terminal knobs of alpha motor neurons and received by postsynaptic receptors (nicotinic acetylcholine receptors) of muscles, thereby relaying the stimulus to contract muscle fibers.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Spinal nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_nerves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_nerves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dorsal root. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/dorsal%20root**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autonomic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/autonomic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intervertebral foramen. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intervertebral%20foramen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Fo...ertebralia.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsal ramus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsal%20ramus. **License**: *CC BY-SA: Attribution-ShareAlike*
- meningeal branches. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/meningeal%20branches. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Fo...ertebralia.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- coccygeal plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/coccygeal%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- brachial plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/brachial%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/nerve%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Fo...ertebralia.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_plexus. **License**: *CC BY: Attribution*
- Brachial plexus 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:B...l_plexus_2.svg. **License**: *Public Domain: No Known Copyright*
- Intercostal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intercostal_nerves. **License**: *CC BY-SA: Attribution-ShareAlike*
- Intercostal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intercostal_nerves. **License**: *CC BY-SA: Attribution-ShareAlike*
- sympathetic trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sympathetic%20trunk. **License**: *CC BY-SA: Attribution-ShareAlike*
- thoracic spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/thoraci...pinal%20nerves. **License**: *CC BY-SA: Attribution-ShareAlike*
- abdominal peritoneum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/abdominal%20peritoneum. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Fo...ertebralia.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_plexus. **License**: *CC BY: Attribution*
- Brachial plexus 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:B...l_plexus_2.svg. **License**: *Public Domain: No Known Copyright*
- Gray821. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray821.png. **License**: *Public Domain: No Known Copyright*
- Dermatome (anatomy). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dermatome_(anatomy). **License**: *CC BY-SA: Attribution-ShareAlike*

- Dermatomes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dermatomes. **License**: *CC BY-SA: Attribution-ShareAlike*
- chickenpox. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chickenpox. **License**: *CC BY-SA: Attribution-ShareAlike*
- shingles. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/shingles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Myotome. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Myotome. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Foraminaintervertebralia.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_plexus. **License**: *CC BY: Attribution*
- Brachial plexus 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brachial_plexus_2.svg. **License**: *Public Domain: No Known Copyright*
- Gray821. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray821.png. **License**: *Public Domain: No Known Copyright*
- Dermatomes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dermatomes. **License**: *Public Domain: No Known Copyright*
- Shingles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Shingles.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...Nervous_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- Peripheral nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Peripheral_nervous_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- peripheral nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/periphe...rvous%20system. **License**: *CC BY-SA: Attribution-ShareAlike*
- mixed spinal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/mixed%20spinal%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Autonomic Nervous System. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Autonomic_nervous_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Foraminaintervertebralia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Foraminaintervertebralia.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_plexus. **License**: *CC BY: Attribution*
- Brachial plexus 2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brachial_plexus_2.svg. **License**: *Public Domain: No Known Copyright*
- Gray821. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Gray821.png. **License**: *Public Domain: No Known Copyright*
- Dermatomes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dermatomes. **License**: *Public Domain: No Known Copyright*
- Shingles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Shingles.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- The Autonomic Nervous System. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:The_Autonomic_Nervous_System.jpg. **License**: *Public Domain: No Known Copyright*
- Spinal nerves. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_nerves. **License**: *Public Domain: No Known Copyright*

This page titled 12.6F: Function and Physiology of the Spinal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7: DISTRIBUTION OF SPINAL NERVES

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

12.7A: Cervical Plexus
12.7B: Brachial Plexus
12.7C: Lumbar Plexus
12.7D: Sacral and Coccygeal Plexuses
12.7E: Sensory and Motor Tracts

This page titled 12.7: Distribution of Spinal Nerves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7A: Cervical Plexus

The cervical plexus is the plexus of the ventral rami of the first four cervical spinal nerves.

Learning Objectives

- Describe the cervical plexus and its function

Key Points

- The cervical plexus describes the plexus of the ventral rami of the first four cervical spinal nerves that arise from the cervical spinal column in the neck.
- The cervical spinal nerves that form the cervical plexus are located laterally (farther from the median line) to the transverse processes of the prevertebral skeletal muscles of the neck from the medial side, and vertebral (closer to the vertebral column) to these muscles from the lateral side.
- The cervical plexus forms an anastomosis, a connection, with the accessory nerve, the hypoglossal nerve, and the sympathetic trunk.
- The cervical plexus is located in the neck, internal to the sternocleidomastoid, an anterior neck muscle.

Key Terms

- **sympathetic trunk**: Also called the sympathetic chain or gangliated cord, these are a paired bundle of nerve fibers that run from the base of the skull to the coccyx.
- **plexus**: A network or interwoven mass, especially of nerves, blood vessels, or lymphatic vessels.
- **platysma**: A superficial muscle that overlaps the sternocleidomastoid.
- **cervical plexus**: A plexus of the ventral rami of the first four cervical spinal nerves that are located from the C1 to C4 cervical segment in the neck. They are located laterally to the transverse processes of the prevertebral muscles from the medial side and vertebral (scalenus, levator scapulae, splenius cervicis muscles) from the lateral side.

Structure and Distribution

The cervical plexus is a plexus of the ventral rami of the first four cervical spinal nerves located from the C1 to C4 cervical segment in the neck. They are located laterally to the transverse processes between prevertebral muscles from the medial side and anteriolateral to the m. scalenus and m. levator scapulae.

There is anastomosis with the accessory nerve, hypoglossal nerve, and sympathetic trunk. It is located deep in the neck, near the sternocleidomastoid muscle.

Nerves formed from the cervical plexus innervate the back of the head, as well as some neck muscles. The branches of the cervical plexus emerge from the posterior triangle at the nerve point, a point that lies midway on the posterior border of the sternocleidomastoid.

Branches and Their Functions

The cervical plexus has two types of branches: cutaneous and muscular.

Cutaneous branches include:

- The lesser occipital nerve, or small occipital nerve, is a cutaneous spinal nerve that arises between the second and third cervical vertebrae, along with the greater occipital nerve. It innervates the scalp in the lateral area of the head posterior to the ear.


- The great auricular nerve originates from the cervical plexus and is composed of branches from spinal nerves C2 and C3. It provides sensory innervation for the skin over the parotid gland and mastoid process, and both surfaces of the outer ear.
- The transverse cervical nerve (superficial cervical or cutaneous cervical) arises from the second and third cervical nerves, turns around the posterior border of the sternocleidomastoideus about its middle, then passes obliquely forward beneath the external jugular vein to the anterior border of the muscle, where it perforates the deep cervical fascia and divides beneath the platysma into ascending and descending branches that are distributed to the antero-lateral parts of the neck.
- The supraclavicular nerves (descending branches) arise from the third and fourth cervical nerves. They emerge beneath the posterior border of the sternocleidomastoideus, and descend in the posterior triangle of the neck beneath the platysma and deep cervical fascia.

Muscular branches include:

- Ansa cervicalis (loop formed from C1–C3), geniohyoid (C1 only), thyrohyoid (C1 only), sternothyroid, sternohyoid, omohyoid: The ansa cervicalis is a loop of nerves that are part of the cervical plexus.
- The phrenic nerve (C3–C5, but primarily C4) is a nerve that originates in the neck and passes down between the lung and heart to reach the diaphragm.
- Segmental branches (C1–C4) innervate the anterior and middle scalenes.

There are two additional branches that are formed by the posterior roots of the spinal nerves:

- Preauricular nerve (from the posterior roots of C2–C3).
- Postauricular nerve (from the posterior roots of C3–C4).



Cervical plexus: Diagram of the cervical plexus.

This page titled 12.7A: Cervical Plexus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7B: Brachial Plexus

The brachial plexus is formed by the four lower cervical spinal nerves and the first thoracic spinal nerve.

Learning Objectives

- Describe the brachial plexus


Key Points

- The nerve bundles of the brachial plexus pass through the cervico-axillary canal to serve the axilla (armpit), brachium (arm), antebrachium (forearm), and hand.
- The brachial plexus provides cutaneous (skin) and muscular innervations for the upper limbs, excluding the trapezius muscle and an area of skin near the axilla that are innervated by the spinal accessory nerve and the intercostobrachial nerve, respectively.
- The brachial plexus is divided into roots, trunks, divisions, cords, and branches.

Key Terms

- **spinal accessory nerve**: A nerve that controls specific muscles of the shoulder and neck.
- **axilla**: The armpit, or the cavity beneath the junction of the arm and shoulder.
- **brachium**: The upper arm.

Formation and Distribution



Right brachial plexus: The right brachial plexus with its short branches, viewed from the front.

The brachial plexus is a network of nerve fibers that run from the spine that are formed by the ventral rami of the lower four cervical and first thoracic nerve roots (C5–C8, T1).


The brachial plexus proceeds through the neck, the axilla (armpit region), and into the arm. It is a collection of nerves passing through the cervico-axillary canal to reach the axilla and innervate the brachium, the antebrachium, and the hand.

The brachial plexus is responsible for cutaneous and muscular innervation of the entire upper limb, with two exceptions: the trapezius muscle is innervated by the spinal accessory nerve (CN XI) and an area of skin near the axilla is innervated by the intercostobrachial nerve. Lesions on the brachial plexus can lead to severe functional impairment.

Branches and Divisions

The brachial plexus is divided into roots, trunks, divisions, cords, and branches. There are five terminal branches and numerous other preterminal or collateral branches that leave the plexus at various points along its length. Its structure includes:

- Five roots: The five anterior rami of the spinal nerves, after they have given off their segmental supply to the muscles of the neck.
- These roots merge to form three trunks: The superior or upper (C5–C6), the middle (C7), and the inferior or lower (C8, T1).
- Each trunk then splits in two, to form six divisions: The anterior divisions of the upper, middle, and lower trunks and the posterior divisions of the upper, middle, and lower trunks.
- These six divisions will regroup to become the three cords. The cords are named by their position with respect to the axillary artery:
    1. The posterior cord is formed from the three posterior divisions of the trunks (C5–C8,T1).
    2. The lateral cord is the anterior divisions from the upper and middle trunks (C5–C7).
    3. The medial cord is simply a continuation of the anterior division of the lower trunk (C8, T1).
    4. Most branches branch from the cords, but a few branch directly from earlier structures. The five on the left are considered terminal branches.



Brachial plexus: Diagram of the brachial plexus and its branches.

This page titled 12.7B: Brachial Plexus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7C: Lumbar Plexus

The lumbar plexus is formed by the subcostal nerve and divisions of the first four lumbar nerves that arise from the middle to lower back.

Learning Objectives

- Describe the lumbar plexus

Key Points

- The lumbar plexus passes through the psoas major muscle and innervates the skin and muscles of the abdominal wall, thigh, and external genitalia.
- The largest nerve that forms part of the lumbar plexus is the femoral nerve, which innervates the anterior thigh muscles and some of the skin distal to the inguinal ligament.
- The ilioinguinal nerve pierces the lateral abdominal wall and runs medially at the level of the inguinal ligament. It supplies motor branches to both transversus abdominis and sensory branches (to the skin over the pubic symphysis and the lateral aspect of the labia majora or scrotum).
- The genital branch of the genitofemoral nerve courses differently in females and males.


Key Terms

- **lumbosacral plexus**: The anterior divisions of the lumbar nerves, sacral nerves, and coccygeal nerve form this plexus, the first lumbar nerve being frequently joined by a branch from the twelfth thoracic nerve.
- **inguinal ligament**: A band running from the pubic tubercle to the anterior superior iliac spine. Its anatomy is very important for operating on hernia patients.
- **subcostal nerve**: The twelfth thoracic nerve, its anterior division is larger than the others; it runs along the lower border of the twelfth rib and often gives a communicating branch to the first lumbar nerve, and then passes under the lateral lumbocostal arch.
- **aponeurosis**: Layers of flat, broad tendons that have a shiny, whitish-silvery color.

Structure and Distribution

The lumbar plexus is a nerve plexus in the lumbar region of the body that forms part of the lumbosacral plexus. It is formed by the ventral divisions of the first four lumbar nerves (L1–L4) and from contributions of the subcostal nerve (T12), which is the last thoracic nerve.


This plexus lies within the psoas major muscle. Nerves of the lumbar plexus serve the skin and the muscles of the lower abdominal wall, the thigh, and external genitals. The largest nerve of the plexus is the femoral nerve and it supplies the anterior muscles of the thigh and a part of skin distal to the inguinal ligament.



Lumbar plexus: An image of the lumbar plexus with its nerves highlighted in yellow.

Branches of the Lumbar Plexus

Iliohypogastric Nerve



Lumbar plexus: Schematic of the lumbar plexus.

Runs anterior to the psoas major on its proximal lateral border to run laterally and obliquely on the anterior side of the quadratus lumborum. Lateral to this muscle, it pierces the transversus abdominis to run above the iliac crest between that muscle and the abdominal internal oblique. It gives off several motor branches to these muscles and a sensory branch to the skin of the lateral hip. Its terminal branch then runs parallel to the inguinal ligament to exit the aponeurosis of the abdominal external oblique above the external inguinal ring, where it supplies the skin above the inguinal ligament (i.e., the hypogastric region) with the anterior cutaneous branch.

Ilioinguinal Nerve

This nerve closely follows the iliohypogastric nerve on the quadratus lumborum but then passes below it to run at the level of the iliac crest. It pierces the lateral abdominal wall and runs medially at the level of the inguinal ligament where it supplies motor branches to both the transversus abdominis and sensory branches through the external inguinal ring to the skin over the pubic symphysis and the lateral aspect of the labia majora in females, and in males, the scrotum.

Genitofemoral Nerve

Pierces the psoas major anteriorly, below the former two nerves to immediately split into two branches that run downward on the anterior side of the muscle. The lateral femoral branch is purely sensory and pierces the vascular lacuna near the saphenous hiatus and supplies the skin below the inguinal ligament. In males, the genital branch runs in the spermatic cord then sends sensory branches to the scrotal skin and supplies motor innervations to the cremaster muscle. In females, it runs in the inguinal canal together with the teres uteri ligament. It then sends sensory branches to the labia majora in females.

Lateral Cutaneous Femoral Nerve

Pierces the psoas major on its lateral side and runs obliquely downward below the iliac fascia. Medial to the anterior superior iliac spine, it leaves the pelvic area through the lateral muscular lacuna and enters the thigh by passing behind the lateral end of the inguinal ligament. In the thigh, it briefly passes under the fascia lata before it breaches the fascia and supplies the skin of the anterior thigh.

Obturator Nerve

Leaves the lumbar plexus and descends behind the psoas major on its medial side, follows the linea terminalis into the lesser pelvis, then finally leaves the pelvic area through the obturator canal. In the thigh, it sends motor branches to obturator externus before dividing into an anterior and a posterior branch, both of which continue distally. These branches are separated by the adductor brevis and supply all thigh adductors with motor innervations. The anterior branch contributes a terminal, sensory branch that passes along the anterior border of gracilis and supplies the skin on the medial, distal part of the thigh.

Femoral Nerve

This is the largest and longest of the plexus' nerves. It gives motor innervation to iliopsoas, pectineus, sartorius, and quadriceps femoris, and sensory innervation to the anterior thigh, posterior lower leg, and hindfoot. In the pelvic area, it runs in a groove between the psoas major and iliacus muscles and gives branches to both. It exits the pelvis through the medial aspect of the muscular lacuna. In the thigh, it divides into numerous sensory and muscular branches and the saphenous nerve, its long sensory terminal branch that continues down to the foot.

This page titled 12.7C: Lumbar Plexus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7D: Sacral and Coccygeal Plexuses

The sacral plexus is the plexus of the three sacral spinal nerves (S2–S4) that arise from the lower back just above the sacrum.

Learning Objectives

- Distinguish between the sacral and coccygeal plexuses


Key Points

- The sacral plexus, together with the lumbar plexus, forms the lumbosacral plexus.
- The largest nerve in the sacral plexus is the sciatic nerve that innervates the thigh, lower leg, and the foot.
- The coccygeal plexus consists of the coccygeal nerve and the fifth sacral nerve, which innervate the skin in the coccygeal region, around the tailbone (called the coccyx).

Key Terms

- **sacral plexus**: A nerve plexus that provides motor and sensory nerves for the posterior thigh, most of the lower leg, the entire foot, and part of the pelvis.
- **coccygeal nerve**: The spinal nerve that corresponds to the coccyx bone.
- **sciatic nerve**: A large nerve that starts in the lower back and runs through the buttock and down the lower limb.

The sacral plexus is a nerve plexus that provides motor and sensory nerves for the posterior thigh, most of the lower leg, the entire foot, and part of the pelvis. It is part of the lumbosacral plexus and emerges from the sacral vertebrae (S2–S4).



The sacral plexus: The nerves of the sacral plexus are shown.

The largest and longest nerve of the human body, the sciatic nerve, is the main branch and gives rami to the motor innervation of the muscles of the foot, the leg, and the thigh.

The sacral plexus is formed by:

- The lumbosacral trunk.


- The anterior division of the first sacral nerve.
- Portions of the anterior divisions of the second and third sacral nerves.

The nerves forming the sacral plexus converge toward the lower part of the greater sciatic foramen and unite to form a flattened band from the anterior and posterior surfaces, from which several branches arise.

The band itself is continued as the sciatic nerve, which splits on the back of the thigh into the tibial nerve and the common fibular nerve. These two nerves sometimes arise separately from the plexus and, in all cases, their independence can be shown by dissection.

Often, the sacral plexus and the lumbar plexus are considered to be one large nerve plexus, the lumbosacral plexus. The lumbosacral trunk connects the two plexuses.

The coccygeal plexus originates from the S4, S5, and Co1 spinal nerves. It is interconnected with the lower part of the sacral plexus. The only nerve in this plexus is the anococcygeal nerve, which serves sensory innervation of the skin in the coccygeal region.



Sacral plexus diagram: Diagram of the sacral plexus showing the various anterior and posterior nerve branches that arise from each lumbar and sacral nerve.

This page titled 12.7D: Sacral and Coccygeal Plexuses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.7E: Sensory and Motor Tracts

The spinothalamic tract is a somatosensory tract and the corticospinal tract is a motor tract.

Learning Objectives

- Distinguish between sensory and motor tracts

Key Points

- The spinothalamic tract is split into the lateral spinothalamic tract that transmits pain and temperature sensation to the thalamus, and the anterior spinothalamic tract that transmits pressure and crude touch sensation to the thalamus.
- The corticospinal tract is a motor tract comprised of mostly motor axons that carry motor information from the cerebral cortex and brainstem to the musculature for voluntary movement.
- The corticospinal tract is split into the lateral and anterior corticospinal tracts, which decussate, or cross, in the medulla oblongata. Therefore, the right brain controls the left side of the body, and the left brain controls the right side of the body.
- Betz cells, the largest pyramidal cells, are only found in the corticospinal tract.

Key Terms

- **spinothalamic tract**: A sensory pathway originating in the spinal cord. It transmits information to the thalamus about pain, temperature, itch, and crude touch.
- **somatosensory tract**: The system that reacts to diverse stimuli using thermoreceptors, nociceptors, mechanoreceptors, and chemoreceptors. The transmission of information from the receptors passes via sensory nerves through tracts in the spinal cord and into the brain.
- **Betz cell**: Giant pyramidal cells (neurons) located within the fifth layer of the grey matter in the primary motor cortex. They have a rapid conduction rate of over 70m/sec, which is the fastest conduction of any signals from the brain to the spinal cord.
- **corticospinal tract**: The nervous system tract that conducts impulses from the brain to the spinal cord. It contains mostly motor axons and is made up of two separate tracts in the spinal cord: the lateral corticospinal tract and the anterior corticospinal tract.

The Somatosensory Tract

The spinothalamic tract is a sensory pathway originating in the spinal cord. It transmits information to the thalamus about pain, temperature, itch, and crude touch. The pathway decussates at the level of the spinal cord.

Somatosensory organization is divided into the dorsal column–medial lemniscus tract (the touch/proprioception/vibration sensory pathway) and the anterolateral system, or ALS (the pain/temperature sensory pathway). Both sensory pathways use three different neurons to get information from sensory receptors at the periphery to the cerebral cortex.

These neurons are designated primary, secondary, and tertiary sensory neurons. In both pathways, primary sensory neuron cell bodies are found in the dorsal root ganglia, and their central axons project into the spinal cord.

Function

The types of sensory information transmitted via the spinothalamic tract are described as affective sensation. This means that the sensation is accompanied by a compulsion to act. For instance, an itch is accompanied by a need to scratch, and a painful stimulus makes us want to withdraw from the pain.

There are two subsystems:

1. Direct (for direct, conscious appreciation of pain).
2. Indirect (for affective and arousal impact of pain).

Indirect projections are further divided into:

- Spino-reticulo-thalamo-cortical (part of the ascending reticular arousal system, also known as ARAS).
- Spino-mesencephalic-limbic (for affective impact of pain).


The Corticospinal Tract

The corticospinal tract conducts impulses from the brain to the spinal cord. It contains mostly motor axons. The corticospinal tract is made up of two separate tracts in the spinal cord: the lateral corticospinal tract and the anterior corticospinal tract.

The corticospinal tract also contains the Betz cell (the largest pyramidal cells) that are not found in any other region of the body. An understanding of these tracts leads to an understanding of why one side of the body is controlled by the opposite side of the brain.

The corticospinal tract is concerned specifically with discrete, voluntary, skilled movements, such as the precise movement of fingers and toes. The brain sends impulses to the spinal cord that relay the message.

This is imperative in understanding that the left hemisphere of the brain controls the RIGHT side of the body, while the right hemisphere of the brain controls the LEFT side of the body. The signals cross in the medulla oblongata, and this process is also known as decussation.



Spinal cord tracts: Spinal cord tracts are identified.

Function

The primary purpose of the corticospinal tract is to maintain voluntary motor control of the body and limbs. However, connections to the somatosensory cortex suggest that the pyramidal tracts are also responsible for modulating sensory information from the body.

Some of these connections cross the midline; therefore, each side of the brain is responsible for controlling muscles for the limbs on opposite sides of the body. However, control of trunk muscles is on the same side of the body.

After a patient's pyramidal tracts are injured, the patient is paralyzed on the corresponding side of the body. Fortunately, they can re-learn some crude, basic motions, but not fine movements. This implies that the connections to these tracts are crucial for fine

movement, and only partial recovery is possible if they are damaged.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cervical plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cervical_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cervical_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cervical plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cervical%20plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plexus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sympathetic trunk. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/sympathetic%20trunk**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical plexus.PNG. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:C...cal_plexus.PNG**. **License**: ***Public Domain: No Known Copyright***
- spinal accessory nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/spinal%...essory%20nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brachial plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brachial_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- axilla. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/axilla**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- brachium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/brachium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical plexus.PNG. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:C...cal_plexus.PNG**. **License**: ***Public Domain: No Known Copyright***
- Gray808.png. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Gray808.png**. **License**: ***Public Domain: No Known Copyright***
- Brachial Plexus Color. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...exus_color.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nervous plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lumbar plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lumbar_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- inguinal ligament. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/inguinal%20ligament**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lumbosacral plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lumbosacral%20plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- subcostal nerve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/subcostal%20nerve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Aponeurosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Aponeurosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical plexus.PNG. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:C...cal_plexus.PNG**. **License**: ***Public Domain: No Known Copyright***
- Gray808.png. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Gray808.png**. **License**: ***Public Domain: No Known Copyright***
- Brachial Plexus Color. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...exus_color.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray823. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray823.png**. **License**: ***Public Domain: No Known Copyright***
- lumbar schem.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/F...bar_plexus.svg**. **License**: ***CC BY: Attribution***
- Nerve plexus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nerve_plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Sacral plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sacral_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- coccygeal nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/coccygeal%20nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- sacral plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sacral%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- sciatic nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sciatic_nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cervical plexus.PNG. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Cervical_plexus.PNG. **License**: *Public Domain: No Known Copyright*
- Gray808.png. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray808.png. **License**: *Public Domain: No Known Copyright*
- Brachial Plexus Color. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...exus_color.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray823. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray823.png. **License**: *Public Domain: No Known Copyright*
- lumbar schem.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F...bar_plexus.svg. **License**: *CC BY: Attribution*
- Gray828.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray828.png. **License**: *Public Domain: No Known Copyright*
- sacral plexus diagram.gif. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...ral_plexus.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinothalamic tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinothalamic_tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Corticospinal tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Corticospinal_tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal tracts. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spinal_tracts. **License**: *CC BY-SA: Attribution-ShareAlike*
- somatosensory tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/somatosensory%20tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- corticospinal tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/corticospinal%20tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- spinothalamic tract. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/spinothalamic%20tract. **License**: *CC BY-SA: Attribution-ShareAlike*
- Betz Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Betz_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cervical plexus.PNG. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Cervical_plexus.PNG. **License**: *Public Domain: No Known Copyright*
- Gray808.png. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray808.png. **License**: *Public Domain: No Known Copyright*
- Brachial Plexus Color. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Brachial_plexus_color.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray823. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray823.png. **License**: *Public Domain: No Known Copyright*
- lumbar schem.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Lumbar_plexus.svg. **License**: *CC BY: Attribution*
- Gray828.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray828.png. **License**: *Public Domain: No Known Copyright*
- sacral plexus diagram.gif. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Sacral_plexus.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal cord tracts. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikiped..._-_English.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 12.7E: Sensory and Motor Tracts is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.8: MOTOR ACTIVITY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

12.8A: Peripheral Motor Endings

12.8B: Overview of Motor Integration

This page titled 12.8: Motor Activity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.8A: Peripheral Motor Endings

A neuromuscular junction exists between the axon terminal and the motor end plate of a muscle fiber where neurotransmitters are released.

Learning Objectives

- Describe the function of peripheral motor endings

Key Points

- A neuromuscular junction is the junction between the axon terminal of a motor neuron and the plasma membrane of the motor end plate of a muscle fiber.
- With the arrival of an action potential to the axon terminal, voltage-dependent calcium channels open, and calcium infuses into the cell. The influx of calcium ions causes the docking of acetylcholine -containing vesicles at the plasma membrane of the neuron and exocytosis into the synaptic cleft.
- Acetylcholine is a neurotransmitter contained in the vesicles of the pre-synaptic neuron. It is released into the synaptic cleft and activates nicotinic acetylcholine receptors on the motor end plate, and causes local motor end plate depolarization, also known as the end plate potential (EPP).
- The end plate potential propagates across the surface of the muscle fiber, causing the fiber to contract and continuing the process of excitation–contraction coupling.

Key Terms

- **axon**: A nerve fiber that is a long, slender projection of a nerve cell that conducts nerve impulses away from the body of the cell to a synapse.
- **voltage-dependent calcium channels**: A group of voltage-gated ion channels found in excitable cells (e.g., muscle, glial cells, neurons, etc.) with a permeability to the ion Ca2+.
- **presynaptic neuron**: The neuron that releases neurotransmitters into the synaptic cleft.
- **nicotinic acetylcholine receptor**: These are cholinergic receptors that form ligand-gated ion channels in the plasma membranes of certain neurons and on the postsynaptic side of the neuromuscular junction.
- **synaptic cleft**: A small space between neurons.
- **excitation-contraction coupling**: This process is fundamental to muscle physiology, whereby the electrical stimulus is usually an action potential and the mechanical response is a contraction.

A neuromuscular junction is the synapse or junction of the axon terminal of a motor neuron with the motor end plate, as shown in Figures 1 and 2. The highly excitable region of muscle fiber plasma membrane is responsible for initiation of action potentials across the muscle's surface, ultimately causing the muscle to contract.

In vertebrates, the signal passes through the neuromuscular junction via the neurotransmitter acetylcholine.




Figure 1. Detailed view of a neuromuscular junction: Detailed view of a neuromuscular junction: 1) Presynaptic terminal; 2) Sarcolemma; 3) Synaptic vesicle; 4) Nicotinic acetylcholine receptor; 5) Mitochondrion.




Figure 2. Neuromuscular junction: Electron micrograph showing a cross section through the neuromuscular junction. T is the axon terminal and M is the muscle fiber. The arrow shows junctional folds with basal lamina. Postsynaptic densities are visible on the tips between the folds. Scale is 0.3 µm.

Upon the arrival of an action potential at the presynaptic neuron terminal, voltage-dependent calcium channels open and Ca^{2+} ions flow from the extracellular fluid into the presynaptic neuron's cytosol. This influx of Ca^{2+} causes neurotransmitter-containing vesicles to dock and fuse to the presynaptic neuron's cell membrane, which results in the emptying of the vesicle's contents (acetylcholine) into the synaptic cleft; this process is known as exocytosis.

Acetylcholine diffuses into the synaptic cleft and binds to the nicotinic acetylcholine receptors located on the motor end plate.

These receptors open to allow sodium ions to flow in and potassium ions to flow out of the muscle's cytosol, producing a local depolarization of the motor end plate, known as an end-plate potential (EPP). This depolarization spreads across the surface of the muscle fiber and continues the excitation–contraction coupling to contract the muscle.

The action potential spreads through the muscle fiber's network of T-tubules, depolarizing the inner portion of the muscle fiber. The depolarization activates L-type, voltage-dependent calcium channels (dihydropyridine receptors) in the T-tubule membrane, which are in close proximity to calcium-release channels (ryanodine receptors) in the adjacent sarcoplasmic reticulum.

As intracellular calcium levels rise, the motor proteins responsible for the contractile response are able to interact, as shown in Figure 3, to form cross-bridges and undergo shortening.

CLINICAL EXAMPLE

Myasthenia gravis is an autoimmune disorder in which circulating antibodies block the nicotinic acetylcholine receptors on the motor end plate of the neuromuscular junction. This blockage of acetylcholine receptors causes muscle weakness, often first

exhibiting drooping eyelids and expanding to include overall muscle weakness and fatigue.

The affects of myasthenia gravis illustrate the importance of effective and functioning neuromuscular junctions for communication between neurons and muscles to allow contraction and relaxation of muscle fibers.




Figure 3. Muscle contraction and actin-myosin interactions: Skeletal muscle contracts following activation by an action potential. The binding of acetylcholine at the motor end plate leads to intracellular calcium release and interactions between myofibrils to elicit contraction.

This page titled 12.8A: Peripheral Motor Endings is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.8B: Overview of Motor Integration

A motor unit is comprised of a single alpha-motor neuron and all the muscle fibers it innervates.

Learning Objectives


- Describe motor integration and how it affects motor activity

Key Points

- Motor units contain muscle fibers of all the same type; these may be many muscle fibers (as in the case of quadriceps) or a few muscle fibers (as in the case of the muscles that control eye movement).
- Groups of motor units often work together to coordinate the contractions of a single muscle; all of the motor units that subserve a single muscle are considered a motor unit pool.
- Motor units are generally recruited in order of smallest to largest (from fewest fibers to most fibers) as contraction increases. This is known as Henneman's Size Principle.
- The smaller the motor unit, the more precise the action of the muscle.

Key Terms

- **Henneman's size principle**: According to this principle, motor unit recruitment is always in the same order from smallest to largest motor unit. Additionally, the motor unit action potential is an all-or-none phenomenon—once the recruitment threshold (the stimulus intensity at which a motor unit begins to fire) is reached, it fires fully.
- **alpha motor neuron**: Alpha motor neurons (α-MNs) are large, lower motor neurons of the brainstem and spinal cord. They innervate the extrafusal muscle fibers of skeletal muscle and are directly responsible for initiating their contraction. Alpha motor neurons are distinct from gamma motor neurons, which innervate the intrafusal muscle fibers of muscle spindles.
- **motor unit**: A neuron with its associated muscle fibers.



Rectus femoris: The rectus femoris muscle is one of the four quadriceps muscles of the human body. These muscles may have as many as a thousand fibers in each motor unit.

A motor unit consists of a single alpha motor neuron and all of the corresponding muscle fibers it innervates; all of these fibers will be of the same type (either fast twitch or slow twitch).

When a motor unit is activated, all of its fibers contract. Groups of motor units often work together to coordinate the contractions of a single muscle. All of the motor units that subserve a single muscle are considered a motor unit pool.


The number of muscle fibers within each unit can vary. Thigh muscles, for example, can have a thousand fibers in each unit, eye muscles might have ten. In general, the number of muscle fibers innervated by a motor unit is a function of a muscle's need for refined motion.

The smaller the motor unit, the more precise the action of the muscle. Muscles requiring more refined motion are innervated by motor units that synapse with fewer muscle fibers.

Motor unit recruitment is the progressive activation of a muscle by the successive recruitment of motor units to accomplish increasing gradations of contractile strength. The activation of more motor neurons will result in more muscle fibers being activated, and therefore a stronger muscle contraction.

Motor unit recruitment is a measure of how many motor neurons are activated in a particular muscle. It is therefore a measure of how many muscle fibers of that muscle are activated. The higher the recruitment, the stronger the muscle contraction will be.

Motor units are generally recruited in order of smallest to largest (from fewest fibers to most fibers) as contraction increases. This is known as Henneman's Size Principle.



The orbicularis oris (eye) muscle: These small motor units may contain only 10 fibers per motor unit. The more precise the action of the muscle, the fewer fibers innervated.

Motor Unit Categories

Motor units are generally categorized based upon the similarities between several factors such as:

Physiological

Contraction speed in isometric contractions:

- Rate of rise of force.
- Time to peak of a twitch contraction (response to a single nerve impulse).

Biochemical

Histochemical (the oldest form of biochemical fiber typing):

- Glycolytic enzyme activity.
- Oxidative enzyme activity.
- Sensitivity of myosin ATPase to acid and alkali.

Immunohistochemical (a more recent form of fiber typing):

- Myosin heavy chain (MHC).
- Myosin light chain—alkali (MLC1).
- Myosin light chain—regulatory (MLC2).

FUNCTIONAL EXAMPLE

The quadriceps muscles contain many thousands of muscle fibers in general, both slow and fast twitch, to produce sufficient force for body movements such as standing, walking, running, and jumping.

The eye muscles, on the other hand, contain few muscle fibers, enabling them to be more exact in movement so that vision is not jumpy, but consequently they produce very little force.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Muscle contraction. **Provided by**: Saylor. **Located at**: http://www.saylor.org/site/wp-conten...ontraction.pdf. **License**: ***CC BY-SA: Attribution-ShareAlike***
- presynaptic neuron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/presynaptic+neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- voltage-dependent calcium channels. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/voltage...ium%20channels. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nicotinic acetylcholine receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/nicotin...ine%20receptor. **License**: ***CC BY-SA: Attribution-ShareAlike***
- excitation-contraction coupling. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/excitat...ion%20coupling. **License**: ***CC BY-SA: Attribution-ShareAlike***
- axon. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/axon. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...synaptic-cleft. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neuromuscular Junction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neuromuscular_junction. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muskel-molekulartranslation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mu...ranslation.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor end plate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_end_plate. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor end plate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_end_plate. **License**: ***Public Domain: No Known Copyright***
- Motor unit recruitment. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_...of_recruitment. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor unit. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Motor_unit. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alpha motor neuron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/alpha%20motor%20neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Henneman's size principle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hennema...ze%20principle. **License**: ***CC BY-SA: Attribution-ShareAlike***
- motor unit. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/motor_unit. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Muskel-molekulartranslation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mu...ranslation.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor end plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_end_plate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor end plate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_end_plate**. **License**: ***Public Domain: No Known Copyright***
- Gray379. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray379.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rectus femoris. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Rectus_femoris.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 12.8B: Overview of Motor Integration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.9: MOTOR PATHWAYS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

12.9A: Organization of Motor Neuron Pathways

12.9B: The Role of the Basal Ganglia in Movement

12.9C: Modulation of Movement by the Cerebellum

12.9D: Functions of the Cerebellum in Integrating Movements

This page titled 12.9: Motor Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.9A: Organization of Motor Neuron Pathways

The motor system is the part of the central nervous system that is involved with movement.

Learning Objectives


- Describe the organization of motor neuron pathways

Key Points

- The pyramidal tract, which includes both the corticospinal and corticobulbar tracts, serves as the motor pathway for upper motor neuronal signals coming from the cerebral cortex and from primitive brainstem motor nuclei.
- Peripheral motor nerves carry the motor impulses from the spinal cord to the voluntary muscles.
- The large majority (90%) of motor neurons cross (decussate) to the contralateral side of the brain at the level of the brainstem.

Key Terms

- **extrapyramidal system**: A biological neural network that is part of the motor system that causes involuntary movements.
- **corticospinal tract**: The nervous system tract that conducts impulses from the brain to the spinal cord. It contains mostly motor axons and is made up of two separate tracts in the spinal cord: the lateral corticospinal tract and the anterior corticospinal tract.
- **motor system**: The part of the central nervous system that is involved with movement. It consists of the pyramidal and extrapyramidal systems.
- **cerebral cortex**: The gray, folded, outermost layer of the cerebrum that is responsible for higher brain processes such as sensation, voluntary muscle movement, thought, reasoning, and memory.



Decussation of the pyramids: A deep dissection, lateral view of a brainstem. The pyramidal tract is visible in red, and pyramidal decussation is labeled at lower right.


The motor system is the part of the central nervous system that is involved with movement. It consists of the pyramidal and extrapyramidal system.

The motor pathway, also called the pyramidal tract or the corticospinal tract, serves as the motor pathway for upper motor neuronal signals coming from the cerebral cortex and from primitive brainstem motor nuclei. There are upper and lower motor neurons in the corticospinal tract.

The motor impulses originate in the giant pyramidal cells (Betz cells) of the motor area, i.e., the precentral gyrus of the cerebral cortex. These are the upper motor neurons of the corticospinal tract. The axons of these cells pass from the cerebral cortex to the midbrain and the medulla oblongata. Peripheral motor nerves carry the motor impulses from the anterior horn to the voluntary muscles.

Cortical upper motor neurons originate from Brodmann areas 1, 2, 3, 4, and 6, then descend into the posterior limb of the internal capsule, through the crus cerebri, down through the pons, and to the medullary pyramids, where about 90% of the axons cross to the contralateral side at the decussation of the pyramids. They then descend as the lateral corticospinal tract.

These axons synapse with lower motor neurons in the ventral horns of all levels of the spinal cord. The remaining 10% of axons descend on the ipsilateral side as the ventral corticospinal tract. These axons also synapse with lower motor neurons in the ventral horns. Most of them will cross to the contralateral side of the cord (via the anterior white commissure) just before synapsing.




Brodmann areas of the brain: This drawing shows the regions of the human cerebral cortex as delineated by Korvinian Brodmann on the basis of cytoarchitecture.

The midbrain nuclei include four motor tracts that send upper motor neuronal axons down the spinal cord to lower motor neurons. These are the rubrospinal tract, the vestibulospinal tract, the tectospinal tract, and the reticulospinal tract.

The function of lower motor neurons can be divided into two different groups: the lateral corticospinal tract and the anterior corticalspinal tract. The lateral tract contains upper motor neuronal axons that synapse on the dorsal lateral lower motor neurons, which are involved in distal limb control.

The anterior corticospinal tract descends ipsilaterally in the anterior column, where the axons emerge and either synapse on ventromedial lower motor neurons in the ventral horn ipsilaterally or descussate at the anterior white commissure where they synapse on ventromedial lower motor neurons contralaterally.

The ventromedial lower motor neurons control the large, postural muscles of the axial skeleton. These lower motor neurons, unlike those of the dorsal lateral, are located in the ventral horn throughout the spinal cord.



Spinal cord tracts: This diagram of spinal cord tracts shows the motor and efferent pathways in red and the sensory and afferent pathways in blue. Included in the diagram are the following motor pathways: corticospinal tracts (pyramidal tract), and extrapyramidal tracts (tectospinal tract not delineated).

This page titled 12.9A: Organization of Motor Neuron Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.9B: The Role of the Basal Ganglia in Movement

The basal ganglia are responsible for voluntary motor control, procedural learning, and eye movement, as well as cognitive and emotional functions.

Learning Objectives

- Describe the role of the basal ganglia in movement

Key Points

- The basal ganglia are studied extensively in the context of two disorders of the basal ganglia: Parksinson's disease and Huntington's disease.
- Hemiballismus, a movement disorder arising from neuronal damage in the subthalamic nucleus, presents with violent movements of the arms and legs.
- Eye movement, a function of the basal ganglia, is influenced by the superior colliculus, a region of the brain that directs eye movement to specific points in space in response to stimuli.
- Basal ganglia are also thought to play a role in motivation.
- In the basal ganglia,
  the majority of the neurons uses gamma-aminobutyric acid (GABA) as the neurotransmitter
  and have inhibitory effects on their
  targets.


Key Terms

- **hemiballismus**: A rare movement disorder with involuntary flinging motions of the extremities.
- **voluntary motor control**: The act of directing motion with intent.
- **forebrain**: The anterior part of the brain, including the cerebrum, thalamus, and hypothalamus.
- **nucleus accumbens**: A region in the basal forebrain rostral to the preoptic area of the hypothalamus. This region and the olfactory tubercle collectively form the ventral striatum.

Location of the Basal Ganglia

The basal ganglia (or basal nuclei) are a group of nuclei of varied origin in the brains of vertebrates that act as a cohesive functional unit. They are situated at the base of the forebrain and are strongly connected with the cerebral cortex, thalamus, and other brain areas.

The basal ganglia are associated with a variety of functions, including voluntary motor control, procedural learning relating to routine behaviors or habits such as bruxism and eye movements, as well as cognitive and emotional functions.



Basal ganglia: Locations of the basal ganglia.

Action Selection

Currently popular theories hold that the basal ganglia play a primary role in action selection. Action selection is the decision of which of several possible behaviors to execute at a given time.

Experimental studies show that the basal ganglia exert an inhibitory influence on a number of motor systems, and that a release of this inhibition permits a motor system to become active. The behavior switching that takes place within the basal ganglia is influenced by signals from many parts of the brain, including the prefrontal cortex, which plays a key role in executive functions.

Movement

The greatest source of insight into the functions of the basal ganglia has come from the study of two neurological disorders, Parkinson's disease and Huntington's disease. For both of these disorders, the nature of the neural damage is well-understood and can be correlated with the resulting symptoms.

Parkinson's disease involves the major loss of dopaminergic cells in the substantia nigra. Huntington's disease involves the massive loss of medium spiny neurons in the striatum.

The symptoms of the two diseases are virtually opposite: Parkinson's disease is characterized by a gradual loss of the ability to initiate movement, whereas Huntington's disease is characterized by an inability to prevent parts of the body from moving unintentionally.

It is noteworthy that, although both diseases have cognitive symptoms, especially in their advanced stages, the most salient symptoms relate to the ability to initiate and control movement. Thus, both are classified primarily as movement disorders.

A different movement disorder, called hemiballismus, may result from damage restricted to the subthalamic nucleus. Hemiballismus is characterized by violent and uncontrollable flinging movements of the arms and legs.

Function in Eye Movement

One of the most intensively studied functions of the basal ganglia is their role in controlling eye movements. Eye movement is influenced by an extensive network of brain regions that converge on a midbrain area called the superior colliculus (SC).

The SC is a layered structure whose layers form two-dimensional retinotopic maps of visual space. A bump of neural activity in

the deep layers of the SC drives eye movement toward the corresponding point in space.

Motivation

Although the role of the basal ganglia in motor control is clear, there are also many indications that it is involved in the control of behavior in a more fundamental way, at the level of motivation. In Parkinson's disease, the ability to execute the components of movement is not greatly affected, but motivational factors such as hunger fail to cause movements to be initiated or switched at the proper times.

The immobility of patients with Parkingson's disease has sometimes been described as a paralysis of the will. These patients have occasionally been observed to show a phenomenon called kinesia paradoxica, in which a person who is otherwise immobile responds to an emergency in a coordinated and energetic way, then lapses back into immobility once the emergency has passed.

The role in motivation of the limbic part of the basal ganglia—the nucleus accumbens (NA), ventral pallidum, and ventral tegmental area (VTA)—is particularly well established. Thousands of experimental studies combine to demonstrate that the dopaminergic projection from the VTA to the NA plays a central role in the brain's reward system.

Numerous things that people find rewarding, including addictive drugs, good-tasting food, and sex, have been shown to elicit activation of the VTA dopamine system. Damage to the NA or VTA can produce a state of profound torpor.


Neurotransmitters

In most regions of the brain, the predominant classes of neurons use glutamate as the neurotransmitter and have excitatory effects on their targets. In the basal ganglia, however, the great majority of neurons uses gamma-aminobutyric acid (GABA) as the neurotransmitter and have inhibitory effects on their targets.

The inputs from the cortex and thalamus to the striatum and subthalamic nucleus are glutamatergic, but the outputs from the striatum, pallidum, and substantia nigra pars reticulata all use GABA. Thus, following the initial excitation of the striatum, the internal dynamics of the basal ganglia are dominated by inhibition and disinhibition.

Other neurotransmitters have important modulatory effects. Dopamine is used by the projection from the substantia nigra pars compacta to the dorsal striatum and also in the analogous projection from the ventral tegmental area to the ventral striatum (nucleus accumbens).

Acetylcholine also plays an important role, as it is used both by several external inputs to the striatum and by a group of striatal interneurons. Although cholinergic cells make up only a small fraction of the total population, the striatum has one of the highest acetylcholine concentrations of any brain structure.



Main circuits of the basal ganglia: This diagram shows the main circuits of the basal ganglia. Two coronal slices have been superimposed to include the involved basal ganglia structures. The + and – signs at the point of the arrows indicate whether the pathway is excitatory or inhibitory, respectively, in effect. Green arrows refer to excitatory glutamatergic pathways, red arrows refer to inhibitory GABAergic pathways and turquoise arrows refer to dopaminergic pathways that are excitatory on the direct pathway and inhibitory on the indirect pathway.

This page titled 12.9B: The Role of the Basal Ganglia in Movement is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.9C: Modulation of Movement by the Cerebellum

The cerebellum is important for motor control—specifically coordination, precision, and timing—as well as some forms of motor learning.

Learning Objectives

- Describe the role of the cerebellum in movement modulation

Key Points

- The cerebellum is a parallel grooved structure at the bottom of the brain containing a highly regular cellular arrangement of Purkinje cells, granule cells, and other cell types.
- The cerebellum adjusts to changes in sensorimotor relationships, possibly functioning as in the Marr-Albus theory: Strong inputs from a single climbing fiber serve as a teaching signal to change the strength of impulses from the corresponding group of parallel fibers.
- Four principles of cerebellum function have been identified. They include: feedforward processing, divergence and convergence, modularity, and plasticity.


Key Terms

- **Purkinje cells**: A class of GABAergic neurons located in the cerebellum.
- **mossy fibers**: One of the major inputs to the cerebellum from sources such as the cerebral cortex.
- **granule cells**: These cells receive excitatory input from mossy fibers that originate from pontine nuclei.

The cerebellum is a region of the brain that plays an important role in motor control. It may also be involved in some cognitive functions such as attention and language, and in regulating fear and pleasure responses, but its movement-related functions are the most solidly established. The cerebellum does not initiate movement, but it contributes to coordination, precision, and accurate timing.


It receives input from sensory systems of the spinal cord and from other parts of the brain, including the cerebral cortex, and integrates these inputs to fine-tune motor activity. Because of this fine-tuning function, damage to the cerebellum does not cause paralysis, but instead produces disorders in fine movement, equilibrium, posture, and motor learning.

The cerebellum differs from most other parts of the brain, especially the cerebral cortex, in regards to the ability of signals to move unidirectionally from input to output. This feedforward mode of operation means that the cerebellum cannot generate self-sustaining patterns of neural activity, in contrast to the cerebral cortex. However, the cerebellum can receive information from the cerebral cortex and processes this information to send motor impulses to the skeletal muscle.



Cerebellum: View of the cerebellum from above and behind.

Anatomy of the Cerebellum



Cerebellum cells: View of the cerebellum from above and behind.

In terms of anatomy, the cerebellum has the appearance of a separate structure attached to the bottom of the brain, tucked underneath the cerebral hemispheres. The surface of the cerebellum is covered with finely spaced parallel grooves, in striking contrast to the broad irregular convolutions of the cerebral cortex. These parallel grooves conceal the fact that the cerebellum is actually a continuous thin layer of tissue (the cerebellar cortex), tightly folded in the style of an accordion.

Within this thin layer are several types of neurons with a highly regular arrangement, the most important being Purkinje cells and granule cells. This complex neural network gives rise to a massive signal-processing capability, but almost all of its output is directed to a set of small, deep cerebellar nuclei lying in the interior of the cerebellum.

Function

Marr-Albus Theory

In addition to its direct role in motor control, the cerebellum is also necessary for several types of motor learning, the most notable one being learning to adjust to changes in sensorimotor relationships.

Several theoretical models have been developed to explain sensorimotor calibration in terms of synaptic plasticity within the cerebellum. Most of them derive from early models formulated by David Marr and James Albus, which were motivated by the observation that each cerebellar Purkinje cell receives two dramatically different types of input.

It receives input from thousands of parallel fibers, each individually very weak. However, each cerebellar Purkinje cell also gets input from one single climbing fiber, which is so strong that a single climbing fiber action potential will reliably cause a target Purkinje cell to fire a burst of action potentials.

The basic concept of the Marr-Albus theory is that the climbing fiber serves as a teaching signal, which induces a long-lasting change in the strength of synchronously activated parallel fiber inputs. Observations of long-term depression in parallel fiber inputs have provided support for theories of this type, but their validity remains controversial.

Insights from Cerebellar Dysfunction

The strongest clues to the function of the cerebellum have come from examining the consequences of damage to it. Animals and humans with cerebellar dysfunction show, above all, problems with motor control. They continue to be able to generate motor activity, but it loses precision, producing erratic, uncoordinated, or incorrectly timed movements.

A standard test of cerebellar function is to reach with the tip of the finger for a target at arm's length. A healthy person will move the fingertip in a rapid straight trajectory, whereas a person with cerebellar damage will reach slowly and erratically, with many mid-course corrections.

Deficits in non-motor functions are more difficult to detect. Thus, the general conclusion reached decades ago is that the basic function of the cerebellum is not to initiate movements, or to decide which movements to execute, but rather to calibrate the detailed form of a movement.

The comparative simplicity and regularity of the cerebellar anatomy led to an early hope that it might imply a similar simplicity of computational function. Although a full understanding of cerebellar function remains elusive, at least four principles are identified as important: 1) feedforward processing, 2) divergence and convergence, 3) modularity, and 4) plasticity.

- Feedforward processing: Refers to the unidirectional movement of signals through the system from input to output, with very little recurrent internal transmission. This means that the cerebellum, in contrast to the cerebral cortex, cannot generate self-sustaining patterns of neural activity. Signals enter the circuit, are processed by each stage in sequential order, and then leave.
- Divergence and convergence: The 1000 or so Purkinje cells belonging to a microzone may receive input from as many as 100 million parallel fibers, and focus their own output down to a group of less than 50 deep nuclear cells. Thus, the cerebellar network receives a modest number of inputs, processes them very extensively through its rigorously structured internal network, and sends out the results via a very limited number of output cells.
- Modularity: The cerebellar system is functionally divided into independent modules. All modules have a similar internal structure, but with different inputs and outputs. The output of one module does not appear to significantly influence the activity of other modules
- Plasticity: The synapses between parallel fibers and Purkinje cells, and the synapses between mossy fibers and deep nuclear cells, are both susceptible to modification of their strength. The influence of each parallel fiber on nuclear cells is adjustable. This arrangement gives tremendous flexibility for fine-tuning the relationship between the cerebellar inputs and outputs.

This page titled 12.9C: Modulation of Movement by the Cerebellum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

12.9D: Functions of the Cerebellum in Integrating Movements

The cerebellum uses feedforward processing and modularity to process information.

Learning Objectives

- Describe the functions of the cerebellum in integrating movments

Key Points

- The function of the cerebellum can be described by the principles of feedforward processing and modularity.
- Feedforward processing means signals move in one direction through the cerebellum, from input to output.
- Modularity describes the modular nature of the cerebellar system, where modules with similar structures function relatively independently. Modules consist of clusters of neurons with common inputs but distinct outputs.

Key Terms

- **purkinje**: Purkinje cells are a class of GABAergic neurons located in the cerebellar cortex. They are some of the largest neurons in the human brain, with an intricately elaborate dendritic arbor characterized by a large number of dendritic spines.
- **feedforward processing**: A property of some neural circuits where signals move unidirectionally through the system from input to output, with very little recurrent internal transmission.
- **microzone**: A group of Purkinje cells that all have the same somatotopic receptive field. Microzones contain on the order of 1,000 Purkinje cells each, arranged in a long, narrow strip, and oriented perpendicular to the cortical folds.

Cerebellar Function


Feedforward Processing

The cerebellum differs from most other parts of the brain in that the signal processing is almost entirely feedforward—that is, signals move unidirectionally through the system from input to output, with very little recurrent internal transmission.

The small amount of recurrence that does exist consists of mutual inhibition; there are no mutually excitatory circuits. This feedforward mode of operation means that the cerebellum, in contrast to the cerebral cortex, cannot generate self-sustaining patterns of neural activity.

Signals enter the circuit, are processed by each stage in sequential order, and then leave. As Eccles, Ito, and Szentágothai wrote,"This elimination in the design of all possibility of reverberatory chains of neuronal excitation is undoubtedly a great advantage in the performance of the cerebellum as a computer, because what the rest of the nervous system requires from the cerebellum is presumably not some output expressing the operation of complex reverberatory circuits in the cerebellum, but rather a quick and clear response to the input of any particular set of information."

Divergence and Convergence



Cells of the Cerebellum: Transverse section of a cerebellar folium, showing its principal cell types and connections.

In the human cerebellum, information from 200 million mossy fiber inputs is expanded to 40 billion granule cells, whose parallel fiber outputs then converge onto 15 million Purkinje cells. Because of the way that they are lined up longitudinally, the 1,000 or so Purkinje cells belonging to a microzone may receive input from as many as 100 million parallel fibers and focus their own output down to a group of less than 50 deep nuclear cells.

Thus, the cerebellar network receives a modest number of inputs, processes them very extensively through its rigorously structured internal network, and sends out the results via a very limited number of output cells.

Modularity


The cerebellar system is functionally divided into more or less independent modules, that probably number in the hundreds to thousands. All modules have a similar internal structure, but different inputs and outputs.

A module (a multizonal microcompartment in the terminology of Apps and Garwicz) consists of a small cluster of neurons in the inferior olivary nucleus, a set of long narrow strips of Purkinje cells in the cerebellar cortex (microzones), and a small cluster of neurons in one of the deep cerebellar nuclei.

Different modules share input from mossy fibers and parallel fibers, but in other respects they appear to function independently. The output of one module does not appear to significantly influence the activity of other modules.

Plasticity

The synapses between parallel fibers and Purkinje cells, and the synapses between mossy fibers and deep nuclear cells, are both susceptible to modification of their strength. In a single cerebellar module, input from as many as a billion parallel fibers converge onto a group of less than 50 deep nuclear cells, and the influence of each parallel fiber on those nuclear cells is adjustable. This arrangement gives tremendous flexibility for fine-tuning the relationships between the cerebellar inputs and outputs.



Zones and microzones in the cerebellum: This schematic illustration of the structure of zones and microzones in the cerebellum shows three levels of magnification. These zones and microzones help explain the modular nature of the cerebellar function. On the left is a simplified illustration of what the cerebellar cortex would look like if all the folds were straightened out—the vertical dimension is the rostro-caudal axis of the cerebellum, the horizontal dimension is the medio-lateral axis. A zone is a longitudinally oriented strip of the cortex, and a microzone is a thin, longitudinally oriented portion of a zone. As the illustration on the right shows, Purkinje cell dendritic trees are flattened in a way that aligns with the microzone length, and parallel fibers cross the microzones at right angles.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Spinal tracts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_...r_organization**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Motor system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Motor_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Upper motor neuron. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Upper_m...ron%23Pathways**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- motor system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/motor%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- corticospinal tract. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/corticospinal%20tract**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pyramidal Tracts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pyramidal_tracts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebral cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cerebral_cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spinal cord tracts - English. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sp..._-_English.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray684. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray684.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brodmann-areas. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Brodmann-areas.png**. **License**: ***Public Domain: No Known Copyright***
- Basal ganglia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Basal_ganglia%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- hemiballismus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hemiballismus. **License**: *CC BY-SA: Attribution-ShareAlike*
- voluntary motor control. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/volunta...otor%20control. **License**: *CC BY-SA: Attribution-ShareAlike*
- forebrain. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/forebrain. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nucleus Accumbens. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nucleus_accumbens. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal cord tracts - English. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sp..._-_English.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray684. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray684.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brodmann-areas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brodmann-areas.png. **License**: *Public Domain: No Known Copyright*
- Basal ganglia circuits. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...a_circuits.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basal Ganglia.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Basal_...salGanglia.png. **License**: *CC BY: Attribution*
- Cerebellum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebellum%23Principles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cerebellum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebellum. **License**: *CC BY-SA: Attribution-ShareAlike*
- cerebellum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cerebellum. **License**: *CC BY-SA: Attribution-ShareAlike*
- Granule Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Granule_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Purkinje Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Purkinje_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mossy Fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mossy_...r_(cerebellum). **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal cord tracts - English. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sp..._-_English.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray684. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray684.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brodmann-areas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brodmann-areas.png. **License**: *Public Domain: No Known Copyright*
- Basal ganglia circuits. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ba...a_circuits.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Basal Ganglia.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Basal_ganglia#/media/File:Blausen_0076_BasalGanglia.png. **License**: *CC BY: Attribution*
- Sobo 1909 653. **Provided by**: Wikipedial Commons. **Located at**: commons.wikimedia.org/wiki/File:Sobo_1909_653.png. **License**: *Public Domain: No Known Copyright*
- Gray706. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gray706.png. **License**: *Public Domain: No Known Copyright*
- Cerebellum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cerebellum%23Principles. **License**: *CC BY-SA: Attribution-ShareAlike*
- purkinje. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/purkinje. **License**: *CC BY-SA: Attribution-ShareAlike*
- microzone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/microzone. **License**: *CC BY-SA: Attribution-ShareAlike*
- feedforward processing. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/feedforward%20processing. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spinal cord tracts - English. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Spinal_cord_tracts_-_English.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray684. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray684.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brodmann-areas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Brodmann-areas.png. **License**: *Public Domain: No Known Copyright*

- Basal ganglia circuits. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ba...a_circuits.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basal Ganglia.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Basal_ganglia#/media/File:Blausen_0076_BasalGanglia.png**. **License**: ***CC BY: Attribution***
- Sobo 1909 653. **Provided by**: Wikipedial Commons. **Located at**: **commons.wikimedia.org/wiki/File:Sobo_1909_653.png**. **License**: ***Public Domain: No Known Copyright***
- Gray706. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Gray706.png**. **License**: ***Public Domain: No Known Copyright***
- Gray706. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray706.png**. **License**: ***Public Domain: No Known Copyright***
- Microzone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Microzone.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 12.9D: Functions of the Cerebellum in Integrating Movements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13: SPECIAL SENSES



- Topic hierarchy

13.1: The Vertebrate Visual System

13.1A: Anatomy of the Eye

13.1B: Transduction of Light

13.1C: Visual Processing

13.2: Taste and Olfaction

13.2A: Tastes and Odors

13.2B: Reception and Transduction

13.3: Hearing and Equilibrium

13.3A: Sound

13.3B: The Vestibular System

13.3C: Reception of Sound

13.3D: Transduction of Sound

This page titled 13: Special Senses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.1: THE VERTEBRATE VISUAL SYSTEM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

13.1A: Anatomy of the Eye

13.1B: Transduction of Light

13.1C: Visual Processing


This page titled 13.1: The Vertebrate Visual System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.1A: Anatomy of the Eye

Learning Objectives

- Explain how eyes have evolved to benefit organisms

The retina, a thin layer of cells located on the inner surface of the back of the eye, consists of photoreceptive cells, which are responsible for the transduction of light into nervous impulses. However, light does not enter the retina unaltered; it must first pass through other layers that process it so that it can be interpreted by the retina.




Retina: (a) The human eye is shown in cross section. The human eye contains structures, such as the cornea, iris, lens, and fovea, that process light so it can be deciphered by the retina. Other structures like the aqueous humor and the vitreous humor help maintain the shape of the eye. (b) A blowup shows the layers of the retina. The retina contains photoreceptive cells. In the retina, light is converted into neural signals sent to the brain.

The cornea, the front transparent layer of the eye, along with the crystalline lens, refract (bend) light to focus the image on the retina. After passing through the cornea, light passes through the aqueous humour, which connects the cornea to the lens. This clear gelatinous mass also provides the corneal epithelium with nutrients and helps maintain the convex shape of the cornea. The iris, which is visible as the colored part of the eye, is a circular muscular ring lying between the lens and the aqueous humour that regulates the amount of light entering the eye. Light passes through the center of the iris, the pupil, which actively adjusts its size to maintain a constant level of light entering the eye. In conditions of high ambient light, the iris contracts, reducing the size of the pupil. In conditions of low light, the iris relaxes and the pupil enlarges.

The main function of the lens is to focus light on the retina and fovea centralis. The lens is a transparent, convex structure located behind the cornea. On the other side of the lens is the vitreous humour, which lets light through without refraction, maintains the shape of the eye, and suspends the delicate lens. The lens focuses and re-focuses light as the eye rests on near and far objects in the visual field. The lens is operated by muscles that stretch it flat or allow it to thicken, changing the focal length of light coming through to focus it sharply on the retina. With age comes the loss of the flexibility of the lens; a form of farsightedness called presbyopia results. Presbyopia occurs because the image focuses behind the retina. It is a deficit similar to a different type of farsightedness, hyperopia, caused by an eyeball that is too short. For both defects, images in the distance are clear, but images nearby are blurry. Myopia (nearsightedness) occurs when an eyeball is elongated and the image focus falls in front of the retina. In this case, images in the distance are blurry, but images nearby are clear.

There are two types of photoreceptors in the retina: rods and cones. Both are named for their general appearance. Rods, strongly photosensitive, are located in the outer edges of the retina. They detect dim light and are used primarily for peripheral and

nighttime vision. Cones, weakly photosensitive, are located near the center of the retina. They respond to bright light; their primary role is in daytime, color vision.



Rods and cones: Rods and cones are photoreceptors in the retina. Rods respond in low light and can detect only shades of gray. Cones respond in intense light and are responsible for color vision.

The fovea is the region in the center back of the eye that is responsible for acute (central) vision. The fovea has a high density of cones. When you bring your gaze to an object to examine it intently in bright light, the eyes orient so that the object's image falls on the fovea. However, when looking at a star in the night sky or other object in dim light, the object can be better viewed by the peripheral vision because it is the rods at the edges of the retina, rather than the cones at the center, that operate better in low light. In humans, cones far outnumber rods in the fovea.

Key Points

- The cornea and the lens bend light to focus the image on the retina; the iris and pupil regulate the amount of light entering the eye.
- The aqueous humour maintains the convex shape of the cornea; the vitreous humour supports the lens and maintains the shape of the entire eye.
- Presbyopia occurs because the image focuses behind the retina; it is similar to hyperopia (farsightedness), which is caused by an eyeball that is too short.
- Myopia (nearsightedness) occurs when an eyeball is elongated; images in the distance appear blurry, but images nearby are clear.
- Rods are used for peripheral and nighttime vision; cones are used for daytime and color vision.
- The fovea is responsible for acute vision because it has a high density of cones.

Key Terms

- **rod**: a rod-shaped cell located in the outer retina of the eye that is extremely sensitive to light
- **retina**: the thin layer of cells at the back of the eyeball where light is converted into neural signals sent to the brain
- **cone**: cell located near the center of the retina that is weakly photosensitive and is responsible for color vision in relatively bright light

This page titled 13.1A: Anatomy of the Eye is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.1B: Transduction of Light

Light is tranduced in rods and cones; visual information is processed in the retina before entering the brain.

Learning Objectives

- Explain retinal processing and the process of transduction of light

Key Points


- When light hits the photoreceptor, the retinal changes shape, which activates the photopigment rhodoposin.
- Primates have full color vision because of the three- cone (trichromatic) system; color is a result of the ratio of activity of the three types of cones.
- There are three types of cones with different photopsins: S cones respond to short waves; M cones respond to medium waves; L cones respond to light to long waves.
- If light is not present, neurons are inhibited by rods and cones; once light is introduced, rods and cones are hyperpolarized, which activates the neurons.
- Activated neurons stimulate ganglion cells, which send action potentials via the optic nerve.
- Horizontal cells can create lateral inhibition, which enhances light and dark contrast in images.

Key Terms


- **tonic activity**: when photoreceptors become slightly active even when not stimulated by light
- **rhodopsin**: a light-sensitive pigment in the rod cells of the retina; it consists of an opsin protein bound to the carotenoid retinal

Transduction of Light

The rods and cones are the site of transduction of light into a neural signal. Both rods and cones contain photopigments, which are pigments that undergo a chemical change when they absorb light. In vertebrates, the main photopigment, rhodopsin, has two main parts: an opsin, which is a membrane protein (in the form of a cluster of α-helices that span the membrane); and retinal, a molecule that absorbs light. When light hits a photoreceptor, it causes a shape change in the retinal, altering its structure from a bent (cis) form of the molecule to its linear (trans) isomer. This isomerization of retinal activates the rhodopsin, starting a cascade of events that ends with the closing of Na^+ channels in the membrane of the photoreceptor. Thus, unlike most other sensory neurons (which become depolarized by exposure to a stimulus), visual receptors become hyperpolarized and are driven away from the threshold.




Hyperpolarized visual receptors: When light strikes rhodopsin, the G-protein transducin is activated, which in turn activates phosphodiesterase. Phosphodiesterase converts cGMP to GMP, thereby closing sodium channels. As a result, the membrane becomes hyperpolarized. The hyperpolarized membrane does not release glutamate to the bipolar cell.



Rhodopsin: (a) Rhodopsin, the photoreceptor in vertebrates, has two parts: the trans-membrane protein opsin and retinal. When light strikes the retinal, it changes shape from (b) a cis to a trans form. The signal is passed to a G-protein called transducin, triggering a series of downstream events.

Trichromatic Coding

There are three types of cones (with different photopsins) that differ in the wavelength to which they are most responsive. Some cones are maximally responsive to short light waves of 420 nm; they are called S cones (“S” for “short”). Other cones (M cones, for “medium”) respond maximally to waves of 530 nm. A third group (L cones, or “long” cones) responds maximally to light of longer wavelengths at 560 nm. With only one type of cone, color vision would not be possible; a two-cone (dichromatic) system has limitations. Primates use a three-cone (trichromatic) system, resulting in full color vision.



Rod and cone cells: Human rod cells and the different types of cone cells each have an optimal wavelength. However, there is considerable overlap in the wavelengths of light detected.

The color we perceive is a result of the ratio of activity of our three types of cones. The colors of the visual spectrum, running from long-wavelength light to short are:

- red (700 nm)
- orange (600 nm)
- yellow (565 nm)
- green (497 nm)
- blue (470 nm)
- indigo (450 nm)
- violet (425 nm).

Humans have very sensitive perception of color and can distinguish about 500 levels of brightness, 200 different hues, and 20 steps of saturation; in all, about 2 million distinct colors.

Retinal Processing

Visual signals leave the cones and rods, travel to the bipolar cells, and then to ganglion cells. A large degree of processing of visual information occurs in the retina itself, before visual information is sent to the brain.

Photoreceptors in the retina continuously undergo tonic activity. That is, they are always slightly active even when not stimulated by light. In neurons that exhibit tonic activity, the absence of stimuli maintains a firing rate at an equilibrium; while some stimuli increase firing rate from the baseline, other stimuli decrease firing rate. In the absence of light, the bipolar neurons that connect rods and cones to ganglion cells are continuously and actively inhibited by the rods and cones. Exposure of the retina to light hyperpolarizes the rods and cones, removing the inhibition of their bipolar cells. The now-active bipolar cells in turn stimulate the ganglion cells, which send action potentials along their axons (which leave the eye as the optic nerve). Thus, the visual system relies on changein retinal activity, rather than the absence or presence of activity, to encode visual signals for the brain. Sometimes horizontal cells carry signals from one rod or cone to other photoreceptors and to several bipolar cells. When a rod or cone stimulates a horizontal cell, the horizontal cell inhibits more-distant photoreceptors and bipolar cells, creating lateral inhibition. This inhibition sharpens edges and enhances contrast in the images by making regions receiving light appear lighter and dark surroundings appear darker. Amacrine cells can distribute information from one bipolar cell to many ganglion cells.

This page titled 13.1B: Transduction of Light is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.1C: Visual Processing

Visual signals are processed in the brain through several different pathways.

Learning Objectives

- Describe the complexity of visual processing in the brain

Key Points

- The magnocellular pathway carries information about form, movement, depth, and differences in brightness; the parvocellular pathway carries information on color and fine detail.
- The optic chiasma allows us to coordinate information between both eyes and is produced by crossing optical information across the brain.
- Visual signals move from the visual cortex to either the parietal lobe or the temporal lobe.
- Some signals move to the thalamus, which sends the visual signals to the primary cortex.
- Visual signals can also travel from the retina to the superior colliculus, where eye movements are coordinated with auditory information.
- Visual signals can move from the retina to the suprachiasmatic nucleus (SCN), the body's internal clock, which is involved in sleep/wake patterns and annual cycles.

Key Terms


- **superior colliculus**: the primary area of the brain where eye movements are coordinated and integrated with auditory information
- **optic chiasma**: found at the base of the brain and coordinates information from both eyes
- **suprachiasmatic nucleus**: cluster of cells that is considered to be the body's internal clock, which controls our circadian (day-long) cycle

Higher Processing

The myelinated axons of ganglion cells make up the optic nerves. Within the nerves, different axons carry different parts of the visual signal. Some axons constitute the magnocellular (big cell) pathway, which carries information about form, movement, depth, and differences in brightness. Other axons constitute the parvocellular (small cell) pathway, which carries information on color and fine detail. Some visual information projects directly back into the brain, while other information crosses to the opposite side of the brain. This crossing of optical pathways produces the distinctive optic chiasma (Greek, for "crossing") found at the base of the brain and allows us to coordinate information from both eyes.

Once in the brain, visual information is processed in several places. Its routes reflect the complexity and importance of visual information to humans and other animals. One route takes the signals to the thalamus, which serves as the routing station for all incoming sensory impulses except smell. In the thalamus, the magnocellular and parvocellular distinctions remain intact; there are different layers of the thalamus dedicated to each. When visual signals leave the thalamus, they travel to the primary visual cortex at the rear of the brain. From the visual cortex, the visual signals travel in two directions. One stream that projects to the parietal lobe, in the side of the brain, carries magnocellular ("where") information. A second stream projects to the temporal lobe and carries both magnocellular ("where") and parvocellular ("what") information.

Another important visual route is a pathway from the retina to the superior colliculus in the midbrain, where eye movements are coordinated and integrated with auditory information. Finally, there is the pathway from the retina to the suprachiasmatic nucleus (SCN) of the hypothalamus. The SCN is a cluster of cells that is considered to be the body's internal clock, which controls our circadian (day-long) cycle. The SCN sends information to the pineal gland, which is important in sleep/wake patterns and annual cycles.



The suprachiasmatic nucleus (SNC): The presence of light and darkness influences circadian rhythms and related physiology and behavior through the SCN.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Ophthalmology/Anatomy of the Eye. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Ophthal...omy_of_the_Eye**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- rod. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/rod**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- retina. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/retina**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_02.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- transduction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/transduction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...tonic-activity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- rhodopsin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/rhodopsin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_02.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_06.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/latest...e_36_05_04.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44761/**

latest...e_36_05_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de.../optic-chiasma. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...smatic-nucleus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...ior-colliculus. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...e_36_05_02.png. **License**: *CC BY: Attribution*
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...e_36_05_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...e_36_05_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...e_36_05_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Vision. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44761/latest...e_36_05_05.jpg. **License**: *CC BY: Attribution*
- Suprachiasmatic nucleus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Suprachiasmatic_nucleus. **License**:

This page titled 13.1C: Visual Processing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.2: TASTE AND OLFACTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

13.2A: Tastes and Odors

13.2B: Reception and Transduction

This page titled 13.2: Taste and Olfaction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.2A: Tastes and Odors

The senses of taste and smell are related because they use the same types of receptors and are stimulated by molecules in solutions or air.

Learning Objectives

- Explain the interaction of taste and odor

Key Points


- Humans can taste sweet, sour, bitter, salty, and umami; umami is the savoriness of certain foods that are commonly high in protein.
- Odors come from molecules in the air that stimulate receptors in the nose; if an organism does not have a receptor for that particular odor molecule, for that organism, the odor has no smell.
- The senses of smell and taste are directly related because they both use the same types of receptors.
- If one's sense of smell is not functional, then the sense of taste will also not function because of the relationship of the receptors.

Key Terms

- **umami**: one of the five basic tastes, the savory taste of foods such as seaweed, cured fish, aged cheeses and meats
- **olfactory**: concerning the sense of smell
- **receptor**: a protein on a cell wall that binds with specific molecules so that they can be absorbed into the cell in order to control certain functions

Tastes and Odors

Both taste and odor stimuli are molecules taken in from the environment. The primary tastes detected by humans are sweet, sour, bitter, salty, and umami. The first four tastes need little explanation. The identification of umami as a fundamental taste occurred fairly recently. It was identified in 1908 by Japanese scientist Kikunae Ikeda while he worked with seaweed broth, but it was not widely accepted as a taste that could be physiologically distinguished until many years later. The taste of umami, also known as savoriness, is attributable to the taste of the amino acid L-glutamate. In fact, monosodium glutamate, or MSG, is often used in cooking to enhance the savory taste of certain foods. The adaptive value of being able to distinguish umami is that savory substances tend to be high in protein.




Uniform Distribution of Taste Receptors: Humans detect taste using receptors called taste buds. Each of these receptors is specially adapted to determine one type of taste sensation. Recent evidence suggests that taste receptors are uniformly distributed across the tongue; thus, the traditional tongue map is no longer valid.

All odors that we perceive are molecules in the air we breathe. If a substance does not release molecules into the air from its surface, it has no smell. If a human or other animal does not have a receptor that recognizes a specific molecule, then that molecule has no smell. Humans have about 350 olfactory receptor subtypes that work in various combinations to allow us to sense about 10,000 different odors. Compare that to mice, for example, which have about 1,300 olfactory receptor types and, therefore, probably sense many more odors.

The senses of smell and taste combine at the back of the throat. When you taste something before you smell it, the smell lingers internally up to the nose causing you to smell it. Both smell and taste use chemoreceptors, which essentially means they are both sensing the chemical environment. This chemoreception in regards to taste, occurs via the presence of specialized taste receptors within the mouth that are referred to as taste cells and are bundled together to form taste buds. These taste buds, located in papillae which are found across the tongue, are specific for the five modalities: salt, sweet, sour, bitter and umami. These receptors are activated when their specific stimulus (i.e. sweet or salt molecules) is present and signals to the brain.

In addition to the activation of the taste receptors, there are similar receptors within the nose that coordinates with activation of the taste receptors. When you eat something, you can tell the difference between sweet and bitter. It is the sense of smell that is used to distinguish the difference. Although humans commonly distinguish taste as one sense and smell as another, they work together to create the perception of flavor. A person's perception of flavor is reduced if he or she has congested nasal passages.



Uniform distribution of taste receptors (the myth of the tongue map): Humans detect taste using receptors called taste buds. Each of these receptors is specially adapted to determine one type of taste sensation. Recent evidence suggests that taste receptors are uniformly distributed across the tongue; thus, this traditional tongue map is no longer valid.

This page titled 13.2A: Tastes and Odors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.2B: Reception and Transduction

Odorants and tastants produce signal molecules received by receptors, which are then processed by the brain to identify smells and tastes.

Learning Objectives

- Describe the process by which tastes and odors are sensed

Key Points


- Odorants are received by receptors in the nose, which send signals to the olfactory bulb of the brain to create an appropriate response; humans have about 12 million receptors.
- Taste results when molecules are dissolved in fluid and reach the gustatory receptors on the tongue; the signals are sent to the brain to determine which flavor (bitter, sour, sweet, salty, umami) is being consumed.
- Taste buds are found on the tongue and contain clusters of gustatory receptors on bumps called papillae; fungiform papillae each contain one to eight taste buds; they also have receptors for pressure and temperature.
- The ability to smell and taste declines with age.

Key Terms

- **tastant**: any substance that stimulates the sense of taste
- **papilla**: a nipple-like anatomical structure
- **odorant**: any substance that has a distinctive smell, especially one added to something (such as household gas) for safety purposes

Reception and Transduction

Odorants (odor molecules) enter the nose and dissolve in the olfactory epithelium, the mucosa at the back of the nasal cavity. The olfactory epithelium is a collection of specialized olfactory receptors in the back of the nasal cavity that spans an area about 5 cm^2 in humans. Recall that sensory cells are neurons. An olfactory receptor, which is a dendrite of a specialized neuron, responds when it binds certain molecules inhaled from the environment by sending impulses directly to the olfactory bulb of the brain. Humans have about 12 million olfactory receptors distributed among hundreds of different receptor types that respond to different odors. Twelve million seems like a large number of receptors, but compare that to other animals: rabbits have about 100 million, most dogs have about 1 billion, and bloodhounds (dogs selectively bred for their sense of smell) have about 4 billion.





Human olfactory system: In the human olfactory system, (a) bipolar olfactory neurons extend from (b) the olfactory epithelium, where olfactory receptors are located, to the olfactory bulb.

Olfactory neurons are bipolar neurons (neurons with two processes from the cell body). Each neuron has a single dendrite buried in the olfactory epithelium; extending from this dendrite are 5 to 20 receptor-laden, hair-like cilia that trap odorant molecules. The sensory receptors on the cilia are proteins. It is the variations in their amino acid chains that make the receptors sensitive to different odorants. Each olfactory sensory neuron has only one type of receptor on its cilia. The receptors are specialized to detect specific odorants, so the bipolar neurons themselves are specialized. When an odorant binds with a receptor that recognizes it, the sensory neuron associated with the receptor is stimulated. Olfactory stimulation is the only sensory information that directly reaches the cerebral cortex, whereas other sensations are relayed through the thalamus.

Taste and Smell

Detecting a taste (gustation) is fairly similar to detecting an odor (olfaction), given that both taste and smell rely on chemical receptors being stimulated by certain molecules. The primary organ of taste is the taste bud. A taste bud is a cluster of gustatory receptors (taste cells) that are located within the bumps on the tongue called papillae (singular: papilla). There are several structurally-distinct papillae. Filiform papillae, which are located across the tongue, are tactile, providing friction that helps the tongue move substances; they contain no taste cells. In contrast, fungiform papillae, which are located mainly on the anterior two-thirds of the tongue, each contain one to eight taste buds; they also have receptors for pressure and temperature. The large circumvallate papillae contain up to 100 taste buds and form a V near the posterior margin of the tongue.





Taste buds: (a) Foliate, circumvallate, and fungiform papillae are located on different regions of the tongue. (b) Foliate papillae are prominent protrusions on this light micrograph.

In humans, there are five primary tastes; each taste has only one corresponding type of receptor. Thus, like olfaction, each receptor is specific to its stimulus (tastant). Transduction of the five tastes happens through different mechanisms that reflect the molecular composition of the tastant. A salty tastant (containing NaCl) provides the sodium ions (Na^+) that enter the taste neurons, exciting them directly. Sour tastants are acids which belong to the thermoreceptor protein family. Binding of an acid or other sour-tasting molecule triggers a change in the ion channel which increases hydrogen ion (H^+) concentrations in the taste neurons; thus, depolarizing them. Sweet, bitter, and umami tastants require a G-protein-coupled receptor. These tastants bind to their respective receptors, thereby exciting the specialized neurons associated with them.

Both tasting abilities and sense of smell change with age. In humans, the senses decline dramatically by age 50 and continue to decline. A child may find a food to be too spicy, whereas an elderly person may find the same food to be bland and unappetizing.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44764/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Smell Fall 2011. **Provided by**: virtualgardneranatphys Wikispace. **Located at**: https://virtualgardneranatphys.wikis...mell+Fall+2011. **License**: ***CC BY-SA: Attribution-ShareAlike***
- receptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/receptor. **License**: ***CC BY-SA: Attribution-ShareAlike***
- olfactory. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/olfactory. **License**: ***CC BY-SA: Attribution-ShareAlike***
- umami. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/umami. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ehumanbiofield - Sense of Taste WikiPage NLL. **Provided by**: Wikispaces. **Located at**: https://ehumanbiofield.wikispaces.co...e+WikiPage+NLL. **License**: ***CC BY-SA: Attribution-ShareAlike***
- The Tongue. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1402_The_Tongue.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44764/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- tastant. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tastant. **License**: ***CC BY-SA: Attribution-ShareAlike***
- papilla. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/papilla. **License**: ***CC BY-SA: Attribution-ShareAlike***
- odorant. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/odorant. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ehumanbiofield - Sense of Taste WikiPage NLL. **Provided by**: Wikispaces. **Located at**: https://ehumanbiofield.wikispaces.co...e+WikiPage+NLL. **License**: ***CC BY-SA: Attribution-ShareAlike***
- The Tongue. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/File:1402_The_Tongue.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Taste and Smell. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44764/latest...e_36_03_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Taste and Smell. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44764/latest...e_36_03_03.jpg. **License**: ***CC BY: Attribution***

This page titled 13.2B: Reception and Transduction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.3: HEARING AND EQUILIBRIUM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

13.3A: Sound

13.3B: The Vestibular System

13.3C: Reception of Sound

13.3D: Transduction of Sound

This page titled 13.3: Hearing and Equilibrium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.3A: Sound

Sound waves, characterized by frequency and amplitude, are perceived uniquely by different organisms.

Learning Objectives

- Describe the relationship of amplitude and frequency of a sound wave to attributes of sound

Key Points

- Sound waves are mechanical pressure waves that must travel through a medium and cannot exist in a vacuum.
- There are four main characteristics of a sound wave: frequency, wavelength, period, and amplitude.
- Frequency is the number of waves per unit of time and is heard as pitch; high-frequency sounds are high-pitched, and low-frequency sounds are low-pitched.
- Most humans can perceive sounds with frequencies between 30 and 20,000 Hz; other animals, such as dolphins, can detect sounds at far higher frequencies.
- Amplitude, the dimension of a wave from peak to trough, is heard as volume; louder sounds have greater amplitudes than those of softer sounds.

Key Terms


- **frequency**: characterized as a periodic vibration that is audible; property of sound that most determines pitch and is measured in hertz
- **amplitude**: measure of a wave from its highest point to its lowest point; heard as volume
- **ultrasound**: sound frequencies above the human detectable ceiling of approximately 20,000 Hz

Sound

Auditory stimuli are sound waves, which are mechanical pressure waves that move through a medium, such as air or water. There are no sound waves in a vacuum since there are no air molecules for the waves to move through. The speed of sound waves differs based on altitude, temperature, and medium. At sea level and a temperature of 20° C (68° F), sound waves travel in the air at about 343 meters per second.

As is true for all waves, there are four main characteristics of a sound wave: frequency, wavelength, period, and amplitude. Frequency is the number of waves per unit of time; in sound, it is heard as pitch. High-frequency (≥15.000Hz) sounds are higher-pitched (short wavelength) than low-frequency (long wavelengths; ≤100Hz) sounds. Frequency is measured in cycles per second. For sound, the most-commonly used unit is hertz (Hz), or cycles per second. Most humans can perceive sounds with frequencies between 30 and 20,000 Hz. Women are typically better at hearing high frequencies, but everyone's ability to hear high frequencies decreases with age. Dogs detect up to about 40,000 Hz; cats, 60,000 Hz; bats, 100,000 Hz; dolphins, 150,000 Hz; and the American shad (*Alosa sapidissima*), a fish, can hear 180,000 Hz. Those frequencies above the human range are called ultrasound.

Amplitude, or the dimension of a wave from peak to trough, in sound is heard as volume. The sound waves of louder sounds have greater amplitude than those of softer sounds. For sound, volume is measured in decibels (dB). The softest sound that a human can hear is the zero point. Humans speak normally at 60 decibels.



Amplitude: For sound waves, wavelength corresponds to pitch. The amplitude of the wave corresponds to volume. The sound wave shown with a dashed line is softer in volume than the sound wave shown with a solid line.

This page titled 13.3A: Sound is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.3B: The Vestibular System

Gravity, acceleration, and deceleration are detected by evaluating the inertia on receptive cells in the vestibular system.

Learning Objectives

- Identify the structures of the vestibular system that respond to gravity

Key Points

- The vestibular system uses hair cells, as does the auditory system, but it excites them in different ways.
- There are five vestibular receptor organs in the inner ear (the vestibular labyrinth): the utricle, the saccule, and three semicircular canals; the utricle and saccule respond to acceleration in a straight line, such as gravity.
- The bending of the stereocilia stimulates specific neurons that signal to the brain that the head is tilted, allowing the maintenance of balance.
- The fluid-filled semicircular canals are tubular loops set at oblique angle, arranged in three spatial planes; the base of each canal contains a cluster of hair cells that monitor angular acceleration and deceleration from rotation.
- Neuronal projections to the temporal cortex account for feelings of dizziness; projections to autonomic nervous system areas in the brainstem account for motion sickness; and projections to the primary somatosensory cortex monitor subjective measurements of the external world and self-movement.


Key Terms

- **vestibulocochlear**: of or pertaining to the vestibular and cochlear nerves
- **vestibular system**: the sensory system in mammals that contributes to movement, sense of balance, and spatial orientation
- **stereocilium**: any of many nonmotile cellular structures resembling long microvilli; those of the inner ear are responsible for auditory transduction

Vestibular Information

The stimuli associated with the vestibular system are linear acceleration (gravity) and angular acceleration/deceleration. Gravity, acceleration, and deceleration are detected by evaluating the inertia on receptive cells in the vestibular system. Gravity is detected through head position, while angular acceleration and deceleration are expressed through turning or tilting of the head.

The vestibular system has some similarities with the auditory system. It utilizes hair cells just like the auditory system, but it excites them in different ways. There are five vestibular receptor organs in the inner ear, all of which help to maintain balance: the utricle, the saccule, and three semicircular canals. Together, they make up what is known as the vestibular labyrinth. The utricle and saccule are most responsive to acceleration in a straight line, such as gravity. The roughly 30,000 hair cells in the utricle and 16,000 hair cells in the saccule lie below a gelatinous layer, with their stereocilia (singular: stereocilium) projecting into the gelatin. Embedded in this gelatin are calcium carbonate crystals, similar to tiny rocks. When the head is tilted, the crystals continue to be pulled straight down by gravity, but the new angle of the head causes the gelatin to shift, thereby bending the stereocilia. The bending of the stereocilia stimulates specific neurons that signal to the brain that the head is tilted, allowing the maintenance of balance. It is the vestibular branch of the vestibulocochlear cranial nerve that deals with balance.



Vestibular labrynth: The structure of the vestibular labyrinth is made up of five vestibular receptor organs in the inner ear: the utricle, the saccule, and three semicircular canals.

The fluid-filled semicircular canals are tubular loops set at oblique angles, arranged in three spatial planes. The base of each canal has a swelling that contains a cluster of hair cells. The hairs project into a gelatinous cap, the cupula, where they monitor angular acceleration and deceleration from rotation. They would be stimulated by driving your car around a corner, turning your head, or falling forward. One canal lies horizontally, while the other two lie at about 45 degree angles to the horizontal axis. When the brain processes input from all three canals together, it can detect angular acceleration or deceleration in three dimensions. When the head turns, the fluid in the canals shifts, thereby bending stereocilia and sending signals to the brain. Upon cessation of acceleration or deceleration, the movement of the fluid within the canals slows or stops. For example, imagine holding a glass of water. When moving forward, water may splash backwards onto the hand; when motion has stopped, water may splash forward onto the fingers. While in motion, the water settles in the glass and does not splash. Note that the canals are not sensitive to velocity itself, but to changes in velocity. In this way, moving forward at 60 mph with your eyes closed would not give the sensation of movement, but suddenly accelerating or braking would stimulate the receptors.

Higher Processing

Hair cells from the utricle, saccule, and semicircular canals also communicate through bipolar neurons to the cochlear nucleus in the medulla. Cochlear neurons send descending projections to the spinal cord and ascending projections to the pons, thalamus, and cerebellum. Connections to the cerebellum are important for coordinated movements. There are also projections to the temporal cortex, which account for feelings of dizziness; projections to autonomic nervous system areas in the brainstem, which account for motion sickness; and projections to the primary somatosensory cortex, which monitors subjective measurements of the external world and self-movement. People with lesions in the vestibular area of the somatosensory cortex see vertical objects in the world as being tilted. Finally, the vestibular signals project to certain optic muscles to coordinate eye and head movements.

This page titled 13.3B: The Vestibular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.3C: Reception of Sound

The outer, middle, and inner structures of the ear collect sound energy, converting it to audible sound.

Learning Objectives

- Explain how animals sense sound

Key Points

- The human ear can be divided into three functional segments: the outer ear, the middle ear, and the inner ear.
- Sound waves are collected by the pinna, travel through the auditory canal, and cause vibration of the tympanum (ear drum).
- The three ossicles of the middle ear (malleus, incus, and stapes) transfer energy from the vibrating ear drum to the inner ear.
- The incus connects the malleus to the stapes, which allows vibrations to reach the inner ear.

Key Terms

- **malleus**: small hammer-shaped bone of the middle ear
- **incus**: small anvil-shaped bone in the middle ear; connects the malleus to the stapes
- **stapes**: small stirrup-shaped bone of the middle ear
- **pinna**: the visible, cartilaginous part of the ear that resides outside of the head and collects sound waves
- **tympanum**: innermost part of the outer ear; the eardrum


Reception of Sound

In order to hear a sound, the auditory system must accomplish three basic tasks. First, it must deliver the acoustic stimulus to the receptors; second, it must convert the stimulus from pressure changes into electrical signals; and third, it must process these electrical signals so that they can efficiently indicate the qualities of the sound source, such as frequency (pitch), amplitude (loudness, volume), and location.

The human ear can be divided into three functional segments:

- the outer ear: collects sound energy from the environment and sends it to the eardrum
- the middle ear: transduces the mechanical pressure signals from the ear drum into electrical signals
- the inner ear: interprets the electrical signals from the middle ear using hair cells

In mammals, sound waves are collected by the external, cartilaginous outer part of the ear called the pinna. They then travel through the auditory canal, causing vibration of the thin diaphragm called the tympanum, or ear drum, the innermost part of the outer ear. Interior to the tympanum is the middle ear, which holds three small bones called the ossicles (“little bones”), that transfer energy from the moving tympanum to the inner ear. The three ossicles are the malleus (also known as the hammer), the incus (the anvil), and stapes (the stirrup). The three ossicles are unique to mammals; each plays a role in hearing. The malleus attaches at three points to the interior surface of the tympanic membrane. The incus attaches the malleus to the stapes. In humans, the stapes is not long enough to reach the tympanum. If we did not have the malleus and the incus, then the vibrations of the tympanum would never reach the inner ear. These bones also function to collect force and amplify sounds. The ear ossicles are homologous to bones in a fish mouth; the bones that support gills in fish are thought to be adapted for use in the vertebrate ear over evolutionary time. Many animals (frogs, reptiles, and birds, for example) use the stapes of the middle ear to transmit vibrations to it.



Human ear: Sound travels through the outer ear to the middle ear, which is bounded on its exterior by the tympanic membrane. The middle ear contains three bones called ossicles that transfer the sound wave to the oval window, the exterior boundary of the inner ear.

This page titled 13.3C: Reception of Sound is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

13.3D: Transduction of Sound

When sound waves reach the ear, the ear transduces this mechanical stimulus (pressure) into a nerve impulse (electrical signal) that the brain perceives as sound.

Learning Objectives

- Describe the transduction of sound and the relevant ear anatomy

Key Points

- The human ear has three distinct functional regions: the outer ear, which collects sound waves; the middle ear, which represents the sound waves as pressure, and the inner ear, which converts those pressure signals into electrical signals that the brain perceives as sound.
- The outer ear involves the pinna (the external shell-shaped structure on the outside of the head), which assists in collecting sound waves; the meatus (the external canal); and the tympanic membrane, also known as the eardrum.
- The middle ear exists between the eardrum and the oval window (the external border with the inner ear) and consists of three separate bones: the malleus, the incus, and the stapes.
- While the middle ear cavity is filled with air, the inner ear is filled with fluid.
- The inner ear exists on the other side of the oval window from the middle ear, by the temple of the human head, and consists of three parts: the semicircular canals, the vestibule, and the cochlea.
- Within the cochlea, the inner hair cells are most important for conveying auditory information to the brain.

Key Terms

- **ossicle**: a small bone (or bony structure), especially one of the three of the middle ear
- **cochlea**: the complex, spirally coiled, tapered cavity of the inner ear in which sound vibrations are converted into nerve impulses
- **transduce**: to convert energy from one form to another

Vibrating objects, such as vocal cords, create sound waves or pressure waves in the air. When these pressure waves reach the ear, the ear transduces this mechanical stimulus (pressure wave) into a nerve impulse (electrical signal) that the brain perceives as sound. The pressure waves strike the tympanum, causing it to vibrate. The mechanical energy from the moving tympanum transmits the vibrations to the three bones of the middle ear. The stapes transmits the vibrations to a thin diaphragm called the oval window, which is the outermost structure of the inner ear.





Diagram of the middle ear: The middle ear exists between the tympanic membrane (the boundary with the outer ear) and the oval window (the boundary with the inner ear) and consists of three bones: the malleus (meaning hammer), the incus (meaning anvil), and the stapes (meaning stirrup).

The structures of the inner ear are found in the labyrinth, a bony, hollow structure that is the most interior portion of the ear. Here, the energy from the sound wave is transferred from the stapes through the flexible oval window and to the fluid of the cochlea. The vibrations of the oval window create pressure waves in the fluid (perilymph) inside the cochlea. The cochlea is a whorled structure, like the shell of a snail, and it contains receptors for transduction of the mechanical wave into an electrical signal. Inside the cochlea, the basilar membrane is a mechanical analyzer that runs the length of the cochlea, curling toward the cochlea's center.



Inner ear: The inner ear can be divided into three parts: the semicircular canals, the vestibule, and the cochlea, all of which are located in the temporal bone.

The mechanical properties of the basilar membrane change along its length, such that it is thicker, tauter, and narrower at the outside of the whorl (where the cochlea is largest), and thinner, floppier, and broader toward the apex, or center, of the whorl (where the cochlea is smallest). Different regions of the basilar membrane vibrate according to the frequency of the sound wave conducted through the fluid in the cochlea. For these reasons, the fluid-filled cochlea detects different wave frequencies (pitches) at different regions of the membrane. When the sound waves in the cochlear fluid contact the basilar membrane, it flexes back and forth in a wave-like fashion. Above the basilar membrane is the tectorial membrane.



Transduction: In the human ear, sound waves cause the stapes to press against the oval window. Vibrations travel up the fluid-filled interior of the cochlea. The basilar membrane that lines the cochlea gets continuously thinner toward the apex of the cochlea. Different thicknesses of membrane vibrate in response to different frequencies of sound. Sound waves then exit through the round window. In the cross section of the cochlea (top right figure), note that in addition to the upper canal and lower canal, the cochlea also has a middle canal. The organ of Corti (bottom image) is the site of sound transduction. Movement of stereocilia on hair cells results in an action potential that travels along the auditory nerve.

The site of transduction is in the organ of Corti (spiral organ). It is composed of hair cells held in place above the basilar membrane like flowers projecting up from soil, with their exposed short, hair-like stereocilia contacting or embedded in the tectorial membrane above them. The inner hair cells are the primary auditory receptors and exist in a single row, numbering approximately 3,500. The stereocilia from inner hair cells extend into small dimples on the tectorial membrane's lower surface. The outer hair cells are arranged in three or four rows. They number approximately 12,000, and they function to fine tune incoming sound waves. The longer stereocilia that project from the outer hair cells actually attach to the tectorial membrane. All of the stereocilia are mechanoreceptors, and when bent by vibrations they respond by opening a gated ion channel (refer to [link]). As a result, the hair cell membrane is depolarized, and a signal is transmitted to the chochlear nerve. Intensity (volume) of sound is determined by how many hair cells at a particular location are stimulated.

The hair cells are arranged on the basilar membrane in an orderly way. The basilar membrane vibrates in different regions, according to the frequency of the sound waves impinging on it. Likewise, the hair cells that lay above it are most sensitive to a specific frequency of sound waves. Hair cells can respond to a small range of similar frequencies, but they require stimulation of greater intensity to fire at frequencies outside of their optimal range. The difference in response frequency between adjacent inner hair cells is about 0.2 percent. Compare that to adjacent piano strings, which are about six percent different. Place theory, which is the model for how biologists think pitch detection works in the human ear, states that high frequency sounds selectively vibrate the basilar membrane of the inner ear near the entrance port (the oval window). Lower frequencies travel farther along the membrane before causing appreciable excitation of the membrane. The basic pitch-determining mechanism is based on the location along the membrane where the hair cells are stimulated. The place theory is the first step toward an understanding of pitch perception. Considering the extreme pitch sensitivity of the human ear, it is thought that there must be some auditory "sharpening" mechanism to enhance the pitch resolution.

When sound waves produce fluid waves inside the cochlea, the basilar membrane flexes, bending the stereocilia that attach to the tectorial membrane. Their bending results in action potentials in the hair cells, and auditory information travels along the neural endings of the bipolar neurons of the hair cells (collectively, the auditory nerve) to the brain. When the hairs bend, they release an excitatory neurotransmitter at a synapse with a sensory neuron, which then conducts action potentials to the central nervous system. The cochlear branch of the vestibulocochlear cranial nerve sends information on hearing. The auditory system is very refined, and there is some modulation or "sharpening" built in. The brain can send signals back to the cochlea, resulting in a change of length in the outer hair cells, sharpening or dampening the hair cells' response to certain frequencies.

Higher Processing

The inner hair cells are most important for conveying auditory information to the brain. About 90 percent of the afferent neurons carry information from inner hair cells, with each hair cell synapsing with 10 or so neurons. Outer hair cells connect to only 10 percent of the afferent neurons, and each afferent neuron innervates many hair cells. The afferent, bipolar neurons that convey auditory information travel from the cochlea to the medulla, through the pons and midbrain in the brainstem, finally reaching the primary auditory cortex in the temporal lobe.

This page titled 13.3D: Transduction of Sound is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14: AUTONOMIC NERVOUS SYSTEM

14.1: Introduction to the Autonomic Nervous System

14.1A: Comparing the Somatic and Autonomic Nervous Systems

14.1B: Divisions of the Autonomic Nervous System

14.2: Structure of the Autonomic Nervous System

14.2A: Preganglionic Neurons

14.2B: Autonomic Ganglia

14.2C: Postganglionic Neurons

14.2D: Autonomic Plexuses

14.2E: Parasympathetic (Craniosacral) Division

14.2F: Autonomic Reflexes

14.2G: Sympathetic Nervous System

14.3: Functions of the Autonomic Nervous System

14.3A: Sympathetic Responses

14.3B: Parasympathetic Responses

14.3C: Autonomic Interactions

14.3D: Control of Autonomic Nervous System Function

14.4: Neurotransmitters and Receptors

14.4A: Cholinergic Neurons and Receptors

14.4B: Adrenergic Neurons and Receptors

14.4C: Agonists, Antagonists, and Drugs

This page titled 14: Autonomic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.1: INTRODUCTION TO THE AUTONOMIC NERVOUS SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

14.1A: Comparing the Somatic and Autonomic Nervous Systems

14.1B: Divisions of the Autonomic Nervous System

This page titled 14.1: Introduction to the Autonomic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.1A: Comparing the Somatic and Autonomic Nervous Systems

The peripheral nervous system includes both a voluntary, somatic branch and an involuntary branch that regulates visceral functions.

Learning Objectives

- Identify the differences between the somatic and autonomic nervous systems

Key Points

- The somatic nervous system (SoNS) is the part of the peripheral nervous system associated with the voluntary control of body movements through the skeletal muscles and mediation of involuntary reflex arcs.
- The autonomic nervous system (ANS) is the part of the peripheral nervous system that controls visceral functions that occur below the level of consciousness.
- The ANS can be subdivided into the parasympathetic nervous system (PSNS) and the sympathetic nervous system (SNS).

Key Terms


- **peripheral nervous system**: Consists of the nerves and ganglia outside of the brain and spinal cord.
- **autonomic**: Acting or occurring involuntarily, without conscious control.
- **somatic nervous system**: The part of the peripheral nervous system that transmits signals from the central nervous system to skeletal muscles, and from receptors of external stimuli, thereby mediating sight, hearing, and touch.

EXAMPLES

Examples of body processes controlled by the ANS include heart rate, digestion, respiratory rate, salivation, perspiration, pupillary dilation, urination, and sexual arousal.

The peripheral nervous system (PNS) is divided into the somatic nervous system and the autonomic nervous system. The somatic nervous system (SoNS) is the part of the peripheral nervous system associated with the voluntary control of body movements via skeletal muscles.

The SoNS consists of efferent nerves responsible for stimulating muscle contraction, including all the non-sensory neurons connected with skeletal muscles and skin. The somatic nervous system controls all voluntary muscular systems within the body, and also mediates involuntary reflex arcs. The somatic nervous system consists of three parts:



The human nervous system: The major organs and nerves of the human nervous system.

1. Spinal nerves are peripheral nerves that carry motor commands and sensory information into the spinal cord.
2. Cranial nerves are the nerve fibers that carry information into and out of the brain stem. They include information related to smell, vision, eyes, eye muscles, the mouth, taste, ears, the neck, shoulders, and the tongue.
3. Association nerves integrate sensory input and motor output; these nerves number in the thousands.

The autonomic nervous system (ANS) is the part of the peripheral nervous system that acts as a control system, functioning largely below the level of consciousness and controlling visceral functions. The ANS affects heart rate, digestion, respiratory rate, salivation, perspiration, pupillary dilation, micturition (urination), and sexual arousal.

Whereas most of its actions are involuntary, some, such as breathing, work in tandem with the conscious mind. The ANS is classically divided into two subsystems: the parasympathetic nervous system (PSNS) and sympathetic nervous system (SNS).

The enteric nervous system is sometimes considered part of the autonomic nervous system, and sometimes considered an independent system.

This page titled 14.1A: Comparing the Somatic and Autonomic Nervous Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.1B: Divisions of the Autonomic Nervous System

The autonomic nervous system (ANS) contains two subdivisions: the parasympathetic (PSNS) and sympathetic (SNS) nervous systems.

Learning Objectives

- Distinguish between the parasympathetic and sympathetic subsystems of the autonomic nervous system

Key Points

- The enteric nervous system is sometimes considered part of the autonomic nervous system, and sometimes considered an independent system.
- Sympathetic and parasympathetic divisions have complementary roles: the sympathetic division functions in actions requiring quick responses (fight or flight) and the parasympathetic division regulates actions that do not require rapid responsiveness (rest and digest).
- The SNS and PSNS can be seen as constantly modulating vital functions, in usually antagonistic fashion, to achieve homeostasis. This includes both cardiovascular and respiratory functions.


Key Terms

- **autonomic**: Acting or occurring involuntarily, without conscious control.
- **fight or flight**: This theory states that animals react to threats with a general discharge of the sympathetic nervous system, priming the animal for fighting or fleeing.
- **vasoconstriction**: The constriction (narrowing) of a blood vessel.

EXAMPLES

Example functions of the SNS include diverting blood flow away from the gastrointestinal (GI) tract and increasing heart rate. Example functions of the PSNS include dilating the blood vessels that lead to the GI tract and stimulating salivary gland secretion.

The autonomic nervous system (ANS) is classically divided into two subsystems: the parasympathetic nervous system (PSNS) and sympathetic nervous system (SNS). The enteric nervous system is sometimes considered part of the autonomic nervous system, and sometimes considered an independent system.



The subdivisions of the autonomic nervous system: In the autonomic nervous system, preganglionic neurons connect the CNS to the ganglion.

Sympathetic and parasympathetic divisions typically function in opposition to each other. This opposition is often viewed as complementary in nature rather than antagonistic. For an analogy, one may think of the sympathetic division as the accelerator and the parasympathetic division as the brake.

The sympathetic division typically functions in actions requiring quick responses. The parasympathetic division functions with actions that do not require immediate reaction. Many think of sympathetic as fight or flight and parasympathetic as rest and digest or feed and breed.

However, many instances of sympathetic and parasympathetic activity cannot be ascribed to fight or rest situations. For example, standing up from a reclining or sitting position would entail an unsustainable drop in blood pressure if not for a compensatory increase in the arterial sympathetic tonus.

Another example is the constant, second-to-second modulation of heart rate by sympathetic and parasympathetic influences as a function of the respiratory cycles. More generally, these two systems should be seen as permanently modulating vital functions, in usually antagonistic fashion, to achieve homeostasis.

Some functions of the SNS include diverting blood flow away from the gastrointestinal (GI) tract and skin via vasoconstriction, enhancing blood flow to skeletal muscles and the lungs, dilating the bronchioles of the lung to allow for greater oxygen exchange, and increasing heart rate.

The PSNS typically functions in contrast to the SNS by dilating the blood vessels leading to the GI tract, causing constriction of the pupil and contraction of the ciliary muscle to the lens to enable closer vision, and stimulating salivary gland secretion, in keeping with the rest and digest functions.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- somatic nervous system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/somatic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peripheral nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Periphe...nervous_system**.

License: ***CC BY-SA: Attribution-ShareAlike***
- Somatic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Somatic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Somatic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Somatic...l_transmission**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Somatic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Somatic...l_transmission**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autonomic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/autonomic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peripheral nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/periphe...rvous%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonom...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vasoconstriction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vasoconstriction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autonomic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/autonomic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fight or flight. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fight%20or%20flight**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Nervous System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 14.1B: Divisions of the Autonomic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2: STRUCTURE OF THE AUTONOMIC NERVOUS SYSTEM

- Topic hierarchy

14.2A: Preganglionic Neurons

14.2B: Autonomic Ganglia

14.2C: Postganglionic Neurons

14.2D: Autonomic Plexuses

14.2E: Parasympathetic (Craniosacral) Division

14.2F: Autonomic Reflexes

14.2G: Sympathetic Nervous System

This page titled 14.2: Structure of the Autonomic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2A: Preganglionic Neurons

In the autonomic nervous system (ANS), nerve fibers that connect the central nervous system to ganglia are known as preganglionic fibers.

Learning Objectives

- Describe characteristics of preganglionic neurons of the autonomic nervous system

Key Points

- All preganglionic fibers of the ANS are cholinergic —meaning they have acetylcholine as their neurotransmitter, and are myelinated for faster transmission.
- Differences between sympathetic and parasympatheic preganglionic fibers include that sympathetic preganglionic fibers tend to be shorter than parasympathetic fibers and sympathetic fibers tend to form more synapses than parasympathetic fibers.
- The parasympathetic division (craniosacral outflow) consists of cell bodies from one of two locations: the brainstem (cranial nerves III, VII, IX, X) or the sacral spinal cord (S2, S3, S4).
- The sympathetic division (thoracolumbar outflow) consists of cell bodies in the lateral horn of the spinal cord (intermediolateral cell columns) from T1 to L2. These cell bodies are GVE (general visceral efferent) neurons and are the preganglionic neurons.

Key Terms

- **cholinergic**: Pertaining to, activated by, producing, or having the same function as acetylcholine.
- **postsynaptic neuron**: The nerve cell that bears receptors for neurotransmitters released into the synaptic cleft by the presynaptic neuron.
- **preganglionic fiber**: In the autonomic nervous system, fibers from the CNS to the ganglion are known as preganglionic fibers.
- **ganglion**: A cluster of interconnecting nerve cells outside the brain.
- **preganglionic neuron**: The nerve fibers that supply a ganglion.

Preganglionic Neuron Properties

In the autonomic nervous system (ANS), fibers from the central nervous system to the ganglion are known as preganglionic fibers. All preganglionic fibers, whether they are in the sympathetic nervous system (SNS) or in the parasympathetic nervous system (PSNS), are cholinergic—that is, these fibers use acetylcholine as their neurotransmitter—and are myelinated.

The ANS is unique in that it requires a sequential two-neuron efferent pathway; the preganglionic neuron must first cross a synapse onto a postganglionic neuron before innervating the target organ. The preganglionic, or first neuron will begin at the outflow and will cross a synapse at the postganglionic, or second neuron's cell body. The postganglionic neuron will then cross a synapse at the target organ.

Sympathetic preganglionic fibers tend to be shorter than parasympathetic preganglionic fibers because sympathetic ganglia are often closer to the spinal cord while parasympathetic preganglionic fibers tend to project to and synapse with the postganglionic fiber close to the target organ.

Outflow Sites

Properties of the SNS and PSNS preganglionic neurons also differ with respect to the spinal cord exit points. The sympathetic division has thoracolumbar outflow, meaning that the neurons begin at the thoracic and lumbar (T1–L2) portions of the spinal


cord. The parasympathetic division has craniosacral outflow, meaning that the neurons begin at the cranial nerves (CN3, CN7, CN9, CN10) and sacral (S2–S4) spinal cord.

The sympathetic division (thoracolumbar outflow) consists of cell bodies in the lateral horn of the spinal cord (intermediolateral cell columns) from T1 to L2. These cell bodies are GVE (general visceral efferent) neurons and are the preganglionic neurons. There are several locations where preganglionic neurons create synapses with their postganglionic neurons:

- The paravertebral ganglia of the sympathetic chain (these run on either side of the vertebral bodies), cervical ganglia, thoracic ganglia, rostral lumbar ganglia, caudal lumbar ganglia, and pelvic ganglia.
- The prevertebral ganglia celiac ganglion, aorticorenal ganglion, superior mesenteric ganglion, inferior mesenteric ganglion.
- The chromaffin cells of the adrenal medulla. This is the one exception to the two-neuron pathway rule: they create a synapse directly onto the target cell bodies.

The parasympathetic division (craniosacral outflow) consists of cell bodies from one of two locations: the brainstem (cranial nerves III, VII, IX, X) or the sacral spinal cord (S2, S3, S4).

These are the preganglionic neurons that synapse with the postganglionic neurons in these locations:



Parasympathetic ganglia of the head: The parasympathetic division has craniosacral outflow, meaning that the neurons begin at the cranial nerves (CN3, CN7, CN9, CN10) and the sacral (S2–S4) spinal cord. Pre- and post-ganglionic fibers and targets are depicted.

- Parasympathetic ganglia of the head (ciliary (CN III)).
- Submandibular (CN VII).
- Pterygopalatine (CN VII).
- Otic (CN IX)).
- In or near the wall of an organ innervated by the vagus (CN X) or sacral nerves (S2, S3, S4).

Divergence

Another major difference between the two ANS systems is divergence, or the number of postsynaptic fibers a single preganglionic fiber creates a synapse with. Whereas in the parasympathetic division there is a divergence factor of roughly 1:4, in the sympathetic division there can be a divergence of up to 1:20.

The site of synapse formation and this divergence for both the sympathetic and parasympathetic preganglionic neurons does, however, occur within ganglia situated within the peripheral nervous system.

This page titled 14.2A: Preganglionic Neurons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2B: Autonomic Ganglia

Autonomic ganglia are clusters of neuron cell bodies that transmit sensory signals from the periphery to the integration centers in the CNS.

Learning Objectives

- Distinguish the differences between sympathetic and parasympathetic ganglia

Key Points

- Autonomic ganglia can be classified as either sympathetic ganglia and parasympathetic ganglia.
- A dorsal root ganglion (or spinal ganglion) is a nodule on a dorsal root of the spine that contains the cell bodies of nerve cells (neurons) that carry signals from sensory organs to the appropriate integration center.
- Sympathetic ganglia deliver information to the body about stress and impending danger, and are responsible for the familiar fight-or-flight response. They contain approximately 20,000–30,000 nerve cell bodies and are located close to and on either side of the spinal cord in long chains.
- The axons of dorsal root ganglion neurons are known as afferents. In the peripheral nervous system, afferents refer to the axons that relay sensory information into the central nervous system (i.e., the brain and the spinal cord).


Key Terms

- **sympathetic ganglion**: The ganglia of the sympathetic nervous system. They deliver information to the body about stress and impending danger, and are responsible for the familiar fight-or-flight response.
- **dorsal root ganglia**: A dorsal root ganglion (or spinal ganglion) is a nodule on a dorsal root of the spine that contains the cell bodies of nerve cells (neurons) that carry signals from sensory organs toward the appropriate integration center.
- **parasympathetic ganglion**: The autonomic ganglia of the parasympathetic nervous system. Most are small terminal ganglia or intramural ganglia, so named because they lie near or within (respectively) the organs they innervate.


Autonomic ganglia are clusters of neuronal cell bodies and their dendrites. They are essentially a junction between autonomic nerves originating from the central nervous system and autonomic nerves innervating their target organs in the periphery.

The dorsal root ganglia lie along the vertebral column by the spine and develop in the embryo from neural crest cells, not neural tube. Therefore, the spinal ganglia can be regarded as gray matter of the spinal cord that became translocated to the periphery.


The two main categories are: sympathetic ganglia and parasympathetic ganglia. An example of parasympathetic ganglion is the ciliary ganglion, involved in pupil constriction and accommodation. A depiction of all the parasympathetic ganglia in the head and neck is shown in the following illustration.



Ciliary ganglion: The pathways of the ciliary ganglion include sympathetic neurons (red), parasympathetic neurons (green), and sensory neurons (blue).



Parasympathetic ganglia of the head: Parasympathetic ganglia of the head (shown as red circles) help supply all parasympathetic innervation to the head and neck.



Anatomy of an autonomic ganglion: The sympathetic connections of the ciliary and superior cervical ganglia are shown in this digram. The postganglionic fibers travel from the ganglion to the effector organ.

Dorsal Root Ganglia

A dorsal root ganglion (or spinal ganglion) is a nodule on a dorsal root of the spine that contains the cell bodies of nerve cells (neurons) that carry signals from the sensory organs towards the appropriate integration center.

Nerves that carry signals towards the brain are known as afferent nerves. The axons of dorsal root ganglion neurons are known as afferents. In the peripheral nervous system, afferents refer to the axons that relay sensory information into the central nervous system (i.e., the brain and the spinal cord).

These neurons are of the pseudo-unipolar type, meaning they have an axon with two branches that act as a single axon, often referred to as a distal process and a proximal process.

Unlike the majority of neurons found in the central nervous system, an action potential in a dorsal root ganglion neuron may initiate in the distal process in the periphery, bypass the cell body, and continue to propagate along the proximal process until reaching the synaptic terminal in the dorsal horn of the spinal cord.

The distal section of the axon may either be a bare nerve ending or encapsulated by a structure that helps relay specific information to a nerve. The nerve endings of dorsal root ganglion neurons have a variety of sensory receptors that are activated by mechanical, thermal, chemical, and noxious stimuli.

In these sensory neurons, a group of ion channels thought to be responsible for somatosensory transduction have been identified. For example, a Meissner's corpuscle or Pacinian corpuscle may encapsulate the nerve ending, rendering the distal process sensitive to mechanical stimulation, such as stroking or vibration, respectively.

Sympathetic Ganglia

Sympathetic ganglia are the ganglia of the sympathetic nervous system. They deliver information to the body about stress and impending danger, and are responsible for the familiar fight-or-flight response. They contain approximately 20,000–30,000 nerve cell bodies and are located close to and on either side of the spinal cord in long chains.

Sympathetic ganglia are the tissue from which neuroblastoma tumors arise. The bilaterally symmetric sympathetic chain ganglia —also called the paravertebral ganglia —are located just ventral and lateral to the spinal cord. The chain extends from the upper neck down to the coccyx, forming the unpaired coccygeal ganglion.

Preganglionic nerves from the spinal cord create a synapse end at one of the chain ganglia, and the postganglionic fiber extends to an effector, typically a visceral organ in the thoracic cavity. There are usually 21 or 23 pairs of these ganglia: three in the cervical region, 12 in the thoracic region, four in the lumbar region, four in the sacral region and a single, unpaired ganglion lying in front of the coccyx called the ganglion impar.

Neurons of the collateral ganglia, also called the prevertebral ganglia, receive input from the splanchnic nerves and innervate organs of the abdominal and pelvic region. These include the celiac ganglia, superior mesenteric ganglia, and inferior mesenteric ganglia.

Parasympathetic Ganglia

Parasympathetic ganglia are the autonomic ganglia of the parasympathetic nervous system. Most are small terminal ganglia or intramural ganglia, so named because they lie near or within (respectively) the organs they innervate. The exceptions are the four paired parasympathetic ganglia of the head and neck.

Efferent parasympathetic nerve signals are carried from the central nervous system to their targets by a system of two neurons. The first neuron in this pathway is referred to as the preganglionic or presynaptic neuron. Its cell body sits in the central nervous system and its axon usually extends to a ganglion somewhere else in the body, where it synapses with the dendrites of the second neuron in the chain.

This second neuron is referred to as the postganglionic or postsynaptic neuron. The axons of presynaptic parasympathetic neurons are usually long. They extend from the CNS into a ganglion that is either very close to or embedded in their target organ. As a result, the postsynaptic parasympathetic nerve fibers are very short.

This page titled 14.2B: Autonomic Ganglia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2C: Postganglionic Neurons

In the autonomic nervous system, fibers from the ganglion to the effector organ are called postganglionic fibers.

Learning Objectives

- Describe sympathetic and parasympathetic postganglionic neurons

Key Points


- Postganglionic fibers in the sympathetic division are adrenergic and use norepinephrine (also called noradrenalin) as a neurotransmitter. By contrast, postganglionic fibers in the parasympathetic division are cholinergic and use acetylcholine as a neurotransmitter.
- In the sympathetic nervous system, the postganglionic neurons of sweat glands release acetylcholine for the activation of muscarinic receptors.
- Chromaffin cells of the adrenal medulla are analogous to post-ganglionic neurons; the adrenal medulla develops in tandem with the sympathetic nervous system and acts as a modified sympathetic ganglion.
- In the sympathetic nervous system, presynaptic nerves ' axons terminate in either the paravertebral ganglia or prevertebral ganglia. In all cases, the axon enters the paravertebral ganglion at the level of its originating spinal nerve.

Key Terms

- **postganglionic fiber**: In the autonomic nervous system, these are the fibers that run from the ganglion to the effector organ.
- **cholinergic**: Pertaining to, activated by, producing, or having the same function as acetylcholine.
- **adrenergic**: Containing or releasing adrenaline.
- **postganglionic neuron**: A nerve cell that is located distal or posterior to a ganglion.

In the autonomic nervous system, fibers from the ganglion to the effector organ are called postganglionic fibers. The post-ganglionic neurons are directly responsible for changes in the activity of the target organ via biochemical modulation and neurotransmitter release.

The neurotransmitters used by postganglionic fibers differ. In the parasympathetic division, they are cholinergic and use acetylcholine as their neurotransmitter. In the sympathetic division, most are adrenergic, meaning they use norepinephrine as their neurotransmitter.



Postganglionic nerve fibers: In the autonomic nervous system, preganglionic fibers (shown in light blue) carry information from the CNS to the ganglion.

The Sympathetic Fibers

At the synapses within the ganglia, the preganglionic neurons release acetylcholine, a neurotransmitter that activates nicotinic acetylcholine receptors on postganglionic neurons. In response to this stimulus, postganglionic neurons—with two important exceptions—release norepinephrine, which activates adrenergic receptors on the peripheral target tissues. The activation of target tissue receptors causes the effects associated with the sympathetic system.

The two exceptions mentioned above are the postganglionic neurons of sweat glands and the chromaffin cells of the adrenal medulla. The postganglionic neurons of sweat glands release acetylcholine for the activation of muscarinic receptors. The chromaffin cells of the adrenal medulla are analogous to post-ganglionic neurons—the adrenal medulla develops in tandem with the sympathetic nervous system and acts as a modified sympathetic ganglion. Within this endocrine gland, the pre-ganglionic neurons create synapses with chromaffin cells and stimulate the chromaffin cells to release norepinephrine and epinephrine directly into the blood.

Presynaptic nerves' axons terminate in either the paravertebral ganglia or prevertebral ganglia. In all cases, the axon enters the paravertebral ganglion at the level of its originating spinal nerve.

After this, it can then either create a synapse in this ganglion, ascend to a more superior ganglion, or descend to a more inferior paravertebral ganglion and make a synapse there, or it can descend to a prevertebral ganglion and create a synapse there with the postsynaptic cell. The postsynaptic cell then goes on to innervate the targeted end effector (i.e., gland, smooth muscle, etc.).

Because paravertebral and prevertebral ganglia are relatively close to the spinal cord, presynaptic neurons are generally much shorter than their postsynaptic counterparts, which must extend throughout the body to reach their destinations.

The Parasympathetic Fibers

The axons of presynaptic parasympathetic neurons are usually long. They extend from the CNS into a ganglion that is either very close to or embedded in their target organ. As a result, the postsynaptic parasympathetic nerve fibers are very short.

In the cranium, preganglionic fibers (cranial nerves III, VII, and IX) usually arise from specific nuclei in the central nervous system (CNS) and create a synapse at one of four parasympathetic ganglia: ciliary, pterygopalatine, otic, or submandibular.

From these four ganglia the postsynaptic fibers complete their journey to target tissues via cranial nerve V (the trigeminal ganglion with its ophthalmic, maxillary, and mandibular branches).

The vagus nerve does not participate in these cranial ganglia, as most of its fibers are destined for a broad array of ganglia on or near the thoracic viscera (esophagus, trachea, heart, lungs) and the abdominal viscera (stomach, pancreas, liver, kidneys). It travels all the way down to the midgut/hindgut junction, which occurs just before the splenic flexure of the transverse colon.

The pelvic splanchnic efferent preganglionic nerve cell bodies reside in the lateral gray horn of the spinal cord at the S2–S4 spinal levels. Their axons continue away from the CNS to synapse at an autonomic ganglion close to the organ of innervation. This differs from the sympathetic nervous system, where synapses between pre- and post-ganglionic efferent nerves in general occur at ganglia that are farther away from the target organ.

The parasympathetic nervous system uses acetylcholine (ACh) as its chief neurotransmitter, although peptides (such as cholecystokinin) may act on the PSNS as a neurotransmitter. The ACh acts on two types of receptors, the muscarinic and nicotinic cholinergic receptors.

Most transmissions occur in two stages: When stimulated, the preganglionic nerve releases ACh at the ganglion, which acts on the nicotinic receptors of the postganglionic neurons. The postganglionic nerve then releases ACh to stimulate the muscarinic receptors of the target organ.

This page titled 14.2C: Postganglionic Neurons is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2D: Autonomic Plexuses

Autonomic plexuses are formed from sympathetic and parasympathetic fibers that innervate and regulate the overall activity of visceral organs.

LEARNING OBJECTIVES

Identify the autonomic plexuses

KEY TAKEAWAYS


Key Points

- The autonomic plexuses include the cardiac plexus, the pulmonary plexus, the esophageal plexus, the abdonimal aortic plexus, and the superior and inferior hypogastric plexuses.
- Autonomic plexuses are formed from sympathetic postganglionic axons, parasympathetic preganglionic axons, and some visceral sensory axons.
- Plexuses provide a complex innervation pattern to the target organs, since most organs are innervated by both divisions of the autonomic nervous system.

Key Terms

- **autonomic plexus**: Any of the extensive networks of nerve fibers and cell bodies associated with the autonomic nervous system that are found in the thorax, abdomen, and pelvis, and that contain sympathetic, parasympathetic, and visceral afferent fibers.
- **abdominal aortic plexus**: This is formed by branches derived, on either side, from the celiac plexus and ganglia, and receives filaments from some of the lumbar ganglia. It is situated upon the sides and front of the aorta, between the origins of the superior and inferior mesenteric arteries.
- **pulmonary plexus**: An autonomic plexus formed from the pulmonary branches of the vagus nerve and the sympathetic trunk. It supplies the bronchial tree and the visceral pleura.

Autonomic plexuses are formed from sympathetic postganglionic axons, parasympathetic preganglionic axons, and some visceral sensory axons. The nerves in the each plexus are close to each other, as in the plexuses of the somatic nervous system, but typically do not interact or synpase together.



Sympathetic trunk: This section of the sympathetic trunk shows both the celiac and the hypogastric plexus.

Instead, they provide a complex innervation pattern to the target organs, since most organs are innervated by both divisions of the autonomic nervous system. The autonomic plexuses include the cardiac plexus, the pulmonary plexus, the esophageal plexus, and abdominal aortic plexus, and the superior and inferior hypogastric plexuses.

Plexuses

Cardiac

The cardiac plexus is a plexus of nerves situated at the base of the heart that innervates the heart. The superficial part of the cardiac plexus lies beneath the arch of the aorta, in front of the right pulmonary artery. It is formed by the superior cardiac branch of the left sympathetic trunk and the lower superior cervical cardiac branch of the left vagus nerve. A small ganglion, the cardiac ganglion of Wrisberg, is occasionally found connected with these nerves at their point of junction.

Pulmonary

The pulmonary plexus is an autonomic plexus formed from pulmonary branches of vagus nerve and the sympathetic trunk. It supplies the bronchial tree and the visceral pleura.

Esophageal

The esophageal plexus is formed by nerve fibers from two sources: the branches of the vagus nerve and the visceral branches of the sympathetic trunk. The esophageal plexus and the cardiac plexus contain the same types of fibers and are both considered thoracic autonomic plexus(es).

Abdominal

The abdominal aortic plexus is formed by branches derived, on either side, from the celiac plexus and ganglia, and receives filaments from some of the lumbar ganglia. It is situated on the sides and front of the aorta, between the origins of the superior and inferior mesenteric arteries.

From this plexus arise parts of the spermatic, the inferior mesenteric, and the hypogastric plexuses; it also distributes filaments to the inferior vena cava.

Superior Hypogastric Plexus

The superior hypogastric plexus (in older texts, hypogastric plexus or presacral nerve) is a plexus of nerves situated on the vertebral bodies below the bifurcation of the abdominal aorta.

Inferior Hypogastric Plexus

The inferior hypogastric plexus (pelvic plexus in some texts) is a plexus of nerves that supplies the viscera of the pelvic cavity. The inferior hypogastric plexus is a paired structure, with each situated on the side of the rectum in the male, and at the sides of the rectum and vagina in the female.

This page titled 14.2D: Autonomic Plexuses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2E: Parasympathetic (Craniosacral) Division

Parasympathetic ganglia are the autonomic ganglia of the parasympathetic nervous system that lie near or within the organs they innervate.

Learning Objectives

- Describe features of the parasympathetic division of the autonomic nervous system

Key Points

- Each PSNS ganglion has three roots: a motor root, a sympathetic root, and a sensory root, as well as a number of exiting branches.
- Most are small terminal ganglia or intramural ganglia, so named because they lie near or within the organs they innervate.
- The parasympathetic system is referred to as having craniosacral outflow because of the location of PSNS fiber origins.

Key Terms


- **lacrimal gland**: One of a pair almond-shaped glands, one for each eye, that secrete the aqueous layer of the tear film.
- **sympathetic root**: This carries postsynaptic sympathetic fibers that traverse the ganglion without crossing a synapse.
- **motor root**: This carries presynaptic parasympathetic nerve fibers that terminate in the ganglion and create synapses for the postsynaptic fibers to travel to their target organs.
- **sensory root**: The proximal end of a dorsal afferent nerve that is attached to the spinal cord.
- **parasympathetic**: Relating to the part of the autonomic nervous system that inhibits or opposes the effects of the sympathetic nervous system.

EXAMPLES

Nerves that supply parasympathetic fibers to the parasympathetic ganglia of the head include the oculomotor nerve (ciliary ganglion); the facial nerve (pterygopalatine ganglion, submandibular ganglion); the glossopharyngeal nerve (otic ganglion); the vagus nerve (no named ganglion); and the pelvic splanchnic nerves (no named ganglion).

Parasympathetic ganglia are the autonomic ganglia of the parasympathetic nervous system, blue fibers). Most are small terminal ganglia or intramural ganglia, so named because they lie near or within (respectively) the organs they innervate.

The exceptions are the four paired parasympathetic ganglia of the head and neck. These paired ganglia supply all parasympathetic innervation to the head and neck: ciliary ganglion (spincter pupillae, ciliary muscle), pterygopalatine ganglion (lacrimal gland, glands of nasal cavity), submandibular ganglion (submandibular and sublingual glands), and otic ganglion (parotid gland).



Nerve innervation of the autonomic nervous system: The parasympathetic nervous system, shown in blue, is a division of the autonomic nervous system.


Each has three roots entering the ganglion (motor, sympathetic, and sensory roots) and a variable number of exiting branches.

1. The motor root carries presynaptic parasympathetic nerve fibers (general visceral efferent fibers) that terminate in the ganglion by creating a synapse for the postsynaptic fibers traveling to target organs.
2. The sympathetic root carries postsynaptic sympathetic fibers (general visceral efferent fibers) that traverse the ganglion without creating a synapse.
3. The sensory root carries general sensory fibers (general somatic afferent fibers) that also do not create a synapse in the ganglion.

Some ganglia also carry special sensory fibers (special visceral afferent) for taste sensation.

The nerves that supply parasympathetic fibers to the parasympathetic ganglia of the head include the oculomotor nerve (ciliary ganglion), the facial nerve (pterygopalatine ganglion, submandibular ganglion), the glossopharyngeal nerve (otic ganglion), the vagus nerve, and the pelvic splanchnic nerves.

Because of its location, the parasympathetic system is commonly referred to as having craniosacral outflow, in contrast to the sympathetic nervous system, which is said to have thoracolumbar outflow.



Parasympathetic ganglia of the head: The parasympathetic division has craniosacral outflow, meaning that the neurons begin at the cranial nerves (CN3, CN7, CN9, CN10) and the sacral spinal cord (S2–S4). Pre- and post-ganglionic fibers and targets are depicted.

This page titled 14.2E: Parasympathetic (Craniosacral) Division is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2F: Autonomic Reflexes

Autonomic reflexes are unconscious motor reflexes relayed from the organs and glands to the CNS through visceral afferent signaling.

Learning Objectives

- Describe autonomic reflexes

Key Points

- While the unconscious reflex arcs are normally undetectable, in certain instances they may trigger pain, typically masked as referred pain.
- The sympathetic nervous system is a quick-response, mobilizing system while the parasympathetic system is a more slowly activated, dampening system—but there are exceptions, such as in sexual arousal and orgasm where both systems play a role.
- Within the brain, the ANS is located in the medulla oblongata in the lower brainstem. The medulla's major ANS functions include respiration, cardiac regulation, vasomotor activity, and certain reflex actions (such as coughing, sneezing, vomiting, and swallowing).

Key Terms

- **reflex arc**: A neural pathway that controls an action reflex. There are two types of reflex arcs: the autonomic reflex arc that affects the inner organs, and the somatic reflex arc that affects muscles.
- **referred pain**: Pain perceived at a location other than the site of the painful stimulus.
- **somatic**: Part of, or relating to, the body of an organism.

EXAMPLES

An example of referred pain from an autonomic reflex arc occurs when the bowel is suddenly distended. In these cases, the body will interpret the afferent pain stimulus as somatic.

The Autonomic Nervous System

The autonomic nervous system (ANS, visceral nervous system, or involuntary nervous system) is the part of the peripheral nervous system that acts as a control system. It functions largely below the level of consciousness, and controls visceral functions.

The ANS affects heart rate, digestion, respiratory rate, salivation, perspiration, pupillary dilation, micturition (urination), and sexual arousal. Most autonomic functions are involuntary but a number of ANS actions can work alongside some degree of conscious control. Everyday examples include breathing, swallowing, and sexual arousal, and in some cases functions such as heart rate.

Functions

Within the brain, the ANS is located in the medulla oblongata in the lower brainstem. The medulla's major ANS functions include respiration (the respiratory control center, or RCC), cardiac regulation (the cardiac control center, or CCC), vasomotor activity (the vasomotor center or VMC), and certain reflex actions (such as coughing, sneezing, vomiting and swallowing).

These then subdivide into other areas and are also linked to ANS subsystems and nervous systems external to the brain. The hypothalamus, just above the brain stem, acts as an integrator for autonomic functions, receiving ANS regulatory input from the limbic system to do so.

Classifications

The ANS is classically divided into two subsystems: the parasympathetic nervous system (PSNS) and sympathetic nervous system (SNS) that operate independently in some functions and interact co-operatively in others. In many cases, the two have opposite actions. When one activates a physiological response, the other inhibits it.


An older simplification of the sympathetic and parasympathetic nervous systems as excitatory and inhibitory was overturned due to the many exceptions found. A more modern characterization is that the sympathetic nervous system is a quick- response, mobilizing system and the parasympathetic is a more slowly activated, dampening system—but there are exceptions, such as in sexual arousal and orgasm where both play a role.

The enteric nervous system is also sometimes considered part of the autonomic nervous system, and sometimes considered an independent system.

The ANS is unique in that it requires a sequential two- neuron efferent pathway; the preganglionic neuron must first creat a synapse to a postganglionic neuron before innervating the target organ. The preganglionic, or first neuron will begin at the outflow and will cross a synapse at the postganglionic, or second neuron's cell body. The postganglionic neuron will then create a synapse at the target organ.

General visceral afferent sensations are mostly unconscious, visceral motor reflex sensations from hollow organs and glands that are transmitted to the CNS (see the following illustration for a depiction of a typical nerve fiber, including general visceral afferent fibers).

While the unconscious reflex arcs are normally undetectable, in certain instances they may send pain sensations to the CNS, masked as referred pain. If the peritoneal cavity becomes inflamed or if the bowel is suddenly distended, the body will interpret the afferent pain stimulus as somatic in origin. This pain is usually non-localized. The pain is usually referred to dermatomes that are at the same spinal nerve level as the visceral afferent synapse.



Typical spinal nerve: This schematic (but not anatomically correct) depiction of a typical spinal nerve indicates (1) somatic efferent fibers, (2) somatic afferent fibers, (3 to 5) sympathetic efferent fibers, and (6 and 7) sympathetic afferent fibers.

This page titled 14.2F: Autonomic Reflexes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.2G: Sympathetic Nervous System

Sympathetic ganglia are the ganglia of the sympathetic nervous system that initiate fight-or-flight, stress-mediated responses.

Learning Objectives


- Describe the features of the sympathetic division of the autonomic nervous system

Key Points

- The bilaterally symmetric sympathetic chain ganglia, also called the paravertebral ganglia, extend from the upper neck down to the coccyx, lateral and slightly ventral to the vertebral column.
- Most sympathetic ganglia are bilaterally symmetric, but an example of an unpaired ganglion, called the ganglion impar is present in front of the coccyx.
- The sympathetic nervous system is said to have thoracolumbar outflow based on the proximity of the ganglia to the corresponding thoracic and lumbar vertebrae.

Key Terms

- **paravertebral ganglia**: Located along the length of the sympathetic trunk, these ganglia are designated as cervical, thoracic, lumbar, and sacral and, except in the neck, closely correspond in number to the vertebrae.
- **fight or flight**: All the coordinated physiological responses that the sympathetic nervous system initiates in response to stress or other emergency situations.
- **sympathetic chain ganglia**: Also called the paravertebral ganglia, these are located just ventral and lateral to the spinal cord. The chain extends from the upper neck down to the coccyx, forming the unpaired coccygeal ganglion.



Nerve innervation of the autonomic nervous system: The parasympathetic nervous system, shown in blue, is a division of the autonomic nervous system.

The sympathetic ganglia are the ganglia of the sympathetic nervous system (the red lines in the diagram below). They deliver information to the body about stress and impending danger, and are responsible for the familiar fight-or-flight response.

This response is also known as the sympathetico-adrenal response because the pre-ganglionic sympathetic fibers that end


in the adrenal medulla—like all sympathetic fibers—secrete acetylcholine. This secretion activates the secretion of adrenaline (epinephrine) and to a lesser extent noradrenaline (norepinephrine) from the adrenal medulla.

Therefore, this response is mediated directly via impulses transmitted through the sympathetic nervous system, and indirectly via catecholamines secreted from the adrenal medulla, and acts primarily on the cardiovascular system.

An example of a sympathetic ganglion in a thoracic nerve is shown in. ganglia contain approximately 20000–30000 nerve cell bodies and are located close to and on either side of the spinal cord in long chains. Sympathetic ganglia are the tissue from which neuroblastoma tumours arise.

The bilaterally symmetric sympathetic chain ganglia, also called the paravertebral ganglia, are located just ventral and lateral to the spinal cord. The chain extends from the upper neck down to the coccyx, forming the unpaired coccygeal ganglion.

Preganglionic nerves from the spinal cord create a synapse at one end of the chain ganglia and the postganglionic fiber extends to an effector, typically a visceral organ, in the thoracic cavity. There are usually 21 or 23 pairs of these ganglia: 3 in the cervical region, 12 in the thoracic region, 4 in the lumbar region, 4 in the sacral region and a single, unpaired ganglion lying in front of the coccyx called the ganglion impar.



Sympathetic ganglion: This intercostal nerve shows the sympathetic ganglion at the top left.

Neurons of the collateral ganglia, also called the prevertebral ganglia, receive input from the splanchnic nerves and innervate organs of the abdominal and pelvic region. These include the celiac ganglia, the superior mesenteric ganglia, and the inferior mesenteric ganglia.

The sympathetic nervous system is said to have thoracolumbar outflow based on its location.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Preganglionic nerve fibers. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Preganglionic_nerve_fibers**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Autonomic_nervous_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- preganglionic fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/preganglionic%20fiber. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cholinergic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cholinergic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ganglion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ganglion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- postsynaptic neuron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/postsynaptic+neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- preganglionic neuron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/preganglionic+neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasympathetic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympath...glia%23Anatomy. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Autonomic_ganglion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasympathetic_ganglia. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_ganglia. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Dorsal root ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dorsal_root_ganglia. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sympathetic ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sympathetic%20ganglion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dorsal root ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dorsal%20root%20ganglia. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parasympathetic ganglion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/parasym...tic%20ganglion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci...n_pathways.png. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: ***Public Domain: No Known Copyright***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasym...nervous_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasym...nervous_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasympathetic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympath...nervous_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- postganglionic fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/postganglionic%20fiber. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...glionic-neuron. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cholinergic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cholinergic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adrenergic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/adrenergic. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci...n_pathways.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: *Public Domain: No Known Copyright*
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: *Public Domain: No Known Copyright*
- Cardiac plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Superior hypogastric plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Superio...gastric_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Abdominal aortic plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Abdominal_aortic_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Esophageal plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Esophageal_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Inferior hypogastric plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Inferio...gastric_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cardiac plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- autonomic plexus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/autonomic+plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- pulmonary plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pulmonary%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- abdominal aortic plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/abdomin...ortic%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci...n_pathways.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: *Public Domain: No Known Copyright*
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: *Public Domain: No Known Copyright*
- Sympathetic trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_trunk. **License**: *Public Domain: No Known Copyright*
- Parasympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasympathetic_ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasym..._head_and_neck. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasym...sical_location. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasym...athetic_fibers. **License**: *CC BY-SA: Attribution-ShareAlike*
- parasympathetic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/parasympathetic. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...n/sensory-root. **License**: *CC BY-SA: Attribution-ShareAlike*
- sympathetic root. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sympathetic%20root. **License**: *CC BY-SA: Attribution-ShareAlike*
- motor root. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/motor%20root. **License**: *CC BY-SA: Attribution-ShareAlike*

- lacrimal gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lacrimal_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci...n_pathways.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: *Public Domain: No Known Copyright*
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: *Public Domain: No Known Copyright*
- Sympathetic trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_trunk. **License**: *Public Domain: No Known Copyright*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pa...ad_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connections of the Parasympathetic Nervous System. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/Fi...ous_System.jpg. **License**: *CC BY: Attribution*
- Sympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Nervous System. **Provided by**: Saylor. **Located at**: http://www.saylor.org/site/wp-conten...ous-System.pdf. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sympathetic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympath...glia%23Anatomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- sympathetic chain ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sympath...hain%20ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- paravertebral ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/paravertebral%20ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- fight or flight. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/fight%20or%20flight. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ciliary_ganglion_pathways.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: *Public Domain: No Known Copyright*
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: *Public Domain: No Known Copyright*
- Sympathetic trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_trunk. **License**: *Public Domain: No Known Copyright*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connections of the Parasympathetic Nervous System. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/Fi...ous_System.jpg. **License**: *CC BY: Attribution*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Gray839. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray839.png. **License**: *Public Domain: No Known Copyright*
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Autonomic_nervous_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Parasympathetic_nervous_system%23Sensation. **License**: *CC BY-SA: Attribution-ShareAlike*
- somatic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somatic. **License**: *CC BY-SA: Attribution-ShareAlike*
- reflex arc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/reflex%20arc. **License**: *CC BY-SA: Attribution-ShareAlike*

- referred pain. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/referred%20pain. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ciliary ganglion pathways. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ciliary_ganglion_pathways.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray840. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray840.png. **License**: *Public Domain: No Known Copyright*
- Postganglionic fibers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Postganglionic_fibers. **License**: *Public Domain: No Known Copyright*
- Sympathetic trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sympathetic_trunk. **License**: *Public Domain: No Known Copyright*
- Parasympathetic head ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parasympathetic_head_ganglia.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Connections of the Parasympathetic Nervous System. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:1503_Connections_of_the_Parasympathetic_Nervous_System.jpg. **License**: *CC BY: Attribution*
- Gray819. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray819.png. **License**: *Public Domain: No Known Copyright*
- Gray839. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray839.png. **License**: *Public Domain: No Known Copyright*
- Gray799. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray799.svg. **License**: *Public Domain: No Known Copyright*

This page titled 14.2G: Sympathetic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.3: FUNCTIONS OF THE AUTONOMIC NERVOUS SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

14.3A: Sympathetic Responses

14.3B: Parasympathetic Responses

14.3C: Autonomic Interactions

14.3D: Control of Autonomic Nervous System Function

This page titled 14.3: Functions of the Autonomic Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.3A: Sympathetic Responses

The sympathetic division of the autonomic nervous system maintains internal organ homeostasis and initiates the stress response.

Learning Objectives

- Describe the sympathetic responses of the autonomic nervous system

Key Points

- The fibers from the sympathetic nervous system (SNS) innervate the tissues in almost every organ system.
- The SNS is best known for mediating the neuronal and hormonal response to stress known as the fight-or-flight response, also known as sympatho-adrenal response.
- The catecholamine hormones adrenaline and noradrenaline are secreted by the adrenal medulla and facilitate physical activity and mobilize the body to respond to threatening environments.
- The primary neurotransmitter of SNS postganglionic fibers is noradrenaline, also called norepinephrine.

Key Terms


- **sympathetic nervous system (SNS)**: One of the three parts of the autonomic nervous system, along with the enteric and parasympathetic systems. Its general action is to mobilize the body's nervous system fight-or-flight response; it is also constantly active at a basal level to maintain homeostasis.
- **sympatho-adrenal response**: Also called the fight-or-flight response, this activates the secretion of adrenaline (epinephrine) and, to a lesser extent, noradrenaline (norepinephrine).
- **stress response**: This halts or slows down various processes, such as sexual responses and digestive systems, to focus on the stressor situation; this usually causes negative effects like constipation, anorexia, difficulty urinating, and difficulty maintaining sexual arousal.

EXAMPLES

Physiological changes induced by the sympathetic nervous system include accelerating the heart rate, widening bronchial passages, decreasing motility of the large intestine, dilating the pupils, and causing perspiration.

Sympathetic Nervous System Physiology

Alongside the other two components of the autonomic nervous system, the sympathetic nervous system aids in the control of most of the body's internal organs. Stress—as in the hyperarousal of the flight-or-fight response—is thought to counteract the parasympathetic system, which generally works to promote maintenance of the body at rest.



Sympathetic nervous system: The sympathetic nervous system extends from the thoracic to lumbar vertebrae and has connections with the thoracic, abdominal aortic, and pelvic plexuses.

The sympathetic nervous system is responsible for regulating many homeostatic mechanisms in living organisms. Fibers from the SNS innervate tissues in almost every organ system and provide physiological regulation over diverse body processes including pupil diameter, gut motility (movement), and urinary output.

The SNS is perhaps best known for mediating the neuronal and hormonal stress response commonly known as the fight-or-flight response, also known as sympatho-adrenal response of the body. This occurs as the preganglionic sympathetic fibers that end in the adrenal medulla secrete acetylcholine, which activates the secretion of adrenaline (epinephrine), and to a lesser extent noradrenaline (norepinephrine).

Therefore, this response is mediated directly via impulses transmitted through the sympathetic nervous system, and also indirectly via catecholamines that are secreted from the adrenal medulla, and acts primarily on the cardiovascular system.

Messages travel through the SNS in a bidirectional flow. Efferent messages can trigger simultaneous changes in different parts of the body.

For example, the sympathetic nervous system can accelerate heart rate, widen bronchial passages, decrease motility of the large intestine, constrict blood vessels, increase peristalsis in the esophagus, cause pupillary dilation, piloerection (goose bumps) and perspiration (sweating), and raise blood pressure.

Afferent messages carry sensations such as heat, cold, or pain. Some evolutionary theorists suggest that the sympathetic nervous system operated in early organisms to maintain survival since the sympathetic nervous system is responsible for priming the body for action. One example of this priming is in the moments before waking, in which sympathetic outflow spontaneously increases in preparation for activity.

The Fight-or-Flight Response

The fight-or-flight response was first described by Walter Bradford Cannon. His theory states that animals react to threats with a general discharge of the sympathetic nervous system, priming the animal for fighting or fleeing. This response was later recognized as the first stage of a general adaptation syndrome that regulates stress responses among vertebrates and other organisms.

Catecholamine hormones, such as adrenaline or noradrenaline, facilitate the immediate physical reactions associated with a preparation for violent muscular action. These include the following:

- Acceleration of heart and lung action.

- Paling or flushing, or alternating between both.
- Inhibition of stomach and upper-intestinal action to the point where digestion slows down or stops.
- General effect on the sphincters of the body.
- Constriction of blood vessels in many parts of the body.
- Liberation of nutrients (particularly fat and glucose) for muscular action.
- Dilation of blood vessels for muscles.
- Inhibition of the lacrimal gland (responsible for tear production) and salivation.
- Dilation of pupil (mydriasis).
- Relaxation of bladder.
- Inhibition of erection.
- Auditory exclusion (loss of hearing).
- Tunnel vision (loss of peripheral vision).
- Disinhibition of spinal reflexes; and shaking.

In prehistoric times, the human fight-or-flight response manifested fight as aggressive, combative behavior and flight as fleeing potentially threatening situations, such as being confronted by a predator.

In current times, these responses persist, but fight-and-flight responses have assumed a wider range of behaviors. For example, the fight response may be manifested in angry, argumentative behavior, and the flight response may be manifested through social withdrawal, substance abuse, and even television viewing.

Males and females tend to deal with stressful situations differently. Males are more likely to respond to an emergency situation with aggression (fight), while females are more likely to flee (flight), turn to others for help, or attempt to defuse the situation (tend and befriend). During stressful times, a mother is especially likely to show protective responses toward her offspring and affiliate with others for shared social responses to threats.

This page titled 14.3A: Sympathetic Responses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.3B: Parasympathetic Responses

The parasympathetic nervous system regulates organ and gland functions during rest and is considered a slowly activated, dampening system.

Learning Objectives

- Describe the parasympathetic responses of the autonomic nervous system


Key Points

- Body functions stimulated by the parasympathetic nervous system (PSNS) include sexual arousal, salivation, lacrimation, urination, digestion, and defecation.
- The PSNS primarily uses acetylcholine as its neurotransmitter.
- Peptides (such as cholecystokinin) may also act on the PSNS as neurotransmitters.

Key Terms

- **acetylcholine**: An organic, polyatomic cation (often abbreviated ACh) that acts as a neurotransmitter in both the peripheral nervous system (PNS) and central nervous system (CNS) in many organisms, including humans.
- **parasympathetic nervous system**: One of the divisions of the autonomic nervous system, based between the brain and the spinal cord, that slows the heart and relaxes muscles.
- **lacrimation**: Shedding tears; crying.

The Parasympathetic Nervous System



Nerve innervation of the autonomic nervous system: The parasympathetic nervous system, shown in blue, is a division of the autonomic nervous system.

The parasympathetic nervous system (PSNS, or occasionally PNS) is one of the two main divisions of the autonomic nervous system (ANS). The autonomic nervous system (ANS, or visceral nervous system, or involuntary nervous system) is the part of the peripheral nervous system that acts as a control system, functioning largely below the level of consciousness and controlling

visceral functions.

The ANS is responsible for regulating the internal organs and glands, which occurs unconsciously. Its roles include stimulation of rest-and-digest activities that occur when the body is at rest, including sexual arousal, salivation, lacrimation (tears), urination, digestion, and defecation.

Its action is described as being complementary to that of one of the other main branches of the ANS, the sympathetic nervous system, which is responsible for stimulating activities associated with the fight-or-flight response.

The sympathetic and parasympathetic divisions typically function in opposition to each other. This natural opposition is better understood as complementary in nature rather than antagonistic.


The sympathetic nervous system can be considered a quick response, mobilizing system; and the parasympathetic system is a more slowly activated, dampening system.

Parasympathetic Nervous System Functions

A useful acronym to summarize the functions of the parasympathetic nervous system is SLUDD (salivation, lacrimation, urination, digestion, and defecation). The parasympathetic nervous system may also be known as the parasympathetic division.

The parasympathetic nervous system uses chiefly acetylcholine (ACh) as its neurotransmitter, although peptides (such as cholecystokinin) may act on the PSNS as neurotransmitters. The ACh acts on two types of receptors, the muscarinic and nicotinic cholinergic receptors.

Most transmission occurs in two stages. When stimulated, the preganglionic nerve releases ACh at the ganglion, which acts on nicotinic receptors of the postganglionic neurons. The postganglionic nerve then releases ACh to stimulate the muscarinic receptors of the target organ.



Nicotinic acetylcholine receptors: Two different subtypes of nicotinic acetylcholine receptors with alpha and beta subunits are shown. The acetylcholine binding sites are indicated by ACh.

This page titled 14.3B: Parasympathetic Responses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.3C: Autonomic Interactions

The sympathetic and parasympathetic autonomic nervous systems cooperatively modulate internal physiology to maintain homeostasis.

Learning Objectives

- Describe the interactions between the sympathetic and parasympathetic divisions of the autonomic nervous system

Key Points

- The sympathetic and parasympathetic divisions typically function in opposition to each other, with one division exciting, triggering, or activating a response that is countered by the alternate system, which serves to relax, decrease, or negatively modulate a process.
- The sympathetic division typically functions in actions requiring quick responses. The parasympathetic division functions with actions that do not require immediate reaction. The sympathetic division initiates the fight-or-flight response and the parasympathetic initiates the rest-and-digest or feed-and-breed responses.
- The sympathetic and parasympathetic nervous systems are important for modulating many vital functions, including respiration and cardiac contractility. For example, the activities of both the sympathetic and parasympathetic systems maintains adequate blood pressure, vagal tone, and heart rate.

Key Terms


- **feed-and-breed**: The parasympathetic nervous system is often colloquially described as the feed-and-breed or rest-and-digest portion of the autonomic nervous system.
- **fight or flight**: All the coordinated physiological responses that the sympathetic nervous system initiates in response to stress or other emergency situations.
- **vital function**: A measure of various physiological states that life depends on, such as recording body temperature, pulse rate (or heart rate), blood pressure, and respiratory rate.

EXAMPLES

Some processes that are modulated by the sympathetic and parasympathetic systems but that are not easily labeled as fight or rest include the maintenance of blood pressure when standing and the maintenance of regular heart rhythms.

Sympathetic and parasympathetic divisions typically function in opposition to each other. However, this opposition is better termed complementary in nature rather than antagonistic. For an analogy, one may think of the sympathetic division as the accelerator and the parasympathetic division as the brake.

The sympathetic division typically functions in actions requiring quick responses. The parasympathetic division functions with actions that do not require immediate reaction. Consider sympathetic as fight or flight and parasympathetic as rest and digest or feed and breed.



The subdivisions of the autonomic nervous system: In the autonomic nervous system, preganglionic neurons connect the CNS to the ganglion.

However, many instances of sympathetic and parasympathetic activity cannot be ascribed to fight or rest situations. For example, standing up from a reclining or sitting position would entail an unsustainable drop in blood pressure if not for a compensatory increase in the arterial sympathetic tonus.

Another example is the constant, second-to-second modulation of heart rate by sympathetic and parasympathetic influences, as a function of the respiratory cycles. More generally, these two systems should be seen as permanently modulating vital functions, usually in an antagonistic fashion, to achieve homeostasis. Some typical actions of the sympathetic and parasympathetic systems are listed below.

The SNS promotes a fight-or-flight response, corresponds with arousal and energy generation, and performs the following functions:

- Inhibits digestion.
- Diverts blood flow away from the gastro-intestinal (GI) tract and skin via vasoconstriction.
- Blood flow to skeletal muscles and the lungs is enhanced (by as much as 1,200% in the case of skeletal muscles).
- Dilates bronchioles of the lung, which allows for greater alveolar oxygen exchange.
- Increases heart rate and the contractility of cardiac cells (myocytes), thereby providing a mechanism for the enhanced blood flow to skeletal muscles.
- Dilates pupils and relaxes the ciliary muscle to the lens, allowing more light to enter the eye and far vision.
- Provides vasodilation for the coronary vessels of the heart.
- Constricts all the intestinal sphincters and the urinary sphincter.
- Inhibits peristalsis.
- Stimulates orgasm.

Conversely, the PSNS promotes a rest-and-digest response, and promotes the following functions:

- Dilates blood vessels leading to the GI tract, increasing blood flow.
- Constricts the bronchiolar diameter when the need for oxygen has diminished.
- Causes constriction of the pupil and contraction of the ciliary muscle to the lens, allowing for closer vision.
- Stimulates salivary gland secretion, and accelerates peristalsis.
- Stimulates sexual arousal.

This page titled 14.3C: Autonomic Interactions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.3D: Control of Autonomic Nervous System Function

The medulla oblongata, in the lower half of the brainstem, is the control center of the autonomic nervous system.

Learning Objectives

- Describe the control of the autonomic nervous system

Key Points


- The medulla contains the cardiac, respiratory, and vasomotor centers.
- The ANS is classically divided into two subdivisions, the sympathetic division and the parasympathetic division.
- As a rule, the SNS functions in actions that require quick responses, while the PSNS is initiated in actions that don't require immediate response.

Key Terms

- **fight or flight**: This theory states that animals react to threats with a general discharge of the sympathetic nervous system, priming the animal for fighting or fleeing.

The autonomic nervous system (ANS) is the part of the peripheral nervous system that controls involuntary functions that are critical for survival. The ANS participates in the regulation of heart rate, digestion, respiratory rate, pupil dilation, and sexual arousal, among other bodily processes.

Within the brain, the ANS is located in the medulla oblongata in the lower brainstem. The medulla's main functions are to control the cardiac, respiratory, and vasomotor centers, to mediate autonomic, involuntary functions, such as breathing, heart rate, and blood pressure, and to regulate reflex actions such as coughing, sneezing, vomiting, and swallowing.



The brain stem with pituitary and pineal glands: The medulla is a subregion of the brainstem and is a major control center for the autonomic nervous system.

The hypothalamus acts to integrate autonomic functions and receives autonomic regulatory feedback from the limbic system to do so. The ANS is classically divided into two subdivisions, the sympathetic division and the parasympathetic division.

The sympathetic division of the ANS is often referred to as the sympathetic nervous system (SNS). The SNS provides

noradrenergic drive to the ANS. It is often referred to as a quick response mobilizing system that initiates the body's fight-or-flight response.

PSNS input to the ANS is responsible for the stimulation of feed-and-breed and rest-and-digest responses, as opposed to the fight-or-flight response initiated by the SNS. The parasympathetic division of the ANS (PSNS) acts to complement and modulate the drive provided by SNS neurotransmission within the ANS.

As a rule, the SNS functions in actions requiring quick responses while the PSNS is initiated in actions that don't require immediate response.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sympath...tem%23Overview**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sympath...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fight-or-flight response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fight-o...ight-or-flight**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sympath...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fight-or-flight response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fight-or-flight_response**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fight-or-flight response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fight-o...e%23Physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sympath...n_transmission**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sympatho-adrenal response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/sympath...nal%20response**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- stress response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/stress%20response**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sympathetic nervous system (SNS). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/sympath...system%20(SNS)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray838. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray838.png**. **License**: ***Public Domain: No Known Copyright***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parasym...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parasym...em%23Receptors**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parasym...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lacrimation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lacrimation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- acetylcholine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acetylcholine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parasympathetic nervous system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/parasy...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray838. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray838.png**. **License**: ***Public Domain: No Known Copyright***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***

- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonom...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- feed-and-breed. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/feed-and-breed**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fight or flight. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fight%20or%20flight**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vital function. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vital%20function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray838. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray838.png**. **License**: ***Public Domain: No Known Copyright***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***
- Sympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sympathetic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autonomic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autonomic_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parasympathetic nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parasym...nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fight or flight. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fight%20or%20flight**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray838. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray838.png**. **License**: ***Public Domain: No Known Copyright***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray839. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray839.png**. **License**: ***Public Domain: No Known Copyright***
- Illu pituitary pineal glands. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...eal_glands.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 14.3D: Control of Autonomic Nervous System Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.4: NEUROTRANSMITTERS AND RECEPTORS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

14.4A: Cholinergic Neurons and Receptors
14.4B: Adrenergic Neurons and Receptors
14.4C: Agonists, Antagonists, and Drugs

This page titled 14.4: Neurotransmitters and Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.4A: Cholinergic Neurons and Receptors

Acetylcholine is a neurotransmitter in the central and peripheral nervous systems that affects plasticity, arousal, and reward.

Learning Objectives

- Identify the cholinergic neurons and receptors in the autonomic system

Key Points

- The neurotransmitter acetylcholine (ACh) is the only neurotransmitter used in the motor division of the somatic nervous system and the principal neurotransmitter at autonomic ganglia.
- In the CNS, the neurons that release and respond to ACh comprise the cholinergic system, which causes anti-excitatory effects.
- ACh plays a role in synaptic plasticity, including learning and short-term memory.
- ACh may bind either muscarinic or nicotinic receptors.
- ACh is synthesized in cholinergic neurons (such as those in the nucleus basalis of Meynert) from choline and acetyl-CoA using an enzyme called choline acetyltransferase.


Key Terms

- **choline acetyltransferase**: Abbreviated as ChAT, this is an enzyme that is synthesized within the body of a neuron. It is then transferred to the nerve terminal via axoplasmic flow. The role of choline acetyltransferase is to join Acetyl-CoA to choline, resulting in the formation of the neurotransmitter acetylcholine.
- **autonomic ganglia**: Clusters of neuronal cell bodies and their dendrites that are a junction between the autonomic nerves originating from the central nervous system and the autonomic nerves innervating their target organs in the periphery.
- **nicotinic receptors**: Also called nAChRs, these are cholinergic receptors that form ligand-gated ion channels in the plasma membranes of certain neurons and on the postsynaptic side of the neuromuscular junction.

Acetylcholine

Acetylcholine (ACh) is an organic, polyatomic ion that acts as a neurotransmitter in both the peripheral nervous system (PNS) and central nervous system (CNS) in many organisms, including humans. Acetylcholine is one of many neurotransmitters in the autonomic nervous system (ANS) and the only neurotransmitter used in the motor division of the somatic nervous system (sensory neurons use glutamate and various peptides at their synapses).

Acetylcholine is also the principal neurotransmitter in all autonomic ganglia. In cardiac tissue, acetylcholine neurotransmission has an inhibitory effect, which lowers heart rate. However, acetylcholine also behaves as an excitatory neurotransmitter at neuromuscular junctions in skeletal muscle.




Acetylcholine: The chemical structure of acetylcholine is depicted.

Acetylcholine was first identified in 1914 by Henry Hallett Dale for its actions on heart tissue. It was confirmed as a neurotransmitter by Otto Loewi, who initially gave it the name Vagusstoff because it was released from the vagus nerve. They jointly received the 1936 Nobel Prize in physiology or medicine for their work. Acetylcholine was also the first neurotransmitter to

be identified.

Functions



Muscarinic acetylcholine receptor M2: This human M2 muscarinic acetylcholine receptor is bound to an antagonist (ACh).

Acetylcholine has functions both in the peripheral nervous system (PNS) and in the central nervous system (CNS) as a neuromodulator. In the peripheral nervous system, acetylcholine activates muscles and is a major neurotransmitter in the autonomic nervous system. In the central nervous system, acetylcholine and its associated neurons form the cholinergic system.

When acetylcholine binds to acetylcholine receptors on skeletal muscle fibers, it opens ligand-gated sodium channels in the cell membrane. Sodium ions then enter the muscle cell, initiating a sequence of steps that finally produce muscle contraction. Although acetylcholine induces contraction of skeletal muscle, it acts via a different type of receptor to inhibit the contraction of cardiac muscle fibers.


In the autonomic nervous system, acetylcholine is released in the following sites: all pre- and post-ganglionic parasympathetic neurons, all pre-ganglionic sympathetic neurons, some post-ganglionic sympathetic fibers, and in the pseudomotor neurons to sweat glands.

In the central nervous system, ACh has a variety of effects as a neuromodulator for plasticity, arousal, and reward. ACh has an important role in the enhancement of sensory perceptions when we wake up and in sustaining attention.

Damage to the cholinergic (acetylcholine-producing) system in the brain has been shown to be plausibly associated with the memory deficits associated with Alzheimer's disease. ACh has also been shown to promote REM sleep.

In the cerebral cortex, tonic ACh inhibits layer 4 neurons, the main targets of thalamocortical inputs while exciting pyramidal cells in layers 2/3 and 5. This filters out weak sensory inputs in layer 4 and amplifies inputs that reach the layers 2/3 and layer 5 excitatory microcircuits.

As a result, these layer-specific effects of ACh might function to improve the signal-to-noise ratio of cortical processing. At the same time, acetylcholine acts through nicotinic receptors to excite certain groups of inhibitory interneurons in the cortex that further dampen cortical activity.



Nicotinic acetylcholine receptors: These schematics describe the heteromeric and homomeric nature of nAChRs. The heteromeric receptors found in the central nervous system are made up of 2 α and 3 β subunits with the binding site at the interface of α and adjacent subunit. Homomeric receptors contain 5 identical subunits and have 5 binding sites located at the interfaces between adjacent subunits.

One well-supported function of ACh in the cortex is an increased responsiveness to sensory stimuli, a form of attention. Phasic increases of ACh during visual, auditory, and somatosensory stimulus presentations have been found to increase the firing rate of neurons in the corresponding primary sensory cortices.

When cholinergic neurons in the basal forebrain are lesioned, animals' ability to detect visual signals was robustly and persistently impaired. In that same study, an animals' ability to correctly reject non-target trials was not impaired, further supporting the interpretation that phasic ACh facilitates responsiveness to stimuli.

ACh has been implicated in reporting expected uncertainty in the environment, based both on the suggested functions listed above and results recorded while subjects perform a behavioral cuing task. Reaction time differences between correctly cued trials and incorrectly cued trials, called the cue validity, was found to vary inversely with ACh levels in primates with pharmacologically and surgically altered levels of ACh. The result was also found in Alzheimer's disease patients and smokers after nicotine (an ACh agonist) consumption.

Creation of ACh

Acetylcholine is synthesized in certain neurons by the enzyme choline acetyltransferase from the compounds choline and acetyl-CoA. Cholinergic neurons are capable of producing ACh.

An example of a central cholinergic area is the nucleus basalis of Meynert in the basal forebrain. The enzyme acetylcholinesterase converts acetylcholine into the inactive metabolites choline and acetate. This enzyme is abundant in the synaptic cleft, and its role in rapidly clearing free acetylcholine from the synapse is essential for proper muscle function.

Certain neurotoxins work by inhibiting acetylcholinesterase, leading to excess acetylcholine at the neuromuscular junction. This results in paralysis of the muscles needed for breathing and stops the beating of the heart.

This page titled 14.4A: Cholinergic Neurons and Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.4B: Adrenergic Neurons and Receptors

Adrenergic receptors are molecules that bind catecholamines. Their activation leads to overall stimulatory and sympathomimetic responses.

Learning Objectives

- Describe adrenergic neurons and receptors in the autonomic nervous system

Key Points


- Adrenergic receptors consist of two main groups, α and β, multiple subgroups (α1, α2, β1, β2, β3), and several subtypes of the α2 subgroup (α2A, α2B, α2C).
- Epinephrine binds both α and β adrenergic receptors to cause vasoconstriction and vasodilation.
- When activated, the α1 receptor triggers smooth muscle contraction in blood vessels in the skin, gastrointestinal tract, kidney, and brain, among other areas.
- When activated, the α2 receptor triggers inhibition of insulin and the induction of glucagon release in the pancreas, contraction of GI tract sphincters, and increased thrombocyte aggregation.
- When activated, the α2 receptor triggers inhibition of insulin and induction of glucagon release in the pancreas, contraction of GI tract sphincters, and increased thrombocyte aggregation.

Key Terms


- **adrenoreceptor**: These are a class of G protein-coupled receptors that are targets of the catecholamines, especially norepinephrine (noradrenaline) and epinephrine (adrenaline). Many cells possess these receptors, and the binding of a catecholamine to the receptor will generally stimulate the sympathetic nervous system.
- **G protein-coupled receptors**: These comprise a large protein family of transmembrane receptors that sense molecules outside the cell and activate inside signal transduction pathways and, ultimately, cellular responses. Any adrenergic effects on cells are generally mediated by G protein-coupled receptors.
- **adrenergic receptor**: Any of several sites in the surface membranes of cells innervated by adrenergic neurons.

The adrenergic receptors (or adrenoceptors) are a class of metabotropic G protein -coupled receptors that are targets of the catecholamines, especially norepinephrine or noradrenaline, and epinephrine (adrenaline). Although dopamine is a catecholamine, its receptors are in a different category.

Many cells possess these receptors, and the binding of an agonist will generally cause a sympathetic (or sympathomimetic) response (e.g., the fight-or-flight response). For instance, the heart rate will increase, pupils will dilate, energy will be mobilized, and blood flow will be diverted from non-essential organs to skeletal muscle.



Adrenaline (epinephrine): The 2D structure of adrenaline (epinephrine) is illustrated.




Noradrenaline (norepinephrine): The 2D structure of noradrenaline (norepinephrine) is illustrated here.

There are two main groups of adrenergic receptors, α and β, with several subtypes. α receptors have the subtypes α1 (a Gq coupled receptor) and α2 (a Gi coupled receptor). Phenylephrine is a selective agonist of the α receptor.

β-receptors have the subtypes β1, β2, and β3. All three are linked to Gs proteins (although β2 also couples to Gi), which in turn are linked to adenylate cyclase. Agonist binding thus causes a rise in the intracellular concentration of the second messenger cAMP. Downstream effectors of cAMP include the cAMP-dependent protein, kinase (PKA), which mediates some of the intracellular events following hormone binding. Isoprenaline is a nonselective agonist.

Adrenaline or noradrenaline are receptor ligands to α1, α2, or β-adrenergic receptors (the pathway is shown in the following diagram).

- α1 couples to Gq, which results in increased intracellular Ca2+ that results in smooth muscle contraction.
- α2, on the other hand, couples to Gi, which causes a decrease of cAMP activity, that results in smooth muscle contraction.
- β receptors couple to Gs, and increases intracellular cAMP activity, resulting in heart muscle contraction, smooth muscle relaxation, and glycogenolysis.



Adrenergic signal transduction: This schematic shows the mechanism of adrenergic receptors. Adrenaline and noradrenaline are ligands to α1, α2, or β-adrenergic receptors. α1-receptors couple to Gq, resulting in increased intracellular Ca2+ and causing smooth muscle contraction. α2 receptors couple to Gi, causing a decrease in cAMP activity and resulting in smooth muscle contraction. β-receptors couple to Gs, increasing intracellular cAMP activity and resulting in heart muscle contraction, smooth muscle relaxation, and glycogenolysis.

Adrenaline (epinephrine) reacts with both α- and β-adrenoceptors, causing vasoconstriction and vasodilation, respectively. Although α receptors are less sensitive to epinephrine, when activated, they override the vasodilation mediated by β-adrenoceptors. The result is that high levels of circulating epinephrine cause vasoconstriction. At lower levels of circulating epinephrine, β-adrenoceptor stimulation dominates, producing an overall vasodilation.

Smooth muscle behavior is variable depending on anatomical location. One important note is the differential effects of increased cAMP in smooth muscle compared to cardiac muscle. Increased cAMP will promote relaxation in smooth muscle, while promoting increased contractility and pulse rate in cardiac muscle.

α-receptors have several functions in common, but also individual effects. Common (or still unspecified) effects include: vasoconstriction of cardiac arteries (coronary artery), vasoconstriction of veins, and decreased motility of smooth muscle in the gastrointestinal tract.

α1-adrenergic receptors are members of the G protein-coupled receptor superfamily. On activation, a heterotrimeric G protein, Gq, activates phospholipase C (PLC).

The PLC cleaves phosphatidylinositol 4,5-bisphosphate (PIP2), which in turn causes an increase in inositol triphosphate (IP3) and diacylglycerol (DAG). The former interacts with calcium channels of the endoplasmic and sarcoplasmic reticulum, thus changing the calcium content in a cell. This triggers all other effects.

Specific actions of the α1-receptor mainly involve smooth muscle contraction. It causes vasoconstriction in many blood vessels, including those of the skin, gastrointestinal system, kidney (renal artery), and brain. Other areas of smooth muscle contraction are:

- Ureter.
- Vas deferens.
- Hair (arrector pili muscles).
- Uterus (when pregnant).
- Urethral sphincter.
- Bronchioles (although minor to the relaxing effect of β2 receptor on bronchioles).
- Blood vessels of ciliary body (stimulation causes mydriasis).

Further effects include glycogenolysis and gluconeogenesis from adipose tissue and the liver, as well as secretion from sweat glands, and Na+ reabsorption from kidney. Antagonists may be used in hypertension.

There are 3 highly homologous subtypes of α2 receptors: α2A, α2B, and α2C.

α2-Receptor Effects

- Inhibition of insulin release in the pancreas.
- Induction of glucagon release from the pancreas.
- Contraction of sphincters of the gastrointestinal tract.
- Negative feedback in the neuronal synapses—presynaptic inhibition of noradrenalin release in CNS.

β1-Receptor Effects

- Increases cardiac output, by raising heart rate (positive chronotropic effect), increasing impulse conduction (positive dromotropic effect), and increasing contraction (positive inotropic effect), thus increasing the volume expelled with each beat (increased ejection fraction).
- Increases renin secretion from the juxtaglomerular cells of the kidney.
- Increases ghrelin secretion from the stomach.

β2-Receptor Effects

- Smooths muscle relaxation, e.g., in bronchi and the GI tract (decreased motility).
- Lipolysis in adipose tissue.
- Anabolism in skeletal muscle.
- Relaxes a non-pregnant uterus.
- Dilates arteries to skeletal muscle.
- Glycogenolysis and gluconeogenesis.
- Stimulates insulin secretion.

- Contracts the sphincters of the GI tract.
- Thickens secretions from the salivary glands.
- Inhibits histamine release from mast cells.
- Increases renin secretion from the kidney.
- Relaxation of bronchioles (salbutamol, a beta-2 agonist, relieves bronchiole constriction).

This page titled 14.4B: Adrenergic Neurons and Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

14.4C: Agonists, Antagonists, and Drugs

Drugs effecting cholinergic neurotransmission may block, hinder, or mimic the action of acetylcholine and alter post-synaptic transmission.

Learning Objectives

- Distinguish between the effects of an agonist versus an antagonist in the autonomic nervous system

Key Points

- Acetylcholine receptor agonists and antagonists have either direct effects on the receptors or act indirectly by affecting the enzyme acetylcholinesterase.
- Agents targeting ACh receptors may target either the nicotinic or muscarinic receptors for ACh.
- Atropine, an antagonist for muscarinic ACh receptors, lowers the parasympathetic activity of muscles and glands in the parasympathetic nervous system.
- Neostigmine is an indirect ACh receptor agonist that inhibits acetylcholinesterase, preventing the breakdown of acetylcholine. It is used in the treatment of myasthenia gravis and to reverse the effects of neuromuscular blockers used for anesthesia.
- Phenylephrine, marketed as a substitute for Sudafed for decongestant purposes, is an α1- adrenergic receptor agonist.
- Beta-blockers, as their name suggests, block the action of epinephrine and norepinephrine on β-adrenergic receptors and are used for the management of cardiac arrhythmias, cardio-protection after a heart attack, and hypertension.

Key Terms

- **acetylcholinesterase**: An enzyme that catalyzes the breakdown of the neurotransmitter acetylcholine.
- **beta-blockers**: Also called beta-adrenergic blocking agents, beta-adrenergic antagonists, beta-adrenoreceptor antagonists, or beta antagonists, these are a class of drugs used for various indications. As beta-adrenergic receptor antagonists, they diminish the effects of epinephrine (adrenaline) and other stress hormones.
- **atropine**: An alkaloid extracted from the plant deadly nightshade (Atropa belladonna) and other sources. It is used as a drug in medicine for its paralytic effects (e.g., in surgery to relax muscles, in dentistry to dry the mouth, in ophthalmology to dilate the pupils), though overdoses are fatal.

Blocking, hindering, or mimicking the action of acetylcholine has many uses in medicine. Drugs that act on the acetylcholine system are either agonists to the receptors that stimulate the system, or antagonists that inhibit it.

Acetylcholine receptor agonists and antagonists can have a direct effect on the receptors or exert their effects indirectly. For example, by affecting the enzyme acetylcholinesterase the receptor ligand is degraded. Agonists increase the level of receptor activation, antagonists reduce it.

Acetylcholine in the ANS

The vagus (parasympathetic) nerves that innervate the heart release acetylcholine (ACh) as their primary neurotransmitter. ACh binds to muscarinic receptors (M2) that are found principally on cells comprising the sinoatrial (SA) and atrioventricular (AV) nodes.


Muscarinic receptors are coupled to the G_i-protein; therefore, vagal activation decreases cAMP. G_i-protein activation also leads to the activation of KACh channels that increase potassium efflux and hyperpolarizes the cells.

Increases in vagal activity to the SA node decreases the firing rate of the pacemaker cells by decreasing the slope of the pacemaker potential and decreasing heart rate. By hyperpolarizing the cells, vagal activation increases the cell's threshold for firing, which contributes to the reduction the firing rate.

Similar electrophysiological effects also occur at the atrioventricular AV node. However, in this tissue, these changes are manifested as a reduction in impulse conduction velocity through the AV node. In the resting state, there is a large degree of vagal tone on the heart, which is responsible for low, resting heart rates.

There is also some vagal innervation of the atrial muscle, and to a much lesser extent, the ventricular muscle. Vagus activation, therefore, results in modest reductions in atrial contractility (inotropy) and even smaller decreases in ventricular contractility.

Muscarinic Antagonists



Atropine: The 2D chemical structure of atropine is illustrated here.

Muscarinic receptor antagonists bind to muscarinic receptors, thereby preventing ACh from binding to and activating the receptor. By blocking the actions of ACh, muscarinic receptor antagonists very effectively block the effects of vagal nerve activity on the heart. By doing so, they increase heart rate and conduction velocity.


Atropine is a naturally occurring tropane alkaloid extracted from deadly nightshade (Atropa belladonna), Jimson weed (Datura stramonium), mandrake (Mandragora officinarum), and other plants of the family Solanaceae. Atropine's pharmacological effects are due to its ability to bind to muscarinic acetylcholine receptors. It is an anti-muscarinic agent.

Working as a nonselective muscarinic acetylcholinergic antagonist, atropine increases firing of the sinoatrial node (SA) and conduction through the atrioventricular node (AV) of the heart, opposes the actions of the vagus nerve, blocks acetylcholine receptor sites, and decreases bronchial secretions. In overdoses, atropine is poisonous.

Nicotinic Agonists

A nicotinic agonist is a drug that mimics, in one way or another, the action of acetylcholine (ACh) at nicotinic acetylcholine receptors (nAChRs). Nicotinic acetylcholine receptors are receptors found in the central nervous system, the peripheral nervous systems, and skeletal muscles.

They are ligand-gated ion channels with binding sites for acetylcholine as well as other agonists. When agonists bind to a receptor it stabilizes the open state of the ion channel allowing an influx of cations.



Nicotinic acetylcholine receptors: NAchR are cholinergic receptors that form ligand-gated ion channels in the plasma membranes of certain neurons and on the postsynaptic side of the neuromuscular junction.

The development of nicotinic acetylcholine receptor agonists began in the early nineties after the discovery of nicotine's positive effects on animal memory. Nicotinic antagonists are mainly used for peripheral muscle paralysis in surgery, the classical agent of this type being tubocurarine, but some centrally acting compounds such as bupropion, mecamylamine, and 18-methoxycoronaridine block nicotinic acetylcholine receptors in the brain and have been proposed for treating drug addiction.

The nicotinic acetylcholine receptor agonists are gaining increasing attention as drug candidates for multiple central nervous system disorders such as Alzheimer's disease, schizophrenia, attention-deficit hyperactivity disorder (ADHD), and nicotine addiction. In 2009 there were at least five drugs on the market that affect the nicotinic acetylcholine receptors.

Most indirect-acting ACh receptor agonists work by inhibiting the enzyme acetylcholinesterase. The resulting accumulation of acetylcholine causes a continuous stimulation of the muscles, glands, and central nervous system.

They are examples of enzyme inhibitors, and increase the action of acetylcholine by delaying degradation; some have been used as nerve agents (sarin and VX nerve gas) or pesticides (organophosphates and the carbamates). In clinical use, they are administered to reverse the action of muscle relaxants, to treat myasthenia gravis, and to treat symptoms of Alzheimer's disease (rivastigmine increases cholinergic activity in the brain).

Beta Receptor Antagonists

Beta blockers (sometimes written as β-blockers) or beta-adrenergic blocking agents, beta-adrenergic antagonists, beta-adrenoreceptor antagonists, or beta antagonists, are a class of drugs used for various indications. They are particularly used for the management of cardiac arrhythmias, cardiac protection after myocardial infarction (heart attack), and hypertension.

As beta-adrenergic receptor antagonists, they diminish the effects of epinephrine (adrenaline) and other stress hormones. Beta blockers block the action of endogenous catecholamines —epinephrine (adrenaline) and norepinephrine (noradrenaline) in particular—on β-adrenergic receptors, part of the sympathetic nervous system that mediates the fight-or-flight response.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Acetylcholine receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acetylcholine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cholinergic. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cholinergic. **License**: *CC BY-SA: Attribution-ShareAlike*
- choline acetyltransferase. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/choline...tyltransferase. **License**: *CC BY-SA: Attribution-ShareAlike*
- nicotinic receptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/nicotinic%20receptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- autonomic ganglia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/autonomic%20ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acetylcholine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine. **License**: *Public Domain: No Known Copyright*
- Muscarinic acetylcholine receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscari...oline_receptor. **License**: *Public Domain: No Known Copyright*
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists. **License**: *CC BY-SA: Attribution-ShareAlike*
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adrenergic_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- G protein-coupled receptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/G%20pro...ed%20receptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- adrenoreceptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adrenoreceptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- adrenergic receptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/adrenergic_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acetylcholine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine. **License**: *Public Domain: No Known Copyright*
- Muscarinic acetylcholine receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscari...oline_receptor. **License**: *Public Domain: No Known Copyright*
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists. **License**: *CC BY-SA: Attribution-ShareAlike*
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adrenergic_receptor. **License**: *Public Domain: No Known Copyright*
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adrenergic_receptor. **License**: *Public Domain: No Known Copyright*
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adrenergic_receptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nicotinic antagonist. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nicotinic_antagonist. **License**: *CC BY-SA: Attribution-ShareAlike*
- Phenylephrine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Phenylephrine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acetylcholine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Beta blockers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Beta_blockers. **License**: *CC BY-SA: Attribution-ShareAlike*
- Atropine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Atropine. **License**: *CC BY-SA: Attribution-ShareAlike*
- acetylcholinesterase. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/acetylcholinesterase. **License**: *CC BY-SA: Attribution-ShareAlike*
- atropine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/atropine. **License**: *CC BY-SA: Attribution-ShareAlike*
- beta-blockers. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/beta-blockers. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acetylcholine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acetylcholine. **License**: *Public Domain: No Known Copyright*

- Muscarinic acetylcholine receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscari...oline_receptor**. **License**: ***Public Domain: No Known Copyright***
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenergic_receptor**. **License**: ***Public Domain: No Known Copyright***
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenergic_receptor**. **License**: ***Public Domain: No Known Copyright***
- Adrenergic receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenergic_receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Drug Discovery and Development: Nicotinic Acetylcholine Receptor Agonists. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Drug_Di...eptor_Agonists**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Atropine. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Atropine**. **License**: ***Public Domain: No Known Copyright***

This page titled 14.4C: Agonists, Antagonists, and Drugs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15: ENDOCRINE SYSTEM



- Topic hierarchy

15.10: The Adrenal (Suprarenal) Glands

15.10A: Overview of the Adrenal Glands

15.10B: Adrenal Cortex

15.10C: Adrenal Medulla

15.11: The Pancreas

15.11A: Overview of Pancreatic Islets

15.11B: Types of Cells in the Pancreas

15.11C: Insulin Secretion and Regulation of Glucagon

15.12: Other Endocrine Tissues and Organs

15.13: Stress

15.1: Overview of the Endocrine System

15.1A: Overview of the Endocrine System

15.1B: Comparing the Nervous and Endocrine Systems

15.2: Hormones

15.2A: Mechanisms of Hormone Action

15.2B: Hormone Receptors

15.2C: Chemistry of Hormones

15.2D: Transport of Hormones

15.3: Mechanisms of Hormone Action

15.3A: Direct Gene Activation and the Second-Messenger System

15.3B: Target Cell Specificity

15.3C: Onset, Duration, and Half-Life of Hormone Activity

15.4: Hormone Interactions

15.4A: Interactions of Hormones at Target Cells

15.5: Hormone Secretion Control

15.5A: Control of Hormone Secretion

15.6: Endocrine Glands

15.6A: Overview of Endocrine Glands

15.7: The Pituitary Gland

15.7A: Overview of the Pituitary Gland

15.7B: Control of the Pituitary Gland by the Hypothalamus

15.7C: The Anterior Pituitary
15.7D: The Posterior Pituitary

15.8: The Thyroid Gland

15.8A: Overview of the Thyroid Gland
15.8B: Formation, Storage, and Release of Thyroid Hormones
15.8C: Action of Thyroid Hormones
15.8D: Control of Thyroid Hormone Release

15.9: The Parathyroid Glands

15.9A: Overview of the Parathyroid Glands
15.9B: Parathyroid Hormone

This page titled 15: Endocrine System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.10: THE ADRENAL (SUPRARENAL) GLANDS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

15.10A: Overview of the Adrenal Glands

15.10B: Adrenal Cortex

15.10C: Adrenal Medulla

This page titled 15.10: The Adrenal (Suprarenal) Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.10A: Overview of the Adrenal Glands

In mammals, the adrenal glands (also known as the suprarenal glands) are endocrine glands that sit atop the kidneys.

Learning Objectives

- Describe the location and function of the adrenal glands

Key Points


- The adrenal glands are are endocrine glands located atop the kidneys.
- They are responsible for releasing three classes of hormones; mineralocorticoids, glucocorticoids, and androgens along with catecholamines.
- Each adrenal gland is composed to two structures: the adrenal medulla and adrenal cortex.

Key Terms

- **adrenal cortex**: The outer portion of the adrenal glands that produces mineralocorticoids, glucocorticoids, and androgens.
- **Adrenal medulla**: The innermost part of the adrenal gland, consisting of cells that secrete adrenaline and noradrenaline.

In mammals, the adrenal glands (also known as the suprarenal glands) are endocrine glands that sit atop the kidneys. They are chiefly responsible for releasing three classes of hormones:

1. Mineralocorticoids (aldosterone)
2. Glucocorticoids (cortisol)
3. Androgens (DHEA)



Adrenal glands: The adrenal glands are triangular-shaped organs on top of the kidneys.

Along with catecholamines (adrenaline), these hormones control a variety of functions including kidney function, metabolism, fight-or-flight response, and sex hormone levels.

In humans, the adrenal glands are found at the level of the 12th thoracic vertebra sitting above and slightly medial to the kidneys, lying within the renal fascia, and separated from the kidneys by a thin layer of connective tissue. In humans, the right adrenal gland is triangular shaped, while the left adrenal gland is semilunar shaped.

Each adrenal gland has two distinct structures, the outer adrenal cortex and the inner medulla—both produce hormones. The cortex mainly produces mineralcorticoids, glucocorticoids, and androgens, while the medulla chiefly produces adrenaline and nor-adrenaline.

This page titled 15.10A: Overview of the Adrenal Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.10B: Adrenal Cortex

The adrenal cortex is devoted to the synthesis of corticosteroid and androgen hormones.

Learning Objectives

- Differentiate among the zones and hormones of the adrenal cortex

Key Points

- Specific cortical cells produce particular hormones, including aldosterone, cortisol, and androgens such as androstenedione.
- The adrenal cortex comprises three zones, or layers: Zona glomerulosa (outer), Zona fasciculata and Zona reticularis.
- The outermost layer, the zona glomerulosa, is the main site for production of mineralocorticoids, mainly aldosterone.
- Zona fasciculata is the layer situated between the glomerulosa and reticularis. This layer is responsible for producing glucocorticoids, such as cortisol.
- Zona reticularis is the inner most cortical layer; the zona reticularis produces androgens, mainly DHEA.

Key Terms

- **adrenal cortex**: The outer portion of the adrenal glands that produces hormones essential to homeostasis.
- **zona glomerulosa**: The outermost layer of the adrenal cortex, responsible for producing mineralocorticoids such as aldosterone.
- **zona fasciculata**: The middle layer of the adrenal cortex, responsible for producing glucocorticoids such as cortisol.
- **zona reticularis**: The inner most layer of the adrenal cortex, responsible for producing androgens such as DHEA.

Zones of the Adrenal Cortex

The cortex is regulated by neuroendocrine hormones secreted by the pituitary gland, which are under the control of the hypothalamus, as well as by the renin- angiotensin system. The adrenal cortex has three zones or layers:

1. Zona glomerulosa (outer)
2. Zona fasciculata
3. Zona reticularis

Zona Glomerulosa


The outermost layer, the zona glomerulosa, is the main site for production of mineralocorticoids, mainly aldosterone, that are largely responsible for the long-term regulation of blood pressure.

Aldosterone exerts its effects on the distal convoluted tubule and collecting duct of the kidney, where it causes increased reabsorption of sodium and increased excretion of both potassium (by principal cells) and hydrogen ions (by intercalated cells of the collecting duct). The major stimulus to produce aldosterone is angiotensin II, as ACTH from the pituitary only produces a transient effect. Angiotensin is stimulated by the juxtaglomerular cells when renal blood pressure drops below 90 mmHg.

Zona Fasciculata

Zona fasciculata is the layer situated between the glomerulosa and reticularis. This layer is responsible for producing glucocorticoids, such as 11-deoxycorticosterone, corticosterone, and cortisol in humans. Cortisol enhances the activity of other hormones including glucagon and catecholamines.

Zona Reticularis



Adrenal Cortex: The three layers of the adrenal cortex are shown, the outermost zona glomerulosa, zona fasciculata and the innermost zona reticularis.

Zona reticularis is the innermost cortical layer, and it produces androgens, mainly dehydroepiandrosterone (DHEA), DHEA sulfate (DHEA-S), and androstenedione (the precursor to testosterone) in humans.

Hormones of the Adrenal Cortex

The different zones of the adrenal cortex produce different hormones.

Mineralocorticoids

These are produced in the zona glomerulosa. The primary mineralocorticoid is aldosterone. Its secretion is regulated by the oligopeptide angiotensin II. Aldosterone is secreted in response to high extracellular potassium levels, low extracellular sodium levels, and low fluid levels and blood volume. Aldosterone secretion affects metabolism in different ways:

- It increases urinary excretion of potassium ions
- It increases interstitial levels of sodium ions
- It increases water retention and blood volume.

Glucocorticoids

These are produced in the zona fasciculata. The primary glucocorticoid released by the adrenal gland in humans is cortisol. Its secretion is regulated by the hormone ACTH from the anterior pituitary gland. Upon binding to its target, cortisol enhances metabolism in several ways:

- It stimulates the release of amino acids from the body
- It stimulates lipolysis, the breakdown of fat
- It stimulates gluconeogenesis, the production of glucose from newly-released amino acids and lipids
- It increases blood glucose levels in response to stress, by inhibiting glucose uptake into muscle and fat cells
- It strengthens cardiac muscle contractions
- It increases water retention
- It has anti-inflammatory and anti-allergic effects.

Androgens

The zona reticularis produces androgens, the most important of which is
DHEA. In general, these hormones do not have an overall effect in the male body, and are converted to more potent androgens such astestosterone and DHT or to estrogens (female sex hormones) in the gonads, acting in this way as a metabolic intermediate.

This page titled 15.10B: Adrenal Cortex is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.10C: Adrenal Medulla

The adrenal cortex is devoted to the synthesis of corticosteroid and androgen hormones.

Learning Objectives

- Describe the adrenal medulla


Key Points

- The adrenal medulla secretes two water-soluble hormones (norepinephrine and epinephrine) that underly the fight-or-flight response.
- To carry out this response, the adrenal medulla receives input from the sympathetic nervous system.

Key Terms

- **adrenal cortex**: The outer portion of the adrenal glands that produces hormones essential to homeostasis.
- **chromaffin**: The cells of the adrenal medulla that secrete adrenaline and noradrenaline in response to nervous stimulation.

The adrenal medulla is the core of the adrenal glands, and is surrounded by the adrenal cortex. The adrenal medulla is responsible for the production of catecholamines, derived from the amino acid tyrosine.



Adrenal gland: The adrenal medulla sits below the three layers of the adrenal cortex and is innervated by nerve fibers.

These water-soluble hormones are the major hormones underlying the fight-or-flight response. The adrenal medulla secretes approximately 20% noradrenaline (norepinephrine) and 80% adrenaline (epinephrine). To carry out this response, the adrenal medulla receives input from the sympathetic nervous system through nerve fibers originating in the thoracic spinal cord from T5–T11.

Chromaffin cells are the neuroendocrine cells found in the medulla; they are modified post-synaptic sympathetic neurons that receive sympathetic input.

When stimulated, chromaffin cells secrete adrenaline and noradrenaline along with enkephalin and enkephalin-containing

peptides into the bloodstream. The secreted adrenaline and noradrenaline play an important role in the fight-or-flight response.

The enkephalins and enkephalin-containing peptides are related to, but also distinct from, the endogenous peptides named endorphins (secreted from the pituitary). All of these peptides bind to opioid receptors and produce analgesic (and other) responses.

Some notable effects of adrenaline and noradrenaline include:

- Increased heart rate and blood pressure.
- Blood vessel constriction in the skin and gastrointestinal tract.
- Smooth muscle dilation.
- Dilating bronchioles and capillaries.
- Increased metabolism.

All of these effects are characteristic of the fight-or-flight response. Receptors for catecholamines are widely distributed throughout the body to allow for a systemic response following secretion.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenal medulla. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal%20medulla**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adrenal cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adrenal_cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland**. **License**: ***Public Domain: No Known Copyright***
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland%23Cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- androgen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/androgen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adrenal cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adrenal_cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland**. **License**: ***Public Domain: No Known Copyright***
- Adrenal cortex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_cortex**. **License**: ***Public Domain: No Known Copyright***
- Chromaffin cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chromaffin_cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pheochromocytoma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pheochromocytoma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenal medulla. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_medulla**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland%23Cortex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Chromaffin cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chromaffin_cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neuroblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neuroblast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neuroendocrine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neuroendocrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adrenal cortex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adrenal_cortex**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland**. **License**: ***Public Domain: No Known Copyright***
- Adrenal cortex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_cortex**. **License**: ***Public Domain: No Known Copyright***
- Adrenal cortex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_cortex**. **License**: ***Public Domain: No Known Copyright***

This page titled 15.10C: Adrenal Medulla is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.11: THE PANCREAS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.11A: Overview of Pancreatic Islets

15.11B: Types of Cells in the Pancreas

15.11C: Insulin Secretion and Regulation of Glucagon

This page titled 15.11: The Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.11A: Overview of Pancreatic Islets

Pancreatic islets, also called the islets of Langerhans, are regions of the pancreas that contain its hormone-producing endocrine cells.

Learning Objectives

- Differentiate among the types of pancreatic islet cells

Key Points


- The pancreatic islets are small islands of cells that produce hormones that regulate blood glucose levels. Hormones produced in the pancreatic islets are secreted directly into the bloodstream by five different types of cells.
- The alpha cells produce glucagon, and make up 15–20% of total islet cells. The beta cells produce insulin and amylin, and make up 65–80% of the total islet cells. The delta cells produce somatostatin, and make up 3–10% of the total islet cells.
- The gamma cells produce pancreatic polypeptide, and make up 3–5% of the total islet cells. The epsilon cells produce ghrelin, and make up less than 1% of the total islet cells.
- The feedback system of the pancreatic islets is paracrine, and is based on the activation and inhibition of the islet cells by the endocrine hormones produced in the islets.

Key Terms

- **endocrine**: Produces internal secretions that are transported around the body by the bloodstream.
- **paracrine**: Describes a hormone or other secretion released from endocrine cells into the surrounding tissue rather than into the bloodstream.
- **exocrine**: Produces external secretions that are released through a duct.

The pancreas serves two functions, endocrine and exocrine. The exocrine function of the pancreas is involved in digestion, and these associated structures are known as the pancreatic acini.


The pancreatic acini are clusters of cells that produce digestive enzymes and secretions and make up the bulk of the pancreas. The endocrine function of the pancreas helps maintain blood glucose levels, and the structures involved are known as the pancreatic islets, or the islets of Langerhans.



Pancreatic Tissue

Pancreatic islets or islets of Langerhans: The islets of Langerhans are the regions of the pancreas that contain its endocrine (hormone-producing) cells.

The pancreatic islets are small islands of cells that produce hormones that regulate blood glucose levels. Hormones produced in the pancreatic islets are secreted directly into the blood flow by five different types of cells.



Pancreatic tissue: The small cells in the middle are beta cells, and the surrounding larger cells are alpha, delta, gamma, and epsilon cells.

The endocrine cell subsets are:

- Alpha cells that produce glucagon and make up 15–20% of total islet cells. Glucagon is a hormone that raises blood glucose levels by stimulating the liver to convert its glycogen into glucose.
- Beta cells that produce insulin and amylin and make up 65–80% of the total islet cells. Insulin lowers blood glucose levels by stimulating cells to take up glucose out of the blood stream. Amylin slows gastric emptying, preventing spikes in blood glucose levels.
- Delta cells that produce somatostatin and make up 3–10% of the total islet cells. Somatostatin is a hormone that suppresses the release of the other hormones made in the pancreas.
- Gamma cells that produce pancreatic polypeptide and make up 3–5% of the total islet cells. Pancreatic polypeptide regulates both the endocrine and exocrine pancreatic secretions.
- Epsilon cells that produce ghrelin and make up less than 1% of the total islet cells. Ghrelin is a protein that stimulates hunger.

The feedback system of the pancreatic islets is paracrine—it is based on the activation and inhibition of the islet cells by the endocrine hormones produced in the islets. Insulin activates beta cells and inhibits alpha cells, while glucagon activates alpha cells, which activates beta cells and delta cells. Somatostatin inhibits the activity of alpha cells and beta cells.

This page titled 15.11A: Overview of Pancreatic Islets is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.11B: Types of Cells in the Pancreas

The islets of Langerhans are the regions of the pancreas that contain many hormone-producing endocrine cells.

Learning Objectives

- Distinguish between the cell types of the pancreas

Key Points

- The pancreas reveals two different types of parenchymal tissue: exocrine acini ducts and the endocrine islets of Langerhans.
- The hormones produced in the islets of Langerhans are insulin, glucagon, somatostatin, pancreatic polypeptide, and ghrelin.
- The pancreatic hormones are secreted by alpha, beta, delta, gamma, and epsilon cells.

Key Terms

- **somatostatin**: A polypeptide hormone, secreted by the pancreas, that inhibits the production of certain other hormones.
- **insulin**: A polypeptide hormone that regulates carbohydrate metabolism.
- **glucagon**: A hormone, produced by the pancreas, that opposes the action of insulin by stimulating the production of sugar.

Pancreatic Cells


The pancreas is a glandular organ that belongs to both the digestive and the endocrine systems of vertebrates. It is an endocrine gland that produces several important hormones, including insulin, glucagon, somatostatin, and pancreatic polypeptide.

It is also a digestive, exocrine organ, that secretes pancreatic juice that contains digestive enzymes to assist with digestion and the absorption of nutrients in the small intestine. These enzymes help to further break down the carbohydrates, proteins, and lipids in the chyme.

Under a microscope, stained sections of the pancreas reveal two different types of parenchymal tissue. The light-stained clusters of cells are called islets of Langerhans, which produce hormones that underlie the endocrine functions of the pancreas.

The dark-stained cells form acini, connected to ducts. Acinar cells belong to the exocrine pancreas and secrete digestive enzymes into the gut via a system of ducts.

Islets of Langerhans



Islets of Langerhans: A porcine islet of Langerhans. On the left is a brightfield image created using hematoxylin stain; nuclei are dark circles and the acinar pancreatic tissue is darker than the islet tissue. The right image is the same section stained by immunofluorescence against insulin, indicating beta cells.

The pancreatic islets are small islands of cells that produce hormones that regulate blood glucose levels. Hormones produced in the pancreatic islets are secreted directly into the blood flow by five different types of cells. The endocrine cell subsets are:

- Alpha cells that produce glucagon, and make up 15–20% of total islet cells. Glucagon is a hormone that raises blood glucose levels by stimulating the liver to convert its glycogen into glucose.
- Beta cells that produce insulin and amylin, and make up 65–80% of the total islet cells. Insulin lowers blood glucose levels by stimulating cells to take up glucose out of the blood stream. Amylin slows gastric emptying, preventing spikes in blood glucose levels.
- Delta cells that produce somatostatin, and make up 3–10% of the total islet cells. Somatostatin is a hormone that suppresses the release of the other hormones made in the pancreas.
- Gamma cells that produce pancreatic polypeptide, and make up 3–5% of the total islet cells. Pancreatic polypeptide regulates both the endocrine and exocrine pancreatic secretions.
- Epsilon cells that produce ghrelin, and make up less than 1% of the total islet cells. Ghrelin is a protein that stimulates hunger.

The islets of Langerhans can influence each other through paracrine and autocrine communication. The paracrine feedback system is based on the following correlations:

- The insulin hormone activates beta cells and inhibits alpha cells.
- The hormone glucagon activates alpha cells which then activate beta cells and delta cells.
- Somatostatin hormone inhibits alpha cells and beta cells.

This page titled 15.11B: Types of Cells in the Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.11C: Insulin Secretion and Regulation of Glucagon

Glucagon is a peptide hormone that works in conjunction with insulin to maintain a stable blood glucose level.

Learning Objectives

- Summarize the relationship between insulin secretion and glucagon regulation in blood glucose homeostasis

Key Points


- Glucagon and insulin are peptide hormones secreted by the pancreas that play a key role in maintaining a stable blood blood glucose level.
- Glucagon is produced by alpha cells in the pancreas and acts to raise blood sugar levels.
- Insulin is produced by beta cells in the pancreas and acts to lower blood sugar levels.

Key Terms

- **insulin**: A polypeptide hormone that regulates carbohydrate metabolism.
- **glycogen**: A polysaccharide that is the main form of carbohydrate storage in animals and also converts to glucose as needed.
- **glucagon**: A hormone, produced by the pancreas, that opposes the action of insulin by stimulating the production of sugar.

Glucagon and insulin are peptide hormones secreted by the pancreas that play a key role in maintaining a stable blood glucose level. The blood glucose level is carefully monitored by cells within the pancreas that respond by secreting key hormones.

Glucagon



Glucagon staining: This is an image from a microscope stained for glucagon.

Glucagon is produced by alpha cells in the pancreas and elevates the concentration of glucose in the blood by promoting gluconeogenesis and glycogenolysis. Glucose is stored in the liver in the form of the polysaccharide glycogen, which is a glucan.

Liver cells have glucagon receptors and when glucagon binds to the liver cells they convert glycogen into individual glucose molecules and release them into the bloodstream—this process is known as glycogenolysis. As these stores become depleted, glucagon then encourages the liver and kidney to synthesize additional glucose by gluconeogenesis. Glucagon also turns off

glycolysis in the liver, causing glycolytic intermediates to be shuttled to gluconeogenesis that can induce lipolysis to produce glucose from fat.

Insulin

Insulin is produced by beta cells in the pancreas and acts to oppose the functions of glucagon. It's main role is to promote the conversion of circulating glucose into glycogen via glycogenesis in the liver and muscle cells.

Insulin also inhibits
gluconeogenesis and promotes the storage of glucose in fat through lipid synthesis and also by inhibiting lipolysis.

In Disease

When control of insulin levels fails, diabetes mellitus can result. As a consequence, insulin is used medically to treat some forms of diabetes mellitus.

Patients with type 1 diabetes depend on external insulin (most commonly injected subcutaneously) for their survival because the hormone is no longer produced internally.

Patients with type 2 diabetes are often insulin resistant and, because of such resistance, they may suffer from a relative insulin deficiency. Some patients with type 2 diabetes may eventually require insulin if other medications fail to control blood glucose levels adequately.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Islets of Langerhans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Islets_of_Langerhans**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pancreatic hormones. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pancreatic_hormones**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- exocrine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/exocrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endocrine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endocrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- paracrine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/paracrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illustration depicting pancreatic islet. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Islets_...aticTissue.png**. **License**: ***CC BY: Attribution***
- Pancreatic tissue. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...sche_Insel.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Islets of Langerhans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Islets_...s%23Cell_types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- somatostatin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/somatostatin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- insulin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/insulin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glucagon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glucagon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illustration depicting pancreatic islet. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Islets_...aticTissue.png**. **License**: ***CC BY: Attribution***
- Pancreatic tissue. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...sche_Insel.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Islets of Langerhans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Islets_of_Langerhans**. **License**: ***Public Domain: No Known Copyright***
- Metabolomics/Hormones/Glucagon. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Metabol...n%23Regulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- insulin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/insulin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycogen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glycogen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glucagon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glucagon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illustration depicting pancreatic islet. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/Islets_...aticTissue.png**. **License**: ***CC BY: Attribution***
- Pancreatic tissue. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/f0/Langerhanssche_Insel.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Islets of Langerhans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Islets_of_Langerhans**. **License**: ***Public Domain: No Known Copyright***
- Glucagon. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Glucagon**. **License**: ***CC BY: Attribution***

This page titled 15.11C: Insulin Secretion and Regulation of Glucagon is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.12: Other Endocrine Tissues and Organs

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.
Topic hierarchy

This page titled 15.12: Other Endocrine Tissues and Organs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.13: Stress

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 15.13: Stress is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.1: OVERVIEW OF THE ENDOCRINE SYSTEM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.1A: Overview of the Endocrine System

15.1B: Comparing the Nervous and Endocrine Systems

This page titled 15.1: Overview of the Endocrine System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.1A: Overview of the Endocrine System

The endocrine system is a system of ductless glands that secrete hormones—chemical messengers that are carried for long distances.

Learning Objectives

- Produce a brief overview of the endocrine system

Key Points

- The endocrine system is a system of ductless glands that secrete hormones directly into the circulatory system to be carried long distances to other target organs that regulate key body and organ functions.
- The major endocrine glands include the pituitary, pineal, ovaries, testes, thyroid, hypothalamus, and adrenal glands.

Key Terms

- **hormone**: A molecule released by a cell or a gland in one part of the body that sends out messages affecting cells in other parts of the organism.
- **endocrine system**: The system of ductless glands that secretes hormones directly into the circulatory system.

The Endocrine System


The endocrine system is a system of ductless glands that secretes hormones directly into the circulatory system to be carried long distances to other target organs regulating key body and organ functions. For example, the pineal gland, located at the base of the brain, secretes the hormone melatonin, responsible for regulating sleep patterns.

Endocrine glands are typically well vascularized and the cells comprising the tissue are typically rich in intracellular vacuoles or granules that store hormones prior to release. Endocrine signaling is typically slow to initiate but is prolonged in response; this provides a counterpoint to the more rapid and short-lived nervous system signals.

The endocrine system is in contrast to the exocrine system, which features ducted glands that secrete substances onto an epithelial surface; for example, a sweat gland. Additionally the endocrine system is differentiated from shorter distance signaling such as autocrine (a cell affecting itself), juxtacrine (a cell affecting it's direct neighbors), and paracrine (a cell affecting other nearby cells) signaling.

Key Endocrine Glands

The major endocrine glands include the pituitary, pineal, ovaries, testes, thyroid, hypothalamus and adrenal glands, additionally other tissues such as the kidney and liver also display secondary adrenal functions.



Endocrine glands of the head and neck: The endocrine systems found in the head and neck include the hypothalamus, pineal, pituitary and thyroid glands.

This page titled 15.1A: Overview of the Endocrine System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.1B: Comparing the Nervous and Endocrine Systems

The nervous system and endocrine system both use chemical messengers to signal cells, but each has a different transmission speed.

Learning Objectives

- Distinguish between the nervous system and the endocrine system

Key Points

- The nervous system can respond quickly to stimuli, through the use of action potentials and neurotransmitters.
- Responses to nervous system stimulation are typically quick but short lived.
- The endocrine system responds to stimulation by secreting hormones into the circulatory system that travel to the target tissue.
- Responses to endocrine system stimulation are typically slow but long lasting.

Key Terms

- **hormone**: A molecule released by a cell or a gland in one part of the body that sends out messages affecting cells in other parts of the organism.
- **neurotransmitters**: Endogenous chemicals that transmit signals from a neuron to a target cell across a synapse.

The body must maintain a constant internal environment, through a process termed homeostasis, while also being able to respond and adapt to external events. The nervous and endocrine systems both work to bring about this adaptation, but their response patterns are different. The nervous system and the endocrine system use chemical messengers to signal cells, but the speed at which these messages are transmitted and the length of their effects differs.

Nervous System

The nervous system responds rapidly to stimuli by sending electrical action potentials along neurons, which in turn transmit these action potentials to their target cells using neurotransmitters, the chemical messenger of the nervous system. The response to stimuli by the nervous system is near instantaneous, although the effects are often short lived. An example is the recoil mechanism of an arm when touching something hot.

Endocrine System

The endocrine system relies on hormones to elicit responses from target cells. These hormones are synthesized in specialized glands at a distance from their target, and travel through the bloodstream or inter-cellular fluid. Upon reaching their target, hormones can induce cellular responses at a protein or genetic level.

This process takes significantly longer than that of the nervous system, as endocrine hormones must first be synthesized, transported to their target cell, and enter or signal the cell. However, although hormones act more slowly than a nervous impulse, their effects are typically longer lasting.

Additionally, the target cells can respond to minute quantities of hormones and are sensitive to subtle changes in hormone concentration. For example, the growth hormones secreted by the pituitary gland are responsible for sustained growth during childhood.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Introduction_To_The_Endocrine_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Introduction_To_The_Endocrine_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Introduction_To_The_Endocrine_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secondary messenger. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/secondary%20messenger**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine Glands in the Head. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Endocrine_central_nervous_en.svg**. **License**:
- Anatomy and Physiology of Animals/Endocrine System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Anatomy_and_Physiology_of_Animals/Endocrine_System%23The_Endocrine_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neurotransmitters. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neurotransmitters**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- action potential. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/action_potential**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine Glands in the Head. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Endocrine_central_nervous_en.svg**. **License**:

This page titled 15.1B: Comparing the Nervous and Endocrine Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.2: HORMONES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.2A: Mechanisms of Hormone Action

15.2B: Hormone Receptors

15.2C: Chemistry of Hormones

15.2D: Transport of Hormones

This page titled 15.2: Hormones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.2A: Mechanisms of Hormone Action

A hormone is a secreted chemical messenger that enables communication between cells and tissues throughout the body.

Learning Objectives

- Summarize the mechanisms of hormone action

Key Points

- Hormones are released into the bloodstream through which they travel to target sites.
- The target cell has receptors specific to a given hormone and will be activated by either a lipid-soluble (permeable to plasma membrane) or water-soluble hormone (binds to a cell-surface receptor).
- Lipid-soluble hormones diffuse through the plasma membrane to enter the target cell and bind to a receptor protein.
- Water-soluble hormones bind to a receptor protein on the plasma membrane of the cell.
- Receptor stimulation results in a change in cell activity, which may send feedback to the original hormone-producing cell.

Key Terms

- **Water-soluble hormone**: A lipophobic hormone that binds to a receptor on, or within, the plasma membrane, to initiate an intracellular signaling cascade.
- **hormone**: A molecule released by a cell or a gland in one part of the body that sends out messages affecting cells in other parts of the organism.
- **Lipid-soluble hormone**: A lipophilic hormone that passes through the plasma membrane of a cell, binds to an intracellular receptor, and changes gene expression.

A hormone is a chemical messenger that enables communication between cells. Hormones are secreted by the glands of the endocrine system and they serve to maintain homeostasis and to regulate numerous other systems and processes, including reproduction and development.

Hormone Signaling


The glands of the endocrine system secrete hormones directly into the extracellular environment. The hormones then diffuse to the bloodstream via capillaries and are transported to the target cells through the circulatory system. This allows hormones to affect tissues and organs far from the site of production or to apply systemic effects to the whole body.

Hormone-producing cells are typically specialized and reside within a particular endocrine gland, such as thryocytes in the thyroid gland. Hormones exit their cell of origin through the process of exocytosis or by other means of membrane transport.

Cellular recipients of a particular hormonal signal may be one of several cell types that reside within a number of different tissues. This is so in the case of insulin, which triggers a diverse range of systemic physiological effects. Different tissue types may also respond differently to the same hormonal signal. As a result, hormonal signaling is elaborate and hard to dissect.

Hormones activate target cells by diffusing through the plasma membrane of the target cells (lipid-soluble hormones) to bind a receptor protein within the cytoplasm of the cell, or by binding a specific receptor protein in the cell membrane of the target cell (water-soluble proteins). In both cases, the hormone complex will activate a chain of molecular events within the cell that will result in the activation of gene expression in the nucleus.


The reaction of the target cells may then be recognized by the original hormone-producing cells, leading to a down-regulation in hormone production. This is an example of a homeostatic negative feedback loop.



Lipid-soluble hormone receptor activation: Nuclear hormone receptors are activated by a lipid-soluble hormone such as estrogen, binding to them inside the cell. Lipid-soluble hormones can cross the plasma membrane.

Steps of Hormonal Signaling

- Biosynthesis of a particular hormone in a particular tissue.
- Storage and secretion of the hormone.
- Transport of the hormone to the target cells, tissues, or organs.
- Recognition of the hormone by an associated cell membrane or an intracellular receptor protein.
- Relay and amplification of the received hormonal signal via a signal transduction process.
- Potential feedback to a hormone-producing cell.



Water-soluble hormone receptor activation: Water-soluble hormones, such as epinephrine, bind to a cell-surface localized receptor, initiating a signaling cascade using intracellular second messengers.

Hormone Classes

Hormones are typically divided into three classes:

1. Peptide: Hormones that are modified amino acids or short (peptide) or long (protein) chains of amino acids. Additionally, they can contain carbohydrate moieties.
2. Lipid: Steroid hormones that contain lipids synthesized from cholesterol and eicosanoids that contain lipids synthesized from the fatty acid chains of phospholipids found in the plasma membrane.

- Monoamine: Hormones derived from aromatic amino acids such as phenylalanine, tyrosine, and tryptophan.

This page titled 15.2A: Mechanisms of Hormone Action is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.2B: Hormone Receptors

Hormones activate a cellular response in the target cell by binding to a specific receptor in the target cell.

Learning Objectives

- Distinguish between the location and function of hydrophilic and lipophilic hormone receptors

Key Points

- For water-soluble proteins, the receptor will be at the plasma membrane of the cell.
- The ligand-bound receptor will trigger a cascade of secondary messengers inside the cell.
- For lipid-soluble hormones, the receptor is typically located within the cytoplasm or nucleus of the cell.
- The binding of the hormone allows the receptor to influence transcription in the nucleus, either alone or in association with other transcription factors.
- The number of hormone molecules is usually the key factor for determining hormone action and it is determined by the concentration of circulating hormones, which in turn is influenced by the rate and level of secretion.
- Another limiting factor for hormone action is the effective concentration of hormone-bound receptor complexes that are formed within the cell. This is determined by the number of hormone/receptor molecules available for complex formation and the binding affinity between the hormone and receptor.

Key Terms

- **secondary messenger**: These are molecules that relay signals from receptors on the cell surface to target molecules inside the cell, in the cytoplasm, or the nucleus.

A hormone receptor is a molecule that binds to a specific hormone. Receptors for peptide hormones tend to be found on the plasma membrane of cells, whereas receptors for lipid-soluble hormones are usually found within the cytoplasm.

Upon hormone binding, the receptor can initiate multiple signaling pathways that ultimately lead to changes in the behavior of the target cells.

The hormone activity within a target cell is dependent on the effective concentration of hormone-receptor complexes that are formed. The number of these complexes is in turn regulated by the number of hormone or receptor molecules available, and the binding affinity between hormone and receptor.

Lipophobic Hormones


Many hormones are composed of polypeptides—such as thyroid -stimulating hormones, follicle-stimulating hormones, luteinizing hormones, and insulin. These molecules are not lipid-soluble and therefore cannot diffuse through cell membranes.

The receptors for these hormones need to be localized to the cells' plasma membranes. Following an interaction with the hormones, a cascade of secondary effects within the cytoplasm of the cell is triggered, often involving the addition or removal of phosphate groups to cytoplasmic proteins, changes in ion channel permeability, or an increase in the concentrations of intracellular molecules that may act as secondary messengers, such as cyclic AMP.

Lipophilic Hormones

Lipophilic hormones—such as steroid or thyroid hormones—are able to pass through the cell and nuclear membrane; therefore receptors for these hormones do not need to be, although they sometimes are, located in the cell membrane.

The majority of lipophilic hormone receptors are transcription factors that are either located in the cytosol and move to the cell nucleus upon activation, or remain in the nucleus waiting for the steroid hormone to enter and activate them.
Upon binding by the hormone the receptor undergoes a conformational change, and the receptor together with the bound hormone influence transcription, either alone or in association with other transcription factors.



Example hormone receptor: The thyroid hormone receptor (TR) heterodimerized to the RXR. In the absence of a ligand, the TR is bound to a corepressor protein. Ligand binding to the TR causes a dissociation of co-repressor and recruitment of co-activator proteins, which in turn recruit additional proteins (such as RNA polymerase) that are responsible for the transcription of downstream DNA into RNA, and eventually into protein that results in a change in cell function.

This page titled 15.2B: Hormone Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.2C: Chemistry of Hormones

There are three classes of hormones: peptide hormones, lipid hormones, and monoamine hormones.

Learning Objectives

- Distinguish between the hydrophilic and lipophilic types of endocrine hormones based on their chemical structures

Key Points

- Peptide hormones are comprised of short (peptides) and long (proteins) chains of amino acids. They are water-soluble but cannot pass through the plasma membrane alone.
- Glyco-protein hormones have a carbohydrate moiety attached to the protein.
- Lipid hormones include steroid and eicosanoid hormones. They are lipid-soluble and can pass through the plasma membrane.
- Steroid hormones are derived from the cholesterol and eicosanoid hormones from fatty acids that compose the plasma membrane.
- The third class of hormones is the monoamines that are derived from aromatic amino acids like phenylalanine, tyrosine, and tryptophan.

A hormone is a chemical released by a cell or a gland in one part of the body that sends out messages that affect cells in other parts of the organism.


There are three classes of hormones:

1. Peptide hormones
2. Lipid-derived hormones
3. Monoamine hormones

Peptide Hormones

Peptide hormones consist of short chains of amino acids, such as vasopressin, that are secreted by the pituitary gland and regulate osmotic balance; or long chains, such as insulin, that are secreted by the pancreas, which regulates glucose metabolism.

Some peptide hormones contain carbohydrate side chains and are termed glyco-proteins, such as the follicle-stimulating hormone. All peptide hormones are hydrophilic and are therefore unable to cross the plasma membrane alone.



Peptide hormone: Representation of the molecular structure of a peptide hormone.

Lipid-Derived Hormones

Lipid and phospholipid-derived hormones are produced from lipids such as linoleic acid and arachidonic acid. Steroid hormones, which form the majority of lipid hormones, are derived from carbohydrates; for example, testosterone is produced primarily in the testes and plays a key role in development of the male reproductive system.

Eicosanoids are also lipid hormones that are derived from fatty acids in the plasma membrane. Unlike other hormones, eicosanoids are not stored in the cell—they are synthesized as required. Both are lipophillic and can cross the plasma membrane.

Monoamine Hormones

Monoamine hormones are derived from single aromatic amino acids like phenylalanine, tyrosine, and tryptophan. For example, the tryptophan-derived melatonin that is secreted by the pineal gland regulates sleep patterns.

This page titled 15.2C: Chemistry of Hormones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.2D: Transport of Hormones

Hormones synthesized by the endocrine glands are transported throughout the body by the bloodstream.

Learning Objectives

- Describe the way in which hormones are transported in the endocrine system

Key Points

- Hormones are typically secreted into systemic circulation. However, some are secreted into portal systems that allow for direct hormone targeting.
- Hormones can exist freely in systemic circulation, but the majority are bound with transport proteins.
- Transport proteins hold hormones inactive in systemic circulation and create a reservoir within the circulation that facilitates an even distribution of hormones throughout the tissue or organ.

Key Terms


- **transport protein**: A protein that binds with a hormone in systemic circulation that facilitates its efficient transport.

The endocrine system is a system of ductless glands that secrete hormones directly into the circulatory system to be carried long distances to other target organs that regulate key body and organ functions.

Some endocrine glands secrete into a portal system rather than the systemic circulation that allows for the direct targeting of hormones. For example, hormones secreted by the pancreas pass into the hepatic portal vein that transports them directly to the liver.

Once within the circulatory system a small proportion of hormones circulate freely, however the majority are bound with a transport protein. Mainly produced in the liver, these transport proteins are hormone specific, such as the sex hormone binding globulin that binds with the sex hormones.

When bound with a transport protein hormones are typically inactive, and their release is often triggered in regions of low hormone concentration or can be controlled by other factors. Therefore, transport proteins can act as a reservoir within the circulatory system and help insure an even distribution of hormones within an organ or tissue.



The endocrine system: The major endocrine glands for men and women (male left, female on the right): 1. Pineal gland 2. Pituitary gland 3. Thyroid gland 4. Thymus 5. Adrenal gland 6. Pancreas 7. Ovary 8. Testis

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...ones_and_Types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...ones_and_Types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hormones. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hormones**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/water-soluble-hormone--2**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/lipid-soluble-hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secondary messenger. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/secondary%20messenger**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hormone%23Interactions_with_receptors**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Type ii nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...tor_action.png**. **License**:
- Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hormone...es_of_hormones**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- prostaglandin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/prostaglandin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- testosterone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/testosterone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cholesterol. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cholesterol**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones...cept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Type ii nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...tor_action.png**. **License**:
- PDB 2dez EBI. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...B_2dez_EBI.jpg**. **License**:
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Hormones_and_Types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Hormones_and_Types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...tion/diffusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones/?ref=/concept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- CK-12 Foundation |. **Provided by**: CK-12. **Located at**: **http://www.ck12.org/concept/Hormones/?ref=/concept/Hormones/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Type ii nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Type_ii_nuclear_receptor_action.png**. **License**:
- PDB 2dez EBI. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...B_2dez_EBI.jpg**. **License**:
- Illu endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/File:Il...ine_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 15.2D: Transport of Hormones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.3: MECHANISMS OF HORMONE ACTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

15.3A: Direct Gene Activation and the Second-Messenger System

15.3B: Target Cell Specificity

15.3C: Onset, Duration, and Half-Life of Hormone Activity

This page titled 15.3: Mechanisms of Hormone Action is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.3A: Direct Gene Activation and the Second-Messenger System

Nuclear receptors function as transcription factors because they can bind to DNA and regulate gene expression.

Learning Objectives

- Distinguish between the hormone mechanisms of direct gene activation and the second-messenger system

Key Points

- Receptors that can directly influence gene expression are termed nuclear receptors.
  Type I nuclear receptors (found in cytosol) are modified to translocate to the nucleus upon hormone binding.
- Type II nuclear receptors remain in the nucleus where they often create a complex with co-repressor proteins, which are released upon hormone binding.
- Secondary messengers relay signals from receptors on the cell surface to the target molecules.
- The secondary messenger systems bind hormones to a receptor that causes a cascade of changes that leads to actions.

Key Terms

- **nuclear receptor**: A class of proteins found within cells that are responsible for sensing steroid and thyroid hormones and certain other molecules, as well as to influence gene expression upon activation.
- **secondary messenger**: Molecules that relay signals from receptors on the cell surface to target molecules inside the cell, in the cytoplasm or nucleus.
- **hormone response element**: A short sequence of DNA within the promoter of a gene that is able to bind a specific hormone receptor complex and therefore regulate gene expression.


Hormones can alter cell activity by binding with a receptor. Receptors can either directly influence gene expression and thus cell activity, or induce a secondary signaling cascade that will in turn influence cell activity.

Direct Gene Activation

Receptors that can directly influence gene expression are termed nuclear receptors. Located within the cytosol or nucleus, nuclear receptors are the target of steroid and thyroid hormones that are able to pass through the cell membrane. Nuclear receptors can bind directly to DNA to regulate specific gene expressions and are, therefore, classified as transcription factors.

Nuclear receptors can be classified into two broad classes according to their mechanism of action and their sub-cellular distribution in the absence of ligand. Type I nuclear receptors are located in the cytosol. Upon binding to a hormone the receptor and hormone translocate into the nucleus, and bind to specific sequences of DNA known as hormone response elements (HREs).

Type II receptors are retained in the nucleus. In the absence of ligand, type II nuclear receptors often form a complex with co-repressor proteins. Hormone binding to the nuclear receptor results in dissociation of the co-repressor and the recruitment of co-activator proteins.



Lipid soluble hormones directly regulate gene expression: This figure depicts the mechanism of a class I nuclear receptor (NR) that, in the absence of ligand, is located in the cytosol. Hormone binding to the NR triggers translocation to the nucleus, where the NR binds to a specific sequence of DNA known as a hormone response element (HRE).


Secondary Messengers

For lipophobic hormones that cannot pass the cellular membrane, activity is mediated and amplified within a cell by the action of second messenger mechanisms (molecules that relay signals from receptors on the cell surface to target molecules inside the cell in the cytoplasm or nucleus).

Most hormone receptors are G protein-coupled receptors. Upon hormone binding, the receptor undergoes a conformational change and exposes a binding site for a G-protein. The G-protein is bound to the inner membrane of the cell and consists of three sub-units: alpha, beta, and gamma.

Upon binding to the receptor, it releases a GTP molecule, at which point the alpha sub-unit of the G-protein breaks free from the beta and gamma sub-units and is able to move along the inner membrane until it contacts another membrane-bound protein: the primary effector.

The primary effector then has an action, which creates a signal that can diffuse within the cell. This signal is called the secondary messenger. The secondary messenger may then activate a secondary effector, whose effects depend on the particular secondary messenger system.



Second messenger mechanisms: General schematic of second messenger generation following activation of membrane bound receptors. 1. The agonist activates the membrane-bound receptor. 2. G-protein is activated and produces an effector. 3. The effector stimulates a second messenger synthesis. 4. The second messenger activates an intercellular process.

This page titled 15.3A: Direct Gene Activation and the Second-Messenger System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.3B: Target Cell Specificity

Hormones target a limited number of cells (based on the presence of a specific receptor) as they circulate in the bloodstream.

Learning Objectives

- Describe target cell specificity in the endocrine system

Key Points

- Target cells are cells that are receptive to a secreted hormone.
- Target cell activation is dependent on three factors; the hormone levels in the blood, the receptor levels on the target cell, and hormone–receptor affinity.

Key Terms

- **target cell**: A cell that is receptive to a secreted hormone.

EXAMPLES

An XY fetus will develop along a female pathway if the target cells fail to respond to androgen. This androgen insensitivity occurs when the receptors on the target cells are unable to accept the hormone due to an impairment in receptor shape.

In endocrinology, target cells can refer to the cells where hormones have an effect. Target cells are capable of responding to hormones because they display receptors to which the circulating hormone can bind. In this way, hormones only affect a limited number of cells even though they are transported in the bloodstream throughout the body.

Target cell activation is dependent on three factors:

1. The levels of hormone in the blood.
2. The relative number of hormone receptors on the target cell.
3. The hormone–receptor affinity.

Modulation of these factors can control target cell response. For example, after receptor stimulation the signaling target cell often sends feedback to the hormone-secreting tissue to down-regulate hormone expression.

Additionally, the target cell can up or down-regulate receptor expression to make it more or less sensitive to the same hormone. Finally, hormone–receptor affinity can be altered by the expression of associated inhibitory or co-activating factors.

In some instances, alterations of receptor structure due to a genetic mutation can lead to a reduction in hormone–receptor affinity, as in the case of androgen insensitivity.

This page titled 15.3B: Target Cell Specificity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.3C: Onset, Duration, and Half-Life of Hormone Activity

A hormone's half-life and duration of activity are limited and vary from hormone to hormone.

Learning Objectives

- Distinguish between a hormone's half-life and its duration of activity

Key Points

- The hormone receptors are dynamic structures that vary in number and sensitivity, that depend on the levels of the stimulating hormone.
- The blood levels of hormones reflect a balance between secretion and degradation/excretion by the liver and kidneys.
- The biological half- life of a hormone is the time it takes for the hormone to lose half of its physiological activity.
- The duration of hormone activity refers to the duration of altered cellular behavior triggered by hormone binding.

Key Terms

- **hormone receptor**: A molecule that binds to a specific hormone that triggers alterations in cell activity.
- **half-life**: The time it takes for a substance (drug, radioactive nuclide, or other) to lose half of its pharmacological, physiological, or radiological activity.

EXAMPLES

Vitamin D is a hormone that has a half-life of one to two months. If one obtains vitamin D solely through sun (UVB) exposure during the summer months, serum vitamin D levels will be critically low by late winter. This is one reason why current recommendations are to take vitamin D supplements in order to maintain serum vitamin D levels throughout the year.

The number of hormone molecules available for complex formation is usually the key factor that determines the level at which signal transduction pathways are activated. The number of hormone molecules that are available is determined by the concentration of circulating hormones.

Half-Life

The blood levels of hormones reflect a balance between synthesis/secretion and degradation/excretion. The liver and kidneys are the major organs that degrade hormones with breakdown products excreted in urine and feces.


A hormone's half-life and duration of activity are limited and vary from hormone to hormone. For instance, the biological half-life of luteinizing hormone is 20 minutes, which is shorter than that of a follicle-stimulating hormone (three to four hours), and of human chorionic gonadotropin (24 hours).

A biological half-life or elimination half-life is the time it takes for a substance such as a hormone or drug to lose half of its pharmacologic or physiologic activity. In a medical context, half-life may also describe the time it takes for the blood plasma concentration of a substance to halve (plasma half-life) its steady-state.

The relationship between the biological and plasma half-lives of a substance can be complex, due to factors including their accumulation in tissues, active metabolites, and receptor interactions.

Duration

The duration of hormone activity refers to the duration of events that were stimulated by hormone-receptor binding. While typically relatively short and measured in minutes or hours, certain events, such as the onset of puberty, are much longer lasting.



Hormone levels during menstrual cycle: This image depicts the levels of certain hormones during the menstrual cycle (B), as they correspond to follicular growth and ovulation (A). 1. Follicle-stimulating hormone 2. Estrogen 3. Luteinizing hormone 4. Progesterone.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Second messenger system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Second_messenger_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nuclear receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nuclear_receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secondary messenger. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/secondary%20messenger**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone response element. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hormone%20response%20element**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nuclear receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/nuclear%20receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Second Messenger Mechanism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Second_Messenger_Mechanism.jpg**. **License**: ***Public Domain: No Known Copyright***
- Nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Nuclear_receptor_action.png**. **License**: ***Public Domain: No Known Copyright***
- Receptor (biochemistry). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Receptor_(biochemistry)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Target cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Target_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endocrine action. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/endocrine%20action**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- paracrine action. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/paracrine%20action**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- target cell. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/target_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Second Messenger Mechanism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Second_Messenger_Mechanism.jpg**. **License**: ***Public Domain: No Known Copyright***
- Nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Nuclear_receptor_action.png**. **License**: ***Public Domain: No Known Copyright***
- Half-life. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Half-life%23Half-life_in_biology_and_pharmacology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Endocrine_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Luteinizing hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Luteinizing_hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- half-life. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/half-life**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone receptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hormone%20receptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Second Messenger Mechanism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Second_Messenger_Mechanism.jpg**. **License**: ***Public Domain: No Known Copyright***
- Nuclear receptor action. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Nuclear_receptor_action.png**. **License**: ***Public Domain: No Known Copyright***
- Hormons level - follicle & corpus luteum. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...pus_luteum.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 15.3C: Onset, Duration, and Half-Life of Hormone Activity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.4: HORMONE INTERACTIONS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.4A: Interactions of Hormones at Target Cells

This page titled 15.4: Hormone Interactions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.4A: Interactions of Hormones at Target Cells

Hormones that act to return body conditions to within acceptable limits from opposite extremes are called antagonistic hormones.

Learning Objectives

- Differentiate among the interactions (permissiveness, antagonism, and synergy) of hormones at target cells

Key Points

- Permissiveness is the situation in which a hormone cannot exert its full effects without the presence of another hormone.
- Synergism occurs when two or more hormones produce the same effects in a target cell and their results are amplified.
- Antagonism occurs when a hormone opposes or reverses the effect of another hormone.

Key Terms

- **antagonism**: When a substance binds to the same site an agonist would bind to without causing activation of the receptor.
- **synergism**: Two or more things functioning together to produce a result not independently obtainable.
- **permissiveness**: A certain relationship between hormones and the target cell when the presence of one hormone, at a certain concentration, is required in order to allow a second hormone to fully affect the target cell.

Permissiveness

In biology, permissiveness is a certain relationship between hormones and the target cell. It can be used to describe situations in which the presence of one hormone, at a certain concentration, is required to allow a second hormone to fully affect the target cell.

For example, thyroid hormones increase the number of receptors available for epinephrine at the latter's target cell, thereby increasing epinephrine's effect at that cell. Without the thyroid hormones, epinephrine would have only a weak effect. Another example is cortisol, which exerts a permissive effect on growth hormones.

Antagonism

Maintaining homeostasis often requires conditions to be limited to a narrow range. When conditions exceed the upper limit of homeostasis, a specific action—usually the production of a hormone—is triggered. When conditions return to normal, hormone production is discontinued.

If conditions exceed the lower limits of homeostasis, a different action, usually the production of a second hormone, is triggered. Hormones that act to return body conditions to within acceptable limits from opposite extremes are called antagonistic hormones. The two glands most responsible for homeostasis are the thyroid and the parathyroid.

The regulation of blood glucose concentration (through negative feedback) illustrates how the endocrine system maintains homeostasis by the action of antagonistic hormones. Bundles of cells in the pancreas, called the islets of Langerhans, contain two kinds of cells: alpha cells and beta cells. These cells control blood glucose concentration by producing the antagonistic hormones insulin and glucagon.


Beta cells secrete insulin. When the concentration of blood glucose rises, such as after eating, beta cells secrete insulin into the blood. Insulin stimulates the liver and most other body cells to absorb glucose.

Liver and muscle cells convert glucose to glycogen, for short-term storage, and adipose cells convert glucose to fat. In response, glucose concentration decreases in the blood, and insulin secretion discontinues through negative feedback from the declining

levels of glucose.

Alpha cells secrete glucagon. When the concentration of blood glucose drops, such as during exercise, alpha cells secrete glucagon into the blood. Glucagon stimulates the liver to release glucose.

The glucose in the liver originates from the breakdown of glycogen. Glucagon also stimulates the production of ketone bodies from amino acids and fatty acids. Ketone bodies are an alternative energy source to glucose for some tissues. When blood glucose levels return to normal, glucagon secretion discontinues through negative feedback.



The glucagon receptor structure: Glucagon is a pancreatic peptide hormone that, as a counter-regulatory hormone for insulin, stimulates glucose release by the liver and maintains glucose homeostasis.

Synergy

Synergism occurs when two or more hormones combine to produce effects greater than the sum of their individual effects. For example, testosterone and follicle – stimulating hormones are required for normal sperm production.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Synergism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Synergi...gical_sciences**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...istic_Hormones**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Permissiveness (biology). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Permissiveness_(biology)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- permissiveness. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/permissiveness**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synergism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/synergism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- antagonism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/antagonism**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- MMDB ID 62600 PDB ID 3C59 Crystal Structure Of The Ligand-Bound Glucagon-Like Peptide- 1 Receptor Extracellular Domain. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...lar_Domain.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 15.4A: Interactions of Hormones at Target Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.5: HORMONE SECRETION CONTROL

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.5A: Control of Hormone Secretion

This page titled 15.5: Hormone Secretion Control is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.5A: Control of Hormone Secretion

A hormone is a molecule released by a cell or a gland in one part of the body that exerts effects via specific receptors at other sites.

Learning Objectives

- Describe the manner in which hormone secretion in the endocrine system is controlled

Key Points

- The endocrine system relies on feedback mechanisms to control the hormone levels in the circulatory system.
- Negative feedback systems prevent deviation from an ideal mean to maintain homoeostasis.
- Positive feedback systems facilitate deviation from the mean.

Key Terms

- **negative feedback**: A system that prevents deviation from a mean value.
- **positive feedback**: A system that promotes deviation from a mean value.

The physiological activity of a hormone depends largely on its concentration within the circulatory system. The effects of too high or too low a concentration of hormones can be damaging—this level must be tightly controlled.

The endocrine system relies on feedback systems to regulate hormone production and secretion.

Negative Feedback

Most endocrine glands are under negative feedback control that acts to maintain homoeostasis, i.e., prevent deviation from an ideal value. A key example of a negative feedback system is the regulation of the thyroid hormone thyroxine, which regulates numerous key metabolic processes.

Briefly, neurons in the hypothalamus secrete thyroid-releasing hormone that stimulate cells in the anterior pituitary to secrete thyroid-stimulating hormone. The thyroid-stimulating hormone then stimulates the release of thyroxine from the thyroid gland.

When the blood concentration of thyroxine rises above the ideal value as detected by sensory neurons, the hypothalamus is signaled to stop thyroid-releasing hormone production, which eventually lowers the levels of thyroxine in the blood. When these drop below the ideal value the hypothalamus is signaled to begin secreting thyroid-releasing hormone again.

Positive Feedback

Positive feedback mechanisms control self-perpetuating events, that is, they encourage deviation from the mean. Positive feedback systems are much less common although they do exist. A key example occurs during childbirth.

The hormone oxytocin is produced by the posterior pituitary that stimulates and enhances contractions during labor. During birth, as the baby moves through the birth canal, pressure receptors within the cervix signal the hypothalamus to stimulate the pituitary to secrete oxytocin.

Oxytocin travels to the uterus through the bloodstream, stimulating the muscles in the uterine wall to contract which in turn increases the activation of the pressure receptors and stimulates further release of oxytocin. The strength of muscle contractions intensifies until the baby is born and the stimulation of the pressure receptors is removed, which stops the release of oxytocin.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Signal cascade. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Signal%20cascade**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endocrine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endocrine**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 15.5A: Control of Hormone Secretion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.6: ENDOCRINE GLANDS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.6A: Overview of Endocrine Glands

This page titled 15.6: Endocrine Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.6A: Overview of Endocrine Glands

Endocrine glands are glands that have no duct and release their secretions directly into the intercellular fluid or into the blood.

Learning Objectives

- Differentiate among the types of endocrine glands (pituitary [posterior pituitary, anterior pituitary], thyroid, parathyroid, adrenal, and pancreas) in the endocrine system


Key Points

- The main endocrine glands are the pituitary (anterior and posterior lobes), thyroid, parathyroid, adrenal (cortex and medulla), pancreas, and gonads.
- The parathyroid glands are four masses of tissue, two embedded posteriorly in each lateral mass of the thyroid gland.
- The pancreas lies along the lower curvature of the stomach, close to where it meets the first region of the small intestine, the duodenum.

Key Terms

- **adrenal gland**: This gland is responsible for releasing hormones in response to stress through the synthesis of corticosteroids, such as cortisol and catecholamines (epinephrine (adrenaline) and norepinephrine), as well as the production of androgens.
- **thyroid**: One of the largest endocrine glands, it is responsible for the secretion of thyroxine which controls how quickly the body uses energy, makes proteins, and is sensitive to other hormones.

Endocrine glands are ductless and release their secretions directly into the intercellular fluid or into the blood. A collection of endocrine glands makes up the endocrine system: the pituitary (anterior and posterior lobes), thyroid, parathyroid, adrenal (cortex and medulla), pancreas and gonads.



Endocrine glands in the human head and neck: The endocrine system is the system of glands, each of which secretes different types of hormones directly into the bloodstream (some of which are transported along nerve tracts) to regulate the body.

Pituitary Gland

The hypothalamus makes up the lower region of the diencephalons and lies just above the brain stem. The pituitary gland (hypophysis) is found in the inferior part of the brain, attached to the bottom of the hypothalamus by a slender stalk called the infundibulum.

The pituitary gland consists of two major regions, the anterior pituitary gland (adenohypophysis) and the posterior pituitary gland (neurohypophysis). The hypothalamus also controls the glandular secretion of the pituitary gland.

Posterior Pituitary Gland

Communication between the hypothalamus and the posterior pituitary occurs through neurosecretory cells that span the short distance between them. Hormones produced by the cell bodies of the neurosecretory cells are packaged in vesicles and transported through the axon and stored in the axon terminals that lie in the posterior pituitary.

When the neurosecretory cells are stimulated, the action potential generated triggers the release of the stored hormones from the axon terminals to a capillary network within the posterior pituitary. Two hormones—oxytocin and antidiuretic hormone (ADH)—are produced and released this way.

Anterior Pituitary Gland

The anterior pituitary is involved in sending hormones that control all other hormones of the body. Its lobe is derived from oral ectoderm and is composed of glandular epithelium.

Communication between the hypothalamus and the anterior pituitary occurs through hormones (releasing hormones and inhibiting hormones) that are produced by the hypothalamus and delivered to the anterior pituitary via a portal network of capillaries. The releasing and inhibiting hormones are produced by specialized neurons of the hypothalamus called neurosecretory cells.

These hormones are released into a capillary network that supplies the anterior pituitary. The hormones then diffuse from this secondary plexus into the anterior pituitary, where they initiate the production of specific hormones by the anterior pituitary.

Thyroid Gland

This is one of the largest endocrine glands in the body. It is positioned on the neck just below the larynx and has two lobes, one on either side of the trachea.

It is involved in the production of the hormones T3 (triiodothyronine) and T4 (thyroxine). These hormones increase the metabolic activity of the body‘s cells.

The thyroid also produces and releases the hormone calcitonin (thyrocalcitonin) that contributes to the regulation of blood calcium levels. Thyrocalcitonin or calcitonin decreases the concentration of calcium in the blood, where most of it is stored in the bones.


Parathyroid

There are four parathyroid glands, all located on the thyroid gland. One of its most important functions is to regulate the body’s calcium and phosphorus levels.

All four glands also secrete parathyroid hormone, or PTH, which causes calcium to be released from the bones back into the extracellular fluid. PTH is released directly into the bloodstream and travels to its target cells, which are often quite far away and found in bone, kidneys, and the gastrointestinal system.

Calcitonin, a hormone produced by the thyroid gland that also regulates ECF calcium levels, serves to counteract the calcium-producing effects of PTH.

Adrenal Glands



Adrenal glands: These are responsible for releasing hormones in response to stress through the synthesis of corticosteroids such as cortisol, and catecholamines such as epinephrine (adrenaline) and norepinephrine.

The adrenal glands are a pair of ductless glands located above the kidneys. Through hormonal secretions, they regulate many essential bodily functions including biochemical balances that influence athletic training and general stress response.

The glucocorticoids include corticosterone, cortisone, and hydrocortisone or cortisol. These hormones serve to stimulate the conversion of amino acids into carbohydrates, a process known as gluconeogenesis, and the formation of glycogen by the liver. They also stimulate the formation of reserve glycogen in the tissues, such as in the muscles.

Pancreas

The pancreas is a very important organ in the digestive and the circulatory systems that helps to maintain our blood sugar levels. It is considered to be part of the gastrointestinal system since it produces digestive enzymes. These are released into the small intestine to aid in reducing food particles to basic elements that can be absorbed by the intestine and used by the body.

It has another, very different function in that it forms insulin, glucagon, and other hormones that are sent into the bloodstream to regulate blood sugar levels and other activities throughout the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- thyroid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/thyroid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The endocrine system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_endocrine_system%23Endocrine_Glands**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/adrenal%20gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine central nervous en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/**

File:Endocrine_central_nervous_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Adrenal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adrenal_gland**. **License**: *Public Domain: No Known Copyright*

This page titled 15.6A: Overview of Endocrine Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.7: THE PITUITARY GLAND

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

15.7A: Overview of the Pituitary Gland

15.7B: Control of the Pituitary Gland by the Hypothalamus

15.7C: The Anterior Pituitary

15.7D: The Posterior Pituitary

This page titled 15.7: The Pituitary Gland is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.7A: Overview of the Pituitary Gland

The pituitary gland is connected to the hypothalamus and secretes nine hormones that regulate body homeostasis.

Learning Objectives

- Summarize the structure and function of the pituitary gland

Key Points

- The pituitary gland, or hypophysis, is an endocrine gland about the size of a pea. Although located at the base of the brain and often considered to be part of the brain, the pituitary gland is in fact a separate organ, and is not part of the brain.
- The pituitary gland is divided into two parts, the anterior pituitary and the posterior pituitary. The anterior pituitary receives signalling molecules from the hypothalamus, and in response, synthesizes and secretes seven hormones.
- The posterior pituitary does not produce any hormones of its own; instead, it stores and secretes two hormones made in the hypothalamus.

Key Terms

- **pituitary gland**: An endocrine gland, about the size of a pea, that sits in a small, bony cavity at the base of the brain whose secretions control the other endocrine glands and influence growth, metabolism, and maturation.
- **hypothalamus**: A region of the forebrain located below the thalamus, forming the basal portion of the diencephalon, and functioning to regulate body temperature, some metabolic processes, and governing the autonomic nervous system.

The pituitary gland, or hypophysis, is an endocrine gland about the size of a pea. Although located at the base of the brain and often considered to be part of the brain, the pituitary gland is in fact a separate organ. It protrudes off the bottom of the hypothalamus at the base of the brain, and rests in a small, bony cavity.




Pituitary location: The location of pituitary gland in the human brain.

The pituitary is functionally connected to the hypothalamus by a small tube called the infundibular stem, or, pituitary stalk. The pituitary gland secretes hormones that regulate homeostasis.

The pituitary gland is divided into two parts, the anterior pituitary and the posterior pituitary.

- The anterior pituitary receives signaling molecules from the hypothalamus, and in response, synthesizes and secretes seven important hormones including thyroid-stimulating hormone and growth hormone.
- The posterior pituitary does not produce any hormones of its own, rather, it stores and secretes two hormones made in the hypothalamus— oxytocin and
  anti-diuretic hormone.



The pituitary gland: In this image, the pituitary gland is referred to by its other name, the hypophysis.

This page titled 15.7A: Overview of the Pituitary Gland is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.7B: Control of the Pituitary Gland by the Hypothalamus

The pituitary gland consists of the anterior pituitary and the posterior pituitary.

Learning Objectives

- Explain how the hypothalamus controls the pituitary gland

Key Points

- While the pituitary gland is known as the master endocrine gland, both of its lobes are under the control of the hypothalamus: the anterior pituitary receives its signals from the parvocellular neurons, and the posterior pituitary receives its signals from the magnocellular neurons.
- The pituitary gland is connected by a system of blood vessels to the hypothalamus. This system of blood vessels is known as the hypophyseal portal system, and it allows endocrine communication between the two structures.
- The mechanism for hormone transport via hypothalamoportal vessels involves cells that are regulated by different nuclei in the hypothalamus; for instance, neurons that release neurotransmitters as hormones in the connective link between the pituitary and the brain.

Key Terms


- **pituitary gland**: An endocrine gland, about the size of a pea, that sits in a small, bony cavity at the base of the brain whose secretions control the other endocrine glands and influence growth, metabolism, and maturation.
- **hypothalamus**: A region of the forebrain located below the thalamus, forming the basal portion of the diencephalon, that regulates body temperature, some metabolic processes, and governs the autonomic nervous system.
- **hypophyseal portal system**: The system of blood vessels that link the hypothalamus and the anterior pituitary in the brain.

The pituitary gland consists of two components: the anterior pituitary and the posterior pituitary, and is functionally linked to the hypothalamus by the pituitary stalk (also named the infundibular stem, or simply the infundibulum).

Whilst the pituitary gland is known as the master endocrine gland, both of the lobes are under the control of the hypothalamus: the anterior pituitary receives its signals from the parvocellular neurons, and the posterior pituitary receives its signals from magnocellular neurons.

The anterior lobe of the pituitary receives
hypothalamic-releasing hormones from the hypothalamus that bind with receptors on endocrine cells in the anterior pituitary that regulate the release of adrenal hormones into the circulatory system. Hormones from the hypothalamus are rapidly degraded in the anterior pituitary, which prevents them from entering the circulatory system.

The posterior lobe of the pituitary gland develops as an extension of the hypothalamus. As such, it is not capable of producing its own hormones; instead, it stores hypothalamic hormones for later release into the systemic circulation.



Pituitary gland: The anterior and posterior lobes of the pituitary (hypophysis) gland are shown.

This page titled 15.7B: Control of the Pituitary Gland by the Hypothalamus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.7C: The Anterior Pituitary

The anterior pituitary secretes seven hormones that regulate several physiological processes, including stress, growth, and reproduction.

Learning Objectives

- Identify the location and the hormones produced by the anterior pituitary

Key Points

- A major organ of the endocrine system, the anterior pituitary, also called the adenohypophysis, is the glandular, anterior lobe of the pituitary gland.
- The anterior pituitary regulates several physiological processes, including stress, growth, reproduction, and lactation.
- The anterior pituitary gland secretes 7 hormones: follicle -stimulating hormone, luteinizing horomone, adrenocorticotropic horomone, thyroid -stimulating horomone, prolactin, endorphins, and growth hormone.

Key Terms

- **anterior pituitary gland**: A major organ of the endocrine system that regulates several physiological processes including stress, growth, reproduction, and lactation.

A major organ of the endocrine system, the anterior pituitary (also called the adenohypophysis) is the glandular, anterior lobe of the pituitary gland. The anterior pituitary regulates several physiological processes including stress, growth, reproduction, and lactation.

Its regulatory functions are achieved through the secretion of various peptide hormones that act on target organs including the adrenal gland, liver, bone, thyroid gland, and gonads. The anterior pituitary itself is regulated by the hypothalamus and by negative feedback from these target organs.

Anatomy of the Anterior Pituitary Gland

The fleshy, glandular anterior pituitary is distinct from the neural composition of the posterior pituitary. The anterior pituitary is composed of multiple parts:

- Pars distalis: This is the distal part that comprises the majority of the anterior pituitary; it is where most pituitary hormone production occurs.
- Pars tuberalis: This is the tubular part that forms a sheath that extends up from the pars distalis and wraps around the pituitary stalk. Its function is poorly understood.
- Pars intermedia: This is the intermediate part that sits between the pars distalis and the posterior pituitary and is often very small in humans.


Major Hormones Secreted by the Anterior Pituitary Gland

- Adrenocorticotropic hormone (ACTH), is a polypeptide whose target is the adrenal gland. The effects of ACTH are upon secretion of glucocorticoid, mineralocorticoids, and sex corticoids.
- Beta-endorphin is a polypeptide that effects the opioid receptor, whose effects include the inhibition of the perception of pain.
- Thyroid-stimulating hormone is a glycoprotein hormone that affects the thyroid gland and the secretion of thyroid hormones.
- Follicle-stimulating hormone is a glycoprotein hormone that targets the gonads and effects the growth of the reproductive system.

- Luteinizing hormone is a glycoprotein hormone that targets the gonads to effect sex-hormone production.
- Growth hormone is a polypeptide hormone that targets the liver and adipose tissue and promotes growth through lipid and carbohydrate metabolism.
- Prolactin is a polypeptide hormone whose target is the ovaries and mammary glands. Prolactin influences the secretion of estrogen/progesterone and milk production.

Regulation

Hormone secretion from the anterior pituitary gland is regulated by hormones secreted by the hypothalamus. Neuroendocrine neurons in the hypothalamus project axons to the median eminence, at the base of the brain. At this site, these neurons can release substances into the small blood vessels that travel directly to the anterior pituitary gland (the hypothalamo-hypophysial portal vessels).



The anterior pituitary: The anterior pituitary, in yellow, is linked to the hypothalamus by a portal system. The hypothalamus releases signaling molecules that incite the anterior pituitary to produce hormones.

This page titled 15.7C: The Anterior Pituitary is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.7D: The Posterior Pituitary

The posterior pituitary secretes two important endocrine hormones—oxytocin and antidiuretic hormone.

Learning Objectives

- Identify the location of the posterior pituitary and the hormones associated with it


Key Points

- The posterior pituitary (or neurohypophysis) comprises the posterior lobe of the pituitary gland and is part of the endocrine system.
- Hormones known as posterior pituitary hormones are synthesized by the hypothalamus, and include oxytocin and antidiuretic hormone.
- The hormones are then stored in neurosecretory vesicles (Herring bodies) before being secreted by the posterior pituitary into the bloodstream.

Key Terms

- **oxytocin**: A hormone that stimulates contractions during labor.
- **posterior pituitary**: The posterior pituitary (or neurohypophysis) comprises the posterior lobe of the pituitary gland and is part of the endocrine system. Despite its name, the posterior pituitary gland is not a true gland; rather, it is largely a collection of axonal projections from the hypothalamus that terminate behind the anterior pituitary gland.
- **Antidiuretic hormone**: A hormone that stimulates water re-absorption in the kidneys.

Posterior Pituitary Gland



Pituitary: Pituitary gland representation.

The posterior pituitary (or neurohypophysis) comprises the posterior lobe of the pituitary gland and is part of the endocrine system. Despite its name, the posterior pituitary gland is not a gland; rather, it is largely a collection of axonal projections from the hypothalamus that terminate behind the anterior pituitary gland.

The posterior pituitary consists mainly of neuronal projections (axons) extending from the supraoptic and paraventricular nuclei of the hypothalamus. These axons release peptide hormones into the capillaries of the hypophyseal circulation. These are then

stored in neurosecretory vesicles (Herring bodies) before being secreted by the posterior pituitary into the systemic bloodstream.

Anatomy of the Posterior Pituitary Gland

The posterior pituitary is derived from the hypothalamus and is distinct from the more fleshy, vascularized anterior lobe. The posterior pituitary is composed of two parts:

- The pars nervosa, also called the neural lobe or posterior lobe, constitutes the majority of the posterior pituitary and is the storage site of oxytocin and vasopressin.
- The infundibular stalk, also known as the infundibulum or pituitary stalk, bridges the hypothalamic and hypophyseal systems.

Major Hormones Secreted by the Posterior Pituitary Gland

The posterior pituitary stores two hormones secreted by the hypothalamus for later release:

- Oxytocin, most of which is released from the paraventricular nucleus in the hypothalamus. Oxytocin is one of the few hormones that create a positive feedback loop.
- Antidiuretic hormone (ADH, also known as vasopressin), the majority of which is released from the supraoptic nucleus in the hypothalamus. ADH acts on the collecting ducts of the kidney to facilitate the reabsorption of water into the blood.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- pituitary gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pituitary_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pituitary_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypothalamus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypothalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pituitary_gland**. **License**: ***Public Domain: No Known Copyright***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Pituitary_gland.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pituitary gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pituitary_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pituitary_gland%23Sections**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypophyseal portal system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypophyseal_portal_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypothalamus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypothalamus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypophyseal portal system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hypophyseal%20portal%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pituitary_gland**. **License**: ***Public Domain: No Known Copyright***
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Pituitary_gland.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pickardt-Fahlbusch-Syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pickard...busch-Syndrome**. **License**: ***Public Domain: No Known Copyright***
- Anterior pituitary. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anterior_pituitary**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- anterior pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/anterio...uitary%20gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pituitary_gland. **License**: *Public Domain: No Known Copyright*
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pituitary_gland.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pickardt-Fahlbusch-Syndrome. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pickard...busch-Syndrome. **License**: *Public Domain: No Known Copyright*
- Pituitary gland et vessel. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._et_vessel.jpg. **License**: *CC BY: Attribution*
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pituita...ohypophysis.29. **License**: *CC BY-SA: Attribution-ShareAlike*
- Posterior pituitary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Posterior_pituitary. **License**: *CC BY-SA: Attribution-ShareAlike*
- vasopressin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasopressin. **License**: *CC BY-SA: Attribution-ShareAlike*
- oxytocin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/oxytocin. **License**: *CC BY-SA: Attribution-ShareAlike*
- posterior pituitary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/posterior%20pituitary. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pituitary_gland. **License**: *Public Domain: No Known Copyright*
- Pituitary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pituitary_gland.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pickardt-Fahlbusch-Syndrome. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pickardt-Fahlbusch-Syndrome. **License**: *Public Domain: No Known Copyright*
- Pituitary gland et vessel. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._et_vessel.jpg. **License**: *CC BY: Attribution*
- Pituitary gland representation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pi...esentation.PNG. **License**: *Public Domain: No Known Copyright*

This page titled 15.7D: The Posterior Pituitary is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.8: THE THYROID GLAND

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

15.8A: Overview of the Thyroid Gland

15.8B: Formation, Storage, and Release of Thyroid Hormones

15.8C: Action of Thyroid Hormones

15.8D: Control of Thyroid Hormone Release

This page titled 15.8: The Thyroid Gland is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.8A: Overview of the Thyroid Gland

The thyroid gland, in the anterior neck, controls body metabolism, protein synthesis, and a body's responsiveness to other hormones.

Learning Objectives

- Summarize the anatomy and purpose of the thyroid gland

Key Points

- The thyroid gland controls how quickly the body uses energy, makes proteins, and controls how sensitive the body is to other hormones. It participates in these processes by producing thyroid hormones, the principal ones being triiodothyronine (T3) and thyroxine (T4).
- Hormones released from the thyroid regulate the rate of metabolism and affect the growth and rate of function of many other systems in the body.
- The thyroid also produces calcitonin, which plays a role in calcium homeostasis.
- Hormonal output from the thyroid is regulated by thyroid-stimulating hormone (TSH) produced by the anterior pituitary, which itself is regulated by thyrotropin-releasing hormone (TRH) produced by the hypothalamus.
- The thyroid gland (the thyroid in vertebrate anatomy) is one of the largest endocrine glands.

Key Terms


- **thyroid-stimulating hormone**: Also known as TSH or thyrotropin, this is a hormone that stimulates the thyroid gland to produce thyroxine (T4), and then triiodothyronine (T3), which stimulates the metabolism of almost every tissue in the body. It is a glycoprotein hormone, synthesized and secreted by thyrotrope cells in the anterior pituitary gland, that regulates the endocrine function of the thyroid gland.
- **thyroxine**: A hormone (an iodine derivative of tyrosine), produced by the thyroid gland, that regulates cell metabolism and growth.

In vertebrates, the thyroid gland is one of the largest endocrine glands. It is found in the neck, below the thyroid cartilage that forms the laryngeal prominence, or Adam's apple. The isthmus (the bridge between the two lobes of the thyroid) is located inferior to the cricoid cartilage.

The thyroid gland controls how quickly the body uses energy, makes proteins, and controls how sensitive the body is to other hormones. It participates in these processes by producing thyroid hormones, the principal ones being triiodothyronine (T3) and thyroxine (sometimes referred to as tetraiodothyronine (T4)).

These hormones regulate the rate of metabolism and affect the growth and rate of function of many other systems in the body. T3 and T4 are synthesized from both iodine and tyrosine.

The thyroid also produces calcitonin, which plays a role in calcium homeostasis. The hormonal output from the thyroid is regulated by thyroid-stimulating hormone (TSH) produced by the anterior pituitary, which itself is regulated by thyrotropin-releasing hormone (TRH) produced by the hypothalamus.



Thyroid system: Thyroid function is regulated by the actions of the hypothalamus and pituitary gland.

Anatomy of the Thyroid Gland

The thyroid gland is a butterfly-shaped organ and is composed of two cone-like lobes or wings, lobus dexter (right lobe) and lobus sinister (left lobe), connected via the isthmus. The organ is situated on the anterior side of the neck, lying against and around the larynx and trachea, reaching posteriorly the oesophagus and carotid sheath.

It starts cranially at the oblique line on the thyroid cartilage (just below the laryngeal prominence, or Adam's Apple), and extends inferiorly to approximately the fifth or sixth tracheal ring. It is difficult to demarcate the gland's upper and lower border with vertebral levels because it moves in position in relation to these structures during swallowing.

This page titled 15.8A: Overview of the Thyroid Gland is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.8B: Formation, Storage, and Release of Thyroid Hormones

Thyroid hormones (T4 and T3) are produced by the follicular cells of the thyroid gland and regulated by thyroid-stimulating hormone (TSH).

Learning Objectives

- Describe the relationship between thyroxine (T4) and triiodothyronine (T3)

Key Points

- Because the effects of T4 in vivo are mediated via T3 (T4 is converted to T3 in target tissues), T3 is three- to five-fold more active than T4.
- Thyroxine is believed to be a pro- hormone and a reservoir for the most active and main thyroid hormone T3. T4 is converted as required in the tissues by iodothyronine deiodinase.
- Thyroid hormones (T4 and T3) are produced by the follicular cells of the thyroid gland and are regulated by thyroid-stimulating hormone secreted by the anterior pituitary gland.

Key Terms

- **thyroid-stimulating hormone**: A hormone that stimulates the thyroid gland to produce thyroxine (T4), and then triiodothyronine (T3), which stimulates the metabolism of almost every tissue in the body.
- **triiodothyronine**: A thyroid hormone also known as T3 that plays a key role in many physiological processes and is the much more active than T4.
- **thyroxine**: A thyroid hormone also known as T4, thought to be a pro-hormone and a reservoir for T3.


The thyroid hormones
thyroxine (T4) and triiodothyronine (T3) are produced from thyroid follicular cells within the thyroid gland, a process regulated by the thyroid-stimulating hormone secreted by the anterior pituitary gland.

Thyroglobulin, the pre-cursor of T4 and T3, is produced by the thyroid follicular cells before being secreted and stored in the follicular lumen. Iodide is actively absorbed from the bloodstream by a process called iodide trapping. In this process, sodium is co-transported with iodide from the basolateral side of the membrane into the cell, and then concentrated in the thyroid follicles to about thirty times its concentration in the blood.

Through a reaction with the enzyme thyroperoxidase, iodine is bound to tyrosine residues in the thyroglobulin molecules to form monoiodotyrosine (MIT) and diiodotyrosine (DIT). Linking two moieties of DIT produces T4. Combining one particle of MIT and one particle of DIT produces T3.

Proteases digest iodinated thyroglobulin, releasing the hormones T4 and T3, the biologically-active agents central to metabolic regulation. T3 is identical to T4, but it has one less iodine atom per molecule.

T4 is believed to be a pro-hormone and a reservoir for the more active and main thyroid hormone T3. T4 is converted as required in the tissues by iodothyronine deiodinase.



Thyroid hormone: Diagrammatic representation of thyroid hormone synthesis in a thyroid follicle.

Effects of Iodine Deficiency

If there is a deficiency of dietary iodine, the thyroid will not be able to make thyroid hormone. A lack of thyroid hormone will lead to decreased negative feedback on the pituitary, which in turn, will lead to increased production of thyroid-stimulating hormone, which causes the thyroid to enlarge (goiter).

This enlarged endemic colloid goiter has the effect of increasing the thyroid's ability to trap more iodide, compensating for the iodine deficiency and allowing it to produce adequate amounts of thyroid hormone.

This page titled 15.8B: Formation, Storage, and Release of Thyroid Hormones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.8C: Action of Thyroid Hormones

The primary function of the thyroid is to produce the hormones triiodothyronine (T3), thyroxine (T4), and calcitonin.

Learning Objectives

- List the actions of thyroid hormones

Key Points

- T4 is converted to T3 by peripheral organs such as the liver, kidney, and spleen.
- Triiodothyronine (T3) is several times more powerful than T4, which is largely a pro- hormone.
- The regulation of actin polymerization by T4 is critical to cell migration in neurons and glial cells and is important for brain development.
- Thyroid hormones play an important role in regulating metabolic rate and body temperature.

Key Terms


- **thyroxine**: A hormone (an iodine derivative of tyrosine) produced by the thyroid gland that regulates cell metabolism and growth.

Triiodothyronine (T3) and thyroxine (T4) are enzymes produced by the thyroid gland. T4 is thought to be a pro-hormone to the more metabolically active T3. T4 is converted to T3 in tissues as required by deiodinase enzymes.

Calcitonin is another hormone released by the thyroid gland that is responsible for modulating blood calcium levels in conjunction with parathyroid hormone, which is released from the parathyroid.

Effect of Thyroid Hormones on Metabolism

The main activity of the thyroid hormones T3 and T4 is to boost the basal metabolic rates of proteins, fats, and carbohydrates as well as vitamins.



Thyroid system: An overview of the thyroid system.

Effect of Thyroid Hormones on Body Temperature

Thyroid hormones affect the dilation of blood vessels, which in turn affects the rate at which heat can escape the body. The more dilated blood vessels are, the faster heat can escape.

A person who suffers from hyperthyroidism (an over-active thyroid) will experience a fever; conversely, a person who suffers from hypothyroidism (a less active thyroid) will experience a decrease in body temperature.

Action of Thyroid Hormones on the Developing Fetus

The cells of the developing brain are a major target for T3 and T4. Thyroid hormones play a particularly crucial role in brain maturation during fetal development by regulating actin polymerization during neuronal development.

Action of Thyroid Hormones in Blood

In the blood, T4 and T3 are partially bound to thyroxine-binding globulin (TBG), transthyretin, and albumin. Only a very small fraction of the circulating hormone is free—T4 0.03% and T3 0.3%. Only the free fraction has hormonal activity.

As with the steroid hormones, thyroid hormones are lipophillic and can cross the cell membrane and bind to intracellular receptors, which act alone as transcription factors or in association with other factors to modulate DNA transcription.

Calcitonin Activity

Calcitonin acts to lower blood calcium levels in three ways:

1. Inhibiting the osteoclast-mediated breakdown of bones.
2. Stimulating osteoblastic activity to produce new bone tissue.
3. Inhibiting re-absorption of calcium in the kidneys.

This page titled 15.8C: Action of Thyroid Hormones is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.8D: Control of Thyroid Hormone Release

The production of thyroxine and triiodothyronine is regulated by thyroid-stimulating hormone (TSH) that is released from the anterior pituitary.

Learning Objectives

- Explain how thyroid hormone release is controlled

Key Points


- Thyroid hormones are released from the thyroid under the control of thyroid stimulating hormone (TSH).
- TSH is secreted by the anterior pituitary gland and is itself under the control of thyrotropin-releasing hormone (TRH).
- TRH is secreted by the hypothalamus.
- Both TSH and TRH secretion are inhibited when elevated thyroid hormone levels are detected in the blood and provide negative feedback to the hypothalamus and anterior pituitary gland.

Key Terms

- **thyroid-stimulating hormone**: A hormone released from the anterior pituitary gland that stimulates the release of thyorid hormones.
- **thyrotropin-releasing hormone**: A hormone released from the hypothalamus that stimulates thyroid-stimulating hormone production from the anterior pituitary gland.

The production of thyroxine (T4) and triiodothyronine (T3) is primarily regulated by thyroid-stimulating hormone (TSH) that is released from the anterior pituitary gland. TSH release, in turn, stimulates the hypothalamus to secrete thyrotropin-releasing hormone (TRH). This results in increased metabolism, growth, development and the activation of numerous other systems controlled by thyroid hormones.

Thyroid hormones also provide negative feedback to the hypothalamus and anterior pituitary gland. When thyroid levels in the blood are elevated TSH and TRH production is reduced. Excessive TRH can also inhibit the production of further TRH.



The thyroid system: Thyroid hormones are produced from the thyroid under the influence of thyroid-stimulating hormone (TSH)

from the anterior pituitary gland, which is itself under the control of thyroptropin-releasing hormone (TRH) secreted by the hypothalamus. Thyroid hormones provide negative feedback, inhibiting secretion of TRH and TSH when blood levels are high.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Thyroid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thyroid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroxine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thyroxine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroid-stimulating hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/thyroid...ting%20hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- triiodothyronine (T3). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/triiodothyronine%20(T3)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thyroid...yroid_hormones**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroxine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thyroxine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroid-stimulating hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/thyroid...ting%20hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid hormone synthesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Th..._synthesis.png**. **License**:
- Thyroid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thyroid...ion_and_action**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroxine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thyroxine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid hormone synthesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Th..._synthesis.png**. **License**:
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyrotropin-releasing hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/thyrotr...sing%20hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thyroid..._T4_regulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- calcitonin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/calcitonin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thyroid follicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/thyroid%20follicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thyroid hormone synthesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_hormone_synthesis.png**. **License**:
- Thyroid system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Thyroid_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 871px-Thyroid_system.svg.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Thyroid_system.svg**. **License**:

This page titled 15.8D: Control of Thyroid Hormone Release is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.9: THE PARATHYROID GLANDS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

15.9A: Overview of the Parathyroid Glands

15.9B: Parathyroid Hormone

This page titled 15.9: The Parathyroid Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.9A: Overview of the Parathyroid Glands

The parathyroid glands are small endocrine glands in the neck that produce parathyroid hormone.

Learning Objectives

- Describe the location and function of the parathyroid glands


Key Points

- The parathyroid glands are four or more small glands, about the size of a grain of rice, located on the posterior surface of the thyroid gland.
- The major function of the parathyroid glands is to maintain the body's calcium level within a very narrow range, so the nervous and muscular systems can function properly.

Key Terms

- **calcitonin**: A hormone that is produced primarily by the parafollicular cells of the thyroid. It acts to reduce blood calcium (Ca2+), opposing the effects of parathyroid hormone.
- **parathyroid gland**: One of four endocrine glands situated in the neck, usually on the posterior surface of the thyroid gland, that produce parathyroid hormone.
- **parathyroid hormone**: A hormone produced by the parathyroid gland that acts to increase blood calcium levels by stimulating osteoclasts to release calcium from the bone.

The parathyroid glands are small endocrine glands —approximately the size of a grain of rice—in the neck that produce parathyroid hormone. Humans usually have four parathyroid glands, which are usually located on the posterior surface of the thyroid gland, or, in rare cases, within the thyroid gland itself or in the chest.



Parathyroid gland: The parathyroid gland in relation to the thyroid gland. They are located on the posterior surface of the lobes.

The two parathyroid glands on each side that are positioned higher are called the superior parathyroid glands, while the lower two are called the inferior parathyroid glands. Occasionally, some individuals may have six, eight, or even more parathyroid glands.

Parathyroid glands control the amount of calcium in the blood and within the bones. The major function of the parathyroid glands is to maintain the body's calcium level within a very narrow range, so that the nervous and muscular systems can function properly.

When blood calcium levels drop below a certain point, calcium-sensing receptors in the parathyroid gland are activated to release parathyroid hormone (PTH) into the blood.

PTH modulates calcium and phosphate homeostasis, as well as bone physiology. PTH has effects antagonistic to those of calcitonin by increasing blood calcium levels by stimulating osteoclasts to break down bone and release calcium. PTH also increases gastrointestinal calcium absorption by activating vitamin D, and promotes calcium conservation by re-absorption in the kidneys.

This page titled 15.9A: Overview of the Parathyroid Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

15.9B: Parathyroid Hormone

Parathyroid hormone maintains the body's calcium levels by increasing the absorption of calcium from the bones, kidneys, and GI tract.

Learning Objectives

- Explain the functioning of parathyroid hormone

Key Points

- Parathyroid hormone works in concert with another hormone, calcitonin, that is produced by the thyroid to maintain blood calcium levels.
- Parathyroid hormone acts to increase blood calcium levels, while calcitonin acts to decrease blood calcium levels.
- When blood calcium levels drop below a certain point, calcium-sensing receptors in the parathyroid gland are activated, and the parathyroid glands release parathyroid hormone into the blood.
- PTH acts on the bone to increase blood calcium levels by stimulating osteoclasts to break down bone and release calcium into the bloodstream; on the GI tract to increase the activity of the enzyme in the intestines that activates vitamin D; and on the kidneys to promote calcium reabsorption.

Key Terms

- **vitamin D**: A fat-soluble vitamin required for normal bone development and that prevents rickets; it can be manufactured in the skin on exposure to sunlight.
- **osteoclast**: A large multinuclear cell associated with the break down and resorption of bone.
- **bone remodeling**: The resorption by osteoclasts and replacement by osteoblasts in bones.


Parathyroid Glands

The parathyroid glands are small, pea-sized endocrine glands located on the rear side of the thyroid gland. The major function of the parathyroid glands is to maintain the body's calcium level within a very narrow range, so that the nervous and muscular systems, which depend on calcium to transmit action potentials, can function properly.

When blood calcium levels drop below a certain point, the calcium-sensing receptors in the parathyroid gland are activated, and the parathyroid glands release parathyroid hormone (PTH) into the blood. PTH is a small protein hormone that is integral to the regulation of the level of calcium in the blood via the bone, kidneys, and intestines.

PTH works in concert with another hormone, calcitonin, that is produced by the thyroid to maintain calcium homoeostasis. Parathyroid hormone acts to increase blood calcium levels, while calcitonin acts to decrease blood calcium levels.

This interaction between parathyroid hormone and calcitonin is also an important part of bone remodeling. This is a lifelong process where mature bone tissue is removed from the skeleton and new bone tissue is formed.



Calcium regulation: Parathyroid hormone regulates the levels of calcium in the blood. to the parathyroid glands.

Parathyroid Hormone Action

- Parathyroid hormone acts on a bone to increase its blood calcium levels by stimulating osteoclasts to break down bone and release calcium into the bloodstream.
- Parathyroid hormone acts on the gastrointestinal tract to increase blood calcium by increasing the activity of the enzyme in the intestines that activates vitamin D.
- It acts on the kidneys to increase blood calcium levels by promoting calcium reabsorption in the nephrons.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Parathyroid gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parathyroid_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- calcitonin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/calcitonin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vitamin D. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vitamin_D**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parathyroid gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/parathyroid_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parathyroid gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parathyroid_gland**. **License**: ***Public Domain: No Known Copyright***
- Parathyroid hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parathyroid_hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteoclast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/osteoclast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bone remodeling. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bone%20remodeling**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vitamin D. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vitamin_D**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Parathyroid gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Parathyroid_gland**. **License**: ***Public***

Domain: No Known Copyright

- Calcium regulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ca...regulation.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 15.9B: Parathyroid Hormone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16: CARDIOVASCULAR SYSTEM - BLOOD

- Topic hierarchy

16.1: Overview of Blood

16.1A: Components of Blood

16.1B: Physical Characteristics and Volume

16.1C: Functions of Blood

16.1D: Blood Plasma

16.2: Red Blood Cells

16.2A: RBC Anatomy

16.2B: RBC Physiology

16.2C: RBC Life Cycle

16.3: White Blood Cells

16.3A: Types of WBCs

16.3B: WBC Function

16.3C: WBC Formation

16.4: Platelets

16.4A: Platelets

16.4B: Platelet Formation

16.5: Hemostasis

16.5A: Overview of Hemostasis

16.5B: Vascular Spasm

16.5C: Platelet Plug Formation

16.5D: Coagulation

16.5E: Role of Vitamin K

16.5F: Clot Retraction and Repair

16.5G: Fibrinolysis

16.6: Transfusions

16.6A: Transfusions of Whole Blood

16.6B: Plasma and Blood Volume Expanders

16.6C: Blood Groups and Blood Types

16.6D: Typing and Cross-Matching for Transfusions

This page titled 16: Cardiovascular System - Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.1: OVERVIEW OF BLOOD



- Topic hierarchy

16.1A: Components of Blood

16.1B: Physical Characteristics and Volume

16.1C: Functions of Blood

16.1D: Blood Plasma

This page titled 16.1: Overview of Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.1A: Components of Blood

Blood is composed of plasma and three types of cells: red blood cells, white blood cells, and platelets.

Learning Objectives

- Differentiate among the blood components of erythrocytes, leukocytes, and thrombocytes

Key Points

- Red blood cells' primary function is to transport oxygen between the lungs and tissues of the body.
- White blood cells, the cells of the immune system, provide defense against pathogens.
- Platelets are involved in clot formation during wound healing.
- Blood is an extracellular matrix tissue in which various blood cells are suspended in the plasma matrix.
- Blood is vital for normal metabolic function due to the transfer of oxygen, carbon dioxide, and glucose to and from the body's tissues. It also transports a number of other cells and molecules throughout the body.

Key Terms

- **plasma**: The straw-colored or pale yellow liquid component of blood in which blood cells are suspended.
- **hemoglobin**: The iron-containing substance in red blood cells that binds to and transports oxygen from the alveoli of the lungs to the tissues of the body. It consists of a protein (globulin) and haem (a porphyrin ring with an atom of iron at its center).



Composition of Blood: Two tubes of EDTA-anticoagulated blood. Left tube: after standing, the RBCs have settled at the bottom of the tube.

Blood is a circulating tissue composed of fluid, plasma, and cells. The cellular components of blood are erythrocytes (red blood cells, or RBCs), leukocytes (white blood cells, or WBCs), and thrombocytes (platelets). By volume, the RBCs constitute about 45% of whole blood, the plasma about 54.3%, and white blood cells about 0.7%. Platelets make up less than 1%. Although it consists of cells suspended in fluid, blood is still considered a tissue as it is technically a type of extracellular matrix.


Blood enables transport of cells and molecules between parts of the body. Oxygen, carbon dioxide, and glucose are among the

most vital molecules transported in blood. Blood cells are essential for normal metabolic and immune system function.

Erythrocytes (RBCs)

Erythrocytes are discs measuring about seven to eight micrometers in diameter. RBCs contain hemoglobin molecules which bind to oxygen so it can be transported to tissues. Mature RBCs lack a nucleus and organelles and have no nuclear DNA. RBCs, endothelial vessel cells, and other blood cells are also marked by glycoproteins that define the different blood types. The ratio of RBCs to blood plasma is referred to as the hematocrit, and is normally about 45%. The combined surface area of all red blood cells of the human body would be roughly 2,000 times greater than the body's exterior surface.

Leukocytes (WBCs)



Components of Blood: From left to right, a diagram of an erythrocyte, thrombocyte, and leukocyte.

Leukocytes are usually larger in size (10–14 micrometers in diameter) than red blood cells. They lack hemoglobin but contain organelles, a nucleus, and nuclear DNA. WBCs are the main functional component of the body's immune system. They destroy and remove old or aberrant cells and cellular debris, as well as attack infectious agents (pathogens) and foreign substances. There are several different types of white blood cells: basophils, eosinophils, neutrophils, monocytes, natural killer cells, B- and T-cell lymphocytes, macrophages, and dendritic cells, all of which perform distinct functions.

Thrombocytes (Platelets)

Thrombocytes measure between one to two micrometers in diameter. These membrane-bound cell fragments lack nuclei and are responsible for blood clotting (coagulation). They result from fragmentation of large cells called megakaryocytes, which are derived from stem cells in the bone marrow. Platelets are produced at a rate of 200 billion per day, a process regulated by the hormone thrombopoietin. Platelets contain mitochondrial DNA, but not nuclear DNA.

The sticky surface of platelets allows them to accumulate at the site of broken blood vessels to form a clot, due in part to the release of clotting factors that occurs during endothelial injury to blood vessels. This process is called hemostatis. Platelets secrete factors that increase local platelet aggregation (e.g., thromboxane A), enhance vasoconstriction (e.g., serotonin), and promote blood coagulation (e.g., thromboplastin, fibrinogen). Platelets are critically important for wound healing, which can only occur once the clot forms and bleeding ceases completely.

This page titled 16.1A: Components of Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.1B: Physical Characteristics and Volume

Blood contains plasma and blood cells, some of which have hemoglobin that makes blood red. The average blood volume in adult is five liters.

Learning Objectives

- Describe the physical characteristics and volume of blood in adults

Key Points

- Blood accounts for 8% of human body weight. The average adult has a blood volume of roughly five liters (1.3 gal).
- By volume, red blood cells constitute about 45% of whole blood, plasma about 54.3%, and white blood cells about 0.7%. Platelets make up less than 1%. Blood also contains proteins such as albumins.
- Hemoglobin is the principal determinant of blood color in vertebrates. Each molecule has four heme groups, and their interaction with various molecules alters the exact color of blood.
- Veins appear to be blue because blue light penetrates the skin better than other forms of light. Deoxygenated blood is not blue.
- Blood volume is a regulated variable that is proportional to blood pressure and a component of homeostasis.
- Injury can cause blood loss. A healthy adult can lose almost 20% of blood volume (1 L) before the first symptom (restlessness) begins, and 40% of volume (2 L) before hypovolemic shock sets in.

Key Terms

- **erythrocyte**: An anucleate cell in the blood involved with the transport of oxygen. Also called a red blood cell because of the red coloring of hemoglobin.
- **hemoglobin**: The iron-containing substance in red blood cells that transports oxygen from the lungs to the rest of the body. It consists of a protein (globulin) and haem (a porphyrin ring with an atom of iron at its center).
- **tissue perfusion**: The amount of blood that can reach the tissues to supply them with oxygen and glucose.

Blood is a specialized bodily fluid in animals that delivers necessary substances, such as nutrients and oxygen, to the cells and transports metabolic waste products away from those same cells. Blood plays many roles in sustaining life and has physical characteristics that distinguish it from other body tissues.

Physical Characteristics

Blood is a fluid that is technically considered a connective tissue. It is an extracellular matrix in which blood cells are suspended in plasma. It normally has a pH of about 7.4 and is slightly denser and more viscous than water. Blood contains red blood cells (RBCs), white blood cells (WBCs), platelets, and other cell fragments, molecules, and debris. Albumin is the main protein found in plasma, and it functions to regulate the colloidal osmotic pressure of blood.

Blood appears red because of the high amount of hemoglobin, a molecule found on RBCs. Each hemoglobin molecule has four heme groups that interact with various molecules, which alters the exact color. In oxygenated blood found in arterial circulation, hemoglobin-bound oxygen creates a distinctive red color.

Deoxygenated blood is a darker shade of red. It is present in veins and can be seen during blood donation or lab tests. Carbon monoxide poisoning causes bright red blood due to the formation of carboxyhemoglobin. In cyanide poisoning, venous blood remains oxygenated, increasing the redness. Under normal conditions, blood can never truly be blue, although most visible veins appear blue because only blue light can can penetrate deeply enough to illuminate veins beneath the skin.

Blood Volume

Blood generally accounts for 8% of the human body weight. The average adult has a blood volume of roughly five liters (1.3 gal). By volume, red blood cells constitute about 45% of whole blood, plasma about 54.3%, and white cells about 0.7%, with platelets making up less than 1%.



Composition of blood: Two tubes of EDTA-anticoagulated blood. Left tube: After standing, the RBCs have settled at the bottom of the tube. Right tube: contains freshly drawn blood.

Blood volume is a regulated variable that is directly proportional to blood pressure through the output of the heart. In order to maintain homeostasis, blood volume and blood pressure must be high enough that blood can reach all of the body's tissues, a process called tissue perfusion. Most tissues can survive without perfusion for a short amount of time, but the brain needs a continuous supply of oxygen and glucose to stay alive.

Many mechanisms exist to regulate blood volume and tissue perfusion, including renal water excretion in the kidney, the pumping activity of the heart, and the abilities of the arteries to constrict or dilate. When blood volume becomes too low, such as from an injury, dehydration, or internal bleeding, the body will enter into a state of hypovolemic shock, in which tissue perfusion decreases too much. A healthy adult can lose almost 20% of blood volume (1 L) before the first symptom, restlessness, begins, and 40% of volume (2 L) before hypovolemic shock sets in. Conversely, higher than normal blood volume may cause hypertension, heart failure, and aneurysms.

This page titled 16.1B: Physical Characteristics and Volume is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.1C: Functions of Blood

The main function of blood is to supply oxygen to tissues and remove carbon dioxide. Other functions include pH regulation and thermoregulation.

Learning Objectives

- Describe the functions blood performs in the body

Key Points

- The main function of blood is to carry oxygen from lungs and deliver it to the body, where it is released and carbon dioxide is consumed.
- The tissues of the body cannot survive without blood perfusion. Without blood, tissues may undergo hypoxia, ischemia, or infarction depending on the severity of the deficiency.
- The blood is involved in the homeostasis of variables such as temperature, blood volume, blood pressure, blood pH, and blood glucose levels.
- Other important blood functions include carbon dioxide transport and hormone signaling.
- The blood is involved in immune system functions such as white blood cell activity and blood clotting.

Key Terms

- **hypoxia**: A condition in which tissues are deprived of an adequate supply of oxygen for metabolic purposes; anoxia.
- **coagulation**: The process by which blood forms solid clots.

Blood performs many functions critical for sustaining metabolic physiological processes in complex organisms. Blood is involved in everything from gas exchange to nutrient transport to immune system and homeostatic functions.

Oxygen and Glucose Transport


Blood's primary function is to transport molecules around the body to support critical metabolic processes. All cells require oxygen and glucose to undergo cellular respiration. Tissues cannot survive very long without these two molecules. Disruption of this process is most dangerous to the brain, which can survive only about two minutes without oxygen and glucose. These terms are used to describe oxygen or blood deficiency to tissues in the body:

- Hypoxia: a state in which the tissues do not receive adequate oxygen supply, generally due to decreased tissue perfusion or decreased oxygen intake.
- Ischemia: a reversible condition in which a tissue does not receive adequate blood supply, usually from an obstructed or ruptured blood vessel.
- Infarction: a usually irreversible condition in which tissues die as a result of prolonged oxygen or blood supply.

Most tissues can survive in a hypoxic or ischemic state for a few hours before infarction sets in. Heart infarction, which often occurs during a heart attack, will cause infarction in other tissues as blood is no longer pumped.

In addition to oxygen and glucose, the blood transports several other important molecules. Carbon dioxide, which travels through the blood mostly as bicarbonate, is transported from tissues as a waste product of cellular respiration to the lungs during gas exchange. Many hormones (chemical messengers) also travel through the blood as a form of communication between interrelated organs, which are often involved in homeostatic control.

Immune System Functions



Cellular Components of Blood: From left, diagrams of an erythrocyte, thrombocyte, and leukocyte.

White blood cells and antibodies circulate through the blood and destroy any foreign invaders (pathogens) that they encounter. Inflammation occurs in blood vessels due to the release of inflammatory mediators in the blood. This causes vasodilation and redness as other white blood cells are drawn to the region through the bloodstream to destroy infectious pathogens. They may also find molecular markers of pathogens called antigens and take them to lymphatic organs to stimulate powerful adaptive immune system responses.

The blood also has the ability to undergo clotting in response to vascular injury such as bleeding. Normally a series of clotting and anti-clotting factors are kept in balance through the blood so that no clotting occurs, but when endothelial cells are injured, the clotting factors are increased and cause blood to clot. Circulating platelets in the blood arrive at the injury site and form a mesh and plug to coagulate the blood and stop the bleeding. Wound healing can only begin after this clotting response occurs.

Homeostatic Functions

Blood is involved in maintaining homeostasis through several ways. Temperature regulation occurs in part as a result of the dilation and constriction of vessels in the blood. Blood pH is a regulated variable of the respiratory system, because the pH of blood is directly proportional to the amount of carbon dioxide dissolved in blood. This makes blood pH an indicator of respiratory homeostasis. Blood glucose levels are regulated by insulin and glucagon secretion. Blood volume and blood pressure are directly proportional regulated variables that are tied to the activity of the heart and the fluid retention of the kidney. If any of these variables are too high or too low, severe problems can occur. For that reason, a number of complex negative feedback mechanisms exist to keep all variables within homeostatic range, despite influences from the internal and external environments.

This page titled 16.1C: Functions of Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.1D: Blood Plasma

Plasma comprises about 55% of total blood volume. It contains proteins and clotting factors, transports nutrients, and removes waste.

LEARNING OBJECTIVES

Describe the features of blood plasma

KEY TAKEAWAYS

Key Points

- The majority of blood volume consists of plasma. This aqueous solution is 92% water. It also contains blood plasma proteins, including serum albumin, blood-clotting factors, and immunoglobulins.
- Plasma circulates respiratory gases, dissolved nutrients, and other materials. It also removes waste products.
- Globulins are a diverse group of proteins that primarily transport other substances and inhibit certain enzymes.
- Albumins maintain osmotic balance between the blood and tissue fluids through exertion of oncotic pressure.
- Fibrinogen is the main clotting protein found in plasma. It is responsible for stopping blood flow during wound healing.

Key Terms

- **platelet**: A small, colorless, disc-shaped particle found in the blood of mammals. It plays an important role in blood clot formation.
- **immunoglobulin**: Any of the glycoproteins in blood serum that respond to invasion by foreign antigens and that protect the host by removing pathogens; an antibody.
- **albumins**: A plasma protein that exerts a high degree of oncotic pressure to pull water and other substances into tissues.

About 55% of blood is blood plasma, a straw-colored liquid matrix in which blood cells are suspended. It is an aqueous solution containing about 90% water, 8% soluble blood plasma proteins, 1% electrolytes, and 1% elements in transit. One percent of the plasma is salt, which helps with pH. Human blood plasma volume averages about 2.7–3.0 liters.

Molecular Contents of Plasma



Composition of Blood: Two tubes of EDTA-anticoagulated blood. Left tube: after standing, the RBCs have settled at the bottom of the tube.

Plasma contains molecules that are transported around the body. Respiratory gases, such as oxygen and carbon dioxide, may be dissolved directly in the plasma. However, most oxygen is hemoglobin bound, and most carbon dioxide is converted to bicarbonate ions in the plasma. Hormones and nutrients such as glucose, amino acids and proteins, lipids and fatty acids, and vitamins are also dissolved in the plasma. Waste products are carried through the plasma during their removal, including urea and ammonia.

Plasma Proteins

The largest group of solutes in plasma contains three important proteins: albumins, globulins, and clotting proteins.

Albumins

Albumins, produced in the liver, make up about two-thirds of the proteins in plasma. Albumins maintain the osmotic balance between the blood and tissue fluids. These proteins exert a force that pulls water towards them, which is called oncotic or osmotic pressure. During inflammation, albumins leave the vascular endothelium and enter the tissues, which transports water and some of the plasma into the interstitial fluid. This is the principal cause of exudate edema, which is the swelling that indicates inflammation.

Albumins also assist in transport of different materials, such as vitamins and certain molecules and drugs (e.g. bilirubin, fatty acids, and penicillin) due to the force exerted by their oncotic pressure. Plasma that is pulled into the tissues by albumin-exerted oncotic pressure becomes interstitial fluid. This gradually drains into the lymphatic system which it turn recirculates it back into the plasma of the circulatory system.

Globulins

Globulins are a diverse group of proteins designated into three groups, gamma, alpha, and beta, based on how far they move during electrophoresis tests. Their main function is to transport various substances in the blood. For example, the beta globulin transferrin can transport iron. Most gamma globulins are antibodies (immunoglobulin), which assist the body's immune system in defense against infections and illness. Alpha globulins are notable for inhibiting certain proteases, while beta globulins often function as enzymes in the body.

Clotting Factors

Clotting proteins are mainly produced in the liver. Twelve proteins known as "clotting factors" participate in the cascade clotting process during endothelial injury. One important clotting factor is fibrinogen. Fibrinogen generates fibrin when activated by the coagulant thrombin, which forms a mesh that clots blood with the assistance of a platelet plug. Normally, anticoagulants and fibrinolytics in the plasma, such as plasmin and heparin, break up fibrin clots and inactivate thrombin. However, during endothelial injury, damaged cells will release tissue factor, another type of clotting factor that causes a cascade of thrombin production that will overpower the anticoagulants and cause a clotting response.

Serum is a term used to describe plasma that has been removed of its clotting factors. Serum still contains albumin and globulins, which are often called serum proteins as a result.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...od_Composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plasma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycoprotein. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glycoprotein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hemoglobin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...od_Composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- erythrocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/erythrocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hemoglobin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plasma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...od_Composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoxia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoxia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- coagulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/coagulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- oxyhemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/oxyhemoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%2...of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Blood_physiology%23Blood_Composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Constituents_of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Red_White_Blood_cells.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood#/media/File:Red_White_Blood_cells.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Blood_physiology%23Plasma_makeup**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- platelet. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/platelet**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- immunoglobulin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/immunoglobulin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- electrolyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/electrolyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Constituents_of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/**

Blood_physiology%23Blood_Composition. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Constituents_of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Red_White_Blood_cells.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood#/media/File:Red_White_Blood_cells.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Constituents_of_human_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 16.1D: Blood Plasma is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.2: RED BLOOD CELLS

16.2A: RBC Anatomy
16.2B: RBC Physiology
16.2C: RBC Life Cycle

This page titled 16.2: Red Blood Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


16.2A: RBC Anatomy

Learning Objectives

- Diagram the anatomy of an erythrocyte (red blood cell, or RBC)

Human erythrocytes or red blood cells (RBCs) are the primary cellular component of blood. They are involved in oxygen transport through the body and have features that distinguish them from every other type of human cell. Adult humans have roughly 20-30 trillion RBCs at any given time, comprising approximately one quarter of the total number of human cells.

External Structure



Red blood cells: Human red blood cells (6–8μm)

RBCs are disc-shaped with a flatter, concave center. This biconcave shape allows the cells to flow smoothly through the narrowest blood vessels. Gas exchange with tissues occurs in capillaries, tiny blood vessels that are only as wide as one cell. Many RBCs are wider than capillaries, but their shape provides the needed flexibility to squeeze through.

A typical human RBC has a disk diameter of 6–8 micrometers and a thickness of 2 micrometers, much smaller than most other human cells. These cells have an average volume of about 90 femtoliters (fL) with a surface area of about 136 square micrometers. They can swell up to a sphere shape containing 150 fL without bursting their cell membrane. When the shape does change, it inhibits their ability to carry oxygen or participate in gas exchange. This occurs in people with spherocytic (sphere-shaped) anemia or sickle-cell anemia.

Internal Structure

Although RBCs are considered cells, they lack a nucleus, nuclear DNA, and most organelles, including the endoplasmic reticulum and mitochondria. RBCs therefore cannot divide or replicate like other labile cells of the body. They also lack the components to express genes and synthesize proteins. While most cells have chemotaxic ways to travel through the body, RBCs are carried through the body by blood flow and pressure alone.

Hemoglobin molecules are the most important component of RBCs. Hemoglobin is a specialized protein that contains a binding site for the transport of oxygen and other molecules. The RBCs' distinctive red color is due to the spectral properties of the binding of hemic iron ions in hemoglobin. Each human red blood cell contains approximately 270 million of these hemoglobin biomolecules, each carrying four heme groups (individual proteins). Hemoglobin comprises about a third of the total RBC volume. This protein is responsible for the transport of more than 98% of the oxygen, while the rest travels as dissolved molecules through the plasma.

Key Points

- The biconcave shape allows RBCs to bend and flow smoothly through the body's capillaries. It also facilitates oxygen transport.
- Red blood cells are considered cells, but they lack a nucleus, DNA, and organelles like the endoplasmic reticulum or mitochondria.
- Red blood cells cannot divide or replicate like other bodily cells. They cannot independently synthesize proteins.
- The blood's red color is due to the spectral properties of the hemic iron ions in hemoglobin.
- Each human red blood cell contains approximately 270 million hemoglobin biomolecules, each carrying four heme groups to which oxygen binds.

Key Terms

- **iron**: A metallic chemical element with atomic number 26 and symbol Fe. Iron-containing enzymes and proteins, often containing heme prosthetic groups, participate in many biological oxidations and in transport.
- **hemoglobin**: The iron-containing substance in RBCs that transports oxygen from the lungs to the rest of the body. It consists of a protein (globulin) and haem (a porphyrin ring with an atom of iron at its center).

This page titled 16.2A: RBC Anatomy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.2B: RBC Physiology

The primary functions of red blood cells (RBCs) include carrying oxygen to all parts of the body, binding to hemoglobin, and removing carbon dioxide.

Learning Objectives

- Discuss the primary function of erythrocytes (red blood cells)

Key Points


- Red blood cells contain hemoglobin,which contains four iron-binding heme groups.
- Oxygen binds the heme groups of hemoglobin. Each hemoglobin molecule can bind four oxygen molecules.
- The binding affinity of hemoglobin for oxygen is cooperative. It is increased by the oxygen saturation of the molecule. Binding of an initial oxygen molecule influences the shape of the other binding sites. This makes binding more favorable for additional oxygen molecules.
- Each hemoglobin molecule contains four iron-binding heme groups which are the site of oxygen binding. Oxygen-bound hemoglobin is called oxyhemoglobin.
- Red blood cells alter blood pH by catalyzing the reversible carbon dioxide to carbonic acid reaction through the enzyme carbonic anhydrase.
- pH is also controlled by carbon dioxide binding to hemoglobin instead of being converted to carbonic acid.

Key Terms

- **carbonic anhydrase**: The enzyme found in RBCs that catalyzes the reaction between carbonic acid and carbon dioxide and water.
- **cooperative binding**: In binding in which multiple molecules can potentially bind to multiple binding sites, when a first molecule is bound to a binding site, the same molecule is favored for the rest of the binding sites through increased binding affinity.

Red blood cells (RBCs) perform a number of human respiratory and cardiovascular system functions. Most of these functions are attributed to hemoglobin content. The main RBC functions are facilitating gas exchange and regulating blood pH.


Gas Exchange



Heme: This is a diagram of the molecular structure of heme.

RBCs facilitate gas exchange through a protein called hemoglobin. The word hemoglobin comes from “hemo” meaning blood and “globin” meaning protein. Hemoglobin is a quaternary structure protein consisting of many smaller tertiary structure proteins composed of amino acid polypeptide chains. Each hemoglobin molecule contains four iron-binding heme groups, which are the site of oxygen (O_2) binding. Oxygen bound hemoglobin is called oxyhemoglobin.


The binding of oxygen is a cooperative process. Hemoglobin bound oxygen causes a gradual increase in oxygen-binding affinity until all binding sites on the hemoglobin molecule are filled. As a result, the oxygen-binding curve of hemoglobin (also called the oxygen saturation or dissociation curve) is sigmoidal, or S-shaped, as opposed to the normal hyperbolic curve associated with noncooperative binding. This curve shows the saturation of oxygen bound to hemoglobin compared to the partial pressure of oxygen (concentration) in blood.



Oxygen saturation curve: Due to cooperative binding, the oxygen saturation curve is S-shaped.

pH Control

RBCs control blood pH by changing the form of carbon dioxide within the blood. Carbon dioxide is associated with blood acidity. That's because most carbon dioxide travels through the blood as a bicarbonate ion, which is the dissociated form of carbonic acid in solution. The respiratory system regulates blood pH by changing the rate at which carbon dioxide is exhaled from the body, which involves the RBC's molecular activity. RBCs alter blood pH in a few different ways.



Quaternary structure: hemoglobin: Hemoglobin is a globular protein composed of four polypeptide subunits (two alpha chains, in blue, and two beta pleated sheets, in red). The heme groups are the green structures nestled among the alpha and beta.

RBCs secrete the enzyme carbonic anhydrase, which catalyzes the conversion of carbon dioxide and water to carbonic acid. This dissociates in solution into bicarbonate and hydrogen ions, the driving force of pH in the blood. This reaction is reversible by the same enzyme. Carbonic anhydrase also removes water from carbonic acid to turn it back into carbon dioxide and water. This process is essential so carbon dioxide can exist as a gas during gas exchange in the alveolar capillaries. As carbon dioxide is converted from its dissolved acid form and exhaled through the lungs, blood pH becomes less acidic. This reaction can occur without the presence of RBCs or carbonic anhydrase, but at a much slower rate. With the catalyst activity of carbonic anhydrase, this reaction is one of the fastest in the human body.

Hemoglobin can also bind to carbon dioxide, which creates carbamino-hemoglobin. When carbon dioxide binds to hemoglobin, it doesn't exist in the form of carbonic acid, which makes the blood less acidic and increases blood pH. However, because of allosteric effects on the hemoglobin molecule, the binding of carbon dioxide decreases the amount of oxygen bound for a given partial pressure of oxygen. This decrease in hemoglobin's affinity for oxygen by the binding of carbon dioxide is known as the Bohr effect, which results in a rightward shift to the O_2-saturation curve. Conversely, when the carbon dioxide levels in the blood decrease (i.e., in the lung capillaries), carbon dioxide and hydrogen ions are released from hemoglobin, increasing the oxygen affinity of the protein. A reduction in the total binding capacity of hemoglobin to oxygen (i.e. shifting the curve down, not just to the right) due to reduced pH is called the Haldane effect.

This page titled 16.2B: RBC Physiology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.2C: RBC Life Cycle

Human erythrocytes are produced through a process called erythropoiesis. They take about seven days to mature.

Learning Objectives

- Outline the life cycle of erythrocytes (red blood cells, or RBCs)


Key Points

- After about 100-120 days, RBCs are removed from circulation through a process called eryptosis.
- Erythropoiesis is the process by which human erythrocytes are produced. It is triggered by erythropoietin, a kidney hormone produced during hypoxia.
- Erythropoiesis takes place in the bone marrow, where hemopoietic stem cells differentiate and eventually shed their nuclei to become reticulocytes. Iron, vitamin B12, and folic acid are required for hemoglobin synthesis and normal RBC maturation.
- Reticulocytes mature into normal, functional RBCs after 24 hours in the bloodstream.
- Following eryptosis, the liver breaks down old hemoglobin into biliverdin and iron. The iron is taken back to the bone marrow for reuse by transferrins, while biliverdin is broken down into bilirubin and excreted through digestive system bile.

Key Terms

- **erythropoietin**: A hormone produced by the kidneys in response to hypoxia, which stimulates erythropoiesis.
- **bilirubin**: A bile pigment that arises when biliverdin is separated from the iron of old hemoglobin molecules in the liver. Bilirubin becomes part of bile salts in the digestive system and is excreted, while the iron content is reused.

Human erythrocytes are produced through a process called erythropoiesis, developing from committed stem cells to mature erythrocytes in about seven days. When matured, these cells circulate in the blood for about 100 to 120 days, performing their normal function of molecule transport. At the end of their lifespan, they degrade and are removed from circulation.



Scanning electron micrograph of blood cells: Shown on the left, the erythrocyte, or red blood cell, has a round, donut-like shape.

Erythopoiesis

Erythropoiesis is the process in which new erythrocytes are produced, which takes about seven days. Erythrocytes are continuously produced in the red bone marrow of large bones at a rate of about 2 million cells per second in a healthy adult. Erythrocytes differentiate from erythrotropietic bone marrow cells, a type of hemopoietic stem cell found in bone marrow. Unlike mature RBCs, bone marrow cells contain a nucleus. In the embryo, the liver is the main site of red blood cell production and bears similar types of stem cells at this stage of development.

Erythropoiesis can be stimulated by the hormone erythropoietin, which is synthesized by the kidney in response to hypoxia (systemic oxygen deficiency). In the last stages of development, the immature RBCs absorb iron, Vitamin B12, and folic acid. These dietary nutrients that are necessary for proper synthesis of hemoglobin (iron) and normal RBC development (B12 and folic acid). Deficiency of any of these nutrients may cause anemia, a condition in which there aren't enough fully functional RBCs carrying oxygen in the bloodstream. Just before and after leaving the bone marrow, the developing cells are known as reticulocytes. These immature RBCs that have shed their nuclei following initial differentiation. After 24 hours in the bloodstream, reticulocytes mature into functional RBCs.

Eryptosis

Eryptosis, a form of apoptosis (programmed cell death), is the aging and death of mature RBCs. As an RBC ages, it undergoes changes in its plasma membrane that make it susceptible to selective recognition by macrophages and subsequent phagocytosis in the reticuloendothelial system (spleen, liver, and bone marrow). This process removes old and defective cells and continually purges the blood. Eryptosis normally occurs at the same rate as erythropoiesis, keeping the total circulating red blood cell count in a state of equilibrium. Many diseases that involve damage to RBCs (hemolytic anemias, sepsis, malaria, pernicious or nutritional anemias) or normal cellular processes that cause cellular damage (oxidative stress) may increase the rate of eryptosis. Conversely, erythropotein and nitric oxide (a vasodilator) will inhibit eryptosis.

Following eryptosis, the hemoglobin content within the RBC is broken down and recirculated throughout the body. The heme components of hemoglobin are broken down into iron ions and a green bile pigment called biliverdin. The biliverdin is reduced to the yellow bile pigment bilirubin, which is released into the plasma and recirculated to the liver, then bound to albumin and stored in the gallbladder. The bilirubin is excreted through the digestive system in the form of bile, while some of the iron is released into the plasma to be recirculated back into the bone marrow by a carrier protein called transferrin. This iron is then reused for erythropoiesis, but additional dietary iron is needed to support healthy RBC life cycles.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Erythrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Erythro...n_erythrocytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Blood_physiology%23Red_Blood_Cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hemoglobin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- iron. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/iron**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- biosynthesis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/biosynthesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Erythrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Erythro...n_erythrocytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Blood_physiology%23Functions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemoglobin%23Structure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- cooperative binding. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cooperative%20binding**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...on/bohr-effect**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- allosteric. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/allosteric**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Erythrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Erythrocyte%23Mammalian_erythrocytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemoglobin**. **License**: ***Public Domain: No Known Copyright***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Erythrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Erythrocyte%23Life_cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- macrophage. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/macrophage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phagocytosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/phagocytosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- xenobiotic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/xenobiotic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Erythrocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Erythrocyte%23Mammalian_erythrocytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemoglobin**. **License**: ***Public Domain: No Known Copyright***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood%23Oxygen_transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Red_White_Blood_cells.jpg. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/24/Red_White_Blood_cells.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 16.2C: RBC Life Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.3: WHITE BLOOD CELLS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

16.3A: Types of WBCs

16.3B: WBC Function

16.3C: WBC Formation

This page titled 16.3: White Blood Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.3A: Types of WBCs

The different types of white blood cells (leukocytes) include neutrophils, basophils, eosinophils, lymphocytes, monocytes, and macrophages.

Learning Objectives

- Distinguish between the two major types of leukocytes (white blood cells): granulocytes and agranulocytes

Key Points

- The two main types of leukocytes are granulocytes and mononuclear leukocytes (agranulocytes).
- Leukocytes arise from hemopoietic stem cells in the bone marrow.
- Leukocytes are involved in pathogen recognition, phagocytosis (ingestion of particles), pathogen destruction, inflammation mediation, and antigen presentation.
- Granulocytes include neutrophils, basophils, eosinophils, and mast cells. Their granules contain enzymes that damage or digest pathogens and release inflammatory mediators into the bloodstream.
- Mononuclear leukocytes include lymphocytes, monocytes, macrophages, and dendritic cells. This group is involved in both innate and adaptive immune system function.


Key Terms

- **endocytosed**: Engulfed during the process by which the plasma membrane of a cell folds inwards to ingest material.
- **antigen**: A substance, usually foreign, that induces an immune response.
- **pathogen**: Any organism or substance, especially a microorganism, capable of causing disease. Examples include bacteria, viruses, protozoa, or fungi. Microorganisms are not considered pathogenic until the population has grown large enough to cause disease.

White blood cells (WBCs), or leukocytes, are immune system cells that defend the body against infectious disease and foreign materials. There are several different types of WBCs. They share commonalities but are distinct in form and function. WBCs are produced in the bone marrow by hemopoeitic stem cells, which differentiate into either lymphoid or myeloid progenitor cells. A major distinguishing feature is the presence of granules; white blood cells are often characterized as granulocytes or agranulocytes.

Granulocytes

Granulocytes, also known as polymorphonuclear (PMN) leukocytes, are characterized by stained granules within their cytoplasm under a microscope. These granules are membrane-bound enzymes that act primarily in the digestion of endocytosed particles. They may also cause granule dependent cell-mediated apoptosis through the release of perforins, granzymes, and proteases. The nucleus contains multiple lobes (polymorphonuclear) as opposed to a single rounded lobe. Granulocytes contain toll-like receptors that allow them to recognize pathogen-associated molecular patterns (PAMPS). All categories except neutrophils contain IgE receptors that implicate them in allergic responses. There are four types of granulocytes:



Granulocytes: From left to right, a neutrophil, an eosinophil, and a basophil.

- Neutrophils defend against bacterial or fungal infection and other very small inflammatory processes. They are usually the first responders to microbial infection. Their activity and death in large numbers from degranulation forms purulent necrosis (pus).
- Eosinophils primarily deal with parasitic infections. They are also the predominant inflammatory cells in allergic reactions.
- Basophils are chiefly responsible for short-term inflammatory response (particularly from allergy or irritation) by releasing the chemical histamine, which causes the vasodilation that occurs with inflammation.
- Mast cells function similarly to basophils in that they often mediate inflammation, but are more common and arise from a different hemopoeitic lineage.

Mononuclear Leukocytes

Mononuclear (MN) leukocytes are characterized by a single round nucleus within the cytoplasm. Some MN leukocytes contain granules while others do not, but the members of this group are sometimes considered agranulocytes by naming convention. MN leukocytes contain lysosomes, small vesicles containing digestive enzymes that break down foreign matter that is endocytosed by the cell during phagocytosis. The cells include:

- Lymphocytes, which come in three types. B-lymphocytes produce antibodies in the humoral immune response. T-lymphocytes participate in the cell-mediated immune response. NK cells are cytotoxic cells that participate in the innate immune response by killing virally infected and tumor cells and mediating fever and long-lasting inflammation. B and T lymphocytes contain MHC antigen receptors and their activity is antigen-specific. Other leukocytes will attack any pathogen but cannot distinguish between different types of pathogens.
- Monocytes are large leukocytes that differentiate into macrophages and dendritic cells under varying conditions, while performing similar functions in phagocytosis and antigen presentation (the process by which molecular components are presented to lymphocytes to stimulate an adaptive immune response). Monocytes and their progeny contain toll-like receptors and granules.
- Macrophages are monocytes that have migrated out of the blood stream and into the internal body tissues. They destroy necrotic cell debris and foreign material including viruses and bacteria, and can present antigens to naive lymphocytes. They typically arrive at the site of inflammation one to three days after the initial neutrophil response to clean up dead neutrophils, cellular debris, and remaining pathogens.
- Dendritic cells are monocytes that have migrated to cells that are in contact with the external environment, such as the skin, intestines, or respiratory epithelium. Their name comes branched projections called dendrites, which increase their surface area. They phagocytize pathogens and present antigens to naive lymphocytes.



A Macrophage: A macrophage phagocytizes two smaller particles, possibly pathogens

This page titled 16.3A: Types of WBCs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.3B: WBC Function

Each type of white blood cell (WBC) has a specific function in defending the body against infections.

Learning Objectives

- Describe the functions of leukocytes (white blood cells)

Key Points

- Leukocyte functions often occur in the bloodstream and may represent either the innate or adaptive immune systems.
- Innate immune system functions are non-specific and include phagocytosis, inflammation, and degranulation.
- Adaptive immune system functions are antigen -specific and involve antigen presentation as well as cell -mediated and humoral -mediated activities.
- Compared to innate immune system functions, adaptive immune system functions take more time to initiate, but work much faster. They have a memory component to prevent reinfection by the same pathogen.

Key Terms


- **macrophage**: A white blood cell that phagocytizes necrotic cell debris and foreign material, including viruses, bacteria, and tattoo ink. It presents foreign antigens on MHC II molecules to lymphocytes. Part of the innate immune system.
- **Inflammation**: An innate immune system function in response to a pathogen or injury. Chemical mediators cause the blood vessels to dilate and become more permeable, which draws neutrophils to the area.
- **cytotoxic**: Any mechanism that can cause the death of a cell (typically without phagocytosis), such as degranulation or cell mediated apoptosis.

Leukocytes (white blood cells) provide a number of functions that are primarily related to defending the body from pathogens (foreign invaders). Much leukocyte activity takes place within the bloodstream, but is not restricted to this area. Many leukocytes are able to perform their functions in tissues or organs during normal transport and in response to injury. Leukocyte functions may be classified as either innate or adaptive based on several characteristics.

Innate Immune System Functions

The innate immune system refers to the body's ability to prevent pathogen entry and destroy pathogens that do enter the body. Its functions are rapid responses that inhibit a pathogen as soon as it is detected in the body. Innate immune system functions involving leukocytes include:

- Phagocytosis of pathogens. This process is performed primarily by neutrophils, macrophages, and dendritic cells, but most other leukocytes can do it as well. It involves the binding of an Fc receptor to a tail on a pathogen. The pathogen is engulfed by the leukocyte and destroyed with enzymes and free radicals.
- Inflammation. This process is performed primarily by mast cells, eosinophils, basophils, and NK cells. When a pathogen is detected or vascular endothelial cells release stress cytokines from injury such as a cut, leukocytes release a variety of inflammatory cytokines such as histamine or TNF-alpha. These cause vasodilation, increase vascular permeability, and promote neutrophil movement to the inflammation site.
- Degranulation. This process is performed by granulocytes like neutrophils. When pathogens are encountered, granule-dependent apoptosis (a mechanism of cytotoxicity) may be induced in the pathogen by releasing perforins, granzymes, and proteases from their granules.



Neutrophils Phagocytizing Bacteria: Here, neutrophils are depicted phagocytizing and completely engulfing bacteria.

Adaptive Immune System Functions

The adaptive immune system is specific to each pathogen on the basis of antigens, molecular components of pathogens used by leukocytes to recognize that specific pathogen. Compared to the innate immune system, adaptive immune functions work much faster and have a memory component that prevents reinfection by the same pathogen. However, more time typically passes before the adpative immune system is functional. Adaptive immune functions of leukocytes include:

- Antigen presentation. This process is primarily performed by macrophages and dendritic cells. Following phagocytosis, protein components (antigens) of the pathogen are expressed on leukocyte MHC molecules and presented to naive T cells (and B cells) in the lymph nodes. The T cells will then start the adaptive immune response by rapidly proliferating and differentiating.
- Cell-mediated activities. This process is performed by T cells. Pathogens that bear the T cell's antigen are destroyed through cytotoxic -induced apoptosis and protease activity.
- Humoral activities. This process is performed by B cells, which secrete antigen-specific antibodies. The antibodies bind to pathogens to opsonize (mark) them for phagocytes to engulf, neutralize, or start a complement cascade in which proteins form a membrane attack complex to lyse the pathogen.
- Memory cell activity. Following antigen presentation, memory B and T cells are created. These rapidly produce new T cells or antibodies if the same pathogen is detected in the future. This prevents that pathogen from reinfecting the organism.

This page titled 16.3B: WBC Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.3C: WBC Formation

Haematopoiesis refers to the formation of blood cells components. It is necessary for vertebrate function.

Learning Objectives

- Describe the formation of leukocytes (white blood cells, or WBCs)

Key Points

- Haematopoietic stem cells are self-renewing and reside in the medulla of the bone (bone marrow).
- All blood cells are divided into two main lineages, produced through lymphoid progenitor cells or myeloid progenitor cells depending on lineage type.
- Lymphoid progenitor cells differentiate into B and T cells and NK cells.
- Myeloid progenitor cells differentiate into myelocytes (granulocytes and monocytes) or non-leukocytes such as erythorocytes and megakaryocytes (which produce platelets).
- Before birth, most blood cell formation occurs in the liver or spleen, which tend to enlarge when used for hematopoiesis. In adults, most blood production occurs in the bone marrow.

Key Terms

- **myelocyte**: A large cell found in bone marrow that becomes a granulocyte or monocyte when mature.
- **differentiation**: The gradual changes that occur when a cell or tissue type changes into a different type. Cells generally become more specialized the more they differentiate, and are considered to be terminally differentiated when they cannot differentiate (and often cannot divide) any further.
- **megakaryocyte**: A large cell found in bone marrow, responsible for the production of platelets.

Haematopoiesis refers to the formation of blood cellular components, including both white and red blood cells. All cellular blood components are derived from haematopoietic stem cells located within the bone marrow. In a healthy adult, approximately 10^{11}–10^{12}new blood cells are produced daily to maintain equilibrium levels in peripheral circulation.

Leukocyte Haematopoiesis


Haematopoietic stem cells (HSCs) reside in the bone marrow and have the unique ability to give rise to all mature blood cell types through differentiation into other progenitor cells. HSCs are self-renewing. When they proliferate, at least some daughter cells remain HSCs, so the pool of stem cells does not become depleted over time. The daughters are the myeloid and lymphoid progenitor cells, which cannot self renew but differentiate into various myeloid leukocytes and lymphocytes respectively. This is one of the body's vital processes.

Leukocyte Lineages

Two different leukocyte lineages and two non-leukocyte lineages arise from the progeny of HSCs. Following this split in differentiation, the subtypes undergo eventual differentiation into terminally-differentiated leukocytes, which typically do not divide independently.

1. The lymphocyte lineage derives from common lymphoid progenitor cells, which in turn become lymphoblasts before differentiating into T cells, B cells, and NK cells.
2. Myelocytes are an offshoot of common myeloid progenitor cells, which also differentiate into the erythropoietic and magakaryotic progenitors. This diverse group differentiates into granulocytes and monocytes. Monocytes further differentiate into macrophages or dendritic cells upon reaching certain tissues.
3. Megakaryocytes (the cells that produce platelets) and erythrocytes (red blood cells) are not formally considered to be

leukocytes, but arise from the common myeloid progenitor cells that produce the other cellular components of blood.



Hematopoiesis in Humans: This diagram shows hematopoiesis as it occurs in humans.

Sites of Haematopoesis in Pre- and Postnatal Periods

In developing embryos, blood formation occurs in aggregates of blood cells in the yolk sac called blood islands. However, most of blood supply comes from the mother through the placenta. As development progresses, blood formation occurs primarily in the spleen, liver, and lymph nodes.

When bone marrow develops, it eventually assumes the task of forming most of the blood cells for the entire organism. However, maturation, activation, and some proliferation of lymphoid cells occurs in lymphoid organs (spleen, thymus, and lymph nodes). In children, haematopoiesis occurs in the marrow of the long bones such as the femur and tibia. In adults, it occurs mainly in the pelvis, cranium, vertebrae, and sternum.

In some cases, the liver, thymus, and spleen may resume their haematopoietic function if necessary. This is called extramedullary haematopoiesis. It may cause these organs to hypertrophy and increase in size substantially. During fetal development, the liver functions as the main haematopoetic organ since bones and marrow develop later. Therefore, the liver is enlarged during development relative to its mature proportions.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Leukocytes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Leukocytes%23Types**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endocytosed. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endocytosed**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pathogen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pathogen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- antigen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/antigen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- White blood cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/White_blood_cell**. **License**:

- Macrophage. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Macrophage.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Leukocytes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Leukocytes. **License**: *CC BY-SA: Attribution-ShareAlike*
- granulocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/granulocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- macrophage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/macrophage. **License**: *CC BY-SA: Attribution-ShareAlike*
- White blood cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/White_blood_cell. **License**:
- Macrophage. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Macrophage.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neutrophils phagocytizing bacteria. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...g_bacteria.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Haematopoiesis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Haematopoiesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//management...ifferentiation. **License**: *CC BY-SA: Attribution-ShareAlike*
- megakaryocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/megakaryocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- myelocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/myelocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- White blood cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/White_blood_cell. **License**:
- Macrophage. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Macrophage.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neutrophils phagocytizing bacteria. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...g_bacteria.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hematopoiesis (human) diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:He...n)_diagram.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 16.3C: WBC Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.4: PLATELETS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

16.4A: Platelets

16.4B: Platelet Formation

This page titled 16.4: Platelets is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.4A: Platelets

Platelets, also called thrombocytes, are membrane-bound cell fragments that are essential for clot formation during wound healing.

Learning Objectives

- Discuss the roles played by platelets in the blood

Key Points

- Platelets, also called thrombocytes, are derived from megakaryocytes, which are derived from stem cells in the bone marrow.
- Platelets circulate in the blood and are involved in hemostasis, leading to the formation of blood clots and blood coagulation.
- Platelets lack a nucleus, but do contain some organelles, such as mitochondria and endoplasmic reticulum fragments.
- If the number of platelets in the blood is too low, excessive bleeding can occur. However, if the number of platelets is too high, blood clots can form (thrombosis), which may obstruct blood vessels.
- Platelets are a natural source of growth factors involved in wound healing, coagulatants, and inflammatory mediators.


Key Terms

- **extracellular matrix**: All the connective tissues and fibers that are not part of a cell, but rather provide support.
- **platelet**: A small, colorless, disc-shaped particle found in the blood of mammals. It plays an important role in the formation of blood clots.

Platelets, also called thrombocytes, are membrane-bound cell fragments derived from the fragmentation of larger precursor cells called megakaryocytes, which are derived from stem cells in the bone marrow. Platelets are important for the blood clotting process, making them essential for wound healing.

Platelet Structure and Distribution

Platelets are irregularly shaped, have no nucleus, and typically measure only 2–3 micrometers in diameter. Platelets are not true cells, but are instead classified as cell fragments produced by megakaryocytes. Because they lack a nucleus, they do not contain nuclear DNA. However, they do contain mitochondria and mitochondrial DNA, as well as endoplasmic reticulum fragments and granules from the megakaryocyte parent cells. Platelets also contain adhesive proteins that allow them to adhere to fibrin mesh and the vascular endothelium, as well as to a microtubule and microfilament skeleton that extends into filaments during platelet activation. Less than 1% of whole blood consists of platelets. They are about $1/10^{th}$ to $1/20^{th}$ as abundant as white blood cells.




Platelet: Image from a light microscope (40×) from a peripheral blood smear surrounded by red blood cells. One platelet can be seen in the upper left side of the image (purple) and is significantly smaller in size than the red blood cells (stained pink) and the two large neutrophils (stained purple).

Platelet Functions

Platelets circulate in blood plasma and are primarily involved in hemostasis (stopping the flow of blood during injury), by causing the formation of blood clots, also known as coagulation. The adhesive surface proteins of platelets allow them to accumulate on the fibrin mesh at an injury site to form a platelet plug that clots the blood. The complex process of wound repair can only begin once the clot has stopped bleeding.

Platelets secrete many factors involved in coagulation and wound healing. During coagulation, they release factors that increase local platelet aggregation (thromboxane A), mediate inflammation (serotonin), and promote blood coagulation through increasing thrombin and fibrin (thromboplastin). They also release wound healing-associated growth factors including platelet-derived growth factor (PDGF), which directs cell movement; TGF beta, which stimulates the deposition of extracellular matrix tissue into a wound during healing; and vascular endothelial growth factor (VEGF), which stimulates angiogenesis, or the regrowth of blood vessels. These growth factors play a significant role in the repair and regeneration of connective tissues. Local application of these platelet-produced healing-associated factors in increased concentrations has been used as an adjunct to wound healing for several decades.



Platelets: A blood slide of platelets aggregating, or, clumping together. The platelets are the small, bright purple fragments.

If the number of platelets is too low, excessive bleeding can occur and wound healing will be impaired. However, if the number of platelets is too high, blood clots can form (thrombosis), which may obstruct blood vessels and result in ischemic tissue damage caused by a stroke, myocardial infarction, pulmonary embolism, or the blockage of blood vessels to other parts of the body. Thrombosis also occurs when blood is allowed to pool, which causes clotting factors and platelets to form a blood clot even in the absence of an injury.

This page titled 16.4A: Platelets is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.4B: Platelet Formation

Platelets are membrane-bound cell fragments derived from megakaryocytes, which are produced during thrombopoiesis.

Learning Objectives

- Describe the process of platelet formation

Key Points

- Megakaryocytes are produced from stem cells in the bone marrow by a process called thrombopoiesis.
- Megaryocytes create platelets by releasing protoplatelets that break up into numerous smaller, functional platelets.
- Thrombopoiesis is stimulated and regulated by the hormone thrombopoietin.
- Platelets have an average life span of five to ten days.
- Old platelets are destroyed by phagocytosis. The spleen holds a reservoir of additional platelets.
- Abnormal numbers of platelets result from problems in thrombopoiesis. This is associated with various disorders and potential pathological complications.

Key Terms

- **megakaryocyte**: A large cell found in bone marrow that is responsible for the production of platelets.
- **Thrombopotein**: A hormone produced by the liver or kidneys that stimulates megakaryocyte differentiation and platelet release.
- **bone marrow**: The fatty vascular tissue that fills the cavities of bones; the place where new blood cells are produced.


Platelets are small, clear, irregularly-shaped cell fragments produced by larger precursor cells called megakaryocytes. Platelets are also called thrombocytes because they are involved in the blood clotting process, which is necessary for wound healing. Platelets are continuously produced as a component product of hematopoiesis (blood cell formation).

Thrombopoiesis

Platelets are produced during hematopoiesis in a sub-process called thromopoiesis, or production of thrombocytes. Thrombopoiesis occurs from common myeloid progenitor cells in the bone marrow, which differentiate into promegakaryocytes and then into megakaryocytes. Megakaryocytes stay in the bone marrow and are thought to produce protoplatelets within their cytoplasm, which are released in cytoplasmic extensions upon cytokine stimulus. The protoplatelets then break up into hundreds of platelets that circulate throughout the bloodstream, while the remaining nucleus of the ruptured megakaryocyte is consumed by macrophages.

Megakaryocyte and platelet production is regulated by thrombopoietin, a hormone produced by the liver and kidneys. Thrombopoietin stimulates differentiation of myeloid progenitor cells into megakaryocytes and causes the release of platelets. Thrombopoietin is regulated by a negative feedback mechanism based on platelet levels in the body so that high levels of platelets result in lower levels of thrombopoietin, while low levels of platelets result in higher levels of thrombopoietin.

Each megakaryocyte produces between 5,000 and 10,000 platelets before its cellular components are fully depleted. Altogether, around 10^11 platelets are produced each day in a healthy adult. The average lifespan of a platelet is just 5 to 10 days. Old platelets are destroyed by macrophage phagocytosis in the spleen and by Kupffer cells in the liver. Up to 40% of platelets are stored in the spleen as a reserve, released when needed by sympathetically-induced splenic muscle contractions during severe injury.



Hematopoiesis: Myeloid progenitor cells differentiate into promegakaryocytes, and megakaryocytes, which release platelets.

Implications of Platelet Formation

Balanced thrombopoiesis is important because it directly influences the amount of platelets in the body and their associated complications. If the number of platelets is too low, excessive bleeding can occur, even from minor or superficial injuries. If the number of platelets is too high, blood clots can form (thrombosis) and travel through the bloodstream, which may obstruct blood vessels and result in ischemic events. These include stroke, myocardial infarction, pulmonary embolism, or infarction of other tissues.

An abnormality or disease of the platelets is called a thrombocytopathy, which could be either a low number of platelets (thrombocytopenia), a decrease in function of platelets (thrombasthenia), or an increase in the number of platelets (thrombocytosis). In any case, issues with the number of circulating platelets is often due to issues in thrombopoietin feedback regulation, but may also be associated with genetic characteristics and certain medications and diseases. For example, thrombocytopenia often occurs in leukemia patents. Cancerous myeloid cells crowd out healthy ones in the bone marrow, causing impaired thrombopoiesis.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- platelet. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/platelet**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...gy%23Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extracellular matrix. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/extracellular_matrix**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Platelets.jpg**. **License**: ***CC BY: Attribution***

- bone marrow. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/bone_marrow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelet. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelet**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...gy%23Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- megakaryocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/megakaryocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- stem cell. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/stem_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Platelets.jpg**. **License**: ***CC BY: Attribution***
- Platelets. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Platelets**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 16.4B: Platelet Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5: HEMOSTASIS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

16.5A: Overview of Hemostasis

16.5B: Vascular Spasm

16.5C: Platelet Plug Formation

16.5D: Coagulation

16.5E: Role of Vitamin K

16.5F: Clot Retraction and Repair

16.5G: Fibrinolysis

This page titled 16.5: Hemostasis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5A: Overview of Hemostasis

Hemostasis is the natural process that stops blood loss when an injury occurs.

Learning Objectives

- Explain the steps involved in hemostasis

Key Points

- Hemostasis is the natural process that stops blood loss when an injury occurs.It involves three steps: (1) vascular spasm (vasoconstriction); (2) platelet plug formation; and (3) coagulation.
- Vasoconstriction is a reflex in which blood vessels narrow to increase blood pressure.
- Next, platelet plug formation involves the activation, aggregation, and adherence of platelets into a plug that serves as a barrier against blood flow.
- Coagulation involves a complex cascade in which a fibrin mesh is cleaved from fibrinogen.
- Fibrin acts as a “molecular glue” during clot formation, holding the platelet plug together.

Key Terms

- **hemostasis**: The process of slowing and stopping the flow of blood to initiate wound healing.
- **coagulation**: The process by which blood forms gelatinous clots.
- **heparin**: A fibrinolytic molecule expressed on endothelial cells or produced as a blood thinner medicine. It prevents activation of platelets and clotting factors.

Hemostasis is the natural process in which blood flow slows and a clot forms to prevent blood loss during an injury, with hemo- meaning blood, and stasis meaning stopping. During hemostasis, blood changes from a fluid liquid to a gelatinous state.

Steps of Hemostasis

Hemostasis includes three steps that occur in a rapid sequence: (1) vascular spasm, or vasoconstriction, a brief and intense contraction of blood vessels; (2) formation of a platelet plug; and (3) blood clotting or coagulation, which reinforces the platelet plug with fibrin mesh that acts as a glue to hold the clot together. Once blood flow has ceased, tissue repair can begin.



Angiogenesis Generates New Blood Vessels: Blood vessel with an erythrocyte (red blood cell) within its lumen, endothelial cells forming its tunica intima or inner layer, and pericytes forming its tunica adventitia (outer layer).

Vasoconstriction

Intact blood vessels are central to moderating blood's clotting tendency. The endothelial cells of intact vessels prevent clotting by expressing a fibrinolytic heparin molecule and thrombomodulin, which prevents platelet aggregation and stops the coagulation cascade with nitric oxide and prostacyclin. When endothelial injury occurs, the endothelial cells stop secretion of coagulation and aggregation inhibitors and instead secrete von Willebrand factor, which causes platelet adherence during the initial formation of a clot. The vasoconstriction that occurs during hemostasis is a brief reflexive contraction that causes a decrease in blood flow to the area.

Platelet Plug Formation

Platelets create the "platelet plug" that forms almost directly after a blood vessel has been ruptured. Within twenty seconds of an injury in which the blood vessel's epithelial wall is disrupted, coagulation is initiated. It takes approximately sixty seconds until the first fibrin strands begin to intersperse among the wound. After several minutes, the platelet plug is completely formed by fibrin.

Contrary to popular belief, clotting of a skin injury is not caused by exposure to air, but by platelets adhering to and being activated by collagen in the blood vessels' endothelium. The activated platelets then release the contents of their granules, which contain a variety of substances that stimulate further platelet activation and enhance the hemostatic process.

When the lining of a blood vessel breaks and endothelial cells are damaged, revealing subendothelial collagen proteins from the extracellular matrix, thromboxane causes platelets to swell, grow filaments, and start clumping together, or aggregating. Von Willebrand factor causes them to adhere to each other and the walls of the vessel. This continues as more platelets congregate and undergo these same transformations. This process results in a platelet plug that seals the injured area. If the injury is small, the platelet plug may be able to form within several seconds.

Coagulation Cascade

If the platelet plug is not enough to stop the bleeding, the third stage of hemostasis begins: the formation of a blood clot. Platelets contain secretory granules. When they stick to the proteins in the vessel walls, they degranulate, thus releasing their products, which include ADP (adenosine diphosphate), serotonin, and thromboxane A2 (which activates other platelets).

First, blood changes from a liquid to a gel. At least 12 substances called clotting factors or tissue factors take part in a cascade of

chemical reactions that eventually create a mesh of fibrin within the blood. Each of the clotting factors has a very specific function. Prothrombin, thrombin, and fibrinogen are the main factors involved in the outcome of the coagulation cascade. Prothrombin and fibrinogen are proteins that are produced and deposited in the blood by the liver.

When blood vessels are damaged, vessels and nearby platelets are stimulated to release a substance called prothrombin activator, which in turn activates the conversion of prothrombin, a plasma protein, into an enzyme called thrombin. This reaction requires calcium ions. Thrombin facilitates the conversion of a soluble plasma protein called fibrinogen into long, insoluble fibers or threads of the protein, fibrin. Fibrin threads wind around the platelet plug at the damaged area of the blood vessel, forming an interlocking network of fibers and a framework for the clot. This net of fibers traps and helps hold platelets, blood cells, and other molecules tight to the site of injury, functioning as the initial clot. This temporary fibrin clot can form in less than a minute and slows blood flow before platelets attach.

Next, platelets in the clot begin to shrink, tightening the clot and drawing together the vessel walls to initiate the process of wound healing. Usually, the whole process of clot formation and tightening takes less than a half hour.



Vasoconstriction: Microvessel showing an erythrocyte (E), a tunica intima of endothelial cells, and a tunica adventitia of pericytes.

This page titled 16.5A: Overview of Hemostasis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5B: Vascular Spasm

Vasoconstriction is the narrowing of the blood vessels, which reduces blood loss during injury.

Learning Objectives

- Describe vascular spasms in hemostasis

Key Points

- Vasoconstriction is the narrowing of the blood vessels, which increases blood pressure but can decrease blood flow and loss.
- Vasoconstriction is mediated by contraction of the smooth muscles lining a blood vessel.
- Vasoconstriction is caused by thromboxane A_2 from activated platelets and injured epithelial cells, nervous system reflexes from pain, and direct injury to vascular smooth muscle.
- Vasopressins are drugs that may induce vasoconstriction and increase blood pressure.
- Vasonstriction only lasts for a few minutes during hemostasis. During inflammation that follows the injury, it is replaced by vasodilation as the healing process begins.

Key Terms

- **endothelial cells**: The endothelium comprises the thin layer of endothelial cells that lines the interior surface of blood and lymphatic vessels, forming an interface between circulating blood or lymph in the lumen and the rest of the vessel wall.
- **vascular**: Of, pertaining to, or containing vessels that conduct or circulate fluids such as blood, lymph, or sap through the body of an animal or plant.
- **inflammation**: A process that occurs during injury and generally follows hemostasis in which vasoconstriction ends and vasodilation begins.

Vasoconstriction is the narrowing of the blood vessels resulting from contraction of the smooth muscle wall of the vessels, particularly in the large arteries and small arterioles. The process is the opposite of vasodilation, the dilation and expansion of blood vessels. During hemostasis, a brief spasm of vasoconstriction occurs, which slows blood flow into the injured area while the clot forms.



Vasoconstriction during hemostasis: Blood vessel experiencing vasoconstriction as its smooth muscle contracts while the blood clot forms.

Mechanisms of Vasoconstriction

The vasoconstriction response is triggered by factors such as a direct injury to vascular smooth muscle, signaling molecules released by injured endothelial cells and activated platelets (such as thromboxane A_2), and nervous system reflexes initiated by local pain receptors. The spasm response becomes more effective as the amount of damage is increased. Vascular spasm is much more effective at slowing the flow of blood in smaller blood vessels. Vasoconstriction also causes an increase in blood pressure for affected blood vessels.

Smooth muscle in the vessel wall goes through intense contractions that constrict the vessel. If the vessels are small, spasms compress the inner walls together and may be able to stop the bleeding completely. If the vessels are medium to large-sized, the spasms slow down immediate outflow of blood, lessening the damage but still preparing the vessel for the later steps of hemostasis. The spasm response becomes stronger and lasts longer in more severe injuries. Vasoconstriction may be induced by drugs called vasopressins, which increase blood pressure and can help treat certain conditions.

Injury and Inflammation

During injury, vasoconstriction is brief, lasting only a few minutes while the platelet plug and coagulation cascade occur. This is because as tissues are damaged during an injury, inflammation occurs as a result of inflammatory mediator release from immune system cells (such as mast cells or NK cells) that receive cell stress cytokines from damaged enothelial cells or vasoactive amines (serotonin) that are secreted by activated platelets. During inflammation, vasodilation occur, along with increased vascular permeability and leukocyte chemotaxis, ending the spasm of vasoconstriction and hemostasis as wound healing begins.

This page titled 16.5B: Vascular Spasm is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5C: Platelet Plug Formation

At the site of vessel injury, platelets stick together to create a plug, which is the beginning of blood clot formation.

Learning Objectives

- Describe the formation of a platelet plug

Key Points


- Platelets adhere to the damaged endothelium to form a platelet plug, temporarily sealing the break in the vessel wall.
- Activated platelets release factors to stimulate further platelet activation, perpetuating plug formation in a positive feedback loop, while other factors stimulate the coagulation cascade and maintain vasoconstriction.
- Platelets adhere to the collagen fibers in the vessel wall
  by becoming adhesive and filamentous
  due to the stimulus of von Willebrand factor.
- During platelet aggregation, platelets bind to von Willebrand factor and fibrinogen to stick together and seal the break in the endothelium.

Key Terms

- **von Willebrand Factor**: The factor responsible for causing platelet adherence and aggregation. It is increased by positive feedback during platelet activation.
- **collagen**: A glycoprotein that forms elongated fibers, usually found in the extracellular matrix of connective tissue such as the matrix beneath the vascular endothelium.

The second critical step in hemostasis, which follows vasoconstriction, is platelet plug formation. The three steps to platelet plug formation are platelet adherence, activation, and aggregation.

Platelet Adherence




Platelets: A blood slide of platelets aggregating or clumping together. The platelets are small, bright purple fragments.

Normally, the endothelial cells express molecules that inhibit platelet adherence and activation while platelets circulate through the blood vessels. These molecules include nitric oxide, prostacylcine (PGI_2) and endothelial ADP-ase.

During an injury, subendothelial collagen from the extracellular matrix beneath the endothelial cells is exposed on the epithelium as the normal epithelial cells are damaged and removed, which releases von Willebrand Factor (VWF). VWF causes the platelets to change form with adhesive filaments (extensions) that adhere to the subendothelial collagen on the endothelial wall.

Platelet Activation

After platelet adherence occurs, the subendothelial collagen binds to receptors on the platelet, which activates it. During platelet activation, the platelet releases a number of important cytokines and chemical mediators via degranulation. The released chemicals include ADP, VWF, thromboxane A2, platelet-derived growth factor (PDGF), vascular endothelial growth factor (VEGF), serotonin, and coagulation factors. The extra ADP and VWF is especially important because it causes nearby platelets to adhere and activate, as well as release more ADP, VWF, and other chemicals. Platelet plug formation is considered a positive feedback process because ADP and VWF levels are successively increased as more and more platelets activate to form the plug.



ADP: The chemical structure of ADP, a molecule that causes platelet activation and is involved in the positive feedback component of platelet activation.

The other factors released during platelet activation perform other important functions. Thromboxane is an arachidonic acid derivative (similar to prostaglandins) that activates other platelets and maintains vasoconstriction. Serotonin is a short-lived inflammatory mediator with a vasoconstrictive effect that contributes to vascular changes associated with inflammation during an injury. PDGF and VEGF are involved in angiogenesis, the growth of new blood vessels and cell cycle proliferation (division) following injury. The coagulation factors include factor V and VIII, which are involved in the coagulation cascade that converts fibrinogen into fibrin mesh after platelet plug formation.

Platelet Aggregation

The final step of platelet plug formation is aggregation of the platelets into a barrier-like plug. Receptors on the platelet bind to VWF and fibrinogen molecules, which hold the platelets together. Platelets may also bind to subendothelial VWF to anchor them to the damaged endothelium. The completed plug will cover the damaged components of the endothelium and will stop blood from flowing out of it, but if the wound is large enough, blood will not coagulate until the fibrin mesh from the coagulation cascade is produced, which strengthens the platelet plug. If the wound is minor, the platelet plug may be enough to stop the bleeding without the coagulation cascade.

This page titled 16.5C: Platelet Plug Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5D: Coagulation

Coagulation is the process by which a blood clot forms to reduce blood loss after damage to a blood vessel.

Learning Objectives

- Outline the process of coagulation in secondary hemostasis


Key Points

- The coagulation cascade is a series of reactions, which is classically divided into three pathways: the contact (also known as the intrinsic) pathway, the tissue factor (also known as the extrinsic pathway), and the common pathway.
- The intrinsic pathway occurs when negatively charged molecule contact causes a cascade of factors that produce factor X. The extrinsic pathway occurs when tissue damage causes the release of tissue factor, creating a smaller cascade that produces factor X. The common pathway merges both pathways as factor X is used to create thrombin from prothrombin.
- Secondary hemostasis involves factors of the coagulation cascade, which collectively strengthen the platelet plug.
- Coagulation can be harmful if blood clots embolize and obstruct other blood vessels. Clots can also occur if blood pools from prolonged immobility.
- A number of anticoagulants exist to inhibit various parts of the coagulation cascade, inactivate thrombin, or degrade fibrin directly.

Key Terms

- **fibrin**: An elastic, insoluble, whitish protein produced by the action of thrombin on fibrinogen and forming an interlacing fibrous network in the coagulation of blood.
- **endothelium**: A thin layer of flat epithelial cells that lines the heart, serous cavities, lymph vessels, and blood vessels.
- **thrombin**: The end product of the coagulation cascade, which cleaves fibrin from fibrinogen.


Coagulation is the process by which a blood clot forms to reduce blood loss after damage to a blood vessel. Several components of the coagulation cascade, including both cellular (e.g. platelets) and protein (e.g. fibrin) components, are involved in blood vessel repair. The role of the cellular and protein components can be categorized as primary hemostasis (the platelet plug) and secondary hemostasis (the coagulation cascade). The coagulation cascade is classically divided into three pathways: the contact (also known as the intrinsic) pathway, the tissue factor (also known as the extrinsic pathway), and the common pathway. Both the contact pathway and the tissue factor feed into and activate the common pathway.



Coagulation Pathway

Secondary Hemostasis

Hemostasis can either be primary or secondary. Primary hemostasis refers to platelet plug formation, which forms the primary clot. Secondary hemostasis refers to the coagulation cascade, which produces a fibrin mesh to strengthen the platelet plug. Secondary hemostasis occurs simultaneously with primary hemostasis, but generally finishes after it. The coagulation factors circulate as inactive enzyme precursors, which, upon activation, take part in the series of reactions that make up the coagulation cascade. The coagulation factors are generally serine proteases (enzymes).



Coagulation Cascade

Intrinsic Pathway

The intrinsic pathway (contact activation pathway) occurs during exposure to negatively charged molecules, such as molecules on bacteria and various types of lipids. It begins with formation of the primary complex on collagen by high-molecular-weight kininogen (HMWK), prekallikrein, and factor XII (Hageman factor). This initiates a cascade in which factor XII is activated, which

then activates factor XI, which activated factor IX, which along with factor VIII activates factor X in the common pathway.

Extrinsic Pathway

The main role of the extrinsic (tissue factor) pathway is to generate a “thrombin burst,” a process by which large amounts of thrombin, the final component that cleaves fibrinogen into fibrin, is released instantly. The extrinsic pathway occurs during tissue damage when damaged cells release tissue factor III. Tissue factor III acts on tissue factor VII in circulation and feeds into the final step of the common pathway, in which factor X causes thrombin to be created from prothrombin.

Common Pathway

In the final common pathway, prothrombin is converted to thrombin. When factor X is activated by either the intrinsic or extrinsic pathways, it activates prothrombin (also called factor II) and converts it into thrombin using factor V. Thrombin then cleaves fibrinogen into fibrin, which forms the mesh that binds to and strengthens the platelet plug, finishing coagulation and thus hemostasis. It also activates more factor V, which later acts as an anticoagulant with inhibitor protein C, and factor XIII, which covalently bonds to fibrin to strengthen its attachment to the platelets.

Coagulation Problems

While the coagulation cascade is critical for hemostasis and wound healing, it can also cause problems. An embolism is any thrombosis (blood clot) that breaks off without being dissolved and travels through the bloodstream to another site. If it obstructs an artery that supplies blood to a tissue or organ, it can cause ischemia and infarcation to those tissues, leading to a pulmonary embolism, stroke, or heart attack).

Coagulation can occur even without injury, as blood pooling from prolonged immobility can cause clotting factors to accumulate and activate a coagulation cascade independently. Additionally, endothelial damage caused by immune system factors like inflammation or hypersensitivity may also cause unnecessary thrombosis and embolism. For example, during severe bacterial infections (septic shock), inflammation-induced tissue damage and the negatively charged molecules of bacteria activate both pathways of the coagulation cascade and cause disseminated intravascular coagulation (DIC), in which many clots form and break off, leading to massive organ failure.

Anticoagulants

Many anticoagulants prevent unnecessary coagulation, and those that genetically lack the ability to produce these molecules will be more susceptible to coagulation. These mechanisms include:

1. Protein C: a vitamin K-dependent serine protease enzyme that degrades Factor V and factor VIII.
2. Antithrombin: a serine protease inhibitor that degrades thrombin, Factor IXa, Factor Xa, Factor XIa, and Factor XIIa.
3. Tissue factor pathway inhibitor (TFPI): limits the action of tissue factor (TF) and the factors it produces.
4. Plasmin: generated by proteolytic cleavage of plasminogen, a potent fibrinolytic that degrades fibrin and destroys clots.
5. Prostacyclin (PGI2): released by the endothelium and inhibits platelet activation.
6. Thrombomodulin: released by the endothelium and converts thrombin into an inactive form.

This page titled 16.5D: Coagulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5E: Role of Vitamin K

Vitamin K is an essential factor of the coagulation cascade.

Learning Objectives

- Describe the role of vitamin K in hemostasis


Key Points

- Vitamin K is involved in the synthesis of many factors of the coagulation cascade.
- Vitamin K is antagonized (inhibited) by the anticoagulant drug warfarin.
- Calcium and phospholipids are needed to activate tenase, which converts prothrombin to thrombin.
- Both calcium and vitamin K are needed to synthesize Protein C, an anticoagulant that prevents excessive coagulation after the coagulation cascade occurs.
- Deficiency of any of these clotting cofactors will cause an impaired ability for blood to coagulate, which can contribute to excessive bleeding and hemorrhage.

Key Terms

- **warfarin**: An anticoagulant medication that is used for the prophylaxis of thrombosis and embolism in many disorders.
- **tenase**: An enzyme activated by a calcium and phospholipid complex that converts prothrombin to thrombin in the common pathway.

Coagulation is a complex cascade that requires many different cofactors and molecules to occur. Vitamin K, calcium, and phospholipids are necessary cofactors for proper coagulation, and people deficient in these substances will be more susceptible to uncontrolled bleeding.



Blood Coagulation Pathways: Blood coagulation pathways in vivo showing the central role played by thrombin.

Vitamin K

Vitamin K is a fat-soluble vitamin necessary for synthesis of coagulation factors involved in the coagulation cascade. Factors II, VII, IX, and X which are all important for the intrinsic and common pathways of coagulation. Vitamin K also synthesizes Protein C, Protein S, and Protein Z, anticoagulant proteins that degrade specific coagulation factors, preventing excessive thrombosis following the initial coagulation cascade.

Vitamin K can be inhibited by the anticoagulant drug warfarin, which acts as an antagonist for vitamin K. Warfarin is used in medicine for those at high risk of thromboembolism to prevent the coagulation cascade by reducing vitamin K dependent synthesis of coagulation factors. Warfarin's effects can be overcome by ingesting more vitamin K to reactivate the coagulation factor synthesis pathway.

Vitamin K deficiency is associated with impaired coagulation function and excessive bleeding and hemorrhage (internal bleeding, often severe). This can be caused by poor diet, malabsorption in the intestines, or liver failure. Those with vitamin K deficiency produce alternative proteins that improperly bind with phospholipids, which also contributes to the lack of coagulant function.

Calcium and Phospholipids

Calcium and phospholipids (a platelet membrane constituent) are required cofactors for prothrombin activation enzyme complexes to function. This enzyme is called tenase, and converts prothrombin to thrombin. Calcium mediates the binding of the tenase enzyme complexes (via the terminal gamma-carboxy residues on FXa and FIXa) to the phospholipid surfaces expressed by platelets, which in turn activates prothrombin to produce thrombin, which then produces fibrin from fibrinogen. Calcium acts as a catalyst for this reaction, speeding up the rate of the reaction to occur within the time frame of the factors involved in the coagulation cascade. Calcium is also required to to synthesize the anticoagulant Protein C (along with vitamin K).

Calcium deficiencies inhibit proper blood coagulation. This can be caused by a nutritional deficiency or acute problems in which calcium is allocated elsewhere in the blood. Phosopholipid deficiency is also associated with thrombocytopenia (platelet deficiency) because the phospholipids involved with clotting come from platelets. Thrombocytopenia causes more severe issues with blood clotting as the platelet plug will not be able to form or activate the coagulation cascade.

This page titled 16.5E: Role of Vitamin K is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5F: Clot Retraction and Repair

Clot retraction is the shrinking of a blood clot facilitated by thrombolytic agents.

Learning Objectives

- Outline the process of clot retraction and repair

Key Points


- Clot retraction is dependent on the release of multiple coagulation factors, specifically Factor XIIIa at the end of the coagulation cascade.
- The formation of blood clots can cause a number of serious diseases. By breaking down the clot, the disease process can be arrested or the complications reduced.
- Clot retraction is the “shrinking” of a blood clot over a number of days. The edges of the blood vessel wall at the point of injury are slowly brought together to repair the damage.
- Clot retraction occurs due to the contraction, knotting, and twisting of the fibrin mesh.
- The steps of wound healing that follow clot retraction include inflammation, tissue proliferation, collagen and granulation tissue deposition, angiogenesis, wound contraction, and epithelialization.

Key Terms

- **Clot retraction**: The shrinking of a blood clot over the day following initial clot formation.
- **angiogenesis**: Growth of new blood vessels during wound healing.
- **thrombus**: Ablood clot formed from platelets and other elements that forms in a blood vessel in a living organism. It may cause thrombosis or obstruction of the vessel at its point of formation or travel to other areas of the body

The blood clots produced in hemostasis are merely the first step in repair and healing that occur after injury. Following a clot, inflammation draws leukocytes to the injury site to eliminate any pathogens that may have entered the body during the initial injury. Then, over the course of the next 24 hours, the clot retracts as tissue healing begins.

Clot Retraction



Thrombus or Blood Clot: Micrograph showing a thrombus (center of image) within a blood vessel of the placenta.

As the healing process occurs following blood clot formation, the clot must be destroyed in order to prevent thromboembolic events, in which clots break off from the endothelium and cause ischemic damage elsewhere in the body. By reducing the size of and breaking down the clot, the disease process can be arrested or the complications reduced.

Clot retraction refers to a regression in size of the blood clot over a number of days. During this process, the edges of the endothelium at the point of injury are slowly brought together again to repair the damage. Clot retraction is dependent on the release of multiple coagulation factors released at the end of the coagulation cascade, most notably factor XIIIa crosslinks. These factors cause the fibrin mesh to contract by forming twists and knots that condense the size of the clot. Clot retraction generally occurs within 24 hours of initial clot formation and decreases the size of the clot by 90%. Following clot retraction, a separate process called fibrinolysis occurs which degrades the fibrin of the clot while macrophages consume the expended platelets, thus preventing possible thromboembolism.

Wound Healing

While the clot retracts, the wound begins to heal. The first step of wound healing is epithelial cell migration, which forms a scab before the clot retracts. This occurs due to the stimulus of platelet-derived growth factor (PDGF). After clot retraction, true repair begins as tissue proliferation starts and collagen from the extracellular matrix is deposited in the wound while granulation tissue forms. Then new blood vessels grow into the healing tissue in a process called angiogenesis, which is stimulated by vascular endothelial growth factor (VEGF). The wound itself contracts, reducing in size. After these steps occur, new epithelial cells grow to cover the wound. If the wound was severe or unevenly shaped, or if healing takes too long, scarring may occur from collagen deposition. Most scarring on the skin is benign, but scarring inside the tissues of organs such as the heart or the lungs can cause health problems.

This page titled 16.5F: Clot Retraction and Repair is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.5G: Fibrinolysis

Fibrinolysis is a process of breaking down clots in order to prevent them from growing and becoming problematic.

Learning Objectives

- Describe the process of fibrinolysis

Key Points

- Fibrinolysis is the breakdown of a fibrin clot.
- Plasmin is the enzyme that breaks down fibrin. It is activated from inactive plasminogen by tissue plasminogen activator (t-PA) and urokinase.
- Tissue plasminogen activator (t-PA) and urokinase are inhibited by plasminogen activator inhibitor-1 and plasminogen activator inhibitor-2 (PAI-1 and PAI-2).
- Many drugs have fibrinolytic properties that can be used to treat uncontrolled clotting and embolism, including streptokinase, synthetic t-PA, aspirin, heparin, warfarin, and citrates.
- Patients suspected to be having a blood clot-induced stroke should be screened for hemorrhage and aneurysm first, since fibrinolytic treatment would help with a stroke but make bleeding conditions much more dangerous.

Key Terms


- **fibrin**: An elastic, insoluble, whitish protein produced by the action of thrombin on fibrinogen and forming an interlacing fibrous network in the coagulation of blood.
- **protease**: An enzyme that cuts or cleaves proteins.
- **Plasmin**: A protease that breaks down plasmin. It is cleaved from inactive plasminogen.

Fibrinolysis is a process that removes clots following hemostasis and clot retraction, preventing uncontrolled thrombosis and embolism. There are two types of fibrinolysis: primary fibrinolysis and secondary fibrinolysis. Primary fibrinolysis is a normal body process, whereas secondary fibrinolysis is the breakdown of clots due to a medication, medical disorder, or other cause.

Mechanisms of Primary Fibrinolysis

Primary fibrinolysis normally occurs following clot retraction, in which the clot has already condensed considerably in size. The main enzyme in primary fibrinolysis is plasmin, a proteolytic enzyme that degrades fibrin mesh. Plasmin cleaves fibrin at various places, leading to the production of circulating fragments that are cleared by other proteases or by the kidneys and liver.

Plasmin is produced in an inactive form, plasminogen, in the liver. Plasminogen cannot cleave fibrin and circulates in the bloodstream. Instead, it is incorporated into the clot when it is formed and then activated into plasmin later. Plasminogen is activated to plasmin by tissue plasminogen activator (t-PA) and urokinase, an enzyme found in the urine.



Fibrinolysis : Blue arrows denote stimulation and red arrows inhibition.

T-PA is released into the blood very slowly by the damaged endothelium of the blood vessels. T-PA and urokinase are themselves inhibited by plasminogen activator inhibitor-1 and plasminogen activator inhibitor-2 (PAI-1 and PAI-2). In contrast, plasmin further stimulates plasmin generation by producing more active forms of both tissue plasminogen activator (tPA) and urokinase. Following fibrin degradation by plasmin, old activated platelets from the platelet plug are phagocytized and destroyed by macrophages.

Alpha 2-antiplasmin and alpha 2-macroglobulin inactivate plasmin. Plasmin activity is also reduced by thrombin -activatable fibrinolysis inhibitor (TAFI), which modifies fibrin to make it more resistant to the tPA-mediated plasminogen. Plasmin operates on a negative feedback process because it is reduced when the fibrin clot is fully degraded.

Mechanisms of Secondary Fibrinolysis

Secondary fibrinolysis generally refers to treatment of pathological thromboembolism. If blood clots embolize to different parts of the body, they can cause tissue death by blocking off blood flow to those tissues. This is a common cause of heart attacks, pulmonary embolism, and strokes. Several medications exist to help treat and prevent these conditions.

Fibrinolytic drugs include synthesized tissue plasminogen activator and streptokinase, a bacterial enzyme that has degrades fibrin directly. Clots may also be prevented or kept from worsening through the use of blood thinners (anticoagulants). Aspirin has anticoagulant properties because it inhibits cyoclo-oxygenase dependent pathways of platelet activation, which can prevent clotting from worsening. Heparin is a fast-acting anticoagulant produced by the body and used as a drug which inhibits the activity of thrombin. Warfarin inhibits vitamin K cofactor activation during the coagulation cascade, and citrates chelate calcium to prevent prothrombin activation into thrombin.

All of these treatments have been shown to have tremendous therapeutic benefit in treating those with thromboembolic diseases; however, they can make injury much more difficult to treat by disrupting the clotting process. For example, patients thought to be suffering from a stroke (obstructed artery in the brain) must be screened through imaging before given aspirin or a fibrinolytic drug, because if they have an aneurysm or hemorrhage (burst blood vessel or bleeding in the brain), administering fibrinolytic treatment would make their condition worse and possibly fatal by inhibiting the clotting that could save their lives.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...or_Clotting.29. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hemostasis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hemostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- hemostasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hemostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- heparin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/heparin. **License**: *CC BY-SA: Attribution-ShareAlike*
- coagulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/coagulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Vasoconstriction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vasoconstriction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...or_Clotting.29. **License**: *CC BY-SA: Attribution-ShareAlike*
- endothelial cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/endothelial%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- vascular. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vascular. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...or_Clotting.29. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hemostasis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hemostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- hemostasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hemostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- collagen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/collagen. **License**: *CC BY-SA: Attribution-ShareAlike*
- serotonin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/serotonin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Adenosine diphosphate ribose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ate_ribose.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Platelets. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Platelets.jpg. **License**: *CC BY: Attribution*
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...or_Clotting.29. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagula...lation_cascade. **License**: *CC BY-SA: Attribution-ShareAlike*
- fibrin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fibrin. **License**: *CC BY-SA: Attribution-ShareAlike*
- endothelium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/endothelium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY:*

Attribution
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Adenosine diphosphate ribose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ate_ribose.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Platelets. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Platelets.jpg. **License**: *CC BY: Attribution*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagula...lation_cascade. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagula...lation_cascade. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagulation%23Cofactors. **License**: *CC BY-SA: Attribution-ShareAlike*
- warfarin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/warfarin. **License**: *CC BY-SA: Attribution-ShareAlike*
- coumarin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/coumarin. **License**: *CC BY-SA: Attribution-ShareAlike*
- oxidized. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/oxidized. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Adenosine diphosphate ribose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Adenosine_diphosphate_ribose.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Platelets. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Platelets.jpg. **License**: *CC BY: Attribution*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagula...lation_cascade. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coagula...lation_cascade. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coagulation in vivo. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Co...on_in_vivo.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- platelet. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/platelet. **License**: *CC BY-SA: Attribution-ShareAlike*
- thrombus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thrombus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clot retraction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clot_retraction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thrombolysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thrombolysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blood clot. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_clot. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blood clot. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_clot. **License**: *CC BY-SA: Attribution-ShareAlike*
- clot. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/clot. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clot retraction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clot%20retraction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*

- Microvessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Microvessel.jpg**. **License**: ***CC BY: Attribution***
- Adenosine diphosphate ribose. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Adenosine_diphosphate_ribose.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Platelets.jpg**. **License**: ***CC BY: Attribution***
- Coagulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coagula...lation_cascade**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coagulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coagulation%23The_coagulation_cascade**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coagulation in vivo. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coagulation_in_vivo.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fetal thrombotic vasculopathy - very high mag. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Fe...y_high_mag.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrinolysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrinolysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- coagulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/coagulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- protease. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/protease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibrin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibrin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Microvessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Microvessel.jpg**. **License**: ***CC BY: Attribution***
- Microvessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Microvessel.jpg**. **License**: ***CC BY: Attribution***
- Microvessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Microvessel.jpg**. **License**: ***CC BY: Attribution***
- Adenosine diphosphate ribose. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Adenosine_diphosphate_ribose.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Platelets. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Platelets.jpg**. **License**: ***CC BY: Attribution***
- Coagulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coagulation%23The_coagulation_cascade**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coagulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coagulation%23The_coagulation_cascade**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coagulation in vivo. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coagulation_in_vivo.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fetal thrombotic vasculopathy - very high mag. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Fetal_thrombotic_vasculopathy_-_very_high_mag.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrinolysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrinolysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 16.5G: Fibrinolysis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.6: TRANSFUSIONS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

16.6A: Transfusions of Whole Blood

16.6B: Plasma and Blood Volume Expanders

16.6C: Blood Groups and Blood Types

16.6D: Typing and Cross-Matching for Transfusions

This page titled 16.6: Transfusions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.6A: Transfusions of Whole Blood

Whole blood refers to human blood transfusion from a standard blood donation.

Learning Objectives

- Describe whole blood in terms of transfusions

Key Points


- Whole blood can be separated into its components: red blood cells, plasma, and platelets.
- Blood can be transfused in either its relatively unprocessed form as whole blood, or through the administration of its processed and separated cellular or plasma componenets.
- Plateletpheresis is a more efficient way to extract platelets than whole blood extraction, because plateletpheresis produces platelets that are more highly concentrated.
- The blood is separated into distinct plasma and cellular layers by centrifugation, by sedimentation, or simply through gravity over a longer period of time.
- Whole blood transfusion has similar risks to a transfusion of red blood cells and must be cross-matched based on blood type to avoid hemolytic transfusion reactions and other complications.

Key Terms

- **buffy coat**: The fraction of an anticoagulated blood sample that contains most of the white blood cells and platelets following density gradient centrifugation of the blood.
- **centrifuge**: A device in which a mixture of denser and lighter materials (normally dispersed in a liquid) is separated by rotation around a central axis at high speed.
- **plasma**: The straw-colored/pale-yellow liquid component of blood that normally holds the blood cells of whole blood in suspension.

Blood transfusions are a key therapeutic component to treating those with excessive blood loss from severe injury or surgery. Whole blood refers to blood drawn directly from the body from which none of the components, such as plasma or platelets, have been removed. The blood is typically combined with an anticoagulant during the collection process, but is otherwise unprocessed. Whole blood may also be altered and processed for use in blood transfusion.

Blood Processing



Centrifuged Blood: Whole blood is a term used in transfusion medicine for human blood from a standard blood donation.

Historically, blood was transfused as whole blood without further processing. Most blood banks now split the whole blood into two or more components, typically red blood cells and a plasma component such as fresh frozen plasma, which is extracted frozen plasma from the blood splitting process.

Platelets for transfusion can also be prepared from the buffy coat of whole blood, which has therapeutic benefits for those with platelet disorders or impaired clotting ability. Some blood banks have replaced this with platelets collected by plateletpheresis, a process in which platelets are extracted during initial blood collection. Plateletpheresis is more efficient because whole blood platelets typically aren't concentrated enough to have a useful effect, while plateletpheresis platelets are highly packed and concentrated. It also minimizes the chance for platelet transplant rejection because a single donor will be able to contribute enough platelets via plateletpheresis.

The collected blood is generally separated into components by one of three laboratory methods:

- Centrifuge quickly separates whole blood into plasma, buffy coat, and red cells by using centrifugal force to drop the cellular components to the bottom of a container.
- Sedimentation, in which whole blood sits overnight, causing the red blood cells and plasma to settle and slowly separate by the force of normal gravity.

Practical Considerations

Whole blood transfusion has similar risks to those of transfusion of red blood cells. It must be cross-matched on the basis of blood type to avoid hemolytic transfusion reactions. Most of the indications for use are identical to those for red blood cells. Whole blood is not used because the extra plasma can contribute to transfusion associated circulatory overload (TACO), a potential complication that can dangerously increase blood pressure, causing pulmonary edema and acute respiratory distress.

Whole blood is sometimes "recreated" from stored red blood cells and fresh frozen plasma for neonatal transfusions. This provides a final product with a very specific hematocrit (percentage of red cells) with type O red cells and type AB plasma to minimize the chance of complications.

Whole blood is typically stored under the same conditions as red blood cells and can be kept up to 35 days if collected with CPDA-1 storage solution or 21 days with other common storage solutions such as CPD. If the blood will be used to make platelets, it is kept at room temperature until the process is complete. This must be done quickly to minimize the warm storage of RBCs in the unit.



Blood Transfusions: Blood transfusions are common during surgeries.

This page titled 16.6A: Transfusions of Whole Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.6B: Plasma and Blood Volume Expanders

A volume expander is a type of intravenous therapy that provides fluid replacement for the circulatory system.

Learning Objectives

- Evaluate the use of blood volume expanders

Key Points

- During blood loss, the amount of oxygen that can be delivered to the tissues is reduced due to lost red blood cells and decreased blood volume, which also causes a decrease in blood pressure.
- Although they cannot replace lost red blood cells, blood volume expanders can help improve oxygen delivery in instances of blood loss by increasing blood volume and blood pressure so that blood can flow to the tissues.
- Survival is possible with low red blood cell and hemoglobin levels as long as blood volume and blood pressure are maintained so blood continues to reach tissues.
- Hypovolemic shock occurs when tissue oxygenation drops due to a decrease in blood volume.
- Crystalloids volume expanders are aqueous solutions of mineral salts or other water-soluble molecules. Although they decrease the osmotic pressure by diluting the red blood cells, they increase both vascular and interstitial volume.
- Colloids volume expanders contain larger insoluble molecules, such as gelatin or hydroxyethyl starch, and theoretically increase the intravascular volume but not interstitial and intracellular volumes.

Key Terms

- **crystalloid**: Aqueous solutions of mineral salts or other water-soluble molecules, such as saline solution.
- **hypovolemic shock**: Shock due to decreased blood volume, such as through severe bleeding or vomiting. It activates dangerous compensatory mechanisms that maintain blood flow to the brain while causing other organs to fail.
- **colloid**: Blood volume expander containing larger insoluble molecules that exert osmotic pressure.

When blood is lost, the greatest immediate need is to stop further blood loss, then lost volume must be replaced. Blood volume is directly proportional to the blood pressure in the body, and when both decrease the flow of blood to important tissues may be inhibited. The remaining red blood cells can still oxygenate body tissue. A volume expander is a type of intravenous therapy that provides blood volume for the circulatory system. It may be used for fluid replacement.

Blood Volume and Oxygen Transport

Normal human blood has a significant excess oxygen transport capability because not all of the hemoglobin molecules are loaded with oxygen under normal conditions. As long as pulmonary function is sufficient for gas exchange and there is enough blood volume to have sufficient blood pressure, very low hemoglobin levels will be enough to sustain the patient. Those with low hemoglobin content will not be able to tolerate situations where a greater amount of oxygen is required (exercise, for example) until their hemoglobin levels are restored.

The body has compensatory feedback mechanisms to deal with lower hemoglobin levels. For instance, the heart pumps more blood with each beat, which increases blood pressure. Blood pressure is detected by the renal system, which increases blood volume and blood pressure by excreting less water during blood filtration. As a result of partial pressure gradient changes, more oxygen is released to the tissues. These adaptations are so effective that if only half of the red blood cells remain, oxygen delivery will still be around 75% of normal. A patient at rest only uses 25% of the oxygen available in their blood. In extreme cases, patients have survived with a hemoglobin level of about 1/7 the normal (i.e. 2 g/dl), although levels this low are very dangerous.

When blood loss is significant, the red blood cell level ultimately drops to a level that is too low for adequate tissue oxygenation. This is marked by hypoxia and hypovolemic shock, a condition in which tissue oxygenation drops from a lack of blood volume and harmful compensatory mechanisms activate, causing more damage. In these situations, the only alternatives are blood

transfusion, packed red blood cells, or oxygen therapy.

Types of Volume Expanders



Saline solution: A bag of saline. Saline can be used to increase blood volume when a blood transfusion is not possible.

There are two main types of volume expanders: crystalloids and colloids. Crystalloids are aqueous solutions of mineral salts or other water-soluble molecules. Colloids contain larger insoluble molecules, such as gelatin; blood itself is a colloid. There are also a few other volume expanders that may be used in certain situations:

- Colloids: These solutions preserve a high-colloid osmotic pressure (protein-exerted pressure) in the blood, while this parameter is decreased by crystalloids due to hemodilution. The higher osmotic pressure from colloids draws fluids inward, preventing it from leaking out into the tissues as easily, which increases intravascular blood volume.
- Crystalloids: The most commonly used crystalloid fluid is normal saline, a solution of sodium chloride at 0.9% concentration, which is close to the concentration in the blood (isotonic). Saline solution is administered intravenously (IV drips) and increases both intravascular and interstitial volume. They decrease osmotic pressure by diluting the blood.
- Dextrose Water: This solution contains dextrose, a form of glucose. It is given to patients who have dangerously low blood sugar levels (important for cellular metabolism) as well as low blood volume.

Another common volume expander includes hydroxyethyl starch (HES/HAES, common trade names: Hespan, Voluven) which is considered a colloid. An intravenous solution of hydroxyethyl starch is used to prevent shock following severe blood loss caused by trauma, surgery, or another problem. It increases the blood volume, allowing red blood cells to continue to deliver oxygen to the body. When tissue blood perfusion is maintained, shock is averted as the dangerous compensatory mechanisms of shock aren't activated.

This page titled 16.6B: Plasma and Blood Volume Expanders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.6C: Blood Groups and Blood Types

Red blood cells have surface-expressed proteins that define the self/not-self nature of the cells.

Learning Objectives

- Evaluate the ABO and Rhesus blood groups in terms of donors and recipients

Key Points

- Surface-expressed proteins called antigens on red blood cells determine an individual's blood type. There are two types of antigen groups: the ABO system antigens and a Rhesus D antigen.
- Exposure to a blood group antigen that is not recognized as self will cause the immune system to make specific antibodies to the new blood group antigen, often leading to destruction of the cells.
- Knowing an individual's antigen type is important to ensure compatibility if a transfusion is needed.
- Blood type is inherited. O type is the most common despite being a recessive gene because it is more highly expressed in the gene pool, while type A and type B are dominant (and type AB is codominant) but are less common because they are less expressed in the gene pool.
- Individuals may also be positive or negative for the rhesus D antigen in addition to their blood type. Rhesus D complications are common during fetal development if the parents differ in rhesus antigen expression.

Key Terms


- **antibodies**: Also known as an immunoglobulin (Ig), a large Y-shaped protein produced by B-cells that is used by the immune system to identify and neutralize foreign objects such as bacteria and viruses.
- **antigen**: A substance that induces an immune response, usually foreign.

Red blood cells have surface-expressed proteins that act as antigens, which are molecules that can illicit an immune system response. Red blood cells belong to different groups on the basis of the type of antigen that they express. Blood type determines compatibility for receiving blood transfusions from other people.

The ABO Blood Group System

If an individual is exposed to a blood group antigen (A or B) that is not recognized as self, the individual can become sensitized to that antigen. This will cause the immune system to make specific antibodies to a particular blood group antigen and form an immunological memory against that antigen. These antibodies can bind to antigens on the surface of transfused red blood cells (or other foreign tissue cells), often leading to destruction of the cells by recruitment of other components of the immune system.

Knowledge of an individual's blood type is important to identify appropriate blood for transfusion or tissue for organ transplantation. There are four blood types that differ based on the antigen expressed by the red blood cell and by the type of associated antibody found in the plasma. The type of antigen determines which blood types that blood type may safely be donated to, while the type of antibody determines which types of antigen (and types of blood) will be rejected by the body.



Blood type classification: Blood type (or blood group) is determined, in part, by the ABO blood group antigens present on red blood cell.

- Blood group A individuals have the A antigen on the surface of their RBCs, and blood serum containing IgM antibodies against the B antigen. Therefore, a group A individual can only receive blood from individuals of groups A or O types, and can donate blood to individuals of groups A or AB.
- Blood group B individuals have the B antigen on their surface of their RBCs, and blood serum containing IgM antibodies against the A antigen. Therefore, a group B individual can only receive blood from individuals of groups B or O, and can donate blood to individuals of groups B or AB.
- Blood Group AB individuals have both A and B antigens on the surface of their RBCs, and their blood serum does not contain any antibodies against either A or B antigen. Therefore, an individual with type AB blood can receive blood from any group, but can only donate blood to another group AB individual. AB blood is also known as “universal receiver.”
- Blood group O individuals do not have either A or B antigens on the surface of their RBCs, but their blood serum contains IgM antibodies against both A and B antigens. Therefore, a group O individual can only receive blood from a group O individual, but they can donate blood to individuals of any ABO blood group (i.e. A, B, O, or AB). O blood is also known as “universal donor.”

Blood types are inherited and represent genetic contributions from both parents. The gene that codes for blood type contains three alelles: IA and IB which give type A and B blood and are dominant, and i, which is recessive and codes type O. Children will have blood types similar to their parents based on inheritance. The i allele is far more commonly expressed in the gene pool than IA and IB, which is why type O blood is the most common type despite being a recessive phenotype. Type AB is the rarest because it is the combination of less commonly expressed alleles, and is the result of codominance between IA and IB alelles.

Rhesus Factor

Many people also have the rhesus D (Rh) antigen expressed by their red blood cells. Those that are have Rh antigens are positive for Rh, while those that don’t have it are Rh negative (ie. type O+ is type O with rhesus, type A- is type A without rhesus). Rh positive individuals do not have the antibodies for the Rh factor, but can make them if exposed to Rh. Besides being a consideration for blood transfusion, parents who differ based on Rh status must be cautious to ensure that maternal antibodies do not destroy their child’s red blood cells during fetal development, which can cause hemolytic anemia.

This page titled 16.6C: Blood Groups and Blood Types is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

16.6D: Typing and Cross-Matching for Transfusions

Blood banks test donor blood to ensure recipient compatibility, reducing the risk of hemolytic reaction, renal failure, and death.

Learning Objectives

- Explain the purposes of typing and of cross-matching blood prior to transfusion

Key Points

- Transfusion medicine is important to treat those with blood loss.
- Given enough time, cross-matching is performed to ensure that donated blood will not cause a transfusion reaction.
- Cross-matching involves mixing a sample of the recipient's serum with a sample of the donor's red blood cells and checking if the mixture agglutinates due to antibody reactivity.
- If a transfusion with non-matched blood occurs, the patient risks red blood cell destruction, renal failure, shock, and death.


Key Terms

- **hemolysis**: The destruction of red blood cells from pathological causes, such as infection or immune system mediated damage.
- **agglutinate**: The act of red blood cells clumping together due to antibody reactivity.

Transfusion medicine is extremely effective at treating those with severe blood loss. Transfusions are often a required component of major surgeries. Due to the different antigen blood types, blood must be cross-matched during processing to avoid potential complications.

The Cross-Matching Process

Much of the routine work of a blood bank involves testing blood from both donors and recipients to ensure that every recipient is given blood that is compatible and is as safe as possible. Several laboratory tests allow cross-matching of compatible blood between donor and recipient. Patients should ideally receive their own blood or type-specific blood products to minimize the chance of a transfusion reaction. Risks can be further reduced by cross-matching blood, but this process isn't always performed if time is short and the need for transfusion has not been anticipated.



Agglutinated RBC: Red blood cells can agglutinate if the serum contains antibodies against the expressed proteins. In this image, the blood serum contains anti-A3 antibodies, which attack and agglutinate type A blood.

Cross-matching involves mixing a sample of the recipient's serum with a sample of the donor's red blood cells and checking if the mixture agglutinates, or forms clumps. These clumps are the result of antibodies binding the red blood cells together. If agglutination is not obvious by direct vision, blood bank technicians check for agglutination with a microscope. If agglutination occurs, that particular donor's blood cannot be transfused to that particular recipient. In a blood bank, it is vital that all blood specimens are correctly identified, so labeling has been standardized using a barcode system known as ISBT 128. The blood group may be included on identification by military personnel in case they need an emergency blood transfusion.

Potential Transfusion Complications

If a patient receives blood during a transfusion that is not compatible with his or her blood type, severe problems can occur. Acute hemolytic transfusion reactions occur if donated blood cells are attacked by matching host antibodies. This can cause shock-like symptoms, such as fever, hypotension, and disseminated intravascular coagulation from immune system mediated endothelial damage. Transfusion reactions are also associated with acute renal failure. Lung injury is common as well, due to pulmonary edema from fluid overload if plasma volume becomes too high or neutrophil activation during a transfusion reaction. If the donated blood is contaminated with bacteria, it may induce septic shock in the patient.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Whole blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Whole_blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- centrifuge. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/centrifuge**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Buffy coat. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Buffy_coat**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plasma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- platelet. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/platelet**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- US Navy 060505-N-2832L-050 Hospital Corpsman Megan Beach, checks the flow of blood through a cell saver, during training for an antilogous blood transfusion. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ransfusion.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood-centrifugation-scheme. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion-scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood plasma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_p...lasma_donation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Volume expander. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Volume_expander**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- isotonic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/isotonic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- crystalloid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/crystalloid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- colloid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/colloid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- US Navy 060505-N-2832L-050 Hospital Corpsman Megan Beach, checks the flow of blood through a cell saver, during training for an antilogous blood transfusion. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ransfusion.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood-centrifugation-scheme. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion-scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Iv1-07 014. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Iv1-07_014.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...O_Group_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- antigen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/antigen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ion/antibodies**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- US Navy 060505-N-2832L-050 Hospital Corpsman Megan Beach, checks the flow of blood through a cell saver, during training for an antilogous blood transfusion. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ransfusion.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood-centrifugation-scheme. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion-scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Iv1-07 014. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Iv1-07_014.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_g..._compatibility**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_g...od_transfusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hemolysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemolysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- agglutinate. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/agglutinate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- complement system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/complement_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- US Navy 060505-N-2832L-050 Hospital Corpsman Megan Beach, checks the flow of blood through a cell saver, during training for an antilogous blood transfusion. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ransfusion.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood-centrifugation-scheme. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion-scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Iv1-07 014. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Iv1-07_014.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anti-HLA agglutinated RBC. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...inated_RBC.png**.

License: *CC BY-SA: Attribution-ShareAlike*

This page titled 16.6D: Typing and Cross-Matching for Transfusions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17: CARDIOVASCULAR SYSTEM: THE HEART

17.1: The Heart

17.1A: Anatomy of the Heart

17.1B: Pericardium

17.1C: Layers of the Heart Walls

17.1D: Chambers of the Heart

17.1E: Great Vessels of the Heart

17.1F: Myocardial Thickness and Function

17.1G: Fibrous Skeleton of the Heart

17.2: Circulation and Heart Valves

17.2A: Heart Circulation

17.2B: Operation of Atrioventricular Valves

17.2C: Operation of Semilunar Valves

17.2D: Systemic and Pulmonary Circulation

17.3: Cardiac Muscle Tissue

17.3A: Mechanism and Contraction Events of Cardiac Muscle Fibers

17.3B: Microscopic Anatomy

17.3C: Energy Requirements

17.4: Physiology of the Heart

17.4A: Electrical Events

17.4B: Electrocardiogram and Correlation of ECG Waves with Systole

17.4C: Heart Sounds

17.4D: Cardiac Cycle

17.4E: Cardiac Output

17.5: Exercise and the Heart

17.5A: Effects of Exercise on the Heart

Section 6:

This page titled 17: Cardiovascular System: The Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1: THE HEART

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

17.1A: Anatomy of the Heart

17.1B: Pericardium

17.1C: Layers of the Heart Walls

17.1D: Chambers of the Heart

17.1E: Great Vessels of the Heart

17.1F: Myocardial Thickness and Function

17.1G: Fibrous Skeleton of the Heart

This page titled 17.1: The Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1A: Anatomy of the Heart

The heart is an organ responsible for pumping blood through the blood vessels using rhythmic contractions of cardiac muscle.

Learning Objectives

- Describe the anatomy of the heart

Key Points

- The heart is a four-chambered muscular organ containing an involuntary conduction system that initiates rhythmic contractions to pump blood throughout the body.
- The heart has its own blood supply and is controlled by self-regulating nerve bundles called nodes.
- The SA and AV nodes send impulses through the Purkinje fibers that cause muscle contractions to the heart.
- The heart is composed of three layers: the epicardium (outer layer) which prevents excess expansion or movement of the heart, the myocardium (middle layer) which initiates contractions driving the cardiac cycle, and the endocardium (inner layer) that lines the cavities and valves.
- The heart contains two upper chambers called atria and two lower chambers called ventricles. The left and right sides of the heart are separated by the septum.
- Valves prevent the backflow of blood and separate the atria from the ventricle and the ventricle from the pulmonary artery or aorta.

Key Terms

- **heart**: A fist-sized muscular organ in the chest that pumps blood through the body using involuntary contractions of cardiac muscle.
- **ventricle**: One of two lower chambers of the heart that receives blood from the atria and pumps it out into pulmonary or systemic circulation, depending on side.
- **myocardium**: The middle of the three layers forming the wall of the heart, containing cardiac muscle tissue. Innervated by the Purkinje fibers.

The human heart is the pump for the circulatory system, and along with the circulatory system is considered to be an organ of the cardiovascular system. It consists of four chambers and pumps blood through both systemic and pulmonary circulation to enable gas exchange and tissue oxygenation. The heart is located in the thoracic cavity between the lungs and protected by the rib cage.

Structure of the Heart

The heart consists of four chambers separated into two sides. Each side contains an atria which receives blood into the heart and flows it into a ventricle, which pumps the blood out of the heart. The atria and ventricle on each side of the heart are linked together by valves that prevent backflow of blood. The wall that separates the left and right side of the heart is called the septum.


The left heart deals with systemic circulation, while the right heart deals with pulmonary circulation. The left side of the heart receives oxygenated blood from the pulmonary vein and pumps it into the aorta, while the right side of the heart receives deoxygenated blood from the vena cava and pumps it into the pulmonary vein. The pulmonary vein and aorta also have valves connecting them to their respective ventricle.

The heart has its own self-sustaining conduction system that sends nervous impulses to cardiac tissue. The sinoatrial (SA) and atrioventricular (AV) nodes are bundles of nerve fibers that form this conduction system. They are located in the left atrial wall of the heart and send nerve impulses to a large, highly specialized set of nerves called the Purkinje fibers, which in turn send those nerve impulses to the cardiac muscle tissue. These nodes can send impulses to the heart without central nervous system stimulation, but may be influenced by nervous stimulation to alter heart rate. The heart also has its own blood supply, the cardiac arteries that provide tissue oxygenation to the heart as the blood within the heart is not used for oxygenation by the heart.

Cardiac Histology

The heart is enclosed in a double-walled protective membrane called the pericardium, which is a mesothelium tissue of the thoracic cavity. The double membrane of pericardium contains pericardial fluid which nourishes the heart and prevents shock. This composite sac protects the heart, anchors it to surrounding structures, and prevents the heart from overfilling with blood.

The wall of the heart is composed of three layers of different tissues. The outer layer is called the epicardium, or visceral pericardium, since it is also the inner wall of the pericardium. The middle layer of the heart, the myocardium, and contains specialized cardiac muscle tissue responsible for contraction. Cardiac muscle tissue is distinct from skeletal or smooth muscle because it pumps involuntarily based on conduction from the AV and SA nodes. The inner layer is called the endocardium and is in contact with the blood that the heart pumps. It also merges with the inner lining of blood vessels and covers heart valves. Cardiac tissue is permanent tissue that does not heal or regenerate when damaged. As a result, is prone to scarring and enlargement due to mechanical stress and injury.



The Mammalian Heart: The position of valves ensures proper directional flow of blood through the cardiac interior. Note the difference in the thickness of the muscled walls of the atrium and the left and right ventricle.

This page titled 17.1A: Anatomy of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1B: Pericardium

The pericardium is a thick, membranous, fluid-filled sac which encloses, protects, and nourishes the heart.

Learning Objectives

- Distinguish between the fibrous and serous layers of the pericardium


Key Points

- The pericardium is a mesothelium tissue of the thoracic cavity which surrounds the heart. The outer layer, the fibrous pericardium, is comprised of dense connective tissue that protects the heart, anchors it to the surrounding walls, and prevents it from overfilling.
- The inner layer of the pericardium, the serous pericardium, is further divided into two layers, an outer parietal layer and an inner visceral layer with the pericardial cavity in between the two layers.
- The serous pericardium functions to lubricate the heart with pericardial fluid, which is stored in the space between the parietal and visceral layers.
- Pericarditis is inflammation of the pericardium, often due to infection, which can constrict and put harmful pressure on the heart.

Key Terms

- **fibrous pericardium**: Composed of dense connective tissue which protects the heart, anchors it to the surrounding walls, and prevents the heart from overfilling with blood.
- **serous pericardium**: Located deeper than the fibrous pericardium, this structure contains two layers that lubricate the heart to prevent friction from occurring during heart activity.
- **pericardium**: A serous membrane that surrounds and protects the heart.

The pericardium is the thick, membranous, fluid-filled sac that surrounds the heart and the roots of the vessels that enter and leave this vital organ, functioning as a protective membrane. The pericardium is one of the mesothelium tissues of the thoracic cavity, along with the pleura which cover the lungs. The pericardium is composed of two layers, an outer fibrous pericardium and an inner serous pericardium.



Membranes of the Thoracic Cavity: A transverse section of the thorax, showing the contents of the middle and the posterior mediastinum. The pleural and pericardial cavities are exaggerated since normally there is no space between parietal and visceral pleura and between pericardium and heart.

Fibrous Pericardium

The fibrous pericardium is the outer layer of the pericardium. It is composed of dense connective tissue which anchors the heart to the mediastinum of the chest wall. It prevents the heart from overfilling with blood and protects it from nearby infections by completely separating it from the rest of the thoracic cavity. It is continuous with the outer fibrous layer of the neighboring great blood vessels.

Serous Pericardium

The serous pericardium, the inner layer of the pericardium, is composed of two different layers. The outer layer, the *parietal layer*, is completely adhered to the fibrous pericardium. The inner layer is known as the *visceral layer*, which covers and protects the great vessels and heart. The space between the parietal and visceral layers is called the pericardial cavity.

The visceral layer is referred to as the *epicardium* in the areas where it is in direct contact with the heart. The space between these two serous layers, the parietal and the visceral, is the *pericardial* cavity, which contains pericardial fluid. The serous pericardium, with its two membranes and the fluid-filled pericardial cavity, provides protection to the heart and a lubricated sliding surface within which the heart can move in response to its own contractions and to the movement of adjacent structures such as the diaphragm and the lungs.

Function of the Pericardium

The pericardium is important because it protects the heart from trauma, shock, stress, and even infections from the nearby lungs. It supports the heart and anchors it to the medastinum so it doesn't move within the body. The pericardium lubricates the heart and prevents it from becoming too large if blood volume is overloaded (though it will not prevent chronic heart enlargement).

Despite these functions, the pericardium is still vulnerable to problems of its own. Pericarditis is the term for inflammation in the pericardium, typically due to infection. Pericarditis is often a severe disease because it can constrict and apply pressure on the heart and work against its normal function. Pericarditis comes in many types depending on which tissue layer is infected.

This page titled 17.1B: Pericardium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1C: Layers of the Heart Walls

The heart wall is comprised of three layers: the outer epicardium, the middle myocardium, and the inner endocardium.

Learning Objectives

- Distinguish between the epicardium, myocardium, and endocardium layers of the heart wall

Key Points


- The epicardium is a thin layer of connective tissue and fat serving as an additional layer of protection for the heart. It is considered a continuation of the serous pericardium.
- The myocardium is the muscle tissue of the heart, composed of cardiac muscle cells called cardiomyocytes that receive nervous stimulation from the sinoatrial (SA) and atrioventricular (AV nodes via the Purkinje fibers.
- Cardiomyocytes are shorter than skeletal myocytes, and contain fewer nuclei. Cardiac muscle is striated.
- The endocardium is composed of endothelial cells that provide a smooth, non-adherent surface for blood collection and pumping and may help regulate contractility.
- An infection or inflammation of the endocardium is called infective endocarditis.

Key Terms

- **Purkinje fibers**: A bundle of nerve fibers located under the endocardium, which supply nervous impulses to the mycardium's cardiac muscle tissues.
- **endocardium**: A thin serous membrane that lines the interior of the heart and valves.
- **cardiomyocyte**: A cardiac muscle cell (or myocyte) in the heart, which makes up the cardiac muscle tissue.

The heart wall is comprised of three layers, the epicardium (outer), myocardium (middle), and endocardium (inner). These tissue layers are highly specialized and perform different functions. During ventricular contraction, the wave of depolarization from the SA and AV nodes moves from within the endocardial wall through the myocardial layer to the epicardial surface of the heart.

Epicardium



The Heart Wall: The wall of the heart is composed of three layers, the thin outer epicardium, the thick middle myocardium, and the very thin inner endocardium. The dark area on the heart wall is scarring from a previous myocardial infarction (heart attack).

The outer layer of the heart wall is the epicardium. The epicardium refers to both the outer layer of the heart and the inner layer of the serous visceral pericardium, which is attached to the outer wall of the heart. The epicardium is a thin layer of elastic connective tissue and fat that serves as an additional layer of protection from trauma or friction for the heart under the pericardium. This layer contains the coronary blood vessels, which oxygenate the tissues of the heart with a blood supply from the coronary arteries.

Myocardium

The middle layer of the heart wall is the myocardium—the muscle tissue of the heart and the thickest layer of the heart wall. It is composed of cardiac muscle cells, or cardiomyocytes. Cardiomyocytes are specialized muscle cells that contract like other muscle cells, but differ in shape. Compared to skeletal muscle cells, cardiac muscle cells are shorter and have fewer nuclei. Cardiac muscle tissue is also striated (forming protein bands) and contains tubules and gap junctions, unlike skeletal muscle tissue. Due to their continuous rhythmic contraction, cardiomyocytes require a dedicated blood supply to deliver oxygen and nutrients and remove waste products such as carbon dioxide from the cardiac muscle tissue. This blood supply is provided by the coronary arteries.

Endocardium

The inner layer of the heart wall is the endocardium, composed of endothelial cells that provide a smooth, elastic, non-adherent surface for blood collection and pumping. The endocardium may regulate metabolic waste removal from heart tissues and act as a barrier between the blood and the heart muscle, thus controlling the composition of the extracellular fluid in which the cardiomyocytes bathe. This in turn can affect the contractility of the heart.

This tissue also covers the valves of the heart and is histologically continuous with the vascular endothelium of the major blood vessels entering and leaving the heart. The Purkinje fibers are located just beneath the endocardium and send nervous impulses from the SA and AV nodes outside of the heart into the myocardial tissues.

The endocardium can become infected, a serious inflammatory condition called infective endocarditis. This and other potential problems with the endocardium may damage the valves and impair the normal flow of blood through the heart.

This page titled 17.1C: Layers of the Heart Walls is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1D: Chambers of the Heart

The heart has four chambers. The two atria receive blood into the heart and the two ventricles pump blood into circulation.

Learning Objectives

- Describe the four chambers of the heart

Key Points

- The right and left atria are the top chambers of the heart and receive blood into the heart. The right atrium receives deoxygenated blood from systemic circulation and the left atrium receives oxygenated blood from pulmonary circulation.
- The atria do not have inlet valves, but are separated from the ventricles by valves. The atria facilitate circulation by allowing uninterrupted blood flow into the heart during ventricular systole.
- The atria have several mechanisms in place to ensure that venous blood flow isn't impaired, such as the lack of valves and the presence of light and brief contractions during atrial systole.
- The right ventricle pumps deoxygenated blood into pulmonary circulation. The left ventricle pumps oxygenated blood into systemic circulation. The left ventricle is thicker because systemic circulation involves a much larger blood volume than pulmonary circulation.
- Systole describes contraction, when blood is pumped from the heart into circulation by the ventricles. It can also describe atrial systole, in which blood moves from the atria to the ventricles during ventricular diastole.
- Diastole describes relaxation, when blood moves into the heart while the ventricles prepare to eject that same blood during systole.

Key Terms

- **diastole**: The relaxation and dilation of the heart chambers between contractions, during which they fill with blood.
- **systole**: The rhythmic contraction of the heart by which blood is ejected from one chamber to the next, or out of the heart and into the arteries.

The heart is the complex pump of the circulatory system, pumping blood throughout the body for the purposes of tissue oxygenation and gas exchange. The heart has four chambers through which blood flows: two sets of each type of chamber (atria and ventricles), one per side, each with distinct functions. The left side of the heart deals with systemic circulation while the right side of the heart deals with pulmonary circulation.

The Atria

The atria are chambers in which blood enters the heart. They are located on the anterior end of the heart, with one atrium on each side. The right atrium receives deoxygenated blood from systemic circulation through the superior vena cava and inferior venae cavae. The left atrium receives oxygenated blood from pulmonary circulation through the left and right pulmonary veins.

Blood passively flows into the atria without passing through valves. The atria relax and dilate (expand) while they fill with blood in a process called atrial diastole. The atria and ventricles are separated by the mitral and tricuspid valves. The atria undergo atrial systole, a brief contraction of the atria that ejects blood from the atria through the valves and into the ventricles. The chordae tendinae are elastic tendons that attach to the valve from the ventricles and relax during atrial systole and ventricular diastole, but contract and close off the valve during ventricular systole.

One of the defining characteristics of the atria is that they do not impede venous flow into the heart. Atria have four essential characteristics that cause them to promote continuous venous flow:

1. There are no atrial inlet valves to interrupt blood flow during atrial systole. The venous blood entering the heart has a very low pressure compared to arterial blood, and valves would require venous blood pressure to build up over a long period of


time to enter the atria.
2. The atrial systole contractions are incomplete and do not block flow from the veins through the atria into the ventricles. During atrial systole, blood not only empties from the atria to the ventricles, but continues to flow uninterrupted from the veins right through the atria into the ventricles.
3. The atrial contractions are slight, preventing significant back pressure that would impede venous flow.
4. The relaxation of the atria is coordinated to begin before the start of ventricular contraction, which also helps prevent the heart from beating too slowly.

Ventricles

The ventricles are located on the posterior end of the heart beneath their corresponding atrium. The right ventricle receives deoxygenated blood from the right atria and pumps it through the pulmonary vein and into pulmonary circulation, which goes into the lungs for gas exchange. The left ventricle receives oxygenated blood from the left atria and pumps it through the aorta into systemic circulation to supply the tissues of the body with oxygen.

The walls of the ventricles are thicker and stronger than those of the atria. The physiologic load on the ventricles, which pump blood throughout the body and lungs, is much greater than the pressure generated by the atria to fill the ventricles. Further, the left ventricle has thicker walls than the right because it pumps blood throughout the body, while the right ventricle pumps only to the lungs, which is a much smaller volume of blood.

During ventricular diastole, the ventricles relax and fill with blood. During ventricular systole, the ventricles contract, pumping blood through the semi-lunar valves into systemic circulation.



Structure of the heart: Structure diagram of a coronal section of the human heart from an anterior view. The two larger chambers are the ventricles.

This page titled 17.1D: Chambers of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1E: Great Vessels of the Heart

Great vessels are the major vessels which directly carry blood into or out of the heart.

Learning Objectives

- Describe the great vessels that carry blood to and from the heart

Key Points


- Five great vessels enter and leave the heart: the superior and inferior vena cava, the pulmonary artery, the pulmonary vein, and the aorta.
- The superior vena cava and inferior vena cava are veins that return deoxygenated blood from circulation in the body and empty it into the right atrium.
- The pulmonary artery carries deoxygenated blood from the right ventricle into the lungs for oxygenation.
- The pulmonary veins carry oxygenated blood from the lungs into the left atrium where it is returned to systemic circulation.
- The aorta is the largest artery in the body. It carries oxygenated blood from the left ventricle of the heart into systemic circulation.
- The aorta has many subdivisions that branch off into smaller arteries. These subdivisions are the ascending and descending aorta, the aortic arch, and the thoracic and abdominal aorta.

Key Terms

- **pulmonary arteries**: The arteries that take deoxygenated blood away from the right side of the heart and into the capillaries of the lungs for the purpose of gas exchange.
- **aorta**: The great artery which carries the blood from the heart into systemic circulation.
- **venae cavae**: The two large vessels, the superior and inferior vena cava, that bring deoxygenated blood from systemic circulation to the heart.

The human circulatory system is a double system, meaning there are two separate systems of blood flow: pulmonary circulation and systemic circulation. The adult human heart consists of two separated pumps, the right side (right atrium and ventricle,) which pumps deoxygenated blood into the pulmonary circulation, and the left side (left atrium and ventricle), which pumps oxygenated blood into the systemic circulation. Great vessels are the major vessels that carry blood into the heart and away from the heart to and from the pulmonary or systemic circuit. The great vessels collect and distribute blood across the body from numerous smaller vessels.

The Venae Cavae



The Systemic Circuit: The venae cavae and the aorta form the systemic circuit, which circulates blood to the head, extremities and abdomen.

The superior and inferior vena cava are collectively called the venae cavae. The venae cavae, along with the aorta, are the great vessels involved in systemic circulation. These veins return deoxygenated blood from the body into the heart, emptying it into the right atrium. The venae cavae are not separated from the right atrium by valves.

Superior Vena Cava

The superior vena cava is a large, short vein that carries deoxygenated blood from the upper half of the body to the right atrium. The right and left subclavian veins, jugular veins, and thyroid veins feed into the superior vena cava. The subclavian veins are significant because the thoracic lymphatic duct drains lymph fluid into the subclavian veins, making the superior vena cava a site of lymph fluid recirculation into the plasma. The superior vena cava begins above the heart.

Inferior Vena Cava

The inferior vena cava is the largest vein in the body and carries deoxygenated blood from the lower half of the body into the heart. The left and right common iliac veins converge to form the inferior vena cava at its lowest point. The inferior vena cava begins posterior to the abdominal cavity and travels to the heart next to the abdominal aorta. Along the way up the body from the iliac veins, the renal and suprarenal veins (kidney and adrenal glands), lumbar veins (from the back), and hepatic veins (from the liver) all drain into the inferior vena cava.

The Aorta

The aorta is the largest of the arteries in systemic circulation. Blood is pumped from the left ventricle through the aortic valve into the aorta. The aorta is a highly elastic artery and is able to dilate and constrict in response to blood pressure and volume. When the left ventricle contracts to force blood through the aortic valve into the aorta, the aorta expands. This expansion provides potential energy to help maintain blood pressure during diastole, when the aorta passively contracts. Blood pressure is highest in the aorta and diminishes through circulation, reaching its lowest points at the end of venous circulation. The difference in pressure between the aorta and right atrium accounts for blood flow in the circulation, as blood flows from areas of high pressure to areas of low pressure.


Components of the Aorta

The aortic arch contains peripheral baroreceptors (pressure sensors) and chemoreceptors (chemical sensors) that relay information concerning blood pressure, blood pH, and carbon dioxide levels to the medulla oblongata of the brain. This information is processed by the brain and the autonomic nervous system mediates the homeostatic responses that involve feedback in the lungs and kidneys. The aorta extends around the heart and travels downward, diverging into the iliac arteries. The five components of the aorta are:

1. The ascending aorta lies between the heart and the arch of aorta. It breaks off into the aortic sinuses, some of which form the coronary arteries.
2. The arch of aorta is the peak of the aorta, which breaks off into the left carotid artery, brachiocephalic trunk, and the left subclavian artery.
3. The descending aorta is the section from the arch of aorta to the point where it divides into the common iliac arteries. It is subdivided into the thoracic and abdominal aorta.
4. The thoracic aorta is the part of the descending aorta above the diaphragm. It branches off into the bronchial, mediastinal, esophageal, and phrenic arteries.
5. The abdominal aorta is the part of the descending aorta below the diaphragm, which divides into the iliac arteries and branches into the renal and suprarenal arteries. This part of the aorta is vulnerable to bursting and hemorrhage (aneurysm) from persistently high blood pressure.

The Pulmonary Arteries

The pulmonary arteries carry deoxygenated blood from the right ventricle into the alveolar capillaries of the lungs to unload carbon dioxide and take up oxygen. These are the only arteries that carry deoxygenated blood, and are considered arteries because they carry blood away from the heart. The short, wide vessel branches into the left and right pulmonary arteries that deliver deoxygenated blood to the respective lungs. Blood first passes through the pulmonary valve as it is ejected into the pulmonary arteries.



Pulmonary circuit: Diagram of pulmonary circulation. Oxygen-rich blood is shown in red; oxygen-depleted blood in blue.

The Pulmonary Veins

The pulmonary veins carry oxygenated blood from the lungs to the left atrium of the heart. Despite carrying oxygenated blood, this great vessel is still considered a vein because it carries blood towards the heart. Four pulmonary veins enter the left atrium. The right pulmonary veins pass behind the right atrium and superior vena cava while the left pass in front of the descending thoracic aorta. The pulmonary arteries and veins are both considered part of pulmonary circulation.

This page titled 17.1E: Great Vessels of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1F: Myocardial Thickness and Function

The myocardium (cardiac muscle) is the thickest section of the heart wall and contains cardiomyocytes, the contractile cells of the heart.

Learning Objectives

- Describe myocardial thickness and function in the heart

Key Points

- Coordinated contraction of cardiomyocytes causes the heart to contract and expel blood into circulation.
- The myocardium is thickest in the left ventricle, as the left ventricle must create a lot of pressure to pump blood into the aorta and throughout systemic circulation.
- Cardiac muscle has a high density of mitochondria and a large blood supply, which keep it functioning continuously.
- Sarcomeres are the basic unit of muscle tissue, composed primarily of actin and myosin myofibrils.
- The heart's thickness may change over time and adapt to compensate for diseases, which often results in heart failure.

Key Terms

- **Cardiac hypertrophy**: An adaptive disease in which the walls of the heart become too thick to pump blood effectively. It is generally a complication of hypertension.
- **sarcomere**: The basic contractile unit of contractile muscle, which contains myofibril filaments made out of myosin and actin, the two proteins that slide past one another to cause a muscle contraction.
- **cardiomyocyte**: A cardiac muscle cell (myocyte) in the heart.


The myocardium, or cardiac muscle, is the thickest section of the heart wall and contains cardiomyocytes, the contractile cells of the heart. As a type of muscle tissue, the myocardium is unique among all other muscle tissues in the human body. The thickness of the myocardium determines the strength of the heart's ability to pump blood.

Myocardial Histology

The structure of cardiac muscle shares some characteristics with skeletal muscle, but has many distinctive features of its own. Cardiomyocytes are shorter than skeletal myocytes and have fewer nuclei. Each muscle fiber connects to the plasma membrane (sarcolemma) with distinctive tubules (T-tubule). At these T-tubules, the sarcolemma is studded with a large number of calcium channels which allow calcium ion exchange at a rate much faster than that of the neuromuscular junction in skeletal muscle. The flux of calcium ions into the muscle cells causes stimulates an action potential, which causes the cells to contract.

Cardiac muscle, like skeletal muscle, is comprised of sarcomeres, the basic, contractile units of muscle. Sarcomeres are composed of long, fibrous proteins that slide past each other when the muscles contract and relax. Two of the important proteins found in sarcomeres are myosin, which forms the thick filament, and actin, which forms the thin filament. Myosin has a long, fibrous tail and a globular head that binds to actin. The myosin head also binds to ATP, the source of energy for cellular metabolism, and is required for the cardiomyocytes to sustain themselves and function normally. Together, myosin and actin form myofibril filaments, the elongated, contractile threads found in muscle tissue. Cardiac muscle and skeletal muscle both contain the protein myoglobin, which stores oxygen.

Cardiac muscle is adapted to be highly resistant to fatigue. Cardiomyocytes have a large number of mitochondria, enabling continuous aerobic respiration. Cardiac muscle also has a large blood supply relative to its size, which provides a continuous stream of nutrients and oxygen while providing ample removal of metabolic waste.




Cardiac Muscle: The tissue structure of cardiac muscle contains sarcomeres that are made of myofibrils with intercalated disks, that contain cardiomyocytes and have many mitocondria.

Myocardial Thickness

The myocardium has variable levels of thickness within the heart. Chambers of the heart with a thicker myocardium are able to pump blood with more pressure and force compared to chambers of the heart with a thinner myocardium. The myocardium is thinnest within the atria, as these chambers primarily fill through passive blood flow. The right ventricle myocardium is thicker than the atrial myocardium, as this muscle must pump all blood returning to the heart into the lungs for oxygenation. The myocardium is thickest in the left ventricle, as this chamber must create substantial pressure to pump blood into the aorta and throughout systemic circulation.

The thickness of the myocardium may change in some individuals as a compensatory adaptation to disease, either thickening and becoming stiff or becoming thinner and flabby. Cardiac hypertrophy is a common result of hypertension (high blood pressure) in which the cells of the myocardium enlarge as an adaptive response to pumping against the higher pressure. Eventually, hypertrophy may become so severe that heart failure occurs when the heart becomes so stiff that it can no longer pump blood. A flabby heart is typically the result of myocardial infections, in which the heart muscle becomes so weak that it cannot efficiently pump blood, which also leads to heart failure.



Right Ventricular Hypertrophy: If the heart adapts to become too thick, it will not be able to pump blood as efficiently, and heart failure may occur.

This page titled 17.1F: Myocardial Thickness and Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.1G: Fibrous Skeleton of the Heart

The cardiac skeleton, also known the heart's fibrous skeleton, consists of dense connective tissue in the heart that separates the atria from the ventricles.

Learning Objectives

- Describe composition and function of the heart's fibrous skeleton

Key Points

- The cardiac skeleton consists of four bands of dense connective tissue, called fibrous rings, that surround the base of the pulmonary trunk, aorta, and mitral and tricuspid valves.
- The heart 's fibrous skeleton stops the flow of electrical currents between the chambers of the heart so that it only flows through the atrioventricular (AV) bundle. This causes a delay in depolarization so that the ventricles contract after they fill with blood.
- The AV bundle is a bundle of electrically-connected cardiomyocytes that transmit impulses from the AV node to the apex of the heart. It is located in the center of the cardiac skeleton.
- The cardiac skeleton consists mainly of the protein collagen, a glycoprotein found in all connective tissues.

Key Terms

- **collagen**: A glycoprotein that forms elongated fibers, usually found in the extracellular matrix of connective tissue.
- **fibrous rings**: Four dense bands of tough elastic tissue that encircle the bases of the valves of the heart.

The cardiac skeleton, or fibrous skeleton of the heart, is the structure of dense connective tissue that separates the atria from the ventricles. The fibrous skeleton provides critical support for the heart and separates the flow of electrical impulses through the heart.

Fibrous Ring Structure



Fibrous Rings of the Heart: Transverse section of the heart showing the fibrous rings surrounding the valves.

The primary structure of cardiac skeleton consists of four dense bands of tough elastic tissue called fibrous rings that encircle the bases of the heart valves. The fibrous skeleton is composed primarily of collagen, a protein found in every type of connective tissue in the human body. There are four fibrous rings:

- The aortic ring encircles the aortic valve. It provides support for the aortic valve so that it is open, yet does not have backflow.
- The pulmonary ring encircles the pulmonary valve. Similar to the aortic ring, it provides structural support for the pulmonary valve.
- The left fibrous ring encircles the bicuspid valve. This ring is the thickest and strongest of all the fibrous rings due to the thickness of the left ventricle, which requires more structural support than the other chambers of the heart. It also surrounds the coronary arteries and AV node.
- The right fibrous ring encircles the tricuspid valve. It also surrounds the coronary arteries and AV node.

The fibrous skeleton provides a great amount of structural and functional support for the valves of the heart by enabling them to stay open and provides a point of attachment for the valves to the wall of the heart.

Electrical Functions

The fibrous skeleton of the heart acts as an insulator for the flow of electrical current across the heart. It stops the flow of electricity between the different chambers of the heart so that electrical impulses do not flow directly between the atria and ventricles. The sinoatrial (SA) node lies on the top of the heart, while the AV node is located at the center of the fibrous skeleton, the only path by which electricity can flow through the heart.

This electrical separation is essential for cardiac function, because electrical impulses flow from the top of the heart to the bottom of the heart. The separation allows the AV node and AV bundle to delay the wave of depolarization so that the atria can contract and assist in ventricular filling before the ventricles themselves depolarize and contract. Without the fibrous skeleton of the heart, the heart's ability to pump blood would be considerably less efficient since the ventricles would contract before filled to capacity. The fibrous skeleton of the heart also protects against cardiac arrhythmias.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human heart. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_heart%23Structure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...em%23The_Heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- heart. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myocardium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myocardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventricle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventricle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Di...%23globalusage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fibrous pericardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fibrous_pericardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Serous pericardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Serous_pericardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pericardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pericardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...%23Pericardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pericardium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pericardium**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- fibrous pericardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/fibrous%20pericardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- serous pericardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/serous%20pericardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Di...%23globalusage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...01/Gray968.png. **License**: *Public Domain: No Known Copyright*
- Epicardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Epicardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Myocardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Myocardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...em%23The_Heart. **License**: *CC BY-SA: Attribution-ShareAlike*
- Endocardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endocardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- epicardium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/epicardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- endocardium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/endocardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardiomyocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cardiomyocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/endothelial%20cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Di...%23globalusage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...01/Gray968.png. **License**: *Public Domain: No Known Copyright*
- Heart ant wall infarction. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...infarction.jpg. **License**: *CC BY: Attribution*
- Atrium (heart). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Atrium_(heart). **License**: *CC BY-SA: Attribution-ShareAlike*
- Ventricle (heart). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventricle_(heart). **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...Heart_Chambers. **License**: *CC BY-SA: Attribution-ShareAlike*
- systole. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/systole. **License**: *CC BY-SA: Attribution-ShareAlike*
- pulsatile. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pulsatile. **License**: *CC BY-SA: Attribution-ShareAlike*
- diastole. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/diastole. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Di...%23globalusage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/01/Gray968.png. **License**: *Public Domain: No Known Copyright*
- Heart ant wall infarction. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Heart_ant_wall_infarction.jpg. **License**: *CC BY: Attribution*
- Heart diagram-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Heart_diagram-en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_cardiovascular_system%23The_Cardiovascular_Pathways. **License**: *CC BY-SA: Attribution-ShareAlike*
- Aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Aorta. **License**: *CC BY-SA: Attribution-ShareAlike*
- Inferior vena cava. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Inferior_vena_cava. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vena cavae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vena_cavae. **License**: *CC BY-SA: Attribution-ShareAlike*

- Pulmonary artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Superior vena cava. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Superior_vena_cava. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary vein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- venae cavae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/venae%20cavae. **License**: *CC BY-SA: Attribution-ShareAlike*
- pulmonary vessels. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pulmonary%20vessels. **License**: *CC BY-SA: Attribution-ShareAlike*
- aorta. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/aorta. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg%23globalusage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/01/Gray968.png. **License**: *Public Domain: No Known Copyright*
- Heart ant wall infarction. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Heart_ant_wall_infarction.jpg. **License**: *CC BY: Attribution*
- Heart diagram-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Heart_diagram-en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_pulmonary_circuit.jpg. **License**: *Public Domain: No Known Copyright*
- Illu systemic circuit. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ic_circuit.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Syncytium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Syncytium. **License**: *CC BY-SA: Attribution-ShareAlike*
- T-tubules. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/T-tubules. **License**: *CC BY-SA: Attribution-ShareAlike*
- Myocardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Myocardium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sarcomere. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sarcomere. **License**: *CC BY-SA: Attribution-ShareAlike*
- Excitation-contraction coupling. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Excitat...ction_coupling. **License**: *CC BY-SA: Attribution-ShareAlike*
- sarcomere. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sarcomere. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardiomyocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cardiomyocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- myoglobin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/myoglobin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg%23globalusage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/01/Gray968.png. **License**: *Public Domain: No Known Copyright*
- Heart ant wall infarction. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Heart_ant_wall_infarction.jpg. **License**: *CC BY: Attribution*
- Heart diagram-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Heart_diagram-en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_pulmonary_circuit.jpg. **License**: *Public Domain: No Known Copyright*
- Illu systemic circuit. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Illu_systemic_circuit.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Right_Ventricular_hypertrophy.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ventricular_hypertrophy#/media/File:Right_Ventricular_hypertrophy.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cardiac_Muscle.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_muscle#/media/File:Cardiac_Muscle.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cardiac skeleton. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_skeleton. **License**: *CC BY-SA: Attribution-ShareAlike*
- collagen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/collagen. **License**: *CC BY-SA: Attribution-*

ShareAlike
- fibrous rings. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fibrous%20rings**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiac skeleton. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cardiac%20skeleton**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Diagram of the human heart (cropped). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg%23globalusage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Transverse Thorax. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/0/01/Gray968.png**. **License**: ***Public Domain: No Known Copyright***
- Heart ant wall infarction. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Heart_ant_wall_infarction.jpg**. **License**: ***CC BY: Attribution***
- Heart diagram-en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Heart_diagram-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pulmonary_circuit.jpg**. **License**: ***Public Domain: No Known Copyright***
- Illu systemic circuit. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Illu_systemic_circuit.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Right_Ventricular_hypertrophy.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ventricular_hypertrophy#/media/File:Right_Ventricular_hypertrophy.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cardiac_Muscle.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cardiac_muscle#/media/File:Cardiac_Muscle.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray495. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray495.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 17.1G: Fibrous Skeleton of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.2: CIRCULATION AND HEART VALVES

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

17.2A: Heart Circulation

17.2B: Operation of Atrioventricular Valves

17.2C: Operation of Semilunar Valves

17.2D: Systemic and Pulmonary Circulation

This page titled 17.2: Circulation and Heart Valves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).


17.2A: Heart Circulation

Learning Objectives

- Describe circulation of blood in the heart

Atherosclerotic plaques in a coronary artery will slowly occlude (block) the vessel. As the vessel diameter narrows, less blood and oxygen will pass through and a region of the myocardium will consequently not receive an adequate supply of oxygen. This could result in angina and ultimately a myocardial infarction.

Coronary circulation is the circulation of blood in the blood vessels of the heart muscle. The vessels that deliver oxygen-rich blood to the myocardium are known as coronary arteries. The vessels that remove the deoxygenated blood from the heart muscle are known as cardiac veins. The blood supply to the heart is greater than that of other body tissues since the heart has a constant metabolic demand that must be satisfied to keep the heart pumping at all times.



Coronary Circulation: Coronary arteries labeled in red text and other landmarks in blue text.

Coronary Artery Structure

The coronary arteries originate from the left side of the heart descending from the aorta. There are multiple coronary arteries derived from the larger right and left coronary arteries. For example, important coronary arteries that branch off from the larger arteries include the left anterior descending (LAD) coronary and the right posterior coronary.

Coronary arteries run both along the surface of the heart and deep within the myocardium, which has the greatest metabolic demands of all the heart tissues due to its muscle content. Epicardial coronary arteries, which run along on the surface of the heart, are capable of autoregulating vasodilation and vasoconstriction to maintain coronary blood flow at appropriate levels to fit the metabolic demands of the heart muscle. These vessels are relatively narrow and thus vulnerable to blockage, which may cause a myocardial infarction. Subendocardial coronary arteries run deep within the myocardium to provide oxygen throughout the muscle tissue of the cardiac wall.

Systole and Diastole

In systole, the ventricular myocardium contracts, generating high intraventricular pressure and compressing the subendocardial

coronary vessels while allowing the epicardial coronary vessels to remain fully open. With the subendocardial coronary vessels compressed, blood flow essentially stops below the surface of the myocardium.

In diastole, the ventricular myocardium contracts, lowering the intraventricular pressure and allowing the subendocardial vessels to become open again. Due to the high pressures generated in the ventricular myocardium during systole, most myocardial tissue perfusion occurs during diastole. Additionally, catecholamines such as norephinephrine, which normally cause vasoconstriction will instead cause vasodilation within the coronary arteries. This mechanism is due to beta-adrenergic receptors in the coronary arteries and helps enable the increased cardiac output associated with fight-or-flight responses.

Myocardial Infarctions

A myocardial infarction (heart attack) may be caused by prolonged ischemia (oxygen deprivation) in the heart, which occurs due to blockage of any of the coronary arteries. Since there is very little unnecessary blood supply to the myocardium, blockage of these vessels can cause serious damage. When these vessels become blocked, the myocardium becomes oxygen-deprived, a condition called ischemia. Brief periods of ischemia in the heart are associated with intense chest pain called angina, which may either be transient if the clot breaks up on its own or stable if it does not. As the time period of ischemia increases, the hypoxic conditions cause muscle tissue to die, causing a myocardial infarction (heart attack).

Myocardial infarction is one of the most common causes of death worldwide. The clots that cause the infarction are usually the result of ruptured atherosclerotic plaques that break off and occlude the coronary arteries, but arterial thrombosis from injury or pooled blood may also cause a heart attack. The tissues of the heart do not regenerate, so those that survive a myocardial infarction will generally have scar tissue in their myocardium and may be more susceptible to other heart problems in the future.

Key Points

- The vessels that supply blood to the myocardium are called coronary arteries and coronary veins.
- The left and right coronary arteries branch off into smaller arteries, such as the important left anterior descending (LAD) coronary artery.
- The vessels that deliver oxygen-rich blood to the myocardium are known as coronary arteries. The vessels that remove the deoxygenated blood from the heart muscle are known as cardiac veins.
- Most tissue perfusion in the heart occurs when the coronary arteries open during diastole.
- Failure of the coronary arteries to provide sufficient blood supply to the heart can lead to ischemia, angina, and myocardial infarction.
- Norepinephrine will cause vasodilation in the coronary arteries but vasoconstriction in the other arteries of the body.
- Myocardial infarctions are the result of ruptured artherosclerotic plaques or arterial thrombosis, which cause the death of heart tissue from prolonged ischemia.

Key Terms

- **myocardial infarction**: Necrosis of heart muscle caused by an interruption to the supply of blood to the heart, often as a result of prolonged ischemia.
- **ischemia**: Oxygen deprivation in tissues due to mechanical obstruction of the blood supply, such as by a narrowed or blocked artery or clot.
- **angina**: Chest pain that indicates ischemia in the heart. It may be either transient (unstable) or stable, and stable anginas typically lead to infarction.


This page titled 17.2A: Heart Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.2B: Operation of Atrioventricular Valves

Learning Objectives

- Describe the operation of the atrioventricular valves: bicuspid (mitral) and tricuspid

A heart valve allows blood flow in only one direction through the heart, and the combination of the atrioventricular and semi-lunar heart valves determines the pathway of blood flow. Valves open or close based on pressure differences across the valve. The atrioventricular (AV) valves separate the atria from the ventricles on each side of the heart and prevent backflow of blood from the ventricles into the atria during systole.



Cross section of heart indicating heart valves: The four valves determine the pathway of blood flow (indicated by arrows) through the heart

Subvalvular Apparatus

The subvalvular apparatus describes the structures beneath the AV valves that prevent the valves prom prolapsing. Valve prolapse means that the valves do not close properly, which may cause regurgitation or backflow of blood from the ventricle back into the atria, which is inefficient. The subvalvular apparatus includes the chordae tendineae and the papillary muscles. The AV valves are anchored to the wall of the ventricle by chordae tendineae (heartstrings), small tendons that prevent backflow by stopping the valve leaflets from inverting. The chordae tendineae are inelastic and attached at one end to the papillary muscles and at the other end to the valve cusps.

Papillary muscles are finger-like projections from the wall of the ventricle that anchor the chordae tendineae. This connection provides tension to hold the valves in place and prevent them from prolapsing into the atria when they close, preventing the risk of regurgitation. The subvalvular apparatus has no effect on the opening and closing of the valves, which is caused entirely by the pressure gradient of blood across the valve as blood flows from high pressure to low pressure areas.

The Mitral Valve

The mitral valve is on the left side of the heart and allows the blood to flow from the left atrium into the left ventricle. It is also known as the bicuspid valve because it contains two leaflets (cusps). The relaxation of the ventricular myocardium and the contraction of the atrial myocardium cause a pressure gradient that allows for rapid blood flow from the left atrium into the

left ventricle across the mitral valve. Atrial systole (contraction) increases the pressure in the atria, while ventricular diastole (relaxation) decreases the pressure in the ventricle, causing pressure-induced flow of blood across the valve. The mitral annulus, a ring around the mitral valve, changes in shape and size during the cardiac cycle to prevent backflow. The ring contracts at the end of atrial systole due to the contraction of the left atrium around it, which aids in bringing the leaflets together to provide firm closure during ventricular systole.

The Tricuspid Valve

The tricuspid valve is the three-leaflet valve on the right side of the heart between the right atrium and the right ventricle and stops the backflow of blood between the two. The tricuspid valve functions similarly to the bicuspid valve except that three chordae tendineae connect the cusps of the valve to three papillary muscles, rather than the pair that connects the bicuspid valve. Blood passes through the tricuspid valve the same as it does through the bicuspid valve, based on a pressure gradient from high pressure to low pressure during systole and diastole.

The reason that the valves have different numbers of leaflets is not fully understood, but may arise from differences in tissue structure and pressure that occur during fetal development.

Key Points

- The atrioventricular valves, the bicuspid (mitral) and tricuspid valves, separate the atria from the ventricles.
- The bicuspid valve is on the left side of the heart and the tricuspid valve is on the right side of the heart.
- Blood flows through an atrioventricular (AV) valve when blood pressure in the atria becomes high during atrial systole and blood pressure in the ventricles becomes low enough during ventricular diastole, creating a blood pressure gradient.
- Papillary muscles, finger-like projections from the wall of the ventricles, connect the chordae tendineae (heartstrings) to the cusps of the atrioventricular valves. This connection prevents the valve from prolapsing under pressure.
- Papillary muscles, together with the chordae tendineae, make up the subvalvular apparatus.

Key Terms

- **Atrioventricular valves**: These valves separate the atria from the ventricles on each side of the heart and prevent backflow from the ventricles into the atria during systole. They include the mitral and tricuspid valves.
- **Subvalvular apparatus**: The papillary muscles and the chordae tendineae, known as the subvalvular apparatus, hold the valves closed so that they do not prolapse.
- **mitral valve**: The bicuspid valve that divides the left atrium and left ventricle of the heart

This page titled 17.2B: Operation of Atrioventricular Valves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.2C: Operation of Semilunar Valves

The semilunar valves allow blood to be pumped into the major arteries while preventing backflow of blood from the arteries into the ventricles.

Learning Objectives

- Describe the operation of the semilunar valves

Key Points

- The semilunar valves act to prevent backflow of blood from the arteries to the ventricles during ventricular diastole and help maintain pressure on the major arteries.
- The aortic semilunar valve separates the left ventricle from the opening of the aorta.
- The aortic and pulmonary valves are semilunar valves which separate the ventricles from the aorta and pulmonary artery, respectively.
- Partial pressure gradient changes during systole and diastole cause the opening and closing of the valves.
- Valve stenosis is when valves narrow and can't open fully, while regurgitation is when they cannot close completely. In both instances, the heart must work harder to compensate for the faulty valves.

Key Terms


- **semilunar valves**: Located at the base of both the trunk of the pulmonary artery and the aorta, and prevent backflow of blood from the arteries into the ventricles.
- **stenosis**: The narrowing of valves, which prevents them from opening completely.

The semilunar valves are located at the connections between the pulmonary artery and the right ventricle, and the aorta and the left ventricle. These valves allow blood to be pumped forward into the arteries, but prevent backflow of blood from the arteries into the ventricles. These valves do not have subvalvular apparatus and are more similar to the semilunar valves in veins and lymphatic vessels than to atrioventricular (AV) valves.

The semilunar valves act in concert with the AV valves to direct blood flow through the heart. When the atrioventricular valves are open, the semi lunar valves are shut and blood is forced into the ventricles. When the AV valves shut, the semilunar valves open, forcing blood into the aorta and pulmonary artery. The mechanism for this process depends on blood pressure gradients in the heart, which provide the force that pushes blood through the semilunar valves.

The Aortic Valve

The aortic valve separates the left ventricle from the aorta and has three cusps. During ventricular systole, pressure rises in the left ventricle. When the pressure in the left ventricle exceeds the pressure in the aorta, the aortic valve opens and blood flows from the left ventricle into the aorta. When ventricular systole ends, pressure in the left ventricle drops rapidly, and the valve closes due to a lack of pressure imposed on them from the left ventricle. Blood pressure within the aorta following systole also causes the closing of the valve. The closing of the aortic valve produces a sound that is a component of the second heart sound.



Heart viewed from above: This anterior view of the heart indicates the semilunar valves, and the aortic and pulmonary valves.

The Pulmonary Valve

The pulmonary valve (also called the pulmonic valve), which also has three cusps, separates the right ventricle from the pulmonary artery. Similar to the aortic valve, the pulmonary valve opens in ventricular systole, when the pressure in the right ventricle exceeds the pressure in the pulmonary artery. When ventricular systole ends, pressure in the right ventricle drops rapidly, and the pressure in the pulmonary artery forces the pulmonary valve to close. The closure of the pulmonary valve also produces a sound, however it is softer than the aortic sound because the blood pressure in the right side of the heart is lower compared to the left side, due to the differences between pulmonary and systemic circulation.

Valve Problems

Valves are vulnerable to several conditions that impair their normal functions. Two of the most common problems with the semilunar valves are stenosis and regurgitation. Valve stenosis refers to narrowing of the valves, which prevents the valve from opening fully, causing an obstruction of blood flow. Valve stenosis is often caused by calcium buildup and scarring from rheumatic fever, and may cause cardiac hypertrophy and heart failure. Valve regurgitation is backflow through the valves that occurs when when they cannot close completely. It is the cause of most heart murmurs, and is generally a minor problem, but if severe enough, it can cause heart failure. Stenosis and regurgitation can occur in both the semilunar and atrioventricular valves.

This page titled 17.2C: Operation of Semilunar Valves is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.2D: Systemic and Pulmonary Circulation

The cardiovascular system has two distinct circulatory paths, pulmonary circulation and systemic circulation.

Learning Objectives

- Distinguish between the systemic and pulmonary circulation circuits

Key Points

- The cardiovascular system is composed of two circulatory paths: pulmonary circulation, the circuit through the lungs where blood is oxygenated, and systemic circulation, the circuit through the rest of the body to provide oxygenated blood.
- In the pulmonary circulation, blood travels through capillaries on the alveoli, air sacs in the lungs which allow for gas exchange.
- As blood flows through circulation, the size of the vessel decreases from artery / vein, to arteriole / venule, and finally to capillaries, the smallest vessels for gas and nutrient exchange.
- Systemic and pulmonary circulation transition to the opposite type of circulation when they return blood to the opposite side of the heart.
- Systemic circulation is a much larger and higher pressure system than pulmonary circulation.

Key Terms


- **alveoli**: Air sacs in the lungs that provide the surface for gas exchange between the air and capillaries.
- **pulmonary circulation**: The part of blood circulation which carries oxygen-depleted blood away from the heart, to the lungs, and returns oxygenated blood back to the heart.
- **systemic circulation**: The part of blood circulation that carries oxygenated blood away from the heart, to the body, and returns deoxygenated blood back to the heart.

The cardiovascular system is composed of two circulatory paths: pulmonary circulation, the circuit through the lungs where blood is oxygenated; and systemic circulation, the circuit through the rest of the body to provide oxygenated blood. The two circuits are linked to each other through the heart, creating a continuous cycle of blood through the body.


Pulmonary Circulation

Pulmonary circulation is the movement of blood from the heart to the lungs for oxygenation, then back to the heart again. Oxygen-depleted blood from the body leaves the systemic circulation when it enters the right atrium through the superior and inferior venae cavae. The blood is then pumped through the tricuspid valve into the right ventricle. From the right ventricle, blood is pumped through the pulmonary valve and into the pulmonary artery. The pulmonary artery splits into the right and left pulmonary arteries and travel to each lung.

At the lungs, the blood travels through capillary beds on the alveoli where gas exchange occurs, removing carbon dioxide and adding oxygen to the blood. Gas exchange occurs due to gas partial pressure gradients across the the alveoli of the lungs and the capillaries interwoven in the alveoli. The oxygenated blood then leaves the lungs through pulmonary veins, which returns it to the left atrium, completing the pulmonary circuit. As the pulmonary circuit ends, the systemic circuit begins.



Alveoli: A diagram of the alveoli, showing the capillary beds where gas exchange with the blood occurs.



Pulmonary circuit: Diagram of pulmonary circulation. Oxygen-rich blood is shown in red; oxygen-depleted blood in blue.

Systemic Circulation

Systemic circulation is the movement of blood from the heart through the body to provide oxygen and nutrients to the tissues of the body while bringing deoxygenated blood back to the heart. Oxygenated blood enters the left atrium from the pulmonary veins. The blood is then pumped through the mitral valve into the left ventricle. From the left ventricle, blood is pumped through the aortic valve and into the aorta, the body's largest artery. The aorta arches and branches into major arteries to the upper body before passing through the diaphragm, where it branches further into the illiac, renal, and suprarenal arteries which supply the lower parts of the body.

The arteries branch into smaller arteries, arterioles, and finally capillaries. Gas and nutrient exchange with the tissues occurs within the capillaries that run through the tissues. Metabolic waste and carbon dioxide diffuse out of the cell into the blood, while oxygen and glucose in the blood diffuses out of the blood and into the cell. Systemic circulation keeps the metabolism of every organ and every tissue in the body alive, with the exception of the parenchyma of the lungs, which are supplied by pulmonary circulation.

The deoxygenated blood continues through the capillaries which merge into venules, then veins, and finally the venae cavae,

which drain into the right atrium of the heart. From the right atrium, the blood will travel through the pulmonary circulation to be oxygenated before returning gain to the system circulation, completing the cycle of circulation through the body. The arterial component of systemic circulation the highest blood pressures in the body. The venous component of systemic circulation has considerably lower blood pressure in comparison, due to their distance from the heart, but contain semi-lunar valves to compensate. Systemic circulation as a whole is a higher pressure system than pulmonary circulation simply because systemic circulation must force greater volumes of blood farther through the body compared to pulmonary circulation.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Coronary circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coronary_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ischemia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ischemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coronary circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Coronary%20circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myocardial infarction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myocardial_infarction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coronary arteries. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coronary_arteries.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Semilunar valves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Semilunar_valves%23Atrioventricular_or_cuspid_valves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mitral valve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mitral_valve%23Normal_physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Atrioventricular valves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Atrioventricular%20valves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Subvalvular apparatus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Subvalvular%20apparatus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mitral valve. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mitral_valve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coronary arteries. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coronary_arteries.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pediatric Cardiothoracic Surgery - Heart Valve Disease. **Provided by**: University of California San Francisco. **Located at**: **http://pediatricct.surgery.ucsf.edu/conditions--procedures/heart-valve-disease.aspx**. **License**: ***Public Domain: No Known Copyright***
- Semilunar valves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Semilunar_valves%23Semilunar_valves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- aortic valve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/aortic%20valve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Semilunar valves. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Semilunar%20valves**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pulmonary valve. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pulmonary%20valve**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coronary arteries. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coronary_arteries.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pediatric Cardiothoracic Surgery - Heart Valve Disease. **Provided by**: University of California San Francisco. **Located at**: **http://pediatricct.surgery.ucsf.edu/conditions--procedures/heart-valve-disease.aspx**. **License**: ***Public Domain: No Known Copyright***
- Gray494. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray494.png**. **License**: ***Public Domain: No Known Copyright***
- Systemic circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Systemic_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pulmonary circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pulmonary_circulation**. **License**: ***CC***

BY-SA: Attribution-ShareAlike
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physics/de...ion/alveoli--2**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- systemic circulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/systemic_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pulmonary circulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pulmonary_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coronary arteries. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Coronary_arteries.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pediatric Cardiothoracic Surgery - Heart Valve Disease. **Provided by**: University of California San Francisco. **Located at**: **http://pediatricct.surgery.ucsf.edu/conditions--procedures/heart-valve-disease.aspx**. **License**: ***Public Domain: No Known Copyright***
- Gray494. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray494.png**. **License**: ***Public Domain: No Known Copyright***
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...ry_circuit.jpg**. **License**: ***Public Domain: No Known Copyright***
- Alveolus diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Al...us_diagram.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 17.2D: Systemic and Pulmonary Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.3: CARDIAC MUSCLE TISSUE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

17.3A: Mechanism and Contraction Events of Cardiac Muscle Fibers
17.3B: Microscopic Anatomy
17.3C: Energy Requirements

This page titled 17.3: Cardiac Muscle Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.3A: Mechanism and Contraction Events of Cardiac Muscle Fibers

Cardiac muscle fibers undergo coordinated contraction via calcium-induced calcium release conducted through the intercalated discs.

Learning Objectives

- Describe the mechanism and contraction events of cardiac muscle fibers

Key Points

- Cardiac muscle fibers contract via excitation-contraction coupling, using a mechanism unique to cardiac muscle called calcium -induced calcium release.
- Excitation-contraction coupling describes the process of converting an electrical stimulus (action potential) into a mechanical response (muscle contraction).
- Calcium-induced calcium release involves the conduction of calcium ions into the cardiomyocyte, triggering further release of ions into the cytoplasm.
- Calcium prolongs the duration of muscle cell depolarization before repolarization occurs.Contraction in cardiac muscle occurs due to the the binding of the myosin head to adenosine triphosphate (ATP), which then pulls the actin filaments to the center of the sarcomere, the mechanical force of contraction.

Key Terms

- **excitation contraction coupling (ECC)**: The physiological process of converting an electrical stimulus to a mechanical response.
- **calcium-induced calcium release (CICR)**: A process whereby calcium can trigger release of further calcium from the muscle sarcoplasmic reticulum.

Cardiomyocytes are capable of coordinated contraction, controlled through the gap junctions of intercalated discs. The gap junctions spread action potentials to support the synchronized contraction of the myocardium. In cardiac, skeletal, and some smooth muscle tissue, contraction occurs through a phenomenon known as excitation contraction coupling (ECC). ECC describes the process of converting an electrical stimulus from the neurons into a mechanical response that facilitates muscle movement. Action potentials are the electrical stimulus that elicits the mechanical response in ECC.

Calcium-Induced Calcium Release

In cardiac muscle, ECC is dependent on a phenomenon called calcium-induced calcium release (CICR), which involves the influx of calcium ions into the cell, triggering further release of ions into the cytoplasm. The mechanism for CIRC is receptors within the cardiomyocyte that bind to calcium ions when calcium ion channels open during depolarization, releasing more calcium ions into the cell.


Similarly to skeletal muscle, the influx of sodium ions causes an initial depolarization; however, in cardiac muscle, the influx of calcium ions sustains the depolarization so that it lasts longer. CICR creates a “plateau phase” in which the cell’s charge stays slightly positive (depolarized) briefly before it becomes more negative as it repolarizes due to potassium ion influx. Skeletal muscle, by contrast, repolarizes immediately.

Pathway of Cardiac Muscle Contraction

The actual mechanical contraction response in cardiac muscle occurs via the sliding filament model of contraction. In the sliding

filament model, myosin filaments slide along actin filaments to shorten or lengthen the muscle fiber for contraction and relaxation. The pathway of contraction can be described in five steps:

1. An action potential, induced by the pacemaker cells in the sinoatrial (SA) and atrioventricular (AV) nodes, is conducted to contractile cardiomyocytes through gap junctions.
2. As the action potential travels between sarcomeres, it activates the calcium channels in the T-tubules, resulting in an influx of calcium ions into the cardiomyocyte.
3. Calcium in the cytoplasm then binds to cardiac troponin-C, which moves the troponin complex away from the actin binding site. This removal of the troponin complex frees the actin to be bound by myosin and initiates contraction.
4. The myosin head binds to ATP and pulls the actin filaments toward the center of the sarcomere, contracting the muscle.
5. Intracellular calcium is then removed by the sarcoplasmic reticulum, dropping intracellular calcium concentration, returning the troponin complex to its inhibiting position on the active site of actin, and effectively ending contraction as the actin filaments return to their initial position, relaxing the muscle.



Sliding Filament Model of Contraction: Muscle fibers in relaxed (above) and contracted (below) positions



Animation of Myosin and Actin: This animation shows myosin filaments (red) sliding along the actin filaments (pink) to contract a muscle cell.

This page titled 17.3A: Mechanism and Contraction Events of Cardiac Muscle Fibers is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.3B: Microscopic Anatomy

Cardiac muscle appears striated due to the presence of sarcomeres, the highly-organized basic functional unit of muscle tissue.

Learning Objectives

- Identify the microscopic anatomy of cardiac muscles

Key Points

- Cardiac muscle, composed of the contractile cells of the heart, has a striated appearance due to alternating thick and thin filaments composed of myosin and actin.
- Actin and myosin are contractile protein filaments, with actin making up thin filaments, and myosin contributing to thick filaments. Together, they are considered myofibrils.
- Myosin and actin adenosine triphosphate (ATP) binding allows for muscle contraction. It is regulated by action potentials and calcium concentrations.
- Adherens junctions, gap junctions, and desmosomes are intercalated discs that connect cardiac muscle cells. Gap junctions specifically allow for the transmission of action potentials within cells.

Key Terms

- **intercalated discs**: Junctions that connect cardiomyocytes together, some of which transmit electrical impulses between cells.
- **sarcomere**: The basic unit of contractile muscle which contains myosin and actin, the two proteins that slide past one another to cause a muscle contraction.


Cardiac muscle, like skeletal muscle, appears striated due to the organization of muscle tissue into sarcomeres. While similar to skeletal muscle, cardiac muscle is different in a few ways. Cardiac muscles are composed of tubular cardiomyocytes, or cardiac muscle cells. The cardiomyocytes are composed of tubular myofibrils, which are repeating sections of sarcomeres. Intercalated disks transmit electrical action potentials between sarcomeres.

Sarcomere Structure


A sarcomere is the basic unit of muscle tissue in both cardiac and skeletal muscle. Sarcomeres appear under the microscope as striations, with alternating dark and light bands. Sarcomeres are connected to a plasma membrane, called a sarcolemma, by T-tubules, which speed up the rate of depolarization within the sarcomere.

Individual sarcomeres are composed of long, fibrous proteins that slide past each other when the muscles contract and relax. The two most important proteins within sarcomeres are myosin, which forms a thick, flexible filament, and actin, which forms the thin, more rigid filament. Myosin has a long, fibrous tail and a globular head which binds to actin. The myosin head also binds to ATP, the source of energy for muscle movement. Actin molecules are bound to the Z-disc, which forms the borders of the sarcomere. Together, myosin and actin form myofibrils, the repeating molecular structure of sarcomeres.

Myofibril activity is required for muscle contraction on the molecular level. When ATP binds to myosin, it separates from the actin of the myofibril, which causes a contraction. Muscle contraction is a complex process regulated by calcium influx and the stimulus of electrical impulses.



Muscle Contraction and Actin-Myosin Interactions: Skeletal muscle contracts following activation by an action potential. Binding of Acetylcholine at the motor end plate leads to intracellular calcium release and interactions between myofibrils, eliciting contraction.



The Sarcomere: A single sarcomere unit with all functional areas labeled, including thick and thin filaments, Z lines, H zone, I bands, and A band.

Intercalated Discs

Intercalated discs are gap junctions that link cardiomyocytes so that electrical impulses (action potentials) can travel between cells. In a more general sense, an intercalated disk is any junction that links cells together between a gap in which no other cells exist, such as an extracellular matrix. In cardiac muscle tissue, they are also responsible for transmission of action potentials and calcium during muscle contraction. In cardiac muscle, intercalated discs connecting cardiomyocytes to the syncytium, a multinucleated muscle cell, to support the rapid spread of action potentials and the synchronized contraction of the myocardium. Intercalated discs consist of three types of cell-cell junctions, most of which are found in other tissues besides cardiac muscle:

1. Adherens junctions, which anchor actin filaments to cytoplasm of cardiomyocytes.
2. Desmosomes, which bind adhesion proteins to the cytoskeleton within cells, thus connecting the cells.
3. Gap junctions, which connect proteins to the cytoplasm of different cells and transmit action potentials between both cells, required for cellular depolarization. It is found primarily in nervous and muscular tissue.

Under light microscopy, intercalated discs appear as thin lines dividing adjacent cardiac muscle cells and running perpendicular to the direction of muscle fibers.

This page titled 17.3B: Microscopic Anatomy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.3C: Energy Requirements

Cardiac cells contain numerous mitochondria, which enable continuous aerobic respiration and production of adenosine triphosphate (ATP) for cardiac function.

Learning Objectives

- Describe the energy requirements of cardiac muscle tissue

Key Points

- The myocardium requires significant energy to contract continually over the human lifetime.
- These energy needs are met through mitochondria, myoglobins, and rich blood supply from the coronary arteries.
- The mitochondria generate ATP for the contraction of cardiomyocytes.
- Myoglobins are oxygen-storing and oxygen-transferring pigments in cardiomyocytes.
- Aerobic metabolism occurs when oxygen is present, while anaerobic respiration occurs when tissue is deprived of oxygen. Aerobic metabolism accounts for nearly all of the metabolic function in the heart, but anerobic metabolism may contribute as well.
- Glucose reservoirs and lactate recycling allow the heart to function even during malnutrition.

Key Terms


- **lactate**: A molecule produced by anaerobic respiration that can be used to produce ATP without oxygen, albeit at lower levels.
- **myoglobin**: A small globular protein containing a heme group that carries oxygen to muscles from the blood and stores reserve oxygen.

The heart muscle pumps continuously throughout life and is adapted to be highly resistant to fatigue. Cardiomyocytes contain large numbers of mitochondria, the powerhouse of the cell, enabling continuous aerobic respiration and ATP production required for mechanical muscle contraction. Cardiac muscle tissue has among the highest energy requirements in the human body (along with the brain) and has a high level of mitochondria and a constant, rich, blood supply to support its metabolic activity.

Aerobic Metabolism

Aerobic metabolism is a necessary component to support the metabolic function of the heart. Oxygen is necessary, and if even a small part of the heart is oxygen-deprived for too long, a myocardial infarction (heart attack) will occur. Coronary circulation branches from the aorta soon after it leaves the heart, and supplies the heart with the nutrients and oxygen needed to sustain aerobic metabolism. Cardiac muscle cells contain larger amounts of mitochondria than other cells in the body, enabling higher ATP production.

The heart derives energy from aerobic metabolism via many different types of nutrients. Sixty percent of the energy to power the heart is derived from fat (free fatty acids and triglycerides), 35% from carbohydrates, and 5% from amino acids and ketone bodies from proteins. These proportions vary widely with available dietary nutrients. Malnutrition will not result in the death of heart tissue in the way that oxygen deficiency will, because the body has glucose reserves that sustain the vital organs of the body and the ability to recycle and use lactate aerobically.



Myoglobin: The heme component of myoglobin, shown in orange, binds oxygen. Myoglobin provides a back-up store of oxygen to muscle cells.

Heart muscle also contains large amounts of a pigment called myoglobin. Myoglobin is similar to hemoglobin in that it contains a heme group (an oxygen binding site). Myoglobin transfers oxygen from the blood to the muscle cell and stores reserve oxygen for aerobic metabolic function in the muscle cell.

Anaerobic Metabolism

While aerobic respiration supports normal heart activity, anaerobic respiration may provide additional energy during brief periods of oxygen deprivation. Lactate, created from lactic acid fermentation, accounts for the anaerobic component of cardiac metabolism. At normal metabolic rates, about 1% of energy is derived from lactate, and about 10% under moderately hypoxic (low oxygen) conditions. Under more severe hypoxic conditions, not enough energy can be liberated by lactate production to sustain ventricular contraction, and heart failure will occur.

Lactate can be recycled by the heart and provides additional support during nutrient deprivation. Recycling lactate is very energy-efficient in the nutrient-deprived myocardium, since one NAD+ is reduced to NADH and H+ (equal to 2.5 or 3 ATP) when lactate is oxidized to pyruvate. The produced pyruvate can then be burned aerobically in the citric acid cycle (also known as the tricarboxylic acid cycle or Krebs cycle), liberating a significant amount of energy.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Myofibrils. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Myofibrils. **License**: *CC BY-SA: Attribution-ShareAlike*
- Myocardium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Myocardium%23Appearance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sarcomeres. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sarcomeres. **License**: *CC BY-SA: Attribution-ShareAlike*
- syncytium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/syncytium. **License**: *CC BY-SA: Attribution-ShareAlike*
- sarcomere. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sarcomere. **License**: *CC BY-SA: Attribution-*

ShareAlike
- intercalated discs. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intercalated%20discs**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muskel-molekulartranslation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muskel-molekulartranslation.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- calcium-induced calcium release (CICR). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/calcium...lease%20(CICR)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myocardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myocardium%23T-Tubules**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Excitation-contraction coupling. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Excitation-contraction_coupling**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myocardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myocardium%23Role_of_calcium_in_contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T-tubule. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/T-tubule**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- excitation contraction coupling (ECC). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/excitat...upling%20(ECC)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muskel-molekulartranslation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muskel-molekulartranslation.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Actin Myosin. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Actin_Myosin.gif**. **License**: ***Public Domain: No Known Copyright***
- Myocardium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myocardium%23Metabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myoglobin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lactate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- basal metabolic rate. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/basal_metabolic_rate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muskel-molekulartranslation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Muskel-molekulartranslation.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Sarcomere.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Actin Myosin. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Actin_Myosin.gif**. **License**: ***Public Domain: No Known Copyright***
- Myoglobin and heme. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Myoglobin_and_heme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 17.3C: Energy Requirements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4: PHYSIOLOGY OF THE HEART

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

17.4A: Electrical Events

17.4B: Electrocardiogram and Correlation of ECG Waves with Systole

17.4C: Heart Sounds

17.4D: Cardiac Cycle

17.4E: Cardiac Output

This page titled 17.4: Physiology of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4A: Electrical Events

Cardiac contraction is initiated in the excitable cells of the sinoatrial (SA) node by both spontaneous depolarization and sympathetic activity.

Learning Objectives

- Describe the electrical events of the heart

Key Points

- The sinoatrial (SA) and atrioventricular (AV) nodes make up the intrinsic conduction system of the heart by setting the rate at which the heart beats.
- The SA node generates action potentials spontaneously.
- The SA node fires at a normal rate of 60–100 beats per minute (bpm), and causes depolarization in atrial muscle tissue and subsequent atrial contraction.
- The AV node slows the impulses from the SA node, firing at a normal rate of 40-60 bpm, and causes depolarization of the ventricular muscle tissue and ventricular contraction.
- Sympathetic nervous stimulation increases the heart rate, while parasympathetic nervous stimulation decreases the heart rate.

Key Terms

- **pacemaker**: A structure that sets the rate at which the heart beats. Under normal conditions, the SA node serves this function for the heart.
- **atrioventricular (AV) node**: The bundle of conducting tissue that receives impulses from the SA node and delays them before stimulating depolarization in the muscles of the ventricles.

The heart's activity is dependent on the electrical impulses from the sinoatrial (SA) node and atrioventricular (AV) node, which form the intrinsic conduction system of the heart. The SA and AV nodes act as a pacemaker for the heart, determining the rate at which it beats, even without signals from the larger nervous system of the human body. The SA and AV nodes initiate the electrical impulses that cause contraction within the atria and ventricles of the heart.

Sinoatrial Node

The SA node is a bundle of nerve cells located on the outer layer of the right atria. These cells are specialized to undergo spontaneous depolarization and generation of action potentials without stimulation from the rest of the nervous system. The SA node nerve impulses travel through the atria and cause direct muscle cell depolarization and contraction of the atria. The SA node stimulates the right atria directly and stimulates the left atria through the Bachmann's bundle. The SA node impulses also travel to the AV node, which stimulates ventricular contraction.


The SA node generates its own action potentials, but may be influenced by the autonomic nervous system. Without autonomic nervous stimulation, the SA node will set the heart rate itself, acting as the primary pacemaker for the heart. The SA node fires to set a heart rate in a range of 60–100 beats per minute (bpm), a normal range that varies from person to person.

Atrioventricular Node

The AV node is a bundle of conducting tissue (not formally classified as nerve tissue) located at the junction between the atria and ventricles of the heart. The AV node receives action potentials from the SA node, and transmits them through the bundle of His, left and right bundle branches, and Purkinje fibers, which cause depolarization of ventricular muscle cells leading to ventricular contraction. The AV node slightly slows the neural impulse from the SA node, which causes a delay between depolarization of the

atria and the ventricles.

The normal firing rate in the AV node is lower than that of the SA node because it slows the rate of neural impulses. Without autonomic nervous stimulation, it sets the rate of ventricular contraction at 40–60 bpm. Certain types of autonomic nervous stimulation alter the rate of firing in the AV node. Sympathetic nervous stimulation still increases heart rate, while parasympathetic nervous stimulation decreases heart rate by acting on the AV node.



The Cardiac Conduction System: The system of nerves that work together to set the heart rate and stimulate muscle cell depolarization within the heart.

This page titled 17.4A: Electrical Events is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4B: Electrocardiogram and Correlation of ECG Waves with Systole

An electrocardiogram, or ECG, is a recording of the heart's electrical activity as a graph over a period of time.

Learning Objectives

- Describe electrocardiograms and their correlation with systole

Key Points


- An ECG is used to measure the rate and regularity of heartbeats as well as the size and position of the chambers, the presence of damage to the heart, and the effects of drugs or devices used to regulate the heart, such as a pacemaker.
- The ECG device detects and amplifies the tiny electrical changes on the skin that are caused when the heart muscle depolarizes during each heartbeat, and then translates the electrical pulses of the heart into a graphic representation.
- A typical ECG tracing of the cardiac cycle (heartbeat) consists of a P wave (atrial depolarization), a QRS complex (ventricular depolarization), and a T wave (ventricular repolarization). An additional wave, the U wave (Purkinje repolarization), is often visible, but not always.
- The ST complex is usually elevated during a myocardial infarction.
- Atrial fibrillation occurs when the P wave is missing and represents irregular, rapid, and inefficient atrial contraction, but is generally not fatal on its own.
- Ventricular fibrillation occurs when all normal waves of an ECG are missing, represents rapid and irregular heartbeats, and will quickly cause sudden cardiac death.

Key Terms

- **fibrillation**: A condition in which parts of the ECG do not appear normally, representing irregular, rapid, disorganized, and inefficient contractions of the atria or ventricles.
- **ST segment**: The line between the QRS complex and the T wave, representing the time when the ventricles are depolarized before repolarization begins.


An electrocardiogram (ECG or EKG) is a recording of the heart's electrical activity as a graph over a period of time, as detected by electrodes attached to the outer surface of the skin and recorded by a device external to the body. The graph can show the heart's rate and rhythm. It can also detect enlargement of the heart, decreased blood flow, or the presence of current or past heart attacks. ECGs are the primary clinical tool to measure electrical and mechanical performance of the heart.

The ECG works by detecting and amplifying tiny electrical changes on the skin that occur during heart muscle depolarization. The output for the ECG forms a graph that shows several different waves, each corresponding to a different electrical and mechanical event within the heart. Changes in these waves are used to identify problems with the different phases of heart activity.



ECG: Illustration of a patient undergoing a 12-lead ECG.

The P Wave



Normal Systole ECG: The U wave is not visible in all ECGs.

The first wave on an ECG is the P wave, indicating atrial depolarization in which the atria contract (atrial systole). The P wave is the first wave on the ECG because the action potential for the heart is generated in the sinoatrial (SA) node, located on the atria, which sends action potentials directly through Bachmann's bundle to depolarize the atrial muscle cells.


Increased or decreased P waves can indicate problems with the potassium ion concentration in the body that will alter nerve activity. A missing P wave indicates atrial fibrillation, a cardiac arrhythmia in which the heart beats irregularly, preventing efficient ventricular diastole. This is generally not fatal on its own.

The QRS Complex

The QRS complex refers to the combination of the Q, R, and S waves, and indicates ventricular depolarization and contraction

(ventricular systole). The Q and S waves are downward waves while the R wave, an upward wave, is the most prominent feature of an ECG. The QRS complex represents action potentials moving from the AV node, through the bundle of His and left and right branches and Purkinje fibers into the ventricular muscle tissue. Abnormalities in the QRS complex may indicate cardiac hypertrophy or myocardial infarctions.

The T Wave and ST Segment



Animation of a Normal ECG Wave: The red lines represent the movement of the electrical signal through the heart.

The T Wave indicates ventricular repolarization, in which the ventricles relax following depolarization and contraction. The ST segment refers to the gap (flat or slightly upcurved line) between the S wave and the T wave, and represents the time between ventricular depolarization and repolarization. An elevated ST segment is the classic indicator for myocardial infarctions, though missing or downward sloping sloping ST segments may indicate myocardial ischemia.

Following the T wave is the U wave, which represents repolarization of the Purkinje fibers. It is not always visible on an ECG because it is a very small wave in comparison to the others.

Ventricular Fibrillation

When ECG output shows no identifiable P waves, QRS complexes, or T waves, it imdicates ventricular fibrillation, a severe arrhythmia. During ventricular fibrillation, the heart beats extremely fast and irregularly and can no longer pump blood, acting as a mass of quivering, disorganized muscle movements. Ventricular fibrillation will cause sudden cardiac death within minutes unless electrical resuscitation (with an AED) is performed immediately. It generally occurs with myocardial infarcations and heart failure, and is thought to be caused by action potentials that re-enter the AV nodes from the muscle tissue and induce rapid, irregular, weak contractions of the heart that fail to pump blood.

This page titled 17.4B: Electrocardiogram and Correlation of ECG Waves with Systole is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4C: Heart Sounds

The two major heart sounds are “lub” (from the closure of AV valves) and “dub: (from the closure of aortic and pulmonary valves).

Learning Objectives

- Describe the sounds the heart makes

Key Points

- The heart tone “lub,” or S1, is caused by the closure of the mitral and tricuspid atrioventricular (AV) valves at the beginning of ventricular systole.
- The heart tone “dub,” or S2 (a combination of A2 and P2), is caused by the closure of the aortic valve and pulmonary valve at the end of ventricular systole.
- The splitting of the second heart tone, S2, into two distinct components, A2 and P2, can sometimes be heard in younger people during inspiration. During expiration, the interval between the two components shortens and the tones become merged.
- Murmurs are a “whoosh” or “slosh” sound that indicate backflow through the valves.
- S3 and S4 are a “ta” sound that indicates ventricles that are either too weak or too stiff to effectively pump blood.

Key Terms

- **dub**: The second heart tone, or S2 (A2 and P2), caused by the closure of the aortic valve and pulmonary valve at the end of ventricular systole.
- **lub**: The first heart tone, or S1, caused by the closure of the atrioventricular valves (mitral and tricuspid) at the beginning of ventricular contraction or systole.
- **Heart murmurs**: A sound made by backflow of blood through either set of valve that cannot close or open properly.

The closing of the heart valves produces a sound. This sound may be described as either a “lub” or a “dub” sound. Heart sounds are a useful indicator for evaluating the health of the valves and the heart as a whole.

S1

The first heart sound, called S1, makes a “lub” sound caused by the closure of the mitral and tricuspid valves as ventricular systole begins. There is a very slight split between the closure of the mitral and tricuspid valves, but it is not long enough to create multiple sounds.

S2


The second heart sound, called S2, makes a “dub” sound caused by the closure of the semilunar (aortic and pulmonary) valves following ventricular systole. S2 is split because aortic valve closure occurs before pulmonary valve closure. During inspiration (breathing in) there is slightly increased blood return to the right side of the heart, which causes the pulmonary valve to stay open slightly longer than the aortic valve. Due to this, the naming convention is to divide the second sound into two second sounds, A2 (aortic), and P2 (pulmonary). The time between A2 and P2 is variable depending on the respiratory rate, but the split is generally only prominent in children during inspiration. In adults and during expiration, the split is usually not long enough to suggest two sounds.

Abnormal Heart Sounds

Abnormal heart sounds may indicate problems with the health of the valves. Heart murmurs sound like a “whoosh” or “slosh” and

indicate regurgitation or backflow of blood through the valves because they cannot close properly. Heart murmurs are common and generally not serious, but some may be more severe and/or caused by severe underlying problems within the heart. Murmurs may also be caused by valve stenosis (improper opening) and cardiac shunts, a severe condition in which a defect in the septum allows blood to flow between both sides of the heart.

Third and fourth heart sounds, S3 and S4, differ from S1 and S2 because they are caused by abnormal contraction and relaxation of the heart instead of the closure of valves and are more often indicative of more severe problems than are heart murmurs. S3 represents a flabby or weak ventricle that fills with more blood than it is able to pump, while S4 represents a stiff ventricle, such as those found in cardiac hypertrophy. S3 makes a “ta” sound after the “lub-dub” while S4 makes a “ta” sound before the “lub-dub.”



Opening and Closing of Heart Valves: The closing of the heart valves generates the “lub, dub” sounds that can be heard though a stethoscope.

This page titled 17.4C: Heart Sounds is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4D: Cardiac Cycle

The cardiac cycle describes the heart's phases of contraction and relaxation that drive blood flow throughout the body.

Learning Objectives

- Describe the cardiac cycle and its three phases

Key Points

- Every single beat of the heart involves three major stages: cardiac diastole, when chambers are relaxed and filling passively; atrial systole when the atria contract leading to ventricular filling; and ventricular systole when blood is ejected into both the pulmonary artery and aorta.
- Pulse is a way of measuring heartbeat, based on the arterial distensions or pulses that occur as blood is pushed through the arteries.
- Resting heart rate typically ranges from 60 to 100 bpm (beats per minute). Athletes often have significantly lower than average heart rates while the sedentary and obese typically have elevated heart rates.
- Systolic blood pressure is the pressure during heart contraction, while diastolic blood pressure is the pressure during heart relaxation.
- The normal range for blood pressure is between 90/60 mmHg and 120/80 mmHg.

Key Terms

- **cardiac cycle**: The term used to describe the relaxation and contraction that occur as a heart works to pump blood through the body.
- **cardiac output**: The volume of blood pumped by the heart each minute, calculated as heart rate (HR) X (times) stroke volume (SV).
- **pulse**: Pressure waves generated by the heart in systole move the arterial walls, creating a palpable pressure wave felt by touch.

The cardiac cycle is the term used to describe the relaxation and contraction that occur as the heart works to pump blood through the body. Heart rate is a term used to describe the frequency of the cardiac cycle. It is considered one of the four vital signs and is a regulated variable. Usually heart rate is calculated as the number of contractions (heartbeats) of the heart in one minute and expressed as "beats per minute" (bpm). When resting, the adult human heart beats at about 70 bpm (males) and 75 bpm (females), but this varies among individuals. The reference range is normally between 60 bpm (lower is termed bradycardia) and 100 bpm (higher is termed tachycardia). Resting heart rates can be significantly lower in athletes and significantly higher in the obese. The body can increase the heart rate in response to a wide variety of conditions in order to increase the cardiac output, the blood ejected by the heart, which improves oxygen supply to the tissues.

Pulse

Pressure waves generated by the heart in systole, or ventricular contraction, move the highly elastic arterial walls. Forward movement of blood occurs when the arterial wall boundaries are pliable and compliant. These properties allow the arterial wall to distend when pressure increases, resulting in a pulse that can be detected by touch. Exercise, environmental stress, or psychological stress can cause the heart rate to increase above the resting rate. The pulse is the most straightforward way of measuring the heart rate, but it can be a crude and inaccurate measurement when cardiac output is low. In these cases (as happens in some arrhythmias), there is little pressure change and no corresponding change in pulse, and the heart rate may be considerably higher than the measured pulse.


Cardiac Cycle

Every single heartbeat includes three major stages: atrial systole, ventricular systole, and complete cardiac diastole.

- Atrial systole is the contraction of the atria that causes ventricular filling.
- Ventricular systole is the contraction of the ventricles in which blood is ejected into the pulmonary artery or aorta, depending on side.
- Complete cardiac diastole occurs after systole. The blood chambers of the heart relax and fill with blood once more, continuing the cycle.

Systolic and Diastolic Blood Pressure

Throughout the cardiac cycle, the arterial blood pressure increases during the phases of active ventricular contraction and decreases during ventricular filling and atrial systole. Thus, there are two types of measurable blood pressure: systolic during contraction and diastolic during relaxation. Systolic blood pressure is always higher than diastolic blood pressure, generally presented as a ratio in which systolic blood pressure is over diastolic blood pressure. For example, 115/75 mmHg would indicated a systolic blood pressure of 115 mmHg and a diastolic blood pressure or 75 mmHG. The normal range for blood pressure is between 90/60 mmHg and 120/80 mmHg. Pressures higher than that range may indicate hypertension, while lower pressures may indicate hypotension. Blood pressure is a regulated variable that is directly related to blood volume, based on cardiac output during the cardiac cycle.



The Cardiac Cycle: Changes in contractility lead to pressure differences in the heart's chambers that drive the movement of blood.

This page titled 17.4D: Cardiac Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.4E: Cardiac Output

Cardiac output (Q or CO) is the volume of blood pumped by the heart, in particular by the left or right ventricle, in one minute.

Learning Objectives

- Describe cardiac output and its function in the cardiovascular system

Key Points

- Cardiac output, a measure of how much blood the heart pumps over the course of a minute, is calculated by multiplying heart rate by stroke volume.
- The heart rate is increased by sympathetic nervous stimulation and decreased by parasympathetic nervous stimulation.
- Stroke volume is end diastolic volume (venous return) minus end systolic volume, the amount of blood left over in the heart after systole.
- The ejection fraction is stroke volume divided by end diastolic volume.
- Mean arterial blood pressure is cardiac output multiplied by total peripheral resistance. A twofold change in vascular size will cause a 16-fold change in resistance in the opposite direction.
- Starling's mechanism states that changes in venous return (preload) to the heart will change cardiac output, which will also change mean arterial blood pressure in the same direction. This means that blood volume and blood pressure are directly related to one another.

Key Terms

- **mean arterial blood pressure**: A measure of blood pressure based on cardiac output and vascular resistance.
- **cardiac output**: The volume of blood pumped by the heart, in particular by the left or right ventricle, in the time interval of one minute.

Cardiac output (CO) is a measure of the heart's performance. While there are many clinical techniques to measure CO, it is best described as a physiological and mathematical relationship between different variables. When one of the variables change, CO as a whole will change as a result. This can also be used to predict other regulated variables, such as blood pressure and blood volume. The mathematical description of CO is that CO=Heart Rate (HR)×Stroke Volume (SV)CO=Heart Rate (HR)×Stroke Volume (SV). Changes in HR, SV, or their components, will change CO.

Heart Rate

The heart rate is determined by spontaneous action potential generation in the sinoatrial (SA) node and conduction in the atrioventricular (AV) node. It refers to the number of heartbeats over the course of a minute. Sympathetic nervous system activation will stimulate the SA and AV nodes to increase the heart rate, which will increase cardiac output. Parasympathetic nervous system activation will conversely act on the SA and AV nodes to decrease the heart rate, which will decrease cardiac output. For the SA node, the rate of depolarization is altered, while the AV node's rate of conduction is altered by autonomic nerve stimulation.

Stroke Volume


Stroke volume refers to the amount of blood ejected from the heart during a single beat. It is a measure of the contractility of the heart based on end diastolic volume (EDV), mathematically described as SV=EDV−ESV (end systolic volumeSV=EDV−ESV (end systolic volume. EDV is the volume of blood in the ventricles at the end of diastole, while ESV is the volume of blood left inside the ventricles at the end of systole, making SV the difference between EDV and ESV. Contractility of the heart refers to the variability in how much blood the heart ejects based on changes in stroke volume rather than than changes in heart rate.

Additionally, another indicator known as the ejection fraction (EF) is used to evaluate stroke volume and contractility. It is described as EF=(SVEDV)×100%EF=(SVEDV)×100% and is a measure of the proportion of blood ejected during systole compared to the amount of blood that was present in the heart. A higher EF suggests more efficient heart activity.

Mean Arterial Pressure

Cardiac output is an indicator of mean arterial blood pressure (MAP), the average measure of blood pressure within the body. It is described as MAP=CO×TPR (total peripheral resistance)MAP=CO×TPR (total peripheral resistance). TPR is a measure of resistance in the blood vessels, which acts as the force by which blood must overcome to flow through the arteries determined by the diameter of the blood vessels. The exact relationship is such that a twofold increase in blood vessel diameter (doubling the diameter) would decrease resistance by 16-fold, and the opposite is true as well. When CO increases, MAP will increase, but if CO decreases, MAP will decrease.

Starling's Law of the Heart



Frank Starling's Law: This chart indicates stroke volume compared to ventricular preload, with labels for preload dependent zone, responsive patient SVV > 10%, and nonresponsive patient SVV < 10 %.

CO can also predict blood pressure based on blood volume. Starling's law of the heart states that the SV of the heart increases in response to an increase in EDV when all other factors remain constant. Essentially, this means that higher venous blood return to the heart (also called the preload) will increase SV, which will in turn increase CO. This is because sarcomeres are stretched further when EDV increases, allowing the heart to eject more blood and keep the same ESV if no other factors change.

The main implication of this law is that increases in blood volume or blood return to the heart will increase cardiac output, which will lead to an increase in MAP. The opposite scenario is true as well. For example, a dehydrated person will have a low blood volume and lower venous return to the heart, which will decrease cardiac output and blood pressure. Those that stand up quickly after lying down may feel light-headed because their venous return to the heart is momentarily impaired by gravity, temporarily decreasing blood pressure and supply to the brain. The adjustment for blood pressure is a quick process, while blood volume is slowly altered. Blood volume itself is another regulated variable, regulated slowly through complex processes in the renal system that alter blood pressure based on the Starling mechanism.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...stem%23Systole**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...tem%23Diastole**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventricle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventricle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- noradrenaline. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/noradrenaline**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gap junction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gap_junction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ConductionsystemoftheheartwithouttheHeart-en.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sinoat...heHeart-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...stem%23The_ECG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Electrocardiography. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Electrocardiography**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sinoatrial node. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/sinoatrial%20node**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- atrioventricular node. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/atrioventricular%20node**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ConductionsystemoftheheartwithouttheHeart-en.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sinoat...heHeart-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ECG principle slow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:EC...ciple_slow.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ECGcolor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ECGcolor.svg**. **License**: ***Public Domain: No Known Copyright***
- Systole QRS Complex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sy...RS_Complex.png**. **License**: ***Public Domain: No Known Copyright***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...3.22Lub-Dub.22**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dub. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/dub**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lub. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lub**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ConductionsystemoftheheartwithouttheHeart-en.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sinoat...heHeart-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ECG principle slow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:EC...ciple_slow.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ECGcolor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ECGcolor.svg**. **License**: ***Public Domain: No Known Copyright***
- Systole QRS Complex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Systole_QRS_Complex.png**. **License**: ***Public Domain: No Known Copyright***
- Apikal4D. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Apikal4D.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_cardiovascular_system%23Cardiac_Cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Electrical conduction system of the heart. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Electrical_conduction_system_of_the_heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pulse. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pulse**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiac cycle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cardiac%20cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/pulse**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiac output. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cardiac%20output**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ConductionsystemoftheheartwithouttheHeart-en.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sinoatrial_node#/media/File:ConductionsystemoftheheartwithouttheHeart-en.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- ECG principle slow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ECG_principle_slow.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- ECGcolor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ECGcolor.svg. **License**: *Public Domain: No Known Copyright*
- Systole QRS Complex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Systole_QRS_Complex.png. **License**: *Public Domain: No Known Copyright*
- Apikal4D. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Apikal4D.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/0d/Cardiac_cycle_pressure_only.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cardiac output. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_output. **License**: *CC BY-SA: Attribution-ShareAlike*
- echocardiography. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/echocardiography. **License**: *CC BY-SA: Attribution-ShareAlike*
- pulse pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pulse%20pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardiac output. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cardiac%20output. **License**: *CC BY-SA: Attribution-ShareAlike*
- Starling's Law. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Frank%E2%80%93Starling_law. **License**: *CC BY-SA: Attribution-ShareAlike*
- ConductionsystemoftheheartwithouttheHeart-en.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sinoatrial_node#/media/File:ConductionsystemoftheheartwithouttheHeart-en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- ECG principle slow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ECG_principle_slow.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- ECGcolor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ECGcolor.svg. **License**: *Public Domain: No Known Copyright*
- Systole QRS Complex. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Systole_QRS_Complex.png. **License**: *Public Domain: No Known Copyright*
- Apikal4D. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Apikal4D.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/0d/Cardiac_cycle_pressure_only.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Frank_Starlingu2019s_curve.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cardiac_output#/media/File:Frank_Starling%E2%80%99s_curve.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 17.4E: Cardiac Output is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.5: EXERCISE AND THE HEART

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

17.5A: Effects of Exercise on the Heart

This page titled 17.5: Exercise and the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

17.5A: Effects of Exercise on the Heart

Aerobic exercise promotes cardiovascular health, while physical inactivity is associated with increased morbidity and mortality.

Learning Objectives

- Describe the effects of exercise on the heart

Key Points

- Exercise causes the heart to pump blood into the circulation more efficiently as a result of more forceful and efficient myocardial contractions, increased perfusion of tissues and organs with blood, and increased oxygen delivery.
- Aerobic exercise trains the heart to become more efficient.
- Aerobic conditioning exercises, such as running and swimming, train the heart and lungs to pump blood more efficiently, allowing more oxygen to get to muscles and organs.
- Exercise is protective against metabolic syndrome, lowers blood pressure, works against blood clotting, and lowers stress, all of which contribute to improved cardiovascular health.
- Too much exercise can be harmful, and may cause cardiac hypertrophy or sudden cardiac death in those with certain genetic characteristics.

Key Terms

- **aerobic exercise**: A process whereby one trains the heart and lungs to pump blood more efficiently, increasing delivery of oxygen to muscles and organs.
- **metabolic syndrome,**: A complex disease, characterized by lack of exercise and high levels of adipose tissue, that turn causes many other cardiovascular and endocrine diseases.

Exercise has been shown to protect against nearly every type of chronic acquired cardiovascular disease. A sedentary and inactive lifestyle is associated with greater risk for hypertension, cardiac hypertrophy, atherosclerosis, and myocardial infarctions, due to the metabolic changes that accompany a sedentary lifestyle. The physiological benefits of exercise on the heart include increased blood filling and pumping ability and improved oxygenation.

Athletic Training and Cardiac Function

Aerobic conditioning is a process by which one trains the heart and lungs to pump blood more efficiently, allowing more oxygen to get to muscles, organs, and the heart itself. Aerobic conditioning and cardiovascular fitness are determining factors in athletic performance in events with duration greater than two minutes. In track competitions this includes all events extending from the 800 meter. Training for these events is done predominantly through cardiovascular exercise like running, swimming, and aerobics. A stronger heart does not pump blood faster but does pump it more forcefully with a greater ejection fraction, suggesting more efficient cardiac output. The heart rates at rest of trained endurance athletes are also significantly lower than those of sedentary individuals, because fewer heartbeats are required to produce the same cardiac output at rest in those with higher stroke volumes. The improved pumping capacity of the heart is due to the increased muscular tone of the myocardium, more efficient delivery of blood and oxygen to the heart itself, and improved filling of and ejection from the heart chambers, causing increased stroke volume.


Protection Against Chronic Disease

Exercise, along with a healthy diet, is effective at preventing chronic acquired cardiovascular diseases for several reasons.

1. Exercise burns fat and works against adipose tissue creation. Higher levels of fat can cause metabolic syndrome, in which chronic inflammation in the blood vessels increases the risk of atherosclerosis and hypertension.

2. Exercise helps prevent blood pooling and clotting, protecting against clots that could cause angina or myocardial infarction.
3. Exercise lowers blood pressure by improving the efficiency of the heart's ability to pump blood.
4. Exercise releases endorphins that reduce stress. Stress causes sympathetic nervous system activation that increases heart rate and blood sugar, as well as the risk for diabetes and metabolic syndrome.

About thirty minutes per day of cardiovascular exercise is recommended for most non-athletes who wish to maintain a healthy lifestyle. Too much exercise can be detrimental to the heart, but generally only in extreme cases in athletes with certain genetic predispositions. For example, some athletes may be at risk for cardiac hypertrophy from too much exercise over long periods of time and sudden cardiac death from exercising to the point that the heart's metabolic demands become too high, causing an arrhythmia.



The mammalian heart: Chronic exercise results in increased pumping efficiency, a greater filling capacity and greater responsiveness of the heart to increased energy demands.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Aerobic conditioning. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Aerobic_conditioning**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Exercise. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Exercise%23Cardiovascular_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anaerobic threshold. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/anaerobic%20threshold**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- aerobic exercise. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/aerobic%20exercise**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Diagram of the human heart (cropped). **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 17.5A: Effects of Exercise on the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

Section 6:

This page titled Section 6: is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18: CARDIOVASCULAR SYSTEM: BLOOD VESSELS

18.10: Circulatory Routes

18.10A: The Aorta and Its Branches

18.10B: Ascending Aorta

18.10C: Arch of the Aorta

18.10D: Thoracic Aorta

18.10E: Abdominal Aorta

18.10F: Arteries of the Pelvis and Lower Limbs

18.1: Blood Vessel Structure and Function

18.1A: Blood Vessel Structure

18.1B: Blood Vessel Function

18.2: Arteries

18.2A: Artery Function

18.2B: Elastic Arteries

18.2C: Muscular Arteries

18.2D: Anastomoses

18.2E: Arterioles

18.2F: Capillaries

18.3: The Venous System

18.3A: Venules

18.3B: Veins

18.4: Physiology of Circulation

18.4A: Introduction to Blood Flow, Pressure, and Resistance

18.4B: Distribution of Blood

18.5: Systemic Blood Pressure

18.5A: Introduction to Blood Pressure

18.5B: Arterial Blood Pressure

18.5C: Venous Blood Pressure

18.6: Control of Blood Pressure

18.6A: Role of the Cardiovascular Center

18.6B: Short-Term Neural Control

18.6C: Checking Circulation

18.6C: Short-Term Chemical Control

18.6D: Long-Term Renal Regulation

18.6E: Pulse

18.6F: Measuring Blood Pressure

18.6G: Extremes in Blood Pressure

18.7: Blood Flow Through the Body

18.7A: Velocity of Blood Flow

18.7B: Local Regulation of Blood Flow

18.7C: Blood Flow in Skeletal Muscle

18.7D: Blood Flow in the Brain

18.7E: Blood Flow in the Skin

18.7F: Blood Flow in the Lungs

18.7G: Blood Flow in the Heart

18.7H: Hepatic Portal Circulation

18.8: Capillary Exchange

18.8A: Capillary Dynamics

18.8B: Transcytosis

18.8C: Bulk Flow: Filtration and Reabsorption

18.9: Circulatory Shock

18.9A: Types of Shock

18.9B: Homeostatic Responses to Shock

18.9C: Signs and Symptoms of Shock

This page titled 18: Cardiovascular System: Blood Vessels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10: CIRCULATORY ROUTES

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.10A: The Aorta and Its Branches

18.10B: Ascending Aorta

18.10C: Arch of the Aorta

18.10D: Thoracic Aorta

18.10E: Abdominal Aorta

18.10F: Arteries of the Pelvis and Lower Limbs

This page titled 18.10: Circulatory Routes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10A: The Aorta and Its Branches

The aorta is the largest artery in the body and is divided into 3 parts: the ascending aorta, arch of the aorta, and descending aorta.

Learning Objectives

- Outline the divisions of the aorta

Key Points

- The blood is pumped from the left ventricle into the aorta and from there branches to all parts of the body.
- The aorta is divided into three parts: the ascending aorta (where the aorta initially leaves the heart and points toward the head), the arch of the aorta (where the aorta changes direction), and the descending aorta (where the aorta points toward the feet).
- The ascending aorta has two small branches, the left and right coronary arteries, that provide blood to the heart muscle.
- The arch of the aorta has three branches: the brachiocephalic artery (which divides into right common carotid artery and the right subclavian artery), the left common carotid artery, and the left subclavian artery. These arteries provide blood to both arms and the head.
- Within the abdomen, the descending aorta branches into the two common iliac arteries; these provide blood to the pelvis and, eventually, the legs.

Key Terms

- **descending aorta**: The region of the aorta that passes inferiorly towards the feet.
- **ascending aorta**: The region of the aorta directly attached to the heart that passes superiorly towards the head.
- **arch of the aorta**: The region of the aorta that changes direction between the ascending and descending aorta.

The aorta is the largest artery in the systemic circulatory system. Blood is pumped from the left ventricle into the aorta and from there branches to all parts of the body. The aorta is an elastic artery, meaning it is able to distend. When the left ventricle contracts to force blood into the aorta, the aorta expands. This stretching generates the potential energy that helps maintain blood pressure during diastole, since during this time the aorta contracts passively.





Diagram of Human Heart: This diagram of the human heart shows all the major vessels, and arrows indicate the direction of flow through the heart.

The aorta is divided into three parts: the ascending aorta, where the aorta initially leaves the heart and points superiorly toward the head; the arch of the aorta where the aorta changes direction; and the descending aorta where the aorta points inferiorly toward the feet.



Ascending Aorta: The aorta has three parts: the ascending, the arch and the descending.

The ascending aorta has two small branches, the left and right coronary arteries. These arteries provide blood to the heart muscle, and their blockage is the cause myocardial infarctions or heart attacks.

The arch of the aorta has three branches: the brachiocephalic artery, which itself divides into right common carotid artery and the right subclavian artery, the left common carotid artery, and the left subclavian artery. These arteries provide blood to both arms and the head.

The descending aorta is the largest artery in the body; it runs from the heart down the length of the chest and abdomen. It is divided into two portions, the thoracic and abdominal, in correspondence with the two great cavities of the trunk in which it sits. Within the abdomen, the descending aorta branches into the two common iliac arteries that provide blood to the pelvis and, eventually, the legs.

This page titled 18.10A: The Aorta and Its Branches is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10B: Ascending Aorta

The ascending aorta is the first portion of the aorta; it includes the aortic sinuses, the bulb of the aorta, and the sinotubular junction.

Learning Objectives

- Describe the role of the ascending aorta in circulation


Key Points

- The ascending aorta passes diagonally upward, forward, and to the right, in the direction of the heart ‘s axis.
- The aortic sinuses end at the sinotubular junction, the point in the ascending aorta where the aorta becomes a tubular structure.
- The ascending aorta has two branching vessels, the left and right coronary arteries, which supply blood to the heart muscle.

Key Terms

- **aortic sinuses**: An aortic sinus is one of the anatomic dilations of the ascending aorta, which occurs just above the aortic valve.
- **sinotubular junction**: The sinotubular junction is the point in the ascending aorta where the aortic sinuses end and the aorta becomes a tubular structure.
- **bulb of the aorta**: At the union of the ascending aorta with the aortic arch, the caliber of the vessel increases with a bulging of its right wall. This dilation is termed the bulb of the aorta, and on transverse section presents a somewhat oval figure.

The ascending aorta is a portion of the aorta beginning at the upper part of the base of the left ventricle, on a level with the lower border of the third costal cartilage behind the left half of the sternum; it passes diagonally upward, forward, and to the right, in the direction of the heart’s axis, as high as the upper border of the second right costal cartilage. Its total length is about five centimeters.



Ascending Aorta: The aorta has three parts: the ascending, the arch and the descending.

The aortic root is the portion of the ascending aorta beginning at the aortic annulus, the fibrous attachment between the heart

and the aorta, and extending to the sinotubular junction. Just above the aortic valve are three small dilations called the aortic sinuses. The two anterior sinuses give rise to the coronary arteries, while the third posterior sinus does not usually give rise to any vessels and so is known as the non-coronary sinus.

The sinotubular junction is the point in the ascending aorta where the aortic sinuses end and the aorta becomes a tubular structure.

At the junction of the ascending aorta with the aortic arch, the caliber of the vessel increases with a bulging of its right wall. This dilatation is termed the “bulb of the aorta.” The ascending aorta is contained within the pericardium. It is enclosed in a tube of the serous pericardium, which also encloses the pulmonary artery.

The ascending aorta is covered at its beginning by the trunk of the pulmonary artery and, higher up, is separated from the sternum by the pericardium, the right pleura, the anterior margin of the right lung, some loose areolar tissue, and the remains of the thymus. Posteriorly, it rests upon the left atrium and right pulmonary artery.

This page titled 18.10B: Ascending Aorta is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10C: Arch of the Aorta

The arch of the aorta follows the ascending aorta and begins at the level of the second sternocostal articulation of the right side.

Learning Objectives


- Describe the function of arch of the aorta

Key Points

- Three vessels come out of the aortic arch: the brachiocephalic artery, the left common carotid artery, and the left subclavian artery.
- These vessels supply blood to the head, neck, thorax and upper limbs.
- In some individuals, the left common carotid artery and left subclavian artery may arise from the brachiocephalic artery rather than the aortic arch.

Key Terms

- **arch of the aorta**: Also called the transverse aortic arch, is continuous with the upper border of the ascending aorta and begins at the level of the upper border of the second sternocostal articulation of the right side.



Aortic arch: This diagram shows the arch of the aorta and its branches.

The arch of the aorta, or the transverse aortic arch, is continuous with the upper border of the ascending aorta and begins at the level of the upper border of the second sternocostal articulation of the right side. The arch of the aorta runs at first upward, backward, and to the left in front of the trachea; it is then directed backward on the left side of the trachea and finally passes downward on the left side of the body of the fourth thoracic vertebra. At the lower border, this vertebra becomes continuous with the descending aorta.

Three vessels come out of the aortic arch: the brachiocephalic artery, the left common carotid artery, and the left subclavian artery. These vessels supply blood to the head, neck, thorax and upper limbs. In approximately 20% of individuals, the left common carotid artery arises from the brachiocephalic artery rather than the aortic arch, and in approximately 7% of individuals the left subclavian artery also arises here.

This page titled 18.10C: Arch of the Aorta is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10D: Thoracic Aorta

The thoracic aorta is the section of the aorta that travels through the thoracic cavity to carry blood to the head, neck, thorax and arms.

Learning Objectives


- Explain the function of the thoracic aorta

Key Points

- The thoracic aorta is contained in the posterior mediastinal cavity, begins at the 4th thoracic vertebra where it is continuous with the aortic arch, and ends in front of the lower border of the twelfth thoracic vertebra.
- Branches from the thoracic aorta include the bronchial arteries, the mediastinal arteries, the esophageal arteries, the pericardial arteries, and the superior phrenic artery.
- The thoracic aorta and the esophagus run parallel for most of its length, with the esophagus lying on the right side of the aorta. At the lower part of the thorax, the esophagus is placed in front of the aorta, situated on its left side close to the diaphragm..

Key Terms

- **mediastinal cavity**: The central compartment of the thoracic cavity. It contains the heart, the great vessels of the heart, the esophagus, the trachea, the phrenic nerve, the cardiac nerve, the thoracic duct, the thymus, and the lymph nodes of the central chest.
- **Thoracic Aorta**: Contained in the posterior mediastinal cavity, it begins at the lower border of the fourth thoracic vertebra where it is continuous with the aortic arch, and ends in front of the lower border of the twelfth thoracic vertebra, at the aortic hiatus in the diaphragm where it becomes the abdominal aorta.



Thoracic aorta: The aorta, highlighted in red, includes the thoracic aorta, the section of the aorta which runs from the lower border of the fourth thoracic vertebra to the diaphragm.

The thoracic aorta forms part of the descending aorta and is continuous with the aortic arch at its origin before becoming the abdominal aorta. Contained within the posterior mediastinal cavity, it begins at the lower border of the fourth thoracic vertebra

where it is continuous with the aortic arch, and ends in front of the lower border of the twelfth thoracic vertebra at the aortic hiatus in the diaphragm. At its commencement, the thoracic aorta is situated on the left of the vertebral column; it approaches the median line as it descends, and at its termination lies directly in front of the column.

The thoracic aorta's relation, from above downward, is as follows: anteriorly with the root of the left lung, the pericardium, the esophagus and the diaphragm; posteriorly with the vertebral column; on the right side with the hemiazygos veins and thoracic duct; and on the left side with the left pleura and lung. The esophagus lies on the right side of the aorta for most of its length, but at the lower part of the thorax is placed in front of the aorta and close to the diaphragm, situated on its left side.

As it descends in the thorax, the aorta gives off several paired branches. In descending order these are the bronchial arteries, the mediastinal arteries, the esophageal arteries, the pericardial arteries, and the superior phrenic artery. The posterior intercostal arteries are branches that originate throughout the length of the posterior aspect of the thoracic aorta.

This page titled 18.10D: Thoracic Aorta is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10E: Abdominal Aorta

The abdominal aorta is the largest artery in the abdominal cavity and supplies blood to most of the abdominal organs.

Learning Objectives


- Describe the abdominal aorta

Key Points

- The abdominal aorta lies slightly to the left of the midline of the body.
- The abdominal aorta has a venous counterpart called the inferior vena cava that travels parallel to it on its right side.
- The abdominal aorta branches into many arteries that supply blood to the abdominal organs.

Key Terms

- **abdominal aorta**: The largest artery in the abdominal cavity. As part of the aorta, it is a direct continuation of the descending aorta (of the thorax).
- **omentum**: Either of two folds of the peritoneum that support the viscera.
- **inferior vena cava**: The large vein which returns blood from the lower extremities and the pelvic and abdominal organs to the right atrium of the heart.



Abdominal Aorta: The aorta, highlighted in red, includes the abdominal aorta which begins at the diaphragm and ends as it branches into the common iliac arteries.

The abdominal aorta is the largest artery in the abdominal cavity. As part of the descending aorta, it is a direct continuation of the thoracic aorta.

It begins at the level of the diaphragm, crossing it via the aortic hiatus. This hole in the diaphragm that allows the passage of the great vessels at the vertebral level of T12. The abdominal aorta travels down the posterior wall of the abdomen, anterior to the vertebral column, following the curvature of the lumbar vertebrae. The abdominal aorta runs parallel to the inferior vena cava, located just to the right of the abdominal aorta.

The abdominal aorta lies slightly to the left of the midline of the body. It is covered anteriorly by the lesser omentum and stomach. Posteriorly, it is separated from the lumbar vertebrae by the anterior longitudinal ligament and left lumbar veins.

The abdominal aorta supplies blood to much of the abdominal cavity through numerous branches that become smaller in diameter as it descends. Terminally it branches into the paired common iliac arteries, which supply the pelvis and lower limbs.

This page titled 18.10E: Abdominal Aorta is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.10F: Arteries of the Pelvis and Lower Limbs

The abdominal aorta divides into the major arteries of the leg: the femoral, popliteal, tibial, dorsal foot, plantar, and fibular arteries.

Learning Objectives


- Outline blood flow in the arteries of the pelvis and lower limbs

Key Points

- The pelvic cavity is supplied by the paired internal iliac arteries.
- The internal iliac artery divides into posterior and anterior trunks which supply the reproductive and other organs of the pelvis with blood.

Key Terms

- **internal iliac arteries**: Formed when the common iliac artery divides the internal iliac artery at the vertebral level L5 descends inferiorly into the lesser pelvis.



Internal Iliac Artery: The division of the internal iliac artery into its posterior and anterior trunks.

The pelvic cavity is largely supplied by the paired internal iliac arteries, formed when the common iliac artery divides the internal iliac artery at the vertebral level L5 descends inferiorly into the lesser pelvis. The external iliac artery passes into the thigh, becoming the femoral artery.

At the most superior border of the greater sciatic foramen, the large opening to the rear of the pelvis, the internal iliac artery divides into anterior and posterior trunks.

The anterior trunk gives rise to numerous arteries that supply the organs of the pelvis and the gluteal and adductor muscles of the leg. Key branches include the obturator artery, the inferior vesical artery in men and the equivalent vaginal artery in females, and the rectal and gluteal arteries.

The posterior trunk gives rise to arteries that supply the posterior pelvic wall and the gluteal region, including the iliolumbar artery that supplies the psoas major muscle, the lateral sacral arteries, and the superior gluteal artery.

This page titled 18.10F: Arteries of the Pelvis and Lower Limbs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.1: BLOOD VESSEL STRUCTURE AND FUNCTION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

18.1A: Blood Vessel Structure

18.1B: Blood Vessel Function

This page titled 18.1: Blood Vessel Structure and Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.1A: Blood Vessel Structure

Blood vessels are flexible tubes that carry blood, associated oxygen, nutrients, water, and hormones throughout the body.

Learning Objectives

- Differentiate among the structure of arteries, veins, and capillaries

Key Points

- Blood vessels consist of arteries, arterioles, capillaries, venules, and veins. Vessel networks deliver blood to all tissues in a directed and regulated manner.
- Arteries and veins are composed of three tissue layers.
- The thick outermost layer of a vessel (tunica adventitia or tunica externa) is made of connective tissue.
- The middle layer (tunica media) is thicker and contains more contractile tissue in arteries than in veins. It consists of circularly arranged elastic fibers, connective tissue, and smooth muscle cells.
- The inner layer (tunica intima) is the thinnest layer, comprised of a single layer of endothelium supported by a subendothelial layer.
- Capillaries consist of a single layer of endothelium and associated connective tissue.

Key Terms

- **tunica intima**: The innermost layer of a blood vessel.
- **tunica externa**: The outermost layer of a blood vessel.
- **capillary**: Any of the small blood vessels that connect arteries to veins.
- **tunica media**: The middle layer of a blood vessel.
- **anastomosis**: The junction between blood vessels.

Blood vessels are key components of the systemic and pulmonary circulatory systems that distribute blood throughout the body. There are three major types of blood vessels: arteries that carry blood away from the heart, branching into smaller arterioles throughout the body and eventually forming the capillary network. The latter facilitates efficient chemical exchange between tissue and blood. Capillaries in turn merge into venules, then into larger veins responsible for returning the blood to the heart. The junctions between vessels are called anastomoses.

Arteries and veins are comprised of three distinct layers while the much smaller capillaries are composed of a single layer.

Tunica Intima


The inner layer (tunica intima) is the thinnest layer, formed from a single continuous layer of endothelial cells and supported by a subendothelial layer of connective tissue and supportive cells. In smaller arterioles or venules, this subendothelial layer consists of a single layer of cells, but can be much thicker in larger vessels such as the aorta. The tunica intima is surrounded by a thin membrane comprised of elastic fibers running parallel to the vessel. Capillaries consist only of the thin endothelial layer of cells with an associated thin layer of connective tissue.

Tunica Media

Surrounding the tunica intima is the tunica media, comprised of smooth muscle cells and elastic and connective tissues arranged circularly around the vessel. This layer is much thicker in arteries than in veins. Fiber composition also differs; veins contain fewer elastic fibers and function to control caliber of the arteries, a key step in maintaining blood pressure.

Tunica Externa

The outermost layer is the tunica externa or tunica adventitia, composed entirely of connective fibers and surrounded by an external elastic lamina which functions to anchor vessels with surrounding tissues. The tunica externa is often thicker in veins to prevent collapse of the blood vessel and provide protection from damage since veins may be superficially located.



Structure of the Artery Wall: This diagram of the artery wall indicates the smooth muscle, external elastic membrane, endothelium, internal elastic membrane, tunica externa, tunica media, and tunica intima.

Valve Function

A major structural difference between arteries and veins is the presence of valves. In arteries, the blood is pumped under pressure from the heart, so backflow cannot occur. However, passing through the capillary network results in a decrease in blood pressure, meaning that backflow of blood is possible in veins. To counteract this, veins contain numerous one-direction valves that prevent backflow.

This page titled 18.1A: Blood Vessel Structure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.1B: Blood Vessel Function

Blood vessels carry nutrients and oxygen throughout the body and aid in gas exchange.

Learning Objectives

- List the functions of blood vessels

Key Points

- Systemic and pulmonary circulatory systems efficiently deliver oxygen to the tissues of the body and remove waste products such as carbon dioxide. Arterial blood (except in the pulmonary artery) is highly saturated with oxygen and supplies oxygen to the body's tissues.
- Venous blood (except in the pulmonary vein) is deoxygenated and returns to the heart to be pumped into the lungs for reoxygenation.
- Nutrients carried in the blood are released to tissues via the permeable endothelium of blood vessels.
- Immune cells move throughout the circulatory system and are able to rapidly permeate the walls of blood vessels to attend sites of injury or infection.
- Blood vessels can increase or decrease blood flow near the surface of the body, either increasing or reducing the amount of heat lost as a means of regulating body temperature.

Key Terms


- **thermoregulation**: The maintenance of a constant internal temperature of an organism independent of the temperature of the environment

Blood plays many critical roles within the body: delivering nutrients and chemicals to tissues, removing waste products, and maintaining homeostasis and health. The circulatory system is transports blood through the body to perform these actions, facilitated by the extensive network of blood vessels.

Gas Transfer

The circulatory system can be split into two sections, systemic and pulmonary. In the systemic circulatory system, highly oxygenated blood (95-100%) is pumped from the left ventricle of the heart and into the arteries of the body. Upon reaching the capillary networks, gas exchange between tissue and blood can occur, facilitated by the narrow walls of the capillaries. Oxygen is released from the blood into the tissues and carbon dioxide, a waste product of respiration, is absorbed. The capillaries merge into venules and then veins, carrying the deoxygenated blood (~75%) back to the right atrium of the heart at the end of the systemic circulatory system.

The much smaller pulmonary system reoxygenates the blood and facilitates the removal of carbon dioxide. After leaving the heart through the right ventricle, the blood passes through the pulmonary artery, the only artery in the body that contains deoxygenated blood, and into the capillary network within the lungs. The close association of the thin-walled alveolae with the equally thin-walled capillaries allows for rapid release of carbon dioxide and uptake of oxygen. After leaving the lungs through the pulmonary vein, the only vein which carries oxygenated blood, the blood enters the left atrium. This completes the pulmonary circulatory system.



The Circulatory System: This simplified diagram of the human circulatory system (anterior view) shows arteries in red and veins in blue.

Additional Functions

Blood vessels also facilitate the rapid distribution and efficient transport of factors such as glucose, amino acids, or lipids into the tissues and the removal of waste products for processing elsewhere, such as lactic acid to the liver or urea to the kidneys. Additionally, blood vessels provide the ideal network for immune system surveillance and distribution. Numerous white blood cells circulate around the body, sensing for infection or injury. Once an injury is detected, they rapidly leave the circulatory system by passing through gaps in vessel walls to reach the affected area while signalling for a larger targeted immune response.

Mechanically the blood vessels, especially those near the skin, play a key role in thermoregulation. Blood vessels can swell to allow greater blood flow, allowing for greater radiant heat loss. Conversely, blood flow through these vessels can be lessened to reduce heat loss in colder climates.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Blood vessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_vessel**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- capillary. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/capillary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anastomosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anastomosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tunica media. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tunica+media**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blausen_0055_ArteryWallStructure.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Blausen_0055_ArteryWallStructure.png**. **License**:
- Blood vessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_vessel**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vein. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vein**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- thermoregulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thermoregulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- artery. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/artery**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blausen_0055_ArteryWallStructure.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Blausen_0055_ArteryWallStructure.png**. **License**:
- Circulatory System en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Circulatory_System_en.svg**. **License**:

This page titled 18.1B: Blood Vessel Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2: ARTERIES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

18.2A: Artery Function

18.2B: Elastic Arteries

18.2C: Muscular Arteries

18.2D: Anastomoses

18.2E: Arterioles

18.2F: Capillaries

This page titled 18.2: Arteries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2A: Artery Function

Arteries are high-pressure blood vessels that carry oxygenated blood away from the heart to all other tissues and organs.

Learning Objectives

- Distinguish the function of the arterial system from that of venous system


Key Points

- Arteries are blood vessels that carry blood away from the heart. This blood is normally oxygenated, with the exception of blood in the pulmonary artery.
- Arteries typically have a thicker tunica media than veins, containing more smooth muscle cells and elastic tissue. This allows for modulation of vessel caliber and thus control of blood pressure.
- The arterial system is the higher-pressure portion of the circulatory system, with pressure varying between the peak pressure during heart contraction (systolic pressure) and the minimum (diastolic) pressure between contractions when the heart expands and refills.
- The increase in arterial pressure during systole, or ventricular contraction, results in the pulse pressure, an indicator of cardiac function.

Key Terms

- **systolic pressure**: The peak arterial pressure during heart contraction.
- **diastolic pressure**: The minimum arterial pressure between contractions, when the heart expands and refills.
- **artery**: An efferent blood vessel from the heart, conveying blood away from the heart regardless of oxygenation status.

Arteries are blood vessels that carry blood away from the heart under pressure. This blood is usually oxygenated, with the exception of that in the pulmonary artery, which carries deoxygenated blood to the lungs.



Arterial system: Simplified diagram of the human arterial system in anterior view.

As with veins, arteries are comprised of three layers: the tunicae intima, media, and externa. In arteries, the tunica media, which contains smooth muscle cells and elastic tissue, is thicker than that of veins so it can modulate vessel caliber and thus control

and maintain blood pressure.

Arterial pressure varies between the peak pressure during heart contraction, called the systolic pressure, and the minimum or diastolic pressure between contractions, when the heart expands and refills. This pressure variation within the artery produces the observable pulse that reflects heart activity. The pressure in the arterial system decreases steadily, highest in the aorta and lowest in the venous system, as blood approaches the heart after delivery of oxygen to tissues in the systemic circulation.

Arteries of the systemic circulation can be subdivided into muscular or elastic types according to the the relative compositions of elastic and muscle tissue in their tunica media. Larger arteries are typically elastic and smaller arteries are more likely to be muscular. These arteries deliver blood to the arterioles, which in turn deliver blood to the capillary networks associated with the body's tissues.

This page titled 18.2A: Artery Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2B: Elastic Arteries

An elastic or conducting artery has a large number of collagen and elastin filaments in the tunica media.

Learning Objectives

- Distinguish the elastic artery from the muscular artery

Key Points


- Elastic arteries include the largest arteries in the body, those closest to the heart. They give rise to medium-sized vessels known as muscular, or distributing, arteries.
- Elastic arteries differ from muscular arteries both in size and in the relative amount of elastic tissue contained within the tunica media.
- Arterial elasticity gives rise to the Windkessel effect, which helps to maintain a relatively constant pressure in the arteries despite the pulsating nature of blood flow.

Key Terms


- **elastic arteries**: An artery with a large number of collagen and elastin filaments, giving it the ability to stretch in response to each pulse.
- **tunica media**: The middle layer of a vein wall with bands of thin smooth muscle.

Elastic arteries contain larger numbers of collagen and elastin filaments in their tunica media than muscular arteries do, giving them the ability to stretch in response to each pulse.

Elastic arteries include the largest arteries in the body, those closest to the heart, and give rise to the smaller muscular arteries. The pulmonary arteries, the aorta, and its branches together comprise the body's system of elastic arteries. In these large arteries, the amount of elastic tissue is considerable and the smooth muscle fiber cells are arranged in 5 to 7 layers in both circular and longitudinal directions.



Anatomy of the Arterial Wall: Arterial wall layers including the tunica intima and the tunica media. In elastic arteries, the tunica media is rich with elastic and connective tissue.



The aorta: The aorta makes up most of the elastic arteries in the body.

Arterial elasticity gives rise to the Windkessel effect, which through passive contraction after expansion helps to maintain a relatively constant pressure in the arteries despite the pulsating nature of the blood flow from the heart.

The Aorta

Due to position as the first part of the systemic circulatory system closest to the heart and the resultant high pressures it will experience, the aorta is perhaps the most elastic artery, featuring an incredibly thick tunica media rich in elastic filaments. The aorta is so thick that it requires its own capillary network to supply it with sufficient oxygen and nutrients to function, the vasa vasorum.

When the left ventricle contracts to force blood into the aorta, the aorta expands. This stretching generates the potential energy that will help maintain blood pressure during diastole, when the aorta contracts passively. Additionally, the elastic recoil helps conserve the energy from the pumping heart and smooth the flow of blood around the body through the Windkessel effect.

This page titled 18.2B: Elastic Arteries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2C: Muscular Arteries

Distributing arteries are medium-sized arteries that draw blood from an elastic artery and branch into resistance vessels.

Learning Objectives


- Distinguish muscular arteries from elastic arteries

Key Points

- In contrast to the mechanism elastic arteries use to store energy generated by the heart ‘s contraction, distributing arteries contain layers of smooth muscle.

Key Terms

- **muscular arteries**: Medium-sized arteries that draw blood from an elastic artery and branch into resistance vessels, including small arteries and arterioles.
- **elastic lamina**: A layer of elastic tissue that forms the outermost part of the tunica intima of blood vessels. It is readily visualized with light microscropy in sections of muscular arteries.
- **arteriole**: One of the small branches of an artery, especially one that connects with capillaries.



Splenic Artery: Transverse section of the human spleen showing the distribution of the splenic artery and its branches

Muscular or distributing arteries are medium-sized arteries that draw blood from an elastic artery and branch into resistance vessels, including small arteries and arterioles. In contrast to the mechanism elastic arteries use to store and dissipate energy generated by the heart’s contraction, muscular arteries contain layers of smooth muscle providing allowing for involuntary control of vessel caliber and thus control of blood flow. Muscular arteries can be identified by the well-defined elastic lamina that lies between the tunicae intima and media.

The splenic artery (lienal artery), the blood vessel that supplies oxygenated blood to the spleen, is an example of a muscular artery. It branches from the celiac artery and follows a course superior to the pancreas. The splenic artery branches off to the stomach and pancreas before reaching the spleen and gives rise to arterioles that directly supply capillaries of these organs.

This page titled 18.2C: Muscular Arteries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2D: Anastomoses

A circulatory anastomosis is a connection or looped interaction between two blood vessels.

Learning Objectives

- Explain the function of arterial anastomoses


Key Points

- Anastomoses occur normally in the body in the circulatory system, serving as backup routes for blood flow if one link is blocked or otherwise compromised.
- Anastomoses between arteries and between veins result in a multitude of arteries and veins, respectively, serving the same volume of tissue.
- Pathological anastomoses result from trauma or disease and are referred to as fistulae.

Key Terms

- **circulatory anastomosis**: A connection between two blood vessels, such as between arteries (arterio-arterial anastomosis), between veins (veno-venous anastomosis), or between an artery and a vein (arterio-venous anastomosis).
- **fistula**: An abnormal connection or passageway between organs or vessels that normally do not connect.

An anastomosis refers to any join between two vessels. Circulatory anastomoses are named based on the vessels they join: two arteries (arterio-arterial anastomosis), two veins (veno-venous anastomosis), or between an artery and a vein (arterio-venous anastomosis).




Anastomoses: The blood vessels of the rectum and anus, showing the distribution and anastomosis on the posterior surface near the termination of the gut.

Anastomoses between arteries and anastomoses between veins result in a multitude of arteries and veins serving the same volume of tissue. Such anastomoses occur normally in the body in the circulatory system, serving as backup routes for blood to flow if one link is blocked or otherwise compromised, but may also occur pathologically.

Examples of Anastomoses

Arterio-arterial anastomoses include actual joins (e.g. palmar arch, plantar arch) and potential ones, which may only function if the normal vessel is damaged or blocked (e.g. coronary arteries and cortical branch of cerebral arteries). Important examples include:

- The circle of Willis in the brain.
  The arrangement of the brain's arteries into the circle of Willis creates redundancies for the cerebral circulation. If one part of the circle becomes blocked or narrowed or one of the arteries supplying the circle is blocked or narrowed, blood flow from the other blood vessels can often preserve the cerebral perfusion well enough to maintain function.
- Joint anastomoses. Almost all joints receive anastomotic blood supply from more than one source. Examples include the knee and geniculate arteries, shoulder and circumflex humeral, and hip and circumflex iliac.
- Coronary artery anastomoses. The coronary arteries are functionally end arteries, so these meetings are referred to as anatomical anastamoses, which lack function. As blockage of one coronary artery generally results in death of the heart tissue due to lack of sufficient blood supply from the other branch, when two arteries or their branches join, the area of the myocardium receives dual blood supply. If one coronary artery is obstructed by an atheroma, a degradation of the arterial walls, the second artery is still able to supply oxygenated blood to the myocardium. However, this can only occur if the atheroma progresses slowly, giving the anastomosis time to form.



The Circle of Willis: Schematic representation of the circle of Willis—arteries of the brain and brain stem. Blood flows up to the brain through the vertebral arteries and through the internal carotid arteries.

Pathological anastomoses result from trauma or disease and are usually referred to as fistulae. They can be very severe if they result in the bypassing of key tissues by the circulatory system.

This page titled 18.2D: Anastomoses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2E: Arterioles

An arteriole is a small diameter blood vessel in the microcirculation system that branches out from an artery and leads to capillaries.

Learning Objectives

- Explain the function of arterioles


Key Points

- Arterioles have muscular walls and are the primary site of vascular resistance, which reduces the pressure and velocity of flow for gas and nutrient exchange to occur within the capillaries.
- Arterioles are innervated and can also respond to other circulating factors to regulate their caliber.

Key Terms

- **microcirculation**: The flow of blood through the smallest vessels: arterioles, capillaries, and venules.
- **arteriole**: One of the small branches of an artery, especially one that connects with capillaries.

An arteriole is a small-diameter blood vessel which forms part of the microcirculation that extends from an artery and leads to capillaries.



Capillary: Arterioles are part of the microcirculation system, along with capillaries, arteries, veins, venules, and tissue cells.

The microcirculation involves the flow of blood in the smallest blood vessels, including arterioles, capillaries, and venules.

Arterioles have muscular walls that usually consist of one or two layers of smooth muscle. They are the primary site of vascular resistance. This reduces the pressure and velocity of blood flow to enable gas and nutrient exchange to occur within the capillaries. Arterioles are innervated and also respond to various circulating hormones and other factors such as pH in order to regulate their caliber, thus modulating the amount of blood flow into the capillary network and tissues.

This page titled 18.2E: Arterioles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.2F: Capillaries

Capillaries, the smallest blood vessels in the body, are part of the microcirculation.

Learning Objectives

- Describe the structure and function of capillaries


Key Points

- Capillaries measure 5-10 μm in diameter and are only one cell thick.
- Capillaries connect arterioles and venules and enable the exchange of water, oxygen, carbon dioxide, and many other nutrients and waste substances between blood and surrounding tissues.
- There are three main types of capillaries: continuous, fenestrated, and sinusoidal.

Key Terms

- **capillary**: Any of the small blood vessels that connect arteries to veins.
- **microcirculation**: The flow of blood through the smallest vessels such as arterioles, capillaries, and venules.

Capillaries, which form part of the micro-circulation, are the smallest of the body's blood vessels at between 5-10 μm in diameter with the endothelial vessel wall of only one cell thick. They are surrounded by a thin basal lamina of connective tissue.



Structure of a capillary: Capillaries are of small diameter with the vessel wall being a single cell thick. Capillaries are surrounded by a thin basal lamina of connective tissue.

Capillary Function

Capillaries form a network through body tissues that connects arterioles and venules and facilitates the exchange of water, oxygen, carbon dioxide, and many other nutrients and waste substances between blood and surrounding tissues.
The thin wall of the capillary and close association with its resident tissue allow for gas and lipophilic molecules to pass through

without the need for special transport mechanisms. This allows bidirectional diffusion depending on osmotic gradients.

Formation of New Capillaries

During embryological development, new capillaries are formed by vasculogenesis, the process of blood vessel formation occurring by de novo production of endothelial cells and their formation into vascular tubes. The term angiogenesis denotes the formation of new capillaries from pre-existing blood vessels.

The Capillary Bed

Capillaries do not function independently. The capillary bed is an interwoven network of capillaries that supplies an organ. The more metabolically active the cells, the more capillaries required to supply nutrients and carry away waste products.

A capillary bed can consist of two types of vessels: true capillaries, which branch mainly from arterioles and provide exchange between cells and the circulation, and vascular shunts, short vessels that directly connect arterioles and venules at opposite ends of the bed, allowing for bypass.

Types of Capillaries

There are three main types of capillaries:

- *Continuous*: Endothelial cells provide an uninterrupted lining, only allowing small molecules like water and ions to diffuse through tight junctions. This leave gaps of unjoined membrane called intercellular clefts.
- *Fenestrated*: Fenestrated capillaries have pores in the endothelial cells (60-80 nanometers in diameter) that are spanned by a diaphragm of radially-oriented fibrils. They allow small molecules and limited amounts of protein to diffuse.
- *Sinusoidal*: Sinusoidal capillaries are a special type of fenestrated capillaries that have larger openings (30–40 μm in diameter) in the endothelium. These types of blood vessels allow red and white blood cells (7.5μm–25μm diameter) and various serum proteins to pass using a process aided by a discontinuous basal lamina. Sinusoid blood vessels are primarily located in the bone marrow, lymph nodes, and adrenal gland. Some sinusoids are special in that they do not have tight junctions between cells. These are called discontinuous sinusoidal capillaries, present in the liver and spleen where greater movement of cells and materials is necessary.

Control of Flow

Capillary beds may control blood flow via autoregulation. This allows an organ to maintain constant flow despite a change in central blood pressure. This is achieved by myogenic response and by tubuloglomerular feedback in the kidney. When blood pressure increases, the arterioles that lead to the capillary bed are stretched and subsequently constrict to counteract the increased tendency for high pressure to increase blood flow. In the lungs, special mechanisms have been adapted to meet the needs of increased necessity of blood flow during exercise. When heart rate increases and more blood must flow through the lungs, capillaries are recruited and are distended to make room for increased blood flow while resistance decreases.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Artery. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Artery**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tolic-pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- artery. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/artery**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tolic-pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:A..._System_en.svg. **License**: *Public Domain: No Known Copyright*
- Internal elastic lamina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Internal_elastic_lamina. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary arteries. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_arteries. **License**: *CC BY-SA: Attribution-ShareAlike*
- Elastic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Elastic_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Tunica media. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Tunica_media. **License**: *CC BY-SA: Attribution-ShareAlike*
- Tunica media. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Tunica_media. **License**: *CC BY-SA: Attribution-ShareAlike*
- Aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Aorta. **License**: *CC BY-SA: Attribution-ShareAlike*
- elastic arteries. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/elastic+arteries. **License**: *CC BY-SA: Attribution-ShareAlike*
- tunica media. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tunica+media. **License**: *CC BY-SA: Attribution-ShareAlike*
- extracellular matrix. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/extracellular_matrix. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:A..._System_en.svg. **License**: *Public Domain: No Known Copyright*
- Endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:En...elial_cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ascending aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ascending_aorta. **License**: *Public Domain: No Known Copyright*
- Radial artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Radial_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Internal elastic lamina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Internal_elastic_lamina. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscular artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscular_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Splenic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenic_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- elastic lamina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/elastic%20lamina. **License**: *CC BY-SA: Attribution-ShareAlike*
- arteriole. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/arteriole. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...cular-arteries. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Arterial_System_en.svg. **License**: *Public Domain: No Known Copyright*
- Endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:En...elial_cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ascending aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ascending_aorta. **License**: *Public Domain: No Known Copyright*
- Splenic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenic_artery. **License**: *Public Domain: No Known Copyright*
- Coronary circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coronar...%23Anastomoses. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory anastomosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circulatory_anastomosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory anastomosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circulatory_anastomosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- fistula. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fistula. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...ry-anastomosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- artery. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Arterial_System_en.svg. **License**:

Public Domain: No Known Copyright
- Endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:En...elial_cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ascending aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ascending_aorta. **License**: *Public Domain: No Known Copyright*
- Splenic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenic_artery. **License**: *Public Domain: No Known Copyright*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci..._Willis_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anastamosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anastamosis. **License**: *Public Domain: No Known Copyright*
- Arterioles. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Arterioles. **License**: *CC BY-SA: Attribution-ShareAlike*
- arteriole. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/arteriole. **License**: *CC BY-SA: Attribution-ShareAlike*
- microcirculation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/microcirculation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Arterial_System_en.svg. **License**: *Public Domain: No Known Copyright*
- Endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:En...elial_cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ascending aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ascending_aorta. **License**: *Public Domain: No Known Copyright*
- Splenic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenic_artery. **License**: *Public Domain: No Known Copyright*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ci..._Willis_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anastamosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anastamosis. **License**: *Public Domain: No Known Copyright*
- Illu capillary. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._capillary.jpg. **License**: *Public Domain: No Known Copyright*
- Capillaries. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Capillaries. **License**: *CC BY-SA: Attribution-ShareAlike*
- microcirculation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/microcirculation. **License**: *CC BY-SA: Attribution-ShareAlike*
- capillary. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/capillary. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial System en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Arterial_System_en.svg. **License**: *Public Domain: No Known Copyright*
- Endothelial cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:En...elial_cell.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ascending aorta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ascending_aorta. **License**: *Public Domain: No Known Copyright*
- Splenic artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Splenic_artery. **License**: *Public Domain: No Known Copyright*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Circle_of_Willis_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anastamosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anastamosis. **License**: *Public Domain: No Known Copyright*
- Illu capillary. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Illu_capillary.jpg. **License**: *Public Domain: No Known Copyright*
- 1280px-Capillary.svg.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Capillary.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 18.2F: Capillaries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.3: THE VENOUS SYSTEM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

18.3A: Venules

18.3B: Veins

This page titled 18.3: The Venous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.3A: Venules

Venules are small blood vessels in the microcirculation that connect capillary beds to veins.

Learning Objectives

- Describe the venules of the venous system


Key Points

- Many venules unite to form a vein.
- Venule walls have three layers: an inner endothelium composed of squamous endothelial cells that act as a membrane, a middle layer of muscle and elastic tissue, and an outer layer of fibrous connective tissue.
- High-endothelial venules are specialized post- capillary venous swellings characterized by plump endothelial cells, in contrast with the thinner endothelial cells found in regular venules.

Key Terms

- **high endothelial venule**: A specialized post-capillary venous swelling of the lymphatic system that allows lymphocytes (white blood cells) to easily exit the circulatory system.
- **venule**: A small blood vessel in the microcirculation that allows deoxygenated blood to return from capillary beds to veins.

A venule is a small blood vessel in the microcirculation that allows deoxygenated blood to return from capillary beds to larger blood vessels called veins. Venules range from 8 to 100μm in diameter and are formed when capillaries come together. Many venules unite to form a vein.



Venule: Venules form when capillaries come together and converging venules form a vein.

Venule walls have three layers: an inner endothelium composed of squamous endothelial cells that act as a membrane, a middle layer of muscle and elastic tissue, and an outer layer of fibrous connective tissue. The middle layer is poorly developed so that venules have thinner walls than arterioles. Venules are extremely porous so that fluid and blood cells can move easily from the bloodstream through their walls.

In contrast to regular venules, high-endothelial venules (HEV) are specialized post-capillary venous swellings. They are characterized by plump endothelial cells as opposed to the usual thinner endothelial cells found in regular venules. HEVs enable lymphocytes (white blood cells) circulating in the blood to directly enter a lymph node by crossing through the HEV.

This page titled 18.3A: Venules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.3B: Veins

Veins are blood vessels that carry blood from tissues and organs back to the heart; they have thin walls and one-way valves.

Learning Objectives

- Distinguish the venous system from the arterial system

Key Points

- The difference between veins and arteries is the direction of blood flow (out of the heart through arteries, returning to the heart through veins).
- Veins differ from arteries in structure and function. For example, arteries are more muscular than veins, veins are often closer to the skin, and veins contain valves to help keep blood flowing toward the heart, while arteries do not have valves and carry blood away from the heart.
- Veins are also called capacitance vessels because they contain 60% of the body's blood volume.
- The return of blood to the heart is assisted by the action of the skeletal- muscle pump. As muscles move, they squeeze the veins running through them. Veins contain a series of one-way valves, and they are squeezed, blood is pushed through the valves, which then close to prevent backflow.

Key Terms


- **venous pooling**: When blood accumulates in the lower extremities, resulting in low venous return to the heart which can result in fainting.
- **skeletal-muscle pump**: Rhythmic contraction of limb muscles that occurs during normal locomotory activity (walking, running, swimming), which promotes venous return by the pumping action on veins within muscles.
- **portal vein**: A short, wide vein that carries blood to the liver from the organs of the digestive system.

Veins are blood vessels that carry blood towards the heart. Most carry deoxygenated blood from the tissues back to the heart, but the pulmonary and umbilical veins both carry oxygenated blood to the heart. The difference between veins and arteries is the direction of blood flow (out of the heart through arteries, back to the heart through veins), not their oxygen content. Veins differ from arteries in structure and function. For example, arteries are more muscular than veins, veins are often closer to the skin, and veins contain valves to help keep blood flowing toward the heart, while arteries do not have valves and carry blood away from the heart. The precise location of veins is much more variable than that of arteries, since veins often display anatomical variation from person to person.

Veins are also called capacitance vessels because they contain 60% of the body's blood volume. In systemic circulation, oxygenated blood is pumped by the left ventricle through the arteries to the muscles and organs of the body, where its nutrients and gases are exchanged at capillaries. The blood then enters venules, then veins filled with cellular waste and carbon dioxide. The deoxygenated blood is taken by veins to the right atrium of the heart, which transfers the blood to the right ventricle, where it is then pumped through the pulmonary arteries to the lungs. In pulmonary circulation the veins return oxygenated blood from the lungs to the left atrium, which empties into the left ventricle, completing the cycle of blood circulation.

Mechanisms to Return Blood

The return of blood to the heart is assisted by the action of the skeletal-muscle pump and by the thoracic pump action of breathing during respiration. As muscles move, they squeeze the veins that run through them. Veins contain a series of one-way valves. As the vein is squeezed, it pushes blood through the valves, which then close to prevent backflow. Standing or sitting for prolonged periods can cause low venous return from venous pooling. In venous pooling, the smooth muscles surrounding the veins become slack and the veins fill with the majority of the blood in the body, keeping blood away from the brain, which can cause unconsciousness.



Venous valve: Venous valves prevent back flow and ensure that blood flows in one direction.

Although most veins take blood back to the heart, portal veins carry blood between capillary beds. For example, the hepatic portal vein takes blood from the capillary beds in the digestive tract and transports it to the capillary beds in the liver. The blood is then drained in the gastrointestinal tract and spleen, where it is taken up by the hepatic veins and blood is taken back into the heart. Since this is an important function in mammals, damage to the hepatic portal vein can be dangerous. Blood clotting in the hepatic portal vein can cause portal hypertension, which results in a decrease of blood fluid to the liver.

Vein Classification

Veins are classified in a number of ways, including superficial vs. deep, pulmonary vs. systemic, and large vs. small:

- Superficial veins: Superficial veins are close to the surface of the body and have no corresponding arteries.
- Deep veins: Deep veins are deeper in the body and have corresponding arteries.
- Communicating veins: Communicating veins (or perforator veins) directly connect superficial veins to deep veins.
- Pulmonary veins: The pulmonary veins deliver oxygenated blood from the lungs to the heart.
- Systemic veins: Systemic veins drain the tissues of the body and deliver deoxygenated blood to the heart.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Venules. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Venules**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- High endothelial venules. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/High_endothelial_venules**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymphocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymphocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- capillary. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/capillary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu capillary. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_capillary.jpg**. **License**: ***Public Domain: No Known Copyright***
- Veins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Veins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...venous-pooling**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- skeletal-muscle pump. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/skeletal-muscle%20pump**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- portal vein. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/portal_vein. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu capillary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_capillary.jpg. **License**: ***Public Domain: No Known Copyright***
- Venous valve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Venous_valve.svg. **License**: ***Public Domain: No Known Copyright***

This page titled 18.3B: Veins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.4: PHYSIOLOGY OF CIRCULATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.4A: Introduction to Blood Flow, Pressure, and Resistance
18.4B: Distribution of Blood

This page titled 18.4: Physiology of Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.4A: Introduction to Blood Flow, Pressure, and Resistance

The circulatory system is the continuous system of tubes that pumps blood to tissues and organs throughout the body.

LEARNING OBJECTIVES

Differentiate between blood flow, blood pressure, and resistance

KEY TAKEAWAYS


Key Points

- The pulmonary circulatory system circulates deoxygenated blood from the heart to the lungs via the pulmonary artery and returns it to the heart via the pulmonary vein.
- The systemic circulatory system circulates oxygenated blood from the heart around the body into the tissues before it is returned to the heart.
- The arteries divide into thin vessels called arterioles, which in turn divide into smaller capillaries that form a network between the cells of the body. The capillaries then join up again to make veins that return the blood to the heart.
- The flow of blood along arteries, arterioles and capillaries is not constant but can be controlled depending upon the body's requirements.
- Vascular resistance generated by the blood vessels must be overcome by blood pressure generated in the heart to allow blood to flow through the circulatory system.

Key Terms

- **vasodilation**: The opening of a blood vessel.
- **flow**: The movement of blood around the body, closely controlled by alterations in resistance and pressure.
- **vasoconstriction**: The closing or tightening of a blood vessel.
- **resistance**: The resistance which must be overcome by pressure to maintain blood flow throughout the body.
- **pressure**: The force which overcomes resistance to maintain blood flow throughout the body.

The circulatory system is the continuous system of tubes through which the blood is pumped around the body. It supplies the tissues with their nutritional requirements and removes waste products. The pulmonary circulatory system circulates deoxygenated blood from the heart to the lungs via the pulmonary artery and returns it to the heart via the pulmonary vein. The systemic circulatory system circulates oxygenated blood from the heart around the body into the tissues before returning deoxygenated blood to the heart.



Pulmonary circulation: Pulmonary circulation is the half of the cardiovascular system that carries oxygen-depleted blood away from the heart to the lungs and returns oxygenated blood back to the heart.

Resistance, Pressure and Flow

Three key factors influence blood circulation.

Resistance

Resistance to flow must be overcome to push blood through the circulatory system. If resistance increases, either pressure must increase to maintain flow, or flow rate must reduce to maintain pressure. Numerous factors can alter resistance, but the three most important are vessel length, vessel radius, and blood viscosity. With increasing length, increasing viscosity, and decreasing radius, resistance is increased. The arterioles and capillary networks are the main regions of the circulatory system that generate resistance, due the small caliber of their lumen. Arterioles in particular are able to rapidly alter resistance by altering their radius through vasodilation or vasoconstriction.

The resistance offered by peripheral circulation is known as systemic vascular resistance (SVR), while the resistance offered by the vasculature of the lungs is known as pulmonary vascular resistance (PVR).

Blood Pressure

Blood pressure is the pressure that blood exerts on the wall of the blood vessels. The pressure originates in the contraction of the heart, which forces blood out of the heart and into the blood vessels. If flow is impaired through increased resistance then blood pressure must increase, so blood pressure is often used as a test for circulatory health. Blood pressure can be modulated through altering cardiac activity, vasoconstriction, or vasodilation.

Blood Flow

Flow is the movement of the blood around the circulatory system. A relatively constant flow is required by the body's tissues, so pressure and resistance are altered to maintain this consistency. A too-high flow can damage blood vessels and tissue, while flow that's too low means tissues served by the blood vessel may not receive sufficient oxygen to function.

This page titled 18.4A: Introduction to Blood Flow, Pressure, and Resistance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.4B: Distribution of Blood

Humans have a closed cardiovascular system, meaning that blood never leaves the network of arteries, veins, and capillaries.

Learning Objectives

- List the components of blood flow distribution


Key Points

- In humans, blood is pumped from the strong left ventricle of the heart through arteries to peripheral tissues and returns to the right atrium of the heart through veins.
- After blood returns to the right atrium, it enters the right ventricle and is pumped through the pulmonary artery to the lungs, then returns to the left atrium through the pulmonary veins. Blood then enters the left ventricle to be circulated through the systemic circulation again.
- The closing of blood vessels is termed vasoconstriction. Vasoconstriction occurs through contraction of the muscular walls of vessels and results in increased blood pressure.
- Vasoconstriction is important for minimizing acute blood loss in the event of hemorrhage as well as retaining body heat and regulating mean arterial pressure.
- Dilation, or opening of blood vessels, is termed vasodilation. Vasodilation occurs through relaxation of smooth muscle cells within vessel walls.
- Vasodilation increases blood flow by reducing vascular resistance. Therefore, dilation of arterial blood vessels (mainly arterioles) causes a decrease in blood pressure.

Key Terms

- **vasoconstriction**: The constriction of the blood vessels.
- **vascular resistance**: The resistance to flow that must be overcome to push blood through the circulatory system. The resistance offered by the peripheral circulation is known as the systemic vascular resistance (SVR), while the resistance offered by the vasculature of the lungs is known as the pulmonary vascular resistance (PVR).
- **vasodilation**: The dilation of the blood vessels.
- **mean arterial pressure**: The average arterial pressure during a single cardiac cycle.


Humans have a closed cardiovascular system, meaning that the blood never leaves the network of arteries, veins, and capillaries. Blood is circulated through blood vessels by the pumping action of the heart, pumped from the left ventricle through arteries to peripheral tissues and returning to the right atrium through veins. It then enters the right ventricle and is pumped through the pulmonary artery to the lungs and returns to the left atrium through the pulmonary veins. Blood then enters the left ventricle to be circulated again.



Pulmonary circuit: Diagram of pulmonary circulation. Oxygen-rich blood is shown in red; oxygen-depleted blood in blue.

Distribution of blood can be modulated by many factors, including increasing or decreasing heart rate and dilation or constriction of blood vessels.

Vasoconstriction



Blood distribution: Oxygenated arterial blood (red) and deoxygenated venous blood (blue) are distributed around the body.

Vasoconstriction is the narrowing of the blood vessels resulting from contraction of the muscular wall of the vessels, particularly the large arteries and small arterioles. The process is the opposite of vasodilation, the widening of blood vessels. The process is particularly important in staunching hemorrhage and acute blood loss. When blood vessels constrict, the flow of blood is restricted or decreased, thus retaining body heat or increasing vascular resistance. This makes the skin turn paler because less blood reaches the surface, reducing the radiation of heat.

On a larger level, vasoconstriction is one mechanism by which the body regulates and maintains mean arterial pressure. Substances causing vasoconstriction are called vasoconstrictors or vasopressors. Generalized vasoconstriction usually results in an increase in systemic blood pressure, but it may also occur in specific tissues, causing a localized reduction in blood flow. The extent of vasoconstriction may be slight or severe depending on the substance or circumstance.

Vasodilation

Vasodilation refers to the widening of blood vessels resulting from relaxation of smooth muscle cells within the vessel walls, particularly in the large veins, large arteries, and smaller arterioles. The process is essentially the opposite of vasoconstriction. When blood vessels dilate, the flow of blood is increased due to a decrease in vascular resistance. Therefore, dilation of arterial blood vessels (mainly the arterioles) causes a decrease in blood pressure. The response may be intrinsic (due to local processes in the surrounding tissue) or extrinsic (due to hormones or the nervous system). Additionally, the response may be localized to a specific organ (depending on the metabolic needs of a particular tissue, as during strenuous exercise), or it may be systemic (seen throughout the entire systemic circulation). Substances that cause vasodilation are termed vasodilators.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Anatomy and Physiology of Animals/Cardiovascular System/Blood circulation. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Anatomy_and_Physiology_of_Animals/Cardiovascular_System/Blood_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tolic-pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tolic-pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypotension. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypotension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...ry_circuit.jpg**. **License**: ***Public Domain: No Known Copyright***
- Circulatory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vasodilation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vasodilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Venoconstriction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Venoconstriction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vasodilation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vasodilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mean arterial pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/mean%20...ial%20pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vascular resistance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vascular%20resistance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...ry_circuit.jpg**. **License**: ***Public Domain: No Known Copyright***
- Blutkreislauf. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Blutkreislauf.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu pulmonary circuit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pulmonary_circuit.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 18.4B: Distribution of Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.5: SYSTEMIC BLOOD PRESSURE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.5A: Introduction to Blood Pressure

18.5B: Arterial Blood Pressure

18.5C: Venous Blood Pressure

This page titled 18.5: Systemic Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.5A: Introduction to Blood Pressure

Blood pressure is a vital sign reflecting the pressure exerted on blood vessels when blood is forced out of the heart during contraction.

Learning Objectives

- Explain the topic of blood pressure

Key Points

- Diastole is the relaxation of the chambers of the heart and systole is the contraction of the heart chambers.
- Blood pressure is composed of systolic and diastolic blood pressure, which correspond to the pressure following contraction of the heart and pressure during relaxation for the heart, respectively. Normal blood pressure should be around 120/80, with the systolic number on top.
- Mean blood pressure decreases as the circulating blood moves away from the heart through arteries, capillaries, and veins due to viscous loss of energy. Mean blood pressure drops during circulation, although most of this decrease occurs along the small arteries and arterioles.


Key Terms

- **blood pressure**: The pressure exerted by the blood against the walls of the arteries and veins; it varies during the heartbeat cycle and according to a person's age, health, and physical condition.
- **systolic pressure**: The peak arterial pressure during heart contraction.
- **diastolic pressure**: The minimum arterial pressure between contractions, when the heart expands and refills.

Blood pressure is the pressure that blood exerts on the wall of the blood vessels. This pressure originates in the contraction of the heart, which forces blood out of the heart and into the blood vessels.

Two mechanisms take place in the heart: diastole and systole. Diastole is the relaxation of the chambers of the heart and systole is the contraction of the heart chambers. Systolic pressure is thus the pressure that your heart emits when blood is forced out of the heart and diastolic pressure is the pressure exerted when the heart is relaxed. This is the main mechanism by which blood pressure operates.

Blood pressure is one of the principal vital signs. During each heartbeat, blood pressure varies between a maximum (systolic) and a minimum (diastolic) pressure. A normal blood pressure should be around 120/80, with the systolic pressure expressed first.



Measurement of vital signs using a sphygmomanometer: Blood pressure and pulse, or the vital signs, are measured as indicators of several aspects of cardiovascular health.

Differences in mean blood pressure are responsible for blood flow from one location to another in circulation. The rate of mean blood flow depends on the resistance to flow presented by the blood vessels. Mean blood pressure decreases as circulating blood moves away from the heart through arteries, capillaries, and veins due to viscous loss of energy. Mean blood pressure decreases during circulation, although most of this decrease occurs along the small arteries and arterioles. Gravity affects blood pressure via hydrostatic forces (for example, during standing) Valves in veins, breathing, and pumping from contraction of skeletal muscles also influence venous blood pressure.

This page titled 18.5A: Introduction to Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.5B: Arterial Blood Pressure

The measurement of blood pressure without further specification usually refers to systemic arterial pressure measured at the upper arm.

Learning Objectives

- Distinguish between arterial blood pressure and venous blood pressure

Key Points

- Systemic blood pressure refers to the pressure exerted on blood vessels in systemic circulation, and is often measured using arterial pressure, or pressure exerted upon arteries during heart contractions.
- Blood pressure (BP), sometimes referred to as arterial blood pressure, is the pressure exerted by circulating blood upon the walls of blood vessels, and is one of the principal vital signs.
- All levels of arterial pressure put mechanical stress on the arterial walls. Higher pressures increase heart workload and progression of unhealthy tissue growth (atheroma) that develops within the walls of arteries.

Key Terms

- **atheroma**: An abnormal fatty deposit that develops within the walls of arteries.
- **arterial blood pressure**: The pressure of the blood within an arterial vessel, typically the brachial artery in the upper arm. Calculated over a cardiac cycle and determined by the cardiac output (CO), systemic vascular resistance (SVR), and central venous pressure (CVP). It can be approximately determined from measurements of the systolic pressure and the diastolic pressure while there is a normal resting heart rate.
- **systemic circulation**: The part of blood circulation that carries oxygenated blood away from the heart to the body, and returns deoxygenated blood back to the heart.

The measurement of blood pressure without further specification usually refers to the systemic arterial pressure, defined as the pressure exerted by circulating blood upon the walls of blood vessels. Pressure is typically measured with a blood pressure cuff (sphygmomanometer) wrapped around a person's upper arm, which measures the pressure in the brachial artery. A person's blood pressure is usually expressed in terms of the systolic pressure over diastolic pressure and is measured in millimeters of mercury (mmHg), for example 140/90.


Blood pressure in the arteries is much higher than in the veins, in part due to receiving blood from the heart after contraction, but also due to their contractile capacity. The tunica media of arteries is thickened compared to veins, with smoother muscle fibers and elastic tissue. Together, these generate of elastic recoil and blood vessel contraction, allowing for the maintenance of a higher pressure.



Typical Tools of Auscultatory Measurement: Shown here are a stethoscope and a sphygmomanometer, used for ascultatory measurement.

Blood Pressure and Cardiovascular Disease

While average values for arterial pressure could be computed for any given population, there is extensive variation from person to person and even from minute to minute for an individual. Additionally, the average arterial pressure of a given population has only a questionable correlation with its general health. However, in a study of 100 human subjects with no known history of hypertension, the average blood pressure of 112/64 mmHg, currently classified as a desirable or “normal” value. Normal values fluctuate through the 24-hour cycle, with the highest readings in the afternoons and lowest readings at night



Changes in Arterial Pressure: Arterial pressures changes across the cardiac cycle.

The risk of cardiovascular disease increases progressively above 115/75 mmHg. In the past, hypertension was only diagnosed if secondary signs of high arterial pressure were present along with a prolonged high systolic pressure reading over several visits. Hypotension is typically diagnosed only if noticeable symptoms are present. Clinical trials demonstrate that people who maintain arterial pressures at the low end of these ranges have much better long-term cardiovascular health. The principal medical debate concerns the aggressiveness and relative value of methods used to lower pressures into this range for those

with high blood pressure. Elevations more commonly seen in older people, though often considered normal, are associated with increased morbidity and mortality.

Arterial Hypertension

Arterial hypertension can be an indicator of other problems and may have long-term adverse effects. Sometimes it can be an acute problem, such as a hypertensive emergency. All levels of arterial pressure put mechanical stress on the arterial walls. Higher pressures increase heart workload and progression of unhealthy tissue growth (atheroma) that develops within the walls of arteries. The higher the pressure, the more stress that is present, the more the atheroma tends to progress, and the more heart muscle may thicken, enlarge, and weaken over time.

Persistent hypertension is one of the risk factors for strokes, heart attacks, heart failure, and arterial aneurysms, and is the leading cause of chronic renal failure. Even moderate elevation of arterial pressure leads to shortened life expectancy. At mean arterial pressures 50% or more above average, a person can expect to live no more than a few years unless appropriately treated.

In the past, most attention was paid to diastolic pressure, but now we know that both high systolic pressure and high pulse pressure (the numerical difference between systolic and diastolic pressures) are also risk factors for disease. In some cases, a decrease in excessive diastolic pressure can actually increase risk, probably due to the increased difference between systolic and diastolic pressures. If systolic blood pressure is elevated (>140) with a normal diastolic blood pressure (<90), it is called “isolated systolic hypertension” and may present a health concern.

This page titled 18.5B: Arterial Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.5C: Venous Blood Pressure

Venous pressure is the vascular pressure in a vein or the atria of the heart, and is much lower than arterial pressure.

Learning Objectives

- Distinguish venous blood pressure from arterial blood pressure


Key Points

- Venous pressure values are commonly 5 mmHg in the right atrium and 8 mmHg in the left atrium.
- Several measurements of venous blood pressure exist in various locations within the heart, including central venous pressure, jugular venous pressure, and portal venous pressure.
- The portal venous pressure is the blood pressure in the portal vein and is normally 5–10 mm Hg.
- Variants of venous pressure include central venous pressure, which is a good approximation of right atrial pressure, which can then be used to calculate right ventricular end diastolic volume.
- Neurogenic and hypovolemic shock can cause fainting. When the smooth muscles surrounding the veins become slack, the veins fill with the majority of the blood in the body, keeping blood away from the brain and causing unconsciousness.

Key Terms

- **central venous pressure**: The pressure of blood in the thoracic vena cava, near the right atrium of the heart, reflecting the amount of blood returning to the heart and the ability of the heart to pump the blood into the arterial system.
- **jugular venous pressure**: The indirectly-observed pressure over the venous system via visualization of the internal jugular vein.
- **venous system**: The portion of the circulatory system composed of veins, which carry blood towards the heart.

Blood pressure generally refers to the arterial pressure in the systemic circulation. However, measurement of pressures in the human venous system and the pulmonary vessels play an important role in intensive care medicine and are physiologically important in ensuring proper return of blood to the heart, maintaining flow in the closed circulatory system.



The Human Venous System: Veins (from the Latin vena) are blood vessels that carry blood towards the heart. Veins differ from arteries in structure and function; arteries are more muscular than veins, while veins are often closer to the skin and contain

valves to help keep blood flowing toward the heart.

Systemic Venous Pressure

Venous pressure is the vascular pressure in a vein or the atria of the heart. It is much lower than arterial pressure, with common values of 5 mmHg in the right atrium and 8 mmHg in the left atrium. Variants of venous pressure include:

1. Central venous pressure, a good approximation of right atrial pressure, which is a major determinant of right ventricular end diastolic volume.
2. Jugular venous pressure (JVP), the indirectly observed pressure over the venous system. It can be useful in differentiating different forms of heart and lung disease.
3. Portal venous pressure or the blood pressure in the portal vein. It is normally 5–10 mmHg.

Vein Structure and Function

In general, veins function to return deoxygenated blood to the heart, and are essentially tubes that collapse when their lumens are not filled with blood. Compared with arteries, the tunica media of veins, which contains smooth muscle or elastic fibers allowing for contraction, is much thinner, resulting in a compromised ability to deliver pressure. The actions of the skeletal-muscle pump and the thoracic pump of breathing during respiration aid in the generation of venous pressure and the return of blood to the heart.

The pressure within the circulatory circuit as a whole is mean arterial pressure (MAP). This value is a function of the cardiac output (total blood pumped) and total peripheral resistance (TPR). TPR is primarily a function of the resistance of the systemic circulation. The resistance to flow generated by veins, due to their minimal ability to contract and reduce their diameter, means that regulation of blood pressure by veins is minimal in contrast to that of muscular vessels, primarily arterioles. The latter can actively contract, reduce diameter, and increase resistance and pressure. In addition, veins can easily distend or stretch. A vein's ability to increase in diameter in response to a given blood volume also contributes to the very low pressures within this segment of the circulatory system.

Pooling and Fainting

Standing or sitting for a prolonged period of time can cause low venous return in the absence of the muscle pump, resulting in venous pooling (vascular) and shock. Fainting can occur, but usually baroreceptors within the aortic sinuses initiate a baroreflex, triggering angiotensin II and norepinephrine release and consequent vasoconstriction and heart rate increases to augment blood flow return.

Neurogenic and hypovolemic shock can also cause fainting. The smooth muscles surrounding the veins become slack and the veins fill with the majority of the blood in the body, keeping blood away from the brain and causing unconsciousness. Jet pilots wear pressurized suits to help maintain their venous return and blood pressure, since high-speed maneuvers increase venous pooling in the legs. Pressure suits specifically squeeze the lower extremities, increasing venous return to the heart. This ensures that end diastolic volumes are maintained and that the brain will receive adequate blood, preventing loss of consciousness.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Arterial blood pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arterial_blood_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- General Anatomy/Regulation of Blood Pressure. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/General...Blood_Pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tolic-pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...tolic-pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- blood pressure. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/blood_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blutdruck. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blutdruck.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- systemic circulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/systemic_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Arterial_blood_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Arterial_blood_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- atheroma. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/atheroma. **License**: *CC BY-SA: Attribution-ShareAlike*
- arterial blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/arteria...ood%20pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blutdruck. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blutdruck.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sphygmomanometer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sp...omanometer.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial-blood-pressure-curve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ar...sure-curve.svg. **License**:
- Veins. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Veins. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_p...enous_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Veins. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Veins. **License**: *CC BY-SA: Attribution-ShareAlike*
- central venous pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/central...ous%20pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- jugular venous pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/jugular...ous%20pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- venous system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/venous%20system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blutdruck. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Blutdruck.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sphygmomanometer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sp...omanometer.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial-blood-pressure-curve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ar...sure-curve.svg. **License**:
- Venous system en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ve..._system_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 18.5C: Venous Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6: CONTROL OF BLOOD PRESSURE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

18.6A: Role of the Cardiovascular Center

18.6B: Short-Term Neural Control

18.6C: Checking Circulation

18.6C: Short-Term Chemical Control

18.6D: Long-Term Renal Regulation

18.6E: Pulse

18.6F: Measuring Blood Pressure

18.6G: Extremes in Blood Pressure

This page titled 18.6: Control of Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6A: Role of the Cardiovascular Center

The cardiovascular system plays a role in body maintenance by transporting hormones and nutrients and removing waste products.

Learning Objectives

- Explain the role of the cardiovascular center in controlling blood pressure

Key Points

- The cardiovascular center is a part of the human brain found in the medulla oblongata, responsible for regulation of cardiac output.
- Numerous receptors in the circulatory system can detect changes in pH or stretch and signal these changes to the cardiovascular center.
- The cardiovascular center can alter heart rate and stroke volume to increase blood pressure and flow.


Key Terms

- **cardiovascular centre**: A region of the brain responsible for nervous control of cardiac output.

The cardiovascular center forms part of the autonomic nervous system and is responsible for regulation of cardiac output. Located in the medulla oblongata, the cardiovascular center contains three distinct components: the cardioaccelerator center, the cardioinhibitor center, and the vasomotor center.

The cardioaccelerator center stimulates cardiac function by regulating heart rate and stroke volume via sympathetic stimulation from the cardiac accelerator nerve. The cardioinhibitor center slows cardiac function by decreasing heart rate and stroke volume via parasympathetic stimulation from the vagus nerve. The vasomotor center controls vessel tone or contraction of the smooth muscle in the tunica media. Changes in diameter affect peripheral resistance, pressure, and flow, which in turn affect cardiac output. The majority of these neurons act via the release of the neurotransmitter norepinephrine from sympathetic neurons. Although each center functions independently, they are not anatomically distinct.

The cardiovascular center can respond to numerous stimuli. Hormones such as epinephrine and norepinephrine or changes in pH such as acidification due to carbon dioxide accumulation in a tissue during exercise are detected by chemoreceptors. Baroreceptors that detect stretch can also signal to the cardiovascular center to alter heart rate.



Human circulatory system: The cardiovascular system is composed largely of the circulatory system, or the system of blood vessels that distributes oxygen from the lungs throughout the body.

This page titled 18.6A: Role of the Cardiovascular Center is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6B: Short-Term Neural Control

Neural regulation of blood pressure is achieved through the role of cardiovascular centers and baroreceptor stimulation.

Learning Objectives

- Describe the role of baroreceptors and cardiovascular centers (ANS) in blood pressure control

Key Points

- The cardioaccelerator center, the cardioinhibitor center, and the vasomotor center form the cardiovascular center, a cluster of neurons that function independently to regulate blood pressure and flow.
- The release of the neurotransmitter norepinephrine from sympathetic neurons directs the majority of neurons associated with the cardiovascular center.
- Baroreceptors respond to the degree of stretch caused by the presence of blood; this stimulates impulses to be sent to the cardiovascular center to regulate blood pressure to achieve homeostasis when needed.

Key Terms

- **autonomic nervous system**: The part of the nervous system that regulates the involuntary activity of the heart, intestines, and glands. These activities include digestion, respiration, perspiration, metabolism, and blood pressure modulation.
- **norepinephrine**: A catecholamine with multiple roles including as a hormone and neurotransmitter. Areas of the body that produce or are affected by this substance are described as noradrenergic.
- **sympathetic**: Of or related to the part of the autonomic nervous system that under stress raises blood pressure and heart rate, constricts blood vessels, and dilates the pupils.
- **baroreceptor**: A nerve ending that is sensitive to changes in blood pressure.
- **parasympathetic**: Of or relating to the part of the autonomic nervous system that inhibits or opposes the effects of the sympathetic nervous system.


The autonomic nervous system plays a critical role in the regulation of vascular homeostasis. The primary regulatory sites include the cardiovascular centers in the brain that control both cardiac and vascular functions.

Neurological regulation of blood pressure and flow depends on the cardiovascular centers located in the medulla oblongata. This cluster of neurons responds to changes in blood pressure as well as blood concentrations of oxygen, carbon dioxide, and other factors such as pH.

Baroreceptor Function

Baroreceptors are specialized stretch receptors located within thin areas of blood vessels and heart chambers that respond to the degree of stretch caused by the presence of blood. They send impulses to the cardiovascular center to regulate blood pressure. Vascular baroreceptors are found primarily in sinuses (small cavities) within the aorta and carotid arteries. The aortic sinuses are found in the walls of the ascending aorta just superior to the aortic valve, whereas the carotid sinuses are located in the base of the internal carotid arteries. There are also low-pressure baroreceptors located in the walls of the venae cavae and right atrium.

When blood pressure increases, the baroreceptors are stretched more tightly and initiate action potentials at a higher rate. At lower blood pressures, the degree of stretch is lower and the rate of firing is slower. When the cardiovascular center in the medulla oblongata receives this input, it triggers a reflex that maintains homeostasis.



Baroreceptor Reflexes: Increased blood pressure results in increased rates of baroreceptor firing, whereas decreased blood pressure results in slower rates of fire, both initiating the homeostatic mechanism to restore blood pressure.

Baroreceptor Reflexes

When blood pressure rises too high, baroreceptors fire at a higher rate and trigger parasympathetic stimulation of the heart. As a result, cardiac output falls. Sympathetic stimulation of the peripheral arterioles will also decrease, resulting in vasodilation. Combined, these activities cause blood pressure to fall.

When blood pressure drops too low, the rate of baroreceptor firing decreases. This triggers an increase in sympathetic stimulation of the heart, causing cardiac output to increase. It also triggers sympathetic stimulation of the peripheral vessels, resulting in vasoconstriction. Combined, these activities cause blood pressure to rise.

The baroreceptors in the venae cavae and right atrium monitor blood pressure as the blood returns to the heart from the systemic circulation. Normally, blood flow into the aorta is the same as blood flow back into the right atrium. If blood is returning to the right atrium more rapidly than it is being ejected from the left ventricle, the atrial receptors will stimulate the cardiovascular centers to increase sympathetic firing and cardiac output until homeostasis is achieved. The opposite is also true. This mechanism is referred to as the atrial reflex.

Other neural mechanisms can also have a significant impact on cardiovascular function. These include the limbic system, which links physiological responses to psychological stimuli, chemoreceptor reflexes, generalized sympathetic stimulation, and parasympathetic stimulation.

This page titled 18.6B: Short-Term Neural Control is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6C: Checking Circulation

Checking circulation involves measurement of blood pressure and pulse through a variety of invasive and noninvasive methods.

Learning Objectives

- Choose the appropriate method for checking circulation

Key Points

- Pulse rate is most commonly measured manually at the wrist by a trained medical professional.
- Arterial catheters and pulse oximetry allow for more accurate and long-term measurement of pulse rate.
- Heart rate can be measured directly by listening to the heart through the chest.
- Electrocardiography, which detects the electrical pattern of the heart muscle through the skin, can be used for more accurate or long-term measurements.
- Arterial pressure is most commonly measured via a sphygmomanometer.
- Blood pressure values are generally reported in millimeters of mercury (mmHg), though aneroid and electronic devices do not use mercury.
- The auscultatory method for determining blood pressure uses a stethoscope and a sphygmomanometer.

Key Terms

- **electrocardiography**: A measure of the electrical output of the heart detected through the skin.
- **sphygmomanometer**: A device used to measure blood pressure.


Circulatory health can be measured in a variety of ways as follows.

Pulse

While a simple pulse rate measurement can be achieved by anyone, trained medical staff are capable of much more accurate measurements. Radial pulse is commonly measured using three fingers: the finger closest to the heart is used to occlude the pulse pressure, the middle finger is used to get a crude estimate of blood pressure, and the finger most distal to the heart is used to nullify the effect of the ulnar pulse as the two arteries are connected via the palmar arches.

Where more accurate or long-term measurements are required, pulse rate, pulse deficits, and much more physiologic data are readily visualized by the use of one or more arterial catheters connected to a transducer and oscilloscope. This invasive technique has been commonly used in intensive care since the 1970's. The rate of the pulse is observed and measured by tactile or visual means on the outside of an artery and recorded as beats per minute (BPM). The pulse may be further indirectly observed under light absorbencies of varying wavelengths with assigned and inexpensively-reproduced mathematical ratios. Applied capture of variances of light signal from the blood component hemoglobin under oxygenated vs. deoxygenated conditions allows the technology of pulse oximetry.

Heart Rate



ECG Graph: ECG graph of a normal heartbeat.

Heart rate can be measured by listening to the heart directly though the chest, traditionally using a stethoscope. For more accurate or long-term measurements, electrocardiography may be used.

During each heartbeat, a healthy heart has an orderly progression of depolarization that starts with pacemaker cells in the sinoatrial (SA) node, spreads out through the atrium, passes through the atrioventricular node down into the bundle of His and into the Purkinje fibers, and down and to the left throughout the ventricles. This organized pattern of depolarization can be detected through electrodes placed on the skin and recorded as the commonly seen ECG tracing. ECG provides a very accurate means to measure heart rate, rhythm, and other factors such as chamber sizing, as well as identifying possible regions of damage.

Blood Pressure

Arterial pressure is most commonly measured via a sphygmomanometer, which historically used the height of a column of mercury to reflect the circulating pressure. Blood pressure values are generally reported in millimeters of mercury (mmHg), though aneroid and electronic devices do not use mercury. For each heartbeat, blood pressure varies between systolic and diastolic pressures. Systolic pressure is peak pressure in the arteries, which occurs near the end of the cardiac cycle when the ventricles are contracting. Diastolic pressure is minimum pressure in the arteries, which occurs near the beginning of the cardiac cycle when the ventricles are filled with blood. An example of normal measured values for a resting, healthy adult human is 120 mmHg systolic and 80 mmHg diastolic.

Hypertension refers to abnormally high arterial pressure, as opposed to hypotension, when it is abnormally low. Along with body temperature, respiratory rate, and pulse rate, blood pressure is one of the four main vital signs routinely monitored by medical professionals and healthcare providers.

Measuring pressure invasively by penetrating the arterial wall to take the measurement is much less common and usually restricted to a hospital setting. The noninvasive auscultatory and oscillometric measurements are simpler and faster than invasive measurements, require less expertise, have virtually no complications, are less unpleasant and painful for the patient. However, noninvasive methods may yield somewhat lower accuracy and small systematic differences in numerical results. Noninvasive measurement methods are more commonly used for routine examinations and monitoring.

The Auscultatory Method



Typical Tools of Auscultatory Measurement: Shown here are a stethoscope and a sphygmomanometer, which are used for ascultatory measurement.

The auscultatory method uses a stethoscope and a sphygmomanometer. This comprises an inflatable cuff placed around the upper arm at roughly the same vertical height as the heart, attached to a mercury or aneroid manometer. The mercury manometer, considered the gold standard, measures the height of a column of mercury, giving an absolute result without need for calibration.

A cuff of appropriate size is fitted smoothly and snugly, then inflated manually by repeatedly squeezing a rubber bulb until the artery is completely occluded. Listening with the stethoscope to the brachial artery at the elbow, the examiner slowly releases the pressure in the cuff. When blood just starts to flow in the artery, the turbulent flow creates a “whooshing” or pounding (first Korotkoff sound). The pressure at which this sound is first heard is the systolic blood pressure. The cuff pressure is further released until no sound can be heard (fifth Korotkoff sound), at the diastolic arterial pressure. The auscultatory method is the predominant method of clinical measurement.

This page titled 18.6C: Checking Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6C: Short-Term Chemical Control

Blood pressure is controlled chemically through dilation or constriction of the blood vessels by vasodilators and vasocontrictors.

Learning Objectives

- Discuss chemical substances to control blood pressure

Key Points

- Constriction or dilation of blood vessels alters resistance, increasing or decreasing blood pressure respectively.
- Generalized vasoconstriction usually results in an increase in systemic blood pressure, but it may also occur in specific tissues, causing a localized reduction in blood flow.
- Vasoconstriction results from increased concentration of calcium (Ca2+) ions within vascular smooth muscle.
- When blood vessels dilate, the flow of blood is increased due to a decrease in vascular resistance. Therefore, dilation of arterial blood vessels (mainly the arterioles) causes a decrease in blood pressure.
- Localized tissues increase blood flow in multiple ways, including releasing vasodilators, primarily adenosine, into the local intersitial fluid, which diffuses to capillary beds provoking local vasodilation.

Key Terms

- **vasodilation**: The dilation (widening) of a blood vessel.
- **vasoconstriction**: The constriction (narrowing) of a blood vessel.

Many physical factors that influence arterial pressure. Each may in turn be influenced by physiological factors such as diet, exercise, disease, drugs or alcohol, stress, and obesity. In practice, each individual's autonomic nervous system responds to and regulates all of these interacting factors so that the actual arterial pressure response varies widely because of both split-second and slow-moving responses of the nervous system and end organs. These responses are very effective in changing the variables and resulting blood pressure from moment to moment.

Chemical Vasoconstriction

Vasoconstriction is the narrowing of blood vessels resulting from contraction of the muscular wall of the vessels, particularly the large arteries and small arterioles. Generalized vasoconstriction usually results in an increase in systemic blood pressure, but may also occur in specific tissues, causing a localized reduction in blood flow.

The mechanism that leads to vasoconstriction results from the increased concentration of calcium (Ca2+ ions) and phosphorylated myosin within vascular smooth muscle cells. When stimulated, a signal transduction cascade leads to increased intracellular calcium from the sarcoplasmic reticulum through IP3 mediated calcium release, as well as enhanced calcium entry across the sarcolemma through calcium channels.

The rise in intracellular calcium interacts with calmodulin, which in turn activates myosin light chain kinase. This enzyme is responsible for phosphorylating the light chain of myosin to stimulate cross-bridge cycling. Once elevated, the intracellular calcium concentration is returned to its basal level through a variety of protein pumps and calcium exchanges located on the plasma membrane and sarcoplasmic reticulum. This reduction in calcium removes the stimulus necessary for contraction allowing for a return to baseline.

Endogenous vasoconstrictors include ATP, epinephrine, and angiotensin II.



Vasoconstriction: Vasoconstriction of a microvessel by pericytes and endothelial cells that encircle an erythrocyte (E).

Chemical Vasodilation

Vasodilation is the widening of blood vessels resulting from relaxation of smooth muscle cells within the vessel walls, particularly in the large veins, large arteries, and smaller arterioles. Generalized vasodilation usually results in a decrease in systemic blood pressure, but may also occur in specific tissues causing a localized increase in blood flow.

The primary function of vasodilation is to increase blood flow in the body to tissues that need it most. This is often in response to a localized need for oxygen, but can occur when the tissue in question is not receiving enough glucose, lipids, or other nutrients. Localized tissues increase blood flow by several methods, including the release of vasodilators, primarily adenosine, into the local interstitial fluid, which diffuses to capillary beds provoking local vasodilation. Some physiologists have suggested the lack of oxygen itself causes capillary beds to vasodilate by the smooth muscle hypoxia of the vessels in the region.

As with vasoconstriction, vasodilation is modulated by calcium ion concentration and myosin phosphorylation within vascular smooth muscle cells. Dephosphorylation by myosin light-chain phosphatase and induction of calcium symporters and antiporters that pump calcium ions out of the intracellular compartment both contribute to smooth muscle cell relaxation and therefore vasodilation. This is accomplished through reuptake of ions into the sarcoplasmic reticulum via exchangers and expulsion across the plasma membrane.

Endogenous vasodilators include arginine and lactic acid.

This page titled 18.6C: Short-Term Chemical Control is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6D: Long-Term Renal Regulation

Consistent and long-term control of blood pressure is determined by the renin-angiotensin system.

Learning Objectives

- Describe the long-term renal regulation of blood pressure

Key Points

- When blood volume is low, renin, excreted by the kidneys, stimulates production of angiotensin I, which is converted into angiotensin II. This substance has many effects, including increase in blood pressure due to its vasoconstrictive properties.
- The cells that excrete renin are called juxtaglomerular cells. When blood volume is low, juxtaglomerular cells in the kidneys secrete renin directly into circulation. Plasma renin then carries out the conversion of angiotensinogen released by the liver to angiotensin I.
- Aldosterone secretion from the adrenal cortex is induced by angiotensin II and causes the tubules of the kidneys to increase the reabsorption of sodium and water into the blood, thereby increasing blood volume and blood pressure.

Key Terms


- **juxtaglomerular cells**: The juxtaglomerular cells (JG cells, or granular cells) are cells in the kidney that synthesize, store, and secrete the enzyme renin.
- **aldosterone**: A mineralocorticoid hormone secreted by the adrenal cortex that regulates the balance of sodium and potassium in the body.
- **adrenal cortex**: The outer portion of the adrenal glands that produces hormones essential to homeostasis.

Along with vessel morphology, blood viscosity is one of the key factors influencing resistance and hence blood pressure. A key modulator of blood viscosity is the renin-angiotensin system (RAS) or the renin-angiotensin-aldosterone system (RAAS), a hormone system that regulates blood pressure and water balance.

When blood volume is low, juxtaglomerular cells in the kidneys secrete renin directly into circulation. Plasma renin then carries out the conversion of angiotensinogen released by the liver to angiotensin I. Angiotensin I is subsequently converted to angiotensin II by the enzyme angiotensin converting enzyme found in the lungs. Angiotensin II is a potent vasoactive peptide that causes blood vessels to constrict, resulting in increased blood pressure. Angiotensin II also stimulates the secretion of the hormone aldosterone from the adrenal cortex.

Aldosterone causes the tubules of the kidneys to increase the reabsorption of sodium and water into the blood. This increases the volume of fluid in the body, which also increases blood pressure. If the renin-angiotensin-aldosterone system is too active, blood pressure will be too high. Many drugs interrupt different steps in this system to lower blood pressure. These drugs are one of the main ways to control high blood pressure (hypertension), heart failure, kidney failure, and harmful effects of diabetes.

It is believed that angiotensin I may have some minor activity, but angiotensin II is the major bioactive product. Angiotensin II has a variety of effects on the body: throughout the body, it is a potent vasoconstrictor of arterioles.



The renin-angiotensin pathway: The figures outlines the origination of the renin-angiotensin pathway molecules, as well as effects on target organs and systems.

This page titled 18.6D: Long-Term Renal Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6E: Pulse

Pulse is a measurement of heart rate by touching and counting beats at several body locations, typically at the wrist radial artery.

Learning Objectives

- Describe the cardiovascular pulse

Key Points

- Physiologically, pulse is the expansion of the artery due to pressure from the heartbeat, and thus is most closely correlated to systolic blood pressure.
- Sometimes the pulse cannot be taken at the wrist and may therefore be taken at the neck against the carotid artery (carotid pulse) or behind the knee (popliteal artery).
- The heart rate may be greater or less than the pulse rate depending upon physiologic demand. In this case, the heart rate is determined by auscultation or audible sounds at the heart apex, not the pulse.
- Pulse rate is recorded as beats per minute (bpm) and varies with age. A newborn or infant can have a heart rate of approximately 130-150 bpm, while an adult pulse rate is between 50 and 80 bpm.

Key Terms

- **popliteal artery**: The popliteal artery is defined as the extension of the superficial femoral artery after passing through the adductor canal and adductor hiatus above the knee.
- **radial artery**: The main artery that enters the wrist on the side of the thumb, it is the most common location for measuring pulse rate.
- **heart rate**: The number of heartbeats per unit of time, usually expressed as beats per minute.
- **pulse rate**: The physical expansion of an artery per unit of time, usually expressed as beats per minute.
- **carotid artery**: Either of a pair of arteries on each side of the neck that branch from the aorta and supply blood to the head.

The pulse is the physical expansion of an artery generated by the increase in pressure associated with systole of the heart. Pulse is often used as an equivalent of heart rate due to the relative ease of measurement; heart rate can be measured by listening to the heart directly through the chest, traditionally using a stethoscope.



Measurement of the pulse: Measurement of the pulse can occur at several locations, including the radial artery shown here.

Pulse rate or velocity is usually measured either at the wrist from the radial artery and is recorded as beats per minute (bpm). Other common measurement locations include the carotid artery in the neck and popliteal artery behind the knee

Pulse varies with age; a newborn or infant can have a heart rate of about 130-150 bpm. A toddler's heart will beat about 100-120 times per minute, an older child's heartbeat is around 60-100 bpm, adolescents around 80-100 bpm, and a healthy adults pulse rate is anywhere between 50 and 80 bpm.

The heart rate may be greater or less than the pulse rate depending upon physiologic demand. In this case, the heart rate is determined by auscultation or audible sounds at the heart apex, not the pulse. The pulse deficit (difference between heartbeats and pulsations at the periphery) is determined by simultaneous palpation at the radial artery and auscultation at the heart apex.

Measurement Techniques

While a simple measurement of pulse rate is achievable by anyone, trained medical staff are capable of much more accurate measurements. Radial pulse is commonly measured using three fingers: the finger closest to the heart used to occlude the pulse pressure, the middle finger used get a crude estimate of blood pressure, and the finger most distal to the heart used to nullify the effect of the ulnar pulse as the two arteries are connected via the palmar arches.

Where more accurate or long-term measurements are required, pulse rate, pulse deficits, and more physiologic data are readily visualized by the use of one or more arterial catheters connected to a transducer and oscilloscope. This invasive technique has been commonly used in intensive care since the 1970's. The rate of the pulse is observed and measured by tactile or visual means on the outside of an artery and is recorded as beats per minute. The pulse may be further indirectly observed under light absorbencies of varying wavelengths with assigned and inexpensively reproduced mathematical ratios. Applied capture of variances of light signal from the blood component hemoglobin under oxygenated vs. deoxygenated conditions allows the technology of pulse oximetry.

This page titled 18.6E: Pulse is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6F: Measuring Blood Pressure

Measurement of blood pressure includes systolic pressure during cardiac contraction and diastolic pressure during cardiac relaxation.

Learning Objectives

- Explain how blood pressure is measured and the ranges of blood pressure readings

Key Points

- The difference between systolic and diastolic pressure is referred to as the pulse pressure. That difference can indicate hypertension or hypotension with a deviation from the norm.
- The measurement of these pressures is now usually done with an aneroid or electronic sphygmomanometer. The classic measurement device is a mercury sphygmomanometer, using a column of mercury measured in millimeters.
- Blood pressures are also taken at other portions of the extremities. These pressures are called segmental blood pressures and are used to evaluate blockage or arterial occlusion in a limb.

Key Terms

- **pulse pressure**: Blood pressure when feeling the pulse, measured by millimeters of mercury (mmHg).
- **diastolic blood pressure**: The lowest pressure within the bloodstream, occurring between heartbeats because of a diastole.
- **systolic blood pressure**: The highest pressure within the bloodstream, occurring during each heartbeat because of the systole.

Blood pressure is the pressure blood exerts on the arterial walls. It is recorded as two readings: the systolic blood pressure (the top number) occurs during cardiac contraction, and the diastolic blood pressure or resting pressure (the bottom number), occurs between heartbeats when the heart is not actively contracting.



A spygmomanometer: A blood pressure cuff and associated monitor used for determining systolic and diastolic pressures within an artery.

A normal blood pressure is about 120 mmHg systolic over 80 mmHg diastolic. Usually the blood pressure is read from the left arm, although blood pressures are also taken at other locations along the extremities. These pressures, called segmental blood

pressures, are used to evaluate blockage or arterial occlusion in a limb (for example, the ankle brachial pressure index).The difference between the systolic and diastolic pressure is called the pulse pressure.

The measurement of these pressures is usually performed with an aneroid or electronic sphygmomanometer. The classic measurement device is a mercury sphygmomanometer, using a column of mercury measured off in millimeters. In the United States and UK, the common form is millimeters of mercury (mm Hg), while elsewhere SI units of pressure are used. There is no natural or normal value for blood pressure, but rather a range of values that are associated with increased risks for disease and health:

- Hypotension: under 90 mmHg systolic and under 60 mmHg diastolic.
- Normal: 90–119 mmHg systolic and 60–79 mmHg diastolic.
- Prehypertensive: 120–139 mmHg systolic and 80–89 mmHg diastolic.
- Hypertensive: 140 mmHg and above systolic and 90 mmHg and above diastolic.

The guidelines for acceptable readings also take into account other cofactors for disease, such as pre-existing health factors. Therefore, hypertension is indicated when the systolic number is persistently over 140–160 mmHg. Low blood pressure, or hypotension, is indicated when the systolic number is persistently below 90 mmHg.

This page titled 18.6F: Measuring Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.6G: Extremes in Blood Pressure

Chronically elevated blood pressure is called hypertension, while chronically low blood pressure is called hypotension.

Learning Objectives

- Distinguish between hypertension and hypotension

Key Points

- Hypertension, the unhealthy elevation of blood pressure, is a major risk factor for stroke, myocardial infarction (heart attacks), heart failure, aneurysms of the arteries, and peripheral arterial disease and a cause of chronic kidney disease.
- Hypertension is classified as either primary or secondary hypertension. The majority of cases are primary hypertension, high blood pressure with no identified cause. The remaining 5–10% of cases (secondary hypertension) are caused by other conditions that affect the organs or endocrine system.
- Dietary and lifestyle changes can improve blood pressure control and decrease the risk of associated health complications, although drug treatment is often necessary in people for whom lifestyle changes prove ineffective or insufficient.
- Hypotension is an abnormally low blood pressure and often indicative of a short-term condition that is not necessarily linked to disease, but rather an altered physiological state.
- For some people who exercise and are in top physical condition, low blood pressure is a sign of good health and fitness.
- For many people, low blood pressure can cause dizziness and fainting or indicate serious heart, endocrine, or neurological disorders.


Key Terms

- **hypertension**: High blood pressure, clinically diagnosed when above 140/90 mmHg.
- **hypotension**: Low blood pressure, clinically diagnosed when below 100/60 mmHg.

In healthy adults, physiological blood pressure should fall between the range of 100-140 mmHg systolic and 60-90 mmHg diastolic. Blood pressures above this are classed as hypertension and those below are hypotension, both considered medical conditions.

Hypertension

Hypertension or high blood pressure, sometimes called arterial hypertension, is a chronic medical condition in which the blood pressure in the arteries is elevated above 140/90 mmHg.



Arterial Blood Pressure over the Cardiac Cycle: Graph showing changes in blood pressure during a single contraction-relaxation cycle of the heart.

Hypertension is classified as either primary (essential) hypertension or secondary hypertension; about 90–95% of cases are categorized as "primary hypertension" which means high blood pressure with no obvious underlying medical cause. The remaining 5–10% of cases (secondary hypertension) are caused by other conditions that affect the kidneys, arteries, heart, or endocrine system.

Hypertension is a major risk factor for stroke, myocardial infarction (heart attacks), heart failure, aneurysms of the arteries (e.g. aortic aneurysm), peripheral arterial disease and a cause of chronic kidney disease. Even moderate elevation of arterial blood pressure is associated with a shortened life expectancy. Dietary and lifestyle changes can improve blood pressure control and decrease the risk of associated health complications, although drug treatment is often necessary in people for whom lifestyle changes prove ineffective or insufficient.

Hypotension

Hypotension is a medical condition in which the blood pressure in the arteries is reduced below 100/60 mmHg. Hypotension is best understood as a physiological state rather than a disease and is often associated with shock, though not necessarily indicative of it. However, blood pressure is considered too low only if noticeable symptoms are present.

For some people who exercise and are in top physical condition, hypotension is a sign of good health and fitness. For many people, low blood pressure can cause dizziness and fainting or indicate serious heart, endocrine, or neurological disorders. Severely low blood pressure can deprive the brain and other vital organs of oxygen and nutrients, leading to a life-threatening condition called shock.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- systemic circulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/systemic_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Homeostasis%23Cardiovascular_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blood pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cardiovascular system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/**

Cardiovascular_system%23Human_cardiovascular_system. License: *CC BY-SA: Attribution-ShareAlike*
- circulatory system. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/circulatory_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- blood pressure. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/blood_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory System en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._System_en.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Homeostatic Regulation of the Vascular System. January 14, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- norepinephrine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/norepinephrine. **License**: *CC BY-SA: Attribution-ShareAlike*
- parasympathetic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/parasympathetic. **License**: *CC BY-SA: Attribution-ShareAlike*
- sympathetic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sympathetic. **License**: *CC BY-SA: Attribution-ShareAlike*
- baroreceptor. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/baroreceptor. **License**: *CC BY-SA: Attribution-ShareAlike*
- autonomic nervous system. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/autonomic_nervous_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory System en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._System_en.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...l_of_Heartbeat. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vasodilator. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vasodilator. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_pressure%23Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vasoconstrictor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vasoconstrictor. **License**: *CC BY-SA: Attribution-ShareAlike*
- vasodilation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasodilation. **License**: *CC BY-SA: Attribution-ShareAlike*
- vasoconstriction. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasoconstriction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory System en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Blood pressure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood_pressure%23Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renin-angiotensin system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renin-angiotensin_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- juxtaglomerular cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/juxtaglomerular%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- aldosterone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/aldosterone. **License**: *CC BY-SA: Attribution-ShareAlike*
- adrenal cortex. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/adrenal_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory System en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Renin-angiotensin-aldosterone system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/

File:Re...one_system.png. License: CC BY-SA: Attribution-ShareAlike
- Blood pressure. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Blood_pressure%23Measurement. License: CC BY-SA: Attribution-ShareAlike
- sphygmomanometer. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/sphygmomanometer. License: CC BY-SA: Attribution-ShareAlike
- palpation. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/palpation. License: CC BY-SA: Attribution-ShareAlike
- Circulatory System en. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. License: CC BY-SA: Attribution-ShareAlike
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. Provided by: OpenStax CNX. Located at: http://cnx.org/content/m46603/latest/. License: CC BY: Attribution
- Microvessel. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Microvessel.jpg. License: CC BY: Attribution
- Renin-angiotensin-aldosterone system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Re...one_system.png. License: CC BY-SA: Attribution-ShareAlike
- SinusRhythmLabels.svg. Provided by: Wikipedia. Located at: commons.wikimedia.org/wiki/F...ythmLabels.svg. License: Public Domain: No Known Copyright
- Sphygmomanometer. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Sp...omanometer.jpg. License: CC BY-SA: Attribution-ShareAlike
- Vital signs. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Vital_s...Blood_pressure. License: CC BY-SA: Attribution-ShareAlike
- Pulse. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Pulse. License: CC BY-SA: Attribution-ShareAlike
- brachial artery. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/brachial%20artery. License: CC BY-SA: Attribution-ShareAlike
- popliteal artery. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/popliteal%20artery. License: CC BY-SA: Attribution-ShareAlike
- pulse rate. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/pulse%20rate. License: CC BY-SA: Attribution-ShareAlike
- heart rate. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/heart_rate. License: CC BY-SA: Attribution-ShareAlike
- carotid artery. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/carotid_artery. License: CC BY-SA: Attribution-ShareAlike
- radial artery. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/radial_artery. License: CC BY-SA: Attribution-ShareAlike
- Circulatory System en. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. License: CC BY-SA: Attribution-ShareAlike
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. Provided by: OpenStax CNX. Located at: http://cnx.org/content/m46603/latest/. License: CC BY: Attribution
- Microvessel. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Microvessel.jpg. License: CC BY: Attribution
- Renin-angiotensin-aldosterone system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Re...one_system.png. License: CC BY-SA: Attribution-ShareAlike
- SinusRhythmLabels.svg. Provided by: Wikipedia. Located at: commons.wikimedia.org/wiki/F...ythmLabels.svg. License: Public Domain: No Known Copyright
- Sphygmomanometer. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Sp...omanometer.jpg. License: CC BY-SA: Attribution-ShareAlike
- Pulse evaluation. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Pu...evaluation.JPG. License: CC BY-SA: Attribution-ShareAlike
- systolic blood pressure. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/systolic_blood_pressure. License: CC BY-SA: Attribution-ShareAlike
- Vital signs. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Vital_signs%23Blood_pressure. License: CC BY-SA: Attribution-ShareAlike
- pulse pressure. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/pulse%20pressure. License: CC BY-SA: Attribution-ShareAlike
- diastolic blood pressure. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/diasto...blood_pressure. License: CC BY-SA: Attribution-ShareAlike
- Circulatory System en. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. License: CC BY-SA: Attribution-ShareAlike

- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Renin-angiotensin-aldosterone system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Renin-angiotensin-aldosterone_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- SinusRhythmLabels.svg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:SinusRhythmLabels.svg. **License**: *Public Domain: No Known Copyright*
- Sphygmomanometer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sp...omanometer.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulse evaluation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pu...evaluation.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Grade 1 hypertension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gr...pertension.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- myocardial infarction (heart attacks). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/myocard...art%20attacks). **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypotension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypotension. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hypertension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypertension. **License**: *CC BY-SA: Attribution-ShareAlike*
- primary (essential) hypertension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/primary...20hypertension. **License**: *CC BY-SA: Attribution-ShareAlike*
- secondary hypertension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/secondary%20hypertension. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory System en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Circulatory_System_en.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Homeostatic Regulation of the Vascular System. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46603/latest/. **License**: *CC BY: Attribution*
- Microvessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Microvessel.jpg. **License**: *CC BY: Attribution*
- Renin-angiotensin-aldosterone system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Renin-angiotensin-aldosterone_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- SinusRhythmLabels.svg. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:SinusRhythmLabels.svg. **License**: *Public Domain: No Known Copyright*
- Sphygmomanometer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sphygmomanometer.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulse evaluation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pu...evaluation.JPG. **License**: *CC BY-SA: Attribution-ShareAlike*
- Grade 1 hypertension. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gr...pertension.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Arterial-blood-pressure-curve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ar...sure-curve.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 18.6G: Extremes in Blood Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7: BLOOD FLOW THROUGH THE BODY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.7A: Velocity of Blood Flow

18.7B: Local Regulation of Blood Flow

18.7C: Blood Flow in Skeletal Muscle

18.7D: Blood Flow in the Brain

18.7E: Blood Flow in the Skin

18.7F: Blood Flow in the Lungs

18.7G: Blood Flow in the Heart

18.7H: Hepatic Portal Circulation

This page titled 18.7: Blood Flow Through the Body is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7A: Velocity of Blood Flow

Blood flow is a pulse wave that moves out from the aorta and through the arterial branches, then is reflected back to the heart.

Learning Objectives

- Explain the velocity of blood flow

Key Points

- Flow is the movement of a liquid or gas over time.
- Flow can be calculated by multiplying velocity, the distance moved by an object over time, with cross-sectional area.
- Within the circulatory system, velocity can be altered by changes in blood pressure, vessel resistance, and blood viscosity.
- Blood vessels can vasoconstrict and vasodilate, which alters cross-sectional area.
- These various factors are under tight regulation to maintain sufficient blood flow to the body's organs and tissues.

Key Terms

- **flow**: The movement of a volume of a liquid or gas over time, e.g. mL/sec.
- **velocity**: The distance moved by an object over time, e.g. cm/sec.

The flow of blood around the circulatory system is modulated by numerous interacting factors. The science dedicated to understanding this flow is called hemodynamics.

Velocity vs. Flow

It's important to understand the different between velocity and flow. Velocity refers to the distance an object moves over time; for example, in blood this measurement is often given as cm/sec. Flow refers to the movement of a *volume* of a liquid or gas over time; for example, in blood this measurement is often given as mL/sec

At its simplest, imagine a perfect, rigid tube with no resistance and with a homogeneous liquid flowing through in a perpendicular manner. Flow can be calculated using the following formula:

$F=v \cdot aF=v \cdot a$

Where F = flow, v = velocity and a = cross-sectional area.

Potential Complications

While the above example is a simple calculation, in reality there are numerous factors that influence velocity and flow.

Velocity and Pressure

Movement of blood throughout the circulatory system is created by differences in pressure generated by the pumping of the heart. Pressure is greatest immediately after exiting the heart and drops as it circulates around the body, particularly through the arterioles and capillary networks. A greater difference in pressure results in a greater velocity assuming all else remains equal, so when increased blood flow is required the heart can pump more quickly and also in larger volume.

Velocity and Resistance

Resistance is the force that must be overcome by pressure in order for flow to occur, and is a factor of vessel length, diameter, surface composition, and the viscosity of the liquid flowing through. As resistance increases the difference in pressure which influences velocity decreases, which in turn reduces flow. For this reason, the narrow arterioles rapidly reduce local blood pressure and slow the flow of blood through the capillaries, a beneficial effect allowing for efficient transfer of chemicals and nutrients. However, pathological changes in blood vessels that result in narrowing or an increase in surface resistance can lead to a reduction in pressure, velocity, and thus flow, which can in turn lead to tissue damage.

Velocity and Viscosity

Blood is a complex liquid formed from plasma and containing numerous cell types. As such, its viscosity is changeable depending on osmotic balance and cell load. Increases in viscosity such as reduced water content lead to increases in resistance and thus reduction in flow.

Vessel Area

Blood vessels are capable of vasodilation and vasoconstriction to alter their diameter. Assuming all else remains equal, a reduction in diameter results in a reduction in flow, whereas an increase in vessel diameter results in an increase in flow.

Regulation

These individual elements are tightly regulated by the body to maintain sufficient flow to the body's organs and tissues

This page titled 18.7A: Velocity of Blood Flow is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7B: Local Regulation of Blood Flow

Blood flow is regulated locally in the arterioles and capillaries using smooth muscle contraction, hormones, oxygen, and changes in pH.

Learning Objectives

- Describe local regulation of blood flow

Key Points

- Regulation of blood flow is managed by adjusting the contraction or relaxation of smooth muscle fibers in the walls of arterioles and capillaries.
- This control can be systemic, affecting the whole circulatory system, or localized to specific tissues or organs.
- Arterioles are the primary blood vessel for local control due to their physical location within tissues and ability to vasodilate and vasocontract to influence blood flow.

Key Terms

- **capillary**: Any of the small blood vessels that connect arteries to veins.
- **arteriole**: One of the small branches of an artery, especially one that connects with capillaries.

The flow of blood along arteries, arterioles, and capillaries is not constant, but can be controlled depending upon the requirements of the body. For example, more blood is directed to the skeletal muscles, brain, or digestive system when they are active, and blood flow to the skin can be reduced or increased to aid with thermoregulation. Blood flow is regulated by vasoconstriction or vasodilation of smooth muscle fibers in the walls of blood vessels, typically arterioles. This regulation can be systemic, affecting the whole of the circulatory system, or localized to specific tissues or organs.

Local Regulation

The greatest change in blood pressure and velocity of blood flow occurs at the transition of arterioles to capillaries. This reduces the pressure and velocity of flow for gas and nutrient exchange to occur within the capillaries. As such arterioles are the main part of the circulatory system in which local control of blood flow occurs.

Arterioles contain smooth muscle fibers in their tunica media, which allows for fine control of their diameter. They are innervated and so can respond to nervous system stimuli and also various circulating hormones. Local responses to stretch, carbon dioxide, pH, and oxygen also influence smooth muscle tone and thus vasoconstriction and vasodilation.

Generally, norepinephrine and epinephrine (hormones secreted by sympathetic nerves and the adrenal gland medulla) are vasoconstrictive, acting on alpha-1-adrenergic receptors. However, the arterioles of skeletal muscle, cardiac muscle, and the pulmonary circulation vasodilate in response to these hormones acting on beta-adrenergic receptors. Generally, stretch and high oxygen tension increase tone, and carbon dioxide and low pH promote vasodilation.

Pulmonary arterioles are a noteworthy exception as they vasodilate in high oxygen. Brain arterioles are particularly sensitive to pH, with reduced pH promoting vasodilation.

A number of hormones influence arteriole tone such as the vasoconstrictive epinephrine, angiotensin II, and endothelin and the vasodilators bradykinin and prostacyclin.

This page titled 18.7B: Local Regulation of Blood Flow is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7C: Blood Flow in Skeletal Muscle

Blood flow to an active muscle changes depending on exercise intensity and contraction frequency and rate.

Learning Objectives

- Summarize the factors involved in blood flow to skeletal muscles

Key Points


- The circulatory system is closely associated with skeletal muscle to provide efficient transfer of oxygen and nutrients required for contraction and the removal of inhibitory waste products.
- At rest, skeletal muscle uses approximately 20% of cardiac output, which can rise to 80% during exercise.
- Return of blood to the heart, especially from the legs, is facilitated by the skeletal muscle pump. Deep-lying veins are compressed by skeletal muscle, forcing blood through the circulatory system back to the heart.
- After repeated stimulation, vascular recruitment can lead to an increase in the number of capillaries present in a muscle tissue, facilitating better supply and more efficient removal of waste products.

Key Terms

- **skeletal muscle pump**: The mechanism whereby skeletal muscles aid the return of blood to the heart by compressing embedded veins.
- **vascular recruitment**: The increase in the number of capillaries in response to a stimulus; for example, repeated exercise results in an increase in the number of capillaries in a skeletal muscle.

Skeletal muscles are important in maintaining posture and controlling locomotion through contraction. For this reason, they receive approximately 20% of cardiac output at rest, which can increase up to a maximum of approximately 80% with exercise. Due to the requirements for large amounts of oxygen and nutrients, muscle vessels are under very tight autonomous regulation to ensure a constant blood flow, and so can have a large impact on the blood pressure of associated arteries.

Blood vessels are closely intertwined with skeletal muscle tissues lying between the fascicles, or bundles of muscle fibers. Each muscle is supplied by many capillaries. This close association reduces the diffusion distances, allowing for the efficient exchange of oxygen and nutrients required for contraction and the rapid removal of inhibitory waste products.



Skeletal Muscle: Skeletal muscle: 1] Bone, 2] Perimysium, 3] Blood vessel, 4] Muscle fiber, 5] Fascicle, 6] Endomysium, and 7] Epimysium Tendon.


Blood Flow During Exercise

Blood flow within muscles fluctuates as they contract and relax. During contraction, the vasculature within the muscle is compressed, resulting in a lower arterial inflow with inflow increased upon relaxation. The opposite effect would be seen if measuring venous outflow.

This rapid increase and decrease in flow is observed over multiple contractions. If the muscle is used for an extended period, mean arterial inflow will increase as the arterioles vasodilate to provide the oxygen and nutrients required for contraction. Following the end of contractions, this increased mean flow remains to resupply the muscle tissue with required nutrients and clear inhibitory waste products, due to the loss of the inhibitory contractile phase.

Skeletal Muscle Pump

Skeletal muscles also play a key role in the movement of blood around the body. Veins embedded within a muscle are compressed during contraction of that muscle, causing an increase in blood pressure due to the presence of one-way valves within the veins. This increase in pressure drives the blood towards the heart. The skeletal muscles of the legs are particularly important skeletal muscle pumps as they prevent pooling of the blood in the feet and calves due to gravity.



Skeletal Muscle Pump: During contraction of the skeletal muscle the vein is compressed which increases blood pressure. Due to the presence of one way valves the blood can pass only in one direction, back towards the heart.

It is unclear whether the action of skeletal muscle pumps influences arterial flow or if this is maintained purely by the pumping of the heart.

Vascular Recruitment

Following repeated stimulus such as through exercise, the number of capillaries present in a muscle tissue can increase. This vascular recruitment increases the capillary surface area within a muscle, allowing for enhanced oxygen exchange with the muscle fibers, prolonging the period of aerobic respiration and thus muscle output, and facilitating a more rapid removal of inhibitory waster factors such as lactic acid, reducing fatigue.

This page titled 18.7C: Blood Flow in Skeletal Muscle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7D: Blood Flow in the Brain

Cerebral circulation is the movement of blood through the network of blood vessels supplying the brain, providing oxygen and nutrients.

Learning Objectives

- Evaluate the results of altered cerebral blood flow

Key Points


- The arteries deliver oxygenated blood, glucose and other nutrients to the brain and the veins carry deoxygenated blood back to the heart, removing carbon dioxide, lactic acid, and other metabolic products.
- In an adult, cerebral blood flow (CBF) is typically 750 milliters per minute or 15% of the cardiac output. CBF is tightly regulated to meet the brain's metabolic demands.
- Too much blood can raise intracranial pressure, which can compress and damage delicate brain tissue. Too little blood flow results in tissue death.
- CBF is determined by a number of factors, such as viscosity of blood, how dilated blood vessels are, and the net pressure of the flow of blood into the brain, which is determined by the body's blood pressure.

Key Terms

- **ischemia**: An inadequate blood supply to an organ or part of the body.
- **cerebral blood flow**: The blood supply to the brain in a given time period.
- **intracranial pressure**: The pressure inside the skull.

Cerebral circulation refers to the movement of blood through the network of blood vessels supplying the brain. The arteries deliver oxygenated blood, glucose, and other nutrients to the brain and the veins carry deoxygenated blood back to the heart, removing carbon dioxide, lactic acid, and other metabolic products. Since the brain is very vulnerable to compromises in its blood supply, the cerebral circulatory system has many safeguards. The circle of Willis, a circulatory anastomosis that supplies blood to the brain and surrounding structures while providing redundancy in case of any interruption, is a key protection. Failure of these safeguards results in cerebrovascular accidents, commonly known as strokes.

The amount of blood that the cerebral circulation carries is known as cerebral blood flow (CBF). In an adult, CBF is typically 750 milliters per minute or 15% of the cardiac output. CBF is tightly regulated to meet the brain's metabolic demands. Too much blood can raise intracranial pressure (ICP), which can compress and damage delicate brain tissue. Too little blood flow (ischemia) results in tissue death. In brain tissue, a biochemical cascade known as the ischemic cascade is triggered when the tissue becomes ischemic, potentially resulting in damage to and death of brain cells. Medical professionals must take steps to maintain proper CBF in patients who have conditions like shock, stroke, and traumatic brain injury.



Brain Blood Flow: Schematic representation of the circle of Willis, arteries of the brain, and brain stem.

This page titled 18.7D: Blood Flow in the Brain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7E: Blood Flow in the Skin

Blood flow to the skin provides nutrition to skin and regulates body heat through the constriction and dilation of blood vessels.

Learning Objectives

- Identify the factors involved in blood flow in the skin

Key Points

- The small blood vessels in the skin contain muscles in their tunica media under the control of the sympathetic nervous system.
- Contraction or relaxation of these muscles leads to vasoconstriction and vasodilation respectively.
- This mechanism is used to alter loss of radiant heat, providing thermoregulation.

Key Terms

- **anastomoses**: The reconnection of two blood vessels that previously branched out.
- **cutaneous**: Of, relating to, existing on, or affecting the exterior skin, especially the cutis.

The skin contains a network of small blood vessels containing muscle fibers in their tunica media. These muscles are under the control of the sympathetic nervous system and provide an efficient means of thermoregulation through vasoconstriction and vasodilation.

When vasoconstricted, blood flow through the skin is reduced, so less core heat is lost. With restricted blood flow, the skin appears paler. When vasodilated, blood flow through the skin is increased, meaning more core heat can be lost through radiation. With increased blood flow, the skin appears red.

Arteriovenous anastomoses can be found in areas of the body exposed to maximal cooling, such as the hands, feet, nose, lips and ears. These richly innervated areas are called apical structures. The anastomoses connect cutaneous arterioles and venules directly, playing an important role in the reduction of blood flow in a cold environment.

This page titled 18.7E: Blood Flow in the Skin is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7F: Blood Flow in the Lungs

Pulmonary circulation in the lungs is responsible for removing carbon dioxide from and replacing oxygen in deoxygenated blood.

Learning Objectives

- Outline the path of pulmonary circulation: blood flow in the lungs

Key Points


- Oxygen-depleted blood is pumped away from the heart via the pulmonary arteries.
- The lung's capillaries line small sacs in the lungs called the alveoli. The alveoli are the site of oxygen and carbon dioxide exchange in the lungs.
- Air is inhaled through the nose or the mouth and fills the lungs. Oxygen passively flows from the air inside the alveoli into the blood in the alveolar capillaries, while carbon dioxide passively flows in the opposite direction. The air, along with the diffused carbon dioxide, is then exhaled.
- The oxygenated blood then leaves the lungs through pulmonary veins, which return it to the left heart, completing the pulmonary cycle.

Key Terms

- **Alveoli**: A small air sac in the lungs where oxygen and carbon dioxide are exchanged with blood.
- **pulmonary vein**: The vein that returns blood to the left atrium of the heart after it has circulated through the lungs.
- **pulmonary artery**: The artery that connects the right ventricle of the heart to the lungs.

The pulmonary circulatory system is the portion of the cardiovascular system in which oxygen-depleted blood is pumped away from the heart to the lungs via the pulmonary artery. Oxygenated blood is then returned to the heart via the pulmonary vein.

From the right ventricle of the heart, blood is pumped through the pulmonary semilunar valve into the left and right pulmonary arteries (one for each lung) and travels through the lungs. The pulmonary arteries carry deoxygenated blood to the lungs, where they release carbon dioxide and pick up oxygen during respiration.



An Alveolus: The alveoli are the site of oxygen and carbon dioxide exchange in the lungs.

The pulmonary arteries divide into thin-walled capillaries closely associated with the alveoli, small air sacs in the lungs where gas exchange occurs. Air is inhaled through the nose or the mouth and fills the lungs. Oxygen passively flows from the air inside the alveoli into the blood in the alveolar capillaries, while carbon dioxide passively flows in the opposite direction. The air, along with the diffused carbon dioxide, is then exhaled.

The oxygenated blood then leaves the lungs through pulmonary veins, which return it to the left atrium of the heart, completing the pulmonary cycle. This blood is pumped through the bicuspid valve into the left ventricle, then distributed to the body through the systemic circulation before returning to the right atrium.

This page titled 18.7F: Blood Flow in the Lungs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7G: Blood Flow in the Heart

The heart pumps oxygenated blood to the body and deoxygenated blood to the lungs.

Learning Objectives

- Sketch the structure of the heart and how blood flows through it

Key Points


- The heart contains four chambers: two atria and two ventricles.
- The blood that is returned to the right atrium is deoxygenated, then passed into the right ventricle to be pumped through the pulmonary artery to the lungs for reoxygenation and removal of carbon dioxide.
- The left atrium receives newly-oxygenated blood from the lungs through the pulmonary veins. This blood is passed into the strong left ventricle to be pumped through the aorta to the different organs of the body.
- The coronary circulation supplies the heart muscle with the oxygen and nutrients it requires to function.

Key Terms

- **Coronary circulation**: The circulation of blood in the blood vessels of the heart muscle.
- **cardiovascular system**: In humans, this system is comprised of the heart, blood, and blood vessels.

The heart is a key organ in the circulatory system responsible for the generation of pressure and thus flow throughout the system and pulmonary circulatory systems.

The heart contains four chambers, two atria and two ventricles. The blood that is returned to the right atrium is deoxygenated and s passed into the right ventricle to be pumped through the pulmonary artery to the lungs for reoxygenation and removal of carbon dioxide. The left atrium receives newly oxygenated blood from the lungs through the pulmonary veins, which is passed into the strong left ventricle to be pumped through the aorta to the different organs of the body.




The Heart: Circulation of blood through the chambers of the heart. Deoxygenated blood is received from the systemic circulation into the right atrium, it is pumped into the right ventricle and then through the pulmonary artery into the lungs. Through association with the alveoli the blood is oxygenated in the lungs and returns to the left atrium through the pulmonary

veins, before passing into the left ventricle and being pumped around the body.

Coronary Blood Supply

The heart has its own circulation system, coronary circulation, which is part of systemic circulation. The coronary arteries derive from the aorta and run along the surface of the heart and within the muscle to deliver oxygen-rich blood to the myocardium. The coronary veins remove deoxygenated blood from the heart muscle, returning it through the coronary sinus into the right atrium.

When healthy, the coronary arteries are capable of autoregulation to maintain blood flow at levels appropriate to the needs of the heart muscle. Blockage of these vessels can lead to poor oxygenation of the heart muscle, impairing its function and in severe cases leading to tissue death, resulting in a myocardial infarction or heart attack.



Coronary Circulation (Anterior)

Coronary Circulation: The coronary circulation supplies the heart muscle with the oxygen and nutrients it requires to function.

This page titled 18.7G: Blood Flow in the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.7H: Hepatic Portal Circulation

The hepatic portal system is responsible for directing blood from parts of the gastrointestinal tract to the liver.

Learning Objectives


- Describe the blood flow of the hepatic portal system

Key Points

- The hepatic portal system is the system of veins comprising the hepatic portal vein and its tributaries.
- The liver consumes about 20% of total body oxygen when at rest, so the total liver blood flow is quite high.
- Blood flow to the liver is unique in that it receives both oxygenated and partially deoxygenated blood.

Key Terms

- **sinusoid**: Any of several channels through which venous blood passes in various organs.
- **hepatic portal system**: In human anatomy, the hepatic portal system is the system of veins comprising the hepatic portal vein and its tributaries.
- **hepatocyte**: Any of the cells in the liver responsible for the metabolism of protein, carbohydrate, and lipids and for detoxification.




Rat Liver Sinusoid: Sinusoid of a rat liver with fenestrated endothelial cells. Fenestrae are approx 100nm diameter, and sinusoidal width 5 microns. Original mag 30,000x. Note the microvilli of hepatocytes in the space of Disse external to the endothelium.

The hepatic portal system is the system of veins comprising the hepatic portal vein and its tributaries. It is responsible for directing blood from the region of the gastrointestinal tract between the esophagus and rectum and also includes venous drainage from the supplementary organs such as the spleen and pancreas. It functions to supply the liver with metabolites and ensures that ingested substances are processed in the liver before reaching the systemic circulation, limiting the damage ingested toxins may cause.

The hepatic portal vein supplies about 75% of the blood the liver requires, with the other 25% supplied by the hepatic artery.

Blood from the hepatic artery is oxygenated but nutrient-poor compared to that supplied by the hepatic portal vein. Blood from either source passes into cavities between the hepatocytes of the liver called sinusoids, which feature a fenestrated, discontinuous endothelium allowing for the effecient transfer and processing of nutrients in the liver. Since blood received from the hepatic portal vein may be contaminated with pathogens such as bacteria, the liver is rich in specialized immune cells called Kupffer cells that detect and destroy foreign organisms. Following processing, blood collects in a central vein that drains into the hepatic vein and finally the inferior vena cava.

The liver consumes about 20% of the total body oxygen when at rest. That is why the total liver blood flow is quite high at about 1 liter a minute and up to two liters a minute. That is about a quarter of the average cardiac output at rest.



Hepatic Portal Circulation: Diagram showing the hepatic portal vein and its territory.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Blood pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blood_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The cardiovascular system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...fter_the_Heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Total peripheral flow resistance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Total_p...low_resistance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Aorta. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Aorta%2...w_and_velocity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- total peripheral resistance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/total%2...l%20resistance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- aorta. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/aorta**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypertension. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and Physiology of Animals/Cardiovascular System/Blood circulation. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Anatomy..._Of_Blood_Flow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vasoconstriction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vasoconstriction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Arterioles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arterioles**. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- Hemodynamics. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemodynamics**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vasodilators. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vasodilators**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- blood flow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/blood%20flow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- arteriole. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/arteriole**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- capillary. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/capillary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Dynamic Medicine. **Provided by**: BioMed Central. **Located at**: **www.biomedcentral.com/1476-5918/content/6/1/5**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Exercise physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Exercis...ology%23Oxygen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Vascular recruitment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vascular_recruitment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Skeletal-muscle pump. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeletal-muscle_pump**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vascular recruitment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vascular%20recruitment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- muscle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/muscle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- perfusion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/perfusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...tal_muscle.png**. **License**: ***Public Domain: No Known Copyright***
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:2..._Vein_Pump.jpg**. **License**: ***CC BY: Attribution***
- Cerebral circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cerebral_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain blood flow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brain_blood_flow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cerebral blood flow. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cerebral%20blood%20flow**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intracranial pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intracranial%20pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ischemia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ischemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...tal_muscle.png**. **License**: ***Public Domain: No Known Copyright***
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:2..._Vein_Pump.jpg**. **License**: ***CC BY: Attribution***
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:C..._Willis_en.svg**. **License**: ***Public Domain: No Known Copyright***
- Blushing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blushing%23Blood_vessels**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plexus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plexus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cutaneous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cutaneous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anastomoses. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anastomoses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...tal_muscle.png**. **License**: ***Public Domain: No Known Copyright***
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:2..._Vein_Pump.jpg**. **License**: ***CC BY: Attribution***
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Circle_of_Willis_en.svg**. **License**: ***Public Domain: No Known Copyright***

- pulmonary vein. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pulmonary_vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circula...ry_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary alveolus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_alveolus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physics/de...ion/alveoli--2. **License**: *CC BY-SA: Attribution-ShareAlike*
- pulmonary artery. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pulmonary_artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a2/Skeletal_muscle.png. **License**: *Public Domain: No Known Copyright*
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:2114_Skeletal_Muscle_Vein_Pump.jpg. **License**: *CC BY: Attribution*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Circle_of_Willis_en.svg. **License**: *Public Domain: No Known Copyright*
- File:Alveolus diagram.svg - Wikimedia Commons. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/w/index...ram.svg&page=1. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coronary circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coronary_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circula...system%23Heart. **License**: *CC BY-SA: Attribution-ShareAlike*
- Coronary circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Coronary%20circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardiovascular system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cardiovascular%20system. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a2/Skeletal_muscle.png. **License**: *Public Domain: No Known Copyright*
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:2114_Skeletal_Muscle_Vein_Pump.jpg. **License**: *CC BY: Attribution*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Circle_of_Willis_en.svg. **License**: *Public Domain: No Known Copyright*
- File:Alveolus diagram.svg - Wikimedia Commons. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/w/index.php?title=File:Alveolus_diagram.svg&page=1. **License**: *CC BY-SA: Attribution-ShareAlike*
- Diagram of the human heart (cropped). **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Blausen_0260_CoronaryVessels_Anterior.png. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0260_CoronaryVessels_Anterior.png. **License**: *CC BY: Attribution*
- Hepatic portal circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hepatic_portal_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver sinusoid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Liver_sinusoid. **License**: *CC BY-SA: Attribution-ShareAlike*
- sinusoid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sinusoid. **License**: *CC BY-SA: Attribution-ShareAlike*
- hepatocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hepatocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- hepatic portal system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/hepatic%20portal%20system. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a2/Skeletal_muscle.png. **License**: *Public Domain: No Known Copyright*
- 2114_Skeletal_Muscle_Vein_Pump.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:2114_Skeletal_Muscle_Vein_Pump.jpg. **License**: *CC BY: Attribution*
- Circle of Willis en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Circle_of_Willis_en.svg. **License**: *Public Domain: No Known Copyright*
- File:Alveolus diagram.svg - Wikimedia Commons. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/w/index.php?title=File:Alveolus_diagram.svg&page=1. **License**: *CC BY-SA: Attribution-ShareAlike*

- Diagram of the human heart (cropped). **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Diagram_of_the_human_heart_(cropped).svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blausen_0260_CoronaryVessels_Anterior.png. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/File:Blausen_0260_CoronaryVessels_Anterior.png**. **License**: ***CC BY: Attribution***
- Hepatic Portal Circulation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/33/Gray591.png**. **License**: ***Public Domain: No Known Copyright***
- Sinusoid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sinusoid.jpeg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 18.7H: Hepatic Portal Circulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.8: CAPILLARY EXCHANGE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.8A: Capillary Dynamics

18.8B: Transcytosis

18.8C: Bulk Flow: Filtration and Reabsorption

This page titled 18.8: Capillary Exchange is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.8A: Capillary Dynamics

Hydrostatic and osmotic pressure are opposing factors that drive capillary dynamics.

Learning Objectives

- Describe hydrostatic pressure and osmotic pressure, the factors of capillary dynamics

Key Points

- Capillary exchange refers to the exchange of material from the blood into the tissues in the capillary.
- There are three mechanisms that facilitate capillary exchange: diffusion, transcytosis and bulk flow.
- Capillary dynamics are controlled by the four Starling forces.
- Oncotic pressure is a form of osmotic pressure exerted by proteins either in the blood plasma or interstitial fluid.
- Hydrostatic pressure is a force generated by the pressure of fluid on the capillary walls either by the blood plasma or interstitial fluid.
- The net filtration pressure is the balance of the four Starling forces and determines the net flow of fluid across the capillary membrane.

Key Terms

- **proteinuria**: Excessive protein in the urine, a condition which can alter the net filtration pressure altering flow of fluid across the capillary wall.
- **hydrostatic pressure**: A pressure generated by fluid on the walls of the capillary, usually forcing water out of the circulatory system.
- **net filtration pressure**: The balance of the four Starling forces that determines the net flow of fluid across the capillary membrane.
- **oncotic pressure**: A form of osmotic pressure exerted by proteins in a fluid that usually tends to pull water into the circulatory system.

Capillary exchange refers to the exchange of material between the blood and tissues in the capillaries. There are three mechanisms that facilitate capillary exchange: diffusion, transcytosis, and bulk flow.

Capillary Exchange Mechanisms

Diffusion, the most widely-used mechanism, allows the flow of small molecules across capillaries such as glucose and oxygen from the blood into the tissues and carbon dioxide from the tissue into the blood. The process depends on the difference of gradients between the interstitium and blood, with molecules moving to low-concentrated spaces from high-concentrated ones.

Transcytosis is the mechanism whereby large, lipid-insoluble substances cross the capillary membranes. The substance to be transported is endocytosed by the endothelial cell into a lipid vesicle which moves through the cell and is then exocytosed to the other side.


Bulk flow is used by small, lipid-insoluble solutes in water to cross the the capillary wall. The movement of materials across the wall is dependent on pressure and is bi-directional depending on the net filtration pressure derived from the four Starling forces that modulate capillary dynamics.

Capillary Dynamics

The four Starling forces modulate capillary dynamics.

- Oncotic or colloid osmotic pressure is a form of osmotic pressure exerted by proteins in the blood plasma or interstitial fluid.
- Hydrostatic pressure is the force generated by the pressure of fluid within or outside of capillary on the capillary wall.

The net filtration pressure derived from the sum of the four forces described above determines the fluid flow into or out of the capillary. Movement from the bloodstream into the interstitium is favored by blood hydrostatic pressure and interstitial fluid oncotic pressure. Alternatively, movement from the interstitium into the bloodstream is favored by blood oncotic pressure and interstitial fluid hydrostatic pressure.



Capillary Dynamics: Oncotic pressure exerted by proteins in blood plasma tends to pull water into the circulatory system.

Due to the pressure of the blood in the capillaries, blood hydrostatic pressure is greater than interstitial fluid hydrostatic pressure, promoting a net flow of fluid from the blood vessels into the interstitium. However, because large plasma proteins, especially albumin, cannot easily cross through the capillary walls, their effect on the osmotic pressure of the capillary interiors will to some extent balance the tendency for fluid to leak from the capillaries.In conditions where plasma proteins are reduced (e.g. from being lost in the urine or from malnutrition), or blood pressure is significantly increased, a change in net filtration pressure and an increase in fluid movement across the capillary result in excess fluid build-up in the tissues (edema).

This page titled 18.8A: Capillary Dynamics is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.8B: Transcytosis

Transcytosis is a process by which molecules are transported into the capillaries.

Learning Objectives

- Describe the process of transcytosis in capillary exchange

Key Points

- Transcytosis is the process by which various macromolecules are transported across the endothelium of the capillaries.
- Due to this function, transcytosis can be a convenient mechanism for pathogens to invade a tissue.

Key Terms

- **transcytosis**: The process whereby macromolecules are transported across the interior of a cell via vesicles.

Transcytosis, or vesicle transport, is one of three mechanisms that facilitate capillary exchange, along with diffusion and bulk flow.

Substances are transported through the endothelial cells themselves within vesicles. This mechanism is mainly used by large molecules, typically lipid-insoluble preventing the use of other transport mechanisms. The substance to be transported is endocytosed by the endothelial cell into a lipid vesicle which moves through the cell and is then exocytosed to the other side. Vesicles are capable of merging, allowing for their contents to mix, and can be transported directly to specific organs or tissues.

Pathology

Due to the function of transcytosis, it can be a convenient mechanism by which pathogens can invade a tissue. Transcytosis has been shown to be critical to the entry of *Cronobacter sakazakii* across the intestinal epithelium and the blood-brain barrier.

Listeria monocytogenes has been shown to enter the intestinal lumen via transcytosis across goblet cells. Shiga toxin secreted by entero-hemorrhagic *E. coli* has been shown to be transcytosed into the intestinal lumen. These examples illustrate that transcytosis is vital to the process of pathogenesis for a variety of infectious agents.

Transcytosis in Pharmaceuticals

Pharmaceutical companies are currently exploring the use of transcytosis as a mechanism for transporting therapeutic drugs across the human blood-brain barrier. Exploiting the body's own transport mechanism can help to overcome the high selectivity of this barrier, which blocks the uptake of most therapeutic antibodies into the brain and central nervous system.

This page titled 18.8B: Transcytosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.8C: Bulk Flow: Filtration and Reabsorption

Capillary fluid movement occurs as a result of diffusion (colloid osmotic pressure), transcytosis, and filtration.

Learning Objectives

- Explain the process of filtration and reabsorption in capillaries

Key Points

- Bulk flow is a process used by small lipid-insoluble proteins to cross the capillary wall.
- Capillary structure plays a large role in the rate of bulk flow, with continuous capillaries limiting flow and discontinuous capillaries facilitating the greatest amount of flow.
- When moving from the blood to the interstitium, bulk flow is termed filtration.
- When moving from the interstitium to the blood, bulk flow is termed re-absorption.
- The kidney is a major site of bulk flow where waste products are filtered from the blood.

Key Terms

- **filtration**: In bulk flow, this refers to the movement of proteins or other large molecules from the blood into the interstitium.
- **reabsorption**: In bulk flow, this refers to the movement of proteins or other large molecules from the interstitium into the blood.

Bulk flow is one of three mechanisms that facilitate capillary exchange, along with diffusion and transcytosis.

Bulk Flow Process

Bulk flow is used by small, lipid-insoluble solutes in water to cross the the capillary wall and is dependent on the physical characteristics of the capillary. Continuous capillaries have a tight structure reducing bulk flow. Fenestrated capillaries permit a larger amount of flow and discontinuous capillaries allow the largest amount of flow.

The movement of materials across the capillary wall is dependent on pressure and is bidirectional depending on the net filtration pressure derived from the four Starling forces.

When moving from the bloodstream into the interstitium, bulk flow is termed filtration, which is favored by blood hydrostatic pressure and interstitial fluid oncotic pressure. When moving from the interstitium into the bloodstream, the process is termed reabsorption and is favored by blood oncotic pressure and interstitial fluid hydrostatic pressure.


Modern evidence shows that in most cases, venular blood pressure exceeds the opposing pressure, thus maintaining a positive outward force. This indicates that capillaries are normally in a state of filtration along their entire length.

The Kidneys and Bulk Flow

The kidney is a major site for bulk flow transport. Blood that enters the kidneys is filtered by nephrons, the functional unit of the kidney. Each nephron begins in a renal corpuscle composed of a glomerulus containing numerous capillaries enclosed in a Bowman's capsule. Proteins and other large molecules are filtered out of the oxygenated blood in the glomerulus and pass into Bowman's capsule and the tubular fluid contained within. Blood continues to flow around the nephron until it reaches another capillary-rich region the peritubular capillaries, where the previously filtered molecules are reabsorbed from the tubule of the nephron.

Tubular reabsorption is the process by which solutes and water are removed from the tubular fluid and transported into the blood.

Reabsorption is a two-step process beginning with the active or passive extraction of substances from the tubule fluid into the renal interstitium, and then the transport of these substances from the interstitium into the bloodstream



Tubular Secretion: Diagram showing the basic physiologic mechanisms of the kidney and the three steps involved in urine formation.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Starling equation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Starling_equation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Colloid osmotic pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Colloid_osmotic_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Microcirculation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Microcirculation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Starling equation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Starling_equation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Capillary Exchange. January 16, 2014. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/contents/797b2238-a9ed-4012-b9af-ce0a25172732@3**. **License**: ***CC BY: Attribution***
- Hydrostatic pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hydrostatic_pressure%23Hydrostatic_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endothelium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endothelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- proteinuria. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/proteinuria**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- oncotic pressure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/oncotic%20pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Capillary Microcirculation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...irculation.jpg**. **License**: ***Public Domain: No Known Copyright***
- Transcytosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Transcytosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Transcytosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Transcytosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- transcytosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/transcytosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Capillary Microcirculation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/0/03/Illu_capillary_microcirculation.jpg**. **License**: ***Public Domain: No Known Copyright***
- Renal physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Renal_p...y%23Mechanisms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Starling equation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Starling_equation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- reabsorption. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/reabsorption**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- filtration. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/filtration**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- active transport. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/active%20transport**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Capillary Microcirculation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/0/03/Illu_capillary_microcirculation.jpg**. **License**: ***Public Domain: No Known Copyright***
- Urinary Excretion. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...of_Nephron.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 18.8C: Bulk Flow: Filtration and Reabsorption is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.9: CIRCULATORY SHOCK

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

18.9A: Types of Shock

18.9B: Homeostatic Responses to Shock

18.9C: Signs and Symptoms of Shock

This page titled 18.9: Circulatory Shock is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.9A: Types of Shock

Circulatory shock is a life-threatening medical condition that occurs due to inadequate substrate for aerobic cellular respiration.

Learning Objectives

- Differentiate among the types of shock


Key Points

- Circulatory shock, commonly known simply as shock, is a life-threatening medical condition that occurs due to the provision of inadequate substrates for cellular respiration.Typical symptoms of shock include elevated but weak heart rate, low blood pressure, and poor organ function, typically observed as low urine output, confusion, or loss of consciousness.
- There are four subtypes of shock with differing underlying causes and symptoms: hypovolemic, cardiogenic, obstructive, and distributive.
- Distributive shock can be further divided into septic, anaphylaxis, and neurogenic shock.

Key Terms

- **shock**: A medical condition that occurs due to an inadequate supply of substrates required for aerobic respiration by the bodies tissues.

Circulatory shock, commonly known simply as shock, is a life-threatening medical condition that occurs due to inadequate substrates for aerobic cellular respiration. In the early stages, this is generally caused by an inadequate tissue level of oxygen. The typical signs of shock are low blood pressure, a rapid heartbeat, and signs of poor end-organ perfusion or decompensation (such as low urine output, confusion, or loss of consciousness). In some people with circulatory shock, blood pressure remains stable.



Shock: The scheme depicts the cell metabolic response as a result of inadequate blood delivery during circulatory shock.

The presentation of shock is variable with some people having only minimal symptoms such as confusion and weakness. While the general signs for all types of shock are low blood pressure, decreased urine output, and confusion, these may not always be present. Specific subtypes of shock may have additional symptoms.

Hypovolemic Shock

Hypovolemic shock, the most common type, is caused by insufficient circulating volume, typically from hemorrhage although severe vomiting and diarrhea are also potential causes.

Hypovolemic shock is graded on a four-point scale depending on the severity of symptoms and level of blood loss. Typical symptoms include a rapid, weak pulse due to decreased blood flow combined with tachycardia, cool, clammy skin, and rapid and shallow breathing.

Cardiogenic Shock

Cardiogenic shock is caused by a failure of the heart to pump correctly, either due to damage to the heart muscle through myocardial infarction or through cardiac valve problems, congestive heart failure, or dysrhythmia.

Obstructive Shock

Obstructive shock is caused by an obstruction of blood flow outside of the heart. This typically occurs due to a reduction in venous return, but may also be caused by blockage of the aorta.

Distributive Shock

Distributive shock is caused by an abnormal distribution of blood to tissues and organs and includes septic, anaphylactic, and neurogenic causes.

Septic

Septic shock is the most common cause of distributive shock and is caused by an overwhelming systemic infection that cannot be cleared by the immune system, resulting in vasodilation and hypotension.

Anaphylactic

Anaphylactic shock is caused by a severe reaction to an allergen, leading to the release of histamine that causes widespread vasodilation and hypotension.

Neurogenic

Neurogenic shock arises due to damage to the central nervous system, which impairs cardiac function by reducing heart rate and loosening the blood vessel tone, resulting in severe hypotension.

This page titled 18.9A: Types of Shock is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.9B: Homeostatic Responses to Shock

An organism responds with numerous reactions during each of the four stages of shock in an attempt to maintain cellular homeostasis.

Learning Objectives

- Describe the homeostatic response to shock


Key Points

- There are four stages of shock. As it is a complex and continuous condition, there is no sudden transition from one stage to the next.
  The initial stage of shock is characterized by hypoxia and anaerobic cell respiration leading to lactic acidosis.
- The compensatory stage is characterized by the employment of neural, hormonal, and biochemical mechanisms in the body's attempt to reverse the condition.
- The progressive stage is the point at which the compensatory mechanisms will begin to fail. If the crisis is not treated successfully, vital organs might be compromised.
- The refractory stage is when vital organs have failed and the shock can no longer be reversed leading to imminent death.

Key Terms

- **hypoperfusion**: Decreased perfusion of blood through an organ.
- **hypoxia**: A condition in which tissues (especially the blood) are deprived of an adequate supply of oxygen; anoxia.

Circulatory shock, commonly known simply as shock, is a life-threatening medical condition that occurs due to inadequate substrate for aerobic cellular respiration. In the early stages this is generally an inadequate level of oxygen in the tissues.



Shock: The scheme depicts the cell metabolic response as a result of inadequate blood delivery during circulatory shock.

There are four stages of shock. As it is a complex and continuous condition there is no sudden transition from one stage to the next.

Initial Stage

During the initial stage, the state of hypoperfusion causes hypoxia. Due to the lack of oxygen, the cells perform anaerobic respiration. Since oxygen is not abundant, the Kreb's cycle is slowed, resulting in lactic acidosis (the accumulation of lactate).

Compensatory Stage

The *compensatory stage* is characterized by the employment of neural, hormonal, and biochemical mechanisms in the body's attempt to reverse the lactic acidosis. The increase in acidity will initiate the Cushing reflex, generating the classic symptoms of shock. The individual will begin to hyperventilate to rid the body of carbon dioxide to raise the blood pH (lower the acidity). As a result, the baroreceptors in the arteries detect the hypotension and initiate the release of epinephrine and norepinephrine to increase heart rate and blood pressure.

Progressive Stage

Should the cause of the crisis not be successfully treated, the shock will proceed to the *progressive stage,* in which the compensatory mechanisms begin to fail. As anaerobic metabolism continues, increasing the body's metabolic acidosis, the arteriolar smooth muscle and precapillary sphincters relax. Blood remains in the capillaries, leading to leakage of fluid and protein into the surrounding tissues. As fluid is lost, blood concentration and viscosity increase, causing blockage of the microcirculation. The prolonged vasoconstriction will also cause the vital organs to be compromised due to reduced perfusion. If the bowel becomes sufficiently ischemic, bacteria may enter the blood stream, resulting in the additional complication of endotoxic shock.

Refractory Stage

At the *refractory stage*, the vital organs have failed and shock can no longer be reversed. Brain damage and cell death are occurring, and death is imminent. Shock is irreversible at this point since a large amount of cellular ATP has been degraded into adenosine in the absence of oxygen as an electron receptor in the mitochondrial matrix. Adenosine easily perfuses out of cellular membranes into extracellular fluid, furthering capillary vasodilation, and then is transformed into uric acid. Because cells can only produce adenosine at a rate of about 2% of the cell's total need per hour, even restoring oxygen is futile at this point because there is no adenosine to phosphorylate into ATP.

This page titled 18.9B: Homeostatic Responses to Shock is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

18.9C: Signs and Symptoms of Shock

The clinical manifestation of shock varies depending on the type of shock and the individual, but there are some general symptoms.

Learning Objectives

- Differentiate among the types of shock and the signs and symptoms associated with each

Key Points

- The general signs for all types of shock are low blood pressure, decreased urine output, and confusion. However, these may not always be present.
- Hypovolemic shock is characterized by loss of effective circulating blood volume, which leads to rapid pulse, cool skin, shallow breathing, hypothermia, thirst, and cold mottled skin.
- Cardiogenic shock is characterized by distended jugular veins, weak or absent pulse, and arrhythmia.
- Distributive shock includes septic shock, characterized by fever or anaphylaxis, and neurogenic shock, characterized by a reduced heart rate and vasodilation of superficial vessels warming the skin.

The presentation of shock varies. Some people presenting only minimal symptoms, such as confusion and weakness. Typical symptoms of shock include elevated but weak heart rate, low blood pressure, and poor organ function, typically observed as low urine output, confusion or loss of consciousness.

While a fast heart rate is common, those on beta blockers and those who are athletic may have a normal or slow heart rate. This also occurs in 30% of cases of shock caused by abdominal bleeding. Specific subtypes of shock may have additional symptoms.

Hypovolemic shock results from the direct loss of effective circulating blood volume. This leads to a rapid, weak pulse due to decreased blood flow combined with tachycardia, stimulation of vasoconstriction, and cool, clammy skin. It also presents with acidosis as well as rapid, shallow breathing due to sympathetic nervous system stimulation. Hypothermia due to decreased perfusion and evaporation of sweat, and thirst and dry mouth due to fluid depletion, may also be present.



Anaphalactic Hives: Hives and flushing on the back of a person with anaphylaxis.

The symptoms of cardiogenic shock are similar to those of hypovolemic shock. Additional symptoms may include arrhythmia of the heart beat and visibly distended jugular veins.

Distributive shock includes septic, anaphylactic, and neurogenic causes. With septic shock, fever may occur and the skin may be warm and sweaty. With anaphylaxis, hives may present on the skin, and there may localized edema, especially around the face,

and weak and rapid pulse. Breathlessness and cough due to narrowed airways and swelling of the throat may also occur. The symptoms of neurogenic shock are distinct from those of classical shock, as the heart rate slows and and superficial vessels vasodilate and warm the skin. These symptoms are caused by neural damage and resultant loss of muscle control.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...s_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neurogenic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neurogenic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anaphylaxis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anaphylaxis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- septic shock. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/septic_shock**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Shock-cell2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Shock-cell2.svg**. **License**: ***Public Domain: No Known Copyright***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...athophysiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Shock (circulatory). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Shock_(circulatory)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circula...athophysiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Pathophysiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Pathophysiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Pathophysiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/shock**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoxia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoxia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoperfusion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoperfusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Shock-cell2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Shock-cell2.svg**. **License**: ***Public Domain: No Known Copyright***
- Shock-cell2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Shock-cell2.svg**. **License**: ***Public Domain: No Known Copyright***
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Signs_and_symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiogenic shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cardiogenic%20shock**. **License**: ***CC BY-***

SA: Attribution-ShareAlike
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Signs_and_symptoms**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Shock (circulatory). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Shock_(circulatory)%23Signs_and_symptoms**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Signs_and_symptoms**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Circulatory shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Circulatory_shock%23Signs_and_symptoms**. **License**: *CC BY-SA: Attribution-ShareAlike*
- distributive shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/distributive%20shock**. **License**: *CC BY-SA: Attribution-ShareAlike*
- anaphylaxis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anaphylaxis**. **License**: *CC BY-SA: Attribution-ShareAlike*
- hypovolemic shock. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hypovolemic%20shock**. **License**: *CC BY-SA: Attribution-ShareAlike*
- septic shock. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/septic_shock**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Shock-cell2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Shock-cell2.svg**. **License**: *Public Domain: No Known Copyright*
- Shock-cell2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Shock-cell2.svg**. **License**: *Public Domain: No Known Copyright*
- BackRed. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:BackRed.JPG**. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 18.9C: Signs and Symptoms of Shock is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19: LYMPHATIC SYSTEM

- Topic hierarchy

19.1: Lymphatic System Structure and Function

19.1A: Structure of the Lymphatic System

19.1B: Functions of the Lymphatic System

19.2: Lymphatic Vessels

19.2A: Lymphatic Vessel Structure

19.2B: Distribution of Lymphatic Vessels

19.2C: Lymph Transport

19.2D: Lymphatic Capillaries

19.2E: Lymph Trunks and Ducts

19.3: Lymph Cells and Tissues

19.3A: Lymphoid Cells

19.3B: Lymphoid Tissue

19.3C: Lymph Nodes

19.4: Other Lymphoid Organs

19.4A: Thymus

19.4B: Spleen

19.4C: Tonsils

This page titled 19: Lymphatic System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.1: LYMPHATIC SYSTEM STRUCTURE AND FUNCTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

19.1A: Structure of the Lymphatic System

19.1B: Functions of the Lymphatic System

This page titled 19.1: Lymphatic System Structure and Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.1A: Structure of the Lymphatic System

The lymphatic system consists of lymphatic vessels and associated lymphoid organs.

Learning Objectives

- Describe the structure and function of the lymphatic system

Key Points

- The lymphatic system is a circulatory system that drains fluid from the blood vessels.
- Lymph vessels are the site of fluid drainage and pump lymph fluid using smooth muscle and skeletal muscle action. The larger vessels contain valves to prevent backflow and pump towards the heart to return lymph fluid to the bloodstream by the subclavian veins.
- A lymph node is an organized collection of lymphoid tissue through which the lymph passes on its way to returning to the blood. Lymph nodes are located at intervals along the lymphatic system.
- Lymphoid tissue contains lymphocytes and other specialized cells and tissues that have immune system functions.

Key Terms


- **lymph node**: Small oval bodies of the lymphatic system, distributed along the lymphatic vessels clustered in the armpits, groin, neck, chest, and abdomen. They filter through lymph fluid.
- **lymph**: A colorless, watery, bodily fluid carried by the lymphatic system, consisting mainly of white blood cells.

The lymphatic system is a collection of structures and vessels that drains lymph from blood and has several other functions. It is a circulatory system for lymph fluid and the site of many key immune system functions.

Lymphatic Vessels

The lymphatic vessels are the lymphatic system equivalent of the blood vessels of the circulatory system and drain fluid from the circulatory system. The network of lymph vessels consists of the initial collectors of lymph fluid, which are small, valveless vessels, and goes on to form the precollector vessels, which have rudimentary valves that are not fully functional. These structures then form increasingly larger lymphatic vessels which form colaterals and have lymph-angions (lymph hearts). The larger lymph vessels contain valves that prevent the backflow of lymph.

The lymphatic system is an active pumping system driven by segments that have a function similar to peristalsis. They lack a central pump (like the heart in the cardio vascular system), so smooth muscle tissue contracts to move lymph along through the vessels. Skeletal muscle contractions also move lymph through the vessels. The lymphatic vessels make their way to the lymph nodes, and from there the vessels form into trunks. In general, the lymph vessels bring lymph fluid toward the heart and above it to the subclavian veins, which enable lymph fluid to re-enter the circulatory system through the vena cava.



The lymphatic system: This diagram shows the network of lymph nodes and connecting lymphatic vessels in the human body.

Lymphatic Tissues and Organs

Lymphoid tissue is found in many organs including the lymph nodes, as well as in the lymphoid follicles in the pharynx such as the tonsils. Lymph nodes are found primarily in the armpits, groin, chest, neck, and abdomen. Lymphoid tissues contain lymphocytes (a type of highly differentiated white blood cell), but they also contain other types of cells for structural and functional support, such as the dendritic cells, which play a key role in the immune system. The system also includes all the structures dedicated to the circulation and production of lymphocytes, including the spleen, thymus, and bone marrow.

This page titled 19.1A: Structure of the Lymphatic System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.1B: Functions of the Lymphatic System

The lymphatic system plays a prominent role in immune function, fatty acid absorption, and removal of interstitial fluid from tissues.

Learning Objectives

- Describe the roles of the lymphatic system

Key Points

- The lymphatic system is a linear network of lymphatic vessels and secondary lymphoid organs. It is the site of many immune system functions as well as its own functions.
- It is responsible for the removal of interstitial fluid from tissues into lymph fluid, which is filtered and brought back into the bloodstream through the subclavian veins near the heart.
- Edema accumulates in tissues during inflammation or when lymph drainage is impaired.
- It absorbs and transports fatty acids and fats as chylomicrons from the digestive system.
- It transports white blood cells and dendritic cells to lymph nodes where adaptive immune responses are often triggered.
- Tumors can spread through lymphatic transport.

Key Terms

- **lacteal**: A lymphatic capillary that absorbs dietary fats in the villi of the small intestine.
- **interstitial fluid**: Also called tissue fluid, a solution that bathes and surrounds the cells of multicellular animals.
- **white blood cell**: A type of blood cell involved with an immune response. Many white blood cells (primarily lymphocytes) are transported by the lymphatic system.


The lymphatic system is the site of many key immune system functions. It is important to distinguish that immune system functions can happen almost anywhere in the body, while the lymphatic system is its own system where many immune system functions take place. Besides immune system function, the lymphatic system has many functions of its own. It is responsible for the removal and filtration of interstitial fluid from tissues, absorbs and transports fatty acids and fats as chyle from the digestive system, and transports many of the cells involved in immune system function via lymph.

Removal of Fluid

Interstitial fluid accumulates in the tissues, generally as a result of the pressure exerted from capillaries (hydrostatic and osmotic pressure) or from protein leakage into the tissues (which occurs during inflammation). These conditions force fluid from the capillaries into the tissues. One of the main functions of the lymphatic system is to drain the excess interstitial fluid that accumulates.

The lymphatic system is a blunt-ended linear flow system, in which tissue fluids, cells, and large extracellular molecules, collectively called lymph, are drained into the initial lymphatic capillary vessels that begin at the interstitial spaces of tissues and organs. They are then transported to thicker collecting lymphatics, which are embedded with multiple lymph nodes, and are eventually returned to the blood circulation through the left and right subclavian veins and into the vena cava. They drain into venous circulation because there is lower blood pressure in veins, which minimizes the impact of lymph cycling on blood pressure. Lymph nodes located at junctions between the lymph vessels also filter the lymph fluid to remove pathogens and other abnormalities.

Fluid removal from tissues prevents the development of edema. Edema is any type of tissue swelling from increased flow of interstitial fluid into tissues relative to fluid drainage. While edema is a normal component of the inflammation process, in some cases it can be very harmful. Cerebral and pulmonary edema are especially problematic, which is why lymph drainage is so important. Abnormal edema can still occur if the drainage components of the lymph vessels are obstructed.



The lymphatic system: A diagram of fluid movement in the lymphatic system.

Fatty Acid Transport

The lymphatic system also facilitates fatty acid absorption from the digestive system. During fat digestion, fatty acids are digested, emulsified, and converted within intestinal cells into a lipoprotein called chylomicrons. Lymph drainage vessels that line the intestine, called lacteals, absorb the chylomicrons into lymph fluid. The lymph vessels then take the chylomicrons into blood circulation, where they react with HDL cholesterols and are then broken down in the liver.

Immune Cell Transport

In addition to tissue fluid homeostasis, the lymphatic system serves as a conduit for transport of cells involved in immune system function. Most notably, highly-specialized white blood cells called lymphocytes and antigen -presenting cells are transported to regional lymph nodes, where the immune system encounters pathogens, microbes, and other immune elicitors that are filtered from the lymph fluid. Much of the adaptive immune system response, which is mediated by dendritic cells, takes place in the lymph nodes. Lymphatic vessels, which uptake various antigens from peripheral tissues, are positively regulated by chemokines/ cytokines secreted by various immune cells during inflammation. This allows antigens to enter lymph nodes, where dendritic cells can present them to lymphocytes to trigger an adaptive immune response.

While the lymphatic system is important for transporting immune cells, its transport capabilities can also provide a pathway for the spread of cancer. Lymph circulation is one of the main ways that tumors can spread to distant parts of the body, which is difficult to prevent.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- lymph node. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lymph_node. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphatic_system%23Lymphoid_tissue. **License**: *CC BY-SA: Attribution-ShareAlike*
- lymphocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lymphocyte. **License**: *CC BY-SA: Attribution-*

ShareAlike
- lymph. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lymph. **License**: *CC BY-SA: Attribution-ShareAlike*
- edema. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/edema. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Lymphatic_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphatic_system%23Functions. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphatic_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- lacteal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lacteal. **License**: *CC BY-SA: Attribution-ShareAlike*
- interstitial fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/interstitial_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- white blood cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/white_blood_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Lymphatic_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and Physiology of Animals/Lymphatic System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy_and_Physiology_of_Animals/Lymphatic_System. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 19.1B: Functions of the Lymphatic System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2: LYMPHATIC VESSELS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

19.2A: Lymphatic Vessel Structure

19.2B: Distribution of Lymphatic Vessels

19.2C: Lymph Transport

19.2D: Lymphatic Capillaries

19.2E: Lymph Trunks and Ducts

This page titled 19.2: Lymphatic Vessels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2A: Lymphatic Vessel Structure

The lymphatic structure is based on that of blood vessels.

Learning Objectives

- Describe lymphatic circulation and the structure of lymphatic vessels

Key Points

- Lymph (or lymphatic) vessels are thin-walled valved structures that carry lymph.
- Lymph vessels are lined by endothelial cells and have a thin layer of smooth muscles and adventitia that bind the lymph vessels to the surrounding tissue.
- Lymph movement occurs despite low pressure due to smooth muscle action, valves, and compression during contraction of adjacent skeletal muscle and arterial pulsation.
- When the pressure inside a lymphangion becomes high enough, lymph fluid will push through the semilunar valve into the next lymphangion, while the valve then closes.
- Lymph vessels are structurally very similar to blood vessels.
- Valves prevent backwards flow of lymph fluid, which allows the lymphatic system to function without a central pump.

Key Terms

- **lymphagion**: The space between two semilunar valves of the lymphatic vessels that forms a distinct functional unit for the forward flow of lymph.
- **adventitia**: The outermost layer of connective tissue encasing a visceral organ or vessel.
- **ISF**: Interstitial (or tissue) fluid, a solution that bathes and surrounds the cells of multicellular animals. It is the main component of extracellular fluid, which also includes plasma and transcellular fluid.
- **endothelial cells**: A thin layer of cells that lines the interior surface of blood and lymphatic vessels, forming an interface between circulating blood or lymph in the lumen and the rest of the vessel wall.

The general structure of lymphatic vessels is similar to that of blood vessels since these are the only two types of vessels in the body. While blood and lymph fluid are two separate substances, both are composed of the same water (plasma or fluid) found elsewhere in the body.

Layers of Lymph Vessels


The endothelium, a general term for the inner layer of a vessel, is composed of an inner lining of single, flattened epithelial cells (simple squamous epithelium). This layer mechanically transports fluid. It sits on a highly permeable basement membrane made out of extracellular matrix that separates the endothelium from the other layers. The endothelium is designed with junctions between cells that allow interstitial fluid to flow into the lumen when pressure becomes high enough (such as from blood capillary hydrostatic pressure), but does not normally allow lymph fluid to leak back out into the interstitial space.

The next layer is *smooth muscles* arranged in a circular fashion around the endothelium that alters the pressure inside the lumen (space) inside the vessel by contracting and relaxing. The activity of smooth muscles allows lymph vessels to slowly pump lymph fluid through the body without a central pump or heart. By contrast, the smooth muscles in blood vessels are involved in vasoconstriction and vasodilation instead of fluid pumping.

The outermost layer is the *adventitia,* consisting of fibrous tissue. It is made primarily out of collagen and serves to anchor the lymph vessels to structures within the body for stability. Larger lymph vessels have many more layers of adventitia than do smaller lymph vessels. The smallest vessels, such as the lymphatic capillaries, may have no outer adventitia. As they proceed forward and integrate into the larger lymph vessels, they develop adventitia and smooth muscle. Blood vessels also have adventitia, sometimes referred to as tunica.

Lymphatic Valves

One of the main structural features of lymph vessels is their valves, which are semilunar structures attached to opposite sides of the lymphatic endothelium. Valves are found in larger lymph vessels and collecting vessels and are absent in the lymphatic capillaries. The valves is to prevent backflow of fluid, so that lymph eventually flows forward instead of falling backwards. When the pressure of lymph fluid increase to a certain point due to filling with more lymph fluid or from smooth muscle contraction, the fluid will be pushed through the valve (opening it) into the next chamber of the vessel (called a lymphangion). As the pressure falls, the open valve then closes so that the lymph fluid cannot flow backwards.



Lymph Vessel: Diagram representing propulsion of lymph through a lymph vessel.

A lymphangion is the term for the space between two semilunar valves in a lymphatic vessel, functional unit of the lymphatic system. Lymph fluid can only flow forward through lymphangions due to the closing of valves after fluid is pushed through by fluid accumulation, smooth muscle contraction, or skeletal muscle contraction.

Without valves, the lymphatic system would be unable to function without a central pump. Smooth muscle contractions only cause small changes in pressure and volume within the lumen of the lymph vessels, so the fluid would just move backwards when the pressure dropped. Blood vessels also have valves, but only in low pressure venous circulation. They function similarly to lymphatic valves, though are comparatively more dependent on skeletal muscle contractions.

This page titled 19.2A: Lymphatic Vessel Structure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2B: Distribution of Lymphatic Vessels

The lymphatic system comprises a network of conduits called lymphatic vessels that carry lymph unidirectionally towards the heart.

Learning Objectives

- Describe the structure of the lymphatic system and its role in the immune system and blood circulation

Key Points


- The lymph system is not a closed system. Lymph flows in one direction toward the heart.
- Lymph nodes are most densely distributed toward the center of the body, particularly around the neck, intestines, and armpits.
- Lymph vessels and nodes are not found within bone or nervous system tissue.
- Afferent lymph vessels flow into lymph nodes, while efferent lymph vessels flow out of them.
- Lymphatic capillaries are the sites of lymph fluid collection, and are distributed throughout most tissues of the body, particularly connective tissue.

Key Terms

- **lymph**: A colorless, watery, bodily fluid carried by the lymphatic system, consisting mainly of white blood cells.
- **plasma**: The straw-colored/pale-yellow liquid component of blood that normally holds the blood cells of whole blood in suspension.
- **Efferent**: A type of vessel that flows out of a structure, such as lymph vessels that leave the spleen or lymph nodes and arterioles that leave the kidney.

The lymphatic system is a circulatory system for lymphatic fluid, comprising a network of conduits called lymphatic vessels that carry the fluid in one direction toward the heart. Its functions include providing sites for certain immune system functions and facilitating plasma circulation in the cardiovascular system. The lymphatic system is composed of many different types of lymph vessels over a wide distribution throughout the body.

Lymph Node Distribution



Lymphatic System: The lymph nodes and lymph vessels in human beings. (Public Domain; NIH via Wikipedia)

Lymphatic vessels are most densely distributed near lymph nodes: bundles of lymphoid tissue that filter the lymph fluid of pathogens and abnormal molecules. Adaptive immune responses usually develop within lymphatic vessels. Large lymphatic vessels can be broadly characterized into two categories based on lymph node distribution.

- Afferent lymphatic vessels flow into a lymph node and carry unfiltered lymph fluid.
- Efferent lymphatic vessels flow out of a lymph node and carry filtered lymph fluid. Lymph vessels that leave the thymus or spleen (which lack afferent vessels) also fall into this category.

Lymph nodes are most densely distributed around the pharynx and neck, chest, armpits, groin, and around the intestines. Afferent and efferent lymph vessels are also most concentrated in these areas so they can filter lymph fluid close to the end of the lymphatic system, where fluid is returned into the cardiovascular system. Conversely, lymph nodes are not found in the areas of the upper central nervous system, where tissue drains into cerebrospinal fluid instead of lymph, though there are some lymph vessels in the meninges. There are few lymph nodes at the ends of the limbs. The efferent lymph vessels in the left and lower side of the body drain into the left subclavian vein through the thoracic duct, while the efferent lymph vessels of the right side of the body drain into the right subclavian vein through the right lymphatic duct.

Flow Through Lymph Vessels

The lymphatic vessels start with the collection of lymph fluid from the interstitial fluid. This fluid is mainly water from plasma that leaks into the intersitial space in the tissues due to pressure forces exerted by capillaries (hydrostatic pressure) or through osmotic forces from proteins (osmotic pressure). When the pressure for interstitial fluid in the interstitial space becomes large enough it leaks into lymph capillaries, which are the site for lymph fluid collection.

Like cardiovascular capillaries, lymph capillaries are well distributed throughout most of the body's tissues, though they are mostly absent in bone or nervous system tissue. In comparison to cardiovascular capillaries, lymphatic capillaries are larger, distributed throughout connective tissues, and have a dead end that completely prevents backflow of lymph. That means the lymphatic system is an open system with linear flow, while the cardiovascular system is a closed system with true circular flow.

Lymph flows in one direction toward the heart. Lymph vessels become larger, with better developed smooth muscle and valves to keep lymph moving forward despite the low pressure and adventia to support the lymph vessels. As the lymph vessels become larger, their function changes from collecting fluid from the tissues to propelling fluid forward. Lymph nodes found closer to the heart filter lymph fluid before it is returned to venous circulation through one of the two lymph ducts.

This page titled 19.2B: Distribution of Lymphatic Vessels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2C: Lymph Transport

Lymph circulates to the lymph node via afferent lymphatic vessels and drains into the lymph node in the subcapsular sinus.

Learning Objectives

- Describe the location of B cells and T cells in lymph nodes and the path of lymph circulation

Key Points

- The sinus space is crisscrossed by the pseudopods of macrophages, which act to trap foreign particles and filter the lymph.
- Lymph then leaves the lymph node via the efferent lymphatic vessel towards either a more central lymph node or for drainage into a central venous subclavian blood vessel.
- Lymphatic transport begins in the lymphatic capillaries, which converge into collecting vessels that flow into afferent vessels, then into lymph nodes.
- The lymph fluid leaves the node through efferent lymph vessels, which converge into lymphatic trunks, which in turn converge into one of the lymphatic ducts that flow lymph back into venous circulation.
- B and T lymphocytes must be transported to different sites within lymph nodes during an adaptive immune response.

Key Terms

- **afferent lymphatic vessels**: These vessels enter into the lymph nodes, flowing into the sinus space below the capsule of the node.
- **lymph**: A colorless, watery bodily fluid carried by the lymphatic system, consisting mainly of white blood cells.
- **germinal centers**: Places within secondary lymph nodes to which B cells migrate to proliferate and differentiate based on an antigen response.

Lymph transport refers to the transport of lymph fluid from the interstitial space inside the tissues of the body, through the lymph nodes, and into lymph ducts that return the fluid to venous circulation.

Transport in the Lymph Capillaries and Vessels

Lymphatic capillaries are the site of lymph fluid collection from the tissues. The fluid accumulates in the interstitial space inside tissues after leaking out through the cardiovascular capillaries. The fluid enters the lymphatic capillaries by leaking through the minivalves located in the junctions of the endothelium. Under ordinary conditions these minivalves prevent the lymph from flowing back into the tissues. In addition to interstitial fluid, pathogens, proteins, and tumor cells may also leak into the lymph capillaries and be transported through lymph.

The lymph capillaries feed into larger lymph vessels. The lymph vessels that receive lymph fluid from many capillaries are called collecting vessels. Semilunar valves work together with smooth muscle contractions and skeletal muscle pressure to slowly push the lymph fluid forward while the valves prevent backflow. The collecting vessels typically transport lymph fluid either into lymph nodes or lymph trunks.


Transport Within Lymph Nodes

Lymph circulates to the lymph node via afferent lymphatic vessels. The lymph fluid drains into the node just beneath the capsule of the node into its various sinus spaces. These spaces are loosely separated by walls, so lymph fluid flows around them throughout the lymph node.

The sinus space is filled with macrophages that engulf foreign particles and pathogens and filter the lymph. The sinuses converge at the hilum of the node, where lymph then leaves the node via an efferent lymphatic vessel toward either a more central lymph

node or a lymph duct for drainage into one of the subclavian veins.

The lymph nodes contain a large number of B and T lymphocytes, which are transported throughout the node during many components of the adaptive immune response. When a lymphocyte is presented with an antigen (such as by an activated helper T cell), B cells become activated and migrate to the germinal centers of the node, where they proliferate and differentiate to be specific to that antigen. When antibody-producing B cells are formed, they migrate to the medullary (central) cords of the node. Stimulation of the lymphocytes by antigens can accelerate the migration process to about ten times normal, resulting in the characteristic swelling of the lymph nodes that is a common symptom of many infections. The lymphocytes are transported through lymph fluid and leave the node through the efferent vessels to travel to other parts of the body to perform adaptive immune response functions.



Flow of Lymph : The lymph flows from the afferent vessels into the sinuses of the lymph node, and then out of the node through the efferent vessels.

The End of Lymphatic Transport

After leaving the lymph node through efferent vessels, lymph travels either to another node further into the body or to a lymph trunk, the larger vessel where many efferent vessels converge. Four pairs of lymph trunks are distributed laterally around the center of the body, along with an unpaired intestinal trunk.

The lymph trunks then converge into the two lymph ducts, the right lymph duct and the thoracic duct. These ducts take the lymph into the right and left subclavian veins, which flow into the vena cava. This is where lymph fluid reaches the end of its journey from the interstitial space of tissues back into blood circulation.

This page titled 19.2C: Lymph Transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2D: Lymphatic Capillaries

Lymph capillaries are tiny, thin-walled vessels, closed at one end and located in the spaces between cells throughout the body.

Learning Objectives

- Describe the location, structure, and role of lymphatic capillaries in maintaining the pressure of the interstitial fluid

Key Points

- Lymph or lymphatic capillaries are tiny thin-walled vessels, closed at one end and located in the spaces between cells throughout the body, except in the central nervous system and non-vascular tissues.
- Lymphatic capillaries are slightly larger in diameter and have greater oncotic pressure than blood capillaries.
- When pressure is greater in the interstitial fluid than in lymph, the minivalve cells separate slightly and interstitial fluid enters the lymphatic capillary. When pressure is greater inside the lymphatic capillary, the cells of the minivalves adhere more closely, and lymph cannot flow back into interstitial fluid.
- Anchoring filaments attach to the minivalves to anchor the capillary to connective tissue, and also pull the capillary open to increase lymph collection when the tissue is swollen.
- Because lymph capillaries have a closed end, lymph is pushed forward into larger vessels as the pressure inside the capillary increases as lymph accumulates from fluid collection.
- Edema can occur when interstitial fluid accumulation in tissues is greater than fluid removal (acute inflammation) or when the lymph vessels are obstructed in some way (elephantiasis).

Key Terms

- **interstitial fluid**: Also called tissue fluid, a solution that bathes and surrounds the cells of multicellular animals.
- **lymph capillaries**: Tiny thin-walled vessels, closed at one end and located in the spaces between cells throughout the body, collect fluid from the tissues.

Lymphatic circulation begins in the smallest type of lymph vessels, the lymph capillaries. These regulate the pressure of interstitial fluid by draining lymph from the tissues.

Structure of Lymphatic Capillaries

Lymph or lymphatic capillaries are tiny thin-walled vessels, closed at one end and located in the spaces between cells throughout the body. These are particularly dense within connective tissue. Lymphatic capillaries are slightly larger in diameter than blood capillaries and contain flap-like “minivalves” that permit interstitial fluid to flow into them but not out, under normal conditions.

Lymphatic capillaries are primarily made out of an endothelium layer that sits on a permeable basement membrane. The flap-like minivalves, located at gap-like junctions in the endothelium, are formed from the overlap of endothelial cells and are normally closed. Attached to the outer opening of the minivalves are anchoring filaments containing elastic fibers. They extend out from the lymphatic capillary, attaching the endothelium to fibroblast cells in the connective tissue. Unlike larger lymphatic vessels, lymphatic capillaries do not contain smooth muscle nor do they have a well developed adventitia, only small elastic filaments that perform a similar function.

Function of Lymphatic Capillaries

The lymph capillaries serve a variety of important functions.

Fluid Pressure Regulation


Lymphatic capillaries collect lymph fluid from the tissues, which allows them to regulate the pressure of interstitial fluid. This fluid is essentially plasma that leaks out of cardiovascular capillaries into the tissues due to the forces of hydrostatic or oncotic pressure. When pressure is greater in the interstitial fluid than in lymph due to accumulation of interstitial fluid, the minivalves separate slightly like the opening of a one-way swinging door so that fluid can enter the lymphatic capillary. When pressure is greater inside the lymphatic capillary, the cells adhere more closely to each other to prevent lymph backflow. The anchoring filaments are also pulled when the tissues are swollen. This opens the lymph capillaries more, increasing their volume and reducing their pressure to further facilitate fluid flow into the capillaries.

Lymph capillaries have a greater oncotic pressure (a pulling pressure exerted by proteins in solution) than blood plasma due to the greater concentration of plasma proteins in lymph. Additionally, the greater size of lymphatic capillaries compared to cardiovascular capillaries allows them to take more fluid proteins into lymph compared to plasma, which is the other reason for their greater levels of oncotic pressure. This also explains why lymph flows into the lymph capillaries easily, since fluid follows proteins that exert oncotic pressure.

Edema Prevention

Under normal conditions, lymph capillaries prevent the accumulation of edema (abnormal swelling) in the tissues. However, edema will still occur during acute inflammation or diseases in which lymph vessels are obstructed. During inflammation, fluid leaks into the tissues at a rate faster than it can be removed by the lymph capillaries due to the increased permeability of cardiovascular capillaries. During lymph vessel obstruction (such as through elephantiasis infection), lymph will be unable to progress normally through the lymphatic system, and pressure within the blocked off lymph capillaries increases to the point where backflow into tissues may occur, while the pressure of interstitial fluid gradually rises.

Drive Lymph Through Lymphatic Vessels



Lymph Capillary: Diagram showing the formation of lymph from interstitial fluid (labeled here as “tissue fluid”). Note: how the tissue fluid is entering the blind ends of lymph capillaries (indicated by deep green arrows).

The lymphatic capillaries bring lymph further into the lymphatic vessels. The capillaries have external valves but no internal valves or smooth muscle, so the pressure of lymph accumulation itself must propel the fluid forward into the larger vessels. Because lymphatic capillaries have a closed end and minivalves normally prevent backflow into tissues, the pressure of lymph becomes higher as more lymph is collected from the tissues, which sends the lymph fluid forward. Multiple capillaries converge in collecting vessels, where the internal valves and smooth muscle start to appear. This moves lymph further along the system despite the fall in pressure that occurs when moving from the higher-pressure capillaries to the lower-pressure collecting vessels.

This page titled 19.2D: Lymphatic Capillaries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.2E: Lymph Trunks and Ducts

The lymph trunks drain into the lymph ducts, which in turn return lymph to the blood by emptying into the respective subclavian veins.

Learning Objectives

- Describe the types of lymph vessels and lymph trunks and their roles

Key Points


- The lymph trunks drain into the lymph ducts, which in turn return lymph to the blood by emptying into the respective subclavian veins.
- There are two lymph ducts in the body: the right lymph duct and the thoracic duct.
- There are four pairs of lymph trunks: jugular lymph trunks, subclavian lymph trunks, bronchomediastinal lymph trunks, and lumbar lymph trunks. In addition, the intestinal lymph trunk is unpaired.
- The intestinal lymph trunk and the thoracic lymph duct contain chyle, a mixture of emulsified fats from the intestines and lymph fluid.

Key Terms

- **thoracic duct**: The lymph duct that drains lymph and chyle from the lower and left halves of the body.
- **subclavian vein**: Two large veins, one on either side of the body, with a diameter similar to that of the smallest finger.
- **lymph**: A colorless, watery body fluid carried by the lymphatic system, consisting mainly of white blood cells.

After filtration by the lymph nodes, efferent lymphatic vessels take lymph to the end of the lymphatic system. The final goal of the lymphatic system is to recirculate lymph back into the plasma of the bloodstream. There are two specialized lymphatic structures at the end of the lymphatic system, called the lymph trunks and ducts.

Lymphatic Trunks



Lymphatic Ducts: The thoracic duct and right lymphatic duct.

A lymphatic trunk is any large lymph vessel that forms from the convergence of many efferent lymph vessels. There are four sets of of lymph trunks that are paired with a right and left half, and one unpaired trunk:

- Jugular lymph trunks, located in the neck, drain lymph fluid from the cervical lymph nodes of the neck.
- Subclavian lymph trunks, located beneath the clavicle, drain lymph fluid from the apical lymph nodes around the armpit, which carry lymph from the arms.
- Bronchomediastinal lymph trunks, located in the chest, drain lymph fluid from the lungs, heart, trachea, mediastinal, and mammary glands.
- Lumbar lymph trunks are the lower pair of lymph trunks that drain lymph fluid from the legs, pelvic region, and kidneys.
- Intestinal lymph trunk is the unpaired lymph trunk that receives chyle (lymph mixed with fats) from the intestines. Chyle typically has a high fatty acid content.

Lymphatic trunks then drain lymph fluid into the lymph ducts, the final part of the lymphatic system.

Lymph Ducts

Two lymph ducts receive lymph from the lymph trunks. These are the largest lymph vessels and contain three layers, similar to those of great veins.

- The thoracic lymph duct, the largest lymph vessel in the body, takes lymph from the lower and left halves of the body. Because the thoracic lymph duct drains the intestinal lymph trunks, it carries a mixture of lymph and emulsified fatty acids called chyle back to the bloodstream.
- The right lymphatic duct receives lymph from the right and upper halves of the body, including the right sides of the jugular, bronchomediastinal, and subclavian lymph trunks.

The thoracic duct drains into to the left subclavian vein while the right duct drains into the right subclavian vein, both at the junction between the respective vein and the jugular vein. The two subclavian veins then merge into the vena cava, the large vein that brings deoxygenated blood to the heart. The lymph ducts each have internal valves at their junction with the subclavian vein. These function similarly to other lymphatic valves and prevent venous blood from flowing into the lymph duct. This point marks the end of lymph fluid's journey through the lymphatic system.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Lymphatic vessels. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphat..._of_Lymphatics**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endothelial cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/endothelial%20cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ISF. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/ISF**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adventitia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adventitia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphatic vessel. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphatic_vessel**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph flow. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...mph_vessel.png**. **License**: ***Public Domain: No Known Copyright***
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphatic_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymph**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plasma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph flow. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...mph_vessel.png**. **License**: ***Public Domain: No Known Copyright***

- Illu lymphatic system. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...tic_system.jpg. **License**: ***Public Domain: No Known Copyright***
- Lymphatic nodules. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphat...ic_circulation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lymph. **License**: ***CC BY-SA: Attribution-ShareAlike***
- afferent lymphatic vessels. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/afferen...atic%20vessels. **License**: ***CC BY-SA: Attribution-ShareAlike***
- subcapsular sinus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/subcapsular%20sinus. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph flow. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...mph_vessel.png. **License**: ***Public Domain: No Known Copyright***
- Illu lymphatic system. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...tic_system.jpg. **License**: ***Public Domain: No Known Copyright***
- Schematic_of_lymph_node_showing_lymph_sinuses.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_...ph_sinuses.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphatic capillaries. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphatic_capillaries. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph capillaries. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lymph%20capillaries. **License**: ***CC BY-SA: Attribution-ShareAlike***
- interstitial fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/interstitial_fluid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph flow. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...mph_vessel.png. **License**: ***Public Domain: No Known Copyright***
- Illu lymphatic system. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...tic_system.jpg. **License**: ***Public Domain: No Known Copyright***
- Schematic_of_lymph_node_showing_lymph_sinuses.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_...ph_sinuses.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu lymph capillary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Il..._capillary.png. **License**: ***Public Domain: No Known Copyright***
- Lymph duct. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_duct. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph vessel. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_vessel. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph trunk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_trunk. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/lymph. **License**: ***CC BY-SA: Attribution-ShareAlike***
- subclavian vein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/subclavian%20vein. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thoracic duct. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/thoracic%20duct. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph flow. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/00/Lymph_vessel.png. **License**: ***Public Domain: No Known Copyright***
- Illu lymphatic system. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...tic_system.jpg. **License**: ***Public Domain: No Known Copyright***
- Schematic_of_lymph_node_showing_lymph_sinuses.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymph_...ph_sinuses.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu lymph capillary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Il..._capillary.png. **License**: ***Public Domain: No Known Copyright***
- Gray599. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray599.png. **License**: ***Public Domain: No Known Copyright***

This page titled 19.2E: Lymph Trunks and Ducts is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.3: LYMPH CELLS AND TISSUES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

19.3A: Lymphoid Cells

19.3B: Lymphoid Tissue

19.3C: Lymph Nodes

This page titled 19.3: Lymph Cells and Tissues is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.3A: Lymphoid Cells

A lymphocyte is a type of white blood cell in the vertebrate immune system.

Learning Objectives

- List the types of lymphocytes in the vertebrate immune system and their roles in innate or humoral immunity

Key Points


- A lymphocyte is a type of white blood cell in the vertebrate immune system.
- NK cells are a part of the innate immune system and play a major role in defending the host from both tumors and virally infected cells.
- T cells are involved in cell-mediated immunity whereas B cells are primarily responsible for humoral immunity (relating to antibodies).
- Helper T-cells coordinate immune responses, while cytotoxic T-cells lyse (break down) pathogens associated with T cell's specific antigen.
- Memory B cells are formed at the end of an adaptive immune response and will produce antibodies more quickly when the antigen is detected again, which is effective at preventing recurrent infections from the same pathogen.
- Sometimes the body will present antigens that aren't harmful (allergy) or antigens from otherwise normally functioning body parts (autoimmunity). The latter can cause severe antibody and T-cell induced immune responses and diseases.

Key Terms

- **humoral immunity**: Immunity to infection due to antibodies that circulate in the blood and lymph and are produced by B cells.
- **antigen**: Any molecule that activates an immune response from a host organism, such as a toxin produced by bacteria or a molecule expressed on the cell wall of a virus -infected cell.
- **lymphocyte**: A type of white blood cell that includes T cells, B cells, and NK cells.

A lymphocyte is a type of white blood cell in the immune system. Lymphocytes develop from lymphoblasts (differentiated blood stem cells) within lymphoid tissue in organs such as the thymus. Lymphocytes are vital for normal immune system function. The three major types of lymphocyte are T cells, B cells, and natural killer cells.

Natural Killer Cells



Lymphocyte: A stained lymphocyte surrounded by red blood cells viewed using a light microscope.

Natural killer (NK) cells are part of the innate immune system and play a major role in defending the host from both tumors and virus-infected cells. NK cells contain receptors for a molecule called MHC (major histocompatibility complex) class I, which allows the NK cell to distinguish between infected cells and tumors from normal and uninfected cells. Normal cells express MHC class I on their cell membranes, while infected or cancerous cells do not express or express reduced amounts of the molecule. Therefore, the molecule acts as an inhibitor of NK cell activity, and NK cells activate and destroy cells on which MHC class I is not detected.

Activated NK cells release cytotoxic (cell-killing) granules that contain perforin and granzyme, which can lyse (break down) cell membranes and induce apoptosis to kill infected or abnormal cells. Cancer cells express much less MHC class I than normal cells, so NK cells are effective at destroying them before they develop into full tumors. If cancer cells evade NK cell detection for long enough, however, they can grow into tumors that are more resistant to NK cell activity.

T Cells and B Cells

T and B lymphocytes are the main forces of adaptive immunity, which includes cell-mediated and humoral immunity. T cells are involved in cell-mediated immunity whereas B cells are primarily responsible for humoral immunity. T cells and B cells irecognize specific “non-self” antigens during a process known as antigen presentation with MHC class II (usually done by dendritic cells). Once they have received an antigen, the cells become specifically tailored to eliminate and inhibit the pathogens or pathogen-infected cells that express that antigen. Sometimes these lymphocytes react to antigens that aren’t harmful (allergy) or will attack antigens expressed from the host’s own body (autoimmunity).

There are two types of T cells involved in adaptive, cell-mediated immunity.


- Helper T cells (CD4s) facilitate the organization of immune responses. They present antigens to B cells, produce cytokines that guide cytotoxic T cells, and activate macrophages.
- Cytotoxic T cells (CD8s) destroy pathogens associated with an antigen. Similar to NK cells, they bind to MHC class I and release granzymes, but will only bind to cells that express their specific antigen. Cytoxic T cells cause much of the damage associated with cell-mediated hypersensitivity, autoimmune disorders, and organ transplant rejection.

B cells are part of the humoral component of adaptive immunity. They respond to pathogens by producing large quantities of antigen-specific antibodies which neutralize foreign objects like bacteria and viruses, and opsonize (mark) them to be more easily recognized by other immune cells.

Following activation, B cells and T cells leave a lasting legacy of the antigens they have encountered in the form of memory

cells. Memory B cells are important for quickly producing antibodies should an antigen be recognized again, which can prevent recurrent infections from the same type of pathogen. This explains why vaccines are so effective, though viruses and bacteria with high mutation rates will express different antigens and thus avoid recognition by memory cells.

Development of Lymphocytes



SEM Lymphocyte: A scanning electron microscope (SEM) image of a single human lymphocyte.

All lymphocytes originate from a common lymphoid progenitor cell known as a lymphoblast, before differentiating into their distinct lymphocyte types. The formation of lymphocytes is known as lymphopoiesis. B cells mature into B lymphocytes in the bone marrow, while T cells migrate to and mature in thymus. Following maturation, the lymphocytes enter the circulation and peripheral lymphoid organs, where they survey for invading pathogens and cancer cells. The lymphocytes involved in adaptive immunity (B and T cells) differentiate further after exposure to an antigen, which occurs in the lymph nodes during antigen presentation from the dendritic cells. The fully differentiated B and T cells are specific to the presented antigen and work to defend the body against pathogens associated with that antigen.

This page titled 19.3A: Lymphoid Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.3B: Lymphoid Tissue

Lymphoid tissue consists of many organs that play a role in the production and maturation of lymphocytes in the immune response.

Learning Objectives

- Describe the structure, development, and location of lymphoid tissue

Key Points

- Lymphoid tissue may be primary or secondary depending upon its stage of lymphocyte development and maturation.
- The secondary lymphoid tissues consist of lymph nodes, tonsils, Peyer's patches, spleen, adenoids, skin, and mucosa-associated lymphoid tissue (MALT). They are responsible for maintaining mature naive lymphocytes and initiating an adaptive immune response.
- The thymus and bone marrow constitute the primary lymphoid tissues that are the sites of lymphocyte generation and maturation.
- Lymphoid tissue develops from venous endothelial tissues after the fifth week of gestation, starting by the end of the lymphatic system (subclavian vein and lymph ducts) and spreading outwards.

Key Terms

- **secondary lymphoid organ**: These organs maintain mature naive lymphocytes and initiate an adaptive immune response through antigen presentation.
- **primary lymphoid organ**: These organs generate lymphocytes from immature progenitor cells and provide an environment in which they mature.

The tissues of lymphoid organs are different than the tissues in most other organ systems in that they vary considerably based on cell cycle proliferation of lymphocytes. The lymphoid tissue may be primary or secondary depending upon its stage of lymphocyte development and maturation. Specialized lymphoid tissue supports proliferation and differentiation of lymphocytes.

Primary Lymphoid Organs


Central or primary lymphoid organs generate lymphocytes from immature progenitor cells such as lymphoblasts. The thymus gland and bone marrow contain primary lymphoid tissue where B and T cells are generated.

Besides generation, primary lymphoid tissue is the site where lymphocytes undergo the early stages of maturation. T cells mature in the thymus, while B cells mature in the bone marrow. T cells born in bone marrow travel to the thymus gland to mature.

Secondary Lymphoid Organs

Secondary or peripheral lymphoid organs maintain mature naive lymphocytes until an adaptive immune response is initiated. During antigen presentation, such as from the dendritic cells, lymphocytes migrate to germinal centers of the secondary lymphoid tissues, where they undergo clonal expansion and affinity maturation. Mature lymphocytes ill then recirculate between the blood and peripheral lymphoid organs until they encounter the specific antigens where they perform their immune response functions.

Secondary lymphoid tissue provides the environment for the antigens to interact with the lymphocytes. It is found mainly in the lymph nodes, but also in the lymphoid follicles in tonsils, Peyer's patches, spleen, adenoids, skin, and other areas associated with the mucosa-associated lymphoid tissue (MALT). In addition to supporting B and T lymphocyte activation, other secondary lymphoid organs perform other unique functions, such as the spleen's ability to filter blood and the tonsil's ability to capture antigens in the upper respiratory tract.



Lymphatic Tissues: The thymus and bone marrow are primary lymphoid tissue, while the lymph nodes, tonsils, and spleen are secondary lymphoid tissue.

Development of Lymphatic Tissue

Lymphatic tissue begins to develop by the end of the fifth week of embryonic development. Lymphatic vessels develop from lymph sacs that arise from developing veins, which are derived from mesoderm, the inner tissue layer of the embryo. Development of lymphatic tissue starts when venous endothelial tissues differentiate into lymphatic endothelial tissues. The lymphatic endothelial cells proliferate into sacs that eventually become lymph nodes, with afferent and efferent vessels that flow out from the lymph nodes. This process begins with he lymph nodes closest to the thoracic and right lymph ducts, which arises from immature subclavian-jugular vein junction. The lymph nodes organized around other lymph trunks, such as those in the abdomen and intestine, develop afterwards from nearby veins. Smaller lymph vessels and lymphatic capillaries develop after that until the lymphatic system is completed at the closed end of each lymphatic capillary.

More specialized primary lymph tissue, such as the thymus, develops from pharyngeal pouches (embryonic structures that differentiate into organs near the pharynx and throat) by the eighth week of gestation.

This page titled 19.3B: Lymphoid Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.3C: Lymph Nodes

Lymph nodes are small oval-shaped balls of lymphatic tissue distributed widely throughout the body and linked by lymphatic vessels.

Learning Objectives

- Describe the structure and function of lymph nodes

Key Points


- Lymph nodes are well-distributed around the chest, armpits, neck, and abdomen.
- Each lymph node is surrounded by a fibrous capsule that encircles the internal cortex and medulla. The cortex is mainly composed of clusters of B and T cells. The medulla contains plasma cells, macrophages, and B cells, as well as sinuses, which are vessel-like spaces that the lymph flows into, and nodules located within the sinuses.
- Lymph nodes contain a hilum beneath the capsule, which brings blood supply to the tissues of the lymph node.
- Antigen presentation by dendritic cells occurs in the lymph nodes, which triggers an adaptive immune response.
- Lymphadenopathy, the swelling of the lymph nodes, can indicate the presence of an infection or cancer.
- Lymph circulates to the lymph node via afferent lymphatic vessels and drains into the efferent lymphatic vessels just beneath the capsule.

Key Terms

- **lymphadenopathy**: Swelling of the lymph nodes that can indicate the presence of an infection or cancer.
- **lymph node**: Small oval bodies of the lymphatic system that act as filters, with an internal honeycomb of connective tissue filled with lymphocytes and macrophages that collect and destroy bacteria, viruses, and foreign matter from lymph.


Lymph nodes are small oval-shaped balls of lymphatic tissue, distributed widely throughout the body and linked by a vast network of lymphatic vessels. Lymph nodes are repositories of B cells, T cells, and other immune system cells, such as dendritic cells and macrophages. They act as filters for foreign particles in the body and are one of the sites where adaptive immune responses are triggered.

Structure of Lymph Nodes



Lymph node structure: This diagram of a lymph node shows the outer capsule, cortex, medulla, hilum, sinus, valve to prevent backflow, nodule, and afferent and efferent vessels.

Lymph nodes are found throughout the body, and are typically 1 to 2 centimeters long. Humans have approximately 500–600 lymph nodes, with clusters found in the underarms, groin, neck, chest, and abdomen. Each lymph node is surrounded by a fibrous capsule that encircles the internal cortex and medulla. The cortex is mainly composed of clusters of B cells in the outer layers and T cells in the inner layers, and may also contain antigen-presenting dendritic cells. The medulla contains plasma cells, macrophages, and B cells as well as sinuses, which are vessel-like spaces that the lymph flows into. Inside each sinus cavity is a nodule, a smaller, denser bundle of lymphoid tissue that usually contains a germinal center, the site of B cell proliferation during antigen presentation. The sinuses are partially divided by capsule tissue, which causes lymph fluid to flow around the nodules in each sinus cavity on their way through the node.



The lymphatic system: This diagram shows the network of lymph nodes and connecting lymphatic vessels in the human body.

Lymph fluid flows into and out of the lymph nodes via the lymphatic vessels, a network of valved vessels that are similar in structure to cardiovascular veins. Each lymph node has an afferent lymph vessel that directs lymph into the node, and an efferent lymph vessel called the hilum that directs lymph out of the node at the concave side of the node. The hilum also contains the blood supply of the lymph node.

Function of Lymph Nodes

Lymph nodes are the primary site for antigen presentation and activation in adaptive immune response in B and T lymphocytes. These lymphocytes are continuously recirculated through the lymph nodes and the bloodstream. Molecules called antigens are found on bacteria cell walls, the cell walls of virus-infected cells, or even chemical substances and toxins secreted from bacteria. These antigens may be taken by cells into the lymph nodes. There, antigen-presenting cells called dendritic cells present the antigen molecule to naive B and T lymphocytes. These undergo cell cycle proliferation into lymphocytes that are able to specifically detect and eliminate pathogens associated with that antigen, through various methods such as cytotoxic action (T cells) and antibody production (B cells).

The lymph nodes also filter the lymph fluid. Macrophages in the sinus spaces phagocytize (engulf) foreign particles such as pathogen, so that lymph fluid that returns to the bloodstream is cleaned of problematic abnormalities. The lymph node is also arranged in such a way that the chance of B and T lymphocytes encountering dendritic cells is quite high, to facilitate antigen presentation.

Lymphadenopathy

Lymphadenopathy describes the clinical condition of swollen lymph nodes. This is usually caused by increased lymph flow into

the nodes. This fluid may carry a higher amount of debris, so inflammation occurs as more neutrophils and later macrophages enter the node to remove debris from the lymph.

Lymphadenopathy is a symptom in conditions from trivial, such as a common cold or a minndor infection, to life-threatening, such as cancer or severe infection. Cancers that are severe and widespread from frequent metastases tend to have lymphadenopathy, so cancer staging criteria includes lymph node involvement. Additionally, cancers like lymphomas that have tumors made out of aberrant lymphocytes nearly always show lymphadenopathy, often an early warning sign for this type of cancer.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Lymphocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymphocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymphocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- humoral immunity. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/humoral_immunity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphocyte2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lymphocyte2.jpg**. **License**: ***Public Domain: No Known Copyright***
- SEM Lymphocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:SEM_Lymphocyte.jpg**. **License**: ***Public Domain: No Known Copyright***
- Lymphoid tissue. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphoi...ymphoid_tissue**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secondary lymphoid organ. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/seconda...mphoid%20organ**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- primary lymphoid organ. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/primary...mphoid%20organ**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph node. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymph_node**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphocyte2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lymphocyte2.jpg**. **License**: ***Public Domain: No Known Copyright***
- SEM Lymphocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:SEM_Lymphocyte.jpg**. **License**: ***Public Domain: No Known Copyright***
- Blausen_0623_LymphaticSystem_Female.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphatic_system#/media/File:Blausen_0623_LymphaticSystem_Female.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymph node. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymph_node**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymphocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymphocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymphadenopathy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymphadenopathy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lymph node. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lymph_node**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphocyte2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lymphocyte2.jpg**. **License**: ***Public Domain: No Known Copyright***
- SEM Lymphocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:SEM_Lymphocyte.jpg**. **License**: ***Public Domain: No Known Copyright***
- Blausen_0623_LymphaticSystem_Female.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphatic_system#/media/File:Blausen_0623_LymphaticSystem_Female.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphatic system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ly...tic_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Illu lymph node structure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il..._structure.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 19.3C: Lymph Nodes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.4: OTHER LYMPHOID ORGANS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

19.4A: Thymus
19.4B: Spleen
19.4C: Tonsils

This page titled 19.4: Other Lymphoid Organs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.4A: Thymus

The thymus is a specialized organ that “educates” T cells or T lymphocytes, which are part of the adaptive immune system.

Learning Objectives

- Describe the, structure, location, and role of the thymus in the immune system

Key Points

- Each T cell is specialized to attack a different antigen, but those that attack self-antigens are destroyed by the thymus during selection processes in lymphocyte proliferation and maturation.
- The organ enlarges during childhood, atrophies at puberty, and is generally replaced with fat. Residual T lymphopoiesis continues through adulthood.
- The thymus is composed of two identical lobes in the upper chest. Its main components are the the cortex, which is the site of lymphocyte maturation, and the medulla, which connects the thymus to venous circulation.
- One of the most important roles of the thymus is the maintenance of central tolerance, which works to prevent autoimmune disorders from occurring.
- T-cells are generated in the bone marrow and travel to the thymus in order to mature.

Key Terms

- **thymus**: A ductless gland consisting mainly of primary lymphatic tissue. It plays an important role in the development of the immune system and produces lymphocytes.
- **Central tolerance**: The ability for T-cells to avoid perceiving normal host molecules as foreign antigens.
- **T lymphocytes**: These cells, also called T cells, belong to a group of white blood cells known as lymphocytes and play a central role in cell-mediated immunity.

EXAMPLES

Thymectomy, the surgical removal of the thymus, is done most often to gain access to the heart in surgeries to correct congenital heart defects performed in the neonatal period. In neonates, the relative size of the thymus obstructs surgical access to the heart. Surprisingly, removal of the thymus does not result in a T cell immunodeficiency. This is because sufficient T cells are generated during fetal life prior to birth. These T cells are long-lived and can proliferate by homeostatic proliferation throughout the patient’s lifetime. However, there is evidence of premature immune aging in patients thymectomized during early childhood.

The thymus is a specialized organ of the immune system. It consists of primary lymphoid tissue, which provides a site for the generation and maturation of T lymphocytes, critical cells of the adaptive immune system.

Structure of the Thymus


The thymus is of a pinkish-gray color, soft, and lobulated on its surfaces. The organ enlarges during childhood into adolescence and begins to atrophy at puberty due to hormonal changes. After puberty, the thymus shrinks rapidly with age, eventually becoming almost indistinguishable from the surrounding fatty tissue.

The thymus consists of two lateral lobes placed in close contact along the middle line situated partly in the thorax, resting in the chest beneath the neck. The two lobes differ slightly in size, may be united or separated, and may be broken down into smaller lobules. It is covered with a capsule of connective tissue that provides structural support.

Histologically, the thymus contains mature lymphocytes, immature lymphocytes, and stroma, while lobule tissues consist of an inner medulla and an outer cortex. The cortex and medulla play different roles in the development of T cells. The cortex is the

site of T cell generation and proliferation, while the medulla connects to the venous bloodstream and allows for transport of mature inactive T cells to the lymph nodes and transport of immature T cells from bone marrow tissue into the thymus cortex for proliferation and maturation.

Function of the Thymus



Thymus: The thymus is the site of T-cell generation and maturation.

The thymus provides an environment for T cells to mature and proliferate, a process called lymphopoeisis. First, immature T cells generated in bone marrow travel to the cortex tissues of the thymus through the bloodstream. Then the immature T cells undergo proliferative expansion, in which they are exposed to growth factors and antigen receptors are formed. Then the T cells are sorted by the thymus so that only T cells that express T-cell receptors (TcRs) and can bind to foreign MHC molecules will survive. The surviving cells will not mistake self-molecules for antigens. Only 2-4% of T cells survive this sorting process. The thymus is most active early in life for building a large reservoir of T cells. Though removal of the thymus in childhood causes severe immunodeficiency, later in life this is not an issue because of the proliferation of thymus activity early in life.

Central tolerance is another function of the thymus. Autoimmune diseases occur when central tolerance is lost, which causes lymphocytes to recognize host molecules as antigens and attack them, even if those tissues otherwise function normally. The thymus sorts T cells so that they will be inactive towards host molecules, though sometimes a few T cells evade this sorting process and may initiate an autoimmune disease. Though the thymus is mostly effective at preventing this occurrence, those with certain genetic characteristics (such as altered MHC complexes) may be more likely to develop autoimmune diseases with the few T-cells that aren't properly selected by the thymus.

This page titled 19.4A: Thymus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.4B: Spleen

The spleen, similar to a large lymph node, acts primarily as a blood filter in the mononuclear phagocyte system of the immune system.

Learning Objectives

- Describe the role of the spleen in the filtration and storage of red blood cells

Key Points

- The primary function of the spleen is to filter old blood cells out of the bloodstream.
- Structurally, the spleen is similar to a massive lymph node; however, it filters blood instead of lymph and therefore contains only efferent lymph vessels.
- Red pulp mechanically filters old blood using macrophage activity. White pulp is responsible for active immune response by synthesizing antibodies.
- The spleen removes some bacteria from the bloodstream, particularly those that cause pneumonia.
- The spleen holds extra blood that can help during hypovolemic shock.
- Survival is possible with removal of the spleen because the lymph nodes and liver can perform most of the same functions.

Key Terms

- **spleen**: A ductless vascular gland that destroys old red blood cells, removes debris from the bloodstream, acts as a reservoir of blood, and produces lymphocytes.
- **white pulp**: The part of the spleen where lymphocytes are maintained in a similar way as in lymph nodes.
- **red pulp**: The site of blood filtration in the spleen.

EXAMPLES

Survival is possible without a spleen. However, retrospective epidemiological studies of World War II veterans found that those who had their spleens removed on the battlefield showed significant mortality risk from pneumonia and a significant excess of mortality from ischemic heart disease, but not from other conditions.

The spleen is the largest distinct organ of the lymphatic system. Similar in structure to a large lymph node, it acts primarily as a blood filter. Despite this important function, a healthy life is possible after removal. The spleen plays important roles in regards to red blood cells and the immune system.


Structure of the Spleen

The spleen is located in the left upper quadrant of the abdomen. It is similar to an enlarged lymph node but is a bit more complex. The spleen is made up of two distinct tissue types:

- Red pulp is the site of blood filtration in the spleen. It is made of connective tissue called the cord of Billroth that can fill with blood and contains many macrophages.
- White pulp is secondary lymphoid tissue that is similar to that in the adenoid tonsils. They contain large amounts of lymphocytes and antigen -presenting cells.

Unlike lymph nodes, the spleen possesses only efferent lymphatic vessels, because it only filters blood instead of lymph fluid. The splenic artery forms its primary blood supply. The spleen is unique in respect to its development within the gut because it is derived from mesenchymal tissue rather than endoderm tissue during embryonic development. However, it still shares the same blood supply as the foregut organs in the abdominal cavity.

Function of the Spleen



Spleen: This diagram of the spleen indicates the vein, artery, white pulp, red pulp, and capsule.

The primary function of the spleen is blood filtration. Blood cells have a lifespan of roughly 120 days. When blood passes through the red pulp of the spleen, healthy blood cells easily pass, while older red blood cells are caught phagocytized by the macrophages within. The macrophages also remove pathogens, denatured hemoglobin, and other cellular debris. Iron from old or damaged hemoglobin content in the blood is filtered out and sent to the liver so that new red blood cells can be created. Antigens are also filtered by the red pulp, which may be presented to naive lymphocytes in the white pulp of the spleen. This stimulates the same type of adaptive immune response that occurs in the lymph nodes.

The spleen is also important for generating new red blood cells early in embryonic development, but this function stops after birth. The spleen may also function as a reservoir of blood and platelets during hypovolemic shock, which occurs when overall tissue perfusion falls due to severe dehydration or severe bleeding or hemorrhage. During hypovolemic shock, the spleen can release up to a cup of extra blood to help mitigate the complications of fluid loss.

The spleen is often removed surgically if it becomes damaged or infected. This causes modest increases in circulating white blood cells and platelets, diminished responsiveness to some vaccines, and increased susceptibility to infection by bacteria and protozoa. In particular, there is an increased risk to infection from gram negative bacteria that cause pneumonia. Besides these increased risks, the loss of the spleen does not cause major immune system impairment, and most people will still live normal and healthy lives because the lymph nodes and liver perform the same functions as the spleen. In particular, those with splenomegaly (an enlarged spleen that could rupture) or splenic cancers are typically better off living without their spleen than living with the risk of severe bleeding from a ruptured spleen or the plethora of symptoms caused by splenic cancer and its metastases.

This page titled 19.4B: Spleen is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

19.4C: Tonsils

The tonsils are one of the immune system's first lines of defense against ingested or inhaled foreign pathogens.

Learning Objectives

- Describe the role of tonsils in the immune system

Key Points

- The four sets of tonsils are the adenoids, palatine tonsils, tubal tonsils,and the lingual tonsils.
- Tonsils consist of epithelial tissue with narrow folds called crypts, secondary lymphoid tissue that contains lymphocytes, and M cells that capture antigens in the respiratory tract.
- Tonsils tend to reach their largest size near puberty and gradually atrophy thereafter.
- Tonsils can become enlarged or inflamed and may be surgically removed in tonsillectomy. Chronic inflammation of the tonsils can cause their cells to increase in size (hypertrophy).
- Enlarged tonsils can make breathing more difficult and disrupt mucus drainage in the pharynx.
- Those with a tonsillectomy show no significant long term difference in immune system function, though minor changes in immune cell and antibody levels do occur.

Key Terms

- **tonsillitis**: Inflammation of the tonsils.
- **tonsillectomy**: The surgical removal of the tonsils, especially the palatine tonsils. Frequently accompanied by an adenoidectomy.
- **squamous cell carcinoma**: a cancer of squamous cell epithelial tissue.
- **tonsils**: Paired masses of secondary lymphoid tissue and epithelial tissue found in the pharynx.

The tonsils are small masses of secondary lymphoid tissue located in the pharynx. They function similarly to other types of secondary lymphoid organs and also capture antigens from respiratory tract pathogens.


Location and Structure of the Tonsils

There are four pairs of tonsils located within the pharynx. They function similarly but have a few structural differences.

- The adenoids are located in the wall of nasopharynx.
- The palatine tonsils are located in the sides of the oropharynx.
- The tubal tonsils are located in the wall of the nasopharynx near the entrance to each Eustachian tube.
- The lingual tonsils are located behind the tongue.

The tonsils are made of secondary lymphoid tissue and covered with an epithelium characteristic of the part of the body where they are located. For example, the adenoids and tubal tonsils are covered with the ciliated psuedostratified columnar epithelium of the nasopharynx, while the palatine and lingual tonsils are made up of the non-keratinized stratified squamous epithelium of the oropharynx. The tonsils also contain very deep and narrow folds in their tissues called crypts. Like the thymus, the tonsils reach their largest size near puberty and gradually atrophy thereafter.

Function of the Tonsils



Tonsils: Palatine tonsils can be seen on the left and right sides at the back of the throat.

The tonsils primarily facilitate adaptive immune responses in the upper respiratory tract, one of the most common pathways for pathogen entry in the body. In a way, the tonsils are the "first line of defense" against potential respiratory pathogens. They contain specialized M cells that collect antigens produced by respiratory tract pathogens. The secondary lymphoid tissue within the tonsils functions like the same type of the tissue in lymph nodes. Captured antigens are presented to B and T cells within the tonsil, then the B cells migrate to germinal centers within the tonsil as an adaptive immune response is initiated. Additionally, evidence exists that suggests that tonsils may play a role in T cell maturation and development like the thymus does, but more research is needed.

Tonsil removal (tonsillectomy) is a common procedure to treat swollen and infected lymph nodes (tonsillitis). It does not appear to cause weakened immune function. Chronic infection of the adenoids can cause adenoid hypertrophy, increases in cell size from repeated damage. Enlarged tonsils can make it more difficult to breath and disrupt normal mucus drainage in the pharynx, so removal is generally recommended in those cases. Squamous cell carcinomas (epithelial tumor) and lymphomas (lymphocyte tumor) can also develop in the tonsillar tissue, and removal is a key treatment. Epidemiological studies show no significant change in immune system function in those that have a tonsillectomy, but minor increases in helper T cell levels and minor decreases in IgA levels (an antibody produced by B cells) were observed.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Thymus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thymus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T lymphocytes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/T%20lymphocytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thymus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thymus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thymocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thymocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu thymus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_thymus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spleen. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spleen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- spleen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/spleen**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- red pulp. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/red%20pulp**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- white pulp. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/white%20pulp**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu thymus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_thymus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikipedia. **Located at**: **Wikipedia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tonsillectomy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tonsillectomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tonsils. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Tonsils**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tonsils. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tonsils**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tonsillitis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tonsillitis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Squamous-cell carcinoma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Squamous-cell_carcinoma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu thymus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_thymus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikipedia. **Located at**: **Wikipedia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tonsils. **Provided by**: Wikipedia. **Located at**: **Wikipedia**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 19.4C: Tonsils is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20: IMMUNE SYSTEM

20.1: Innate Immunity

20.1A: Skin and Mucosae (Surface Barriers)

20.1B: Phagocytes

20.1C: Natural Killer Cells

20.1D: Inflammation

20.1E: Antimicrobial Proteins

20.1F: Fever

20.2: Stress and Immunity

20.2A: Stress and Immunity

20.3: Adaptive Immunity

20.3A: Overview of Adaptive Immunity

20.3B: Types of Adaptive Immunity

20.3C: Maturation of T Cells

20.3D: Lymphocytes

20.3E: Antigen-Presenting Cells

20.4: Cytokines

20.4A: Types of Cytokines Participating in Immune Response

20.5: Antigens

20.5A: Antigens and Antigen Receptors

20.5B: Complete Antigens and Haptens

20.5C: Antigenic Determinants and Processing Pathways

20.6: Humoral Immune Response

20.6A: Clonal Selection and B-Cell Differentiation

20.6B: Structure and Function of Antibodies

20.6C: Role of the Complement System in Immunity

20.6D: Immunological Memory

20.6E: Major Histocompatibility Complex Antigens (Self-Antigens)

20.7: Cell-Mediated Immune Response

20.7A: Clonal Selection and T-Cell Differentiation

20.7B: Specific T-Cell Roles

20.7C: Active and Passive Humoral Immunity

This page titled 20: Immune System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1: INNATE IMMUNITY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

20.1A: Skin and Mucosae (Surface Barriers)

20.1B: Phagocytes

20.1C: Natural Killer Cells

20.1D: Inflammation

20.1E: Antimicrobial Proteins

20.1F: Fever

This page titled 20.1: Innate Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1A: Skin and Mucosae (Surface Barriers)

In mammals, the skin and mucosae constitute complex protective barriers that guard against infection and injury.

Learning Objectives

- Describe how the skin and mucosae serve as a protective barrier which guards against infection and injury

Key Points

- The skin consists of the epidermis, dermis, and basement membrane.
- The epidermis is the outermost layer of skin and forms a protective barrier over the body's surface. Its layers continually grow outward as older layers shed away.
- The dermis is below the epidermis and is made of connective tissue that cushions the body from stress and strain. It contains mechanoreceptors, blood vessels, and sweat glands.
- The mucosae are linings covered in epithelium and involved in absorption and secretion. They line cavities that are exposed to the external environment and internal organs.
- The barrier immune system is part of the innate immune system, and consists of anything that the skin, mucosae, and chemical secretions of the body do to prevent pathogens from invading.
- The barrier system can fail when the skin breaks or when pathogens invade the mucosal epithelium, so other innate and adaptive immune system functions exist to destroy pathogens in these cases.

Key Terms

- **mucous membranes**: Specialized epithelium for internal and semi-internal structures that usually secretes mucus and provides some barrier immune system function.
- **barrier immune system**: A component of the innate immune system that refers to the physical and chemical barriers that prevent pathogens from entering and infecting the body.

The skin is one of the most important body parts because it interfaces with the environment and is the first line of defense from external factors. While it performs a wide range of functions, including sensation, heat regulation, control of evaporation, storage, synthesis, absorption, and water resistance, but its innate immune system functions as the barrier immune system are the most critical.

In humans, the outer covering of the body consists of the skin and mucosae, which together make up the barrier immune system.

Components of the Skin

The skin is made up of several layers that together protect the body, regulate temperature, keep water inside the body, and have sensory function. The skin is the largest organ in the body, and consists of three components that differ greatly in structure and function:

- The epidermis comprises the outermost layers of the skin. It forms a protective barrier over the body's surface, responsible for keeping water in the body and preventing pathogens from entering. It is made of stratified squamous epithelium tissues, composed of proliferating basal and differentiated suprabasal keratinocytes that form an extracellular matrix that continually divides as the older outer layers of the epidermis shed. The epidermis also helps the skin regulate body temperature through sweat pores that connect to underlying sweat glands in the dermis.
- The basement membrane is a thin sheet of fibers called the basement membrane that separates the dermis and epidermis. It controls the traffic of cells and molecules between the dermis and epidermis and the release of cytokines and growth factors during wound healing.
- The dermis is the layer of skin beneath the epidermis that consists of connective tissue and cushions the body from stress and strain. The dermis provides strength and elasticity to the skin through an extracellular matrix composed of

collagen fibrils, microfibrils, and elastic fibers, embedded in proteoglycans. It harbors many mechanoreceptors (sensory nerve endings) that provide the sensations of touch and heat. It also contains the hair follicles, sweat glands, sebaceous glands, apocrine glands, lymphatic vessels, and blood vessels. The blood vessels in the dermis provide nourishment and waste removal for their own cells and for the epidermis.

Mucous Membranes

The mucous membranes (or mucosae; singular mucosa) are linings of mostly endodermal origin, covered in various types of epithelium, that are involved in absorption and secretion. They line cavities that are exposed to the external environment and internal organs. They attached to skin at the nostrils, mouth, lips, eyelids, and genital area, but are also located within the body cavities, such as in the stomach, anus, trachea, and ears.


Most mucous membranes secrete a sticky, thick fluid called mucus, which facilitates several barrier immune system functions and provides a moist environment for internal and semi-internal structures. The mucosae are highly specialized in each organ to deal with different conditions. The most variation is seen in the epithelium. In the esophagus and oropharynx, the epithelium is stratified, squamous and non-keratinising, to protect these areas from harsh or acidic foods. In the stomach it is columnar and organized into gastric pits and glands to secrete acids and pepsin. The small intestine epithelium is specialized for absorption, organized into simple columnar epithelium on protruding villi with narrow crypts that have a high surface area. The mucosal epithelium in the nasopharynx is psuedostratified and ciliated, which helps accumulate and remove mucus.

The Barrier Immune System

Together, the skin and mucosae form the the barrier immune system, technically considered a component of the innate immune system. These structures form physical barriers to infection that prevent pathogens from entering the body through a variety of methods. While the skin simply prevents pathogen entry, more specialized structures like the mucociliary escalator in the trachea trap pathogens in mucus secretions and use cilia to push them out of the trachea to prevent entry into the lungs.

The barrier system also includes chemical barriers that prevent pathogen entry. Notable examples include stomach acidity which kills most microbes, antimicrobial peptides on mucosal epithelial tissue, and even the flow of urine that flushes pathogens out of the urethra.

The barrier system is the first line of defense against pathogen invasion, though it is not perfect. The skin can be broken through cuts or abrasions that expose the bloodstream to the environment. Not every pathogen is caught nor inhibited in mucus, and some may infect the mucosal epithelium directly. Smoking and alcohol abuse damage the muciliary escalator and make it easier for pathogens to invade the lungs. Fortunately, other mechanisms of the innate and adaptive immune systems defend the body when the barrier system fails.



Dermis: A diagrammatic view of a skin section.

This page titled 20.1A: Skin and Mucosae (Surface Barriers) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1B: Phagocytes

Phagocytes are the white blood cells that protect the body by ingesting harmful foreign particles and help initiate an immune response.

Learning Objectives

- Describe the types of phagocytes and their roles in initiating an immune response

Key Points

- Many white blood cells and other cells in the body use phagocytosis to engulf and kill cells.
- Phagocytosis occurs over several steps, which include binding to an opsonized pathogen with a receptor and killing it using an oxidative burst.
- Monocytes are phagocytes that can differentiate into macrophages and dendritic cells based on conditions within the body.
- Macrophages are the cleanup crew for the innate immune system. They remove debris, pathogens, and dead neutrophils after an inflammatory response.
- Neutrophils are polymorphonuclear (PMN) granulocytes that are the first responders to an inflammatory response. They kill pathogens through phagocytosis and degranulation, but die as a result.
- Mast cells are circulation PMN granulocytes that trigger an immune response by releasing an inflammatory mediator when they detect an antigen with their toll-like receptors.

Key Terms

- **oxidative burst**: A chemical reaction that occurs in phagocytes in which an engulfed pathogen is destroyed by exposure to oxidative stress from reactive oxygen species.
- **PMN granulocyte**: A type of phagocyte that contains PMN granules, most notably neutrophils and mast cells, but also basophiles and eosinophils.

Any cell that undergoes phagocytosis, a process in which pathogens and other foreign particles and debris are engulfed by a cell to be destroyed, is considered a phagocyte. Most phagocytes are types of white blood cells that use phagocytosis to perform basic innate immune system function within the body.

The Mechanism of Phagocytosis

Phagocytosis is the process by which a phagocyte engulfs a pathogen or debris. It can occur in almost any tissue, most often in the bloodstream and interstitial space but also the alveoli of the lungs and the parenchyma of most other major organs in the body. Typical phagocytosis occurs over the course of a few steps:

1. A receptor on the phagocyte's cell membrane binds to a foreign particle, such as a pathogenic microbe or a toxin. The Fc receptor is typically the receptor of use, which binds to antibodies that have opsonized (marked) a pathogen or toxin.
2. The cytoplasm surrounds and engulfs the bound pathogen through endocytosis.
3. The engulfed pathogen is kept in a vacuole called a phagosome, which then binds to the lysosomes inside the cell.
4. A series of chemical reactions called an oxidative burst occurs, which uses reactive oxygen species and NADPH oxidase to damage and kill the pathogen through oxidative stress. Oxidative stress can kill a cell through DNA, cell membrane, or mitochondrial damage.
5. The remains of the pathogen are expelled by exocytosis.

These are the general mechanisms used by phagocytosis to engulf and kill pathogens, but some variations can occur. For instance, other receptors may be used to engulf pathogens, and other non-oxidative methods (such as lysozyme) exist to kill the phagocytized pathogen.

Types of Phagocytes

There are many classes of phagocytes within the body, each with different specialized functions involving phagocytosis. Most phagocytes are derived from stem cells in the bone marrow. The main types of phagocytes are monocytes, macrophages, neutrophils, tissue dendritic cells, and mast cells. Other cells, such as epithelial cells and fibroblasts, may also engage in phagocytosis, but lack receptors to detect opsonized pathogens and are not primarily immune system cells.


Monocytes

Monocytes develop in the bone marrow and reach maturity in the blood. Mature monocytes have large, smooth, lobed nuclei and an abundant cytoplasm that contains granules, but are not technically considered granulocytes. Monocytes ingest foreign or dangerous substances and present antigens to other cells of the immune system. Monocytes form two groups: a circulating group and a marginal group that remains in other tissues (approximately 70% are in the marginal group). Most monocytes leave the blood stream after 20–40 hours to travel to tissues and organs; during this process, they differentiate into macrophages or dendritic cells depending on the signals they receive.

Macrophages

Mature macrophages are derived from monocytes, granulocyte stem cells, or the cell division of pre-existing macrophages. Macrophages do not have granules, but contain many lysosomes. They are found throughout the body in almost all tissues and organs, but are rarely found in the bloodstream. Macrophages cause inflammation through the production of interleukin-1, interleukin-6, and TNF-alpha. Macrophages are activated in a number of ways, including by T cells, cytokines such as IFN-gamma, or pathogen-derived compounds such as LPS toxins from bacteria. During inflammation, they enter about 72 hours after the initial response to clean up debris and dead neutrophils.

Dendritic Cells




Dendritic cell: Dendritic cell characterized by membranous projections that resemble spines.

Dendritic cells are specialized antigen-presenting cells that have long outgrowths called dendrites, which help to engulf microbes and other invaders. They express MHC class II molecules, which makes them the ideal antigen-presenting cell. Dendritic cells are present in the tissues that are in contact with the external environment, mainly the skin, the inner lining of the nose, the lungs, the stomach, and the intestines.

Once activated, they mature and migrate to the lymphoid tissues, where they present antigens to T and B cells to initiate the adaptive immune response. This involves deriving T and B cells that are specific towards a single antigen from naive lymphocytes.

Neutrophils


Neutrophils are a type of PMN granulocyte normally found in the bloodstream. They are the most abundant type of phagocyte and the first responder during inflammation. Once they have received the appropriate chemokine signals, neutrophils leave the bloodstream and reach the site of an infection through adhering to the vascular endothelium to squeeze into the tissues. There, they rapidly engulf invaders coated with antibodies, damaged cells, or cellular debris. They also degranulate to release perforin, granzyme, proteases, and other chemicals to cause cytotoxic damage to pathogens (and occasionally normal bodily tissues as well). Neutrophils die after phagocytosis, becoming pus that is later cleaned up by macrophages.



Extravasion of Neutrophils: Neutrophils move through the blood to the site of infection by rolling onto the vascular endothelium and adhering to it to slip through small gaps into the tissues during an inflammatory response.

Mast Cells

Mast cells are PMN granulocytes with toll-like receptors that tend to trigger inflammatory responses. Mast cells express MHC class II molecules and can participate in antigen presentation; however, the mast cell's role in antigen presentation is not well-understood. Mast cells can consume, kill, and process their antigens. In addition to these functions, mast cells produce cytokines kept in their granules, such as histamine, that induce an inflammatory response when a pathogen is detected. Because of this function, allergic inflammatory responses occur when a mast cell is sensitized to an antigen that it normally wouldn't react to.



Leukocyte Differentiation: Phagocytes derive from stem cells in the bone marrow. Monocytes differentiate into dendritic cells and macrophages, while mast cells and neutrophils are in a separate group of PMN granuolcytes as well.

This page titled 20.1B: Phagocytes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1C: Natural Killer Cells

Natural killer (NK) cells are cytotoxic lymphocytes critical for the innate immune system.

Learning Objectives

- Describe the role of natural killer cells in the innate immune system

Key Points

- NK cells differentiate from lymphocyte progenitor cells and are a critical part of the innate immune system.
- NK cells recognize abnormal or infected cells with activating receptors and inhibitory receptors.
- All normal cells in the body express MHC I to signal that those cells are part of the body.
- Inhibitory receptors recognize MHC class I alleles, which inhibits the NK killing response and explains why NK cells will kill cells possessing few or no MHC class I molecules.
- Activating receptors recognize antigens, antibodies, or other opsonins on a cell's surface and activate a killing response.
- NK cells are cytotoxic; small granules in their cytoplasm contain proteins such as perforin and proteases known as granzymes that trigger either apoptosis or cell lysis in an abnormal cell.
- NK cells may release two potent immune system cytokines, IFN-gamma and TNF-alpha, when certain receptors are activated.

Key Terms

- **Major Histocompatability Complex I**: A molecule expressed on cells to signal to immune system cells that they are normal cells of that organism's body. Abbreviated as MHC I.
- **apoptosis**: A response in which a cell undergoes programmed cell death and its DNA and other components are destroyed completely. It is a mechanism to stop viral infections and cancer development and is a result of cellular stress.

Natural killer (NK) cells are cytotoxic lymphocytes critical to the innate immune system. The role of NK cells is similar to that of cytotoxic T cells in the adaptive immune response. NK cells provide rapid responses to virus-infected cells and respond to tumor formation by destroying abnormal and infected cells. NK cells use wo cytolytic granule-mediated apoptosis to destroy abnormal and infected cells.

Natural Killer Overview

Typically, immune cells detect major histocompatibility complex (MHC) presented on cell surfaces, triggering cytokine release and lysis or apoptosis in cells that do not express MHC I or express much less of it than normal cells. Unlike phagocytes, NK cells do not need their targets to be opsonized (marked) by antibodies before they can act, allowing for a much faster immune reaction. However, opsonins do speed up the process.

These cells named "natural killers" because they were thought to work without cytokine or chemokine activation. However, later research proved that cytokines play a role in guiding NK cells to stressed cells that may need to be destroyed.


NK cells are large granular lymphocytes derived from the common lymphoid progenitor cells (lymphoblasts), which also generate B and T lymphocytes. NK cells differentiate and mature in the bone marrow, lymph nodes, spleen, tonsils and thymus, where they then enter into the bloodstream.

MHC I Recognition

In order for NK cells to defend the body against viruses and pathogens, they require mechanisms to determine whether a cell is infected. The exact mechanisms remain the subject of current investigation, but recognition of an "altered self" state is thought to be involved. To control their cytotoxic activity, NK cells possess two types of surface receptors: activating receptors and inhibitory

receptors. Most of these receptors are also present in certain T cells. These receptors recognize major histocompatability complex I (MHC I), a molecule expressed on every cell to signal that the cell belongs to the body.

When the NK cell recognizes MHC I on a cell using an inhibitory receptor, its killing response is inhibited. When the NK cell does not recognize MHC I on the cell with an inhibitory receptor, or detects an antigen with an activating receptor, the killing response is activated. Virus-infected cells and foreign pathogens such as bacteria and fungi will not express the MHC I specific to the host organism, which will fail to inhibit the NK cell's killing responses. If both types of receptors are being stimulated, the receptor that experiences a higher degree of relative stimulation will determine the NK cell behavior. Some tumor cells may still express MHC I in low amounts, so they may evade NK cell destruction based on the balance of activating and inhibiting stimuli.



Complementary Activities of Cytotoxic T-cells and NK cells: Schematic diagram indicating the complementary activities of cytotoxic T-cells and NK cells. T-cells are activated by recognizing antigens, while NK cells are activated by not recognizing MHC I.

Mechanisms of Cytotoxicity

The granules of NK cells contain proteins such as perforin and proteases known as granzymes. Upon binding to a cell slated for killing, perforin forms pores in the cell membrane of the target cell, creating an aqueous channel through which the granzymes and associated molecules can enter, inducing either apoptosis or osmotic cell lysis (a form of cell necrosis). Defensins, an antimicrobial secreted by NK cells, directly kills bacteria by disrupting its cell walls.

Apoptosis is a form of "programmed cell death" in which the cell is stimulated by the cytotoxic mechanisms to destroy itself. Unlike with lysis, apoptosis does not degrade DNA, and cells are destroyed cleanly and completely on their own. Cellular lysis causes necrosis of that cell, in which the DNA and cell components degrade into debris that must be phagocytized by macrophages. This distinction has many important implications. Virus-infected cells destroyed by cell lysis release their replicated virus particles into the body, which infects other cells. In apoptosis, these virus particles are destroyed. However, cancer cells often develop genetic mechanisms to prevent apoptosis signals from occurring, so cell lysis is generally more effective.

Cells that are osponized with antibodies are easier for NK cells to detect and destroy. Antibodies that bind to antigens can be recognized by FcγRIII (CD16) receptors (a type of activating receptor), resulting in NK activation, release of cytolytic granules, and consequent cell apoptosis.

Cytokines in NK Cell Activity

Cytokines play a role in NK cell activation. Many cells release cytokines as a result of cellular stress when infected with a virus. Cytokines involved in NK activation include IL-12, IL-15, IL-18, IL-2, and CCL5. NK cells are activated in response to interferons or macrophage-derived cytokines. They serve to contain viral infections while the adaptive immune response generates antigen-

specific cytotoxic T cells that can clear the infection.

NK cells also secrete their own cytokines of their own to help facilitate immune responses, generally upon NK cell activation. NK cells work to control viral infections by secreting IFNγ (interferon gamma) and TNFα (tumor necrosis factor alpha). IFNγ activates macrophages for phagocytosis and lysis while TNFα acts to promote direct NK tumor cell killing. It is also a potent inflammatory mediator that causes long-lasting inflammatory responses and fever in response to more severe infections. Patients deficient in NK cells prove to be more susceptible to most infections than people with normal levels of NK cells, due to a loss of innate immune system function and efficiency.

This page titled 20.1C: Natural Killer Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1D: Inflammation

Inflammation is part of the biological response of vascular tissues to harmful stimuli.

Learning Objectives

- Describe the biological mechanisms of inflammation and its role in innate immunity

Key Points

- Acute inflammation occurs due to infection, injury, or irritation, and is an essential part of the healing process to remove pathogens and start the wound-healing process.
- During inflammation, vasodilation occurs, the endothelium becomes more permeable as exudate leaks into the tissues, and neutrophils migrate to the site of inflammation.
- Acute inflammation is initiated by cells already present in all tissues, such as mast cells that recognize pathogen-associated molecular patterns (PAMPs) with toll-like receptors, but other cells like natural killer (NK) cells can trigger inflammation as well.
- Acute inflammation is characterized by pain, redness, immobility (loss of function), swelling, and heat.
- Neutrophils migrate to inflamed tissues by rolling onto the endothelium with selectins, adhering to it with integrins, and sliding through its gaps with PECAM-1. Then they follow chemokines to the tissues to find pathogens to destroy.
- Repeated bouts of acute inflammation lead to chronic inflammation, a process of constant healing and inflammation-induced damage as the initial problem is never truly healed.
- Allergic reactions are the result of an inappropriate immune response that triggers inflammation.

Key Terms

- **extravasion**:
- **exudate**: Protein-rich edema caused by proteins flowing into the tissues during inflammation due to increased vascular permeability and oncotic pressure.
- **inflammatory mediator**: Any chemical released from cells that stimulates the vasodilation and increased permeability that occur during acute inflammation.

Inflammation is part of the complex biological response of vascular tissues to harmful stimuli, such as pathogens, injury or trauma, and irritants. Inflammation is a protective attempt by the organism to remove injurious stimuli and initiate the healing process. Inflammation is not a synonym for infection, even in cases where inflammation is caused by infection. Rather, it refers to the response of the body to try and fight the infection. While inflammation is an important mechanism of innate immunity, it can harm the body in cases of allergy, autoimmunity, and infections in tissues with poor regenerative capacity such as the heart.

Functions and Components of Inflammatory Response

The main function of inflammation is to trigger an immune response in an area of the body that needs it to fight off pathogens that may cause an infection or to help heal an injury. The main symptoms of acute inflammation are swelling, redness, pain, loss of function, and heat. Three components to the basic acute inflammatory response occur every time: vasodilation, increased vascular permeability, and migration of leukocytes to the affected tissues.

Vasodilation

An inflammatory response can be caused by any of numerous inflammatory mediators released from innate immune system cells. The most common short term mediators are histamine and seratonin from mast cells, but bradykinin, complement proteins, some interleukins, prostaglandins, and TNF-alpha may also trigger inflammation from other types of cells. Circulating mast cells contain toll-like receptors, which can detect pathogen-associated molecular patterns (PAMPS) on the surface of pathogens and release

an inflammatory mediator such as histamine in response. Alternatively, mast cells may release inflammatory mediators due to signals from damaged cells (which will release clotting factors) during trauma or injury.

After an inflammatory mediator is released in the bloodstream, a period of transient vasoconstriction, lasting only a few seconds, occurs. Then blood vessels expand to undergo vasodilation from the stimulus of the vasoactive inflammatory mediator, which increases blood flow to the area. This causes slowing and stasis of red blood cells, which can be involved in the clotting response needed to stop bleeding in the case of injury. Vasodilation is the reason for the redness, heat, and pain associated with inflammation.

Increased Vascular Permeability

The next step of acute inflammation is an increase in vascular permeability due to inflammatory mediator activity, which causes the blood vessels to become more permeable. Normally only water and small compounds can exit the bloodstream into the tissues, but during inflammation, large proteins in the bloodstream, such as serum albumins, can leak out and into the tissues. Water follows these proteins due to the force of oncotic pressure that the proteins exert. This is called exudate, a form of edema. As exudate accumulates within the tissues, they become swollen. The exudate may carry antimicrobial proteins and antibodies into the tissues, and stimulates lymphatic drainage.

Leukocyte Migration to the Tissues

The next step of the acute inflammatory response is chemotaxis migration of neutrophils to the affected area. Neutrophils are recruited to the site of inflammation by various cytokines. Other inflammatory mediators, such as TNF-alpha and IL-1, increase the expression of adhesion molecules on vascular endothelial cells. The neutrophils loosely attach to the endothelial cells through use of selectins, a process called *rolling*. Then integrins firmly attach to the adhesion molecules on the endothelial cells, which is called *adhesion*.Together, rolling and adhesion are referred to as *margination*, the accumulation of leukocytes on the endothelium.

The next step is for neutrophils to squeeze through the gaps in the endothelium into the tissues through binding with PECAM-1 expressed on the endothelium, a process called *extravasion*. Then the neutrophils follow a chemotactic gradient to the site of infection of injury in the tissues, where they will degranulate and phagocytize pathogens. Later, macrophages enter the tissues through a similar process to clean up dead neutrophils and cellular debris.

Outcomes of Acute Inflammation



Inflammation: Toes inflamed by chilblains

When acute inflammation ends (typically by release of anti-inflammatory mediators such as IL-10 or an end to the release of inflammatory mediators) resolution will occur if the problem is alleviated. Resolution involves physiological responses that are part of the healing process, such as wound healing. If the problem is not resolved, acute inflammation could occur again. Repeated bouts of acute inflammation, known as chronic inflammation, leads to a progressive shift in the type of cells present at the site of inflammation and is characterized by simultaneous destruction and healing of the tissue from the inflammatory process. In particular, fibrosis (scarring) and tissue necrosis are common outcomes of chronic inflammation.

This page titled 20.1D: Inflammation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1E: Antimicrobial Proteins

Antimicrobial peptides are an evolutionarily-conserved component of the innate immune response found among all classes of life.

Learning Objectives

- Describe the role of antimicrobial peptides in the innate immune system

Key Points

- Antimicrobial peptides include a net positive charge and hydrophilic and hydrophobic ends, which allow them to adhere to the lipid membranes of bacterial cell membranes.
- The positive charge makes antimicrobial peptides selective, so they only adhere to negatively-charged bacterial cell membranes instead of host cell membranes.
- The modes of action by which antimicrobial peptides kill bacteria are varied and include disrupting membranes, interfering with metabolism, and targeting cytoplasmic components.
- Many bacteria have developed antimicrobial resistance, in which a component of their cell membranes or enzyme secretions is altered to prevent the peptides from binding to the bacteria, or by inhibiting the peptides directly.

Key Terms

- **antimicrobial resistance**: Any mechanism that enables bacteria to evade or inhibit antimicrobial action.
- **peptide**: A class of organic compounds consisting of various numbers of amino acids in which the amine of one is reacted with the carboxylic acid of the next to form anamide bond.
- **amphipathic**: A molecule with both hydrophobic and hydrophilic groups that allow it to adhere to lipid structures more easily.

Antimicrobial peptides (also called host defense peptides) are an evolutionarily-conserved component of the innate immune response found among all known species. These peptides are found in many of the mucus membranes across the human body and are therefore considered to be part of the barrier immune system. The function of these peptides is to kill microbial pathogens and prevent them from entering the body.

Structure


Antimicrobial peptides are a unique and diverse group of molecules. As peptides, they consist of chains of amino acids that determine their composition and structure. These peptides have a stronger positive than negative charge, which is an important component of their selectivity. They also include hydrophobic and hydrophilic groups that enable them to latch onto other molecules (often lipids) through intermolecular forces, such as the lipid bilayer that forms cell membranes.

The secondary structures of these molecules follow four themes, including i) α-helical, ii) β-stranded due to the presence of two or more disulfide bonds, iii) β-hairpin or loop due to the presence of a single disulfide bond and/or cyclization of the peptide chain, and iv) extended. Many of these peptides are unstructured and inactive in free solution, and fold into their final configuration upon reaching mucus membranes. The amphipathicity (hydrophilic and hydrophobic ends) and positive charge of peptides are their defining structural features.


Antimicrobial Action

The modes of action by which antimicrobial peptides kill bacteria are varied and include disrupting cell membranes, interfering with metabolism, and damaging organelles. The initial contact between the peptide and target organism is electrostatic due to the force of negative and positive ionic charge. The amino acid composition, amphipathicity, cationic charge, and size allow them to attach to and insert into membrane bilayers to form pores by barrel-stave, carpet, or toroidal-pore mechanisms.

The peptides are selective and thus more likely to adhere to bacterial cell membranes than to cell membranes of the host cells. The peptides have a greater positive charge than negative charge, while bacterial cell membranes have a greater negative charge than host cell membranes. This causes the peptide to bind to bacterial membranes instead of host cell membranes.



Mechanism of Selectivity of Antimicrobial Peptides: Cell membranes of bacteria are different from the cell membranes of plants and animals and are preferentially targeted by the antimicrobial proteins.



Modes of Action by Antimicrobial Peptides: Antimicrobial peptides multiple various modes of action.

Other antimicrobial mechanisms include intracellular binding models. These involve inhibition of cell wall synthesis, alteration of the cytoplasmic membrane, activation of autolysin, inhibition of DNA, RNA, and protein synthesis, and inhibition of certain bacterial enzymes. However, in many cases, the exact killing mechanism is unknown. In general, the antimicrobial activity of these peptides is determined by measuring the minimal inhibitory concentration (MIC), the lowest concentration of drug that inhibits bacterial growth and an indicator of the antimicrobial strength of that peptide.

Antimicrobial Resistance

Despite the efficiency of antimicrobial peptides to inhibit and kill bacteria, they are still able to get inside the body and cause

infections. Bacteria can develop resistance to antimicrobial peptides (as well as separate resistances to antibiotics and other antimicriobials). Bacteria like *staphylococcus aureas*, which forms the highly resistant MRSA strain, can reduce the negativity of the charge of its cell membrane by bringing amino acids from the cytoplasm into its cell membrane so antimicrobial peptides won't bind to it. Other forms of antimicrobial resistance include producing enzymes that inhibit the antimicrobial peptides, altering the hydrophobic forces on the cell membrane, and capturing antimicrobial peptides in vesicles on the cell membrane to remove them from the bacterium.

Additionally, commensal bacteria have developed antimicrobial resistance to peptides, but they are normal flora of the body. Most never act as pathogens, though some may be opportunistic pathogens or only act as pathogens in people with certain genetic characteristics.

This page titled 20.1E: Antimicrobial Proteins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.1F: Fever

Fever is an elevation of body temperature above the regulatory set point, mediated through the release of prostaglandin E2.

Learning Objectives

- Describe how fever, a common symptom of medical conditions, is induced by endogenous and exogenous pyrogens


Key Points

- Temperature is ultimately regulated in the hypothalamus. A trigger of the fever, called a pyrogen, causes a release of prostaglandin E2 (PGE2). PGE2 then acts on the hypothalamus, which raises the temperature set point so that the body temperature increases through heat generation and vasoconstriction.
- Fever may be useful as an innate immune response to fight off infections by killing bacteria and viruses.
- Aspirin is a potent anti-fever drug because it inhibits COX-2 production, which inhibits PGE2 release.
- A pyrogen is a substance that induces fever and can be either internal (endogenous) or external (exogenous) to the body.
- During severe infections, fever can be more harmful than helpful as the body's cells are injured in addition to the bacterial cells, which can cause more problems for the innate immune system to handle.

Key Terms

- **pyrogen**: Any substance that produces fever or a rise in body temperature through the arachidonic acid pathway.
- **arachidonic acid pathway**: The pathway by which the fever regulator prostaglandin E-2 and several inflammatory mediators are produced by pyrogen activity with phospolipids and COX-2, usually in the brain or liver.

Fever (also known as pyrexia) is a physiological process of the innate immune response against many infections and diseases, characterized by an elevation of temperature above the normal range of 36.5–37.5 °C (98–100 °F) due to an increase in the body temperature regulatory set-point. Although the person's temperature increases, there is often a feeling of cold. Once the new temperature is reached, there is a feeling of warmth. A fever can be caused by many conditions ranging from benign to potentially serious. Fevers are helpful in fighting infections, but can also cause damage in the body.



Performance of the Various Types of Fever: Performance of the various types of fever: a) Fever continues b) Fever continues to abrupt onset and remission c) Fever remittent d) Intermittent fever e) Undulant fever f) Relapsing fever

Fever Pathways

Temperature is ultimately regulated in the hypothalamus. The primary fever mediator in the human body is prostaglandin E2 (PGE2), which acts on the hypothalamus to raise the temperature set point. PGE2 release comes from the arachidonic acid pathway, which also produces inflammatory mediators such as thromboxane and leukotriene.

This pathway is mediated by the enzymes phospholipase A2 (PLA2), cyclooxygenase-2 (COX-2), and prostaglandin E2 synthase. These enzymes ultimately mediate the synthesis and release of PGE2. Therefore, COX-2 inhibitors such as aspirin are commonly used to reduce fever, although treatments designed to inhibit pyrogens are also effective.

The hypothalamus is the thermostat of the body, in that it alters the temperature set point during temperature feedback and fevers. During a fever, the set point is raised, which causes the body to increase its temperature through both actively generating and retaining heat (vasoconstriction). If these measures are insufficient to make the blood temperature in the brain match the new setting in the hypothalamus, then shivering begins so those muscle movements produce more heat. When the fever stops (when PGE2 release ends), the temperature set point is lowered to normal, and the reverse of these processes (vasodilation, end of shivering, and nonshivering heat production) as well as sweating are used to cool the body to the new, lower setting.

Pyrogens

A pyrogen is a substance that induces fever and can be either internal (endogenous) or external (exogenous) to the body. Pyrogenicity can vary: in extreme examples, bacterial pyrogens known as superantigens can cause rapid and dangerous fevers. Depyrogenation may be achieved through filtration, distillation, chromatography, or inactivation.

Exogenous factors s lipopolysaccharide toxin (from gram negative bacteria) which can activate a number of innate immune activation pathways. These pathways induce the expression of endogenous pyrogens, including a variety of cytokines such as IL1α, IL1β, IL6, TNFα, TNFβ, IFNα, INFβ, and INFγ. For example, if an NK cell detects lipopolysaccharide from a pathogen, it will release TNFα, which will travel through the bloodstream to induce a number of long-lasting inflammatory changes including fever. When TNFα or any of these cytokine factors bind to cells in phospolipids in the brain, the arachidonic acid pathway is activated and PGE2 released to act on the hypothalamus and cause the fever response.

Problems with Fever

Fever is normally a beneficial immune process since increased body temperature can kill off bacteria and viruses and denature bacterial enzymes. But when the body temperature climbs too high, fever is often more harmful than helpful. High fevers also denature the body's own proteins, which can alter normal cell metabolism, leading to cell injury and death. Persistent high body temperature can also trigger apoptosis. Treatments for severe fevers include antipyrogens and aspirin, which also helps to stop blood clots that may coincide with severe fever.

High fevers (more than 104 degrees Fahrenheit) are a symptom of severe infections. While fevers typically aren't the direct cause of death in these cases, they do tend to worsen the prognosis. For example, septic shock is a severe bacterial infection in which bacterial toxins stimulate pyrogen and inflammatory mediator activity causes high fever. The fever makes it harder for the body to stop the systemic organ failure that occurs from the compensatory mechanisms in septic shock. Organs fail as blood is pulled away from them to fight the infection (compensatory mechanisms), the damage caused by the fever results in even more compensatory mechanism activity. While septic shock is one of the worst possible examples of fever, it illustrates an important concept in pathophysiology: that normal immune functions are as easily able to hurt us as help us.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Mucosal immune system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mucosal_immune_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Skin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Skin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mucous membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mucous_membrane. **License**: *CC BY-SA: Attribution-ShareAlike*
- Intestinal mucosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intesti...ucosa%23Mucosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- integumentary system. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/integumentary_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:G...ng%23filelinks. **License**: *Public Domain: No Known Copyright*
- Phagocyte. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Phagocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- epithelial cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/epithelial%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- mesenchymal cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/mesenchymal%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- chemotaxis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chemotaxis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:G...ng%23filelinks. **License**: *Public Domain: No Known Copyright*
- Dendritic cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Dendritic_cell.JPG. **License**: *CC BY: Attribution*
- Myeloid cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Myeloid_cells.png. **License**: *Public Domain: No Known Copyright*
- NeutrophilerAktion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:NeutrophilerAktion.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Natural killer cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Natural_killer_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cytotoxic T cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cytotoxic%20T%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:G...ng%23filelinks. **License**: *Public Domain: No Known Copyright*
- Dendritic cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Dendritic_cell.JPG. **License**: *CC BY: Attribution*
- Myeloid cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Myeloid_cells.png. **License**: *Public Domain: No Known Copyright*
- NeutrophilerAktion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:NeutrophilerAktion.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Missingself. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Missingself.svg. **License**: *Public Domain: No Known Copyright*
- Inflammation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Inflammation. **License**: *CC BY-SA: Attribution-ShareAlike*
- pattern recognition receptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pattern%20recognition%20receptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray940.png%23filelinks. **License**: *Public Domain: No Known Copyright*
- Dendritic cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Dendritic_cell.JPG. **License**: *CC BY: Attribution*
- Myeloid cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Myeloid_cells.png. **License**: *Public Domain: No Known Copyright*
- NeutrophilerAktion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:NeutrophilerAktion.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Missingself. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Missingself.svg. **License**: *Public Domain: No Known Copyright*
- Wintertenen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Wintertenen.jpg. **License**: *Public Domain: No Known Copyright*
- Antimicrobial peptides. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Antimicrobial_peptides. **License**: *CC BY-SA: Attribution-ShareAlike*
- amphipathic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amphipathic. **License**: *CC BY-SA: Attribution-*

ShareAlike
- peptide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/peptide. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray940.png%23filelinks. **License**: ***Public Domain: No Known Copyright***
- Dendritic cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Dendritic_cell.JPG. **License**: *CC BY: Attribution*
- Myeloid cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Myeloid_cells.png. **License**: *Public Domain: No Known Copyright*
- NeutrophilerAktion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:NeutrophilerAktion.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Missingself. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Missingself.svg. **License**: *Public Domain: No Known Copyright*
- Wintertenen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Wintertenen.jpg. **License**: *Public Domain: No Known Copyright*
- Modes of action. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Modes_of_action.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mechanim of Selectivity of Antimicrobial Peptides. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mechanim_of_Selectivity_of_Antimicrobial_Peptides.jpg. **License**: *Public Domain: No Known Copyright*
- Fever. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Fever. **License**: *CC BY-SA: Attribution-ShareAlike*
- thermoregulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thermoregulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- pyrogen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pyrogen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray940. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray940.png%23filelinks. **License**: *Public Domain: No Known Copyright*
- Dendritic cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Dendritic_cell.JPG. **License**: *CC BY: Attribution*
- Myeloid cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Myeloid_cells.png. **License**: *Public Domain: No Known Copyright*
- NeutrophilerAktion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:NeutrophilerAktion.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Missingself. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Missingself.svg. **License**: *Public Domain: No Known Copyright*
- Wintertenen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Wintertenen.jpg. **License**: *Public Domain: No Known Copyright*
- Modes of action. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Modes_of_action.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mechanim of Selectivity of Antimicrobial Peptides. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Mechanim_of_Selectivity_of_Antimicrobial_Peptides.jpg. **License**: *Public Domain: No Known Copyright*
- Febbre. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Febbre.gif. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 20.1F: Fever is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.2: STRESS AND IMMUNITY

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

20.2A: Stress and Immunity

This page titled 20.2: Stress and Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.2A: Stress and Immunity

Evidence shows that stress has a negative effect on the body's immune system.

Learning Objectives

- Describe the effects of stress on the immune system

Key Points

- Cortisol is released in the adrenocortex by the neuroendocrine activity of the hypothalamus and anterior pituitary gland during acute stress responses, which involve sympathetic nervous system stimulation to cause a fight or flight response.
- Chronic stress is a state of prolonged tension from internal or external stressors, which may cause various physical manifestations— asthma, back pain, arrhythmias, fatigue, headaches, high blood pressure, irritable bowel syndrome, ulcers, high blood sugar, diabetes, heart problems, and Cushing's syndrome.
- Cortisol, a hormone released during stressful situations, affects the immune system by preventing the production of inflammatory mediators. During chronic stress, cortisol is overproduced, causing fewer receptors to be produced in immune cells and inducing chronic inflammation.
- Managing stress is important for maintaining a healthy lifestyle. Exercise, meditation, hobbies, social activities, and listening to music are all effective ways to manage stress.

Key Terms

- **stress**: Emotional pressure suffered by a human being or other animal.
- **Cortisol**: The hormone that facilitates the physiological response to stress by raising blood sugar to increase cellular metabolism, activating the sympathetic nervous system and suppressing the inflammatory response.

EXAMPLES

In one study, individuals caring for spouses with dementia, representing the stress group, saw a significant decrease in immune response to an influenza-virus vaccine compared to a non-stressed control group.

Stress is the body's reaction to any stimuli that disturb its equilibrium. Stress responses release a hormone called cortisol in the adrenocortex through a complex neuroendocrine pathway that is controlled by the hypothalamus and anterior pituitary gland.

Effects of Acute Stress on the Immune System

Cortisol has numerous effects on the body, such as sympathetic nervous system activation, increasing blood sugar for energy purposes, and anti-inflammatory effects that include the inhibition of certain inflammatory mediators that are important in innate immunity. For example, expression of IL-1 IFN-gamma and TNF-alpha from immune cells are all inhibited by cortisol, while cytokines that control helper T-cell activity are increased. Evolutionary biologists believe that this stress mechanism is intended to protect the immune system from becoming overactive to facilitate the fight-or-flight response, which would be weakened by inflammation. Other studies show that the weakened inflammatory effect makes pathogen-caused diseases more likely to lead to infection.

Research studies in which participants were subjected to a variety of viruses showed that stress has a negative effect on the immune system. A study was conducted using a rhinovirus, the causative agent of the common cold. Participants were infected with the virus and given a stress index. Results showed that an increase in score on the stress index correlated with greater severity of cold symptoms. Studies with HIV have also shown that stress speeds up viral progression. Men with HIV were two to three times more likely to develop AIDS when under above-average stress.

Effects of Chronic Stress

Normally, stress responses are beneficial for the body, provided they are moderate and cortisol returns to normal levels after the stressful situation ends. Chronic stress occurs when the body's stress levels remain too high for too long and do not return to normal levels for long enough. Chronic stress is associated with many chronic diseases, including asthma, back pain, arrhythmia, fatigue, headaches, high blood pressure, irritable bowel syndrome, ulcers, high blood sugar and diabetes.

Chronic stress takes a more significant toll on the body than does acute stress. It can raise blood pressure enough to cause hypertension, increase the risk of heart attack, stroke, and cardiac hypertrophy, increase vulnerability to anxiety and depression, contribute to infertility, and hasten the aging process. Cushing's syndrome is a chronic disease caused by persistent and excessive cortisol levels (from stress and often tumors that secrete cortisol). It typically causes weight gain, hair growth, and increased risk for diabetes. Over time, immune cells may become desensitized to cortisol and express fewer cortisol receptors, so chronic inflammation can develop as the anti-inflammatory effect of cortisol is weakened.



Skin Ulcer: Erythema nodosum—lesions that occur in some patients suffering from inflammatory bowel syndrome.

Dealing with Stress

Proper stress management is important for good overall health because it can prevent the plethora of chronic diseases and short-term illnesses incurred by a stressful lifestyle. Common effective ways to manage stress include exercise, meditation, therapy, and participating in hobbies, music, and social activities. Exercise relieves stress through endorphins, which have an inhibitory effect on cortisol secretion. Other activities like meditation and listening to music have been found to calm the mind, which can relax the parts of the brain responsible for the stress response (such as the anterior pituitary gland). There are many other ways to deal with stress, and they all have the potential to improve both mental and physiological health.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Stress (biology). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stress...mmune_response. **License**: *CC BY-SA: Attribution-ShareAlike*
- stress. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/stress. **License**: *CC BY-SA: Attribution-ShareAlike*
- lymphoproliferative response. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lymphop...ive%20response.

License: ***CC BY-SA: Attribution-ShareAlike***

- Ulcer (dermatology). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ulcer_(dermatology)**. **License**: ***Public Domain: No Known Copyright***

This page titled 20.2A: Stress and Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3: ADAPTIVE IMMUNITY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

20.3A: Overview of Adaptive Immunity

20.3B: Types of Adaptive Immunity

20.3C: Maturation of T Cells

20.3D: Lymphocytes

20.3E: Antigen-Presenting Cells

This page titled 20.3: Adaptive Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3A: Overview of Adaptive Immunity

The adaptive immune system is composed of highly-specialized systemic cells and processes that eliminate or prevent pathogenic growth.

Learning Objectives

- Describe adaptive immunity

Key Points

- The adaptive immune response provides the vertebrate immune system with the ability to recognize and remember specific pathogens to generate immunity, and mount stronger attacks each time the pathogen is encountered.
- The cells of the adaptive immune system are a type of leukocyte called a lymphocyte. B cells and T cells are the major types of lymphocytes involved in adaptive immunity.
- B and T cells can create memory cells to defend against future attacks by the same pathogen by mounting a stronger and faster adaptive immune response against that pathogen before it can even cause symptoms of infection.
- Antigen -presenting cells present captured antigens to immature lymphocytes, which then mature to be specific to that antigen and work to destroy pathogens that express that antigen.
- Hypersensitivity disorders (allergies) may occur when an adaptive immune response forms against antigens that aren't associated with pathogens, such as pollen. More complex hypersensitivity disorders may involve cytotoxic T cells and cause chronic inflammation and damage to the body's own tissues.

Key Terms

- **antigen**: A substance that induces an immune response, usually a molecule found on a pathogen such as a toxin or molecule expressed by the pathogen or pathogen-infected cells.
- **hypersensitivity**: A disorder in which an adaptive immune response forms memory cells specific to antigens that aren't associated with pathogens.
- **antigen presenting cell**: A cell that presents captured antigens to immature T-cells. Dendritic cells and macrophages are the best examples, but several other cells can present antigens as well.

The adaptive immune system, also known as the specific immune system, is composed of highly-specialized systemic cells and processes that eliminate or prevent pathogenic growth. The adaptive immune system works to protect and heal the body when the innate immune system fails. It provides the body with the ability to recognize and remember specific pathogens through their antigens. This mechanism allows the immune system to mount stronger attacks each time the pathogen is encountered, thus preparing itself for future challenges and preventing reinfection by the same pathogen.

Functions of the Adaptive Immune System


The adaptive immune system starts to work after the innate immune system is activated. If an infection progresses despite the inflammation, fever, natural killer (NK) cell and phagocyte activity of the innate immune system, a more coordinated response is required in order to destroy the pathogen. The adaptive immune response occurs a few days after the innate immune response is initiated. The major functions of the adaptive immune system include:

- The recognition of specific "non-self" antigens in the presence of "self" during the process of antigen presentation
- The generation of responses that are tailored to maximally eliminate specific pathogens or pathogen-infected cells
- The development of immunological memory in which each pathogen is "remembered" by a signature antibody, which can then be called upon to quickly eliminate a pathogen should subsequent infections occur.

Formation of Adaptive Immunity

Adaptive immunity is triggered when a pathogen evades the innate immune system for long enough to generate a threshold level of an antigen. An antigen is any molecule that induces an immune response, such as a toxin or molecular component of a pathogen cell membrane, and is unique to each species of pathogen. A typical adaptive immune response includes several steps:


1. The antigen for the pathogen is taken up by an antigen-presenting cell (APC), such as a dendritic cell or macrophage, through phagocytosis.
2. The APC travels to a part of the body that contains immature T and B cells, such as a lymph node.
3. The antigen is processed by the APC and bound to MHC class II receptors and MHC class I receptors on the cell membrane of the APC.
4. The antigen is presented to immature helper T cells and cytotoxic T cells through binding the MHC II (helper T) or MHC I (cytotoxic T) to T-cell receptors.
5. These T lymphocytes mature and proliferate. Helper T cells activate B cells, which proliferate and produce antibodies specific to the antigen, while cytotoxic T cells destroy pathogens that bear the antigen that was presented to them by the APCs.
6. Memory B and T cells are formed after the infection ends.



Antigen Presentation: Antigen presentation stimulates T cells to become either “cytotoxic” CD8+ cells or “helper” CD4+ cells. Cytotoxic cells directly attack cells carrying certain foreign or abnormal molecules on their surfaces. Helper T cells, or Th cells, coordinate immune responses by communicating with other cells. In most cases, T cells only recognize an antigen if it is carried on the surface of a cell by one of the body’s own MHC, or major histocompatibility complex, molecules.

Immunological Memory

When B cells and T cells are activated, some become memory cells. Throughout the lifetime of an animal, these memory cells form a database of effective B and T lymphocytes. Upon interaction with a previously-encountered antigen, the appropriate memory cells are selected and activated. In this manner, the second and subsequent exposures to an antigen produce a stronger and faster immune response. This is “adaptive” because the body’s immune system prepares itself for future challenges, which can stop an infection by the same pathogen before it can even cause symptoms.



Antibody: An antibody is made up of two heavy chains and two light chains. The unique variable region allows an antibody to recognize its matching antigen

Immunological memory can either be in the form of passive short-term memory or active long-term memory. Passive memory is usually short-term, lasting between a few days and several months, and is particularly important for newborn infants, who are given passive memory from maternal antibodies and immune cells before birth. Active immunity is generally long-term and can be acquired by infection followed by B cells and T cells activation, or artificially acquired by vaccines in a process called immunization.

The memory system does have a few flaws. Pathogens that undergo mutation often have different antigens than those known by memory B and T cells, meaning that different strains of the same pathogen can avoid the memory-enhanced immune response. Additionally, the memory cell function enables the development of hypersensitivity disorders, such as allergies and many chronic diseases (like multiple sclerosis or myasthenia gravis). In these cases, memory cells form for an antigen that elicits an immune response without actually being caused by a pathogen, which leads to immune system mediated-damage to the body from mast cell, antibody, or T-cell mediated activities and inflammation.

This page titled 20.3A: Overview of Adaptive Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3B: Types of Adaptive Immunity

The adaptive immune response is mediated by B and T cells and creates immunity memory.

Learning Objectives

- Distinguish between the types of adaptive immunity

Key Points

- B cells and T cells, the major types of lymphocytes, are very important in the adaptive immune system.
- B cells, type 2 helper T cells, antibodies, mast cells, and eosinophils are involved in the humoral immune response.
- Type 1 helper T cells and cytoxic T-cells are involved in cell-mediated immune response.
- Cytotoxic T cells kill pathogens through release of perforin, granzymes, and proteases, which cause the target cell to undergo apoptosis.
- Antibodies bind to pathogens to opsonize them, neutralize pathogen toxins, and activate the complement complex system. IgE also alerts circulating mast cells and eosinophils of known antigens, which causes a rapid inflammatory response.

Key Terms

- **cell-mediated immunity**: Adaptive immunity that is not controlled by antibodies and is instead mediated directly by immune cells themselves, most notably type 1 helper T cells and cytotoxic T-cells.
- **humoral immunity**: Adaptive immunity that refers to antigen-specific components flowing through the plasma, such as antibodies, their function, and the cells that produce them.

The adaptive immune system mounts a stronger, antigen-specific immune response after the innate immune response fails to prevent a pathogen from causing an infection. There are two subdivisions of the adaptive immune system: cell-mediated immunity and humoral immunity.

Cell-Mediated Immunity

Cell mediated immunity is controlled by type 1 helper T cells (T_h1) and cytotoxic T cells. These cells are activated by antigen-presenting cells, which causes them to rapidly mature into forms specific to that antigen. T cells then circulate through the body to destroy pathogens in several ways. Helper T cells facilitate the immune response by guiding cytotoxic T cells to pathogens or pathogen-infected cells, which they will then destroy.

Cytotoxic T cells kill pathogens in several ways, including the release of granules that contain the cytotoxins perforin and granzyme, which lyse small pores in the membrane of a pathogen. Then T-cell produced proteases enter the pathogen and induce an apoptosis response within the cell. Helper T cells secrete cytokines such as interferon-gamma, which can activate cytotoxic T cells and macrophages.


Humoral Immunity

Humoral immunity refers to the component of the adaptive immune response that is caused by B cells, antibodies, and type 2 helper T cells (T_h2), as well as circulating mast cells and eosinophils to a lesser extent. Its name comes from the idea that blood is one of the humors of the body, since antibodies provide passive or active immunity through circulation in the bloodstream.

Type 2 helper T cells are included in the humoral immune system because they present antigens to immature B-cells, which undergo proliferation to become specific to the presented antigen. The B cells then rapidly produce a large number of antibodies that circulate through the body's plasma.

Antibodies provide a number of functions in humoral immunity. Six different classes of antibodies provide distinct functions and interact with different cells in the immune system. All antibodies bind to pathogens to opsonize them, which makes it easier for phagocytic cells to bind to and destroy the pathogen. They also neutralize the toxins produced by certain pathogens and provide complement pathway activation, in which circulating proteins are combined in a complex cascade that forms a membrane attack complex on a pathogen cell membrane, which lyses the cell.

Mast cells and eosinophils are considered part of the humoral immune system because they can be sensitized towards certain antigens through circulating immunoglobin E (IgE), a specific type of antibody produced by B cells. IgE binds to the mast cells and eosinophils when an antigen is detected, using a type of Fc receptor on the mast cell or eosinophil that has a high-binding affinity with IgE. This binding will cause degranulation and release of inflammatory mediators that start an immune response against the antigen. This process is the reason why memory B cells can cause hypersensitivity (allergy) formation, as circulating IgE from those memory cells will activate a rapid inflammatory and immune response.



Types of Adaptive Immunity: This diagram of adaptive immunity indicates the flow from antigen to APC, MHC2, CD4+, T helper cells, B cells, antibodies, macrophages, and killer T cells.

This page titled 20.3B: Types of Adaptive Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3C: Maturation of T Cells

T cells originate from hematopoietic stem cells in the bone marrow and undergo positive and negative selection in the thymus to mature.

Learning Objectives

- Describe T cell maturation

Key Points

- Hematopoietic progenitors derived from hematopoietic stem cells populate the thymus and expand by cell division to generate a large population of immature thymocytes. After the thymus becomes inactive later in life, existing immature T cells will proliferate through clonal expansion.
- About 98% of thymocytes die during the development processes in the thymus by failing either positive selection or negative selection, while the other 2% survive and leave the thymus to become mature immunocompetent T cells.
- During positive selection, only T cells that can bind to MHC are kept alive. The rest are killed by an apoptotic signal so that non-functional T cells don't get into the body and crowd out functional T cells.
- During negative selection, most T cells that bind too easily to self antigens are killed. Some are kept alive and differentiate into T reg cells, which help prevent overactive cell mediated immune responses.
- Autoimmune diseases may be caused either by antibodies or T cells that can bind to self antigens, causing damage to self organs and tissues. They may be caused by failed negative selection and often have a genetic component.

Key Terms

- **thymus**: A ductless gland consisting mainly of primary lymphatic tissue that is the site of lymphocyte maturation and selection.
- **negative selection**: The process by which T cells are screened so that those with a high affinity for binding to self antigens (and potentially causing autoimmunity) are destroyed.
- **Positive selection**: The process by which T cells are screened so that only those capable of binding to MHC are kept alive.

T cells belong to a group of white blood cells known as lymphocytes and play a central role in the cell-mediated branch of the adaptive immune system. They are distinguished from other lymphocytes, such as B cells and natural killer cells (NK cells), by the presence of a T cell receptor (TCR) on the cell surface. T cells are produced in the bone marrow but travel to the thymus to mature. T cells can be either helper T cells or cytoxic T cells based on whether they express CD4 (helper) or CD8 (cytotoxic) glycoprotein.

Maturation of T Cells

All T cells originate from hematopoietic stem cells in the bone marrow, which are capable of differentiating into any type of white blood cell. Immature T cells that migrate to the thymus are called thymocytes. The earliest thymocytes express neither CD4 nor CD8, and are therefore classed as double-negative (CD4-CD8-) cells. As they progress through their development they become double-positive thymocytes (CD4+CD8+) and finally mature to single-positive (CD4+CD8- or CD4-CD8+) thymocytes that are released from the thymus to peripheral tissues. Typically, these mature thymocytes are still referred to as either "immature" or "naive" because they have not been presented with an antigen. They travel to sites that contain secondary lymphoid tissue, such as the lymph nodes and tonsils, where antigen presentation. This facilitates the development of antigen-specific adaptive immunity.

The thymus contributes fewer cells as a person ages. As its functional mass shrinks by about 3% a year throughout middle age, there is a corresponding fall in the thymic production of naive T cells, leaving clonal expansion of immature T cells to play a greater role in protecting older subjects. The thymus is thus thought to be important in building a large stock of naive T cells soon after birth that can later function without thymus support.


Positive Selection of T Cells

During thymocyte maturation, 98% of T cells are discarded by selection, thich is a mechanism designed to ensure that T cells function without major problems. Positive selection designates T cells capable of interacting with MHC. Double-positive thymocytes (CD4+/CD8+) move deep into the thymic cortex tissue where they are presented with self-antigens. These are expressed by thymic cortical epithelial cells that express both MHC I and MHC II molecules on the surface of cortical cells. Only those thymocytes that interact with MHC I or MHC II will receive a vital “survival signal.” Thosethat can’t interact will undergo apoptosis (cell death). This insures T cell functionality since T cells with non-functional receptors cannot receive antigens and are thus useless to the immune system. If non-functional T cells were allowed into circulation, they would crowd out functional T cells and slow down the rate at which adaptive immune responses are formed. The vast majority of thymocytes die during this process.

A thymocyte’s differentiation into either a helper or cytotoxic version is also determined during positive selection. Double-positive cells (CD4+/CD8+) that are positively selected on MHC class II molecules will eventually become CD4+ helper T cells, while cells positively selected on MHC class I molecules mature into CD8+ cytotoxic T cells. A T cell is then signaled by the thymus to become a CD4+ cell by reducing expression of its CD8 cell surface receptors. If the cell does not lose its signal, it will continue reducing CD8 and become a CD4+, single positive cell. But if there is a signal interruption, it will instead reduce CD4 molecules, eventually becoming a CD8+, single positive cell. This process does not remove thymocytes that may become sensitized against self-antigens, which causes autoimmunity. The potentially autoimmune cells are removed by the process of negative selection.

Negative Selection of T Cells

Negative selection removes thymocytes that are capable of strongly binding with self-antigens presented by MHC. Thymocytes that survive positive selection migrate towards the boundary of the thymic cortex and thymic medulla (the part of the thymus where T cells enter circulation). While in the medulla, they are again presented with self-antigen in complex with MHC molecules on thymic epithelial cells. Thymocytes that interact too strongly with the antigen receive an apoptotic signal that leads to cell death.



Blood cells: Scanning electron micrograph of T lymphocyte (right), a platelet (center), and a red blood cell (left).

However, some cells are selected to become T-reg cells, which retain their ability to bind to self-antigens in order to suppress overactive immune responses. These cells may be protective against autoimmunity. The remaining cells exit the thymus as mature naive T cells. This process is an important component of central tolerance, a process that prevents the formation of self-reactive T cells that are capable of inducing autoimmune diseases in the host. Autoimmune diseases reflect a loss of central tolerance in which the body’s own B and T cells become sensitized towards self-antigens. Many autoimmune disorders are primarily antibody-mediated, but some are T cell mediated. One example of the latter is Crohn’s disease, in which T cells attack the colon. These autoimmune disorders may be caused by problems in negative selection and tend to have genetic components.

This page titled 20.3C: Maturation of T Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3D: Lymphocytes

A lymphocyte is a type of white blood cell in the vertebrate immune system.

Learning Objectives

- Describe the role of lymphocytes

Key Points

- The main types of T cells are helper T cells, cytotoxic T cells, memory T cells, and regulatory T cells.
- The main types of B cells are plasma cells, plasmablasts, memory B cells, and regulatory B cells.
- T cells are involved in cell-mediated immunity, whereas B cells are primarily responsible for humoral (antibody -related) immunity.
- If an antigen is detected again after the initial adaptive immune response, memory T cells create new helper and cytotoxic T cells, while memory B cells create new antibodies.
- Regulatory T and B cells suppress immune responses at the end of an infection and suppress T and B cells involved in autoimmunity.

Key Terms

- **Memory T cells**: A type of T cell that rapidly differentiates into helper and cytotoxic T cells if its associated antigen is detected.
- **Plasma Cell**: A type of B cell that produces most of the antibodies during development of adaptive immune response.

A lymphocyte is a type of white blood cell in the immune system, including both the B and T cells of the adaptive immune system and natural killer (NK) cells of the innate immune system.

B and T cells and their various subdivisions perform many adaptive immune functions.


T Cells

T cells mature in the thymus and contain T cell receptors (TCRs) that allow them to bind to antigens on MHC complexes. T cells are a major component in cell-mediated adaptive immunity because they provide a pathway for the direct killing of pathogens. There are two main types of T cells that express either CD4 or CD8 depending on signals that occur during T cell maturation, as well as less common types:

- Helper T cells (CD4s) facilitate the organization of immune responses, and can bind to MHC class II. Subtype 2 helper T cells present antigens to B cells. Subtype 1 helper T cells produce cytokines that guide cytotoxic T cells to pathogens and activate macrophages.
- Cytotoxic T cells (CD8s) destroy pathogens associated with an
antigen. They function similarly to NK cells by binding to
MHC class I and releasing perforin, granzymes, and proteases to induce apoptosis in a pathogen. They are different from NK cells because they only bind to
cells that express their specific antigen, and are not large or granular like NK cells.
- Suppressor T cells (T-reg cells) retain some of their ability to bind to self-cells. They have an immunosuppressive effect that inhibits cell-mediated immunity at the end of a response and destroys autoimmune T cells that aren't filtered out by negative selection in the thymus.
- Memory T cells are created after an adaptive immune response subsides, retaining the presented antigen. They rapidly proliferate and differentiate into helper and cytotoxic T cells that are specific to that antigen should it be detected in the body again.

While these are the main categories of T lymphocytes, there are other subtypes within these categories as well as additional categories that are not fully understood.

B Cells



Lymphocyte: A scanning electron microscope (SEM) image of a single human lymphocyte.

B cells are involved in humoral adaptive immunity, producing the antibodies that circulate through the plasma. They are produced and mature in bone marrow tissues and contain B cell receptors (BCRs) that bind to antigens. While in the bone marrow, B cells are sorted through positive and negative selection in a manner somewhat similiar to T cell maturation in the thymus, with the same process of killing B cells that are nonreactive to antigens or reactive to self-antigens. Instead of apoptosis, though, defective B cells are killed through other mechanisms such as clonal deletion. Mature B cells leave the thymus and travel to secondary lymphoid tissue such as the lymph nodes.

During antigen presentation, antigen-presenting cells first present antigens to T cells. Then mature helper T cells bind their antigen to naive B cells through BCRs. After antigen presentation, the naive B cells migrate together to germinal centers within the lymphoid tissue, where they undergo extensive proliferation and differentiation into different types of mature B cells. Some of the major categories of B cells that arise include:

- Plasma cell and long-lived B cells that are the main source of antibodies. They do not have the ability to proliferate and are considered terminally-differentiated.
- Plasmablasts are short-lived B cells produced early in an infection. Their antibodies have a weaker binding affinity than those of plasma cells.
- Regulatory B cells (B reg cells) are immunosuppresive B cells that secrete anti-inflammatory cytokines (such as IL-10) to inhibit autoimmune lymphocytes.
- Memory B cells are dormant B cells with the same BCR as the B cell from which they differentiated. They are specific to the antigen presented to that BCR and rapidly secrete large amounts of antigen-specific antibodies to prevent reinfection if that antigen is detected again.

Besides antibody production, B cells may also function in antigen presentation, though not to the degree of macrophages or dendritic cells. B cells are important to adaptive immune function but can cause problems as well. Autoreactive B cells may cause autoimmune disease that involves antibody-induced damage and inflammation. Certain B cells may undergo malignant tranformation into cancer cells such as lymphoma, in which they continually divide and form solid tumors.

This page titled 20.3D: Lymphocytes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.3E: Antigen-Presenting Cells

Antigen presentation is a process by which immune cells capture antigens and then enable their recognition by T cells.

Learning Objectives

- Describe the role of antigen-presenting cells

Key Points

- The host's cells express "self" antigens that identify them as such. These antigens are different from those in bacteria ("non-self" antigens) and in virus-infected host cells ("missing-self").
- Antigen presentation consists of pathogen recognition, phagocytosis of the pathogen or its molecular components, processing of the antigen, and then presentation of the antigen to naive T cells.
- The T cell receptor is restricted to recognizing antigenic peptides only when bound to appropriate molecules of the major histocompatibility complex (MHC), also known in humans as human leukocyte antigen (HLA).
- Helper T cells recieve antigens from MHC II on an APC, while cytotoxic T cells recieve antigens from MHC I. Helper T cells present their antigen to B cells as well.Dendritic cells, B cells, and macrophages play a major role in the innate response, and are the primary antigen-presenting cells (APC).
- APCs use toll-like receptors to identify PAMPS and DAMPs, which are signs of an infection and may be processed into antigen peptides if phagocytized. Most APCs cannot tell the difference between different types of antigens like B and T cells can.

Key Terms

- **damage-associated molecular pattern**: Protein or nucleic acid based signs of pathogen induced damage. Protein DAMPs may be phagocytized and processed for antigen presentation.
- **cytotoxic**: A population of T cells specialized for inducing the deaths of other cells.

Antigen presentation is a process in the body's immune system by which macrophages, dendritic cells and other cell types capture antigens, then present them to naive T-cells. The basis of adaptive immunity lies in the capacity of immune cells to distinguish between the body's own cells and infectious pathogens. The host's cells express "self" antigens that identify them as belonging to the self. These antigens are different from those in bacteria ("non-self" antigens) or in virally-infected host cells ("missing-self"). Antigen presentation broadly consists of pathogen recognition, phagocytosis of the pathogen or its molecular components, processing of the antigen, and then presentation of the antigen to naive (mature but not yet activated) T cells. The ability of the adaptive immune system to fight off pathogens and end an infection depends on antigen presentation.

Antigen Presenting Cells

Antigen Presenting Cells (APCs) are cells that capture antigens from within the body, and present them to naive T-cells. Many immune system cells can present antigens, but the most common types are macrophages and dendritic cells, which are two types of terminally differentiated leukocytes that arise from monocytes. Both of these APCs perform many immune functions that are important for both innate and adaptive immunity, such as removing leftover pathogens and dead neutrophils after an inflammatory response. Dendritic cells (DCs) are generally found in tissues that have contact with the external environment (such as the skin or respiratory epithelium) while macrophages are found in almost all tissues. Some types of B cells may also present antigens as well, though it is not their primary function.

APCs phagocytize exogenous pathogens such as bacteria, parasites, and toxins in the tissues and then migrate, via chemokine signals, to lymph nodes that contain naive T cells. During migration, APCs undergo a process of maturation in which they digest phagocytized pathogens and begin to express the antigen in the form of a peptide on their MHC complexes, which enables them to present the antigen to naive T cells. The antigen digestion phase is also called "antigen processing," because it prepares the antigens for presentation. This MHC:antigen complex is then recognized by T cells passing through the lymph node. Exogenous antigens are usually displayed on MHC Class II molecules, which interact with CD4+ helper T cells.


This maturation process is dependent on signaling from other pathogen-associated molecular pattern (PAMP) molecules (such as a toxin or component of a cell membrane from a pathogen) through pattern recognition receptors (PRRs), which are received by Toll-like receptors on the DC's body. They may also recognize damage-associated molecular pattern (DAMP) molecules, which include degraded proteins or nucleic acids released from cells that undergo necrosis. PAMPs and DAMPS are not technically considered antigens themselves, but instead are signs of pathogen presence that alert APCs through Toll-like receptor binding. However if a DC phagocytzes a PAMP or DAMP, it could be used as an antigen during antigen presentation. APCs are unable to distinguish between different types of antigens themselves, but B and T cells can due to their specificity.

Antigen Presentation

T cells must be presented with antigens in order to perform immune system functions. The T cell receptor is restricted to recognizing antigenic peptides only when bound to appropriate molecules of the MHC complexes on APCs, also known in humans as Human leukocyte antigen (HLA).

Several different types of T cell can be activated by APCs, and each type of T cell is specially equipped to deal with different pathogens, whether the pathogen is bacterial, viral or a toxin. The type of T cell activated, and therefore the type of response generated, depends on which MHC complex the processed antigen-peptide binds to.

MHC Class I molecules present antigen to CD8+ cytotoxic T cells, while MHC class II molecules present antigen to CD4+ helper T cells. With the exception of some cell types (such as erythrocytes), Class I MHC is expressed by almost all host cells. Cytotoxic T cells (also known as TC, killer T cell, or cytotoxic T-lymphocyte (CTL)) are a population of T cells that are specialized for inducing the death of other cells. Recognition of antigenic peptides through Class I by CTLs leads to the killing of the target cell, which is infected by virus, intracytoplasmic bacterium, or are otherwise damaged or dysfunctional. Additionally, some helper T cells will present their antigen to B cells, which will activate their proliferation response.



Antigen presentation: In the upper pathway; foreign protein or antigen (1) is taken up by an antigen-presenting cell (2). The antigen is processed and displayed on a MHC II molecule (3), which interacts with a T helper cell (4). In the lower pathway; whole foreign proteins are bound by membrane antibodies (5) and presented to B lymphocytes (6), which process (7) and present antigen on MHC II (8) to a previously activated T helper cell (10), spurring the production of antigen-specific antibodies (9).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- antigen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antigen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Adaptive immune system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Adaptive_immune_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- antibody. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antibody. **License**: *CC BY-SA: Attribution-ShareAlike*
- macrophage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/macrophage. **License**: *CC BY-SA: Attribution-ShareAlike*
- Antibody. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antibody.jpg. **License**: *Public Domain: No Known Copyright*
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antigen_presentation.svg. **License**: *Public Domain: No Known Copyright*
- Immune system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immune_system%23Innate_immune_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- antigen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antigen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Antibody. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antibody.jpg. **License**: *Public Domain: No Known Copyright*
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antigen_presentation.svg. **License**: *Public Domain: No Known Copyright*
- Lymphocyte_activation_simple.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immune_system#/media/File:Lymphocyte_activation_simple.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- T cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/T_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- B cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/B_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Immune cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immune_cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- thymus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thymus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Antibody. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antibody.jpg. **License**: *Public Domain: No Known Copyright*
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antigen_presentation.svg. **License**: *Public Domain: No Known Copyright*
- Lymphocyte_activation_simple.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immune_system#/media/File:Lymphocyte_activation_simple.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Red White Blood cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Red_White_Blood_cells.jpg. **License**: *Public Domain: No Known Copyright*
- Lymphocyte. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- B cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/B%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- natural killer (NK) cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/natural...20(NK)%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- T cells. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/T_cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- Antibody. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antibody.jpg. **License**: *Public Domain: No Known Copyright*
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Antigen_presentation.svg. **License**: *Public Domain: No Known Copyright*
- Lymphocyte_activation_simple.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immune_system#/media/File:Lymphocyte_activation_simple.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Red White Blood cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Red_White_Blood_cells.jpg. **License**: *Public Domain: No Known Copyright*
- Lymphocyte. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lymphocyte. **License**: *Public Domain: No Known Copyright*
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Antigen_presentation. **License**: *CC BY-SA: Attribution-ShareAlike*
- cytotoxic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cytotoxic. **License**: *CC BY-SA: Attribution-*

ShareAlike
- Antibody. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antibody.jpg**. **License**: ***Public Domain: No Known Copyright***
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antigen_presentation.svg**. **License**: ***Public Domain: No Known Copyright***
- Lymphocyte_activation_simple.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Immune_system#/media/File:Lymphocyte_activation_simple.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Red White Blood cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Red_White_Blood_cells.jpg**. **License**: ***Public Domain: No Known Copyright***
- Lymphocyte. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphocyte**. **License**: ***Public Domain: No Known Copyright***
- Antigen presentation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Antigen_presentation**. **License**: ***Public Domain: No Known Copyright***

This page titled 20.3E: Antigen-Presenting Cells is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.4: CYTOKINES

20.4A: Types of Cytokines Participating in Immune Response

This page titled 20.4: Cytokines is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.4A: Types of Cytokines Participating in Immune Response

Cytokines are small cell-signaling protein molecules secreted by numerous cells.

Learning Objectives

- Distinguish between the types of cytokines participating in immune response

Key Points

- Cytokines are immune system regulatory agents that work at both systemic and local levels. Cytokines are also involved in several developmental processes during embryogenesis.
- Each cytokine has a matching cell -surface receptor. Upon binding, intracellular signaling and gene expression regulations are altered, leading to the production of other cytokines, surface receptors, or feedback inhibition.
- Interleukins principally target leukocytes. They include common inflammatory and anti- inflammatory mediators and lymphokines.
- Chemokines mediate chemoattraction (chemotasis) between cells.
- Interferons have antiviral function and can act as a pyrogen.
- Tumor necrosis factor causes long-lasting inflammatory effects and fever during systemic immune responses. It stimulates the acute phase reaction in the liver and is responsible for much of the immune system-caused damage in severe infections.

Key Terms

- **chemokine**: Any of various cytokines produced during inflammation that organize the leukocytes by providing a stimuli for chemotaxis.
- **IL-10**: Also known as human cytokine synthesis inhibitory factor (CSIF), an anti-inflammatory cytokine.
- **cytokine**: Any of various small regulatory proteins or glycoproteins that regulate the cells of the immune system.

Cytokines are small cell-signaling protein molecules secreted by numerous cells and used extensively in intercellular communication. Cytokines can be classified as proteins, peptides, or glycoproteins. They provide the signaling pathways that orchestrate the complex immune responses of the human body. Cytokines are similar to hormones, which are also chemical messengers, but hormones have considerably more variation in molecular structure and are involved more in tissue signaling than cellular signaling.

Each cytokine has a matching cell-surface receptor. Subsequent cascades of intracellular signalling then alter cell functions. This may include the upregulation (increased expression) and/or downregulation (decreased expression) of several genes and their transcription factors resulting in the production of other cytokines, an increase in the number of surface receptors for other molecules, or the suppression of their own effects by feedback inhibition.

Interleukins



Interleukin: Crystallographic structure of human interleukin.

Interleukins are a class of cytokines primarily expressed by leukocytes. They are glycoproteins involved in the signaling of many types of immune system functions. There are 17 different families of interleukins. Some of the more important ones include inflammatory mediators such as IL-1, IL-4, and IL-6, the potent anti-inflammatory IL-10, and other interleukins involved with T and B cell signaling following antigen presentation. Many interleukins are also considered lymphokines, interleukins released by helper T cells to organize immune responses.

Interferons

Interferons are protein cytokines that have antiviral functions. They can activate macrophages and natural killer (NK) cells to attack and lyse virus-infected cells. One common interferon is IFN-gamma, a pyrogen involved in inflammatory response and macrophage and NK cell activation. IFN-gamma is produced by T cells (both CD4 and CD8) and NK cells.

Chemokines

Chemokines are protein cytokines that are mainly involved in facilitating chemotaxis (chemical-stimulated movement) in immune cells. Leukocytes travel along chemotactic gradients that guide them to sites of injury, infection, or inflammation. By definition, inflammatory mediators in other classes of cytokines are also considered chemokines. This category also includes cytokines that are only involved in leukocyte migration, such as CCL2 which causes monocyte chemotaxis and stimulates its differentiation into macrophages inside of tissues.

Tumor Necrosis Factor

Tumor necrosis factor (TNF) are cytokines that induce apoptosis in abnormal cells such as tumor cells. It is a protein released by NK cells, macrophages, and helper T cells, typically in systemic immune responses. TNF-alpha is the most notable example. This long-lasting inflammatory mediator and pyrogen can cause fever and inflammation for up to 24 hours. It also stimulates acute phase reaction in the liver, a component of systemic immune system activation where the liver makes proteins involved in immune system response such as complement proteins. TNF-alpha is released in very high amounts in response to lipopolysaccharide (infection with gram negative bacteria), which facilitates much of the self-destructive immune response in septic shock. In these cases, TNF-alpha can cause organ failure from tissue hypoperfusion, caused by damage and blood clotting from an overactive immune response.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cytokines. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cytokines**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lymphokines. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lymphokines**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Interleukins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Interleukins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cytokine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cytokine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chemokine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chemokine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- IL-10. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/IL-10**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Interleukins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Interleukins**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 20.4A: Types of Cytokines Participating in Immune Response is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.5: ANTIGENS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

20.5A: Antigens and Antigen Receptors
20.5B: Complete Antigens and Haptens
20.5C: Antigenic Determinants and Processing Pathways

This page titled 20.5: Antigens is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.5A: Antigens and Antigen Receptors

Antigens are molecules that initiate the immune response and can be bound by antibodies.

Learning Objectives

- Distinguish between antigens and antigen receptors

Key Points

- An antigen is a molecule that initiates the production of an antibody and causes an immune response.
- Antigens are typically proteins, peptides, or polysaccharides. Lipids and nucleic acids can combine with those molecules to form more complex antigens, like lipopolysaccharide, a potent bacterial toxin.
- An epitope is a molecular surface feature of an antigen that can be bound by an antibody. A paratope is the molecular surface feature of an antibody that binds to an epitope.
- Antigens are classified as exogenous (entering from outside) endogenous (generated within cells), an autoantigen, a tumor antigen, or a native antigen.
- Antigenic specificity is the ability of host cells to recognize an antigen by its unique molecular structure, such as the relationship between antigen epitopes and antibody paratopes.

Key Terms

- **antigen**: A substance that induces an immune response, usually foreign, but self antigens and internally produced antigens exist as well.
- **autoantigen**: Any antigen that stimulates auto antibodies in the organism that produced it. These are “self” antigens that are involved in autoimmune disease pathogenesis.

EXAMPLES

Fluorescein, along with other haptens such as biotin, is used in various cell and molecular biological techniques. Fluorescein is often conjugated to a protein to allow scientists to examine its location using a fluorescent microscope.

In immunology, an antigen is a substance that evokes an immune response. Formally they are defined as a substance that causes the production of antibodies specific to that antigen, however they also cause T cell mediated immune responses, and may lead to an inflammatory response. The substance may be from the external environment or formed within the body. The immune system will try to destroy or neutralize any antigen that is recognized as a foreign and potentially harmful invader.”Self” antigens are usually tolerated by the immune system; whereas “non-self” antigens can be identified as invaders and can be attacked by the immune system.

Molecular Structure of Antigens

At the molecular level, an antigen is characterized by its ability to be “bound” at the antigen-binding site of an antibody. Antibodies tend to discriminate between the specific molecular structures presented on the surface of the antigen. Antigens are usually either proteins, peptides, or polysaccharides. This includes parts (coats, capsules, cell walls, flagella, fimbrae, and toxins) of bacteria, viruses, and other microorganisms. Lipids and nucleic acids are antigenic only when combined with proteins and polysaccharides. For example, the combination of lipids and polysaccharides are lipopolysaccharides (LPS), which are the primary component of gram negative bacterial endotoxin. LPS forms the cell wall of gram negative bacteria and causes a powerful immune response when bound. Cells present their immunogenic-antigens to the immune system via a major histocompatibility (MHC) molecule. Depending on the antigen presented and the type of the histocompatibility molecule, several types of immune cells can become activated due to an antigen.

Antigens have several structural components of interaction that may be bound by different classes of antibodies. Each of these distinct structural components is considered to be an epitope, also called an antigenic determinent. Therefore, most antigens have the potential to be bound by several distinct antibodies, each of which is specific to a particular epitope. The antigen binding receptor on an antibody is called a paratope, and is specific to the epitope of the antigen. Using the “lock and key” metaphor, the antigen itself can be seen as a string of keys – any epitope being a “key” – each of which can match a different lock.

Types of Antigens

Antigens are categorized into broad classes of antigens based on their origin. So many different molecules can function as an antigen in the body, and there is considerable diversity even within these categories.

These are the main classes of antigens that are involved in immune system activation. Their diversity is analogous to the immense diversity of the diseases that the immune system works to overcome.

Exogenous Antigens

Exogenous antigens are antigens that have entered the body from the outside, for example by inhalation, ingestion, or injection. Exogenous antigens are the most common kinds of antigens, and includes pollen or foods that may cause allergies, as well as the molecular components of bacteria and other pathogens that could cause an infection.

Endogenous Antigens

Endogenous antigens are that have been generated within previously-normal cells as a result of normal cell metabolism or because of viral or intracellular bacterial infection (which both change cells from the inside in order to reproduce). The fragments are then presented on the surface of the infected cells in the complex with MHC class I molecules.

Autoantigens

Autoantigens are normal “self” protein or complex of proteins or nucleic acid that is attacked by the host’s immune system, causing an autoimmune disease. These antigens should, under normal conditions, not be the target of the immune system, but due to mainly genetic and environmental factors, the normal immunological tolerance for such an antigen has been lost.

Tumor Antigens (Neoantigens)

These antigens are presented by MHC I or MHC II molecules on the surface of tumor cells.These antigens result from a tumor-specific mutation during malignant transformation of normal cells into cancer cells. Despite expressing this antigen, many tumors have developed ways to evade antigen recognition and immune system killing.

Native Antigens

A native antigen is an antigen that is not yet processed by an APC to smaller parts. T cells cannot bind native antigens, but require that they be digested and processed by APCs, whereas B cells can be activated by native ones without prior processing.

This page titled 20.5A: Antigens and Antigen Receptors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.5B: Complete Antigens and Haptens

Haptens are molecules that create an immune response when attached to proteins.

Learning Objectives

- Describe haptens and complete antigens

Key Points

- Haptens are incomplete antigens that do not cause an immune response upon binding because they cannot bind to MHC complexes.
- Haptens may bind with a carrier protein to form an adduct, which is also a complete antigen.
- While haptens don't directly cause immune responses, they may sensitize the body towards hypersensitivity and autoimmune responses.
- Haptens may inhibit antibody immune responses by binding with antibodies in place of the actual antigen until there aren't enough antibodies left to bind to the complete antigen.

Key Terms


- **adduct**: A complex molecule formed by the combination of two or more molecules, such as a complete antigen created by a hapten and a carrier.
- **hapten**: Any small molecule that can elicit an immune response only when attached to a large carrier such as a protein.

Antigens are basic molecules that induce an immune response when detected by immune system cells. Antigens may be either complete or incomplete based on the nuances of their molecule structure.

Haptens

A hapten is essentially an incomplete antigen. These small molecules can elicit an immune response only when attached to a large carrier such as a protein; the carrier typically does not illicit an immune response by itself. Many hapten carriers are normal molecules that circulate through the body. When haptens and carriers combine, the resulting molecule is called an adduct, the combination of two or more molecules. Haptens cannot independently bind to MHC complexes, so they cannot be presented to T cells.

The first haptens used were aniline and its carboxyl derivatives (o-, m-, and p-aminobenzoic acid). One well-known hapten is urushiol, the toxin found in poison ivy and a common cause of cell-mediated contact dermatitis. When absorbed through the skin from a poison ivy plant, urushiol undergoes oxidation in the skin cells to generate the actual hapten, a reactive molecule called a quinone, which then reacts with skin proteins to form hapten adducts. Usually, the first exposure causes only sensitization, in which there is a proliferation of helper and cytotoxic T cells. After a second exposure, the proliferated T cells can become activated, generating an immune reaction and producing the characteristic blisters of poison ivy exposure.



Fluorescein Molecule: Fluorescein is an example of a hapten used in molecular biology.

Some haptens induce autoimmune disease. An example is hydralazine, a blood pressure-lowering drug that occasionally causes lupus erythematosus (an autoimmune inflammatory disorder) in certain individuals with genetic predispositions to the disease. This also appears to be the mechanism by which the anesthetic gas halothane can cause life-threatening hepatitis and penicillin-class drugs cause autoimmune hemolytic anemia. Other haptens, such as flourescein, detect proteins with which they form adducts. This makes them a common part of molecular biology lab techniques.

Complete Antigens

A complete antigen is essentially a hapten-carrier adduct. Once the body has generated antibodies to a hapten-carrier adduct, the small-molecule hapten may also be able to bind to the antibody, but will usually not initiate an immune response. In most cases this can only be elicited by theonly the hapten-carrier adduct. Sometimes the small-molecule hapten can block immune response to the complete antigen by preventing the adduct from binding to the antibody, a process called hapten inhibition. In this case, the hapten acts as the epitope for the antigen, which binds to the antibodies without causing a response. If this happens with enough haptens, there will not be enough antibodies left to bind to the complete antigen, thus inhibiting the antibody response.

This page titled 20.5B: Complete Antigens and Haptens is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.5C: Antigenic Determinants and Processing Pathways

Antigen epitopes make it possible for the immune system to recognize pathogens.

Learning Objectives

- Describe antigenic determinants and pathways of processing

Key Points

- An epitope (also known as an antigenic determinant) is part of an antigen that is recognized by the immune system, specifically by antibodies and B and T cells. Other immune cells like APCs cannot recognize epitopes (only PAMPS and DAMPS).
- Antigenic determinants (epitopes) are divided into conformational epitopes and linear epitopes.
- Antigen processing occurs within a cell and results in fragmentation of proteins, association of the fragments with MHC molecules, and expression of the peptide -MHC molecules at the cell surface where they can be recognized by the T cell receptor on a T cell.
- Antigen processing may be done through either the endogenous pathway (viral proteins from within an infected cell) or through the exogenous pathway (engulfing a pathogen and isolating its antigen from within the APC).
- The endogenous pathway uses MHC class I and binds to cytotoxic T cells, while the exogenous pathway uses MHC class II and binds to helper T cells.
- Some viruses can prevent antigen processing by disrupting movement of MHC within the cell.

Key Terms

- **Linear epitopes**: These consist of the primary amino acid structure of a protein that makes up the larger antigen.
- **The Exogenous Pathway**: Phagocytized pathogens are broken down from within the cell and their broken-down antigens bind with MHC II, which then is expressed on the surface of the antigen-presenting cell.

An epitope, also known as an antigenic determinant, is the part of an antigen that is recognized by the immune system, specifically by antibodies, B cells, and T cells. The latter can use epitopes to distinguish between different antigens, and only binds to their specific antigen. In antibodies, the binding site for an epitope is called a paratope. Although epitopes are usually derived from non-self proteins, sequences derived from the host that can be recognized are also classified as epitopes. Epitopes determine how antigen binding and antigen presentation occur.

Types of Antigenic Determinants


The epitopes of protein antigens are divided into two categories based on their structures and interaction with the paratope.

- A conformational epitope is composed of discontinuous sections of the antigen's amino acid sequence. These epitopes interact with the paratope based on the 3-D surface features and tertiary structure (overall shape) of the antigen. Most epitopes are conformational.
- Linear epitopes interact with the paratope based on their primary structure (shape of the protein's components). A linear epitope is formed by a continuous sequence of amino acids from the antigen, which creates a "line" of sorts that builds the protein structure.

Antigenic determinants recognized by B cells and the antibodies secreted by B cells can be either conformational or linear epitopes. Antigenic determinants recognized by T cells are typically linear epitopes. T cells do not recognize polysaccharide or nucleic acid antigens. This is why polysaccharides are generally T-independent antigens and proteins are generally T-dependent antigens. The determinants need not be located on the exposed surface of the antigen in its original form, since recognition of the determinant by T cells requires that the antigen be first processed by antigen presenting cells. Free peptides flowing through the body are not recognized by T cells, but the peptides associate with molecules coded for by the major histocompatibility complex

(MHC). This combination of MHC molecules and peptide is recognized by T cells.

Antigen-Processing Pathways



Antigen-Binding Site of an Antibody: Antigen-binding sites can recognize different epitopes on an antigen.

In order for an antigen-presenting cell (APC) to present an antigen to a naive T cell, it must first be processed so itacan be recognized by the T cell receptor. This occurs within an APC that phagocytizes an antigen and then digests it through fragmentation (proteolysis) of the antigen protein, association of the fragments with MHC molecules, and expression of the peptide-MHC molecules at the cell surface. There, they are recognized by the T cell receptor on a T cell during antigen presentation. MHC molecules must move between the cell membrane and cytoplasm in order for antigen processing to occur properly. However, the pathway leading to the association of protein fragments with MHC molecules differs between class I and class II MHC, which are presented to cytotoxic or helper T cells respectively. There are two different pathways for antigen processing:

- The endogenous pathway occurs when MHC class I molecules present antigens derived from intracellular (endogenous) proteins in the cytoplasm, such as the proteins produced within virus-infected cells. Generally, proteosomes are used to break up the viral proteins and combine them with MHC I.
- The exogenous pathway occurs when MHC class II molecules present fragments derived from extracellular (exogenous) proteins that are located within the cell. First, pathogens are phagocytized, then endosomes within the cell break down antigens with proteases, which then combine with MHC II.

Some viral pathogens have developed ways to evade antigen processing. For example, cytomegalovirus and HIV-infected cells sometimes disrupt MHC movement through the cytoplasm, which may prevent them from binding to antigens or from moving back to the cell membrane after binding with an antigen.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Antigen. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Antigen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- antigen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/antigen**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- autoantigen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/autoantigen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hapten. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hapten**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hapten. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hapten**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fluorescein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fluorescein**. **License**: ***Public Domain: No Known Copyright***
- Antigenic determinants. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Antigenic_determinants**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T cell epitopes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/T%20cell%20epitopes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- major histocompatibility complex. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/major+...bility+complex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fluorescein. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fluorescein**. **License**: ***Public Domain: No Known Copyright***
- Antigen. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Antigen**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 20.5C: Antigenic Determinants and Processing Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6: HUMORAL IMMUNE RESPONSE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

20.6A: Clonal Selection and B-Cell Differentiation

20.6B: Structure and Function of Antibodies

20.6C: Role of the Complement System in Immunity

20.6D: Immunological Memory

20.6E: Major Histocompatibility Complex Antigens (Self-Antigens)

This page titled 20.6: Humoral Immune Response is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6A: Clonal Selection and B-Cell Differentiation

B cells mature in the bone marrow, where they undergo VDJ recombination to produce unique receptors that do not react to self-antigens.

Learning Objectives

- Describe clonal selection and B cell differentiation

Key Points

- In the bone marrow, central tolerance of B cells is produced through negative selection. Immature B cells are tested for auto-reactivity before leaving the bone marrow. The immature B cells whose receptors (BCRs) bind too strongly to self antigens will not be killed.
- When the B cell receptor on the cell surface matches its cognate antigen in the body, the B cell proliferates and secretes a free form of those receptors (antibodies), with identical binding sites as on the original cell surface.
- B cells that have not been exposed to antigen, also known as naïve B cells, can be activated in a T cell-dependent or independent manner.
- There are two types of T cell independent activation: Type 1 T cell-independent (polyclonal) activation, and type 2 T cell-independent activation.
- During clonal selection, random mutations during clonal expansion cause the production of B cells with increased antibody-binding affinity for their antigens.
- The clonal selection hypothesis may explain why secondary immune responses are so effective at preventing reinfection by the same pathogen.

Key Terms

- **Clonal selection**: The theory that lymphocytes bear antigen receptors before activation and that random mutations during clonal expansion cause the development of lymphocytes with high binding affinities for their antigens.
- **humoral**: Of or relating to the body fluids or humors.
- **memory B cell**: A B cell subtype formed following primary infection in which the cell recognizes a specific epitope.

B cells are lymphocytes that play a large role in the humoral immune response (as opposed to the cell-mediated immune response, which is governed by T cells). B cells primarily function to make antibodies against antigens, act as antigen-presenting cells (APCs), and eventually develop into memory B cells to provide long-term immunity. B cells undergo clonal selection and develop similarly to T cells with some notable differences.

B Cell Development


Immature B cells are produced in the bone marrow of most mammals. Their development occurs through several stages, each representing a change in the genome content at the antibody loci. An antibody is composed of two identical light (L) and two identical heavy (H) chains, and the genes specifying them are found in the V (variable) region and C (constant) region. The heavy-chain V region has three segments, V, D and J. These segments recombine randomly in a process called VDJ recombination to produce a unique variable domain in the immunoglobulin of each individual B cell.

Similar rearrangements occur for the light-chain V region but with only two segments involved: V and J. When the B cell fails in any step of the maturation process, it will die by apoptosis, here called clonal deletion. This is a form of positive selection. B cells are also tested for autoreactivity through negative selection. If these B cells have high affinity for binding to self-antigens, they will die by clonal deletion or another pathway such as anergy.

B Cell Activation

B cell activation refers to the differentiation and clonal expansion of B cells.When the B cell receptor on the cell surface matches its cognate antigen in the body, the B cell proliferates and secretes a free form of those receptors (antibodies) in the body, with binding sites identical to those on the original cell surface. However, B cell recognition of antigens is not the only element necessary for B cell activation. B cells that have not been exposed to antigen, also known as naïve B cells, can be activated in a T cell-dependent or independent manner.

- T cell dependent activation is activation of B cells by type 2 helper T cells in the lymph nodes.
- T cell independent activation occurs when antigens directly bind to B cell themselves, usually through cross-linking the antigen to the B cell receptor or receiving the antigen with a toll-like receptor.



T-dependent B cell activation: T cell-dependent B cell activation, showing a TH2-cell (left), B cell (right), and several interaction molecules.

B Cell Differentiation and Clonal Expansion

After activation, the B cell undergoes differentiation and clonal expansion, which usually involves migration to germinal centers if the activation takes place in a lymph node. B cell differentiation is the process by which B cells change into different types, such as plasma cells and plasma blasts. Clonal expansion is the process by which daughter cells arise from a parent cell. During B cell clonal expansion, many copies of that B cell are produced that share affinity with and specificity of the same antigen.

Clonal Selection

Clonal selection is a theory stating that B cells express antigen-specific receptors before antigens are ever encountered in the body. After B cell activation, the B cells clone themselves through clonal expansion, but during each cellular division, random mutations occur that gradually increase the binding affinity for B cell-produced antibodies to antigens.

For example, memory B cells that differentiate after an adaptive immune response are thought to undergo clonal selection so that antibodies produced by newer memory B cells have considerably higher binding affinities to their antigens. This theory may explain why secondary immune responses from memory cells are so effective that repeated infections by the same pathogen are stopped before symptoms even develop. Following the initial infection, random mutations during clonal selection could produce memory B cells that can more easily bind to antigens than can the original B cells.

This page titled 20.6A: Clonal Selection and B-Cell Differentiation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6B: Structure and Function of Antibodies

An antibody is a Y-shaped protein produced by B cells to identify and neutralize antigens in the body.

Learning Objectives

- Describe antibody structure and function

Key Points

- An antibody, also known as an immunoglobulin, is a large Y-shaped protein produced by B- cells and used by the immune system to identify and neutralize foreign objects such as bacteria and viruses.
- Each tip of the "Y" of an antibody contains a paratope (a structure analogous to a lock) that is specific for one particular epitope (similarly analogous to a key) on an antigen, allowing these two structures to bind together with precision.
- Though the general structure of antibodies is very similar, a small region at the tip of the protein is extremely variable, allowing millions of antibodies with different antigen binding sites to exist. This region is known as the hypervariable region.
- Five isotypes of antibodies are found in different locations and perform different specific functions.
- The base of the Y plays a role in modulating immune cell activity. This region is called the Fc region, and phagocytes may bind to it to initiate phagocytosis.
- Antibodies that bind to surface antigens on a bacterium attract the first component of the complement cascade with their Fc region and initiate activation of the classical complement system.


Key Terms

- **epitope**: Part of a biomolecule (such as a protein) that is the target of an immune response.
- **paratope**: Part of the molecule of an antibody that binds to an antigen.
- **isotype**: A marker corresponding to an antigen found in all members of a subclass of a specific class of immunoglobulins.

An antibody (formally called immunoglobulin) is a large Y-shaped glycoprotein produced by B-cells and used by the immune system to identify and neutralize pathogens. Antibodies are produced by B cells, and are either secreted into circulation or remain expressed on the surface of the B cell.

Structure of Antibodies

The antibody recognizes a unique part of an antigen (foreign object). Each tip of the "Y" of an antibody contains a paratope (a structure analogous to a lock) that is specific for one particular epitope (similarly analogous to a key) on an antigen, allowing these two structures to bind together with precision. Using this binding mechanism, an antibody can neutralize its target directly or tag it for attack by other parts of the immune system.



Antibody: Each antibody binds to a specific antigen, an interaction similar to a lock and key.

Antibodies are glycoproteins belonging to the immunoglobulin superfamily, typically made of basic structural units each with two large heavy chains and two small light chains. Most antibodies exist as a monomer, in which they have a single “Y” shaped sub-unit, but some antibodies can exist as dimers (two subunits) or pentamers (five subunits). The paratope is considered a hypervariable region and has the same specificity and antigen-binding affinity as the B cell receptor of the B cell that created the antibody. In some isotypes, the tail end of the antibody is called the constant region and faces away from the “Y-shaped” paratobe ends, functioning as an Fc tail to which phagocytes can bind.

Isotypes

Five different isotypes of antibodies each perform different functions and are generally found in different parts of the body.

- IgA: A dimer secreted into mucosal surfaces, such as the gut, respiratory tract, and urogenital tract, that prevents mucosal invasion into the body by pathogens. It is resistant to the proteolytic enzymes found in the gastrointestinal mucosae.
- IgD: Functions mainly as an antigen receptor on B cells that have not been exposed to antigens. It has been shown to activate basophils and mast cells to produce antimicrobial factors.
- IgE: Found in circulation and binds to allergens, triggering histamine release from mast cells and basophils. Also protects against parasitic worms.
- IgG: Has four different forms and provides the majority of antibody-based immunity against invading pathogens as the best opsonin of any type of antibody. This is because it expresses a tail for Fc receptors on phagocytes to bind to, which activates phagocytosis. It is the only antibody capable of crossing the placenta to give passive immunity to fetus, and can activate the classical complement system.
- IgM: Expressed on the surface of B cells (monomer) and in a secreted pentamer with very high avidity. Eliminates pathogens in the early stages of B cell-mediated (humoral) immunity before there is sufficient IgG. Like IgG, it can also activate the classical complement system.

Function of Antibodies

Circulating antibodies are produced by clonal B cells that specifically respond to only one antigen. Antibodies contribute to immunity in three ways: preventing pathogens from entering or damaging cells by binding to them (neutralization); stimulating removal of pathogens by macrophages and other cells by coating the pathogen (opsonization); and triggering destruction of pathogens by stimulating other immune responses such as the complement pathway. The complement system starts a long cascade of protein productions that either opsonize a pathogen for phagocytosis or lyse it directly by forming a membrane attack complex. During opsonization, the antibody expresses the tail for an Fc receptor on a macrophage, neutrophil, or natural killer cell. The immune cell will then bind to the antibody’s Fc tail instead of the pathogen itself, which speeds up the process of finding pathogens to phagocytize. Additionally, because antibodies have two or more paratopes, they can sometimes link pathogens

together, making phagocytosis more efficient.

This page titled 20.6B: Structure and Function of Antibodies is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6C: Role of the Complement System in Immunity

The complement system is the ability of antibodies and phagocytic cells to remove pathogens from an organism.

Learning Objectives

- Describe the role of the complement system in immunity

Key Points

- The complement system helps antibodies and phagocytic cells clear pathogens from an organism.
- The complement system consists of a number of small proteins produced by the acute phase reaction in the liver during inflammation.
- The complement system might play a role in diseases with an immune component and those of the central nervous system. Complement protein deficiency is a form of primary immunodeficiency.
- The classical complement pathway starts with antibody binding, which causes a cascade reaction of complement proteins that gradually form a membrane attack complex.
- The alternative complement pathway is usually stimulated by pathogen antigens or toxins rather than antibodies, and cleaves C3 until there is enough to continue the steps of the classical complement pathway from the C5 convertase step.
- The lectin pathway is homologous to the classical pathway, but with the opsonin, mannose-binding lectin (MBL), and ficolins, instead of C1 from the antibody. This pathway uses proteases on the MBL to form C3 convertase, which continues the steps of the classical complement pathway from the C3 convertase step.
- The complement system is regulated by complement control proteins, such as decay accelerating pathway, which prevent complement proteins from forming MAC on the body's cells.

Key Terms

- **C5a**: A complement protein that is an acute phase inflammatory mediator, causing vasodilation and neutrophil chemotaxis.
- **membrane attack complex**: The final complex of all complement system pathways that lyses the pathogen. It is composed of C5b, C6, C7, C8, and C9.
- **mannan-binding-lectin**: A protein that binds to carbohydrates on pathogens to activate the lectin complement pathway.

The complement system or "complements" the ability of antibodies and phagocytic cells to clear pathogens from an organism. The complement system consists of a number of small proteins found in the blood, generally synthesized by the liver as a part of the acute phase reaction during systemic inflammation (from TNF-alpha release). When stimulated by one of several triggers, proteases in the system cleave specific proteins to release cytokines and initiate an amplifying cascade of further cleavages. The end result of this activation cascade is massive amplification of the response and activation of the cell-killing membrane attack complex (also called MAC). There are three different pathways by which the complement system may occur.


Classical Complement Pathway

The classical complement pathway is the main pathway by which the complement system occurs. It is comprised of a cascade of many steps with complement proteins cleaving one another in a sequential order:

1. The antibody binds to an antigen on the surface of a pathogen, activating the C1 complement protein.
2. C1 acts a protease and cleaves C2 and C4 to form C4b2b.
3. C42b converts C3 into C3a and C3b, which forms a C5 convertase.
4. C5 convertase cleaves C5 into C5a and C5b.
5. C5b forms a complex with C6, C7, and then C8, and C9, which becomes the membrane attack complex that lyses the pathogen.

Note that C5a has a number of other functions in the immune system, such as causing vasodilation during inflammation and

stimulating neutrophil chemotaxis. Additionally, the body's cells express a glycoprotein called decay accelerating factor, which decays C3 and C5 convertase on the body's cells. This factor prevents membrane attack complexes from forming on the body's cells under normal conditions.



The Classical and Alternative Complement Pathways: The classical and alternative complement pathways start off differently, but end in the same cascade of complement proteins that combine to form a membrane attack complex.

The Alternative Complement Pathway

The alternative pathway may be a leftover evolutionary precursor to the classical pathway. Unlike the classical pathway, the alternative pathway is generally activated by microbial inflammatory mediators instead of antibodies. For example, lipopolysaccharide, the toxin of gram-negative bacteria, may activate this pathway. The steps for the alternative pathway are:

1. The pathogenic antigen (such as LPS) activates C3 so it creates a C3B complex
2. Factor D cleaves the C3B complex so that C3bBb is created.
3. C3bBb is a C3 convertase, which converts more C3 into C3a and C3b.
4. Similarly to the classical pathway, C3b forms a C42b complex, and the rest of the steps are essentially the same as the classical pathway, ending with C5b forming a membrane attack complex with C6, C7, C8, and C9.

Lectin Pathway

The lectin pathway is not caused by antibody binding, but by a carbohydrate -binding-protein called mannan-binding-lectin (MBL). It is an acute phase reactant produced in the liver and binds to the carbohydrates on the surfaces of many pathogens. The steps for the lectin pathway are:

1. MBL binds to the carbohydrates on a pathogen.
2. Proteases bound on the other side of the MBL cleaves C4 into C4a and C4b.
3. C4b creates C3 convertase, and the rest of the steps happen identically to the classical pathway from the C3 convertase step.

Problems with the Complement System

The complement system might play a role in diseases with an immune component, such as Barraquer-Simons Syndrome, asthma, lupus erythematosus, glomerulonephritis, various forms of arthritis, autoimmune heart disease, multiple sclerosis, inflammatory bowel disease, ischemia-reperfusion injuries, and rejection of transplanted organs. The complement system is also becoming increasingly implicated in diseases of the central nervous system such as Alzheimer's disease and other neurodegenerative

conditions such as spinal cord injuries. Additionally, deficiencies in complement proteins produced in the liver can lead to a form of primary (congenital) immunodeficiency, in which the body is more susceptible to disease, particularly autoimmune diseases and severe bacterial infections.

This page titled 20.6C: Role of the Complement System in Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6D: Immunological Memory

Immunological memory refers to the ability of B and T cells to produce long-lived memory cells that defend against specific pathogens.

Learning Objectives

- Describe immunological memory of the immune system


Key Points

- When B and T cells begin to replicate, some offspring will become long-lived memory cells.
- Memory cells remember all specific pathogens encountered during the animal's lifetime and can thus call forth a strong response if the pathogen ever invades the body again.
- Passive immunity comes from IgG antibodies given through the mother during fetal development and through breast milk. This memory is short term, but protects the infant until its own adaptive immune system is functional.
- During a secondary immune response, memory B and T cells work to rapidly eliminate the pathogen, preventing reinfection by the same pathogen.
- During a vaccination, the antigen of a pathogen is introduced into the body through a weakened form of the pathogen that cannot cause an infection. This stimulates the immune system to develop a specific immunity against that pathogen without actually causing the disease that the pathogen brings.
- Vaccines do not exist for every pathogen due to frequent strain mutations and challenges in producing an immunization strong enough to work, but not strong enough to cause an infection.

Key Terms

- **secondary immune response**: The act of exposure to the same pathogen after the initial immune response. Memory B and T cells work to rapidly eliminate the pathogen to prevent reinfection.
- **vaccination**: Inoculation with the weakened form of a pathogen to protect against a particular disease or strain of disease by stimulating the development of immunological memory against that pathogen.

When B and T cells begin to replicate during an adaptive immune response, some offspring become long-lived memory cells. These memory cells remember all specific pathogens encountered during the animal's lifetime and can thus call forth a stronger response, called the secondary immune response, if the pathogen ever invades the body again. The adaptive immune system is so-named because it is a result of an adaptation to an infection. Immunological memory can either exist in active long-term memory or passive short-term memory.



Immune response: When B and T cells begin to replicate, some of the offspring that they produce will end up becoming long-lived memory cells. These memory cells will remember all specific pathogens encountered during the animal's lifetime and can thus call forth a strong response if the pathogen ever invades the body again.

Passive Memory

Newborn infants are particularly vulnerable to infections since they have no prior exposure to pathogens. Thus, the mother protects the infant through several layers of passive protection. During pregnancy, IgG, a certain isotype of antibody, is transported to the baby from the mother through the placenta, so even babies have high levels of antibodies with similar antigen specificities as the mother. Even breast milk contains antibodies that are transferred to the infant's gastrointestinal tract and protect against bacterial infections until the baby is capable of making its own antibodies. Since the fetus isn't making any memory cells or antibodies, this is called passive immunity. Passive immunity is short-lived, ranging from a couple days to a couple months.

As the infant matures, their thymus and bone marrow work to raise a stock of mature lymphocytes that form the foundation for the infant's personal adaptive immune system. Because the passive memory comes from antibodies instead of B cells themselves, infants do not inherit long-term immunological memory from the mother. Even if the infant receives antibodies specific to certain diseases from its mother, the infant wouldn't be able to bolster a long-term memory that would direct antigen exposure and presentation.

Active Memory and Immunization

Following an infection, long-term active memory is acquired by activation of B and T cells. Memory cells derive from their parent B and T cells, and undergo clonal selection following infection, which increases antigen-binding affinity. Following reinfection, the secondary immune response typically eliminates the pathogen before symptoms of an infection can occur. During the secondary immune response, memory T cells rapidly proliferate into active helper and cytotoxic T cells specific to that antigen, while memory B cells rapidly produce antibodies to neutralize the pathogen. Long-term active memory consists of rapid response and form permanent immunological memory so long as those memory cells survive.

Vaccinations take advantage of memory lymphocyte development by artificially-generating active immunity, a process called immunization. During a vaccination, the antigen of a pathogen is introduced into the body and stimulates the immune system to develop a specific immunity against that pathogen. It doesn't cause the disease that the pathogen brings because the vaccine uses an attenuated form of the pathogen that contains the same antigen but doesn't have the capacity for replication. This deliberate introduction of the pathogen is successful since it exploits the immune system's natural specificity and inducibility. Vaccination is an extremely effective manipulation of the immune system that helps fight diseases. Over the course of vaccine development, they have saved countless lives, and diseases like rubella and polio are not the widespread causes of disability they once were.

Despite the effectiveness of vaccines, methods do not yet exist to develop vaccines for every pathogen. Many pathogens undergo mutations that change the expression of their antigens, making immunization attempts fruitless for diseases like the common cold or norovirus. Many parasitic pathogens, such as the *plasmodium* protist that causes malaria, haven't successfully been vaccinated against because it is challenging to develop a vaccine that is strong enough to stimulate an immune response (sufficient immunogenicity) without causing a live infection.

This page titled 20.6D: Immunological Memory is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.6E: Major Histocompatibility Complex Antigens (Self-Antigens)

The major histocompatibility complex (MHC) is a cell surface molecule that regulates interactions between white blood cells and other cells.

Learning Objectives

- Describe major histocompatibility complex antigens

Key Points

- Major histocompatibility complex (MHC) is a cell -surface molecule that mediates interactions between white blood cells and other leukocytes or cells.
- The MHC is divided into class I, class II, and class III subgroups, all encoded by the same gene.
- MHC antigen presentation gains its diversity from the high degree of polymorphism of the MHC protein itself.
- Differing HLA between members of the same species is the main reason why transplanted organs are rejected by the organ recipient.

Key Terms

- **human leukocyte antigen (HLA)**: The name of the major histocompatibility complex (MHC) in humans.


Major histocompatibility complex (MHC) is a cell-surface molecule encoded by a large gene family in all vertebrates. MHC molecules mediate interactions of leukocytes with other leukocytes or body cells. MHC determines compatibility of donors for organ transplant as well as one's susceptibility to an autoimmune disease via cross-reacting immunization. In humans, MHC is also called human leukocyte antigen (HLA), because MHC can often act as an antigen for human leukocytes.

Types of MHC


MHC is a polymorphic protein attached to the surface of cells. Three different classes of MHC differ in structure and function in the immune system, but are all encoded by the same set of gene sequences.

- MHC I is presented on all cells of the body. It contains an epitope that forms the structural binding site for an antigen. MHC I interacts with natural killer (NK) cells and cytotoxic T cells to signal whether a cell is self or non-self, and whether it contains an antigen specific to that T cell.
- MHC II is presented mainly on macrophages, dendritic cells, and helper T cells, which are all involved in antigen presentation. It has a longer helical region than MHC I, which allows it to bind to CD4 (helper T cells) during antigen presentation.
- MHC III is a secreted enzyme that is neither membrane-bound nor involved in antigen presentation like MHC I and II. It is merely included as an MHC protein because it is encoded by the same set of genes. It is involved in production of complement proteins and inflammatory cytokines.


During antigen processing prior to presentation, protein peptides from pathogens and MHC (I or II) travel through the cytoplasm and then to the cell membrane surface. A complex series of vesicles enables MHC transport from the cell, and the endoplasmic reticulum and Golgi bodies facilitate this transport during antigen processing. After processing, the MHC can present the peptide antigen bound to it to naive T cells.



MHC Class I processing: MHC class I pathway: proteins in the cytosol are degraded by the proteasome, liberating peptides internalized by TAP channel in the endoplasmic reticulum, there associating with MHC-I molecules freshly synthesized. MHC-I/ peptide complexes enter Golgi apparatus, are glycosylated, enter secratory vesicles, fuse with the cell membrane, and externalize on the cell membrane interacting with T lymphocytes.



MHC Class I: MHC class I protein molecule.



MHC Class II: MHC class II protein molecule.

HLA and Organ Rejection

Organ transplantation is a complex procedure that can potentially cure many chronic diseases and acute injuries. However, surgically-replaced organs are often rejected by the body's immune system. MHC is also called human leukocyte antigen (HLA) and varies considerably among different members of the same species. If the T and B cells of the body recognize the HLA of the graft as foreign, they will attack the organ graft. The damage in organ rejection can be acute or chronic, cell-mediated or antibody -mediated, and often involves diffuse damage of the graft that cause necrosis and infarction (tissue death from lack of oxygen) to the graft tissue by attacking its vascular components.

In nearly all cases, immunosuppressive chemotherapy is a requirement for successful organ transplantation. These drugs can stop acute organ rejection after the procedure, but will not stop chronic organ rejection, in which gradual vascular lesions and endothelial thickening slowly kill the graft. If an organ donor has HLA similar to that of the recipient, the risk of organ rejection is reduced. However this isn't feasible for heart, liver, or lung transplants because there generally isn't enough time in these cases to find a matching organ donor.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- B cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/B_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clonal selection. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clonal_selection. **License**: *CC BY-SA: Attribution-ShareAlike*
- apoptosis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/apoptosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- memory B cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/memory%20B%20cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- humoral. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/humoral. **License**: *CC BY-SA: Attribution-ShareAlike*
- T-dependent B cell activation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:T-dependent_B_cell_activation.png. **License**:
- Antibodies. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Antibodies. **License**: *CC BY-SA: Attribution-*

ShareAlike
- epitope. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/epitope**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- paratope. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/paratope**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- isotype. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/isotype**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T-dependent B cell activation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:T-dependent_B_cell_activation.png**. **License**:
- Antibody. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antibody.svg**. **License**:
- complement system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/complement_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Complement system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Complement_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- innate immune system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/innate_immune_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T-dependent B cell activation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:T-dependent_B_cell_activation.png**. **License**:
- Antibody. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antibody.svg**. **License**:
- Complement_pathway.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Complement_component_4#/media/File:Complement_pathway.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Structural Biochemistry/Genetic code/Immune System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structural_Biochemistry/Genetic_code/Immune_System%23Immunological_Memory**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vaccination. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vaccination**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T-dependent B cell activation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:T-dependent_B_cell_activation.png**. **License**:
- Antibody. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antibody.svg**. **License**:
- Complement_pathway.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Complement_component_4#/media/File:Complement_pathway.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Immune response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Immune_response.jpg**. **License**:
- Major histocompatibility complex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Major_histocompatibility_complex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- human leukocyte antigen (HLA). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/human%2...ntigen%20(HLA)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T-dependent B cell activation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:T-dependent_B_cell_activation.png**. **License**:
- Antibody. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Antibody.svg**. **License**:
- Complement_pathway.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Complement_component_4#/media/File:Complement_pathway.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Immune response. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Immune_response.jpg**. **License**:
- MHC Class 1. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MHC_Class_1.svg**. **License**:
- MHC Class I processing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MHC_Class_I_processing.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- MHC Class 2. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MHC_Class_2.svg**. **License**:

This page titled 20.6E: Major Histocompatibility Complex Antigens (Self-Antigens) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.7: CELL-MEDIATED IMMUNE RESPONSE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

20.7A: Clonal Selection and T-Cell Differentiation

20.7B: Specific T-Cell Roles

20.7C: Active and Passive Humoral Immunity

This page titled 20.7: Cell-Mediated Immune Response is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.7A: Clonal Selection and T-Cell Differentiation

Antigens are selected to form clones of themselves, both memory and effector.

Learning Objectives

- Describe clonal selection and T cell differentiation

Key Points

- All T cells originate from hematopoietic stem cells in the bone marrow and generate a large population of immature thymocytes. The thymocytes progress from double negative cells to become double-positive thymocytes (CD4+CD8+), and finally mature to single-positive (CD4+CD8- or CD4-CD8+).
- Clonal selection is used during negative selection to destroy lymphocytes that may be able to bind with self antigens.
- Clonal selection is the theory that specific antigen receptors exist on lymphocytes before they are presented with an antigen due to random mutations during initial maturation and proliferation. After antigen presentation, selected lymphocytes undergo clonal expansion because they have the needed antigen receptor.
- Clonal selection may explain why memory cells can initiate secondary immune responses more quickly than the primary immune response, due to increased binding affinity from clonal expansion.
- During T cell differentiation, the naive T cell becomes a blast cell that proliferates by clonal expansion and differentiates into memory and effector T cells.
- Many subsets of helper T cells are created during T cell differentiation and perform vastly different functions for the immune system.

Key Terms

- **Clonal selection**: The idea that lymphocytes have antigen-specific binding receptors before they encounter with an antigen, and are selected to proliferate because they have the specific antigen receptor needed during an adaptive immune response.

Clonal selection is an theory that attempts to explain why lymphocytes are able to respond to so many different types of antigens. T and B cells are able to respond to nearly all of the world's vast variety of antigens upon presentation. Clonal selection assumes that lymphocytes are selected during antigen presentation because they already have receptors for that antigen.


Clonal Selection

In clonal selection, an antigen is presented to many circulating naive B and (via MHC) T cells, and the lymphocytes that match the antigen are selected to form both memory and effector clones of themselves. This mass production is termed "clonal expansion," in which daughter cells proliferate into several generations of clones of the original parent cells. The theoretical basis of clonal selection is the assumption that lymphocytes bearing an antigen receptor for an antigen exist long before antigen presentation occurs, explained by the idea of random mutations (VDJ recombination) that occur during lymphocyte maturation. During antigen presentation, pre-existing lymphocytes that bear that antigen receptor are merely selected because they can bind with that antigen. It is also assumed that most lymphocytes never encounter the antigen for which they bear a receptor.

Clonal selection may also be used during negative selection during T cell maturation. Here, the body's own epitopes are presented to the infant lymphocytes; those that react are recognized as auto-reactive and destroyed before they (and their future cloned daughter cells) can leave and wreak havoc in the body. This assumes that random mutations resulted in lymphocytes that were autoreactive instead of reactive to non-self antigens.

Following an adaptive immune response, memory cells are able to respond to a new infection of the same pathogen much more quickly than the original effector T cells during the formation of the adaptive immune response. Clonal selection is thought to cause mutations of antigen-binding affinity in memory cells during clonal expansion so that memory cells have greatly increased antigen-binding affinity than the effector cells during the first response. The increased binding affinity may be why memory cells

can eliminate a pathogen more rapidly than the original generation of effector cells. This idea is still only a theory, but explains many of the nuances of the adaptive immune system.



Clonal selection of lymphocytes: A hematopoietic stem cell undergoes differentiation and genetic rearrangement to produce lymphocytes in the immune system. Clonal selection of lymphocytes: 1) A hematopoietic stem cell undergoes differentiation and genetic rearrangement to produce 2) immature lymphocytes with many different antigen receptors. Those that bind to 3) antigens from the body's own tissues are destroyed, while the rest mature into 4) inactive lymphocytes. Most of these will never encounter a matching 5) foreign antigen, but those that do are activated and produce 6) many clones of themselves.

T Cell Differentiation

Following T cell maturation, naive T cells circulate through the circulatory and lymphatic systems of the body until presented with an antigen for which they bear the receptor. T cells are sorted to be either helper, cytotoxic, or regulatory variants during maturation, but may differentiate into subsets following T cell activation. Following antigen presentation, the T cell is activated and begins to differentiate. T cell differentiation happens via the following steps:

1. The activated T cell becomes a large blast cell.
2. The blast cell proliferates by clonal expansion.
3. Cloned daughter cells differentiate into either effector T cells or memory T cells.
4. Cytotoxic effector T cells are finished, but helper T cells continue to differentiate into individual subsets of helper T cells.

Many different subsets of helper T cells perform various functions. The most common subsets are Th1, which mediates cyotoxic T cell activity through cytokine release, and Th2, which presents antigens to B cells. Additionally, Th17, which only differentiates from effector cells if certain cytokines are present, is important in regulating and inhibiting T-reg cell activity. The effector cells are short-lived for the duration of the adaptive immune response while memory cells are long-lived and are the basis of the secondary immune response.

This page titled 20.7A: Clonal Selection and T-Cell Differentiation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.7B: Specific T-Cell Roles

T helper cells assist the maturation of B cells and memory B cells while activating cytotoxic T cells and macrophages.

Learning Objectives

- Distinguish between the different types of T cell roles

Key Points

- Helper T cells secrete small proteins called cytokines that regulate or assist in the active immune response by activating other immune cells. They also present antigens to B cells.
- Cytotoxic T cells (TC cells, or CTLs) destroy virus-infected cells and tumor cells, and are implicated in transplant rejection and autoimmune disease.
- Memory T cells persist long-term after an infection has resolved. They quickly expand to large numbers of effector T cells upon re-exposure to their antigens, thus providing the immune system with “memory” against past infections.
- Regulatory T cells are crucial for the maintenance of immunological tolerance, because they play a role in suppressing overactive immune responses.
- Natural killer (NK) T cells bridge the adaptive immune system with the innate immune system by producing cytokines and binding to non-MHC or protein bound antigens, such as glycolipids and lipids.

Key Terms

- **Natural Killer T cells**: A heterogeneous group of T cells that shares properties of both T cells and natural killer (NK) cells, and recognizes the non-polymorphic CD1d molecule, an antigen-presenting molecule that binds self- and foreign lipids and glycolipids instead of MHC.

Many different categories and subsets of T cells perform various roles for the immune system. Differentiation for most categories of T cells occurs during the the T cell maturation, but memory cell and helper T subset differentiation occurs after maturation following antigen presentation. The different categories of T cells are the basis for cell-mediated immune system activity.

Helper T Cells

Helper T cells assist other white blood cells in immunologic processes by facilitating cytokines that activate and direct other immune cells. Their primary functions include antigen presentation and activation of B cells, and activation of cytotoxic T cells and macrophages. These cells are also known as CD4+ T cells because they express the CD4 protein on their surfaces. Helper T cells become activated when presented with peptide antigens by MHC class II molecules, which are expressed on the surface of antigen-presenting cells. Once activated, they divide rapidly and secrete regulatory cytokines such as IFN-gamma and certain interleukins. These cells can differentiate into one of several subtypes, including TH1, TH2, TH3, TH17, or TFH, which secrete different cytokines to facilitate a different type of immune response. Differentiation into helper T cell subtypes occurs during clonal selection following T cell activation of naive T cells.

Cytotoxic T cells

Cytotoxic T cells (TC cells, or CTLs) destroy virus-infected cells and tumor cells, and cause much of the damage in in transplant rejection and autoimmune diseases. These cells are also known as CD8+ T cells since they express the CD8 glycoprotein at their surfaces. They recognize their targets by binding to antigens associated with MHC class I, which is present on the surface of nearly every cell of the body. Cytotoxic T cells recognize their antigen on pathogens through their T cell receptor, and kill the pathogen through degranulation and cell-mediated apoptosis. The cytotoxic enzymes and proteases travel to their target cells through a microtubule cytoskeleton. Through IL-10, adenosine and other anti-inflammatory cytokines secreted by regulatory T cells, the CD8+ cells can be inactivated to an anergic state, which can prevent or reduce the severity of autoimmune diseases.

Memory T Cells


Memory T cells are a subset of antigen-specific T cells that persist long after an infection has resolved. They rapidly proliferate to large numbers of effector T cells upon re-exposure to their antigens, thus providing the immune system with “memory” against past infections. The secondary immune response mediated by memory T cells is much faster and more effective at eliminating pathogens compared to the initial immune response. Memory T cells comprise two subtypes: central memory T cells (TCM cells) and effector memory T cells (TEM cells), which have different properties and release different cytokines. Effector memory cells may be either CD4+ or CD8+, and produce either helper or cytotoxic T cells in a secondary immune response.

Regulatory T Cells

Regulatory T cells (Treg cells), also known as suppressor T cells, are crucial for the maintenance of immunological tolerance. Their major role is to shut down T cell-mediated immunity toward the end of an immune reaction and suppress auto-reactive T cells that escaped the process of negative selection in the thymus. Most Treg cells are CD4+ and arise in the thymus. Naturally-occurring Treg cells can be distinguished from other T cells by the presence of an intracellular molecule called FoxP3. Mutations of the FOXP3 gene can prevent regulatory T cell development, causing the fatal autoimmune disease IPEX.

Natural Killer T Cells

Natural killer T cells (NKT cells – not to be confused with natural killer cells) bridge the adaptive immune system with the innate immune system. Unlike conventional T cells that recognize peptide antigens presented by major histocompatibility complex (MHC) molecules, NKT cells recognize glycolipid antigen presented by a molecule called CD1d. Once activated, these cells perform functions ascribed to both Th and Tc cells (i.e., cytokine production and release of cytolytic/cell killing molecules). They are also able to recognize and eliminate some tumor cells and cells infected with herpes viruses. They are among the least common type of T cell in the body and are found in the highest density in the liver. There is an association between NKT cell deficiency and development of autoimmune diseases and chronic inflammatory diseases like asthma, but the exact mechanism of this association is not fully understood.



T cell Activation: T cells become activated upon encountering a pathogen and can become either cytoxic T or helper T cells.

This page titled 20.7B: Specific T-Cell Roles is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

20.7C: Active and Passive Humoral Immunity

The humoral immune response is the aspect of immunity mediated by secreted antibodies.

Learning Objectives

- Distinguish between active and passive humoral immunity

Key Points

- Passive immunity is the transfer of active humoral immunity in the form of ready-made antibodies from one individual to another.
- Naturally-acquired passive immunity includes antibodies given from the mother to her child during fetal development or through breast milk after birth.
- Artificially-acquired passive immunity is a short-term immunization achieved by the transfer of antibodies, and can be administered in several forms.
- Active immunity is long-lasting immunity produced by the body's own immune system and involves the production of long-lasting memory cells.
- Active immunity can either be natural, such as from an infection, or artificial, such as from vaccination.

Key Terms

- **Artificially acquired passive immunity**: A short-lived form of immunity gained from transfer of antibodies produced by another organism.

The humoral immune response (HIR) is the aspect of immunity mediated by secreted antibodies produced by B cells. Secreted antibodies bind to antigens on the surfaces of invading pathogens, which flag them for destruction. Humoral immunity is so named because it involves substances found in the humors, or body fluids. There are two types of humoral immunity: active and passive.

Active Humoral Immunity

Active humoral immunity refers to any form of immunity that occurs as a result of the formation of an adaptive immune response from the body's own immune system. Active immunity is long term (sometimes lifelong) because memory cells with antigen-binding affinity maturation are produced during the lymphocyte differentiation and proliferation that occurs during the formation of an adaptive immune response. It also refers to the effector functions of antibodies, which include pathogen and toxin neutralization, classical complement activation, and opsonin promotion of phagocytosis and pathogen elimination.

Active immunity can either be naturally-occurring or passive. Natural active immunity generally occurs as a result of infection with a pathogen, in which memory cells that remember the antigen of the infectious agent remain in the body. Artificial active immunity is the result of vaccination. During vaccination, the body is exposed to a weakened form of a pathogen that contains the same antigens as the live pathogen, but cannot mount an infection against the body in its weakened state. Vaccinations have become an effective form of disease prevention that is especially useful in preventing diseases that would normally have a high risk of mortality during an infection, where relying on natural active immunity would prove dangerous. However, active immunity does not work to protect against all pathogens, because many can mutate and change their antigen structure over time, which enables them to evade the defenses of immunological memory.


Passive Immunity

Passive immunity is the transfer of active humoral immunity in the form of ready-made antibodies from one individual to another. Passive immunization is used when there is a high risk of infection and insufficient time for the body to develop its own immune

response, or to reduce the symptoms of ongoing or immunosuppressive diseases. Unlike active immunity, passive immunity is short-lived (often only for a few months), because it does not involve the production and upkeep of memory cells.

Passive immunity can occur naturally or artificially. Maternal passive immunity is a type of naturally-acquired passive immunity, and refers to antibody-mediated immunity conveyed to a fetus by its mother during pregnancy. IgG is passed through the placenta to the developing fetus, and is the only antibody isotype that can pass through the placenta. Because passive immunity is short-lived, vaccination is often required shortly following birth to prevent diseases such as tuberculosis, hepatitis B, polio, and pertussis; however, maternal antibodies can inhibit the induction of protective vaccine responses throughout the first year of life. This effect is usually overcome by secondary responses to booster immunization. Passive immunity is also provided through the transfer of IgA antibodies found in breast milk, which are transferred to the gastrointestinal tract of the infant, protecting against bacterial infections until the newborn has produced enough matured B cells to synthesize its own antibodies.

Artificially-acquired passive immunity is a short-term immunization achieved by the transfer of antibodies, and can be administered in several forms: as human or animal blood (usually horse) plasma or serum, as pooled human immunoglobulin for intravenous (IVIG) or intramuscular (IG) use, and as monoclonal antibodies (MAb). Passive transfer is used to help treat those with immunodeficiency and for several types of severe acute infections that have no vaccine, such as the Ebola virus. Immunity derived from passive immunization lasts for only a short period of time, and there is potential risk for hypersensitivity reactions and serum sickness, especially from gamma globulin of non-human origin. Passive immunity provides immediate protection, but the body does not develop memory; therefore, the patient is at risk of being infected by the same pathogen later.



Innate and adaptive immunity in the immune system: This chart depicts the different divisions of immunity, including adaptive, innate, natural, artificial, passive (maternal), active (infection), passive (antibody transfer) and active (immunization).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- T cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/T_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Immunology/Introduction. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Immunology/Introduction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thymocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thymocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Clonal selection. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Clonal_selection**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- T cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/T_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Natural Killer T cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Natural...er%20T%20cells**. **License**: ***CC***

BY-SA: Attribution-ShareAlike
- Clonal selection. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clonal_selection. **License**: *CC BY-SA: Attribution-ShareAlike*
- T cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/T_cell. **License**: *Public Domain: No Known Copyright*
- Passive immunity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Passive_immunity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Humoral immunity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Humoral_immunity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Artificially acquired passive immunity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Artific...ive%20immunity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clonal selection. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Clonal_selection. **License**: *CC BY-SA: Attribution-ShareAlike*
- T cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/T_cell. **License**: *Public Domain: No Known Copyright*
- Immunity (medical). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Immunity_(medical). **License**: *Public Domain: No Known Copyright*

This page titled 20.7C: Active and Passive Humoral Immunity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21: RESPIRATORY SYSTEM

21.10: Respiration Control

21.10A: Neural Mechanisms (Respiratory Center)

21.10B: Neural Mechanisms (Cortex)

21.10C: Chemoreceptor Regulation of Breathing

21.10D: Proprioceptor Regulation of Breathing

21.11: Respiratory Adjustments

21.11A: Adjustments During Exercise

21.11B: Adjustments at High Altitude

21.1: Overview of the Respiratory System

21.1A: The Reason for Breathing

21.1B: Functional Anatomy of the Respiratory System

21.2: Conducting Zone

21.2A: Nose and Paranasal Sinuses

21.2B: Pharynx

21.2C: Larynx

21.2D: Structures Used in Voice Production

21.2E: Trachea

21.3: Respiratory Zone

21.3A: Bronchi and Subdivisions

21.3B: Alveoli

21.4: The Lungs and Pleurae

21.4A: Lungs

21.4B: Lobes, Fissures, and Lobules

21.4C: Blood Supply to the Lungs

21.5: Mechanics of Breathing

21.5A: Pressure Changes During Pulmonary Ventilation

21.5B: Inspiration

21.5C: Expiration

21.5D: Breathing Patterns

21.6: Factors Affecting Pulmonary Ventilation

21.6A: Factors Affecting Pulmonary Ventilation: Surface Tension of Alveolar Fluid

21.6B: Factors Affecting Pulmonary Ventilation: Compliance of the Lungs

21.6C: Factors Affecting Pulmonary Ventilation: Airway Resistance

21.7: Nonrespiratory Lung Functions

21.7A: Lung Capacity and Volume

21.7B: Nonrespiratory Air Movements

21.8: Gas Laws

21.8A: Dalton's Law of Partial Pressure

21.8B: Henry's Law

21.9: Gas Exchange

21.1A: External Respiration

21.9B: Internal Respiration

21.9C: Oxygen Transport

21.9D: Carbon Dioxide Transport

This page titled 21: Respiratory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.10: RESPIRATION CONTROL

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.10A: Neural Mechanisms (Respiratory Center)

21.10B: Neural Mechanisms (Cortex)

21.10C: Chemoreceptor Regulation of Breathing

21.10D: Proprioceptor Regulation of Breathing

This page titled 21.10: Respiration Control is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.10A: Neural Mechanisms (Respiratory Center)

The medulla and the pons are involved in the regulation of the ventilatory pattern of respiration.


Learning Objectives

- Describe the neural mechanism of the respiratory center in respiration control

Involuntary respiration is any form of respiratory control that is not under direct, conscious control. Breathing is required to sustain life, so involuntary respiration allows it to happen when voluntary respiration is not possible, such as during sleep. Involuntary respiration also has metabolic functions that work even when a person is conscious.

The Respiratory Centers

Involuntary respiration is controlled by the respiratory centers of the upper brainstem (sometimes termed the lower brain, along with the cerebellum). This region of the brain controls many involuntary and metabolic functions besides the respiratory system, including certain aspects of cardiovascular function and involuntary muscle movements (in the cerebellum).



Anatomy of the brainstem: The brainstem, which includes the pons and medulla.

The respiratory centers contain chemoreceptors that detect pH levels in the blood and send signals to the respiratory centers of the brain to adjust the ventilation rate to change acidity by increasing or decreasing the removal of carbon dioxide (since carbon dioxide is linked to higher levels of hydrogen ions in blood).

There are also peripheral chemoreceptors in other blood vessels that perform this function as well, which include the aortic and carotid bodies.

The Medulla

The medulla oblongata is the primary respiratory control center. Its main function is to send signals to the muscles that control respiration to cause breathing to occur. There are two regions in the medulla that control respiration:

- The ventral respiratory group stimulates expiratory movements.

- The dorsal respiratory group stimulates inspiratory movements.

The medulla also controls the reflexes for nonrespiratory air movements, such as coughing and sneezing reflexes, as well as other reflexes, like swallowing and vomiting.

The Pons

The pons is the other respiratory center and is located underneath the medulla. Its main function is to control the rate or speed of involuntary respiration. It has two main functional regions that perform this role:

- The apneustic center sends signals for inspiration for long and deep breaths. It controls the intensity of breathing and is inhibited by the stretch receptors of the pulmonary muscles at maximum depth of inspiration, or by signals from the pneumotaxic center. It increases tidal volume.
- The pneumotaxic center sends signals to inhibit inspiration that allows it to finely control the respiratory rate. Its signals limit the activity of the phrenic nerve and inhibits the signals of the apneustic center. It decreases tidal volume.

The apneustic and pneumotaxic centers work against each other together to control the respiratory rate.

Key Points

- The ventral respiratory group controls voluntary forced exhalation and acts to increase the force of inspiration.
- The dorsal respiratory group (nucleus tractus solitarius) controls mostly inspiratory movements and their timing.
- Ventilatory rate (minute volume) is tightly controlled and determined primarily by blood levels of carbon dioxide as determined by metabolic rate.
- Chemoreceptors can detect changes in blood pH that require changes in involuntary respiration to correct.The apneustic (stimulating) and pneumotaxic (limiting) centers of the pons work together to control rate of breathing.
- The medulla sends signals to the muscles that initiate inspiration and expiration and controls nonrespiratory air movement reflexes, like coughing and sneezing.

Key Terms

- **respiratory control centers**: The medulla which sends signals to the muscles involved in breathing, and the pons which controls the rate of breathing.
- **chemorecepters**: These are receptors in the medulla and in the aortic and carotid bodies of the blood vessels that detect changes in blood pH and signal the medulla to correct those changes.

This page titled 21.10A: Neural Mechanisms (Respiratory Center) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.10B: Neural Mechanisms (Cortex)

Learning Objectives

- Describe the mechanism of the neural cortex in respiration control

Voluntary respiration is any type of respiration that is under conscious control. Voluntary respiration is important for the higher functions that involve air supply, such as voice control or blowing out candles. Similarly to how involuntary respiration's lower functions are controlled by the lower brain, voluntary respiration's higher functions are controlled by the upper brain, namely parts of the cerebral cortex.

The Motor Cortex

The primary motor cortex is the neural center for voluntary respiratory control. More broadly, the motor cortex is responsible for initiating any voluntary muscular movement.

The processes that drive its functions aren't fully understood, but it works by sending signals to the spinal cord, which sends signals to the muscles it controls, such as the diaphragm and the accessory muscles for respiration. This neural pathway is called the ascending respiratory pathway.


Different parts of the cerebral cortex control different forms of voluntary respiration. Initiation of the voluntary contraction and relaxation of the internal and external intercostal muscles takes place in the superior portion of the primary motor cortex.

The center for diaphragm control is posterior to the location of thoracic control (within the superior portion of the primary motor cortex). The inferior portion of the primary motor cortex may be involved in controlled exhalation.

Activity has also been seen within the supplementary motor area and the premotor cortex during voluntary respiration. This is most likely due to the focus and mental preparation of the voluntary muscular movement that occurs when one decides to initiate that muscle movement.

Note that voluntary respiratory nerve signals in the ascending respiratory pathway can be overridden by chemoreceptor signals from involuntary respiration. Additionally, other structures may override voluntary respiratory signals, such as the activity of limbic center structures like the hypothalamus.

During periods of perceived danger or emotional stress, signals from the hypothalamus take over the respiratory signals and increase the respiratory rate to facilitate the fight or flight response.



Topography of the primary motor cortex: Topography of the primary motor cortex, on an outline drawing of the human brain. Each part of the primary motor cortex controls a different part of the body.

Nerves Used in Respiration

There are several nerves responsible for the muscular functions involved in respiration. There are three types of important respiratory nerves:

- The phrenic nerves: The nerves that stimulate the activity of the diaphragm. They are composed of two nerves, the right and left phrenic nerve, which pass through the right and left side of the heart respectively. They are autonomic nerves.
- The vagus nerve*:* Innervates the diaphragm as well as movements in the larynx and pharynx. It also provides parasympathetic stimulation for the heart and the digestive system. It is a major autonomic nerve.
- The posterior thoracic nerves: These nerves stimulate the intercostal muscles located around the pleura. They are considered to be part of a larger group of intercostal nerves that stimulate regions across the thorax and abdomen. They are somatic nerves.

These three types of nerves continue the signal of the ascending respiratory pathway from the spinal cord to stimulate the muscles that perform the movements needed for respiration.

Damage to any of these three respiratory nerves can cause severe problems, such as diaphragm paralysis if the phrenic nerves are damaged. Less severe damage can cause irritation to the phrenic or vagus nerves, which can result in hiccups.

Key Points

- The motor cortex within the cerebral cortex of the brain controls voluntary respiration (the ascending respiratory pathway).
- Voluntary respiration may be overridden by aspects of involuntary respiration, such as chemoreceptor stimulus, and hypothalamus stress response.
- The phrenic nerves, vagus nerves, and posterior thoracic nerves are the major nerves involved in respiration.
- Voluntary respiration is needed to perform higher functions, such as voice control.

Key Terms

- **The Phrenic Nerves**: A set of two nerves that brings nerve impulses from the spinal cord to the diaphragm.
- **primary motor cortex**: The region in the brain that initiates all voluntary muscular movement, including those for respiration.

This page titled 21.10B: Neural Mechanisms (Cortex) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.10C: Chemoreceptor Regulation of Breathing

Chemoreceptors detect the levels of carbon dioxide in the blood by monitoring the concentrations of hydrogen ions in the blood.

Learning Objectives

- Describe the role of chemoreceptors in the regulation of breathing

Key Points

- An increase in carbon dioxide concentration leads to a decrease in the pH of blood due to the production of H^+ ions from carbonic acid.
- In response to a decrease in blood pH, the respiratory center (in the medulla) sends nervous impulses to the external intercostal muscles and the diaphragm, to increase the breathing rate and the volume of the lungs during inhalation.
- Hyperventilation causes alakalosis, which causes a feedback response of decreased ventilation (to increase carbon dioxide), while hypoventilation causes acidosis, which causes a feedback response of increased ventilation (to remove carbon dioxide).
- Any situation with hypoxia (too low oxygen levels) will cause a feedback response that increases ventilation to increase oxygen intake.
- Vomiting causes alkalosis and diarrhea causes acidosis, which will cause an appropriate respiratory feedback response.

Key Terms

- **hypoxia**: A system-wide deficiency in the levels of oxygen that reach the tissues.
- **central chemoreceptors**: Located within the medulla, they are sensitive to the pH of their environment.
- **peripheral chemoreceptors**: The aoritic and carotid bodies, which act principally to detect variation of the oxygen concentration in the arterial blood, also monitor arterial carbon dioxide and pH.

Chemoreceptor regulation of breathing is a form of negative feedback. The goal of this system is to keep the pH of the blood stream within normal neutral ranges, around 7.35.

Chemoreceptors

A chemoreceptor, also known as chemosensor, is a sensory receptor that transduces a chemical signal into an action potential. The action potential is sent along nerve pathways to parts of the brain, which are the integrating centers for this type of feedback. There are many types of chemoreceptors in the body, but only a few of them are involved in respiration.

The respiratory chemoreceptors work by sensing the pH of their environment through the concentration of hydrogen ions. Because most carbon dioxide is converted to carbonic acid (and bicarbonate) in the bloodstream, chemoreceptors are able to use blood pH as a way to measure the carbon dioxide levels of the bloodstream.

The main chemoreceptors involved in respiratory feedback are:

1. Central chemoreceptors: These are located on the ventrolateral surface of medulla oblongata and detect changes in the pH of spinal fluid. They can be desensitized over time from chronic hypoxia (oxygen deficiency) and increased carbon dioxide.
2. Peripheral chemoreceptors: These include the aortic body, which detects changes in blood oxygen and carbon dioxide, but not pH, and the carotid body which detects all three. They do not desensitize, and have less of an impact on the respiratory rate compared to the central chemoreceptors.

Chemoreceptor Negative Feedback

Negative feedback responses have three main components: the sensor, the integrating sensor, and the effector. For the

respiratory rate, the chemoreceptors are the sensors for blood pH, the medulla and pons form the integrating center, and the respiratory muscles are the effector.

Consider a case in which a person is hyperventilating from an anxiety attack. Their increased ventilation rate will remove too much carbon dioxide from their body. Without that carbon dioxide, there will be less carbonic acid in blood, so the concentration of hydrogen ions decreases and the pH of the blood rises, causing alkalosis.


In response, the chemoreceptors detect this change, and send a signal to the medulla, which signals the respiratory muscles to decrease the ventilation rate so carbon dioxide levels and pH can return to normal levels.

There are several other examples in which chemoreceptor feedback applies. A person with severe diarrhea loses a lot of bicarbonate in the intestinal tract, which decreases bicarbonate levels in the plasma. As bicarbonate levels decrease while hydrogen ion concentrations stays the same, blood pH will decrease (as bicarbonate is a buffer) and become more acidic.

In cases of acidosis, feedback will increase ventilation to remove more carbon dioxide to reduce the hydrogen ion concentration. Conversely, vomiting removes hydrogen ions from the body (as the stomach contents are acidic), which will cause decreased ventilation to correct alkalosis.

Chemoreceptor feedback also adjusts for oxygen levels to prevent hypoxia, though only the peripheral chemoreceptors sense oxygen levels. In cases where oxygen intake is too low, feedback increases ventilation to increase oxygen intake.

A more detailed example would be that if a person breathes through a long tube (such as a snorkeling mask) and has increased amounts of dead space, feedback will increase ventilation.



Respiratory feedback: The chemoreceptors are the sensors for blood pH, the medulla and pons form the integrating center, and the respiratory muscles are the effector.

This page titled 21.10C: Chemoreceptor Regulation of Breathing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.10D: Proprioceptor Regulation of Breathing

The Hering–Breuer inflation reflex prevents overinflation of the lungs.

Learning Objectives

- Evaluate the effect of proprioception (the sense of the relative position of the body and effort being employed in movement) on breathing

Key Points


- Pulmonary stretch receptors present in the smooth muscle of the airways and the pleura respond to excessive stretching of the lung during large inspirations.
- The Hering–Breuer inflation reflex is initiated by stimulation of stretch receptors. The deflation reflex is initiated by stimulation of the compression receptors (called proprioceptors) or deactivation of stretch receptors when the lungs deflate.
- Activation of the pulmonary stretch receptors (via the vagus nerve) results in inhibition of the inspiratory stimlus in the medulla, and thus inhibition of inspiration and initiation of expiration.
- An increase in pulmonary stretch receptor activity leads to an elevation of heart rate (tachycardia).
- A cyclical, elevated heart rate from inspiration is called sinus arrhythmia and is a normal response in youth. Inhibition of inspiration is important to allow expiration to occur.

Key Terms

- **sinus arryhthmia**: A normal cyclical heart rate change in which an increase in heart rate occurs during inspiration, but returns to normal during expiration.
- **pulmonary stretch receptors**: A sensory receptor that sends an action potential when it detects pressure, tension, stretch, or distortion.

The lungs are a highly elastic organ capable of expanding to a much larger volume during inflation. While the volume of the lungs is proportional to the pressure of the pleural cavity as it expands and contracts during breathing, there is a risk of over-inflation of the lungs if inspiration becomes too deep for too long. Physiological mechanisms exist to prevent over-inflation of the lungs.

The Hering–Bauer Reflex



Cardiac and respiratory branches of the vagus nerve: The vagus nerve is the neural pathway for stretch receptor regulation of breathing.

The Hering–Breuer reflex (also called the inflation reflex) is triggered to prevent over-inflation of the lungs. There are many stretch receptors in the lungs, particularly within the pleura and the smooth muscles of the bronchi and bronchioles, that activate when the lungs have inflated to their ideal maximum point.

These stretch receptors are mechanoreceptors, which are a type of sensory receptor that specifically detects mechanical pressure, distortion, and stretch, and are found in many parts of the human body, especially the lungs, stomach, and skin. They do not detect fine-touch information like most sensory receptors in the human body, but they do create a feeling of tension or fullness when activated, especially in the lungs or stomach.

When the lungs are inflated to their maximum volume during inspiration, the pulmonary stretch receptors send an action potential signal to the medulla and pons in the brain through the vagus nerve.

The pneumotaxic center of the pons sends signals to inhibit the apneustic center of the pons, so it doesn't activate the inspiratory area (the dorsal medulla), and the inspiratory signals that are sent to the diaphragm and accessory muscles stop. This is called the inflation reflex.

As inspiration stops, expiration begins and the lung begins to deflate. As the lungs deflate the stretch receptors are deactivated (and compression receptors called proprioreceptors may be activated) so the inhibitory signals stop and inhalation can begin again—this is called the deflation reflex.

Early physiologists believed this reflex played a major role in establishing the rate and depth of breathing in humans. While this may be true for most animals, it is not the case for most adult humans at rest. However, the reflex may determine the breathing rate and depth in newborns and in adult humans when tidal volume is more than 1 L, such as when exercising.

Additionally, people with emphysema have an impaired Hering–Bauer reflex due to a loss of pulmonary stretch receptors from the destruction of lung tissue, so their lungs can over-inflate as well as collapse, which contributes to shortness of breath.

Sinus Arrhythmia

As the Hering–Bauer reflex uses the vagus nerve as its neural pathway, it also has a few cardiovascular system effects because the vagus nerve also innervates the heart.

During stretch receptor activation, the inhibitory signal that travels through the vagus nerve is also sent to the sinus-atrial node

of the heart. Its stimulation causes a short-term increase in resting heart rate, which is called tachycardia.

The heart rate returns to normal during expiration when the stretch receptors are deactivated. When this process is cyclical it is called a sinus arrhythmia, which is a generally normal physiological phenomenon in which there is short-term tachycardia during inspiration.

Sinus arryhthmias do not occur in everyone, and are more common in youth. The sensitivity of the sinus-atrial node to the inflation reflex is lost over time, so sinus arryhthmias are less common in older people.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Control of respiration. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Control_of_respiration**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Control of respiration. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Control_of_respiration**. **License**: *CC BY-SA: Attribution-ShareAlike*
- dorsal respiratory group. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/dorsal%...ratory%20group**. **License**: *CC BY-SA: Attribution-ShareAlike*
- respiratory control centers. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/respir...ontrol+centers**. **License**: *CC BY-SA: Attribution-ShareAlike*
- aortic bodies. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/aortic+bodies**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain sagittal section stem highlighted. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Brain_sagittal_section_stem_highlighted.svg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Control of respiration. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Control_of_respiration**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Exhalation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Exhalation**. **License**: *CC BY-SA: Attribution-ShareAlike*
- neuron. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neuron**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/external-internal-costals**. **License**: *CC BY-SA: Attribution-ShareAlike*
- glottis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glottis**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain sagittal section stem highlighted. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Brain_sagittal_section_stem_highlighted.svg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human motor cortex topography. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_motor_cortex_topography.png**. **License**: *Public Domain: No Known Copyright*
- Chemoreceptor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chemoreceptor**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/central-chemoreceptors**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/peripheral-chemoreceptors**. **License**: *CC BY-SA: Attribution-ShareAlike*
- external intercostal muscles. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/external%20intercostal%20muscles**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brain sagittal section stem highlighted. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Brain_sagittal_section_stem_highlighted.svg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human motor cortex topography. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_motor_cortex_topography.png**. **License**: *Public Domain: No Known Copyright*
- 2327_Respiratory_Centers_of_the_Brain.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Respiratory_center#/media/File:2327_Respiratory_Centers_of_the_Brain.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Heringu2013Breuer reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hering%E2%80%93Breuer_reflex%23Rate_and_depth_of_breathing**. **License**: *CC BY-SA: Attribution-ShareAlike*

- Heringu2013Breuer reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hering%E2%80%93Breuer_reflex%23Rate_and_depth_of_breathing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tachycardia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tachycardia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/nucleus-ambiguus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- proprioceptor. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/proprioceptor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain sagittal section stem highlighted. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Brain_sagittal_section_stem_highlighted.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human motor cortex topography. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_motor_cortex_topography.png**. **License**: ***Public Domain: No Known Copyright***
- 2327_Respiratory_Centers_of_the_Brain.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Respiratory_center#/media/File:2327_Respiratory_Centers_of_the_Brain.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray793. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray793.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 21.10D: Proprioceptor Regulation of Breathing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.11: RESPIRATORY ADJUSTMENTS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

21.11A: Adjustments During Exercise

21.11B: Adjustments at High Altitude

This page titled 21.11: Respiratory Adjustments is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.11A: Adjustments During Exercise

Aerobic and anaerobic exercise work to increase the mechanical efficiency of the heart.

Learning Objectives

- Describe respiratory adjustments during exercise

Key Points

- There is tremendous variation in an individual's response to excercise.
- The beneficial effects of physical activity include improved cognition from increased flow of blood and oxygen to the brain, and reduced stress response.
- Anaerobic respiration happens during exercise if the body does not get enough oxygen to meet its metabolic demands.
- Extremely intense, long-term cardiovascular exercise can be deleterious and has been associated with asthma and sudden cardiac death.
- Sudden cardiac death from exercise has a genetic component that is worsened by cardiac hypertrophy.

Key Terms

- **anaerobic exercise**: This occurs during exercise when the body's oxygen intake doesn't meet oxygen demands to produce enough ATP.
- **asthma**: A long-term respiratory condition, marked by airway obstruction from bronchospams. It is often in response to an allergen, cold air, exercise, or emotional stress.

During exercise, the human body needs a greater amount of oxygen to meet the increased metabolic demands of the muscle tissues. Various short-term respiratory changes must occur in order for those metabolic demands to be reached. Eventually exercise can induce long term cardiovascular and respiratory changes, which can be both healthy and unhealthy.

Short-Term Changes

During exercise, carbon dioxide levels (the metabolic waste) rise in arterial blood. Carbon dioxide induces vasodilation in the arteries while the heart rate increases, which leads to better blood flow and tissue perfusion, and better oxygen delivery to the tissues.

In particular, the blood flow to the brain and heart is increased, while increased blood flow to the muscles makes exercise easier. Additionally, the respiratory rate increases as a result of higher carbon dioxide levels (through chemoreceptor regulation), which allows the body to release more carbon dioxide while increasing oxygen intake.

If exercise is too intense for oxygen demands to be satisfied in the short term, anaerobic respiration will be used to make up for the ATP deficit in the muscles. This can cause a buildup of lactic acid in the muscles, which is the byproduct of lactic acid fermentation (the most common anaerobic respiration process in the human body).

This is one reason why muscles may become sore during exercise, though the lactic acid is eventually removed through conversion to glucose in the liver.

Beneficial Long-Term Changes

In the long run, exercise results in increased levels of arterial oxygen levels at rest, due to chemoreceptor desensitization to carbon dioxide levels and a lack of oxygen supply relative to oxygen demands during exercise. Over time, the elevated respiratory rate past what is needed to restore normal blood pH levels following exercise causes a long-term increase in arterial oxygen

levels.

Increased oxygen levels in the body are especially important to the long-term health of the brain and heart, two organs that are vital to sustain life and that require large amounts of oxygen to function well. Brain plasticity and cognition also improves as a result.

Exercise also has beneficial effects for reducing stress responses in the body due to increased endorphin production in the brain from exercise. In long-term exercise of appropriate intensity, the volume and strength of the heart are improved, which makes additional exercise easier.

Adverse Long-Term Changes

When exercise is performed at too intense levels for too long or without adequate rest, it can cause long term, adverse health effects. Muscle tissue repair is impaired in those that exercise too frequently. Exercise induced asthma is another common complication from too much exercise.


Normally asthma is caused by an allergic response within the lungs, but exercise can induce a similar response from too much intake of dry and cold air during the increased respiratory rate from exercise. It is most common in those that do more cardiac-oriented exercise.

The air in the lungs is meant to be moistened and humidified before entering the lungs, but if it is not adequately treated in the upper airways, it can induce bronchospams in the brionchioles, which cause the wheezing and coughing that occurs during an asthma attacks. Asthma treatments (such as medicines from an inhaler) can help prevent exercise-induced asthma, though it is only a particular risk in those who already have allergic asthma, or who simply exercise at too much intensity for what their body is capable of handling.

Sudden cardiac death is notable for occurring in otherwise healthy and young athletes who train too much for long-distance running. While many factors that can lead to sudden cardiac death in athletes are genetic (such as inherited problems with heart rhythm or coronary artery blood supply), many of these deaths are caused by cardiac hypertrophy, in which the heart becomes too thick from damage and scarring from too-intense exercise over long periods of time.

Initially, hypertrophy improves blood flow due to increases in the strength of the heart, but it eventually leads to heart failure as the tissues become too thick to pump normally.

Athletes with genetic susceptibilities are more likely to experience sudden cardiac death as a response to their hypertrophied heart, which can contribute to development of a severe arrhythmia (such as ventricular fibrillation). Exercising at appropriate intensities significantly reduces the risk of sudden cardiac death, though those with genetic susceptibilities should take more caution.



Asthma: Obstruction of the lumen of the bronchiole by mucoid exudate, goblet cell metaplasia, epithelial basement membrane thickening, and severe inflammation of bronchiole.

This page titled 21.11A: Adjustments During Exercise is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.11B: Adjustments at High Altitude

The human body can adapt to high altitude through immediate and long-term acclimatization processes.

Learning Objectives

- Evaluate the respiratory adjustments to high altitude

Key Points

- At high altitude, in the short term, the lack of oxygen is sensed by the peripheral chemoreceptors, which causes an increase in ventilation. An increase in heart rate and decrease in stroke volume also occurs.
- During acclimatization over a few days to weeks, the body produces more red blood cells to counteract the lower oxygen saturation in blood in high altitudes.
- Full adaptation to high altitude is achieved when the increase of red blood cells reaches a plateau and stops.
- Increased red blood cell levels remain for about two weeks after acclimatization, which makes it a popular training regimen for athletes.

Key Terms

- **acclimatization**: Long-term adjustment to high altitude, which is primarily due to increased red blood cell production and capillary tissue perfusion.
- **hematocrit**: This is the amount of red blood cells in a given volume of blood.

The human body can adapt to high altitude through immediate and long-term acclimatization. At high altitude there is lower air pressure compared to a lower altitude or sea-level altitude.

Due to Boyle's law, at higher altitude the partial pressure of oxygen in the air is lower, and less oxygen is breathed in with every breath. The partial pressure gradients for gas exchange are also decreased, along with the percentage of oxygen saturation in hemoglobin.

Humans can survive at high altitudes with impaired short-term functions that eventually adjust in the long term. Some altitudes are too high for acclimatization to work, and can cause death if people stay there for too long.

Short-Term Adjustments

At high altitude, in the short term, the lack of oxygen is sensed by the peripheral chemoreceptors, which causes an increase in breathing rate (hyperventilation). However, hyperventilation also causes the adverse effect of alkalosis due to increasing the rate by which carbon dioxide is removed from the body, which inhibits the respiratory center from enhancing the respiratory rate to meet the oxygen demands.

Additionally, the peripheral chemoreceptors cause sympathetic nervous system stimulation, which causes the heart rate to increase while stroke volume decreases, and digestion is impaired. Shortness of breath is common, and urination increases.

Along with alkalosis, these effects make up the symptoms of altitude sickness, which become worse during exercise at high altitudes (which involves more anaerobic respiration than at lower altitudes), but falls off during acclimatization.

Acclimatization

Acclimatization to high altitude requires days, or even weeks. Gradually, the body compensates for the respiratory alkalosis by kidney excretion of bicarbonate, which allows adequate respiration to provide oxygen without risking alkalosis. It takes about four

days at any given altitude, and can be enhanced by drugs such as acetazolamide (which decreases fluid retention).

Staying hydrated during acclimatization is important to minimize altitude sickness symptoms and to counteract increased urination. The heart rate and ventilation rate at rest both remain elevated despite the acclimatization, while heart rate at maximum activity level will be reduced.

The main difference brought about by acclimatization that explain why it makes high altitudes more comfortable for the body is increased levels of circulating red blood cells, which improve the carrying capacity of oxygen by hemoglobin in the body. This is an adaptive response due to erythropotein secretions in the kidneys (from lack of oxygen in the tissues) that act on the liver to increase erythrocyte (red blood cell) production.

Blood volume decreases, which also increases the hematocrit, which is the concentration of hemoglobin in blood. This increase in red blood cells remains for a few weeks after one returns to lower altitude, so those who acclimatize to high altitude will experience improved athletic performance at lower altitudes. Capillary density and tissue perfusion also increase.

These physiological changes make high-altitude athletic training popular for athletes, such as Olympic athletes. Full hematological adaptation to high altitude is achieved when the increase of red blood cells reaches a plateau and stops.

The length of full hematological adaptation can be approximated by multiplying the altitude in kilometers by 11.4 days. For example, to adapt to 4,000 meters (13,000 ft.) of altitude would require 45.6 days.

The upper altitude limit of this linear relationship has not been fully established, in part because extremely high altitudes have such little oxygen content that they would be fatal regardless of acclimatization.



An athletic training camp in the Swiss Alps

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Physical exercise. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Physical_exercise%23Cardiovascular_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Physical exercise. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Physical_exercise%23Cardiovascular_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Physical exercise. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Physical_exercise%23Cardiovascular_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- asthma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/asthma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/anaerobic-exercise**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Asthma_.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Effects of high altitude on humans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Effects_of_high_altitude_on_humans**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lung volume. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lung_volume**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alkalosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hematocrit. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hematocrit**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Asthma_.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Swiss_Olympic_training_base.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Effects_of_high_altitude_on_humans#/media/File:Swiss_Olympic_training_base.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 21.11B: Adjustments at High Altitude is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.1: OVERVIEW OF THE RESPIRATORY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.1A: The Reason for Breathing

21.1B: Functional Anatomy of the Respiratory System

This page titled 21.1: Overview of the Respiratory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.1A: The Reason for Breathing

Breathing allows for the delivery oxygen to internal tissues and cells where it is needed, and allows for the removal of CO2.

Learning Objectives

- List the main reasons for breathing

Key Points

- Breathing is the process that moves air in and out of the lungs of terrestrial vertebrates, to take in oxygen and remove carbon dioxide.
- Aerobic organisms require oxygen to release energy via respiration, in the form of metabolized, energy-rich molecules such as glucose.
- Another key role of respiration is to maintain proper blood pH —too much carbon dioxide causes acidosis, and too little carbon dioxide causes causes alkalosis.
- Once these dissolved gases are in the blood, the circulatory system transports them around the body, thereby bringing oxygen to the tissues, and carbon dioxide to the lungs.

Key Terms

- **passive diffusion**: Net movement of material from an area of high concentration to an area of lower concentration without any energy input.
- **alkalosis**: When blood pH becomes alkaline due to too few hydrogen ions and too little carbon dioxide.
- **acidosis**: When blood pH becomes acidic due to too many hydrogen ions and too much carbon dioxide.

The Purpose of Breathing

Breathing is the physiological process that moves air in and out of the lungs in terrestrial vertebrates. Respiration is often referred to as breathing, but it can also mean cellular respiration, which is the main reason why breathing is important.

Cells require oxygen from the air to extract energy from glucose through respiration, which produces carbon dioxide and water as a waste product. Therefore, oxygen is vital for every part of normal cellular function, and oxygen deficiency can have severe pathological consequences.

The respiratory system facilitates breathing. In the alveoli tissue of the lungs, the exchange of oxygen and carbon dioxide molecules between the air and the bloodstream occurs by passive transport, so that oxygen is taken in and carbon dioxide and water are removed.

Passive diffusion (also called bulk flow) is the term for the movement of these gases between the air and bloodstream based on their relative concentration, with the gas with the greater concentration moving across to the area with the lower concentration. This process consumes no energy.

The circulatory system is deeply connected with the respiratory system because it distributes the dissolved oxygen to the tissues of the body and the waste carbon dioxide to the lungs.


Breathing Controls Blood Chemistry

Another key role of respiration is maintaining proper blood pH. The concentration of hydrogen ions in blood is partially determined by the amount of dissolved carbon dioxide in the blood, so that more carbon dioxide results in more hydrogen, causing the blood to have a lower pH and be more acidic.

When the blood becomes acidic, respiratory acidosis occurs, which can cause tissue damage if too severe. Acidosis can be caused by hypoventilation (too little breathing), which reduces the removal rate of carbon dioxide, causing it to build up in the bloodstream along with hydrogen. There are many symptoms of acidosis, such as headache, confusion, increased heart rate, and muscle weakness.

Respiratory alkalosis happens when the opposite effect occurs. When the blood pH becomes too high, from too few hydrogen ions because of too little carbon dioxide, the blood will become alkaline, which is also harmful to the body. Alkalosis can happen from hyperventilation (too much breathing) which removes too much carbon dioxide from the bloodstream.

Thankfully, negative feedback mechanisms exist so that hyperventilation and hypoventilation can be corrected. These feedback mechanisms can fail in people with chronic respiratory diseases like emphysema and bronchitis, or from the side effects of certain drugs, in which acidosis and alkalosis will occur regardless.



Some Symptoms of Acidosis and Alkalosis: One of the primary reasons for breathing is to regulate blood pH so that respiratory acidosis and alkalosis don't occur.

This page titled 21.1A: The Reason for Breathing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.1B: Functional Anatomy of the Respiratory System

The respiratory system include lungs, airways and respiratory muscles. Ventilation is the rate at which gas enters or leaves the lung.

Learning Objectives

- Summarize the the functional anatomy of the respiratory system

Key Points


- Ventilation occurs under the control of the autonomic nervous system from parts of the brain stem—the medulla oblongata and the pons —that together form the respiration regulatory center.
- The three types of ventilation are minute ventilation, alevolar ventilation, and dead space ventilation.
- Inhalation is initiated by the diaphragm and supported by the external intercostal muscles. Additional accessory muscles include sternocleidomastoid, platysma, the scalene muscles of the neck, pectoral muscles, and the latissimus dorsi.
- When the diaphragm contracts, the ribcage expands and the contents of the abdomen are moved downward, resulting in a larger thoracic volume and negative pressure (with respect to atmospheric pressure) inside the chest.
- Exhalation is generally a passive process since the lungs have a natural elasticity; they recoil from the stretch of inhalation and air flows back out until the pressures in the chest and the atmosphere reach equilibrium.
- Gas exchange occurs at the alveoli, the tiny sacs that are the basic functional component of the lungs. The alveoli are interwoven with capillaries that connect to the larger bloodstream.

Key Terms

- **elastic recoil**: The lungs' rebound from the stretch of inhalation that passively removes air from the lungs during exhalation.
- **Dead space**: Any space in the airways that is not involved in alveolar gas exhange, such as the conducting zones.
- **ventilation**: The bodily process of breathing, the inhalation of air to provide oxygen, and the exhalation of spent air to remove carbon dioxide.

The Respiratory System

The primary function of the respiratory system is gas exchange between the external environment and an organism's circulatory system. In humans and other mammals, this exchange balances oxygenation of the blood with the removal of carbon dioxide and other metabolic wastes from the circulation.



Bronchial anatomy: The pulmonary alveoli are the terminal ends of the respiratory tree, outcropping from either alveolar sacs or alveolar ducts, which are both sites of gas exchange with the blood.

As gas exchange occurs, the acid-base balance of the body is maintained as part of homeostasis. If proper ventilation is not maintained, two opposing conditions could occur: respiratory acidosis (a life threatening condition) and respiratory alkalosis.

At the molecular level, gas exchange occurs in the alveoli—tiny sacs which are the basic functional component of the lungs. The alveolar epithelial tissue is extremely thin and permeable, allowing for gas exchange between the air inside the lungs and the capillaries of the blood stream. Air moves according to pressure differences, in which air flows from areas of high pressure to areas of low pressure.

The Ventilation Rate


In respiratory physiology, ventilation rate is the rate at which gas enters or leaves the lung. There are several different terms used to describe the nuances of the ventilation rate.

- Minute Ventilation (V_E): The amount of air entering the lungs per minute. It can be defined as tidal volume (the volume of air inhaled in a single breath) times the amount of breaths in a minute.
- Alveolar Ventilation (V_A): The amount of gas per unit of time that reaches the alveoli (the functional part of the lungs where gas exchange occurs). It is defined as tidal volume minus dead space (the space in the lungs where gas exchange does not occur) times the respiratory rate.
- Dead Space Ventilation (V_D): The amount of air per unit of time that doesn't reach the alveoli. It is defined as volume of dead space times the respiratory rate.

Dead space is any space that isn't involved in alveolar gas exchange itself, and it typically refers to parts of the lungs that are conducting zones for air, such as the trachea and bronchioles.

If someone breathes through a snorkeling mask, the length of their conducting zones increases, which increases dead space and reduces on alveolar ventilation. Feedback mechanisms increase the ventilation rate in such a case, but if dead space becomes too great, they won't be able to counteract the effect.

The ventilation rate is controlled by several centers of the autonomic nervous system in the brain, primarily the medulla and the pons.



The human respiratory system: A complete, schematic view of the human respiratory system with its parts and functions.

Mechanisms of Inhalation

Inhalation is initiated by the activity of the diaphragm and supported by the external intercostal muscles. A normal human respiratory rate is 10 to 18 breaths per minute.

During vigorous inhalation (at rates exceeding 35 breaths per minute), or in approaching respiratory failure, accessory muscles—such as the sternocleidomastoid, platysma, and the scalene muscles of the neck—are recruited to help sustain the increased respiratory rate. Pectoral muscles and latissimus dorsi are also accessory muscles for the activity of the lungs.

Under normal conditions, the diaphragm is the primary driver of inhalation. When the diaphragm contracts, the rib cage expands and the contents of the abdomen are moved downward, resulting in a larger thoracic volume and negative pressure (with respect to atmospheric pressure) inside the thorax.

As air moves from zones of high pressure to zones of low pressure, the contraction of the diaphragm allows the air to enter the conducting zone (such as the trachea, bronchioles, etc.), where it is filtered, warmed, and humidified as it flows to the lungs.

Mechanisms of Exhalation

Exhalation is generally a passive process. The lungs have high degree of elastic recoil, so they rebound from the stretch of inhalation and air flows out until the pressures in the lungs and the atmosphere reach equilibrium.

The reason for the elastic recoil of the lung is the surface tension from water molecules on the epithelium of the lungs. A molecule called surfactant (secreted by the alveoli) prevents the surface tension from becoming too great and collapsing the lungs.

Active or forced exhalation is achieved by the abdominal and the internal intercostal muscles. During this process, air is forced or exhaled out. During forced exhalation, as when blowing out a candle, the expiratory muscles, including the abdominal muscles and internal intercostal muscles, generate abdominal and thoracic pressure that force air out of the lungs.

Forced exhalation is often used as an indicator to measure airway health, as people with obstructive lung diseases (such as emphysema, asthma, and bronchitis) will not be able to actively exhale as much as a healthy person because of obstruction in the conducting zones from inhlation, or from a loss of elastic recoil of the lungs.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Breathing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Breathing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- passive diffusion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/passive%20diffusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 2716_Symptoms_of_Acidosis_Alkalosis.jpg. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/wikiped..._Alkalosis.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Respira...ogy_in_mammals**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- apneustic center. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/apneustic%20center**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pneumotaxic center. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pneumotaxic%20center**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventilation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 2716_Symptoms_of_Acidosis_Alkalosis.jpg. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/wikiped..._Alkalosis.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory system complete en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...omplete_en.svg**. **License**: ***Public Domain: No Known Copyright***
- Alveoli. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alveoli**. **License**: ***CC BY: Attribution***

This page titled 21.1B: Functional Anatomy of the Respiratory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2: CONDUCTING ZONE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.2A: Nose and Paranasal Sinuses
21.2B: Pharynx
21.2C: Larynx
21.2D: Structures Used in Voice Production
21.2E: Trachea

This page titled 21.2: Conducting Zone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2A: Nose and Paranasal Sinuses

The shape of the nose is determined by the ethmoid bone and the nasal septum.

Learning Objectives

- Describe the anatomy of the nose and paranasal sinuses

Key Points


- The shape of the nose is determined by the ethmoid bone and the nasal septum, which consists mostly of cartilage and separates the nostrils.
- The nose and paranasal sinuses are part of the upper respiratory tract.
- The functions of the nose include the sense of smell and conditioning of inhaled air by warming it and making it more humid.
- Hairs inside the nose prevent large particles from entering the lungs.
- Nasal mucosa and cilia help prevent pathogens and dust from reaching the lungs. Sneezing helps remove foreign particles that irritate the nasal mucosa.
- The paranasal sinuses are air-filled spaces around the nasal cavity that have many possible functions.
- The mucosa of the upper respiratory tract contain antimicrobial proteins that are a barrier component of the innate immune system.

Key Terms

- **nostril**: Either of the two orifices located on the nose (or on the beak of a bird); used as a passage for air and other gases to travel the nasal passages.
- **paranasal sinuses**: Four air-filled spaces around the nasal cavity that perform many functions, such as draining mucus from the nose.

The nose and paranasal sinuses form much of the upper respiratory tract, along with the pharynx. The upper respiratory tract is the entrance to the respiratory system, where air first enters the body. The overall function of the upper respiratory tract is to provide a pathway for air to reach the lower respiratory tract, where gas exchange occurs.

Anatomy and Physiology of the Nose



Internal diagram of the human nose: Air flows in through the nasal passage on the right and out through the nasopharynx on the posterior side.

The external part of the human nose is the protruding part of the face that bears the nostrils. The shape of the nose is determined by the ethmoid bone and the nasal septum.

The ethmoid bone is the bone that separates the nose from the brain, and supports the shape and structure of the nasal and orbital cavities. The nasal septum is wall of cartilage that separates the right and left nostril chambers from each other. On average, the nose of a male is larger than that of a female, due to differences in facial bone structure between genders.


The interior of the nasal cavity is lined with mucous membranes, nasal hairs, and cilia (microscopic hairs), that perform many of the specialized functions of the nose. The macroscopic nasal hairs prevent large particles from reaching the lungs, while the cilia and mucus trap pathogens and dust to take them to the pharynx, where they can be destroyed by digestion.

Another function of the nose is the conditioning of inhaled air, warming it and making it more humid. Sneezing occurs from irritation of the nasal mucus, which expels foreign particles, but can also spread microbial and viral infections between humans.

Finally, the nose has an area of specialized cells that are responsible for smelling, which is considered a nervous system function rather than a respiratory system function.

Anatomy and Physiology of the Paranasal Sinuses

The paranasal sinuses are a group of four, paired, air-filled spaces, lined with respiratory epithelium (ciliated columnar epithelium). These are named according to the bones within which the sinuses lie: surrounding the nasal cavity (maxillary sinuses), above the eyes (frontal sinuses), between the eyes (ethmoid sinuses), and behind the ethmoid bone (sphenoid sinuses).



Locations of the paranasal sinuses: The paranasal sinuses are four airspaces around the nasal cavity.

The functions of the sinuses are not fully understood, but there are many possible functions. The most important function is the sinuses' role in draining mucus from the nasal cavity to the nasopharynx, which helps regulate pressure inside the nasal cavity. This may be a component of the barrier defenses of the innate immune system because of antimicrobial proteins found in the mucosa.

Other possible sinus functions include giving resonance to the voice, supporting the structure of the skull and facial bones, heating and humidifying inhaled air, and protecting the face from injury.

This page titled 21.2A: Nose and Paranasal Sinuses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2B: Pharynx

The human pharynx is part of the digestive system and also the respiratory system.

Learning Objectives

- Describe the respiratory anatomy of the pharynx


Key Points

- The human pharynx (plural: pharynges) is part of the digestive system and also the respiratory system. It is situated immediately posterior to (behind) the mouth and nasal cavity, and superior to (above) the esophagus and larynx.
- The human pharynx is conventionally divided into three sections: the nasopharynx (epipharynx), the oropharynx (mesopharynx), and the laryngopharynx (hypopharynx).
- The Eustachian tubes connect the middle ear to the nasopharynx, and serve to equalize the barometric pressure in the middle ear with that of the ambient atmosphere.
- Because both food and air pass through the pharynx, a flap of connective tissue called the epiglottis closes over the glottis when food is swallowed to prevent food from getting into the lungs.
- The laryngopharynx includes three major sites: the pyriform sinus, postcricoid area, and the posterior pharyngeal wall.
- Tonsils (lymphoid tissue) exist in the pharynx. Two of the major sets of tonsils are the adenoids in the nasopharynx, and the palatine tonsils in the oropharyx.
- The oropharynx is the middle chamber of the pharynx that passes food from the mouth into the laryngopharynx. The nasopharynx opens above it as well.
- The laryngopharynx is the bottom part of the pharynx that marks the branching pathway between the digestive and respiratory systems.

Key Terms

- **nasopharynx**: The upper part of the pharynx that connects the nasal cavity to the throat.
- **tonsils**: Masses of lymphoid tissue found in the pharynx that play a small role in immune system function.
- **laryngopharynx**: The lower part of the pharynx above the larynx and below the oropharynx.
- **oropharynx**: The middle part of the pharynx that connects to the oral cavity and the other two chambers of the pharynx.

The Pharynx




The three main sections of the pharynx: This figure illustrates the three main subdivisions of the pharynx.

The human pharynx (plural: pharynges) is the part of the throat situated immediately posterior to the mouth and nasal cavity, and superior to the esophagus and larynx.

The human pharynx is divided into three sections: the nasopharynx (epipharynx), the oropharynx (mesopharynx), and the laryngopharynx (hypopharynx), which are all innervated by the pharyngeal plexus.

The pharynx is part of both the digestive system and the respiratory system. As a component of the upper respiratory tract, the pharynx is part of the conducting zone for air into the lungs. Therefore, one of its primary functions is to warm and humidify air before it reaches the lungs.

The Nasopharynx



The Pharynx: This is a detailed diagram of the pharynx from Gray's Anatomy, showing the major structures in each part of the pharynx.

The nasopharynx is the upper region of the pharynx. It extends from the base of the skull to the upper surface of the soft palate above the oral cavity. The nasophaynx connects the nasal cavity with the throat.

The nasopharynx connects to the eustachian tubes of the middle ear, which allows the nasopharynx to help balance pressure within the ear. However, it also allows infections to spread easily between the nasopharynx and ear. The nasopharynx contains psuedo-stratified squamous cell epithelia tissue that is ciliated (covered in tiny hairs that move mucus).

The adenoids (pharyngeal tonsils) are a mass of lymphatic tissue found in the roof of the nasopharynx. The adenoids play a minor role in embyonic development and have a minor role in producing T-lymphocytes for the immune system after birth.

The adenoids are often removed in childhood due to infection or hypertrophy (enlargement of the cells in its tissues), which can obstruct the flow of air from the nose to the lung if left untreated. While loss of the adenoids does not make a significant difference in immune system function, the procedure occasionally has complications.

The lateral walls of the nasopharynx are made of the pharyngeal ostia (bone) of the auditory tube, and supported by the torus tubarus, a mound of cartilage tissue from the auditory tube. Two folds arise from the cartilaginous opening of the auditory tube.

The salpingopharyngeal fold is a vertical fold of mucous membrane extending from the inferior part of the torus and is made up of salpingopharyngeus muscle. The salpingopalatine fold is a smaller fold extending from the superior part of the torus to the palate; it contains the levator veli palatini muscle.

Behind the bone of the auditory tube is a deep recess, the pharyngeal recess. Above the adenoid, in the midline, is an irregular flask-shaped depression of the mucous membrane called the pharyngeal bursa.

The Oropharynx

The orpharnyx (mesopharynx) is the middle portion of the pharynx. It lies between the oral cavity, below the nasopharynx, and above the laryngopharynx, and has an opening to each of these other cavities. The anterior wall of the oropharynx consists of the base of the tongue and the superior wall consists of the bottom surface of the soft palate and the uvula.

The oropharynx is lined by non-keratinized squamous stratified epithelium, which is thicker than the epithelium found in other parts of the respiratory tract in order to prevent damage from food, but not as thick as skin as it lacks keratin.

The epiglottis lies between the oropharynx and the laryngopharynx, and it is a flap of elastic cartilage that closes during swallowing to ensure food enters the esophagus rather than the trachea.

The oropharynx contains the palatine tonsils, which are masses of lymphoid tissue found on the lateral walls of the oropharynx. Compared to the adenoids of the nasopharynx, the palatine tonsils contain many folds (called crypts), and aren't ciliated like the adenoids are. These tonsils are also occasionally removed in people with infection or enlargement.

The Laryngopharynx

The laryngopharynx or hypopharynx is the caudal part of the pharynx; it is the part of the throat that connects to the esophagus and trachea. It lies inferior to the epiglottis and marks the division between the respiratory and digestive system pathways.

During swallowing, the epiglottis closes over the trachea and air passage temporarily stops. The laryngopharynx naturally continues into the esophagus tissue and is made up of a similar type of stratified squamous epithelium tissue.

The layngopharynx itself has a few important demarcations and regions. The formal superior boundary that separates the laryngopharynx from the oropharynx is at the level of the hyoid bone.

The laryngopharynx includes three major regions: the pyriform sinus, the postcricoid area, and the posterior pharyngeal wall, which are separated by small folds of cartilage. Unlike the nasopharynx and oropharynx, there are no tonsils in the laryngopharynx.

This page titled 21.2B: Pharynx is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2C: Larynx

The larynx is an organ in the neck involved in breathing, sound production, and protecting the trachea against food aspiration.

Learning Objectives

- Describe the anatomic structure of the larynx


Key Points

- In adult humans, the larynx is found in the anterior neck at the level of the C3–C6 vertebrae and consists of nine cartilages: three single (epiglottic, thyroid, and cricoid) and three paired (arytenoid, corniculate, and cuneiform).
- Its interior can be divided in supraglottis, glottis, and subglottis.
- The larynx houses the vocal folds—commonly but improperly termed the vocal cords. They are situated just below where the tract of the pharynx splits into the trachea and the esophagus; they are essential for phonation.
- The vocal folds are closed together by adducting the arytenoid cartilages, so that they vibrate (see phonation).
- The vocal folds are controlled by the action of the vagus nerve.
- The larynx closes and rises during swallowing to move the epiglottis over the trachea.
- The larynx closes during a coughing reflex to protect the lungs from inhaling something that could damage it, and to remove foreign material from the trachea and lungs.

Key Terms

- **vocal cords**: Two folds of tissue located in the larynx that vibrate when air passes over them, producing the sound waves associated with talking and singing.
- **cough reflex**: This occurs when the larynx closes as it forces air out of the lungs to protect the lungs from aspirated materials.
- **larynges**: Part of the respiratory tract between the pharynx and the trachea, having walls of cartilage and muscle and containing the vocal cords enveloped in folds of mucous membrane.

The larynx (plural: larynges), commonly called the voice box, is an organ in the neck of humans and most animals that is involved in breathing, sound production, coughing, and protecting the trachea against food aspiration during eating.



External view of the larynx: This figure is a detailed view of the external aspect of the larynx.

Anatomy of the Larynx

In adult humans, the larynx is found in the anterior neck at the level of the C3–C6 vertebrae in the backbone. It connects the inferior part of the pharynx (laryngopharynx) with the trachea. The laryngeal skeleton consists of three single cartilages (thyroid, epiglottic, and cricoid).

1. The thyroid cartilage is particularly notable for forming the Adam's apple, the visible bulge made by the larynx when looking at the throat, and protects the larynx from injury.
2. The epiglottic cartilage is the body of the epiglottis itself that connects to the larynx from above.
3. The cricoid cartilage connects the larynx to the trachea from below.

There are also three sets of cartilages that are paired on either side of the larynx (arytenoid, corniculate, and cuneiform) that allow the position of the larynx to move during voice production.

The larynx connects to the hyoid bone (the bone that forms the floor of the mouth) from above. The larynx extends vertically from the tip of the epiglottis to the border of the cricoid cartilage that marks the formal beginning of the trachea.

The interior of the larynx consists of three regions, the supraglottis, glottis, and subglottis. The glottis is the midsection that contains the vocal folds (folds of muscular epithelium), while the supraglottis and subglottis are the areas of the larynx that are above and below the glottis respectively. In newborn infants, the larynx is initially at the level of the C2–C3 vertebrae, but descends as the child grows.

The glottis consists of two pairs of mucosal folds. These folds are false vocal folds (vestibular folds) and true vocal folds (folds). The false vocal folds are covered by respiratory epithelium, while the true vocal folds are covered by stratified squamous epithelium.

The false vocal folds are not responsible for sound production, but rather for resonance. These false vocal folds do not contain muscle, while the true vocal folds do have skeletal muscle. The two sets of folds are separated by the vocal ligament, with the false vocal folds above, and the true vocal cords below the ligament. The true vocal folds are often referred to as the vocal cords, however the folds technically aren't cords.

Physiology of the Larynx

The most notable and unique function of the larynx is phonation (voice production). The vocal folds of the larynx have two positions, open and closed. During breathing the folds remain open, but they close during swallowing or phonation.

When air from the lungs passes through closed folds during exhalation, the folds vibrate and create sound. The pitch produced depends on the length and tightness of the vocal folds.

The vagus nerves innervate the larynx and signal the muscles and paired cartilage (the arytenoid) of the larynx to work together to open and close the vocal folds as well as change their length and tension to alter pitch. Longer vocal folds have a lower pitch, which is part of the reason why men have deeper voices compared to women, because their larger larynxes have longer vocal folds.

Besides phonation, there are a few other important functions of the larynx. The folds of the larynx close and move upwards during swallowing, which causes the epiglottis to close off the trachea. This helps prevent aspiration of food into the lungs or choking from a blockage of food in the trachea.

The larynx closes off during coughing to help prevent harmful gasses from entering the lungs. During a cough reflex, air is forced out of the lungs, which can remove accumulated mucus, fluid, or blood from the lungs during injury, infection, or cancer of the lungs, as well as food or objects in the trachea during choking.

Finally, the larynx can be signaled to open its folds wider than usual to increase the flow of air into and out of the lungs during heavy breathing when the body requires more oxygen.

This page titled 21.2C: Larynx is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2D: Structures Used in Voice Production

Voices produce sounds through a steady flow of air through the larynx, which causes vibrations and creates fluctuations in air pressure.

Learning Objectives

- Describe the anatomy of voice production structures in the respiratory system

Key Points

- The three basic mechanisms of voice production are air supply, vibration, and resonance.
- Pressure and air-flow speed through the larynx determine the strength and volume of voice.
- The articulation of consonants involves parts of the vocal tract obstructing phonation, and can be active or passive.
- A vowel is any articulation that comes from an open vocal tract.
- Vowel articulation depends mainly on the shape of the lips, position of the tongue, but the shape of the vocal folds are involved as well.

Key Terms

- **Resonance**: The amplication of vibration by the structures of the upper respiratory tract, which can also influence the quality or tone of the sound.
- **Articulation**: The process by which raw phonation from the vocal cords is refined into specific sounds, such as consonants and vowels.
- **glottis**: An organ of speech located in the larynx and consisting of the true vocal cords and the opening between them.

Voice production is a complex process with many different layers and intricacies. The three basic mechanisms of voice production are air supply, vibration, and resonance.

Passive and active articulation shapes and refines phonation (vocal sound production) into the sounds and words used in communication. Voice production is an important physiological process because it enables complex communication between humans.

While the brain is responsible for higher organization and understanding language, the structures of the respiratory system are largely responsible for the production of sound itself.

Basic Mechanisms of Voice Production

Sound is produced by a combination of different structures of the respiratory system working together to create and resonate a sound. There are three basic mechanisms by which the human body produces a voice.

1. Air Supply: In order for voice to be produced, air must flow through the vocal folds. The supply of air for phonation comes from the lungs, and the speed and pressure by which is flows through the vocal folds is determined by the diaphragm. The speed of air flow also determines the strength and loudness of the voice.
2. Vibration: The vocal folds in the glottis of the larynx vibrate as air passes through them. The vibration creates changes in air pressure that manifest as audible sound waves. They only vibrate if the vocal folds are in the closed position, when the folds are held together by the movement of ayrtenoid cartilage. The pitch of the vibration depends on the length and tension of the vocal folds, which can be altered by muscle action.
3. Resonance: The structures of the upper respiratory tract—particularly the soft palate of the mouth, the nasopharynx, and the paranasal sinuses —resonate and amplify the vibration of the vocal folds, making the sound louder and changing its tone. It works similarly to the way the sounding board of a guitar amplifies the vibration of the strings.

These basic mechanisms work together to create the voice. If they are altered, the produced voice will also be altered as well.

For example, during loud voice production, such as shouting or singing, a greater air supply and greater pressure for the flow of air through the vocal folds is required to produce the louder sound. The diaphragm must contract harder to support this greater flow of air compared to normal speech.

Similarly, whispering takes less air compared to normal speech, because the sound produced during whispering is much weaker in comparison.

Articulation

Articulation is the process by which phonation is refined into the specific consonants and vowels used to form words. The articulation of consonants occurs at a point of either active or passive articulation, which is a place in the vocal tract where an obstruction stops the sound.

After the sound is obstructed, the pressure from the air builds based on the shape of that obstruction, which changes the sound into the form it is vocalized as. Vowels are articulated sounds that do not come from obstruction, and instead come from an open vocal tract.

Passive Place of Articulation

The passive place of articulation is the place on the more stationary part of the vocal tract where the articulation occurs. It can be anywhere from the lips, upper teeth, gums, roof of the mouth, or the back of the throat. These areas are passive because no specific action or activity is involved within that area to pronounce the consonant.

Passive articulation is considered a continuum because the obstruction of many different places is needed to produce most of the consonants. There are also several different combinations of areas that can produce the same consonant; for example, many languages may distinguish consonants by articulating them in different areas. Passive places of articulation include:

- The upper lip (labial).
- The upper teeth, either on the edge of the teeth or inner surface (dental).
- The alveolar ridge, the gum line just behind the teeth (alveolar).
- The back of the alveolar ridge (post-alveolar).
- The hard palate on the roof of the mouth (palatal).
- The soft palate further back on the roof of the mouth (velar).
- The uvula hanging down at the entrance to the throat (uvular).
- The throat itself, also known as the pharynx (pharyngeal).
- The epiglottis at the entrance to the windpipe, above the voice box (epiglottal).

Active Place of Articulation

The articulatory gesture of the active place of articulation involves the more mobile part of the vocal tract. This is typically some part of the tongue or lips. It is considered active because these areas change the consonant pronounced by moving or changing.


The active places of articulation are not considered a continuum (unlike passive articulation) because they work independently of each other, but they have the capacity to work together for certain consonants. Active places of articulation include:

- The lower lip (labial).
- Various parts of the front of the tongue.
- The back of the tongue. The aryepiglottic folds at the entrance to the larynx (also epiglottal).
- The glottis (laryngeal).

Vowels

A vowel is a sound that comes from an open vocal tract, and does involve strict obstruction of the sound as with consonants. Therefore, there is more variation in the mechanisms used to create vowels compared to consonants. Vowels are mainly

articulated by the shape of the lips, the position of the tongue (both vertical and horizontal), and by the phonation of the larynx itself.



Places of articulation for voice production: Places of articulation (active and passive): 1. Exo-labial (outer part of lip), 2. Endo-labial (inner part of lip), 3. Dental (teeth), 4. Alveolar (front part of alveolar ridge), 5. Post-alveolar (rear part of alveolar ridge & slightly behind it), 6. Pre-palatal (front part of hard palate that arches upward), 7. Palatal (hard palate), 8. Velar (soft palate), 9. Uvular (a.k.a. Post-velar; uvula), 10. Pharyngeal (pharyngeal wall), 11. Glottal (a.k.a. Laryngeal; vocal folds), 12. Epiglottal (epiglottis), 13. Radical (tongue root), 14. Postero-dorsal (back of tongue body), 15. Antero-dorsal (front of tongue body), 16. Laminal (tongue blade), 17. Apical (apex or tongue tip), and 18. Sub-laminal (also known as sub-apical; underside of tongue)

This page titled 21.2D: Structures Used in Voice Production is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.2E: Trachea

The trachea, or windpipe, is a tube that connects the pharynx or larynx to the lungs, allowing the passage of air.

Learning Objectives


- Describe the anatomic structure of the trachea

Key Points

- The trachea is lined with pseudostratified ciliated columnar epithelium cells with goblet cells that produce mucus.
- There are about 15 to 20 C-shaped cartilaginous rings that reinforce the anterior and lateral sides of the trachea to protect and maintain the airway, leaving a membranous wall (pars membranacea) dorsally without cartilage where the C-shape is open.
- The cartilaginous rings are C-shaped to allow the trachea to collapse slightly at the opening so that food can pass down the esophagus.
- The trachealis muscle connects the ends of the open part of the C-shaped rings and contracts during coughing, reducing the size of the lumen of the trachea to increase the air flow rate.
- The esophagus lies posteriorly to the trachea.
- The mucociliary escalator helps prevent pathogens from entering the lungs.
- The trachea is part of the conducting zone and contributes to anatomical dead space.

Key Terms

- **cilia**: Tiny, hair-like projections from a cell.
- **mucociliary escalator**: The ladder formed by mucus and cilia in the trachea that pushes mucus up the trachea and into the pharynx to prevent mucus pathogens from entering the lungs.
- **anatomical dead space**: The space in the respiratory tract that isn't involved in alveolar ventilation and is part of the normal conducting zone of the respiratory system.



The trachea: This is the trachea in relation to the rest of the respiratory system.

The trachea, or windpipe, is a tube that connects the pharynx or larynx to the lungs, allowing the passage of air. It is lined with


pseudostratified ciliated columnar epithelium cells with goblet cells that produce mucus. The trachea is part of the conducting zone for air into and out of the lungs.

Anatomy of the Trachea

The trachea is a long tube that extends from the pharynx and larynx to the bronchi of the lungs. It typically has an inner diameter of about 25.4 millimeters (1.00 in) and a length of about 10 to 16 centimeters.

The trachea commences at the lower border of the larynx, level with the sixth cervical vertebra, and bifurcates into the primary bronchi at the vertebral level of thoracic vertebra T5, or up to two vertebrae lower or higher, depending on breathing.

At the top of the trachea and bottom of the larynx is the cricoid cartilage, the only complete ring of cartilage in the trachea. Extending downward throughout the length of the tube are about fifteen to 20 C-shaped cartilaginous rings that reinforce the outer structure and shape of the trachea—the open part of each C-shaped ring reveals a membranous wall on the inside of the trachea.



Histology of the Trachea: A cross section of the trachea, showing the hyaline cartilage, mucus glands, and ciliated epithelium.

The cartilage of the trachea is considered hyaline cartilage: simple, transparent, and made primarily of collagen. The trachealis muscle connects the open ends of the C-shaped rings of cartilage and contracts during coughing, reducing the size of the lumen of the trachea to increase the air flow rate.

The esophagus lies behind the trachea. The C-shaped cartilaginous rings allow the trachea to collapse slightly at its opening, so food can pass down the esophagus after swallowing.

The epiglottis closes the opening to the larynx during swallowing to prevent swallowed matter from entering the trachea.

Physiology of the Trachea

This mucus and cilia of the trachea form the mucociliary escalator, which lines the cells of the trachea with mucus to trap inhaled foreign particles. The cilia then waft upward toward the larynx and the pharynx, where it can be either swallowed into the stomach (and destroyed by acid) or expelled as phlegm.

The mucociliary escalator is one of the most important functions of the trachea and is also considered a barrier component of the immune system due its role in preventing pathogens from entering the lungs. The epithelium and the mucociliary ladder can be damaged by smoking tobacco and alcohol consumption, which can make pneumonia (an infection of the alveoli of the lungs) from bacteria in the upper respiratory tract more likely to occur due to the loss of barrier function.

As a part of the conducting zone of the lungs, the trachea is important in warming and moistening air before it reaches the lungs. The trachea is also considered a part of normal anatomical dead space (space in the airway that isn't involved in alveolar gas exchange) and its volume contributes to calculations of ventilation and physiological (total) dead space. It is not considered alveolar dead space, a term that refers to alveoli that don't partake in gas exchange due to damage or lack of blood supply.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human nose. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_nose**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Paranasal sinuses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Paranasal_sinuses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- frontal sinuses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/frontal%20sinuses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- maxillary sinuses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/maxillary%20sinuses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nostril. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nostril**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- TE-Nose diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:TE-Nose_diagram.svg**. **License**: ***CC BY: Attribution***
- Illu nose nasal cavities. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...l_cavities.jpg**. **License**: ***Public Domain: No Known Copyright***
- Human pharynx. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_pharynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...atory-membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- laryngopharynx. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/laryngopharynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glottis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glottis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pharynges. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pharynges**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- TE-Nose diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:TE-Nose_diagram.svg**. **License**: ***CC BY: Attribution***
- Illu nose nasal cavities. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Il...l_cavities.jpg**. **License**: ***Public Domain: No Known Copyright***
- Illu pharynx. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pharynx.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gray994. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray994.png**. **License**: ***Public Domain: No Known Copyright***
- Larynx. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Larynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- larynges. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/larynges**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- voice box. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/voice_box**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vocal cords. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/vocal_cords**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- TE-Nose diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:TE-Nose_diagram.svg**. **License**: ***CC BY: Attribution***
- Illu nose nasal cavities. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_nose_nasal_cavities.jpg**. **License**: ***Public Domain: No Known Copyright***
- Illu pharynx. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Illu_pharynx.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gray994. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray994.png**. **License**: ***Public Domain: No Known Copyright***

- Larynx external en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:La...xternal_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Voice production. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Voice_production. **License**: *CC BY-SA: Attribution-ShareAlike*
- glottis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glottis. **License**: *CC BY-SA: Attribution-ShareAlike*
- cartilage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cartilage. **License**: *CC BY-SA: Attribution-ShareAlike*
- TE-Nose diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:TE-Nose_diagram.svg. **License**: *CC BY: Attribution*
- Illu nose nasal cavities. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_nose_nasal_cavities.jpg. **License**: *Public Domain: No Known Copyright*
- Illu pharynx. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_pharynx.jpg. **License**: *Public Domain: No Known Copyright*
- Gray994. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray994.png. **License**: *Public Domain: No Known Copyright*
- Larynx external en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Larynx_external_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Places of articulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Pl...ticulation.svg. **License**: *Public Domain: No Known Copyright*
- Vertebrate trachea. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vertebrate_trachea. **License**: *CC BY-SA: Attribution-ShareAlike*
- windpipe. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/windpipe. **License**: *CC BY-SA: Attribution-ShareAlike*
- cilia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cilia. **License**: *CC BY-SA: Attribution-ShareAlike*
- TE-Nose diagram. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:TE-Nose_diagram.svg. **License**: *CC BY: Attribution*
- Illu nose nasal cavities. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_nose_nasal_cavities.jpg. **License**: *Public Domain: No Known Copyright*
- Illu pharynx. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Illu_pharynx.jpg. **License**: *Public Domain: No Known Copyright*
- Gray994. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray994.png. **License**: *Public Domain: No Known Copyright*
- Larynx external en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Larynx_external_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Places of articulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Places_of_articulation.svg. **License**: *Public Domain: No Known Copyright*
- Illu conducting passages. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Il...g_passages.svg. **License**: *Public Domain: No Known Copyright*
- Gray964.png. **Provided by**: Wikipedia Commons. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f7/Gray964.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 21.2E: Trachea is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.3: RESPIRATORY ZONE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

21.3A: Bronchi and Subdivisions

21.3B: Alveoli

This page titled 21.3: Respiratory Zone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.3A: Bronchi and Subdivisions

Learning Objectives

- Illustrate the anatomical structure of the bronchi and their subdivisions

A bronchus (plural bronchi, adjective bronchial) is a passage of airway in the respiratory tract that conducts air into the lungs. The bronchus branches into smaller tubes called bronchioles. The bronchi and bronchioles are considered anatomical dead space, like the trachea and upper respiratory tract, because no gas exchange takes place within this zone.

Anatomy of the Bronchi

The human trachea divides into two main bronchi (also called mainstem bronchi), that extend laterally (but not symmetrically) into the left and right lung respectively, at the level of the sternum. The point where the trachea divides into the bronchi is called the **carina**. The right main bronchus is wider, shorter than the left main bronchus, which is thinner and longer. The right main bronchus subdivides into three lobar bronchi, while the left main bronchus divides into two. The lobar bronchi (also called secondary bronchi) divide into tertiary bronchi, each of which supplies air to a different bronchopulmonary segment.

A bronchopulmonary segment is a distinct region of the lung separated from the rest of the lung by connective tissue. Each bronchopulmonary segment forms a discrete functional unit in the lung that is independent of the other segments. This property allows a bronchopulmonary segment to be surgically removed without affecting other segments.

There are 10 segments in the right lung and 8 to 9 segments in the left lung due to anatomical differences. The segmental bronchi divide into many primary bronchioles that divide into terminal bronchioles. Each terminal bronchiole then gives rise to several respiratory bronchioles, which go on to divide into two to 11 alveolar ducts. There are five or six alveolar sacs associated with each alveolar duct. The alveolus is the smallest anatomical unit of the lung, and the site of gas exchange between the lung and the bloodstream.

Histology

The histology of the bronchi are largely similar to that of the trachea. There is hyaline (transparent and consisting of collagen) cartilage present in the bronchi, in rings that are more irregular than those in the trachea. There are also small plates and islands of hyaline cartilage in the primary and terminal bronchioles. Smooth muscle is present continuously around the bronchi (similar to the trachealis muscle of the trachea) and is innervated with the parasympathetic nervous system.

The amount of bronchial smooth muscle increases as the amount of hyaline cartilage decreases as the bronchi become smaller further into the lungs. The mucous membrane lining the bronchi also undergoes a transition—from ciliated pseudostratified columnar epithelium to simple cuboidal epithelium to simple squamous epithelium further into the lungs.


Physiology of the Bronchi

Like the trachea, the bronchi and bronchioles are part of the conducting zone, so they moisten and warm air and contribute to the volume of anatomical dead space. The bronchi and bronchioles are also part of the mucociliary escalator that removes mucus and pathogens from the lungs.

A unique characteristic of the bronchi and bronchioles is **bronchoconstriction**, in which the smooth muscle of the bronchi or bronchioles tightens. This leads to coughing, wheezing, and dyspnea (shortness of breath).

It is caused by activation of the parasympathetic nervous system and release of acetylcholine in the bronchi, as well as by overproduction of mucus or allergic reactions and inflammation. It is a symptom of diseases such as bronchitis (chronic inflammation and mucus production in the bronchi) and asthma (an acute attack of bronchoconstriction, often allergic). Both cause obstruction of the airways and make it more difficult to breathe.

Bronchoconstriction is treated with anti-inflammatory drugs, such as corticosteroids, and prevented by maintaining lung health, such as through avoiding smoking, air pollution, and airborne allergens.



The complete respiratory system: This figure details the respiratory system including the bronchi and its many subdivisions.

Key Points

- The human trachea (windpipe) divides into two main bronchi (also called mainstem bronchi), at the anatomical point known as the carina.
- The right main bronchus is wider and shorter than the left main bronchus. The right main bronchus subdivides into three lobar bronchi and the left main bronchus divides into two.
- The lobar bronchi divide into tertiary bronchi, also known as segmentalinic bronchi, each of which supplies a bronchopulmonary segment.
- The segmental bronchi divide into many primary bronchioles that divide into terminal bronchioles, each of which then gives rise to several respiratory bronchioles, which go on to divide into and terminate in tiny air sacs called alveoli.
- The mucous membrane of the primary bronchi is initially lined by ciliated pseudostratified columnar epithelium, but eventually the lining transitions to simple cuboidal epithelium, and then to simple squamous epithelium.
- The bronchi are part of the conducting zone and contribute to anatomical dead space.
- Bronchoconstriction is the tightening of the smooth muscle of the bronchi from a variety of causes, which makes it more difficult to breathe.

Key Terms

- **bronchus**: Either of the two airways that are the primary branches of the trachea, leading directly into the lungs.
- **bronchoconstriction**: The tightening of the smooth muscle of the bronchi due to parasympathetic nervous system stimulation, excess mucus production, inflammation, or allergic reactions.
- **bronchopulmonary segment**: A distinct functional region of the lung that is separated from the rest of the lung by connective tissue.

This page titled 21.3A: Bronchi and Subdivisions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.3B: Alveoli

Alveoli are hollow cavities in the lung that perform gas exchange with the blood.

Learning Objectives

- Describe the alveoli of lungs

Key Points

- An alveolus is an anatomical structure that has the form of a hollow cavity and is found in the lung parenchyma (tissue inside the lung).
- The pulmonary alveoli are the terminal ends of the respiratory tree that outcrop from either alveolar sacs or alveolar ducts; both are sites of gas exchange.
- The alveolar membrane is the gas-exchange surface. Carbon-dioxide-rich blood is pumped from the rest of the body into the alveolar blood vessels where, through diffusion, it releases its carbon dioxide and absorbs oxygen.
- The alveoli consist of an epithelial layer and an extracellular matrix surrounded by capillaries. In some alveolar walls there are pores between alveoli called pores of Kohn.
- Great alveolar cells secrete pulmonary surfactant to lower the surface tension of water to maintain the lungs elastic recoil.
- Insufficient pulmonary surfactant in the alveoli can contribute to the collapse of a lung.
- Alveolar dead space occurs when an alveolus can't participate in gas exchange due to damage, injury, scarring, or infection. It is a common symptom of many lung diseases.

Key Terms


- **alveolar dead space**: Alveolar space that can no longer contribute to gas exchange due to damage or scarring.
- **extracellular matrix**: A fluid space secreted by cells (including alveolar cells) for structural and functional support.
- **Pulmonary surfactant**: Pulmonary surfactant is a surface-active lipoprotein complex formed by type II alveolar cells that maintains the elastic recoil of the lung.

An alveolus is an anatomical structure that has the form of a hollow cavity. Its plural is alveoli, from the Latin alveolus, meaning little cavity.

Found in the lung parenchyma, the pulmonary alveoli are the terminal ends of the respiratory tree that outcrop from either alveolar sacs or alveolar ducts; both are sites of gas exchange with blood.

The alveolar membrane is the gas-exchange surface. Carbon-dioxide-rich blood is pumped from the rest of the body into the alveolar blood vessels where, through passive diffusion, it releases its carbon dioxide and absorbs oxygen into the blood vessels.

Anatomy of the Alveoli



Pulmonary alveolus: A diagram of the pulmonary alveolus.

The alveoli are located in the respiratory zone of the lungs, at the distal termination of the alveolar ducts. These air sacs are at the end points of the respiratory tract.

There are approximately 700 million alveoli in the lungs, covering a total surface area of about 70 m^2, which is a considerably larger surface area relative to volume. The large surface area makes gas exchange with the bloodstream more efficient.

The alveoli are highly elastic, so the alveoli can stretch as they are filled with air during inhalation. They then spring back during exhalation in order to expel the carbon-dioxide-rich air.

Histology

The alveoli consist of an extremely thin epithelial layer and an extracellular matrix (a fluid space made of collagen and elastin that contains no cells); it is surrounded by many capillaries, the tiniest type of blood vessel.

The fluid extracellular matrix supports the structure of the alveoli and allows gases to dissolve in water, making passive diffusion of those gases with the capillaries possible. In some alveolar walls there are pores between alveoli called the pores of Kohn, that connect alveoli in order to equalize air pressure between the different sacs of an alveolus.

There are two major types of epithelial cells found in alveoli (pneumocytes):

- Type I (Squamous Alveolar) cells: These form the structure of an alveolar wall. They are extremely thin, and permeable, which facilitates gas exhange with the capillaries. They can't undergo mitosis, making them vulnerable to injury.
- Type II (Great Alveolar) cells: These are the site of surfactant production in the lungs, making them critical for maintaining the elastic recoil of the lung. They are more common than type I cells and can undergo mitosis, and may even proliferate into new type I cells when necessary.

Besides these epithelium cells, there are many macrophages found in the alveoli that provide immune system defense of the alveoli from pathogens and foreign material.

Physiology of the Alveoli

The surfactant produced by type II epithelial cells is very important for maintaining the elastic recoil of the lungs. It is a lipoprotein with hydrophilic and hydrophobic ends that reduce the amount of surface tension from water in the lungs. Without

surfactant, the surface tension would cause the lungs to collapse during exhalation, making normal breathing impossible.

Surfactant is first produced by human lungs between 24 and 28 weeks in the womb, and many infants born prematurely do not have enough surfactant to breathe on their own after birth. Surfactant replacement therapy is necessary to save the lives of these premature births.

The alveoli are the site of alveolar ventilation, and are not normally considered dead space. However, alveoli that are injured and can no longer contribute to gas exchange become alveolar dead space.

This is a common occurrence in people with lung diseases like COPD (chronic pulmonary obstructive disorder, i.e., emphysema and bronchitis) or restrictive lung diseases like pulmonary fibrosis, in which scarring of the lung tissue hinders gas exchange in the alveoli, or lung infections like pnuemonia.

Physiological dead space is the sum of normal anatomical dead space and alveolar dead space, and can be used to determine the rate of ventilation (gas exchange) in the lungs. When any type of dead space increases, the rate of ventilation in the lungs will decrease.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Bronchi. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bronchi**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tertiary bronchi. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/tertiary%20bronchi**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bronchus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/bronchus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bronchopulmonary segment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/broncho...nary%20segment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory system complete numbered. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Re...e_numbered.svg**. **License**: ***Public Domain: No Known Copyright***
- Alveoli. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alveoli**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ung-parenchyma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...ary-surfactant**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology.../pores-of-kohn**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory system complete numbered. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Re...e_numbered.svg**. **License**: ***Public Domain: No Known Copyright***
- Alveoli. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alveoli**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 21.3B: Alveoli is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.4: THE LUNGS AND PLEURAE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

21.4A: Lungs

21.4B: Lobes, Fissures, and Lobules

21.4C: Blood Supply to the Lungs

This page titled 21.4: The Lungs and Pleurae is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.4A: Lungs

The lungs are an essential organ which is necessary for the exchange of oxygen and carbon dioxide from the body.

Learning Objectives

- Describe the structure, principal function, and location of the lungs

Key Points

- The principal function of the lungs is to transport oxygen from the atmosphere into the bloodstream and to release carbon dioxide from the bloodstream into the atmosphere.
- Each alveolus is tightly wrapped in blood vessels and it is here that gas exchange actually occurs.
- The diaphragm contracts during inhalation, and relaxes during exhaltion.
- The pleural cavity is the fluid-filled space between the two layers of the pluera, and provides a space for the lung to expand against during inhalation.
- Too little ventilation indicates a shunt (lack of air supply relative to perfusion), while too little alveolar perfusion indicates alveolar dead space from damage or disease. These are both considered V/Q ratio mismatches.

Key Terms


- **pleura**: The double-layered membranous lining of the thoracic cavity that covers the lungs.
- **V/Q ratio**: The ratio of ventilation and blood perfusion of the lungs. Its balance is an important indicator of lung function and efficiency.

The lung is the essential organ of the respiratory system. In mammals and more complex life forms, the two lungs are located near the backbone on either side of the heart.

Their principal function is to transport oxygen from the atmosphere into the bloodstream, and to release carbon dioxide from the bloodstream into the atmosphere. Air travels through the mouth or nose into the pharynx, larynx, trachea, and bronchi and bronchioles in order to reach the lungs.

The exchange of gases is performed by the alveoli, the functional units of the lungs.

Anatomy of the Lungs



The human lungs flank the heart and great vessels in the chest cavity: The human lungs flank the heart and great vessels in the chest cavity.


The lungs are found in the thoracic cavity, and extend laterally into the right and left halves around the heart. The rib cage is a structure of bones that surrounds and protects the thoracic cavity, with 12 ribs protecting each of the two lungs.

The vertebral column is posteriorly attached to the ribcage and the lungs. The lungs are cone-shaped—the apex refers to the top of the lung, while the base refers to the bottom of the lung. The base is curved inward to allow it to rest on the diaphragm.

The diaphragm, a sheet of skeletal muscle that lies beneath the lungs, plays an important role in inhalation. The right lung is larger than the left lung, and the left lung contains the cardiac notch, a concave impression that the heart lies against.

The outer layer of the lungs are the pleura, a type of mesothelium (membrane tissue) that surrounds the lung and attaches it to the thoracic cavity. The pleura contain two layers, the outer parietal pleura (attached to the thoracic cavity), and the inner visceral pleura (covers the lungs).

The pleural cavity is the fluid-filled space between the parietal and visceral pleura, and provides room for the lung to expand during inhalation. The fluid inside the pleural cavity protects against irritation during inhalation as well.



Interior of the thoracic cavity: This is a transverse view of the thoracic cavity, showing the pleural cavity and the major structures around the lungs.

The interior of the lung contains the alveoli, where gas exchange occurs. The alveoli branch off from the bronchioles and bronchi that connect to the trachea and allow air to pass into the lungs.

The lungs are divided into lobes by fissures on the outer surface of the lung, and divide further into segments and finally into hexagonal lobules, the smallest divisions of the lungs.

Physiology of the Lungs

Breathing is largely driven by the muscular diaphragm at the bottom of the thorax. Contraction of the diaphragm pulls the bottom of the cavity in which the lung is enclosed downward, increasing volume and thus decreasing pressure, causing air to flow into the airways and into the lungs.

During normal breathing, expiration is passive— no muscles are contracted—and the diaphragm relaxes from its contracted state. The elastic recoil of the lungs automatically pulls the lungs inward during exhalation. The rib cage itself is also able to expand and contract to some degree during breathing, through the action of other respiratory and accessory respiratory muscles. As a result, air is transported into or expelled out of the lungs.

The major function of the lungs is gas exchange, which occurs in the alveoli of the lung. Oxygen dissolves through the extracellular matrix of the alveoli that lets the gas diffuse into the capillaries based on the relative partial pressures of the gasses (gasses flow from areas of high pressure to low pressure). Oxygen passively diffuses into the deoxygenated blood of the capillaries while carbon dioxide passively diffuses out of the deoxygenated blood and into the airways.

Only a relatively small proportion of alveoli in the lungs are perfused with blood and actually take part in gas exchange. The ratio of ventilation in the lungs and perfusion of the lungs (the air and blood supply of the alveoli respectively) is called the V/Q ratio (with Q being perfusion); it is an important indicator of efficiency in the lungs.

The V/Q in a healthy individual is variable but balanced, usually around.8, but can be higher or lower depending on the region of the lung sampled. An imbalance in V/Q ratio is called a mismatch, and indicates a severe problem.

Too low perfusion (and a higher ratio) indicates alveolar dead space, while too low ventilation (and a lower ratio) indicates a shunt, which is a lack of air supply relative to perfusion. V/Q mismatches occur during many different kinds of lung failure, and have many different causes.

This page titled 21.4A: Lungs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.4B: Lobes, Fissures, and Lobules

The lungs are located on either side of the heart and are separated by fissures into lobes, three in the right and two lobes in the left.

Learning Objectives

- Distinguish between the right and left lungs based on their lobes, fissures, and lobules

Key Points

- Human lungs are located in two cavities on either side of the heart and are separated into lobes by fissures.
- The two lungs are not identical. The right lung has three lobes and left has two lobes. They are further divided into segments and then into lobules.
- Lobules are hexagonal divisions of the lungs that are the smallest subdivision visible to the naked eye.
- The right lung is divided by the oblique fissure, which separates the inferior lobe from the middle and superior lobes, and the horizontal fissure, which separates the superior from the middle lobe.
- The human left lung is divided into two lobes, an upper and a lower, by the oblique fissure. It has a cardiac notch, a concave impression molded to accommodate the shape of the heart.
- The lingula is not technically a lobe, but is the left lung equivalent of the right lung's middle lobe.
- The hilium is the root of the lung and contains the structures involved in pulmonary circulation, as well as the pulmonary nerves and lymph vessels.

Key Terms


- **hilium**: The root of the lung that contains the pulmonary veins and arteries that supply blood to the lungs.
- **cardiac notch**: A concave impression molded into the left lung to accommodate the shape of the heart.

The lungs are located in two chambers of the thoracic cavity on either side of the heart. Though similar in appearance, the two lungs are not identical, nor wholly symmetrical.

Fissures are double folds of pleura that divide the lung into lobes. There are three lobes in the right lung and two in the left lung. The lobes are further divided into segments and then into lobules, which are hexagonal divisions of the lungs that are the smallest visible subdivision.

The lobes are further divided into segments and then into lobules, hexagonal divisions of the lungs that are the smallest subdivision visible to the naked eye.

The Right Lung



The lobes of the lungs: The right lung has three lobes and the left lung has two. (Copyright Public Domain)

The right lung is five centimeters shorter than the left lung to accommodate the diaphragm, which rises higher on the right side over the liver; it is also broader. The volume, the total capacity, and the weight of the right lung is greater than that of the left. The right lung is divided into three lobes.

The Upper Lobe (Right Lung)

The upper lobe is the largest lobe of the right lung. It extends from the apex of the lung down to the horizontal and oblique fissures. It bears apical, anterior, and posterior bronchopulmonary segments.


The Middle Lobe (Right Lung)

The middle lobe is the smallest lobe of the right lung, located between the horizontal and oblique fissures. It bears medial and lateral bronchopulmonary segments.

The Lower Lobe (Right Lung)

The lower lobe is the bottom lobe of the right lung. It lies beneath the oblique fissure. It bears medial, lateral, superior, anterior, and posterior bronchopulmonary segments.

The Left Lung



The Left Lung: This has a concave depression that accommodates the shape of the heart, called the cardiac notch.

The human left lung is smaller and narrower that the right lung, and is divided into two lobes, an upper and a lower, by the oblique fissure. The left lung has only two formal lobes because of the space taken up in the left side of the chest cavity by the heart, though it does have the lingula, which is similar to a lobe.

The left lung has a depression on the medial side of its surface called the cardiac notch, a concave impression molded to accommodate the shape of the heart.

The Upper Lobe (Left Lung)

The upper lobe of the left lung contains anterior and apicoposterior bronchopulmonary segments. It is above the oblique fissure.

The Lower Lobe (Left Lung)

The lower lobe of the left lung contains superior, anterior, posterior, medial, and lateral bronchopulmonary segments.

The Lingula

The lingula is not formally considered to be a lobe. It is a small, tongue-like projection of the left lung that is analogous to the middle lobe of the right lung. It contains superior and inferior bronchopulmonary segments.

The Hilium

Above and behind the cardiac impression is a triangular depression named the hilum. The hilum is the root of the lung where that contains structures that supply the lungs with blood, lymph fluid, and innervation, such as the pulmonary vein, pulmonary artery, pulmonary nerves, and lymphatic vessels.

These structures are enclosed by pleura. There is a hilium for each of the lungs found in the mediastinum (backside) of the lungs. The hilium is thinner in the left lung compared to the right lung because it lies between the cardiac notch and the groove for the aorta.

The hilium is important because it is the primary way in which the respiratory system links with the cardiovascular and nervous systems.

This page titled 21.4B: Lobes, Fissures, and Lobules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.4C: Blood Supply to the Lungs

Pulmonary circulation transports oxygen-depleted blood away from the heart to the lungs and returns oxygenated blood back to the heart.

Learning Objectives

- Distinguish between pulmonary and systemic circulation of blood

Key Points

- Pulmonary circulation is the portion of the cardiovascular system that carries oxygen-depleted blood to the lungs from the heart, and returns oxygenated (oxygen-rich) blood back to the heart.
- Systemic circulation is the portion of the cardiovascular system that brings oxygen to the tissues of the body. De-oxygenated blood enters the right atrium. Blood then moves to the right ventricle that pumps blood from the heart to the lungs where it releases carbon dioxide and picks up oxygen.
- Oxygenated blood leaves the lungs through pulmonary veins, completing the pulmonary cycle. This blood enters the left atrium and is then transferred to the left ventricle, which pumps the newly oxygenated blood back into systemic circulation.
- The pulmonary arteries carry deoxygenated blood to the lungs, where it releases carbon dioxide and picks up oxygen during respiration.
- A pulmonary embolism can occur if blood pools in veins of the legs and forms a blood clot due to immobilization. The resulting blood clot can block off the pulmonary artery and cause the alveoli inside the lung to die.

Key Terms

- **pulmonary embolism**: A blockage of the blood supply the lungs by a blood clot.
- **atrium**: An upper chamber of the heart that receives blood from the veins and forces it into a ventricle.
- **ventricle**: The lower chamber of the heart that receives blood from the atrium and pumps it into the arteries.

There are two primary types of circulation in the human.

1. Pulmonary circulation refers to blood supply to the lungs for the purpose of gas exchange.
2. Systemic circulation refers to blood supply to the rest of the body, for the purpose of supplying oxygen to the tissues.

Bronchial circulation (by the bronchial arteries) supplies blood to the tissues of the bronchi and the pleura, and is considered part of systemic circulation.

Pulmonary Circulation

The right side of the heart deals with pulmonary circulation. At the end of systemic circulation, the veins take blood back to the heart through the vena cava.

The vena cava fills the right atrium with blood, which then ejects blood into the right ventricle by passing through the tricuspid valve. After blood fills in the right ventricle, it contracts and pumps the blood through the pulmonary valve, and into the pulmonary arteries.


There are two pulmonary arteries (one for each lung) that bring the deoxygenated blood to the lungs through the hilium. The arteries branch into the capillaries of the alveoli. Capillaries are the thinnest and smallest type of blood vessel, and they supply oxygen to individual tissues everywhere in the human body.

Gas exchange occurs by passive diffusion in the alveoli, so that dissolved oxygen enters the capillaries, while carbon dioxide leaves pulmonary circulation. The oxygenated blood then leaves the lungs through pulmonary veins (also contained in the hilium),

which return the blood to the left side of the heart, completing the cycle of pulmonary circulation.

This blood then enters and fills inside the left atrium, which pumps it through the mitral valve (also called bicuspid) into the left ventricle. The blood fills inside the left ventricle and is then pumped through the aortic valve into the aorta, which marks the beginning of systemic circulation.

Systemic circulation and pulmonary circulation form the overall cycle of the circulatory system: transporting oxygen throughout the body.



Pulmonary circuit: Diagram of pulmonary circulation. Oxygen-rich blood is shown in red; oxygen-depleted blood in blue.

Problems in Pulmonary Circulation

While the cycle of pulmonary and systemic circulation is a well designed and effective system, it is not immune to certain problems. The most serious issue in pulmonary circulation is a pulmonary embolism, which is where a blood clot travels to the lung and causes an infarction of the lung (tissue death from lack of oxygen).

These blood clots typically originate in the deep veins of the legs (part of systemic circulation) as a result of blood pooling from injury or immobilization. As the veins of the leg are on their way to the right side of the heart, the clots are less likely to break up before they reach pulmonary circulation.

When the clot reaches the pulmonary artery, it obstructs the flow of blood into the lung, which causes the alveoli in the effected lung to die as a result. This results in an increase in aveolar dead space and decreased perfusion, (leading to shortness of breath and chest pain) and can be fatal if not treated in time by fibrinolytics (medications that dissolve the clot).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Lung. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lung. **License**: *CC BY-SA: Attribution-ShareAlike*
- bronchi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/bronchi. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray968.png. **Provided by**: Wikipedia Commons. **Located at**: en.Wikipedia.org/wiki/Pleura...le:Gray968.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lung. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lung. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardiac notch. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cardiac%20notch. **License**: *CC BY-SA: Attribution-ShareAlike*
- Right lung. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Right_lung. **License**: *CC BY-SA: Attribution-ShareAlike*
- Oblique fissure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Oblique_fissure. **License**: *CC BY-SA: Attribution-ShareAlike*

- Left lung. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Left_lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lung. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Right Lung. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Right+Lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Left Lung. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Left+Lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray968.png. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/Pleural_cavity%23/media/File:Gray968.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lung. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray973. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray973.png**. **License**: ***Public Domain: No Known Copyright***
- Lung and diaphragm. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Lung_and_diaphragm.jpg**. **License**: ***Public Domain: No Known Copyright***
- Pulmonary circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pulmonary_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- atrium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/atrium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventricle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventricle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray968.png. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/Pleural_cavity%23/media/File:Gray968.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lung. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lung**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray973. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray973.png**. **License**: ***Public Domain: No Known Copyright***
- Lung and diaphragm. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Lung_and_diaphragm.jpg**. **License**: ***Public Domain: No Known Copyright***
- Pulmonary circulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pulmonary_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 21.4C: Blood Supply to the Lungs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.5: MECHANICS OF BREATHING

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.5A: Pressure Changes During Pulmonary Ventilation
21.5B: Inspiration
21.5C: Expiration
21.5D: Breathing Patterns

This page titled 21.5: Mechanics of Breathing is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.5A: Pressure Changes During Pulmonary Ventilation

Ventilation is the rate at which gas enters or leaves the lung.

LEARNING OBJECTIVE

Differentiate among the types of pulmonary ventilation: minute, alveolar, dead space

Key Takeaways

Key Points

- Ventilation is the rate at which gas enters or leaves the lung.
- The three types of ventilation are minute ventilation, alveolar ventilation, and dead space ventilation.
- The alveolar ventilation rate changes according to the frequency of breath, tidal volume, and amount of dead space.
- PA refers to alveolar partial pressure of a gas, while Pa refers to the partial pressure of that gas in arterial blood.
- Gas exchange occurs from passive diffusion because PAO2 is greater than PaO2 in deoxygenated blood.

Key Terms

- **ventilation**: The bodily process of breathing, the inhalation of air to provide oxygen, and the exhalation of spent air to remove carbon dioxide.
- **partial pressure**: The pressure exerted by a gas, either in air or dissolved, that indicates the concentration of that gas.

The Types of Ventilation Rates

In respiratory physiology, the ventilation rate is the rate at which gas enters or leaves the lung. Ventilation is generally expressed as volume of air times a respiratory rate.

The volume of air can refer to tidal volume (the amount inhaled in an average breath) or something more specific, such as the volume of dead space in the airways. The three main types of ventilation rates used in respiratory physiology are:

1. Minute ventilation (VE): The amount of air entering the lungs per minute. It can be defined as VE=Tidal Volume×Breaths Per MinuteVE=Tidal Volume×Breaths Per Minute
2. Alveolar ventilation (VA): The amount of gas per unit of time that reaches the alveoli and becomes involved in gas exchange. It is defined as VA=(Tidal Volume−Dead Space Volume)×Respiratory RateVA=(Tidal Volume−Dead Space Volume)×Respiratory Rate
3. Dead space ventilation (VD): The amount of air per unit of time that is not involved in gas exchange, such as the air that remains in the conducting zones. It is defined as VD=Dead Space Volume×Respiratory RateVD=Dead Space Volume×Respiratory Rate.


Additionally, minute ventilation can be described as the sum of alveolar and dead space ventilation, provided that the respiratory rate used to derive them is in terms of breaths per minute.

The three types of ventilation are mathematically linked to one another, so changes in one ventilation rate can cause the change of the other. This is most apparent in changes of the dead space volume. Breathing through a snorkeling tube and having a pulmonary embolism both increase the amount of dead space volume (through anatomical versus alveolar dead space respectively), which will reduce alveolar ventilation.

Alveolar ventilation is the most important type of ventilation for measuring how much oxygen actually gets into the body, which can initiate negative feedback mechanisms to try and increase alveolar ventilation despite the increase in dead space. In particular, the body will generally attempt to combat increased dead space by raising the frequency of breaths to try and maintain sufficient

levels of alveolar ventilation.

Partial Pressure of Gasses



Gaseous Exchange in the Lungs: Diagram of gas exchange in the lungs.

When gasses dissolve in the bloodstream during ventilation, they are generally described by the partial pressure of the gasses. Partial pressure more specifically refers to the relative concentration of those gasses by the pressure they exert in a dissolved state.

In respiratory physiology, PAO_2 and $PACO_2$,refer to the partial pressures of oxygen and carbon dioxide in the alveoli.

PaO_2 and $PaCO_2$ refer to the partial pressures of oxygen and carbon dioxide within arterial blood. Differences in partial pressures of gasses between the alveolar air and the blood stream are the reason that gas exchange occurs by passive diffusion.

Under normal conditions, PAO_2 is about 100 mmHg, while PaO_2 is 80–100 mmHg in systemic arteries, but 40–50 mmHg in the deoxygenated blood of the pulmonary artery going to the lungs.

Recall that gasses travel from areas of high pressure to areas of low pressure, so the greater pressure of oxygen in the alveoli compared to that of the deoxygenated blood explains why oxygen can passively diffuse into the bloodstream during gas exchange.

Conversely, $PACO_2$ is 35 mmHg, while $PaCO_2$ is about 40–45 mmHG in systemic arteries and 50 mmHg in the pulmonary artery. The partial pressure, and thus concentration of carbon dioxide, is greater in the in the capillaries of the alveoli compared to the alveolar air, so carbon dioxide will passively diffuse from the bloodstream into the alveoli during gas exchange.

Additionally, because $PaCO_2$ is an indicator of the concentration of carbon dioxide in arterial blood, it can be used to measure blood pH and identify cases of respiratory acidosis and alkalkosis.

This page titled 21.5A: Pressure Changes During Pulmonary Ventilation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.5B: Inspiration

Inhalation is the flow of air into an organism that is due to a pressure difference between the atmosphere and alveolus.

LEARNING OBJECTIVE

Outline the mechanics of inspiration

Key Takeaways

Key Points

- In humans, inspiration is the flow of air into an organism from the external environment, through the airways, and into the alveoli.
- Inhalation begins with the onset of a contraction of the diaphragm, which results in expansion of the thoracic and pleural cavities and a decrease in pressure (also called an increase in negative pressure).
- There are many accessory muscles involved in inhalation—such as external intercostal muscles, scalene muscles, the sternocleidomastoid muscle, and the trapezius muscle.
- Breathing only with the accessory muscles instead of the diaphragm is considered inefficient, and provides much less air during inhalation.
- The negative pressure in the pleural cavity is enough to hold the lungs open in spite of the inherent elasticity of the tissue. The thoracic cavity increases in volume causing a drop in the pressure (a partial vacuum) within the lung itself.
- As long as pressure within the alveoli is lower than atmospheric pressure, air will continue to move inwardly, but as soon as the pressure is stabilized air movement stops.

Key Terms

- **inspiration**: The drawing of air into the lungs, accomplished in mammals by elevation of the chest walls and flattening of the diaphragm.
- **accessory muscles**: Muscles that help expand small parts of the thoracic cavity, either working in addition to the diaphragm or substituting for it if the diaphragm becomes injured.
- **intrapleural pressure**: The pressure inside the pleural cavity, which is negative compared to outside air and becomes even more negative during inspiration.

Inspiration refers to inhalation—it is the flow of the respiratory current into an organism. In humans it is the movement of ambient air through the airways and into the alveoli of the lungs.

The Process of Inspiration


Inspiration begins with the contraction of the diaphragm, which results in expansion of the thoracic cavity and the pleural cavity. The pleural cavity normally has a lower pressure compared to ambient air (–3 mmHg normally and typically –6 mmHg during inspiration), so when it expands, the pressure inside the lungs drops.

Pressure and volume are inversely related to each other, so the drop in pressure inside the lung increases the volume of air inside the lung by drawing outside air into the lung. As the volume of air inside the lung increases, the lung pushes back against the expanded pleural cavity as a result of the drop in intrapleural pressure (pressure inside the pleural cavity).

The force of the intrapleural pressure is even enough to hold the lungs open during inpiration despite the natural elastic recoil of the lung. The alveolar sacs also expand as a result of being filled with air during inspiration, which contributes to the expansion inside the lung.

Eventually, the pressure inside the lung becomes less negative as the volume inside the lung increases and, when pressure and

volume stabilize, air movement stops, inspiration ends, and expiration (exhalation) will begin. Deeper breaths have higher tidal volumes and require a greater drop in intrapleural pressure compared to shallower breaths.



Respiratory System: Resistance in any part of the respiratory tract can cause problems.

Accessory Muscles of Inspiration

The diaphragm is the primary muscle involved in breathing, however several other muscles play a role in certain circumstances. These muscles are referred to as accessory muscles of inhalation.

- External intercostal muscles: Muscles located between the ribs that help the thoracic cavity and pleural cavity expand during quiet and forced inspiration.
- Scalene muscles: Muscles in the neck that lift the upper ribs (and thoracic cavity around the upper ribs) to help with breathing. They provide a mechanism for inspiration when the diaphragm is injured and can't contract normally.
- Sternocleidomastoid muscle: Muscles that connect the sternum to the neck and allow for rotation and turning of the head. They can lift the upper ribs like the scalene muscles can.
- Trapezius muscle: Muscles in the shoulders that retract the scapula and expand the upper part of the thoracic cavity.

The accessory muscles assist breathing by expanding the thoracic cavity in a similar way to the diaphragm. However, they expand a much smaller part of the thoracic cavity compared to diaphragm. Therefore they should not be used as the primary mechanism of inhalation, because they take in much less air compared to the diaphragm resulting in a much lower tidal volume.

For example, singers need a lot of air to support the powerful voice production needed for singing. A common problem in novice singers is breathing with the accessory muscles of the neck, shoulder, and ribs instead of the diaphragm, which gives them a much smaller air supply than what is needed to sing properly.

This page titled 21.5B: Inspiration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.5C: Expiration

Exhalation (or expiration) is the flow of the respiratory current out of the organism.

LEARNING OBJECTIVE

Outline the mechanics of expiration

Key Takeaways


Key Points

- In humans, exhalation is the movement of air out of the bronchial tubes, through the airways, to the external environment during breathing.
- Exhalation is a passive process because of the elastic properties of the lungs.
- During forced exhalation, internal intercostal muscles which lower the rib cage and decrease thoracic volume while the abdominal muscles push up on the diaphragm which causes the thoracic cavity to contract.
- Relaxation of the thoracic diaphragm causes contraction of the pleural cavity which puts pressure on the lungs to expel the air.
- Brain control of exhalation can be broken down into voluntary control and involuntary control.

Key Terms

- **Intercostal muscles**: Intercostal muscles are several groups of muscles that run between the ribs, and help form and move the chest wall.
- **exhalation**: The act or process of exhaling, or sending forth in the form of steam or vapor; evaporation.

Expiration, also called exhalation, is the flow of the respiratory current out of the organism. The purpose of exhalation is to remove metabolic waste, primarily carbon dioxide from the body from gas exchange. The pathway for exhalation is the movement of air out of the conducting zone, to the external environment during breathing.



Respiratory System: As the diaphragm relaxes, the pleural cavity contracts, which exerts pressure on the lungs, which reduces

the volume of the lungs as air is passively pushed out of the lungs.

The Process of Expiration

Expiration is typically a passive process that happens from the relaxation of the diaphragm muscle (that contracted during inspiration). The primary reason that expiration is passive is due to the elastic recoil of the lungs. The elasticity of the lungs is due to molecules called elastins in the extracellular matrix of lung tissues and is maintained by surfactant, a chemical that prevents the elasticity of the lungs from becoming too great by reducing surface tension from water. Without surfactant the lungs would collapse at the end of expiration, making it much more difficult to inhale again. Because the lung is elastic, it will automatically return to its smaller size as air leaves the lung.

Exhalation begins when inhalation ends. Just as the plueral cavity's increased negative pressure leads to air uptake during inhalation, the pleural cavity will contract during the exhalation (due to relaxation of the diaphragm),which exerts pressure on the lungs and causes the pressure inside the cavity to be less negative. An increase in pressure leads to a decrease in volume inside the lung, and air is pushed out into the airways as the lung returns to its smaller size. The pleural cavity is so important to breathing because it's pressure changes the volume of the lungs, and it provides a friction-less space for the lung to expand and contract against during breathing.

While expiration is generally a passive process, it can also be an active and forced process. There are two groups of muscles that are involved in forced exhalation.

- Internal Intercostal Muscles: Muscles of the ribcage that help lower the ribcage, which pushes down on the thoracic cavity, causing forced exhalation. *Note that these are not the same as the external intercostal muscles involved in inspiration*.
- Abdominal Muscles: Any number of muscles in the abdomen that exert pressure on the diaphragm from below to expand it, which in turn contracts the thoracic cavity, causing forced exhalation.

This happens due to elastic properties of the lungs, as well as the internal intercostal muscles that lower the rib cage and decrease thoracic volume. As the thoracic diaphragm relaxes during exhalation it causes the tissue it has depressed to rise superiorly and put pressure on the lungs to expel the air.

Control of Expiration

Expiration can be either voluntary or involuntary in order to serve different purposes for the body. These two types of expiration are controlled by different centers within the body.

Voluntary expiration is actively controlled. It is generally defined by holding air in the lungs and releasing it at a fixed rate, which enables control over when and how much air to exhale. It is required for voice production during speech or singing, which requires very specific control over air, or even simpler activities, like blowing out a candle on one's birthday. The nervous system component that controls voluntary expiration is the motor cortex (the ascending respiratory pathway), because it controls muscle movements, but this pathway isn't fully understood, and there are many other possible sites in the brain that may also be involved.

Involuntary expiration is not under conscious control, and is an important component for metabolic function. Examples include breathing during sleep or meditation. Changes in breathing patterns may also occur for metabolic reasons, such as through increased breathing rate in people with acidosis from negative feedback. The principle neural control center for involuntary expiration consists of the medulla oblongata and the pons, which are located in the brainstem directly beneath the brain. While these two structures are involved in neural respiratory control, they also have other metabolic regulatory functions for other body systems, such as the cardiovascular system.

This page titled 21.5C: Expiration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.5D: Breathing Patterns

Breathing is an autonomic process that moves air in and out of the lungs.

LEARNING OBJECTIVE

Describe the process of breathing in humans

Key Takeaways

Key Points

- Breathing patterns consist of tidal volume and respiratory rate in an individual.
- An average breathing pattern is 12 breaths per minute and 500 mL per breath.
- Eupnea is normal breathing at rest.
- There are types of altered breathing patterns that are symptoms of many diseases.
- Altered breathing patterns refer to changes in respiratory rate or amount of air exchanged during breathing, and do not always indicate changes in alveolar ventilation.
- The mechanism of generation of the ventilatory pattern involves the integration of neural signals by respiratory control centers in the medulla and pons.


Key Terms

- **altered breathing patterns**: Abnormal breathing patterns that indicate typically indicate either too fast or too slow respiratory rate or too much or too little tidal volume.
- **tidal volume**: The amount of air displaced or exchanged in a single breath.

Breathing patterns refer to the respiratory rate, which is defined as the frequency of breaths over a period of time, as well as the amount of air cycled during breathing (tidal volume). Breathing patterns are an important diagnostic criteria for many diseases, including some which involve more than the respiratory system itself.

Characteristics of the Breathing Patterns

The respiratory rate is frequency of breaths over time. The time period is variable, but usually expressed in breaths per minute because it that time period allows for estimation of minute ventilation. During normal breathing, the volume of air cycled through inhalation and exhalation is called tidal volume (VT), and is the amount of air exchanged in a single breath. Tidal volume multiplied by the respiratory rate is minute ventilation, which is one of the most important indicators of lung function. In an average human adult, the average respiratory rate is 12 breaths per minute, with a tidal volume of .5 liters and a minute ventilation of 6 liters per minute, though these numbers vary from person to person. Infants and children have considerably higher respiratory rates than adults.



Spirometry curve: The normal respiratory rate refers to the cyclical inhalation and exhalation of tidal volume (VT).

The respiratory rate is controlled by involuntary processes of the autonomic nervous system. In particular, the respiratory centers of the medulla and the pons control the overall respiratory rate based on a variety of chemical stimuli from within the body. The hypothalamus can also influence the respiratory rate during emotional and stress responses.

Normal and Altered Breathing Patterns

Eupnea is the term for the normal respiratory rate for an individual at rest. Several other terms describe abnormal breathing patterns that are indicative of symptoms of many diseases, many of which aren't mainly respiratory diseases. Some of the more common terms for altered breathing patterns include:

- Dyspnea: commonly called shortness of breath. It describes dramatically decreased tidal volume and sometimes increased respiratory rate, leading to a sensation of breathlessness. It is a common symptom of anxiety attacks, pulmonary embolisms, heart attacks, and emphysema, among other things.
- Hypernea: refers to increased volume of air cycled to meet the body's metabolic needs, which may or may not involve a change in frequency of breathing. It is a symptom of exercise and adjustment to high altitude, which are generally not problematic, but can also be seen in those with anemia or septic shock, which is problematic.
- Tachypnea: describes increased respiratory rate. Often a symptom of carbon monoxide poisoning or pnuemonia.
- Bradypnea: describes decreased respiratory rate. Often a symptom of hypertension, heart arryhmias, or slow metabolic rate from hypothyroidism.
- Apnea: Transient stopped breathing that begins again soon afterwards. It is the main symptom of sleep apnea, in which breathing temporarily stops during sleep.

These terms all describe an altered breathing pattern through increased or decreased (or stopped) tidal volume or respiratory rate. It is important to distinguish these terms from hyperventilation and hypoventilation, which refer to abnormalities in alveolar gas exchange (and thus blood pH) instead of an altered breathing pattern, but they may be associated with an altered breathing pattern. For example dyspnea or tachypnea often occur together with hyperventilation during anxiety attacks, though not always.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Pulmonary ventilation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pulmonary_ventilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Modes of mechanical ventilation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Modes_of_mechanical_ventilation%23Pressure_controlled**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ventilation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ventilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Respiratory_physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Inhalation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Inhalation**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- negative pressure. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/negative_pressure. **License**: *CC BY-SA: Attribution-ShareAlike*
- inspiration. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/inspiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- The respiratory system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_respiratory_system. **Project**: Human Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Respiratory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Respiratory_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Exhalation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Exhalation. **License**: *CC BY-SA: Attribution-ShareAlike*
- exhalation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/exhalation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Respiratory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Respiratory_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thoracic diaphragm. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thoracic_diaphragm. **License**: *CC BY-SA: Attribution-ShareAlike*
- Breathing. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Breathing. **License**: *CC BY-SA: Attribution-ShareAlike*
- Control of respiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Control_of_respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- breathing. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/breathing. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lung Volume. **Authored by**: Vihsadas. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:LungVolume.jpg. **License**: *Public Domain: No Known Copyright*
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 21.5D: Breathing Patterns is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.6: FACTORS AFFECTING PULMONARY VENTILATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.6A: Factors Affecting Pulmonary Ventilation: Surface Tension of Alveolar Fluid
21.6B: Factors Affecting Pulmonary Ventilation: Compliance of the Lungs
21.6C: Factors Affecting Pulmonary Ventilation: Airway Resistance

This page titled 21.6: Factors Affecting Pulmonary Ventilation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.6A: Factors Affecting Pulmonary Ventilation: Surface Tension of Alveolar Fluid

The surface tension of alveolar fluid is regulated by pulmonary surfactant, allowing efficient respiration.

Learning Objectives

- Describe surfactant and its role in alveolar fluid surface tension

Key Points

- Type II avleolar epithelial cells secrete pulmonary surfactant to lower the surface tension of water, which helps prevent airway collapse.
- Reinflation of the alveoli following exhalation is made easier by pulmonary surfactant.
- The surfactant reduces surface tension within all alveoli through hydrophilic and hydrophobic forces.
- Insufficient pulmonary surfactant in the alveoli can contribute to atelectasis (collapse of part or all of the lung).
- Premature infants often don't have the capacity to produce enough surfactant to survive on their own.

Key Terms

- **atelectasis**: The collapse of a part of or the whole lung caused by inner factors, rather than a pneumothorax.
- **surfactant**: A lipoprotein in the tissues of the lung that reduces surface tension and permits more efficient gas transport.
- **Surface tension**: The inward force created by films of molecules that can reduce the area of a surface.

EXAMPLES

Elective cesarean sections are becoming more common. One unfortunate consequence of this is that many of the infants delivered by this method are actually slightly physiologically premature. They lack sufficient surfactant to initiate proper breathing, and therefore, go into respiratory distress.

The alveoli are highly elastic structures in the parenchyma of the lungs that are the functional site of gas exchange. As the alveoli fill with air during inhalation they expand, and as air leaves the lung with exhalation, the alvoli return to their non-inflated size. The reason for the elasticity of the alveoli is a protein found in the extracellular matrix of the alveoli, called elastin, as well as the surface tension of water molecules on the alveoli themselves.

Surface Tension in the Lung

Surface tension is the force exerted by water molecules on the surface of the lung tissue as those water molecules pull together. Water (H_2O) is a highly polar molecule, so it forms strong covalent bonds with other water molecules. The force of these covalent bonds effectively creates an inward force on surfaces, such as lung tissue, with the effect of lowering the surface area of that surface as the tissue is pulled together. As the air inside the lungs is moist, there is considerable surface tension within the tissue of the lungs. Because the alveoli of the lungs are highly elastic, they do not resist surface tension on their own, which allows the force of that surface tension to deflate the alveoli as air is forced out during exhalation by the contraction of the pleural cavity.

Pulmonary Surfactant

The force of surface tension in the lungs is so great that without something to reduce the surface tension, the airways would collapse after exhalation, making re-inflation during inhalation much more difficult and less effective. Collapse of the lungs is called alectasis. Fortunately, the type II epithelial cells of the alveoli continually secrete a molecule called surfactant that solves

this problem.

Surfactant is a lipoprotein molecule that reduces the force of surface tension from water molecules on the lung tissue. The main reason that surfactant has this function is due to a lipid called dipalmitoylphosphatidylcholine (DPPC) which contains hydophilic and hydrophobic ends. The hydrophilic ends are water soluable and attach to the water molecules on the surface of the lungs. The hydrophilic ends are water insoluable and face towards the air and pull away from the water. The net result is that the surface tension of the lungs from water is reduced so that the lungs can still inflate and deflate properly without the possibility of collapse from surface tension alone.

As unborn humans grow and develop in the womb, they receive oxygen from the mother, so their lungs aren't fully functional right away. Of particular importance is the fact that they don't produce surfactant until 24 weeks of development and usually don't have enough built up to prevent lung collapse until 35 weeks of development. Therefore prematurely born infants are at a high risk of respiratory distress syndrome from airway collapse, which can cause death if untreated. It is treated through pulmonary surfactant replacement therapy and mechanical ventilator treatment until the infant's lungs are old enough to secrete enough surfactant to survive on their own. Other diseases may cause atelectasis, such as COPD, or any sort of lung trauma and inflammation that involves extensive damage to the pleural cavity or the lung parenchyma.




Diagram of an Alveoli: An alveoli with both cross-section and external views

This page titled 21.6A: Factors Affecting Pulmonary Ventilation: Surface Tension of Alveolar Fluid is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.6B: Factors Affecting Pulmonary Ventilation: Compliance of the Lungs

Lung compliance refers to the magnitude of change in lung volume as a result of the change in pulmonary pressure.

Learning Objectives

- Evaluate the factors involved in lung compliance

Key Points

- A low lung compliance would mean that the lungs would need a greater-than-average change in intrapleural pressure to change the volume of the lungs.
- A high lung compliance would indicate that little pressure difference in intrapleural pressure is needed to change the volume of the lungs.
- Persons with low lung compliance due to obstructive lung diseases tend to take rapid shallow breaths and sit hunched over to make exhalation less difficult.
- Persons with high lung compliance due to restrictive lung diseases tend to have difficulty expanding and deflating the lungs.
- Two factors determine lung compliance: elasticity of the lung tissue and surface tensions at air water interfaces.
- Two factors determine lung compliance – elasticity of the lung tissue and surface tensions at air water interfaces.

Key Terms

- **Lung compliance**: The ability of the lungs and pleural cavity to change in volume based on changes in pressure.

EXAMPLES


Low lung compliance can be the result of interstitial lung diseases resulting from the inhalation of particulate substances such as asbestos (asbestosis) and silicon (silicosis).

Compliance is the ability of lungs and pleural cavity to expand and contract based on changes in pressure. Lung compliance is defined as the volume change per unit of pressure change across the lung, and is an important indicator of lung health and function. Measurements of lung volumes differ at the same pressure between inhalation and exhalation, meaning that lung compliance differs between inhalation and exhalation. Lung compliance can either be measured as static or dynamic based on whether only volume and pressure (static) is measured or if their changes over time are measured as well (dynamic).

Compliance and Elastic Recoil of the Lung

Compliance depends on the elasticity and surface tension of the lungs. Compliance is inversely related to the elastic recoil of the lungs, so thickening of lung tissue will decrease lung compliance. The lungs must also be able to overcome the force of surface tension from water on lung tissue during inflation in order to be compliant, and greater surface tension causes lower lung compliance. Therefore, surfactant secreted by type II epithelial cells increases lung compliance by reducing the force of surface tension.

A low lung compliance means that the lungs are “stiff” and have a higher than normal level of elastic recoil. A stiff lung would need a greater-than-average change in pleural pressure to change the volume of the lungs, and breathing becomes more difficult as a result. Low lung compliance is commonly seen in people with restrictive lung diseases, such as pulmonary fibrosis, in which scar tissue deposits in the lung making it much more difficult for the lungs to expand and deflate, and gas exchange is impaired. Pulmonary fibrosis is caused by many different types of inhalation exposures, such as silica dust.



Pulmonary Fibrosis: Pulmonary fibrosis stiffens the lungs through deposits of scar tissue, decreasing low compliance and making it more difficult for the lungs to inflate and deflate.

A high lung compliance means that the lungs are too pliable and have a lower than normal level of elastic recoil. This indicates that little pressure difference in pleural pressure is needed to change the volume of the lungs. Exhalation of air also becomes much more difficult because the loss of elastic recoil reduces the passive ability of the lungs to deflate during exhalation. High lung compliance is commonly seen in those with obstructive diseases, such of emphysema, in which destruction of the elastic tissue of the lungs from cigarette smoke exposure causes a loss of elastic recoil of the lung. Those with emphysema have considerable difficulty with exhaling breaths and tend to take fast shallow breaths and tend to sit in a hunched-over position in order to make exhalation easier.

This page titled 21.6B: Factors Affecting Pulmonary Ventilation: Compliance of the Lungs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.6C: Factors Affecting Pulmonary Ventilation: Airway Resistance

Airway resistance refers to resistance in the respiratory tract to airflow.

Learning Objectives

- Describe airway resistance and how it affects pulmonary ventilation

Key Points

- Airway resistance is a concept in respiratory physiology that describes the resistance of the respiratory tract to airflow during inspiration and expiration.
- Airway resistance can be indirectly measured with body plethysmography.
- A two-fold change in the radius/diameter of an airway causes a 16-fold change in air resistance in the opposite direction (an inverse relationship).
- Diseases affecting the respiratory tract can increase airway resistance.
- Laminar flow is orderly and has low resistance while turbulent flow is disorganized and has high resistance.

Key Terms

- **Airway resistance**: Airway Resistance is a concept in respiratory physiology that describes the resistance of the respiratory tract to airflow during inspiration and expiration.
- **plethysmography**: The diagnostic use of a plethysmograph to measure changes in volume within an organ or whole body.
- **Turbulent flow**: Air with disorganized layers that has higher resistance. It is often located in areas where the airways branch or diverge.

EXAMPLES

Airway resistance can change over time, especially during an asthma attack when the airways constricts causing an increase in airway resistance.

Airway Resistance

Airway resistance is the resistance to flow of air caused by friction with the airways, which includes the conducting zone for air, such as the trachea, bronchi and bronchioles. The main determinants of airway resistance are the size of the airway and the properties of the flow of air itself.

Size of the Airway

Resistance in an airway is inversely proportional to the radius of the airway. However the ratio for this relationship is not 1:1. Below is the equation for calculating airway resistance (R).

$R = 8(\text{Length} \times \text{Gas Viscosity})(\pi r^4)$

The most important part of this formula is radius of the airway (*r*). A common example is that if one were to double the diameter of an airway (thus doubling the radius as well) the resistance of the airway would drop by a factor of 16. This mathematical property between radius and resistance is consistent for all tubes, and is often applied to the blood vessels in the cardiovascular system.


The radius of the airways of the conducting zone become smaller as air goes deeper into the lungs. Therefore the resistance to air in the bronchi is greater than the resistance to air in the trachea. The number of airways also plays a large role in the resistance to air, with more airways reducing resistance because there are more paths for the air to flow into. Therfore, despite the fact that the terminal bronchioles are the smallest airway in terms of radius, their high number compared to the larger airways means that the bronchi actually have greater resistance because there are less of them compared to the terminal bronchioles. Another important fact is that airway resistance is inversely related to lung volumes because the airways expand a bit as they inflate, so the airways in a fully inflated lung will have lower resistance than a lung after exhalation.

Airway resistance can be indirectly measured with body plethysmography, which is an instrument used for measuring changes in volume within a structure, such as the airways. The resistance of the airways is an important indicator of lung health and function and can be used to diagnose lung diseases.

The size of the airways, and thus the resistance can change based on the health and conditions of the lungs. Most lung diseases increase airway resistance in many different ways. For example, in asthma attacks the bronchioles spasm and constrict, which increases resistance. Emphysema also increases airway resistance because the lung tissue becomes too pliable and it the airways become more difficult to hold open by the flow of air.

Flow of Air

The air that flows through the lungs varies considerably in the properties of the flow of air. The air flow can either be turbulant, transitional or laminar based on the airway. Laminar flow involves an orderly and concentric distribution of layers of air particles and tends to occur in smaller airways, and has lower resistance. Turbulent flow is disorganized distribution of the layers of air and tends to occur in larger airways and places where the airways branch, and has a higher resistance. Transitional flow occurs in places that branch within smaller airways, in which the air flow becomes in between laminar and turbulent flow and has moderate resistance. The relationship between resistance and type of airflow is difficult to measure and apply, but some mathematical models (such as the Reynold's number) can provide a rough estimate.



Laminar and Turbulent Flow: Laminar flow (a) has orderly layers and low resistance. Turbulent flow (b) has disorganized layers and high resistance.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Alveolar. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Alveolar. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary surfactant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- atelectasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/atelectasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- great alveolar cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/great%20alveolar%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- surfactant. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary surfactant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The respiratory system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...%23Inspiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary surfactant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmona...t%23Compliance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...ung-compliance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary surfactant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ipf_NIH 1.jpg. **Provided by**: Wikipedia . **Located at**: en.Wikipedia.org/wiki/Idiopa...le:Ipf_NIH.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Airway resistance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Airway_resistance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Plethysmograph. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Plethysmograph. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...iratory-system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...way-resistance. **License**: *CC BY-SA: Attribution-ShareAlike*
- plethysmography. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/plethysmography. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pulmonary surfactant. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pulmonary_surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Ipf_NIH 1.jpg. **Provided by**: Wikipedia . **Located at**: en.Wikipedia.org/wiki/Idiopa...le:Ipf_NIH.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- 753px-Laminar_and_turbulent_flows.svg.png. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hydrod...lent_flows.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 21.6C: Factors Affecting Pulmonary Ventilation: Airway Resistance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.7: NONRESPIRATORY LUNG FUNCTIONS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.7A: Lung Capacity and Volume

21.7B: Nonrespiratory Air Movements

This page titled 21.7: Nonrespiratory Lung Functions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.7A: Lung Capacity and Volume

Lung volumes and capacities refer to phases of the respiratory cycle; lung volumes are directly measured while capacities are inferred.

LEARNING OBJECTIVE

Differentiate among tidal volume, inspiratory reserve volume, expiratory reserve volume, and vital capacity of lungs

Key Takeaways

Key Points

- Lung capacity is a measure of lung volume inferred from the exhaled during the various cycles of breathing.
- There is residual air leftover in the lungs during normal breathing.
- Vital capacity is used to diagnose restrictive diseases, while the FEV1/FVC ratio is used to diagnose obstructive diseases.
- FEV1/FVC ratio declines as someone ages, but declines faster in those who smoke due to damage caused by smoking.

Key Terms

- **FEV1/FVC ratio**: The ratio between forced expiratory volume and forced vital capacity, which is used to measure the level of obstruction in the lungs.
- **vital capacity**: The maximum volume of air that can be discharged from the lungs following maximum inspiration.

EXAMPLES

- Pulmonary function tests (PFTs) may be used to help diagnose different pulmonary diseases. The two most often used measurements are FVC (forced vital capacity) and FEV1 (forced expiratory volume in one second).
- An FEV1/FVC ratio of >80% indicates a restrictive lung disease like pulmonary fibrosis or infant respiratory distress syndrome.
- An FEV1/FVC ratio of <70% indicates an obstructive lung disease like asthma or COPD.

Lung Capacity

Lung capacity generally refers to the total amount of air inside the lungs at certain phases of the respiratory cycle. It is usually measured as the amount of air that is exhaled after inhalation; this is measured with a device called a spirometer.

There are many different types and terms for the different components of lung capacity that all have different characteristics. In general, measuring lung capacity is important because it serves as the best indicator of lung health by quantifying the functional ability of the lungs to cycle air.

Vital Capacity

Vital capacity (VC) is the maximum amount of air that a person can exhale after inhaling as much air as possible. It is also the sum of tidal volume and the inspiratory and expiratory reserve volumes, which capture the differences between normal breathing and maximal breathing.

The inspiratory reserve volume is the extra space for air after a normal inspiration, and the expiratory reserve volume is the extra air that can be exhalaed after a normal expiration. VC tends to be decreased in those with restrictive lung diseases, such as pulmonary fibrosis, making VC a good diagnostic indicator of restrictive lung diseases.

Other important lung volumes related to lung capacity are residual volume (RV) and total lung capacity (TLC).


- RV: The amount of air left in the lungs after a maximal expiration.
- TLC: The volume of the lungs at maximal inflation, which is the sum of VC and RV.

FEV1/FVC Ratio

The most widely used diagnostic application for lung capacities is the ratio between forced expiratory volume (FEV1) and forced vital capacity (FVC).

- FEV1: The volume of air exhaled in one second of forced expiration.
- FVC: The total volume exhaled air during a forced expiration.

The FEV1/FVC ratio is an important indicator of lung health and is the standard approach for diagnosing COPD (chronic pulmonary obstructive disease), which includes emphysema and bronchitis, which are both caused by smoking. An FEV1/FVC ratio that is greater than .8 indicates a normal lung with generally healthy function, however a ratio below .8 indicates a significant degree of airway obstruction and suggests COPD.



Lung Capacity: Lung capacity at the various stages of the respiratory cycle, which is one inhalation followed by an exhalation.

The obstruction becomes worse the lower the ratio becomes, which increases the likelihood of respiratory failure and death. The FEV1/FVC ratio naturally falls as humans age, however smoking (the cause of COPD) will cause much larger decreases in FEV1/FVC ratio than what is normal.

Smoking causes this damage by initiating an inflammatory response in the lungs. Those who quit smoking will not experience a regain the FEV1/FVC ratio lost from smoking, however their rate of FEV1/FVC ratio decline will slow to normal, and their life expectancy will be less impacted.

Those with asthma, an acute form obstructive lung disease, will show a low FEV1/FVC ratio during an asthma attack, which returns to normal after the attack is over. Therefore, to diagnose asthma, many clinicians expose patients to methacholine or histamine to trigger mild asthma attacks to measure FEV1/FVC ratios.

This page titled 21.7A: Lung Capacity and Volume is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.7B: Nonrespiratory Air Movements

The lungs have a number of metabolic functions in addition to their functions in gas exchange.

LEARNING OBJECTIVE

Discuss the non-respiratory air movements of the respiratory system

Key Takeaways

Key Points

- The lungs have a number of metabolic functions, such as the secretion of ACE (angiotensin converting enzyme), which converts angiotensin I to angiotensin II to stimulate changes in the renal system.
- Higher levels of ACE lead to higher blood pressure. ACE inhibitors are used to treat hypertension by reducing ACE to reduce blood pressure.
- Airway epithelial cells can secrete a variety of molecules—immunoglobulins (IgA), proteases, reactive oxygen species, and antimicrobial peptides—that all help protect the lungs and body from pathogens.
- Non-respiratory air movements are mechanical functions that aren't involved in gas exchange, such as voice production and coughing.

Key Terms

- **ACE**: Angiotensin converting enzyme, which is secreted in the lungs and helps to increase blood pressure in the body through renal system feedback loops.
- **Airway epithelial cells**: Airway epithelial cells can secrete a variety of molecules that aid in the immune system defense of lungs.

EXAMPLE


Non-respiratory air movements do not involve gas exchange. Examples are: sneezing, coughing, burping, laughing, singing, and talking.

While the primary function of the lungs is gas exchange, they have several other functions, which are both metabolic and mechanical. These include the secretion of many enzymes and proteins involved in other body systems and nonrespiratory air movements.

Metabolic Functions

The lungs secrete many enzymes and proteins that serve non-respiratory metabolic functions.

ACE (angiotensin converting enzyme) is an enzyme secreted by the endothelial cells of the capillaries in the lungs. ACE converts angiotensin I into angiotensin II, which are two important hormones in the renin-angiotensin feedback loop of the renal system.



The renin-angiotensin-aldosterone system: The renin-angiotensin-aldosterone system is dependent on ACE from the lungs to regulate blood pressure. ACE activity results in increased blood pressure.

This system works to regulate blood pressure and blood volume by changing the amount of water retained by the kidneys. In general, more ACE leads to more angiotensin II, which leads to more aldosterone, which leads to more retained water through sodium reabsorption in the kidney, which leads to increased blood volume and blood pressure.

ACE inhibitors are a common treatment for those with hypertension, as it will reduce the amount of ACE, which will cause the kidney to excrete more water, which lowers blood volume and blood pressure.

The epithelial cells and macrophages of the lungs secrete many molecules that have immune system functions. In general these molecules have anti-microbial functions.

- Immunoglobin A (IgA): An antibody that can attack pathogens and mark them for phagocytosis from macrophages and neutrophils.
- Protease: Secreted from lung macrophages and neutrophils during inflammatory response to damage pathogens. A fibrinolytic that can break up thrombosis (blood clots) in the lungs.
- Reactive oxygen species (ROS): Free radicals, which are any substance with an unpaired electron in the valence shell, can cause oxidative stress (damage) in cells. They are used to kill pathogens after being engulfed (phagocytized) by immune cells.
- Anti-microbial peptides: Various chemokines and proteins that are secreted by the mucus membranes of the airways. They can damage and inhibit pathogens and are considered a barrier component of the immune system.

Mechanical Functions

There are several types of non-respiratory air movements that have important functions that are not primarily related to gas exchange. One example is voice production for speaking and singing, which involves fine control over the direction and flow of the air as it passes into the upper respiratory tract.

Other mechanical functions include sneezing and coughing, which protect the lungs and airways from irritants that could potentially cause damage. Coughing is a result of constriction from nervous stimulation in the trachea and larynx and also serves to dislodge mucus trapped inside the lungs.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Lung volume. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lung_volume. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tidal volume. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tidal_volume. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vital capacity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vital_capacity. **License**: ***CC BY-SA: Attribution-ShareAlike***
- partial pressure. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/partial_pressure. **License**: ***CC BY-SA: Attribution-ShareAlike***

- The respiratory system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...iratory_system. **Project**: Human Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Respiratory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Respiratory_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- syrinx. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/syrinx. **License**: *CC BY-SA: Attribution-ShareAlike*
- surfactant. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/surfactant. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renin angiotensin aldosterone system. **Authored by**: A. Rad. **Located at**: commons.wikimedia.org/wiki/F...one_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 21.7B: Nonrespiratory Air Movements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.8: GAS LAWS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

21.8A: Dalton's Law of Partial Pressure

21.8B: Henry's Law

This page titled 21.8: Gas Laws is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.8A: Dalton's Law of Partial Pressure

Dalton's law of partial pressures states that the pressure of a mixture of gases is the sum of the pressures of the individual components.

Learning Objectives

- Infer from Dalton's law of partial pressure the sum of partial pressures in alveoli

Key Points

- This empirical law was observed by John Dalton in 1801 and is related to the ideal gas laws.
- Atmospheric air is a mixture of nitrogen, water, oxygen, carbon dioxide, and other minor gasses. The relative concentrations of a gasses don't change even as the pressure and volume of the total gasses change.
- Gasses flow from areas of high to low pressure, so the partial pressures of inhaled and alveolar air determine why oxygen goes into the alveoli, and why carbon dioxide leaves the alveoli.
- Dalton's law is only completely accurate for ideal gasses.

Key Terms

- **Dalton's law**: The total pressure of a mixture of gases is the sum of the partial pressures of each gas in the mixture; it is only true for ideal gases.

Dalton's law states that the total pressure exerted by the mixture of inert (non-reactive) gases is equal to the sum of the partial pressures of individual gases in a volume of air. This empirical law was observed by John Dalton in 1801 and is related to the ideal gas laws.

Dalton's Law in Respiration

The air in the atmosphere is a mixture of many different gases, that vary in concentration. Dalton's law states that at any given time, the percentage of each of these gasses in the air we breathe makes its contribution to total atmospheric pressure, and this contribution will depend on how much of each gas is in the air we breathe.

Dalton's law also implies that the relative concentration of gasses (their partial pressures) does not change as the pressure and volume of the gas mixture changes, so that air inhaled into the lungs will have the same relative concentration of gasses as atmospheric air. In the lungs, the relative concentration of gasses determines the rate at which each gas will diffuse across the alveolar membranes.

Mathematically, the pressure of a mixture of gases can be defined as the sum of the partial pressures of each of the gasses in air.

$$P_{total}=P1+P2+P3+\cdots+Pn=n\sum i=1PiPtotal=P1+P2+P3+\cdots+Pn=\sum i=1nPi$$

DALTON'S LAW

In regards to atmospheric air, Dalton's law becomes:

$$Atm=PN2+PO2+PCO2+PH2O+P(other\ gasses)Atm=PN2+PO2+PCO2+PH2O+P(other\ gasses)$$

For the purposes of gas exchange, O_2 and CO_2 are mainly considered due to their metabolic importance in gas exchange. Because gasses flow from areas of high pressure to areas of low pressure, atmospheric air has higher partial pressure of oxygen

than alveolar air (P_{O_2}= 159mm Hg compared to PA_{O_2}= 100mm Hg).

Similarly, atmospheric air has a much lower partial pressure for carbon dioxide compared to alveolar air (P_{CO_2}= .3mm Hg compared to PA_{CO_2}= 40mm Hg). These pressure differences explain why oxygen flows into the alveoli and why carbon dioxide flows out of the alveoli through passive diffusion (just as a similar process explains alveolar and arterial gas exchange).

While inhaled air is similar to atmospheric air due to Dalton's law, exhaled air will have relative concentrations that are in between atmospheric and alveolar air due to the passive diffusion of gasses during gas exchange.

Dalton's law is only truly applicable in every situation to ideal gasses. Therefore most gasses will not follow it exactly, especially in conditions of extremely high pressure, or in situations where intermolecular forces act to keep the gasses together.

This page titled 21.8A: Dalton's Law of Partial Pressure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.8B: Henry's Law

Henry's law states that the amount of a gas that dissolves in a liquid is directly proportional to the partial pressure of that gas.

Learning Objectives

- Explain the way in which Henry's law relates to gas exchange in the respiratory system

Key Points

- At a constant temperature, the amount of a given gas that dissolves in a given type and volume of liquid is directly proportional to the partial pressure of that gas in equilibrium with that liquid.
- Gasses with a higher solubility will have more dissolved molecules than gasses with a lower solubility if they have the same partial pressure.
- Henry's law explains how gasses dissolve across the alveoli – capillary barrier.
- Henry's law predicts how gasses behave during gas exchange based on the partial pressure gradients and solubility of oxygen and carbon dioxide.


Key Terms

- **Henry's law**: At a constant temperature, the amount of a given gas that dissolves in a given type and volume of liquid is directly proportional to the partial pressure of that gas in equilibrium with that liquid.
- **partial pressure gradient**: The difference between the partial pressures (and thus concentration) of gasses between gaseous and dissolved forms.

EXAMPLES

An everyday example of Henry's law is given by carbonated soft drinks. Before the bottle or can is opened, the gas above the drink is almost pure carbon dioxide at a pressure slightly higher than atmospheric pressure. The drink itself contains dissolved carbon dioxide. When the bottle or can is opened, some of this gas escapes, giving the characteristic hiss (or pop in the case of a sparkling wine bottle). Because the pressure above the liquid is now lower, some of the dissolved carbon dioxide comes out of solution as bubbles. If a glass of the drink is left in the open, the concentration of carbon dioxide in solution will come into equilibrium with the carbon dioxide in the air, and the drink will go flat.

Henry's law states that at a constant temperature, the amount of a gas that dissolves in a liquid is directly proportional to the partial pressure of that gas in equilibrium with that liquid. It was formulated by William Henry in 1803.



Henry's law: Henry's law states that when a gas is in contact with the surface of a liquid, the amount of the gas which will go into solution is proportional to the partial pressure of that gas.

The practical description for the law is that the solubility (i.e., equilibrium) of a gas in a liquid is directly proportional to the partial pressure of that gas. In addition, the partial pressure is able to predict the tendency to dissolve simply because the gasses with higher partial pressures have more molecules and will bounce into the solution they can dissolve into more often than gasses with lower partial pressures.

Henry's law also applies to the solubility of other substances that aren't gaseous, such as the equilibrium of organic pollutants in water being based on the relative concentration of that pollutant in the media its suspended in.

Henry's law can be put into mathematical terms (at constant temperature):

$p=kHcp=kHc$

Where p is the partial pressure of the solute in the gas above the solution, c is the concentration of the solute, the solubility of the substance is k, and the Henry's law constant (H), which depends on the solute, the solvent, and the temperature.

The solubility captures the tendency of a substance to go towards equilibrium in a solution, which explains why gasses that have the same partial pressure may have different tendencies to dissolve.

Henry's Law in Respiration

The main application of Henry's law in respiratory physiology is to predict how gasses will dissolve in the alveoli and bloodstream during gas exchange. The amount of oxygen that dissolves into the bloodstream is directly proportional to the partial pressure of oxygen in alveolar air.

The partial pressure of oxygen is greater in alveolar air than in deoxygenated blood, so oxygen has a high tendency to dissolve into deoxygenated blood. Conversely the opposite is true for carbon dioxide, which has a greater partial pressure in deoxygenated blood than in the alveolar air, so it will diffuse out of the solution and back into gaseous form.

Recall that the difference in partial pressures between the bloodstream and alveoli (the partial pressure gradient) are much smaller for carbon dioxide compared to oxygen. Carbon dioxide has much higher solubility in the plasma of blood than oxygen (roughly 22 times greater), so more carbon dioxide molecules are able to diffuse across the small pressure gradient of the capillary and alveoli.

Oxygen has a larger partial pressure gradient to diffuse into the bloodstream, so it's lower solubility in blood doesn't hinder it

during gas exchange. Therefore, based on the properties of Henry's law, both the partial pressure and solubility of the oxygen and carbon dioxide determine how they will behave during gas exchange.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Dalton's law. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Dalton's_law**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gas Laws. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gas_Laws**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Dalton's law. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Dalton's_law**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gas Laws. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gas_Laws**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Henry's law. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Henry's_law**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gas Laws. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gas_Laws**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Henry's law. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Henry's_law**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gas Laws. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gas_Laws**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Amazon Web Services. **Located at**: **s3.amazonaws.com/figures.boun.../henryslaw.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 21.8B: Henry's Law is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.9: GAS EXCHANGE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

21.1A: External Respiration

21.9B: Internal Respiration

21.9C: Oxygen Transport

21.9D: Carbon Dioxide Transport

This page titled 21.9: Gas Exchange is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.1A: External Respiration

Respiration is the transport of oxygen to the cells within tissues and the transport of carbon dioxide in the opposite direction.

Learning Objectives

- Describe the four stages of external respiraton

Key Points

- External respiration describes the exchange of gasses between the external environment and the bloodstream.
- The components of external respiration include alveolar surface area, ventilation and perfusion matching, and partial pressure gradients.
- Partial pressure gradients allow gasses to flow from areas of high pressure to areas of lower pressure.
- Ventilation and perfusion in the alveoli must be balanced to maintain efficient gas exchange.

Key Terms

- **passive diffusion**: The net movement of material from an area of high concentration to an area of lower concentration without any energy input.
- **perfusion**: The blood supply to the alveolar capillaries that must be balanced with ventilation.
- **Equilibrium**: The rate of change of the gasses in the alveoli and capillary become equal to eachother at the end of gas exchange.

External Respiration

External respiration is the formal term for gas exchange. It describes both the bulk flow of air into and out of the lungs and the transfer of oxygen and carbon dioxide into the bloodstream through diffusion. While the bulk flow of air from the external environment happens due to pressure changes in the lungs, the mechanisms of alveolar gas exchange are more complicated. The primary three components of external respiration are the surface area of the alveolar membrane, the partial pressure gradients of the gasses, and the matching of perfusion and ventilation.


Surface Area

The alveoli have a very high surface area to volume ratio that allows for efficient gas exchange. The alveoli are covered with a high density of capillaries that provide many sites for gas exchange.

The walls of the alveolar membrane are thin and covered with a fluid, extra-cellular matrix that provides a surface for gas molecules in the air of the lungs to diffuse into, from which they can then diffuse into the capillaries.

Partial Pressure Gradients

Partial pressure gradients (differences in partial pressure) allow the loading of oxygen into the bloodstream and the unloading of carbon dioxide out of the bloodstream. These two processes occur at the same time.



Gas exchange in the alveolus: External respiration is a result of partial pressure gradients, alveolar surface area, and ventilation and perfusion matching.

Oxygen has a partial pressure gradient of about 60 mmHg (100 mmHg in alveolar air and 40 mmHg in deoxygenated blood) and diffuses rapidly from the alveolar air into the capillary.

Equilibrium between the alveolar air and capillaries is reached quickly, within the first third of the length of the capillary within a third of a second. The partial pressure of oxygen in the oxygenated blood of the capillary after oxygen loading is about 100 mmHg.

The process is similar in carbon dioxide. The partial pressure gradient for carbon dioxide is much smaller compared to oxygen, being only 5 mmHg (45 mmHg in deoxygenated blood and 40 mmHg in alveolar air).

Based on Henry's law, the greater solubility of carbon dioxide in blood compared to oxygen means that diffusion will still occur very rapidly despite the lower partial pressure gradient. Equilibrium between the alveolar air and the capillaries for carbon dioxide is reached within the first half of the length of the capillaries within half a second. The partial pressure of carbon dioxide in the blood leaving the capillaries is 40 mmHg.

Ventilation and Perfusion Matching

The exchange of gas and blood supply to the lungs must be balanced in order to facilitate efficient external respiration. While a severe ventilation–perfusion mismatch indicates severe lung disease, minor imbalances can be corrected by maintaining air flow that is proportional to capillary blood flow, which maintains the balance of ventilation and perfusion.

Perfusion in the capillaries adjusts to changes in PAO_2. Constriction in the airways (such as from the bronchospasms in an asthma attack) lead to decreased PAO_2 because the flow of air into the lungs is slowed.

In response, the arteries being supplied by the constricted airway undergo vasocontriction, reducing the flow of blood into those alveoli so that the perfusion doesn't become much greater relative to the decreased ventilation (a type of ventilation–perfusion mismatch called a shunt).

Alternatively, breathing in higher concentrations of oxygen from an oxygen tank will cause vasodilation and increased blood perfusion in the capillaries.

Ventilation adjusts from changes in $PACO_2$. When airflow becomes higher relative to perfusion, $PACO_2$ decreases, so the bronchioles will constrict in order to maintain to the balance between airflow (ventilation) and perfusion. When airflow is reduced, $PACO_2$ increases, so the bronchioles will dilate in order to maintain the balance.

This page titled 21.1A: External Respiration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.9B: Internal Respiration

Cellular respiration is the metabolic process by which an organism obtains energy through the reaction of oxygen with glucose.

Learning Objectives

- Describe internal (cellular) respiration

Key Points

- Cellular respiration is the metabolic process by which an organism obtains energy by reacting oxygen with glucose to give water, carbon dioxide, and adenosine triphosphate (energy).
- The 3 steps of cellular respiration are glycolysis, the Krebs cycle, and oxidative phosphorylation.
- Carbon dioxide is a waste product of cellular respiration that comes from the carbon in glucose and the oxygen used in cellular respiration.
- Internal respiration involves gas exchange between the bloodstream and tissues, and cellular respiration.
- Gas exchange in tissues is a result of tissue surface area, gas partial pressure gradients, and the blood perfusion of those tissues.

Key Terms

- **Cellular respiration**: The cellular process of producing adenosine triphosphate (ATP), water, and carbon dioxide from glucose and oxygen.
- **Oxidative Phosphorylation**: The final step of cellular phosphorylation in which oxygen is used to a create an electron transport chain that produces a large amount of ATP from nicotinamide adenine dinucleotide (NADH).

Internal respiration refers to two distinct processes. The first is the exchange of gasses between the bloodstream and the tissues. The second is the process of cellular respiration, from which cells utilize oxygen to perform basic metabolic functions.

Gas Exchange with Tissues

Gas exchange occurs in the alveoli so that oxygen is loaded into the bloodstream and carbon dioxide is unloaded from the bloodstream. Afterwards, oxygen is brought to the left side of the heart via the pulmonary vein, which pumps it into systemic circulation.


Red blood cells carry the oxygen into the capillaries of the tissues of the body. Oxygen diffuses into the cells of the tissues, while carbon dioxide diffuses out of the cells of the tissues and into the bloodstream.

The factors that influence tissue gas exchange are similar to the factors of alveolar gas exchange, and include partial pressure gradients between the blood and the tissues, the blood perfusion of those tissues, and the surface areas of those tissues. Each of those factors generally increase gas exchange as those factors are increased (i.e., more oxygen diffusion in tissues with more blood perfusion).

Regarding the partial pressure gradients in systemic capillaries, they have a PaO_2 of 100mmHg and a $PaCO_2$ of 40mmHG within the capillary and a PaO_2 of 40 mmHg and $PaCO_2$ of 45 mmHg inside issue cells, which allows gas exchange to occur.

Cellular Respiration

Cellular respiration is the metabolic process by which an organism obtains energy through the reaction of oxygen with glucose to produce water, carbon dioxide and ATP, which is the functional source of energy for the cell. The oxygen supply for cellular respiration comes from the external respiration of the respiratory system.



Overview of cellular respiration: A diagram of cellular respiration including glycolysis, the Krebs cycle (also called the citric acid cycle), and the electron transport chain.

Cellular respiration includes three major steps, and occurs mainly in the cytoplasm of the cell and within the mitochondria of the cell. The net formula for cellular respiration is:

1. Glycolysis: The breakdown of glucose into pyruvate, ATP, H_2O, and heat.
2. Krebs Cycle: Produces NADH from pyruvate.
3. Oxidative Phosphorylation: Produces ATP from NADH, oxygen, and H+. The oxygen plays the role of electron receptor in an electron transport chain to produce ATP.

The net formula for cellular respiration is:

Glucose+6 Oxygen→6 Carbon Dioxide+6 Water+38 ATPGlucose+6 Oxygen→6 Carbon Dioxide+6 Water+38 ATP

The carbon dioxide waste is the result of the carbon from glucose ($C_6H_{12}O_6$) being broken down to produce the pyruvate and NADH intermediates needed to produce ATP at the end of respiration. The energy stored in ATP can then be used to drive processes that require energy, including biosynthesis, locomotion, or transportation of molecules across cell membranes.

Cellular respiration can occur anaerobically without oxygen, such as through lactic acid fermentation. Human cells may use lactic acid fermentation in muscle tissue during strenuous exercise when there isn't enough oxygen to power the tissues. This process is very inefficient compared to aerobic respiration, as without oxidative phosphorylation, the cell cannot produce nearly as much ATP (2 ATP compared to 38 during cellular respiration).

This page titled 21.9B: Internal Respiration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.9C: Oxygen Transport

Hemoglobin is the primary transporter of oxygen with an oxygen binding capacity between 1.36 and 1.37 ml O_2 per gram Hgb.

Learning Objectives

- Describe the process of oxygen transport

Key Points

- About 98.5% of the oxygen in a sample of arterial blood in a healthy human breathing air at sea-level pressure is bound to the hemoglobin in blood.
- Each molecule of hemoglobin is made up of four heme groups.
- The oxygen dissociation curve shows the amount of oxygen saturated in hemoglobin for a given partial pressure of oxygen.
- The curve shifts right when oxygen's affinity for binding to hemoglobin is decreased, and shifts left when its affinity for binding is increased.
- A rightward shift occurs during exercise, decreasing blood pH and increasing temperature, or during anemia (with a downward shift as well).
- A leftward shift occurs during increased blood pH, decreased temperature, or carbon monoxide exposure.


Key Terms

- **hemoglobin**: A protein found in red blood cells that provides a binding site for oxygen to transport it to the tissues of the body.
- **dissociation curve**: The oxygen–hemoglobin dissociation curve plots the proportion of hemoglobin saturated with oxygen on the vertical axis against the partial pressure of oxygen on the horizontal axis.

Hemoglobin

About 98.5% of the oxygen in a sample of arterial blood in a healthy human breathing air at sea-level pressure is bound to the hemoglobin in blood (Hgb). Hemoglobin is a protein found in red blood cells (also called erythrocytes).

There are roughly 270 million hemoglobin molecules in a single red blood cell, and each contains 4 heme groups. The function of Hgb is to provide a binding site for oxygen to carry oxygen throughout the bloodstream to the systemic tissues for cellular respiration.



Hemoglobin: Hemoglobin is the iron-containing, oxygen-transport metalloprotein in the red blood cells of all vertebrates.

About 1.5% of oxygen is physically dissolved in the other blood liquids and not connected to Hgb. It has an oxygen binding capacity between 1.36 and 1.37 ml O_2 per gram Hgb.


Oxyhemoglobin Dissociation Curve

The percentage of oxygen that is saturated in the hemoglobin of blood is generally represented by a curve that shows the relationship between PaO_2 and O_2 saturation. Saturation of O_2 in hemoglobin is an indicator for how much O_2 is able to reach the tissues of the body.

Higher PaO_2 means higher saturation of oxygen in blood. Under normal conditions the PaO_2 in systemic blood is equal to 50%, about 26.6 mmHg,; this is called the P_{50}.

The curve starts to plateau at PaO_2 higher than 60 mmHG, meaning that increases in PaO_2 after that point won't significantly increase saturation. This also means that the approximate carrying capacity for oxygen in hemoglobin has been reached and excess oxygen won't go into hemoglobin.

The carrying capacity can be increased if more hemoglobin is added to the system, such as through greater red blood cell generation in high altitude, or from blood transfusions. The lower areas of the curve show saturation when oxygen is unloaded into the tissues.



The oxyhaemoglobin dissociation curve: The oxygen–hemoglobin dissociation curve plots the percent hemoglobin saturation (y-axis) against the partial pressure of oxygen in the blood (PO_2). The blue curve is standard curve, while the red and green curves are right and leftward shifts respectively.

The oxyhemoglobin dissociation curve can shift in response to a variety of factors. A change in the P_{50} of the curve is a sign that the dissociation curve as a whole has shifted. Shifts indicate a change in affinity for oxygen's binding to hemoglobin, which changes the ability of oxygen to bind to hemoglobin and stay bound to it (i.e., not be released from it).

Rightward shifts indicate a decreased affinity for the binding of hemoglobin, so that less oxygen binds to hemoglobin, and more oxygen is unloaded from it into the tissues. The curve shifts right during decreased blood pH (called the Bohr effect), increased temperature, and during exercise among other things.

Anemia (a disorder marked by a decreased red blood cell count and less hemoglobin) also causes a rightward shift, but also changes the shape of the curve so that it moves downward as well as a result of the reduced levels of hemoglobin.

Leftward shifts indicate an increased affinity for the binding of hemoglobin, so that more oxygen binds to hemoglobin, but less oxygen is unloaded from it into the tissues. Causes of leftward shifts include increased blood pH, decreased temperature, and carbon monoxide exposure. Carbon monoxide binds to hemoglobin in place of oxygen, so that less oxygen reaches the tissues; this can be fatal if severe enough.

This page titled 21.9C: Oxygen Transport is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

21.9D: Carbon Dioxide Transport


Learning Objectives

- Describe the process of carbon dioxide transport

Carbon Dioxide Transport

Carbon dioxide is the product of cellular respiration, and is transported from the cells of tissues in the body to the alveoli of the lungs through the bloodstream. Carbon dioxide is carried in the blood through three different ways.

About 5% of carbon dioxide is transported in the plasma of the blood as dissolved CO_2 molecules that aren't bound to anything else. Carbon dioxide has a much higher solubility than oxygen, which explains why a relatively greater amount of carbon dioxide is dissolved in the plasma compared to oxygen. While oxygen binds to the iron content in the heme of hemoglobin, carbon dioxide can bind to the amino acid chains on hemoglobin. When carbon dioxide clings to hemoglobin it forms carbanimohemoglobin. About 10% of carbon dioxide in the human body is transported this way. Carbanimohemoglobin gives red blood cells a bluish color, which is one of the reasons why the veins that carry deoxygenated blood appear to be blue.



Structure of human hemoglobin: Hemoglobin is a tetramer of alpha (red) and beta (blue) subunits with iron containing heme groups (green).

A property of hemoglobin called the **Haldane effect** states that deoxygenated blood has an increased capacity to carry carbon dioxide, while oxygenated blood has a decreased capacity to carry carbon dioxide.

This property means that hemoglobin will primarily carry oxygen in systemic circulation until it unloads that oxygen and is able to carry a relatively higher amount of carbon dioxide. This is due to deoxygenated blood's increased capacity to carry carbon dioxide, and from the carbon dioxide loaded from the tissues during tissue gas exchange.

Bicarbonate Ions

The majority (85%) of carbon dioxide travels in the blood stream as bicarbonate ions. The reaction that describes the formation of bicarbonate ions in the blood is:

$$CO_2 + H_2O \longrightarrow H_2CO_3 \longrightarrow H^+ + HCO_3^- \quad (21.9D.1)$$

This means that carbon dioxide reacts with water to form carbonic acid, which dissociates in solution to form hydrogen ions and bicarbonate ions. The main implication of this process is that the pH of blood becomes a way of determining the amount of carbon dioxide in blood. This is because if carbon dioxide increases in the body, it will manifest as increased concentrations of bicarbonate and increased concentrations of hydrogen ions that reduce blood pH and make the blood more acidic. Conversely, if carbon dioxide levels are reduced, there will be less bicarbonate and less hydrogen ions dissolved in the blood, so pH will increase and blood will become more basic. Bicarbonate ions act as a buffer for the pH of blood so that blood pH will be neutral as long as bicarbonate and hydrogen ions are balanced.

This connection explains how ventilation rate and blood chemistry are related, as hyperventilation will cause alkalosis, and hypoventilation will cause acidosis, due to the changes in carbon dioxide levels that they cause. Bicarbonate is also carried in the fluids of tissues besides the blood vessels, especially in the duodenum and intestine, so problems in those organs can cause a respiratory system response.

Transport to the Alveoli

After carbon dioxide travels through the bloodstream to the capillaries covering the alveoli of the lungs through any of the 3 methods listed above, it must return to dissolved carbon dioxide form in order to diffuse across the capillary into the alveolus. Dissolved carbon dioxide is already able to diffuse into the alveolus, while hemoglobin-bound carbon dioxide is unloaded into the plasma.

For carbon dioxide stored in bicarbonate, it undergoes a reaction reversal. Bicarbonate ions dissolved in the plasma enter the red blood cells by diffusing across a chloride ion gradient (replacing chloride inside the cell), and combining with hydrogen to form carbonic acid.

Next, the action of carbonic anhydrase breaks carbonic acid down into carbon dioxide in water, which leaves the cell by diffusion. The dissolved carbon dioxide is then able to diffuse into the alveolus.

Key Points

- Carbon dioxide is transported through the bloodstream either dissolved in blood, bound the hemoglobin, or converted to bicarbonate ions.
- The Haldane effect is the decreased binding of carbon dioxide in hemoglobin due to increased oxygen levels and increased binding of carbon dioxide to hemoglobin from decreased oxygen levels.
- $CO_2 + H_2O \longrightarrow H_2CO_3 \longrightarrow H^+ + HCO_3^-$ is the bicarbonate formation reaction in the plasma.
- Increased carbon dioxide means increased blood acidity and conversely while decreased carbon dioxide means decreased blood acidity.
- The bicarbonate reaction is reversible, which allows bicarbonate to turn back into dissolved carbon dioxide to diffuse from the alveolus.

Key Terms

- **red blood cell**: A type of cell in the blood of vertebrates that contains hemoglobin and transports oxygen from the lungs to the tissues; an erythrocyte.
- **Bicarbonate**: A negatively charged ion that accumulates in plasma when carbon dioxide dissolves in water and reacts with it. It also acts as a buffer for blood pH.
- **carbonic anhydrase**: An enzyme that catalyzes the bicarbonate reaction in either direction.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- passive diffusion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/passive%20diffusion. **License**: *CC BY-SA: Attribution-ShareAlike*
- External respiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/External_respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...costal-muscles. **License**: *CC BY-SA: Attribution-ShareAlike*
- gas transport. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/gas%20transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gas_exchange_in_the_aveolus_simple_en.svg 1.png. **Provided by**: Wikimedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gas_exchange_in_the_aveolus.svg. **License**: *CC BY: Attribution*
- Cellular respiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cellular_respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gas exchange. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gas_exchange. **License**: *CC BY-SA: Attribution-ShareAlike*
- External respiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/External_respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- respiration. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- metabolite. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/metabolite. **License**: *CC BY-SA: Attribution-ShareAlike*
- transport. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gas_exchange_in_the_aveolus_simple_en.svg 1.png. **Provided by**: Wikimedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gas_exchange_in_the_aveolus.svg. **License**: *CC BY: Attribution*
- CellRespiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:CellRespiration.svg. **License**: *Public Domain: No Known Copyright*
- Oxygen transport. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Oxygen_transport%23Oxygen_transport. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/dissociation-curve. **License**: *CC BY-SA: Attribution-ShareAlike*
- acidity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/acidity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/binding-capacity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hemoglobin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Oxygenu2013haemoglobin dissociation curve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Oxygen%E2%80%93haemoglobin_dissociation_curve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gas_exchange_in_the_aveolus_simple_en.svg 1.png. **Provided by**: Wikimedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Gas_exchange_in_the_aveolus.svg. **License**: *CC BY: Attribution*
- CellRespiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:CellRespiration.svg. **License**: *Public Domain: No Known Copyright*
- 1904%20Hemoglobin. **Provided by**: Wikimedia%20Commons. **Located at**: commons.wikimedia.org/wiki/File:1904_Hemoglobin.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Oxyhaemoglobin dissociation curve. **Provided by**: Wikimedia Commons. **Located at**: en.Wikipedia.org/wiki/File:Oxyhaemoglobin_dissociation_curve.png. **License**: *Public Domain: No Known Copyright*
- Carbaminohemoglobin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Carbaminohemoglobin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hemoglobin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Carbon dioxide. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Carbon_dioxide%23Transport_in_the_blood. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/carbamino. **License**: *CC BY-SA: Attribution-ShareAlike*
- enzyme. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/enzyme. **License**: *CC BY-SA: Attribution-ShareAlike*
- red blood cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/red_blood_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gas_exchange_in_the_aveolus_simple_en.svg 1.png. **Provided by**: Wikimedia Commons. **Located at**:

commons.wikimedia.org/wiki/File:Gas_exchange_in_the_aveolus.svg. **License**: ***CC BY: Attribution***
- CellRespiration. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:CellRespiration.svg**. **License**: ***Public Domain: No Known Copyright***
- 1904%20Hemoglobin. **Provided by**: Wikimedia%20Commons. **Located at**: **commons.wikimedia.org/wiki/File:1904_Hemoglobin.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Oxyhaemoglobin dissociation curve. **Provided by**: Wikimedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Oxyhaemoglobin_dissociation_curve.png**. **License**: ***Public Domain: No Known Copyright***
- Hemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemoglobin**. **License**: ***Public Domain: No Known Copyright***

This page titled 21.9D: Carbon Dioxide Transport is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22: DIGESTIVE SYSTEM

22.1: Overview of the Digestive System

22.1A: Anatomy of the Digestive System

22.1B: Processes and Functions of the Digestive System

22.1C: Organs of the Digestive System

22.2: Nervous System of the Digestive System

22.2A: Enteric Nervous System

22.2B: Gastrointestinal Reflex Pathways

22.3: The Peritoneum

22.3: The Peritoneum

22.4: The Alimentary Canal

22.4A: Mouth

22.4B: Pharynx

22.4C: Esophagus

22.5: Layers of the Alimentary Canal

22.5A: Mucosa

22.5B: Submucosa

22.5C: Muscularis

22.5D: Serosa

22.7: The Liver

22.7A: The Liver

22.7B: Anatomy of the Liver

22.7C: Histology of the Liver

22.7D: Blood Supply to the Liver

22.7E: Liver Function

22.7F: Bile

22.8: The Gallbladder

22.8A: The Gallbladder

22.8B: Anatomy of the Gallbladder

22.9: The Pancreas

22.9A: Pancreas

22.9B: Anatomy of the Pancreas

22.9C: Histology of the Pancreas

22.9D: Pancreatic Juice

22.10: The Small Intestine

22.10A: Anatomy of the Small Intestine

22.10B: Histology of the Small Intestine
22.10C: Digestive Processes of the Small Intestine

22.11: The Large Intestine
22.11A: Anatomy of the Large Intestine
22.11B: Histology of the Large Intestine
22.11C: Bacterial Flora
22.11D: Digestive Processes of the Large Intestine
22.11E: Absorption and Feces Formation in the Large Intestine
22.11F: Defecation Reflex

22.12: Chemical Digestion
22.12A: Mechanisms of Chemical Digestion
22.12B: Chemical Digestion of Carbohydrates, Proteins, Lipids, and Nucleic Acids

22.13: Absorption
22.13A: Absorption in the Small Intestine
22.13B: Absorption of Monosaccharides, Amino Acids, Dipeptides, Tripeptides, Lipids, Electrolytes, Vitamins, and Water

22.14: Phases of Digestion
22.14A: Cephalic Phase
22.14B: Gastric Phase

22.6: The Stomach
22.6A: Gross Anatomy of the Stomach
22.6B: Microscopic Anatomy of the Stomach
22.6C: Digestive Properties of the Stomach

Index

This page titled 22: Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.1: OVERVIEW OF THE DIGESTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.1A: Anatomy of the Digestive System
22.1B: Processes and Functions of the Digestive System
22.1C: Organs of the Digestive System

This page titled 22.1: Overview of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.1A: Anatomy of the Digestive System

The human gastrointestinal tract refers to the stomach and intestine, and sometimes to all the structures from the mouth to the anus.

Learning Objectives


- Outline the anatomical organization of the digestive system

Key Points

- The major organs of the digestive system are the stomach and intestine.
- The upper gastrointestinal tract consists of the esophagus, stomach, and duodenum.
- The lower gastrointestinal tract includes the small intestine and the large intestine.
- Digestive juices are produced by the pancreas and the gallbladder.
- The small intestine includes the duodenum, jejunum, and ileum.
- The large intestine includes the cecum, colon, rectum, and anus.

Key Terms

- **upper gastrointestinal tract**: This tract consists of the esophagus, stomach, and duodenum.
- **lower gastrointestinal tract**: This tract includes most of the small intestine and all of the large intestine.



Upper and lower gastrointestinal tract: The major organs of the human gastrointestinal system.

The human gastrointestinal tract refers to the stomach and intestine, and sometimes to all the structures from the mouth to the anus.

Upper Gastrointestinal Tract

The upper gastrointestinal tract consists of the esophagus, stomach, and duodenum. The exact demarcation between upper and lower can vary. Upon gross dissection, the duodenum may appear to be a unified organ, but it is often divided into two parts

based upon function, arterial supply, or embryology.


The upper gastrointestinal tract includes the:

- Esophagus, the fibromuscular tube that food passes through—aided by peristaltic contractions—the pharynx to the stomach.
- Stomach, which secretes protein -digesting enzymes called proteases and strong acids to aid in food digestion, before sending the partially digested food to the small intestines.
- Duodenum, the first section of the small intestine that may be the principal site for iron absorption.

Lower Gastrointestinal Tract

The lower gastrointestinal tract includes most of the small intestine and all of the large intestine. According to some sources, it also includes the anus.

The small intestine has three parts:



Small intestine: This image shows the position of the small intestine in the gastrointestinal tract.

- Duodenum: Here the digestive juices from the pancreas (digestive enzymes) and the gallbladder (bile) mix together. The digestive enzymes break down proteins and bile and emulsify fats into micelles. The duodenum contains Brunner's glands that produce bicarbonate, and pancreatic juice that contains bicarbonate to neutralize hydrochloric acid in the stomach.
- Jejunum: This is the midsection of the intestine, connecting the duodenum to the ileum. It contains the plicae circulares and villi to increase the surface area of that part of the GI tract.
- Ileum: This has villi, where all soluble molecules are absorbed into the blood (through the capillaries and lacteals).

The large intestine has four parts:

1. Cecum, the vermiform appendix that is attached to the cecum.
2. Colon, which includes the ascending colon, transverse colon, descending colon, and sigmoid flexure. The main function of the colon is to absorb water, but it also contains bacteria that produce beneficial vitamins like vitamin K.
3. Rectum.
4. Anus.

The ligament of Treitz is sometimes used to divide the upper and lower GI tracts.

This page titled 22.1A: Anatomy of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.1B: Processes and Functions of the Digestive System

Digestion is necessary for absorbing nutrients from food and occurs through two processes: mechanical and chemical digestion.

Learning Objectives

- Describe the processes and functions of the digestive system

Key Points

- Two important functions of the digestive system are digestion and absorption.
- The nutrients that come from food are derived from proteins, fats, carbohydrates, vitamins, and minerals. These complex macromolecules must be broken down and absorbed in the gastrointestinal (GI) tract.
- Mechanical digestion starts in the mouth, with the physical processing of food by the teeth, and continues in the stomach.
- Chemical digestion starts with the release of enzymes in saliva, and continues in the stomach and intestines.
- During absorption, the nutrients that come from food pass through the wall of the small intestine and into the bloodstream.

Key Terms

- **mastication**: The process of mechanical breakdown by the teeth; also known as chewing.
- **bolus**: Moistened and mechanically manipulated food.
- **mechanical digestion**: The breaking down of food into digestible chunks, normally using the teeth.
- **chemical digestion**: A process that involves the action of enzymes to break down food into components that can be absorbed by the small intestine.
- **gastrointestinal tract**: This tract consists of the stomach and intestine, and sometimes includes all the structures from the mouth to the anus. The digestive system is a broader term that includes other structures, including the accessory organs of digestion, such as the liver, gallbladder, and pancreas.

The Digestive System

The proper functioning of the gastrointestinal (GI) tract is imperative for our well being and life -long health. A non-functioning or poorly-functioning GI tract can be the source of many chronic health problems that can interfere with your quality of life.

Here is a look at the importance of two main functions of the digestive system: digestion and absorption.

Digestion

The gastrointestinal tract is responsible for the breakdown and absorption of the various foods and liquids needed to sustain life. Many different organs have essential roles in the digestion of food, from the mechanical breakdown of food by the teeth to the creation of bile (an emulsifier) by the liver.

Bile production plays a important role in digestion: it is stored and concentrated in the gallbladder during fasting stages, and discharged to the small intestine. Pancreatic juices are excreted into the digestive system to break down complex molecules such as proteins and fats.

Absorption

Absorption occurs in the small intestines, where nutrients directly enter the bloodstream.

Each component of the digestive system plays a special role in these complimentary processes. The structure of each component highlights the function of that particular organ, providing a seamless anatomy to keep our body fueled and healthy.

Components of the Digestive System

The digestive system is comprised of the alimentary canal, or the digestive tract, and other accessory organs that play a part in digestion—such as the liver, the gallbladder, and the pancreas. The alimentary canal and the GI tract are terms that are sometimes used interchangeably.

The alimentary canal is the long tube that runs from the mouth (where the food enters) to the anus (where indigestible waste leaves). The organs in the alimentary canal include the mouth (the site of mastication), the esophagus, the stomach, the small and large intestines, the rectum, and the anus. From mouth to anus, the average adult digestive tract is about thirty feet (30′) long.

Processes of Digestion

Food is the body's source of fuel. The nutrients in food give the body's cells the energy they need to operate. Before food can be used it has to be mechanically broken down into tiny pieces, then chemically broken down so nutrients can be absorbed.

In humans, proteins need to be broken down into amino acids, starches into sugars, and fats into fatty acids and glycerol. This mechanical and chemical breakdown encompasses the process of digestion.

To recap these twin processes:

1. Mechanical digestion: Larger pieces of food get broken down into smaller pieces while being prepared for chemical digestion; this process starts in the mouth and continues into the stomach.
2. Chemical digestion: Several different enzymes break down macromolecules into smaller molecules that can be absorbed. The process starts in the mouth and continues into the intestines.

Moistening and Breakdown of Food

Digestion begins in the mouth. A brain reflex triggers the flow of saliva when we see or even think about food. Enzymes in saliva then begin the chemical breakdown of food; teeth aid in the mechanical breakdown of larger food particles.

Saliva moistens the food, while the teeth masticate the food and make it easier to swallow. To accomplish this moistening goal, the salivary glands produce an estimated three liters of saliva per day.

Amylase, the digestive enzyme found in saliva, starts to break down starch into simple sugars before the food even leaves the mouth. The nervous pathway involved in salivary excretion requires stimulation of receptors in the mouth, sensory impulses to the brain stem, and parasympathetic impulses to salivary glands. Once food is moistened and rolled and ready to swallow, it is known as a bolus.

Swallowing and the Movement of Food


For swallowing to happen correctly a combination of 25 muscles must all work together at the same time. Swallowing occurs when the muscles in your tongue and mouth move the bolus into your pharynx.

The pharynx, which is the passageway for food and air, is about five inches (5″) long—a remarkably small space. A small flap of skin called the epiglottis closes over the pharynx to prevent food from entering the trachea, which would cause choking. Instead, food is pushed into the muscular tube called the esophagus. Waves of muscle movement, called peristalsis, move the bolus down to the stomach.

While in the digestive tract, the food is really passing through the body rather than being in the body. The smooth muscles of the tubular digestive organs move the food efficiently along as it is broken down into easily absorbed ions and molecules.

Large-scale Breakdown in the Stomach

Once the bolus reaches the stomach, gastric juices mix with the partially digested food and continue the breakdown process. The bolus is converted into a slimy material called chyme.



Major digestive hormones: There are at least five major digestive hormones in the gut of mammals that help process food through chemical digestion in the gall bladder, duodenum, stomach, and pancrease. These hormones are cholecystokinin, gastric inhibitory polypeptide, motilin, secretin, and gastrin.

The stomach is a muscular bag that maneuvers food particles, mixing highly acidic gastric juice and powerful digestive enzymes with the chyme to prepare for nutrient absorption in the small intestine. Stimulatory hormones such as gastrin and motilin help the stomach pump gastric juice and move chyme. The complex network of hormones eventually prepares chyme for entry into the duodenum, the first segment of the small intestine.

Absorption in the Small Intestine

During absorption, the nutrients that come from food (such as proteins, fats, carbohydrates, vitamins, and minerals) pass through the wall of the small intestine and into the bloodstream. In this way nutrients can be distributed throughout the rest of the body. The small intestine increases surface area for absorption through tiny interior projections, like small fingers, called villi.

Waste Compaction in the Large Intestine

In the large intestine there is resorption of water and absorption of certain minerals as feces are formed. Feces are the waste parts of the food that the body passes out through the anus.

This page titled 22.1B: Processes and Functions of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.1C: Organs of the Digestive System

The organs of the digestive system can be divided into upper and lower digestive tracts. The upper digestive tract consists of the esophagus, stomach, and the small intestine; the lower tract includes all of the large intestine, the rectum, and anus.

Learning Objectives

- Outline the relationship, structure, and function of the digestive organs

Key Points


- The gastrointestinal tract is made up of upper and lower tracts.
- Food moves from the mouth to the stomach via the esophagus.
- The small intestine has three parts: the duodenum, jejunum, and ileum.
- The large intestine has four parts: the cecum, colon, rectum, and anus.

Key Terms

- **small intestine**: A winding, digestive tube and the site of large scale nutrient absorption comprised of the duodenum, jejunum, and ileum.
- **esophagus**: An organ in vertebrates that is a muscular tube through which food passes from the pharynx to the stomach.
- **stomach**: An organ in animals that stores and breaks down food in the process of digestion.
- **large intestine**: The second to last part of the digestive system comprised of the cecum and colon.

The human body uses a variety of mental and physiological cues to initiate the process of digestion. Throughout our gastrointestinal (GI) tract, each organ serves a specific purpose to bring our food from the plate to a digestible substance from which nutrients can be extracted.

The Digestive Tube



The organs of the gastrointestinal tract: This diagram shows the relationship between the various organs of the digestive system. It shows how the oral cavity connects to the esophagus and descends into the stomach and then the small intestine. It

then connects to the large intestine, then the rectum, and, finally, the anus.

Our digestive system is like a long tube, with different segments doing different jobs. The major organs within our digestive system can be split into two major segments of this tube: the upper gastrointestinal tract, and the lower gastrointestinal tract.

The Upper Gastrointestinal Tract

The upper gastrointestinal, or GI, tract is made up of three main parts:

1. The esophagus.
2. The stomach.
3. The small intestine.

The Lower Gastrointestinal Tract

The lower GI tract contains the remainder of the system:

1. The large intestine.
2. The rectum.
3. The anus.

The exact dividing line between upper and lower tracts can vary, depending on which medical specialist is examining the GI tract.

Food Breakdown and Absorption: The Upper GI Tract

When we take a bite of food, the food material gets chewed up and processed in the mouth, where saliva begins the process of chemical and mechanical breakdown. The chewing process is also known as mastication.

When we mix up food with saliva, the resulting mushy wad is called a bolus. The bolus gets swallowed, and begins its journey through the upper gastrointestinal tract.

The Esophagus

The upper GI tract begins with the esophagus, the long muscular tube that carries food to the stomach. The throat cavity in which our esophagus originates is known as the pharynx. As we swallow, the bolus moves down our esophagus, from the pharynx to the stomach, through waves of muscle movement known as peristalsis. Next the bolus reaches the stomach itself.

The Stomach

The stomach is a muscular, hollow bag that is an important part of the upper GI tract. Many organisms have a variety of stomach types, with many segments or even multiple stomachs. As humans, we have only one stomach.

Here our bolus gets mixed with digestive acids, furthering breakdown of the bolus, and turning the bolus material into a slimy mess called chyme. The chyme moves on into the small intestine, where nutrients are absorbed.

The Small Intestine

The small intestine is an impressive digestive tube, spanning an average of 20 feet in length. The twists and turns of the small intestine, along with tiny interior projections known as villi, help to increase the surface area for nutrient absorption.

This snaking tube is made up of three parts, in order from the stomach:

1. The duodenum.
2. The jejunum.
3. The ileum.

As the chyme makes its way through each segment of the small intestine, pancreatic juices from the pancreas start to break down proteins. Soapy bile from the liver, stored in the gallbladder, gets squirted into the small intestine to help emulsify—or break apart—fats.

Now thoroughly digested, with its nutrients absorbed along the path of the small intestine, what remains of our food gets passed into the lower GI tract.

Waste Compaction and Removal: The Lower Gastrointestinal Tract

The Large Intestine (Colon)

Following nutrient absorption, the food waste reaches the large intestine, or colon. The large intestine is responsible for compacting waste material, removing water, and producing feces —our solid-waste product.

Accessory organs like the cecum and appendix, which are remnants of our evolutionary past, serve as special pockets at the beginning of the large intestine. The compacted and dried-out waste passes to the rectum, and out of the body through the anus. Healthy gut bacteria in the large intestine also help to metabolize our waste as it finishes its journey.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23The_Digestive_System**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Digestive System. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digestive_System**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Digestive System. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digestive_System%23Upper_gastrointestinal_tract**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/upper-gastrointestinal-tract**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/lower-gastrointestinal-tract**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Duodenum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Duodenum**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Esophagus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Esophagus**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Stomach**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...um_diagram.svg**. **License**: *Public Domain: No Known Copyright*
- Digestive system diagram edit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Digestive_system_diagram_edit.svg**. **License**: *Public Domain: No Known Copyright*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23The_Digestive_System**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23Introduction**. **License**: *CC BY-SA: Attribution-ShareAlike*
- chemical digestion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chemical+digestion**. **License**: *CC BY-SA: Attribution-ShareAlike*
- mechanical digestion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mechanical_digestion**. **License**: *CC BY-SA: Attribution-ShareAlike*

- gastrointestinal tract. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gastrointestinal_tract**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Small intestine. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...um_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Digestive system diagram edit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Digestive_system_diagram_edit.svg**. **License**: ***Public Domain: No Known Copyright***
- Digestive hormones. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Digestive_hormones.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Digestive System. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digestive_System%23Upper_gastrointestinal_tract**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- stomach. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/stomach**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/esophagus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- large intestine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/large_intestine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Small intestine. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...um_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Digestive system diagram edit. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Digestive_system_diagram_edit.svg**. **License**: ***Public Domain: No Known Copyright***
- Digestive hormones. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Digestive_hormones.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Digestive system diagram en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Digestive_system_diagram_en.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 22.1C: Organs of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.2: NERVOUS SYSTEM OF THE DIGESTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.2A: Enteric Nervous System

22.2B: Gastrointestinal Reflex Pathways

This page titled 22.2: Nervous System of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.2A: Enteric Nervous System

The enteric nervous system (ENS) is a subdivision of the autonomic nervous system (ANS) that directly controls the gastrointestinal system.

Learning Objectives

- Describe the structure and function of the enteric nervous system (ENS)

Key Points

- The enteric nervous system (ENS), which is embedded in the lining of the gastrointestinal system, can operate independently of the brain and the spinal cord.
- The ENS consists of two plexuses, the submucosal and the myenteric. The myenteric plexus increases the tone of the gut and the velocity and intensity of contractions. The submucosal plexus is involved with local conditions and controls local secretion, absorption, and muscle movements.
- While described as a second brain, the enteric nervous system normally communicates with the central nervous system (CNS) through the parasympathetic (via the vagus nerve) and sympathetic (via the prevertebral ganglia) nervous systems, but can still function when the vagus nerve is severed.
- The ENS includes efferent neurons, afferent neurons, and interneurons, all of which make the ENS capable of carrying reflexes and acting as an integrating center in the absence of CNS input.
- The ENS contains support cells, which are similar to the astroglia of the brain, and a diffusion barrier around the capillaries surrounding the ganglia, which is similar to the blood –brain barrier of cerebral blood vessels.

Key Terms

- **enteric nervous system**: A subdivision of the peripheral nervous system that directly controls the gastrointestinal system.

EXAMPLES

The second brain of the enteric nervous system is the reason we get butterflies in our stomach or need to use the restroom more frequently when we are nervous and/or under stress.

The gastrointestinal (GI) system has its own nervous system, the enteric nervous system (ENS). Neurogastroenterology is the study of the enteric nervous system, a subdivision of the autonomic nervous system (ANS) that directly controls the gastrointestinal system. The ENS is capable of autonomous functions such as the coordination of reflexes.

Although it receives considerable innervation from the autonomic nervous system, it can and does operate independently of the brain and the spinal cord. The ENS consists of some 100 million neurons, one-thousandth of the number of neurons in the brain, and about one-tenth the number of neurons in the spinal cord. The enteric nervous system is embedded in the lining of the gastrointestinal system.

Ganglia of the ENS

The neurons of the ENS are collected into two types of ganglia:


1. The myenteric (Auerbach's) plexus, located between the inner and outer layers of the muscularis externa
2. The submucosal (Meissner's) plexus, located in the submucosa

The Myenteric Plexus

The myenteric plexus is mainly organized as a longitudinal chains of neurons. When stimulated, this plexus increases the tone of the gut as well as the velocity and intensity of its contractions. This plexus is concerned with motility throughout the whole gut. Inhibition of the myenteric system helps to relax the sphincters —the muscular rings that control the flow of digested food or food waste.

The Submucosal Plexus

The submucosal plexus is more involved with local conditions and controls local secretion and absorption, as well as local muscle movements. The mucosa and epithelial tissue associated with the submucosal plexus have sensory nerve endings that feed signals to both layers of the enteric plexus. These tissues also send information back to the sympathetic pre-vertebral ganglia, the spinal cord, and the brain stem.



Neural control of the gut: An illustration of neural control of the gut wall by the autonomic nervous system and the enteric nervous system.

Function and Structure of the ENS

The enteric nervous system has been described as a second brain. There are several reasons for this. For instance, the enteric nervous system can operate autonomously. It normally communicates with the central nervous system (CNS) through the parasympathetic (e.g., via the vagus nerve) and sympathetic (e.g., via the prevertebral ganglia) nervous systems. However, vertebrate studies show that when the vagus nerve is severed, the enteric nervous system continues to function.

In vertebrates, the enteric nervous system includes efferent neurons, afferent neurons, and interneurons, all of which make the enteric nervous system capable of carrying reflexes and acting as an integrating center in the absence of CNS input. For instance, the sensory neurons report mechanical and chemical conditions, while the motor neurons control peristalsis and the churning of intestinal contents through the intestinal muscles. Other neurons control the secretion of enzymes.

The enteric nervous system also makes use of more than 30 neurotransmitters, most of which are identical to the ones found in the CNS, such as acetylcholine, dopamine, and serotonin. More than 90% of the body's serotonin is in the gut, as well as about 50% of the body's dopamine, which is currently being studied to further our understanding of its utility in the brain.

The enteric nervous system has the capacity to alter its response depending on factors such as bulk and nutrient composition. In addition, the ENS contains support cells that are similar to the astroglia of the brain, as well as a diffusion barrier around the capillaries that surround the ganglia, which is similar to the blood–brain barrier of the cerebral blood vessels.

Regulation of ENS Function

The parasympathetic nervous system is able to stimulate the enteric nerves in order to increase enteric function. The parasympathetic enteric neurons function in defecation and provide a rich nerve supply to the sigmoid colon, the rectum, and the anus.

Conversely, stimulation of the enteric nerves by the sympathetic nervous system will inhibit enteric function and capabilities. Neurotransmitter secretion and direct inhibition of the enteric plexuses cause this stall in function. If the gut tract is irritated or distended, afferent nerves will send signals to the medulla of the brain for further processing.

This page titled 22.2A: Enteric Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.2B: Gastrointestinal Reflex Pathways

The digestive system functions via a system of long reflexes, short reflexes, and extrinsic reflexes from gastrointestinal (GI) peptides that work together.

Learning Objectives

- Differentiate among the gastrointestinal reflex pathways

Key Points

- Long reflexes to the digestive system involve a sensory neuron that sends external or internal digestive information to the brain. This type of reflex includes reactions to food, emotion, or danger.
- Short reflexes to the digestive system provide shortcuts for the enteric nervous system (ENS) to act quickly and effectively, and form a sort of digestive brain. It reacts to digestive movement and chemical changes.
- The enterogastric reflex is stimulated by the senses. This reflex releases acid in the duodenum or in the stomach, and suppresses the release of digestive proteins.
- The gastrocolic reflex increases movement in the gastrointestinal tract, and reacts to stretches in the stomach walls as well as in the colon. It is responsible for the urge to defecate, the movement of digested material in the small intestine, and it makes room for more food within the stomach.
- The gastroileal reflex works with the gastrocolic reflex to stimulate the urge to defecate. It does so by opening the ileocecal valve and moving the digested contents from the ileum of the small intestine into the colon for compaction.
- GI peptides act on a variety of tissues including the brain, the digestive accessory organs, and the GI tract.

Key Terms

- **gastrocolic reflex**: One of the three extrinsic physiological reflexes that control the motility or peristalsis of the gastrointestinal tract; it involves an increase in the motility of the colon, creates the urge to defecate along with the gastroileal reflex, and helps make room for food in the stomach.
- **enterogastric reflex**: One of the three extrinsic reflexes of the gastrointestinal tract that is stimulated by the presence of acid levels in the duodenum or in the stomach. It releases acids and controls the release of stomach proteins such as gastrin.
- **gastroileal reflex**: One of the three extrinsic reflexes of the gastrointestinal tract that works with the gastrocolic reflex to stimulate the urge to defecate. This reflex is stimulated by the opening of the ileocecal valve and moves the digested contents from the ileum of the small intestine into the colon for compaction.

EXAMPLES

The gastrocolic reflex can cause irritable bowel syndrome. This can lead to abdominal pain, diarrhea, or constipation.

Food in the Digestive System

The digestive system has a complex system of food movement and secretion regulation, which are vital for its proper function. Movement and secretion are regulated by long reflexes from the central nervous system (CNS), short reflexes from the enteric nervous system (ENS), and reflexes from the gastrointestinal system (GI) peptides that work in harmony with each other.

In addition, there are three overarching reflexes that control the movement, digestion, and defecation of food and food waste:

1. The enterogastric reflex
2. The gastrocolic reflex
3. The gastroileal reflex

Long and Short Reflexes

Long reflexes to the digestive system involve a sensory neuron that sends information to the brain. This sensory information can come from within the digestive system, or from outside the body in the form of emotional response, danger, or a reaction to food.

These alternative sensory responses from outside the digestive system are also known as feedforward reflexes. Emotional responses can also trigger GI responses, such as the butterflies in the stomach feeling when nervous.


Control of the digestive system is also maintained by enteric nervous system (ENS), which can be thought of as a digestive brain that helps to regulate motility, secretion, and growth. The enteric nervous system can act as a fast, internal response to digestive stimuli. When this occurs, it is called a short reflex.

Three Main Types of Gastrointestinal Reflex

The Enterogastric Reflex

The enterogastric reflex is stimulated by the presence of acid levels in the duodenum at a pH of 3–4 or in the stomach at a pH of 1.5. When this reflex is stimulated, the release of gastrin from G- cells in the antrum of the stomach is shut off. In turn, this inhibits gastric motility and the secretion of gastric acid (HCl). Enterogastric reflex activation causes decreased motility.

The Gastrocolic Reflex



Peristalis: The gastrocolic reflex is one of a number of physiological reflexes that control the motility, or peristalsis, of the gastrointestinal tract.

The gastrocolic reflex is the physiological reflex that controls the motility, or peristalsis, of the gastrointestinal tract. It involves an increase in motility of the colon in response to stretch in the stomach and the byproducts of digestion in the small intestine. Thus, this reflex is responsible for the urge to defecate following a meal. The small intestine also shows a similar motility response. The gastrocolic reflex also helps make room for food in the stomach.

The Gastroileal Reflex

The gastroileal reflex is a third type of gastrointestinal reflex. It works with the gastrocolic reflex to stimulate the urge to defecate. This urge is stimulated by the opening of the ileocecal valve and the movement of the digested contents from the ileum of the small intestine into the colon for compaction.

GI Peptides that Contribute to Gastrointestinal Signals

GI peptides are signal molecules that are released into the blood by the GI cells themselves. They act on a variety of tissues that include the brain, the digestive accessory organs, and the GI tract.

The effects range from excitatory or inhibitory effects on motility and secretion, to feelings of satiety or hunger when acting on the brain. These hormones fall into three major categories:

1. The gastrin family
2. The secretin family
3. A third family that is composed of the hormones that do not fit into either of these two families

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Autonomic Nervous System (ANS). **Provided by**: Boundless. **Located at**: **www.boundless.com/users/3718...ans-1069-2395/**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medical Physiology/Gastrointestinal Physiology/Principles of GI function. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Medical...Nervous_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Enteric nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Enteric_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Enteric nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Enteric...tem%23Function**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- enteric nervous system. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/enteric_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural control gut. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ontrol_gut.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medical Physiology/Gastrointestinal Physiology/Principles of GI function. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Medical...tinal_Reflexes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gastrocolic reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gastrocolic_reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pharyngeal reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pharyngeal_reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Enterogastric reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Enterogastric_reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gastrointestinal physiology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gastroi...y%23Regulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gastrocolic reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/gastrocolic%20reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- enterogastric reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/enterogastric%20reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural control gut. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Neural_control_gut.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peristalsis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Peristalsis.gif**. **License**: ***CC BY: Attribution***

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Medical Physiology/Gastrointestinal Physiology/Digestion & Absorption. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Medical...rate_Digestion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hydrolysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hydrolysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- amylase. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amylase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peptidase. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/peptidase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amino acid4. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...mino_acid4.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amylase hydrolysisl 1-4. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...lysisl_1-4.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Catabolism schematic.svg - Wikipedia, the free encyclopedia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/w/index.php?...tic.svg&page=1**. **License**: ***Public Domain: No Known Copyright***
- Chemical digestion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chemica...tein_digestion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- zymogen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/zymogen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactose intolerance. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lactose_intolerance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amino acid4. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...mino_acid4.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amylase hydrolysisl 1-4. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Amylase_hydrolysisl_1-4.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Catabolism schematic.svg - Wikipedia, the free encyclopedia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/w/index.php?title=File:Catabolism_schematic.svg&page=1**. **License**: ***Public Domain: No Known Copyright***
- Carbohydrate digestion. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/4/41/Small_intestine_sugar_enzymes.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lipid digestion. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/be/Micelle_fat_absorption.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 22.2B: Gastrointestinal Reflex Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.3: THE PERITONEUM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

22.3: The Peritoneum

This page titled 22.3: The Peritoneum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.3: The Peritoneum

The peritoneum, the serous membrane that forms the lining of the abdominal cavity, covers most of the intra-abdominal organs.

Learning Objectives

- Differentiate among the digestive organs and their location relative to the peritoneum

Key Points

- The peritoneum supports the abdominal organs and serves as a conduit for their blood and lymph vessels and nerves.
- There are two layers of the peritoneum: the outer layer, called the parietal peritoneum, is attached to the abdominal wall; the inner layer, the visceral peritoneum, is wrapped around the internal organs that are located inside the intraperitoneal cavity. The mesentery is the double layer of visceral peritoneum.
- The potential space between these two layers, the peritoneal cavity, is filled with a small amount of slippery serous fluid that allows the two layers to slide freely over each other.
- The structures in the abdomen are classified as intraperitoneal, retroperitoneal, or infraperitoneal, depending on whether they are covered with visceral peritoneum and are attached by mesenteries.
- There are two main regions of the peritoneum connected by the epiploic foramen: the greater sac or general cavity of the abdomen, and the lesser sac or omental bursa.
- Intraperitoneal organs and retroperitoneal organs weave in and out of these membranes, and serve varying functions. Retroperitoneal structures tend to be more static than intraperitoneal ones.

Key Terms

- **greater omentum**: A large fold of visceral peritoneum that hangs down from the stomach.
- **retroperitoneal**: Located outside of the peritoneum.
- **serous membrane**: A thin membrane that secretes serum that lines an internal body cavity, such as the peritoneum, the pericardium, and the pleura.
- **peritoneum**: In mammals, the serous membrane that lines the cavity of the abdomen and that is folded over the viscera.
- **intraperitoneal**: Within the cavity of the peritoneum.
- **mesentery**: The membrane that attaches the intestines to the wall of the abdomen and maintains their position in the abdominal cavity to supply them with blood vessels, nerves, and lymphatics.


EXAMPLES

In one form of dialysis, called peritoneal dialysis, a glucose solution is sent through a tube into the peritoneal cavity. The fluid is left there for a prescribed amount of time to absorb waste products, and then removed through the tube. This form of dialysis is effective because of the the high number of arteries and veins in the peritoneal cavity which, through the mechanism of diffusion, remove waste products from the blood.

The peritoneum is the serous membrane that forms the lining of the abdominal cavity or the coelom. It covers most of the intra-abdominal, or coelomic, organs. It is composed of a layer of mesothelial tissue, supported by a thin layer of connective tissue.

The peritoneum provides support and protection for the abdominal organs, and is the main conduit for the associated lymph vessels, nerves, and abdominal arteries and veins.

The abdominal cavity is the open space surrounded by the vertebrae, abdominal muscles, diaphragm, and pelvic floor. Remember not to confuse the abdominal cavity with the intraperitoneal space, which is located within the abdominal cavity and wrapped in peritoneum tissue. For example, a kidney is inside the abdominal cavity, but is retroperitoneal—located outside the peritoneum.



The peritoneum and the kidney: A sagittal section through the posterior abdominal wall, showing the kidney residing outside the peritoneum.

Although they ultimately form one continuous sheet, there are two layers of peritoneum and potential space between those layers.


- The outer layer, called the parietal peritoneum, is attached to the abdominal wall.
- The inner layer, the visceral peritoneum, is wrapped around the internal organs that are located inside the intraperitoneal cavity.
- The potential space between these two layers is the peritoneal cavity. It is filled with a small amount of slippery serous fluid that allows the two layers to slide freely over each other.

The term mesentery is often used to refer to a double layer of visceral peritoneum. There are generally blood vessels, nerves, and other structures between these layers. The space between the two layers is technically outside of the peritoneal sac, and thus not in the peritoneal cavity.

The Regions of the Peritoneum

There are two main regions of the peritoneum, connected by the epiploic foramen (also known as the omental foramen or foramen of Winslow). The first is the greater sac or general cavity of the abdomen. The second is the lesser sac or omental bursa.

The lesser sac is divided into two omenta: the gastrohepatic and the gastrocolic. The gastrohepatic omentum is attached to the lesser curvature of the stomach and the liver. The gastrocolic omentum hangs from the greater curve of the stomach and loops down in front of the intestines before curving upwards to attach to the transverse colon. Like a curtain of tissue, it is draped in front of the intestines to insulate and protect them.



Substructures of the peritoneum: This is a midsagittal, cross-section drawing of the epiploic foramen, the greater sac or general cavity (red), and the lesser sac or omental bursa (blue).

Abdomen Structures

The structures in the abdomen are classified as intraperitoneal, retroperitoneal, or infraperitoneal depending on whether they are covered with visceral peritoneum and are attached by mesenteries, such as the mensentery and mesocolon.

Intraperitoneal Structures

Intraperitoneal organs include the stomach, the first five centimeters and the fourth part of the the duodenum, the jejunum, the ileum, the cecum, the appendix, the transverse colon, the sigmoid colon, and the upper third of the rectum.


Other organs located in the intraperitoneal space are the liver, spleen, and the tail of the pancreas. In women, the uterus, fallopian tubes, ovaries, and gonadal blood vessels are located in the intraperitoneum.

Retroperitoneal Structures

Retroperitoneal structures include the rest of the duodenum, the ascending colon, the descending colon, the middle third of the rectum, and the remainder of the pancreas. Other organs located in the retroperitoneal space are the kidneys, adrenal glands, proximal ureters, and renal vessels. Organs located below the peritoneum in the subperitoneal space include the lower third of the rectum and the urinary bladder.

Intraperitoneal Structures

Structures that are intraperitoneal are generally mobile, while those that are retroperitoneal are relatively fixed in their location. Some structures, such as the kidneys, are primarily retroperitoneal, while others such as the majority of the duodenum, are secondarily retroperitoneal, meaning that structure developed intraperitoneally, but lost its mesentery and thus became retroperitoneal.



Peritoneum: The peritoneum illustrated, indicated by blue.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Peritoneum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peritoneum%23Structure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peritoneum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Periton...nal_structures**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peritoneum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peritoneum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Peritoneum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Periton...inical_aspects**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Greater omentum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Greater_omentum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mesentery. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/mesentery**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peritoneum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/peritoneum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intraperitoneal. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/intraperitoneal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- serous membrane. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/serous_membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1040. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1040.png**. **License**: ***Public Domain: No Known Copyright***
- Gray1125. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1125.png**. **License**: ***Public Domain: No Known Copyright***
- Gray1035. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1035.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 22.3: The Peritoneum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.4: THE ALIMENTARY CANAL

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

22.4A: Mouth

22.4B: Pharynx

22.4C: Esophagus

This page titled 22.4: The Alimentary Canal is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.4A: Mouth

Learning Objectives


- Describe the features of the mouth that play a role in digestion

The mouth has a variety of roles in human anatomy and sociology. While its primary function is to begin the process of mechanically and chemically digesting food, the mouth is also the beginning of the alimentary canal—a larger digestive tube. Without the human mouth, expressions of the lips and language of the tongue and throat would be impossible.

The mouth is the first portion of the alimentary canal. It receives food and moistens the food with saliva, while the food is mechanically processed (mastication) by the teeth. The mouth is also known as the oral cavity, and within the oral cavity sits the tongue, the soft and hard palate, the uvula, and numerous salivary glands.

The oral mucosa is the mucous membrane epithelial tissue that lines the inside of the mouth. This membrane maintains a moist and lubricated environment within the mouth to prepare the digestive system for the entry of food.

The Mouth as a Communication and Breathing Tool



Inside of the mouth: An illustration of the inside of a human mouth. The cheeks have been omitted in the drawing and the lips pulled back for an unobstructed view of the teeth, tongue, jaw bones, uvula, and alimentary canal.

In addition to its primary function as the beginning of the digestive system, the mouth also plays a significant role in human communication and breathing. The primary features of human voice are produced in the throat, but the tongue, lips, and jaw also work together to produce the range of sounds we see in human language.

Air is drawn in through the mouth to the trachea and lungs, and the lips and tongue form words. The lips mark the transition from the mucous membrane to the outer epithelial skin that covers most of the body. Lips are remarkably sensitive and often serve as an infant's second hands with which to explore the world.

Mechanical Food Breakdown by Teeth

In the digestive process, the mouth's purpose is to prepare food for further digestion in the stomach and the small intestine. This

process begins with the mechanical breakdown of food by the teeth, which fit into the alveolar arches. The front teeth (incisors and canines) are used to cut and tear food, while the teeth further back (bicuspids and molars) crush and grind.

Food Lubrication and Chemical Digestion By Saliva

Saliva is projected from three main pairs of salivary glands: the large parotid glands near the cheeks, the submandibular glands beneath the mandible, and the sublingual glands beneath the tongue.

Saliva keeps the mouth moist and lubricates the food, helping the tongue form the food into a soft wad, called a bolus. The fluid of saliva also contains several enzymes, notably lysozyme—an antibacterial agent—and amylase, which catalyzes large starch molecules into simpler sugars via hydrolysis.



***Cross section of the head and neck**: A cross section of the head and neck in mid-sagittal view, showing the structures of the mouth and throat.*

Once properly chewed and lubricated, food and drink are swallowed into the esophagus, the tube that leads to the stomach.

The Structures of the Lips and External Mouth

Infant humans are born with an instinctual sucking reflex, by which they know how to gain nourishment using their lips and jaw. The philtrum, or bow of the lip, is the vertical groove or dip just below the nose. The nasolabial folds are the deep creases of tissue that extend from the nose to the sides of the mouth. One of the first signs of age on the human face is the increase in prominence of the nasolabial folds.

Key Points

- The mouth is also known as the oral cavity. Its purpose is to mechanically break down food, moisten it with saliva, and swallow the food into the esophagus and the stomach.
- While vocal sounds are primarily produced in the throat, the tongue, lips, and jaw are also needed to produce the range of sounds included in human language.
- Saliva is produced by three main pairs of salivary glands: the parotid, the submandibular, and sublingual. When food is chewed and mixed with this saliva, the resulting wad is known as a bolus.

Key Terms

- **mastication**: The process of physical and mechanical breakdown of food; chewing.
- **mucous membrane**: A membrane that secretes mucus. It forms the lining of various body passages that communicate with the air, such as the respiratory, genitourinary, and alimentary tracts.

- **mouth**: The opening of a organism through which food is ingested.
- **saliva**: A clear, slightly alkaline liquid secreted into the mouth by the salivary glands and mucous glands that consists of water, mucin, protein, and enzymes. It moistens the mouth, lubricates ingested food, and begins the breakdown of starches.
- **uvula**: A soft, punching-bag-like piece of tissue that hangs at the back of the mouth and functions in closing the air passages during swallowing, in conjunction with the epiglottis of the trachea.
- **hard palate**: The bony roof of the mouth, located ventrally to the soft palate.
- **alimentary canal**: The organs of a human or a non-human animal through which food passes.
- **alveolar arch**: The part of the upper or lower jawbones in which the teeth are set.

This page titled 22.4A: Mouth is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.4B: Pharynx

The pharynx is part of the digestive and respiratory systems and consists of three main parts: the nasopharynx, oropharynx, and laryngopharynx.

Learning Objectives

- Describe the pharynx's role in digestion


Key Points

- The human pharynx (plural: pharynges) is the part of the throat situated immediately inferior to (below) the mouth and nasal cavity, and superior to the esophagus and larynx.
- The human pharynx is conventionally divided into three sections: the nasopharynx (epipharynx), the oropharynx (mesopharynx), and the laryngopharynx (hypopharynx).
- The nasopharynx extends from the base of the skull to the upper surface of the soft palate. Adenoids are lymphoid tissue structures located in the posterior wall of the nasopharynx. The nasopharynx communicates with the middle ear, nasal cavities, and auditory tube.
- The oropharynx lies behind the oral cavity and extends from the uvula to the level of the hyoid bone. A flap of connective tissue called the epiglottis closes over the glottis when food is swallowed to prevent aspiration.
- The laryngopharynx is the caudal part of the pharynx; it is the part of the throat that connects to the esophagus. It includes three major sites: the pyriform sinus, the postcricoid area, and the posterior pharyngeal wall.

Key Terms

- **adenoid**: One of the two folds of lymphatic tissue covered by ciliated epithelium. They are found in the roof and posterior wall of the nasopharynx at the back of the throat behind the uvula. They may obstruct normal breathing and make speech difficult when swollen—a condition often called adenitis.
- **epiglottis**: A cartilaginous organ in the throat of terrestrial vertebrates that covers the glottis when swallowing to prevent food and liquid from entering the trachea. In Homo sapiens it is also a speech organ.
- **uvula**: The fleshy appendage that hangs from the back of the palate and closes the nasopharynx during swallowing.
- **pharynx**: The part of the alimentary canal that extends from the mouth and nasal cavities to the larynx, where it becomes continuous with the esophagus.

The human pharynx (plural: pharynges) is the part of the throat situated immediately inferior to (below) the mouth and nasal cavity, and superior to the esophagus and larynx.



Head and neck overview: The human pharynx is situated immediately below the mouth and nasal cavity, and above the esophagus and larynx.

The human pharynx is conventionally divided into three sections: the nasopharynx (epipharynx), the oropharynx (mesopharynx), and the laryngopharynx (hypopharynx). The pharynx is part of the digestive system and also the respiratory system, as well as an important part in vocalization.

Nasopharynx

The nasopharynx is the most cephalad (toward the head) portion of the pharynx. It extends from the base of the skull to the upper surface of the soft palate. It includes the space between the internal nares and the soft palate, and lies superior to the oral cavity.

The pharyngeal tonsils, more commonly referred to as the adenoids, are lymphoid tissue structures located in the posterior wall of the nasopharynx. Polyps or mucus can obstruct the nasopharynx, as can congestion due to an upper respiratory infection.

The eustachian tubes connect the middle ear to the pharynx and open into the nasopharynx. The opening and closing of the eustachian tubes serves to equalize the barometric pressure in the middle ear with that of the ambient atmosphere.

The anterior portion of the nasopharynx connects with the nasal cavities through openings known as choanae. The nasopharynx and its associated nasal tissues are lined with ciliated pseudostratified columnar epithelium, which is excellent for sweeping debris from the nasal passages.

The Pharyngeal Ostia

On the lateral walls of the nasopharynx are the pharyngeal ostia of the auditory tube—triangle-shaped openings bound from behind by a firm prominence called the torus tubarius or cushion.

This binding is formed by a cartilaginous tube-like opening. Two folds arise from the cartilaginous opening:

1. The salpingopharyngeal fold, a vertical fold of mucous membrane that extends from the inferior part of the torus.
2. The salpingopalatine fold, a smaller fold that extends from the superior part of the torus to the palate.

Behind the ostia of the auditory tube is a deep recess known as the pharyngeal recess (or fossa of Rosenmüller).

The posterior wall of the nasopharynx holds the pharyngeal tonsils, which can be especially marked in childhood. Superior to the pharyngeal tonsil, in the midline, an irregular flask-shaped depression of the mucous membrane sometimes extends upward; it is known as the pharyngeal bursa.

Oropharynx

The oropharynx or mesopharynx lies behind the oral cavity and extends from the uvula to the level of the hyoid bone. It opens anteriorly, through the isthmus faucium, into the mouth, and contains the palatine tonsil—another grouping of adenoid tissue.

The anterior wall consists of the base of the tongue and the the epiglottis tissue. The lateral walls are made up of the tonsil and associated tonsilar tissues. The superior wall consists of the inferior surface of the soft palate and the uvula.

Because both food and air pass through the pharynx, a flap of connective tissue called the epiglottis closes over the glottis (tracheal opening) when food is swallowed to prevent accidental inhalation. The oropharynx is lined by non-keratinized stratified squamous epithelium.

Laryngopharynx

The hypopharynx or laryngopharynx is the caudal (most inferior) part of the pharynx; it is the part of the throat that connects to the esophagus. It lies inferior to the epiglottis and extends to the location where this common pathway diverges into the respiratory (larynx) and digestive (esophagus) pathways.

At that point, the laryngopharynx is continuous with the esophagus posteriorly. The esophagus conducts food and fluids to the stomach; air enters the larynx anteriorly. During swallowing, food has the right of way and air passage temporarily stops.

The laryngopharynx includes three major sites:

1. The pyriform sinus.
2. The postcricoid area.
3. The posterior pharyngeal wall.

Like the oropharynx above it, the laryngopharynx serves as a passageway for food and air and is lined with a stratified squamous epithelium.

This page titled 22.4B: Pharynx is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.4C: Esophagus

The esophagus is a muscular tube that moves food from the pharynx to the stomach via peristalsis.

Learning Objectives

- Describe the role of the esophagus in digestion

Key Points

- The esophagus is the muscular tube that moves food material from the pharynx to the stomach via waves of muscle movement known as peristalsis. The junction between the esophagus and the stomach is known as the gastroesophageal junction or GE junction.
- The entry to the esophagus opens only when swallowing or vomiting due to specialized muscles that control the opening.


Key Terms

- **esophagus**: The esophagus is an organ in vertebrates that consists of a muscular tube through which food passes from the pharynx to the stomach.
- **peristalsis**: The rhythmic, wave-like contraction of both longitudinal and circular smooth muscle fibers within the digestive tract that forces food through it.
- **mucus**: A slippery secretion from the lining of the mucous membranes.

EXAMPLES

Swallowing is a voluntary act that utilizes the muscles of the mouth and tongue to push food into the esophagus. Once food material is pushed into the throat, or pharynx, the trachea (windpipe) is blocked by a flap of tissue known as the epiglottis to prevent the aspiration of food. Food then moves down the esophageal tube through waves of muscle movement, or peristalsis, until it reaches the stomach.

The esophagus is an organ in vertebrates that consists of a muscular tube through which food passes from the pharynx to the stomach. It is a major component of the upper digestive system.



Esophagus: The location of the esophagus within the greater digestive system in humans.

The word esophagus is derived from the Latin œsophagus, which derives from the Greek word oisophagos, meaning entrance for eating. It is lined with mucus to aid in the passage of food.

Length and Location

In humans the esophagus is continuous with the laryngeal part of the pharynx within the neck, and it passes through the thorax diaphragm and into abdomen to reach the cardiac orifice of the stomach. It is usually about 10–50 cm long depending on an individual's height. Due to the inferior pharyngeal constrictor muscle, the entry to the esophagus opens only when swallowing or vomiting.

Layers of Tissue

The esophageal tube in humans is comprised of two main layers of smooth muscle, though striated muscle comprises the tube near the pharynx. This combination of muscle tissue allows peristalsis to push food downward, and aids in regurgitation at the pharynx.

The innermost layer of smooth muscle is arranged in a series of concentric rings, while the outermost layer is arranged longitudinally.

In much of the gastrointestinal tract, smooth muscles contract in sequence to produce a peristaltic wave which forces a ball of food (called a bolus) from the pharynx to the stomach.

The Gastroesophageal Junction

The junction between the esophagus and the stomach (the gastroesophageal junction or GE junction) is not actually considered a valve in the anatomical sense, although it is sometimes called the cardiac sphincter.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- mucous membrane. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mucous_membrane. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human mouth. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_mouth. **License**: *CC BY-SA: Attribution-ShareAlike*
- Philtrum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Philtrum. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mouth. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mouth. **License**: *CC BY-SA: Attribution-ShareAlike*
- alveolar arch. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/alveolar%20arch. **License**: *CC BY-SA: Attribution-ShareAlike*
- saliva. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/saliva. **License**: *CC BY-SA: Attribution-ShareAlike*
- mouth. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mouth. **License**: *CC BY-SA: Attribution-ShareAlike*
- alimentary canal. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/alimentary_canal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mouth. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mouth. **License**: *CC BY-SA: Attribution-ShareAlike*
- Esophagus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Esophagus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human pharynx. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_pharynx. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pharynx. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pharynx. **License**: *CC BY-SA: Attribution-*

ShareAlike
- adenoid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/adenoid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- uvula. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/uvula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pharynx. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pharynx**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epiglottis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/epiglottis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mouth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mouth**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Esophagus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Esophagus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu01 head neck. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Illu01_head_neck.jpg**. **License**: ***Public Domain: No Known Copyright***
- peristalsis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/peristalsis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/esophagus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Esophagus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Esophagus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mucus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mucus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mouth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mouth**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Esophagus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Esophagus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Illu01 head neck. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Illu01_head_neck.jpg**. **License**: ***Public Domain: No Known Copyright***
- Digestive_system_diagram_edit.svg.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digestion**. **License**: ***Public Domain: No Known Copyright***

This page titled 22.4C: Esophagus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.5: LAYERS OF THE ALIMENTARY CANAL

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.5A: Mucosa
22.5B: Submucosa
22.5C: Muscularis
22.5D: Serosa

This page titled 22.5: Layers of the Alimentary Canal is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.5A: Mucosa

The mucosa, composed of simple epithelium cells, is the innermost layer of the gastrointestinal (GI) tract. It is the absorptive and secretory layer of the GI tract.

Learning Objectives

- Describe the structure and function of the mucosa of the GI tract

Key Points

- The mucosa is the innermost layer of the GI tract. It is made up of three layers: the epithelium, lamina propria, and muscularis mucosae.
- The mucosa surrounds the lumen, or open space within the digestive tube. This layer comes in direct contact with digested food (chyme).
- The epithelium of the mucosa is particularly specialized, depending on the portion of the digestive system.

Key Terms


- **lumen**: The inside space of a tubular structure, such as an artery or intestine.

Layers of GI Tissue

The GI tract is composed of four layers. Each layer has different tissues and functions. From the inside out they are called: mucosa, submucosa, muscularis, and serosa.

The mucosa is the innermost layer, and functions in absorption and secretion. It is composed of epithelium cells and a thin connective tissue.


The mucosa contains specialized goblet cells that secrete sticky mucus throughout the GI tract. On the mucosa layer, small finger-like projections called villi and microvilli help to increase surface area for nutrient absorption.



Layers of GI tissue: Note the mucosa, located at the innermost layer.

Layers of Tissue Within the Mucosa

Since the mucosa is the innermost layer within the GI tract, it surrounds an open space known as the lumen. Food, mucus, and digestive juices pass through the lumen, and the mucosa comes in direct contact with digested food (chyme).



General structure of the gut wall: This cross section shows the mucosa in relation to the interior space, or lumen.

The mucosa is made up of three layers:

1. The epithelium is the innermost layer and it is responsible for most digestive, absorptive, and secretory processes.
2. The lamina propria is a layer of connective tissue that is unusually cellular compared to most connective tissue.
3. The muscularis mucosae is a thin layer of smooth muscle and its function is still under debate.

The mucosae (singular: mucosa) are highly specialized in each organ of the gastrointestinal tract in order to deal with different digestive tract conditions. The most variation is seen in the epithelium tissue layer of the mucosa.

- In the esophagus, the epithelium is stratified, squamous, and non-keratinizing, for protective purposes.
- In the stomach. the epithelium is simple columnar, and is organized into gastric pits and glands to deal with secretion.
- In the small intestine, the epithelium (particularly the ileum) is specialized for absorption, with villi and microvilli increasing surface area.

This page titled 22.5A: Mucosa is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.5B: Submucosa

The submucosa is a dense, irregular layer of connective tissue with large blood vessels, lymphatics, and nerves that supports the mucosa.

Learning Objectives

- Describe the submucosa of the GI tract

Key Points

- The absorbed elements that pass through the mucosa are picked up from the blood vessels of the submucosa.
- In the gastrointestinal tract, the submucosa is the layer of dense, irregular connective tissue or loose connective tissue that supports the mucosa, as well as joins the mucosa to the bulk of underlying smooth muscle (fibers that run circularly within a layer of longitudinal muscle).
- Tiny parasympathetic ganglia are scattered around to form the submucosal plexus (or Meissner's plexus) where preganglionic parasympathetic neurons create synapses with postganglionic nerve fibers that supply the muscularis mucosae.

Key Terms

- **parasympathetic ganglia**: The autonomic ganglia of the parasympathetic nervous system; they lie near or within (respectively) the organs they innervate.
- **lymphatic**: The system that carries a clear fluid called lymph that is formed from interstitial fluid collected through the capillaries.
- **nerve**: A bundle of neurons with their connective tissue sheaths, blood vessels, and lymphatics.

The Layers of the GI tract


The GI tract is composed of four layers. Each layer has different tissues and functions. From the inside out they are called:

1. Mucosa
2. Submucosa
3. Muscularis
4. Serosa

The Submucosa

The submucosa is relatively thick, highly vascular, and serves the mucosa. The absorbed elements that pass through the mucosa are picked up from the blood vessels of the submucosa.


The submucosa also has glands and nerve plexuses. The submucosa lies under the mucosa and consists of fibrous connective tissue, separating the mucosa from the next layer, the muscularis externa.



Layers of stomach lining: Stomach. The serosa is labeled at far right, and is colored yellow.

The Muscularis

The muscularis in the stomach differs from that of other GI organs in that it has three layers of muscle instead of two. Under these muscle layers is the adventitia—layers of connective tissue that are continuous with the omenta.



General structure of the gut wall: The general structure of the gut wall is illustrated.

The submucosa consists of a dense irregular layer of connective tissue with large blood vessels, lymphatics, and nerves that branch into the mucosa and muscularis externa. It contains Meissner's plexus, an enteric nervous plexus, situated on the inner surface of the muscularis externa.

In the gastrointestinal tract, the submucosa is the layer of dense irregular connective tissue or loose connective tissue that supports the mucosa. It also joins the mucosa to the bulk of underlying smooth muscle (fibers running circularly within layer of longitudinal muscle).

Blood vessels, lymphatic vessels, and nerves (all supplying the mucosa) will run through here. Tiny parasympathetic ganglia are scattered around forming the submucosal plexus (or Meissner's plexus) where preganglionic parasympathetic neurons create synapses with the postganglionic nerve fibers that supply the muscularis mucosae.

This page titled 22.5B: Submucosa is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.5C: Muscularis

The muscularis is responsible for the segmental contractions and peristaltic movements in the gastrointestinal (GI) tract.

Learning Objectives

- Identify the function of muscularis in the GI tract

Key Points


- The muscularis, or muscularis externa, consists of an inner circular muscular layer and a longitudinal outer muscular layer. The coordinated contractions of these layers is called peristalsis, which propels the food through the GI tract.
- Between the two muscle layers is the myenteric or Auerbach's plexus, which controls peristalsis.
- In the colon, the muscularis externa is much thicker because the feces are large and heavy, requiring more force to push along.
- The stomach has a third layer of muscularis externa: the inner oblique layer. This helps churn the chyme in the stomach.
- Peristaltic activity in the muscularis externa is regulated by the enteric nervous system and the autonomic nervous system.

Key Terms

- **muscularis externa**: A region of muscle in many organs in the vertebrate body, adjacent to the submucosa membrane. It is responsible for gut movement such as peristalsis.
- **oblique layer**: This layer is responsible for creating the motion that churns and physically breaks down the food.
- **tiniae coli**: These are three, separate longitudinal ribbons of smooth muscle on the outside of the ascending, transverse, descending, and sigmoid colons.

The gastrointestinal (GI) tract is composed of four layers of tissue, known as tunics. Each layer has different structures and functions. From the inside out they are called the mucosa, submucosa, muscularis externa, and serosa.

Structure of the Muscularis Externa



Muscularis mucosa of the submucosa: The muscularis mucosa is adjacent to the submucosa, and should not be confused


with the muscularis externa.

The muscularis externa is responsible for segmental contractions and peristaltic movement in the GI tract. These muscles cause food to move and churn together with digestive enzymes down the GI tract. The muscularis externa consists of an inner circular layer and a longitudinal outer muscular layer. It should not be confused with a thin layer of muscle known as the muscularis mucosa, which lies within the submucosa, a layer of tissue adjacent to the muscularis externa. The muscularis mucosa is made up of smooth muscle, and is most prominent in the stomach.

Within the muscularis externa, the circular muscle layer prevents food from traveling backward, while the longitudinal layer shortens the tract. The layers are not truly longitudinal or circular, rather the layers of muscle are helical with different pitches. The inner circular is helical with a steep pitch and the outer longitudinal is helical with a much shallower pitch.

The coordinated contractions of these layers is called peristalsis. Between the two muscle layers is the myenteric or Auerbach's plexus, which controls peristalsis. Peristaltic activity is regulated by these nerve cells, and the rate of peristalsis can be modulated by the rest of the autonomic nervous system.

The thickness of muscularis externa varies in each part of the tract. In the colon, for example, the muscularis externa is much thicker because the feces are large and heavy, and require more force to push along. The outer longitudinal layer of the colon thins out into three discontinuous longitudinal bands known as tiniae coli (bands of the colon). This is one of the three features helping to distinguish between the large and small intestine.



General Structure of the gut wall: General structure of the gut wall—the muscularis externa is labeled circular muscle and longitudinal muscle here.

Occasionally in the large intestine (two to three times a day), there will be mass contraction of certain segments, moving a lot of feces along. This is generally when one gets the urge to defecate.

The pylorus of the stomach has a thickened portion of the inner circular layer: the pyloric sphincter. Alone among the GI tract, the stomach has a third layer of muscularis externa. This is the inner oblique layer, and helps churn the chyme in the stomach.

This page titled 22.5C: Muscularis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.5D: Serosa

Serosa consists of a secretory epithelial layer and a thin connective tissue layer that reduce the friction from muscle movements.

Learning Objectives

- Diagram the structure of the serosa

Key Points

- The serous layer provides a partition between the internal organs and the abdominal cavity.
- Cells of the serous layer secrete a serous fluid that provides lubrication to reduce friction.
- The connective tissue layer provides blood vessels and nerves.
- The three serous cavities within the human body are the pericardial cavity (surrounding the heart), the pleural cavity (surrounding the lungs), and peritoneal cavity (surrounding most organs of the abdomen).
- The serous membrane covers the heart; it has an inner layer (the parietal pericardium) and an outer layer (the visceral pericardium).
- The serosa of the uterus is called the perimetrium.


Key Terms

- **pleural cavity**: The body cavity that surrounds the lungs and is enclosed by the pleura.
- **coelom**: A fluid-filled cavity within the body of an animal. The digestive system is suspended within the cavity that is lined by a tissue called the peritoneum.
- **serosa**: A membrane that lines an internal cavity to protect the contents and secretes serum.

The Serous Membrane

In anatomy, the serous membrane (or serosa) is a smooth membrane that consists of a thin connective tissue layer and a thin layer of cells that secrete serous fluid. Serous membranes line and enclose several body cavities, known as serous cavities, where they secrete a lubricating fluid to reduce friction from muscle movements.

Serosa is not to be confused with adventitia, a connective tissue layer that binds together structures rather than reduces friction between them.



Layers of stomach lining: The serosa is labeled at far right, and is colored yellow.

Each serous membrane is composed of a secretory epithelial layer and a connective tissue layer underneath. The epithelial layer, known as mesothelium, consists of a single layer of avascular flat nucleated cells (simple squamous epithelium) that produce the lubricating serous fluid. This fluid has a consistency similar to thin mucus.

These cells are bound tightly to the underlying connective tissue. The connective tissue layer provides the blood vessels and nerves for the overlying secretory cells, and also serves as the binding layer that allows the whole serous membrane to adhere to organs and other structures.

For the heart, the surrounding serous membranes include: the outer, inner, parietal pericardium, and visceral pericardium (epicardium). Other parts of the body may also have specific names for these structures. For example, the serosa of the uterus is called the perimetrium.

The pericardial cavity (surrounding the heart), pleural cavity (surrounding the lungs) and peritoneal cavity (surrounding most organs of the abdomen) are the three serous cavities within the human body. While serous membranes have a lubricative role to play in all three cavities, in the pleural cavity it has a greater role to play in the function of breathing.

The serous cavities are formed from the intraembryonic coelom and are basically an empty space within the body surrounded by a serous membrane. Early in embryonic life, visceral organs develop adjacent to a cavity and invaginate into the bag-like coelom.

Therefore each organ becomes surrounded by a serous membrane—they do not lie within the serous cavity. The layer in contact with the organ is known as the visceral layer, while the parietal layer is in contact with the body wall.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gastrointestinal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lumen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lumen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Serosa. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Serosa**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/parasympathetic-ganglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Submucosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Submucosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/lymphatic. **License**: *CC BY-SA: Attribution-ShareAlike*
- nerve. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serous membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serous_membrane. **License**: *Public Domain: No Known Copyright*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscularis mucosae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscularis_mucosae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/oblique-layer. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology/definition/tiniae-coli. **License**: *CC BY-SA: Attribution-ShareAlike*
- muscularis externa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/muscularis%20externa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serous membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serous_membrane. **License**: *Public Domain: No Known Copyright*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscularis mucosae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscularis_mucosae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- coelom. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/coelom. **License**: *CC BY-SA: Attribution-ShareAlike*
- serosa. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- pleural cavity. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pleural_cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*

- Serous membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serous_membrane. **License**: *Public Domain: No Known Copyright*
- Gastrointestinal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrointestinal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscularis mucosae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscularis_mucosae. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serous membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serous_membrane. **License**: *Public Domain: No Known Copyright*

This page titled 22.5D: Serosa is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7: THE LIVER

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.7A: The Liver

22.7B: Anatomy of the Liver

22.7C: Histology of the Liver

22.7D: Blood Supply to the Liver

22.7E: Liver Function

22.7F: Bile

This page titled 22.7: The Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7A: The Liver

The liver makes bile, which is essential for the digestion of fats.

Learning Objectives

- Summarize the roles of the liver in digestion

Key Points


- The liver is a vital organ with a wide range of functions, including detoxification, protein synthesis, and the production of bile, which is necessary for digestion.
- The bile produced by the liver is essential for the digestion of fats. Bile is formed in the liver and either stored in the gallbladder or released directly into the small intestine.

Key Terms

- **liver**: A large organ in the body that stores and metabolizes nutrients, destroys toxins, and produces bile. It is responsible for thousands of biochemical reactions.
- **bile**: A bitter, brownish-yellow or greenish-yellow secretion produced by the liver, stored in the gallbladder, and discharged into the duodenum, where it aids the process of digestion.

The Role of the Liver

The liver normally weighs between 1.3—3.0 kilograms and is a soft, pinkish-brown organ. It is the second-largest organ in the body, and is located on the right side of the abdomen.



Human liver: Photo of recently removed human liver.

The liver plays a major role in metabolism and has a number of functions in the body, including glycogen storage, plasma protein synthesis, and drug detoxification. It also produces bile, which is important for digestion.

The liver is supplied by two main blood vessels on its right lobe: the hepatic artery and the portal vein. The portal vein brings venous blood from the spleen, pancreas, and small intestine so that the liver can process the nutrients and byproducts of food digestion.

Bile

The bile produced in the liver is essential for the digestion of fats. Bile is formed in the liver, and it is stored in the gallbladder or released directly into the small intestine. After being stored in the gallbladder, the bile becomes more concentrated than when it left the liver; this increases its potency and intensifies its effect in digesting fats.

This page titled 22.7A: The Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7B: Anatomy of the Liver

The liver is located in the abdomen and has four lobes.

Learning Objectives

- Detail the anatomy of the liver

Key Points

- A human liver normally weighs 1.44–1.66 kg (3.2–3.7 lb), and is a soft, pinkish-brown, triangular organ.
- The liver is both the largest internal organ (the skin being the largest organ overall) and the largest gland in the human body.
- The liver is connected to two large blood vessels: the hepatic artery and the portal vein.
- Traditionally, the liver is divided into four lobes: left, right, caudate, and quadrate. The lobes are further divided into lobules, the functional units of the liver.
- Each lobule is made up of millions of hepatic cells that are the basic metabolic cells of the liver.


Key Terms

- **lobule**: A subdivision of the four main liver lobes, the basic functional unit of the liver.

The Liver

The human liver is both the largest internal organ (the skin being the largest organ overall) and the largest gland in the human body. It is a soft, pinkish-brown, triangular organ normally weighing 1.44–1.66 kg (3.2–3.7 lb).

The liver has a wide range of functions including detoxification, protein synthesis, and the production of the biochemicals necessary for digestion. It is located in the right upper quadrant of the abdominal cavity, resting just below the diaphragm. The liver lies to the right of the stomach and overlies the gall bladder.



The position of the liver: The spatial relationship between the liver, stomach, gall bladder, and pancreas. The liver is seen above the stomach, gall bladder, and pancreas.

The liver is connected to two large blood vessels, the hepatic artery and the portal vein. The hepatic artery carries blood from the aorta to the liver, whereas the portal vein carries blood containing the digested nutrients from the entire gastrointestinal tract, and also from the spleen and pancreas to the liver. These blood vessels subdivide into capillaries that then lead to a lobule.

Lobes of the Liver

Traditionally, the liver is divided into four lobes: left, right, caudate, and quadrate. The lobes are further divided into lobules, the functional units of the liver. Each lobule is made up of millions of hepatic cells that are the basic metabolic cells of the liver.

This page titled 22.7B: Anatomy of the Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7C: Histology of the Liver

Hepatocytes are the main tissue cells of the liver. The gallbladder contains the mucosa, muscularis, perimuscular, and serosa layers.

Learning Objectives

- Describe the histology of the liver


Key Points

- A hepatocyte is the main tissue cell of the liver and makes up 70–80% of the liver's cytoplasmic mass.
- Hepatocytes contain large amounts of rough endoplasmic reticulum and free ribosomes.
- Hepatocytes are involved in: protein synthesis; protein storage; the transformation of carbohydrates,; the synthesis of cholesterol, bile salts, and phospholipids; and detoxification, modification, and excretion of exogenous and endogenous substances.
- Hepatocytes are unique in that they are one of the few types of cells in the human body that are capable of regeneration.
- There are several different layers of the gallbladder: the mucosa (epithelium and lamina propria), the muscularis, the perimuscular, and the serosa.

Key Terms

- **hepatocyte**: Any of the cells in the liver responsible for the metabolism of proteins, carbohydrates, and lipids, and for detoxification.

The Liver



Hepatocytes: A cross-section of a human liver that shows hepatocytes.

A hepatocyte is the main tissue cell of the liver and makes up 70–80% of the liver's cytoplasmic mass. Hepatocytes contain large amounts of rough endoplasmic reticulum and free ribosomes. Hepatocytes are involved in:

- Protein synthesis.
- Protein storage.
- The transformation of carbohydrates.
- The synthesis of cholesterol, bile salts, and phospholipids.
- The detoxification, modification, and excretion of exogenous and endogenous substances.

Hepatocytes also initiate the formation and secretion of bile. Hepatocytes are organized into plates separated by vascular channels (sinusoids) for blood vessels. The hepatocyte plates are one cell thick in mammals.

Hepatocytes are unique in that they are one of the few types of cell in the human body that are capable of regeneration. Hepatocytes are derived from hepatoblasts, the precursor stem cell of the liver that divides to produce new hepatocytes. The liver is capable of complete regeneration from as little as 25% of the original organ.

This page titled 22.7C: Histology of the Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7D: Blood Supply to the Liver

In the hepatic portal system, the liver receives a dual blood supply from the hepatic portal vein and the hepatic arteries.

Learning Objectives

- Outline the blood flow to and from the liver

Key Points


- The hepatic portal vein supplies 75% of the blood to the liver, while the hepatic arteries supply the remaining 25%.
- Approximately half of the liver's oxygen demand is met by the hepatic portal vein, and half is met by the hepatic arteries.
- The hepatic portal system connects the capillaries of the gastrointestinal tract with the capillaries in the liver. Nutrient-rich blood leaves the gastrointestinal tract and is first brought to the liver for processing before being sent to the heart.

Key Terms

- **hepatic arteries**: A blood vessel that supplies oxygenated blood to the liver.
- **hepatic portal vein**: A vessel located in the abdominal cavity that is formed by the union of the superior mesenteric and splenic veins that channel blood from the gastrointestinal tract and spleen to the capillary beds in the liver.
- **cofactors**: A substance, especially a coenzyme or a metal, that must be present for an enzyme to function.

In the hepatic portal system, the liver receives a dual blood supply from the hepatic portal vein and hepatic arteries. The hepatic portal vein carries venous blood drained from the spleen, gastrointestinal tract and its associated organs; it supplies approximately 75% of the liver's blood. The hepatic arteries supply arterial blood to the liver and account for the remainder of its blood flow.

Oxygen is provided from both sources; approximately half of the liver's oxygen demand is met by the hepatic portal vein, and half is met by the hepatic arteries. Blood flows through the liver tissue and empties into the central vein of each lobule. The central veins coalesce into hepatic veins that collect the blood leaving the liver and bring it to the heart.




Hepatic veins: An image of a liver with the hepatic veins labeled. They are located in the inferior vena cava.

A portal system is a venous structure that enables blood from one set of capillary beds to drain into another set of capillary beds, without first returning this blood to the heart. The majority of capillaries in the body drain directly into the heart, so portal systems are unusual.

The hepatic portal system connects the capillaries of the gastrointestinal tract with the capillaries in the liver. Nutrient-rich blood leaves the gastrointestinal tract and is first brought to the liver for processing before being sent to the heart. Here, carbohydrates and amino acids can be stored or used to make new proteins and carbohydrates.

The liver also removes vitamins and cofactors from the blood for storage, as well as filters any toxins that may have been absorbed along with the food. When any of these stored substances are needed, the liver releases them back into circulation through the hepatic veins.



Hepatic portal circulation: A diagram that shows the hepatic portal vein and its territory.

This page titled 22.7D: Blood Supply to the Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7E: Liver Function

The liver is thought to be responsible for up to 500 separate functions.

Learning Objectives

- List some of the functions of the liver

Key Points


- The liver is thought to be responsible for up to 500 separate functions, usually in combination with other systems and organs.
- The various functions of the liver are carried out by the liver cells or hepatocytes.
- Excessive alcohol consumption can cause liver disease.
- The liver tissue of an alcoholic may become clogged with fats and adversely affect liver function.

Key Terms

- **liver disease**: Also called hepatic disease, this is an umbrella term referring to damage to or disease of the liver.
- **IGF**: A hormone similar in molecular structure to insulin. It plays an important role in childhood growth and continues to have anabolic effects in adults. A synthetic analog of IGF-1, mecasermin is used for the treatment of growth failure.


Functions of the Liver

The human liver is thought to be responsible for up to 500 separate functions, usually in combination with other systems and organs. The various functions of the liver are carried out by the liver cells or hepatocytes. Currently, there is no artificial organ or device capable of emulating all the functions of the liver.



The liver: The liver, or hepar, is a vital organ present in vertebrates and some other animals. It has a wide range of functions including detoxification, protein synthesis, and the production of the biochemicals necessary for digestion.

The liver is the mainstay of protein metabolism— it synthesizes as well as degrades. It performs several roles in carbohydrate and lipid metabolism. The bulk of the lipoproteins are synthesized in the liver.



Liver tissue of an alcoholic: A healthy liver can break down alcohol. However, the overstressed liver of an alcoholic may become clogged with fats that adversely affect liver function. This type of tissue is most common in alcoholic hepatitis (a prevalence of 65%) and alcoholic cirrhosis (a prevalence of 51%).

In the first- trimester fetus, the liver is the main site of red blood cell production. By the 32nd week of gestation, the bone marrow has almost completely taken over that task.

The liver also produces the insulin -like growth factor 1 (IGF-1), a polypeptide protein hormone that plays an important role in childhood growth and continues to have anabolic effects in adults.

The liver stores a multitude of substances, including glucose (in the form of glycogen), vitamin A (1–2 years' supply), vitamin D (1–4 months' supply), vitamin B12 (1–3 years' supply), iron, and copper. The liver is responsible for immunological effects, acting as a sieve for the antigens that are carried to it via the portal system.

The liver synthesizes angiotensinogen, a hormone that is responsible for raising blood pressure when the angiotensinogen is activated by renin, an enzyme that is released when the kidney senses low blood pressure.

The liver breaks down or modifies toxic substances, such as alcohol and most medicinal products, in a process called drug metabolism. This sometimes results in toxication, when the metabolite is more toxic than its precursor.

Preferably, the toxins are conjugated to avail excretion in bile or urine. The liver breaks down insulin and other hormones.

This page titled 22.7E: Liver Function is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.7F: Bile

Bile is a fluid produced by the liver that aids the process of digestion and the absorption of lipids in the small intestine.

Learning Objectives


- Summarize the composition and function of bile

Key Points

- Bile is a composition of the following materials: water (85%), bile salts (10%), mucus and pigments (3%), fats (1%), inorganic salts (0.7%), and cholesterol (0.3%).
- Bile can either drain directly into the duodenum or be temporarily stored in the gallbladder.
- Bile, which is alkaline, also has the function of neutralizing any excess stomach acid in the small intestine.

Key Terms

- **bile**: A bitter, brownish-yellow or greenish-yellow secretion produced by the liver, stored in the gallbladder, and discharged into the duodenum, where it aids the process of digestion.



Bile: Micrograph of bile (yellow material) in a liver biopsy.

Bile, or gall, is a bitter-tasting, dark-green to yellowish-brown fluid produced by the liver that aids the process of digestion of lipids in the small intestine. Bile is stored in the gallbladder, and upon eating is discharged into the duodenum through the bile duct. Bile is a composition of the following materials: water (85%), bile salts (10%), mucus and pigments (3%), fats (1%), inorganic salts (0.7%), and cholesterol (0.3%).


Bile acts as a surfactant, helping to emulsify the fats in the food, in the same way that soap emulsifies fat. The bile salts are ionically charged, with a hydrophobic end and a hydrophillic end.

When exposed to water mixed with fat, such as in the small intestine, the bile salts congregate around a fat droplet with their hydrophobic side pointing towards the fat and their hydrophillic side pointing towards the water. This increases the surface area of the fat and allows greater access by the pancreatic enzymes that break down fats.

Since bile increases the absorption of fats, it is an important part of the absorption of the fat-soluble vitamins, such as the vitamins D, E, K, and A.

Besides its digestive function, bile serves also as the route of excretion for bilirubin, a waste byproduct of red blood cells that is recycled by the liver. The alkaline bile also has the function of neutralizing any excess stomach acid before it enters the ileum, the final section of the small intestine.

Bile salts also act as bactericides, destroying many of the microbes that may be present in the food.



Bile salt action on lipids: Bile salts congregate around fat and separate them into small droplets called micelles.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23Pancreas.2C_Liver.2C_and_Gallbladder**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Liver**. **License**: *CC BY-SA: Attribution-ShareAlike*
- bile. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/bile**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/gallbladder**. **License**: *CC BY-SA: Attribution-ShareAlike*
- liver. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/liver**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg**. **License**: *Public Domain: No Known Copyright*
- Liver. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Liver%23Anatomy**. **License**: *CC BY-SA: Attribution-ShareAlike*
- lobule. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lobule**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg**. **License**: *Public Domain: No Known Copyright*
- Anatomy of liver and gall bladder. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Anatomy_of_liver_and_gall_bladder.png**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatocytes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hepatocytes**. **License**: *CC BY-SA: Attribution-ShareAlike*

- Regeneration (biology). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Regeneration_(biology)%23Liver. **License**: *CC BY-SA: Attribution-ShareAlike*
- hepatocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hepatocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg. **License**: *Public Domain: No Known Copyright*
- Anatomy of liver and gall bladder. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Anatomy_of_liver_and_gall_bladder.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatocytes. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fc/Human_liver.jpg. **License**: *Public Domain: No Known Copyright*
- Liver. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Liver%23Blood_flow. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatic portal vein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hepatic_portal_vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- cofactors. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cofactors. **License**: *CC BY-SA: Attribution-ShareAlike*
- hepatic arteries. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/hepatic%20arteries. **License**: *CC BY-SA: Attribution-ShareAlike*
- hepatic portal vein. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hepatic_portal_vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg. **License**: *Public Domain: No Known Copyright*
- Anatomy of liver and gall bladder. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Anatomy_of_liver_and_gall_bladder.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatocytes. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fc/Human_liver.jpg. **License**: *Public Domain: No Known Copyright*
- Hepatic veins. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/e/e5/Liver_veins.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatic portal circulation. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/33/Gray591.png. **License**: *Public Domain: No Known Copyright*
- Liver. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Liver%23Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- liver disease. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/liver%20disease. **License**: *CC BY-SA: Attribution-ShareAlike*
- IGF. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/IGF. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg. **License**: *Public Domain: No Known Copyright*
- Anatomy of liver and gall bladder. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Anatomy_of_liver_and_gall_bladder.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatocytes. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fc/Human_liver.jpg. **License**: *Public Domain: No Known Copyright*
- Hepatic veins. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/e/e5/Liver_veins.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatic portal circulation. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/33/Gray591.png. **License**: *Public Domain: No Known Copyright*
- Gray1224. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Gray1224.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/0b/Mallory_body_high_mag_cropped.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bile. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bile. **License**: *CC BY-SA: Attribution-ShareAlike*
- bile. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/bile. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/6f/Human_Hepar.jpg. **License**: *Public Domain: No Known Copyright*
- Anatomy of liver and gall bladder. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Anatomy_of_liver_and_gall_bladder.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Hepatocytes. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fc/Human_liver.jpg. **License**: *Public Domain: No Known Copyright*
- Hepatic veins. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/e/e5/Liver_veins.jpg.

License: *CC BY-SA: Attribution-ShareAlike*
- Hepatic portal circulation. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/33/Gray591.png. **License**: *Public Domain: No Known Copyright*
- Gray1224. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Gray1224.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Liver. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/0b/Mallory_body_high_mag_cropped.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bile. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/31/Cholestasis_2_high_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Lipid and bile salts. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Lipid_and_bile_salts.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 22.7F: Bile is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.8: THE GALLBLADDER

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

22.8A: The Gallbladder

22.8B: Anatomy of the Gallbladder

This page titled 22.8: The Gallbladder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.8A: The Gallbladder

The gallbladder stores bile produced by the liver.

Learning Objectives


- Describe the gallbladder

Key Points

- The gallbladder has a muscular wall that contracts in response to cholecystokinin, a peptide hormone that is synthesized by the small intestine.
- When food containing fat enters the digestive tract, the secretion of cholecystokinin is stimulated and the gall bladder releases the bile into the small intestine.

Key Terms

- **gallbladder**: In vertebrates, a small organ that aids mainly in fat digestion and concentrates bile produced by the liver.



Gallbladder: The gallbladder as depicted in Gray's Anatomy.

The gallbladder is a pear-shaped organ that stores about 50 ml of the bile produced by the liver until the body needs it for digestion. It is about 7–10cm long in humans and is dark green in color.

The gallbladder has a muscular wall that contracts in response to cholecystokinin, a peptide hormone that is synthesized by the small intestine.

Bile and the Gallbladder

When food containing fat enters the digestive tract, the secretion of cholecystokinin (CCK) is stimulated, and the gallbladder releases the bile into the small intestine. The bile emulsifies fats and neutralizes acids in partly digested food. After being stored in the gallbladder, the bile becomes more concentrated to increase its potency and intensify its effect in fats.

This page titled 22.8A: The Gallbladder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.8B: Anatomy of the Gallbladder

The gallbladder, a hollow organ that stores bile, is located under the liver.

Learning Objectives

- Detail the anatomy of the gallbladder

Key Points


- The gallbladder is a hollow organ that sits beneath the liver and stores bile that is made in the liver.
- The gallbladder is divided into three sections: the fundus, body, and neck. The neck connects to a system of ducts.

Key Terms

- **gallstones**: Small, hard objects, in the shape of a pebble, that sometimes form in the gallbladder or bile duct that are composed of cholesterol, bile pigments, and calcium salts.

Anatomy of the Gallbladder

The gallbladder is a hollow organ that sits beneath the liver and stores bile made in the liver. In adults, the gallbladder measures approximately eight centimeters (3.1 in) in length and four centimeters (1.6 in) in diameter when fully distended.



The gallbladder, labeled: An illustration of the gallbladder from Gray's Anatomy with each section labeled.

The gallbladder is divided into three sections:

1. The fundus.
2. The body.
3. The neck.


The neck tapers and connects to the biliary tree via the cystic duct, which then joins the common hepatic duct to become the

common bile duct. At the neck of the gallbladder is a mucosal fold where gallstones commonly get stuck.

Layers of the Gallbladder

There are several different layers of the gallbladder: the mucosa (epithelium and lamina propria), the muscularis, the perimuscular, and the serosa.

- The epithelium is a thin sheet of cells that is closest to the inside of the gallbladder.
- The lamina propria is a thin layer of loose connective tissue, which together with the epithelium, forms the mucosa.
- The muscularis is a layer of smooth muscular tissue that helps the gallbladder contract and squirt its bile into the bile duct.
- The perimuscular (meaning around the muscle) is a fibrous connective tissue layer that surrounds the muscularis.
- The serosa is a smooth membrane that is the outer covering of the gallbladder.



Layers of the gallbladder: A micrograph of the layers of a gallbladder.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- The gastrointestinal system: Liver and gallbladder. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23Pancreas.2C_Liver.2C_and_Gallbladder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gallbladder. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gallbladder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gallbladder.png. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/6c/Gray1095-gall_bladder.png**. **License**: ***Public Domain: No Known Copyright***
- Gallbladder. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gallbladder%23Human_anatomy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gallstone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gallstone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gallbladder.png. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/6/6c/Gray1095-gall_bladder.png**. **License**: ***Public Domain: No Known Copyright***
- Gray1095_parts. **Provided by**: Wikimedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Gray1095_parts.png**.

License: *Public Domain: No Known Copyright*

- Gallbladder - intermed mag. **Provided by**: Wikimedia Commons. **Located at**: **upload.wikimedia.org/Wikipedia/commons/b/bd/Gallbladder_-_intermed_mag.jpg**. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 22.8B: Anatomy of the Gallbladder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.9: THE PANCREAS



- Topic hierarchy

22.9A: Pancreas

22.9B: Anatomy of the Pancreas

22.9C: Histology of the Pancreas

22.9D: Pancreatic Juice

This page titled 22.9: The Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.9A: Pancreas

The pancreas is a gland organ in the digestive and endocrine systems.

Learning Objectives

- Describe the function of the pancreas

Key Points

- The pancreas is a gland organ in the digestive and endocrine systems.
- As an endocrine gland, the pancreas produces several important hormones that include insulin, glucagon, somatostatin, and pancreatic polypeptide.
- As a digestive organ, the pancreas secretes pancreatic juice that contains digestive enzymes that assist the absorption of nutrients and digestion in the small intestine.
- These enzymes help to further break down the carbohydrates, proteins, and lipids in the chyme.

Key Terms

- **pancreas**: A gland near the stomach that secretes a fluid into the duodenum to help with food digestion.

Anatomy of the Pancreas

The pancreas is a gland organ in the digestive and endocrine systems. As an endocrine gland, the pancreas produces several important hormones that include insulin, glucagon, somatostatin, and pancreatic polypeptide.

As a digestive organ, the pancreas secretes pancreatic juice that contains digestive enzymes that assist the absorption of nutrients and digestion in the small intestine. These enzymes help to further break down the carbohydrates, proteins, and lipids in the chyme.

Location

The pancreas is located posterior to the stomach and next to the duodenum. The pancreas functions as both an exocrine and endocrine gland. The exocrine function of the pancreas is essential for digestion as it produces many of the enzymes that break down the protein, carbohydrates, and fats in digestible foods.

Composition

The pancreas is composed of pancreatic exocrine cells, whose ducts are arranged in clusters called acini. The cells are filled with secretory granules containing the inactivated digestive enzymes, mainly trypsinogen, chymotrypsinogen, pancreatic lipase, and amylase, that are secreted into the lumen of the acini.

Glandular Function of the Pancreas

The pancreas is a dual-function gland, having features of both endocrine and exocrine glands.

Exocrine Function

The pancreas synthesizes its enzymes in the inactive form, known as zymogens, to avoid digesting itself. The enzymes are

activated once they reach the small intestine. The pancreas also secretes bicarbonate ions from the ductal cells to neutralize the acidic chyme that the stomach churns out.

The exocrine function of the pancreas is controlled by the hormones gastrin, cholecystokinin, and secretin, which are hormones secreted by cells in the stomach and duodenum in response to food.

The two major proteases that the pancreas synthesizes are trypsinogen and chymotrypsinogen. These zymogens are inactivated forms of trypsin and chymotrypsin.

Once released in the intestine, the enzyme enterokinase, which is produced by the intestinal mucosa, activates trypsinogen by cleaving it to form trypsin. The free trypsin then cleaves the rest of the trypsinogen and chymotrypsinogen to their active forms. Pancreatic secretions accumulate in small ducts that drain to the main pancreatic duct that drains directly into the duodenum.


Endocrine Function

The part of the pancreas with endocrine function is made up of approximately a million cell clusters called the islets of Langerhans. Four main cell types exist in the islets. They are relatively difficult to distinguish using standard staining techniques, but they can be classified by their secretions:

1. α cells secrete glucagon (increase glucose in blood).
2. β cells secrete insulin (decrease glucose in blood).
3. Delta cells secrete somatostatin (regulates/stops α and β cells).
4. PP cells or gamma cells, secrete pancreatic polypeptide.

The Islets of Langerhans

The islets are a compact collection of endocrine cells arranged in clusters and cords that are crisscrossed by a dense network of capillaries. The capillaries of the islets are lined by layers of endocrine cells that are in direct contact with blood vessels, either by cytoplasmic processes or by direct apposition.



Pancreas: This image shows the location of the pancreas relative to other organs. The pancreas is seen positioned with the duodenum slightly on top of it and next to the right kidney. The pancreas crosses above the left kidney.

This page titled 22.9A: Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.9B: Anatomy of the Pancreas

The pancreas lies in the epigastrium or upper central region of the abdomen and can vary in shape.

Learning Objectives

- Outline the anatomy of the pancreas

Key Points

- The pancreas lies in the epigastrium or upper central region of the abdomen.
- The pancreas is composed of a head, uncinate process, neck, body, and tail.
- A number of blood vessels connect the pancreas to the duodenum, spleen, and liver.

Key Terms

- **epigastrium**: The upper middle region of the abdomen, between the umbilical and hypochondriac regions.

Variation

Pancreatic tissue is present in all vertebrate species, but its precise form and arrangement varies widely. There may be up to three separate pancreases, two of which arise from ventral buds, and the other dorsally. In most species (including humans), these fuse in the adult, but there are several exceptions.

Even when a single pancreas is present, two or three pancreatic ducts may persist, each draining separately into the duodenum (or an equivalent part of the foregut). Birds, for example, typically have three such ducts.

In teleosts, and a few other species (such as rabbits), there is no discrete pancreas at all, with pancreatic tissue being distributed diffusely across the mesentery and even within other nearby organs, such as the liver or spleen.


Anatomy of the Pancreas

The pancreas lies in the epigastrium or upper central region of the abdomen. It is composed of several parts.

- The head lies within the concavity of the duodenum.
- The uncinate process emerges from the lower part of head, and lies deep to superior mesenteric vessels.
- The neck is the constricted part between the head and the body.
- The body lies behind the stomach.
- The tail is the left end of the pancreas. It lies in contact with the spleen.

The superior pancreaticoduodenal artery from the gastroduodenal artery and the inferior pancreaticoduodenal artery from the superior mesenteric artery run in the groove between the pancreas and the duodenum and supply the head of pancreas.

The pancreatic branches of the splenic artery also supply the neck, body, and tail of the pancreas. The body and neck of the pancreas drain into the splenic vein; the head drains into the superior mesenteric and portal veins. Lymph is drained via the splenic, celiac, and superior mesenteric lymph nodes.



Parts of a pancreas: 1: Head of pancreas 2: Uncinate process of pancreas 3: Pancreatic notch 4: Body of the pancreas 5: Anterior surface of the pancreas 6: Inferior surface of the pancreas 7: Superior margin of the pancreas 8: Anterior margin of the pancreas 9: Inferior margin of the pancreas 10: Omental tuber 11: Tail of the pancreas 12: Duodenum.

This page titled 22.9B: Anatomy of the Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.9C: Histology of the Pancreas

The pancreas serves digestive and endocrine functions, and it is composed of two types of tissue: islets of Langerhans and acini.

Learning Objectives

- Describe the histology of the pancreas

Key Points

- Under a microscope, the stained sections of the pancreas reveal two different types of parenchymal tissue.
- The light-stained clusters of cells are called islets of Langerhans. These produce hormones that underlie the endocrine functions of the pancreas.
- The dark-stained cells form acini that are connected to ducts. Acinar cells belong to the exocrine pancreas and secrete digestive enzymes into the gut via a system of ducts.

Key Terms

- **islets of Langerhans**: Regions in the pancreas that contain its endocrine cells.
- **acini**: An acinus (adjective: acinar; plural: acini) refers to any cluster of cells that resembles a many-lobed berry, such as a raspberry (acinus is Latin for berry).

The pancreas is a glandular organ in the digestive system and endocrine system of vertebrates. It is both an endocrine gland that produces several important hormones—including insulin, glucagon, somatostatin, and pancreatic polypeptide—as well as a digestive organ that secretes pancreatic juice that contain digestive enzymes to assist the absorption of nutrients and digestion in the small intestine. These enzymes also help to further break down the carbohydrates, proteins, and lipids in the chyme.

Anatomy of the Pancreas

Under a microscope, stained sections of the pancreas reveal two different types of parenchymal tissue. Light-stained clusters of cells are called islets of Langerhans. These produce hormones that underlie the endocrine functions of the pancreas.

The dark-stained cells form acini that are connected to ducts. Acinar cells belong to the exocrine pancreas and secrete digestive enzymes into the gut via a system of ducts.

The pancreas is a dual-function gland that has the features of endocrine and exocrine glands.

The part of the pancreas with endocrine function is made up of approximately a million cell clusters called islets of Langerhans. Four main cell types exist in the islets. They are relatively difficult to distinguish using standard staining techniques, but they can be classified by their secretion

1. α cells secrete glucagon (increase glucose in blood).
2. β cells secrete insulin (decrease glucose in blood).
3. Delta cells secrete somatostatin (regulates/stops α and β cells).
4. PP cells or gamma cells, secrete pancreatic polypeptide.

The islets are a compact collection of endocrine cells arranged in clusters and cords and are crisscrossed by a dense network of capillaries. The capillaries of the islets are lined by layers of endocrine cells in direct contact with vessels, and most endocrine cells are in direct contact with blood vessels, either by cytoplasmic processes or by direct apposition.

This page titled 22.9C: Histology of the Pancreas is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.9D: Pancreatic Juice

Pancreatic fluid contains digestive enzymes that help to further break down the carbohydrates, proteins, and lipids in the chyme.

Learning Objectives

- Describe the nature and function of pancreatic juice

Key Points

- Pancreatic fluid or juice contains digestive enzymes that pass to the small intestine where they help to further break down the carbohydrates, proteins, and lipids (fats) in the chyme.
- Pancreatic fluid is alkaline in nature due to its high concentration of bicarbonate ions that neutralize the gastric acid and allow effective enzymic action.
- Pancreatic juice secretion is regulated by the hormones secretin and cholecystokinin. It is produced by the walls of the duodenum upon detection of acid food, proteins, fats, and vitamins.


Key Terms

- **pancreatic fluid**: A liquid secreted by the pancreas that contains a variety of enzymes, including trypsinogen, chymotrypsinogen, elastase, carboxypeptidase, pancreatic lipase, and amylase.

The pancreas is a glandular organ in the digestive system and endocrine system of vertebrates. It is both an endocrine gland that produces several important hormones—including insulin, glucagon, somatostatin, and pancreatic polypeptide—and a digestive organ that secretes pancreatic juice that has digestive enzymes that assist the absorption of nutrients and digestion in the small intestine. These enzymes help to further break down the carbohydrates, proteins, and lipids in the chyme.

Pancreatic Juice

Pancreatic juice is a liquid secreted by the pancreas that contains a variety of enzymes, including trypsinogen, chymotrypsinogen, elastase, carboxypeptidase, pancreatic lipase, nucleases, and amylase.



Schematic Diagram of
Pancreatic Acini and ducts

Pancreatic fluid: A schematic diagram that shows pancreatic acini and the ducts where fluid is created and released.

Pancreatic juice is alkaline in nature due to its high concentration of bicarbonate ions that neutralize the gastric acid and allow effective enzymic action.

Pancreatic juice secretion is regulated by the hormones secretin and cholecystokinin. It is produced by the walls of the duodenum upon detection of acid food, proteins, fats, and vitamins. Pancreatic secretion consists of an aqueous bicarbonate component from the duct cells and an enzymatic component from the acinar cells.

Because the pancreas is a sort of storage depot for digestive enzymes, injury to the pancreas is potentially fatal. A puncture of the pancreas generally requires prompt and experienced medical intervention.

A variety of factors cause a high pressure within pancreatic ducts. Pancreatic duct rupture and pancreatic juice leakage cause pancreatic self-digestion.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The gastrointestinal system.. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_gastrointestinal_system%23The_Digestive_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- pancreas. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a3/Pancreas.jpg. **License**: *Public Domain: No Known Copyright*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- epigastrium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/epigastrium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a3/Pancreas.jpg. **License**: *Public Domain: No Known Copyright*
- Pancreas. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fe/Illu_pancreas_duodenum.jpg. **License**: *Public Domain: No Known Copyright*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas%23Histology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- acini. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/acini. **License**: *CC BY-SA: Attribution-ShareAlike*
- islets of Langerhans. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/islets_of_Langerhans. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreas. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a3/Pancreas.jpg. **License**: *Public Domain: No Known Copyright*
- Pancreas. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/fe/Illu_pancreas_duodenum.jpg. **License**: *Public Domain: No Known Copyright*
- Pancreas. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreas. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pancreatic juice. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pancreatic_juice. **License**: *CC BY-SA:*

Attribution-ShareAlike
- Pancreas. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pancreas**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pancreatic fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/pancreatic%20fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pancreas. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/a/a3/Pancreas.jpg**. **License**: ***Public Domain: No Known Copyright***
- Pancreas. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fe/Illu_pancreas_duodenum.jpg**. **License**: ***Public Domain: No Known Copyright***
- Pancreatic. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/85/Pancreas_secretions2.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 22.9D: Pancreatic Juice is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.10: THE SMALL INTESTINE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

22.10A: Anatomy of the Small Intestine

22.10B: Histology of the Small Intestine

22.10C: Digestive Processes of the Small Intestine

This page titled 22.10: The Small Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.10A: Anatomy of the Small Intestine

The small intestine is the part of the gastrointestinal tract where much of the digestion and absorption of food takes place.

Learning Objectives

- Diagram the anatomy of the small intestine

Key Points


- The small intestine is the part of the gastrointestinal tract that follows the stomach, which is in turn followed by the large intestine.
- The average length of the small intestine in an adult human male is 6.9 m (22 feet, 6 inches), and in the adult female 7.1 m (23 feet, 4 inches).
- The small intestine is divided into the duodenum, jejunum, and ileum.
- Much of the small intestine is covered in projections called villi that increase the surface area of the tissue available to absorb nutrients from the gut contents.

Key Terms

- **duodenum**: The first part of the small intestine that starts at the lower end of the stomach and extending to the jejunum.
- **ileum**: The last, and usually the longest, division of the small intestine; the part between the jejunum and large intestine.
- **small intestine**: The upper part of the intestine, between the stomach and the large intestine, that is divided into the duodenum, the jejunum, and the ileum.
- **jejunum**: The central of the three divisions of the small intestine that lies between the duodenum and the ileum.

The Small Intestine

The small intestine is the part of the gastrointestinal tract that follows the stomach, which is in turn followed by the large intestine. The small intestine is the site where almost all of the digestion and absorption of nutrients and minerals from food takes place.



Small intestine: An illustration of the small intestine with the duodenum, jejunum, and ileum labeled.

The average length of the small intestine in an adult human male is 6.9 m (22 feet, 6 inches), and in the adult female 7.1 m (23 feet,

4 inches). It can vary greatly, from as short as 4.6 m (15 feet) to as long as 9.8 m (32 feet). The small intestine is approximately 2.5–3 cm in diameter, and is divided into three sections:

1. The duodenum is the first section of the small intestine and is the shortest part of the small intestine. It is where most chemical digestion using enzymes takes place.
2. The jejunum is the middle section of the small intestine. It has a lining which is designed to absorb carbohydrates and proteins. The inner surface of the jejunum, its mucous membrane, is covered in projections called villi, which increase the surface area of tissue available to absorb nutrients from the gut contents. The epithelial cells which line these villi possess even larger numbers of microvilli. The transport of nutrients across epithelial cells through the jejunum includes the passive transport of some carbohydrates and the active transport of amino acids, small peptides, vitamins, and most glucose. The villi in the jejunum are much longer than in the duodenum or ileum.
3. The ileum is the final section of the small intestine. The function of the ileum is mainly to absorb vitamin B12, bile salts, and any products of digestion that were not absorbed by the jejunum. The wall itself is made up of folds, each of which has many tiny finger-like projections known as villi on its surface. The ileum has an extremely large surface area both for the adsorption of enzyme molecules and for the absorption of products of digestion.

The Villi

The villi contain large numbers of capillaries that take the amino acids and glucose produced by digestion to the hepatic portal vein and the liver. Lacteals are the small lymph vessels that are present in villi. They absorb fatty acids and glycerol, the products of fat digestion, into direct circulation.

Layers of circular and longitudinal smooth muscle enable the digested food to be pushed along the ileum by waves of muscle contractions called peristalsis. The undigested food (waste and water) are sent to the colon.

This page titled 22.10A: Anatomy of the Small Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.10B: Histology of the Small Intestine

The small intestine wall has four layers: the outermost serosa, muscularis, submucosa, and innermost mucosa.

Learning Objectives

- Describe the histology of the small intestine


Key Points

- The outermost layer of the intestine, the serosa, is a smooth membrane consisting of a thin layer of cells that secrete serous fluid, and a thin layer of connective tissue.
- The muscularis is a region of muscle adjacent to the submucosa membrane. It is responsible for gut movement (also called peristalsis). It usually has two distinct layers of smooth muscle: circular and longitudinal.
- The submucosa is the layer of dense irregular connective tissue or loose connective tissue that supports the mucosa; it also joins the mucosa to the bulk of underlying smooth muscle.
- The mucosa is the innermost tissue layer of the small intestines and is a mucous membrane that secretes digestive enzymes and hormones. The intestinal villi are part of the mucosa.
- The three sections of the small intestine look similar to each other at a microscopic level, but there are some important differences. The jejunum and ileum do not have Brunner's glands in the submucosa, while the ileum has Peyer's patches in the mucosa, but the duodenum and jejunum do not.

Key Terms

- **Brunner's glands**: Compound, tubular, submucosal glands found in that portion of the duodenum that is above the hepatopancreatic sphincter (sphincter of Oddi).
- **Peyer's patches**: Patches of lymphoid tissue or lymphoid nodules on the walls of the ileum in the small intestine.
- **intestinal wall**: The wall of the small intestine is composed of four layers, from the outside to the inside: serosa, muscularis, submucosa, and mucosa.

The Small Intestine's Layers



Section of duodenum: This image shows the layers of the duodenum: the serosa, muscularis, submucosa, and mucosa.

The small intestine has four tissue layers:

1. The serosa is the outermost layer of the intestine. The serosa is a smooth membrane consisting of a thin layer of cells that secrete serous fluid, and a thin layer of connective tissue. Serous fluid is a lubricating fluid that reduces friction from the movement of the muscularis.
2. The muscularis is a region of muscle adjacent to the submucosa membrane. It is responsible for gut movement, or peristalsis. It usually has two distinct layers of smooth muscle: circular and longitudinal.
3. The submucosa is the layer of dense, irregular connective tissue or loose connective tissue that supports the mucosa, as well as joins the mucosa to the bulk of underlying smooth muscle.
4. The mucosa is the innermost tissue layer of the small intestines, and is a mucous membrane that secretes digestive enzymes and hormones. The intestinal villi are part of the mucosa.

The three sections of the small intestine look similar to each other at a microscopic level, but there are some important differences. The jejunum and ileum do not have Brunner's glands in the submucosa, while the ileum has Peyer's patches in the mucosa, but the duodenum and jejunum do not.

Brunner's Glands


Brunner's glands (or duodenal glands) are compound tubular submucosal glands found in the duodenum. The main function of these glands is to produce a mucus-rich, alkaline secretion (containing bicarbonate) in order to neutralize the acidic content of chyme that is introduced into the duodenum from the stomach, and to provide an alkaline condition for optimal intestinal enzyme activity, thus enabling absorption to take place and lubricate the intestinal walls.

Peyer's Patches

Peyer's patches are organized lymph nodules. They are aggregations of lymphoid tissue that are found in the lowest portion of the small intestine, which differentiate the ileum from the duodenum and jejunum.

Because the lumen of the gastrointestinal tract is exposed to the external environment, much of it is populated with potentially pathogenic microorganisms. Peyer's patches function as the immune surveillance system of the intestinal lumen and facilitate the generation of the immune response within the mucosa.

Intestinal Villi



Micrograph of the small intestine: A low-magnification micrograph of small intestinal mucosa that shows villi.

Intestinal villi (singular: villus) are tiny, finger-like projections that protrude from the epithelial lining of the mucosa. Each villus is approximately 0.5–1.6 mm in length and has many microvilli (singular: microvillus), each of which are much smaller than a single villus.

Villi increase the internal surface area of the intestinal walls. This increased surface area allows for more intestinal wall area to be available for absorption. An increased absorptive area is useful because digested nutrients (including sugars and amino acids) pass into the villi, which is semi-permeable, through diffusion, which is effective only at short distances.

In other words, the increased surface area (in contact with the fluid in the lumen) decreases the average distance traveled by the nutrient molecules, so the effectiveness of diffusion increases.

The villi are connected to blood vessels that carry the nutrients away in the circulating blood.

This page titled 22.10B: Histology of the Small Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.10C: Digestive Processes of the Small Intestine

The small intestine uses different enzymes and processes to digest proteins, lipids, and carbohydrates.

Learning Objectives

- Describe the small intestine's role in the digestive process

Key Points

- The small intestine is where most chemical digestion in the human body takes place.
- Most of the digestive enzymes in the small intestine are secreted by the pancreas and enter the small intestine via the pancreatic duct.
- The three major classes of nutrients that undergo digestion are proteins, lipids (fats), and carbohydrates.

Key Terms

- **digestive enzymes**: Enzymes that break down polymeric macromolecules into their smaller building blocks to facilitate their absorption by the body.

Chemical Digestion in the Small Intestine

The small intestine is where most chemical digestion takes place. Most of the digestive enzymes in the small intestine are secreted by the pancreas and enter the small intestine via the pancreatic duct.

These enzymes enter the small intestine in response to the hormone cholecystokinin, which is produced in response to the presence of nutrients. The hormone secretin also causes bicarbonate to be released into the small intestine from the pancreas to neutralize the potentially harmful acid coming from the stomach.

The three major classes of nutrients that undergo digestion are proteins, lipids (fats), and carbohydrates.

Proteins

Proteins are degraded into small peptides and amino acids before absorption. Their chemical breakdown begins in the stomach and continues through the large intestine.

Proteolytic enzymes, including trypsin and chymotrypsin, are secreted by the pancreas and cleave proteins into smaller peptides. Carboxypeptidase, a pancreatic brush border enzyme, splits one amino acid at a time. Aminopeptidase and dipeptidase free the end amino acid products.

Lipids

Lipids (fats) are degraded into fatty acids and glycerol. Pancreatic lipase breaks down triglycerides into free fatty acids and monoglycerides. Pancreatic lipase works with the help of the salts from bile secreted by the liver and the gallbladder.

Bile salts attach to triglycerides and help to emulsify them; this aids access by pancreatic lipase because the lipase is water-soluble, but the fatty triglycerides are hydrophobic and tend to orient toward each other and away from the watery intestinal surroundings.

The bile salts act to hold the triglycerides in their watery surroundings until the lipase can break them into the smaller components

that are able to enter the villi for absorption.

Carbohydrates

Some carbohydrates are degraded into simple sugars, or monosaccharides (e.g., glucose, galactose) and are absorbed by the small intestine. Pancreatic amylase breaks down some carbohydrates (notably starch) into oligosaccharides. Other carbohydrates pass undigested into the large intestine, where they are digested by intestinal bacteria.

Brush border enzymes take over from there. The most important brush border enzymes are dextrinase and glucoamylase, which further break down oligosaccharides. Other brush border enzymes are maltase, sucrase, and lactase.

Lactase is absent in most adult humans and for them lactose, like most poly-saccharides, is not digested in the small intestine. Some carbohydrates, such as cellulose, are not digested at all, despite being made of multiple glucose units. This is because the cellulose is made out of beta-glucose that makes the inter-monosaccharidal bindings different from the ones present in starch, which consists of alpha-glucose. Humans lack the enzyme for splitting the beta-glucose-bonds—that is reserved for herbivores and bacteria in the large intestine.

Major Digestive Enzymes			
Enzyme	**Produced In**	**Site of Release**	**pH Level**
Carbohydrate Digestion			
Salivary amylase	Salivary glands	Mount	Neutral
Pancreatic amylase	Pancreas	Small intestine	Basic
Maltase	Small intestine	Small intestine	Basic
Protein Digestion			
Pepsin	Gastric glands	Stomach	Acidic
Trypsin	Pancreas	Small intestine	Basic
Peptidases	Small intestine	Small intestine	Basic
Nucleic Acid Digestion			
Nuclease	Pancreas	Small intestine	Basic
Nucleosidases	Pancreas	Small intestine	Basic
Fat Digestion			
Lipase	Pancreas	Small intestine	Basic

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Ileum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ileum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Duodenum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Duodenum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Small intestine. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Small_intestine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Jejunum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Jejunum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- jejunum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/jejunum**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- duodenum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/duodenum. **License**: *CC BY-SA: Attribution-ShareAlike*
- ileum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ileum. **License**: *CC BY-SA: Attribution-ShareAlike*
- small intestine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/small_intestine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._intestine.jpg. **License**: *Public Domain: No Known Copyright*
- Brunner's glands. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Brunner's_glands. **License**: *CC BY-SA: Attribution-ShareAlike*
- Submucosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Submucosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Muscularis externa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Muscularis_externa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Small_i...ne%23Histology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Serosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Serosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Intestinal villi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intestinal_villi. **License**: *CC BY-SA: Attribution-ShareAlike*
- Peyer's patches. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Peyer's_patches. **License**: *CC BY-SA: Attribution-ShareAlike*
- Brunner's glands. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Brunner's%20glands. **License**: *CC BY-SA: Attribution-ShareAlike*
- intestinal wall. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/intestinal%20wall. **License**: *CC BY-SA: Attribution-ShareAlike*
- Peyer's patches. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Peyer's_patches. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/3d/Illu_small_intestine.jpg. **License**: *Public Domain: No Known Copyright*
- Duodenum. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/5/54/Gray1058.png. **License**: *Public Domain: No Known Copyright*
- Small intestinal mucosa. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/38/Small_intestine_low_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Small_intestine. **License**: *CC BY-SA: Attribution-ShareAlike*
- digestive enzymes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/digestive%20enzymes. **License**: *CC BY-SA: Attribution-ShareAlike*
- Small intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/3d/Illu_small_intestine.jpg. **License**: *Public Domain: No Known Copyright*
- Duodenum. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/5/54/Gray1058.png. **License**: *Public Domain: No Known Copyright*
- Small intestinal mucosa. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/38/Small_intestine_low_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Table of Digestive Enzymes. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/6f/Major_digestive_enzymes.png. **License**: *Public Domain: No Known Copyright*

This page titled 22.10C: Digestive Processes of the Small Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11: THE LARGE INTESTINE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.11A: Anatomy of the Large Intestine

22.11B: Histology of the Large Intestine

22.11C: Bacterial Flora

22.11D: Digestive Processes of the Large Intestine

22.11E: Absorption and Feces Formation in the Large Intestine

22.11F: Defecation Reflex

This page titled 22.11: The Large Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11A: Anatomy of the Large Intestine

The large intestine absorbs water from the remaining indigestible food matter and compacts feces prior to defecation.

Learning Objectives

- Describe the function and anatomy of the large intestine

Key Points


- The large intestine starts in the right iliac region of the pelvis, just at or below the right waist, where it is joined to the bottom end of the small intestine. It is about 4.9 feet (1.5 m) long, which is about one-fifth of the whole length of the intestinal canal.
- The appendix is attached to its inferior surface of the cecum. It contains the least lymphoid tissue, and it is a part of mucosa-associated lymphoid tissue that gives it an important role in immunity.
- On the surface of the large intestine, three bands of longitudinal muscle fibers called taeniae coli, each about 0.2 inches wide, can be identified. They start at the base of the appendix and extend from the cecum to the rectum.

Key Terms

- **appendix**: An inner organ without any known use that can become inflamed.
- **cecum**: A pouch, usually peritoneal, that is considered to be the beginning of the large intestine.
- **colon**: The part of the large intestine that is the final segment of the digestive system, after (distal to) the ileum and before (proximal to) the anus.

Function and Form of the Large Intestine

The function of the large intestine (or large bowel) is to absorb water from the remaining indigestible food matter, and then to pass the useless waste material from the body. The large intestine consists of the cecum and colon.



Large intestine: A schematic of the large intestine, with the colon marked as follows: cecum; 1) ascending colon; 2) transverse colon; 3) descending colon; 4) sigmoid colon, rectum, and anus.

It starts in the right iliac region of the pelvis, just at or below the right waist, where it is joined to the bottom end of the small intestine (cecum). From here it continues up the abdomen (ascending colon), then across the width of the abdominal cavity (transverse colon), and then it turns down (descending colon), continuing to its endpoint at the anus (sigmoid colon to rectum to anus). The large intestine is about 4.9 feet (1.5 m) long—about one-fifth of the whole length of the intestinal canal.

Differences Between Large and Small Intestine

The large intestine differs in physical form from the small intestine in several ways. The large intestine is much wider, and the longitudinal layers of the muscularis are reduced to three, strap-like structures known as the taeniae coli.

The wall of the large intestine is lined with simple columnar epithelium. Instead of having the evaginations of the small intestine (villi), the large intestine has invaginations (the intestinal glands).

While both the small intestine and the large intestine have goblet cells, they are more abundant in the large intestine.

Additional Structures

The appendix is attached to its inferior surface of the cecum. It contains the least lymphoid tissue, and it is a part of mucosa-associated lymphoid tissue, which gives it an important role in immunity.

Appendicitis is the result of a blockage that traps infectious material in the lumen. The appendix can be removed with no apparent damage or consequence to the patient.

On the surface of the large intestine, bands of longitudinal muscle fibers called taeniae coli, each about 0.2 inches wide, can be identified. There are three bands, starting at the base of the appendix and extending from the cecum to the rectum.

Along the sides of the taeniae, tags of peritoneum filled with fat, called epiploic appendages (or appendices epiploicae) are found. The sacculations, called haustra, are characteristic features of the large intestine, and distinguish it from the small intestine.

This page titled 22.11A: Anatomy of the Large Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11B: Histology of the Large Intestine

The large intestine has taeniae coli and invaginations (the intestinal glands), unlike the small intestines.

Learning Objectives

- Describe the histology of the large intestine

Key Points

- The longitudinal layer of the muscularis of the large intestine is reduced to three, strap-like structures known as the taeniae coli—bands of longitudinal muscle fibers, each about 1/5 in wide.
- The bands of longitudinal muscle fibers start at the base of the appendix and extend from the cecum to the rectum.
- The wall of the large intestine is lined with simple columnar epithelium.
- Both the small intestine and the large intestine have goblet cells, but they are abundant in the large intestine.

Key Terms

- **goblet cell**:
- **columnar epithelium**: Epithelial cells whose heights are at least four times their width.
- **mucin**: A family of high molecular weight, heavily glycosylated proteins (glycoconjugates) produced by the epithelial tissues in most metazoans.
- **goblet cells**: Glandular, simple, columnar epithelial cells whose sole function is to secrete mucin, which dissolves in water to form mucus.

Histology of the Large Intestine



Colon biopsy: Micrograph of a colon biopsy.


The large intestine, or large bowel, is the last part of the digestive system in vertebrate animals. Its function is to absorb water from the remaining indigestible food matter, and then to pass the useless waste material from the body. The large intestine consists of the cecum, colon, rectum, and anal canal.

It starts in the right iliac region of the pelvis, just at or below the right waist, where it is joined to the bottom end of the small intestine. From here it continues up the abdomen, across the width of the abdominal cavity, and then it turns downward, continuing to its endpoint at the anus.

The large intestine differs in physical form from the small intestine in being much wider. The longitudinal layer of the muscularis is reduced to three strap-like structures known as the taeniae coli—bands of longitudinal muscle fibers, each about 1/5 in wide. These three bands start at the base of the appendix and extend from the cecum to the rectum.

Along the sides of the taeniae are tags of peritoneum filled with fat; these are called epiploic appendages, or appendices epiploicae. The wall of the large intestine is lined with simple columnar epithelium.

Instead of having the evaginations of the small intestine (villi), the large intestine has invaginations (the intestinal glands). While both the small intestine and the large intestine have goblet cells that secrete mucin to form mucus in water, they are abundant in the large intestine.



Sigmoid colon: A photograph of the large bowel (sigmoid colon) that shows multiple diverticula on either side of the longitudinal muscle bundle (Taenia coli).

In histology, an intestinal crypt—called the crypt of Lieberkühn—is a gland found in the epithelial lining of the small intestine and colon. The crypts and intestinal villi are covered by epithelium that contains two types of cells: goblet cells that secrete mucus and enterocytes that secrete water and electrolytes.

The enterocytes in the mucosa contain digestive enzymes that digest specific food while they are being absorbed through the epithelium. These enzymes include peptidases, sucrase, maltase, lactase and intestinal lipase. This is in contrast to the stomach, where the chief cells secrete pepsinogen. In the intestine, the digestive enzymes are not secreted by the cells of the intestine.

Also, new epithelium is formed here, which is important because the cells at this site are continuously worn away by the passing food. The basal portion of the crypt, further from the intestinal lumen, contains multipotent stem cells.

During each mitosis, one of the two daughter cells remains in the crypt as a stem cell, while the other differentiates and migrates up the side of the crypt and eventually into the villus. Goblet cells are among the cells produced in this fashion. Many genes have been shown to be important for the differentiation of intestinal stem cells.

The loss of proliferation control in the crypts is thought to lead to colorectal cancer.

This page titled 22.11B: Histology of the Large Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11C: Bacterial Flora

The largest bacteria ecosystem in the human body is in the large intestine, where it plays a variety of important roles.

Learning Objectives

- Describe other factors about gut/bacterial flora

Key Points

- The large intestine absorbs some of the products formed by the bacteria that inhabit this region, such as short-chain fatty acids that are metabolized from undigested polysaccharides (fiber).
- Other bacterial products of undigested polysaccharide fermentation include gas (flatus), which consists primarily of nitrogen and carbon dioxide.
- These bacteria also produce large amounts of vitamins, especially vitamin K and biotin (a B vitamin), for absorption into the blood.
- Factors that disrupt the microorganism population of the large intestine include antibiotics, stress, and parasites.

Key Terms


- **passive diffusion**: The net movement of material from an area of high concentration to an area of low concentration without any energy input.
- **bacterial flora**: A community of bacteria that exists on or in the body, and possesses a unique ecological relationship with the host.
- **colitis**: An inflammation of the colon or the large intestine.

Bacterial Flora

The large intestine houses over 700 species of bacteria that perform a wide variety of functions; it is the largest bacterial ecosystem in the human body. The large intestine absorbs some of the products formed by the bacteria that inhabit this region.

For example, undigested polysaccharides (fiber) are metabolized to short-chain fatty acids by the bacteria in the large intestine, and then are absorbed by passive diffusion. The bicarbonate that the large intestine secretes helps to neutralize the increased acidity that results from the formation of these fatty acids.

Bacteria and Vitamins



Bacterial flora: Escherichia coli is one of the many species of bacteria present in the human gut.

These bacteria also produce large amounts of vitamins, especially vitamin K and biotin (a B vitamin), for absorption into the blood. Although this source of vitamins, in general, provides only a small part of the daily requirement, it makes a significant contribution when dietary vitamin intake is low.

An individual who depends just on the absorption of vitamins formed by bacteria in the large intestine may become vitamin deficient if treated with antibiotics that inhibit other species of bacteria, as well as the disease-causing bacteria.

Other bacterial products include gas (flatus), which is a mixture of nitrogen and carbon dioxide, with small amounts of hydrogen, methane, and hydrogen sulphide. These are produced as a result of the bacterial fermentation of undigested polysaccharides. The normal flora is also essential for the development of certain tissues, including the cecum and lymphatics.

Bacteria and Antibodies

Bacterial flora is also involved in the production of cross-reactive antibodies. These are antibodies produced by the immune system against the normal flora, that are also effective against related pathogens, and prevent infection or invasion.

The most prevalent bacteria are the bacteroides, which have been implicated in the initiation of colitis and colon cancer. Bifidobacteria are also abundant, and are often described as friendly bacteria.

A mucus layer protects the large intestine from attacks from colonic commensal bacteria. Some factors that disrupt the microorganism population of the large intestine include antibiotics, stress, and parasites.

This page titled 22.11C: Bacterial Flora is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11D: Digestive Processes of the Large Intestine

In the large intestine, a host of microorganisms known as gut flora help digest the remaining food matter and create vitamins.

Learning Objectives

- Summarize the digestive processes of the gut flora of the large intestine


Key Points

- The large intestine takes about 16 hours to finish the remaining processes of the digestive system.
- The colon absorbs vitamins created by the colonic bacteria. Gut flora consists of microorganisms that live in the digestive tracts of animals; the digestive tract is the largest reservoir of human flora.
- The colon compacts feces and stores fecal matter in the rectum until it can be defecated.
- The gut flora performs many useful functions, such as fermenting unused energy substrates, training the immune system, preventing the growth of pathogenic bacteria, regulating the development of the gut, producing vitamins for the host, and producing hormones to direct the host to store fats.

Key Terms

- **saccharolytic**: The breakdown of carbohydrates for energy.
- **gut flora**: The microorganisms that normally live in the digestive tract of animals.
- **vitamin**: Any of a specific group of organic compounds essential in small quantities for healthy human growth, metabolism, development, and body function that are found in minute amounts in plant and animal foods or sometimes produced synthetically; deficiencies of specific vitamins produce specific disorders.

Overview of the Large Intestine



Digestive processes in large intestine: This image shows the relationship of the colon to the other parts of the digestive system.

The large intestine takes about 16 hours to finish up the remaining processes of the digestive system. Food is no longer broken

down at this stage of digestion. The colon absorbs vitamins created by the colonic bacteria—such as vitamin K (especially important as the daily ingestion of vitamin K is not normally enough to maintain adequate blood coagulation), vitamin B12, thiamine, and riboflavin. It also compacts feces, and stores fecal matter in the rectum until it can be defecated.

Gut Flora

Gut flora consists of microorganisms that live in the digestive tracts of animals—the gut is the largest reservoir of human flora. The human body, which consists of about 10 trillion cells, carries about ten times as many microorganisms in the intestines.


The metabolic activities performed by these bacteria resemble those of an organ, leading some to liken gut bacteria to a forgotten organ. It is estimated that these gut flora have around a hundred times as many genes in aggregate as there are in the human genome.

Bacteria make up most of the flora in the colon and up to 60 percent of the dry mass of feces. Somewhere between 300 and 1000 different species live in the gut, with most estimates at about 500. Ninety-nine percent of the bacteria probably come from about 30 or 40 species.

Research suggests that the relationship between gut flora and humans is not merely commensal (a non-harmful coexistence), but rather a mutualistic relationship. Though people can survive without gut flora, the microorganisms perform a host of useful functions, such as:

- Fermenting unused energy substrates.
- Training the immune system.
- Preventing growth of harmful, pathogenic bacteria.
- Regulating the development of the gut.
- Producing vitamins for the host (such as biotin and vitamin K).
- Producing hormones to direct the host to store fats.

Gut Flora and Specialized Digestion



Bacterial flora: Escherichia coli, one of the many species of bacteria present in the human gut.

Without gut flora, the human body would be unable to utilize some of the undigested carbohydrates it consumes. Some types of gut flora have enzymes that human cells lack for breaking down certain polysaccharides. Carbohydrates that need bacterial assistance for digestion include:

- Certain starches.

- Fiber.
- Oligosaccharides and sugars like lactose (in the case of lactose intolerance) and sugar alcohols.
- Mucus produced by the gut.
- Various proteins.

Fermentation

Bacteria turn the carbohydrates they ferment into short-chain fatty acids (SCFAs) by a form of fermentation called saccharolytic fermentation. These SCFAs include acetic acid, propionic acid, and butyric acid.

SCFAs can be used by host cells as a major source of useful energy and nutrients for humans. They also help the body absorb essential dietary minerals such as calcium, magnesium, and iron. Evidence indicates that bacteria enhance the absorption and storage of lipids and produce and aid the absorption of needed vitamins, such as vitamin K.

This page titled 22.11D: Digestive Processes of the Large Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11E: Absorption and Feces Formation in the Large Intestine

The large intestine absorbs water from the chyme and stores feces until they can be defecated.

Learning Objectives

- Describe the process of absorption and feces formation in the large intestine

Key Points

- Partially digested food passes from the small intestine to the large intestine or colon.
- Within the colon, digestion is retained long enough to allow fermentation via gut bacteria that break down some of the substances that remain after processing in the small intestine.
- The large intestine houses over 700 species of bacteria that metabolize polysaccharides into short-chain fatty acids that produce large amounts of vitamins —especially vitamin K and biotin—and gas.
- The normal flora of bacteria in the large intestine is essential in the development of certain tissues, including the cecum and lymphatics.

Key Terms

- **feces**: Digested waste material that is discharged from the bowels; excrement.
- **polysaccharide**: A polymer made of many saccharide units that are linked by glycosidic bonds.
- **anal sphincter**: A ring muscle that surrounds the anus (anal orifice).

After the food has been passed through the small intestine, it enters the large intestine. Within the large intestine, digestion is retained long enough to allow fermentation via gut bacteria that break down some of the substances that remain after processing in the small intestine.

Some of the breakdown products are absorbed. In humans, these include most complex saccharides (at most, three disaccharides are digestible by humans).

Intestinal Bacteria

The large intestine houses over 700 species of bacteria that perform a variety of functions. The large intestine absorbs some of the products formed by the bacteria that inhabit this region.

Undigested polysaccharides (fiber) are metabolized into short-chain fatty acids by bacteria in the large intestine and get absorbed by passive diffusion. The bicarbonate that the large intestine secretes helps to neutralize the increased acidity from the formation of fatty acids.

Intestinal bacteria also produce large amounts of vitamins, especially vitamin K and biotin (a B vitamin), which are absorbed into the blood. Although this source of vitamins provides only a small part of the daily requirement, it makes a significant contribution when dietary vitamin intake is low. An individual that depends on absorption of vitamins formed by bacteria in the large intestine may become vitamin-deficient if treated with antibiotics that inhibit other species of bacteria while targeting the disease-causing bacteria.

Other bacterial products include gas (flatus)—a mixture of nitrogen and carbon dioxide, with small amounts of the gases hydrogen, methane, and hydrogen sulphide. The bacterial fermentation of undigested polysaccharides produces these gases.

Intestinal flora are also essential for the development of certain tissues, including the cecum and lymphatics.

Water and Cellulose

The large intestine absorbs water from the chyme and stores feces until it can be defecated. Food products that cannot go through the villi, such as cellulose (dietary fiber), are mixed with other waste products from the body and become hard and concentrated feces.

The feces is stored in the rectum for a certain period and then the stored feces is eliminated from the body due to the contraction and relaxation of the anus. The exit of this waste material is regulated by the anal sphincter.

This page titled 22.11E: Absorption and Feces Formation in the Large Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.11F: Defecation Reflex

Defecation is a combination of voluntary and involuntary processes that create enough force to remove waste material from the digestive system.

Learning Objectives

- Describe the defecation reflex

Key Points

- The rectal ampulla acts as a temporary storage facility for the unneeded digestive material.
- A sufficient increase in fecal material in the rectum causes stretch receptors from the nervous system that are located in the rectal walls to trigger the contraction of rectal muscles, the relaxation of the internal anal sphincter, and an initial contraction of the skeletal muscle of the external sphincter.
- The relaxation of the internal anal sphincter causes a signal to be sent to the brain indicating an urge to defecate.
- If defecation is delayed for a prolonged period, the fecal matter may harden and autolyze, and result in constipation.
- Once the voluntary signal to defecate is sent back from the brain, the ano-rectal angle decreases, becoming almost straight, and the external anal sphincter relaxes. The rectum contracts and shortens in peristaltic waves, forcing fecal material out of the rectum and down through the anal canal.

Key Terms

- **defecation**: The act or process of voiding feces from the bowels.
- **rectum**: The terminal part of the large intestine through which feces pass.
- **anal canal**: The terminal part of the large intestine, situated between the rectum and anus.
- **constipation**: A state of the bowels in which the evacuations are infrequent and difficult, or the intestines become filled with hardened feces.
- **autolyze**: To destroy itself; to be destroyed by its own enzymes.


EXAMPLES

Constipation is uncomfortable, but it may be a signal that your diet is deficient in fibrous foods (eat more fruits and vegetables) and that you are not drinking enough water, so laxatives are generally not the best way to treat it.

Defecation

For the adult human, the process of defecation is normally a combination of both voluntary and involuntary processes that create enough force to remove waste material from the digestive system.

The rectal ampulla acts as a temporary storage facility for the unneeded material. As additional fecal material enters the rectum, the rectal walls expand. A sufficient increase in fecal material in the rectum causes the stretch receptors from the nervous system, located in the rectal walls, to trigger the contraction of rectal muscles, the relaxation of the internal anal sphincter, and an initial contraction of the skeletal muscle of the external sphincter. The relaxation of the internal anal sphincter causes a signal to be sent to the brain indicating an urge to defecate.



Defecation reflex: The conscious and parasympathetic pathways of the defecation reflex.

If this urge is not acted upon, the material in the rectum is often returned to the colon by reverse peristalsis where more water is absorbed, thus temporarily reducing pressure and stretching within the rectum. The additional fecal material is stored in the colon until the next mass peristaltic movement of the transverse and descending colon. If defecation is delayed for a prolonged period, the fecal matter may harden and autolyze, resulting in constipation.

Once the voluntary signal to defecate is sent back from the brain, the final phase begins. The abdominal muscles contract (straining), causing the intra-abdominal pressure to increase. The perineal wall is lowered and causes the anorectal angle to decrease from 90 degrees to less than 15 degrees (almost straight), and the external anal sphincter relaxes.

The rectum now contracts and shortens in peristaltic waves, thus forcing fecal material out of the rectum and down through the anal canal. The internal and external anal sphincters, along with the puborectalis muscle, allow the feces to be passed by pulling the anus up and over the exiting feces in shortening and contracting actions.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Large Intestine. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Large_Intestine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large intestine. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Large_intestine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- appendix. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/appendix**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- colon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/colon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/cecum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...lon_scheme.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- columnar epithelium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/columnar%20epithelium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intestinal gland. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intestinal_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Large intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Large_intestine. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Large_I...o_other_organs. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mucin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/mucin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- goblet cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/goblet%20cell. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...n/goblet-cells. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...lon_scheme.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Bowel. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...moid_colon.jpg. **License**: ***Public Domain: No Known Copyright***
- Colon biopsy. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...i_high_mag.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Large_I...acterial_flora. **License**: ***CC BY-SA: Attribution-ShareAlike***
- passive diffusion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/passive%20diffusion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bacterial flora. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/bacterial%20flora. **License**: ***CC BY-SA: Attribution-ShareAlike***
- colitis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/colitis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...lon_scheme.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Bowel. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...moid_colon.jpg. **License**: ***Public Domain: No Known Copyright***
- Colon biopsy. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...i_high_mag.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Coli_NIAID.jpg. **License**: ***Public Domain: No Known Copyright***
- Gut flora. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gut_flora. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Large_I...o_other_organs. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gut flora. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/gut+flora. **License**: ***CC BY-SA: Attribution-ShareAlike***
- saccharolytic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/saccharolytic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vitamin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vitamin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...lon_scheme.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large Bowel. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...moid_colon.jpg. **License**: ***Public Domain: No Known Copyright***
- Colon biopsy. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...i_high_mag.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Coli_NIAID.jpg. **License**: ***Public Domain: No Known Copyright***
- Colon. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ne-diagram.svg. **License**: ***Public Domain: No Known Copyright***
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Coli_NIAID.jpg. **License**: ***Public Domain: No Known Copyright***
- Digestion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Digesti...arge_intestine. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Large intestine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Large_intestine. **License**: ***CC BY-SA: Attribution-ShareAlike***
- polysaccharide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/polysaccharide. **License**: ***CC BY-SA: Attribution-ShareAlike***

- feces. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/feces. **License**: *CC BY-SA: Attribution-ShareAlike*
- anal sphincter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/anal_sphincter. **License**: *CC BY-SA: Attribution-ShareAlike*
- Large Intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...lon_scheme.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Large Bowel. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...moid_colon.jpg. **License**: *Public Domain: No Known Copyright*
- Colon biopsy. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...i_high_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Coli_NIAID.jpg. **License**: *Public Domain: No Known Copyright*
- Colon. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ne-diagram.svg. **License**: *Public Domain: No Known Copyright*
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Coli_NIAID.jpg. **License**: *Public Domain: No Known Copyright*
- Defecation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Defecation. **License**: *CC BY-SA: Attribution-ShareAlike*
- defecation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/defecation. **License**: *CC BY-SA: Attribution-ShareAlike*
- autolyze. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/autolyze. **License**: *CC BY-SA: Attribution-ShareAlike*
- rectum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/rectum. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...n/constipation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anal canal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anal_canal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Large Intestine. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/6/62/Colon_scheme.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Large Bowel. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/8/83/Diverticula,_sigmoid_colon.jpg. **License**: *Public Domain: No Known Copyright*
- Colon biopsy. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/e/ef/Melanosis_coli_high_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/32/EscherichiaColi_NIAID.jpg. **License**: *Public Domain: No Known Copyright*
- Colon. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/1/11/Intestine-diagram.svg. **License**: *Public Domain: No Known Copyright*
- E. Coli. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/32/EscherichiaColi_NIAID.jpg. **License**: *Public Domain: No Known Copyright*
- Defecation reflex. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ion_reflex.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 22.11F: Defecation Reflex is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.12: CHEMICAL DIGESTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.12A: Mechanisms of Chemical Digestion

22.12B: Chemical Digestion of Carbohydrates, Proteins, Lipids, and Nucleic Acids

This page titled 22.12: Chemical Digestion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.12A: Mechanisms of Chemical Digestion

Chemical digestion is the enzyme-mediated, hydrolysis process that breaks down large macronutrients into smaller molecules.

Learning Objectives

- Differentiate among the methods used to chemically break down food molecules

Key Points

- Carbohydrates are mainly taken in the form of amylose and glycogen. Amylases hydrolyze the long carbohydrate chains that break amylose down into disaccharides, and glycogen into polysaccharides. The enzymes in the small intestine then break these down to monosaccharides.
- Proteins are digested by hydrolysis of the carbon–nitrogen (C–N) bond. Peptidases are secreted in an inactive form, to prevent auto-digestion. Endopeptidases cleave the polypeptides at the interior peptide bonds, and the exopeptidases cleave the terminal amino acids.
- Fats are digested by lipases that hydrolyze the glycerol fatty acid bonds. Bile salts emulsify the fats to allow for their solution as micelles in the chyme and to increase the surface area for the pancreatic lipases to operate.
- RNA and DNA are hydrolized by the pancreatic enzymes (ribonucleases, deoxyribonucleases) into nucleic acids, which are further broken down to purine and pyrimidine bases and pentoses, by enzymes in the intestinal mucosa (nucleases).
- Once the voluntary signal to defecate is sent back from the brain, the final phase begins.


Key Terms

- **peptidase**: Any enzyme that catalyzes the hydrolysis of peptides into amino acids; a protease.
- **amylase**: Any of a class of digestive enzymes that are present in saliva and that break down complex carbohydrates, such as starch, into simple sugars, such as glucose.
- **hydrolysis**: The degradation of certain biopolymers (proteins, complex sugars) by the chemical process that results in smaller polymers or monomers, such as amino acids or monosaccharides.

EXAMPLES

Because amylase turns some potato or rice starch into sugar, these foods taste slightly sweet.

Digestion is a form of catabolism: a breakdown of large food molecules (i.e., polysaccharides, proteins, fats, nucleic acids) into smaller ones (i.e., monosaccharides, amino acids, fatty acids, nucleotides).




Catabolism: A simplified outline of the catabolism of proteins, carbohydrates, and fats.

Carbohydrates are taken in mainly in the form of plant carbohydrate (amylose) and animal carbohydrate (glycogen) together with some sugars, mainly disaccharides. About 80% of the western diet is in the form of amylose. Amylose is not highly branched and consists mainly of long chains of glucose linked by α1:4 linkages.

Cellulose, the most abundant starch in nature, is formed of β1:4 linkages and cannot be digested in humans, although the bacterial action in the colon does break down a minute amount of it.

Glycogen is a multi-branched starch with linkages at the 1:4 and 1:6 position. This creates very large granules of multi-branched starch. Both the parotid and pancreatic amylases hydrolyse the 1:4 link, but not the terminal 1:4 links or the 1:6 links. This breaks amylose down into mainly disaccharides, and glycogen with its 1:6 linkages into polysaccharides.

The net result of these actions are numerous disaccharides and polysaccharides. Enzymes attached to the enterocycytes of the small intestine break these down to monosaccharides.



Hydrolysis of 1:4α Linkage


Hydrolysis by amylase: Both the parotid and pancreatic amylases hydrolyse the 1:4 link, but not the terminal 1:4 links or the 1:6 links.

Proteins and polypeptides are digested by hydrolysis of the carbon–nitrogen (C–N) bond. The proteolytic enzymes are all secreted in an inactive form, to prevent auto-digestion, and are activated in the lumen of the gut. Activation is caused by HCl in the case of the stomach enzyme pepsinogen, and by enteropeptidase and trypsin in the case of the pancreatic enzymes.

Final digestion takes place by small intestine enzymes that are embedded in the brush border of the small intestine. The enzymes are divided into endo- and exo-peptidases.

- The endopeptidases cleave the polypeptide at the interior peptide bonds, while the exopeptidases cleave the terminal amino acid.
- Exopeptidases are further subclassified into aminopeptidases—that cleave off the terminal amino acid at the amine end of the chain—and carboxypeptidases that cleave off the terminal amino acid at the carboxyl end of the chain.

Stomach pepsin cleaves the interior bonds of the amino acids, and is particularly important for its ability to digest collagen. This is a major constituent of the connective tissue of meat. In the absence of stomach pepsin, digestion in the small intestine proceeds with difficulty. Stomach pepsin digests about 20% of the proteins, and the rest are digested by the pancreatic and small intestine enzymes.



Hydrolysis of Peptide Bond

Hydrolysis of peptide bond: Proteins and polypeptides are digested by hydrolysis of the C–N bond.

Fats are digested by lipases that hydrolyze the glycerol fatty acid bonds. Of particular importance in fat digestion and absorption are the bile salts, which emulsify the fats to allow for their solution as micelles in the chyme, and increase the surface area for the pancreatic lipases to operate.

Lipases are found in the mouth, the stomach, and the pancreas. Because the lingual lipase is inactivated by stomach acid, it is formally believed to be mainly present for oral hygiene and for its anti-bacterial effect in the mouth. However, it can continue to operate on food stored in the fundus of the stomach, and as much as 30% of the fats can be digested by this lipase.

Gastric lipase is of little importance in humans. Pancreatic lipase accounts for the majority of fat digestion and operates in conjunction with the bile salts.

RNA and DNA are hydrolized by the pancreatic enzymes (ribonucleases, deoxyribonucleases) into nucleic acids, which are further broken down to purine and pyrimidine bases and pentoses, by enzymes in the intestinal mucosa (nucleases).

This page titled 22.12A: Mechanisms of Chemical Digestion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.12B: Chemical Digestion of Carbohydrates, Proteins, Lipids, and Nucleic Acids

The chemical breakdown of the macromolecules contained in food is completed by various enzymes produced in the digestive system.

Learning Objectives

- Summarize chemical digestion

Key Points

- Protein digestion occurs in the stomach and the duodenum through the action of three main enzymes: pepsin, secreted by the stomach, and trypsin and chymotrypsin, secreted by the pancreas.
- During carbohydrate digestion the bonds between glucose molecules are broken by salivary and pancreatic amylase.
- The digestion of certain fats begins in the mouth, where short-chain lipids break down into diglycerides because of lingual lipase. The fat present in the small intestine stimulates the release of lipase from the pancreas, and bile from the liver enables the breakdown of fats into fatty acids.
- DNA and RNA are broken down into mononucleotides by the nucleases deoxyribonuclease and ribonuclease (DNase and RNase) that are released by the pancreas.

Key Terms

- **zymogen**: A proenzyme, or enzyme precursor, that requires a biochemical change (i.e., hydrolysis) to become an active form of the enzyme.
- **lactose intolerance**: The inability to fully metabolize lactose.

Chemical Digestion

Protein digestion occurs in the stomach and the duodenum through the action of three primary enzymes:

1. Pepsin, secreted by the stomach.
2. Trypsin, secreted by the pancreas.
3. Chymotrypsin, secreted by the pancreas.

These enzymes break down food proteins into polypeptides that are then broken down by various exopeptidases and dipeptidases into amino acids. The digestive enzymes, however, are secreted mainly as their inactive precursors, the zymogens.

Thus, trypsin is secreted by the pancreas in the form of trypsinogen, which is activated in the duodenum by enterokinase to form trypsin. Trypsin then cleaves proteins into smaller polypeptides.

In humans, dietary starches are composed of glucose units arranged in long chains of polysaccharide called amylose. During digestion, the bonds between glucose molecules are broken by salivary and pancreatic amylase, and result in progressively smaller chains of glucose. This process produces the simple sugars glucose and maltose (two glucose molecules) that can be absorbed by the small intestine.


Sucrase is an enzyme that breaks down disaccharide sucrose, commonly known as table sugar, cane sugar, or beet sugar. Sucrose digestion yields the sugars fructose and glucose, which are readily absorbed by the small intestine.

Lactase is an enzyme that breaks down the disaccharide lactose into its component parts, glucose and galactose, that are absorbed by the small intestine. Approximately half the adult population produces only small amounts of lactase and are therefore

unable to eat milk-based foods. This condition is commonly known as lactose intolerance.


The digestion of certain fats begins in the mouth, where lingual lipase breaks down short chain lipids into diglycerides. The presence of fat in the small intestine produces hormones that stimulate the release of pancreatic lipase from the pancreas, and bile from the liver, to enable the breakdown of fats into fatty acids. The complete digestion of one molecule of fat (a triglyceride) results in three fatty acid molecules and one glycerol molecule.

DNA and RNA are broken down into mononucleotides by the nucleases deoxyribonuclease and ribonuclease (DNase and RNase), which are released by the pancreas.



Action of Small Intestine enzymes on various sugar substrates

Carbohydrate digestion: A diagram of the action of the oligosaccharide-cleaving enzymes in the small intestine.



Lipid digestion: Lipid digestion involves the formation of micelles in the presence of bile salts, and the passage of micelles and fatty acids through the unstirred layer. The diagram depicts dietary fat at the top, with pancreatic lipase and bile salts forming micelles that will pass through the unstirred layer at the bottom of the diagram.

This page titled 22.12B: Chemical Digestion of Carbohydrates, Proteins, Lipids, and Nucleic Acids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.13: ABSORPTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.13A: Absorption in the Small Intestine

22.13B: Absorption of Monosaccharides, Amino Acids, Dipeptides, Tripeptides, Lipids, Electrolytes, Vitamins, and Water

This page titled 22.13: Absorption is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.13A: Absorption in the Small Intestine

The absorption of nutrients occurs partially by diffusion through the wall of the small intestine.

Learning Objectives

- Describe the role played by the small intestine in the absorption of nutrients

Key Points

- Digested food is able to pass into the blood vessels in the wall of the small intestine through the process of diffusion.
- The inner wall, or mucosa, of the small intestine is covered in wrinkles or folds called plicae circulares that project microscopic finger-like pieces of tissue called villi, which in turn have finger-like projections known as microvilli.
- The function of the plicae circulares, the villi, and the microvilli is to increase the amount of surface area available for the absorption of nutrients.
- Each villus transports nutrients to a network of capillaries and fine lymphatic vessels called lacteals close to its surface.

Key Terms


- **villi**: Tiny, finger-like projections that protrude from the epithelial lining of the intestinal wall.
- **plicae circulares**: These circular folds (known as the valves of Kerckring or the valvulae conniventes) are large, valvular flaps that project into the lumen of the bowel.
- **diffusion**: The act of diffusing or dispersing something, or the property of being diffused or dispersed; dispersion.

EXAMPLES

Examples of nutrients absorbed by the small intestine include carbohydrates, lipids, proteins, iron, vitamins, and water.

The Small Intestine

The small intestine is the part of the gastrointestinal tract between the stomach and the large intestine where much of the digestion of food takes place. The primary function of the small intestine is the absorption of nutrients and minerals found in food.



Intestinal villus: An image of a simplified structure of the villus. The thin surface layer appear above the capillaries that are connected to a blood vessel. The lacteal is surrounded by the capillaries.

Digested nutrients pass into the blood vessels in the wall of the intestine through a process of diffusion. The inner wall, or mucosa, of the small intestine is lined with simple columnar epithelial tissue.

Structurally, the mucosa is covered in wrinkles or folds called plicae circulares—these are permanent features in the wall of the organ. They are distinct from the rugae, which are non-permanent features that allow for distention and contraction.


From the plicae circulares project microscopic finger-like pieces of tissue called villi (Latin for shaggy hair). The individual epithelial cells also have finger-like projections known as microvilli. The function of the plicae circulares, the villi, and the microvilli is to increase the amount of surface area available for the absorption of nutrients.

Each villus has a network of capillaries and fine lymphatic vessels called lacteals close to its surface. The epithelial cells of the villi transport nutrients from the lumen of the intestine into these capillaries (amino acids and carbohydrates) and lacteals (lipids).

The absorbed substances are transported via the blood vessels to different organs of the body where they are used to build complex substances, such as the proteins required by our body. The food that remains undigested and unabsorbed passes into the large intestine.

Absorption of the majority of nutrients takes place in the jejunum, with the following notable exceptions:

- Iron is absorbed in the duodenum.
- Vitamin B12 and bile salts are absorbed in the terminal ileum.
- Water and lipids are absorbed by passive diffusion throughout the small intestine.
- Sodium bicarbonate is absorbed by active transport and glucose and amino acid co-transport.
- Fructose is absorbed by facilitated diffusion.



Section of duodenum: Section of duodenum with villi at the top layer.

This page titled 22.13A: Absorption in the Small Intestine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.13B: Absorption of Monosaccharides, Amino Acids, Dipeptides, Tripeptides, Lipids, Electrolytes, Vitamins, and Water

Glucose, amino acids, fats, and vitamins are absorbed in the small intestine via the action of hormones and electrolytes.

Learning Objectives

- Describe the process of absorption of nutrients in the small intestine

Key Points

- Proteins are degraded into small peptides and amino acids (di- and tripeptides) before their absorption by proteolytic and digestive enzymes such as trypsin.
- Lipids (fats) are degraded into fatty acids and glycerol by pancreatic lipase.
- Carbohydrates are degraded into monosaccharide or oligosaccharide sugars by the action of amylase. Carbohydrates, such as cellulose, pass through the human intestinal tract undigested.
- Water and some water-soluble vitamins are absorbed by diffusion. Some electrolytes and water non-soluble vitamins require an active uptake mechanism.

Key Terms

- **trypsin**: A digestive enzyme that cleaves peptide bonds (a serine protease).
- **lipase**: Any of a group of enzymes that catalyses the hydrolysis of lipids.
- **amylase**: Any of a class of digestive enzymes that are present in saliva and that break down complex carbohydrates, such as starch, into simpler sugars, such as glucose.


EXAMPLES

During breastfeeding, the lactase enzyme breaks down lactose (milk sugar). However, lactase production ceases after weaning in most populations, so adults in those populations experience gastric discomfort or distress when eating dairy products.

Digestive Enzymes and the Small Intestine

The small intestine is where most chemical digestion occurs. Most of the digestive enzymes that act in the small intestine are secreted by the pancreas and enter the small intestine via the pancreatic duct.

The enzymes enter the small intestine in response to the hormone cholecystokinin, which is produced in the small intestine in response to the presence of nutrients. The hormone secretin also causes bicarbonate to be released into the small intestine from the pancreas in order to neutralize the potentially harmful acid that comes from the stomach.



Small intestine: This image shows the position of the small intestine in the gastrointestinal tract.

The three major classes of nutrients that undergo digestion are:

1. Proteins. These are degraded into small peptides and amino acids before absorption. Their chemical breakdown begins in the stomach and continues in the large intestine. Proteolytic enzymes, including trypsin and chymotrypsin, are secreted by the pancreas and cleave proteins into smaller peptides. Carboxypeptidase, which is a pancreatic brush border enzyme, splits one amino acid at a time. Aminopeptidase and dipeptidase free the final amino acid products.
2. Lipids (fats). These are degraded into fatty acids and glycerol. Pancreatic lipase breaks down the triglycerides into free fatty acids and monoglycerides. Pancreatic lipase works with the help of the salts from the bile secreted by the liver and the gall bladder. Bile salts attach to triglycerides to help emulsify them and aid access by pancreatic lipase. This occurs because the lipase is water soluble, but the fatty triglycerides are hydrophobic and tend to orient towards each other and away from the watery intestinal surroundings. The bile salts are the main thing that holds the triglycerides in their watery surroundings until the lipase can break them into the smaller components that can enter the villi for absorption.
3. Carbohydrates. Some carbohydrates are degraded into simple sugars, or monosaccharides (e.g., glucose). Pancreatic amylase breaks down some carbohydrates (notably starch) into oligosaccharides. Other carbohydrates pass undigested into the large intestine for further handling by intestinal bacteria.

Brush Border Enzymes

Brush border enzymes take over from there. The most important brush border enzymes are dextrinase and glucoamylase that further break down oligosaccharides. Other brush border enzymes are maltase, sucrase, and lactase.

Lactase is absent in most adult humans and so lactose, like most polysaccharides, is not digested in their small intestine. Some carbohydrates, such as cellulose, are not digested at all despite being made of multiple glucose units. This is because the cellulose is made out of beta-glucose that makes the inter-monosaccharidal bindings different from the ones present in starch, which consists of alpha-glucose. Humans lack the enzyme for splitting the beta-glucose-bonds, something reserved for herbivores and the bacteria from the large intestine.

The fat-soluble vitamins A, D, and E are absorbed in the upper small intestine. The factors that cause the malabsorption of fat can also affect the absorption of these vitamins. Vitamin B12 is absorbed in the ilium and must be bound to intrinsic factor, a protein secreted in the stomach, in order to be absorbed. If intrinsic factor is missing, then Vitamin B12 is not absorbed and pernicious anemia results.

Of the water-soluble vitamins, the transport of folate and B12 across the apical membrane are independent from sodium (Na+), but the other water-soluble vitamins are absorbed by Na+ co-transporters. In physiology, the primary ions of electrolytes are sodium (Na+), potassium (K+), calcium (Ca2+), magnesium (Mg2+), chloride (Cl–), hydrogen phosphate (HPO42–), and hydrogen carbonate (HCO3–).

Sodium is the main electrolyte found in extracellular fluid and is involved in fluid balance and blood pressure control. Electrolyte

balance is regulated by hormones, generally with the kidneys flushing out excess levels. In humans, electrolyte homeostasis is regulated by hormones such as the antidiuretic, aldosterone, and parathyroid hormones.

Serious electrolyte disturbances, such as dehydration and over-hydration, may lead to cardiac and neurological complications that, unless they are rapidly resolved, will result in a medical emergency. Water and minerals are reabsorbed back into the blood in the colon (large intestine) where the pH is slightly acidic—about 5.6 to 6.9.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Absorption (small intestine). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Absorp...)%23Absorption**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...tion/diffusion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- villi. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/villi**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plicae circulares. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/plicae%20circulares**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1058. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1058.png**. **License**: ***Public Domain: No Known Copyright***
- Intestinal villus simplified. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:In...simplified.svg**. **License**: ***Public Domain: No Known Copyright***
- Medical Physiology/Gastrointestinal Physiology/Digestion & Absorption. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Medical...ion%23Vitamins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Digestion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Digestion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Electrolytes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Electrolytes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Small Intestine. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Small_Intestine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/definition/lipase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- amylase. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amylase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- trypsin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trypsin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1058. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1058.png**. **License**: ***Public Domain: No Known Copyright***
- Intestinal villus simplified. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:In...simplified.svg**. **License**: ***Public Domain: No Known Copyright***
- Small intestine. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...um_diagram.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 22.13B: Absorption of Monosaccharides, Amino Acids, Dipeptides, Tripeptides, Lipids, Electrolytes, Vitamins, and Water is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.14: PHASES OF DIGESTION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

22.14A: Cephalic Phase

22.14B: Gastric Phase

This page titled 22.14: Phases of Digestion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.14A: Cephalic Phase

The cephalic phase of gastric secretion occurs before food enters the stomach due to neurological signals.

Learning Objectives

- Outline the cephalic phase of digestion

Key Points

- The cephalic phase of gastric secretion is initiated by the sight, smell, thought or taste of food.
- Neurological signals originate from the cerebral cortex and in the appetite centers of the amygdala and hypothalamus.
- This enhanced secretory activity is a conditioned reflex.
- This phase of secretion normally accounts for about 20 percent of the gastric secretion associated with eating a meal.

Key Terms

- **conditioned reflex**: A response, to a stimulus, that has been acquired by operant conditioning.
- **cephalic phase**: This occurs before food enters the stomach, especially while it is being eaten.

The cephalic phase of gastric secretion occurs before food enters the stomach, especially while it is being eaten. It results from the sight, smell, thought, or taste of food; and the greater the appetite, the more intense is the stimulation.

Neurogenic signals that initiate the cephalic phase of gastric secretion originate from the cerebral cortex, and in the appetite centers of the amygdala and hypothalamus. They are transmitted through the dorsal motor nuclei of the vagi, and then through the vagus nerve to the stomach.

This phase of secretion normally accounts for about 20% of the gastric secretions that are associated with eating a meal. Since this enhanced secretory activity is brought on by the thought or sight of food it is a conditioned reflex—it only occurs when we like or want food. When one's appetite is depressed this part of the cephalic reflex is inhibited.

The cephalic phase causes ECL cells to secrete histamine and increase HCl acid in the stomach. There will also be an influence on G cells to increase gastrin circulation.

Chain of Events for the Nervous System and Hormone System

1. Thinking of food (i.e., smell, sight) stimulates the cerebral cortex.
2. The cerebral cortex sends messages to the hypothalamus, the medulla, and the parasympathetic nervous system via the vagus nerve, and to the stomach via the gastric glands in the walls of the fundus and the body of stomach.
3. The gastric glands secrete gastric juice.
4. When food enters the stomach, the stomach stretches and activates stretch receptors.
5. The stretch receptors send a message to the medulla and then back to the stomach via the vagus nerve.
6. The gastric glands secrete more gastric juice.
7. Chemical stimuli (i.e., partially digested proteins, caffeine) directly activate G cells (enteroendocrine cells) that are located in the pyloric region of the stomach to secrete gastrin; this in turn stimulates the gastric glands to secrete gastric juice.

This page titled 22.14A: Cephalic Phase is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.14B: Gastric Phase

Learning Objectives

- Outline the gastric phase of digestion

The gastric phase is a period in which swallowed food and semi-digested protein (peptides and amino acids) activate gastric activity. About two-thirds of gastric secretion occurs during this phase. Ingested food stimulates gastric activity in two ways: by stretching the stomach and by raising the pH of its contents. Stretching activates two reflexes: a short reflex is mediated through the myenteric nerve plexus; and a long reflex is mediated through the vagus nerves and brainstem.

Gastric Secretion


Gastric secretion is stimulated chiefly by three chemicals:

1. Acetylcholine (ACh). This is secreted by the parasympathetic nerve fibers of both the short and long reflex pathways.
2. Histamine. This is a paracrine secretion from the enteroendocrine cells in the gastric glands.
3. Gastrin. This is a hormone produced by enteroendocrine G cells in the pyloric glands.

All three of these stimulate parietal cells to secrete hydrochloric acid and intrinsic factor. The chief cells secrete pepsinogen in response to gastrin and especially ACh, and ACh also stimulates mucus secretion.

As dietary protein is digested, it breaks down into smaller peptides and amino acids that directly stimulate the G cells to secrete even more gastrin: this is a positive feedback loop that accelerates protein digestion.

Small peptides also buffer the stomach acid so the pH does not fall excessively low. As digestion continues and these peptides empty from the stomach, the pH drops lower and lower. Below pH of 2, stomach acid inhibits the parietal cells and G cells: this is a negative feedback loop that winds down the gastric phase as the need for pepsin and HCl declines.



The gastric phase of digestion: During the gastric phase, gastrin is secreted. The stomach stretches and churns while enzymes break down proteins.

Key Points

- The gastric phase accounts for about two-thirds of gastric secretions.
- Ingested food stimulates gastric activity by stretching the stomach and raising the pH of its contents; this causes a cascade of events that leads to the release of hydrochloric acid by the parietal cells that lower the pH and break apart the food.
- Gastric secretion is stimulated chiefly by three chemicals: acetylcholine (ACh), histamine, and gastrin.
- Below pH of 2, stomach acid inhibits the parietal cells and G cells; this is a negative feedback loop that winds down the gastric phase as the need for pepsin and HCl declines.

Key Terms

- **gastric phase**: The second phase of digestion that follows mastication (chewing) and takes place in the stomach.

This page titled 22.14B: Gastric Phase is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.6: THE STOMACH

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

22.6A: Gross Anatomy of the Stomach

22.6B: Microscopic Anatomy of the Stomach

22.6C: Digestive Properties of the Stomach

This page titled 22.6: The Stomach is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.6A: Gross Anatomy of the Stomach

The stomach is divided into four sections, each of which has different cells and functions.

Learning Objectives

- Describe the gross anatomy of the stomach


Key Points

- The stomach is divided into four sections: the cardiac region, the fundus, the body, and the pylorus or atrium.
- The stomach is lined by a mucous membrane that contains glands (with chief cells) that secrete gastric juices.
- Two smooth muscle valves, or sphincters, keep the contents of the stomach contained: the cardiac or esophageal sphincter and the pyloric sphincter.
- The arteries supplying the stomach are the left gastric, the right gastric, and the right gastroepiploic branches of the hepatic, and the left gastroepiploic and short gastric branches of the lineal.
- Lymphatics consist of a superficial and a deep set, and pass to the lymph glands found along the two curvatures of the organ.
- The nerves are the terminal branches of the right and left urethra and other parts; the former are distributed upon the back, and the latter upon the front, part of the organ.

Key Terms

- **chief cell**: A cell located in the stomach and parathyroid gland that secretes precursor enzymes.
- **pylorus**: The opening in a vertebrate, including humans, at the lower end of the stomach that opens into the duodenum.
- **sphincter**: A ringlike band of muscle that surrounds a bodily opening and constricts and relaxes as required for normal physiological functioning.

The stomach is a thick, walled organ that lies between the esophagus and the first part of the small intestine (the duodenum). It is on the left side of the abdominal cavity, the fundus of the stomach lying against the diaphragm. Lying beneath the stomach is the pancreas. The greater omentum hangs from the greater curvature.



Sections of the stomach: This diagram of the stomach shows the cardiac region, fundus, body, and pylorus.

A mucous membrane lines the stomach that contains the glands (with chief cells) that secrete gastric juices. Up to three quarts of this digestive fluid is produced daily. The gastric glands begin secreting before food enters the stomach due to the parasympathetic impulses of the vagus nerve, that also make the stomach a storage vat for that acid.

The stomach is divided into four sections, each of which has different cells and functions. The sections are:

1. The cardiac region, where the contents of the esophagus empty into the stomach.
2. The fundus, which is formed by the upper curvature of the organ.
3. The body, the main central region.
4. The pylorus or atrium, the lower section of the organ that facilitates the emptying of the contents into the small intestine.

Two smooth muscle valves, or sphincters, keep the contents of the stomach contained. They are the:

1. Cardiac or esophageal sphincter that divides the tract above.
2. Pyloric sphincter or pyloric orifice that divides the stomach from the small intestine.

The arteries that supply the stomach are the left gastric, the right gastric, and right gastroepiploic branches of the hepati;, and the left gastroepiploic and short gastric branches of the lineal. They supply the muscular coat, ramify in the submucous coat, and are finally distributed to the mucous membrane.

The arteries break up at the base of the gastric tubules into a plexus of fine capillaries that run upward between the tubules. They anatomize with each other and end in a plexus of larger capillaries that surround the mouths of the tubes and also form hexagonal meshes around the ducts.

From these the veins arise, and pursue a straight course downward, between the tubules, to the submucous tissue; they end in the lineal and superior mesenteric veins or directly in the portal vein.

The lymphatics are numerous. They consist of a superficial and a deep set, and pass to the lymph glands found along the two curvatures of the organ.

The nerves are the terminal branches of the right and left urethra and other parts, the former being distributed upon the back, and the latter upon the front part of the organ. A great number of branches from the celiac plexus of the sympathetic are also distributed to it.

Nerve plexuses are found in the submucous coat and between the layers of the muscular coat as in the intestine. Fibrils are distributed from these plexuses to the muscular tissue and the mucous membrane.

This page titled 22.6A: Gross Anatomy of the Stomach is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.6B: Microscopic Anatomy of the Stomach

The layers of the stomach produce mucus to protect itself, enyzmes to break down the food for digestion, and muscles to churn the food.

Learning Objectives

- Diagram the microscopic anatomy of the stomach

Key Points

- The stomach walls are made of the following layers (inside to outside): mucosa, muscularis mucosa, submucosa, and muscularis externa.
- The epithelium of the stomach forms deep pits (fundic or oxyntic glands) where chief cells produce pepsinogen, an inactive precursor of pepsin that degrades proteins. The secretion of pepsinogen prevents self-digestion of the stomach cells.
- Gastric acid kills most of the bacteria in food, stimulates hunger, and activates pepsinogen into pepsin; it also denatures the complex protein molecule as a precursor to protein digestion.
- Goblet cells produce mucus that protects the stomach from self-digestion.
- The muscularis externa has three layers of smooth muscle. The innermost obliquely-oriented layer is responsible for creating the motion that churns and physically breaks down the food, and is unique to the stomach. The other layers are present as in other parts of the GI tract.

Key Terms

- **muscularis externa**: A region of muscle in many organs in the vertebrate body that is adjacent to the submucosa membrane. It is responsible for gut movements, such as peristalsis.
- **muscularis mucosae**: Also called the lamina muscularis mucosae, this is the thin layer of smooth muscle found in most parts of the gastrointestinal tract. It is located outside the lamina propria mucosae and separates it from the submucosa.
- **submucosa**: A layer of connective tissue beneath a mucous membrane.

EXAMPLES

If there is low or no gastric acid in the stomach, this could lead to problems as the disinfectant properties of the gastric lumen are decreased. In such conditions, there is a greater risk of infections in the digestive tract (such as infection with Vibrio or Helicobacter bacteria).

Anatomy of the Stomach

Like the other parts of the gastrointestinal tract, the stomach walls are made of a number of layers. From the inside to the outside, the first main layer is the mucosa.

Mucosa



The stomach wall: A micrograph that shows a cross section of the stomach wall, in the body portion of the stomach.

This consists of an epithelium, the lamina propria underneath, and a thin bit of smooth muscle called the muscularis mucosae. The submucosa lies under this and consists of fibrous connective tissue that separate the mucosa from the next layer, the muscularis externa.

The muscularis in the stomach differs from that of other GI organs in that it has three layers of muscle instead of two. Under these muscle layers is the adventitia, layers of connective tissue continuous with the omenta.

The epithelium of the stomach forms deep pits, called fundic or oxyntic glands. Different types of cells are at different locations down the pits. The cells at the base of these pits are chief cells that are responsible for the production of pepsinogen, an inactive precursor of pepsin, which degrades proteins. The secretion of pepsinogen prevents self-digestion of the stomach cells.

Further up the pits, parietal cells produce gastric acid and a vital substance, intrinsic factor. The function of gastric acid is two fold:

1. It kills most of the bacteria in food, stimulates hunger, and activates pepsinogen into pepsin.
2. It denatures the complex protein molecule as a precursor to protein digestion through enzyme action in the stomach and small intestines.

Near the top of the pits, closest to the contents of the stomach, there are mucus-producing cells called goblet cells that help protect the stomach from self-digestion.

The Muscularis Externa

The muscularis externa is made up of three layers of smooth muscle.

- The innermost layer is obliquely-oriented: this is not seen in other parts of the digestive system. This layer is responsible for creating the motion that churns and physically breaks down the food.
- The next layers are the square and then the longitudinal, which are present as in other parts of the GI tract.
- The pyloric antrum has thicker skin cells in its walls and performs more forceful contractions than the fundus. The pylorus is surrounded by a thick circular muscular wall that is normally tonically constricted, forming a functional (if not anatomically discrete) pyloric sphincter that controls the movement of chyme.

This page titled 22.6B: Microscopic Anatomy of the Stomach is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

22.6C: Digestive Properties of the Stomach

The movement and flow of chemicals into the stomach is controlled by the autonomic nervous system and various digestive system hormones.

Learning Objectives

- List the various digestive properties of the stomach

Key Points

- Digestive hormones include: gastrin, somatostatin, cholecystokinin, secretin, gastric inhibitory peptide, and enteroglucagon.
- Gastrin causes an increase in the secretion of HCl from the parietal cells, an increase in pepsinogen from the chief cells in the stomach, and an increase in stomach motility.
- Somatostatin inhibits the activity of gastrin. Cholecystokinin (CCK) causes gall bladder contractions; it also decreases gastric emptying and increases the release of pancreatic juice that is alkaline and neutralizes the chyme. Secretin diminishes acid secretion in the stomach.
- Gastric inhibitory peptide (GIP) and enteroglucagon decrease gastric acid release and motility. They also turn off stomach action when the gall bladder and liver have not yet finished absorption.
- Epidermal growth factor (EGF), produced by the salivary glands, heals oral and gastroesophageal ulcers, inhibits gastric acid secretion, and stimulates DNA synthesis as well as mucosal protection.
- The stomach can sense, independently to the tongue and oral taste receptors, glucose, carbohydrates, proteins, fats, and sodium glutamate via the vagus nerve, which allows the brain to link the nutritional value of foods to their tastes.

Key Terms

- **gastrin**: A hormone that stimulates the production of gastric acid in the stomach.
- **antrum**: A bodily cavity, especially one having bony walls, especially in the sinuses.
- **enteroglucagon**: A peptide hormone derived from preproglucagon that is secreted from the mucosal cells, primarily of the colon and terminal ileum, that follows ingestion of a mixed meal to delay gastric emptying.
- **somatostatin**: A polypeptide hormone, secreted by the pancreas, that inhibits the production of certain other hormones.

EXAMPLES

Gastric ulcers can be healed by supplemental daily doses of epidermal growth factor (EGF). EGF inhibits bacterial colonization and speeds healing just as well as antibiotics.

Stomach Hormones



Fundic gland: The fundic gland, found in the stomach, secretes gastrin and other hormones.

The movement and the flow of chemicals into the stomach are controlled by both the autonomic nervous system and the various digestive system hormones.

The hormone gastrin causes an increase in the secretion of HCl from the parietal cells and pepsinogen from the chief cells in the stomach. It also causes increased motility (movement, peristalsis) in the stomach.

Gastrin is released by G-cells in the stomach, via the base cells of the pyloric, cardiac, and fundic glands, in response to distension of the antrum, and digestive products (especially large quantities of incompletely digested proteins). It is inhibited by a pH normally less than four (high acid), as well as the hormone somatostatin.

Cholecystokinin (CCK) primarily effects the gall bladder and causes it to contract, but it also decreases gastric emptying and increases release of pancreatic juice, which is alkaline and neutralizes the chyme.

In a different and rare manner, secretin that is produced in the small intestine primarily effects the pancreas, but it will also diminish acid secretion in the stomach. Gastric inhibitory peptide (GIP) decreases both gastric acid release and motility. Enteroglucagon decreases both gastric acid and motility.



Cardiac gland: The cardiac gland, found in the stomach, secretes gastrin and other hormones.

Other than gastrin, these hormones all act to turn off the stomach's action. This is in response to food products in the liver and gall bladder that have not yet been absorbed. The stomach needs to push food into the small intestine only when it is not busy. While the intestine is full and still digesting food, the stomach will act as storage for food.

Epidermal Growth Factor

Epidermal growth factor (EGF) results in cellular proliferation, differentiation, and survival. EGF is a low-molecular-weight polypeptide that was first purified from a mouse's submandibular gland, but has since been found in many human tissues including the submandibular and parotid glands.



Pyloric gland: The pyloric gland, found in the stomach, secretes gastrin and other hormones.

Salivary EGF, which seems to be regulated by dietary inorganic iodine, plays an important physiological role in the maintenance of oroesophageal and gastric tissue integrity. The biological effects of salivary EGF include the healing of oral and gastroesophageal ulcers, the inhibition of gastric acid secretion, and the stimulation of DNA synthesis, as well as providing mucosal protection from intraluminal injurious factors—such as gastric acid, bile acids, pepsin, and trypsin, and to physical, chemical, and bacterial agents.

The stomach can taste sodium glutamate using glutamate receptors. This information is passed to the lateral hypothalamus and limbic system in the brain as a palatability signal through the vagus nerve. The stomach can also sense, independently from the tongue and oral taste receptors, glucose, carbohydrates, proteins, and fats. This allows the brain to link the nutritional value of foods to their tastes.

Although absorption is the primary function of the small intestine, some absorption of certain small molecules does nevertheless occur in the stomach through its lining. This includes water if the body is too dehydrated; medication, like aspirin; and amino acids (e.g., whey protein shake).

There are many different gastric glands that secrete many different chemicals. Parietal cells secrete hydrochloric acid and intrinsic factor

- Chief cells secrete pepsinogen.
- Goblet cells secrete mucus.
- Argentaffin cells secrete serotonin and histamine.
- G cells secrete the hormone gastrin.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...stem%23Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- pylorus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pylorus. **License**: *CC BY-SA: Attribution-ShareAlike*
- sphincter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sphincter. **License**: *CC BY-SA: Attribution-ShareAlike*
- chief cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chief_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...stem%23Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gastric acid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastric_acid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- muscularis mucosae. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/muscularis%20mucosae. **License**: *CC BY-SA: Attribution-ShareAlike*
- submucosa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/submucosa. **License**: *CC BY-SA: Attribution-ShareAlike*
- muscularis externa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/muscularis%20externa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The gastrointestinal system. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...stem%23Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach%23Glands. **License**: *CC BY-SA: Attribution-ShareAlike*
- antrum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antrum. **License**: *CC BY-SA: Attribution-ShareAlike*
- somatostatin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somatostatin. **License**: *CC BY-SA: Attribution-ShareAlike*
- enteroglucagon. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/enteroglucagon. **License**: *CC BY-SA: Attribution-ShareAlike*
- gastrin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/gastrin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*
- Stomach. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stomach. **License**: *CC BY-SA: Attribution-ShareAlike*

PUBLIC DOMAIN CONTENT

- Gray1046.svg. **Authored by**: Henry Gray. **Located at**: commons.wikimedia.org/wiki/File:Gray1046.svg. **License**: *Public Domain: No Known Copyright*

This page titled 22.6C: Digestive Properties of the Stomach is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23: NUTRITION AND METABOLISM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.10: Diet, Digestion, and Energy Storage Regulation

23.10A: Hormonal Responses to Food

23.10B: Neural Responses to Food

23.10C: Food Energy and ATP

23.1: Nutrition

23.1A: Digestive Systems

23.1B: Carbohydrates: Sources, Uses in the Body, and Dietary Requirements

23.1C: Lipids: Sources, Uses in the Body, and Dietary Requirements

23.1D: Proteins: Sources, Uses in the Body, and Dietary Requirements

23.1E: Food Requirements and Essential Nutrients

23.2: Carbohydrates

23.2A: Carbohydrate Molecules

23.2B: Importance of Carbohydrates

23.3: Lipids

23.3A: Lipid Molecules

23.3B: Phospholipids

23.3C: Steroids

23.4: Proteins

23.4A: Types and Functions of Proteins

23.4B: Amino Acids

23.4C: Protein Structure

23.5: Metabolic Reactions

23.5A: Breakdown of Pyruvate

23.5B: Citric Acid Cycle

23.5C: Importance of Glycolysis

23.5D: Electron Transport Chain

23.5E: ATP Yield

23.5F: Control of Catabolic Pathways

23.5G: Transforming Chemical Energy

23.6: Metabolism of Nutrients

23.6A: Connecting Other Sugars to Glucose Metabolism

23.6B: Connecting Proteins to Glucose Metabolism

23.7: Metabolic Body States

23.7A: Catabolic-Anabolic Steady State

23.7B: Absorptive State

23.7C: Postabsorptive State

23.8: Metabolism and the Liver

23.8A: Cholesterol Metabolism

23.8B: Regulation of Blood Cholesterol Levels

23.9: Metabolic Rate and Thermoregulation

23.9A: Homeostatic Process

23.9B: Homeostasis: Thermoregulation

23.9C: Heat Conservation and Dissipation

This page titled 23: Nutrition and Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.10: DIET, DIGESTION, AND ENERGY STORAGE REGULATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

23.10A: Hormonal Responses to Food
23.10B: Neural Responses to Food
23.10C: Food Energy and ATP

This page titled 23.10: Diet, Digestion, and Energy Storage Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.10A: Hormonal Responses to Food

The endocrine system controls the release of hormones responsible for starting, stopping, slowing, and quickening digestive processes.

Learning Objectives

- Describe hormonal responses to food

Key Points

- The presence and absence of hormones that are released into the bloodstream generate specific digestive responses; they either stimulate or discontinue digestive processes.
- In hormone control, a negative feedback mechanism takes place when the stomach is empty and its acidic environment does not need to be maintained; as a result, a hormone is released to stop the release of hydrochloric acid, which was previously activated to aid digestion.
- In some cases, hormones work in tandem and sequentially to achieve important digestive functions, such as in the breakdown of acidic chyme, where hormones act in releasing the appropriate secretions in the appropriate stages of digestion.
- When digesting certain types of foods, such as ones high in fat, hormones can control the speed of food digestion and, therefore, absorption.

Key Terms


- **endocrine system**: a control system of ductless glands that secrete hormones which circulate via the bloodstream to affect cells within specific organs
- **chyme**: the thick semifluid mass of partly digested food that is passed from the stomach to the duodenum
- **secretin**: a peptide hormone, secreted by the duodenum, that serves to regulate its acidity
- **cholecystokinin**: any of several peptide hormones that stimulate the digestion of fat and protein
- **somatostatin**: a polypeptide hormone, secreted by the pancreas, that inhibits the production of certain other hormones
- **gastrin**: a hormone that stimulates the production of gastric acid in the stomach

Hormonal Responses to Food

The endocrine system controls the response of the various glands in the body and the release of hormones at the appropriate times. The endocrine system's effects are slow to initiate, but prolonged in their response, lasting from a few hours up to weeks. The system is made of a series of glands that produce chemicals called hormones. These hormones are chemical mediators released from endocrine tissue into the bloodstream where they travel to target tissue and generate a response.

One of the important factors under hormonal control is the stomach acid environment. During the gastric phase, the hormone gastrin is secreted by G cells in the stomach in response to the presence of proteins. Gastrin stimulates the release of stomach acid, or hydrochloric acid (HCl), which aids in the digestion of the majority of proteins. However, when the stomach is emptied, the acidic environment need not be maintained and a hormone called somatostatin stops the release of hydrochloric acid. This is controlled by a negative feedback mechanism.

In the duodenum, digestive secretions from the liver, pancreas, and gallbladder play an important role in digesting chyme during the intestinal phase. In order to neutralize the acidic chyme, a hormone called secretin stimulates the pancreas to produce alkaline bicarbonate solution and deliver it to the duodenum. Secretin acts in tandem with another hormone called cholecystokinin (CCK). Not only does CCK stimulate the pancreas to produce the requisite pancreatic juices, it also stimulates the gallbladder to release bile into the duodenum.



Digestive endocrine system: Hormones, such as secretin and cholecystokinin, play important roles in digestive processes. These hormones are released from endocrine tissue to generate specific controls in the digestion of chyme. As seen in the image, hormones are vital players in several bodily functions and processes.

Another level of hormonal control occurs in response to the composition of food. Foods high in lipids (fatty foods) take a long time to digest. A hormone called gastric inhibitory peptide is secreted by the small intestine to slow down the peristaltic movements of the intestine to allow fatty foods more time to be digested and absorbed.

Understanding the hormonal control of the digestive system is an important area of ongoing research. Scientists are exploring the role of each hormone in the digestive process and developing ways to target these hormones. Advances could lead to knowledge that may help to battle the obesity epidemic.

This page titled 23.10A: Hormonal Responses to Food is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.10B: Neural Responses to Food

All three phases of digestive responses to food (the cephalic, gastric, and intestinal stages) are managed through enzymatic neural control.

Learning Objectives

- Summarize the neural responses to food

Key Points

- The cephalic phase is controlled by sight, sense, and smell, which trigger neural responses, including salivation and hydrochloric acid production, before food has even reached the mouth.
- Once food reaches the stomach, gastric acids and enzymes process the ingested materials in the gastric phase, which involves local, hormonal, and neural responses.
- The intestinal phase controls the rate of gastric emptying and the release of hormones needed to digest chyme in the small intestine.

Key Terms

- **neural**: of, or relating to the nerves, neurons or the nervous system
- **salivary gland**: any of several exocrine glands that produce saliva to break down carbohydrates in food enzymatically
- **peristaltic**: of, or pertaining to the rhythmic, wave-like contraction of the digestive tract that forces food through it

Neural Responses to Food

In reaction to the smell, sight, or thought of food, the first hormonal response is that of salivation. The salivary glands secrete more saliva in response to the stimulus presented by food in preparation for digestion. Simultaneously, the stomach begins to produce hydrochloric acid to digest the food. Recall that the peristaltic movements of the esophagus and other organs of the digestive tract are under the control of the brain. The brain prepares these muscles for movement as well. When the stomach is full, the part of the brain that detects satiety signals fullness. There are three overlapping phases of gastric control: the cephalic phase, the gastric phase, and the intestinal phase. Each requires many enzymes and is under neural control as well.



Salivation: Seeing a plate of food triggers the secretion of saliva in the mouth and the production of hydrochloric acid in the stomach.

Digestive Phases

The response to food begins even before food enters the mouth. The first phase of ingestion, called the cephalic phase, is controlled by the neural response to the stimulus provided by food. All aspects, such as sight, sense, and smell, trigger the neural responses resulting in salivation and secretion of gastric juices. The gastric and salivary secretion in the cephalic phase can also take place at the thought of food. Right now, if you think about a piece of chocolate or a crispy potato chip, the increase in salivation is a cephalic phase response to the thought. The central nervous system prepares the stomach to receive food.

The gastric phase begins once the food arrives in the stomach. It builds on the stimulation provided during the cephalic phase. Gastric acids and enzymes process the ingested materials. The gastric phase is stimulated by (1) distension of the stomach, (2) a decrease in the pH of the gastric contents, and (3) the presence of undigested material. This phase consists of local, hormonal, and neural responses. These responses stimulate secretions and powerful contractions.

The intestinal phase begins when chyme enters the small intestine, triggering digestive secretions. This phase controls the rate of gastric emptying. In addition to gastric emptying, when chyme enters the small intestine, it triggers other hormonal and neural events that coordinate the activities of the intestinal tract, pancreas, liver, and gallbladder.

This page titled 23.10B: Neural Responses to Food is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.10C: Food Energy and ATP


Learning Objectives

- Summarize the ways in which animals obtain, store, and use food energy

Animals need food to obtain energy and maintain homeostasis. Homeostasis is the ability of a system to maintain a stable internal environment even in the face of external changes to the environment. For example, the normal body temperature of humans is 37°C (98.6°F). Humans maintain this temperature even when the external temperature is hot or cold. The energy it takes to maintain this body temperature is obtained from food.

The primary source of energy for animals is carbohydrates, primarily glucose: the body's fuel. The digestible carbohydrates in an animal's diet are converted to glucose molecules and into energy through a series of catabolic chemical reactions.

Adenosine triphosphate, or ATP, is the primary energy currency in cells. ATP stores energy in phosphate ester bonds, releasing energy when the phosphodiester bonds are broken: ATP is converted to ADP and a phosphate group. ATP is produced by the oxidative reactions in the cytoplasm and mitochondrion of the cell, where carbohydrates, proteins, and fats undergo a series of metabolic reactions collectively called cellular respiration.



ATP Production Pathways

***ATP production pathways**: ATP is the energy molecule of the cell. It is produced through various pathways during the cellular respiration process, with each making different amounts of energy.*

ATP is required for all cellular functions. It is used to build the organic molecules that are required for cells and tissues. It also provides energy for muscle contraction and for the transmission of electrical signals in the nervous system. When the amount of ATP available is in excess of the body's requirements, the liver uses the excess ATP and excess glucose to produce molecules called glycogen (a polymeric form of glucose) that is stored in the liver and skeletal muscle cells. When blood sugar drops, the liver releases glucose from stores of glycogen. Skeletal muscle converts glycogen to glucose during intense exercise. The process of converting glucose and excess ATP to glycogen and the storage of excess energy is an evolutionarily-important step in helping animals deal with mobility, food shortages, and famine.

Key Points

- Animals obtain energy from the food they consume, using that energy to maintain body temperature and perform other metabolic functions.

- Glucose, found in the food animals eat, is broken down during the process of cellular respiration into an energy source called ATP.
- When excess ATP and glucose are present, the liver converts them into a molecule called glycogen, which is stored for later use.

Key Terms

- **glucose**: a simple monosaccharide (sugar) with a molecular formula of C6H12O6; it is a principal source of energy for cellular metabolism
- **adenosine triphosphate**: a multifunctional nucleoside triphosphate used in cells as a coenzyme, often called the "molecular unit of energy currency" in intracellular energy transfer
- **phosphodiester**: any of many biologically active compounds in which two alcohols form ester bonds with phosphate

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44744/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Endocrine system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endocrine_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- secretin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/secretin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- somatostatin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somatostatin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cholecystokinin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cholecystokinin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gastrin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/gastrin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chyme. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/chyme. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endocrine system. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/endocrine_system. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine Alimentary system en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._system_en.svg. **License**:
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44744/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- neural. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neural. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peristaltic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/peristaltic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- salivary gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/salivary_gland. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endocrine Alimentary system en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._system_en.svg. **License**:
- OpenStax College, Digestive System Regulation. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44744/latest...e_34_04_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44740/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- phosphodiester. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/phosphodiester. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glucose. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glucose. **License**: ***CC BY-SA: Attribution-ShareAlike***
- adenosine triphosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adenosine%20triphosphate.

License: *CC BY-SA: Attribution-ShareAlike*
- Endocrine Alimentary system en. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Endocrine_Alimentary_system_en.svg. **License**:
- OpenStax College, Digestive System Regulation. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44744/latest...e_34_04_01.jpg. **License**: *CC BY: Attribution*
- ATP Production Pathways. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...n_Pathways.jpg. **License**: *CC BY: Attribution*

This page titled 23.10C: Food Energy and ATP is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1: NUTRITION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.1A: Digestive Systems

23.1B: Carbohydrates: Sources, Uses in the Body, and Dietary Requirements

23.1C: Lipids: Sources, Uses in the Body, and Dietary Requirements

23.1D: Proteins: Sources, Uses in the Body, and Dietary Requirements

23.1E: Food Requirements and Essential Nutrients

This page titled 23.1: Nutrition is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1A: Digestive Systems

Animals use the organs of their digestive systems to extract important nutrients from food they consume, which can later be absorbed.

Learning Objectives

- Summarize animal nutrition and the digestive system

Key Points

- Animals obtain lipids, proteins, carbohydrates, essential vitamins, and minerals from the food they consume.
- The digestive system is composed of a series of organs, each with a specific, yet related function, that work to extract nutrients from food.
- Organs of the digestive system include the mouth, esophagus, stomach, small intestine, and the large intestine.
- Accessory organs, such as the liver and pancreas, secrete digestive juices into the gastrointestinal tract to assist with food breakdown.

Key Terms

- **digestion**: the process, in the gastrointestinal tract, by which food is converted into substances that can be utilized by the body
- **macromolecule**: a very large molecule, especially used in reference to large biological polymers (e.g. nucleic acids and proteins)
- **alimentary canal**: the organs of a human or an animal through which food passes; the digestive tract

All living organisms need nutrients to survive. While plants can obtain the molecules required for cellular function through the process of photosynthesis, most animals obtain their nutrients by the consumption of other organisms. At the cellular level, the biological molecules necessary for animal function are amino acids, lipid molecules, nucleotides, and simple sugars. The food consumed consists of protein, fat, and complex carbohydrates, but the requirements of each are different for each animal.



Balanced human diet: For humans, fruits and vegetables are important in maintaining a balanced diet. Both of these are an important source of vitamins and minerals, as well as carbohydrates, which are broken down through digestion for energy.


Animals must convert these macromolecules into the simple molecules required for maintaining cellular functions, such as assembling new molecules, cells, and tissues. The conversion of the food consumed to the nutrients required is a multi-step

process involving digestion and absorption. During digestion, food particles are broken down to smaller components which will later be absorbed by the body.

Digestive System

The digestive system is one of the largest organ systems in the human body. It is responsible for processing ingested food and liquids. The cells of the human body all require a wide array of chemicals to support their metabolic activities, from organic nutrients used as fuel to the water that sustains life at the cellular level. The digestive system not only effectively chemically reduces the compounds in food into their fundamental building blocks, but also acts to retain water and excrete undigested materials. The functions of the digestive system can be summarized as follows: ingestion (eat food), digestion (breakdown of food), absorption (extraction of nutrients from the food), and defecation (removal of waste products).

The digestive system consists of a group of organs that form a closed tube-like structure called the gastrointestinal tract (GI tract) or the alimentary canal. For convenience, the GI tract is divided into upper GI tract and lower GI tract. The organs that make up the GI tract include the mouth, the esophagus, the stomach, the small intestine, and the large intestine. There are also several accessory organs that secrete various enzymes into the GI tract. These include the salivary glands, the liver, and the pancreas.



Generalized animal digestive system: This diagram shows a generalized animal digestive system, detailing the different organs and their functions.

Challenges to Human Nutrition

One of the challenges in human nutrition is maintaining a balance between food intake, storage, and energy expenditure. Imbalances can have serious health consequences. For example, eating too much food while not expending much energy leads to obesity, which in turn will increase the risk of developing illnesses such as type-2 diabetes and cardiovascular disease. The recent rise in obesity and related diseases means that understanding the role of diet and nutrition in maintaining good health is more important than ever.

This page titled 23.1A: Digestive Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1B: Carbohydrates: Sources, Uses in the Body, and Dietary Requirements

Carbohydrates, which break down to glucose, are a major source of energy for humans, but are not an essential nutrient.

Learning Objectives

- Evaluate the sources and uses of carbohydrates in the body

Daily food intake that includes 8–10 fruit and vegetable servings (not starchy potatoes or grains, such as corn and rice) will not only provide plenty of energy but will also keep glucose levels in balance.




Carbohydrates are a class of macromolecule: Grain products are rich sources of carbohydrates. (Public Domain; US Department of Agriculture via Wikipedia)

Foods high in carbohydrate include fruits, sweets, soft drinks, breads, pastas, beans, potatoes, bran, rice, and cereals. Carbohydrates are a common source of energy in living organisms, however, a carbohydrate is not an essential nutrient in humans.

Carbohydrates are not necessary building blocks of other molecules, and the body can obtain all its energy from protein and fats. The brain and neurons generally cannot burn fat for energy but use solely glucose or ketones. Humans can synthesize some glucose (in a set of processes known as “gluconeogenesis”) from specific amino acids or from the glycerol backbone in triglycerides and, in some cases, from fatty acids. Carbohydrate and protein contain 4 kilocalories per gram, while fats contain 9 kilocalories per gram. In the case of protein, this is somewhat misleading as only some amino acids are able to undergo conversion into useful energy forms.

Organisms typically cannot metabolize all types of carbohydrate to yield energy. Glucose is a nearly universal and accessible source of calories. Many organisms also have the ability to metabolize other monosaccharides and disaccharides, though glucose is preferred. Polysaccharides are also common sources of energy. Even though these complex carbohydrates are not very digestible, they may comprise important dietary elements for humans. Called “dietary fiber,” these carbohydrates enhance digestion, among other benefits.



Glucose Molecule: Image of a glucose molecule containing a fixed ratio of carbon, hydrogen and oxygen.

Based on the effects on risk of heart disease and obesity, the Institute of Medicine (IOM) recommends that American and Canadian adults get between 45–65% of dietary energy from carbohydrates. The Food and Agriculture Organization (FAO) and World Health Organization (WHO) jointly recommend that national dietary guidelines set a goal of 55–75% of total energy from carbohydrates, but only 10% directly from sugars (their term for simple carbohydrates).

Key Points

- Carbohydrates include such items as fruits, grains, beans, and potatoes, along with sugars and sugared foods.
- While fat is a better source of energy, the brain cannot burn fat and instead requires glucose.
- Polysaccharides (complex carbs) are difficult for humans to breakdown, but are useful as fiber to enhance the digestive process.
- Government agencies recommend a dietary intake of 45–65% or 55–75% of carbohydrates to meet daily energy needs.
- Of daily carbohydrate intake, only 10% should be simple carbs or sugars.

Key Terms

- **glucose**: A simple monosaccharide (sugar) with a molecular formula of C6H12O6; it is a principle source of energy for cellular metabolism.
- **carbohydrate**: A sugar, starch, or cellulose that is a food source of energy for an animal or plant; a saccharide.
- **saccharide**: The unit structure of carbohydrates, of general formula CnH2nOn. Either the simple sugars or polymers such as starch and cellulose. The saccharides exist in either a ring or short chain conformation, and typically contain five or six carbon atoms.

This page titled 23.1B: Carbohydrates: Sources, Uses in the Body, and Dietary Requirements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1C: Lipids: Sources, Uses in the Body, and Dietary Requirements

Fats store energy, facilitate absorption of fat-soluble vitamins, aid brain growth and development, and protect against many diseases.

Learning Objectives

- Evaluate the sources and uses of lipids in the body

Key Points

- Vitamins A, D, E, and K should be taken with some dietary fat in order to facilitate their absorption and activity.
- Humans cannot synthesize omega-6 and omega-3 fatty acids, so these fats must be obtained from the diet.
- Omega-6 fatty acids are found in many foods, while omega-3 fatty acids are found in walnuts and are especially abundant in fatty fish.
- Omega-3 fatty acids have many positive health benefits including reduced rates of cancer, cardiovascular disease, mental illness, and dementia.
- Studies of dietary fat intake have found no link between percentage of calories obtained from fats and risk of cancer, heart disease, or obesity.
- Trans fats, such as those present in partially hydrogenated vegetable oils, are a risk factor for cardiovascular disease.

Key Terms


- **phospholipids**: Phospholipids are a class of lipids that are a major component of all cell membranes as they can form lipid bilayers.
- **trans fats**: Trans fats are unsaturated fats generated by physical agents such as heat or pressure that can lead to a variety of health problems.
- **fatty acid**: Fatty acids can be saturated or unsaturated and are usually derived from triglycerides or phospholipids.

EXAMPLES

College students require optimal brain function, which is supported by fatty fish and walnuts.

Most of the fats found in food are triglycerides, cholesterol, and phospholipids. Some dietary fat is necessary for the absorption of fat-soluble vitamins (A, D, E, and K) and carotenoids. Humans and other mammals require fatty acids such as linoleic acid (an omega-6 fatty acid) and alpha-linolenic acid (an omega-3 fatty acid), because they cannot be synthesized from simple precursors in the diet.

Both omega-6 and omega-3 are 18-carbon polyunsaturated fatty acids that differ in the number and position of their double bonds. Most vegetable oils (safflower, sunflower, and corn oils) are rich in linoleic acid. Alpha-linolenic acid is found in the green leaves of plants, selected seeds, nuts, and legumes, and particularly in flax, rapeseed, walnut, and soy. Fish oils are especially rich in the longer-chain omega-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Numerous studies have shown that the consumption of omega-3 fatty acids has positive benefits in terms of infant development, cancer, cardiovascular disease, and mental illnesses such as depression, attention-deficit hyperactivity disorder, and dementia. In contrast, the consumption of trans fats, such as those present in partially hydrogenated vegetable oils, are now known to be a risk factor for cardiovascular disease.



Omega-3 Fatty Acid: Docosahexaenoic acid (DHA) is an omega-3 fatty acid. It is essential for proper functioning of the brain in both adults and infants. DHA concentrations in breast milk range from 0.07-1.0% of total fatty acids and are influenced by the amount of fatty fish in the mother's diet. In the U.S., infant formula has been supplemented with DHA since 2001. Research suggests that DHA contributes to numerous nervous system functions such as visual acuity, neurogenesis, and synaptogenesis, and that it lowers the risk for cardiovascular disease. It is highly concentrated in the brain and eye.

Several studies have suggested that total dietary fat intake is linked to obesity and diabetes. However, influential studies like the Women's Health Initiative Dietary Modification Trial (an eight-year study of 49,000 women), as well as the Nurses' Health Study and the Health Professionals Follow-up Study, have revealed no such link between percentage of calories from fat and risk of cancer, heart disease, or weight gain. The Nutrition Source, a website maintained by the Department of Nutrition at the Harvard School of Public Health, summarizes the current evidence on the impact of dietary fat as follows: "Detailed research—much of it done at Harvard—shows that the total amount of fat in the diet isn't really linked with weight or disease."



Salmon: Salmon is an excellent source of omega-3 fatty acids.

This page titled 23.1C: Lipids: Sources, Uses in the Body, and Dietary Requirements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1D: Proteins: Sources, Uses in the Body, and Dietary Requirements

Proteins are composed of 20 different amino acids, about half of which are essential, meaning they must be obtained from the diet.

Learning Objectives

- Evaluate protein sources and uses in the body


Key Points

- Protein -based foods (plant and animal) provide amino acids; however, the best source of essential amino acids is animal.
- If the diet does not provide adequate protein, the body will obtain what it needs from itself, especially from its own muscles.
- While adequate protein is required for building skeletal muscle and other tissues, there is ongoing debate regarding the use and necessity of high-protein diets in anaerobic exercise, in particular weight training and bodybuilding.
- This use of protein as a fuel is particularly important under starvation conditions as it allows the body's own proteins to be used to support life.

Key Terms


- **amino acid**: Any of the twenty naturally occurring α-amino acids (having the amino, and carboxylic acid groups on the same carbon atom), and a variety of side chains, that combine, via peptide bonds, to form proteins.
- **denaturation**: Denaturation is a process in which proteins or nucleic acids lose their tertiary and secondary structure which is present in their native state, by application of some external stress or compound such as a strong acid or base, a concentrated inorganic salt, an organic solvent (e.g., alcohol or chloroform), or heat.
- **protein**: Any of numerous large, complex naturally-produced molecules composed of one or more long chains of amino acids, in which the amino acid groups are held together by peptide bonds.

Most microorganisms and plants can biosynthesize all 20 standard amino acids, while animals (including humans) must obtain some of the amino acids from the diet. The amino acids that an organism cannot synthesize on its own are referred to as essential amino acids. Key enzymes that synthesize certain amino acids are not present in animals. One example is aspartokinase, which catalyzes the first step in the synthesis of lysine, methionine, and threonine from aspartate.



Amino Acid: Ball-and-stick model of the cystine molecule, an amino acid formed from two cysteine molecules.

In animals, amino acids are obtained through the consumption of foods containing protein. Ingested proteins are then broken down into amino acids through digestion, which typically involves denaturation of the protein through exposure to acid and hydrolysis by enzymes called proteases. Some ingested amino acids are used for protein biosynthesis, while others are converted to glucose through gluconeogenesis, or fed into the citric acid cycle. This use of protein as a fuel is particularly important under starvation conditions as it allows the body's own proteins to be used to support life, particularly those found in muscle. Amino acids are also an important dietary source of nitrogen.



Protein: Muscle meat, such as steak, is an excellent source of the essential amino acids the body needs to create all necessary proteins.

A high-protein diet is often recommended by bodybuilders and nutritionists to help efforts to build muscle and lose fat. It should not be confused with low-carb diets, such as the Atkins Diet, which are not calorie-controlled and which often contain large amounts of fat. While adequate protein is required for building skeletal muscle and other tissues, there is ongoing debate regarding the use and necessity of high-protein diets in anaerobic exercise in particular weight training and bodybuilding. Extreme protein intake (in excess of 200g per day), coupled with inadequate intake of other calorie sources (fat or carbohydrates), can cause a form of metabolic disturbance and death commonly known as rabbit starvation.

Relatively little evidence has been gathered regarding the effect of long-term high intake of protein on the development of chronic diseases. Increased load on the kidney is a result of an increase in reabsorption of NaCl. This causes a decrease in the sensitivity of tubuloglomerular feedback, which, in turn, results in an increased glomerular filtration rate. This increases pressure in glomerular capillaries. When added to any additional renal disease, this may cause permanent glomerular damage.

This page titled 23.1D: Proteins: Sources, Uses in the Body, and Dietary Requirements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.1E: Food Requirements and Essential Nutrients

Essential nutrients are those that cannot be created by an animal's metabolism and need to be obtained from the diet.

Learning Objectives

- Describe the essential nutrients required for cellular function that cannot be synthesized by the animal body


Key Points

- The animal diet needs to be well-balanced in order to ensure that all necessary vitamins and minerals are being obtained.
- Vitamins are important for maintaining bodily health, making bones strong, and seeing in the dark.
- Water-soluble vitamins are not stored by the body and need to be consumed more regularly than fat-soluble vitamins, which build up within body tissues.
- Essential fatty acids need to be consumed through the diet and are important building blocks of cell membranes.
- Nine of the 20 amino acids cannot be synthesized by the body and need to be obtained from the diet.

Key Terms

- **nutrient**: a source of nourishment, such as food, that can be metabolized by an organism to give energy and build tissue
- **catabolism**: destructive metabolism, usually including the release of energy and breakdown of materials
- **vitamin**: any of a specific group of organic compounds essential in small quantities for healthy human growth, metabolism, development, and body function

Food Requirements



A balanced diet: For humans, a balanced diet includes fruits, vegetables, grains, and protein. Each of these food sources provides different nutrients the body cannot make for itself. These include vitamins, omega 3 fatty acids, and some amino acids.

What are the fundamental requirements of the animal diet? The animal diet should be well balanced and provide nutrients required for bodily function along with the minerals and vitamins required for maintaining structure and regulation necessary for good health and reproductive capability.

Organic Precursors

The organic molecules required for building cellular material and tissues must come from food. Carbohydrates or sugars are the primary source of organic carbons in the animal body. During digestion, digestible carbohydrates are ultimately broken down into glucose and used to provide energy through metabolic pathways. The excess sugars in the body are converted into glycogen and stored in the liver and muscles for later use. Glycogen stores are used to fuel prolonged exertions, such as long-distance running, and to provide energy during food shortage. Excess digestible carbohydrates are stored by mammals in order to survive famine and aid in mobility.

Another important requirement is that of nitrogen. Protein catabolism provides a source of organic nitrogen. Amino acids are the building blocks of proteins and protein breakdown provides amino acids that are used for cellular function. The carbon and nitrogen derived from these become the building block for nucleotides, nucleic acids, proteins, cells, and tissues. Excess nitrogen must be excreted, as it is toxic. Fats add flavor to food and promote a sense of satiety or fullness. Fatty foods are also significant sources of energy because one gram of fat contains nine calories. Fats are required in the diet to aid the absorption of fat-soluble vitamins and the production of fat-soluble hormones.

Essential Nutrients

While the animal body can synthesize many of the molecules required for function from the organic precursors, there are some nutrients that need to be consumed from food. These nutrients are termed essential nutrients: they must be eaten as the body cannot produce them.

Vitamins and minerals are substances found in the food we eat. Your body needs them to be able to work properly and for growth and development. Each vitamin has its own special role to play. For example, vitamin D (added to whole milk or naturally-occurring in sardines), helps make bones strong, while vitamin A (found in carrots) helps with night vision. Vitamins fall into two categories: fat soluble and water soluble. The fat-soluble vitamins dissolve in fat and can be stored in your body, whereas the water-soluble vitamins need to dissolve in water before your body can absorb them; therefore, the body cannot store them.

Fat-soluble vitamins are found primarily in foods that contain fat and oil, such as animal fats, vegetable oils, dairy foods, liver, and fatty fish. Your body needs these vitamins every day to enable it to work properly. However, you do not need to eat foods containing these every day. If your body does not need these vitamins immediately, they will be stored in the liver and fat tissues for future use. This means that stores can build up; if you have more than you need, fat soluble vitamins can become harmful. Some fat-soluble vitamins include vitamin A, vitamin K, vitamin D, and vitamin E. Unlike the other fat-soluble vitamins, vitamin D is difficult to obtain in adequate quantities in a normal diet; therefore, supplementation may be necessary.


Water-soluble vitamins are not stored in the body; therefore, you need to have them more frequently. If you have more then you need, the body rids itself of the extra vitamins during urination. Because the body does not store these vitamins, they are generally not harmful. Water-soluble vitamins are found in foods that include fruits, vegetables, and grains. Unlike fat-soluble vitamins, they can be destroyed by heat. This means that sometimes these vitamins can often be lost during cooking. This is why it is better to steam or grill these foods rather then boil them. Some water-soluble vitamins include vitamin B6, vitamin B12, vitamin C, biotin, folic acid, niacin, and riboflavin.



Sea buckthorn seed oil: Sea buckthorn seed oil contains many vital nutrients.

The omega-3 alpha-linolenic acid and the omega-6 linoleic acid are essential fatty acids needed to synthesize some membrane phospholipids. Many people take supplements to ensure they are obtaining all the essential fatty acids they need. Sea buckthorn contains many of these fatty acids and is also high in vitamins. Sea buckthorn can be used to treat acne and promote weight loss and wound healing.

Minerals are inorganic essential nutrients that must also be obtained from food. Among their many functions, minerals help in cell structure and regulation; they are also considered co-factors. In addition to vitamins and minerals, certain amino acids must also be procured from food and cannot be synthesized by the body. These amino acids are the “essential” amino acids. The human body can synthesize only 11 of the 20 required amino acids. The rest must be obtained from food.



Amino Acids: There are 20 known amino acids. Animals can make only 11, so the others must be obtained through the diet. Meats are the best source of amino acids, although some amino acids can also be obtained from vegetables and grains.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Digestive System. **Provided by**: sciencemasters Wikispace. **Located at**: http://sciencemasters.wikispaces.com/Digestive+System. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- macromolecule. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/macromolecule. **License**: ***CC BY-SA: Attribution-ShareAlike***
- digestion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/digestion. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alimentary canal. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/alimentary_canal. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and physiology of animals Summary of the main functions of the different regions of the gut. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...of_the_gut.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...e_34_00_01.jpg. **License**: ***CC BY: Attribution***
- Carbohydrate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Carbohydrate%23Nutrition. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glucose. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glucose. **License**: ***CC BY-SA: Attribution-ShareAlike***
- saccharide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/saccharide. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carbohydrate. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/carbohydrate. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and physiology of animals Summary of the main functions of the different regions of the gut. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...of_the_gut.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...e_34_00_01.jpg. **License**: ***CC BY: Attribution***
- Starchy-foods.. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Starchy-foods..jpg. **License**: ***Public Domain: No Known Copyright***
- Glucose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Glucose.PNG. **License**: ***Public Domain: No Known Copyright***
- Lipid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lipid%2...ion_and_health. **License**: ***CC BY-SA: Attribution-ShareAlike***
- trans fats. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/trans%20fats. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phospholipids. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/phospholipids. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fatty acid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/fatty_acid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and physiology of animals Summary of the main functions of the different regions of the gut. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...of_the_gut.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...e_34_00_01.jpg. **License**: ***CC BY: Attribution***
- Starchy-foods.. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Starchy-foods..jpg. **License**: ***Public Domain: No Known Copyright***
- Glucose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Glucose.PNG. **License**: ***Public Domain: No Known Copyright***
- Salmon. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Salmon.JPG. **License**: ***Public Domain: No Known Copyright***
- DHA Molecule. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...A_Molecule.gif. **License**: ***Public Domain: No Known Copyright***
- Protein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Protein%23Nutrition. **License**: ***CC BY-SA: Attribution-ShareAlike***
- High-protein diet. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/High-protein_diet. **License**: ***CC BY-SA: Attribution-ShareAlike***
- protein. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/protein. **License**: ***CC BY-SA: Attribution-ShareAlike***

- denaturation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/denaturation. **License**: *CC BY-SA: Attribution-ShareAlike*
- amino acid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amino_acid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals Summary of the main functions of the different regions of the gut. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...of_the_gut.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Introduction. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...e_34_00_01.jpg. **License**: *CC BY: Attribution*
- Starchy-foods.. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Starchy-foods..jpg. **License**: *Public Domain: No Known Copyright*
- Glucose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Glucose.PNG. **License**: *Public Domain: No Known Copyright*
- Salmon. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Salmon.JPG. **License**: *Public Domain: No Known Copyright*
- DHA Molecule. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...A_Molecule.gif. **License**: *Public Domain: No Known Copyright*
- N2 steak. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:N2_steak.jpg. **License**: *CC BY: Attribution*
- Cystine 3D ball. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ne_3D_ball.png. **License**: *Public Domain: No Known Copyright*
- Vitamins and Minerals.. **Provided by**: healthyeatingfoods Wikispace. **Located at**: http://healthyeatingfoods.wikispaces...s+and+Minerals.. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44740/latest...ol11448/latest. **License**: *CC BY: Attribution*
- catabolism. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/catabolism. **License**: *CC BY-SA: Attribution-ShareAlike*
- vitamin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vitamin. **License**: *CC BY-SA: Attribution-ShareAlike*
- nutrient. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nutrient. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals Summary of the main functions of the different regions of the gut. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Summary_of_the_main_functions_of_the_different_regions_of_the_gut.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Introduction. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44734/latest...e_34_00_01.jpg. **License**: *CC BY: Attribution*
- Starchy-foods.. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Starchy-foods..jpg. **License**: *Public Domain: No Known Copyright*
- Glucose. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Glucose.PNG. **License**: *Public Domain: No Known Copyright*
- Salmon. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Salmon.JPG. **License**: *Public Domain: No Known Copyright*
- DHA Molecule. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...A_Molecule.gif. **License**: *Public Domain: No Known Copyright*
- N2 steak. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:N2_steak.jpg. **License**: *CC BY: Attribution*
- Cystine 3D ball. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ne_3D_ball.png. **License**: *Public Domain: No Known Copyright*
- OpenStax College, Nutrition and Energy Production. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44740/latest...e_34_02_01.jpg. **License**: *CC BY: Attribution*
- Sea buckthorn seed oil gel. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...d_oil_gel.JPEG. **License**: *CC BY: Attribution*
- Amino Acids. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...mino_Acids.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 23.1E: Food Requirements and Essential Nutrients is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.2: CARBOHYDRATES

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.2A: Carbohydrate Molecules

23.2B: Importance of Carbohydrates

This page titled 23.2: Carbohydrates is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.2A: Carbohydrate Molecules

Carbohydrates are essential macromolecules that are classified into three subtypes: monosaccharides, disaccharides, and polysaccharides.

Learning Objectives

- Describe the structure of mono-, di-, and poly-saccharides

Key Points

- Monosaccharides are simple sugars made up of three to seven carbons, and they can exist as a linear chain or as ring-shaped molecules.
- Glucose, galactose, and fructose are monosaccharide isomers, which means they all have the same chemical formula but differ structurally and chemically.
- Disaccharides form when two monosaccharides undergo a dehydration reaction (a condensation reaction); they are held together by a covalent bond.
- Sucrose (table sugar) is the most common disaccharide, which is composed of the monomers glucose and fructose.
- A polysaccharide is a long chain of monosaccharides linked by glycosidic bonds; the chain may be branched or unbranched and can contain many types of monosaccharides.


Key Terms

- **isomer**: Any of two or more compounds with the same molecular formula but with different structure.
- **dehydration reaction**: A chemical reaction in which two molecules are covalently linked in a reaction that generates H2O as a second product.
- **biopolymer**: Any macromolecule of a living organism that is formed from the polymerization of smaller entities; a polymer that occurs in a living organism or results from life.

Carbohydrates can be represented by the stoichiometric formula $(CH_2O)_n$, where n is the number of carbons in the molecule. Therefore, the ratio of carbon to hydrogen to oxygen is 1:2:1 in carbohydrate molecules. The origin of the term “carbohydrate” is based on its components: carbon (“carbo”) and water (“hydrate”). Carbohydrates are classified into three subtypes: monosaccharides, disaccharides, and polysaccharides.

Monosaccharides

Monosaccharides (mono- = “one”; sacchar- = “sweet”) are simple sugars. In monosaccharides, the number of carbons usually ranges from three to seven. If the sugar has an aldehyde group (the functional group with the structure R-CHO), it is known as an aldose, and if it has a ketone group (the functional group with the structure RC(=O)R’), it is known as a ketose. Depending on the number of carbons in the sugar, they also may be known as trioses (three carbons), pentoses (five carbons), and or hexoses (six carbons). Monosaccharides can exist as a linear chain or as ring-shaped molecules; in aqueous solutions they are usually found in ring forms.



Monosaccharides: Monosaccharides are classified based on the position of their carbonyl group and the number of carbons in the backbone. Aldoses have a carbonyl group (indicated in green) at the end of the carbon chain, and ketoses have a carbonyl group in the middle of the carbon chain. Trioses, pentoses, and hexoses have three, five, and six carbon backbones, respectively.


Common Monosaccharides

Glucose ($C_6H_{12}O_6$) is a common monosaccharide and an important source of energy. During cellular respiration, energy is released from glucose and that energy is used to help make adenosine triphosphate (ATP). Plants synthesize glucose using carbon dioxide and water, and glucose, in turn, is used for energy requirements for the plant.

Galactose (a milk sugar) and fructose (found in fruit) are other common monosaccharides. Although glucose, galactose, and fructose all have the same chemical formula ($C_6H_{12}O_6$), they differ structurally and stereochemically. This makes them different molecules despite sharing the same atoms in the same proportions, and they are all isomers of one another, or isomeric monosaccharides. Glucose and galactose are aldoses, and fructose is a ketose.

Disaccharides

Disaccharides (di- = “two”) form when two monosaccharides undergo a dehydration reaction (also known as a condensation reaction or dehydration synthesis). During this process, the hydroxyl group of one monosaccharide combines with the hydrogen of another monosaccharide, releasing a molecule of water and forming a covalent bond. A covalent bond formed between a carbohydrate molecule and another molecule (in this case, between two monosaccharides) is known as a glycosidic bond. Glycosidic bonds (also called glycosidic linkages) can be of the alpha or the beta type.



Disaccharides: Sucrose is formed when a monomer of glucose and a monomer of fructose are joined in a dehydration reaction to form a glycosidic bond. In the process, a water molecule is lost. By convention, the carbon atoms in a monosaccharide are numbered from the terminal carbon closest to the carbonyl group. In sucrose, a glycosidic linkage is formed between carbon 1 in glucose and carbon 2 in fructose.

Common Disaccharides

Common disaccharides include lactose, maltose, and sucrose. Lactose is a disaccharide consisting of the monomers glucose and galactose. It is found naturally in milk. Maltose, or malt sugar, is a disaccharide formed by a dehydration reaction between two glucose molecules. The most common disaccharide is sucrose, or table sugar, which is composed of the monomers glucose and fructose.

Polysaccharides


A long chain of monosaccharides linked by glycosidic bonds is known as a polysaccharide (poly- = “many”). The chain may be branched or unbranched, and it may contain different types of monosaccharides. Starch, glycogen, cellulose, and chitin are primary examples of polysaccharides.

Plants are able to synthesize glucose, and the excess glucose is stored as starch in different plant parts, including roots and seeds. Starch is the stored form of sugars in plants and is made up of glucose monomers that are joined by α1-4 or 1-6 glycosidic bonds. The starch in the seeds provides food for the embryo as it germinates while the starch that is consumed by humans is broken down by enzymes into smaller molecules, such as maltose and glucose. The cells can then absorb the glucose.

Common Polysaccharides

Glycogen is the storage form of glucose in humans and other vertebrates. It is made up of monomers of glucose. Glycogen is the animal equivalent of starch and is a highly branched molecule usually stored in liver and muscle cells. Whenever blood glucose levels decrease, glycogen is broken down to release glucose in a process known as glycogenolysis.

Cellulose is the most abundant natural biopolymer. The cell wall of plants is mostly made of cellulose and provides structural support to the cell. Cellulose is made up of glucose monomers that are linked by β 1-4 glycosidic bonds. Every other glucose monomer in cellulose is flipped over, and the monomers are packed tightly as extended long chains. This gives cellulose its rigidity and high tensile strength—which is so important to plant cells.



Polysaccharides: In cellulose, glucose monomers are linked in unbranched chains by β 1-4 glycosidic linkages. Because of the way the glucose subunits are joined, every glucose monomer is flipped relative to the next one resulting in a linear, fibrous structure.

Carbohydrate Function

Carbohydrates serve various functions in different animals. Arthropods have an outer skeleton, the exoskeleton, which protects their internal body parts. This exoskeleton is made of chitin, which is a polysaccharide-containing nitrogen. It is made of repeating units of N-acetyl-β-d-glucosamine, a modified sugar. Chitin is also a major component of fungal cell walls.

This page titled 23.2A: Carbohydrate Molecules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.2B: Importance of Carbohydrates

Carbohydrates are a major class of biological macromolecules that are an essential part of our diet and provide energy to the body.

Learning Objectives

- Describe the benefits provided to organisms by carbohydrates

Key Points


- Carbohydrates provide energy to the body, particularly through glucose, a simple sugar that is found in many basic foods.
- Carbohydrates contain soluble and insoluble elements; the insoluble part is known as fiber, which promotes regular bowel movement, regulates the rate of consumption of blood glucose, and also helps to remove excess cholesterol from the body.
- As an immediate source of energy, glucose is broken down during the process of cellular respiration, which produces ATP, the energy currency of the cell.
- Since carbohydrates are an important part of the human nutrition, eliminating them from the diet is not the best way to lose weight.

Key Terms

- **carbohydrate**: A sugar, starch, or cellulose that is a food source of energy for an animal or plant; a saccharide.
- **glucose**: a simple monosaccharide (sugar) with a molecular formula of C6H12O6; it is a principal source of energy for cellular metabolism
- **ATP**: A nucleotide that occurs in muscle tissue, and is used as a source of energy in cellular reactions, and in the synthesis of nucleic acids. ATP is the abbreviation for adenosine triphosphate.

Benefits of Carbohydrates

Biological macromolecules are large molecules that are necessary for life and are built from smaller organic molecules. One major class of biological macromolecules are carbohydrates, which are further divided into three subtypes: monosaccharides, disaccharides, and polysaccharides. Carbohydrates are, in fact, an essential part of our diet; grains, fruits, and vegetables are all natural sources of carbohydrates. Importantly, carbohydrates provide energy to the body, particularly through glucose, a simple sugar that is a component of starch and an ingredient in many basic foods.



Carbohydrates: Carbohydrates are biological macromolecules that are further divided into three subtypes: monosaccharides, disaccharides, and polysaccharides. Like all macromolecules, carbohydrates are necessary for life and are built from smaller organic molecules.

Carbohydrates in Nutrition

Carbohydrates have been a controversial topic within the diet world. People trying to lose weight often avoid carbs, and some diets completely forbid carbohydrate consumption, claiming that a low-carb diet helps people to lose weight faster. However, carbohydrates have been an important part of the human diet for thousands of years; artifacts from ancient civilizations show the presence of wheat, rice, and corn in our ancestors' storage areas.

Carbohydrates should be supplemented with proteins, vitamins, and fats to be parts of a well-balanced diet. Calorie-wise, a gram of carbohydrate provides 4.3 Kcal. In comparison, fats provide 9 Kcal/g, a less desirable ratio. Carbohydrates contain soluble and insoluble elements; the insoluble part is known as fiber, which is mostly cellulose. Fiber has many uses; it promotes regular bowel movement by adding bulk, and it regulates the rate of consumption of blood glucose. Fiber also helps to remove excess cholesterol from the body. Fiber binds and attaches to the cholesterol in the small intestine and prevents the cholesterol particles from entering the bloodstream. Then cholesterol exits the body via the feces. Fiber-rich diets also have a protective role in reducing the occurrence of colon cancer. In addition, a meal containing whole grains and vegetables gives a feeling of fullness. As an immediate source of energy, glucose is broken down during the process of cellular respiration, which produces adenosine triphosphate (ATP), the energy currency of the cell. Without the consumption of carbohydrates, the availability of "instant energy" would be reduced. Eliminating carbohydrates from the diet is not the best way to lose weight. A low-calorie diet that is rich in whole grains, fruits, vegetables, and lean meat, together with plenty of exercise and plenty of water, is the more sensible way to lose weight.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- dehydration reaction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/dehydration_reaction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- isomer. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/isomer**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- biopolymer. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/biopolymer**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_07.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_04.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- ATP. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ATP**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carbohydrate. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/carbohydrate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glucose. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glucose**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_07.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_04.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Carbohydrates. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44400/latest...e_03_02_01.jpg**. **License**: ***CC BY: Attribution***
- Wikimedia. **Provided by**: Wikimedia. **Located at**: **www.wikimedia.org**. **License**:

This page titled 23.2B: Importance of Carbohydrates is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.3: LIPIDS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.3A: Lipid Molecules

23.3B: Phospholipids

23.3C: Steroids

This page titled 23.3: Lipids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.3A: Lipid Molecules

Fats and oils, which may be saturated or unsaturated, can be unhealthy but also serve important functions for plants and animals.

Learning Objectives

- Differentiate between saturated and unsaturated fatty acids

Key Points

- Fats provide energy, insulation, and storage of fatty acids for many organisms.
- Fats may be saturated (having single bonds) or unsaturated (having double bonds).
- Unsaturated fats may be cis (hydrogens in same plane) or trans (hydrogens in two different planes).
- Olive oil, a monounsaturated fat, has a single double bond whereas canola oil, a polyunsaturated fat, has more than one double bond.
- Omega-3 fatty acid and omega-6 fatty acid are essential for human biological processes, but they must be ingested in the diet because they cannot be synthesized.


Key Terms

- **hydrogenation**: The chemical reaction of hydrogen with another substance, especially with an unsaturated organic compound, and usually under the influence of temperature, pressure and catalysts.
- **ester**: Compound most often formed by the condensation of an alcohol and an acid, by removing water. It contains the functional group carbon-oxygen double bond joined via carbon to another oxygen atom.
- **carboxyl**: A univalent functional group consisting of a carbonyl and a hydroxyl functional group (-CO.OH); characteristic of carboxylic acids.

Fats have important functions, and many vitamins are fat soluble. Fats serve as a long-term storage form of fatty acids and act as a source of energy. They also provide insulation for the body.

Glycerol and Fatty Acids

A fat molecule consists of two main components: glycerol and fatty acids. Glycerol is an alcohol with three carbons, five hydrogens, and three hydroxyl (OH) groups. Fatty acids have a long chain of hydrocarbons with a carboxyl group attached and may have 4-36 carbons; however, most of them have 12-18. In a fat molecule, the fatty acids are attached to each of the three carbons of the glycerol molecule with an ester bond through the oxygen atom. During the ester bond formation, three molecules are released. Since fats consist of three fatty acids and a glycerol, they are also called triacylglycerols or triglycerides.




Triacylglycerols: Triacylglycerol is formed by the joining of three fatty acids to a glycerol backbone in a dehydration reaction. Three molecules of water are released in the process.

Saturated vs. Unsaturated Fatty Acids

Fatty acids may be saturated or unsaturated. In a fatty acid chain, if there are only single bonds between neighboring carbons in the hydrocarbon chain, the fatty acid is said to be saturated. Saturated fatty acids are saturated with hydrogen since single bonds increase the number of hydrogens on each carbon. Stearic acid and palmitic acid, which are commonly found in meat, are examples of saturated fats.

When the hydrocarbon chain contains a double bond, the fatty acid is said to be unsaturated. Oleic acid is an example of an unsaturated fatty acid. Most unsaturated fats are liquid at room temperature and are called oils. If there is only one double bond in the molecule, then it is known as a monounsaturated fat; e.g. olive oil. If there is more than one double bond, then it is known as a polyunsaturated fat; e.g. canola oil. Unsaturated fats help to lower blood cholesterol levels whereas saturated fats contribute to plaque formation in the arteries.

Unsaturated fats or oils are usually of plant origin and contain cis unsaturated fatty acids. Cis and trans indicate the configuration of the molecule around the double bond. If hydrogens are present in the same plane, it is referred to as a cis fat; if the hydrogen atoms are on two different planes, it is referred to as a trans fat. The cis double bond causes a bend or a “kink” that prevents the fatty acids from packing tightly, keeping them liquid at room temperature.




Fatty Acids: Saturated fatty acids have hydrocarbon chains connected by single bonds only. Unsaturated fatty acids have one or more double bonds. Each double bond may be in a cis or trans configuration. In the cis configuration, both hydrogens are on the same side of the hydrocarbon chain. In the trans configuration, the hydrogens are on opposite sides. A cis double bond causes a kink in the chain.

Trans Fats

In the food industry, oils are artificially hydrogenated to make them semi-solid and of a consistency desirable for many processed food products. During this hydrogenation process, gas is bubbled through oils to solidify them, and the double bonds of the cis-conformation in the hydrocarbon chain may be converted to double bonds in the trans-conformation.

Margarine, some types of peanut butter, and shortening are examples of artificially-hydrogenated trans fats. Recent studies have shown that an increase in trans fats in the human diet may lead to an increase in levels of low-density lipoproteins (LDL), or “bad” cholesterol, which in turn may lead to plaque deposition in the arteries, resulting in heart disease. Many fast food restaurants have recently banned the use of trans fats, and food labels are required to display the trans fat content.

Essential Fatty Acids



Omega Fatty Acids: Alpha-linolenic acid is an example of an omega-3 fatty acid. It has three cis double bonds and, as a result, a curved shape. For clarity, the carbons are not shown. Each singly bonded carbon has two hydrogens associated with it, also not shown.

Essential fatty acids are fatty acids required for biological processes, but not synthesized by the human body. Consequently, they have to be supplemented through ingestion via the diet and are nutritionally very important. Omega-3 fatty acid, or alpha-linoleic acid (ALA), falls into this category and is one of only two fatty acids known to be essential for humans (the other being omega-6 fatty acid, or linoleic acid). These polyunsaturated fatty acids are called omega-3 because the third carbon from the end of the hydrocarbon chain is connected to its neighboring carbon by a double bond. Salmon, trout, and tuna are good sources of omega-3 fatty acids.

Research indicates that omega-3 fatty acids reduce the risk of sudden death from heart attacks, reduce triglycerides in the blood, lower blood pressure, and prevent thrombosis by inhibiting blood clotting. They also reduce inflammation and may help reduce the risk of some cancers in animals.

This page titled 23.3A: Lipid Molecules is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.3B: Phospholipids

Phospholipids are amphipathic molecules that make up the bilayer of the plasma membrane and keep the membrane fluid.

Learning Objectives

- Describe phospholipids and their role in cells


Key Points

- Phospholipids consist of a glycerol molecule, two fatty acids, and a phosphate group that is modified by an alcohol.
- The phosphate group is the negatively-charged polar head, which is hydrophilic.
- The fatty acid chains are the uncharged, nonpolar tails, which are hydrophobic.
- Since the tails are hydrophobic, they face the inside, away from the water and meet in the inner region of the membrane.
- Since the heads are hydrophilic, they face outward and are attracted to the intracellular and extracellular fluid.
- If phospholipids are placed in water, they form into micelles, which are lipid molecules that arrange themselves in a spherical form in aqueous solutions.

Key Terms

- **micelle**: Lipid molecules that arrange themselves in a spherical form in aqueous solutions.
- **amphipathic**: Describing a molecule, such as a detergent, which has both hydrophobic and hydrophilic groups.

Defining Characteristics of Phospholipids



Phospholipid Molecule: A phospholipid is a molecule with two fatty acids and a modified phosphate group attached to a glycerol backbone. The phosphate may be modified by the addition of charged or polar chemical groups. Two chemical groups that may modify the phosphate, choline and serine, are shown here. Both choline and serine attach to the phosphate group at the position labeled R via the hydroxyl group indicated in green.

Phospholipids are major components of the plasma membrane, the outermost layer of animal cells. Like fats, they are composed of fatty acid chains attached to a glycerol backbone. Unlike triglycerides, which have three fatty acids, phospholipids have two


fatty acids that help form a diacylglycerol. The third carbon of the glycerol backbone is also occupied by a modified phosphate group. However, just a phosphate group attached to a diacylglycerol does not qualify as a phospholipid. This would be considered a phosphatidate (diacylglycerol 3-phosphate), the precursor to phospholipids. To qualify as a phospholipid, the phosphate group should be modified by an alcohol. Phosphatidylcholine and phosphatidylserine are examples of two important phospholipids that are found in plasma membranes.

Structure of a Phospholipid Molecule

A phospholipid is an amphipathic molecule which means it has both a hydrophobic and a hydrophilic component. A single phospholipid molecule has a phosphate group on one end, called the “head,” and two side-by-side chains of fatty acids that make up the lipid “tails.” The phosphate group is negatively charged, making the head polar and hydrophilic, or “water loving.” The phosphate heads are thus attracted to the water molecules in their environment.

The lipid tails, on the other hand, are uncharged, nonpolar, and hydrophobic, or “water fearing.” A hydrophobic molecule repels and is repelled by water. Some lipid tails consist of saturated fatty acids and some contain unsaturated fatty acids. This combination adds to the fluidity of the tails that are constantly in motion.

Phospholipids and Biological Membranes




Phospholipid Bilayer: The phospholipid bilayer consists of two adjacent sheets of phospholipids, arranged tail to tail. The hydrophobic tails associate with one another, forming the interior of the membrane. The polar heads contact the fluid inside and outside of the cell.

The cell membrane consists of two adjacent layers of phospholipids, which form a bilayer. The fatty acid tails of phospholipids face inside, away from water, whereas the phosphate heads face the outward aqueous side. Since the heads face outward, one layer is exposed to the interior of the cell and one layer is exposed to the exterior. As the phosphate groups are polar and hydrophilic, they are attracted to water in the intracellular fluid.

Because of the phospholipds’ chemical and physical characteristics, the lipid bilayer acts as a semipermeable membrane; only lipophilic solutes can easily pass the phospholipd bilayer. As a result, there are two distinct aqueous compartments on each side of the membrane. This separation is essential for many biological functions, including cell communication and metabolism.

Membrane Fluidity



Micelles: An example of micelles in water.

A cell's plasma membrane contain proteins and other lipids (such as cholesterol) within the phospholipid bilayer. Biological membranes remain fluid because of the unsaturated hydrophobic tails, which prevent phospholipid molecules from packing together and forming a solid.

If a drop of phospholipids is placed in water, the phospholipids spontaneously form a structure known as a micelle, with their hydrophilic heads oriented toward the water. Micelles are lipid molecules that arrange themselves in a spherical form in aqueous solution. The formation of a micelle is a response to the amphipathic nature of fatty acids, meaning that they contain both hydrophilic and hydrophobic regions.

This page titled 23.3B: Phospholipids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.3C: Steroids

Steroids, like cholesterol, play roles in reproduction, absorption, metabolism regulation, and brain activity.

Learning Objectives

- Describe some functions of steroids

Key Points


- Steroids are lipids because they are hydrophobic and insoluble in water, but they do not resemble lipids since they have a structure composed of four fused rings.
- Cholesterol is the most common steroid and is the precursor to vitamin D, testosterone, estrogen, progesterone, aldosterone, cortisol, and bile salts.
- Cholesterol is a component of the phospholipid bilayer and plays a role in the structure and function of membranes.
- Steroids are found in the brain and alter electrical activity in the brain.
- Because they can tone down receptors that communicate messages from neurotransmitters, steroids are often used in anesthetic medicines.

Key Terms


- **neurotransmitter**: any substance, such as acetylcholine or dopamine, responsible for sending nerve signals across a synapse between two neurons
- **osmoregulation**: the homeostatic regulation of osmotic pressure in the body in order to maintain a constant water content
- **hormone**: any substance produced by one tissue and conveyed by the bloodstream to another to affect physiological activity

Structure of Steroid Molecules

Unlike phospholipids and fats, steroids have a fused ring structure. Although they do not resemble the other lipids, they are grouped with them because they are also hydrophobic and insoluble in water. All steroids have four linked carbon rings, and many of them, like cholesterol, have a short tail. Many steroids also have the –OH functional group, and these steroids are classified as alcohols called sterols.



Cholesterol



Cortisol

Steroid Structures: Steroids, such as cholesterol and cortisol, are composed of four fused hydrocarbon rings.

Cholesterol

Cholesterol is the most common steroid and is mainly synthesized in the liver; it is the precursor to vitamin D. Cholesterol is also a precursor to many important steroid hormones like estrogen, testosterone, and progesterone, which are secreted by the gonads and endocrine glands. Therefore, steroids play very important roles in the body's reproductive system. Cholesterol also plays a role in synthesizing the steroid hormones aldosterone, which is used for osmoregulation, and cortisol, which plays a role in metabolism.

Cholesterol is also the precursor to bile salts, which help in the emulsification of fats and their absorption by cells. It is a component of the plasma membrane of animal cells and the phospholipid bilayer. Being the outermost structure in animal cells, the plasma membrane is responsible for the transport of materials and cellular recognition; and it is involved in cell-to-cell communication. Thus, steroids also play an important role in the structure and function of membranes.

It has also been discovered that steroids can be active in the brain where they affect the nervous system, These neurosteroids alter electrical activity in the brain. They can either activate or tone down receptors that communicate messages from neurotransmitters. Since these neurosteroids can tone down receptors and decrease brain activity, steroids are often used in anesthetic medicines.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- hydrogenation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hydrogenation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ester. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ester. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carboxyl. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/carboxyl. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_02.jpg. **License**: ***CC BY: Attribution***

- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_05.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- Structural Biochemistry/Lipids/Micelles. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structu...ipids/Micelles. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/definition/micelle. **License**: *CC BY-SA: Attribution-ShareAlike*
- amphipathic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amphipathic. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_05.jpg. **License**: *CC BY: Attribution*
- Structural Biochemistry/Lipids/Micelles. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structu...ipids/Micelles. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest..._05_01_03a.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Hormones. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46667/latest/. **License**: *CC BY: Attribution*
- Metabolomics/Metabolites/Lipids/Steroids. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Metabol...ipids/Steroids. **License**: *CC BY-SA: Attribution-ShareAlike*
- hormone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hormone. **License**: *CC BY-SA: Attribution-ShareAlike*
- osmoregulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osmoregulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- neurotransmitter. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurotransmitter. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_05.jpg. **License**: *CC BY: Attribution*
- Structural Biochemistry/Lipids/Micelles. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structural_Biochemistry/Lipids/Micelles. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest..._05_01_03a.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Cell Membrane. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46021/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Lipids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44401/latest...e_03_03_10.jpg. **License**: *CC BY: Attribution*

This page titled 23.3C: Steroids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.4: PROTEINS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.4A: Types and Functions of Proteins

23.4B: Amino Acids

23.4C: Protein Structure

This page titled 23.4: Proteins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.4A: Types and Functions of Proteins

Proteins perform many essential physiological functions, including catalyzing biochemical reactions.

Learning Objectives

- Differentiate among the types and functions of proteins

Key Points

- Proteins are essential for the main physiological processes of life and perform functions in every system of the human body.
- A protein's shape determines its function.
- Proteins are composed of amino acid subunits that form polypeptide chains.
- Enzymes catalyze biochemical reactions by speeding up chemical reactions, and can either break down their substrate or build larger molecules from their substrate.
- The shape of an enzyme's active site matches the shape of the substrate.
- Hormones are a type of protein used for cell signaling and communication.

Key Terms


- **amino acid**: Any of 20 naturally occurring α-amino acids (having the amino, and carboxylic acid groups on the same carbon atom), and a variety of side chains, that combine, via peptide bonds, to form proteins.
- **polypeptide**: Any polymer of (same or different) amino acids joined via peptide bonds.
- **catalyze**: To accelerate a process.

EXAMPLES

Salivary amylase is an enzyme in the mouth that breaks down starch (a long carbohydrate chain) into amylose (a short chain of glucose molecules). The longer you chew a cracker, the sweeter the cracker will taste because your taste buds are receptive to the glucose molecules exposed by the amylase.

Proteins perform essential functions throughout the systems of the human body. These long chains of amino acids are critically important for:

- catalyzing chemical reactions
- synthesizing and repairing DNA
- transporting materials across the cell
- receiving and sending chemical signals
- responding to stimuli
- providing structural support



Human Hemoglobin: Structure of human hemoglobin. The proteins' α and β subunits are in red and blue, and the iron-containing heme groups in green. From the protein data base.

Proteins (a polymer) are macromolecules composed of amino acid subunits (the monomers). These amino acids are covalently attached to one another to form long linear chains called polypeptides, which then fold into a specific three-dimensional shape. Sometimes these folded polypeptide chains are functional by themselves. Other times they combine with additional polypeptide chains to form the final protein structure. Sometimes non-polypeptide groups are also required in the final protein. For instance, the blood protein hemogobin is made up of four polypeptide chains, each of which also contains a heme molecule, which is ring structure with an iron atom in its center.


Proteins have different shapes and molecular weights, depending on the amino acid sequence. For example, hemoglobin is a globular protein, which means it folds into a compact globe-like structure, but collagen, found in our skin, is a fibrous protein, which means it folds into a long extended fiber-like chain. You probably look similar to your family members because you share similar proteins, but you look different from strangers because the proteins in your eyes, hair, and the rest of your body are different.

Because form determines function, any slight change to a protein's shape may cause the protein to become dysfunctional. Small changes in the amino acid sequence of a protein can cause devastating genetic diseases such as Huntington's disease or sickle cell anemia.

Enzymes

Enzymes are proteins that catalyze biochemical reactions, which otherwise would not take place. These enzymes are essential for chemical processes like digestion and cellular metabolism. Without enzymes, most physiological processes would proceed so slowly (or not at all) that life could not exist.

Because form determines function, each enzyme is specific to its substrates. The substrates are the reactants that undergo the chemical reaction catalyzed by the enzyme. The location where substrates bind to or interact with the enzyme is known as the active site, because that is the site where the chemistry occurs. When the substrate binds to its active site at the enzyme, the enzyme may help in its breakdown, rearrangement, or synthesis. By placing the substrate into a specific shape and microenvironment in the active site, the enzyme encourages the chemical reaction to occur. There are two basic classes of enzymes:



Enzyme reaction: A catabolic enzyme reaction showing the substrate matching the exact shape of the active site.

- Catabolic enzymes: enzymes that break down their substrate
- Anabolic enzymes: enzymes that build more complex molecules from their substrates


Enzymes are essential for digestion: the process of breaking larger food molecules down into subunits small enough to diffuse through a cell membrane and to be used by the cell. These enzymes include amylase, which catalyzes the digestion carbohydrates in the mouth and small intestine; pepsin, which catalyzes the digestion of proteins in the stomach; lipase, which catalyzes reactions need to emulsify fats in the small intestine; and trypsin, which catalyzes the further digestion of proteins in the small intestine.

Enzymes are also essential for biosynthesis: the process of making new, complex molecules from the smaller subunits that are provided to or generated by the cell. These biosynthetic enzymes include DNA Polymerase, which catalyzes the synthesis of new strands of the genetic material before cell division; fatty acid synthetase, which the synthesis of new fatty acids for fat or membrane lipid formation; and components of the ribosome, which catalyzes the formation of new polypeptides from amino acid monomers.

Hormones

Some proteins function as chemical-signaling molecules called hormones. These proteins are secreted by endocrine cells that act to control or regulate specific physiological processes, which include growth, development, metabolism, and reproduction. For example, insulin is a protein hormone that helps to regulate blood glucose levels. Other proteins act as receptors to detect the concentrations of chemicals and send signals to respond. Some types of hormones, such as estrogen and testosterone, are lipid steroids, not proteins.

Other Protein Functions



Tubulin: The structural protein tubulin stained red in mouse cells.

Proteins perform essential functions throughout the systems of the human body. In the respiratory system, hemoglobin (composed of four protein subunits) transports oxygen for use in cellular metabolism. Additional proteins in the blood plasma and lymph carry nutrients and metabolic waste products throughout the body. The proteins actin and tubulin form cellular structures, while keratin forms the structural support for the dead cells that become fingernails and hair. Antibodies, also called immunoglobins, help recognize and destroy foreign pathogens in the immune system. Actin and myosin allow muscles to contract, while albumin nourishes the early development of an embryo or a seedling.

This page titled 23.4A: Types and Functions of Proteins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.4B: Amino Acids

An amino acid contains an amino group, a carboxyl group, and an R group, and it combines with other amino acids to form polypeptide chains.

Learning Objectives

- Describe the structure of an amino acid and the features that confer its specific properties

Key Points


- Each amino acid contains a central C atom, an amino group (NH2), a carboxyl group (COOH), and a specific R group.
- The R group determines the characteristics (size, polarity, and pH) for each type of amino acid.
- Peptide bonds form between the carboxyl group of one amino acid and the amino group of another through dehydration synthesis.
- A chain of amino acids is a polypeptide.

Key Terms

- **amino acid**: Any of 20 naturally occurring α-amino acids (having the amino, and carboxylic acid groups on the same carbon atom), and a variety of side chains, that combine, via peptide bonds, to form proteins.
- **R group**: The R group is a side chain specific to each amino acid that confers particular chemical properties to that amino acid.
- **polypeptide**: Any polymer of (same or different) amino acids joined via peptide bonds.

Structure of an Amino Acid


Amino acids are the monomers that make up proteins. Each amino acid has the same fundamental structure, which consists of a central carbon atom, also known as the alpha (α) carbon, bonded to an amino group (NH_2), a carboxyl group (COOH), and to a hydrogen atom. In the aqueous environment of the cell, the both the amino group and the carboxyl group are ionized under physiological conditions, and so have the structures $-NH_3^+$ and $-COO^-$, respectively. Every amino acid also has another atom or group of atoms bonded to the central atom known as the R group. This R group, or side chain, gives each amino acid proteins specific characteristics, including size, polarity, and pH.



Amino acid structure: Amino acids have a central asymmetric carbon to which an amino group, a carboxyl group, a hydrogen atom, and a side chain (R group) are attached. This amino acid is unionized, but if it were placed in water at pH 7, its amino group would pick up another hydrogen and a positive charge, and the hydroxyl in its carboxyl group would lose and a hydrogen and gain a negative charge.

Types of Amino Acids

The name “amino acid” is derived from the amino group and carboxyl-acid-group in their basic structure. There are 21 amino acids present in proteins, each with a specific R group or side chain. Ten of these are considered essential amino acids in humans because the human body cannot produce them and they must be obtained from the diet. All organisms have different essential amino acids based on their physiology.



Types of amino acids: There are 21 common amino acids commonly found in proteins, each with a different R group (variant group) that determines its chemical nature. The 21st amino acid, not shown here, is selenocysteine, with an R group of -CH_2-SeH.


Characteristics of Amino Acids

Which categories of amino acid would you expect to find on the surface of a soluble protein, and which would you expect to find in the interior? What distribution of amino acids would you expect to find in a protein embedded in a lipid bilayer?

The chemical composition of the side chain determines the characteristics of the amino acid. Amino acids such as valine, methionine, and alanine are nonpolar (hydrophobic), while amino acids such as serine, threonine, and cysteine are polar (hydrophilic). The side chains of lysine and arginine are positively charged so these amino acids are also known as basic (high pH) amino acids. Proline is an exception to the standard structure of an amino acid because its R group is linked to the amino group, forming a ring-like structure.

Amino acids are represented by a single upper case letter or a three-letter abbreviation. For example, valine is known by the letter V or the three-letter symbol val.

Peptide Bonds



Peptide bond formation: Peptide bond formation is a dehydration synthesis reaction. The carboxyl group of one amino acid is linked to the amino group of the incoming amino acid. In the process, a molecule of water is released.

The sequence and the number of amino acids ultimately determine the protein's shape, size, and function. Each amino acid is attached to another amino acid by a covalent bond, known as a peptide bond. When two amino acids are covalently attached by a peptide bond, the carboxyl group of one amino acid and the amino group of the incoming amino acid combine and release a molecule of water. Any reaction that combines two monomers in a reaction that generates H_2O as one of the products is known as a dehydration reaction, so peptide bond formation is an example of a dehydration reaction.

Polypeptide Chains

The resulting chain of amino acids is called a polypeptide chain. Each polypeptide has a free amino group at one end. This end is called the N terminal, or the amino terminal, and the other end has a free carboxyl group, also known as the C or carboxyl terminal. When reading or reporting the amino acid sequence of a protein or polypeptide, the convention is to use the N-to-C direction. That is, the first amino acid in the sequence is assumed to the be one at the N terminal and the last amino acid is assumed to be the one at the C terminal.

Although the terms polypeptide and protein are sometimes used interchangeably, a polypeptide is technically any polymer of amino acids, whereas the term protein is used for a polypeptide or polypeptides that have folded properly, combined with any additional components needed for proper functioning, and is now functional.

This page titled 23.4B: Amino Acids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.4C: Protein Structure

Each successive level of protein folding ultimately contributes to its shape and therefore its function.

Learning Objectives

- Summarize the four levels of protein structure

Key Points

- Protein structure depends on its amino acid sequence and local, low-energy chemical bonds between atoms in both the polypeptide backbone and in amino acid side chains.
- Protein structure plays a key role in its function; if a protein loses its shape at any structural level, it may no longer be functional.
- Primary structure is the amino acid sequence.
- Secondary structure is local interactions between stretches of a polypeptide chain and includes α-helix and β-pleated sheet structures.
- Tertiary structure is the overall the three-dimension folding driven largely by interactions between R groups.
- Quarternary structures is the orientation and arrangement of subunits in a multi-subunit protein.


Key Terms

- **antiparallel**: The nature of the opposite orientations of the two strands of DNA or two beta strands that comprise a protein's secondary structure
- **disulfide bond**: A bond, consisting of a covalent bond between two sulfur atoms, formed by the reaction of two thiol groups, especially between the thiol groups of two proteins
- **β-pleated sheet**: secondary structure of proteins where N-H groups in the backbone of one fully-extended strand establish hydrogen bonds with C=O groups in the backbone of an adjacent fully-extended strand
- **α-helix**: secondary structure of proteins where every backbone N-H creates a hydrogen bond with the C=O group of the amino acid four residues earlier in the same helix.

The shape of a protein is critical to its function because it determines whether the protein can interact with other molecules. Protein structures are very complex, and researchers have only very recently been able to easily and quickly determine the structure of complete proteins down to the atomic level. (The techniques used date back to the 1950s, but until recently they were very slow and laborious to use, so complete protein structures were very slow to be solved.) Early structural biochemists conceptually divided protein structures into four "levels" to make it easier to get a handle on the complexity of the overall structures. To determine how the protein gets its final shape or conformation, we need to understand these four levels of protein structure: primary, secondary, tertiary, and quaternary.


Primary Structure

A protein's primary structure is the unique sequence of amino acids in each polypeptide chain that makes up the protein. Really, this is just a list of which amino acids appear in which order in a polypeptide chain, not really a structure. But, because the final protein structure ultimately depends on this sequence, this was called the primary structure of the polypeptide chain. For example, the pancreatic hormone insulin has two polypeptide chains, A and B.



Primary structure: The A chain of insulin is 21 amino acids long and the B chain is 30 amino acids long, and each sequence is unique to the insulin protein.

The gene, or sequence of DNA, ultimately determines the unique sequence of amino acids in each peptide chain. A change in nucleotide sequence of the gene's coding region may lead to a different amino acid being added to the growing polypeptide chain, causing a change in protein structure and therefore function.




Sickle cell disease: Sickle cells are crescent shaped, while normal cells are disc-shaped.

The oxygen-transport protein hemoglobin consists of four polypeptide chains, two identical α chains and two identical β chains. In sickle cell anemia, a single amino substitution in the hemoglobin β chain causes a change the structure of the entire protein. When the amino acid glutamic acid is replaced by valine in the β chain, the polypeptide folds into an slightly-different shape that creates a dysfunctional hemoglobin protein. So, just one amino acid substitution can cause dramatic changes. These dysfunctional hemoglobin proteins, under low-oxygen conditions, start associating with one another, forming long fibers made from millions of aggregated hemoglobins that distort the red blood cells into crescent or "sickle" shapes, which clog arteries. People affected by the disease often experience breathlessness, dizziness, headaches, and abdominal pain.

Secondary Structure

A protein's secondary structure is whatever regular structures arise from interactions between neighboring or near-by amino acids as the polypeptide starts to fold into its functional three-dimensional form. Secondary structures arise as H bonds form between local groups of amino acids in a region of the polypeptide chain. Rarely does a single secondary structure extend throughout the polypeptide chain. It is usually just in a section of the chain. The most common forms of secondary structure are the α-helix and β-pleated sheet structures and they play an important structural role in most globular and fibrous proteins.




Secondary structure: The α-helix and β-pleated sheet form because of hydrogen bonding between carbonyl and amino groups in the peptide backbone. Certain amino acids have a propensity to form an α-helix, while others have a propensity to form a β-pleated sheet.

In the α-helix chain, the hydrogen bond forms between the oxygen atom in the polypeptide backbone carbonyl group in one amino acid and the hydrogen atom in the polypeptide backbone amino group of another amino acid that is four amino acids farther along the chain. This holds the stretch of amino acids in a right-handed coil. Every helical turn in an alpha helix has 3.6 amino acid residues. The R groups (the side chains) of the polypeptide protrude out from the α-helix chain and are not involved in the H bonds that maintain the α-helix structure.

In β-pleated sheets, stretches of amino acids are held in an almost fully-extended conformation that "pleats" or zig-zags due to the non-linear nature of single C-C and C-N covalent bonds. β-pleated sheets never occur alone. They have to held in place by other β-pleated sheets. The stretches of amino acids in β-pleated sheets are held in their pleated sheet structure because hydrogen bonds form between the oxygen atom in a polypeptide backbone carbonyl group of one β-pleated sheet and the hydrogen atom in a polypeptide backbone amino group of another β-pleated sheet. The β-pleated sheets which hold each other together align parallel or antiparallel to each other. The R groups of the amino acids in a β-pleated sheet point out perpendicular to the hydrogen bonds holding the β-pleated sheets together, and are not involved in maintaining the β-pleated sheet structure.

Tertiary Structure

The tertiary structure of a polypeptide chain is its overall three-dimensional shape, once all the secondary structure elements have folded together among each other. Interactions between polar, nonpolar, acidic, and basic R group within the polypeptide chain create the complex three-dimensional tertiary structure of a protein. When protein folding takes place in the aqueous environment of the body, the hydrophobic R groups of nonpolar amino acids mostly lie in the interior of the protein, while the hydrophilic R groups lie mostly on the outside. Cysteine side chains form disulfide linkages in the presence of oxygen, the only covalent bond forming during protein folding. All of these interactions, weak and strong, determine the final three-dimensional shape of the protein. When a protein loses its three-dimensional shape, it will no longer be functional.




Tertiary structure: The tertiary structure of proteins is determined by hydrophobic interactions, ionic bonding, hydrogen bonding, and disulfide linkages.

Quaternary Structure

The quaternary structure of a protein is how its subunits are oriented and arranged with respect to one another. As a result, quaternary structure only applies to multi-subunit proteins; that is, proteins made from more than one polypeptide chain. Proteins made from a single polypeptide will not have a quaternary structure.

In proteins with more than one subunit, weak interactions between the subunits help to stabilize the overall structure. Enzymes often play key roles in bonding subunits to form the final, functioning protein.

For example, insulin is a ball-shaped, globular protein that contains both hydrogen bonds and disulfide bonds that hold its two polypeptide chains together. Silk is a fibrous protein that results from hydrogen bonding between different β-pleated chains.



Four levels of protein structure: The four levels of protein structure can be observed in these illustrations.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- amino acid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amino_acid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Proteins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Proteins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Structural Biochemistry/Proteins/Amino Acids. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...ns/Amino_Acids**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- catalyze. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/catalyze**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- polypeptide. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/polypeptide**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- BPAE cells | Flickr - Photo Sharing!. **Provided by**: Flickr. **Located at**: **http://www.flickr.com/photos/6345024...-dtyzGH-duUj14**. **License**: ***CC BY: Attribution***
- 1GZX Haemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1G...aemoglobin.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Induced fit diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:In...it_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- amino acid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amino_acid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/definition/r-group**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- polypeptide. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/polypeptide**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- BPAE cells | Flickr - Photo Sharing!. **Provided by**: Flickr. **Located at**: **http://www.flickr.com/photos/6345024...-dtyzGH-duUj14**. **License**: ***CC BY: Attribution***
- 1GZX Haemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1G...aemoglobin.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Induced fit diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:In...it_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...e_03_04_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...e_03_04_02.png**. **License**: ***CC BY: Attribution***
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...e_03_04_01.jpg**. **License**: ***CC BY: Attribution***
- hydrogen bond. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hydrogen_bond**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44402/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...n/antiparallel**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- disulfide bond. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/disulfide_bond**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- BPAE cells | Flickr - Photo Sharing!. **Provided by**: Flickr. **Located at**: **http://www.flickr.com/photos/6345024...-dtyzGH-duUj14**. **License**: ***CC BY: Attribution***
- 1GZX Haemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:1G...aemoglobin.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Induced fit diagram. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Induced_fit_diagram.svg**. **License**:

Public Domain: No Known Copyright

- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_03.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_02.png. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_09.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_07.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_04.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_06.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Proteins. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44402/latest...e_03_04_08.jpg. **License**: *CC BY: Attribution*

This page titled 23.4C: Protein Structure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5: METABOLIC REACTIONS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.5A: Breakdown of Pyruvate

23.5B: Citric Acid Cycle

23.5C: Importance of Glycolysis

23.5D: Electron Transport Chain

23.5E: ATP Yield

23.5F: Control of Catabolic Pathways

23.5G: Transforming Chemical Energy

This page titled 23.5: Metabolic Reactions is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5A: Breakdown of Pyruvate

After glycolysis, pyruvate is converted into acetyl CoA in order to enter the citric acid cycle.

Learning Objectives

- Explain why cells break down pyruvate

Key Points


- In the conversion of pyruvate to acetyl CoA, each pyruvate molecule loses one carbon atom with the release of carbon dioxide.
- During the breakdown of pyruvate, electrons are transferred to NAD+ to produce NADH, which will be used by the cell to produce ATP.
- In the final step of the breakdown of pyruvate, an acetyl group is transferred to Coenzyme A to produce acetyl CoA.

Key Terms

- **acetyl CoA**: a molecule that conveys the carbon atoms from glycolysis (pyruvate) to the citric acid cycle to be oxidized for energy production

Breakdown of Pyruvate

In order for pyruvate, the product of glycolysis, to enter the next pathway, it must undergo several changes to become acetyl Coenzyme A (acetyl CoA). Acetyl CoA is a molecule that is further converted to oxaloacetate, which enters the citric acid cycle (Krebs cycle). The conversion of pyruvate to acetyl CoA is a three-step process.



Breakdown of Pyruvate: Each pyruvate molecule loses a carboxylic group in the form of carbon dioxide. The remaining two carbons are then transferred to the enzyme CoA to produce Acetyl CoA.

Step 1. A carboxyl group is removed from pyruvate, releasing a molecule of carbon dioxide into the surrounding medium. (Note: carbon dioxide is one carbon attached to two oxygen atoms and is one of the major end products of cellular respiration.) The

result of this step is a two-carbon hydroxyethyl group bound to the enzyme pyruvate dehydrogenase; the lost carbon dioxide is the first of the six carbons from the original glucose molecule to be removed. This step proceeds twice for every molecule of glucose metabolized (remember: there are two pyruvate molecules produced at the end of glycolysis); thus, two of the six carbons will have been removed at the end of both of these steps.

Step 2. The hydroxyethyl group is oxidized to an acetyl group, and the electrons are picked up by NAD^+, forming NADH (the reduced form of NAD+). The high- energy electrons from NADH will be used later by the cell to generate ATP for energy.

Step 3. The enzyme-bound acetyl group is transferred to CoA, producing a molecule of acetyl CoA. This molecule of acetyl CoA is then further converted to be used in the next pathway of metabolism, the citric acid cycle.

This page titled 23.5A: Breakdown of Pyruvate is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5B: Citric Acid Cycle

The citric acid cycle is a series of reactions that produces two carbon dioxide molecules, one GTP/ATP, and reduced forms of NADH and FADH2.

Learning Objectives

- List the steps of the Krebs (or citric acid) cycle

Key Points


- The four-carbon molecule, oxaloacetate, that began the cycle is regenerated after the eight steps of the citric acid cycle.
- The eight steps of the citric acid cycle are a series of redox, dehydration, hydration, and decarboxylation reactions.
- Each turn of the cycle forms one GTP or ATP as well as three NADH molecules and one FADH2 molecule, which will be used in further steps of cellular respiration to produce ATP for the cell.

Key Terms

- **citric acid cycle**: a series of chemical reactions used by all aerobic organisms to generate energy through the oxidization of acetate derived from carbohydrates, fats, and proteins into carbon dioxide
- **Krebs cycle**: a series of enzymatic reactions that occurs in all aerobic organisms; it involves the oxidative metabolism of acetyl units and serves as the main source of cellular energy
- **mitochondria**: in cell biology, a mitochondrion (plural mitochondria) is a membrane-enclosed organelle, often described as “cellular power plants” because they generate most of the ATP

Citric Acid Cycle (Krebs Cycle)

Like the conversion of pyruvate to acetyl CoA, the citric acid cycle takes place in the matrix of the mitochondria. Almost all of the enzymes of the citric acid cycle are soluble, with the single exception of the enzyme succinate dehydrogenase, which is embedded in the inner membrane of the mitochondrion. Unlike glycolysis, the citric acid cycle is a closed loop: the last part of the pathway regenerates the compound used in the first step. The eight steps of the cycle are a series of redox, dehydration, hydration, and decarboxylation reactions that produce two carbon dioxide molecules, one GTP/ATP, and reduced forms of NADH and FADH2. This is considered an aerobic pathway because the NADH and FADH2 produced must transfer their electrons to the next pathway in the system, which will use oxygen. If this transfer does not occur, the oxidation steps of the citric acid cycle also do not occur. Note that the citric acid cycle produces very little ATP directly and does not directly consume oxygen.



The citric acid cycle: In the citric acid cycle, the acetyl group from acetyl CoA is attached to a four-carbon oxaloacetate molecule to form a six-carbon citrate molecule. Through a series of steps, citrate is oxidized, releasing two carbon dioxide molecules for each acetyl group fed into the cycle. In the process, three NAD+ molecules are reduced to NADH, one FAD molecule is reduced to FADH2, and one ATP or GTP (depending on the cell type) is produced (by substrate-level phosphorylation). Because the final product of the citric acid cycle is also the first reactant, the cycle runs continuously in the presence of sufficient reactants.

Steps in the Citric Acid Cycle

Step 1. The first step is a condensation step, combining the two-carbon acetyl group (from acetyl CoA) with a four-carbon oxaloacetate molecule to form a six-carbon molecule of citrate. CoA is bound to a sulfhydryl group (-SH) and diffuses away to eventually combine with another acetyl group. This step is irreversible because it is highly exergonic. The rate of this reaction is controlled by negative feedback and the amount of ATP available. If ATP levels increase, the rate of this reaction decreases. If ATP is in short supply, the rate increases.

Step 2. Citrate loses one water molecule and gains another as citrate is converted into its isomer, isocitrate.

Steps 3 and 4. In step three, isocitrate is oxidized, producing a five-carbon molecule, α-ketoglutarate, together with a molecule of $CO_2$and two electrons, which reduce NAD+ to NADH. This step is also regulated by negative feedback from ATP and NADH and by a positive effect of ADP. Steps three and four are both oxidation and decarboxylation steps, which release electrons that reduce NAD^+ to NADH and release carboxyl groups that form CO_2 molecules. α-Ketoglutarate is the product of step three, and a succinyl group is the product of step four. CoA binds the succinyl group to form succinyl CoA. The enzyme that catalyzes step four is regulated by feedback inhibition of ATP, succinyl CoA, and NADH.

Step 5. A phosphate group is substituted for coenzyme A, and a high- energy bond is formed. This energy is used in substrate-level phosphorylation (during the conversion of the succinyl group to succinate) to form either guanine triphosphate (GTP) or ATP. There are two forms of the enzyme, called isoenzymes, for this step, depending upon the type of animal tissue in which they are found. One form is found in tissues that use large amounts of ATP, such as heart and skeletal muscle. This form produces ATP. The second form of the enzyme is found in tissues that have a high number of anabolic pathways, such as liver. This form produces GTP. GTP is energetically equivalent to ATP; however, its use is more restricted. In particular, protein synthesis primarily uses GTP.

Step 6. Step six is a dehydration process that converts succinate into fumarate. Two hydrogen atoms are transferred to FAD, producing $FADH_2$. The energy contained in the electrons of these atoms is insufficient to reduce NAD^+ but adequate to reduce FAD. Unlike NADH, this carrier remains attached to the enzyme and transfers the electrons to the electron transport chain directly. This process is made possible by the localization of the enzyme catalyzing this step inside the inner membrane of the mitochondrion.

Step 7. Water is added to fumarate during step seven, and malate is produced. The last step in the citric acid cycle regenerates oxaloacetate by oxidizing malate. Another molecule of NADH is produced.

Products of the Citric Acid Cycle

Two carbon atoms come into the citric acid cycle from each acetyl group, representing four out of the six carbons of one glucose molecule. Two carbon dioxide molecules are released on each turn of the cycle; however, these do not necessarily contain the most recently-added carbon atoms. The two acetyl carbon atoms will eventually be released on later turns of the cycle; thus, all six carbon atoms from the original glucose molecule are eventually incorporated into carbon dioxide. Each turn of the cycle forms three NADH molecules and one $FADH_2$ molecule. These carriers will connect with the last portion of aerobic respiration to produce ATP molecules. One GTP or ATP is also made in each cycle. Several of the intermediate compounds in the citric acid cycle can be used in synthesizing non-essential amino acids; therefore, the cycle is amphibolic (both catabolic and anabolic).

This page titled 23.5B: Citric Acid Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5C: Importance of Glycolysis

Glycolysis is the first step in the breakdown of glucose to extract energy for cellular metabolism.

Learning Objectives

- Explain the importance of glycolysis to cells

Key Points


- Glycolysis is present in nearly all living organisms.
- Glucose is the source of almost all energy used by cells.
- Overall, glycolysis produces two pyruvate molecules, a net gain of two ATP molecules, and two NADH molecules.

Key Terms

- **glycolysis**: the cellular metabolic pathway of the simple sugar glucose to yield pyruvic acid and ATP as an energy source
- **heterotroph**: an organism that requires an external supply of energy in the form of food, as it cannot synthesize its own

Nearly all of the energy used by living cells comes to them from the energy in the bonds of the sugar glucose. Glucose enters heterotrophic cells in two ways. One method is through secondary active transport in which the transport takes place against the glucose concentration gradient. The other mechanism uses a group of integral proteins called GLUT proteins, also known as glucose transporter proteins. These transporters assist in the facilitated diffusion of glucose. Glycolysis is the first pathway used in the breakdown of glucose to extract energy. It takes place in the cytoplasm of both prokaryotic and eukaryotic cells. It was probably one of the earliest metabolic pathways to evolve since it is used by nearly all of the organisms on earth. The process does not use oxygen and is, therefore, anaerobic.

Glycolysis is the first of the main metabolic pathways of cellular respiration to produce energy in the form of ATP. Through two distinct phases, the six-carbon ring of glucose is cleaved into two three-carbon sugars of pyruvate through a series of enzymatic reactions. The first phase of glycolysis requires energy, while the second phase completes the conversion to pyruvate and produces ATP and NADH for the cell to use for energy. Overall, the process of glycolysis produces a net gain of two pyruvate molecules, two ATP molecules, and two NADH molecules for the cell to use for energy. Following the conversion of glucose to pyruvate, the glycolytic pathway is linked to the Krebs Cycle, where further ATP will be produced for the cell's energy needs.



Cellular Respiration: Glycolysis is the first pathway of cellular respiration that oxidizes glucose molecules. It is followed by the Krebs cycle and oxidative phosphorylation to produce ATP.

This page titled 23.5C: Importance of Glycolysis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5D: Electron Transport Chain

The electron transport chain uses the electrons from electron carriers to create a chemical gradient that can be used to power oxidative phosphorylation.

Learning Objectives

- Describe how electrons move through the electron transport chain

Key Points

- Oxidative phosphorylation is the metabolic pathway in which electrons are transferred from electron donors to electron acceptors in redox reactions; this series of reactions releases energy which is used to form ATP.
- There are four protein complexes (labeled complex I-IV) in the electron transport chain, which are involved in moving electrons from NADH and $FADH_2$ to molecular oxygen.
- Complex I establishes the hydrogen ion gradient by pumping four hydrogen ions across the membrane from the matrix into the intermembrane space.
- Complex II receives $FADH_2$, which bypasses complex I, and delivers electrons directly to the electron transport chain.
- Ubiquinone (Q) accepts the electrons from both complex I and complex II and delivers them to complex III.
- Complex III pumps protons through the membrane and passes its electrons to cytochrome c for transport to the fourth complex of proteins and enzymes.
- Complex IV reduces oxygen; the reduced oxygen then picks up two hydrogen ions from the surrounding medium to make water.

Key Terms


- **prosthetic group**: The non-protein component of a conjugated protein.
- **complex**: A structure consisting of a central atom, molecule, or protein weakly connected to surrounding atoms, molecules, or proteins.
- **ubiquinone**: A lipid soluble substance that is a component of the electron transport chain and accepts electrons from complexes I and II.

Oxidative phosphorylation is a highly efficient method of producing large amounts of ATP, the basic unit of energy for metabolic processes. During this process electrons are exchanged between molecules, which creates a chemical gradient that allows for the production of ATP. The most vital part of this process is the electron transport chain, which produces more ATP than any other part of cellular respiration.

Electron Transport Chain

The electron transport chain is the final component of aerobic respiration and is the only part of glucose metabolism that uses atmospheric oxygen. Electron transport is a series of redox reactions that resemble a relay race. Electrons are passed rapidly from one component to the next to the endpoint of the chain, where the electrons reduce molecular oxygen, producing water. This requirement for oxygen in the final stages of the chain can be seen in the overall equation for cellular respiration, which requires both glucose and oxygen.

A complex is a structure consisting of a central atom, molecule, or protein weakly connected to surrounding atoms, molecules, or proteins. The electron transport chain is an aggregation of four of these complexes (labeled I through IV), together with associated mobile electron carriers. The electron transport chain is present in multiple copies in the inner mitochondrial membrane of eukaryotes and the plasma membrane of prokaryotes.



The electron transport chain: The electron transport chain is a series of electron transporters embedded in the inner mitochondrial membrane that shuttles electrons from NADH and $FADH_2$ to molecular oxygen. In the process, protons are pumped from the mitochondrial matrix to the intermembrane space, and oxygen is reduced to form water.

Complex I

To start, two electrons are carried to the first complex aboard NADH. Complex I is composed of flavin mononucleotide (FMN) and an enzyme containing iron-sulfur (Fe-S). FMN, which is derived from vitamin B_2 (also called riboflavin), is one of several prosthetic groups or co-factors in the electron transport chain. A prosthetic group is a non-protein molecule required for the activity of a protein. Prosthetic groups can be organic or inorganic and are non-peptide molecules bound to a protein that facilitate its function.

Prosthetic groups include co-enzymes, which are the prosthetic groups of enzymes. The enzyme in complex I is NADH dehydrogenase, a very large protein containing 45 amino acid chains. Complex I can pump four hydrogen ions across the membrane from the matrix into the intermembrane space; it is in this way that the hydrogen ion gradient is established and maintained between the two compartments separated by the inner mitochondrial membrane.

Q and Complex II

Complex II directly receives $FADH_2$, which does not pass through complex I. The compound connecting the first and second complexes to the third is ubiquinone (Q). The Q molecule is lipid soluble and freely moves through the hydrophobic core of the membrane. Once it is reduced to QH_2, ubiquinone delivers its electrons to the next complex in the electron transport chain. Q receives the electrons derived from NADH from complex I and the electrons derived from $FADH_2$ from complex II, including succinate dehydrogenase. This enzyme and $FADH_2$ form a small complex that delivers electrons directly to the electron transport chain, bypassing the first complex. Since these electrons bypass, and thus do not energize, the proton pump in the first complex, fewer ATP molecules are made from the $FADH_2$ electrons. The number of ATP molecules ultimately obtained is directly proportional to the number of protons pumped across the inner mitochondrial membrane.

Complex III

The third complex is composed of cytochrome b, another Fe-S protein, Rieske center (2Fe-2S center), and cytochrome c proteins; this complex is also called cytochrome oxidoreductase. Cytochrome proteins have a prosthetic heme group. The heme molecule is similar to the heme in hemoglobin, but it carries electrons, not oxygen. As a result, the iron ion at its core is reduced and oxidized as it passes the electrons, fluctuating between different oxidation states: Fe^{2+} (reduced) and Fe^{3+} (oxidized). The heme molecules in the cytochromes have slightly different characteristics due to the effects of the different proteins binding them,

which makes each complex. Complex III pumps protons through the membrane and passes its electrons to cytochrome c for transport to the fourth complex of proteins and enzymes. Cytochrome c is the acceptor of electrons from Q; however, whereas Q carries pairs of electrons, cytochrome c can accept only one at a time.

Complex IV

The fourth complex is composed of cytochrome proteins c, a, and a_3. This complex contains two heme groups (one in each of the cytochromes a and a_3) and three copper ions (a pair of Cu_A and one Cu_B in cytochrome a_3). The cytochromes hold an oxygen molecule very tightly between the iron and copper ions until the oxygen is completely reduced. The reduced oxygen then picks up two hydrogen ions from the surrounding medium to produce water (H_2O). The removal of the hydrogen ions from the system also contributes to the ion gradient used in the process of chemiosmosis.

This page titled 23.5D: Electron Transport Chain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5E: ATP Yield

The amount of energy (as ATP) gained from glucose catabolism varies across species and depends on other related cellular processes.

Learning Objectives

- Describe the origins of variability in the amount of ATP that is produced per molecule of glucose consumed

Key Points


- While glucose catabolism always produces energy, the amount of energy (in terms of ATP equivalents) produced can vary, especially across different species.
- The number of hydrogen ions the electron transport chain complexes can pump through the membrane varies between species.
- NAD^+ provides more ATP than FAD^+ in the electron transport chain and can lead to variance in ATP production.
- The use of intermediates from glucose catabolism in other biosynthetic pathways, such as amino acid synthesis, can lower the yield of ATP.

Key Terms

- **catabolism**: Destructive metabolism, usually including the release of energy and breakdown of materials.

ATP Yield


In a eukaryotic cell, the process of cellular respiration can metabolize one molecule of glucose into 30 to 32 ATP. The process of glycolysis only produces two ATP, while all the rest are produced during the electron transport chain. Clearly, the electron transport chain is vastly more efficient, but it can only be carried out in the presence of oxygen.



Cellular respiration in a eukaryotic cell: Glycolysis on the left portion of this illustration can be seen to yield 2 ATP molecules, while the Electron Transport Chain portion at the upper right will yield the remaining 30-32 ATP molecules under the presence

of oxygen.

The number of ATP molecules generated via the catabolism of glucose can vary substantially. For example, the number of hydrogen ions the electron transport chain complexes can pump through the membrane varies between species. Another source of variance occurs during the shuttle of electrons across the membranes of the mitochondria. The NADH generated from glycolysis cannot easily enter mitochondria. Thus, electrons are picked up on the inside of mitochondria by either NAD^+ or FAD^+. These FAD^+ molecules can transport fewer ions; consequently, fewer ATP molecules are generated when FAD^+ acts as a carrier. NAD^+ is used as the electron transporter in the liver, and FAD^+ acts in the brain.



Adenosine triphosphate: ATP is the main source of energy in many living organisms.

Another factor that affects the yield of ATP molecules generated from glucose is the fact that intermediate compounds in these pathways are used for other purposes. Glucose catabolism connects with the pathways that build or break down all other biochemical compounds in cells, but the result is not always ideal. For example, sugars other than glucose are fed into the glycolytic pathway for energy extraction. Moreover, the five-carbon sugars that form nucleic acids are made from intermediates in glycolysis. Certain nonessential amino acids can be made from intermediates of both glycolysis and the citric acid cycle. Lipids, such as cholesterol and triglycerides, are also made from intermediates in these pathways, and both amino acids and triglycerides are broken down for energy through these pathways. Overall, in living systems, these pathways of glucose catabolism extract about 34 percent of the energy contained in glucose.

This page titled 23.5E: ATP Yield is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5F: Control of Catabolic Pathways

Catabolic pathways are controlled by enzymes, proteins, electron carriers, and pumps that ensure that the remaining reactions can proceed.

Learning Objectives

- Explain how catabolic pathways are controlled

Key Points

- Glycolysis, the citric acid cycle, and the electron transport chain are catabolic pathways that bring forth non-reversible reactions.
- Glycolysis control begins with hexokinase, which catalyzes the phosphorylation of glucose; its product is glucose-6-phosphate, which accumulates when phosphofructokinase is inhibited.
- The citric acid cycle is controlled through the enzymes that break down the reactions that make the first two molecules of NADH.
- The rate of electron transport through the electron transport chain is affected by the levels of ADP and ATP, whereas specific enzymes of the electron transport chain are unaffected by feedback inhibition.

Key Terms


- **phosphofructokinase**: any of a group of kinase enzymes that convert fructose phosphates to biphosphate
- **glycolysis**: the cellular metabolic pathway of the simple sugar glucose to yield pyruvic acid and ATP as an energy source
- **kinase**: any of a group of enzymes that transfers phosphate groups from high-energy donor molecules, such as ATP, to specific target molecules (substrates); the process is termed phosphorylation

Control of Catabolic Pathways

Enzymes, proteins, electron carriers, and pumps that play roles in glycolysis, the citric acid cycle, and the electron transport chain tend to catalyze non-reversible reactions. In other words, if the initial reaction takes place, the pathway is committed to proceeding with the remaining reactions. Whether a particular enzyme activity is released depends upon the energy needs of the cell (as reflected by the levels of ATP, ADP, and AMP).

Glycolysis

The control of glycolysis begins with the first enzyme in the pathway, hexokinase. This enzyme catalyzes the phosphorylation of glucose, which helps to prepare the compound for cleavage in a later step. The presence of the negatively-charged phosphate in the molecule also prevents the sugar from leaving the cell. When hexokinase is inhibited, glucose diffuses out of the cell and does not become a substrate for the respiration pathways in that tissue. The product of the hexokinase reaction is glucose-6-phosphate, which accumulates when a later enzyme, phosphofructokinase, is inhibited.



Glycolysis: The glycolysis pathway is primarily regulated at the three key enzymatic steps (1, 2, and 7) as indicated. Note that the first two steps that are regulated occur early in the pathway and involve hydrolysis of ATP.


Phosphofructokinase is the main enzyme controlled in glycolysis. High levels of ATP, citrate, or a lower, more acidic pH decrease the enzyme's activity. An increase in citrate concentration can occur because of a blockage in the citric acid cycle. Fermentation, with its production of organic acids like lactic acid, frequently accounts for the increased acidity in a cell; however, the products of fermentation do not typically accumulate in cells.

The last step in glycolysis is catalyzed by pyruvate kinase. The pyruvate produced can proceed to be catabolized or converted into the amino acid alanine. If no more energy is needed and alanine is in adequate supply, the enzyme is inhibited. The enzyme's activity is increased when fructose-1,6-bisphosphate levels increase. (Recall that fructose-1,6-bisphosphate is an intermediate in the first half of glycolysis.) The regulation of pyruvate kinase involves phosphorylation, resulting in a less-active enzyme. Dephosphorylation by a phosphatase reactivates it. Pyruvate kinase is also regulated by ATP (a negative allosteric effect).

If more energy is needed, more pyruvate will be converted into acetyl CoA through the action of pyruvate dehydrogenase. If either acetyl groups or NADH accumulate, there is less need for the reaction and the rate decreases. Pyruvate dehydrogenase is also regulated by phosphorylation: a kinase phosphorylates it to form an inactive enzyme, and a phosphatase reactivates it. The kinase and the phosphatase are also regulated.

Citric Acid Cycle


The citric acid cycle is controlled through the enzymes that catalyze the reactions that make the first two molecules of NADH. These enzymes are isocitrate dehydrogenase and α-ketoglutarate dehydrogenase. When adequate ATP and NADH levels are available, the rates of these reactions decrease. When more ATP is needed, as reflected in rising ADP levels, the rate increases. α-Ketoglutarate dehydrogenase will also be affected by the levels of succinyl CoA, a subsequent intermediate in the cycle, causing a decrease in activity. A decrease in the rate of operation of the pathway at this point is not necessarily negative as the increased levels of the α-ketoglutarate not used by the citric acid cycle can be used by the cell for amino acid (glutamate) synthesis.



Citric Acid Cycle: Enzymes, isocitrate dehydrogenase and α-ketoglutarate dehydrogenase, catalyze the reactions that make the first two molecules of NADH in the citric acid cycle. Rates of the reaction decrease when sufficient ATP and NADH levels are reached.

Electron Transport Chain

Specific enzymes of the electron transport chain are unaffected by feedback inhibition, but the rate of electron transport through the pathway is affected by the levels of ADP and ATP. Greater ATP consumption by a cell is indicated by a buildup of ADP. As ATP usage decreases, the concentration of ADP decreases: ATP begins to build up in the cell. This change in the relative concentration of ADP to ATP triggers the cell to slow down the electron transport chain.



Electron Chain Transport: Levels of ADP and ATP affect the rate of electron transport through this type of chain transport.

This page titled 23.5F: Control of Catabolic Pathways is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.5G: Transforming Chemical Energy

Cellular respiration is the process of transforming chemical energy into forms usable by the cell or organism.

Learning Objectives

- Discuss the importance of cellular respiration

Key Points

- Organisms ingest organic molecules like the carbohydrate glucose to obtain the energy needed for cellular functions.
- The energy in glucose can be extracted in a series of chemical reactions known as cellular respiration.
- Cellular respiration produces energy in the form of ATP, which is the universal energy currency for cells.

Key Terms

- **aerobic respiration**: the process of converting the biochemical energy in nutrients to ATP in the presence of oxygen
- **adenosine triphosphate**: a multifunctional nucleoside triphosphate used in cells as a coenzyme, often called the "molecular unit of energy currency" in intracellular energy transfer
- **cellular respiration**: the set of the metabolic reactions and processes that take place in the cells of organisms to convert biochemical energy from nutrients into adenosine triphosphate (ATP)
- **catabolism**: the breakdown of large molecules into smaller ones usually accompanied by the release of energy

Introduction: Cellular Respiration

An electrical energy plant converts energy from one form to another form that can be more easily used. For example, geothermal energy plants start with underground thermal energy (heat) and transform it into electrical energy that will be transported to homes and factories.



Energy Plant: This geothermal energy plant transforms thermal energy from deep in the ground into electrical energy, which can be easily used.

Like a generating plant, living organisms must take in energy from their environment and convert it into to a form their cells can use. Organisms ingest large molecules, like carbohydrates, proteins, and fats, and convert them into smaller molecules like carbon dioxide and water. This process is called cellular respiration, a form of catabolism, and makes energy available for the cell to use. The energy released by cellular respiration is temporarily captured by the formation of adenosine triphosphate (ATP) within the cell. ATP is the principle form of stored energy used for cellular functions and is frequently referred to as the energy currency of the cell.

The nutrients broken down through cellular respiration lose electrons throughout the process and are said to be oxidized. When oxygen is used to help drive the oxidation of nutrients the process is called aerobic respiration. Aerobic respiration is common among the eukaryotes, including humans, and takes place mostly within the mitochondria. Respiration occurs within the cytoplasm of prokaryotes. Several prokaryotes and a few eukaryotes use an inorganic molecule other than oxygen to drive the oxidation of their nutrients in a process called anaerobic respiration. Electron acceptors for anaerobic respiration include nitrate, sulfate, carbon dioxide, and several metal ions.

The energy released during cellular respiration is then used in other biological processes. These processes build larger molecules that are essential to an organism's survival, such as amino acids, DNA, and proteins. Because they synthesize new molecules, these processes are examples of anabolism.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- acetyl CoA. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/acetyl%20CoA. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: ***CC BY: Attribution***
- Krebs cycle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Krebs_cycle. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- mitochondria. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/mitochondria. **License**: ***CC BY-SA: Attribution-ShareAlike***
- citric acid cycle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/citric%20acid%20cycle. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44432/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- heterotroph. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/heterotroph. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycolysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glycolysis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Carbohydrate Metabolism. November 10, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46451/latest/. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...ol11448/latest. **License**: ***CC BY: Attribution***

- ubiquinone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/ubiquinone. **License**: *CC BY-SA: Attribution-ShareAlike*
- complex. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/complex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Oxidative phosphorylation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Oxidative_phosphorylation. **License**: *CC BY-SA: Attribution-ShareAlike*
- prosthetic group. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/prosthetic_group. **License**: *CC BY-SA: Attribution-ShareAlike*
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Carbohydrate Metabolism. November 10, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46451/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Oxidative Phosphorylation. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...e_07_04_01.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...ol11448/latest. **License**: *CC BY: Attribution*
- catabolism. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/catabolism. **License**: *CC BY-SA: Attribution-ShareAlike*
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Carbohydrate Metabolism. November 10, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46451/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Oxidative Phosphorylation. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...e_07_04_01.jpg. **License**: *CC BY: Attribution*
- ATP-3D-vdW. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ATP-3D-vdW.png. **License**: *Public Domain: No Known Copyright*
- Cellular Respiration in a Eukaryotic Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cellul...espiration.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44442/latest...ol11448/latest. **License**: *CC BY: Attribution*
- phosphofructokinase. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/phosphofructokinase. **License**: *CC BY-SA: Attribution-ShareAlike*
- kinase. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/kinase. **License**: *CC BY-SA: Attribution-ShareAlike*
- glycolysis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glycolysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...cetylCoA-L.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Carbohydrate Metabolism. November 10, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46451/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Oxidative Phosphorylation. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...e_07_04_01.jpg. **License**: *CC BY: Attribution*
- ATP-3D-vdW. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ATP-3D-vdW.png. **License**: *Public Domain: No Known Copyright*
- Cellular Respiration in a Eukaryotic Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cellul...espiration.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Regulation of Cellular Respiration. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44442/latest...e_07_07_02.jpg. **License**: *CC BY: Attribution*
- Citric acid cycle. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...acid_cycle.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Electron transport chain simple. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ain_simple.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- adenosine triphosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adenosine%20triphosphate.

License: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44430/latest...ol11448/latest. **License**: *CC BY: Attribution*
- cellular respiration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cellular%20respiration. **License**: *CC BY-SA: Attribution-ShareAlike*
- photosynthesis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/photosynthesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- 09 10PyruvateToAcetylCoA-L. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:09_10PyruvateToAcetylCoA-L.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Oxidation of Pyruvate and the Citric Acid Cycle. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44433/latest...e_07_03_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Carbohydrate Metabolism. November 10, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46451/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Oxidative Phosphorylation. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44437/latest...e_07_04_01.jpg. **License**: *CC BY: Attribution*
- ATP-3D-vdW. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ATP-3D-vdW.png. **License**: *Public Domain: No Known Copyright*
- Cellular Respiration in a Eukaryotic Cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cellul...espiration.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Regulation of Cellular Respiration. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44442/latest...e_07_07_02.jpg. **License**: *CC BY: Attribution*
- Citric acid cycle. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...acid_cycle.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Electron transport chain simple. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ain_simple.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44430/latest...e_07_00_01.jpg. **License**: *CC BY: Attribution*

This page titled 23.5G: Transforming Chemical Energy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.6: METABOLISM OF NUTRIENTS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

23.6A: Connecting Other Sugars to Glucose Metabolism
23.6B: Connecting Proteins to Glucose Metabolism

This page titled 23.6: Metabolism of Nutrients is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.6A: Connecting Other Sugars to Glucose Metabolism

Sugars, such as galactose, fructose, and glycogen, are catabolized into new products in order to enter the glycolytic pathway.

Learning Objectives

- Identify the types of sugars involved in glucose metabolism

Key Points


- When blood sugar levels drop, glycogen is broken down into glucose -1-phosphate, which is then converted to glucose-6-phosphate and enters glycolysis for ATP production.
- In the liver, galactose is converted to glucose-6-phosphate in order to enter the glycolytic pathway.
- Fructose is converted into glycogen in the liver and then follows the same pathway as glycogen to enter glycolysis.
- Sucrose is broken down into glucose and fructose; glucose enters the pathway directly while fructose is converted to glycogen.

Key Terms

- **disaccharide**: A sugar, such as sucrose, maltose, or lactose, consisting of two monosaccharides combined together.
- **glycogen**: A polysaccharide that is the main form of carbohydrate storage in animals; converted to glucose as needed.
- **monosaccharide**: A simple sugar such as glucose, fructose, or deoxyribose that has a single ring.

You have learned about the catabolism of glucose, which provides energy to living cells. But living things consume more than glucose for food. How does a turkey sandwich end up as ATP in your cells? This happens because all of the catabolic pathways for carbohydrates, proteins, and lipids eventually connect into glycolysis and the citric acid cycle pathways.


Metabolic pathways should be thought of as porous; that is, substances enter from other pathways, and intermediates leave for other pathways. These pathways are not closed systems. Many of the substrates, intermediates, and products in a particular pathway are reactants in other pathways. Like sugars and amino acids, the catabolic pathways of lipids are also connected to the glucose catabolism pathways.



Glycogen Pathway: Glycogen from the liver and muscles, hydrolyzed into glucose-1-phosphate, together with fats and proteins, can feed into the catabolic pathways for carbohydrates.

Glycogen, a polymer of glucose, is an energy-storage molecule in animals. When there is adequate ATP present, excess glucose


is shunted into glycogen for storage. Glycogen is made and stored in both the liver and muscles. The glycogen is hydrolyzed into the glucose monomer, glucose-1-phosphate (G-1-P), if blood sugar levels drop. The presence of glycogen as a source of glucose allows ATP to be produced for a longer period of time during exercise. Glycogen is broken down into G-1-P and converted into glucose-6-phosphate (G-6-P) in both muscle and liver cells; this product enters the glycolytic pathway.



Glycogen Structure: Schematic two-dimensional cross-sectional view of glycogen: A core protein of glycogenin is surrounded by branches of glucose units. The entire globular granule may contain around 30,000 glucose units.

Galactose is the sugar in milk. Infants have an enzyme in the small intestine that metabolizes lactose to galactose and glucose. In areas where milk products are regularly consumed, adults have also evolved this enzyme. Galactose is converted in the liver to G-6-P and can thus enter the glycolytic pathway.

Fructose is one of the three dietary monosaccharides (along with glucose and galactose) which are absorbed directly into the bloodstream during digestion. Fructose is absorbed from the small intestine and then passes to the liver to be metabolized, primarily to glycogen. The catabolism of both fructose and galactose produces the same number of ATP molecules as glucose.



Fructose Metabolism: Although the metabolism of fructose and glucose share many of the same intermediate structures, they have very different metabolic fates in human metabolism.

Sucrose is a disaccharide with a molecule of glucose and a molecule of fructose bonded together with a glycosidic linkage. The catabolism of sucrose breaks it down to monomers of glucose and fructose. The glucose can directly enter the glycolytic pathway while fructose must first be converted to glycogen, which can be broken down to G-1-P and enter the glycolytic pathway as described above.

This page titled 23.6A: Connecting Other Sugars to Glucose Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.6B: Connecting Proteins to Glucose Metabolism

Excess amino acids are converted into molecules that can enter the pathways of glucose catabolism.

Learning Objectives

- Describe the role played by proteins in glucose metabolism

Key Points


- Amino acids must be deaminated before entering any of the pathways of glucose catabolism: the amino group is converted to ammonia, which is used by the liver in the synthesis of urea.
- Deaminated amino acids can be converted into pyruvate, acetyl CoA, or some components of the citric acid cycle to enter the pathways of glucose catabolism.
- Several amino acids can enter the glucose catabolism pathways at multiple locations.

Key Terms

- **catabolism**: Destructive metabolism, usually including the release of energy and breakdown of materials.
- **keto acid**: Any carboxylic acid that also contains a ketone group.
- **deamination**: The removal of an amino group from a compound.

Metabolic pathways should be thought of as porous; that is, substances enter from other pathways and intermediates leave for other pathways. These pathways are not closed systems. Many of the substrates, intermediates, and products in a particular pathway are reactants in other pathways. Proteins are a good example of this phenomenon. They can be broken down into their constituent amino acids and used at various steps of the pathway of glucose catabolism.

Proteins are hydrolyzed by a variety of enzymes in cells. Most of the time, the amino acids are recycled into the synthesis of new proteins or are used as precursors in the synthesis of other important biological molecules, such as hormones, nucleotides, or neurotransmitters. However, if there are excess amino acids, or if the body is in a state of starvation, some amino acids will be shunted into the pathways of glucose catabolism.



Connection of Amino Acids to Glucose Metabolism Pathways: The carbon skeletons of certain amino acids (indicated in

boxes) are derived from proteins and can feed into pyruvate, acetyl CoA, and the citric acid cycle.

Each amino acid must have its amino group removed (deamination) prior to the carbon chain's entry into these pathways. When the amino group is removed from an amino acid, it is converted into ammonia through the urea cycle. The remaining atoms of the amino acid result in a keto acid: a carbon chain with one ketone and one carboxylic acid group. In mammals, the liver synthesizes urea from two ammonia molecules and a carbon dioxide molecule. Thus, urea is the principal waste product in mammals produced from the nitrogen originating in amino acids; it leaves the body in urine. The keto acid can then enter the citric acid cycle.

When deaminated, amino acids can enter the pathways of glucose metabolism as pyruvate, acetyl CoA, or several components of the citric acid cycle. For example, deaminated asparagine and aspartate are converted into oxaloacetate and enter glucose catabolism in the citric acid cycle. Deaminated amino acids can also be converted into another intermediate molecule before entering the pathways. Several amino acids can enter glucose catabolism at multiple locations.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44441/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Medical Physiology/Basic Biochemistry/Sugars. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Medical...ose_Metabolism. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycogen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glycogen. **License**: ***CC BY-SA: Attribution-ShareAlike***
- monosaccharide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/monosaccharide. **License**: ***CC BY-SA: Attribution-ShareAlike***
- disaccharide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/disaccharide. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medical Physiology/Basic Biochemistry/Sugars. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Medical...ose_Metabolism. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glycogen Structure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glycog..._structure.svg. **License**:
- Glycogen pathway.jpg. **Provided by**: OpenStax CNX. **Located at**: https://cnx.org/contents/ZP457F64@7/Connections-of-Carbohydrate-Pr. **License**: ***CC BY-SA: Attribution-ShareAlike***
- keto acid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/keto_acid. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44441/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- deamination. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/deamination. **License**: ***CC BY-SA: Attribution-ShareAlike***
- catabolism. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/catabolism. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Medical Physiology/Basic Biochemistry/Sugars. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Medical...ose_Metabolism. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glycogen Structure. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glycogen%23mediaviewer/File:Glycogen_structure.svg. **License**:
- Glycogen pathway.jpg. **Provided by**: OpenStax CNX. **Located at**: https://cnx.org/contents/ZP457F64@7/Connections-of-Carbohydrate-Pr. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Connections of Carbohydrate, Protein, and Lipid Metabolic Pathways. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44441/latest...e_07_06_01.jpg. **License**: ***CC BY: Attribution***

This page titled 23.6B: Connecting Proteins to Glucose Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.7: METABOLIC BODY STATES

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

23.7A: Catabolic-Anabolic Steady State
23.7B: Absorptive State
23.7C: Postabsorptive State

This page titled 23.7: Metabolic Body States is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.7A: Catabolic-Anabolic Steady State

Catabolic reactions that break complex molecules provide the energy needed by anabolic reactions to produce complex molecules.

Learning Objectives

- Differentiate between anabolism and catabolism

Key Points

- Catabolism is the process of transforming chemical fuels such as glucose into ATP (energy). Anabolism, the process of cell differentiation and growth, requires energy (ATP).
- Anabolism takes a few, basic raw materials and produces a wide variety of products such as peptides, proteins, polysaccharides, lipids, and nucleic acids.
- Catabolism and anabolism have separate metabolic pathways controlled by a distinct set of hormones.
- Growth hormone, testosterone, and estrogen are anabolic hormones. Adrenaline, cortisol, and glucagon are catabolic hormones.
- Glucose metabolism fluctuates with an individual's circadian rhythms which regulate anabolism and catabolism.
- Adrenaline, cortisol, and glucagon are catabolic hormones.
- Glucose metabolism fluctuates with an individual's circadian rhythms which regulate anabolism and catabolism.


Key Terms

- **anabolism**: The constructive metabolism of the body, as distinguished from catabolism.
- **circadian rhythms**: A circadian rhythm is any biological process that displays an endogenous, entrainable oscillation of about 24 hours.
- **catabolism**: The destructive metabolism, usually including the release of energy and breakdown of materials.

EXAMPLES

Babies experience a tremendous amount of growth during their first years, requiring that enough fuel be converted to the energy needed to facilitate this growth. Hence the reason that when most babies aren't sleeping, they are usually eating.

Anabolic reactions require energy. The chemical reaction where ATP changes to ADP supplies energy for this metabolic process. Cells can combine anabolic reactions with catabolic reactions that release energy to form an efficient energy cycle. The catabolic reactions transform chemical fuels into cellular energy, which is then used to initiate the energy-requiring anabolic reactions. ATP, a high energy molecule, couples anabolism by the release of free energy. This energy does not come through the breakage of phosphate bonds; instead, it is released from the hydration of the phosphate group.



Anabolism and Catabolism: Catabolic reactions release energy, while anabolic reactions use up energy.

Anabolism is the opposite of catabolism. For example, synthesizing glucose is an anabolic process, whereas the breaking down of glucose is a catabolic process. Anabolism requires the input of energy, described as an energy intake ("uphill") process. Catabolism is a "downhill" process where energy is released as the organism uses up energy. Anabolism and catabolism must be regulated to avoid the two processes occurring simultaneously. Each process has its own set of hormones that switch these processes on and off. Anabolic hormones include growth hormone, testosterone and estrogen. Catabolic hormones include adrenaline, cortisol and glucagon. The balance between anabolism and catabolism is also regulated by circadian rhythms, with processes such as glucose metabolism fluctuating to match an animal's normal periods of activity throughout the day.

Anabolism can be viewed as a set of metabolic processes in which the synthesis of complex molecules is initiated by energy released through catabolism. These complex molecules are produced through a systematic process from small and simple precursors. For example, an anabolic reaction can begin with relatively simple precursor molecules (created previously by catabolic reactions) and end with fairly complex products such as sugar, certain lipids, or even DNA, which has an extremely complex physical structure. The increased complexity of the products of anabolic reactions also means they are more energy-rich than their simple precursors.

Anabolic reactions constitute divergent processes. That is, relatively few types of raw materials are used to synthesize a wide variety of end products, resulting in an increase in cellular size, complexity, or both. Anabolic processes are responsible for cell differentiation and increases in body size. Bone mineralization and muscle mass are attributed to these processes. Anabolic processes produce peptides, proteins, polysaccharides, lipids and nucleic acids. These molecules comprise all the materials of living cells such as membranes and chromosomes, as well as specialized products of specific types of cells, such as enzymes, antibodies, hormones and neurotransmitters.

This page titled 23.7A: Catabolic-Anabolic Steady State is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.7B: Absorptive State

When the gastrointestinal tract is full, anabolism exceeds catabolism; this is the absorptive state.

Learning Objectives

- Differentiate among the nutrients in the absorptive state

Key Points

- During the absorptive state, anabolic processes use glucose in a variety of ways.
- In the liver, glucose is converted to glycogen or fat, which store energy for future use. Fat is also stored in adipose tissue and glycogen in muscle tissue.
- Glucose is also carried in the bloodstream to cells where it will be used to provide energy for cellular processes.
- Also during the absorptive state, chylomicrons, the product of fat digestion, are reconstituted to fat and stored in adipose tissue or, in a low carb environment, are used as an energy source.
- The liver deaminates amino acids to keto acids which can be used in the krebs cycle to produce ATP, or can be converted to fat, or can be used by other body cells to create proteins.

Key Terms

- **absorptive state**: The period during digestion when anabolism exceeds catabolism.
- **Kreb's cycle**: The Kreb's cycle is a series of chemical reactions used by all aerobic organisms to generate energy through the oxidization of acetate derived from carbohydrates, fats and proteins into carbon dioxide.
- **glycogen**: A polysaccharide that is the main form of carbohydrate storage in animals; converted to glucose as needed.

EXAMPLES

The baby who has finished nursing has a full tummy and now will probably fall asleep. During this sleep period, anabolic processes are busy building up stores of fats and glycogen that will be needed in the future to provide energy for the growing baby.

Absorptive state is the period in which the gastrointestinal tract is full and the anabolic processes exceed catabolism. The fuel used for this process is glucose.

Carbohydrates


Simple sugars are sent to the liver where they are converted to glucose. The glucose then travels to the blood or is converted to glycogen and fat (triglyceride) for energy storage. The glycogen and fat will be stored in the liver and adipose tissue, respectively, as reserves for the post-absorptive state. The remaining glucose is taken in for use by body cells or stored in skeletal muscle as glycogen.

Triglycerides

Chylomicrons are lipoprotein particles that consist of triglycerides (85-92%), phospholipids (6-12%), cholesterol (1-3%) and proteins (1–2%). This main product of fat digestion is first broken down to fatty acids and glycerol through hydrolysis using lipoprotein lipase. This allows them to pass freely through capillary walls. Most of this will be reconstituted as triglycerides and stored in adipose tissue. The rest will be used for energy in adipose cells, skeletal muscle and hepatocytes. In a low carb environment, other cells of the body will also begin to use triglycerides as energy sources.

Amino acids

The liver deaminates amino acids to keto acids to be used in the Kreb's cycle in order to generate energy in the form of ATP. They may also be converted to fat for energy storage. Some are used to make plasma proteins, but most leave through liver sinusoids to be used by body cells to construct proteins.



Glucose Metabolism: Glucose metabolism and various forms of it in the process.

This page titled 23.7B: Absorptive State is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.7C: Postabsorptive State

The post-absorptive state occurs around three to five hours after a meal has been completely digested and absorbed.

Learning Objectives

- Describe the postabsorptive state

Key Points

- Once a meal has been completely absorbed (typically three to five hours after a meal), the metabolism changes to a fasting state, which is synonymous with " post-absorptive state," in contrast to the "post-prandial" state of ongoing digestion.
- Post-absorptive plasma glucose concentration has been discovered to be physiologically maintained within the range of 70 mg/dl [3.9 mmol/l] to 110 mg/dl [6.1 mmol/l] in humans. This is accomplished via increased glucose levels from glucagon and decreased glucose levels from insulin.
- Chronic insulin and glucagon deficiencies have been proven to cause hyperglycemia and, therefore, suggesting that insulin is the predominant factor of postabsorptive glucose levels.

Key Terms

- **post-absorptive state**: The metabolic state achieved after complete digestion and absorption of a meal.

EXAMPLES

Three to five hours after nursing, the baby wakes up ready to nurse again.


In a physiological context, fasting may refer to:

1. The metabolic status of a person who has not eaten overnight.
2. The metabolic state achieved after complete digestion and absorption of a meal.

Several metabolic adjustments occur during fasting, and some diagnostic tests are used to determine a fasting state. For example, a person is assumed to be fasting after 8–12 hours. Metabolic changes toward the fasting state begin after absorption of a meal (typically three to five hours after a meal); "post-absorptive state" is synonymous with this usage, in contrast to the "post-prandial" state of ongoing digestion. A diagnostic fast refers to prolonged fasting (from 8–72 hours depending on age) conducted under observation for investigation of a problem, usually hypoglycemia. Finally, extended fasting has been recommended as therapy for various conditions by health professionals of most cultures, throughout history, from ancient to modern.

During fasting, post-absorptive state, fatty acid oxidation contributes proportionately more to energy expenditure than does carbohydrate oxidation. This phenomenon is due largely to greater lipid and lower carbohydrate availability, as plasma non-esterified fatty acid (NEFA) concentrations rise in response to lower insulin and higher counter-regulatory hormone concentrations.

Post-absorptive plasma glucose concentration has been discovered to be physiologically maintained within the range of 70 mg/dl [3.9 mmol/l] to 110 mg/dl [6.1 mmol/l] in humans. This is accomplished via increased glucose levels from glucagon and decreased glucose levels from insulin. However, there has not yet been any convincing evidence of the involvement of glucagon in post-absorptive plasma glucose concentration maintenance.



The flactuations of glucose and insulin in human during the course of a day: The fluctuation of blood sugar (red) and the sugar-lowering hormone insulin (blue) in humans during the course of a day with three meals. One of the effects of a sugar-rich vs a starch-rich meal is highlighted.



Postabsorptive State: Resting after absorption of meal.

Combined deficiency of insulin and glucagon results in an initial drop in plasma glucose levels, but is followed by an increase in plasma glucose levels. This indicates that there is support of post-absorptive plasma glucose concentrations from glucagon, when in concert with insulin. Changes in plasma glucose concentrations also result from changes in glucose production, but not from glucose utilization. Furthermore, during insulin and partial glucagon deficiency, and the exclusive partial deficiency of glucagon, the rate of glucose appearance increases to a point greater than the rate of glucose disappearance. This rate increase seems to be even larger than during insulin and glucagon deficiency, as well as when glucagon is made exclusively deficient. Both scenarios result in much higher plasma glucose concentrations.

Increases in plasma glucose levels are ultimately followed by plateaus. These plateaus occur within a postabsorptive physiological range, and after octreotide-induced suppression of insulin and glucagon secretion. It has been determined that hormones and additional factors are involved in postabsorptive glucose level maintenance, after short periods of time. However, chronic insulin and glucagon deficiencies still remain victims of diabetes. Therefore, insulin has been proven to contribute to the maintenance of postabsorptive plasma glucose concentrations, while high levels of glucagon are not required to onset diabetes.

These findings do not distinguish the individual roles of insulin and of glucagon. However, chronic insulin and glucagon deficiencies have been proven to cause hyperglycemia and, therefore, strongly suggest that insulin is the predominant factor of postabsorptive glucose levels.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Structural Biochemistry/Anabolism. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...stry/Anabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anabolism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- catabolism. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/catabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anabolism. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- circadian rhythms. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/circadian%20rhythms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Biochemistry/Metabolism and energy. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Biochem...and_catabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Chylomicrons. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chylomicrons**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Absorptive state. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Absorptive_state**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glycogen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glycogen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Kreb's cycle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Kreb's%20cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- absorptive state. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/absorptive%20state**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Biochemistry/Metabolism and energy. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Biochem...and_catabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glucose metabolism. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...metabolism.png**. **License**: ***Public Domain: No Known Copyright***
- Metabolomics/Hormones/Glucagon. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Metabol...mones/Glucagon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nutrition & Metabolism | Full text | Does prior acute exercise affect postexercise substrate oxidation in response to a high carbohydrate meal?. **Provided by**: Nutrition and Metabolism Journal. **Located at**: **http://www.nutritionandmetabolism.com/content/5/1/2**. **License**: ***CC BY: Attribution***
- Fasting. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fasting**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- post-absorptive state. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/post-absorptive%20state**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Biochemistry/Metabolism and energy. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Biochem...and_catabolism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Glucose metabolism. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...metabolism.png**. **License**: ***Public Domain: No Known Copyright***
- Suckale08 fig3 glucose insulin day. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Su...nsulin_day.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Baby sleeping. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...y_sleeping.jpg**. **License**: ***CC BY: Attribution***

This page titled 23.7C: Postabsorptive State is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.8: METABOLISM AND THE LIVER

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

23.8A: Cholesterol Metabolism

23.8B: Regulation of Blood Cholesterol Levels

This page titled 23.8: Metabolism and the Liver is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.8A: Cholesterol Metabolism

One quarter of body cholesterol is produced by the liver, and 50% of this is reabsorbed back into the circulation via the small intestine.

Learning Objectives

- Describe cholesterol metabolism

Key Points

- Cholesterol is also synthesized in the adrenal glands and reproductive organs.
- Cholesterol is kept in balance by homeostatic mechanisms: higher dietary intake leads to reduced synthesis in the body.
- Dietary cholesterol is poorly absorbed; therefore, intake of cholesterol has little effect on blood cholesterol levels.
- Cholesterol forms part of the bile acids excreted from the liver.
- Crystallized cholesterol is the primary component of gallstones.
- Cholesterol in the colon is metabolized by colonic bacteria which convert it to a nonabsorbable sterol that is excreted in the feces.


Key Terms

- **gallbladder**: In vertebrates, a small organ that aids mainly in fat digestion and concentrates bile produced by the liver.
- **cholesterol**: A sterol lipid synthesized by the liver and transported in the bloodstream to the membranes of all animal cells; it plays a central role in many biochemical processes and, as a lipoprotein that coats the walls of blood vessels, is associated with cardiovascular disease.
- **bile**: A bitter brownish-yellow or greenish-yellow secretion produced by the liver, stored in the gall bladder, and discharged into the duodenum, where it aids the process of digestion.

EXAMPLES


Eggs are an excellent, low-calorie source of protein, but they also are fairly high in cholesterol, so, in the past, it was thought that eggs should be limited in the diet. However, as we now know that cholesterol is kept in balance in the body, and that food intake has little effect on blood cholesterol concentrations, it appears that eggs can be a healthy part of one's diet.

About 20–25% of total daily cholesterol production occurs in the liver. Other sites of higher synthesis rates include the intestines, adrenal glands, and reproductive organs. Biosynthesis of cholesterol is directly regulated by the cholesterol levels present, although the homeostatic mechanisms involved are only partly understood. A higher intake from food leads to a net decrease in endogenous production, whereas lower intake from food has the opposite effect. However, most ingested cholesterol is esterified; esterified cholesterol is poorly absorbed. The body also compensates for any absorption of additional cholesterol by reducing cholesterol synthesis. For these reasons, cholesterol intake in food has little, if any, effect on total body cholesterol content or concentrations of cholesterol in the blood.



Cholesterol: A molecule of cholesterol consisting of four linked hydrocarbon rings forming the bulky steroid structure. There is a hydrocarbon tail linked to one end of the steroid and a hydroxyl group linked to the other end.

Cholesterol is recycled. The liver excretes it in a non-esterified form (via bile) into the digestive tract. Typically about 50% of the excreted cholesterol is reabsorbed by the small bowel back into the bloodstream.



Cholesterol Crystals: Microscopic appearance of cholesterol crystals in water. Photo taken under polarized light.

Cholesterol is oxidized by the liver into a variety of bile acids. These, in turn, are conjugated with glycine, taurine, glucuronic acid, or sulfate. A mixture of conjugated and nonconjugated bile acids, along with cholesterol itself, is excreted from the liver into the bile. Approximately 95% of the bile acids are reabsorbed from the intestines, and the remainder is lost in the feces. The excretion and reabsorption of bile acids forms the basis of the enterohepatic circulation, which is essential for the digestion and absorption of dietary fats. Under certain circumstances, when more concentrated, as in the gallbladder, cholesterol crystallizes and is the major constituent of most gallstones. Lecithin and bilirubin gallstones also occur, but less frequently. Every day, up to one gram of cholesterol enters the colon. This cholesterol originates from the diet, bile, and desquamated intestinal cells; it can be metabolized by the colonic bacteria. Cholesterol is converted mainly into coprostanol, a nonabsorbable sterol that is excreted in the feces. A cholesterol-reducing bacterium origin has been isolated from human feces.

This page titled 23.8A: Cholesterol Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.8B: Regulation of Blood Cholesterol Levels

Cholesterol is transported through the blood by lipoproteins which direct cholesterol to where it is needed.

Learning Objectives

- Differentiate among the cholesterol transport lipoproteins

Key Points

- There are several types of lipoproteins that are classified by their density or ratio of lipid to protein. The lower the protein, the less dense it is.
- LDL lipoproteins contain the highest percentage of cholesterol and are the major carriers through the bloodstream.
- LDL and its receptor form a vesicle which fuses with a lysosome that releases the cholesterol for use in membrane biosynthesis or cellular storage.
- HDL lipoproteins transport cholesterol back to the liver for excretion or reprocessing.
- High levels of HDL lipoproteins are associated with better health outcomes.
- LDL lipoproteins contain the highest percentage of cholesterol and are the major carriers through the bloodstream.
- LDL and its receptor form a vesicle which fuses with a lysosome that releases the cholesterol to be used in membrane biosynthesis or to be stored in the cell.
- When a cell has sufficient cholesterol LDL receptors are blocked, but when the cell is deficient, more LDL receptors are created.
- If the system breaks down, LDL lipoproteins accumulate in the blood, are engulfed by macrophages, and may become trapped on the walls of blood vessels to form plaques.
- HDL lipoproteins transport cholesterol back to the liver for excretion or reprocessing.
- High levels of HDL lipoproteins are associated with better health outcomes.

Key Terms


- **lysosome**: An organelle found in all types of animal cells which contains a large range of digestive enzymes capable of splitting most biological macromolecules.
- **lipoprotein**: Any of a large group of complexes of protein and lipid with many biochemical functions.
- **chylomicrons**: Chylomicrons are one of the five major groups of lipoproteins (chylomicrons, VLDL, IDL, LDL, HDL) that enable fats and cholesterol to move within the water-based solution of the bloodstream.

EXAMPLES

A cholesterol test will measure your HDL, LDL, VLDL, and triglyceride levels. The ratio of HDL to LDL is a useful screening method to determine your possible risk for heart disease or stroke.

Blood Cholesterol Levels and Transport

Cholesterol is only slightly soluble in water; it can dissolve and travel in the water-based bloodstream at exceedingly small concentrations. Since cholesterol is insoluble in blood, it is transported in the circulatory system within lipoproteins, which are complex discoidal particles that have an exterior composed of amphiphilic proteins and lipids whose outward-facing surfaces are water-soluble and inward-facing surfaces are lipid-soluble; triglycerides and cholesterol esters are carried internally. Phospholipids and cholesterol, being amphipathic, are transported in the surface monolayer of the lipoprotein particle.



Lipid vesicles are aqueous compartments enclosed by a lipid bilayer. One layer of "hydrophilic heads" faces outside the cell attracted to the water in the environment, whereas another layer of "heads" faces inside attracted to water inside the cell. "Hydrophobic tails" of one layer face the "tails" of the other layer , lining up to form a surface away from the aqueous environment.

liposome structure: Lipid vesicles (or liposomes) are small, intracellular, membrane-enclosed bubbles of liquid within a cell. They are formed because of the properties of lipid membranes, where the hydrophobic chains are packed together so they are not in contact with the aqueous solution.

In addition to providing a soluble means for transporting cholesterol through the blood, lipoproteins have cell-targeting signals that direct the lipids they carry to certain tissues. For this reason, there are several types of lipoproteins within the blood called, in order of increasing density:

1. chylomicrons
2. very-low-density lipoprotein (VLDL)
3. intermediate-density lipoprotein (IDL)
4. low-density lipoprotein (LDL), and
5. high-density lipoprotein (HDL).

The more lipid and less protein a lipoprotein has, the less dense it is. The cholesterol within all the various lipoproteins is identical, although some cholesterol is carried as the “free” alcohol and some is carried as fatty acyl esters referred to as cholesterol esters. However, the different lipoproteins contain apolipoproteins, which serve as ligands for specific receptors on cell membranes. In this way, the lipoprotein particles are molecular addresses that determine the start and endpoints for cholesterol transport.

Chylomicrons, the least dense type of cholesterol transport molecules, contain apolipoprotein B-48, apolipoprotein C, and apolipoprotein E in their shells. Chylomicrons are the transporters that carry fats from the intestine to muscle and other tissues that need fatty acids for energy or fat production. Cholesterol that is not used by muscles remains in more cholesterol-rich chylomicron remnants, which are taken up from here to the bloodstream by the liver.

VLDL molecules are produced by the liver and contain excess triacylglycerol and cholesterol that is not required by the liver for synthesis of bile acids. These molecules contain apolipoprotein B100 and apolipoprotein E in their shells. During transport in the bloodstream, the blood vessels cleave and absorb more triacylglycerol from IDL molecules, which contain an even higher percentage of cholesterol. The IDL molecules have two possible fates: half are metabolized by HTGL, taken up by the LDL receptor on the liver cell surfaces and the other half continue to lose triacylglycerols in the bloodstream until they form LDL molecules, which have the highest percentage of cholesterol within them.

LDL molecules, therefore, are the major carriers of cholesterol in the blood, and each one contains approximately 1,500 molecules of cholesterol ester. The shell of the LDL molecule contains just one molecule of apolipoprotein B100, which is recognized by the LDL receptor in peripheral tissues. Upon binding of apolipoprotein B100, many LDL receptors become localized in clathrin-coated pits. Both the LDL and its receptor are internalized by endocytosis to form a vesicle within the cell. The vesicle then fuses with a lysosome, which has an enzyme called lysosomal acid lipase that hydrolyzes the cholesterol esters. Now within the cell, the cholesterol can be used for membrane biosynthesis or esterified and stored within the cell, so as to not interfere with cell membranes.

Synthesis of the LDL receptor is regulated by SREBP, the same regulatory protein as was used to control synthesis of cholesterol de novo in response to cholesterol presence in the cell. When the cell has abundant cholesterol, LDL receptor synthesis is blocked so new cholesterol in the form of LDL molecules cannot be taken up. On the converse, more LDL receptors are made when the cell is deficient in cholesterol. When this system is deregulated, many LDL molecules appear in the blood without receptors on the peripheral tissues. These LDL molecules are oxidized and taken up by macrophages, which become engorged and form foam cells. These cells often become trapped in the walls of blood vessels and contribute to artherosclerotic plaque formation. Differences in cholesterol homeostasis affect the development of early atherosclerosis (carotid intima-media thickness). These plaques are the main causes of heart attacks, strokes, and other serious medical problems, leading to the association of so-called LDL cholesterol (actually a lipoprotein) with “bad” cholesterol.

Also, HDL particles are thought to transport cholesterol back to the liver for excretion or to other tissues that use cholesterol to synthesize hormones in a process known as reverse cholesterol transport (RCT). Having large numbers of large HDL particles correlates with better health outcomes. In contrast, having small numbers of large HDL particles is independently associated with atheromatous disease progression within the arteries.



Cholesterol Blood Tests: Reference ranges for blood tests, showing usual, as well as optimal, levels of HDL, LDL and total cholesterol in mass and molar concentrations, is found in orange color at right, that is, among the blood constituents with the highest concentration.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholest...23Biosynthesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholest..._and_excretion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bile. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/bile**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...on/gallbladder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cholesterol. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cholesterol**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholesterol**. **License**: ***CC BY: Attribution***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholesterol**. **License**: ***CC BY: Attribution***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholest..._of_absorption**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lipoprotein. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lipoprotein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lysosome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lysosome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chylomicrons. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/chylomicrons**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholesterol**. **License**: ***CC BY: Attribution***
- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholesterol**. **License**: ***CC BY: Attribution***
- Tina Liposome. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...a_Liposome.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Cholesterol. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cholest..._of_absorption**. **License**: ***Public Domain: No Known Copyright***

This page titled 23.8B: Regulation of Blood Cholesterol Levels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.9: METABOLIC RATE AND THERMOREGULATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

23.9A: Homeostatic Process
23.9B: Homeostasis: Thermoregulation
23.9C: Heat Conservation and Dissipation

This page titled 23.9: Metabolic Rate and Thermoregulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.9A: Homeostatic Process

Learning Objectives

- Give an example and describe a homeostatic process.

The human organism consists of trillions of cells working together for the maintenance of the entire organism. While cells may perform very different functions, the cells are quite similar in their metabolic requirements. Maintaining a constant internal environment with everything that the cells need to survive (oxygen, glucose, mineral ions, waste removal, etc.) is necessary for the well-being of individual cells and the well-being of the entire body. The varied processes by which the body regulates its internal environment are collectively referred to as homeostasis.


Homeostasis

Homeostasis, in a general sense, refers to stability, balance, or equilibrium. Physiologically, it is the body's attempt to maintain a constant and balanced internal environment, which requires persistent monitoring and adjustments as conditions change. Adjustment of physiological systems within the body is called homeostatic regulation, which involves three parts or mechanisms: (1) the receptor, (2) the control center, and (3) the effector.

The receptor receives information that something in the environment is changing. The control center or integration center receives and processes information from the receptor. The effector responds to the commands of the control center by either opposing or enhancing the stimulus. This ongoing process continually works to restore and maintain homeostasis. For example, during body temperature regulation, temperature receptors in the skin communicate information to the brain (the control center) which signals the effectors: blood vessels and sweat glands in the skin. As the internal and external environment of the body are constantly changing, adjustments must be made continuously to stay at or near a specific value: the set point.

Purpose of Homeostasis

The ultimate goal of homeostasis is the maintenance of equilibrium around the set point. While there are normal fluctuations from the set point, the body's systems will usually attempt to revert to it. A change in the internal or external environment (a stimulus) is detected by a receptor; the response of the system is to adjust the deviation parameter toward the set point. For instance, if the body becomes too warm, adjustments are made to cool the animal. If the blood's glucose rises after a meal, adjustments are made to lower the blood glucose level by moving the nutrient into tissues in the command center that require it, or to store it for later use.



***Blood glucose homeostasis**: An example of how homeostasis is achieved by controlling blood sugar levels after a meal.*

Key Points

- Homeostasis is the body's attempt to maintain a constant and balanced internal environment, which requires persistent monitoring and adjustments as conditions change.
- Homeostatic regulation is monitored and adjusted by the receptor, the command center, and the effector.
- The receptor receives information based on the internal environment; the command center, receives and processes the information; and the effector responds to the command center, opposing or enhancing the stimulus.

Key Terms

- **homeostasis**: the ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium
- **effector**: any muscle, organ etc. that can respond to a stimulus from a nerve

This page titled 23.9A: Homeostatic Process is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.9B: Homeostasis: Thermoregulation

Animals use different modes of thermoregulation processes to maintain homeostatic internal body temperatures.

Learning Objectives

- Outline the various types of processes utilized by animals to ensure thermoregulation.

Key Points

- In response to varying body temperatures, processes such as enzyme production can be modified to acclimate to changes in the temperature.
- Endotherms regulate their own internal body temperature, regardless of fluctuating external temperatures, while ectotherms rely on the external environment to regulate their internal body temperature.
- Homeotherms maintain their body temperature within a narrow range, while poikilotherms can tolerate a wide variation in internal body temperature, usually because of environmental variation.
- Heat can be exchanged between environment and animals via radiation, evaporation, convection, or conduction processes.

Key Terms

- **ectotherm**: An animal that relies on external environment to regulate its internal body temperature.
- **endotherm**: An animal that regulates its own internal body temperature through metabolic processes.
- **homeotherm**: An animal that maintains a constant internal body temperature, usually within a narrow range of temperatures.
- **poikilotherm**: An animal that varies its internal body temperature within a wide range of temperatures, usually as a result of variation in the environmental temperature.

Thermoregulation to Maintain Homeostasis

Internal thermoregulation contributes to animal's ability to maintain homeostasis within a certain range of temperatures. As internal body temperature rises, physiological processes are affected, such as enzyme activity. Although enzyme activity initially increases with temperature, enzymes begin to denature and lose their function at higher temperatures (around 40-50 C for mammals). As internal body temperature decreases below normal levels, hypothermia occurs and other physiological process are affected. There are various thermoregulation mechanisms that animals use to regulate their internal body temperature.

Types of Thermoregulation (Ectothermy vs. Endothermy)

Thermoregulation in organisms runs along a spectrum from endothermy to ectothermy. Endotherms create most of their heat via metabolic processes, and are colloquially referred to as "warm-blooded." Ectotherms use external sources of temperature to regulate their body temperatures. Ectotherms are colloquially referred to as "cold-blooded" even though their body temperatures often stay within the same temperature ranges as warm-blooded animals.

Ectotherm




Ectotherm: The Common frog is an ecotherm and regulates its body based on the temperature of the external environment.

An ectotherm, from the Greek (ektós) “outside” and (thermós) “hot,” is an organism in which internal physiological sources of heat are of relatively small or quite negligible importance in controlling body temperature. Since ectotherms rely on environmental heat sources, they can operate at economical metabolic rates. Ectotherms usually live in environments in which temperatures are constant, such as the tropics or ocean. Ectotherms have developed several behavioral thermoregulation mechanisms, such as basking in the sun to increase body temperature or seeking shade to decrease body temperature.

Endotherms

In contrast to ectotherms, endotherms regulate their own body temperature through internal metabolic processes and usually maintain a narrow range of internal temperatures. Heat is usually generated from the animal’s normal metabolism, but under conditions of excessive cold or low activity, an endotherm generate additional heat by shivering. Many endotherms have a larger number of mitochondria per cell than ectotherms. These mitochondria enables them to generate heat by increasing the rate at which they metabolize fats and sugars. However, endothermic animals must sustain their higher metabolism by eating more food more often. For example, a mouse (endotherm) must consume food every day to sustain high its metabolism, while a snake (ectotherm) may only eat once a month because its metabolism is much lower.

Homeothermy vs. Poikilothermy




Homeotherm vs. Poikilotherm: Sustained energy output of an endothermic animal (mammal) and an ectothermic animal (reptile) as a function of core temperature. In this scenario, the mammal is also a homeotherm because it maintains its internal body temperature in a very narrow range. The reptile is also a poikilotherm because it can withstand a large range of temperatures.

A poikilotherm is an organism whose internal temperature varies considerably. It is the opposite of a homeotherm, an organism which maintains thermal homeostasis. Poikilotherm's internal temperature usually varies with the ambient environmental temperature, and many terrestrial ectotherms are poikilothermic. Poikilothermic animals include many species of fish, amphibians, and reptiles, as well as birds and mammals that lower their metabolism and body temperature as part of hibernation or torpor. Some ectotherms can also be homeotherms. For example, some species of tropical fish inhabit coral reefs that have such stable ambient temperatures that their internal temperature remains constant.

Means of Heat Transfer

Heat can be exchanged between an animal and its environment through four mechanisms: radiation, evaporation, convection, and conduction. Radiation is the emission of electromagnetic "heat" waves. Heat radiates from the sun and from dry skin the same manner. When a mammal sweats, evaporation removes heat from a surface with a liquid. Convection currents of air remove heat from the surface of dry skin as the air passes over it. Heat can be conducted from one surface to another during direct contact with the surfaces, such as an animal resting on a warm rock.



Mechanisms for heat exchange: Heat can be exchanged by four mechanisms: (a) radiation, (b) evaporation, (c) convection, or (d) conduction.

This page titled 23.9B: Homeostasis: Thermoregulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

23.9C: Heat Conservation and Dissipation

Animals have processes that allow for heat conservation and dissipation in order to maintain a homeostatic internal body temperature.

Learning Objectives

- Describe some of the changes animals use in order to maintain body temperature

Key Points

- Heat conservation is characterized by the ability to ensure blood remains in the core by undergoing vasoconstriction, reducing blood flow to the periphery (also known as peripheral vasoconstriction).
- Heat dissipation is characterized by the ability to undergo vasodilation which increases blood flow to the periphery, resulting in evaporative heat loss.
- Endothermic animals are defined by their ability to utilize both vasoconstriction and vasodilation to maintain internal body temperature.
- Ectothermic animals are defined by their change in behavior (lying in sunlight to warm up, hiding in shade to cool down) to regulate body temperature.


Key Terms

- **endotherm**: a warm-blooded animal that maintains a constant body temperature
- **ectotherm**: a cold-blooded animal that regulates its body temperature by exchanging heat with its surroundings

Heat Conservation and Dissipation

Animals conserve or dissipate heat in a variety of ways. In certain climates, endothermic animals have some form of insulation, such as fur, fat, feathers, or some combination thereof. Animals with thick fur or feathers create an insulating layer of air between their skin and internal organs. Polar bears and seals live and swim in a subfreezing environment, yet they maintain a constant, warm, body temperature. The arctic fox uses its fluffy tail as extra insulation when it curls up to sleep in cold weather. Mammals have a residual effect from shivering and increased muscle activity: arrector pili muscles create “goose bumps,” causing small hairs to stand up when the individual is cold; this has the intended effect of increasing body temperature. Mammals use layers of fat to achieve the same end; the loss of significant amounts of body fat will compromise an individual’s ability to conserve heat.

Endotherms use their circulatory systems to help maintain body temperature. For example, vasodilation brings more blood and heat to the body surface, facilitating radiation and evaporative heat loss, which helps to cool the body. However, vasoconstriction reduces blood flow in peripheral blood vessels, forcing blood toward the core and the vital organs found there, conserving heat. Some animals have adaptions to their circulatory system that enable them to transfer heat from arteries to veins, thus, warming blood that returns to the heart. This is called a countercurrent heat exchange; it prevents the cold venous blood from cooling the heart and other internal organs. This adaption, which can be shut down in some animals to prevent overheating the internal organs, is found in many animals, including dolphins, sharks, bony fish, bees, and hummingbirds. In contrast, similar adaptations (as in dolphin flukes and elephant ears) can help cool endotherms when needed.



Control of body temperature: In endotherms, the circulatory system is used to help maintain body temperature, either by vasodilation or vasoconstriction.

Many animals, especially mammals, use metabolic waste heat as a heat source. When muscles are contracted, most of the energy from the ATP used in muscle actions is wasted energy that translates into heat. In cases of severe cold, a shivering reflex is activated that generates heat for the body. Many species also have a type of adipose tissue called brown fat that specializes in generating heat.

Ecothermic animals use changes in their behavior to help regulate body temperature. For example, a desert ectothermic animal may simply seek cooler areas during the hottest part of the day in the desert to keep from becoming too warm. The same animals may climb onto rocks to capture heat during a cold desert night. Some animals seek water to aid evaporation in cooling them, as seen with reptiles. Other ectotherms use group activity, such as the activity of bees to warm a hive to survive winter.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44733/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...gy/Homeostasis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homeostasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/homeostasis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- effector. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/effector. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 25, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44733/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 26, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44733/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- pyrogen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/pyrogen. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vasoconstriction. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasoconstriction. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vasodilation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasodilation. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- Ectotherm. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ectotherm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thermoregulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thermoregulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Endotherm. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Endotherm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 25, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44733/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Sustained%2525252520energy%2525252520output%2525252520of%2525252520an%2525252520endothermic%25252 **Provided by**: Wikimedia%2525252520commons. **Located at**: **en.Wikipedia.org/wiki/Endothe...kilothermy.png**. **License**: ***Public Domain: No Known Copyright***
- data-attribution-url=cnx.org/content/m44733/latest...e_33_03_03.jpg. **Provided by**: Connexions. **License**: ***CC BY: Attribution***
- Theu00a0Common frogu00a0is a poikilotherm and is able to function over a wide range of body core temperatures.. **Provided by**: Wikimedia commons. **Located at**: **en.Wikipedia.org/wiki/Poikilo...frosch_ies.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44733/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 23, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44733/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- ectotherm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ectotherm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endotherm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endotherm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 25, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44733/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Sustained%2525252520energy%2525252520output%2525252520of%2525252520an%2525252520endothermic%25252 **Provided by**: Wikimedia%2525252520commons. **Located at**: **en.Wikipedia.org/wiki/Endothe...kilothermy.png**. **License**: ***Public Domain: No Known Copyright***
- data-attribution-url=cnx.org/content/m44733/latest...e_33_03_03.jpg. **Provided by**: Connexions. **License**: ***CC BY: Attribution***
- Theu00a0Common frogu00a0is a poikilotherm and is able to function over a wide range of body core temperatures.. **Provided by**: Wikimedia commons. **Located at**: **en.Wikipedia.org/wiki/Poikilo...frosch_ies.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Homeostasis. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44733/latest...e_33_03_04.png**. **License**: ***CC BY: Attribution***

This page titled 23.9C: Heat Conservation and Dissipation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24: URINARY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

24.1: Overview of the Urinary System
24.1A: Overview of the Urinary System

24.2: The Kidneys
24.2A: Location and External Anatomy of the Kidneys
24.2B: Internal Anatomy of the Kidneys
24.2C: Supply of Blood and Nerves to the Kidneys
24.2D: Nephron, Parts, and Histology

24.3: Physiology of the Kidneys
24.3A: Overview of Urine Formation
24.3B: Glomerular Filtration
24.3C: Regulation of Glomerular Filtration Rate
24.3D: Tubular Reabsorption
24.3E: Tubular Secretion

24.4: Urine
24.4A: Physical Characteristics of Urine
24.4B: Chemical Composition of Urine
24.4C: Regulation of Urine Concentration and Volume
24.4D: Urinalysis
24.4E: Renal Clearance

24.5: Urine Transport, Storage, and Elimination
24.5A: Overview of Urine Transport, Storage, and Elimination
24.5B: Ureters
24.5C: Urinary Bladder
24.5D: Urethra
24.5E: Micturition and the Micturition Reflex
24.5F: Waste Management in Other Body Systems

This page titled 24: Urinary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.1: OVERVIEW OF THE URINARY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

24.1A: Overview of the Urinary System

This page titled 24.1: Overview of the Urinary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.1A: Overview of the Urinary System

The urinary system maintains blood homeostasis by filtering out excess fluid and other substances from the bloodstream and secreting waste.

Learning Objectives

- Review the urinary system

Key Points

- The renal system eliminate wastes from the body, controls levels of electrolytes and metabolites, controls the osmoregulation of blood volume and pressure, and regulates blood pH.
- The renal system organs include the kidneys, ureter, bladder, and urethra. Nephrons are the main functional component of the kidneys.
- The respiratory and cardiovascular systems have certain functions that overlap with renal system functions.
- Metabolic wastes and excess ions are filtered out of the blood, combined with water, and leave the body in the form of urine.
- A complex network of hormones controls the renal system to maintain homeostasis.


Key Terms

- **ureter**: These are two long, narrow ducts that carry urine from the kidneys to the urinary bladder.
- **osmoregulation**: The most important function of the renal system, in which blood volume, blood pressure, and blood osmolarity (ion concentration) is maintained in homeostasis.

The Renal System

The renal system, which is also called the urinary system, is a group of organs in the body that filters out excess fluid and other substances from the bloodstream. The purpose of the renal system is to eliminate wastes from the body, regulate blood volume and pressure, control levels of electrolytes and metabolites, and regulate blood pH.

The renal system organs include the kidneys, ureters, bladder, and urethra. Metabolic wastes and excess ions are filtered out of the blood, along with water, and leave the body in the form of urine.



Components of the renal system: Here are the major organs of the renal system.

Renal System Functions

The renal system has many functions. Many of these functions are interrelated with the physiological mechanisms in the cardiovascular and respiratory systems.

1. Removal of metabolic waste products from the body (mainly urea and uric acid).
2. Regulation of electrolyte balance (e.g., sodium, potassium, and calcium).
3. Osmoregulation controls the blood volume and body water contents.
4. Blood pressure homeostasis: The renal system alters water retention and thirst to slowly change blood volume and keep blood pressure in a normal range.
5. Regulation of acid-base homeostasis and blood pH, a function shared with the respiratory system.

Many of these functions are related to one another as well. For example, water follows ions via an osmotic gradient, so mechanisms that alter sodium levels or sodium retention in the renal system will alter water retention levels as well.

Organs of the Renal System

Kidneys and Nephrons

Kidneys are the most complex and critical part of the urinary system. The primary function of the kidneys is to maintain a stable internal environment (homeostasis) for optimal cell and tissue metabolism. The kidneys have an extensive blood supply from the renal arteries that leave the kidneys via the renal vein.

Nephrons are the main functional component inside the parenchyma of the kidneys, which filter blood to remove urea, a waste product formed by the oxidation of proteins, as well as ions like potassium and sodium. The nephrons are made up of a capsule capillaries (the glomerulus) and a small renal tube.

The renal tube of the nephron consists of a network of tubules and loops that are selectively permeable to water and ions. Many hormones involved in homeostasis will alter the permeability of these tubules to change the amount of water that is retained by the body.

Ureter

Urine passes from the renal tube through tubes called ureters and into the bladder.

Bladder

The bladder is flexible and is used as storage until the urine is allowed to pass through the urethra and out of the body.

Urethra

The female and male renal system are very similar, differing only in the length of the urethra.

Human Osmoregulation

The kidneys play a very large role in human osmoregulation by regulating the amount of water reabsorbed from the glomerular filtrate in kidney tubules, which is controlled by hormones such as antidiuretic hormone (ADH), renin, aldosterone, and angiotensin I and II.

A basic example is that a decrease in water concentration of blood is detected by osmoreceptors in the hypothalamus, which stimulates ADH release from the pituitary gland to increase the permeability of the wall of the collecting ducts and tubules in the nephrons. Therefore, a large proportion of water is reabsorbed from fluid to prevent a fair proportion of water from being excreted.

The extent of blood volume and blood pressure regulation facilitated by the kidneys is a complex process. Besides ADH secretion, the renin-angiotensin feedback system is critically important to maintain blood volume and blood pressure homeostasis.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Introduction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- urinary system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/urinary%20system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ureter. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ureter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human%20Physiology/The%20Urinary%20System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Introduction**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 24.1A: Overview of the Urinary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.2: THE KIDNEYS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

24.2A: Location and External Anatomy of the Kidneys

24.2B: Internal Anatomy of the Kidneys

24.2C: Supply of Blood and Nerves to the Kidneys

24.2D: Nephron, Parts, and Histology

This page titled 24.2: The Kidneys is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.2A: Location and External Anatomy of the Kidneys

The kidneys are located at the rear wall of the abdominal cavity and they are protected by the ribcage.

Learning Objectives

- Describe the external anatomy and location of the kidneys

Key Points

- The kidneys lie in the lower abdominal cavity, on its rear wall.
- Adhering to the surface of each kidney are two layers of fat to help cushion them.
- The kidneys are located at the rear wall of the abdominal cavity just above the waistline and are protected by the ribcage. They are considered retroperitoneal, which means that they lie behind the peritoneum.
- The renal artery connects the kidneys to the aorta, while the renal vein connects the kidneys to the inferior vena cava.
- The adrenal glands lie superior to the kidneys.

Key Terms

- **adrenal gland**: Two glands that are responsible for releasing hormones in response to stress, such as catecholamines, and that also produce aldosterone, a renal system hormone.

The Kidneys


The kidneys are the primary functional organ of the renal system. They are essential in homeostatic functions such as the regulation of electrolytes, maintenance of acid–base balance, and the regulation of blood pressure (by maintaining salt and water balance). They serve the body as a natural filter of the blood and remove wastes that are excreted through the urine.

They are also responsible for the reabsorption of water, glucose, and amino acids, and will maintain the balance of these molecules in the body. In addition, the kidneys produce hormones including calcitriol, erythropoietin, and the enzyme renin, which are involved in renal and hemotological physiological processes.

Anatomical Location

The kidneys are a pair of bean-shaped, brown organs about the size of your fist. They are covered by the renal capsule, which is a tough capsule of fibrous connective tissue. Adhering to the surface of each kidney are two layers of fat to help cushion them.

The asymmetry within the abdominal cavity caused by the liver typically results in the right kidney being slightly lower than the left, and left kidney being located slightly more medial than the right. The right kidney sits just below the diaphragm and posterior to the liver, the left below the diaphragm and posterior to the spleen.



The kidneys: Human kidneys viewed from behind with the spine removed.

Resting on top of each kidney is an adrenal gland (adrenal meaning on top of renal), which are involved in some renal system processes despite being a primarily endocrine organ. The upper parts of the kidneys are partially protected by lower ribs, and each whole kidney and adrenal gland are surrounded by two layers of fat (the perirenal and pararenal fat) and the renal fascia.

The kidneys are located at the rear wall of the abdominal cavity just above the waistline and are protected by the ribcage. They are considered retroperitoneal, which means that they lie behind the peritoneum, the membrane lining of the abdominal cavity.

There are a number of important external structures connecting the kidneys to the rest of the body. The renal artery branches off from the lower part of the aorta and provides the blood supply to the kidneys. Renal veins take blood away from the kidneys into the inferior vena cava. The ureters are structures that come out of the kidneys, bringing urine downward into the bladder.

This page titled 24.2A: Location and External Anatomy of the Kidneys is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.2B: Internal Anatomy of the Kidneys

The cortex and medulla make up two of the internal layers of a kidney and are composed of individual filtering units known as nephrons.

Learning Objectives

- Distinguish between the cortex and medulla in the internal anatomy of the kidney

Key Points

- The renal cortex, renal medulla, and renal pelvis are the three main internal regions found in a kidney.
- Nephrons, masses of tiny tubules, are largely located in the medulla and receive fluid from the blood vessels in the renal cortex.
- The renal cortex produces erythropotein.
- The kidneys are made up by three external layers, which include the renal fascia (the outermost layer), the perirenal fat capsule, and lastly, the innermost layer, the renal capsule, which then surround the space of the renal cortex.
- The Bowman's or renal capsule, located at one end of each nephron in the cortex of the kidney, is the blood-filtering region of the nephron.
- The renal pelvis contains a hilium —the concave site in which the renal artery and vein and nerves enter the kidney and the ureter leaves the kidney.

Key Terms

- **renal medulla**: The inner-most region of the kidney, arranged into pyramid-like structures, that consists of the bulk of nephron structure.
- **renal cortex**: The outer region of the kidney, between the renal capsule and the renal medulla, that consists of a space that contains blood vessels that connect to the nephrons.
- **nephron**: The basic structural and functional unit of the kidney that filters the blood in order to regulate chemical concentrations and produce urine.

There are three major regions of the kidney:


1. Renal cortex
2. Renal medulla
3. Renal pelvis

The renal cortex is a space between the medulla and the outer capsule. The renal medulla contains the majority of the length of nephrons, the main functional component of the kidney that filters fluid from blood. The renal pelvis connects the kidney with the circulatory and nervous systems from the rest of the body.

Renal Cortex

The kidneys are surrounded by a renal cortex, a layer of tissue that is also covered by renal fascia (connective tissue) and the renal capsule. The renal cortex is granular tissue due to the presence of nephrons—the functional unit of the kidney—that are located deeper within the kidney, within the renal pyramids of the medulla.

The cortex provides a space for arterioles and venules from the renal artery and vein, as well as the glomerular capillaries, to perfuse the nephrons of the kidney. Erythropotein, a hormone necessary for the synthesis of new red blood cells, is also produced in the renal cortex.



Kidney structure: The kidney is made up of three main areas: the outer cortex, a medulla in the middle, and the renal pelvis.

Renal Medulla

The medulla is the inner region of the parenchyma of the kidney.
The medulla consists of multiple pyramidal tissue masses, called the renal pyramids, which are triangle structures that contain a dense network of nephrons.

At one end of each nephron, in the cortex of the kidney, is a cup-shaped structure called the Bowman's capsule. It surrounds a tuft of capillaries called the glomerulus that carries blood from the renal arteries into the nephron, where plasma is filtered through the capsule.

After entering the capsule, the filtered fluid flows along the proximal convoluted tubule to the loop of Henle and then to the distal convoluted tubule and the collecting ducts, which flow into the ureter. Each of the different components of the nephrons are selectively permeable to different molecules, and enable the complex regulation of water and ion concentrations in the body.

Renal Pelvis

The renal pelvis contains the hilium. The hilum is the concave part of the bean-shape where blood vessels and nerves enter and exit the kidney; it is also the point of exit for the ureters—the urine-bearing tubes that exit the kidney and empty into the urinary bladder. The renal pelvis connects the kidney to the rest of the body.

This page titled 24.2B: Internal Anatomy of the Kidneys is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.2C: Supply of Blood and Nerves to the Kidneys

The renal veins drain the kidney and the renal arteries supply blood to the kidney.

Learning Objectives

- Summarize the blood and nerve supply of the kidney

Key Points

- The renal arteries branch off of the abdominal aorta and supply the kidneys with blood. The arterial supply of the kidneys varies from person to person, and there may be one or more renal arteries to supply each kidney.
- The renal veins are the veins that drain the kidneys and connect them to the inferior vena cava.
- The kidney and the nervous system communicate via the renal plexus. The sympathetic nervous system will trigger vasoconstriction and reduce renal blood flow, while parasympathetic nervous stimulation will trigger vasodilation and increased blood flow.
- Afferent arterioles branch into the glomerular capillaries, while efferent arterioles take blood away from the glomerular capillaries and into the interlobular capillaries that provide oxygen to the kidney.

Key Terms

- **renal artery**: These arise off the side of the abdominal aorta, immediately below the superior mesenteric artery, and supply the kidneys with blood.
- **renal vein**: The veins that drain the kidney and connect the kidney to the inferior vena cava.

Because the kidney filters blood, its network of blood vessels is an important component of its structure and function. The arteries, veins, and nerves that supply the kidney enter and exit at the renal hilum.

Renal Arteries

The renal arteries branch off of the abdominal aorta and supply the kidneys with blood. The arterial supply of the kidneys is variable from person to person, and there may be one or more renal arteries supplying each kidney.

Due to the position of the aorta, the inferior vena cava, and the kidneys in the body, the right renal artery is normally longer than the left renal artery. The renal arteries carry a large portion of the total blood flow to the kidneys—up to a third of the total cardiac output can pass through the renal arteries to be filtered by the kidneys.

Renal blood supply starts with the branching of the aorta into the renal arteries (which are each named based on the region of the kidney they pass through) and ends with the exiting of the renal veins to join the inferior vena cava. The renal arteries split into several segmental arteries upon entering the kidneys, which then split into several arterioles.

These afferent arterioles branch into the glomerular capillaries, which facilitate fluid transfer to the nephrons inside the Bowman's capsule, while efferent arterioles take blood away from the glomerulus, and into the interlobular capillaries, which provide tissue oxygenation to the parenchyma of the kidney.


Renal Veins

The renal veins are the veins that drain the kidneys and connect them to the inferior vena cava. The renal vein drains blood from venules that arise from the interlobular capillaries inside the parenchyma of the kidney.

Renal Plexus

The renal plexus are the source of nervous tissue innervation within the kidney, which surround and primarily alter the size of the arterioles within the renal cortex. Input from the sympathetic nervous system triggers vasoconstriction of the arterioles in the kidney, thereby reducing renal blood flow into the glomerulus.

The kidney also receives input from the parasympathetic nervous system, by way of the renal branches of the vagus nerve (cranial nerve X), which causes vasodilation and increased blood flow of the afferent arterioles. Due to this mechanism, sympathetic nervous stimulation will decrease urine production, while parasympathetic nervous stimulation will increase urine production.



Blood supply to the kidneys: The renal arteries branch off of the abdominal aorta and supply the kidneys with blood.

This page titled 24.2C: Supply of Blood and Nerves to the Kidneys is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.2D: Nephron, Parts, and Histology

The nephron of the kidney is involved in the regulation of water and soluble substances in blood.

Learning Objectives

- Identify the parts and histology of kidney nephrons

Key Points


- The glomerulus is the site in the nephron where fluid and solutes are filtered out of the blood to form a glomerular filtrate.
- The proximal and distal tubules, the loop of Henle, and the collecting ducts are sites for the reabsorption of water and ions.
- All of the glucose in the blood is reabsorbed by the proximal convoluted tubule through ion cotransport.
- The loop of Henle (sometimes known as the nephron loop) is a U-shaped tube that consists of a descending limb and ascending limb, which differ in permeability.
- The collecting duct and distal convoluted tubule are normally impermeable to water, but this is altered due to hormone stimulus during homeostasis.

Key Terms

- **Bowman's capsule**: A cup-like sac at the beginning of the tubular component of a nephron in the mammalian kidney.
- **proximal tubule**: The first and primary sort of water and ion reabsorption in the kidney, where all glucose in the blood is reabsorbed.
- **glomerulus**: A small, intertwined group of capillaries within the nephrons of the kidney that filter the blood to make urine.

A Nephron

A nephron is the basic structural and functional unit of the kidneys that regulates water and soluble substances in the blood by filtering the blood, reabsorbing what is needed, and excreting the rest as urine. Its function is vital for homeostasis of blood volume, blood pressure, and plasma osmolarity. It is regulated by the neuroendocrine system by hormones such as antidiuretic hormone, aldosterone, and parathyroid hormone.



The basic physiology of a nephron within a kidney: The labels are: 1. Glomerulus, 2. Efferent arteriole, 3. Bowman's capsule, 4. Proximal tube, 5. Cortical collecting tube, 6. Distal tube, 7. Loop of Henle, 8. Collecting duct, 9. Peritubular capillaries, 10. Arcuate vein, 11. Arcuate artery, 12. Afferent arteriole, and 13. Juxtaglomerular apparatus.

The Glomerulus

The glomerulus is a capillary tuft that receives its blood supply from an afferent arteriole of the renal circulation. Here, fluid and solutes are filtered out of the blood and into the space made by Bowman's capsule.

A group of specialized cells known as juxtaglomerular apparatus (JGA) are located around the afferent arteriole where it enters the renal corpuscle. The JGA secretes an enzyme called renin, due to a variety of stimuli, and it is involved in the process of blood volume homeostasis.

The Bowman's capsule (also called the glomerular capsule) surrounds the glomerulus. It is composed of visceral (simple squamous epithelial cells; inner) and parietal (simple squamous epithelial cells; outer) layers. The visceral layer lies just beneath the thickened glomerular basement membrane and only allows fluid and small molecules like glucose and ions like sodium to pass through into the nephron.

Red blood cells and large proteins, such as serum albumins, cannot pass through the glomerulus under normal circumstances. However, in some injuries they may be able to pass through and can cause blood and protein content to enter the urine, which is a sign of problems in the kidney.

Proximal Convoluted Tubule

The proximal tubule is the first site of water reabsorption into the bloodstream, and the site where the majority of water and salt reabsorption takes place. Water reabsorption in the proximal convoluted tubule occurs due to both passive diffusion across the basolateral membrane, and active transport from Na^+/K^+/ATPase pumps that actively transports sodium across the basolateral membrane.

Water and glucose follow sodium through the basolateral membrane via an osmotic gradient, in a process called co-transport. Approximately 2/3rds of water in the nephron and 100% of the glucose in the nephron are reabsorbed by cotransport in the proximal convoluted tubule.

Fluid leaving this tubule generally is unchanged due to the equivalent water and ion reabsorption, with an osmolarity (ion concentration) of 300 mOSm/L, which is the same osmolarity as normal plasma.

The Loop of Henle

The loop of Henle is a U-shaped tube that consists of a descending limb and ascending limb. It transfers fluid from the proximal to the distal tubule. The descending limb is highly permeable to water but completely impermeable to ions, causing a large amount of water to be reabsorbed, which increases fluid osmolarity to about 1200 mOSm/L. In contrast, the ascending limb of Henle's loop is impermeable to water but highly permeable to ions, which causes a large drop in the osmolarity of fluid passing through the loop, from 1200 mOSM/L to 100 mOSm/L.

Distal Convoluted Tubule and Collecting Duct

The distal convoluted tubule and collecting duct is the final site of reabsorption in the nephron. Unlike the other components of the nephron, its permeability to water is variable depending on a hormone stimulus to enable the complex regulation of blood osmolarity, volume, pressure, and pH.

Normally, it is impermeable to water and permeable to ions, driving the osmolarity of fluid even lower. However, anti-diuretic hormone (secreted from the pituitary gland as a part of homeostasis) will act on the distal convoluted tubule to increase the permeability of the tubule to water to increase water reabsorption. This example results in increased blood volume and increased blood pressure. Many other hormones will induce other important changes in the distal convoluted tubule that fulfill the other homeostatic functions of the kidney.

The collecting duct is similar in function to the distal convoluted tubule and generally responds the same way to the same hormone stimuli. It is, however, different in terms of histology. The osmolarity of fluid through the distal tubule and collecting duct is highly variable depending on hormone stimulus. After passage through the collecting duct, the fluid is brought into the ureter, where it leaves the kidney as urine.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Kidneys. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Kidneys%23Location. **License**: *CC BY-SA: Attribution-ShareAlike*
- Kidneys. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Kidneys%23Location. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Kidneys_And_Their_Structure. **License**: *CC BY-SA: Attribution-ShareAlike*
- adrenal gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/adrenal%20gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- retroperitoneal. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/retroperitoneal. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal pelvis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/renal_pelvis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1123. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1123.png. **License**:
- Anatomy and Physiology of Animals/Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy_and_Physiology_of_Animals/Urinary_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- Kidney. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Kidney. **License**: *CC BY-SA: Attribution-ShareAlike*
- Robert Bear and David Rintoul, Urinary System. January 14, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m47445/latest/. **License**: *CC BY: Attribution*
- nephron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nephron. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal medulla. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/renal_medulla. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal cortex. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/renal_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1123. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1123.png. **License**:
- Robert Bear and David Rintoul, Urinary System. January 14, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m47445/latest/. **License**: *CC BY: Attribution*
- Kidney. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Kidney. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Renal_Vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal vein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/renal%20vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal plexus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/renal%20plexus. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal artery. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/renal%20artery. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1123. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1123.png. **License**:
- Robert Bear and David Rintoul, Urinary System. January 14, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m47445/latest/. **License**: *CC BY: Attribution*
- Kidneys. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Kidneys.jpg%23file. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Nephrons. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Nephrons. **License**: *CC BY-SA: Attribution-ShareAlike*

- nephron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nephron. **License**: *CC BY-SA: Attribution-ShareAlike*
- glomerulus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/glomerulus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Bowman's_capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1123. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1123.png. **License**:
- Robert Bear and David Rintoul, Urinary System. January 14, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m47445/latest/. **License**: *CC BY: Attribution*
- Kidneys. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Kidneys.jpg%23file. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nephron illustration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Nephron_illustration.svg. **License**:

This page titled 24.2D: Nephron, Parts, and Histology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3: PHYSIOLOGY OF THE KIDNEYS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

24.3A: Overview of Urine Formation

24.3B: Glomerular Filtration

24.3C: Regulation of Glomerular Filtration Rate

24.3D: Tubular Reabsorption

24.3E: Tubular Secretion

This page titled 24.3: Physiology of the Kidneys is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3A: Overview of Urine Formation

Urine is formed in three steps: filtration, reabsorption, and secretion.

Learning Objectives

- Summarize the steps in urine formation

Key Points

- Filtration involves the transfer of soluble components, such as water and waste, from the blood into the glomerulus.
- Reabsorption involves the absorption of molecules, ions, and water that are necessary for the body to maintain homeostasis from the glomerular filtrate back into the blood.
- Secretion involves the transfer of hydrogen ions, creatinine, drugs, and urea from the blood into the collecting duct, and is primarily made of water.
- Blood and glucose are not normally found in urine.

Key Terms

- **urine**: A liquid excrement consisting of water, salts, and urea, which is made in the kidneys then released through the urethra.
- **glomerulus**: A small, intertwined group of capillaries within nephrons of the kidney that filter the blood to make urine.

Urine is a waste byproduct formed from excess water and metabolic waste molecules during the process of renal system filtration. The primary function of the renal system is to regulate blood volume and plasma osmolarity, and waste removal via urine is essentially a convenient way that the body performs many functions using one process.
Urine formation occurs during three processes:

1. Filtration
2. Reabsorption
3. Secretion

Filtration

During filtration, blood enters the afferent arteriole and flows into the glomerulus where filterable blood components, such as water and nitrogenous waste, will move towards the inside of the glomerulus, and nonfilterable components, such as cells and serum albumins, will exit via the efferent arteriole. These filterable components accumulate in the glomerulus to form the glomerular filtrate.

Normally, about 20% of the total blood pumped by the heart each minute will enter the kidneys to undergo filtration; this is called the filtration fraction. The remaining 80% of the blood flows through the rest of the body to facilitate tissue perfusion and gas exchange.


Reabsorption

The next step is reabsorption, during which molecules and ions will be reabsorbed into the circulatory system. The fluid passes through the components of the nephron (the proximal/distal convoluted tubules, loop of Henle, the collecting duct) as water and ions are removed as the fluid osmolarity (ion concentration) changes. In the collecting duct, secretion will occur before the fluid leaves the ureter in the form of urine.

Secretion

During secretion some substances±such as hydrogen ions, creatinine, and drugs—will be removed from the blood through the peritubular capillary network into the collecting duct. The end product of all these processes is urine, which is essentially a collection of substances that has not been reabsorbed during glomerular filtration or tubular reabsorbtion.

Urine is mainly composed of water that has not been reabsorbed, which is the way in which the body lowers blood volume, by increasing the amount of water that becomes urine instead of becoming reabsorbed. The other main component of urine is urea, a highly soluble molecule composed of ammonia and carbon dioxide, and provides a way for nitrogen (found in ammonia) to be removed from the body. Urine also contains many salts and other waste components. Red blood cells and sugar are not normally found in urine but may indicate glomerulus injury and diabetes mellitus respectively.



Normal kidney physiology: This illustration demonstrates the normal kidney physiology, showing where some types of diuretics act, and what they do.

This page titled 24.3A: Overview of Urine Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3B: Glomerular Filtration

Glomerular filtration is the renal process whereby fluid in the blood is filtered across the capillaries of the glomerulus.

Learning Objectives

- Explain the process of glomerular filtration in the kidneys

Key Points

- The formation of urine begins with the process of filtration. Fluid and small solutes are forced under pressure to flow from the glomerulus into the capsular space of the glomerular capsule.
- The Bowman's capsule is the filtration unit of the glomerulus and has tiny slits in which filtrate may pass through into the nephron. Blood entering the glomerulus has filterable and non-filterable components.
- Filterable blood components include water, nitrogenous waste, and nutrients that will be transferred into the glomerulus to form the glomerular filtrate.
- Non-filterable blood components include blood cells, albumins, and platelets, that will leave the glomerulus through the efferent arteriole.
- Glomerular filtration is caused by the force of the difference between hydrostatic and osmotic pressure (though the glomerular filtration rate includes other variables as well).


Key Terms

- **glomerulus**: A small, intertwined group of capillaries within nephrons of the kidney that filter the blood to make urine.
- **hydrostatic pressure**: The pushing force exerted by the pressure in a blood vessel. It is the primary force that drives glomerular filtration.

Glomerular filtration is the first step in urine formation and constitutes the basic physiologic function of the kidneys. It describes the process of blood filtration in the kidney, in which fluid, ions, glucose, and waste products are removed from the glomerular capillaries.

Many of these materials are reabsorbed by the body as the fluid travels through the various parts of the nephron, but those that are not reabsorbed leave the body in the form of urine.

Glomerulus Structure



Glomerulus structure: A diagram showing the afferent and efferent arterioles bringing blood in and out of the Bowman's capsule, a cup-like sac at the beginning of the tubular component of a nephron.

Blood plasma enters the afferent arteriole and flows into the glomerulus, a cluster of intertwined capillaries. The Bowman's capsule (also called the glomerular capsule) surrounds the glomerulus and is composed of visceral (simple squamous epithelial cells—inner) and parietal (simple squamous epithelial cells—outer) layers.

The visceral layer lies just beneath the thickened glomerular basement membrane and is made of podocytes that form small slits in which the fluid passes through into the nephron. The size of the filtration slits restricts the passage of large molecules (such as albumin) and cells (such as red blood cells and platelets) that are the non-filterable components of blood.

These then leave the glomerulus through the efferent arteriole, which becomes capillaries meant for kidney–oxygen exchange and reabsorption before becoming venous circulation. The positively charged podocytes will impede the filtration of negatively charged particles as well (such as albumins).

The Mechanisms of Filtration

The process by which glomerular filtration occurs is called renal ultrafiltration. The force of hydrostatic pressure in the glomerulus (the force of pressure exerted from the pressure of the blood vessel itself) is the driving force that pushes filtrate out of the capillaries and into the slits in the nephron.

Osmotic pressure (the pulling force exerted by the albumins) works against the greater force of hydrostatic pressure, and the difference between the two determines the effective pressure of the glomerulus that determines the force by which molecules are filtered. These factors will influence the glomeruluar filtration rate, along with a few other factors.

This page titled 24.3B: Glomerular Filtration is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3C: Regulation of Glomerular Filtration Rate

Regulation of GFR requires both a mechanism of detecting an inappropriate GFR as well as an effector mechanism that corrects it.

Learning Objectives

- List the conditions that can affect the glomerular filtration rate (GFR) in kidneys and the manner of its regulation

Key Points

- Glomerular filtration is occurs due to the pressure gradient in the glomerulus.
- Increased blood volume and increased blood pressure will increase GFR.
- Constriction in the afferent arterioles going into the glomerulus and dilation of the efferent arterioles coming out of the glomerulus will decrease GFR.
- Hydrostatic pressure in the Bowman's capsule will work to decrease GFR.
- Normally, the osmotic pressure in the Bowman's space is zero, but it will become present and decrease GFR if the glomerulus becomes leaky.
- Low GFR will activate the renin–angiotensin feedback system that will address the low GFR by increasing blood volume.

Key Terms

- **Bowman's capsule**: A cup-like sac at the beginning of the tubular component of a nephron in the mammalian kidney.
- **osmotic pressure**: The pressure exerted by proteins that attracts water. Water tends to follow proteins based on an osmotic pressure gradient.

Glomerular Filtration Rate

Glomerular filtration rate (GFR) is the measure that describes the total amount of filtrate formed by all the renal corpuscles in both kidneys per minute. The glomerular filtration rate is directly proportional to the pressure gradient in the glomerulus, so changes in pressure will change GFR.

GFR is also an indicator of urine production, increased GFR will increase urine production, and vice versa.

The Starling equation for GFR is:

GFR=Filtration Constant × (Hydrostatic Glomerulus Pressure–Hydrostatic Bowman's Capsule Pressure)–(Osmotic Glomerulus Pressure+Osmotic Bowman's Capsule Pressure)

The filtration constant is based on the surface area of the glomerular capillaries, and the hydrostatic pressure is a pushing force exerted from the flow of a fluid itself; osmotic pressure is the pulling force exerted by proteins. Changes in either the hydrostatic or osmotic pressure in the glomerulus or Bowman's capsule will change GFR.

Hydrostatic Pressure Changes

Many factors can change GFR through changes in hydrostatic pressure, in terms of the flow of blood to the glomerulus. GFR is most sensitive to hydrostatic pressure changes within the glomerulus. A notable body-wide example is blood volume.

Due to Starling's law of the heart, increased blood volume will increase blood pressure throughout the body. The increased blood volume with its higher blood pressure will go into the afferent arteriole and into the glomerulus, resulting in increased GFR. Conversely, those with low blood volume due to dehydration will have a decreased GFR.

Pressure changes within the afferent and efferent arterioles that go into and out of the glomerulus itself will also impact GFR. Vasodilation in the afferent arteriole and vasconstriction in the efferent arteriole will increase blood flow (and hydrostatic pressure) in the glomerulus and will increase GFR. Conversely, vasoconstriction in the afferent arteriole and vasodilation in the efferent arteriole will decrease GFR.


The Bowman's capsule space exerts hydrostatic pressure of its own that pushes against the glomerulus. Increased Bowman's capsule hydrostatic pressure will decrease GFR, while decreased Bowman's capsule hydrostatic pressure will increase GFR.

An example of this is a ureter obstruction to the flow of urine that gradually causes a fluid buildup within the nephrons. An obstruction will increase the Bowman's capsule hydrostatic pressure and will consequently decrease GFR.

Osmotic Pressure Changes

Osmotic pressure is the force exerted by proteins and works against filtration because the proteins draw water in. Increased osmotic pressure in the glomerulus is due to increased serum albumin in the bloodstream and decreases GFR, and vice versa.

Under normal conditions, albumins cannot be filtered into the Bowman's capsule, so the osmotic pressure in the Bowman's space is generally not present, and is removed from the GFR equation. In certain kidney diseases, the basement membrane may be damaged (becoming leaky to proteins), which results in decreased GFR due to the increased Bowman's capsule osmotic pressure.



Glomeruluar filtration: The glomerulus (red) filters fluid into the Bowman's capsule (blue) that sends fluid through the nephron (yellow). GFR is the rate at which is this filtration occurs.

GFR Feedback

GFR is one of the many ways in which homeostasis of blood volume and blood pressure may occur. In particular, low GFR is one of the variables that will activate the renin–angiotensin feedback system, a complex process that will increase blood volume, blood pressure, and GFR. This system is also activated by low blood pressure itself, and sympathetic nervous stimulation, in addition to low GFR.

This page titled 24.3C: Regulation of Glomerular Filtration Rate is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3D: Tubular Reabsorption

Tubular reabsorption is the process by which solutes and water are removed from the tubular fluid and transported into the blood.

Learning Objectives

- Describe the process of tubular reabsorption in kidney physiology

Key Points

- Proper function of the kidney requires that it receives and adequately filters blood.
- Reabsorption includes passive diffusion, active transport, and cotransport.
- Water is mostly reabsorbed by the cotransport of glucose and sodium.
- Filtrate osmolarity changes drastically throughout the nephron as varying amounts of the components of filtrate are reabsorbed in the different parts of the nephron.
- The normal osmolarity of plasma is 300 mOsm/L, which is the same osmolarity within the proximal convoluted tubule.


Key Terms

- **NA+/K+ ATPase**: An ATPase pump that consumes ATP to facilitate the active transport of ions in filtrate of the nephron.
- **peri-tubular capillaries**: The capillaries through which components of filtrate are reabsorbed from the lumen of the nephron.

Filtrate

The fluid filtered from blood, called filtrate, passes through the nephron, much of the filtrate and its contents are reabsorbed into the body. Reabsorption is a finely tuned process that is altered to maintain homeostasis of blood volume, blood pressure, plasma osmolarity, and blood pH. Reabsorbed fluids, ions, and molecules are returned to the bloodstream through the peri-tubular capillaries, and are not excreted as urine.

Mechanisms of Reabsorption



Tubular secretion: Diagram showing the basic physiologic mechanisms of the kidney and the three steps involved in urine formation. Namely filtration, reabsorption, secretion, and excretion.

Reabsorption in the nephron may be either a passive or active process, and the specific permeability of the each part of the nephron varies considerably in terms of the amount and type of substance reabsorbed. The mechanisms of reabsorption into the peri-tubular capillaries include:

- Passive diffusion—passing through plasma membranes of the kidney epithelial cells by concentration gradients.
- Active transport—membrane-bound ATPase pumps (such as NA^+/K^+ ATPase pumps) with carrier proteins that carry substances across the plasma membranes of the kidney epithelial cells by consuming ATP.
- Cotransport—this process is particularly important for the reabsorption of water. Water can follow other molecules that are actively transported, particularly glucose and sodium ions in the nephron.


These processes involve the substance passing though the luminal barrier and the basolateral membrane, two plasma membranes of the kidney epithelial cells, and into the peri-tubular capillaries on the other side. Some substances can also pass through tiny spaces in between the renal epithelial cells, called tight junctions.

Osmolarity Changes

As filtrate passes through the nephron, its osmolarity (ion concentration) changes as ions and water are reabsorbed. The filtrate entering the proximal convoluted tubule is 300 mOsm/L, which is the same osmolarity as normal plasma osmolarity.

In the proximal convoluted tubules, all the glucose in the filtrate is reabsorbed, along with an equal concentration of ions and water (through cotransport), so that the filtrate is still 300 mOsm/L as it leaves the tubule. The filtrate osmolarity drops to 1200 mOsm/L as water leaves through the descending loop of Henle, which is impermeable to ions. In the ascending loop of Henle, which is permeable to ions but not water, osmolarity falls to 100–200 mOsm/L.

Finally, in the distal convoluted tubule and collecting duct, a variable amount of ions and water are reabsorbed depending on hormonal stimulus. The final osmolarity of urine is therefore dependent on whether or not the final collecting tubules and ducts are permeable to water or not, which is regulated by homeostasis.



Reabsorption throughout the nephron: A diagram of the nephron that shows the mechanisms of reabsorption.

This page titled 24.3D: Tubular Reabsorption is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.3E: Tubular Secretion

Hydrogen, creatinine, and drugs are removed from the blood and into the collecting duct through the peritubular capillary network.

Learning Objectives

- Describe the purpose of tubular secretion in kidney physiology

Key Points

- The substance that remains in the collecting duct of the kidneys following reabsorption is better known as urine.
- Secreted substances largely include hydrogen, creatinine, ions, and other types of waste products, such as drugs. Tubular secretion is the transfer of materials from peritubular capillaries to the renal tubular lumen and occurs mainly by active transport and passive diffusion.
- It is the tubular secretion of H+ and NH_4+ from the blood into the tubular fluid that helps to keep blood pH at its normal level—this is also a respiratory process.
- Urine leaves the kidney though the ureter following secretion.

Key Terms

- **collecting duct**: A system of the kidneys that consists of a series of tubules and ducts that connect the nephrons to the ureter.
- **peritubular capillaries**: Tiny blood vessels that travel alongside nephrons, allowing reabsorption and secretion between blood and the inner lumen of the nephron.
- **lumen**: The inside space of a tubular structure, such as an artery or intestine.

Tubular secretion is the transfer of materials from peritubular capillaries to the renal tubular lumen; it is the opposite process of reabsorption. This secretion is caused mainly by active transport and passive diffusion.

Usually only a few substances are secreted, and are typically waste products. Urine is the substance leftover in the collecting duct following reabsorption and secretion.


Mechanisms of Secretion

The mechanisms by which secretion occurs are similar to those of reabsorption, however these processes occur in the opposite direction.

- Passive diffusion—the movement of molecules from the peritubular capillaries to the intersitial fluid within the nephron.
- Active transport—the movement of molecules via ATPase pumps that transport the substance through the renal epithelial cell into the lumen of the nephron.

Renal secretion is different from reabsorption because it deals with filtering and cleaning substances from the blood, rather than retaining them. The substances that are secreted into the tubular fluid for removal from the body include:

- Potassium ions (K+)
- Hydrogen ions (H+)
- Ammonium ions (NH_4+)
- Creatinine
- Urea
- Some hormones
- Some drugs (e.g., penicillin)



Tubular secretion: Diagram showing the basic physiologic mechanisms of the kidney and the three steps involved in urine formation.

Many pharmaceutical drugs are protein-bound molecules thatDiagram showing the basic physiologic mechanisms of the kidney and the three steps involved in urine formation. amely filtration, reabsorption, secretion, and excretion. are easily secreted, which is why urine testing can detect the exposure to many types of drugs. Tubular secretion occurs throughout the different parts of the nephron, from the proximal convoluted tubule to the collecting duct at the end of the nephron.

Hydrogen Ion Secretion

The tubular secretion of H+ and NH_4+ from the blood into the tubular fluid is involved in blood pH regulation. The movement of these ions also helps to conserve sodium bicarbonate ($NaHCO_3$). The typical pH of urine is about 6.0, while it is ideally 7.35 to 7.45 for blood.

pH regulation is primarily a respiratory system process, due to the exchange of carbon dioxide (a component of carbonic acid in blood), however tubular secretion assists in pH homeostasis as well.

Following Secretion

Urine that is formed via the three processes of filtration, reabsorption, and secretion leaves the kidney through the ureter, and is stored in the bladder before being removed through the urethra. At this final stage it is only approximately one percent of the originally filtered volume, consisting mostly of water with highly diluted amounts of urea, creatinine, and variable concentrations of ions.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Filtration**. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Filtration**. **License**: *CC BY-SA: Attribution-ShareAlike*

- urine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- glomerular filtrate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/glomerular%20filtrate. **License**: *CC BY-SA: Attribution-ShareAlike*
- glomerulus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/glomerulus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal Diuretics. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._Diuretics.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...m%23Glomerulus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...an.27s_Capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- glomerulus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/glomerulus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Bowman's_capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal Diuretics. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Renal_Diuretics.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bowman's_capsule. **License**: *Public Domain: No Known Copyright*
- Bowman's capsule. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Bowman's_capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...an.27s_Capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...an.27s_Capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal function. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renal_function. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...an.27s_Capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- macula densa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/macula%20densa. **License**: *CC BY-SA: Attribution-ShareAlike*
- clearance rate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/clearance%20rate. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal Diuretics. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Renal_Diuretics.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bowman's_capsule. **License**: *Public Domain: No Known Copyright*
- Renal_corpuscle.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glomerulus_(kidney)#/media/File:Renal_corpuscle.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Bowman's_capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...23Reabsorption. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal physiology. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renal_physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- urine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- tubular fluid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/tubular%20fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal Diuretics. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Renal_Diuretics.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bowman's capsule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bowman's_capsule. **License**: *Public Domain: No Known Copyright*
- Renal_corpuscle.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glomerulus_(kidney)#/media/File:Renal_corpuscle.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Urine formation. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/2/2b/Physiology_of_Nephron.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Kidney_nephron_molar_transport_diagram.svg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Collecting_duct_system#/media/File:Kidney_nephron_molar_transport_diagram.svg. **License**: *CC BY-SA: Attribution-*

ShareAlike

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Secretion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lumen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lumen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peritubular capillaries. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/peritubular%20capillaries**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- collecting duct. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/collecting%20duct**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Renal Diuretics. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Renal_Diuretics.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bowman's capsule. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bowman's_capsule**. **License**: ***Public Domain: No Known Copyright***
- Renal_corpuscle.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Glomerulus_(kidney)#/media/File:Renal_corpuscle.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Urine formation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/2b/Physiology_of_Nephron.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Kidney_nephron_molar_transport_diagram.svg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Collecting_duct_system#/media/File:Kidney_nephron_molar_transport_diagram.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Urine formation. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/2b/Physiology_of_Nephron.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 24.3E: Tubular Secretion is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4: URINE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

24.4A: Physical Characteristics of Urine

24.4B: Chemical Composition of Urine

24.4C: Regulation of Urine Concentration and Volume

24.4D: Urinalysis

24.4E: Renal Clearance

This page titled 24.4: Urine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4A: Physical Characteristics of Urine

Urine is a sterile waste product composed of water soluble nitrogen products.

Learning Objectives

- List the physical characteristics of urine

Key Points

- Urine color is an indicator for hydration.
- Urine pH is often influenced by diet.
- Urine smell indicates age of the urine and may indicate the prescence of glucose and ketones.
- Urine turbidity may indicate urinary tract infection or obstruction.
- Urinalysis is the process of analyzing and detecting chemicals excreted in urine.
- Physical characteristics of urine include color, smell, pH, density and turbidity

Key Terms

- **urine**: A liquid excrement consisting of water, salts, and urea, which is made in the kidneys then released through the urethra.
- **urinalysis**: A urinalysis (UA), also known as Routine and Microscopy (R&M), is an array of tests performed on urine, and one of the most common methods of medical diagnosis.

Urine, a typically sterile liquid by-product of the body, is secreted by the kidneys through a process called urination and excreted through the urethra. Urine is often used as a diagnostic feature for many disease conditions. These may b based on either physical or chemical components, that may give insight to processes within the body, often through urinalysis, a common clinical analysis of urine.

Physical Characteristics

Physical characteristics that can be applied to urine include color, turbidity (transparency), smell (odor), pH (acidity – alkalinity) and density. Many of these characteristics are notable and identifiable by by vision alone, but some require laboratory testing.

- Color: Typically yellow-amber, but varies according to recent diet and the concentration of the urine. Drinking more water generally tends to reduce the concentration of urine, and therefore causes it to have a lighter color. Dark urine may indicate dehydration. Red urine indicates red blood cells within the urine, a sign of kidney damage and disease.
- Smell: The smell of urine may provide health information. For example, urine of diabetics may have a sweet or fruity odor due to the presence of ketones (organic molecules of a particular structure) or glucose. Generally fresh urine has a mild smell but aged urine has a stronger odor similar to that of ammonia.
- The pH of normal urine is generally in the range 4.6 – 8, with a typical average being around 6.0. Much of the variation occurs due to diet. For example, high protein diets result in more acidic urine, but vegetarian diets generally result in more alkaline urine (both within the typical range of 4.6 – 8).
- Density: Density is also known as “specific gravity.” This is the ratio of the weight of a volume of a substance compared with the weight of the same volume of distilled water. The density of normal urine ranges from 0.001 to 0.035.
- Turbidity: The turbidity of the urine sample is gauged subjectively and reported as clear, slightly cloudy, cloudy, opaque or flocculent. Normally, fresh urine is either clear or very slightly cloudy. Excess turbidity results from the presence of suspended particles in the urine, the cause of which can usually be determined by the results of the microscopic urine sediment examination. Common causes of abnormal turbidity include: increased cells, urinary tract infections or obstructions.

Abnormalities in any of these of physical characteristics may indicate disease or metabolic imbalances. These problems may seem superficial or minor on their own, but can actually be the symptoms for more serious diseases, such as diabetes mellitus, or a damaged glomerulus.

This page titled 24.4A: Physical Characteristics of Urine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4B: Chemical Composition of Urine

Normal urine consists of water, urea, salts, and pigments.

Learning Objectives

- Describe how normal urine consists of water, urea, salts and pigment

Key Points

- Urine is a liquid by-product of the body secreted by the kidneys through a process called urination and excreted through the urethra.
- Urine is an aqueous solution of greater than 95% water. Other constituents include urea, chloride, sodium, potassium, creatinine and other dissolved ions, and inorganic and organic compounds.
- Urea is a non-toxic molecule made of toxic ammonia and carbon dioxide. Any abnormal constituents found in urine are an indication of disease.
- The presence of red blood cells in urine is referred to as haematuria.
- The presence of proteins, which are normally too large to pass through the tubules, can be an indication of damage to the tubules, and is called proteinuria.

Key Terms

- **urine**: A liquid excrement that consists of water, salts, and urea, and is made in the kidneys then released through the urethra.
- **diabetes**: A group of endocrine diseases whereby a person has high blood sugar due to an inability to produce, metabolize, or respond to the hormone insulin.

Urine is a liquid byproduct of the body secreted by the kidneys through a process called urination and excreted through the urethra. The normal chemical composition of urine is mainly water content, but it also includes nitrogenous molecules, such as urea, as well as creatinine and other metabolic waste components.

Other substances may be excreted in urine due to injury or infection of the glomeruli of the kidneys, which can alter the ability of the nephron to reabsorb or filter the different components of blood plasma.

Normal Chemical Composition of Urine

Urine is an aqueous solution of greater than 95% water, with a minimum of these remaining constituents, in order of decreasing concentration:

- Urea 9.3 g/L.
- Chloride 1.87 g/L.
- Sodium 1.17 g/L.
- Potassium 0.750 g/L.
- Creatinine 0.670 g/L.
- Other dissolved ions, inorganic and organic compounds (proteins, hormones, metabolites).

Urine is sterile until it reaches the urethra, where epithelial cells lining the urethra are colonized by facultatively anaerobic gram-negative rods and cocci. Urea is essentially a processed form of ammonia that is non-toxic to mammals, unlike ammonia, which can be highly toxic. It is processed from ammonia and carbon dioxide in the liver.

Abnormal Types of Urine

There are several conditions that can cause abnormal components to be excreted in urine or present as abnormal characteristics of urine. They are mostly referred to by the suffix -uria. Some of the more common types of abnormal urine include:

- Proteinuria—Protein content in urine, often due to leaky or damaged glomeruli.
- Oliguria—An abnormally small amount of urine, often due to shock or kidney damage.
- Polyuria—An abnormally large amount of urine, often caused by diabetes.
- Dysuria—Painful or uncomfortable urination, often from urinary tract infections.
- Hematuria—Red blood cells in urine, from infection or injury.
- Glycosuria— Glucose in urine, due to excess plasma glucose in diabetes, beyond the amount able to be reabsorbed in the proximal convoluted tubule.

This page titled 24.4B: Chemical Composition of Urine is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4C: Regulation of Urine Concentration and Volume

Antidiuretic hormone (ADH) is produced by the pituitary gland to control the amount of water that is reabsorbed through the collecting ducts.

Learning Objectives

- Describe how regulating the amount of water excreted in urine is an essential component of homeostasis that is regulated by the antidiuretic hormone (ADH)

Key Points

- Urine volume and concentration is regulated through the same processes that regulate blood volume.
- Antidiuretic hormone (ADH)—produced by the posterior pituitary gland —increases the amount of water reabsorbed in the distal convoluted tubule and collecting duct.
- Osmoreceptors in the hypothalamus signal the posterior pituitary gland to increase ADH secretion when plasma osmolarity becomes too high.
- ADH causes decreased urine volume and decreased plasma osmolarity.
- A diuretic increases urine volume and increases plasma osmolarity.
- Common diuretics include alcohol, water, caffeine, and many medications, and they generally function as diuretics via different mechanisms.

Key Terms

- **diuretic**: A substance that increases urine volume and increases plasma osmolarity, often by inhibiting ADH secretion to prevent water reabsorption in the nephron.
- **antidiuretic hormone**: A hormone secreted by the posterior pituitary gland that increases water retention to decrease urine volume and decrease plasma osmolarity.

Urine is produced not only to eliminate many cellular waste products, but also to control the amount of water in the body. In a way, urine volume regulation is part of homeostasis, in that it directly regulates blood volume, because greater amounts of urine will reduce the volume of waters in blood.

There are a few complex systems involved in regulating blood volume and urine production, such as the intricate renin–angiotensin system, and the simpler anti-diuretic hormone (ADH) feedback system.

Anti-Diuretic Hormone Feedback

An anti-diruetic is a substance that decreases urine volume, and ADH is the primary example of it within the body. ADH is a hormone secreted from the posterior pituitary gland in response to increased plasma osmolarity (i.e., increased ion concentration in the blood), which is generally due to an increased concentration of ions relative to the volume of plasma, or decreased plasma volume.

The increased plasma osmolarity is sensed by osmoreceptors in the hypothalamus, which will stimulate the posterior pituitary gland to release ADH. ADH will then act on the nephrons of the kidneys to cause a decrease in plasma osmolarity and an increase in urine osmolarity.

ADH increases the permeability to water of the distal convoluted tubule and collecting duct, which are normally impermeable to water. This effect causes increased water reabsorption and retention and decreases the volume of urine produced relative to its ion content.


After ADH acts on the nephron to decrease plasma osmolarity (and leads to increased blood volume) and increase urine

osmolarity, the osmoreceptors in the hypothalamus will inactivate, and ADH secretion will end. Due to this response, ADH secretion is considered to be a form of negative feedback.

Diuretics

A diuretic is any substance that has the opposite effect of ADH— they increase urine volume, decrease urine osmolarity, lead to an increased plasma osmolarity, and often reduced blood volume. Many substances can act as diuretics, albeit with different mechanisms.

A common example is alcohol and water ingestion, which directly inhibit ADH secretion in the pituitary gland. Alternatively caffeine is a diuretic because it interferes with sodium reabsorption (reducing the amount of water reabsorbed by sodium cotransport) and increases the glomerular filtration rate by temporarily increasing blood pressure. Many medications are diuretics because they inhibit the ATPase pumps, thus slowing water reabsorption further.



Summary of the process of urine formation: As the fluid flows along the proximal convoluted tubule useful substances like glucose, water, salts, potassium ions, calcium ions, and amino acids are reabsorbed into the blood capillaries that form a network around the tubules. Many of these substances are transported by active transport and energy is required.

This page titled 24.4C: Regulation of Urine Concentration and Volume is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4D: Urinalysis

Urinalysis is the process of analyzing urine for target parameters of health and disease.

Learning Objectives

- Describe how urinalysis can be used as a method of diagnosis in medicine

Key Points

- The characteristics that can be detected in urine include cells, substances, and properties, such as specific gravity or pH.
- Urinalysis can be performed on test strips (routine) by light microscopy.
- The numbers and types of cells and/or material, such as urinary casts, can yield a great detail of information and may suggest a specific diagnosis.
- Urinary casts include hyaline casts, granular casts, white blood cell casts, red blood cell casts, epithelial cell casts, or bacterial cell casts, which all indicate different abnormalities within urine.

Key Terms

- **urinary cast**: Tiny structures formed from bound abnormal cells and molecules within the nephrons that are excreted in urine.
- **urinalysis**: Also known as routine and microscopy (R&M), this is an array of tests performed on urine, and one of the most common methods of medical diagnosis.

Urinalysis

A urinalysis (UA), also known as routine and microscopy (R&M), is an array of tests performed on urine, and one of the most common methods of medical diagnosis. Urinalysis means the analysis of urine, and it is used to diagnose several diseases.

The target parameters that are measured or quantified in urinalysis include many substances and cells, as well as other properties, such as specific gravity. A part of a urinalysis can be performed by using urine test strips, in which the test results can be read as the strip's color changes. Another method is light microscopy of urine samples.

When doctors order a urinalysis, they will request either a routine urinalysis or a routine and microscopy (R&M) urinalysis; the difference being that a routine urinalysis does not include microscopy or culture. R&M is used specifically for culturing bacteria found in urine, which can make it an important tool for diagnosing specific urinary tract infections.

Test Strip Urinalysis

Test strip urinalysis exposes urine to strips that react if the urine contains certain cells or molecules. Test strip urinalysis is the most common technique used in routine urinalysis. A urine test strip can identify:


- Leukocytes—their presence in urine is known as leukocyturia.
- Nitrites—their presence in urine is known as nitrituria.
- Proteins —their presence in urine is known as proteinuria, albuminuria, or microalbuminuria.
- Blood—its presence in urine is known as hematuria.
- pH—the acidity of urine is easily quantified by test strips, which can identify cases of metabolic acidosis or alkalosis.

Urine Microscopy

The numbers and types of cells and/or material, such as urinary casts, can yield a great detail of information and may suggest a specific diagnosis. A urinary cast is any tiny structure found in urine that consists of multiple molecules or cells bound together.

Casts form within the nephron when abnormal cells and molecules are filtered from blood, and are excreted as the bound structures in urine. Microscopy can identify casts in urine and use them to diagnose kidney diseases, by characterizing symptoms such as:

- Red blood cell casts are associated with glomerulonephritis, vasculitis, or malignant hypertension.
- White blood cell casts are associated with acute interstitial nephritis, exudative glomerulonephritis, or severe pyelonephritis.
- Epithelial cell casts are associated with toxin-induced, acute tubular necrosis, hepatitis, and cytomegalovirus.
- (Heme) granular casts are associated with acute tubular necrosis, and are often composed of proteins, especially antibodies.
- Hyaline casts are associated with dehydration; it is the most common type of cast.
- Bacterial casts are associated with urinary tract infection; the cast may be cultured in order to identify the causative organism of the cast.



Urinalysis: White blood cells seen under a microscope from a urine sample.

This page titled 24.4D: Urinalysis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.4E: Renal Clearance

Clearance is a measurement of the renal excretion ability.

Learning Objectives

- Describe how clearance is a measure of the renal excretion ability

Key Points

- Each substance has a specific clearance that depends on its filtration characteristics, size, and molecular structure.
- Clearance is dependent upon glomerular filtration, secretion, and reabsorption.
- Clearance may be either constant or variable over time.
- Many drugs can either be bound to plasma proteins or unbound in plasma; however, they will not be cleared from the body when bound to proteins.
- Renal clearance is the main form of clearance in the body, and when combined with other routes of clearance in the body it accounts for most of the total body clearance.


Key Terms

- **nephron**: The basic structural and functional unit of the kidney that filters the blood in order to regulate chemical concentrations and thereby produce urine.
- **Renal clearance**: The rate at which a substance is removed from the plasma via renal system activity over a unit of time.

Clearance

In renal physiology, clearance is a measurement of the renal excretion ability, which measures the amount of plasma from which a substance is removed from the body over an interval of time. Each substance has its own specific clearance that depends on its unique filtration characteristics.

Clearance is a function of glomerular filtration, secretion from the peritubular capillaries to the nephron, and reabsorption from the nephron back to the peritubular capillaries. Clearance can be either a constant or variable component over time, depending on the type of substance.



Physiology of the nephron: Diagram showing the basic physiologic mechanisms of the kidney.

Clearance Mechanisms

Renal clearance depends mainly on GFR, tubular absorption, and tubular secretion. If any of those variables change, the renal clearance rate of a substance will change as well. These variables alter clearance through the following rules:

- Increased GFR will increase clearance, while decreased GFR will decrease clearance.
- Increased tubular secretion will increase clearance, while decreased tubular secretion will decrease clearance. This variable is sometimes altered through changes in the expression of ATPase pumps involved in active transport.
- Increased tubular reabsorption will decrease clearance, while increased tubular reabsorption will increase clearance.

Additionally, the characteristics of the substance of interest will also determine some components of clearance. For example, certain pharmaceuticals have the tendency to bind to plasma proteins or exist unbound in plasma. Only those that are unbound will be filtered and cleared from the body. Size and molecular structure will also alter the clearance rate.

It is also important to note that renal clearance is not the only form of clearance that occurs for the substances within the plasma of the body. The other types of clearance are

- Biliary (through bile).
- Salivary.
- Pulmonary clearance (removed during alveolar gas exchange).

These types of clearance may also excrete certain molecules from the bloodstream based on their size and molecular structure; however, these forms of clearance are generally relatively minor compared to renal clearance.

These types of clearance all add up to a summation known as total body clearance, which refers to the removal of a substance from the plasma over time, incorporating all routes of removal in the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Urine. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urine%23Characteristics. **License**: *CC BY-SA: Attribution-ShareAlike*
- micturition. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/micturition. **License**: *CC BY-SA: Attribution-ShareAlike*
- urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- urine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: cms.boundlesslearning.com/mod...b02abfc1e84c12. **License**: *CC BY-SA: Attribution-ShareAlike*
- diabetes. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/diabetes. **License**: *CC BY-SA: Attribution-ShareAlike*
- urine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- red blood cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/red_blood_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- antidiuretic hormone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/antidiuretic_hormone. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and Physiology of Animals/Urinary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Anatomy_and_Physiology_of_Animals/Urinary_System%23The_Production_Of_Concentrated_Urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- capsule. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/capsule. **License**: *CC BY-SA: Attribution-ShareAlike*
- glomerulus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/glomerulus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals Summary of the processes involved in the formation of urine. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Summary_of_the_processes_involved_in_the_formation_of_urine.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- urine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urine. **License**: *CC BY-SA: Attribution-ShareAlike*
- urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals Summary of the processes involved in the formation of urine. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Summary_of_the_processes_involved_in_the_formation_of_urine.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal clearance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renal_clearance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal clearance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renal_clearance. **License**: *CC BY-SA: Attribution-ShareAlike*
- nephron. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nephron. **License**: *CC BY-SA: Attribution-ShareAlike*
- kidney. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/kidney. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anatomy and physiology of animals Summary of the processes involved in the formation of urine. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Summary_of_the_processes_involved_in_the_formation_of_urine.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Urinalysis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urinalysis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Physiology of Nephron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Physiology_of_Nephron.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 24.4E: Renal Clearance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5: URINE TRANSPORT, STORAGE, AND ELIMINATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

24.5A: Overview of Urine Transport, Storage, and Elimination

24.5B: Ureters

24.5C: Urinary Bladder

24.5D: Urethra

24.5E: Micturition and the Micturition Reflex

24.5F: Waste Management in Other Body Systems

This page titled 24.5: Urine Transport, Storage, and Elimination is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5A: Overview of Urine Transport, Storage, and Elimination

The urinary organs include the kidneys, ureters, bladder, and urethra.

Learning Objectives

- Outline the process of urine transport, storage, and elimination


Key Points

- Urine collects from the nephrons and flows into the ureters.
- The ureters use smooth muscle contractions to facilitate the flow of urine.
- The urinary bladder is a hollow, muscular, and elastic organ that stores urine.
- Urine exits the bladder and the body through the urethra.
- The kidneys, ureters, bladder, and urethra make up the urinary tract, the pathway through which urine flows and is eliminated from the body.

Key Terms

- **ureter**: These are two long, narrow ducts that carry urine from the kidneys to the urinary bladder.
- **urinary bladder**: An elastic, muscular sac situated in the pelvic cavity, into which urine from the kidneys is stored prior to disposal by urination. Urine enters the bladder via the ureters and exits via the urethra.

The Urinary System



Urinary tract: The transport and removal of urine from the body follows the urinary tract.

The organs, tubes, muscles, and nerves that work together to create, store, and carry urine are referred to as the urinary system, which is another name for the renal system. The renal system filters the plasma of blood and regulates blood volume by excreting excess water in the form of urine. Urine transport follows a path through the kidneys, ureters, bladder, and urethra, which are collectively known as the urinary tract.

Urine Transport

Urine is essentially water, ions, and secreted molecules that leave the collecting duct of the many nephrons of the kidney and flow into the ureters. The ureters are two tubes that drain urine from the kidneys to the bladder.

Each ureter is a muscular tube that drains into the bladder. Smooth muscle contractions in the walls of the ureters, over time, send the urine in small spurts into the bladder, the organ where urine is stored before it can be eliminated.

Urine Storage

The bladder is a hollow muscular organ shaped like a balloon. It sits in the pelvis and is held in place by ligaments attached to other organs and the pelvic bones. The bladder stores urine until enough of it accumulates for removal from the body. It swells into a round shape when it is full and gets smaller when empty.

If the urinary system is healthy, the bladder can hold up to 16 ounces (2 cups) of urine comfortably for 2 to 5 hours. Circular muscles called sphincters help keep urine from leaking. The sphincter muscles close tightly, like a rubber band, around the opening of the bladder into the urethra, the tube that allows urine to pass outside the body.

Urine Elimination

Nerves in the bladder are stimulated as the bladder fills with urine and becomes larger, which in turn stimulates the need to urinate. When you urinate, the brain signals the bladder muscles to tighten, squeezing urine out of the bladder.

At the same time, the brain signals the sphincter muscles to relax. As these muscles relax, urine exits the bladder through the urethra, and leaves the body through an opening in the genital region that contains the urethra. When all the signals occur in the correct order, normal urination occurs, removing urine from the body.

This page titled 24.5A: Overview of Urine Transport, Storage, and Elimination is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5B: Ureters

The ureters are two tubes that drain urine from each of the kidneys into the bladder.

Learning Objectives

- Describe the role of ureters in the urinary system and their structure

Key Points

- The ureter contains transitional epithelium and an additional smooth muscle layer in the more distal one-third to assist with peristalsis.
- The ureters are usually 25–30 cm (10–12 in) long and 3–4 mm in diameter.
- Muscles in the walls of the ureters send the urine in small spurts into the bladder.
- Kidney stones are a common disease of the ureter, in which a stone made of minerals obstructs the ureter.
- The ureter has a few points of constriction where kidney stones are more commonly found.


Key Terms

- **transitional epithelial tissues**: Epithelial tissues that can change between columnar and squamous forms that are found in the ureters and the cervix.
- **peristalsis**: The rhythmic, wave-like contraction of smooth muscle tissue that can propel substances through the many muscular tubes of the body.

The ureters are tubes made of smooth muscle fibers that propel urine from the kidneys to the urinary bladder. In the adult, the ureters are usually 25–30 cm (10–12 in) long and 3–4 mm in diameter. The ureter is one of the essential organs of urinary tract that controls urine transport.

Ureter Structure and Function

The ureters are two tubes that are made out of smooth muscle and transitional epithelial tissues, which are a type of epithelial tissue that may either be columnar or squamous. Each kidney has its own ureter through which urine drains into.



Human urinary system: 1) Human urinary system. 2) Kidney. 3) Renal pelvis. 4) Ureter. 5) Urinary bladder. 6) Urethra (left side with frontal section). 7) Adrenal gland vessels. 8) Renal artery and vein. 9) Inferior vena cava. 10) Abdominal aorta. 11) Common iliac artery and vein with transparency. 12) Liver. 13) Large intestine. 14) Pelvis.

The ureters are long tubes that have a few points of constriction, where obstructions are more common. The ureters receive a blood supply from a few different major arteries including the renal and illiac artery derivatives, and have a relatively dense nerve supply as well.

Muscles in the walls of the ureters send the urine in small spurts into the bladder, in a process called peristalsis. After the urine enters the bladder from the ureters, small folds in the bladder mucosa act like valves to prevent the backward flow of the urine; these are called the ureteral valves. The ureteral valves function similarly to the semilunar valves in the veins of the body, but are structurally different, consisting of transverse mucosal epithelial folds.

Ureter Pathology

Kidney stones and cancer are common diseases of the ureter. A kidney stone can move from the kidney and become lodged inside the ureter, which can block the flow of urine, as well as cause a sharp cramp in the back, side, or lower abdomen. The affected kidney could then develop hydronephrosis, should a part of the kidney become swollen due to blocked flow of urine.

Kidney stones are very common and are usually clumps of aggregated minerals that are most often found at the constriction points in the ureter. Ureter cancer is often due to a malignant transformation of of the transitional epithelial tissue, which is more vulnerable to developing cancer cells compared to other tissues.

This page titled 24.5B: Ureters is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5C: Urinary Bladder

The urinary bladder is a hollow, muscular, and distendible or elastic organ that sits on the pelvic floor.

Learning Objectives

- Summarize the features and function of the urinary bladder

Key Points

- The bladder is made of transitional epithelium and has a trigone.
- A normal bladder empties completely upon a complete discharge, otherwise it is a sign that its elasticity is compromised.
- When the bladder becomes half full, stretch receptors send nerve impulses to the spinal cord, which then sends a reflex nerve impulse back to the sphincter (muscular valve) at the neck of the bladder, causing it to relax and allow the flow of urine into the urethra.
- The internal urethral sphincter is involuntary and the external sphincter is voluntary.
- The bladder plays a minor role in temperature regulation.
- The bladder has both mechanical and sensory innervation through the sympathetic and parasympathetic nervous systems.

Key Terms

- **trigone**: A triangle-shaped part of an organ, such as the bottom of the bladder.
- **urine**: A liquid excrement consisting of water, salts, and urea that is made in the kidneys then released through the urethra.
- **reflex**: An automatic response to a simple stimulus that does not require mental processing.

The Urinary Bladder


The urinary bladder is a urine storage organ that is a part of the urinary tract. The bladder is a hollow, muscular, and elastic organ that sits on the pelvic floor. The bladder expands and fills with urine before it is discharged into the urethra during urination.

Bladder Anatomy

The bladder is a hollow, sac-like organ made of transitional epithelium, similar to the ureter that feeds into it. The ureters enter the bladder diagonally from its dorsolateral floor in
an area called the trigone, which is a triangle-shaped anatomical region. The urethra exits at the
lowest point of the triangle of the trigone.

There are two sphincters, or muscular valves, that separate the bladder from the urethra. The sphincters must open before the urine can flow into the urethra. The internal sphincter is under involuntary control and the external sphincter is under voluntary control.

Bladder Physiology



The urinary bladder: The urinary bladder is composed of several layers of tissue that facilitate urine storage and expulsion. The associated structures of the urinary and male reproductive tract are labelled.

When the bladder fills with urine stretch receptors send nerve impulses to the spinal cord, which then sends a reflex nerve impulse back to the internal sphincter valve at the neck of the bladder that causes it to relax and allow the flow of urine into the urethra. The internal urethral sphincter is involuntary and controlled by the autonomic nerves.

The bladder has a minor temperature regulation function since some heat may leave the body in the form of urine. A normal bladder empties completely upon a complete discharge, otherwise it is a sign that its elasticity is compromised; when it becomes completely void of fluid, it may cause a chilling sensation due to the rapid change of body temperature.

The urinary bladder usually holds 300–350 ml of urine. As urine accumulates, the walls of the bladder thin as it stretches, allowing the bladder to store larger amounts of urine without a significant rise in internal pressure of the bladder.

The bladder receives motor innervation from both sympathetic fibers, most of which arise from the hypogastric plexuses and nerves, and parasympathetic fibers, which come from the pelvic splanchnic nerves and the inferior hypogastric plexus. Sensation from the bladder is transmitted to the central nervous system (CNS) via general visceral afferent fibers.

This page titled 24.5C: Urinary Bladder is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5D: Urethra

The urethra is a muscular tube that connects the bladder with the outside of the body and removes urine from the body.

LEARNING OBJECTIVES

Describe the location and function of the urethra as part of the urinary system

KEY TAKEAWAYS

Key Points

- The external urethral sphincter is a striated muscle that allows voluntary control over urination.
- Since the urethra is much shorter in females, they are much more susceptible to bacterial infections or a urinary tract infection.
- The urethra has smooth muscle tissue, erectile tissue, and mucous membrane tissue. In males, the urethra travels through the penis and carries semen as well as urine.
- Semen is carried into the urethra through the vas deferens.


Key Terms

- **vagina**: A fibromuscular tubular tract that is the female sex organ and has two main functions: sexual intercourse and childbirth.
- **penis**: The male sexual organ for copulation and urination; also the tubular portion of the male genitalia (excluding the scrotum).

The Urethra

The urethra is a tube that connects the urinary bladder to the genitals for the removal of fluids from the body. The external urethral sphincter is a striated muscle that allows voluntary control over urination by controlling the flow of urine from the bladder into the urethra.

The Female Urethra



Female urethra: A depiction of a female urethra.


In females, the urethra is shorter relative to males, and emerges above the vaginal opening. Its lining is composed primarily of non-keratinized stratified squamous epithelium that becomes transitional near the bladder. The urethra consists of three layers of tissues:

1. The muscular layer is a continuation of the smooth muscle of the bladder.
2. The erectile urethral tissue is specialized tissue that may become engorged with blood during sexual arousal.
3. The mucuous urethral tissue are mucous membranes that line the interior of the urethra. The mucuous membranes are especially important for protecting the urethral tissues from the corrosive effects of urine.

The urethral sphincter separates the bladder from the urethra. Somatic (conscious) innervation of the external urethral sphincter is supplied by the pudendal nerve, which allows the sphincter to open and close. As the urethra is shorter in females relative to males, they are more vulnerable to bacterial urinary tract infections.

The Male Urethra

In males, the urethra travels through the penis and carries semen as well as urine. Semen does not flow through the bladder or the rest of the urinary tract, instead it is a fluid made of sperm cells and other fluids that passes through a few different glands from the testes to the urethra through the vas deferens. Aside from semen, and the male urethra's greater length, it is structurally similar to the female urethra.



Male urethra: A detailed view of the male urethra.

This page titled 24.5D: Urethra is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5E: Micturition and the Micturition Reflex

Micturition is the ejection of urine from the urinary bladder through the urethra to the outside of the body.

Learning Objectives

- Summarize the process of micturition via the micturition reflex

Key Points

- In infants, elderly individuals, and those with neurological injury, urination may occur as an involuntary reflex.
- Physiologically, micturition involves coordination among the central, autonomic, and somatic nervous systems that stem from different centers of the brain.
- The brain centers that regulate urination include the pontine micturition center, periaqueductal gray, and the cerebral cortex.
- Micturition consists of a storage phase and a voiding phase. Stretch receptors in the bladder increase their firing rate as the bladder becomes more full. This causes the micturition reflex, and increases urinary urge, and can even cause involuntary urination.

Key Terms

- **micturition**: Also known as urination, this is the ejection of urine from the urinary bladder through the urethra to the outside of the body.
- **urine**: A liquid excrement consisting of water, salts, and urea that is made in the kidneys then released through the urethra.


Micturition, also known as urination, is the ejection of urine from the urinary bladder through the urethra to the outside of the body. In healthy humans the process of urination is under voluntary control. In infants, elderly individuals, and those with neurological injury, urination may occur as an involuntary reflex.

Physiology of Micturition

Physiologically, micturition involves the coordination of the central, autonomic, and somatic nervous systems. The brain centers that regulate urination include the pontine micturition center, the periaqueductal gray, and the cerebral cortex, which cause both involuntary and voluntary control over micturition.

In males, urine is ejected through the penis, and in females through the urethral opening. Due to sexual dimorphism, and the positions where the urethra ends, males and females often use different techniques for urination. Micturition consists of two phases:

- The storage phase: A relaxed bladder in which urine slowly fills the bladder.
- The voiding phase: A contracted bladder that forces the external sphincter open and discharges urine through the urethra.



Location of urethral orifice in females: Location of the external urethral orifice in adult females. between the vestibule and the vaginal orifice.

The muscles controlling micturition are controlled by the autonomic and somatic nervous systems, which open the two sphincters during the voiding phase of micturition. During the storage phase the internal urethral sphincter is tense and the detrusor muscle is relaxed by sympathetic stimulation. During the voiding phase of micturition, parasympathetic stimulation causes the internal urethral sphincter to relax. The external urethral sphincter (sphincter urethrae) is under somatic control and is consciously relaxed (and thus opened) during micturition.

Many males prefer to urinate standing. In females, the urethra opens straight into the vulva. Because of this, the urine often does not exit at a distance from the body and is therefore seen as harder to control.

The Micturition Reflex

The state of the micturition reflex system is dependent on both a conscious signal from the brain and the firing rate of sensory stretch fibers from the bladder and urethra. At low bladder volumes, the afferent firing of the stretch receptors is low, and results in relaxation of the bladder. At high bladder volumes, the afferent firing of the stretch receptors increases, and causes a conscious sensation of urinary urge. This urge becomes stronger as the bladder becomes more full.

The micturition reflex causes bladder contraction during voiding, through a neural pathway. This reflex may lead to involuntary micturition in individuals that may not be able to feel the sensation of urinary urge, due to the firing of the stretch receptors themselves.

This page titled 24.5E: Micturition and the Micturition Reflex is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

24.5F: Waste Management in Other Body Systems

In addition to the kidneys, the liver, skin, and lungs also have important roles in the excretion of waste from the body.

Learning Objectives

- Differentiate among the non-urinary waste management systems


Key Points

- The lungs, liver, and skin are the main non- renal system organs involved in waste excretion.
- The liver is a vital organ with a wide range of functions, including detoxification, protein synthesis, and the production of the biochemicals necessary for digestion.
- The liver converts waste into other substances, but does not remove it from the body directly.
- The skin has sweat glands that secrete a fluid waste called perspiration, or sweat, which is a pathway for water and ion removal from the body, among other things.
- The lungs diffuse gaseous wastes, such as carbon dioxide, from the bloodstream as a normal part of respiration.

Key Terms

- **bilirubin**: A bile pigment that is a product of the breakdown of the heme portion of hemoglobin, that is processed into bile by the liver.
- **urea**: A water-soluble organic compound, $CO(NH2)2$, formed by the metabolism of proteins and excreted in the urine. The liver converts ammonia to urea.

Besides the renal system, many other organs and body systems are directly involved in the excretion of waste products. These other systems are responsible for the elimination of the waste products of the metabolism, as well as other liquid and gaseous wastes, but also provide other critical functions.



The lungs: The lungs are responsible for the removal of gaseous waste from the body.

Similar to renal clearance, these other organs have a clearance rate of their own, and also forms a part of total body clearance. The skin, lungs, and liver are the main waste removal organs outside of the non-renal system; however, their ability to remove

wastes is generally less than that of the renal system.

Liver Waste Removal

The liver has many vital functions, including a role in digestion, protein synthesis, and nutrient and vitamin storage. The liver also plays an important function in the management of body waste. It acts as a detoxification system for the body, processing and neutralizing drugs and toxins.

The liver is involved in the breakdown and recycling of red blood cells, including the removal of bilirubin from the body by secreting it into the bile, which then functions as a fat emulsifier in the digestive system.

Additionally, the liver is involved in the removal of toxic ammonia from the body by converting it to non-toxic urea, which is then excreted by the kidneys and put into urine during tubular secretion. While the liver may breakdown or neutralize waste in a different and safer form for the body, it doesn't directly remove the waste from the body.

Skin Waste Removal

Skin has sweat glands that secrete a fluid waste called perspiration. Perspiration, or sweat, is a fluid consisting primarily of water, as well as various dissolved solid wastes, that is excreted by the sweat glands. The sweat glands remove a bit of excess water and salts, and also serve the function of cooling the body during thermoregulation.

Respiratory Waste Removal

The lungs are responsible for the removal of gaseous waste from the body. Carbon dioxide is a toxic byproduct of cellular respiration. The lungs remove carbon dioxide from the bloodstream during ventilation, exchanging it for oxygen. This function is essentially important for regulating blood pH, since blood acidity is directly related to the concentration of carbon dioxide dissolved in carbonic acid in the blood stream. Excreting more carbon dioxide waste will reduce the acidity of blood.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...ation_of_Urine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ureter. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ureter**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- urinary bladder. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/urinary_bladder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Urinary_tract_en.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Urinar...y_tract_en.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...stem%23Ureters**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- transitional. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/transitional**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- urine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/urine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peristalsis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/peristalsis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Urinary_tract_en.png. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Urinary_system#/media/File:Urinary_tract_en.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Urinary system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Urinary_system.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/**

Human_P...stem%23Ureters. License: CC BY-SA: Attribution-ShareAlike
- vagina. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/vagina. License: CC BY-SA: Attribution-ShareAlike
- reflex. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/reflex. License: CC BY-SA: Attribution-ShareAlike
- urine. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/urine. License: CC BY-SA: Attribution-ShareAlike
- Urinary_tract_en.png. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Urinary_system#/media/File:Urinary_tract_en.png. License: CC BY-SA: Attribution-ShareAlike
- Urinary system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Urinary_system.svg. License: CC BY-SA: Attribution-ShareAlike
- Illu bladder. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/Fi...lu_bladder.jpg. License: CC BY-SA: Attribution-ShareAlike
- Human Physiology/The Urinary System. Provided by: Wikibooks. Located at: en.wikibooks.org/wiki/Human_P...stem%23Urethra. License: CC BY-SA: Attribution-ShareAlike
- penis. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/penis. License: CC BY-SA: Attribution-ShareAlike
- urine. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/urine. License: CC BY-SA: Attribution-ShareAlike
- vagina. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/vagina. License: CC BY-SA: Attribution-ShareAlike
- Urinary_tract_en.png. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Urinary_system#/media/File:Urinary_tract_en.png. License: CC BY-SA: Attribution-ShareAlike
- Urinary system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Urinary_system.svg. License: CC BY-SA: Attribution-ShareAlike
- Illu bladder. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Illu_bladder.jpg. License: CC BY-SA: Attribution-ShareAlike
- Human Physiology/The Urinary System. Provided by: Wikibooks. Located at: en.wikibooks.org/wiki/Human_P...stem%23Urethra. License: CC BY-SA: Attribution-ShareAlike
- File:Gray1142.png - Wikibooks, open books for an open world. Provided by: Wikibooks. Located at: en.wikibooks.org/w/index.php?...20051111182253. License: CC BY-SA: Attribution-ShareAlike
- Micturition reflex. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Micturi...x%23Physiology. License: CC BY-SA: Attribution-ShareAlike
- Micturition. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Micturition. License: CC BY-SA: Attribution-ShareAlike
- micturition. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/micturition. License: CC BY-SA: Attribution-ShareAlike
- urine. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/urine. License: CC BY-SA: Attribution-ShareAlike
- cerebral cortex. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/cerebral_cortex. License: CC BY-SA: Attribution-ShareAlike
- Urinary_tract_en.png. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Urinary_system#/media/File:Urinary_tract_en.png. License: CC BY-SA: Attribution-ShareAlike
- Urinary system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Urinary_system.svg. License: CC BY-SA: Attribution-ShareAlike
- Illu bladder. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Illu_bladder.jpg. License: CC BY-SA: Attribution-ShareAlike
- Human Physiology/The Urinary System. Provided by: Wikibooks. Located at: en.wikibooks.org/wiki/Human_P...stem%23Urethra. License: CC BY-SA: Attribution-ShareAlike
- File:Gray1142.png - Wikibooks, open books for an open world. Provided by: Wikibooks. Located at: en.wikibooks.org/w/index.php?title=File:Gray1142.png&filetimestamp=20051111182253. License: CC BY-SA: Attribution-ShareAlike
- Micturition reflex. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Micturi...x%23Physiology. License: CC BY-SA: Attribution-ShareAlike
- Excretory system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Excretory_system. License: CC BY-SA: Attribution-ShareAlike
- Liver. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Liver. License: CC BY-SA: Attribution-ShareAlike
- bilirubin. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/bilirubin. License: CC BY-SA: Attribution-ShareAlike
- urea. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/urea. License: CC BY-SA: Attribution-ShareAlike
- Urinary_tract_en.png. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Urinary_system#/media/File:Urinary_tract_en.png. License: CC BY-SA: Attribution-ShareAlike
- Urinary system. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Urinary_system.svg. License: CC BY-SA: Attribution-ShareAlike
- Illu bladder. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Illu_bladder.jpg. License: CC BY-SA: Attribution-ShareAlike

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/The_Urinary_System%23Urethra**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Gray1142.png - Wikibooks, open books for an open world. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/w/index.php?title=File:Gray1142.png&filetimestamp=20051111182253**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Micturition reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Micturition_reflex%23Physiology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lungs (animated). **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Lungs_(animated).gif**. **License**: ***Public Domain: No Known Copyright***

This page titled 24.5F: Waste Management in Other Body Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25: BODY FLUIDS AND ACID-BASE BALANCE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

25.1: Water Balance

25.1A: Regulation of Water Intake

25.1B: Regulation of Water Output

25.1C: Nitrogenous Waste in Terrestrial Animals: The Urea Cycle

25.1D: Water Balance Disorders

25.2: Body Fluids

25.2A: Water Content in the Body

25.2B: Fluid Compartments

25.2C: Body Fluid Composition

25.2D: Movement of Fluid Among Compartments

25.3: Electrolyte Balance

25.3A: Sodium, Electrolytes, and Fluid Balance

25.3B: Sodium Balance Regulation

25.3C: Potassium Balance Regulation

25.3D: Calcium and Phosphate Balance Regulation

25.3E: Anion Regulation

25.4: Acid-Base Balance

25.4A: pH, Buffers, Acids, and Bases

25.4B: Chemical Buffer Systems

25.4C: Regulation of H+ by the Lungs

25.4D: The Role of the Kidneys in Acid-Base Balance

This page titled 25: Body Fluids and Acid-Base Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.1: WATER BALANCE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

25.1A: Regulation of Water Intake

25.1B: Regulation of Water Output

25.1C: Nitrogenous Waste in Terrestrial Animals: The Urea Cycle

25.1D: Water Balance Disorders

This page titled 25.1: Water Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.1A: Regulation of Water Intake

Fluid can enter the body as preformed water, ingested food and drink, and, to a lesser extent, as metabolic water.

Learning Objectives

- Describe the regulation of water intake in the body

Key Points

- A constant supply of water is needed to replenish the fluids lost through normal physiological activities, such as respiration, sweating, and urination.
- Thirst is a sensation created by the hypothalamus that drives organisms to ingest water.
- Increased osmolarity in the blood acts on osmoreceptors that either stimulate the hypothalamus directly or cause the release of angiotensin II to stimulate the hypothalamus to cause thirst.
- The renin –angiotensin system increases thirst as a way to increase blood volume. It is activated by high plasma osmolarity, low blood volume, low blood pressure, and stimulation of the sympathetic nervous system.

Key Terms

- **thirst**: The sensation that drives organisms to ingest water. It is considered a basic survival instinct.
- **osmoreceptors**: Sensory receptors that are primarily found in the hypothalamus or macula densa that detect changes in the solute concentration of blood.

Water Intake

Fluid can enter the body as preformed water, ingested food and drink, and, to a lesser extent, as metabolic water that is produced as a by-product of aerobic respiration and dehydration synthesis. A constant supply is needed to replenish the fluids lost through normal physiological activities, such as respiration, sweating, and urination.

Water generated from the biochemical metabolism of nutrients provides a significant proportion of the daily water requirements for some arthropods and desert animals, but it provides only a small fraction of a human's necessary intake. In the normal resting state, the input of water through ingested fluids is approximately 2500 ml/day.

Body water homeostasis is regulated mainly through ingested fluids, which, in turn, depends on thirst. Thirst is the basic instinct or urge that drives an organism to ingest water.

Thirst is a sensation created by the hypothalamus, the thirst center of the human body. Thirst is an important component of blood volume regulation, which is slowly regulated by homeostasis.


Hypothalamus-Mediated Thirst

An osmoreceptor is a sensory receptor that detects changes in osmotic pressure and is primarily found in the hypothalamus of most homeothermic organisms. Osmoreceptors detect changes in plasma osmolarity (that is, the concentration of solutes dissolved in the blood).

When the osmolarity of blood changes (it is more or less dilute), water diffusion into and out of the osmoreceptor cells changes. That is, the cells expand when the blood plasma is more dilute and contract with a higher concentration.

When the osmoreceptors detect high plasma osmolarity (often a sign of a low blood volume), they send signals to the hypothalamus, which creates the biological sensation of thirst. Osmoreceptors also stimulate vasopressin (ADH) secretion, which

starts the events that will reduce plasma osmolarity to normal levels.



The hypothalamus: The hypothalamus is the thirst center of the human body.

Renin–Angiotensin System-Mediated Thirst

Another way through which thirst is induced is through angiotensin II, one of the hormones involved in the renin–angiotensin system. The renin–angiotensin system is a complex homeostatic pathway that deals with blood volume as a whole, as well as plasma osmolarity and blood pressure.

The macula densa cells in the walls of the ascending loop of Henle of the nephron is another type of osmoreceptor; however it stimulates the juxtaglomerular apparatus (JGA) instead of the hypothalamus. When the macula densa is stimulated by high osmolarity, The JGA releases renin into the bloodstream, which cleaves angiotensinogen into angiotensin I. Angiotensin I is converted into angiotensin II by ACE in the lungs. ACE is a hormone that has many functions.

Angiotensin II acts on the hypothalamus to cause the sensation of thirst. It also causes vasoconstriction, and the release of aldosterone to cause increased water reabsorption in a mechanism that is very similar to that of ADH.

Note that the renin–angiotensin system, and thus thirst, can be caused by other stimuli besides increased plasma osmolarity or a decrease in blood volume. For example, stimulation of the sympathetic nervous system and low blood pressure in the kidneys (decreased GFR) will stimulate the renin–angiotensin system and cause an increase in thirst.

This page titled 25.1A: Regulation of Water Intake is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.1B: Regulation of Water Output

Fluid can leave the body in three ways: urination, excretion (feces), and perspiration (sweating).

Learning Objectives

- Describe the regulation of water output in humans

Key Points

- The majority of fluid output occurs from urination. Some fluid is lost through perspiration (part of the body's temperature control mechanism) and as water vapor in expired air.
- The body's homeostatic control mechanisms ensure that a balance between fluid gain and fluid loss is maintained. The hormones ADH (antidiuretic hormone, also known as vasopressin) and aldosterone play a major role in this.
- If the body is becoming fluid deficient, increased plasma osmolarity is sensed by the osmoreceptors. This results in an increase in the secretion of ADH that causes fluid to be retained by the kidneys and urine output to be reduced.
- Aldosterone is the major end-product of the renin – angiotensin system, and increases the expression of ATPase pumps in the nephron that causes an increase in water reabsorption through sodium cotransport.
- ADH increases water reabsorption by increasing the nephron's permeability to water, while aldosterone works by increasing the reabsorption of both sodium and water.

Key Terms

- **osmoreceptors**: Sensory receptors, primarily found in the hypothalamus, that detect changes in plasma osmolarity and contribute to the fluid-balance regulation in the body.
- **anti-diuretic hormone**: A neurohypophysial hormone found in most mammals that is responsible for increasing water absorption in the collecting ducts of the kidney nephrons.
- **aldosterone**: A corticoid hormone that is secreted by the adrenal cortex that regulates the balance of sodium and potassium and thus the water-balance levels in the body.

Water Output

Fluid can leave the body in three ways:

1. Urination
2. Excretion (feces)
3. Perspiration (sweating)

The majority of fluid output occurs from urination, at approximately 1500 ml/day (approximately 1.59 qt/day) in a normal adult at resting state. Some fluid is lost through perspiration (part of the body's temperature control mechanism) and as water vapor in expired air; however these fluid losses are considered to be very minor.

The body's homeostatic control mechanisms maintain a constant internal environment to ensure that a balance between fluid gain and fluid loss is maintained. The hormones ADH (anti-diuretic hormone, also known as vasopressin) and aldosterone, a hormone created by the renin–angiotensin system, play a major role in this balance.

If the body is becoming fluid deficient, there will be an increase in the secretion of these hormones that causes water to be retained by the kidneys through increased tubular reabsorption and urine output to be reduced. Conversely, if fluid levels are excessive, the secretion of these hormones is suppressed and results in less retention of fluid by the kidneys and a subsequent increase in the volume of urine produced, due to reduced fluid retention.

ADH Feedback

When blood volume becomes too low, plasma osmolarity will increase due to a higher concentration of solutes per volume of water. Osmoreceptors in the hypothalamus detect the increased plasma osmolarity and stimulate the posterior pituitary gland to secrete ADH.


ADH causes the walls of the distal convoluted tubule and collecting duct to become permeable to water—this drastically increases the amount of water that is reabsorbed during tubular reabsorption. ADH also has a vasoconstrictive effect in the cardiovascular system, which makes it one of the most important compensatory mechanisms during hypovolemic shock (shock from excessive fluid loss or bleeding).

Aldosterone Feedback

Aldosterone is a steroid hormone (corticoid) produced at the end of the renin–angiotensin system. To review the renin–angiotensin system, low blood volume activates the juxtaglomerular apparatus in a variety of ways to make it secrete renin. Renin cleaves angiotensin I from the liver -produced angiotensinogen. Angiotensin converting enzyme (ACE) in the lungs converts angiotensin I into angiotensin II. Angiotensin II has a variety of effects (such as increasing thirst) but it also causes release of aldosterone from the adrenal cortex.

Aldosterone has a number of effects that are involved in the regulation of water output. It acts on mineral corticoid receptors in the epithelial cells of the distal convoluted tubule and collecting duct to increase their expression of Na^+/K^+ ATPase pumps and to activate those pumps. This causes greatly increased reabsorption of sodium and water (which follows sodium osmotically by cotransport), while causing the secretion of potassium into urine.

Aldosterone increases water reabsorption; however, it involves an exchange of sodium and potassium that ADH reabsoption regulation does not involve. Aldosterone will also cause a similar ion -balancing effect in the colon and salivary glands as well.



A schematic diagram of the renin–angiotensin system: Overview of the renin–angiotensin system that regulates blood pressure and plasma osmolarity.

This page titled 25.1B: Regulation of Water Output is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.1C: Nitrogenous Waste in Terrestrial Animals: The Urea Cycle

Urea, a nitrogenous waste material, is the end product excreted in urine when ammonia is metabolized by animals, such as mammals.

Learning Objectives

- Discuss the urea cycle

Key Points

- Ureotelic animals, which includes mammals, produce urea as the main nitrogenous waste material.
- $2\ NH_3 + CO_2 + 3\ ATP + H_2O \rightarrow H_2N\text{-}CO\text{-}NH_2 + 2\ ADP + 4\ Pi + AMP$ is the chemical reaction by which toxic ammonia is converted to urea.
- The urea cycle involves the multi-step conversion (carried out by five different enzymes) of the amino acid L- ornithine into different intermediates before being regenerated.

Key Terms

- **ureotelic**: animals that secrete urea as the primary nitrogenous waste material
- **ornithine**: an amino acid, which acts as an intermediate in the biosynthesis of urea
- **urea**: a water-soluble organic compound, CO(NH2)2, formed by the metabolism of proteins and excreted in the urine


Nitrogenous Waste in Terrestrial Animals: The Urea Cycle

Mammals, including humans, are the primary producers of urea. Because they secrete urea as the primary nitrogenous waste product, they are called ureotelic animals. Urea serves an important role in the metabolism of nitrogen-containing compounds by animals. It is the main nitrogen-containing substance in the urine of mammals. Urea is a colorless, odorless solid, highly soluble in water, and practically non-toxic. Dissolved in water, it is neither acidic nor alkaline. The body uses it in many processes, the most notable one being nitrogen excretion. Urea is widely used in fertilizers as a convenient source of nitrogen. It is also an important raw material for the chemical industry.

Apart from mammals, urea is also found in the urine of amphibians, as well as some fish. Interestingly, tadpoles excrete ammonia, but shift to urea production during metamorphosis. In humans, apart from being a carrier of waste nitrogen, urea also plays a role in the countercurrent exchange system of the nephrons, which allows for re-absorption of water and critical ions from the excreted urine. This mechanism, controlled by an anti-diuretic hormone, allows the body to create hyperosmotic urine, which has a higher concentration of dissolved substances than the blood plasma. This mechanism is important to prevent the loss of water, to maintain blood pressure, and to maintain a suitable concentration of sodium ions in the blood plasmas.

The urea cycle is the primary mechanism by which mammals convert ammonia to urea. Urea is made in the liver and excreted in urine. The overall chemical reaction by which ammonia is converted to urea is $2\ NH_3$ (ammonia) $+ CO_2 + 3\ ATP + H_2O \rightarrow H_2N\text{-}CO\text{-}NH_2$ (urea) $+ 2\ ADP + 4\ P_i + AMP$.

The urea cycle utilizes five intermediate steps, catalyzed by five different enzymes, to convert ammonia to urea. The amino acid L-ornithine is converted into different intermediates before being regenerated at the end of the urea cycle. Hence, the urea cycle is also referred to as the ornithine cycle. The enzyme ornithine transcarbamylase catalyzes a key step in the urea cycle. Its deficiency can lead to accumulation of toxic levels of ammonia in the body. The first two reactions occur in the mitochondria, while the last three reactions occur in the cytosol.



Urea Cycle: The urea cycle converts ammonia to urea in five steps that include the catalyzation of five different enzymes.

This page titled 25.1C: Nitrogenous Waste in Terrestrial Animals: The Urea Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.1D: Water Balance Disorders

Dehydration is the excessive loss of body fluid.

Learning Objectives

- Describe disorders due to water balance problems

Key Points

- There are three types of dehydration: hypotonic or hyponatremic, hypertonic or hypernatremic, and isotonic or isonatremic.
- Hypotonic dehydration is primarily a loss of electrolytes, sodium in particular.
- Hypertonic dehydration is primarily a loss of water.
- Isotonic dehydration is an equal loss of water and electrolytes.
- Hypvolemia is a loss of blood volume, and may cause hypovolemic shock. In humans, the most common type of dehydration by far is isotonic (isonatraemic) dehydration.
- Water balance disorders are generally treated by increasing water intake and reducing or stopping fluid loss.

Key Terms

- **isotonic**: When comparing solutions, an isotonic solution has the same osmolarity (ion concentration) as the solution it is being compared to.
- **plasma**: The straw-colored/pale-yellow, liquid component of blood that normally holds the blood cells of whole blood in suspension.
- **electrolyte**: Any of the various ions (such as sodium or chloride) that regulate the electric charge on cells and the flow of water across their membranes.

Water Balance Disorders

In physiology and medicine, dehydration (hypohydration) is defined as the excessive loss of body fluid. It is literally the removal of water from an object. However, in physiological terms, it entails a deficiency of fluid within an organism.

Much of the physiological effects of dehydration is due to the changes in ion concentration that may occur as a result of the dehydration. Alternatively, hypovolemia may occur due to loss of blood volume itself.

Dehydration

There are three types of dehydration that differ based on the type of change in ion concentrations:

1. Hypotonic—primarily a loss of electrolytes, sodium in particular. Hypotonic dehydration causes decreased plasma osmolarity.
2. Hypertonic—primarily a loss of water. Hypertonic dehydration causes increased plasma osmolarity.
3. Isotonic—an equal loss of water and electrolytes. Isotonic dehydration will not change plasma osmolarity, but it will reduce overall plasma volume. Isotonic dehydration is the most common type of dehydration.

Further complications may also occur. In hypotonic dehydration, intravascular water shifts to the extravascular space and exaggerates intravascular volume depletion for a given amount of total body water loss.

Neurological complications can occur in hypotonic and hypertonic states. The former can lead to seizures, while the latter can lead to osmotic cerebral edema upon rapid rehydration.

Hypovolemia

Hypovolemia is specifically a decrease in the volume of blood plasma. Furthermore, hypovolemia defines water deficiency in terms of blood volume rather than the overall water content of the body.



IV fluid and electrolyte administration: Intravenous administration of fluid is one effective treatment of dehydration in humans.

Hypovolemia is a cause of hypovolemic shock. Shock is any condition in which the body's fluids are unable to properly circulate and oxygenate the major organs of the human body; this causes compensatory mechanisms to activate that cause further bodily harm as the body's metabolism is maintained for a while longer.

In the case of hypovolemic shock, the tissue metabolism is impaired due to a lack of blood volume and makes it difficult for red blood cells to reach all of the tissues of the body. It is most often caused by severe vomiting, diarrhea, blood loss, or hemorrhage. Other forms of shock with similar symptoms may be due to problems in the heart (cardiogenic) or bacterial infection (septic).

Treatment Options

To treat minor dehydration water intake must be increased, while the source of fluid loss must be reduced or stopped altogether. Plain water restores only the volume of the blood plasma and inhibits the thirst mechanism before solute levels can be replenished.

Solid foods can contribute to fluid loss from vomiting and diarrhea. In more severe cases, correction of a dehydrated state is accomplished by the replenishment of necessary water and electrolytes through oral rehydration therapy or fluid replacement by intravenous therapy (an IV drip).

As oral rehydration is easier to provide, it is the treatment of choice for mild dehydration. Solutions used for intravenous rehydration must be isotonic or hypotonic. Pure water injected into the veins will cause the breakdown (lysis) of red blood cells that could cause other problems.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Free water clearance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Free_water_clearance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Osmoreceptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Osmoreceptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- Fluid balance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Fluid_balance. **License**: *CC BY-SA: Attribution-ShareAlike*
- dehydration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/dehydration. **License**: *CC BY-SA: Attribution-ShareAlike*
- thirst. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thirst. **License**: *CC BY-SA: Attribution-ShareAlike*
- osmoreceptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/osmoreceptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1806_The_Hypothalamus-Pituitary_Complex.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoth...ry_Complex.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Fluid balance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Fluid_balance. **License**: *CC BY-SA: Attribution-ShareAlike*
- aldosterone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/aldosterone. **License**: *CC BY-SA: Attribution-ShareAlike*
- anti-diuretic hormone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/anti-diuretic%20hormone. **License**: *CC BY-SA: Attribution-ShareAlike*
- osmoreceptors. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/osmoreceptors. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1806_The_Hypothalamus-Pituitary_Complex.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoth...ry_Complex.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renin-angiotensin system in man shadow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Re...man_shadow.svg. **License**: *Public Domain: No Known Copyright*
- OpenStax College, Biology. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44811/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Urea. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Urea. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. December 2, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44811/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...tion/ureotelic. **License**: *CC BY-SA: Attribution-ShareAlike*
- ornithine. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ornithine. **License**: *CC BY-SA: Attribution-ShareAlike*
- urea. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/urea. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1806_The_Hypothalamus-Pituitary_Complex.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoth...ry_Complex.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renin-angiotensin system in man shadow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Renin-angiotensin_system_in_man_shadow.svg. **License**: *Public Domain: No Known Copyright*
- OpenStax College, Nitrogenous Wastes. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44811/latest...e_41_04_01.jpg. **License**: *CC BY: Attribution*
- Dehydration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Dehydration. **License**: *CC BY-SA: Attribution-ShareAlike*
- isotonic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/isotonic. **License**: *CC BY-SA: Attribution-ShareAlike*
- electrolyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/electrolyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- plasma. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/plasma. **License**: *CC BY-SA: Attribution-ShareAlike*
- 1806_The_Hypothalamus-Pituitary_Complex.jpg. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypothalamus#/media/File:1806_The_Hypothalamus-Pituitary_Complex.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renin-angiotensin system in man shadow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Renin-angiotensin_system_in_man_shadow.svg. **License**: *Public Domain: No Known Copyright*
- OpenStax College, Nitrogenous Wastes. October 17, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44811/latest/Figure_41_04_01.jpg. **License**: *CC BY: Attribution*
- ICU IV 1. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:ICU_IV_1.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 25.1D: Water Balance Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.2: BODY FLUIDS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

25.2A: Water Content in the Body

25.2B: Fluid Compartments

25.2C: Body Fluid Composition

25.2D: Movement of Fluid Among Compartments

This page titled 25.2: Body Fluids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.2A: Water Content in the Body

A significant percentage of the human body is water, which includes intracellular and extracellular fluids.

Learning Objectives

- Describe the characteristics of water content in the body

Key Points


- On average, body water can account for 50% of the total human body weight and it is significantly higher in newborns. Obesity decreases the percentage of water in the body.
- Body water is regulated by hormones, including anti-diuretic hormone (ADH), aldosterone, and atrial natriuretic peptide.
- Water content in the body can be evaluated using bioelectrical impedance and mass spectrometry.
- Important functions of water in the body include supporting the cellular metabolism, molecular transport, biochemical reactions, and the physical properties of water, such as surface tension.

Key Terms

- **hydrolysis**: A biochemical reaction in which water molecules are used to break down a molecule into smaller molecules.
- **bioelectrical impedance analysis**: A commonly used method for estimating body composition, by measuring resistance to the flow of electricity in the body, which is associated with hydration levels.

Water Content

In physiology, body water is the water content of the human body. It makes up a significant percentage of the total composition of a body. Water is a necessary component to support life for many reasons. All cells in the human body are made mostly of water content in their cytoplasm.



Water molecule: A 3-dimensional model of hydrogen bonds (labeled 1) between molecules of water.

Water also provides a fluid environment for extracellular communication and molecular transport throughout the body. Water itself

is also a key component of biochemical reactions involved in physiology, such as hydrolysis. Many organ systems depend on the physical properties of water, such as the surface tension of water in the alveoli of the lungs.

Overall Water Content

The total amount of water in a human of average weight (70 kilograms) is approximately 40 liters, averaging 57 percent of his total body weight. In a newborn infant, this may be as high as 79 percent of the body weight, but it progressively decreases from birth to old age, with most of the decrease occurring during the first 10 years of life.

Also, obesity decreases the percentage of water in the body, sometimes to as low as 45 percent. The water in the body is distributed among various fluid compartments that are interspersed in the various cavities of the body through different tissue types. In diseased states where body water is affected, the fluid compartments that have changed can give clues to the nature of the problem.

Water Content Regulation and Measurement

Body water is regulated largely by the renal and neuro-endocrine systems. Water content regulation is one of the most important parts of homeostasis due to its influence on blood pressure and cardiac output. Much of this regulation is mediated by hormones, including anti-diuretic hormone (ADH), renin, angiotensin II, aldosterone, and atrial natriuretic peptide (ANP).

These hormones act as messengers between the kidneys and the hypothalamus; however, the lungs and heart are also involved in the secretion of some of these hormones, such as angiotensin converting enzyme (ACE) and ANP respectively.

There are many clinical methods to determine body water. One way to get an uncertain estimate is by calculation based on body weight and urine output. Another way to measure body water is through dilution and equilibration using mass spectrometry, which measures the abundance of water in breath samples from an individual.

In bioelectrical impedance analysis, a person's hydration level is calculated from high-precision measurements of the opposition to the flow of an electric current through body tissues. Since water conducts electricity, a lower hydration level will cause a greater amount of resistance to electrical flow through the body.

This page titled 25.2A: Water Content in the Body is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.2B: Fluid Compartments

The major body-fluid compartments includ: intracellular fluid and extracellular fluid (plasma, interstitial fluid, and transcellular fluid).

Learning Objectives

- Distinguish between intracellular and extracellular fluids

Key Points

- The intracellular fluid of the cytosol or intracellular fluid (or cytoplasmic matrix) is the liquid found inside cells.
- The cytosol is a complex mixture of substances that include proteins, ions, and organelles dissolved in water.
- Extracellular fluid (ECF) or extracellular fluid volume (ECFV) usually denotes all body fluid outside of cells, and consists of plasma, interstitial, and transcellular fluid.
- An extracellular matrix is an extracellular fluid space containing cell-excreted molecules, and they vary in their type and function.
- Plasma also serves as an extracellular matrices (ECM) for the cells and molecules of the blood.
- Interstitial fluid (or tissue fluid) is a solution that bathes and surrounds the cells of multicellular animals.
- Transcellular fluid is the portion of total body water contained within epithelial -lined spaces.

Key Terms

- **intracellular fluid**: The liquid found inside cells, between the endomembrane and the membrane-bound organelles.
- **interstitial fluid**: A solution that bathes and surrounds the cells of multicellular animals; also called tissue fluid.
- **plasma**: The straw-colored/pale-yellow, liquid component of blood that normally holds the blood cells of whole blood in suspension.

Fluid Compartments


The fluids of the various tissues of the human body are divided into fluid compartments. Fluid compartments are generally used to compare the position and characteristics of fluid in relation to the fluid within other compartments.

While fluid compartments may share some characteristics with the divisions defined by the anatomical compartments of the body, these terms are not one in the same. Fluid compartments are defined by their position relative to the cellular membrane of the cells that make up the body's tissues.

Intracellular Fluid

The intracellular fluid of the cytosol or intracellular fluid (or cytoplasm) is the fluid found inside cells. It is separated into compartments by membranes that encircle the various organelles of the cell. For example, the mitochondrial matrix separates the mitochondrion into compartments.

The contents of a eukaryotic cell within the cell membrane, excluding the cell nucleus and other membrane-bound organelles (e.g., mitochondria, plastides, lumen of endoplasmic reticulum, etc.), is referred to as the cytoplasm.



The cytosol: The cytosol (11) is the fluid within the plasma membrane of a cell and contains the organelles. The cytosol includes dissolved molecules and water.

The cytosol is a complex mixture of substances dissolved in water. Although water forms the large majority of the cytosol, it mainly functions as a fluid medium for intracellular signaling (signal transduction) within the cell, and plays a role in determining cell size and shape.


The concentrations of ions, such as sodium and potassium, are generally lower in the cytosol compared to the extracellular fluid; these differences in ion levels are important in processes such as osmoregulation and signal transduction. The cytosol also contains large amounts of macromolecules that can alter how molecules behave, through macromolecular crowding.

Extracellular Fluid

Extracellular fluid (ECF) or extracellular fluid volume (ECFV) usually denotes all the body fluid that is outside of the cells. The extracellular fluid can be divided into two major subcompartments: interstitial fluid and blood plasma.

The extracellular fluid also includes the transcellular fluid; this makes up only about 2.5% of the ECF. In humans, the normal glucose concentration of extracellular fluid that is regulated by homeostasis is approximately 5 mm. The pH of extracellular fluid is tightly regulated by buffers and maintained around 7.4.

The volume of ECF is typically 15L (of which 12L is interstitial fluid and 3L is plasma). The ECF contains extracellular matrices (ECMs) that act as fluids of suspension for cells and molecules inside the ECF.



Extracellular matrix: Spatial relationship between the blood vessels, basement membranes, and interstitial space between structures.

Blood Plasma

Blood plasma is the straw-colored/pale-yellow, liquid component of blood that normally holds the blood cells in whole blood in suspension, making it a type of ECM for blood cells and a diverse group of molecules. It makes up about 55% of total blood volume.

It is the intravascular fluid part of the extracellular fluid. It is mostly water (93% by volume) and contains dissolved proteins (the major proteins are fibrinogens, globulins, and albumins), glucose, clotting factors, mineral ions (Na+, Ca++, Mg++, HCO3- Cl-, etc.), hormones, and carbon dioxide (plasma is the main medium for excretory product transportation). It plays a vital role in intravascular osmotic effects that keep electrolyte levels balanced and protects the body from infection and other blood disorders.

Interstitial Fluid

Interstitial fluid (or tissue fluid) is a solution that bathes and surrounds the cells of multicellular animals. The interstitial fluid is found in the interstitial spaces, also known as the tissue spaces.

On average, a person has about 11 liters (2.4 imperial gallons or about 2.9 U.S. gal) of interstitial fluid that provide the cells of the body with nutrients and a means of waste removal. The majority of the interstitial space functions as an ECM, a fluid space consisting of cell-excreted molecules that lies between the basement membranes of the interstitial spaces. The interstitial ECM contains a great deal of connective tissue and proteins (such as collagen) that are involved in blood clotting and wound healing.

Transcellular Fluid

Transcellular fluid is the portion of total body water contained within the epithelial-lined spaces. It is the smallest component of extracellular fluid, which also includes interstitial fluid and plasma. It is often not calculated as a fraction of the extracellular fluid, but it is about 2.5% of the total body water.

Examples of this fluid are cerebrospinal fluid, ocular fluid, joint fluid, and the pleaural cavity that contains fluid that is only found in their respective epithelium-lined spaces.

The function of transcellular fluid is mainly lubrication of these cavities, and sometimes electrolyte transport.

This page titled 25.2B: Fluid Compartments is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.2C: Body Fluid Composition

The composition of tissue fluid depends upon the exchanges between the cells in the biological tissue and the blood.

Learning Objectives

- Describe the composition of intracellular and extracellular fluid in the body

Key Points

- The cytosol or intracellular fluid consists mostly of water, dissolved ions, small molecules, and large, water-soluble molecules (such as proteins).
- Enzymes in the cytosol are important for cellular metabolism.
- The extracellular fluid is mainly cations and anions.
- Plasma is mostly water and dissolved proteins, but also contains metabolic blood gasses, hormones, and glucose.
- The composition of transcellular fluid varies, but some of its main electrolytes include sodium ions, chloride ions, and bicarbonate ions.

Key Terms


- **electrolyte**: Any of the various ions (such as sodium or chloride) that regulate the electric charge on cells and the flow of water across their membranes.
- **transcellular fluid**: The portion of total body water contained within epithelial-lined spaces, such as the cerebrospinal fluid, and the fluid of the eyes and joints.
- **ion**: An atom or molecule in which the total number of electrons is not equal to the total number of protons, giving it a net positive or negative electrical charge.

Body Fluid Composition

The composition of tissue fluid depends upon the exchanges between the cells in the biological tissue and the blood. This means that fluid composition varies between body compartments.

Intracellular Fluid Composition

The cytosol or intracellular fluid consists mostly of water, dissolved ions, small molecules, and large, water-soluble molecules (such as proteins). This mixture of small molecules is extraordinarily complex, as the variety of enzymes that are involved in cellular metabolism is immense.



Ions: Ions in solution.

These enzymes are involved in the biochemical processes that sustain cells and activate or deactivate toxins. Most of the cytosol is water, which makes up about 70% of the total volume of a typical cell. The pH of the intracellular fluid is 7.4. The cell membrane separates cytosol from extracellular fluid, but can pass through the membrane via specialized channels and pumps during passive and active transport.

The concentrations of the other ions in cytosol or intracellular fluid are quite different from those in extracellular fluid. The cytosol also contains much higher amounts of charged macromolecules, such as proteins and nucleic acids, than the outside of the cell.

In contrast to extracellular fluid, cytosol has a high concentration of potassium ions and a low concentration of sodium ions. The reason for these specific sodium and potassium ion concentrations are Na+/K ATPase pumps that facilitate the active transport of these ions. These pumps transport ions against their concentration gradients to maintain the cytosol fluid composition of the ions.

Extracellular Fluid Composition

The extracellular fluid is mainly cations and anions. The cations include: sodium (Na+ = 136-145 mEq/L), potassium (K+ = 3.5-5.5 mEq/L) and calcium (Ca2+ = 8.4-10.5 mEq/L). Anions include: chloride (mEq/L) and hydrogen carbonate (HCO3- 22-26 mM). These ions are important for water transport throughout the body.

Plasma is mostly water (93% by volume) and contains dissolved proteins (the major proteins are fibrinogens, globulins, and albumins), glucose, clotting factors, mineral ions (Na+, Ca++, Mg++, HCO3- Cl- etc.), hormones and carbon dioxide (plasma being the main medium for excretory product transportation). These dissolved substances are involved in many varied physiological processes, such as gas exchange, immune system function, and drug distribution throughout the body.

Transcellular Fluid Composition

Due to the varying locations of transcellular fluid, the composition changes dramatically. Some of the electrolytes present in the transcellular fluid are sodium ions, chloride ions, and bicarbonate ions.

Cerebrospinal fluid is similar in composition to blood plasma, but lacks most proteins, such as albumins, because they are too large to pass through the blood–brain barrier. Ocular fluid in the eyes contrasts with cerebrospinal fluid by containing high concentrations of proteins, including antibodies.

This page titled 25.2C: Body Fluid Composition is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.2D: Movement of Fluid Among Compartments

How fluid moves through compartments depends on several variables described by Starling's equation.

Learning Objectives

- Describe the movement of fluid between extracellular compartments

Key Points

- Interstitial fluid is formed when hydrostatic pressure generated by the heart pushes water out of the capillaries. The water passes from a high concentration outside of the vessels to a low concentration inside of the vessels, but equilibrium is never reached because the constant blood flow.
- Osmotic pressure works opposite to hydrostatic pressure to hold water and substances in the capillaries.
- Hydrostatic pressure is stronger in the arterial ends of the capillaries, while osmotic pressure is stronger at the venous ends of the capillaries.
- Interstitial fluid is removed through the surrounding lymph vessels, and eventually ends up rejoining the blood. Sometimes the removal of tissue fluid does not function correctly and there is a buildup, called edema.
- The Starling equation describes the pressure gradients that drive the movement of water across fluid compartments.

Key Terms

- **Starling equation**: An equation that illustrates the role of hydrostatic and oncotic forces in the movement of fluid across capillary membranes.
- **interstitial fluid**: A solution that bathes and surrounds the cells of multicellular animals.

Fluid Movement

Extracellular fluid is separated among the various compartments of the body by membranes. These membranes are hydrophobic and repel water; however, there a few ways that fluids can move between body compartments. There are small gaps in membranes, such as the tight junctions, that allow fluids and some of their contents to pass through membranes by way of pressure gradients.

Formation of Interstitial Fluid

Hydrostatic pressure is generated by the contractions of the heart during systole. It pushes water out of the small tight junctions in the capillaries. The water potential is created due to the ability of the small solutes to pass through the walls of capillaries.

This buildup of solutes induces osmosis. The water passes from a high concentration (of water) outside of the vessels to a low concentration inside of the vessels, in an attempt to reach an equilibrium. The osmotic pressure drives water back into the vessels. Because the blood in the capillaries is constantly flowing, equilibrium is never reached.

The balance between the two forces differs at different points on the capillaries. At the arterial end of a vessel, the hydrostatic pressure is greater than the osmotic pressure, so the net movement favors water and other solutes being passed into the tissue fluid.

At the venous end, the osmotic pressure is greater, so the net movement favors substances being passed back into the capillary. This difference is created by the direction of the flow of blood and the imbalance in solutes created by the net movement of water that favors the tissue fluid.

Removal of Interstitial Fluid

The lymphatic system plays a part in the transport of tissue fluid by preventing the buildup of tissue fluid that surrounds the cells in the tissue. Tissue fluid passes into the surrounding lymph vessels and eventually rejoins the blood.

Sometimes the removal of tissue fluid does not function correctly and there is a buildup, which is called edema. Edema is responsible for the swelling that occurs during inflammation, and in certain diseases where the lymphatic drainage pathways are obstructed.


Starling Equation

image

The Starling model: Note the concentration of interstitial solutes (orange) increases proportionally to the distance from the arteriole.

Capillary permeability can be increased by the release of certain cytokines, anaphylatoxins, or other mediators (such as leukotrienes, prostaglandins, histamine, bradykinin, etc.) that are released by cells during inflammation. The Starling equation defines the forces across a semipermeable membrane to calculate the net flux.

The solution to the equation is known as the net filtration or net fluid movement. If positive, fluid will tend to leave the capillary (filtration). If negative, fluid will tend to enter the capillary (absorption). This equation has a number of important physiologic implications, especially when disease processes grossly alter one or more of the variables.



Capillary dynamics: Oncotic pressure exerted by the proteins in blood plasma tends to pull water into the circulatory system.

This is a diagram of the Starling model. Note how the concentration of interstitial solutes increases proportionally to the distance from the arteriole.

According to Starling's equation, the movement of fluid depends on six variables:

1. Capillary hydrostatic pressure (Pc)
2. Interstitial hydrostatic pressure (Pi)
3. Capillary oncotic pressure (πz)
4. Interstitial oncotic pressure (πi)
5. Filtration coefficient (Kf)
6. Reflection coefficient (σ)

The Starling Equation is mathematically described as Flux=Kf[(Pc-Pi)-σ (πz-πi)].

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Bioelectrical Impedance Analysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bioelectrical_Impedance_Analysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Body water. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Body_water**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bioelectrical impedance analysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bioelec...nce%20analysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/extracellular%20fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intracellular%20fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Water. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Water**. **License**: ***Public Domain: No Known Copyright***
- Extracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extracellular_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Body water. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Body_water**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Interstitial fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Interstitial_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intracellular_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- plasma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/plasma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intracellular%20fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- interstitial fluid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/interstitial_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Water. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Water**. **License**: ***Public Domain: No Known Copyright***
- Extracellular Matrix. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ex...lar_Matrix.png**. **License**: ***Public Domain: No Known Copyright***
- Biological cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Biological_cell.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Extracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extracellular_fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Interstitial fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Interstitial_fluid%23Composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intracellular fluid. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intracellular_fluid%23Properties_and_composition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ion. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/transcellular-fluid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- electrolyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/electrolyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Water. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Water**. **License**: ***Public Domain: No Known Copyright***
- Extracellular Matrix. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Extracellular_Matrix.png**. **License**: ***Public Domain: No Known Copyright***
- Biological cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Biological_cell.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Experiment #11: Testing Water Filters. **Provided by**: Chemistry Land. **Located at**: **http://www.chemistryland.com/CHM107Lab/Lab5/Filters/Lab5Exp2filters.html**. **License**: ***Public Domain: No Known Copyright***

- interstitial fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/interstitial_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Capillary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Capillary. **License**: *CC BY-SA: Attribution-ShareAlike*
- Starling equation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Starling_equation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Interstitial fluid. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Interstitial_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Capillary. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Capillary. **License**: *CC BY-SA: Attribution-ShareAlike*
- Starling equation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Starling%20equation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Water. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Water. **License**: *Public Domain: No Known Copyright*
- Extracellular Matrix. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Extracellular_Matrix.png. **License**: *Public Domain: No Known Copyright*
- Biological cell. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Biological_cell.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Experiment #11: Testing Water Filters. **Provided by**: Chemistry Land. **Located at**: http://www.chemistryland.com/CHM107Lab/Lab5/Filters/Lab5Exp2filters.html. **License**: *Public Domain: No Known Copyright*
- Capillary dynamics. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...irculation.jpg. **License**: *Public Domain: No Known Copyright*

This page titled 25.2D: Movement of Fluid Among Compartments is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3: ELECTROLYTE BALANCE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

25.3A: Sodium, Electrolytes, and Fluid Balance

25.3B: Sodium Balance Regulation

25.3C: Potassium Balance Regulation

25.3D: Calcium and Phosphate Balance Regulation

25.3E: Anion Regulation

This page titled 25.3: Electrolyte Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3A: Sodium, Electrolytes, and Fluid Balance

Electrolytes play a vital role in maintaining homeostasis within the body.

Learning Objectives

- Identify the importance of sodium and fluid/electrolyte balance

Key Points

- Electrolytes help to regulate myocardial and neurological functions, fluid balance, oxygen delivery, acid–base balance, and much more.
- The most serious electrolyte disturbances involve abnormalities in the levels of sodium, potassium, and/or calcium.
- Kidneys work to keep the electrolyte concentrations in the blood constant despite changes in the body.

Key Terms

- **homeostasis**: The ability of a system or living organism to adjust its internal environment to maintain a stable equilibrium; such as the ability of warm-blooded animals to maintain a constant temperature.
- **electrolyte**: Any of the various ions (such as sodium or chloride) that regulate the electric charge on cells and the flow of water across their membranes.
- **sodium**: A chemical element with symbol Na (from Latin: natrium) and atomic number 11. It is a soft, silvery white, highly reactive metal and is a member of the alkali metals.

Importance of Electrolyte Balance

Electrolytes play a vital role in maintaining homeostasis within the body. They help regulate myocardial and neurological function, fluid balance, oxygen delivery, acid-base balance, and other biological processes.

Electrolytes are important because they are what cells (especially those of the nerve, heart, and muscle) use to maintain voltages across their cell membranes and to carry electrical impulses (nerve impulses, muscle contractions) across themselves and to other cells.

Electrolyte imbalances can develop from excessive or diminished ingestion and from the excessive or diminished elimination of an electrolyte. The most common cause of electrolyte disturbances is renal failure. The most serious electrolyte disturbances involve abnormalities in the levels of sodium, potassium, and/or calcium.

Other electrolyte imbalances are less common, and often occur in conjunction with major electrolyte changes. Chronic laxative abuse or severe diarrhea or vomiting (gastroenteritis) can lead to electrolyte disturbances combined with dehydration. People suffering from bulimia or anorexia nervosa are especially at high risk for an electrolyte imbalance.

Kidneys work to keep the electrolyte concentrations in blood constant despite changes in your body. For example, during heavy exercise electrolytes are lost through sweating, particularly sodium and potassium, and sweating can increase the need for electrolyte (salt) replacement. It is necessary to replace these electrolytes to keep their concentrations in the body fluids constant.

Dehydration

There are three types of dehydration:

1. Hypotonic or hyponatremic (primarily a loss of electrolytes, sodium in particular).
2. Hypertonic or hypernatremic (primarily a loss of water).


3. Isotonic or isonatremic (an equal loss of water and electrolytes).

In humans, the most common type of dehydration by far is isotonic (isonatraemic) dehydration; which effectively equates with hypovolemia; but the distinction of isotonic from hypotonic or hypertonic dehydration may be important when treating people with dehydration.

Physiologically, and despite the name, dehydration does not simply mean loss of water, as both water and solutes (mainly sodium) are usually lost in roughly equal quantities as to how they exist in blood plasma. In hypotonic dehydration, intravascular water shifts to the extravascular space and exaggerates the intravascular volume depletion for a given amount of total body water loss.

Neurological complications can occur in hypotonic and hypertonic states. The former can lead to seizures, while the latter can lead to osmotic cerebral edema upon rapid rehydration.

In more severe cases, the correction of a dehydrated state is accomplished by the replenishment of necessary water and electrolytes (through oral rehydration therapy or fluid replacement by intravenous therapy). As oral rehydration is less painful, less invasive, less expensive, and easier to provide, it is the treatment of choice for mild dehydration. Solutions used for intravenous rehydration must be isotonic or hypotonic.



Cell electrolytes: This diagram illustrates the mechanism for the transportation of water and electrolytes across the epithelial cells in the secretory glands.

This page titled 25.3A: Sodium, Electrolytes, and Fluid Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3B: Sodium Balance Regulation

Sodium is an important cation that is distributed primarily outside the cell.

Learning Objectives

- Describe the mechanisms by which sodium balance is regulated

Key Points

- The body has a potent sodium -retaining mechanism: the rennin– angiotensin system.
- In states of sodium depletion, aldosterone levels increase; in states of sodium excess, aldosterone levels decrease.
- The major physiological controller of aldosterone secretion is the plasma angiotensin II level that increases aldosterone secretion.

Key Terms

- **sodium**: A chemical element with symbol Na (from Latin: natrium) and atomic number 11. It is a soft, silvery white, highly reactive metal and is a member of the alkali metals.
- **aldosterone**: A mineralocorticoid hormone that is secreted by the adrenal cortex and regulates the balance of sodium and potassium in the body.
- **angiotensin**: Any of several polypeptides that narrow the blood vessels and regulate arterial pressure.

Sodium Regulation


Sodium is an important cation that is distributed primarily outside the cell. The cell sodium concentration is about 15 mmol/l, but it varies in different organs; it has an intracellular volume of 30 liters and about 400 mmol are inside the cell.

The plasma and interstitial sodium is about 140 mmol/l with an extracellular volume of about 13 liters, 1,800 mmol are in the extracellular space. The total body sodium, however, is about 3,700 mmol as there is about 1,500 mmol stored in bones.

The body has potent sodium-retaining mechanisms and even if a person is on five mmol Na+/day they can maintain sodium balance. Extra sodium is lost from the body by reducing the activity of the renin –angiotensin system that leads to increased sodium loss from the body. Sodium is lost through the kidneys, sweat, and feces.

In states of sodium depletion, the aldosterone levels increase. In states of sodium excess, aldosterone levels decrease. The major physiological controller of aldosterone secretion is the plasma angiotensin II level that increases aldosterone secretion.

A high plasma potassium level also increases aldosterone secretion because, besides retaining Na+, high plasma aldosterone causes K+ loss by the kidney. Plasma Na+ levels have little effect on aldosterone secretion.



Renin–angiotensin system: The regulation of sodium via the hormones renin, angiotensin, and aldosterone. In states of sodium depletion, the aldosterone levels increase, and in states of sodium excess, the aldosterone levels decrease.

A low renal perfusion pressure stimulates the release of renin, which forms angiotensin I that is converted to angiotensin II. Angiotensin II will correct the low perfusion pressure by causing the blood vessels to constrict, and increase sodium retention by its direct effect on the proximal renal tubule and by an effect operated through aldosterone. The perfusion pressure to the adrenal gland has little direct effect on aldosterone secretion and the low blood pressure operates to control aldosterone via the renin–angiotensin system.

Aldosterone also acts on the sweat ducts and colonic epithelium to conserve sodium. When aldosterone is activated to retain sodium the plasma sodium tends to rise. This immediately causes the release of ADH, which causes water to be retained, thus balancing Na+ and H2O in the right proportion to restore plasma volume.

In addition to aldosterone and angiotensin II, other factors influence sodium excretion.

- Atrial peptide causes the loss of sodium by the kidneys: it is secreted from the heart in high sodium states due to excess intake or cardiac disease.
- Elevated blood pressure will also cause Na+ loss, and a low blood pressure usually leads to sodium retention.

This page titled 25.3B: Sodium Balance Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3C: Potassium Balance Regulation

Potassium is mainly an intracellular ion.

Learning Objectives

- Describe the mechanisms of potassium balance regulation

Key Points

- Most of the total body potassium is inside the cells and the next largest proportion is in the bones.
- In an unprocessed diet, potassium is much more plentiful than sodium and it is present as an organic salt, while sodium is added as NaCl.
- High potassium intake can potentially increase the extracellular K+ level two times before the kidney can excrete the extra potassium.
- A high plasma potassium increases aldosterone secretion and this increases the potassium loss from the body to restore balance.

Key Terms

- **alkalotic**: A condition that reduces the hydrogen ion concentration of arterial blood plasma (alkalemia). Generally, alkalosis is said to occur when the blood pH exceeds 7.45.
- **Potassium**: A chemical element with the symbol K and the atomic number 19. Elemental potassium is a soft, silvery white, alkali metal that oxidizes rapidly in the air and is very reactive with water—it can generate sufficient heat to ignite the hydrogen emitted in the reaction.
- **acidosis**: An increase in acidity of the blood and other body tissue (i.e., an increased hydrogen ion concentration). If not further qualified, it usually refers to the acidity of the blood plasma.

Potassium Balance

Potassium is predominantly an intracellular ion. Most of the total body potassium of about 4,000 mmol is inside the cells, and the next largest proportion (300–500 mmol) is in the bones. Cell K+ concentration is about 150 mmol/l but varies in different organs. Extracellular potassium is about 4.0 mmol/l, with an extracellular value of about 13 liters, 52 mmol (i.e., less than 1.5%) is present here and only 12 mmol is in the plasma.

In an unprocessed diet potassium is much more plentiful than sodium. It is present as an organic salt, while sodium is added as NaCl. In a hunter-gatherer, K+ intake may be as much as 400 mmol/d while in the Western diet it is 70 mmol/d or less if a person has a minimal amount of fresh fruit and vegetables.

The processing of foods replaces K+ with NaCl. While the body can excrete a large K+ load, it is unable to conserve K+. On a zero K+ intake, or in a person with K+ depletion, there will still be a loss of K+ of 30–50 mmol/d in the urine and feces.

Acid– Base Status Control

If there is a high potassium intake, for example, 100 mmol, this would potentially increase the extracellular K+ level two times before the kidney could excrete the extra potassium. The body buffers the extra potassium by equilibrating it within the cells.

The acid–base status controls the distribution between plasma and cells. A high pH (i.e., alkalosis >7.4) favors the movement of K+ into the cells, and a low pH (i.e., acidosis) causes movement out of the cell. A high plasma potassium level increases aldosterone secretion and this increases the potassium loss from the body to restore balance.

This change of distribution with the acid–base status means that the plasma K+ may not reflect the total body content. Therefore, a person with an acidosis (pH 7.1) and a plasma K+ of 6.5 mmol/l could be depleted of total body potassium. This occurs in diabetic acidosis. Conversely, a person who is alkalotic with a plasma K+ of 3.4 mmol/l may have a normal level of total body potassium.

This page titled 25.3C: Potassium Balance Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3D: Calcium and Phosphate Balance Regulation

Calcium is a key electrolyte: 99% is deposited in the bones and the remainder is associated with hormone release and cell signaling.

Learning Objectives

- Describe calcium and phosphate balance regulation

Key Points

- Calcium absorption is controlled by vitamin D, and calcium excretion is controlled by the parathyroid hormones.
- There is a constant loss of calcium by the kidney even if there is none in the diet.
- Calcium in plasma exists in three forms: ionized, nonionized and protein bound.

Key Terms

- **calcium**: A chemical element, atomic number 20, that is an alkaline earth metal and occurs naturally as carbonate in limestone and as silicate in many rocks.
- **parathyroid hormone**: A polypeptide hormone that is released by the chief cells of the parathyroid glands and is involved in raising the levels of calcium ions in the blood.
- **vitamin D**: A fat-soluble vitamin that is required for normal bone development and that prevents rickets; it can be manufactured in the skin on exposure to sunlight.

Calcium is a very important electrolyte. Ninety-nine percent or more is deposited in the bones and the remainder plays a vital role in nerve conduction, muscle contraction, hormone release, and cell signaling.

The plasma concentration of Ca++ is 2.2 mmol/l, and phosphate is 1.0 mmol/l. The solubility product of Ca and P is close to saturation in plasma. The concentration of Ca++ in the cytoplasm is < 10–6 mmol/l but the concentration of Ca++ in the cell is much higher as calcium is taken up (and is able to be released from) cell organelles.

In the typical Australian diet, there is about 1200 mg/d of calcium. Even if it was all soluble it is not all absorbed as it combines with phosphates in the intestinal secretions. In addition, absorption is regulated by the active vitamin D; increased amounts of vitamin D increase Ca++ absorption.

Absorption is controlled by vitamin D while excretion is controlled by parathyroid hormones. However, the distribution from bone to plasma is controlled by both the parathyroid hormones and vitamin D.

There is also a constant loss of calcium via the kidneys even if there is none in the diet. This excretion of calcium by the kidneys and its distribution between bone and the rest of the body is primarily controlled by the parathyroid hormone.


The calcium in plasma exists in three forms:

1. Ionized.
2. Nonionized.
3. Protein bound.

It is the ionized calcium concentration that is monitored by the parathyroid gland —if it is low, parathyroid hormone secretion is increased. This increases the ionized calcium levels by increasing bone re-absorption, decreasing renal excretion, and acting on the kidney to increase the rate of formation of active vitamin D, thereby increasing the gut's absorption of calcium.

The usual amount of phosphate in the diet is about 1 g/d but not all of it is absorbed. Any excess is excreted by the kidney and this excretion is increased by the parathyroid hormone.

This hormone also causes phosphate to leach out of the bones. Plasma phosphate has no direct effect on parathyroid hormone secretion; however, if it is elevated it combines with Ca++ to decrease ionized Ca++ in plasma, and thereby increase parathyroid hormone secretion.



Calcium regulation: This is an illustration of how parathyroid hormone regulates the levels of calcium in the blood.

This page titled 25.3D: Calcium and Phosphate Balance Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.3E: Anion Regulation

The anions chloride, bicarbonate, and phosphate have important roles in maintaining the balances and neutrality of vital body mechanisms.

Learning Objectives

- Describe anion regulation in the body

Key Points

- Chloride is needed to maintain proper hydration, as well as to balance cations, and maintain the electrical neutrality of the extracellular fluid.
- Bicarbonate ‘s main role is to maintain the body’s acid–base balance through a buffer system.
- Phosphate is a major constituent of the intracellular fluid, and it is important in the regulation of metabolic processes and as a buffering agent in animal cells.
- The kidneys regulate the salt balance in the blood by controlling the excretion and the reabsorption of various ions.

Key Terms

- **anion**: An negatively charged ion.
- **hyperphosphatemia**: An elevated amount of phosphate in the blood.
- **hypochloremia**: An electrolyte disturbance caused by an abnormally depleted level of chloride ions in the blood.
- **hypophosphatemia**: An electrolyte disturbance caused by an abnormally low level of phosphate in the blood.

Anion Regulation

The excretion of ions occurs mainly through the kidneys, with lesser amounts of ions being lost in sweat and in feces. In addition, excessive sweating may cause a significant loss, especially of the anion chloride. Severe vomiting or diarrhea will also cause a loss of chloride and bicarbonate ions.

Adjustments in the respiratory and renal functions allow the body to regulate the levels of these ions in the extracellular fluid (ECF).

Chloride

Chloride is the predominant extracellular anion and it is a major contributor to the osmotic pressure gradient between the intracellular fluid (ICF) and extracellular fluid (ECF). Chloride maintains proper hydration and functions to balance the cations in the ECF to keep the electrical neutrality of this fluid. The paths of secretion and reabsorption of chloride ions in the renal system follow the paths of sodium ions.

Hypochloremia, or lower-than-normal blood chloride levels, can occur because of defective renal tubular absorption. Vomiting, diarrhea, and metabolic acidosis can also lead to hypochloremia.

In contrast, hyperchloremia, or higher-than-normal blood chloride levels, can occur due to dehydration, excessive intake of dietary salt (NaCl) or the swallowing of sea water, aspirin intoxication, congestive heart failure, and the hereditary, chronic lung disease cystic fibrosis. In people who have cystic fibrosis, the chloride levels in their sweat are two to five times those of normal levels; therefore, analysis of their sweat is often used to diagnose the disease.

Bicarbonate


Bicarbonate is the second-most abundant anion in the blood. Its principal function is to maintain your body's acid–base balance by being part of buffer systems.

Bicarbonate ions result from a chemical reaction that starts with the carbon dioxide (CO_2) and water (H_2O) molecules that are produced at the end of aerobic metabolism. Only a small amount of CO_2 can be dissolved in body fluids; thus, over 90 percent of the CO_2 is converted into bicarbonate ions, HCO_3-, through the following reactions:

$$CO_2 + H_2O \leftrightarrow H_2CO_3 \leftrightarrow H_2CO_3^- + H^+$$

The bidirectional arrows indicate that the reactions can go in either direction depending on the concentrations of the reactants and products. Carbon dioxide is produced in large amounts in tissues that have a high metabolic rate, and is converted into bicarbonate in the cytoplasm of the red blood cells through the action of an enzyme called carbonic anhydrase.

Bicarbonate is transported in the blood and once in the lungs, the reactions reverse direction, and CO_2 is regenerated from the bicarbonate to be exhaled as metabolic waste.



Bicarbonate as a buffering system: In the lungs, CO_2 is produced from bicarbonate and removed as metabolic waste through the reverse reaction of the bicarbonate bidirectional equation.

Phosphate

Phosphate is present in the body in three ionic forms:

1. $H_2PO_4^-$
2. HPO_4^{2-}
3. PO_4^{3-}

The addition and removal of phosphate from the proteins in all cells is a pivotal strategy in the regulation of metabolic processes. Phosphate is useful in animal cells as a buffering agent, and the most common form is HPO^{2-}_4. Bone and teeth bind up 85 percent of the body's phosphate as part of calcium phosphate salts. In addition, phosphate is found in phospholipids, such as those that make up the cell membrane, and in ATP, nucleotides, and buffers.

Hypophosphatemia, or abnormally low phosphate blood levels, occurs with the heavy use of antacids, during alcohol withdrawal, and during malnourishment. In the face of phosphate depletion, the kidneys usually conserve phosphate, but during starvation,

this conservation is impaired greatly.

Hyperphosphatemia, or abnormally increased levels of phosphates in the blood, occurs if there is decreased renal function or in cases of acute lymphocytic leukemia. Additionally, because phosphate is a major constituent of the ICF, any significant destruction of cells can result in the dumping of phosphate into the ECF.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Water-electrolyte imbalance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Water-e...lyte_imbalance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...3Water_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...3Water_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Water-electrolyte imbalance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Water-e...lyte_imbalance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sodium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sodium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homeostasis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homeostasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- electrolyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/electrolyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cell electrolytes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ectrolytes.png**. **License**: ***CC BY: Attribution***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...Sodium_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...Sodium_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sodium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sodium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- angiotensin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/angiotensin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- aldosterone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/aldosterone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cell electrolytes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ectrolytes.png**. **License**: ***CC BY: Attribution***
- Renin-angiotensin system. **Provided by**: Wikimedia commons. **Located at**: **en.Wikipedia.org/wiki/File:Re...one_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...assium_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- acidosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...n/potassium--2**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...tion/alkalotic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cell electrolytes. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ectrolytes.png**. **License**: ***CC BY: Attribution***
- Renin-angiotensin system. **Provided by**: Wikimedia commons. **Located at**: **en.Wikipedia.org/wiki/File:Re...one_system.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...sphate_Balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Human Physiology/Homeostasis. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...sphate_Balance. **License**: *CC BY-SA: Attribution-ShareAlike*
- calcium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/calcium. **License**: *CC BY-SA: Attribution-ShareAlike*
- vitamin D. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vitamin_D. **License**: *CC BY-SA: Attribution-ShareAlike*
- parathyroid hormone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/parathyroid_hormone. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cell electrolytes. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ectrolytes.png. **License**: *CC BY: Attribution*
- Renin-angiotensin system. **Provided by**: Wikimedia commons. **Located at**: en.Wikipedia.org/wiki/File:Re...one_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Calcium regulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ca...regulation.png. **License**:
- OpenStax College, Electrolyte Balance. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46414/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Electrolyte Balance. January 15, 2014. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m46414/latest/. **License**: *CC BY: Attribution*
- Phosphate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Phosphate. **License**: *CC BY-SA: Attribution-ShareAlike*
- hypophosphatemia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypophosphatemia. **License**: *CC BY-SA: Attribution-ShareAlike*
- hyperphosphatemia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hyperphosphatemia. **License**: *CC BY-SA: Attribution-ShareAlike*
- anion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/anion. **License**: *CC BY-SA: Attribution-ShareAlike*
- hypochloremia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hypochloremia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cell electrolytes. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...ectrolytes.png. **License**: *CC BY: Attribution*
- Renin-angiotensin system. **Provided by**: Wikimedia commons. **Located at**: en.Wikipedia.org/wiki/File:Re...one_system.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Calcium regulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Ca...regulation.png. **License**:
- Riassorbimento bicarbonati e respirazione cellulare. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._cellulare.svg. **License**:

This page titled 25.3E: Anion Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.4: ACID-BASE BALANCE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

25.4A: pH, Buffers, Acids, and Bases

25.4B: Chemical Buffer Systems

25.4C: Regulation of H+ by the Lungs

25.4D: The Role of the Kidneys in Acid-Base Balance

This page titled 25.4: Acid-Base Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.4A: pH, Buffers, Acids, and Bases

Acids dissociate into H^+ and lower pH, while bases dissociate into OH^- and raise pH; buffers can absorb these excess ions to maintain pH.

Learning Objectives

- Explain the composition of buffer solutions and how they maintain a steady pH

Key Points

- A basic solution will have a pH above 7.0, while an acidic solution will have a pH below 7.0.
- Buffers are solutions that contain a weak acid and its a conjugate base; as such, they can absorb excess H^+ ions or OH^-ions, thereby maintaining an overall steady pH in the solution.
- pH is equal to the negative logarithm of the concentration of H^+ ions in solution: $pH = -\log[H^+]$.

Key Terms

- **alkaline**: having a pH greater than 7; basic
- **acidic**: having a pH less than 7
- **buffer**: a solution composed of a weak acid and its conjugate base that can be used to stabilize the pH of a solution

Self-Ionization of Water

Hydrogen ions are spontaneously generated in pure water by the dissociation (ionization) of a small percentage of water molecules into equal numbers of hydrogen (H^+) ions and hydroxide (OH^-) ions. The hydroxide ions remain in solution because of their hydrogen bonds with other water molecules; the hydrogen ions, consisting of naked protons, are immediately attracted to un-ionized water molecules and form hydronium ions (H_3O^+). By convention, scientists refer to hydrogen ions and their concentration as if they were free in this state in liquid water.

2H2O⇌H3O++OH−2H2O⇌H3O++OH−


The concentration of hydrogen ions dissociating from pure water is 1×10^{-7} moles H^+ ions per liter of water. The pH is calculated as the negative of the base 10 logarithm of this concentration:

$pH = -\log[H^+]$

The negative log of 1×10^{-7} is equal to 7.0, which is also known as neutral pH. Human cells and blood each maintain near-neutral pH.

pH Scale

The pH of a solution indicates its acidity or basicity (alkalinity). The pH scale is an inverse logarithm that ranges from 0 to 14: anything below 7.0 (ranging from 0.0 to 6.9) is acidic, and anything above 7.0 (from 7.1 to 14.0) is basic (or alkaline). Extremes in pH in either direction from 7.0 are usually considered inhospitable to life. The pH in cells (6.8) and the blood (7.4) are both very close to neutral, whereas the environment in the stomach is highly acidic, with a pH of 1 to 2.



The pH scale: The pH scale measures the concentration of hydrogen ions (H^+) in a solution.

Non-neutral pH readings result from dissolving acids or bases in water. Using the negative logarithm to generate positive integers, high concentrations of hydrogen ions yield a low pH, and low concentrations a high pH.

An acid is a substance that increases the concentration of hydrogen ions (H^+) in a solution, usually by dissociating one of its hydrogen atoms. A base provides either hydroxide ions (OH^-) or other negatively-charged ions that react with hydrogen ions in solution, thereby reducing the concentration of H^+ and raising the pH.


Strong Acids and Strong Bases

The stronger the acid, the more readily it donates H^+. For example, hydrochloric acid (HCl) is highly acidic and completely dissociates into hydrogen and chloride ions, whereas the acids in tomato juice or vinegar do not completely dissociate and are considered weak acids; conversely, strong bases readily donate OH^- and/or react with hydrogen ions. Sodium hydroxide (NaOH) and many household cleaners are highly basic and give up OH^- rapidly when placed in water; the OH^- ions react with H^+ in solution, creating new water molecules and lowering the amount of free H^+ in the system, thereby raising the overall pH. An example of a weak basic solution is seawater, which has a pH near 8.0, close enough to neutral that well-adapted marine organisms thrive in this alkaline environment.

Buffers

How can organisms whose bodies require a near-neutral pH ingest acidic and basic substances (a human drinking orange juice, for example) and survive? Buffers are the key. Buffers usually consist of a weak acid and its conjugate base; this enables them to readily absorb excess H^+ or OH^-, keeping the system's pH within a narrow range.

Maintaining a constant blood pH is critical to a person's well-being. The buffer that maintains the pH of human blood involves carbonic acid (H_2CO_3), bicarbonate ion (HCO_3^-), and carbon dioxide (CO_2). When bicarbonate ions combine with free hydrogen ions and become carbonic acid, hydrogen ions are removed, moderating pH changes. Similarly, excess carbonic acid can be converted into carbon dioxide gas and exhaled through the lungs; this prevents too many free hydrogen ions from building up in the blood and dangerously reducing its pH; likewise, if too much OH^- is introduced into the system, carbonic acid will combine with it to create bicarbonate, lowering the pH. Without this buffer system, the body's pH would fluctuate enough to jeopardize survival.



Buffers in the body: This diagram shows the body's buffering of blood pH levels: the blue arrows show the process of raising pH as more CO2 is made; the purple arrows indicate the reverse process, lowering pH as more bicarbonate is created.

Antacids, which combat excess stomach acid, are another example of buffers. Many over-the-counter medications work similarly to blood buffers, often with at least one ion (usually carbonate) capable of absorbing hydrogen and moderating pH, bringing relief to those that suffer "heartburn" from stomach acid after eating.

This page titled 25.4A: pH, Buffers, Acids, and Bases is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.4B: Chemical Buffer Systems

Chemical buffers, such as bicarbonate and ammonia, help keep the blood's pH in the narrow range that is compatible with life.

Learning Objectives

- Distinguish between buffer solutions, ventilation, and renal function as buffer systems to control acid–base balance

Key Points

- The body's acid– base balance is tightly regulated to keep the arterial blood pH between 7.38 and 7.42. Buffer solutions keep the pH constant in a wide variety of chemical actions.
- A buffer solution is a mixture of a weak acid and its conjugate base, or a weak base and its conjugate acid.
- The bicarbonate buffering system maintains optimal pH levels and regulates the carbon dioxide concentration that, in turn, shifts any acid–base imbalance.
- Renal physiology controls pH levels through several powerful mechanisms that excrete excess acid or base.

Key Terms

- **bicarbonate**: An alkaline, vital component of the pH buffering system of the human body that maintains acid–base homeostasis.
- **buffer**: A solution used to stabilize the pH (acidity) of a liquid.
- **pH**: In chemistry, a measure of the activity of the hydrogen ion concentration.

EXAMPLES

Anything that adversely affects an individual's bloodstream will have a negative impact on that individual's health since the blood acts as a chemical buffer solution to keep all the body's cells and tissues properly balanced.

Acid–Base Homeostasis

Acid–base homeostasis concerns the proper balance between acids and bases; it is also called body pH. The body is very sensitive to its pH level, so strong mechanisms exist to maintain it. Outside an acceptable range of pH, proteins are denatured and digested, enzymes lose their ability to function, and death may occur.

Buffer Solution

A buffer solution is an aqueous solution of a weak acid and its conjugate base, or a weak base and its conjugate acid. Its pH changes very little when a small amount of strong acid or base is added to it. Buffer solutions are used as a means of keeping pH at a nearly constant value in a wide variety of chemical applications.

Many life forms thrive only in a relatively small pH range, so they utilize a buffer solution to maintain a constant pH. One example of a buffer solution found in nature is blood. The body's acid–base balance is normally tightly regulated, keeping the arterial blood pH between 7.38 and 7.42.

Several buffering agents that reversibly bind hydrogen ions and impede any change in pH exist. Extracellular buffers include bicarbonate and ammonia, whereas proteins and phosphates act as intracellular buffers.

The bicarbonate buffering system is especially key, as carbon dioxide (CO_2) can be shifted through carbonic acid (H_2CO_3) to hydrogen ions and bicarbonate (HCO^{3-}):


H2O+CO2⇌H2CO3⇌H++CO−3H2O+CO2⇌H2CO3⇌H++CO3−

Acid–base imbalances that overcome the buffer system can be compensated in the short term by changing the rate of ventilation. This alters the concentration of carbon dioxide in the blood and shifts the above reaction according to Le Chatelier's principle, which in turn alters the pH.

Renal Physiology

The kidneys are slower to compensate, but renal physiology has several powerful mechanisms to control pH by the excretion of excess acid or base. In response to acidosis, the tubular cells reabsorb more bicarbonate from the tubular fluid, and the collecting duct cells secrete more hydrogen and generate more bicarbonate, and ammoniagenesis leads to an increase of the NH_3 buffer.

In its responses to alkalosis, the kidneys may excrete more bicarbonate by decreasing hydrogen ion secretion from the tubular epithelial cells, and lower the rates of glutamine metabolism and ammonium excretion.



pH range: Buffering agents keep blood pH between 7.38 and 7.42.

This page titled 25.4B: Chemical Buffer Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.4C: Regulation of H+ by the Lungs

Acid–base imbalances in the blood's pH can be altered by changes in breathing to expel more CO_2 and raise pH back to normal.

Learning Objectives

- Describe the regulation of hydrogen ions by the lungs

Key Points

- Hydrogen ions (H+) are carried in the blood along with oxygen and carbon dioxide.
- Sixty percent of the carbon dioxide is carried as dissolved bicarbonate.
- A small amount of carbon dioxide is carried on the hemoglobin as carbaminohemoglobin, which is transported to the lungs for removal.
- Following Le Chatelier's principle, an imbalance in pH is returned to normal by increasing the rate of ventilation in the lungs.
- To compensate for acidemia, more CO_2 is expelled, while the opposite occurs for alkalemia.

Key Terms

- **carbaminohemoglobin**: A compound of hemoglobin and carbon dioxide. It is one of the forms in which carbon dioxide exists in the blood.
- **Le Chatelier's principle**: A principle that states that if a chemical system at equilibrium experiences a change in concentration, temperature, or total pressure, the equilibrium will shift in order to minimize that change.

EXAMPLES


Since maintaining normal pH is vital for life, and since the lungs play a critical role in maintaining normal pH, smokers have yet another reason to quit smoking.

Acid–base imbalance occurs when a significant insult causes the blood pH to shift out of its normal range (7.35 to 7.45). An excess of acid in the blood is called acidemia and an excess of base is called alkalemia.

The process that causes the imbalance is classified based on the etiology of the disturbance (respiratory or metabolic) and the direction of change in pH (acidosis or alkalosis). There are four basic processes and one or a combination may occur at any given time.

1. Metabolic acidosis
2. Respiratory acidosis
3. Metabolic alkalosis
4. Respiratory alkalosis

Blood carries oxygen, carbon dioxide, and hydrogen ions (H+) between tissues and the lungs. The majority of CO_2 transported in the blood is dissolved in plasma (60% is dissolved bicarbonate).



Expiration: When blood pH drops too low, the body compensates by increasing breathing to expel more carbon dioxide.

A smaller fraction is transported in the red blood cells that combine with the globin portion of hemoglobin as carbaminohemoglobin. This is the chemical portion of the red blood cell that aids in the transport of oxygen and nutrients around the body, but, this time, it is carbon dioxide that is transported back to the lung.

Acid–base imbalances that overcome the buffer system can be compensated in the short term by changing the rate of ventilation. This alters the concentration of carbon dioxide in the blood, shifting the above reaction according to Le Chatelier's principle, which in turn alters the pH. The basic reaction governed by this principle is as follows:

H2O+CO2⇌H2CO3⇌H++CO−3H2O+CO2⇌H2CO3⇌H++CO3−

When the blood pH drops too low (acidemia), the body compensates by increasing breathing to expel more CO_2; this shifts the above reaction to the left such that less hydrogen ions are free; thus, the pH will rise back to normal. For alkalemia, the opposite occurs.

This page titled 25.4C: Regulation of H+ by the Lungs is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

25.4D: The Role of the Kidneys in Acid-Base Balance

The kidneys help maintain the acid–base balance by excreting hydrogen ions into the urine and reabsorbing bicarbonate from the urine.

Learning Objectives

- Describe the role of the kidneys to maintain the acid–base balance

Key Points

- The kidneys maintain homeostasis through the excretion of waste products.
- Acidosis causes more bicarbonate to be reabsorbed from the tubular fluid, while the collecting ducts secrete more hydrogen to generate more bicarbonate, and more NH_3 buffer is formed.
- Alkalosis causes the kidney to excrete more bicarbonate as there is a reduced secretion of hydrogen ions and more ammonium is excreted.

Key Terms

- **base**: Any of a class of generally water-soluble compounds, that have a bitter taste, turn red litmus paper blue, and react with acids to form salts.
- **renal**: Pertaining to the kidneys.

EXAMPLES

Urine testing is important because it can detect acid–base imbalances. For instance, uncontrolled diabetes results in highly acidic urine. If the diabetes remains uncontrolled, the kidneys could become over-stressed and malfunction, which could lead to coma or death.

Within the human body, fluids such as blood must be maintained within the narrow range of 7.35 to 7.45, making it slightly alkaline. Outside that range, pH becomes incompatible with life; proteins are denatured and digested, enzymes lose their ability to function, and the body is unable to sustain itself.

To maintain this narrow range of pH the body has a powerful buffering system. Acid–base imbalances that overcome this system are compensated in the short term by changing the rate of ventilation.

Kidneys and Acid–Base Balance

The kidneys have two very important roles in maintaining the acid–base balance:

1. They reabsorb bicarbonate from urine.
2. They excrete hydrogen ions into urine.

The kidneys are slower to compensate than the lungs, but renal physiology has several powerful mechanisms to control pH by the excretion of excess acid or base. The major, homeostatic control point for maintaining a stable pH balance is renal excretion.

Bicarbonate (HCO_{3-}) does not have a transporter, so its reabsorption involves a series of reactions in the tubule lumen and tubular epithelium. In response to acidosis, the tubular cells reabsorb more bicarbonate from the tubular fluid, and the collecting duct cells secrete more hydrogen and generate more bicarbonate, and ammoniagenesis leads to an increase in the formation of the NH_3 buffer.

In response to alkalosis, the kidneys may excrete more bicarbonate by decreasing hydrogen ion secretion from the tubular epithelial cells, and lowering the rates of glutamine metabolism and ammonium excretion.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest/?collection=col11448/latest**. **License**: ***CC BY: Attribution***
- acidic. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/acidic**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- buffer. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/buffer**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alkaline. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/alkaline**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_07.jpg**. **License**: ***CC BY: Attribution***
- Buffering solution. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Buffering_solution**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Acid-base homeostasis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-ba...is%23Mechanism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Acid-base homeostasis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-base_homeostasis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Acid-base homeostasis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-ba...is%23Mechanism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bicarbonate. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bicarbonate**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- buffer. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/buffer**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- pH. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/pH**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_07.jpg**. **License**: ***CC BY: Attribution***
- Power of Hydrogen (pH) chart. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...(pH)_chart.svg**. **License**: ***Public Domain: No Known Copyright***
- Acid-base balance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-base_balance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Acid-base homeostasis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-ba...is%23Mechanism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gas exchange. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gas_exchange**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carbaminohemoglobin. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/carbaminohemoglobin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Le Chatelier's principle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Le_Cha...er's_principle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44392/latest...e_02_02_07.jpg**. **License**: ***CC BY: Attribution***
- Power of Hydrogen (pH) chart. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...(pH)_chart.svg**. **License**: ***Public Domain: No Known Copyright***
- Expiration diagram. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...on_diagram.svg**. **License**: ***Public Domain: No Known Copyright***
- Acid-base balance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acid-base_balance**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Blood. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Blood%2...e_of_pH_values. **License**: *CC BY-SA: Attribution-ShareAlike*
- Acid-base balance. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acid-base_balance. **License**: *CC BY-SA: Attribution-ShareAlike*
- Kidney. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Kidney%...se_homeostasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Renal physiology. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Renal_physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- renal. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/renal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//chemistry/definition/base. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_08.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Water. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44392/latest/Figure_02_02_07.jpg. **License**: *CC BY: Attribution*
- Power of Hydrogen (pH) chart. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...(pH)_chart.svg. **License**: ***Public Domain: No Known Copyright***
- Expiration diagram. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...on_diagram.svg. **License**: ***Public Domain: No Known Copyright***

This page titled 25.4D: The Role of the Kidneys in Acid-Base Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26: THE REPRODUCTIVE SYSTEM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

26.1: Overview of the Reproductive System

26.1A: Overview of the Male and Female Reproductive Systems

26.2: Meiosis

26.2A: Introduction to Meiosis

26.2B: Comparing Meiosis and Mitosis

26.3: The Male Reproductive System

26.3A: Anatomy of the Male Reproductive System

26.3B: Scrotum

26.3C: Testes

26.3D: Penis

26.3E: Testes Ducts

26.3F: Accessory Sex Glands

26.4: Physiology of the Male Reproductive System

26.4A: Male Sexual Response

26.4B: Sperm

26.4C: Spermatogenesis

26.4D: Semen

26.4E: Hormonal Regulation of the Male Reproductive System

26.5: The Female Reproductive System

26.5A: Anatomy of the Female Reproductive System

26.5B: Ovaries

26.5C: Uterus

26.5D: Female Duct System

26.5E: Vagina

26.5F: Vulva

26.5G: Perineum

26.5H: Mammary Glands

26.6: Physiology of the Female Reproductive System

26.6A: Oogenesis

26.6B: Ovarian Cycle
26.6C: Uterine (Menstrual) Cycle
26.6D: Hormonal Regulation of the Female Reproductive Cycle
26.6E: Extrauterine Effects of Estrogens and Progesterone
26.6F: Female Sexual Response

This page titled 26: The Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.1: OVERVIEW OF THE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

26.1A: Overview of the Male and Female Reproductive Systems

This page titled 26.1: Overview of the Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.1A: Overview of the Male and Female Reproductive Systems

The human reproductive system functions to produce human offspring, with the male providing sperm and the female providing the ovum.

Learning Objectives

- Summarize the reproductive systems of men and women

Key Points

- The male reproductive system consists of external organs. The testes in the scrotum produce the male gamete, sperm, which is ejaculated in seminal fluid by the penis.
- The female reproductive system primarily consists of internal organs. The female gamete, ovum, is produced in the ovaries and is released monthly to travel to the uterus via the Fallopian tubes.
- Fertilization can occur if the penis is inserted through the vulva into the vagina and sperm is ejaculated towards the cervix. If an ovum is currently in the uterus, it can then be fertilized by sperm that manage to enter the cervix.
- Once fertilized, an ovum becomes a zygote and if all goes well, develops into a fetus in the uterus.
- Natural birth occurs when the fetus is pushed from the vagina after nine months in the uterus.

Key Terms

- **fallopian tubes**: The Fallopian tubes, also known as oviducts, uterine tubes, and salpinges (singular salpinx) are two very fine tubes lined with ciliated epithelia leading from the ovaries of female mammals into the uterus, via the utero-tubal junction.
- **penis**: The male sexual organ for copulation and urination; the tubular portion of the male genitalia (excluding the scrotum).
- **vagina**: A fibromuscular tubular tract which is the female sex organ and has two main functions: sexual intercourse and childbirth.

EXAMPLES

While the ultimate purpose of the human reproductive system is to produce offspring, the proximate purpose is to produce pleasure and induce bonding. This can be seen in our closest relatives, the bonobo chimpanzees, who have sex for a wide variety of reasons including pleasure, bonding, and alleviating tension in addition to producing offspring.


The reproductive system or genital system is a set of organs within an organism that work together to produce offspring. Many non-living substances, such as fluids, hormones, and pheromones, are important accessories to the reproductive system. Unlike most organ systems, the sexes of differentiated species often have significant differences. These differences allow for a combination of genetic material between two individuals and thus the possibility of greater genetic fitness of the offspring.

The Reproductive Process

Human reproduction takes place as internal fertilization by sexual intercourse. During this process, the erect penis of the male is inserted into the female's vagina until the male ejaculates semen, which contains sperm, into the vagina. The sperm travels through the vagina and cervix into the uterus for potential fertilization of an ovum. Upon successful fertilization and implantation, gestation of the fetus occurs within the female's uterus for approximately nine months (pregnancy). Gestation ends with labor resulting in birth. In labor, the uterine muscles contract, the cervix dilates, and the baby passes out through the vagina. Human babies and children are nearly helpless and require high levels of parental care for many years. One important type of parental care is the use of the mammary glands in the female breasts to nurse the baby.

The Male Reproductive System

The human male reproductive system is a series of organs located outside of the body and around the pelvic region. The primary direct function of the male reproductive system is to provide the male gamete or spermatozoa for fertilization of the ovum. The major reproductive organs of the male can be grouped into three categories. The first category is sperm production and storage. Production takes place in the testes, housed in the temperature-regulating scrotum. Immature sperm then travel to the epididymis for development and storage. The second category, the ejaculatory fluid-producing glands, includes the seminal vesicles, prostate, and vas deferens. The final category, used for copulation and deposition of the spermatozoa (sperm) within the female, includes the penis, urethra, vas deferens, and Cowper's gland.




The Human Male Reproductive System: Cross-sectional diagram of the male reproductive organs.

Only our species has a distinctive mushroom-capped glans, which is connected to the shaft of the penis by a thin tissue of frenulum (the delicate tab of skin just beneath the urethra). One of the most significant features of the human penis is the coronal ridge underneath the gland around the circumference of the shaft. Magnetic imaging studies of heterosexual couples having sex reveal that during coitus, the typical penis expands to fill the vaginal tract, and with full penetration can even reach the woman's cervix and lift her uterus. This combined with the fact that human ejaculate is expelled with great force and considerable distance (up to two feet if not contained), suggests that men are designed to release sperm into the uppermost portion of the vagina. This may be an evolutionary adaptation to expel the semen left by other males while at the same time increasing the possibility of fertilization with the current male's semen.

The Female Reproductive System

The human female reproductive system is a series of organs primarily located inside the body and around the pelvic region. It contains three main parts: the vagina, which leads from the vulva, the vaginal opening, to the uterus; the uterus, which holds the developing fetus; and the ovaries, which produce the female's ova. The breasts are also a reproductive organ during parenting, but are usually not classified as part of the female reproductive system. The vagina meets the outside at the vulva, which also includes the labia, clitoris, and urethra. During intercourse, this area is lubricated by mucus secreted by the Bartholin's glands. The vagina is attached to the uterus through the cervix, while the uterus is attached to the ovaries via the Fallopian tubes. At certain intervals, approximately every 28 days, the ovaries release an ovum that passes through the Fallopian tube into the uterus. If the ova is fertilized by sperm, it attaches to the endometrium and the fetus develops. In months when fertilization does not occur, the lining of the uterus, called the endometrium, and unfertilized ova are shed each cycle through a process known as menstruation.



The Human Female Reproductive System: The female reproductive system is largely internal.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Reproductive system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Reproductive_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- penis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/penis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vagina. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vagina**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fallopian tubes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fallopian%20tubes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Female Reproductive System. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...em_lateral.png**. **License**: ***Public Domain: No Known Copyright***
- Male Reproductive System. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...le_anatomy.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 26.1A: Overview of the Male and Female Reproductive Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.2: MEIOSIS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

26.2A: Introduction to Meiosis

26.2B: Comparing Meiosis and Mitosis

This page titled 26.2: Meiosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.2A: Introduction to Meiosis

Meiosis is the nuclear division of diploid cells into haploid cells, which is a necessary step in sexual reproduction.

Learning Objectives

- Describe the importance of meiosis in sexual reproduction

Key Points

- Sexual reproduction is the production of haploid cells and the fusion of two of those cells to form a diploid cell.
- Before sexual reproduction can occur, the number of chromosomes in a diploid cell must decrease by half.
- Meiosis produces cells with half the number of chromosomes as the original cell.
- Haploid cells used in sexual reproduction, gametes, are formed during meiosis, which consists of one round of chromosome replication and two rounds of nuclear division.
- Meiosis I is the first round of meiotic division, while meiosis II is the second round.

Key Terms

- **haploid**: of a cell having a single set of unpaired chromosomes
- **gamete**: a reproductive cell, male (sperm) or female (egg), that has only half the usual number of chromosomes
- **diploid**: of a cell, having a pair of each type of chromosome, one of the pair being derived from the ovum and the other from the spermatozoon

Introduction: Meiosis and Sexual Reproduction

The ability to reproduce *in kind* is a basic characteristic of all living things. In kind means that the offspring of any organism closely resemble their parent or parents. Sexual reproduction requires fertilization: the union of two cells from two individual organisms. Haploid cells contain one set of chromosomes. Cells containing two sets of chromosomes are called diploid. The number of sets of chromosomes in a cell is called its ploidy level. If the reproductive cycle is to continue, then the diploid cell must somehow reduce its number of chromosome sets before fertilization can occur again or there will be a continual doubling in the number of chromosome sets in every generation. Therefore, sexual reproduction includes a nuclear division that reduces the number of chromosome sets.



(a) (b) (c)

Offspring Closely Resemble Their Parents: *In kind* means that the offspring of any organism closely resemble their parent or

parents. The hippopotamus gives birth to hippopotamus calves (a). Joshua trees produce seeds from which Joshua tree seedlings emerge (b). Adult flamingos lay eggs that hatch into flamingo chicks (c).

Sexual reproduction is the production of haploid cells (gametes) and the fusion (fertilization) of two gametes to form a single, unique diploid cell called a zygote. All animals and most plants produce these gametes, or eggs and sperm. In most plants and animals, through tens of rounds of mitotic cell division, this diploid cell will develop into an adult organism.

Haploid cells that are part of the sexual reproductive cycle are produced by a type of cell division called meiosis. Meiosis employs many of the same mechanisms as mitosis. However, the starting nucleus is always diploid and the nuclei that result at the end of a meiotic cell division are haploid, so the resulting cells have half the chromosomes as the original. To achieve this reduction in chromosomes, meiosis consists of one round of chromosome duplication and two rounds of nuclear division. Because the events that occur during each of the division stages are analogous to the events of mitosis, the same stage names are assigned. However, because there are two rounds of division, the major process and the stages are designated with a “I” or a “II.” Thus, meiosis I is the first round of meiotic division and consists of prophase I, prometaphase I, and so on. Meiosis II, the second round of meiotic division, includes prophase II, prometaphase II, and so on.

This page titled 26.2A: Introduction to Meiosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.2B: Comparing Meiosis and Mitosis

Mitosis and meiosis share some similarities, but also some differences, most of which are observed during meiosis I.

Learning Objectives

- Compare and contrast mitosis and meiosis

Key Points


- For the most part, in mitosis, diploid cells are partitioned into two new diploid cells, while in meiosis, diploid cells are partitioned into four new haploid cells.
- In mitosis, the daughter cells have the same number of chromosomes as the parent cell, while in meiosis, the daughter cells have half the number of chromosomes as the parent.
- The daughter cells produced by mitosis are identical, whereas the daughter cells produced by meiosis are different because crossing over has occurred.
- The events that occur in meiosis but not mitosis include homologous chromosomes pairing up, crossing over, and lining up along the metaphase plate in tetrads.
- Meiosis II and mitosis are not reduction division like meiosis I because the number of chromosomes remains the same; therefore, meiosis II is referred to as equatorial division.
- When the homologous chromosomes separate and move to opposite poles during meiosis I, the ploidy level is reduced from two to one, which is referred to as a reduction division.

Key Terms

- **reduction division**: the first of the two divisions of meiosis, a type of cell division
- **ploidy**: the number of homologous sets of chromosomes in a cell
- **equatorial division**: a process of nuclear division in which each chromosome divides equally such that the number of chromosomes remains the same from parent to daughter cells

Comparing Meiosis and Mitosis

Mitosis and meiosis are both forms of division of the nucleus in eukaryotic cells. They share some similarities, but also exhibit distinct differences that lead to very different outcomes. The purpose of mitosis is cell regeneration, growth, and asexual reproduction,while the purpose of meiosis is the production of gametes for sexual reproduction. Mitosis is a single nuclear division that results in two nuclei that are usually partitioned into two new daughter cells. The nuclei resulting from a mitotic division are genetically identical to the original nucleus. They have the same number of sets of chromosomes, one set in the case of haploid cells and two sets in the case of diploid cells. In most plants and all animal species, it is typically diploid cells that undergo mitosis to form new diploid cells. In contrast, meiosis consists of two nuclear divisions resulting in four nuclei that are usually partitioned into four new haploid daughter cells. The nuclei resulting from meiosis are not genetically identical and they contain one chromosome set only. This is half the number of chromosome sets in the original cell, which is diploid.



						OUTCOME
PROCESS	DNA synthesis	Synapsis of homologous chromosomes	Crossover	Homologous chromosomes line up at metaphase plate	Sister chromatids line up at metaphase plate	Number and genetic composition of daughter cells
MEIOSIS	Occurs in S phase of interphase	During prophase I	During prophase I	During metaphase I	During metaphase II	Four haploid cells at the end of meiosis II
MITOSIS	Occurs in S phase of interphase	Does not occur in mitosis	Does not occur in mitosis	Does not occur in mitosis	During metaphase	Two diploid cells at the end of mitosis

Comparing Meiosis and Mitosis: Meiosis and mitosis are both preceded by one round of DNA replication; however, meiosis includes two nuclear divisions. The four daughter cells resulting from meiosis are haploid and genetically distinct. The daughter cells resulting from mitosis are diploid and identical to the parent cell.

The main differences between mitosis and meiosis occur in meiosis I. In meiosis I, the homologous chromosome pairs become associated with each other and are bound together with the synaptonemal complex. Chiasmata develop and crossover occurs between homologous chromosomes, which then line up along the metaphase plate in tetrads with kinetochore fibers from opposite spindle poles attached to each kinetochore of a homolog in a tetrad. All of these events occur only in meiosis I.

When the tetrad is broken up and the homologous chromosomes move to opposite poles, the ploidy level is reduced from two to one. For this reason, meiosis I is referred to as a reduction division. There is no such reduction in ploidy level during mitosis.

Meiosis II is much more similar to a mitotic division. In this case, the duplicated chromosomes (only one set, as the homologous pairs have now been separated into two different cells) line up on the metaphase plate with divided kinetochores attached to kinetochore fibers from opposite poles. During anaphase II and mitotic anaphase, the kinetochores divide and sister chromatids, now referred to as chromosomes, are pulled to opposite poles. The two daughter cells of mitosis, however, are identical, unlike the daughter cells produced by meiosis. They are different because there has been at least one crossover per chromosome. Meiosis II is not a reduction division because, although there are fewer copies of the genome in the resulting cells, there is still one set of chromosomes, as there was at the end of meiosis I. Meiosis II is, therefore, referred to as equatorial division.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44468/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, The Process of Meiosis. October 28, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44469/latest/**. **License**: ***CC BY: Attribution***
- haploid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/haploid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diploid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diploid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gamete. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gamete**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/**

m44468/latest...7_00_02abc.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44469/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...orial-division. **License**: *CC BY-SA: Attribution-ShareAlike*
- ploidy. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ploidy. **License**: *CC BY-SA: Attribution-ShareAlike*
- reduction division. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/reduction_division. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44468/latest...7_00_02abc.jpg. **License**: *CC BY: Attribution*
- OpenStax College, The Process of Meiosis. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44469/latest...e_11_01_06.jpg. **License**: *CC BY: Attribution*

This page titled 26.2B: Comparing Meiosis and Mitosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3: THE MALE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

26.3A: Anatomy of the Male Reproductive System

26.3B: Scrotum

26.3C: Testes

26.3D: Penis

26.3E: Testes Ducts

26.3F: Accessory Sex Glands

This page titled 26.3: The Male Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3A: Anatomy of the Male Reproductive System

The male reproductive system includes external (penis, scrotum, epididymus, and testes) and internal (accessory) organs.

Learning Objectives

- Distinguish among the parts and functions of the male reproductive system

Key Points

- The functions of the male reproductive system include producing and transporting sperm, ejaculating sperm into the female reproductive tract, and producing and secreting male hormones.
- Most of the male reproductive system is located outside of the body. These external structures are the penis, scrotum, epididymis, and testes.
- The internal organs of the male reproductive system are called accessory organs. They include the vas deferens, seminal vesicles, prostate gland, and bulbourethral glands.

Key Terms


- **semen**: Contains spermatozoa, proteolytic and other enzymes, and fructose that promotes spermatozoa survival. It also provides a medium for sperm motility.
- **spermatogenesis**: The process of sperm production within the seminiferous tubules in the testes.
- **testosterone**: Steroid hormone produced primarily in the male testes and responsible for the development of male secondary sex characteristics.

The organs of the male reproductive system are specialized for three primary functions:

1. To produce, maintain, transport, and nourish sperm (the male reproductive cells), and protective fluid (semen).
2. To discharge sperm within the female reproductive tract.
3. To produce and secrete male sex hormones.

External Male Sex Organs

Most of the male reproductive system is located outside of the man’s body. These external structures are the penis, scrotum, epididymis, and testes.



Male Reproductive System: Lateral view of male reproductive system with organs labeled.

The penis is the male organ for sexual intercourse and urination. Semen and urine leave the penis through the urethra. The scrotum is a loose, pouch-like sack of skin that hangs behind the penis, containing the testes.

The scrotum has a protective function, including the maintenance of optimal temperatures for sperm survival and function. For sperm development, the testes must maintain a temperature slightly cooler than normal body temperature. Special muscles in the wall of the scrotum contract and relax in order to move the testes near the body.

The epididymus is located at the back of the testis and connects it to the vas deferens. Its function is to store and carry sperm. The testis is the location for testosterone production. The coiled collection of tubes within the testes are the seminiferous tubules. Within these tubules, spermatogenesis takes place.

Accessory Sex Organs

The internal organs of the male reproductive system are called accessory organs. They include the vas deferens, seminal vesicles, prostate gland, and bulbourethral (Cowper's) glands.

- Vas deferens: Transports mature sperm to the urethra in preparation for ejaculation.
- Seminal vesicles: Sac-like pouches that attach to the vas deferens near the base of the bladder. The vesicles produce molecules such as fructose that serve as energy sources for sperm. The seminal vesicle fluid makes up most of the volume of a man's ejaculate.
- Prostate gland: A walnut-sized structure located below the urinary bladder in front of the rectum. It contributes additional fluid to the ejaculate that serves as nourishment for sperm.
- Bulbourethral (Cowper's) glands: Pea-sized structures located on the sides of the urethra just below the prostate gland. These glands produce a clear, slippery fluid that empties directly into the urethra. Fluid produced by these glands lubricates the urethra and neutralizes acidity associated with residual urine.

This page titled 26.3A: Anatomy of the Male Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3B: Scrotum

The purpose of the scrotum is to provide the testes with a chamber of appropriate temperature for optimal sperm production.

Learning Objectives

- Describe the functions of the scrotum

Key Points

- Moving the testes away from the abdomen and increasing the exposed surface area allow a faster dispersion of excess heat.
- If testes were in the abdomen, constant pressure from abdominal muscles would possibly empty the testes and epididymis before sperm were sufficiently mature for fertillization.
- The function of the scrotum appears to be to keep the temperature of the testes slightly lower than that of the rest of the body.

Key Terms

- **scrotum**: The bag of skin and muscle that contains the testes in mammals.
- **testosterone**: A steroid hormone that plays a key role in male reproductive development including the promotion of secondary sexual characteristics.
- **epididymis**: A narrow, tightly-coiled tube where sperm are stored during maturation. It connects the efferent ducts from the rear of each testicle to its vas deferens.

Anatomical Considerations


The scrotum is a dual-chambered suspended sack of skin and smooth muscle that contains the testes, and is homologous to the labia majora in females. It is an extension of the perineum, and is located between the penis and anus. In humans and some other mammals, increased testosterone secretion during puberty causes the darkening of the skin and development of pubic hair on the scrotum. The left testis is usually lower than the right, which may function to avoid compression in the event of impact. This asymmetry may also allow more effective cooling of the testes.

Function

The function of the scrotum appears to be to keep the temperature of the testes slightly lower than that of the rest of the body. For human beings, the temperature should be one or two degrees Celsius below body temperature (around 35 degrees Celsius or 95 degrees Fahrenheit); higher temperatures may be damaging to sperm count.

The temperature is controlled by scrotal movement of the testes away or towards the body depending on the environmental temperatures. Moving the testes away from the abdomen and increasing the exposed surface area allow a faster dispersion of excess heat. This is done by means of contraction and relaxation of the cremaster muscle and the dartos fascia in the scrotum.

However, temperature regulation may not be the only function of the scrotum. It has been suggested that if testes were situated within the abdominal cavity, they would be subjected to the regular changes in abdominal pressure that are exerted by the abdominal muscles, resulting in the more rapid emptying of the testes and epididymis of sperm before the spermatozoa were matured sufficiently for fertilization. Some mammals (elephants and marine mammals, for example) do keep their testes within the abdomen where there may be mechanisms to prevent this inadvertent emptying.



The scrotum: Image of the external, muscle, and deep tissue views of the scrotum.

This page titled 26.3B: Scrotum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3C: Testes

The testis is homologous to the ovary in that it produces the male gamete (sperm) while the ovary produces the female gamete (egg).

Learning Objectives


- Describe the function of testes and the development of sperm

Key Points

- The testes produce the hormones testosterone and other androgens.
- Sperm are produced within seminiferous tubules.
- Leydig cells produce and secrete male hormones.
- Sertoli cells help in the process of spermatogenesis.

Key Terms

- **Leydig cells**: Also known as interstitial cells of Leydig, these are found adjacent to the seminiferous tubules in the testicle and produce testosterone in the presence of luteinizing hormone.
- **follicle-stimulating hormone**: Stimulates the growth and recruitment of immature ovarian follicles in females. In males, it is critical for spermatogenesis as it stimulates primary spermatocytes to form secondary spermatocytes.
- **sertoli cells**: Part of the seminiferous tubule that helps in the process of spermatogenesis.
- **Luteinizing hormone**: A hormone produced by gonadotropic cells of the anterior pituitary gland. It triggers ovulation and development of the corpus luteum in females and stimulates Leydig cell production of testosterone in males.
- **testes**: Also referred to as testicles, the male gonads in animals.



Testicle: A diagram of the major components of an adult human testis, including the following numbered items: 1. Tunica albuginea, 2. Septula testis, 3. Lobulus testis, 4. Mediastinum testis, 5. Tubuli seminiferi contorti, 6. Tubuli seminiferi recti, 7. Rete testis, 8. Ductuli efferentes testis, 9a. Head of epididymis, 9b. Body of epididymis, 9.c Tail of epididymis,10. Vas deferens, 11a. Tunica vaginalis (parietal lamina), 11b. Tunica vaginalis (visceral lamina), and 12. Cavity of tunica vaginalis.


The testis is the male gonad in animals. Like the ovaries to which they are homologous, testes are components of both

the reproductive system and the endocrine system. Thee testes produce sperm (spermatogenesis) and androgens, primarily testosterone. Both functions of the testis are influenced by gonadotropic hormones produced by the anterior pituitary gland. Luteinizing hormone results in testosterone release. The presence of both testosterone and follicle-stimulating hormone (FSH) is needed to support spermatogenesis.

Almost all healthy male vertebrates have two testes. In mammals, the testes are often contained within an extension of the abdomen called the scrotum. In mammals with external testes, it is most common for one testicle to hang lower than the other. While the size of the testis varies, it is estimated that 21.9% of men have one higher-positioned testis, while 27.3% of men have reported equally-positioned testicles.

The tough membranous shell called the tunica albuginea contains very fine coiled tubes called seminiferous tubules. These are lined with a layer of germ cells that develop into sperm cells (also known as spermatozoa or male gametes) from puberty into old age. The developing sperm travels through the seminiferous tubules to the rete testis located in the mediastinum testis, to the efferent ducts, and then to the epididymis where newly-created sperm cells mature. The sperm moves into the vas deferens and is eventually expelled through the urethra, via the urethral orifice through muscular contractions.

Leydig cells located between seminiferous tubules produce and secrete testosterone and other androgens important for sexual development and puberty, including secondary sexual characteristics such as facial hair and sexual behavior. They also support libido, spermatogenesis, and erectile function. In addition, testosterone controls testicular volume. The sertoli cells are the testes' somatic cells, necessary for testis development and spermatogenesis.



Inside the Human Testes: Diagram illustrates the scrotum with a portion of the covering removed to display the testis.

This page titled 26.3C: Testes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3D: Penis

In human males, the penis serves as both a reproductive organ and as a urinal duct.

Learning Objectives

- Describe the structure and function of the penis


Key Points

- The major structure of the penis is formed by columns of corpus cavernosum and spongiosum tissue.
- The head of the penis, called the glans, contains the opening for the urethral duct, the passage for urine and seminal fluid.
- The penis and clitoris (found in women) are homologous organs.
- Blood engorgement of penile tissue causes the penis to become erect, facilitating sexual intercourse.
- Ejaculation is the release of sperm, which propels the sperm into the vaginal canal when it occurs during intercourse.

Key Terms

- **glans**: The vascular body which forms the apex of the penis.
- **penis**: The male sexual organ for copulation and urination; the tubular portion of the male genitalia (excluding the scrotum).
- **intromittent organ**: A term for a male external organ that delivers sperm during copulation.
- **clitoral glans**: Highly innervated part of the clitoris that exists at the tip of the clitoral body as a fibrovascular cap.
- **ejaculation**: The forcible ejection of semen from the mammalian urethra, a reflex in response to sexual stimulation.

The penis is an intromittent organ of male animals with reproductive and urinary functions. Unlike many other species, the human penis has no baculum, or erectile bone. Instead, it relies entirely on engorgement with blood to achieve erection. The human penis cannot be withdrawn into the groin, and
is larger than that of any other primate, particularly in regards to proportion to body mass.



Human Penis: This is a diagram of a human penis with its parts labeled.

Parts of the Penis


The human penis is made up of three columns of tissue: two corpora cavernosa that lie next to each other on the dorsal side and a corpus spongiosum that lies between the corpora cavernosa on the ventral side. The glans penis is the bulbous end of the penis formed by the corpus spongiosum. It supports the foreskin (prepuce) that retracts to expose the glans. The area on the underside of the penis, where the foreskin is attached, is called the frenum (or frenulum). The rounded base of the glans is called the corona. The perineal raphe is the noticeable line along the underside of the penis.

The urethra is the last part of the urinary tract and traverses the corpus spongiosum. The urethral opening is called the meatus and lies on the tip of the glans penis. It serves as a passage both for urine and semen. Sperm are produced in the testes and stored in the attached epididymis. Sperm are propelled through the vas deferens during ejaculation. Fluids are added by the seminal vesicles before the vas deferens carries the sperm to the ejaculatory ducts, which join the urethra inside the prostate gland. The prostate as well as the bulbourethral glands add further secretions, then the semen is expelled through the penis. The raphe is the noticeable ridge between the halves of the penis. It is located on the ventral aspect of the penis and runs from the meatus and across the scrotum to the perineum (area between scrotum and anus).

Homology to the Clitoris

The various parts of the male penis are homologous to parts of the female clitoris:

- Glans of the penis: homologous to the clitoral glans
- Corpora cavernosa: homologous to the body of the clitoris
- Corpus spongiosum: homologous to vestibular bulbs beneath the labia minora
- Scrotum: homologous to the labia minora and labia majora
- Foreskin: homologous to the clitoral hood



Penile and Clitoral Structure: This diagram compares the structure of the penis to the clitoris.

The Penis As a Sexual Organ

An erection is the stiffening and rising of the penis that facilitates sexual arousal, though it can also happen in nonsexual situations. The primary physiological mechanism that brings about an erection is the autonomic dilation of arteries supplying blood to the penis. This allows more blood to fill the three spongy erectile tissue chambers in the penis, causing it to lengthen and stiffen.

The engorged erectile tissue presses against and constricts the veins that carry blood away from the penis. More blood enters

than leaves the penis until an equilibrium is reached where an equal volume of blood flows into the dilated arteries and out of the constricted veins; a constant erectile size is achieved at this equilibrium. Although many erect penises point upwards, it is common and normal for the erect penis to point nearly vertically upwards, nearly vertically downwards, or even horizontally forward, depending on the tension of the suspensory ligament that holds it in position.

Ejaculation is the ejection of semen from the penis and is usually accompanied by orgasm. A series of muscular contractions delivers semen, containing sperm cells or spermatozoa, from the penis. It is usually the result of sexual stimulation, including prostate stimulation. Ejaculation may occur spontaneously during sleep (known as a nocturnal emission or “wet dream”).

Ejaculation has two phases: emission and ejaculation proper. The emission phase of the ejaculatory reflex is under control of the sympathetic nervous system, while the ejaculatory phase is under the control of a spinal reflex at the level of spinal nerves S2 to S4 via the pudendal nerve. A refractory period succeeds the ejaculation, and sexual stimulation precedes it. Anejaculation is the condition of being unable to ejaculate.

This page titled 26.3D: Penis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3E: Testes Ducts

Testes ducts, which include the seminiferous tubules and vas deferens, are involved in the creation or transportation of sperm.

Learning Objectives


- Differentiate among the ducts of the testes

Key Points

- Creation of sperm occurs in the seminiferous tubules of the testes.
- The efferent ducts maintain proper fluid concentration in the testes and propel sperm to the epididymus.
- The vas deferens carry the sperm from the epididymus to the urethra.
- A vasectomy cuts the vas deferens to prevent sperm from entering the urethra and being ejaculated.

Key Terms

- **vasectomy**: The surgical removal of all or part of the vas deferens, usually as a means of male sterilization.
- **peristalsis**: A radially symmetrical contraction and relaxation of muscles that propagates in a anterograde (forward) wave down a tube.
- **vas deferens**: The duct in the testes that carries semen from the epididymis to the ejaculatory duct.
- **rete testis**: An anastomosing network of tubules located in the hilum of the testicle (mediastinum testis) that carries sperm from the seminiferous tubules to the efferent ducts.



Inside the Human Testes: Diagram of cross-section of testes: 1: Testicular septa 2: Convoluted seminiferous tubules 3: Testicular lobules 4: Straight seminiferous tubules 5: Efferent ductules 6: Rete testis.


Seminiferous tubules, located in the testes, are where meiosis occurs and the gametes (spermatozoa) are created. The seminiferous tubules are formed from primitive sex cords from the gonadal ridge. The epithelium of the tubule consists of tall, columnar cells called Sertoli cells. Between the Sertoli cells are spermatogenic cells, which differentiate through meiosis to become sperm cells. There are two types of seminiferous tubules: convoluted, located toward the lateral side, and straight, as the tubule comes medially to form ducts that will exit the testis.

Efferent Ducts

The efferent ducts connect the rete testis with the initial section of the epididymis. There are two basic types of efferent ductule structure.

1. Multiple entries into the epididymis: This type is seen in most large mammals. In humans and other large mammals, there are approximately 15–20 efferent ducts which occupy nearly one-third of the head of the epididymis.
2. Single entry: This type is seen in most small animals such as rodents. This is characterized by three to six ductules that merge into a single small ductule prior to entering the epididymis.

The ductuli are unilaminar and composed of columnar ciliated and nonciliated (absorptive) cells. The ciliated cells stir the luminal fluids, which may help ensure homogeneous absorption of water from the fluid produced by the testis. This results in an increase in the concentration of luminal sperm. The epithelium is surrounded by a band of smooth muscle that helps to propel the sperm toward the epididymis.



Histiological Representation of Seminferious Tubules: Seminiferous tubule in cross-section (large tubular structure: center of image) with sperm (black, tiny, ovoid bodies furthest from the outer edge of the tubular structure).

The ductus (vas) deferens, also called the sperm duct, extend from the epididymis on each side of the scrotum into the abdominal cavity through the inguinal canal, an opening in the abdominal wall for the spermatic cord. The spermatic cord is a connective tissue sheath that contains the vas deferens, testicular blood vessels, and nerves.

The smooth muscle layer of the vas deferens contracts in waves of peristalsis during ejaculation. Two ducts connect the left and right epididymis to the ejaculatory ducts in order to move sperm. In humans, each tube is about 30 centimeters (a foot) long, 3 to 5 mm in diameter, and surrounded by smooth muscle. The sperm are transferred from the vas deferens into the urethra, collecting secretions from the male accessory sex glands such as the seminal vesicles, prostate gland, and bulbourethral glands, which provide the bulk of semen.

The procedure of deferentectomy, also known as a vasectomy, is a method of contraception in which the vas deferens are permanently cut, though in some cases it can be reversed. A modern procedure that does not include cutting the ducts involves injecting an obstructive material into the ductus to block the flow of sperm. Research in male contraception has focused primarily on the vas deferens with the use of the intra-vas device and reversible inhibition of sperm under guidance.

This page titled 26.3E: Testes Ducts is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.3F: Accessory Sex Glands

The accessory sex glands produce seminal fluid and clean and lubricate the urethra.

Learning Objectives

- Describe the functions of male accessory sex glands

Key Points

- Seminal glands contain seminal vesicles that produce 50–70% of the seminal fluid.
- The excretory duct of the seminal gland opens into the vas deferens as it enters the prostate gland.
- Sperm are not in contact with the seminal fluid produced by the seminal vesicles, possibly to block the progress of sperm from other males.
- The prostate secretes 20–30% of the seminal fluid which carries the sperm in the ejaculate.
- The prostate surrounds the urethra below the bladder and can be felt in a rectal exam.
- Bulbourethral glands produce a pre-ejaculate secretion which lubricates and flushes out the urethra in preparation for the sperm.

Key Terms

- **bulbourethral gland**: An exocrine gland which secretes a clear fluid upon sexual arousal as pre-ejaculate (or Cowper's fluid).
- **Lipofuscin**: Yellow-brown pigment granules composed of lipid-containing residues of lysosomal digestion.
- **prostate**: A compound tubuloalveolar exocrine gland of the male reproductive system in most mammals.
- **seminal gland**: A pair of simple tubular glands located within the pelvis that secrete fluid that partly composes semen.

The accessory sex glands, including the seminal, prostate glands, and bulbourethral glands, produce seminal fluid and clean and lubricate the urethra.

Seminal Gland Anatomy

Each seminal gland forms as an outward growth of the wall of ampulla of each vas deferens. They are curled and folded within the gland and can spread out to approximately 5 cm, but the unfolded length is approximately 10 cm. The excretory duct of the seminal gland opens into the vas deferens as it enters the prostate gland.

Seminal Gland Physiology and Function


The seminal vesicles secrete a significant proportion of the fluid that ultimately becomes semen. Lipofuscin granules from dead epithelial cells give the secretion its yellowish color. About 50–70% of seminal fluid in humans originates from the seminal vesicles, but is not expelled in the first ejaculate fractions which are dominated by spermatozoa and zinc-rich prostatic fluid. Seminal vesicle fluid is alkaline, resulting in human semen with a mildly alkaline pH. This helps neutralize the acidity of the vaginal tract, prolonging the lifespan of sperm. Acidic ejaculate (pH <7.2) may be associated with ejaculatory duct obstruction. The vesicle produces a substance that causes the semen to become sticky after ejaculation, thought to help keep the semen near the cervix.

The thick seminal vesicle secretions contain proteins (including enzymes), mucus, fructose, vitamin C, flavins, phosphorylcholine, and prostaglandins. The high fructose concentrations provide nutrient energy for the spermatozoa when stored in semen in the laboratory. Spermatozoa ejaculated into the vagina are not likely to have contact with seminal vesicular fluid, but transfer directly from the prostatic fluid into the cervical mucus as the first step on their travel through the female reproductive system.

Seminal vesicle fluid is expelled under sympathetic contraction of the muscularis muscle coat. In vitro studies have shown that

sperm expelled together with seminal vesicular fluid show poor motility and survival and less-protected chromatin. Thus, the exact physiological importance of seminal vesicular fluid is unclear. It may be a developmental rest, such as in some rodents where the last part of the ejaculate forms a spermicidal plug to reduce the chances for sperm from a later-arriving male to proceed to the oocyte.

Prostate Gland Anatomy



Prostate: Prostate with seminal vesicles and seminal ducts, viewed from the front and above, including the urethra, seminal vesicle, vas deferens, ampulla, ejaculatory duct, and isthmus.


The prostate surrounds the urethra just below the urinary bladder and can be felt during a rectal exam. It is the only exocrine organ located in the midline in humans and similar animals. Within the prostate, the urethra coming from the bladder is called the prostatic urethra and merges with the two ejaculatory ducts. The prostate is sheathed in the muscles of the pelvic floor, which contract during the ejaculatory process.

Prostate Gland Physiology and Function

The prostate secretes a slightly acidic fluid, milky or white in appearance, that usually constitutes 20–30% of the volume of the semen along with spermatozoa and seminal vesicle fluid. The prostatic fluid is expelled in the first ejaculate fractions, together with most of the spermatozoa. In comparison with the few spermatozoa expelled in seminal vesicular fluid, those expelled in prostatic fluid have better motility, longer survival, and better protection of the genetic material. The prostate also contains some smooth muscles that help expel semen during ejaculation.

To work properly, the prostate needs male hormones (e.g., testosterone), which are produced mainly by the testes. Some male hormones are produced in small amounts by the adrenal glands. However, dihydrotestosterone regulates the prostate. A healthy human prostate is slightly larger than a walnut in adult males, with a weight ranging between 7 and 16 grams.

Bulbourethral Glands



Bulbourethral Gland: Image shows internal view of penis and male sexual anatomy. The bulbourethral gland is labeled at center left.

Bulbourethral glands are located posterior and lateral to the membranous portion of the urethra at the base of the penis, between the two layers of the fascia of the urogenital diaphragm in the deep perineal pouch. They are enclosed by transverse fibers of the sphincter urethrae membranacea muscle. The bulbourethral glands are compound tubulo-alveolar glands, each approximately the size of a pea. They are composed of several lobules held together by a fibrous covering. Each lobule opens into a duct that joins with the ducts of other lobules to form a single excretory duct. This duct is approximately 2.5 cm long and opens into the urethra at the base of the penis. The glands gradually diminish in size with advancing age.

During sexual arousal, each gland produces a clear, salty, viscous secretion known as pre-ejaculate. This fluid helps to lubricate the urethra for spermatozoa to pass through, neutralizes traces of acidic urine in the urethra, and helps flush out any residual urine or foreign matter. It is possible for this fluid to pick up sperm remaining in the urethral bulb from previous ejaculations and carry them out prior to the next ejaculation.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Epididymis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epididymis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Semen. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Semen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Male Reproductive System. **Provided by**: brookingsapbiology Wikispace. **Located at**: **http://brookingsapbiology.wikispaces.com/Male+Reproductive+System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- spermatogenesis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/spermatogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- testosterone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/testosterone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...anatomy_en.svg**. **License**: ***Public Domain: No Known Copyright***
- Scrotum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Scrotum**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- epididymis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/epididymis**. License: ***CC BY-SA: Attribution-ShareAlike***
- scrotum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/scrotum**. License: ***CC BY-SA: Attribution-ShareAlike***
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...anatomy_en.svg**. License: ***Public Domain: No Known Copyright***
- Figure 28 01 02.JPG. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...e_28_01_02.JPG**. License: ***CC BY: Attribution***
- Sertoli Cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sertoli_cell**. License: ***CC BY-SA: Attribution-ShareAlike***
- Luteinizing Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Luteinizing_hormone**. License: ***CC BY-SA: Attribution-ShareAlike***
- Follicle-Stimulating Hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Follic...cts_in_females**. License: ***CC BY-SA: Attribution-ShareAlike***
- Testicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Testicle**. License: ***CC BY-SA: Attribution-ShareAlike***
- Leydig cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Leydig%20cells**. License: ***CC BY-SA: Attribution-ShareAlike***
- testicle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/testicle**. License: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...inition/testes**. License: ***CC BY-SA: Attribution-ShareAlike***
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...anatomy_en.svg**. License: ***Public Domain: No Known Copyright***
- Figure 28 01 02.JPG. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...e_28_01_02.JPG**. License: ***CC BY: Attribution***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...le-Revised.jpg**. License: ***CC BY-SA: Attribution-ShareAlike***
- Gray1144. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray1144.png**. License: ***Public Domain: No Known Copyright***
- Intromittent organ. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Intromittent_organ**. License: ***CC BY-SA: Attribution-ShareAlike***
- Clitoris. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Clitoris#Glans_and_body**. License: ***CC BY-SA: Attribution-ShareAlike***
- Penis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Penis**. License: ***CC BY-SA: Attribution-ShareAlike***
- Human penis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_penis**. License: ***CC BY-SA: Attribution-ShareAlike***
- ejaculation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ejaculation**. License: ***CC BY-SA: Attribution-ShareAlike***
- glans. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glans**. License: ***CC BY-SA: Attribution-ShareAlike***
- penis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/penis**. License: ***CC BY-SA: Attribution-ShareAlike***
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...anatomy_en.svg**. License: ***Public Domain: No Known Copyright***
- Figure 28 01 02.JPG. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...e_28_01_02.JPG**. License: ***CC BY: Attribution***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...le-Revised.jpg**. License: ***CC BY-SA: Attribution-ShareAlike***
- Gray1144. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Gray1144.png**. License: ***Public Domain: No Known Copyright***
- Penis/Clitoris. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi..._Structure.JPG**. License: ***Public Domain: No Known Copyright***
- Penis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...c/Gray1158.png**. License: ***Public Domain: No Known Copyright***
- Peristalsis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peristalsis**. License: ***CC BY-SA: Attribution-ShareAlike***
- Vas deferens. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Vas_deferens**. License: ***CC BY-SA: Attribution-ShareAlike***

- Seminiferous tubules. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Seminiferous_tubules. **License**: *CC BY-SA: Attribution-ShareAlike*
- Efferent ductules. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Efferent_ductules. **License**: *CC BY-SA: Attribution-ShareAlike*
- vasectomy. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vasectomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- vas deferens. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/vas_deferens. **License**: *CC BY-SA: Attribution-ShareAlike*
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/d/d1/Male_anatomy_en.svg. **License**: *Public Domain: No Known Copyright*
- Figure 28 01 02.JPG. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_01_02.JPG. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...le-Revised.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1144. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Gray1144.png. **License**: *Public Domain: No Known Copyright*
- Penis/Clitoris. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._Structure.JPG. **License**: *Public Domain: No Known Copyright*
- Penis. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/7c/Gray1158.png. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/4/49/Seminiferous_tubule_and_sperm_low_mag.jpg/504px-Seminiferous_tubule_and_sperm_low_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu testis schematic. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Illu_testis_schematic.jpg. **License**: *Public Domain: No Known Copyright*
- Lipfuscin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Lipofuscin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Seminal vesicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Seminal_vesicle. **License**: *CC BY-SA: Attribution-ShareAlike*
- Bulbourethral gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bulbourethral_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Prostate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Prostate. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology.../seminal-gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- prostate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/prostate. **License**: *CC BY-SA: Attribution-ShareAlike*
- bulbourethral gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/bulbourethral_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- male anatomymale anatomyKevin (tsaitgaist) REDON. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/d/d1/Male_anatomy_en.svg. **License**: *Public Domain: No Known Copyright*
- Figure 28 01 02.JPG. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_01_02.JPG. **License**: *CC BY: Attribution*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...le-Revised.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray1144. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Gray1144.png. **License**: *Public Domain: No Known Copyright*
- Penis/Clitoris. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/4/44/Penile-Clitoral_Structure.JPG. **License**: *Public Domain: No Known Copyright*
- Penis. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/7c/Gray1158.png. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/4/49/Seminiferous_tubule_and_sperm_low_mag.jpg/504px-Seminiferous_tubule_and_sperm_low_mag.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Illu testis schematic. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Illu_testis_schematic.jpg. **License**: *Public Domain: No Known Copyright*
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/9/97/Gray1160.png. **License**: *Public Domain: No Known Copyright*

- Prostate. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/2b/Gray543.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 26.3F: Accessory Sex Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4: PHYSIOLOGY OF THE MALE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

26.4A: Male Sexual Response

26.4B: Sperm

26.4C: Spermatogenesis

26.4D: Semen

26.4E: Hormonal Regulation of the Male Reproductive System

This page titled 26.4: Physiology of the Male Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4A: Male Sexual Response

Physiological changes occur to male genitalia during sexual arousal.

LEARNING OBJECTIVE

Outline the process of male sexual response

Key Takeaways

Key Points

- Male sexual response is demonstrated by vasodilation and blood engorgement of the penis, leading to an erection.
- The testes rise and grow larger and warmer as blood pressure rises.
- The muscles of the pelvic floor, the vesicles, and the prostrate contract, injecting sperm into the urethra of the penis and resulting in the onset of orgasm.
- Ejaculation continues with orgasm.
- Following orgasm, there is a gradual loss of erection and a feeling of relaxation known as the refractory period.
- Cognitive factors involving visual stimuli and high levels of activity in the amygdala and hypothalamus contribute to sexual arousal and sexual response in males.

Key Terms

- **erection**: The physiological process by which a penis becomes engorged with blood.
- **sexual arousal**: Changes that occur during or in anticipation of sexual activity.
- **refractory period**: The period after excitation, during which a membrane recovers its polarization and is not able to respond to a second stimulus.
- **sex flush**: Increased blood flow leading to reddening of the skin in response to sexual arousal or orgasm.
- **tumescence**: The normal engorgement of the erectile tissue with blood.
- **genitalia**: Sex organs.

The erect penis is commonly correlated with male sexual arousal.

Physiological Effects of Arousal

Physical and/or psychological stimulation leads to vasodilation and subsequent increased blood flow into the three spongy areas that run along the length of the penis (the two corpora cavernosa and the corpus spongiosum). The penis grows enlarged and firm, the skin of the scrotum is pulled tighter, and the testes are pulled up against the body.

As sexual arousal and stimulation continues, the glans of the erect penis will swell wider. As the genitals become further engorged with blood, their color deepens and the testes can grow up to 50% larger. As the testes continue to rise, a feeling of warmth may develop around them and the perineum. With further sexual stimulation, the heart rate increases, blood pressure rises, and breathing becomes more rapid. The increase in blood flow in the genitals and other regions may lead, in some men, to a sex flush.

The muscles of the pelvic floor, the ductus deferens (between the testes and the prostate), the seminal vesicles, and the prostate gland may begin to contract in a way that forces sperm and semen into the urethra inside the penis. This is the onset of orgasm and once this has started, the man likely will continue to ejaculate and orgasm fully, with or without further stimulation. If sexual stimulation stops before orgasm, the physical effects of the stimulation, including the vasocongestion, will subside in a short time. Repeated or prolonged stimulation without orgasm and ejaculation can lead to discomfort in the testes that is sometimes called “blue balls.”

Nonarousal-Related Erection

The relationship between erection and arousal is not one-to-one. Some men older than age 40 report that they do not always have an erection when sexually aroused. A male erection can occur during sleep (nocturnal penile tumescence) without conscious sexual arousal or due to mechanical stimulation (e.g. rubbing against a bed sheet) alone. A young man or one with a strong sexual drive may experience enough sexual arousal for an erection with a passing thought or just the sight of a passerby. Once erect, his penis may gain enough stimulation from contact with the inside of his clothing to maintain the erection for more time.

Post-Orgasm Response

After orgasm and ejaculation, a refractory period usually ensues, characterized by loss of erection, a decline in any sex flush, decreased interest in sex, and a feeling of relaxation associated with the action of the neurohormones oxytocin and prolactin. The intensity and duration of the refractory period can be very short in a highly aroused young man in a highly-arousing situation, perhaps without even a noticeable loss of erection. It can be as long as a few hours or days in mid-life and older men.

Hormonal Responses

Several hormones affect sexual arousal, including testosterone, cortisol, and estradiol. However, the specific roles of these hormones are not clear. Testosterone is the most commonly-studied hormone involved with sexuality, and it plays a key role in sexual arousal in males, with strong effects on central arousal mechanisms.

This page titled 26.4A: Male Sexual Response is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4B: Sperm

Sperm are the male “seeds,” germ cells, or gametes.

LEARNING OBJECTIVE

Describe the anatomy and function of sperm

Key Takeaways

Key Points

- Sperm fertilize the oocyte, donate the paternal chromatin, and provide the centrosome that maintains the zygote’s microtubule system.
- Sperm have three parts: a head, which holds the chromatin, a midpiece filled with mitochondria to provide energy, and a flagellum or tail to move the sperm from the vagina to the oocyte.
- Sperm with one tail, such as human sperm, are referred to as spermatozoa.
- Sperm quality and quantity decrease with age.

Key Terms


- **anisogamy**: The form of sexual reproduction that involves the union or fusion of two gametes that differ in size and/or form.
- **spermatozoa**: A motile sperm cell or moving form of the haploid cell that is the male gamete.
- **acrosome**: A caplike structure over the anterior half of the sperm’s head.
- **ATP**: An acronym for adenosine triphosphate, which transports chemical energy within cells for metabolism.
- **oogamy**: A form of anisogamy (heterogamy) in which the female gamete (oocyte) is significantly larger than the male gamete (sperm) and is non-motile. The male gametes are highly motile and compete for the fertilization of the immotile oocyte.

EXAMPLE

The term sperm is derived from the Greek word for seed and refers to the male reproductive cells. In the types of sexual reproduction known as anisogamy and oogamy, there are marked differences in the size of the gametes, with the smaller termed the “male” or sperm cells. Sperm cells cannot divide and have a limited lifespan. After fusion with egg cells during fertilization, a new organism forms, beginning as a totipotent zygote. The human sperm cell is haploid so that its 23 chromosomes can join the 23 chromosomes of the female egg to form a diploid cell. During fertilization, the sperm provides the following three essential parts to the oocyte:

1. A signalling or activating factor that causes the metabolically dormant oocyte to activate
2. The haploid paternal genome
3. The centrosome, which is responsible for maintaining the microtubule system

Sperm Anatomy




Closeup of Mammalian Fertilization: Micrograph of a sperm poised to enter an ovum

Sperm develop in the testes and consist of a head, a midpiece, and a tail. The head contains the nucleus with densely coiled chromatin fibers, surrounded anteriorly by an acrosome that contains enzymes for penetrating the female egg. The midpiece has a central filamentous core with many mitochondria spiraled around it.

Sperm Physiology and Function

In animals, most of the energy (ATP) for sperm motility is derived from the metabolism of fructose carried in the seminal fluid. This takes place in the mitochondria located in the sperm's midpiece. This energy is used for the journey through the female cervix, uterus, and uterine tubes.

Motile sperm cells typically move via flagella and require a water medium in order to swim toward the egg for fertilization.These cells cannot swim backwards due to the nature of their propulsion. The uniflagellated sperm cells (with one flagellum) of animals are referred to as spermatozoa.



Human Sperm: Detailed and labeled diagram of a human spermatozoa

Fertility Factors

Sperm quantity and quality are the main parameters in semen quality, a measure of the ability of semen to accomplish fertilization. The genetic quality of sperm, as well as its volume and motility, all typically decrease with age.

This page titled 26.4B: Sperm is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4C: Spermatogenesis

Male gametes (sperm cells) are haploid cells produced via spermatogenesis.

LEARNING OBJECTIVE

Describe the process of spermatogenesis

Key Takeaways

Key Points

- Spermatogenesis begins with a diploid spermatogonium in the seminiferous tubules, which divides mitotically to produce two diploid primary spermatocytes.
- The primary spermatocyte then undergoes meiosis I to produce two haploid secondary spermatocytes.
- The haploid secondary spermatocytes undergo meiosis II to produce four haploid spermatids.
- Each spermatid begins to grow a tail and a mitochondrial-filled midpiece, while the chromatin is tightly packaged into an acrosome at the head.
- Maturation removes excess cellular material, turning spermatids into inactive, sterile spermatozoa that are transported via peristalis to the epididymus.
- The spermatozoa gain motility in the epididymus, but do not use that ability until they are ejaculated into the vagina.
- Spermatogenesis requires optimal environmental conditions.

Key Terms

- **spermatozoa**: A motile sperm cell, or moving form of the haploid cell that is the male gamete.
- **spermatocyte**: A male gametocyte, from which a spermatozoon develops.
- **axoneme**: Cytoskeletal inner core structure of eukaryotic flagella.
- **spermatid**: A haploid cell produced by meiosis of a spermatocyte that develops into a spermatozoon.
- **spermatogonium**: Any of the undifferentiated cells in the male gonads that become spermatocytes.

Spermatogenesis is the process by which male primary sperm cells undergo meiosis and produce a number of cells calls spermatogonia, from which the primary spermatocytes are derived. Each primary spermatocyte divides into two secondary spermatocytes and each secondary spermatocyte into two spermatids or young spermatozoa. These develop into mature spermatozoa, also known as sperm cells. Thus, the primary spermatocyte gives rise to two cells, the secondary spermatocytes, which in turn produce four spermatozoa.

Male Gametogenesis


Spermatozoa are the mature male gametes in many sexually reproducing organisms. Spermatogenesis is the male version of gametogenesis and results in the formation of spermatocytes possessing half the normal complement of genetic material.

In mammals, it occurs in the male testes and epididymis in a stepwise fashion that takes approximately 64 days.

Spermatogenesis, essential for sexual reproduction is highly dependent upon optimal conditions to occur correctly. DNA methylation and histone modification have been implicated in the regulation of this process. It starts at puberty and usually continues uninterrupted until death, although a slight decrease can be discerned in the quantity of sperm produced with increase in age.

Steps in Spermatogenesis

Step 1: Spermatocytogenesis



Spermatocytogenesis: Diagram of the steps of spermatocytogenesis, including type Ad spermatogonium, type Ap Spermatogonium, type B spermatogonium, primary spermatocyte, and secondary spermatocyte.

Mitotic division of a diploid spermatogonium that resides in the basal compartment of the seminiferous tubules, resulting in two diploid intermediate cells called primary spermatocytes.

Each primary spermatocyte then moves into the adluminal compartment of the seminiferous tubules, duplicates its DNA, and subsequently undergoes meiosis I to produce two haploid secondary spermatocytes.

Secondary spermatocytes later divide into haploid spermatids. During this division, random inclusion of either parental chromosome and chromosomal crossover both increase the genetic variability of the gamete.

Each cell division from a spermatogonium to a spermatid is incomplete; the cells remain connected to one another by bridges of cytoplasm to allow synchronous development. Not all spermatogonia divide to produce spermatocytes; otherwise, the supply would run out. Instead, certain types of spermatogonia divide to produce copies of themselves, thereby ensuring a constant supply of gametogonia to fuel spermatogenesis.

Step 2: Spermatidogenesis

The creation of spermatids from secondary spermatocytes. Secondary spermatocytes produced earlier rapidly enter meiosis II and divide to produce haploid spermatids. The brevity of this stage means that secondary spermatocytes are rarely seen in histological preparations.


Step 3: Spermiogenesis

At this stage, each spermatid begins to grow a tail and develop a thickened midpiece where the mitochondria gather and form an axoneme. Spermatid DNA also undergoes packaging, becoming highly condensed. The DNA is packaged with specific nuclear basic proteins, which are subsequently replaced with protamines during spermatid elongation. The resultant tightly packed chromatin is transcriptionally inactive. The Golgi apparatus surrounds the now condensed nucleus, becoming the acrosome. One of the centrioles of the cell elongates to become the tail of the sperm.

The non-motile spermatozoa are transported to the epididymis in testicular fluid secreted by the Sertoli cells with the aid of

peristaltic contraction. While in the epididymis, the spermatozoa gain motility and become capable of fertilization. However, transport of the mature spermatozoa through the remainder of the male reproductive system is achieved via muscle contraction rather than the spermatozoon's recently acquired motility.


Physiology of Spermatogenesis



Seminiferous Tubule: Micrograph showing seminiferous tubule with maturing sperm.

Maturation takes place under the influence of testosterone, which removes the remaining unnecessary cytoplasm and organelles. The excess cytoplasm, known as residual bodies, is phagocytosed by surrounding Sertoli cells in the testes. The resulting spermatozoa are now mature but lack motility, rendering them sterile. The mature spermatozoa are released from the protective Sertoli cells into the lumen of the seminiferous tubule in a process called spermiation.

Spermatogenesis is highly sensitive to fluctuations in the environment, particularly hormones and temperature. Seminiferous epithelium is sensitive to elevated temperature in humans and is adversely affected by temperatures as high as normal body temperature. Consequently, the testes are located outside the body in a sack of skin called the scrotum. The optimal temperature is maintained at 2 °C below body temperature in human males. This is achieved by regulation of blood flow and positioning towards and away from the heat of the body by the cremaster muscle and the dartos smooth muscle in the scrotum. Dietary deficiencies (such as vitamins B, E, and A), anabolic steroids, metals (cadmium and lead), x-ray exposure, dioxin, alcohol, and infectious diseases will also adversely affect the rate of spermatogenesis.



Spermatozoon: Diagram of parts of a spermatozoon, including the acrosome, plasma membrane, nucleus, centriole, mitochondria, terminal disc, axial filament, tail, endpiece, midpiece, and head.

This page titled 26.4C: Spermatogenesis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4D: Semen

Semen is a fluid produced by the seminal vesicles.

LEARNING OBJECTIVE

Describe semen

Key Takeaways

Key Points

- Seminal fluid mixes with fluids produced by the prostate and bulbourethral glands.
- The seminal fluid provides nutrition and protection for sperm during its journey through the female reproductive tract.
- Semen initially coagulates in the vagina, then liquefies to allow the sperm to move.

Key Terms

- **seminal vesicle**: One of two simple tubular glands located behind the male urinary bladder, responsible for the production of about sixty percent of the fluid that ultimately becomes semen.
- **seminal fluid**: Semen is a fluid that helps in promoting the survival of spermatozoa and provides a medium through which they can move.
- **semen**: The fluid produced in male reproductive organs of an animal that contains the reproductive cells.

Semen is an organic fluid, also known as seminal fluid, that may contain spermatozoa. It is secreted by the gonads (sexual glands) and can fertilize female ova. In humans, seminal fluid contains several components besides spermatozoa, including enzymes (proteolytic and others) and fructose. These elements promote the survival of spermatozoa and provide a medium for motility. Semen is produced and originates from the seminal vesicles, located in the pelvis. The process that results in the discharge of semen is called ejaculation.

Semen Production and Secretion

During the process of ejaculation, sperm pass through the ejaculatory ducts and mix with fluids from the seminal vesicle, the prostate, and the bulbourethral glands to form semen. The seminal vesicles produce a yellowish viscous fluid rich in fructose, amino acids, and other substances that make up about 70% of human semen. The prostatic secretion, influenced by dihydrotestosterone, is a whitish (sometimes clear), thin fluid containing proteolytic enzymes, citric acid, acid phosphatase, and lipids. The bulbourethral glands secrete a clear fluid to lubricate the lumen of the urethra.

Sperm Protection and Transport

Sertoli cells, which nurture and support developing spermatocytes, secrete a fluid into seminiferous tubules that helps transport sperm to the genital ducts. The ductuli efferentes possess cuboidal cells with microvilli and lysosomal granules that modify the semen by reabsorbing some fluid. Once the semen enters the ductus epididymis, the principal cells (which contain pinocytotic vessels indicating fluid reabsorption) secrete glycerophosphocholine, which most likely inhibits premature capacitation.

The seminal plasma provides a nutritive and protective medium for the spermatozoa during their journey through the female reproductive tract. The normal environment of the vagina is a hostile one for sperm cells, as it is acidic (from the native microflora producing lactic acid), viscous, and patrolled by immune cells. The components in the seminal plasma attempt to compensate for this hostile environment. Basic amines such as putrescine, spermine, spermidine, and cadaverine are responsible for the smell and flavor of semen. These alkaline bases counteract the acidic environment of the vaginal canal and protect DNA inside the sperm from acidic denaturation.

Characteristics of Ejaculate

According to the World Health Organization, normal human semen has a volume of 2 ml or greater, pH of 7.2 to 8.0, sperm concentration of 20×106 spermatozoa/ml or more, sperm count of 40×106 spermatozoa per ejaculate or more, and motility of 50% or more within 60 minutes of ejaculation. After ejaculation, the latter part of the semen coagulates immediately, forming globules. After about 15–30 minutes, a prostate-specific antigen present in the semen causes the decoagulation of the seminal coagulum. It is postulated that the initial clotting helps keep the semen in the vagina, while liquefaction frees the sperm to make their journey to the ova.

Semen quality is a measure of the ability of semen to accomplish fertilization and thus a measure of a man's fertility. Semen can be preserved for long-term storage by cryopreservation. For human sperm, the longest reported successful storage with this method is 21 years.

This page titled 26.4D: Semen is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.4E: Hormonal Regulation of the Male Reproductive System

The male reproductive system is regulated by the production, stimulation, and feedback of specific hormones.

LEARNING OBJECTIVE

Differentiate among the hormones of the male reproductive system

Key Takeaways

Key Points

- GnRH is made in the hypothalamus and travels to the pituitary where it stimulates FSH and LH secretion.
- FSH is necessary for sperm maturation.
- LH binds to Leydig cells to stimulate testosterone secretion and androgen production.
- Testosterone stimulates sex drive.
- Inhibin acts as negative feedback to slow the release of FSH and GnRH.

Key Terms

- **GnRH**: Gonadotropin-releasing hormone is a trophic peptide hormone responsible for the release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the anterior pituitary.
- **LH**: Luteinizing hormone is produced by the anterior pituitary gland and in males causes the synthesis and secretion of testosterone and androgen.
- **FSH**: Follicle-stimulating hormone stimulates both the production of androgen-binding protein by Sertoli cells and the formation of the blood-testis barrier.

Hormonal control of spermatogenesis varies among species. In humans, the mechanisms are not completely understood. However, it is known that initiation of spermatogenesis occurs at puberty due to the interaction of the hypothalamus, pituitary gland, and Leydig cells. If the pituitary gland is removed, spermatogenesis can still be initiated by follicle-stimulating hormone (FSH) and testosterone.

The Role of Hormones in Male Reproduction

Studies from rodent models suggest that gonadotropin hormones (both LH and FSH) support the process of spermatogenesis by suppressing the proapoptotic signals and thus promoting spermatogenic cell survival. The Sertoli cells themselves mediate parts of spermatogenesis through hormone production. They are capable of producing the hormones estradiol and inhibin. The Leydig cells are also capable of producing estradiol in addition to their main product, testosterone.

Gonadotropin-Releasing Hormone

Gonadotropin-releasing hormone (GnRH) is mainly made in the preoptic area of the hypothalamus before traveling to the pituitary gland. There it stimulates the synthesis and secretion of the gonadotropins, FSH and luteinizing hormone (LH).

Follicle-Stimulating Hormone

Follicle-stimulating hormone (FSH) is released by the anterior pituitary gland. Its presence in males is necessary for the maturation of spermatozoa. Follicle-stimulating hormone stimulates both the production of androgen-binding protein by Sertoli

cells and the formation of the blood-testis barrier. Increasing the levels of FSH increases the production of spermatozoa by preventing the apoptosis of type A spermatogonia.

Luteinizing Hormone


Luteinizing hormone (LH) is released by the anterior pituitary gland. In the testes, LH binds to receptors on Leydig cells, which stimulates the synthesis and secretion of testosterone.

Testosterone

Testosterone is made in the interstitial cells of the testes. It stimulates the sex drive and is associated with aggression. Androgen-binding protein is essential to concentrating testosterone in levels high enough to initiate and maintain spermatogenesis, which can be 20-50 times higher than the concentration found in blood. The sequestering of testosterone in the testes is initiated by FSH, and only testosterone is required to maintain spermatogenesis.

Inhibin

Inhibin is secreted by the Sertoli cells and acts to decrease the levels of FSH. The hormone is released into the circulation when the sperm count is too high.



Hormonal Influence: This flowchart details the steps involved in hormonal control of male reproduction.

This diagram depicts the hormonal regulation of male reproduction, including the following steps: hypothalamus, GnHR secretion, anterior pituitary, FSH and LH secretion, negative feedback, Leydig cells, Sertoli cells, testosterone secretion, inhibit secretion, spermatogenesis, various target tissues, maintenance of accessory reproductive organs and secondary sex characteristics, sex drive, protein synthesis in skeletal muscle, and bone growth in adolescents.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- erection. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/erection. **License**: ***CC BY-SA: Attribution-ShareAlike***
- refractory period. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/refractory_period. **License**: ***CC BY-SA: Attribution-ShareAlike***
- genitalia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/genitalia. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Sexual arousal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sexual_arousal. **License**: *CC BY-SA: Attribution-ShareAlike*
- Oogamy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Oogamy. **License**: *CC BY-SA: Attribution-ShareAlike*
- spermatozoa. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/spermatozoa. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anisogamy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anisogamy. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sperm. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sperm. **License**: *CC BY: Attribution*
- acrosome. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/acrosome. **License**: *CC BY-SA: Attribution-ShareAlike*
- sperm. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sperm. **License**: *CC BY-SA: Attribution-ShareAlike*
- Spermatogenesis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Spermatogenesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- spermatid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/spermatid. **License**: *CC BY-SA: Attribution-ShareAlike*
- spermatocyte. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/spermatocyte. **License**: *CC BY-SA: Attribution-ShareAlike*
- spermatogonium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/spermatogonium. **License**: *CC BY-SA: Attribution-ShareAlike*
- Seminiferous_tubule_and_sperm. **Authored by**: Nephron. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/F..._and_sperm.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- seminal vesicle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/seminal_vesicle. **License**: *CC BY-SA: Attribution-ShareAlike*
- seminal fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/seminal_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Semen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Semen. **License**: *CC BY-SA: Attribution-ShareAlike*
- semen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/semen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Male Reproductive System. **Provided by**: brookingsapbiology. **Located at**: http://brookingsapbiology.wikispaces...ductive+System. **Project**: Wikispace. **License**: *CC BY-SA: Attribution-ShareAlike*
- GnRH. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/GnRH. **License**: *CC BY-SA: Attribution-ShareAlike*
- LH. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/LH. **License**: *CC BY-SA: Attribution-ShareAlike*
- FSH. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/FSH. **License**: *CC BY-SA: Attribution-ShareAlike*
- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

PUBLIC DOMAIN CONTENT

- Sperm-egg. **Authored by**: Unknown. **Located at**: en.Wikipedia.org/wiki/File:Sperm-egg.jpg. **License**: *Public Domain: No Known Copyright*
- Sperm. **Authored by**: LadyofHats. **Located at**: commons.wikimedia.org/wiki/F...matozoa_en.svg. **License**: *Public Domain: No Known Copyright*
- Male reproductive hormone chart. **Authored by**: Stephaniegreenwood. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...mone_chart.gif. **License**: *Public Domain: No Known Copyright*

This page titled 26.4E: Hormonal Regulation of the Male Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5: THE FEMALE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

26.5A: Anatomy of the Female Reproductive System

26.5B: Ovaries

26.5C: Uterus

26.5D: Female Duct System

26.5E: Vagina

26.5F: Vulva

26.5G: Perineum

26.5H: Mammary Glands

This page titled 26.5: The Female Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5A: Anatomy of the Female Reproductive System

The human female reproductive system contains two main parts: the uterus and the ovaries, which produce a woman's egg cells.

Learning Objectives

- Outline the anatomy of the female reproductive system from external to internal

Key Points

- An female's internal reproductive organs are the vagina, uterus, fallopian tubes, cervix, and ovary.
- External structures include the mons pubis, pudendal cleft, labia majora and minora, vulva, Bartholin's gland, and the clitoris.
- The female reproductive system contains two main parts: the uterus, which hosts the developing fetus, produces vaginal and uterine secretions, and passes the anatomically male sperm through to the fallopian tubes; and the ovaries, which produce the anatomically female egg cells.

Key Terms

- **ovary**: A female reproductive organ, often paired, that produces ova and in mammals secretes the hormones estrogen and progesterone.
- **oviduct**: A duct through which an ovum passes from an ovary to the uterus or to the exterior (called fallopian tubes in humans).
- **vulva**: The consists of the female external genital organs.
- **oogenesis**: The formation and development of an ovum.

The human female reproductive system (or female genital system) contains two main parts:


1. Uterus
    - Hosts the developing fetus
    - Produces vaginal and uterine secretions
    - Passes the anatomically male sperm through to the fallopian tubes
2. Ovaries
    - Produce the anatomically female egg cells.
    - Produce and secrete estrogen and progesterone

These parts are internal; the vagina meets the external organs at the vulva, which includes the labia, clitoris, and urethra. The vagina is attached to the uterus through the cervix, while the uterus is attached to the ovaries via the fallopian tubes. At certain intervals, the ovaries release an ovum, which passes through the fallopian tube into the uterus.

If, in this transit, it meets with sperm, the sperm penetrates and merges with the egg, fertilizing it. The fertilization usually occurs in the oviducts, but can happen in the uterus itself. The zygote then implants itself in the wall of the uterus, where it begins the process of embryogenesis and morphogenesis. When developed enough to survive outside the womb, the cervix dilates and contractions of the uterus propel the fetus through the birth canal (vagina).

The ova are larger than sperm and have formed by the time an anatomically female infant is born. Approximately every month, a process of oogenesis matures one ovum to be sent down the fallopian tube attached to its ovary in anticipation of fertilization. If not fertilized, this egg is flushed out of the system through menstruation.

An anatomically female's internal reproductive organs are the vagina, uterus, fallopian tubes, cervix, and ovary.
The external components include the mons pubis, pudendal cleft, labia majora, labia minora, Bartholin's glands, and clitoris.



The Female Reproductive System

Female Repro: Illustrated sagittal view of the female reproductive system.

This page titled 26.5A: Anatomy of the Female Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5B: Ovaries

The ovaries are the ovum-producing organs of the internal female reproductive system.

Learning Objectives

- Describe the location and function of the ovaries

Key Points

- In addition to producing ova, the ovaries are endocrine organs and produce hormones that act during the female menstrual cycle and pregnancy.
- Ovaries secrete estrogen and progesterone.
- Each ovary is located in the lateral wall of the pelvis in a region called the ovarian fossa.
- The ovaries are attached to the uterus via the ovarian ligament (which runs in the broad ligament).
- Usually, the ovaries take turns releasing eggs every month; however, if one ovary is absent or dysfunctional then the other ovary releases eggs every month.
- There are two extremities to the ovary, the tubal extremity and the uterine extremity.

Key Terms

- **intraperitoneal**: Located within the inner layer of the peritoneum (serous membrane that forms the lining of the abdominal cavity).
- **corpus luteum**: A temporary endocrine structure in female ovaries that is essential for establishing and maintaining pregnancy.
- **libido**: A person's overall sexual drive.
- **follicle**: A spheroid cellular aggregation found in the ovaries that secretes hormones that influence the stages of the menstrual cycle.
- **ovary**: A female reproductive organ, often paired, that produces ova and in mammals secretes the hormones estrogen and progesterone.


The ovary is an ovum-producing reproductive organ, typically found in pairs as part of the vertebrate female reproductive system. Ovaries in females are analogous to testes in males in that both are gonads and endocrine glands. Ovaries secrete both estrogen and progesterone. Estrogen is responsible for the appearance of secondary sex characteristics of females at puberty and for the maturation and maintenance of the reproductive organs in their mature functional state. Progesterone functions with estrogen by promoting menstrual cycle changes in the endometrium.

Anatomical Features

The ovaries are located in the lateral wall of each side of the pelvis in a region called the ovarian fossa. The fossa usually lies beneath the external iliac artery and in front of the ureter and internal iliac artery.

In humans, the paired ovaries lie within the pelvic cavity on either side of the uterus, to which they are attached via a fibrous cord called the ovarian ligament. The ovaries are tethered to the body wall via the suspensory ligament of the ovary. The part of the broad ligament of the uterus that covers the ovary is known as the mesovarium. The ovary is the only organ in the human body which is totally invaginated into the peritonium, making it the only intraperitoneal organ.

There are two extremities to the ovary, the tubal extremity and the uterine extremity. The tubal extremity is the end to which the Fallopian tube attaches via the infundibulopelvic ligament. The uterine extremity points downward and is attached to the uterus via the ovarian ligament.



Ovary: A pictorial illustration of the female reproductive system

Physiology and Function

The ovaries are the site of egg cell production and also have specific endocrine function.

Oogenesis

The ovaries are the site of gamete (egg cell, oocyte) production. The developing egg cell (or oocyte) grows within the environment provided by ovarian follicles. Follicles are composed of different types and number of cells according to their maturation stage, which can be determined by their size. When oocyte maturation is completed, a luteinizing hormone (LH) surge secreted by the pituitary gland stimulates follicle rupture and oocyte release.

This oocyte development and release process is referred to as ovulation. The follicle remains functional and transforms into a corpus luteum, which secretes progesterone to prepare the uterus for possible embryo implantation. Usually each ovary takes turns releasing eggs each month. However, this alternating egg release is random. When one ovary is absent or dysfunctional, the other ovary will continue to release eggs each month.

Endocrine Function

Ovaries secrete estrogen, progesterone, and testosterone. Estrogen is responsible for the secondary sex characteristics of females at puberty. It is also crucial for the maturation and maintenance of the mature and functional reproductive organs. Progesterone prepares the uterus for pregnancy and the mammary glands for lactation. The co-actions of progesterone and estrogen promote menstrual cycle changes in the endometrium. In women, testosterone is important for the development of muscle mass, muscle and bone strength, and for optimal energy level. It also has a role in libido in women.

This page titled 26.5B: Ovaries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5C: Uterus

The uterus is the largest and major organ of the female reproductive tract that is the site of fetal growth and is hormonally responsive.

Learning Objectives

- Describe the structure and function of the uterus

Key Points

- The body of the uterus is connected to the ovaries via the fallopian tubes, and opens into the vagina via the cervix.
- Two Müllerian ducts
  usually form initially in a female fetus, but in humans they completely fuse into a single uterus during gestation.
- The uterus is essential in sexual response by directing blood flow to the pelvis and to the external genitalia, including the ovaries, vagina, labia, and clitoris.
- The reproductive function of the uterus is to accept a fertilized ovum which passes through the utero-tubal junction from the fallopian tube.
- The lining of the uterine cavity is called the endometrium.


Key Terms

- **linea terminalis**: Part of the pelvic brim, which is the edge of the pelvic inlet.
- **adenomyosis**: A condition characterized by the breaking through of the endometrium into the muscle wall of the uterus.
- **uterus**: An organ of the female reproductive system in which the young are conceived and develop until birth; the womb.
- **endometrium**: The mucous membrane that lines the uterus in mammals and in which fertilized eggs are implanted.
- **fallopian tubes**: The fallopian tubes, also known as oviducts, uterine tubes, and salpinges (singular salpinx) are two very fine tubes lined with ciliated epithelia, and lead from the ovaries of female mammals into the uterus via the utero-tubal junction.

The uterus or womb is a major female hormone -responsive reproductive sex organ of most mammals including humans. One end, the cervix, opens into the vagina, while the other is connected to one or both fallopian tubes, depending on the species. It is within the uterus that the fetus develops during gestation, usually developing completely in placental mammals such as humans.

Two Müllerian ducts usually form initially in a female fetus and, in humans, they completely fuse into a single uterus depending on the species. The uterus consists of a body and a cervix. The cervix protrudes into the vagina. The uterus is held in position within the pelvis by condensations of endopelvic fascia, which are called ligaments. These ligaments include the pubocervical, transverse, cervical, cardinal, and uterosacral ligaments. It is covered by a sheet-like fold of peritoneum, the broad ligament.


The uterus is essential in sexual response by directing blood flow to the pelvis and to the external genitalia, including the ovaries, vagina, labia, and clitoris. The reproductive function of the uterus is to accept a fertilized ovum which passes through the utero-tubal junction from the fallopian tube. It implants into the endometrium, and derives nourishment from blood vessels which develop exclusively for this purpose.



Uterus: Vessels of the uterus and its appendages, rear view.

The fertilized ovum becomes an embryo, attaches to a wall of the uterus, creates a placenta, and develops into a fetus (gestates) until childbirth. Due to anatomical barriers such as the pelvis, the uterus is pushed partially into the abdomen due to its expansion during pregnancy. Even during pregnancy, the mass of a human uterus amounts to only about a kilogram (2.2 pounds).

The uterus is located inside the pelvis immediately dorsal (and usually somewhat rostral) to the urinary bladder and ventral to the rectum. The human uterus is pear-shaped and about three inches (7.6 cm) long. The uterus can be divided anatomically into four segments: The fundus, corpus, cervix and the internal os.



Ovary: A pictorial illustration of the female reproductive system.

The uterus is in the middle of the pelvic cavity in frontal plane (due to ligamentum latum uteri). The fundus does not surpass the linea terminalis. The fundus of the uterus is the top, rounded portion, opposite from the cervix. The vaginal part of the cervix does not extend below interspinal line. The uterus is mobile and moves under the pressure of the full bladder or full rectum anteriorly, whereas if both are full it moves upwards. Increased intra-abdominal pressure pushes it downwards. The mobility is conferred to it by musculo-fibrous apparatus that consists of a suspensory and sustentacular part. Under normal circumstances the suspensory part keeps the uterus in anteflexion and anteversion (in 90% of women) and keeps it “floating” in the pelvis. In cases where the uterus is “tipped,” also known as retroverted uterus, women may have symptoms of pain during sexual intercourse, pelvic

pain during menstruation, minor incontinence, urinary tract infections, difficulty conceiving, and difficulty using tampons. A pelvic examination by a doctor can determine if a uterus is tipped.

The lining of the uterine cavity is called the endometrium. It consists of the functional endometrium and the basal endometrium from which the former arises. Damage to the basal endometrium results in adhesion formation and/or fibrosis (Asherman's syndrome). In all placental mammals, including humans, the endometrium builds a lining periodically which is shed or reabsorbed if no pregnancy occurs. Shedding of the functional endometrial lining is responsible for menstrual bleeding (known colloquially as a "period" in humans, with a cycle of approximately 28 days, +/- 7 days of flow and +/- 21 days of progression) throughout the fertile years of a female and for some time beyond.

Depending on the species and attributes of physical and psychological health, weight, environmental factors of circadian rhythm, photoperiodism (the physiological reaction of organisms to the length of day or night), the effect of menstrual cycles to the reproductive function of the uterus is subject to hormone production, cell regeneration, and other biological activities. The menstrual cycles may vary from a few days to six months, but can vary widely even in the same individual, often stopping for several cycles before resuming.

The uterus mostly consists of smooth muscle, known as myometrium. The innermost layer of myometrium is known as the junctional zone, which becomes thickened in adenomyosis. The parametrium is the loose connective tissue around the uterus. The perimetrium is the peritoneum covering of the fundus and ventral and dorsal aspects of the uterus. The uterus is primarily supported by the pelvic diaphragm, perineal body, and the urogenital diaphragm. Secondarily, it is supported by ligaments and the peritoneum (broad ligament of uterus).

This page titled 26.5C: Uterus is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5D: Female Duct System

The Fallopian tubes, or oviducts, connect the ovaries to the uterus.

Learning Objectives

- Describe the structures of the female duct system

Key Points

- The Fallopian tube allows passage of the egg from the ovary to the uterus.
- The lining of the Fallopian tubes are ciliated and have several segments, including the infundibulum, ampullary, isthmus, and interstitial regions.
- Interspersed between the ciliated cells are peg cells, which contain apical granules and produce the tubular fluid that contains nutrients for spermatozoa, oocytes, and zygotes.
- Occasionally, the embryo implants into the Fallopian tube instead of the uterus, creating an ectopic pregnancy.

Key Terms


- **oviduct**: A duct through which an ovum passes from an ovary to the uterus or to the exterior.
- **fallopian tubes**: Also known as oviducts, uterine tubes, and salpinges (singular salpinx), two very fine tubes lined with ciliated epithelia, leading from the ovaries of female mammals into the uterus via the uterotubal junction.
- **ovarian follicle**: The basic units of female reproductive biology, each composed of roughly spherical aggregations of cells found in the ovary.

The Fallopian tubes, also known as oviducts, uterine tubes, and salpinges (singular salpinx), are two very fine tubes lined with ciliated epithelia, leading from the ovaries of female mammals into the uterus via the uterotubal junction. In non-mammalian vertebrates, the equivalent structures are the oviducts. These tubes allows passage of the egg from the ovary to the uterus.

The different segments of the fallopian tube are (lateral to medial):

- The infundibulum with associated fimbriae near the ovary
- The ampullary region that represents the major portion of the lateral tube
- The isthmus, which is the narrower part of the tube that links to the uterus
- The interstitial (intramural) part that transverses the uterine musculature

The tubal ostium is the point at which the tubal canal meets the peritoneal cavity, while the uterine opening of the Fallopian tube is the entrance into the uterine cavity, the uterotubal junction.



Uterine Segments: Illustrative drawing of the anterior view of the uterus showing the uterine segments

There are two types of cells within the simple columnar epithelium of the Fallopian tube. Ciliated cells predominate throughout the tube, but are most numerous in the infundibulum and ampulla. Estrogen increases the production of cilia on these cells.

Interspersed between the ciliated cells are peg cells, which contain apical granules and produce the tubular fluid. This fluid contains nutrients for spermatozoa, oocytes, and zygotes. The secretions also promote capacitation of the sperm by removing glycoproteins and other molecules from the plasma membrane of the sperm. Progesterone increases the number of peg cells, while estrogen increases their height and secretory activity. Tubal fluid flows against the action of the ciliae, toward the fimbrated end.

When an ovum is developing in an ovary, it is encapsulated in a sac known as an ovarian follicle. On maturation, the follicle and the ovary's wall rupture, allowing the ovum to escape. The egg is caught by the fimbriated end and travels to the ampulla where typically the sperm are met and fertilization occurs. The fertilized ovum, now a zygote, travels towards the uterus aided by the tubal cilia and tubal muscle. After about five days, the new embryo enters the uterine cavity and implants about a day later. Occasionally, the embryo implants into the Fallopian tube instead of the uterus, creating an ectopic pregnancy.

This page titled 26.5D: Female Duct System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5E: Vagina

The vagina is the female reproductive tract and has two primary functions: sexual intercourse and childbirth.

LEARNING OBJECTIVES

Describe the vagina and its functions

KEY TAKEAWAYS

Key Points


- The vagina is situated between the cervix of the uterus and the external genitalia, primarily the vulva.
- Although there is wide anatomical variation, the length of the unaroused vagina of a woman of child-bearing age is approximately 6 to 7.5 cm (2.5 to 3 in) across the anterior wall (front), and 9 cm (3.5 in) long across the posterior wall (rear).
- During sexual arousal the vagina expands in both length and width.
- A series of ridges produced by the folding of the wall of the outer third of the vagina is called the vaginal rugae.
- Vaginal lubrication is provided by the Bartholin's glands near the vaginal opening and the cervix.
- The hymen is a membrane of tissue that surrounds or partially covers the external vaginal opening.

Key Terms

- **vulva**: The vaginal opening to the uterus.
- **clitoris**: A small, sensitive, and elongated erectile organ at the anterior part of the vulva in female mammals, homologous with the penis.
- **Skene's glands**: Glands located on the anterior wall of the vagina, around the lower end of the urethra, that drain into the urethra and near the urethral opening. These may be near or part of the G-spot.
- **vagina**: A fibromuscular tubular tract which is the female sex organ and has two main functions, sexual intercourse and childbirth.

The vagina, a female sex organ, is a fibromuscular tubular tract that has two main functions: sexual intercourse and childbirth. In humans, this passage leads from the opening of the vulva to the uterus, but the vaginal tract ends at the cervix.

Anatomy of the Vagina



Vagina: The vagina is the most immediate internal female reproductive organ. This diagram also indicates the ovaries, uterus, and cervix.

The vaginal opening is much larger than the urethral opening. During arousal, the vagina gets moist to facilitate the entrance of the penis. The inner texture of the vagina creates friction for the penis during intercourse.

The vaginal opening is at the caudal end of the vulva behind the opening of the urethra. The upper quarter of the vagina is separated from the rectum by the rectouterine pouch. The vagina and the inside of the vulva are a reddish-pink color, as are most healthy internal mucous membranes in mammals. A series of ridges produced by the folding of the wall of the outer third of the vagina is called the vaginal rugae. These transverse epithelial ridges and provide the vagina with increased surface area for extension and stretching.

Vaginal lubrication is provided by the Bartholin's glands near the vaginal opening and the cervix. The membrane of the vaginal wall also produces moisture, although it does not contain any glands. Before and during ovulation, the cervix's mucus glands secrete different variations of mucus, which provides an alkaline environment in the vaginal canal that is favorable to the survival of sperm.

The hymen is a membrane of tissue that surrounds or partially covers the external vaginal opening. The tissue may or may not be ruptured by vaginal penetration. It can also be ruptured by childbirth, a pelvic examination, injury, or sports. The absence of a hymen may not indicate prior sexual activity. Similarly, its presence may not indicate a lack of prior sexual activity.

Function of the Vagina

The vagina's primary functions are sexual arousal and intercourse as well as childbirth.

Sexual Arousal and Intercourse

The concentration of the nerve endings close to the entrance of a woman's vagina (the lower third) can provide pleasurable sensation during sexual activity when stimulated. Ninety percent of the vagina's nerve endings are in this area. However, the vagina as a whole has insufficient nerve endings for sexual stimulation and orgasm; this lack of nerve endings makes childbirth significantly less painful.

Research indicates that clitoral tissue extends considerably into the vulva and vagina. During sexual arousal, and particularly clitoral stimulation, the vaginal walls lubricate to reduce friction caused by sexual activity. With arousal, the vagina lengthens rapidly to an average of about 4 in. (10 cm), and can continue to lengthen in response to pressure. As the woman becomes fully

aroused, the vagina tents (expands in length and width), while the cervix retracts. The walls of the vagina are composed of soft elastic folds of mucous membrane which stretch or contract (with support from pelvic muscles) to the size of the inserted penis or other object, stimulating the penis and helping the male to experience orgasm and ejaculation, thus enabling fertilization.

An erogenous zone commonly referred to as the G-Spot (also known as the Gräfenberg Spot) is located at the anterior wall of the vagina, about five centimeters in from the entrance. Some women experience intense pleasure if the G-Spot is stimulated appropriately during sexual activity. A G-Spot orgasm may be responsible for female ejaculation, leading some doctors and researchers to believe that G-Spot pleasure comes from the Skene's glands, a female homologue of the prostate, rather than any particular spot on the vaginal wall. Other researchers consider the connection between the Skene's glands and the G-Spot to be weak. They contend that the Skene's glands do not appear to have receptors for touch stimulation and that there is no direct evidence for their involvement. The G-Spot's existence as a distinct structure, is still under dispute, as its location can vary from woman to woman and is sometimes nonexistent.

The Vagina and Childbirth

The vagina provides the channel to deliver the baby from the uterus to its independent life outside the mother's body. During birth, the elasticity of the vagina allows it to stretch to many times its normal diameter. The vagina is often referred to as the birth canal in the context of pregnancy and childbirth.

This page titled 26.5E: Vagina is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5F: Vulva

Learning Objectives

- Describe the structures and functions of the vulva


The vulva consists of the external genital organs of the female mammal. Its development occurs during several phases, chiefly during the fetal and pubertal periods. As the outer portal of the human uterus or womb, the vulva protects its opening with a "double door": the labia majora (large lips) and the labia minora (small lips). The vulva also contains the opening of the female urethra, and thus serves the vital function of passing urine.

In human beings, major structures of the vulva are:

- The mons pubis
- The labia majora and the labia minora
- The external portion of the clitoris and the clitoral hood
- The vulval vestibule
- The pudendal cleft
- The frenulum labiorum pudendi or fourchette
- The opening (or urinary meatus) of the urethra
- The opening (or introitus) of the vagina
- The hymen

Other notable structures include:

- The perineum
- The sebaceous glands on labia majora
- The vaginal glands (Bartholin's glands and paraurethral or Skene's, glands)



Vulva: Labeled image of a vulva, showing external and internal views.

The soft mound at the front of the vulva, the mons pubis, is formed by fatty tissue covering the pubic bone. The mons pubis separates into two folds of skin called the labia majora, literally "major (or large) lips." The cleft between the labia majora is called the pudendal cleft, or cleft of Venus, and it contains and protects the other, more delicate structures of the vulva. The labia majora meet again at the perineum, a flat area between the pudendal cleft and the anus. The color of the outside skin of the labia majora

is usually close to the individual's overall skin color although there is considerable variation.

The inside skin and mucus membrane are often pink or brownish. After the onset of puberty, the mons pubis and the labia majora become covered by pubic hair. This hair sometimes extends to the inner thighs and perineum, but the density, texture, color, and extent of pubic hair coverage vary considerably due to both individual variation and cultural practices of hair modification or removal. The labia minora are two soft folds of skin within the labia majora.

The clitoris is located at the front of the vulva where the labia minora meet. The visible portion of the clitoris is the clitoral glans, roughly the size and shape of a pea. The clitoral glans is highly sensitive, containing as many nerve endings as the analogous organ in males, the glans penis. The point where the labia minora attach to the clitoris is called the frenulum clitoridis. A prepuce, the clitoral hood, normally covers and protects the clitoris; however, in women with particularly large clitorises or small prepuces, the clitoris may be partially or wholly exposed. The clitoral hood is the female equivalent of the male foreskin and may be partially hidden inside of the pudendal cleft.

The area between the labia minora is called the vulval vestibule, and it contains the vaginal and urethral openings. The urethral opening (meatus) is located below the clitoris and just in front of the vagina. This is where urine passes from the urinary bladder.

The opening of the vagina is located at the bottom of the vulval vestibule toward the perineum. The term introitus is more technically correct than "opening," since the vagina is usually collapsed, with the opening closed unless something is inserted. The introitus is sometimes partly covered by a membrane called the hymen. The hymen will rupture during the first episode of vigorous sex, and the blood produced by this rupture has been traditionally seen as a sign of virginity. However, the hymen may also rupture spontaneously during exercise or be stretched by normal activities such as use of tampons. Slightly below and to the left and right of the vaginal opening are two Bartholin glands which produce a waxy, pheromone-containing substance, the purpose of which is not yet fully known.

Key Points

- Major structures of the vulva include the labia major and minora, mons pubis, clitoris, bulb of vestibule, vulva vestibule, vestibular glands, and the genital orifice (or opening of the vagina).
- The vulva is rich in nerves that are stimulated during sexual activity and arousal.
- The vulva also contains the opening of the female urethra and thus serves the vital function of passing urine.

Key Terms

- **labia minora**: The two inner folds of skin within the cleft of the labia majora.
- **vulva**: The vaginal opening to the uterus.
- **mons pubis**: A fleshy protuberance over the pubic bones that becomes covered with hair during puberty.
- **labia majora**: The two outer rounded folds of adipose tissue that lie on either side of the opening of the vagina.

This page titled 26.5F: Vulva is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5G: Perineum

The perineum is the region between the genitals and the anus, including the perineal body and surrounding structures.

Learning Objectives

- Describe the perineum and its functions


Key Points

- The perineum refers to both external and deep structures.
- Perineal tears and episiotomy often occur in childbirth with first-time deliveries, but the risk of these injuries can be reduced by preparing the perineum through massage.
- The perineum is an erogenous zone for both males and females.

Key Terms

- **lower rabbus**: The term for perineum often used in the UK.
- **perineum**: The region of the body inferior to the pelvic diaphragm and between the legs. It is a diamond-shaped area on the inferior surface of the trunk which includes the anus and, in females, the vagina.
- **episiotomy**: A surgical incision through the perineum made to enlarge the vagina and assist childbirth.
- **perineal body**: A pyramid-shaped fibromuscular mass in the middle line of the perineum at the junction between the urogenital triangle and the anal triangle.

In human anatomy, the perineum is the surface region between the pubic symphysis and coccyx in both males and females, including the perineal body and surrounding structures. The boundaries vary in classification but generally include the genitals and anus. It is an erogenous zone for both males and females.



Perineum Illustration: Illustrated drawing of the muscles of the female perineum.

The term perineum may refer to only the superficial structures in this region or be used to include both superficial and deep structures. The term lower rabbus is used colloquially in the UK to describe this structure. Perineal tears and episiotomy often occur in childbirth with first-time deliveries, but the risk of these injuries can be reduced by preparing the perineum through

massage.

The perineum corresponds to the outlet of the pelvis. Its deep boundaries are:

- The pubic arch and the arcuate ligament of the pubis
- The tip of the coccyx
- he inferior rami of the pubis and ischial tuberosity, and the sacrotuberous ligament

The perineum includes two distinct regions separated by the pelvic diaphragm. Its structures include:

- Superficial and deep perineal pouches
- Ischioanal fossa, a fat-filled space at the lateral sides of the anal canal bounded laterally by obturator internus muscle, medially by pelvic diaphragm and the anal canal.
- Anal canal
- Pudendal canal, which contains internal pudendal artery and the pudendal nerve

This page titled 26.5G: Perineum is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.5H: Mammary Glands

A mammary gland is an organ in female mammals that produces milk to feed young offspring.

Learning Objectives

- Describe the function and structure of mammary glands

Key Points

- Mammary glands are not associated with the female reproductive tract, but develop as secondary sex characteristics in reproductive-age females.
- The basic components of a mature mammary gland are the alveoli, hollow cavities, a few millimeters large lined with milk-secreting cuboidal cells and surrounded by myoepithelial cells.
- Alveoli join up to form groups known as lobules, and each of which has a lactiferous duct that drains into openings in the nipple.
- Secretory alveoli develop mainly in pregnancy, when rising levels of prolactin, estrogen, and progesterone cause further branching, together with an increase in adipose tissue and a richer blood flow.


Key Terms

- **Wnts**: Morphogenic signaling proteins that regulate cell-cell interactions.
- **beta-1 integrin**: One of the regulators of mammary epithelial cell growth and differentiation.
- **mammary gland**: A gland that secretes milk for suckling an infant or offspring.
- **lactiferous duct**: The components that form a branched system connecting the lobules of the mammary gland to the tip of the nipple.

A mammary gland is an organ in female mammals that produces milk to feed young offspring.

Anatomy of the Mammary Gland

The basic components of a mature mammary gland are the alveoli, hollow cavities, a few millimeters large lined with milk-secreting cuboidal cells and surrounded by myoepithelial cells. These alveoli join to form groups known as lobules, and each lobule has a lactiferous duct that drains into openings in the nipple. The myoepithelial cells can contract under the stimulation of oxytocin, excreting milk secreted from alveolar units into the lobule lumen toward the nipple where it collects in sinuses of the ducts. As the infant begins to suck, the hormonally (oxytocin) mediated “let-down reflex” ensues, and the mother’s milk is secreted into the baby’s mouth.



Mammary Gland: Cross-section of the mammary-gland. 1. Chest wall 2. Pectoralis muscles 3. Lobules 4. Nipple 5. Areola 6. Milk duct 7. Fatty tissue 8. Skin EndFragment

All the milk-secreting tissue leading to a single lactiferous duct is called a simple mammary gland; a complex mammary gland is all the simple mammary glands serving one nipple. Humans normally have two complex mammary glands, one in each breast, and each complex mammary gland consists of 10–20 simple glands. The presence of more than two nipples is known as polythelia, and the presence of more than two complex mammary glands as polymastia.

Development of the Mammary Glands

Mammary glands develop during different growth cycles. They exist in both sexes during the embryonic stage, forming only a rudimentary duct tree at birth. In this stage, mammary gland development depends on systemic (and maternal) hormones, but is also under the local regulation of paracrine communication between neighboring epithelial and mesenchymal cells by parathyroid hormone-related protein. This locally-secreted factor gives rise to a series of outside-in and inside-out positive feedback between these two types of cells, so that mammary bud epithelial cells can proliferate and sprout down into the mesenchymal layer until they reach the fat pad to begin the first round of branching.

Lactiferous duct development occurs in females in response to circulating hormones, first during pre- and postnatal stages and later during puberty. Estrogen promotes branching differentiation, which is inhibited by testosterone in males. A mature duct tree reaching the limit of the fat pad of the mammary gland is formed by bifurcation of duct terminal end buds, secondary branches sprouting from primary ducts and proper duct lumen formation.

The Process of Milk Production

Secretory alveoli develop mainly in pregnancy, when rising levels of prolactin, estrogen, and progesterone cause further branching, together with an increase in adipose tissue and a richer blood flow. In gestation, serum progesterone remains at a high concentration so signaling through its receptor is continuously activated. As one of the transcribed genes, Wnts secreted from mammary epithelial cells act paracrinely to induce branching of neighboring cells. When the lactiferous duct tree is almost ready, alveoli are differentiated from luminal epithelial cells and added at the end of each branch. In late pregnancy and for the first few days after giving birth, colostrum is secreted.

Milk secretion (lactation) begins a few days after birth, caused by reduction in circulating progesterone and the presence of prolactin, which mediates further alveologenesis and milk protein production and regulates osmotic balance and tight junction function.
The binding of laminin and collagen in the myoepithelial basement membrane with beta-1 integrin on the epithelial surface insures correct placement of prolactin receptors on basal lateral side of alveoli cells and directional secretion of milk into lactiferous ducts. Suckling of the baby causes release of hormone oxytocin which stimulates contraction of the myoepithelial cells.

With combined control from the extracellular matrix (ECM) and systemic hormones, milk secretion can be reciprocally amplified to provide enough nutrition for the baby.

During weaning, decreased prolactin, lack of mechanical stimulation through suckling, and changes in osmotic balance caused by milk stasis and leaking of tight junctions cause cessation of milk production. In some species there is complete or partial involution of alveolar structures after weaning; however, in humans there is only partial involution, which widely varies among individuals. Shrinkage of the mammary duct tree and ECM remodeling by various proteinase is under the control of somatostatin and other growth-inhibiting hormones and local factors. This structure change leads loose fat tissue to fill the empty space. However, a functional lactiferous duct tree can be reformed when a female is pregnant again.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Female Reproductive System. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Female_...ductive_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ovary. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ovary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- oogenesis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/oogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vulva. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vulva**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- oviduct. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/oviduct**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...oSystem_02.png**. **License**: ***CC BY: Attribution***
- Corpus Luteum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Corpus_luteum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ovarian follicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ovarian_follicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ovary. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ovary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ovary. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ovary**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...oSystem_02.png**. **License**: ***CC BY: Attribution***
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F..._PID-Sites.png**. **License**: ***CC BY: Attribution***
- Linea Terminalis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Linea_terminalis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Uterus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Uterus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fundus (uterus). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fundus_(uterus)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- uterus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/uterus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endometrium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endometrium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fallopian tubes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fallopian%20tubes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...oSystem_02.png**. **License**: ***CC BY: Attribution***
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F..._PID-Sites.png**. **License**: ***CC BY: Attribution***
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/F..._PID-Sites.png**. **License**: ***CC BY: Attribution***
- Uterus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Uterus**. **License**: ***Public Domain: No Known Copyright***
- Fallopian tube. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fallopian_tube**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- oviduct. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/oviduct. **License**: *CC BY-SA: Attribution-ShareAlike*
- fallopian tubes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/fallopian%20tubes. **License**: *CC BY-SA: Attribution-ShareAlike*
- ovarian follicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/ovarian%20follicle. **License**: *CC BY-SA: Attribution-ShareAlike*
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...oSystem_02.png. **License**: *CC BY: Attribution*
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F..._PID-Sites.png. **License**: *CC BY: Attribution*
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F..._PID-Sites.png. **License**: *CC BY: Attribution*
- Uterus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Uterus. **License**: *Public Domain: No Known Copyright*
- Ovaries, Uterine Tubes, and Uterus. **Provided by**: OpenStax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Skene Gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Skene%27s_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vagina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vagina. **License**: *CC BY-SA: Attribution-ShareAlike*
- vulva. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vulva. **License**: *CC BY-SA: Attribution-ShareAlike*
- clitoris. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/clitoris. **License**: *CC BY-SA: Attribution-ShareAlike*
- vagina. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vagina. **License**: *CC BY-SA: Attribution-ShareAlike*
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0400_FemaleReproSystem_02.png. **License**: *CC BY: Attribution*
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- Uterus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Uterus. **License**: *Public Domain: No Known Copyright*
- Ovaries, Uterine Tubes, and Uterus. **Provided by**: OpenStax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Scheme female reproductive system-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_female_reproductive_system-en.svg. **License**: *Public Domain: No Known Copyright*
- Vulva. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vulva. **License**: *CC BY-SA: Attribution-ShareAlike*
- vulva. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vulva. **License**: *CC BY-SA: Attribution-ShareAlike*
- labia minora. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/labia_minora. **License**: *CC BY-SA: Attribution-ShareAlike*
- labia majora. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/labia_majora. **License**: *CC BY-SA: Attribution-ShareAlike*
- mons pubis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mons_pubis. **License**: *CC BY-SA: Attribution-ShareAlike*
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0400_FemaleReproSystem_02.png. **License**: *CC BY: Attribution*
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- Uterus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Uterus. **License**: *Public Domain: No Known Copyright*
- Ovaries, Uterine Tubes, and Uterus. **Provided by**: OpenStax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Scheme female reproductive system-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_female_reproductive_system-en.svg. **License**: *Public Domain: No Known Copyright*
- File:Figure_28_02_02. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_02.jpg. **License**: *CC BY: Attribution*
- Perineum. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Perineum. **License**: *CC BY-SA: Attribution-ShareAlike*

- lower rabbus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lower%20rabbus. **License**: *CC BY-SA: Attribution-ShareAlike*
- perineal body. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/perineal%20body. **License**: *CC BY-SA: Attribution-ShareAlike*
- episiotomy. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/episiotomy. **License**: *CC BY-SA: Attribution-ShareAlike*
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0400_FemaleReproSystem_02.png. **License**: *CC BY: Attribution*
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- Uterus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Uterus. **License**: *Public Domain: No Known Copyright*
- Ovaries, Uterine Tubes, and Uterus. **Provided by**: OpenStax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Scheme female reproductive system-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_female_reproductive_system-en.svg. **License**: *Public Domain: No Known Copyright*
- File:Figure_28_02_02. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_02.jpg. **License**: *CC BY: Attribution*
- File:Gray408. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Gray408.png. **License**: *Public Domain: No Known Copyright*
- Mammary gland. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mammary_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- lactiferous duct. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lactiferous%20duct. **License**: *CC BY-SA: Attribution-ShareAlike*
- mammary gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mammary_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- FemaleReproSystem. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0400_FemaleReproSystem_02.png. **License**: *CC BY: Attribution*
- Blausen_0732_PID-Sites. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- /File:Blausen_0732_PID-Sites. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Blausen_0732_PID-Sites.png. **License**: *CC BY: Attribution*
- Uterus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Uterus. **License**: *Public Domain: No Known Copyright*
- Ovaries, Uterine Tubes, and Uterus. **Provided by**: OpenStax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Scheme female reproductive system-en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Scheme_female_reproductive_system-en.svg. **License**: *Public Domain: No Known Copyright*
- File:Figure_28_02_02. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_02.jpg. **License**: *CC BY: Attribution*
- File:Gray408. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Gray408.png. **License**: *Public Domain: No Known Copyright*
- File:Breast_anatomy_normal_scheme. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Breast_anatomy_normal_scheme.png. **License**: *CC BY: Attribution*

This page titled 26.5H: Mammary Glands is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6: PHYSIOLOGY OF THE FEMALE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

26.6A: Oogenesis

26.6B: Ovarian Cycle

26.6C: Uterine (Menstrual) Cycle

26.6D: Hormonal Regulation of the Female Reproductive Cycle

26.6E: Extrauterine Effects of Estrogens and Progesterone

26.6F: Female Sexual Response

This page titled 26.6: Physiology of the Female Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6A: Oogenesis

Oogenesis is the maturation of the female gametes through meiotic division.

Learning Objectives

- Differentiate among the stages of oogenesis and the actions of hormones during each stage

Key Points

- Oogenesis starts with the process of developing oogonia via the transformation of primordial follicles into primary oocytes, a process called oocytogenesis.
- Oocytogenesis is complete either before or shortly after birth in humans. During the menstrual cycle primary oocytes complete maturation through further meiotic divisions.
- The first meiotic division is coordinated by hormones: follicle stimulating hormone (FSH), estrogen, luteinizing hormone (LH), and progesterone.
- The oocyte is arrested in cell division prior to the second meiotic division, which only occurs after fertilization.

Key Terms

- **polar body**: A small haploid cell formed concomitantly as an egg cell during oogenesis, but which does not have the ability to be fertilized.
- **ootid**: A nearly mature ovum that results from meiotic division of a secondary oocyte.
- **lutenizing hormone**: A hormone produced by the anterior pituitary gland. In females, an acute rise of LH triggers ovulation and development of the corpus luteum.
- **follicle**: A small cavity or sac that is the site of oocyte development in the ovary.
- **follicle stimulating hormone**: A gonadotropic glycoprotein hormone, secreted in part of the pituitary gland that stimulates the growth of ovarian follicles in female mammals and induces spermatogenesis in male mammals.

The menstrual cycle begins with the maturation of oocytes through the process of oogenesis, as well as concurrent follicle development that stimulates ovulation. Oogenesis starts with the process of developing oogonia via the transformation of primordial follicles into primary oocytes, a process called oocytogenesis. Oocytogenesis is complete either before or shortly after birth in humans. During the menstrual cycle primary oocytes complete maturation through further meiotic divisions.


Follicle development signals the beginning of the menstrual cycle. At the start of the menstrual cycle, some 12-20 primary follicles begin to develop under the influence of elevated levels of follicle-stimulating hormone (FSH) to form secondary follicles. The primary follicles form from primordial follicles, which develop in the ovary as a fetus during conception and are arrested in the prophase state of the cellular cycle.

By around day 9 of the menstrual cycle, only one healthy secondary follicle remain. The rest are reabsorbed into the ovary. The remaining follicle, called the dominant follicle, is responsible for producing large amounts of estrogen during the late follicular phase.

On day 14 of the cycle, a luteinizing hormone surge is triggered by the positive feedback of estrogen. This causes the secondary follicle to develop into a tertiary follicle, which then leaves the ovary 24–36 hours later. An important event in the development of the tertiary follicle occurs when the primary oocyte completes the first meiotic division, resulting in the formation of a polar body and a secondary oocyte. The empty follicle then forms a corpus luteum which later releases progesterone to maintain a potential pregnancy.

Immediately after meiosis I, the haploid secondary oocyte initiates meiosis II. However, this process is also halted at the metaphase II stage until fertilization occurs. When meiosis II has completed, an ootid and another polar body is created.

Both polar bodies disintegrate at the end of meiosis II, leaving only the ootid, which eventually develops into a mature ovum. The formation of polar bodies serves to discard the extra haploid sets of chromosomes that have resulted as a consequence of meiosis.



The Oogenesis Process: Diagram showing maturation of the ovum.

This page titled 26.6A: Oogenesis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6B: Ovarian Cycle

The menstrual cycle is the physiological process that fertile women undergo for the purposes of reproduction and fertilization.

Learning Objectives

- Differentiate among the phases of the menstrual cycle

Key Points

- The ovarian cycle refers to the series of changes in the ovary during which the follicle matures, the ovum is shed, and the corpus luteum develops.
- The follicular phase describes the development of the follicle in response to follicle stimulation hormone (FSH). As luteinizing hormone (LH) and FSH levels increase they stimulate ovulation, or the release of a mature oocyte into the fallopian tubes.
- In the luteal phase, the corpus luteum forms on the ovary and secretes many hormones, most significantly progesterone, which makes the endometrium of the uterus ready for implantation of an embryo.
- If implantation does not occur, the corpus luteum will be degraded, resulting in menstruation.
- If implantation occurs the corpus luteum is maintained.

Key Terms

- **ischemic phase**: The final part of the secretory phase. The endometrium becomes pale and arteries constrict due to lower hormone release by the disintegrating corpus luteum.
- **granulosa cells**: These cells produce hormones and growth factors that interact with the oocyte during its development.
- **menstrual cycle**: The recurring cycle of physiological changes in the females of some animal species that is associated with reproductive fertility.
- **luteal phase**: The latter part of the menstrual cycle that occurs after ovulation, in which the corpus luteum secretes progesterone to prepare the endometrium for the implantation of an embryo.
- **follicular phase**: The phase of the estrous cycle that involves follicular maturation within the ovary and, controlled by the hormone estradiol.

The menstrual cycle is the scientific term for the physiological changes that occur in fertile women for the purpose of sexual reproduction.The menstrual cycle is controlled by the endocrine system and commonly divided into three phases: the follicular phase, ovulation, and the luteal phase. However, some sources define these phases as menstruation, proliferative phase, and secretory phase. Menstrual cycles are counted from the first day of menstrual bleeding.

The Follicular Phase

The follicular phase (or proliferative phase) is the phase of the menstrual cycle in humans and great apes during which follicles in the ovary mature, ending with ovulation. The main hormone controlling this stage is estradiol. During the follicular phase, follicle-stimulating hormone (FSH) is secreted by the anterior pituitary gland. FSH levels begin to rise in the last few days of the previous menstrual cycle and peak during the first week of the follicular phase. The rise in FSH levels recruits five to seven tertiary-stage ovarian follicles (also known as Graafian or antral follicles) for entry into the menstrual cycle. These follicles compete with each other for dominance.

FSH induces the proliferation of granulosa cells in the developing follicles and the expression of luteinizing hormone (LH) receptors on these granulosa cells. Two or three days before LH levels begin to increase, usually by day seven of the cycle, one or occasionally two of the recruited follicles emerges as dominant. Many endocrinologists believe that estrogen secretion of the

dominant follicle increases to a level that indirectly lowers the levels of LH and FSH. This slowdown in LH and FSH production leads to the atresia (death) of most of the recruited follicles, though the dominant follicle continues to mature.

These high estrogen levels initiate the formation of a new layer of endometrium in the uterus. Crypts in the cervix are also stimulated to produce fertile cervical mucus that reduces the acidity of the vagina, creating a more hospitable environment for sperm. In addition, basal body temperature may lower slightly under the influence of high estrogen levels. Ovulation normally occurs 36 (± 2) hours after the beginning of the LH surge (when LH is first detectable in urine).

Ovulation


Ovulation is the phase in which a mature ovarian follicle ruptures and discharges an ovum (also known as an oocyte, female gamete, or egg). Ovulation also occurs in the estrous cycle of other female mammals, which differs in many fundamental ways from the menstrual cycle. The time immediately surrounding ovulation is referred to as the ovulatory phase or the periovulatory period.

The Luteal Phase

The luteal phase (or secretory phase) is the latter part of the menstrual or estrous cycle. It begins with the formation of the corpus luteum and ends in either pregnancy or luteolysis. The main hormone associated with this stage is progesterone, which is significantly higher during the luteal phase than in other phases of the cycle. Some sources define the end of the luteal phase as a distinct ischemic phase.

After ovulation, the pituitary hormones FSH and LH cause the remaining parts of the dominant follicle to transform into the corpus luteum. It continues to grow for some time after ovulation and produces significant amounts of hormones, particularly progesterone, and to a lesser extent, estrogen. Progesterone plays a vital role in making the endometrium receptive to implantation of the blastocyst and supportive of the early pregnancy. It also raises the woman's basal body temperature. The hormones produced by the corpus luteum suppress production of the FSH and LH, causing the corpus luteum will atrophy. The death of the corpus luteum results in falling levels of progesterone and estrogen. This in turn causes increased levels of FSH, leading to recruitment of follicles for the next cycle. Continued drops in estrogen and progesterone levels trigger the end of the luteal phase, menstruation, and the beginning of the next cycle.

The loss of the corpus luteum can be prevented by implantation of an embryo. After implantation, human embryos produce human chorionic gonadotropin (hCG), which is structurally similar to LH and can preserve the corpus luteum. Because the hormone is unique to the embryo, most pregnancy tests look for the presence of hCG. If implantation occurs, the corpus luteum will continue to produce progesterone (and maintain high basal body temperatures) for eight to twelve weeks, after which the placenta takes over this function.



The ovarian cycle: The ovarian cycle is the series of changes that occur in the ovary during the menstrual cycle that cause maturation of a follicle, ovulation, and development of the corpus luteum.

This page titled 26.6B: Ovarian Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6C: Uterine (Menstrual) Cycle

The uterine cycle describes a series of changes that occur to the lining of the uterus, or endometrium, during a typical menstrual cycle.

Learning Objectives

- Outline the process of the uterine cycle

Key Points

- The uterine cycle includes the increase in the endometrium in preparation for implantation and the shedding of the lining following lack of implantation, termed menstruation.
- Menstrual cycles are counted from the first day of menstrual bleeding.
- Endometrial thickening is stimulated by the increasing amount of estrogen in the follicular phase.
- If implantation does not occur, progesterone and estrogen levels drop, which stimulates menstruation.

Key Terms


- **endometrium**: The mucous membrane that lines the uterus in mammals, in which fertilized eggs are implanted.
- **menstrual cycle**: In the females of some animal species, the recurring cycle of physiological changes associated with reproductive fertility.
- **decidua**: A mucous membrane that lines the uterus: it is shed during menstruation and modified during pregnancy.

Several changes to the uterine lining (endometrium) occur during the menstrual cycle, also called the uterine cycle. The endometrium is the innermost glandular layer of the uterus. During the menstrual cycle, the endometrium grows to a thick, blood vessel-rich tissue lining, representing an optimal environment for the implantation of a blastocyst upon its arrival in the uterus. Menstrual cycles are counted from the first day of menstrual bleeding and are typically 28 days long.

During menstruation, the body begins to prepare for ovulation again. The levels of estrogen gradually rise, signalling the start of the follicular, or proliferation, phase of the menstrual cycle. The discharge of blood slows and then stops in response to rising hormone levels and the lining of the uterus thickens, or proliferates. Ovulation is triggered by a surge in luteinizing hormone. The sudden change in hormones at the time of ovulation sometimes causes minor changes in the endometrium and light midcycle blood flow.

After ovulation, under the influence of progesterone, the endometrium changes to a secretory lining in preparation for the potential implantation of an embryo to establish a pregnancy. If a blastocyst implants, then the lining remains as the decidua. This becomes part of the placenta and provides support and protection for the embryo during gestation.

If implantation does not occur within approximately two weeks, the progesterone-producing corpus luteum in the ovary will recede, causing sharp drops in levels of both progesterone and estrogen. This hormone decrease causes the uterus to shed its lining and the egg in menstruation. The cessation of menstrual cycles at the end of a woman's reproductive period is termed menopause. The average age of menopause in women is 52 years, but it can occur anytime between 45 and 55.



The Uterine Cycle: High estrogen and progesterone levels stimulate increased endometrial thickness, but following their decline from a lack of implantation, the endometrium is shed and menstruation occurs.

Normal menstrual flow can occur although ovulation does not occur. This is referred to as an anovulatory cycle. Follicular development may start but not be completed although estrogen will still stimulate the uterine lining. Anovulatory flow that results from a very thick endometrium caused by prolonged, continued high estrogen levels is called estrogen breakthrough bleeding.However, if it is triggered by a sudden drop in estrogen levels, it is called withdrawal bleeding. Anovulatory cycles commonly occur before menopause and in women with polycystic ovary syndrome.

This page titled 26.6C: Uterine (Menstrual) Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6D: Hormonal Regulation of the Female Reproductive Cycle

The menstrual cycle is controlled by a series of changes in hormone levels, primarily estrogen and progesterone.

LEARNING OBJECTIVES

Differentiate among the phases of the menstrual cycle

KEY TAKEAWAYS


Key Points

- The follicular phase begins with an increase in follicle -stimulation hormone (FSH), which causes increases in luteinizing hormone (LH) and gonadotropin-releasing hormone (GnRH). Concurrent increases in estrogen levels cause increases in progesterone, stimulating proliferation of the endometrium.
- A spike in LH and FSH (“LH surge”) causes ovulation, following a suppression of GnRH.
- Estrogen levels continue to rise following ovulation and the corpus luteum forms, which secretes progesterone in significant levels and causes decreases in LH and FSH levels.
- Without implantation, estrogen and progesterone levels will fall and the corpus luteum will degrade.

Key Terms

- **estrogen**: A hormone responsible for the appearance of secondary sex characteristics of human females at puberty and the maturation and maintenance of the reproductive organs in their functional state.
- **cumulus**: A cluster of cells that surround the oocyte both in the ovarian follicle and after ovulation. These cells coordinate follicular development and oocyte maturation.
- **progesterone**: A steroid hormone secreted by the ovaries that prepares the uterus for the implantation of a fertilized ovum and subsequent pregnancy.
- **theca cells**: A group of endocrine cells in the ovary made up of connective tissue surrounding the follicle. They provide androgen synthesis and signal transduction between granulosa cells and oocytes during development.
- **luteinizing hormone surge**: Acute rise of LH levels that triggers ovulation and development of the corpus luteum.

The menstrual cycle is the physiological change that occurs under the control of the endocrine system in fertile women for the purposes of sexual reproduction and fertilization.



The Menstrual Cycle: The menstrual cycle is controlled by the endocrine system, with distinct phases correlated to changes in hormone concentrations.

Phases of the Menstrual Cycle

The menstrual cycle is divided into three stages: follicular phase, ovulation, and the luteal phase.

Follicular Phase

During the follicular phase (or proliferative phase), follicles in the ovary mature under the control of estradiol. Follicle-stimulating hormone (FSH) is secreted by the anterior pituitary gland beginning in the last few days of the previous menstrual cycle. Levels of FSH peak during the first week of the follicular phase. The rise in FSH recruits tertiary-stage ovarian follicles (antral follicles) for entry into the menstrual cycle.

Follicle-stimulating hormone induces the proliferation of granulosa cells in the developing follicles and the expression of luteinizing hormone (LH) receptors on these cells. Under the influence of FSH, granulosa cells begin estrogen secretion. This increased level of estrogen stimulates production of gonadotropin-releasing hormone (GnRH), which increases production of LH. LH induces androgen synthesis by theca cells, stimulates proliferation and differentiation, and increases LH receptor expression on granulosa cells.

Throughout the entire follicular phase, rising estrogen levels in the blood stimulate growth of the endometrium and myometrium of the uterus. This also causes endometrial cells to produce receptors for progesterone, which helps prime the endometrium to the late proliferative phase and the luteal phase. Two or three days before LH levels begin to increase, one or occasionally two of the recruited follicles emerge as dominant. Many endocrinologists believe that the estrogen secretion of the dominant follicle lowers the levels of LH and FSH, leading to the atresia (death) of most of the other recruited follicles. Estrogen levels will continue to increase for several days.

High estrogen levels initiate the formation of a new layer of endometrium in the uterus, the proliferative endometrium. Crypts in the cervix are stimulated to produce fertile cervical mucus that reduces the acidity of the vagina, creating a more hospitable environment for sperm. In addition, basal body temperature may lower slightly under the influence of high estrogen levels.

Ovulation

Estrogen levels are highest right before the LH surge begins. The short-term drop in steroid hormones between the beginning of the LH surge and ovulation may cause mid-cycle spotting or bleeding. Under the influence of the preovulatory LH surge, the first meiotic division of the oocytes is completed. The surge also initiates luteinization of theca and granulosa cells. Ovulation normally occurs 30 (± 2) hours after the beginning of the LH surge.

Ovulation is the process in a female's menstrual cycle by which a mature ovarian follicle ruptures and discharges an ovum (oocyte). The time immediately surrounding ovulation is referred to as the ovulatory phase or the periovulatory period. In the preovulatory phase of the menstrual cycle, the ovarian follicle undergoes cumulus expansion stimulated by FSH. The ovum then leaves the follicle through the formed stigma. Ovulation is triggered by a spike in the amount of FSH and LH released from the pituitary gland.

Luteal Phase

The luteal phase begins with the formation of the corpus luteum stimulated by FSH and LH and ends in either pregnancy or luteolysis. The main hormone associated with this stage is progesterone, which is produced by the growing corpus luteum and is significantly higher during the luteal phase than other phases of the cycle. Progesterone plays a vital role in making the endometrium receptive to implantation of the blastocyst and supportive of the early pregnancy. It also raises the woman's basal body temperature.

Several days after ovulation, the increasing amount of estrogen produced by the corpus luteum may cause one or two days of fertile cervical mucus, lower basal body temperatures, or both. This is known as a secondary estrogen surge. The hormones produced by the corpus luteum suppress production of the FSH and LH, which leads to its atrophy. The death of the corpus luteum results in falling levels of progesterone and estrogen, which triggers the end of the luteal phase. Increased levels of FSH start recruiting follicles for the next cycle.

Alternatively, the loss of the corpus luteum can be prevented by implantation of an embryo: after implantation, human embryos produce human chorionic gonadotropin (hCG). Human chorionic gonadotropin is structurally similar to LH and can preserve the corpus luteum. Because the hormone is unique to the embryo, most pregnancy tests look for the presence of hCG. If implantation occurs, the corpus luteum will continue to produce progesterone (and maintain high basal body temperatures) for eight to 12 weeks, after which the placenta takes over this function.

This page titled 26.6D: Hormonal Regulation of the Female Reproductive Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6E: Extrauterine Effects of Estrogens and Progesterone

Estrogen and progesterone have several effects beyond their immediate roles in the menstrual cycle, pregnancy, and labor.

Learning Objectives

- Describe the functions of progesterone and estrogen

Key Points


- Estrogen promotes female secondary sex characteristics and has structural and metabolic functions, including bone formation, salt and water retention, and increased cortisol levels.
- The effects of progesterone are largely amplified by estrogen and include increases in core temperature, anti-inflammatory effects, increased use of fat for energy, and regulation of insulin release.
- Progesterone is sometimes called the " hormone of pregnancy" as it has many roles relating to the fetal development, including preparing the uterus for implantation, increasing the amount of cervical mucus, and decreasing the maternal immune response.

Key Terms

- **estrogen**: A hormone responsible for the appearance of secondary sex characteristics of human females at puberty and for the maturation and maintenance of the reproductive organs in their mature functional state.
- **aldosterone**: A mineralocorticoid hormone secreted by the adrenal cortex that regulates the balance of sodium and potassium in the body.
- **progesterone**: A steroid hormone secreted by the ovaries that prepares the uterus for the implantation of a fertilized ovum and subsequent pregnancy.

Both estrogens and progesterone serve functions in the body beyond their roles in menstruation, pregnancy, and childbirth.

Estrogens Overview




Estradiol: One of the estrogens produced in the human body, predominant during a woman's reproductive years.

Estrogens are a group of compounds named for their importance in the estrous cycle of humans and other animals. They are the primary female sex hormones, although they are found in males as well. The three major naturally occurring forms of estrogen in women are estrone (E1), estradiol (E2), and estriol (E3). Estetrol (E4) is produced only during pregnancy.

Natural estrogens are steroid hormones, while some synthetic versions are non-steroidal. Estrogens are synthesized in all

vertebrates as well as some insects, and their presence in both suggests that they have an ancient evolutionary history. Like all steroid hormones, estrogen readily diffuses across the cell membrane. Once inside the cell, it binds to and activates estrogen receptors which in turn modulate the expression of many genes.

Functions of Estrogens



Estriol: Another one of the three main estrogens produced in humans.

While estrogens are present in both men and women, they are usually at significantly higher levels in women of reproductive age. They promote the development of female secondary sexual characteristics, such as breasts, pubic hair, and female fat distribution. They are also involved in the thickening of the endometrium and other aspects of menstrual cycle regulation.

Other functions of and structural changes induced by estrogen include:

- Formation of female secondary sex characteristics
- Accelerating metabolism
- Increasing fat stores
- Stimulating endometrial growth
- Increasing uterine growth
- Increasing vaginal lubrication
- Thickening the vaginal wall
- Maintaining blood vessels and skin
- Reducing bone resorption, increasing bone formation
- Reducing muscle mass


Effect on Libido

Sex drive is dependent on androgen levels only in the presence of estrogen. Without estrogen, free testosterone levels actually decrease sexual desire, as demonstrated in women who have hypoactive sexual desire disorder. The sexual desire in these women can be restored by administration of estrogen through oral contraceptives.

Mental Health

Estrogen plays a significant role in women's mental health. Sudden estrogen withdrawal, fluctuating estrogen, and periods of sustained low levels of estrogen correlate with significant mood changes. Restoration or stabilization of estrogen levels is clinically effective for recovery from postpartum, perimenopause, and postmenopause depression.

Progesterone Overview



Progesterone: Belongs to the progestogen class of hormones and is the predominant example in the human body.

Progesterone is a steroid hormone involved in the female menstrual cycle, pregnancy (supports gestation), and embryogenesis of humans and other species.

Progesterone belongs to a class of hormones called progestogens and is the major naturally-occurring human form in this category. Progesterone exerts its primary action through the intracellular progesterone receptor, although a distinct, membrane-bound progesterone receptor has also been postulated.

Functions of Progesterone

Progesterone has a number of physiological effects that are amplified in the presence of estrogen. Estrogen, through estrogen receptors, upregulates the expression of progesterone receptors. Also, elevated levels of progesterone potently reduce the sodium-retaining activity of aldosterone, resulting in natriuresis and a reduction in extracellular fluid volume. Progesterone withdrawal, on the other hand, is associated with a temporary increase in sodium retention (reduced natriuresis, with an increase in extracellular fluid volume) due to the compensatory increase in aldosterone production. This combats the blockade of the mineralocorticoid receptor by the previously-elevated level of progesterone.

Progesterone has key effects via non-genomic signalling on human sperm as they migrate through the female tract before fertilization occurs, though the receptor(s) as yet remain unidentified. Detailed characterization of the events occurring in sperm in response to progesterone has shed light on intracellular calcium transients, maintained changes, and slow calcium oscillations, now thought to possibly regulate motility.

Progesterone is sometimes called the “hormone of pregnancy” and has many roles relating to fetal development. It converts the endometrium to its secretory stage to prepare the uterus for implantation. At the same time, it affects the vaginal epithelium and cervical mucus, making them thick and impenetrable to sperm.

If pregnancy does not occur, progesterone levels will decrease, leading to menstruation. Normal menstrual bleeding is progesterone-withdrawal bleeding. If ovulation does not occur and the corpus luteum does not develop, its levels may be low, leading to anovulatory dysfunctional uterine bleeding. During implantation and gestation, progesterone appears to decrease the maternal immune response to allow for the acceptance of the pregnancy and decrease contractility of the uterine smooth muscle.

In addition, progesterone inhibits lactation during pregnancy. A drop in its levels is facilitates the onset of labor. Another drop following delivery is one of the triggers for milk production. The fetus metabolizes placental progesterone in the production of adrenal steroids.

This page titled 26.6E: Extrauterine Effects of Estrogens and Progesterone is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

26.6F: Female Sexual Response

Female sexual arousal causes physiological changes including increased blood flow to the genitals and enlargement and lubrication of the vagina.

Learning Objectives

- Explain the process of female sexual response

Key Points


- Responses specific to females include engorgement of several tissues including the nipples, vulva, clitoris, and vaginal walls.
- Further changes include changes to the shape of the vagina and the positioning of the uterus in the pelvis, as well as increases in blood pressure and heart rate.
- Experienced by males and females, orgasms are controlled by the involuntary or autonomic nervous system.
- Age-related changes in sexual responsiveness may be associated with changes in estrogen.

Key Terms

- **vulva**: The vaginal opening to the uterus.
- **estrogen**: A hormone responsible for the appearance of secondary sex characteristics of females at puberty and for the maturation and maintenance of the reproductive organs in their mature functional state.
- **clitoris**: A small, sensitive, and elongated erectile organ at the anterior part of the vulva in females and homologous with the penis.

Sexual arousal is caused by sexual desire during or in anticipation of sexual activity. A number of physiological changes occur in the body and mind in preparation for sex and continue during the act. For women, these changes include increased blood flow to the nipples, vulva, clitoris, and vaginal walls, and increased vaginal lubrication.

Physiological Response



Features of the vulva: The clitoris and labial folds are labelled.

The beginnings of sexual arousal in a woman's body is usually marked by vaginal lubrication, engorgement of the external genitals, and internal enlargement of the vagina. Further stimulation can lead to more vaginal wetness and further engorgement and swelling of the clitoris and the labia, along with increased redness or darkening of the skin in these areas. Changes also occur to the internal shape of the vagina and to the position of the uterus within the pelvis.

Other bodily changes include an increase in heart rate and blood pressure, as well as flushing across the chest and upper body. If sexual stimulation continues, then sexual arousal may peak into orgasm, resulting in rhythmic muscular contractions in the pelvic region characterized by an intense sensation of pleasure. Experienced by males and females, orgasms are controlled by the involuntary or autonomic nervous system.

As women age, estrogen levels decrease. Reduced estrogen levels may be associated with increased vaginal dryness and less clitoral erection when aroused, but are not directly related to other aspects of sexual interest or arousal. In older women, decreased pelvic muscle tone may prolong the time to reach orgasm, diminish the intensity of orgasms, and cause more rapid resolution. In some women, the uterine contracts that occur during orgasm may cause pain or discomfort.

Psychological Response

Mental and physical stimuli such as touch and the internal fluctuation of hormones influence sexual arousal. Cognitive factors like sexual motivation, perceived gender role expectations, and sexual attitudes play important roles in women's self-reported levels of sexual arousal. Basson suggests that women's need for intimacy prompts them to engage with sexual stimuli, leading to an experience of sexual desire and psychological sexual arousal.

Research by Goldey and van Anders showed that sexual cognition impacts hormone levels in women. For instance, sexual thoughts result in a rapid increase in testosterone in women who were not using hormonal contraception. Inconsistent study results indicate that, although testosterone is involved in libido and sexuality of some women, its effects can be obscured by the coexistence of psychological factors in others.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Immature Ovum. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Immature_ovum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Oogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Oogenes...uman_Oogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lutenizing hormone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lutenizing%20hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- follicle. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/follicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- follicle stimulating hormone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/follicle_stimulating_hormone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/F...e_28_02_03.JPG**. **License**: ***CC BY: Attribution***
- Granulosa Cell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Granulosa_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Luteal phase. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Luteal_phase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ovulation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ovulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Follicular phase. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Follicular_phase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Menstrual cycle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Menstrual_cycle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Anatomy and Physiology of Animals/Reproductive System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/**

Anatomy..._Ovarian_Cycle. License: CC BY-SA: Attribution-ShareAlike
- menstrual cycle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/menstrual_cycle. **License**: CC BY-SA: Attribution-ShareAlike
- follicular phase. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/follicular_phase. **License**: CC BY-SA: Attribution-ShareAlike
- luteal phase. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/luteal_phase. **License**: CC BY-SA: Attribution-ShareAlike
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F...e_28_02_03.JPG. **License**: CC BY: Attribution
- Anatomy and physiology of animals Ovarian cycle showing from top left clockwise. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._clockwise.jpg. **License**: CC BY-SA: Attribution-ShareAlike
- Endometrium. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endometrium. **License**: CC BY-SA: Attribution-ShareAlike
- Menstrual cycle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Menstrual_cycle. **License**: CC BY-SA: Attribution-ShareAlike
- decidua. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/decidua. **License**: CC BY-SA: Attribution-ShareAlike
- endometrium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/endometrium. **License**: CC BY-SA: Attribution-ShareAlike
- menstrual cycle. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/menstrual_cycle. **License**: CC BY-SA: Attribution-ShareAlike
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/F...e_28_02_03.JPG. **License**: CC BY: Attribution
- Anatomy and physiology of animals Ovarian cycle showing from top left clockwise. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._clockwise.jpg. **License**: CC BY-SA: Attribution-ShareAlike
- MenstrualCycle2 en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Me...lCycle2_en.svg. **License**: CC BY-SA: Attribution-ShareAlike
- Theca of Follicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Theca_of_follicle. **License**: CC BY-SA: Attribution-ShareAlike
- Cumulus Oophorus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cumulus_oophorus. **License**: CC BY-SA: Attribution-ShareAlike
- Luteal phase. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Luteal_phase. **License**: CC BY-SA: Attribution-ShareAlike
- Follicular phase. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Follicular_phase. **License**: CC BY-SA: Attribution-ShareAlike
- Menstrual cycle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Menstrual_cycle. **License**: CC BY-SA: Attribution-ShareAlike
- Ovulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Ovulation. **License**: CC BY-SA: Attribution-ShareAlike
- estrogen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/estrogen. **License**: CC BY-SA: Attribution-ShareAlike
- progesterone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/progesterone. **License**: CC BY-SA: Attribution-ShareAlike
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//physiology...-hormone-surge. **License**: CC BY-SA: Attribution-ShareAlike
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_03.JPG. **License**: CC BY: Attribution
- Anatomy and physiology of animals Ovarian cycle showing from top left clockwise. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi..._clockwise.jpg. **License**: CC BY-SA: Attribution-ShareAlike
- MenstrualCycle2 en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Me...lCycle2_en.svg. **License**: CC BY-SA: Attribution-ShareAlike
- Menstrual Cycle. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...le2_en.svg.png. **License**: CC BY-SA: Attribution-ShareAlike
- Progesterone. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Progesterone%23Effects. **License**: CC BY-SA: Attribution-ShareAlike
- Estrogen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Estrogen%23Function. **License**: CC BY-SA: Attribution-ShareAlike
- estrogen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/estrogen. **License**: CC BY-SA: Attribution-

ShareAlike
- progesterone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/progesterone. **License**: *CC BY-SA: Attribution-ShareAlike*
- aldosterone. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/aldosterone. **License**: *CC BY-SA: Attribution-ShareAlike*
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_03.JPG. **License**: *CC BY: Attribution*
- Anatomy and physiology of animals Ovarian cycle showing from top left clockwise. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Ovarian_cycle_showing_from_top_left_clockwise.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- MenstrualCycle2 en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Me...lCycle2_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Menstrual Cycle. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/2a/MenstrualCycle2_en.svg/1000px-MenstrualCycle2_en.svg.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Progesteron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Progesteron.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Estradiol2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Estradiol2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Estriol v2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Estriol_v2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sexual arousal. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sexual_arousal. **License**: *CC BY-SA: Attribution-ShareAlike*
- estrogen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/estrogen. **License**: *CC BY-SA: Attribution-ShareAlike*
- vulva. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vulva. **License**: *CC BY-SA: Attribution-ShareAlike*
- clitoris. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/clitoris. **License**: *CC BY-SA: Attribution-ShareAlike*
- File:Figure_28_02_03. **Provided by**: Wikipedia. **Located at**: commons.wikimedia.org/wiki/File:Figure_28_02_03.JPG. **License**: *CC BY: Attribution*
- Anatomy and physiology of animals Ovarian cycle showing from top left clockwise. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Anatomy_and_physiology_of_animals_Ovarian_cycle_showing_from_top_left_clockwise.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- MenstrualCycle2 en. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:MenstrualCycle2_en.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Menstrual Cycle. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/thumb/2/2a/MenstrualCycle2_en.svg/1000px-MenstrualCycle2_en.svg.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Progesteron. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Progesteron.svg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Estradiol2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Estradiol2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Estriol v2. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Estriol_v2.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Clitoris outer anatomy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Clitoris_outer_anatomy.gif. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 26.6F: Female Sexual Response is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27: HUMAN DEVELOPMENT AND PREGNANCY

- Topic hierarchy

27.10: Changes to the Mother's Body During Pregnancy

27.10A: Anatomical Changes

27.10B: Metabolic Changes

27.10C: Physiological Changes

27.10D: Exercise and Pregnancy

27.1: Overview of Pregnancy and Human Development

27.1A: Introduction to Pregnancy and Human Development

27.2: First Week of Development

27.2A: Fertilization

27.2B: Cleavage of the Zygote

27.2C: Blastocyst Formation

27.2D: Implantation

27.3: Second Week of Development

27.3A: Trophoblast Development

27.3B: Bilaminar Embryonic Disc Development

27.3C: Amnion Development

27.3D: Yolk Sac Development

27.3E: Sinusoid Development

27.3F: Development of the Extraembryonic Coelom

27.4: Third Week of Development

27.4A: Gastrulation

27.4B: Neurulation

27.4C: Somite Development

27.4D: Development of the Cardiovascular System

27.4E: Chorionic Villi and Placental Development

27.5: Fourth Week of Development

27.5A: Fourth Week of Development

27.6: Fifth Through Eighth Weeks of Development

27.6A: Fifth Through Eighth Weeks of Development

27.7: Birth (Parturition)

27.7A: Onset of Labor

27.7B: Stages of Labor

27.7C: Adjustments of the Infant at Birth

27.8: Lactation

27.8A: Physiology of Lactation

27.9: The Fetal Period

27.9A: Fetal Development

27.9B: Teratogens

27.9C: Prenatal Diagnostic Tests

This page titled 27: Human Development and Pregnancy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.10: CHANGES TO THE MOTHER'S BODY DURING PREGNANCY

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.10A: Anatomical Changes
27.10B: Metabolic Changes
27.10C: Physiological Changes
27.10D: Exercise and Pregnancy

This page titled 27.10: Changes to the Mother's Body During Pregnancy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.10A: Anatomical Changes

Women undergo many physical changes during pregnancy due to hormonal fluctuations and the need to accommodate a growing fetus.

Learning Objectives

- Describe the anatomic changes to a pregnant woman that occur over the course of the pregnancy

Key Points

- As the fetus grows and develops, several anatomical changes must occur to the female body to accommodate the growing fetus, including placental development, weight gain, abdominal extension, breast enlargement, glandular development, and posture changes.
- During the second trimester, morning sickness subsides, the uterus expands up to 20 times its normal size, breasts enlarge, and movements of the fetus may be felt.
- During the third trimester the fetus grows most rapidly and final weight gain occurs. The abdomen drops and fetal movement can become quite strong. The woman feels ready to give birth.

Key Terms

- **implantation**: The attachment of the fertilized ovum to the uterine wall.
- **decidua**: A mucous membrane that lines the uterus that is shed during menstruation and modified during pregnancy.
- **pelvic cavity**: A body cavity that is bounded by the bones of the pelvis. Its oblique roof is the pelvic inlet (the superior opening of the pelvis). Its lower boundary is the pelvic floor.

Physical Changes During Pregnancy

Pregnancy begins when the developing embryo implants in the endometrial lining of a woman's uterus. Most pregnant women do not have any specific signs or symptoms after implantation, although it is not uncommon to experience minimal bleeding.

After implantation, the uterine endometrium is called the decidua. The placenta, which is partly formed from the decidua and partly from outer layers of the embryo, connects the developing embryo to the uterine wall to allow nutrient uptake, waste elimination, and gas exchange via the mother's blood supply.

The umbilical cord connects the embryo or fetus to the placenta. The developing embryo undergoes tremendous growth and changes during the process of fetal development.

Most pregnant women experience a number of symptoms that can signify pregnancy. The symptoms include nausea and vomiting, excessive tiredness and fatigue, cravings for certain foods that are not normally sought out, and frequent urination, particularly during the night.

A number of early medical signs are associated with pregnancy. These signs typically appear, if at all, within the first few weeks after conception. Not all of these signs are universally present, nor are all of them diagnostic by themselves; taken together, however, they may make a presumptive diagnosis of pregnancy.

These signs include:

- The presence of human chorionic gonadotropin (hCG) in the blood and urine.
- Missed menstrual period.
- Implantation bleeding (occurs at implantation of the embryo in the uterus during the third or fourth week after last menstrual period).

- Increased basal body temperature sustained for over 2 weeks after ovulation.
- Chadwick's sign (darkening of the cervix, vagina, and vulva).
- Goodell's sign (softening of the vaginal portion of the cervix).
- Hegar's sign (softening of the uterus isthmus).
- Pigmentation of the linea alba (called linea nigra), which is darkening of the skin in a midline of the abdomen. This darkening is caused by hyperpigmentation resulting from hormonal changes, usually appearing around the middle of pregnancy.

Breast tenderness is common during the first trimester. Shortly after conception, the nipples and areolas begin to darken due to a temporary increase in hormones. This process continues throughout the pregnancy.

Despite all the signs, some women may not realize they are pregnant until they are far along in pregnancy. In some cases, a few have not been aware of their pregnancy until they begin labor. This can be caused by many factors, including irregular periods (quite common in teenagers), certain medications (not related to conceiving children), and obese women who disregard the pregnancy-related weight gain. Others may be in denial of their situation.

The First Trimester

The first 12 weeks of pregnancy are known as the first trimester. During this trimester, fetal development can be divided into different stages.

At the beginning of the fetal stage, the risk of miscarriage decreases sharply. All major structures, including the head, brain, hands, feet, and other organs have been formed by the fetal stage. Once pregnancy moves into the second trimester, the risks of miscarriage and birth defects drops drastically.

The Second Trimester

Weeks 13 to 28 of the pregnancy are called the second trimester. Most women feel more energized in this period. They put on weight as the symptoms of morning sickness subside and eventually fade away.

By the end of the second trimester, the expanding uterus has created a visible baby bump. Although the breasts have been developing internally since the beginning of the pregnancy, most of the visible changes appear after this point.

The uterus can expand up to 20 times its normal size during pregnancy. Although the fetus begins to move and takes a recognizable human shape during the first trimester, it is not until the second trimester that movement of the fetus, often referred to as quickening, can be felt by the woman.

The Third Trimester

Final weight gain takes place during the third trimester, and it is the most weight gain throughout the pregnancy. The fetus will be growing most rapidly during this stage, gaining up to 28g per day.



A pregnant woman: A pregnant woman's abdomen drops due to the fetus turning in a downward position ready for birth.

The woman's abdomen will transform in shape as the abdomen drops due to the fetus turning in a downward position ready for birth, and the woman will be able to lift her belly up and down. The woman's navel will sometimes become convex—popping out—due to her expanding abdomen. This period of her pregnancy can be uncomfortable and cause symptoms like weak bladder control and backache.

The body's posture changes as the pregnancy progresses. The pelvis tilts and the back arches to help keep balance. Poor posture occurs naturally from the stretching of the woman's abdominal muscles as the fetus grows. These muscles are less able to contract and keep the lower back in proper alignment.

The pregnant woman has a different gait. The step lengthens as the pregnancy progresses due to weight gain and changes in posture. In addition, the increased body weight of pregnancy, fluid retention, and weight gain lowers the arches of the foot, further adding to the foot's length and width.

The influences of increased hormones such as estrogen and relaxin initiate the remodeling of soft tissues, cartilage, and ligaments. Certain skeletal joints (e.g., the pubic symphysis and sacroiliac) widen or have increased laxity.

This page titled 27.10A: Anatomical Changes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.10B: Metabolic Changes

Protein and carbohydrate metabolisms are affected during pregnancy and maternal insulin resistance can lead to gestational diabetes.

Learning Objectives

- Analyze the metabolic factors involved in gestational diabetes

Key Points

- During pregnancy, the metabolism shifts to provide the growing fetus with more nutrients as well as to ensure the development of the uterine lining and breast glandular tissue.
- Hormonal changes during pregnancy increase nutrient requirements and fat deposition.
- Insulin resistance can develop and lead to gestational diabetes.

Key Terms

- **cortisol**: A steroid hormone (also called hydrocortisone) produced by the adrenal cortex, that regulates the metabolism of carbohydrates and maintains blood pressure.
- **gestational diabetes**: Also called gestational diabetes mellitus (GDM), it is a condition in which women without previously diagnosed diabetes exhibit high blood glucose levels during pregnancy (especially during the third trimester).
- **lactogen**: Human placental lactogen (HPL), also called human chorionic somatomammotropin (HCS), is a polypeptide placental hormone. Its structure and function is similar to that of human growth hormone. It modifies the metabolic state of the mother during pregnancy to facilitate the energy supply of the fetus.

Nutrient Metabolism

During pregnancy, both protein metabolism and carbohydrate metabolism are affected. One kilogram of extra protein is deposited, with half going to the fetus and placenta, and another half going to uterine contractile proteins, breast glandular tissue, plasma protein, and hemoglobin.


An increase in nutrients is required for fetal growth and fat deposition. Changes are caused by steroid hormones, lactogen, and cortisol. Increased liver metabolism is also seen, with increased gluconeogenesis that leads to increased maternal glucose levels. Maternal insulin resistance can lead to gestational diabetes.

Gestational Diabetes

Gestational diabetes (or gestational diabetes mellitus, GDM) is a condition in which women without previously diagnosed diabetes exhibit high blood glucose levels during pregnancy (especially during the third trimester). There is some question whether the condition is natural during pregnancy.

Maternal insulin resistance can lead to gestational diabetes. This type of diabetes is caused when the insulin receptors do not function properly. This is likely due to pregnancy-related factors, such as the presence of human placental lactogen that interferes with susceptible insulin receptors. This in turn causes inappropriately elevated blood sugar levels.

Gestational diabetes generally has few symptoms and it is most commonly diagnosed by screening during pregnancy. Diagnostic tests detect inappropriately high levels of glucose in blood samples. Gestational diabetes affects 3–10% of pregnancies, depending on the population studied, so it may be a natural phenomenon.



Effect of insulin on glucose uptake and metabolism: Insulin binds to its receptor (1) on the cell membrane which in turn starts many protein activation cascades (2). These include: the translocation of the glut-4 transporter to the plasma membrane and an influx of glucose (3), glycogen synthesis (4), glycolysis (5) and fatty acid synthesis (6).

Babies born to mothers with untreated gestational diabetes are typically at an increased risk of problems, such as being large for gestational age (which may lead to delivery complications), low blood sugar, and jaundice. If untreated, it can also cause seizures or still birth.

Gestational diabetes is a treatable condition and women who have adequate control of glucose levels can effectively decrease these risks.

This page titled 27.10B: Metabolic Changes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.10C: Physiological Changes

Maternal physiological changes in pregnancy are entirely normal and serve as adaptations to better accommodate embryonic/fetal development.

Learning Objectives

- Describe the physiological changes a woman undergoes during pregnancy

Key Points

- Women undergo several changes during pregnancy, including cardiovascular, hematologic, metabolic, renal, and respiratory changes that provide adequate nutrition and gas exchange for the developing fetus.
- Progesterone and estrogen levels rise continually through pregnancy, together with blood sugar, breathing rate, and cardiac output.
- The body's posture changes during pregnancy to accommodate the growing fetus and the mother will experience weight gain.
- Breasts grow and change in preparation for lactation once the infant is born. Once lactation begins, the woman's breasts swell significantly and can feel achy, lumpy, and heavy (engorgement). This is relieved by nursing the infant.
- Plasma and blood volume increase over the course of the pregnancy and lead to changes in heart rate and blood pressure. Women may also have a higher risk of blood clots, especially in the weeks following labor.

Key Terms

- **human placental lactogen**: Also called human chorionic somatomammotropin, this is a polypeptide placental hormone. Its structure and function is similar to that of human growth hormone. It modifies the metabolic state of the mother during pregnancy to facilitate the energy supply of the fetus.
- **human chorionic gonadotropin**: A peptide hormone produced during pregnancy that prevents the breakdown of the corpus luteum and maintains progesterone production.
- **progesterone**: A steroid hormone, secreted by the ovaries, whose function is to prepare the uterus for the implantation of a fertilized ovum and to maintain pregnancy.

Maternal physiological changes in pregnancy are the normal adaptations that a woman undergoes during pregnancy to better accommodate the embryo or fetus, and include cardiovascular, hematologic, metabolic, renal, and respiratory changes. The female body must change its physiological and homeostatic mechanisms in pregnancy to ensure proper fetal development. Increases in blood sugar, breathing, and cardiac output are all required.

Hormonal Changes

Pregnant women experience adjustments in their endocrine system. Levels of progesterone and estrogens rise continuously throughout pregnancy to suppress the hypothalamic axis and, subsequently, the menstrual cycle.

Estrogen produced by the placenta is associated with fetal well being. Women also experience an increase in human chorionic gonadotropin (β-hCG), which is produced by the placenta and maintains progesterone production by the corpus luteum.

The increase in progesterone production primarily functions to relax smooth muscles. Prolactin levels increase due to maternal pituitary gland enlargement that mediate a change in the structure of the mammary gland from ductal to lobular-alveolar.

Parathyroid hormone increases and leads to increased calcium uptake in the gut and reabsorption by the kidney. Adrenal hormones such as cortisol and aldosterone also increase.

Human placental lactogen (HPL) is produced by the placenta, stimulating lipolysis and fatty acid metabolism by the woman and

conserving blood glucose for use by the fetus. It can also decrease maternal tissue sensitivity to insulin and result in gestational diabetes.

Weight Changes

One of the most noticeable alterations in pregnancy is the gain in weight. The enlarging uterus, the growing fetus, the placenta and liquor amnii, and the acquisition of fat and water retention, all contribute to weight gain.

The weight gain varies and can be anywhere from five pounds (2.3 kg) to over 100 pounds (45 kg). In the U.S., the doctor-recommended weight gain range is 25 pounds (11 kg) to 35 pounds (16 kg), less if the woman is overweight, more (up to 40 pounds 18 kg) if the woman is underweight.



Pregnancy: During pregnancy, a woman gains weight and her breasts enlarge.

A woman's breasts grow during pregnancy, usually one to two cup sizes, but possibly larger. A woman who wore a C cup bra prior to her pregnancy may need to buy an F cup or larger bra while nursing. A women's torso also grows and her bra band size may increase one or two sizes.

Once the baby is born (about 50 to 73 hours after birth), the mother will experience her breasts filling with milk, at which point changes in the breast happen very quickly. Once lactation begins, the woman's breasts swell significantly and can feel achy, lumpy, and heavy (engorgement). Her breasts may increase again in size. Individual breast size can vary daily or for longer periods depending on how much the infant nurses from each breast.

Circulatory Changes

Plasma and blood volume slowly increase by 40–50% over the course of the pregnancy (due to increased aldosterone) to accommodate the changes, resulting in an increase in heart rate (15 beats/min more than usual), stroke volume, and cardiac output. Cardiac output increases by about 50%, primarily during the first trimester.

The systemic vascular resistance also drops due to the smooth muscle relaxation and overall vasodilation caused by elevated progesterone, leading to a fall in blood pressure. Diastolic blood pressure consequently decreases between 12–26 weeks, and increases again to pre-pregnancy levels by 36 weeks.

Edema (swelling) of the feet is common during pregnancy, partly because the enlarging uterus compresses veins and lymphatic drainage from the legs.

A pregnant woman will also become hypercoagulable, leading to increased risk for developing blood clots and embolisms due

to increased liver production of coagulation factors. Women are at highest risk for developing clots (thrombi) during the weeks following labor.

Clots usually develop in the left leg or the left iliac venous system because the left iliac vein is crossed by the right iliac artery. The increased flow in the right iliac artery after birth compresses the left iliac vein leading to an increased risk for thrombosis (clotting) that is exacerbated by a lack of ambulation (walking) following delivery. Both underlying thrombophilia and caesarean section can further increase these risks.

This page titled 27.10C: Physiological Changes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.10D: Exercise and Pregnancy

In the absence of complications, pregnant women should continue aerobic and strength training exercise for the duration of gestation.

Learning Objectives

- Evaluate the types of exercise appropriate for pregnant women

Key Points

- Moderate aerobic exercise and strength training improve the health of pregnant women while having no adverse consequences on the developing fetus.
- A variety of exercise activities are appropriate, with the exception of those with a high risk for abdominal trauma, such as horseback riding, skiing, soccer, or hockey.
- Contraindications of exercise include: vaginal bleeding, dyspnea before exertion, dizziness, headache, chest pain, muscle weakness, preterm labor, decreased fetal movement, amniotic fluid leakage, and calf pain or swelling (to rule out thrombophlebitis).

Key Terms

- **contraindication**: A factor or symptom that makes the prescribed treatment inadvisable.
- **aerobic exercise**: Physical exercise of low to high intensity that depends primarily on the aerobic energy-generating process.
- **strength-conditioning**: The use of resistance to muscular contraction to build the strength, anaerobic endurance, and size of skeletal muscles. There are many different methods of strength training, the most common being the use of gravity or elastic/hydraulic forces to oppose muscle contraction.
- **thrombophlebitis**: Phlebitis (vein inflammation) related to a thrombus (blood clot).



Exercising while pregnant: A strong, healthy woman will generally have a good pregnancy outcome. Physicians recommend moderate exercise during pregnancy, including strength-training.

Regular aerobic exercise during pregnancy appears to improve (or maintain) physical fitness. Although an upper level of safe exercise intensity has not been established, women who were regular exercisers before pregnancy and who have uncomplicated,

healthy pregnancies should be able to engage in high-intensity exercise programs (e.g., jogging and aerobics) for less than 45 minutes with no adverse effects. They just need to be mindful of the possibility that they may need to increase their energy intake, and are careful to not become overheated.

In the absence of either medical or obstetric complications, doctors advise an accumulation of 30 minutes a day of exercise on most if not all days of the week. The Clinical Practice Obstetrics Committee of Canada recommends that "all women without contraindications should be encouraged to participate in aerobic and strength-conditioning exercises as part of a healthy lifestyle during their pregnancy."

In general, participation in a wide range of recreational activities appears to be safe. Pregnant women just need to avoid those with a high risk of falling such as horseback riding or skiing, or those that carry a risk of abdominal trauma, such as soccer or hockey.

In the past, the main concerns of exercise in pregnancy were focused on the fetus and any potential maternal benefit was thought to be offset by potential risks to the fetus. However, more recent information suggests that in the uncomplicated pregnancy, fetal injuries are highly unlikely.

Contraindications for exercise include, vaginal bleeding, dyspnea before exertion, dizziness, headache, chest pain, muscle weakness, preterm labor, decreased fetal movement, amniotic fluid leakage, and calf pain or swelling (to rule out thrombophlebitis).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Pregnancy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pregnancy%23Physiology. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pregnancy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pregnan...Physical_signs. **License**: *CC BY-SA: Attribution-ShareAlike*
- decidua. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/decidua. **License**: *CC BY-SA: Attribution-ShareAlike*
- pelvic cavity. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/pelvic%20cavity. **License**: *CC BY-SA: Attribution-ShareAlike*
- implantation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/implantation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dropped abdomen. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/7c/Swanger_vrou2.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Maternal physiological changes in pregnancy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Maternal_physiological_changes_in_pregnancy%23Metabolic. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gestational diabetes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gestational_diabetes. **License**: *CC BY-SA: Attribution-ShareAlike*
- cortisol. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cortisol. **License**: *CC BY-SA: Attribution-ShareAlike*
- gestational diabetes. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/gestational%20diabetes. **License**: *CC BY-SA: Attribution-ShareAlike*
- lactogen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/lactogen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Dropped abdomen. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/7c/Swanger_vrou2.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Insulin glucose metabolism. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Insulin_glucose_metabolism.jpg. **License**: *Public Domain: No Known Copyright*
- human chorionic gonadotropin. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/human_chorionic_gonadotropin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Maternal physiological changes in pregnancy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/

Maternal_physiological_changes_in_pregnancy%23Musculoskeletal. License: CC BY-SA: Attribution-ShareAlike
- human placental lactogen. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/human%20placental%20lactogen. License: CC BY-SA: Attribution-ShareAlike
- progesterone. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/progesterone. License: CC BY-SA: Attribution-ShareAlike
- Dropped abdomen. Provided by: Wikimedia. Located at: upload.wikimedia.org/Wikipedia/commons/7/7c/Swanger_vrou2.jpg. License: CC BY-SA: Attribution-ShareAlike
- Insulin glucose metabolism. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Insulin_glucose_metabolism.jpg. License: Public Domain: No Known Copyright
- Pregnant woman black and white shadows. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Pregnant_woman_black_and_white_shadows.jpg. License: CC BY-SA: Attribution-ShareAlike
- contraindication. Provided by: Wiktionary. Located at: en.wiktionary.org/wiki/contraindication. License: CC BY-SA: Attribution-ShareAlike
- strength-conditioning. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/strength-conditioning. License: CC BY-SA: Attribution-ShareAlike
- Thrombophlebitis. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Thrombophlebitis. License: CC BY-SA: Attribution-ShareAlike
- Aerobic Exercise. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/Aerobic_exercise. License: CC BY-SA: Attribution-ShareAlike
- Dropped abdomen. Provided by: Wikimedia. Located at: upload.wikimedia.org/Wikipedia/commons/7/7c/Swanger_vrou2.jpg. License: CC BY-SA: Attribution-ShareAlike
- Insulin glucose metabolism. Provided by: Wikipedia. Located at: en.Wikipedia.org/wiki/File:Insulin_glucose_metabolism.jpg. License: Public Domain: No Known Copyright
- Pregnant woman black and white shadows. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Pregnant_woman_black_and_white_shadows.jpg. License: CC BY-SA: Attribution-ShareAlike
- Pregnant Woman With Dumbells. Provided by: Wikimedia. Located at: commons.wikimedia.org/wiki/File:Pregnant_Woman_With_Dumbells.JPG. License: Public Domain: No Known Copyright

This page titled 27.10D: Exercise and Pregnancy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.1: OVERVIEW OF PREGNANCY AND HUMAN DEVELOPMENT

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

27.1A: Introduction to Pregnancy and Human Development

This page titled 27.1: Overview of Pregnancy and Human Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.1A: Introduction to Pregnancy and Human Development

Pregnancy is the period of gestation from the fertilization of an egg, through development of a fetus, and ending at birth.

Learning Objectives

- Explain pregnancy

Key Points


- A pregnancy can be multiple gestations, as in the case of twins or triplets. Childbirth usually occurs about 38 weeks after conception, or approximately 40 weeks from the start of the last normal menstrual period.
- Pregnancy is divided into three trimesters that roughly approximate specific developmental stages. The risk of spontaneous abortion (miscarriage) is higher during the first trimester and lessens in the second and third trimesters.
- The growth and development of the fetus is more easily monitored during the second trimester, while a fetus is generally viable (although it possibly requires medical intervention) in the third trimester.
- Gravidity (gravid = heavy) is another term for pregnancy as a pregnant woman generally gains a significant amount of weight during gestation. A woman who has never been pregnant is a nulligravida; a woman who is pregnant for the first time is a primigravida; and a woman in subsequent pregnancies is a multigravida.

Key Terms

- **gravidity**: Another term for pregnancy.
- **conceptus**: The prenatal, developing offspring and its associated membranes.
- **prenatal**: During pregnancy, before birth.
- **gestation**: The carrying of an embryo or fetus inside female viviparous (having live births) animals, including humans.
- **trimester**: A period of approximately three months.
- **miscarriage**: The spontaneous natural termination of a pregnancy that expels a fetus from the womb before term.

Pregnancy is the state of fertilization and development for one or more offspring within a woman's uterus. The prenatal offspring (also called the conceptus) is referred to as an embryo or fetus.

The term embryo is used primarily for developing humans up to eight weeks after fertilization (to the 10th week of gestation). After that, the term fetus is used.



Pregnancy: Drawing of the lateral (side) view of a pregnant woman's abdomen revealing the developing fetus.

In a pregnancy, there can be multiple gestations, as in the case of twins or triplets. Childbirth usually occurs about 38 weeks after conception. In women who have a menstrual-cycle length of four weeks, this is approximately 40 weeks from the start of their last normal menstrual period.

In many societies' medical or legal definitions, human pregnancy is somewhat arbitrarily divided into three trimester periods as a means to simplify reference to the different stages of prenatal development. The first trimester carries the highest risk of miscarriage (natural death of embryo or fetus). During the second trimester, the development of the fetus is more easily monitored. The beginning of the third trimester often approximates the point of viability, or the ability of the fetus to survive, with or without medical help, outside of the uterus.

The Carnegie stages is a standardized system of 23 stages used to provide a unified description of the developmental maturation of the vertebrate embryo. This system bases stages on the development of structures instead of days of development or the size of the conceptus. This staging method is used only for the first 56 days in humans (prior to the fetal period).



Ultrasound scan: Ultrasound is used to monitor the age and health of the fetus during pregnancy.

One scientific term for the state of pregnancy is gravidity, which is Latin for heavy. A pregnant female is sometimes referred to as

a gravida.

Similarly, the term parity (abbreviated as para) is used for the number of times a female has given birth; parity counts twins and other multiple births as one pregnancy. Medically, a woman who has never been pregnant is referred to as a nulligravida, and a woman in subsequent pregnancies is referred to as a multigravida.

A woman who has had more than one live birth is referred to as multiparous, and a woman who has never given birth is referred to as nulliparous.

During a second pregnancy, a woman would be described as gravida 2, para 1 and, upon live delivery, as gravida 2, para 2. An in-progress pregnancy, as well as abortions, miscarriages, or stillbirths, account for parity values being less than the gravida number. In the case of twins, triplets, and so on, the gravida number and parity value are increased by one only.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Pregnancy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pregnancy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pregnancy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pregnancy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gravidity. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gravidity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- trimester. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trimester**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- miscarriage. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/miscarriage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Conceptus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Conceptus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gestation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gestation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Carnegie Stages. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Carnegie_stages**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pregnancy.png. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Blausen_0747_Pregnancy.png**. **License**: ***CC BY: Attribution***
- Scan12weeks. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...can12weeks.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 27.1A: Introduction to Pregnancy and Human Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.2: FIRST WEEK OF DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.2A: Fertilization
27.2B: Cleavage of the Zygote
27.2C: Blastocyst Formation
27.2D: Implantation

This page titled 27.2: First Week of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.2A: Fertilization

Fertilization occurs when a sperm and an egg have fused together to form a zygote, which begins to divide as it moves towards the uterus.

Learning Objectives

- Describe human fertilization

Key Points

- Fertilization is commonly known as conception. Once the fertilized gamete (ovum) implants itself in the uterine lining, pregnancy begins.
- The fusion of male and female gametes (sperm and ovum, respectively) usually occurs following the act of sexual intercourse. However, artificial insemination and in vitro fertilization have made achieving pregnancy possible without engaging in sexual intercourse.
- The process of fertilization occurs in several steps and the interruption of any of them can lead to failure.
- Prior to fertilization, sperm undergo a process of capacitation in response to conditions in the female reproductive tract, which include increases in motility and destabilization of the cell membrane that allows the head of the sperm to penetrate the egg.


Key Terms

- **fertilization**: The act of fecundating or impregnating animal or vegetable gametes.
- **capacitation**: A step spermatozoa undergo in the female reproductive track that renders them capable of fertilizing an oocyte.
- **implantation**: The embedding of the fertilized ovum into the uterine wall.
- **Nondisjunction**: Nondisjunction is the failure of homologous chromosomes or sister chromatids to separate properly during cell division.
- **zygote**: A fertilized egg cell.

If pregnancy is considered to begin at the point of implantation, the process leading to pregnancy occurs earlier as the result of the female gamete, or oocyte, merging with the male gamete, or spermatozoon. In medicine, this process is referred to as fertilization; in lay terms, it is more commonly known as conception.

After the point of fertilization the fused product of the female and male gamete is referred to as a zygote or fertilized egg. For species that undergo internal fertilization, such as humans, the fusion of male and female gametes usually occurs following the act of sexual intercourse.

However, the advent of artificial insemination and in vitro fertilization have made achieving pregnancy possible without engaging in sexual intercourse. This approach may be undertaken as a voluntary choice or due to infertility.



Human fertilization: The sperm and ovum unite through fertilization, creating a zygote that (over the course of 8–9 days) will implant in the uterine wall, where it will reside over the course of 9 months.

The process of fertilization occurs in several steps and the interruption of any of them can lead to failure. At the beginning of the process, the sperm undergoes a series of changes, as freshly ejaculated sperm is unable or poorly able to fertilize.

The sperm must undergo capacitation in the female's reproductive tract over several hours, which increases its motility and destabilizes its membrane. By destabilizing the membrane, the sperm prepares for the acrosome reaction, the enzymatic penetration of the egg's tough membrane, the zona pellucida. The sperm and the egg cell (which has been released from one of the female's two ovaries) unite in one of the two fallopian tubes.

The fertilized egg, known as a zygote, then moves toward the uterus, a journey that can take up to a week to complete until implantation occurs. Through fertilization, the egg is activated to begin its developmental process (progressing through meiosis II), and the haploid nuclei of the two gametes come together to form the genome of a new diploid organism.

Nondisjunction during the completion of meiosis or problems with early cell division in the zygote to blastula stages can lead to problems with implantation and pregnancy failure.

This page titled 27.2A: Fertilization is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.2B: Cleavage of the Zygote

The process of cleavage is the step of embryogenesis where the zygote divides to produce a cluster of cells known as the morula.

Learning Objectives

- Describe cleavage of the zygote

Key Points


- Following fertilization a series of rapid cell divisions occur that decrease the cells' size with each subsequent division—this eventually produces a morula. The different cells derived from cleavage, up to the blastula stage, are called blastomeres.
- For species such as humans, there is little yolk in eggs, and the divisions are relatively symmetrical, or holoblastic.
- For other species, such as birds and reptiles, the presence of yolk dictates uneven meroblastic divisions that produce cells of uneven size and distribution.
- The end of cleavage is known as the midblastula transition and coincides with the onset of zygotic transcription.
- The cells of the morula are at first closely aggregated, but quickly become arranged into an outer or peripheral layer, the trophoblast, which does not contribute to the formation of the embryo proper, and an inner cell mass from which the embryo develops.

Key Terms

- **cleavage**: In embryology, this is the division of cells in the early embryo.
- **trophoblast**: The membrane of cells that forms the wall of a blastocyst during early pregnancy and provides nutrients to the embryo, and later develops into part of the placenta.
- **zygote**: A fertilized egg cell.
- **morula**: A spherical mass of blastomeres that forms following the splitting of a zygote; it becomes the blastula.

Cell division with no significant growth that produces a cluster of cells that is the same size as the original zygote, is called cleavage. At least four initial cell divisions occur, resulting in a dense ball of at least sixteen cells called the morula.

The different cells derived from cleavage up to the blastula stage are called blastomeres. Depending mostly on the amount of yolk in the egg, the cleavage can be holoblastic (total) or meroblastic (partial).



Cell cleavage: Early development is characterized by cleavage of the zygote, which refers to cell divisions that are not associated with significant growth of the embryo.

Holoblastic cleavage occurs in animals with little yolk in their eggs. These species, such as humans and other mammals, receive nourishment as embryos from the mother via the placenta or milk after birth.

On the other hand, meroblastic cleavage occurs in animals whose eggs have more yolk, such as birds and oviparous reptiles (although some viviparous reptiles also exist). Since cleavage is impeded by the vegetal pole, there is a very uneven distribution and size of cells. Cells are more numerous and smaller at the animal pole of the zygote than at the vegetal pole.

In holoblastic eggs, the first cleavage always occurs along the vegetal–animal axis of the egg, and the second cleavage is perpendicular to the first. From here, the spatial arrangement of blastomeres can follow various patterns, due to different planes of cleavage in various organisms.The end of cleavage is known as the midblastula transition and coincides with the onset of zygotic transcription.

In amniotes, the cells of the morula are at first closely aggregated. However, they quickly become arranged into an outer or peripheral layer, the trophoblast, and an inner cell mass. The trophoblast does not contribute to the formation of the embryo proper; the embryo develops from the inner cell mass.

Fluid collects between the trophoblast and the greater part of the inner cell mass, and thus the morula, is converted into the blastodermic vesicle (also called the blastocyst or blastula). The inner cell mass remains in contact with the trophoblast at one pole of the ovum. This is named the embryonic pole, since it indicates the location where the future embryo will develop.

In the case of monozygotic twins (derived from one zygote), a zygote divides into two separate cells (embryos) at the first cleavage division. Monozygotic twins can also develop from two inner cell masses.

A rare occurrence is the division of a single inner cells mass giving rise to twins. However, if one inner cell mass divides incompletely, the result is conjoined twins. Dizygotic twins is the development of two embryos from two different zygotes.

This page titled 27.2B: Cleavage of the Zygote is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.2C: Blastocyst Formation

The blastocyst forms early in embryonic development and has two layers that form the embryo and placenta.

Learning Objectives

- Outline the process of blastocyst formation

Key Points

- The human blastocyst possesses an inner cell mass (ICM), or embryoblast, which subsequently forms the embryo, and an outer layer of cells, or trophoblast, which later forms the placenta.
- The trophoblast surrounds the inner cell mass and a fluid-filled, blastocyst cavity known as the blastocoele or the blastocystic cavity. The trophoblast combines with the maternal endometrium to form the placenta in eutherian mammals.
- Before gastrulation, the cells of the trophoblast become differentiated into two strata: the ectoderm of the chorion plays a role in the development of the placenta, and the endoderm differentiates and quickly assumes the form of a small sac, called the yolk sac.
- The embryoblast is the source of embryonic stem cells and gives rise to all later structures of the adult organism.
- The floor of the amniotic cavity is formed by the embryonic disk, which is composed of a layer of prismatic cells, and the embryonic ectoderm, which is derived from the inner cell mass and lies in opposition to the endoderm.


Key Terms

- **embryonic disk**: The floor of the amniotic cavity is formed by the embryonic disk (or disc) that is composed of a layer of prismatic cells called the embryonic ectoderm. It is the part of the inner cell mass from which the embryo is developed.
- **blastocyst**: An early form in the development of an embryo that consists of a spherical layer of cells filled with fluid.
- **gastrulation**: The stage of embryonic development when a gastrula is formed from the blastula by the inward migration of cells.
- **eutherian**: Refers to all species of which the female gives birth to live young that receive prenatal nourishment via the placenta.

In humans, the blastocyst is formed approximatelyy five days after fertilization. This stage is preceded by the morula. The morula is a solid ball of about 16 undifferentiated, spherical cells. As cell division continues in the morula, the blastomeres change their shape and tightly align themselves against each other. This is called compaction and is likely mediated by cell surface adhesion glycoproteins.

The blastocyst possesses an inner cell mass (ICM), or embryoblast, which subsequently forms the embryo, and an outer layer of cells, or trophoblast, which later forms the placenta. The trophoblast surrounds the inner cell mass and a fluid-filled, blastocyst cavity known as the blastocoele or the blastocystic cavity.

The embryoblast is the source of embryonic stem cells and gives rise to all later structures of the adult organism. The trophoblast combines with the maternal endometrium to form the placenta in eutherian mammals.



Blastocyst: The blastocyst possesses an inner cell mass from which the embryo will develop, and an outer layer of cells, called the trophoblast, which will eventually form the placenta.

Before gastrulation, the cells of the trophoblast become differentiated into two strata. The outer stratum forms a syncytium, which is a layer of protoplasm studded with nuclei that shows no evidence of subdivision into cells (termed the syncytiotrophoblast).

The inner layer, the cytotrophoblast or layer of Langhans, consists of well-defined cells. As already stated, the cells of the trophoblast do not contribute to the formation of the embryo proper; they form the ectoderm of the chorion and play an important part in the development of the placenta.

On the deep surface of the inner cell mass, a layer of flattened cells, called the endoderm, is differentiated and quickly assumes the form of a small sac, called the yolk sac. Spaces appear between the remaining cells of the mass and, by the enlargement and coalescence of these spaces, a cavity called the amniotic cavity is gradually developed.

The floor of this cavity is formed by the embryonic disk, which is composed of a layer of prismatic cells called the embryonic ectoderm. This layer is derived from the inner cell mass and lies in opposition to the endoderm.

This page titled 27.2C: Blastocyst Formation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.2D: Implantation

Implantation is the very early stage of pregnancy at which the embryo adheres to the wall of the uterus and begins to form the placenta.

Learning Objectives

- Outline the process of implantation in pregnancy

Key Points

- At this stage of prenatal development the embryo is a blastocyst. In humans, implantation of a fertilized ovum occurs between 6 to 12 days after ovulation.
- In preparation for implantation, the blastocyst sheds its outside layer, the zona pellucida, and is replaced by a layer of underlying cells called the trophoblast. The trophoblast will give rise to the placenta after implantation.
- During implantation, the trophoblast differentiates into two distinct layers: the inner cytotrophoblast, and the outer syncytiotrophoblast. The syncytiotrophoblast then implants the blastocyst into the endometrium by forming finger-like projections into the uterine wall called chorionic villi.

Key Terms


- **endometrium**: The mucous membrane that lines the uterus in mammals and in which fertilized eggs are implanted.
- **trophoblast**: The membrane of cells that forms the wall of a blastocyst during early pregnancy and provides nutrients to the embryo and later develops into part of the placenta.
- **human chorionic gonadotropin (hCG)**: In molecular biology, human chorionic gonadotropin (hCG) is a hormone produced during pregnancy that is made by the developing placenta after conception, and later by the placental component.
- **implantation**: The embedding of the blastocyst to the uterine wall.

Implantation is the very early stage of pregnancy during which the embryo embeds into the wall of the uterus. At this stage of prenatal development, the embryo is a blastocyst.

It is by this adhesion that the fetus receives oxygen and nutrients from the mother to be able to grow. In humans, implantation of a blastocyst occurs between 6 to 12 days after ovulation.

In preparation for implantation, the blastocyst sheds its outside layer, the zona pellucida, which binds sperm during fertilization. The zona pellucida degenerates and decomposes, and is replaced by a layer of underlying cells called the trophoblast.

The trophoblast will give rise to the placenta after implantation. During implantation, the trophoblast differentiates into two distinct layers: the inner cytotrophoblast, and the outer syncytiotrophoblast.



Chorionic villi: During implantation, extensions of the trophoblast, the syncytiotrophoblasts, embed within the endometrium and form chorionic villi.

The syncytiotrophoblast then implants the blastocyst into the endometrium of the uterus by forming finger-like projections into the uterine wall called chorionic villi. The chorionic villi grow outwards until they come into contact with the maternal blood supply.

The chorionic villi will be the border between maternal and fetal blood during the pregnancy, and the location of gas and nutrient exchange between the fetus and the mother. The creation of chorionic villi is assisted by hydrolytic enzymes that erode the uterine epithelium.

The syncytiotrophoblast also produces human chorionic gonadotropin (hCG), a hormone that notifies the mother's body that she is pregnant and prevents menstruation by sustaining the function of the progesterone-producing corpus luteum within the ovary.

Human chorionic gonadotropin is the hormone that is detected by pregnancy tests, as it is found in the maternal bloodstream and urine.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Pregnancy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pregnancy%23Initiation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- capacitation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/capacitation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- implantation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/implantation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- zygote. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/zygote**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fertilization. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fertilization**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nondisjunction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nondisjunction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Fertilization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Hu...tilization.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Embryogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryog...age_and_morula**. **License**: ***CC BY-***

SA: Attribution-ShareAlike
- trophoblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trophoblast**. License: *CC BY-SA: Attribution-ShareAlike*
- morula. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/morula**. License: *CC BY-SA: Attribution-ShareAlike*
- cleavage. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cleavage**. License: *CC BY-SA: Attribution-ShareAlike*
- zygote. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/zygote**. License: *CC BY-SA: Attribution-ShareAlike*
- Human Fertilization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_Fertilization.png**. License: *CC BY-SA: Attribution-ShareAlike*
- Gray9. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray9.png**. License: ***Public Domain: No Known Copyright***
- Blastocyst. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blastocyst**. License: *CC BY-SA: Attribution-ShareAlike*
- Embryogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryog...f_the_blastula**. License: *CC BY-SA: Attribution-ShareAlike*
- embryonic disk. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/embryonic%20disk**. License: *CC BY-SA: Attribution-ShareAlike*
- blastocyst. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/blastocyst**. License: *CC BY-SA: Attribution-ShareAlike*
- gastrulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gastrulation**. License: *CC BY-SA: Attribution-ShareAlike*
- Human Fertilization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_Fertilization.png**. License: *CC BY-SA: Attribution-ShareAlike*
- Gray9. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray9.png**. License: ***Public Domain: No Known Copyright***
- Blastocyst English. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Blastocyst_English.svg**. License: *CC BY-SA: Attribution-ShareAlike*
- Human embryogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Human_e...23Implantation**. License: *CC BY-SA: Attribution-ShareAlike*
- Implantation (human embryo). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Implantation_(human_embryo)**. License: *CC BY-SA: Attribution-ShareAlike*
- endometrium. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/endometrium**. License: *CC BY-SA: Attribution-ShareAlike*
- implantation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/implantation**. License: *CC BY-SA: Attribution-ShareAlike*
- human chorionic gonadotropin (hCG). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/human%2...tropin%20(hCG)**. License: *CC BY-SA: Attribution-ShareAlike*
- trophoblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trophoblast**. License: *CC BY-SA: Attribution-ShareAlike*
- Human Fertilization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Human_Fertilization.png**. License: *CC BY-SA: Attribution-ShareAlike*
- Gray9. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray9.png**. License: *Public Domain: No Known Copyright*
- Blastocyst English. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Blastocyst_English.svg**. License: *CC BY-SA: Attribution-ShareAlike*
- Gray36. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray36.png**. License: *Public Domain: No Known Copyright*

This page titled 27.2D: Implantation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3: SECOND WEEK OF DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.3A: Trophoblast Development
27.3B: Bilaminar Embryonic Disc Development
27.3C: Amnion Development
27.3D: Yolk Sac Development
27.3E: Sinusoid Development
27.3F: Development of the Extraembryonic Coelom

This page titled 27.3: Second Week of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3A: Trophoblast Development

Trophoblasts are the outer layer of cells that provide nutrients to the embryo and form part of the placenta.

Learning Objectives

- Describe the trophoblast

Key Points

- The trophoblast is subdivided into two layers, the inner cytotrophoblast, which proliferates upon implantation, and the outer syncytiotrophoblast, which supports cytotrophoblast proliferation through contact with the maternal blood.
- The cytotrophoblast, at the edges of the villi in the placenta, further differentiates into extravillous trophoblasts, which penetrate into the uterus and attach to the placenta and the mother to promote placental vasculature.
- Some trophoblasts replace endothelial cells in the uterine spiral arteries and remodel them into wide-bore conduits that are independent of maternal vasoconstriction. This ensures the fetus receives a steady supply of blood, and that the placenta is not sensitive to damaging fluctuations in oxygen.

Key Terms

- **decidualization**: The changes in the endometrial lining after ovulation, characterized by its transformation into a secretory lining in preparation to accept an embryo.
- **extravillous trophoblast**: These grow out from the placenta and penetrate into the decidualized uterus.
- **trophoblast**: The membrane of cells that forms the wall of a blastocyst during early pregnancy, and also provides nutrients to the embryo and later develops into part of the placenta.

Trophoblasts

Trophoblasts (from the Greek words trephein, to feed, and blastos, germinator) are cells that form the outer layer of a blastocyst. These cells provide nutrients to the embryo and develop into a large part of the placenta.

They are formed during the first stage of pregnancy and are the first cells to differentiate from the fertilized egg. This layer of trophoblasts is also collectively referred to as the trophoblast, or, after gastrulation, the trophectoderm, as it is then contiguous with the ectoderm of the embryo.

Trophoblast Function

Trophoblasts play an important role in embryo implantation and interaction with the endometrium of the maternal uterus following decidualization. The trophoblast is composed of two layers: an inner cytotrophoblast and an outer syncytiotrophoblast.


The syncytiotrophoblast is non-proliferative and thus relies on fusion of the underlying cytotrophoblast cells to expand. The syncytiotrophoblast are the cells in direct contact with the maternal blood that reaches the placental surface, and thus facilitates the exchange of nutrients, wastes, and gases between the maternal and fetal systems.

Cytotrophoblast in the tips of villi can differentiate into another type of trophoblast called the extravillous trophoblast. Extravillous trophoblasts grow out from the placenta and penetrate into the decidualized uterus.

This process is essential not only for physically attaching the placenta to the mother, but also for altering the vasculature in the uterus to allow it to provide an adequate blood supply to the growing fetus as pregnancy progresses.

Some of the trophoblast even replaces the endothelial cells in the uterine spiral arteries as they remodel these vessels into wide

bore conduits that are independent of maternal vasoconstriction. This ensures the fetus receives a steady supply of blood, and the placenta is not sensitive to fluctuations in oxygen that could cause it damage.



Extraembryonic coelom or chorionic cavitiy: A blastocyst embedded in the uterine decidua. (am. Amniotic cavity, b.c. Blood clot., b.s. Body stalk., etc Embryonic ectoderm, ent. Entoderm., mes. Mesoderm, m.v. Maternal vessels., tr. Trophoblast., u.e. Uterine epithelium, u.g. Uterine glands., y.s. Yolk-sac.)

Clinical Example

Intrauterine growth restriction (IUGR) can result in an undersized fetus with poor organ development or even death. The primary factor in IUGR is placental dysfunction caused by a failure of the extravillous trophoblasts to penetrate and modify the uterine spiral arteries. The result is a poorly-oxygenated environment and reduced fetal growth.

This page titled 27.3A: Trophoblast Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3B: Bilaminar Embryonic Disc Development

The floor of the amniotic cavity is formed by the embryonic disc.

Learning Objectives

- Identify the bilaminar embryonic disc

Key Points

- In humans, the formation of the embryonic disc occurs after implantation and prior to embryonic folding (between about day 14 to day 21 post-fertilization).
- The embryonic disc is derived from the epiblast layer, which lies between the hypoblast layer and the amnion. The epiblast layer is derived from the inner cell mass.
- The formation of the bilaminar embryonic disc precedes gastrulation.


Key Terms

- **optic disc**: The location where ganglion cell axons exit the eye to form the optic nerve.
- **bilaminar**: Formed of, or having, two laminae, or thin plates.

The floor of the amniotic cavity is formed by the embryonic disc, which is composed of a layer of prismatic cells and the embryonic ectoderm. It is derived from the inner cell mass and lies adjacent to the endoderm.

In humans, the formation of the embryonic disc occurs after implantation and prior to embryonic folding (between about day 14 to day 21 post-fertilization). The embryonic disc is derived from the epiblast layer, which lies between the hypoblast layer and the amnion. The epiblast layer is derived from the inner cell mass.

Embryonic Disc Development



Surface view of a rabbit embryo: A drawing of a rabbit embryo that identifies the embryonic disc and the primitive streak (arg—embryonic disc; pr—primitive streak).

The embryonic disc forms during early development. By the blastocyst stage, the embryo is a hollow ball of cells with the inner cell mass (embryoblast) off to one side, and the blastocystic cavity fills the remainder of the sphere.

As the embryo progresses in implantation, a small space appears in the embryoblast and forms the amniotic cavity. Simultaneously, morphological changes occur in the embryoblast that result in the formation of a flat, almost circular bilaminar plate of cells that include the epiblast and the hypoblast—the embryonic disk.

The epiblast forms the floor of the amniotic cavity and is continuous with the amnion. The hypoblast forms the roof of the exocoelomic cavity and is continuous with the thin exocoelomic membrane.

The formation of the bilaminar embryonic disc precedes gastrulation. As gastrulation progresses, the embryonic disc becomes trilaminar and the notochord is formed. Through the process of neurulation, the notochord induces the formation of the central nervous system in the embryonic disc.

This page titled 27.3B: Bilaminar Embryonic Disc Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3C: Amnion Development

The amnion contains the fluid that cushions and protects the fetus.

Learning Objectives


- Describe the development of the amnion

Key Points

- In humans, the amnion is present in the earliest observed embryonic stage, appearing as a cavity within a mass of cells.
- The amniotic cavity is roofed in by a single stratum of flattened, ectodermal cells, called the amniotic ectoderm, while the floor is composed of the prismatic ectoderm of the embryonic disk.
- Outside the amniotic ectoderm, a thin layer of mesoderm is connected by the body stalk with the mesodermal lining of the chorion.
- By approximately the fourth to fifth week, amniotic fluid (liquor amnii) begins to accumulate in the amnion, which increases in quantity and expands to contact the chorion.
- During the later stages of pregnancy, the amniotic fluid allows easier and more fluid movements of the fetus and diminishes the risk of injury.

Key Terms

- **amniotic fluid**: In placental mammals, a fluid contained within the amnion membrane that surrounds a developing embryo or fetus (also called liquor amnii).
- **embryonic disk**: The floor of the amniotic cavity is formed by the embryonic disk (or disc), which is composed of a layer of prismatic cells.
- **amnion**: The innermost membrane of the fetal membranes of reptiles, birds, and mammals; the sac in which the embryo is suspended.
- **chorion**: One of the membranes that exist during pregnancy between the developing fetus and mother.



Amnion: The human fetus is enclosed within the amnion.

The amnion is a closed sac appearing in the inner cell mass as a cavity. This cavity is roofed in by a single stratum of flattened,

ectodermal cells called the amniotic ectoderm. Its floor consists of the prismatic ectoderm of the embryonic disk.

The continuity between the roof and the floor is established at the margin of the embryonic disk. Outside the amniotic ectoderm is a thin layer of mesoderm (continuous with that of the somatopleure), which is connected by the body stalk with the mesodermal lining of the chorion.

When first formed, the amnion is in contact with the body of the embryo, but by about the fourth or fifth week, amniotic fluid (liquor amnii) begins to accumulate within it. This fluid increases in quantity, causing the amnion to expand and ultimately to adhere to the inner surface of the chorion so that the extra-embryonic part of the coelom is obliterated.

This increase continues up to the sixth or seventh month of pregnancy, after which it diminishes somewhat. At the end of pregnancy, it amounts to about one liter.

The amniotic fluid allows some free movement for the fetus during the later stages of pregnancy and also diminishes the risk of injury. It contains less than two percent solids, and consists mainly of urea and other extractives, inorganic salts, a small amount of protein, and, frequently, a trace of sugar.

Note: That some of the liquor amnii is swallowed by the fetus is proved by the fact that epidermal debris and hairs have been found among the contents of the fetal alimentary canal.

This page titled 27.3C: Amnion Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3D: Yolk Sac Development

The yolk sac is vascularized and contributes nutrients to the embryo.

Learning Objectives

- Discuss the location and purpose of the yolk sac

Key Points


- The yolk sac is the earliest visible portion of the gestational sac and is situated on the ventral portion of the embryo, arising from extra-embryonic endoderm.
- The blood to and from the yolk sac is conveyed via the primitive aorta and vitelline circulation.
- At approximately the end of the fourth week, the yolk sac is connected to the primitive digestive system, which allows the yolk sac to contribute nutrients to the embryo.

Key Terms

- **yolk sac**: A membranous sac attached to an embryo that provides early nourishment in the form of yolk in bony fishes, sharks, reptiles, birds, and mammals. It functions as the developmental circulatory system of the human embryo before internal circulation begins.
- **vitelline circulation**: The system of blood flowing from the embryo to the yolk sac and back again.
- **gestational sac**: The gestational sac (or gestation sac) is the only available intrauterine structure that can be used to determine if an intrauterine pregnancy (IUP) exists until the embryo is identified.
- **Heuser's membrane**: Also called the exocoelomic membrane, it is a short lived combination of hypoblast cells and extracellular matrix.
- **mesenchyme**: A type of tissue characterized by loosely associated cells that lack polarity and are surrounded by a large extracellular matrix.

The yolk sac is the first element seen in the gestational sac during pregnancy. In humans, it is usually visible at five weeks of gestation. Identifying a true gestation sac is a critical landmark, and is reliably seen in early pregnancy through ultrasound. The yolk sac, situated on the ventral aspect of the embryo, is lined by extra-embryonic endoderm, outside of which is a layer of extra-embryonic mesenchyme derived from the mesoderm.

Blood is conveyed to the wall of the sac by the primitive aorta. After circulating through a wide-meshed capillary plexus, it is returned by the vitelline veins to the tubular heart of the embryo. This vitelline circulation absorbs nutritive material from the yolk sac that is conveyed to the embryo.



Yolk sac: The yolk sac is a membranous sac attached to the embryo that provides nourishment in the form of yolk.

At the end of the fourth week, the yolk sac has the appearance of a small, pear-shaped vesicle (umbilical vesicle) opening into the digestive tube by a long, narrow tube, the vitelline duct. The vesicle can be seen in the afterbirth as a small, somewhat oval-shaped body whose diameter varies from 1 mm to 5 mm. It is situated between the amnion and the chorion and may lie on or at a varying distance from the placenta.

As a rule, the vitelline duct undergoes complete obliteration during the seventh week. In about 2% of the cases, its proximal portion persists as a diverticulum from the small intestine (Meckel's diverticulum), which is situated about 60 cm proximal to the ileocecal valve. It may be attached by a fibrous cord to the abdominal wall at the umbilicus. Sometimes, a narrowing of the lumen of the ileum is seen opposite the site of attachment of the duct.

The yolk sac starts forming during the second week of embryonic development, at the same time of the shaping of the amniotic sac. The hypoblast starts proliferating laterally and descending. In the meantime Heuser's membrane, located on the opposite pole of the developing vesicle, starts its upward proliferation and meets the hypoblast.

This page titled 27.3D: Yolk Sac Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3E: Sinusoid Development

The sinusoids are capillaries that develop after implantation to allow the exchange of gas and nutrients with the mother.

Learning Objectives

- Describe the development of embryonic sinusoids

Key Points


- The vitelline veins drain blood from the yolk sac. Portions of the veins above the upper ring become interrupted by the developing liver and broken up by it into a plexus of small capillary -like vessels termed sinusoids.
- Sinusoids are found in the liver, lymphoid tissue, endocrine organs, and hematopoietic organs such as the bone marrow and the spleen. Sinusoids found within the terminal villi of the placenta are not comparable to these because they possess a continuous endothelium and complete basal lamina.
- The invasion of endometrial sinusoids by the trophoblast allows maternal blood flow into the trophoblastic lacunae; this forms uteroplacental blood circulation.

Key Terms

- **sinusoid**: Any of several channels through which venous blood passes in various organs.
- **trophoblastic lacunae**: Spaces in the early syncytiotrophoblastic layer of the chorion present prior to the development of the villi.
- **vitelline veins**: Veins that drain blood from the yolk sac.

The vitelline veins drain blood from the yolk sac. At first, they run upward in front on either side of the intestinal canal and, subsequently, unite on the ventral aspect of the canal. Beyond this, they are connected to one another by two anastomotic branches, one on the dorsal and the other on the ventral aspect of the duodenal portion of the intestine.

Thus, it is encircled by two venous rings. The superior mesenteric vein opens into the middle or dorsal anastomosis. The portions of the veins above the upper ring become interrupted by the developing liver and broken up by it into a plexus of small, capillary-like vessels termed sinusoids.



Vitelline veins: The liver and veins of a human embryo, 24 or 25 days old, as seen from the ventral surface. (The vitelline veins are visible at the center bottom.)

A sinusoid is a small blood vessel that is a type of capillary similar to a fenestrated endothelium. Sinusoids are actually classified as a type of open pore capillary (that is, discontinuous) as opposed to fenestrated.

Fenestrated capillaries have diaphragms that cover the pores, whereas open pore capillaries lack a diaphragm, having just an open pore. The open pore endothelial cells greatly increase their permeability.

Permeability is also increased by large intercellular clefts and fewer tight junctions. The level of permeability can allow small and medium-sized proteins, such as albumin, to readily enter and leave the blood stream.

Sinusoids are found in the liver, lymphoid tissue, endocrine organs, and hematopoietic organs, such as the bone marrow and the spleen. Sinusoids found within the terminal villi of the placenta are not comparable to these because they possess a continuous endothelium and complete basal lamina.

The invasion of endometrial sinusoids by the trophoblast allows maternal blood flow into the trophoblastic lacunae; this forms the uteroplacental blood circulation.

This page titled 27.3E: Sinusoid Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.3F: Development of the Extraembryonic Coelom

The extra-embryonic coelom is a cavity that contains the chorion. It is located between Heuser's membrane and the trophoblast.

Learning Objectives

- Describe the development of the extraembryonic coelom


Key Points

- The extra-embryonic coelom develops at the same time as the primitive yolk sac through the proliferation and differentiation of hypoblast cells into mesenchymal cells that fill the area between Heuser's membrane and the trophoblast. The entire structure is enclosed by the chorionic plate.
- The extra-embryonic mesoderm is subdivided into two layers: the extra-embryonic splanchnopleuric mesoderm, which is outside the primitive yolk sac; and the extra-embryonic somatopleuric mesoderm, which is adjacent to the cytotrophoblast.
- The chorion is one of the membranes that exist during pregnancy between the developing fetus and mother.

Key Terms

- **chorion**: The protective and nutritive membrane that attaches higher vertebrate fetuses to the uterus.
- **Heuser's membrane**: Heuser's membrane (or the exocoelomic membrane) is a short-lived combination of hypoblast cells and extracellular matrix.
- **coelum**: Also called the chorionic cavity, this is a portion of the conceptus that consists of a cavity between Heuser's membrane and the trophoblast.

Development of the Extra-Embryonic Coelom



Gestational sac: An artificially colored image of the contents in the cavity of the uterus seen at approximately 5 weeks of gestational age by obstetric ultrasonography.

The extra-embryonic coelom (or chorionic cavity) is a portion of the conceptus consisting of a cavity between Heuser's membrane and the trophoblast. During the formation of the primitive yolk sac, some of the migrating hypoblast cells transdifferentiate into mesenchymal cells that fill the space between Heuser's membrane and the trophoblast to form the extra-embryonic mesoderm.

As development progresses, small lacunae begin to form within the extra-embryonic mesoderm that become larger and form the extra-embryonic coelom.

The extra-embryonic mesoderm is divided into two layers: the extra-embryonic splanchnopleuric mesoderm, which lies adjacent to Heuser's membrane around the outside of the primitive yolk sac; and the extra-embryonic somatopleuric mesoderm, which lies adjacent to the cytotrophoblast layer of the embryo.

The extra-embryonic coelomic cavity is also called the chorionic cavity—it is enclosed by the chorionic plate. The chorionic plate is composed of an inner layer of somatopleuric mesoderm and an outer layer of trophoblast cells. It is the fetal aspect of the placenta that gives rise to chorionic villi.

Chorion Development

The chorion is one of the membranes that exist during pregnancy between the developing fetus and the mother. It consists of an extra-embryonic mesoderm and two layers of trophoblast and surrounds the embryo and other membranes.

Villi emerge from the chorion, which invade the endometrium, destroy the uterine decidua, and allow the transfer of nutrients from maternal blood to fetal blood.

Chorionic villi are at first small and nonvascular, and consist of only trophoblast, but they increase in size and branch. Blood is carried to the villi by the paired umbilical arteries, which branch into chorionic arteries and enter the chorionic villi as cotyledon arteries.

After circulating through the capillaries of the villi, the blood is returned to the embryo by the umbilical veins. Until about the end of the second month of pregnancy, the villi cover the entire chorion, and are almost uniform in size; but, after this stage, they develop unequally.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Trophoblast. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Trophoblast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extravillous trophoblast. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/extravi...%20trophoblast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- trophoblast. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trophoblast**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Decidualization. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Decidualization**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray32. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray32.png**. **License**: ***Public Domain: No Known Copyright***
- optic disc. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/optic_disc**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Development of Bilaminar Embryo. **Provided by**: scientia Wikispace. **Located at**: **http://scientia.wikispaces.com/Devel...laminar+Embryo**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Embryonic disk. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryonic_disk**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bilaminar. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/bilaminar**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray32. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray32.png**. **License**: ***Public Domain: No Known Copyright***
- Embryonic disc. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryonic_disc**. **License**: ***Public Domain: No Known Copyright***

- Amnion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Amnion%23In_humans. **License**: *CC BY-SA: Attribution-ShareAlike*
- amnion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amnion. **License**: *CC BY-SA: Attribution-ShareAlike*
- embryonic disk. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/embryonic%20disk. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chorion. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chorion. **License**: *CC BY-SA: Attribution-ShareAlike*
- amniotic fluid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/amniotic_fluid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray32. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray32.png. **License**: *Public Domain: No Known Copyright*
- Embryonic disc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Embryonic_disc. **License**: *Public Domain: No Known Copyright*
- Gray30. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray30.png. **License**: *Public Domain: No Known Copyright*
- Yolk sac. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Yolk_sac%23In_humans. **License**: *CC BY-SA: Attribution-ShareAlike*
- Endodermal sinus tumor. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Endodermal_sinus_tumor. **License**: *CC BY-SA: Attribution-ShareAlike*
- vitelline circulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vitelline%20circulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- gestational sac. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/gestational%20sac. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mesenchyme. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Mesenchyme. **License**: *CC BY-SA: Attribution-ShareAlike*
- yolk sac. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/yolk_sac. **License**: *CC BY-SA: Attribution-ShareAlike*
- Heuser's Membrane. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Heuser%27s_membrane. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray32. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray32.png. **License**: *Public Domain: No Known Copyright*
- Embryonic disc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Embryonic_disc. **License**: *Public Domain: No Known Copyright*
- Gray30. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray30.png. **License**: *Public Domain: No Known Copyright*
- Yolk sac. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Yolk_sac. **License**: *Public Domain: No Known Copyright*
- Implantation (human embryo). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Implantation_(human_embryo). **License**: *CC BY-SA: Attribution-ShareAlike*
- Sinusoid (blood vessel). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sinusoid_(blood_vessel). **License**: *CC BY-SA: Attribution-ShareAlike*
- Vitelline veins. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vitelline_veins. **License**: *CC BY-SA: Attribution-ShareAlike*
- vitelline veins. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/vitelline%20veins. **License**: *CC BY-SA: Attribution-ShareAlike*
- sinusoid. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sinusoid. **License**: *CC BY-SA: Attribution-ShareAlike*
- Gray32. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray32.png. **License**: *Public Domain: No Known Copyright*
- Embryonic disc. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Embryonic_disc. **License**: *Public Domain: No Known Copyright*
- Gray30. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray30.png. **License**: *Public Domain: No Known Copyright*
- Yolk sac. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Yolk_sac. **License**: *Public Domain: No Known Copyright*
- Gray475. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray475.png. **License**: *Public Domain: No Known Copyright*

- Extra-embryonic coelom. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Extra-embryonic_coelom**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Chorion. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chorion%23Growth**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/definition/chorion**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Heuser's membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Heuser's%20membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- extraembryonic coelom. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/extraembryonic%20coelom**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray32. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray32.png**. **License**: ***Public Domain: No Known Copyright***
- Embryonic disc. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryonic_disc**. **License**: ***Public Domain: No Known Copyright***
- Gray30. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray30.png**. **License**: ***Public Domain: No Known Copyright***
- Yolk sac. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Yolk_sac**. **License**: ***Public Domain: No Known Copyright***
- Gray475. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray475.png**. **License**: ***Public Domain: No Known Copyright***
- Ultrasound of embryo at 5 weeks, colored. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:Ultrasound_of_embryo_at_5_weeks,_colored.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 27.3F: Development of the Extraembryonic Coelom is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4: THIRD WEEK OF DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.4A: Gastrulation

27.4B: Neurulation

27.4C: Somite Development

27.4D: Development of the Cardiovascular System

27.4E: Chorionic Villi and Placental Development

This page titled 27.4: Third Week of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4A: Gastrulation

During gastrulation, the embryo develops three germ layers (endoderm, mesoderm, and ectoderm) that differentiate into distinct tissues.

Learning Objectives

- Describe gastrulation and germ-layer formation


Key Points

- Gastrulation takes place after cleavage and the formation of the blastula. Formation of the primitive streak is the beginning of gastrulation. It is followed by organogenesis—when individual organs develop within the newly-formed germ layers.
- The ectoderm layer will give rise to neural tissue, as well as the epidermis.
- The mesoderm develops into somites that differentiate into skeletal and muscle tissues, the notochord, blood vessels, dermis, and connective tissues.
- The endoderm gives rise to the epithelium of the digestive and respiratory systems and the organs associated with the digestive system, such as the liver and pancreas.

Key Terms

- **somite**: One of the paired masses of mesoderm, distributed along the sides of the neural tube, that will eventually become dermis, skeletal muscle, or vertebrae.
- **gastrulation**: The stage of embryonic development at which a gastrula is formed from the blastula by the inward migration of cells.
- **notochord**: A structure found in the embryos of vertebrates from which the spine develops.
- **epiboly**: One of many movements in the early embryo that allow for dramatic physical restructuring and is characterized by a thinning and spreading of cell layers.

Gastrulation is a phase early in the embryonic development of most animals during which the single-layered blastula is reorganized into a trilaminar (three-layered) structure known as the gastrula. These three germ layers are known as the ectoderm, mesoderm, and endoderm.



Gastrulation: Formation of the three primary germ layers occurs during the first two weeks of development. The embryo at this stage is only a few millimeters in length.

Gastrulation takes place after cleavage and the formation of the blastula and the primitive streak. It is followed by organogenesis, when individual organs develop within the newly-formed germ layers. Each layer gives rise to specific tissues and organs in the developing embryo.

In amniotes such as humans, gastrulation occurs in the following sequence:

1. The embryo becomes asymmetric.
2. The primitive streak forms.
3. Cells from the epiblast at the primitive streak undergo an epithelial to mesenchymal transition and ingress at the primitive streak to form the germ layers.

The ectoderm gives rise to the epidermis, and also to the neural crest and other tissues that will later form the nervous system. The mesoderm is found between the ectoderm and the endoderm, giving rise to somites.

The somites form muscle, the cartilage of the ribs and vertebrae, the dermis, the notochord, blood and blood vessels, bone, and connective tissue.

The endoderm gives rise to the epithelium of the digestive and respiratory systems, and the organs associated with the digestive system, such as the liver and pancreas. Following gastrulation, the cells in the body are either organized into sheets of connected cells (as in epithelia), or as a mesh of isolated cells, such as mesenchyme.

The molecular mechanism and timing of gastrulation is different in different organisms. However, some common features of gastrulation across triploblastic organisms include:

- A change in the topological structure of the embryo, from a simply connected surface (sphere-like), to a non-simply connected surface (torus-like)
- The differentiation of cells into one of three types (endodermal, mesodermal, or ectodermal).
- The digestive function of a large number of endodermal cells.

Although gastrulation patterns exhibit enormous variation throughout the animal kingdom, they are unified by the five basic types of cell movements that occur during gastrulation:

1. Invagination
2. Involution
3. ingression
4. Delamination

5. Epiboly

This page titled 27.4A: Gastrulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4B: Neurulation

Following gastrulation, the neurulation process develops the neural tube in the ectoderm, above the notochord of the mesoderm.

Learning Objectives

- Outline the process of neurulation

Key Points

- The notochord stimulates neurulation in the ectoderm after its development.
- The neuronal cells running along the back of the embryo form the neural plate, which folds outward to become a groove.
- During primary neurulation, the folds of the groove fuse to form the neural tube. The anterior portion of the tube forms the basal plate, the posterior portion forms the alar plate, and the center forms the neural canal.
- The ends of the neural tube close at the conclusion of the fourth week of gestation.

Key Terms

- **basal plate**: In the developing nervous system, this is the region of the neural tube ventral to the sulcus limitans. It extends from the rostral mesencephalon to the end of the spinal cord and contains primarily motor neurons.
- **neurulation**: The process that forms the vertebrate nervous system in embryos.
- **alar plate**: The alar plate (or alar lamina) is a neural structure in the embryonic nervous system, part of the dorsal side of the neural tube, that involves the communication of general somatic and general visceral sensory impulses. The caudal part later becomes the sensory axon part of the spinal cord.
- **notochord**: Composed of cells derived from the mesoderm, this provides signals to the surrounding tissue during development.

Neurulation is the formation of the neural tube from the ectoderm of the embryo. It follows gastrulation in all vertebrates. During gastrulation cells migrate to the interior of the embryo, forming the three germ layers: the endoderm (the deepest layer), the mesoderm (the middle layer), and the ectoderm (the surface layer) from which all tissues and organs will arise.


In a simplified way, it can be said that the ectoderm gives rise to skin and the nervous system, the endoderm to the intestinal organs, and the mesoderm to the rest of the organs.

After gastrulation, the notochord—a flexible, rod-shaped body that runs along the back of the embryo—is formed from the mesoderm. During the third week of gestation the notochord sends signals to the overlying ectoderm, inducing it to become neuroectoderm.

This results in a strip of neuronal stem cells that runs along the back of the fetus. This strip is called the neural plate, and it is the origin of the entire nervous system.

The neural plate folds outwards to form the neural groove. Beginning in the future neck region, the neural folds of this groove close to create the neural tube (this form of neurulation is called primary neurulation).

The anterior (ventral or front) part of the neural tube is called the basal plate; the posterior (dorsal or rear) part is called the alar plate. The hollow interior is called the neural canal. By the end of the fourth week of gestation, the open ends of the neural tube (the neuropores) close off.




Neurulation: Transverse sections that show the progression of the neural plate into the neural tube.

Secondary neurulation of vertebrates occurs when primary neurulation terminates. It is the process by which the neural tube at the lower levels and the caudal to the mid-sacral region is formed.

In general, it entails the cells of the neural plate forming a cord-like structure that migrates inside the embryo and hollows to form the tube. Each organism uses primary and secondary neurulation to varying degrees (except fish, which use only secondary neurulation).

CLINICAL EXAMPLE

Spina bifida is a developmental congenital disorder caused by the incomplete closing of the neural tube during neurulation.



Spina bifida: An illustration of a child with spina bifida

This page titled 27.4B: Neurulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4C: Somite Development

Somites develop from the paraxial mesoderm and participate in the facilitation of multiple developmental processes.

Learning Objectives

- Describe the functions of somites

Key Points

- The paraxial mesoderm is distinct from the mesoderm found more internally in the embryo.
- Alongside the neural tube, the mesoderm develops distinct paired structures called somites that develop into dermis, skeletal muscle, and vertebrae.
- Each somite has four compartments: the sclerotome, myotome, dermatome, and the syndetome. Each becomes a specific tissue during development.

Key Terms

- **neural crest cells**: A transient, multipotent, migratory cell population that gives rise to a diverse cell lineage including melanocytes, craniofacial cartilage, bone, smooth muscle, peripheral and enteric neurons, and glia.
- **conceptus**: The fetus or embryo, including all the surrounding tissues protecting and nourishing it during pregnancy.
- **somite**: One of the paired masses of mesoderm, distributed along the sides of the neural tube, that will eventually become dermis, skeletal muscle, or vertebrae.

Intraembryonic Coelom Development


In the development of the human embryo the intraembryonic coelom (or somatic coelom) is a portion of the conceptus that forms in the mesoderm. During the second week of development the lateral mesoderm splits into a dorsal somatic mesoderm (somatopleure) and a ventral splanchnic mesoderm (splanchnopleure).

By the third week of development, this process gives rise to a cavity between the somatopleure and splanchnopleure referred to as the intraembryonic celom. This space later gives rise to both the thoracic and abdominal cavities.

Somite Development

In the developing vertebrate embryo, *somites* are masses of mesoderm that can be found distributed along the two sides of the neural tube. They will eventually become dermis (dermatome), skeletal muscle (myotome), vertebrae (sclerotome), and tendons and cartilage (syndetome).

The mesoderm found lateral to the neural tube is called the paraxial mesoderm. It is separate from the chordamesoderm underneath the neural tube. The paraxial mesoderm is initially called the unsegmented
mesoderm in vertebrates, but is called the segmented mesoderm in chick embryos.



Somites: A dorsal view of a human embryo. The repetitive somites are marked with the older term primitive segments.

As the primitive streak regresses and the neural folds gather preceding the formation of the neural tube, the paraxial mesoderm divides into blocks called somites. Somites play a critical role in early development by participating in the specification of the migration paths of neural crest cells and spinal nerve axons.

Later in development, somites separate into four compartments:

Dermatome

The dermatome is the dorsal portion of the paraxial mesoderm somite. In the human embryo it arises in the third week of embryogenesis.

The dermatomes contribute to the skin, fat, and connective tissue of the neck and of the trunk, though most of the skin is derived from the lateral plate mesoderm.

Myotome

The myotome is that part of a somite that forms the muscles. Each myotome divides into an epaxial part (epimere), at the back, and a hypaxial part (hypomere) at the front.

The myoblasts from the hypaxial division form the muscles of the thoracic and anterior abdominal walls. The epaxial muscle mass loses its segmental character to form the extensor muscles of the neck and trunk of mammals.

Sclerotome

The sclerotome forms the vertebrae and the rib cartilage and part of the occipital bone. It forms the musculature of the back, the ribs, and the limbs.

Syndetome

The syndetome forms the tendons and some blood vessels.

This page titled 27.4C: Somite Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4D: Development of the Cardiovascular System

The circulatory system develops initially via vasculogenesis, with the arterial and venous systems developing from distinct embryonic areas.

Learning Objectives

- Outline the development of the cardiovascular system

Key Points


- The aortic arches are a series of six, paired, embryological vascular structures that give rise to several major arteries. The first and second arches disappear early. The third arch becomes the carotid artery.
- The fourth right arch forms the right subclavian artery, while the fourth left arch forms the arch of the aorta. The fifth arch disappears on both sides.The proximal part of the sixth right arch persists as the proximal right pulmonary artery. The sixth left arch gives off the left pulmonary artery.
- Approximately 30 posterolateral branches arise off the dorsal aortae and will form the intercostal arteries, the upper and lower extremity arteries, the lumbar arteries, and the lateral sacral arteries. The lateral branches of the aorta form the definitive renal, suprarenal, and gonadal arteries.
- The ventral branches consist of the vitelline and umbilical arteries. The vitelline arteries form the celiac, and superior and inferior mesenteric arteries of the gastrointestinal tract. After birth, the umbilical arteries will form the internal iliac arteries.
- The venous system develops from the vitelline veins, umbillical veins, and the cardinal veins, all of which empty into the sinus venosus.

Key Terms

- **sinus venosus**: A large quadrangular cavity that precedes the atrium on the venous side of the chordate heart. In humans, it exists distinctly only in the embryonic heart, where it is found between the two venae cavae.
- **aortic arches**: Also known as pharyngeal arch arteries, this is a series of six, paired, embryological vascular structures that give rise to several major arteries. They are ventral to the dorsal aorta.
- **cardinal vein**: The precardinal veins or anterior cardinal veins contribute to the formation of the internal jugular veins and, together with the common cardinal vein, form the superior vena cava. In an anastomosis by anterior cardinal veins, the left brachiocephalic vein is produced.

Vasculogenesis

The human arterial system originates from the aortic arches and from the dorsal aortae starting from week 4 of embryonic life. The development of the circulatory system initially occurs by the process of vasculogenesis, the formation of new blood vessels when there are no preexisting ones.




Embryonic cardiovascular system: A profile view of a human embryo estimated at twenty or twenty-one days old.

Vasculogenesis is when endothelial precursor cells (angioblasts) migrate and differentiate in response to local cues (such as growth factors and extracellular matrix) to form new blood vessels. The human arterial and venous systems develop from different embryonic areas.

Aortic Arches

The aortic arches—or pharyngeal arch arteries—are a series of six, paired, embryological vascular structures that give rise to several major arteries. They are ventral to the dorsal aorta and arise from the aortic sac.

Arches 1 and 2



Aortic arches: A schematic of the aortic arches and their arterial destinations.

The first and second arches disappear early, but the dorsal end of the second gives origin to the stapedial artery, a vessel that atrophies in humans, but persists in some mammals. It passes through the ring of the stapes and divides into supraorbital,

infraorbital, and mandibular branches that follow the three divisions of the trigeminal nerve.

The infraorbital and mandibular branches arise from a common stem, the terminal part of which anastomoses with the external carotid. On the obliteration of the stapedial artery, this anastomosis enlarges and forms the internal maxillary artery; the branches of the stapedial artery are now branches of this vessel.

The common stem of the infraorbital and mandibular branches passes between the two roots of the auriculotemporal nerve and becomes the middle meningeal artery. The original supraorbital branch of the stapedial artery is represented by the orbital branches of the middle meningeal artery.

Arches 3 and 4

The third aortic arch constitutes the commencement of the internal carotid artery, and is named the carotid arch. The fourth right arch forms the right subclavian artery as far as the origin of its internal mammary branch. The fourth left arch constitutes the arch of the aorta between the origin of the left carotid artery and the termination of the ductus arteriosus.

Arches 5 and 6

The fifth arch disappears on both sides.The proximal part of the sixth right arch persists as the proximal part of the right pulmonary artery, while the distal section degenerates. The sixth left arch gives off the left pulmonary artery and forms the ductus arteriosus.

This duct remains during fetal life, but closes within the first few days after birth due to increased O_2 concentration. This causes the production of bradykinin which causes the ductus to constrict, occluding all flow. Within one to three months, the ductus is obliterated and becomes the ligamentum arteriosum.

Aortic Branches

The dorsal aortae are initially bilateral and then fuse to form the definitive dorsal aorta. Approximately 30 posterolateral branches arise off the aorta and will form the intercostal arteries, upper and lower extremity arteries, lumbar arteries, and the lateral sacral arteries.

The lateral branches of the aorta form the definitive renal, suprarenal, and gonadal arteries. Finally, the ventral branches of the aorta consist of the vitelline arteries and umbilical arteries.

The vitelline arteries form the celiac, and superior and inferior mesenteric arteries of the gastrointestinal tract. After birth, the umbilical arteries will form the internal iliac arteries.

The human venous system develops mainly from the vitelline, umbilical, and cardinal veins, all of which empty into the sinus venosus. The venous system arises during the fourth to eighth weeks of human development.

CLINICAL EXAMPLE

Most defects of the great arteries arise as a result of the persistence of aortic arches that normally should regress or due to the regression of arches that normally should not.

A double aortic arch occurs with the development of an abnormal right aortic arch, in addition to the left aortic arch, forming a vascular ring around the trachea and esophagus, which usually causes difficulty breathing and swallowing.

Occasionally, the entire right dorsal aorta abnormally persists and the left dorsal aorta regresses. In this case, the right aorta will have to arch across from the esophagus, causing difficulty breathing or swallowing.

This page titled 27.4D: Development of the Cardiovascular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.4E: Chorionic Villi and Placental Development

In the placenta, chorionic villi develop to maximize surface-area contact with the maternal blood for nutrient and gas exchange.

Learning Objectives

- Summarize the development of the chorionic villi and placenta

Key Points


- Chorionic villi invade and destroy the uterine decidua while at the same time they absorb nutritive materials from it to support the growth of the embryo.
- The villi begin primary development in the fourth week, becoming fully vascularized between the fifth and sixth weeks.
- Placental development begins with implantation of the blastocyst; this leads to its differentiation into several layers that allow nutrient, gas, and waste exchange to the developing embryo and fetus —as well as forming a protective barrier.

Key Terms

- **chorion**: The protective and nutritive membrane that attaches higher vertebrate fetuses to the uterus.
- **uterine decidua**: The term for the uterine lining (endometrium) during a pregnancy, which forms the maternal part of the placenta. It is formed under the influence of progesterone and forms highly characteristic cells.
- **chorionic villi**: These sprout from the chorion in order to give a maximum area of contact with the maternal blood.
- **placenta**: A vascular organ present only in the female during gestation. It supplies food and oxygen from the mother to the fetus, and passes back waste. It is implanted in the wall of the uterus and links to the fetus through the umbilical cord. It is expelled after birth.

Chorionic Villi

Chorionic villi sprout from the chorion after their rapid proliferation in order to give a maximum area of contact with the maternal blood. These villi invade and destroy the uterine decidua while at the same time they absorb nutritive materials from it to support the growth of the embryo.



Chorionic artery: An image showing the chorionic villi and the maternal vessels.

During the primary stage (the end of fourth week), the chorionic villi are small, nonvascular, and contain only the trophoblast. During the secondary stage (the fifth week), the villi increase in size and ramify, while the mesoderm grows into them; at this point the villi contain trophoblast and mesoderm.

During the tertiary stage (fifth to sixth week), the branches of the umbilical vessels grow into the mesoderm; in this way, the chorionic villi are vascularized. At this point, the villi contain trophoblast, mesoderm, and blood vessels.

Embryonic blood is carried to the villi by the branches of the umbilical arteries. After circulating through the capillaries of the villi, it is returned to the embryo by the umbilical veins. Chorionic villi are vital in pregnancy from a histomorphologic perspective and are, by definition, products of conception.


Placenta

The placenta is a fetally derived organ that connects the developing fetus to the uterine wall to allow nutrient uptake, waste elimination, and gas exchange via the mother's blood supply. The placenta begins to develop upon implantation of the blastocyst into the maternal endometrium.

The placenta functions as a fetomaternal organ with two components: the fetal placenta (chorion frondosum), which develops from the same blastocyst that forms the fetus; and the maternal placenta (decidua basalis), which develops from the maternal uterine tissue.

The outer layer of the blastocyst becomes the trophoblast, which forms the outer layer of the placenta. This layer is divided into two further layers: the underlying cytotrophoblast layer and the overlying syncytiotrophoblast layer.

The latter is a multinucleated, continuous cell layer that covers the surface of the placenta. It forms as a result of the differentiation and fusion of the underlying cytotrophoblast cells, a process that continues throughout placental development. The syncytiotrophoblast (otherwise known as syncytium) thereby contributes to the barrier function of the placenta.



Placenta: Image illustrating the placenta and chorionic villi. The umbilical cord is seen connected to the fetus and the placenta.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Gastrulation. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Gastrulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- somite. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somite. **License**: *CC BY-SA: Attribution-ShareAlike*
- gastrulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/gastrulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- notochord. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/notochord. **License**: *CC BY-SA: Attribution-ShareAlike*
- Epiboly. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Epiboly. **License**: *CC BY-SA: Attribution-ShareAlike*
- Germ Layers. **Provided by**: Open Stax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Neural development in humans. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neural_...%23Neurulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- basal plate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/basal%20plate. **License**: *CC BY-SA: Attribution-ShareAlike*
- alar plate. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/alar%20plate. **License**: *CC BY-SA: Attribution-ShareAlike*
- neurulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- Neural Tube. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neural_tube. **License**: *CC BY-SA: Attribution-ShareAlike*
- Germ Layers. **Provided by**: Open Stax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Neurulation. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F....jpg#filelinks. **License**: *CC BY: Attribution*
- Spina_bifida-web.jpg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F...bifida-web.jpg. **License**: *Public Domain: No Known Copyright*
- Somite. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Somite%23In_vertebrates. **License**: *CC BY-SA: Attribution-ShareAlike*
- Somite. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Somite%23In_vertebrates. **License**: *CC BY-SA: Attribution-ShareAlike*
- somite. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somite. **License**: *CC BY-SA: Attribution-ShareAlike*
- neural crest cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/neural%20crest%20cells. **License**: *CC BY-SA: Attribution-ShareAlike*
- conceptus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/conceptus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Paraxial Mesoderm. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Paraxial_mesoderm. **License**: *CC BY-SA: Attribution-ShareAlike*
- Germ Layers. **Provided by**: Open Stax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Neurulation. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/F....jpg#filelinks. **License**: *CC BY: Attribution*
- Spina_bifida-web.jpg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Spina_bifida-web.jpg. **License**: *Public Domain: No Known Copyright*
- Gray20. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray20.png. **License**: *Public Domain: No Known Copyright*
- Circulatory system. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Circula...%23Development. **License**: *CC BY-SA: Attribution-ShareAlike*
- Aortic arches. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Aortic_arches. **License**: *CC BY-SA: Attribution-ShareAlike*
- Vasculogenesis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Vasculogenesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- aortic arches. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/aortic%20arches. **License**: *CC BY-SA: Attribution-ShareAlike*
- cardinal vein. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cardinal%20vein. **License**: *CC BY-SA: Attribution-ShareAlike*
- sinus venosus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sinus%20venosus. **License**: *CC BY-SA: Attribution-ShareAlike*
- Germ Layers. **Provided by**: Open Stax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Neurulation. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/

File:2912_Neurulation-02.jpg#filelinks. License: CC BY: Attribution
- Spina_bifida-web.jpg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Spina_bifida-web.jpg. **License**: *Public Domain: No Known Copyright*
- Gray20. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray20.png. **License**: *Public Domain: No Known Copyright*
- Gray472. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray472.png. **License**: *Public Domain: No Known Copyright*
- Gray473. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray473.png. **License**: *Public Domain: No Known Copyright*
- Hydatidiform mole. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hydatidiform_mole. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chorionic villi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chorionic_villi. **License**: *CC BY-SA: Attribution-ShareAlike*
- Chorionic villi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chorionic_villi%23Development. **License**: *CC BY-SA: Attribution-ShareAlike*
- Placenta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Placenta. **License**: *CC BY-SA: Attribution-ShareAlike*
- Placenta. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Placenta%23Development. **License**: *CC BY-SA: Attribution-ShareAlike*
- chorionic villi. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/chorionic%20villi. **License**: *CC BY-SA: Attribution-ShareAlike*
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/definition/chorion. **License**: *CC BY-SA: Attribution-ShareAlike*
- placenta. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/placenta. **License**: *CC BY-SA: Attribution-ShareAlike*
- uterine decidua. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/uterine%20decidua. **License**: *CC BY-SA: Attribution-ShareAlike*
- Germ Layers. **Provided by**: Open Stax College. **Located at**: http://openstaxcollege.org. **License**: *CC BY: Attribution*
- Neurulation. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:2912_Neurulation-02.jpg#filelinks. **License**: *CC BY: Attribution*
- Spina_bifida-web.jpg. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Spina_bifida-web.jpg. **License**: *Public Domain: No Known Copyright*
- Gray20. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray20.png. **License**: *Public Domain: No Known Copyright*
- Gray472. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray472.png. **License**: *Public Domain: No Known Copyright*
- Gray473. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray473.png. **License**: *Public Domain: No Known Copyright*
- Chorionic Vessels. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Chorionic_vessels. **License**: *Public Domain: No Known Copyright*
- Placenta. **Provided by**: Wikipedia Commons. **Located at**: commons.wikimedia.org/wiki/File:Placenta.png. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 27.4E: Chorionic Villi and Placental Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.5: FOURTH WEEK OF DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.5A: Fourth Week of Development

This page titled 27.5: Fourth Week of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.5A: Fourth Week of Development

The fourth week of gestation is characterized by the flexion of the superior portion of the neural tube to create the mesencephalon.

Learning Objectives

- Describe the fourth week of embryonic development

Key Points

- Neural tube flexion patterns determine neural development, including the positioning and differentiation of the prosencephalon and rhombencephalon, as well as the optical vesicle.
- Important changes occur to the embryonic heart as well, including development of the pharyngeal arches.
- At the end of the fourth week, the yolk sac presents the appearance of a small pear-shaped vesicle (the umbilical vesicle) that opens into the digestive tube by a long, narrow tube—the vitelline duct.

Key Terms

- **mesencephalon**: A part of the brain located rostral to the pons and caudal to the thalamus and the basal ganglia, composed of the tectum (dorsal portion) and the tegmentum (ventral portion).
- **mandibular arch**: The first pharyngeal arch, also called the mandibular arch, is the first of six aortic arches that develops in fetal life during the fourth week of development.
- **optical vesicle**: The eyes begin to develop as a pair of diverticula from the lateral aspects of the forebrain. These diverticula make their appearance before the closure of the anterior end of the neural tube; after the closure of the tube they are known as the optic vesicles.


Placentation

Trophoblast cells surrounding the embryonic cells proliferate and invade deeper into the uterine lining. They will eventually form the placenta and embryonic membranes. At the end of the fourth week the yolk sac presents the appearance of a small pear-shaped vesicle (the umbilical vesicle) opening into the digestive tube by a long narrow tube, the vitelline duct.

The chorion undergoes rapid proliferation and forms numerous processes. The chorionic villi, which invade and destroy the uterine decidua and at the same time absorb from it nutritive materials for the growth of the embryo. Until about the end of the second month of pregnancy the villi cover the entire chorion, and are almost uniform in size, but after this they develop unequally.

Mesencephalon Development

Late in the fourth week of gestation, the superior part of the neural tube flexes at the level of the future midbrain, the mesencephalon. Superior to the mesencephalon is the prosencephalon (future forebrain) and inferior to it is the rhombencephalon (future hindbrain). The optical vesicle (which will eventually become the optic nerve, retina, and iris) forms at the basal plate of the prosencephalon.




Embryo brain at four weeks: At four weeks the embryo's brain can be differentiated into the proscephalon, mesencephalon, and rhombencephalon. A white circle represents the area of the optical vesicle.

Pharyngeal Arch Development

The embryonic heart attains functionality and starts beating at 22 days after conception (about five weeks after the last menstrual period). It can sometimes be seen as flickering in the embryonic chest by an ultrasound performed during the fourth week after conception.

During the fourth week of development, the first pharyngeal arch, also called the mandibular arch, is the first of six pharyngeal arches that begins to develop. It is located between the stomodeum and the first pharyngeal groove.



Developing fetus: A schematic of a developing fetus with the first, second, and third arches labeled.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- First pharyngeal arch. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/First_pharyngeal_arch**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Prenatal development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Prenatal_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural development in humans. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neural_development_in_humans%23Formation_of_the_brain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Yolk sac. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Yolk_sac**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Embryonic heart. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryonic_heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mesencephalon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mesencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mandibular arch. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/mandibular%20arch**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- optical vesicle. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/optical%20vesicle**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pharyngeal Arch. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pharyngeal_arch**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Placentation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Placentation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray41.png. **Provided by**: Wikipedia Commons. **Located at**: **en.Wikipedia.org/wiki/File:Gray41.png**. **License**:
- 4 week embryo brain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:4_week_embryo_brain.jpg**. **License**:

This page titled 27.5A: Fourth Week of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.6: FIFTH THROUGH EIGHTH WEEKS OF DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.6A: Fifth Through Eighth Weeks of Development

This page titled 27.6: Fifth Through Eighth Weeks of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.6A: Fifth Through Eighth Weeks of Development

Weeks five to eight of gestation develops the major organs, including the circulatory, nervous, and gastrointestinal systems.

Learning Objectives

- Outline embryonic development between five and eight weeks

Key Points

- Being susceptible to the effects of teratogens is high during embryonic development.
- At week five, the brain, spinal cord, vertebrae, heart, vasculature, and gastrointestinal tract begin to develop.
- During weeks six to seven, the embryo grows from 4 mm in length to 9 mm and begins to curve into a C-shape. The fetal heart bulges, develops further, and begins to beat in a regular rhythm. Rudimentary blood begins to move through the blood vessels. The neural tube, which forms the brain, closes.
- During weeks six and seven, the limb buds form. The eyes, mouth, and ear structures begin to form. The initial differentiation of the tissues that will become the spleen, gallbladder, pancreas, liver, kidneys, stomach, and lungs occurs.
- By week eight of gestation, the lungs begin to form, as well as the lymphatic system. The main development of the external genitalia begins, and the brain continues to develop. The arms and legs have grown longer, and the foot and hand areas can be clearly distinguished.

Key Terms

- **chorionic membrane**: One of the membranes that exist during pregnancy between the developing fetus and mother. The chorionic villi emerge from the chorion, invade the endometrium, and allow transfer of nutrients from maternal blood to fetal blood.
- **Gestational age**: The time that has passed since the onset of the last menstruation.
- **Embryonic age**: Measures the actual age of the embryo or fetus from the time of fertilization.
- **teratogens**: An agent, such as a virus, a drug, or radiation, that causes a malformation of an embryo or fetus.


Weeks five through eight of gestation are characterized by the development of the major organ systems, including the circulatory, nervous, lymphatic, and gastrointestinal systems. During this time, the embryo is extremely susceptible to the effects of teratogens.

Gestational age is the time that has passed since the onset of the last menstruation, which occurs two weeks before the actual fertilization. Embryonic age measures the actual age of the embryo or fetus from the time of fertilization. Thus, the first week of embryonic age is already week three counting with gestational age. The number of the week (used here) is one more than the actual age of the embryo/fetus. For example, the embryo is 0 whole weeks old during the first week after fertilization.

Week 5

At week five, the brain, spinal cord, vertebrae, heart, vasculature, and gastrointestinal tract begin to develop.

Week 6–7



Human embryo at seven weeks: An embryo from an ectopic pregnancy, still in the oviduct. This embryo is about five weeks old (or from the seventh week of menstrual age). The heart is the dark spot at the center of the image, bulging out of the embryo.

During weeks six and seven, the embryo grows from four millimeters in length to nine millimeters and begins to curve into a C-shape. The fetal heart bulges, develops further, and begins to beat in a regular rhythm. Rudimentary blood begins to move through the main embryonic blood vessels, connecting to the yolk sac and the chorionic membrane of the placenta.

The arm and leg buds, which will grow into the full limbs over the rest of development, become visible. The neural tube, which forms the brain, closes. The brain then develops into five areas and some cranial nerves are visible.

The eyes, mouth, and ear structures begin to form. The initial differentiation of the tissues that will become the spleen, gallbladder, pancreas, liver, kidneys, stomach, and lungs occurs.

Week 8

By week eight of gestation, the embryo measures 13 millimeters in length. The lungs begin to form, as well as the lymphatic system. The main development of the external genitalia begins, and the brain continues to develop.

The arms and legs have grown longer, and the foot and hand areas can be clearly distinguished. The hands and feet have fingers and toes, but may still be webbed.



Human embryo from an ectopic pregnancy: An embryo from an ectopic pregnancy, located in the part of the uterus to which the fallopian tube is attached. The features are consistent with a developmental age of seven weeks (the ninth week of menstrual age).

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Fetal development: MedlinePlus Medical Encyclopedia. **Provided by**: National Insitute of Health. **Located at**: **http://www.nlm.nih.gov/medlineplus/e...cle/002398.htm**. **License**: ***Public Domain: No Known Copyright***
- teratogens. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/teratogens**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chorionic membrane. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/chorionic%20membrane**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Prenatal Development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Prenatal_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Tubal Pregnancy with embryo. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Tu...ith_embryo.jpg**. **License**: ***Public Domain: No Known Copyright***
- Human_Embryo_from_Ectopic_Pregnancy.jpg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:9-Week_Human_Embryo_from_Ectopic_Pregnancy.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 27.6A: Fifth Through Eighth Weeks of Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.7: BIRTH (PARTURITION)

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.7A: Onset of Labor
27.7B: Stages of Labor
27.7C: Adjustments of the Infant at Birth

This page titled 27.7: Birth (Parturition) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.7A: Onset of Labor

Prodromal labor, which includes the latent phase of labor, marks the initial stages of parturition.

Learning Objectives

- Describe prodromal labor

Key Points

- Prodromal labor, the body's preparation for real labor, may include an increase in blood volume, Braxton Hicks contractions, the presence of colostrum in the breasts, and the dislodging of the mucus plug.
- In contrast to Braxton Hicks contractions, the latent phase is marked by the point at which the woman perceives regular uterine contractions; cervical effacement will reach completion at the end of this stage.
- The latent phase ends with the onset of active first stage, which is marked by advanced dilation.
- The duration of labor varies widely, but its active phase averages some 20 hours for women giving birth to their first child (primiparae) and 8 hours for women who have already given birth (multiparae).
- In a caesarean section, the removal of the neonate is through a surgical incision in the abdomen, rather than through vaginal birth.

Key Terms

- **caesarean section**: A caesarean section, also known as C-section, is a surgical procedure in which one or more incisions are made through a mother's abdomen (laparotomy) and uterus (hysterotomy) to deliver one or more babies, or, rarely, to remove a dead fetus.
- **Braxton Hicks contractions**: Known as false labor or practice contractions. They are sporadic uterine contractions that sometimes start around six weeks but are not usually felt until the second or third trimester of pregnancy.
- **cervical effacement**: The thinning and stretching of the cervix.
- **parturition**: The act or process of bearing or bringing forth offspring from the uterus; giving birth.

Pre-labor (First Stage of Labor)

Pre-labor, also called prodromal labor, consists of the early contractions and labor signs before actual labor starts. It is the body's preparation for real labor.

Prodromal labor, often misnamed false labor, begins much as traditional labor but does not progress to the birth of the baby. Not everyone feels this stage of labor although it does always occur. However, this does not mean that every woman will experience every symptom.

The term is used to describe a cluster of physical changes that may take place in a pregnant woman before she goes into actual labor. These changes can include:

- An increase in blood volume (sometimes resulting in edema).
- Braxton Hicks contractions.
- The presence of colostrum in the breasts.
- The dislodging of the mucous plug that sealed the cervix during the pregnancy.

The term false labor is sometimes used to describe a cluster of Braxton Hicks contractions that are mistaken for real labor. The terms false labor and false pains are sometimes considered equivalent.

Latent Phase

The latent phase is generally defined as beginning at the point at which the woman perceives regular uterine contractions. In contrast, Braxton Hicks contractions should be infrequent, irregular, and involve only mild cramping.

Cervical effacement (the thinning and stretching of the cervix and cervical dilation) occurs during the closing weeks of pregnancy and is usually complete, or near complete, by the end of the latent phase.

When the contractions of labor begin, the walls of the uterus start to contract through stimulation by the release of the pituitary hormone oxytocin. The contractions cause the cervix to widen and begin to open.

As labor progresses the amniotic sac can rupture and cause a slow or a fast gush of fluids. Labor usually begins within a 24-hour period after the amniotic sac ruptures. As contractions become closer and stronger the cervix will gradually start to dilate.

The degree of cervical effacement may be felt during a vaginal examination. A long cervix implies that effacement has not yet occurred. At this stage, the cervix may dilate from 1 to 4 cm. The latent phase ends with the onset of the active phase, which is marked by an accelerated cervical dilation.

This page titled 27.7A: Onset of Labor is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.7B: Stages of Labor

Vaginal delivery childbirth has three distinct phases: dilation of the cervix, delivery of the infant, and delivery of the placenta.

Learning Objectives

- Differentiate among the stages of labor

Key Points

- The first stage of labor begins when the effaced (thinned) cervix is 3 cm dilated. During effacement, the cervix is pulled upward during contractions. Full dilation occurs when the cervix is wide enough to allow passage of the baby's head, about 10 cm for a full-term infant.
- The duration of labor varies widely, but the active phase averages some 20 hours for women giving birth to their first child (primiparae) and 8 hours for women who have previously given birth (multiparae).
- The second stage begins when the cervix is fully dilated, ending when the baby is born. Fetal movement through the birth canal is assisted by the additional maternal efforts of bearing down or pushing. The fetal head is seen to crown as the labia part, and the baby is born.
- The period from just after the fetus is expelled until just after the placenta is expelled is called the third stage of labor, which lasts, on average, 10–12 minutes.
- A fourth stage of labor may refer to the hour immediately after delivery of the placenta, or may refer to the weeks following delivery. It is the time in which the mother's body returns to a nonpregnant state.

Key Terms

- **multiparae**: Women who have given birth to more than one viable fetus.
- **Ferguson reflex**: An example of positive feedback and the female body's response to pressure application in the cervix or vaginal walls. Upon application of pressure, oxytocin is released and uterine contractions are stimulated (which will in turn increase oxytocin production, and hence, increase contractions even more), until the baby is delivered.
- **primiparae**: Women who have given birth to only one child, or who are giving birth for the first time.

The infant's head and shoulders must go through a specific sequence of maneuvers in order to pass through the ring of the mother's pelvis. The six phases of a typical vertex (head-first presentation) delivery are:

1. Engagement of the fetal head in the transverse position.
2. Descent and flexion of the fetal head.
3. Internal rotation of the fetal head so that the baby's face is towards the mother's rectum.
4. Delivery by extension (the fetal head passes out of the birth canal).
5. Restitution: The fetal head turns through 45 degrees to restore its normal relationship with the shoulders, which are still at an angle.
6. External rotation: The shoulders repeat the corkscrew movements of the head, which can be seen in the final movements of the fetal head.

The Bishop score defines several factors that midwives and physicians use to assess the laboring mother's progress.
The score is used to predict whether the mother is likely to spontaneously progress into the second stage of labor and whether induction of labor will be required. It has also been used to assess the odds of spontaneous preterm delivery.

Stages of Labor

First Stage (Active Phase)

The first stage of labor classically starts when the effaced (thinned) cervix is 3 cm dilated, although there is variation as some women may or may not have active contractions prior to reaching this point. The onset of actual labor is defined when the cervix begins to progressively dilate. Rupture of the membranes or a bloody discharge may or may not occur at or around this stage.

Tterine muscles form opposing spirals from the top of the upper segment of the uterus to its junction with the lower segment. During effacement, the cervix becomes incorporated into the lower segment of the uterus. During a contraction, these muscles shorten the upper segment, drawing upwards the lower segment in a gradual expulsive motion.

The presenting fetal part is then permitted to descend. Full dilation is reached when the cervix has widened enough to allow passage of the baby's head, around 10 cm dilation for a term baby. The duration of labor varies widely, but the active phase averages some 20 hours for women giving birth to their first child (primiparae), and 8 hours for women who have already given birth (multiparae).

Active phase arrest is defined, in a primigravid woman, as the failure of the cervix to dilate at a rate of 1.2 cm/hr over a period of at least two hours. This definition is based on Friedman's Curve, which plots the typical rate of cervical dilation and fetal descent during active labor. Some practitioners may diagnose failure to progress, and consequently, perform a caesarean.

Second Stage (Fetal Expulsion)

The second stage begins when the cervix is fully dilated and ends when the baby is born. As pressure on the cervix increases, the Ferguson reflex increases uterine contractions.

At the beginning of the normal second stage, the head is fully engaged in the pelvis: the widest diameter of the head has passed below the level of the pelvic inlet. The fetal head then continues descending into the pelvis, below the pubic arch, and out through the vagina.

This is assisted by the additional maternal efforts of bearing down or pushing. The fetal head is seen to crown as the labia part. At this point, the woman may feel a burning or stinging sensation. The complete expulsion of the baby signals the successful completion of the second stage of labor.

The second stage of birth will vary by factors including parity, fetal size, anesthesia, or the presence of infection. Longer labors are associated with declining rates of spontaneous vaginal delivery and increasing rates of infection, perineal laceration, obstetric hemorrhage, as well as need for intensive care of the neonate.

Third Stage (Placental Delivery)

The third stage of labor is the period from just after the fetus is expelled until just after the placenta is expelled. The average time from delivery of the baby until complete expulsion of the placenta is estimated to be 10–12 minutes.


Placental expulsion can be managed actively, by giving a uterotonic, such as oxytocin, along with appropriate cord traction and fundal massage to assist in delivering the placenta by a skilled birth attendant. Alternatively, it can be managed expectantly, allowing the placenta to be expelled without medical assistance. The umbilical cord is routinely clamped and cut in this stage, but it would normally close naturally even if not clamped and cut.

When the amniotic sac has not ruptured during labour or pushing, the infant can be born with the membranes intact. This is referred to as being born in the caul. The caul is harmless and its membranes are easily broken and wiped away. With the advent of modern interventive obstetrics, the artificial rupture of the membranes has become common, so babies are rarely born in the caul (en-caul birth).

Fourth Stage (Postpartum Period)

The fourth stage of labor is a term used in two different senses:

1. It can refer to the immediate puerperium, or the hours immediately after delivery of the placenta.
2. It can be used in a more metaphorical sense to describe the weeks following delivery.



Birth.: A sequence of images showing the stages of ordinary childbirth.

This page titled 27.7B: Stages of Labor is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.7C: Adjustments of the Infant at Birth

Post birth, an infant's physiology must adapt to breathing independently, changes in blood flow and energy access, and a cold environment.

Learning Objectives

- Analyze the perinatal adjustments of the infant

Key Points

- Following birth, the newborn's lungs inflate and respiration begins to occur through the pulmonary system. The subsequent changes to oxygenated blood flow that occur follow the adult pattern of circulation.
- The energy metabolism of the infant is also quickly converted from a continuous placental supply to intermittent feeding. Glycogen stores created during the second and third trimesters of pregnancy are relied upon for energy during the first few days following birth.
- Physiological changes occur via thermoregulation to maintain core temperature including: vasoconstriction (a decrease of blood flow to skin), maintaining the fetal position (a decrease of the surface area of exposure), jittery large muscle activity (to generate muscular heat), and non-shivering thermogenesis.

Key Terms

- **respiratory distress syndrome**: Infant respiratory distress syndrome (IRDS), also called neonatal respiratory distress syndrome or respiratory distress syndrome (previously called hyaline membrane disease), is a syndrome in premature infants caused by the developmental insufficiency of surfactant production and structural immaturity in the lungs. It can also result from a genetic problem with the production of surfactant-associated proteins.
- **surfactant**: A lipoprotein in the tissues of the lung that reduces surface tension and permits more efficient gas transport.
- **non-shivering thermogenesis**: The process of heat production in organisms. Non-shivering thermogenesis usually occurs in brown adipose tissue (brown fat) that is present in human infants and hibernating mammals.

The first challenge of a newborn is to perfuse its body by breathing independently instead of utilizing placental oxygen. At birth, the baby's lungs are filled with fetal lung fluid (which is not amniotic fluid) and are not inflated.

As the newborn is expelled from the birth canal, its central nervous system reacts to the sudden change in temperature and environment. This triggers it to take the first breath within about 10 seconds of delivery.

With the first breaths, there is a fall in pulmonary vascular resistance and an increase in the surface area available for gas exchange. Over the next 30 seconds, the pulmonary blood flow increases and is oxygenated as it flows through the alveoli of the lungs. Oxygenated blood now reaches the left atrium and ventricle and, through the descending aorta, reaches the umbilical arteries.

Respiration and Circulation

Oxygenated blood now stimulates constriction of the umbilical arteries resulting in a reduction in placental blood flow. As the pulmonary circulation increases, there is an equivalent reduction in the placental blood flow that normally ceases completely after about three minutes. These two changes result in a rapid redirection of blood flow into the pulmonary vascular bed, from approximately 4% to 100% of cardiac output.

The increase in pulmonary venous return results in left atrial pressure being slightly higher than right atrial pressure, which closes the foramen ovale. This change in the pattern of flow results in a drop in blood flow across the ductus arteriosus.

The higher blood oxygen content of blood within the aorta stimulates the constriction and ultimately the closure of this fetal

circulatory shunt. All of these cardiovascular system changes result in the adaptation from fetal circulation patterns to an adult circulation pattern.

During this transition, some types of congenital heart diseases that were not symptomatic in utero during fetal circulation will present with cyanosis or respiratory signs.

Following birth, the expression and re-uptake of surfactant, which begins production at 20 weeks gestation, is accelerated. Expression of surfactant into the alveoli is necessary to prevent alveolar closure.

At this point, rhythmic breathing movements also commence. If there are any problems with breathing, management can include stimulation, bag and mask ventilation, intubation, and ventilation. Cardiorespiratory monitoring is essential to track potential problems.

Pharmacological therapy, such as caffeine, can also be given to treat apnea in premature newborns. A positive airway pressure should be maintained and neonatal sepsis must be ruled out. Potential neonatal respiratory problems include apnea, transient tachypnea of the newborn (TTNB), respiratory distress syndrome (RDS), meconium aspiration syndrome (MAS), airway obstruction, and pneumonia.

Energy Requirements

The energy metabolism of the fetus must be converted from a continuous placental supply of glucose to intermittent feeding. While the fetus is dependent on maternal glucose as the main source of energy, it can use lactate, free fatty acids, and ketone bodies under some conditions. Plasma glucose is maintained by glycogenolysis.

Glycogen synthesis in the liver and muscle begins in the late second trimester of pregnancy, and storage is completed in the third trimester. Glycogen stores are maximal at term, but even then, the fetus only has enough glycogen available to meet its energy needs for 8–10 hours, which can be depleted even more quickly if the demand is high.

Newborns will then rely on gluconeogenesis for energy. Fat stores are the largest storage source of energy. At 27 weeks gestation, only 1% of a fetus' body weight is fat. At 40 weeks, that number increases to 16%.

Inadequate glucose substrate can lead to hypoglycemia, fetal growth restriction, preterm delivery, or other problems. Similarly, an excess substrate can also lead to problems, such as infant of a diabetic mother (IDM), hypothermia, or neonatal sepsis.

Anticipating potential problems is the key to managing most neonatal problems of energy metabolism. For example, early feeding in the delivery room or as soon as possible may prevent hypoglycemia. If the blood glucose is still low, then an intravenous (IV) bolus of glucose may be delivered, with continuous infusion if necessary. Rarely, steroids or glucagon may have to be employed.

Environmental Temperature



Infant in incubator: Infants, especially preterm infants, have trouble regulating their body temperature. Placement in an incubator allows them to adjust without draining their energy stores.

Newborns come from a warm environment to the cold and fluctuating temperatures of this world. They are naked, wet, and have a large surface area to mass ratio, with variable amounts of insulation, limited metabolic reserves, and a decreased ability to shiver.

Physiologic mechanisms for preserving core temperature include vasoconstriction (a decrease of blood flow to the skin), maintaining the fetal position (a decrease of the surface area exposed to the environment), jittery large muscle activity (to generate muscular heat), and non-shivering thermogenesis.

Non-shivering thermogenesis occurs in brown fat. Brown fat is specialized adipose tissue with a high concentration of mitochondria designed to rapidly oxidize fatty acids in order to generate metabolic heat. The newborn's capacity to maintain these mechanisms is limited, especially in premature infants.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Pre-labor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pre-labor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Childbirth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Childbirth**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Pregnancy and birth. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P...abor_and_Birth**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Braxton Hicks contractions. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Braxton...20contractions**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parturition. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/parturition**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cervical effacement. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cervical%20effacement**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- caesarean section. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/caesarean%20section**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- primiparae. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/primiparae**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- Childbirth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Childbi...tage:_dilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ferguson reflex. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ferguson%20reflex**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- multiparae. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/multiparae**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bishop Score. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bishop_score**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- birth.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Childbirth#/media/File:2920_Stages_of_Childbirth-02.jpg**. **License**: ***CC BY: Attribution***
- Adaptation to extrauterine life. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Adaptat...nd_circulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- non-shivering thermogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/non-shi...0thermogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- surfactant. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/surfactant**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- respiratory distress syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/respira...ess%20syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- birth.jpg. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Childbirth#/media/File:2920_Stages_of_Childbirth-02.jpg**. **License**: ***CC BY: Attribution***
- Human Infant in Incubator. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi..._Incubator.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 27.7C: Adjustments of the Infant at Birth is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.8: LACTATION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

27.8A: Physiology of Lactation

This page titled 27.8: Lactation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.8A: Physiology of Lactation

Lactation is the secretion of milk from specialized glands (mammary glands) to provide nourishment to offspring.

Learning Objectives

- Outline the process of lactation

Key Points

- Lactation is a hallmark feature of female mammals.
- Lactation is under endocrine control. The two main hormones involved are prolactin and oxytocin.
- Lactogenesis, or the process of changes to the mammary glands to begin producing milk, begins during the late stages of pregnancy. The delivery of the placenta and the resulting dramatic reduction in progesterone, estrogen, and human placental lactogen levels stimulate milk production.
- Colostrum is the first milk a breastfed baby receives. It contains higher amounts of white blood cells and antibodies than mature milk and is especially high in immunoglobulin A. This immunoglobulin coats the lining of the baby's immature intestines, helping to prevent pathogens from invading the baby's system.

Key Terms

- **witch's milk**: Witch's milk or neonatal milk is milk secreted from the breasts of some newborn human infants of either sex. Neonatal milk secretion is considered a normal physiological occurrence and no treatment or testing is necessary.
- **mammary gland**: A gland that secretes milk for suckling an infant or offspring.
- **lactation**: 1. The secretion of milk from the mammary gland of a female mammal. 2. The process of providing the milk to the young, such as breastfeeding. 3. The period of time that a mother lactates to feed her young; the lactation period.
- **human placental lactogen**: A hormone closely associated with prolactin that is instrumental in breast, nipple, and areola growth before birth.
- **colostrum**: A form of milk produced by the mammary glands in late pregnancy and the few days after giving birth. Human and bovine colostrum is thick and yellowish. In humans, it has high concentrations of nutrients and antibodies, but it is small in quantity.

Overview of Lactation

Lactation describes the secretion of milk from the mammary glands and the period of time that a mother lactates to feed her young. The process occurs in all female mammals, although it predates the origin of mammals.

In humans the process of feeding milk is called breastfeeding or nursing.
The chief function of lactation is to provide nutrition and immune protection to the young after birth. In almost all mammals, lactation induces a period of infertility, which serves to provide the optimal birth spacing for survival of the offspring.

In most species, milk comes out of the mother's nipples; however, the platypus (a non-placental mammal) releases milk through ducts in its abdomen. In only one species of mammal, the dayak fruit bat, is milk production a normal male function.

In some other mammals, the male may produce milk as the result of a hormone imbalance. This phenomenon may also be observed in newborn infants as well (for instance, witch's milk).

Galactopoiesis is the maintenance of milk production. This stage requires prolactin and oxytocin.

Preparation for Lactation

By the fifth or sixth month of pregnancy, the breasts are ready to produce milk. During the latter part of pregnancy, the woman's breasts enter into the lactogenesis I stage. This is when the breasts make colostrum, a thick, sometimes yellowish fluid.


At this stage, high levels of progesterone inhibit most milk production. It is not a medical concern if a pregnant woman leaks any colostrum before her baby's birth, nor is it an indication of future milk production.

At birth, prolactin levels remain high, while the delivery of the placenta results in a sudden drop in progesterone, estrogen, and human placental lactogen levels. This abrupt withdrawal of progesterone in the presence of high prolactin levels stimulates the copious milk production of the lactogenesis II stage.

When the breast is stimulated, prolactin levels in the blood rise and peak in about 45 minutes, then return to the pre-breastfeeding state about three hours later. The release of prolactin triggers the cells in the alveoli to make milk.

Colostrum

Colostrum is the first milk a breastfed baby receives. It contains higher amounts of white blood cells and antibodies than mature milk, and is especially high in immunoglobulin A (IgA), which coats the lining of the baby's immature intestines, and helps to prevent pathogens from invading the baby's system. Secretory IgA also helps prevent food allergies. Over the first two weeks after the birth, colostrum production slowly gives way to mature breast milk.



Lactation: A positive feedback loop ensures continued milk production as long as the infant continues to breastfeed.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- mammary gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mammary_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lactation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lactati...uman_lactation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lactation**. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- colostrum. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/colostrum**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Witch's Milk. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Witch%27s_milk**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Lactation. **Provided by**: Opens Stax College. **Located at**: **http://openstaxcollege.org**. **License**: ***CC BY: Attribution***

This page titled 27.8A: Physiology of Lactation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.9: THE FETAL PERIOD

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

27.9A: Fetal Development
27.9B: Teratogens
27.9C: Prenatal Diagnostic Tests

This page titled 27.9: The Fetal Period is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.9A: Fetal Development

At the end of the 10th week of gestation, the fetal period begins.

Learning Objectives

- Outline the progression of fetal development from 11 weeks to 40 weeks

Key Points

- Although fetuses can be adversely affect by environmental toxicants, they are less sensitive to them than embryos because the precursors for most organs are already formed.
- At the start of the fetal stage, the fetus is typically about 30 millimeters in length from crown to rump, and weighs about eight grams. The head makes up nearly half of the fetus' size.
- The precursors of all the major organs are created by the time the fetal period begins. Therefore, the fetal period is described both by organ and a list of the changes by weeks of gestational age.

Key Terms

- **meconium**: The earliest stool of a mammalian infant. Unlike later feces, meconium is composed of materials ingested during the time the infant spends in the uterus (intestinal epithelial cells, lanugo, mucus, amniotic fluid, bile, and water).
- **gestational age**: This relates to the age of an embryo or fetus (or newborn infant). In human obstetrics, this age is often defined as the time elapsed since 14 days prior to fertilization; this is approximately the duration since the woman's last menstrual period began.
- **lanugo**: Soft down or fine hair that covers the human fetus.

In humans, the fetal stage of prenatal development starts at the beginning of the 11th week in gestational age, which is the ninth week after fertilization. Since the precursors of all the major organs are created by this time, the fetal period is described both by organ and a list of changes by weeks of gestational age.

All major structures are already formed in the fetus, but they continue to grow and develop. Therefore, the fetus is not as sensitive to damage from environmental exposures as the embryo, though toxic exposures often cause physiological abnormalities, growth retardation, or minor congenital malformations.

At the start of the fetal stage, the fetus is typically about 30 millimeters in length from crown to rump and weighs about eight grams. The head makes up nearly half of the fetus' size. The four-chamber heart is finishing development, and the embryonic tail goes away. The breathing-like movement of the fetus is necessary for stimulation of lung development, rather than for obtaining oxygen.

- **Week 10:** Finger nails and hair start to grow. The heart, hands, feet, brain, and other organs are present, but are only at the beginning of development and have minimal operation.
- **Week 11:** Nearly all structures and organs are formed. Fingers and toes are separated and the genitals begin to take on the proper gender characteristics.
- **Week 12:** The digestive system and liver function. The pancreas makes insulin.
- **Week 13:** The fetus begins to get its nourishment from the placenta and the veins and organs are visible through the skin.
- **Week 14:** The kidneys produce urine, and the liver makes bile. In boys, the prostate gland develops, while in girls the ovaries move from the abdomen to the pelvis.
- **Weeks 15 to 16:** The heart pumps out 25 quarts of blood a day and the fetal structures are looking more normal.
- **Week 17:** The fetus start to move its joints and the retina becomes sensitive to light. It weighs about five ounces.
- **Week 18:** The fetus starts to hear and is startled by noise. Its skin is starting to grow a protective, wax-like layer and tiny air spaces begin to form in the lungs and the vocal cords.
- **Week 19:** The brain is designating areas for the five senses.
- **Week 20:** The fetus weighs about 10 ½ oz, swallows more, and produces meconium.

- **Week 21:** The eyebrows and lids are present, and for the female fetus, the vagina begins to form.
- **Week 22:** Tiny tooth buds beneath the gums develops. The eyes form, but the irises lack pigment.
- **Week 23:** The fetus weighs over a pound. It can feel movements and hear sounds. Blood vessels in his/her lungs are developing to prepare for breathing.
- **Week 24:** The fetus is almost a foot long. The lungs are developing branches of the respiratory tree as well as cells that produce surfactant, a substance that will help the air sacs inflate once born.
- **Week 25:** The amount of body fat rapidly increases. The lungs are not fully mature. The bones are fully developed, but are still soft and pliable. Thalamic brain connections form that mediate sensory input. Iron, calcium, and phosphorus become more abundant. The fingernails reach the end of the fingertips. The lanugo, or fine hair, begins to disappear, until it is gone (except on the upper arms and shoulders). Small breast buds are present on both sexes. Head hair becomes coarse and thicker.

Birth is imminent and occurs around the 40th week. The fetus is considered full-term between weeks 37 and 40, which means that the fetus is considered sufficiently developed for life outside the uterus.

This page titled 27.9A: Fetal Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.9B: Teratogens

Learning Objectives

Differentiate among teratogens that could negatively impact fetal development

A teratogen is a compound that permanently deforms the function or structure of a developing embryo or fetus in utero. In general, the degree of teratogenicity depends on:


- The potency of the drug as a mutagen.
- The susceptibility of the fetus to teratogenesis.
- The dose of the teratogen.
- The duration of teratogen exposure.
- The time of exposure.
- The degree of transfer from maternal to fetal circulation.

The global average of all live births complicated by malformation is 6% (Environmental Health Perspectives, (NIH), October 2009). The majority of these complications are due to unknown factors. The vast majority of recognized etiologies are genetic, with only 10% being attributed to environmental etiologies such as maternal health, infection, and toxicants. In general, the central nervous and skeletal systems are the most affected.

Women may encounter a number of teratogens. Smoking is most likely to cause growth retardation, but has also been implicated in the prelabor rupture of the membranes, preterm labor, abruption of the placenta, spontaneous abortion, perinatal morbidity and mortality, and sudden infant death syndrome. Smoking may exert its effects through competitive binding of carbon monoxide with hemoglobin and/or through the various other components found in cigarettes that cause adverse biological effects.

Examples

Alcohol use in pregnancy may result in fetal alcohol syndrome (FAS), which occurs in approximately 1% of all births. Children with FAS present with a flattened and thin upper lip, small palpebral fissures, epicanthal folds, flattened nasal bridge, and a short nose. They may also exhibit microcephaly, mental retardation, and have learning disabilities. It is not clear if there is any safe amount of alcohol consumption in pregnancy.



Fetal alcohol syndrome: *Alcohol is a teratogen. When consumed in pregnancy, it can result in mothers giving birth to children*

with fetal alcohol syndrome. The facial characteristics highlighted in the photograph are small eye opening, smooth philtrum, and a thin upper lip.

Cocaine generally produces growth restriction, preterm delivery, microcephaly, spontaneous abortion, placental abruption, limb anomalies, and central nervous system abnormalities. Cocaine appear to exert a number of its effects through peripheral vasoconstriction that leads to fetal hypoxia.

Women with indications for warfarin therapy should either abstain from pregnancy or switch to low molecular weight heparins. Warfarin typically produces mental retardation, growth restriction, nasal hypoplasia, and opthalmic abnormalities.

Angiotensin converting enzyme (ACE) inhibitors will cause fetal renal failure and oligohydramnios that lead to pulmonary hypoplasia and limb contracture. Fetal cranial bone abnormalities are also common.

Isotretinoin (Accutane), used to treat acne, may cause cardiac, oral, otological, thymic, and central nervous system abnormalities. In one quarter of cases, it causes mental retardation.



Thalidomide effects: A photograph of the limbs of baby born to a mother who took thalidomide while pregnant.

Other teratogenic substance classes and conditions include

- Various prescription drugs and nutrient deficiencies (e.g., insufficient folic acid).
- Chemical compounds such as methyl iodide (used in pesticides) and bisphenol A (used in plastics) are suspected teratogens.
- Thalidomide (a sedative previously marketed in Europe to prevent morning sickness) is a classic teratogen that caused limb defects in babies born to women who took this drug in the 1960s.

Key Points

- The effects of a teratogen on the fetus depend on several factors: the potency of the teratogen, the susceptibility of the fetus to the teratogen, the dose and duration of teratogen exposure, the degree of transfer from maternal to fetal circulation, and when during development the exposure occurs.
- Approximately 10% of congenital malformations are attributed to environmental factors, and 20% are due to genetic or hereditary factors. The rest have unknown causes or are due to a mix of different factors.
- The central nervous and skeletal systems tend to be most affected by teratogens.
- Cigarette components, alcohol, cocaine, warfarin, ACE inhibitors, and Accutane are all teratogens that affect fetal development.

Key Terms

- **teratogen**: Any agent or substance that can cause embryonic malformations or birth defects.
- **fetal alcohol syndrome**: Any of a spectrum of birth defects that result from excessive alcohol consumption by the mother during pregnancy.
- **microcephaly**: A neurological disorder in which the person affected has an abnormally small head due to a failure of brain

growth.

This page titled 27.9B: Teratogens is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

27.9C: Prenatal Diagnostic Tests

Prenatal diagnosis is a way to screen a fetus for diseases and/or conditions that may increase its morbidity and/or mortality.

Learning Objectives

- Infer the type of prenatal diagnostic test that should be used in a particular case

Key Points

- Prenatal diagnosis allows: 1) timely medical treatment of a condition before or after birth; 2) parents to make decisions regarding whether to abort a fetus with a diagnosed condition; 3) parents to prepare psychologically, socially, financially, and medically for a baby with a health problem or disability.
- The types of conditions commonly screened for include neural tube defects, chromosome abnormalities (e.g., Down syndrome, also called trisomy 21), and genetic abnormalities (e.g., Tay-Sachs disease, sickle cell anemia, and cystic fibrosis).
- Diagnostic screening can be invasive (for example, amniocentesis) or noninvasive (such as ultrasound). Risks are associated with invasive measures. It is often recommended that women who are at high risk of pregnancy loss due to advanced age or who previously had children with birth defects receive invasive testing.
- Early ultrasound scans are done at around six weeks; they are used to confirm the gestational age of the embryo and check for a single or twin pregnancy.
- Around weeks 10–11, a nuchal thickness scan (NT) may be offered that can be combined with blood tests that correlate with chromosomal abnormalities. The results of the blood tests are combined with the NT ultrasound measurements, maternal age, and gestational age of the fetus to yield a risk score.
- There are ethical issues associated with prenatal tests, including whether parents should elect to abort fetuses with detected abnormalities, the high rate of false positives and false negatives, and whether prenatal screening will lead to the rise of designer babies.

Key Terms

- **amniocentesis**: A procedure for obtaining amniotic fluid from a pregnant female by inserting a hollow needle through the abdominal wall and into the amniotic sac. It is used to diagnose possible genetic defects and/or obstetric complications.
- **chorionic villus sampling**: A form of prenatal diagnosis in which part of the placental tissue is tested for genetic abnormalities.
- **nuchal thickness scan**: A sonographic prenatal screening scan (ultrasound) to help identify any high risks of chromosomal defects, including Down's Syndrome in a fetus, particularly for older women who have higher risks of such pregnancies.


Prenatal Screening Overview

Prenatal diagnosis and prenatal screening are methods for testing for diseases or conditions in a fetus or embryo before it is born. There are three purposes of prenatal diagnosis:

1. To enable timely medical or surgical treatment of a condition before (fetal therapy) or after birth.
2. To give the parents the chance to decide to abort a fetus with the diagnosed condition.
3. To give parents the chance to prepare psychologically, socially, financially, and medically for a baby with a health problem or disability, or for the likelihood of a stillbirth.

Having this information in advance of the birth means that healthcare staff, as well as parents, can better prepare themselves for the delivery of a child with a health problem.

For example, Down Syndrome is associated with cardiac defects that may need intervention immediately upon birth. The other types of conditions commonly screened for include neural tube defects, chromosome abnormalities, genetic diseases, spina bifida, cleft palate, Tay-Sachs disease, sickle cell anemia, thalassemia, cystic fibrosis, and fragile X syndrome.



Prenatal diagnosis: Karyotype: This karyotype indicates that the fetus has Down Syndrome as it has three of chromosome 21 instead of two.

Fetal screening is also done to determine characteristics that are not generally considered birth defects, such as determining the sex of the offspring. The primary method for sex determination is prenatal ultrasound. Cell-free fetal DNA testing is also possible, wherein the maternal blood is analyzed to detect the small amount of free circulating fetal DNA.


Invasive and Noninvasive Prenatal Screening

Diagnostic prenatal testing can be done by invasive or noninvasive methods. An invasive method involves probes or needles being inserted into the uterus, e.g., amniocentesis (that can be done from about 14 weeks gestation up to about 20 weeks), and chorionic villus sampling (that can be done earlier: between 9.5 and 12.5 weeks gestation).

Chorionic villus sampling is associated with slightly more risk to the fetus. Since chorionic villus sampling is performed earlier in the pregnancy than amniocentesis, typically during the first trimester, it can reasonably be expected that there will be a higher rate of miscarriage after chorionic villus sampling than after amniocentesis.

Non-invasive techniques include examinations of the woman's womb through ultrasonography and maternal serum screens. Blood tests for select trisomies based on detecting fetal DNA present in maternal blood are available. Technologies to perform prenatal testing in a noninvasive manner reduce the risk of pregnancy loss that can occur with other methods.

These include the use of maternal blood and DNA sequencing methods such as next-generation sequencing to determine the presence of genetic defects. If an elevated risk of chromosomal or genetic abnormality is indicated by a noninvasive screening test, a more invasive technique may be employed to gather more information. In the case of neural tube defects, a detailed ultrasound can non-invasively provide a definitive diagnosis.



Amniocentesis: An illustration of the amniocentesis procedure.

Because of the risk of miscarriage and fetal damage associated with amniocentesis and chorionic villus sampling procedures, many women prefer to first undergo screening so they can find out if the fetus' risk of birth defects is high enough to justify the risks of invasive testing. Screening tests yield a risk score that represents the chance that a baby has the birth defect.

The most common threshold for high risk is 1:270. A risk score of 1:300 would be considered low-risk by many physicians. However, the trade-off between the risk of a birth defect and the risk of complications from invasive testing is relative and subjective. Some parents may decide that even a 1:1,000 risk of birth defects warrants an invasive test, while others would not opt for an invasive test even if they had a 1:10 risk score.

The American Congress of Obstetricians and Gynecologists guidelines currently recommend that all pregnant women, regardless of age, be offered invasive testing to obtain a definitive diagnosis of certain birth defects. Therefore, most physicians offer diagnostic testing to all their patients, with or without prior screening, and let the patient decide. Invasive testing is warranted in the following cases:

- Women over the age of 35.
- Women who have previously had premature babies or babies with a birth defect.
- Women who have high blood pressure, lupus, diabetes, asthma, or epilepsy.
- Women who have family histories or ethnic backgrounds prone to genetic disorders, or whose partners have these.
- Women who are pregnant with twins or more.
- Women who have miscarried.

Determination of Gestational Age

At about 6 weeks of pregnancy, an ultrasound dating scan may be offered to help confirm the gestational age of the embryo and check if a single or twin pregnancy exists, but such a scan is unable to detect common abnormalities. Around weeks 10–11, a nuchal thickness scan (NT) may be offered, which can be combined with blood tests that correlate with chromosomal abnormalities.

The results of the blood tests are then combined with the NT ultrasound measurements, maternal age, and gestational age of the fetus to yield a risk score for Down syndrome, trisomy 18, and trisomy 13. Other testing may combine a first trimester blood test with one done in the second trimester to determine if further action needs to be taken.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Fetal development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fetal_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Prenatal development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Prenata...23Fetal_period**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fetus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fetus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lanugo. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lanugo**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gestational age. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/gestational%20age**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Meconium. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Meconium**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Obstetrics and Gynecology/Teratology and Associated Outcomes. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Obstetr..._to_Teratology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- teratogen. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/teratogen**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- microcephaly. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/microcephaly**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fetal alcohol syndrome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fetal_alcohol_syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Teratology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Teratology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fetal alcohol syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fetal_alcohol_syndrome**. **License**: ***Public Domain: No Known Copyright***
- thalidomide.jpg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:NCP14053.jpg**. **License**: ***CC BY: Attribution***
- Genetic counseling. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Genetic_counseling**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nuchal thickness scan. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/nuchal%...ickness%20scan**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- amniocentesis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amniocentesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chorionic villus sampling. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chorio...illus_sampling**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Prenatal Sex Discernment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Prenat...ex_discernment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Fetal alcohol syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Fetal_alcohol_syndrome**. **License**: ***Public Domain: No Known Copyright***
- thalidomide.jpg. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:NCP14053.jpg**. **License**: ***CC BY: Attribution***
- Amniocentesis.png. **Provided by**: Wikipedia Commons. **Located at**: **commons.wikimedia.org/wiki/F...iocentesis.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Down Syndrome Karyotype. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Do..._Karyotype.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 27.9C: Prenatal Diagnostic Tests is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28: REPRODUCTION, CHROMOSOMES, AND MEIOSIS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.10: Exceptions to Mendelian Inheritance

28.10A: Mendel's Laws of Heredity

28.1: The Link Between Genotype and Phenotype

28.1A: The Process and Purpose of Gene Expression Regulation

28.1B: The Relationship Between Genes and Proteins

28.1C: Chromosomal Theory of Inheritance

28.2: Inherited Variation

28.2A: Advantages and Disadvantages of Sexual Reproduction

28.3: Chromosomal Behavior in Meiosis and Fertilization

28.3A: The Structure and Sequence of DNA

28.4: DNA

28.4A: DNA and RNA

28.4B: The DNA Double Helix

28.4C: DNA Packaging

28.4D: Types of RNA

28.5: Cell Division

28.5A: The Role of the Cell Cycle

28.5B: Genomic DNA and Chromosomes

28.5C: Eukaryotic Chromosomal Structure and Compaction

28.6: Sources of Genetic Variation

28.6A: Genetic Variation

28.6B: Mendel's Law of Segregation

28.6C: Mendel's Law of Independent Assortment

28.7: Complex Inheritance Patterns

28.3C: Mendel's Model System

28.7A: The Punnett Square Approach for a Monohybrid Cross

28.7B: Phenotypes and Genotypes

28.8: Patterns of Inheritance

28.8A: Lethal Inheritance Patterns
28.8B: Sex-Linked Traits

28.9: Genetic Disequilibrium

28.9A: Genetic Linkage and Distances
28.9B: Chromosomal Structural Rearrangements

This page titled 28: Reproduction, Chromosomes, and Meiosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.10: EXCEPTIONS TO MENDELIAN INHERITANCE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.10A: Mendel's Laws of Heredity

This page titled 28.10: Exceptions to Mendelian Inheritance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.10A: Mendel's Laws of Heredity

Mendel formed the Laws of Heredity (the Law of Segregation and the Law of Independent Assortment) from his pea plant experiments.

Learning Objectives

- Discuss the methods Mendel utilized in his research that led to his success in understanding the process of inheritance

Key Points

- By crossing purple and white pea plants, Mendel found the offspring were purple rather than mixed, indicating one color was dominant over the other.
- Mendel's Law of Segregation states individuals possess two alleles and a parent passes only one allele to his/her offspring.
- Mendel's Law of Independent Assortment states the inheritance of one pair of factors (genes) is independent of the inheritance of the other pair.
- If the two alleles are identical, the individual is called homozygous for the trait; if the two alleles are different, the individual is called heterozygous.
- Mendel cross-bred dihybrids and found that traits were inherited independently of each other.

Key Terms

- **homozygous**: of an organism in which both copies of a given gene have the same allele
- **heterozygous**: of an organism which has two different alleles of a given gene
- **allele**: one of a number of alternative forms of the same gene occupying a given position on a chromosome


Introduction

Mendelian inheritance (or Mendelian genetics or Mendelism) is a set of primary tenets relating to the transmission of hereditary characteristics from parent organisms to their children; it underlies much of genetics. The tenets were initially derived from the work of Gregor Mendel published in 1865 and 1866, which was "re-discovered" in 1900; they were initially very controversial, but they soon became the core of classical genetics.

The laws of inheritance were derived by Gregor Mendel, a 19th century monk conducting hybridization experiments in garden peas (*Pisum sativum*). Between 1856 and 1863, he cultivated and tested some 28,000 pea plants. From these experiments, he deduced two generalizations that later became known as Mendel's Laws of Heredity or Mendelian inheritance. He described these laws in a two part paper,"Experiments on Plant Hybridization," which was published in 1866.

Mendel's Laws

Mendel discovered that by crossing true-breeding white flower and true-breeding purple flower plants, the result was a hybrid offspring. Rather than being a mix of the two colors, the offspring was purple flowered. He then conceived the idea of heredity units, which he called "factors," one of which is a recessive characteristic and the other dominant. Mendel said that factors, later called genes, normally occur in pairs in ordinary body cells, yet segregate during the formation of sex cells. Each member of the pair becomes part of the separate sex cell. The dominant gene, such as the purple flower in Mendel's plants, will hide the recessive gene, the white flower. After Mendel self-fertilized the F1 generation and obtained an F2 generation with a 3:1 ratio, he correctly theorized that genes can be paired in three different ways for each trait: AA, aa, and Aa. The capital A represents the dominant factor while the lowercase a represents the recessive.



Mendel's Pea Plants: In one of his experiments on inheritance patterns, Mendel crossed plants that were true-breeding for violet flower color with plants true-breeding for white flower color (the P generation). The resulting hybrids in the F1 generation all had violet flowers. In the F2 generation, approximately three-quarters of the plants had violet flowers, and one-quarter had white flowers.

Mendel stated that each individual has two alleles for each trait, one from each parent. Thus, he formed the "first rule," the Law of Segregation, which states individuals possess two alleles and a parent passes only one allele to his/her offspring. One allele is given by the female parent and the other is given by the male parent. The two factors may or may not contain the same information. If the two alleles are identical, the individual is called homozygous for the trait. If the two alleles are different, the individual is called heterozygous. The presence of an allele does not promise that the trait will be expressed in the individual that possesses it. In heterozygous individuals, the only allele that is expressed is the dominant. The recessive allele is present, but its expression is hidden. The genotype of an individual is made up of the many alleles it possesses. An individual's physical appearance, or phenotype, is determined by its alleles as well as by its environment.

Mendel also analyzed the pattern of inheritance of seven pairs of contrasting traits in the domestic pea plant. He did this by cross-breeding dihybrids; that is, plants that were heterozygous for the alleles controlling two different traits. Mendel then crossed these dihybrids. If it is inevitable that round seeds must always be yellow and wrinkled seeds must be green, then he would have expected that this would produce a typical monohybrid cross: 75 percent round-yellow; 25 percent wrinkled-green. But, in fact, his mating generated seeds that showed all possible combinations of the color and texture traits. He found 9/16 of the offspring were round-yellow, 3/16 were round-green, 3/16 were wrinkled-yellow, and 1/16 were wrinkled-green. Finding in every case that each of his seven traits was inherited independently of the others, he formed his "second rule," the Law of Independent Assortment, which states the inheritance of one pair of factors (genes) is independent of the inheritance of the other pair. Today we know that this rule holds only if the genes are on separate chromosomes

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Le Dinh Luong, Genetics in Clasical Understanding. October 30, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m26573/latest/**. **License**: ***CC BY: Attribution***
- heterozygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/heterozygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- homozygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homozygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- allele. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/allele**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Mendelu2019s Experiments and the Laws of Probability. October 30, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44476/latest/**. **License**: ***CC BY: Attribution***

This page titled 28.10A: Mendel's Laws of Heredity is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.1: THE LINK BETWEEN GENOTYPE AND PHENOTYPE

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

28.1A: The Process and Purpose of Gene Expression Regulation

28.1B: The Relationship Between Genes and Proteins

28.1C: Chromosomal Theory of Inheritance

This page titled 28.1: The Link Between Genotype and Phenotype is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.1A: The Process and Purpose of Gene Expression Regulation

Gene expression is a highly complex, regulated process that begins with DNA transcribed into RNA, which is then translated into protein.

Learning Objectives

- Discuss how the genome and proteome contribute to the specialization of a cell


Key Points

- Every cell within an organism shares the same genome (with exceptions, i.e. mature red blood cells), but has variation between its proteomes.
- Gene expression involves the process of transcribing DNA into RNA and then translating RNA into proteins.
- Gene expression is a highly complex and tightly-regulated process.

Key Terms

- **somatic**: part of, or relating to the body of an organism
- **genome**: the cell's complete genetic information packaged as a double-stranded DNA molecule
- **proteome**: the complete set of proteins encoded by a particular genome

Each somatic cell in the body generally contains the same DNA. A few exceptions include red blood cells, which contain no DNA in their mature state, and some immune system cells that rearrange their DNA while producing antibodies. In general, however, the genes that determine whether you have green eyes, brown hair, and how fast you metabolize food are the same in the cells in your eyes and your liver, even though these organs function quite differently. If each cell has the same DNA, how is it that cells or organs are different? Why do cells in the eye differ so dramatically from cells in the liver ?



Gene Expression: The genetic content of each somatic cell in an organism is the same, but not all genes are expressed in every cell. The control of which genes are expressed dictates whether a cell is (a) an eye cell or (b) a liver cell. It is the differential gene expression patterns that arise in different cells that give rise to (c) a complete organism.

Whereas each cell shares the same genome and DNA sequence, each cell does not turn on, or express, the same set of genes. Each cell type needs a different set of proteins to perform its function. Therefore, only a small subset of proteins is expressed in a cell that constitutes its proteome. For the proteins to be expressed, the DNA must be transcribed into RNA and the RNA must be translated into protein. In a given cell type, not all genes encoded in the DNA are transcribed into RNA or translated into protein because specific cells in our body have specific functions. Specialized proteins that make up the eye (iris, lens, and cornea) are only expressed in the eye, whereas the specialized proteins in the heart (pacemaker cells, heart muscle, and valves) are only expressed in the heart. At any given time, only a subset of all of the genes encoded by our DNA are expressed and translated into proteins. The expression of specific genes is a highly-regulated process with many levels and stages of control. This complexity ensures the proper expression in the proper cell at the proper time.

In this section, you will learn about the various methods of gene regulation and the mechanisms used to control gene expression, such as: epigenetic, transcriptional, post-transcriptional, translational, and post-translational controls in eukaryotic gene expression, and transcriptional control in prokaryotic gene expression.

This page titled 28.1A: The Process and Purpose of Gene Expression Regulation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.1B: The Relationship Between Genes and Proteins

Proteins, encoded by individual genes, orchestrate nearly every function of the cell.

Learning Objectives

- Describe transcription and translation

Key Points

- Genes are composed of DNA arranged on chromosomes.
- Some genes encode structural or regulatory RNAs. Other genes encode proteins.
- Replication copies DNA; transcription uses DNA to make complementary RNAs; translation uses mRNAs to make proteins.
- In eukaryotic cells, replication and transcription take place within the nucleus while translation takes place in the cytoplasm.
- In prokaryotic cells, replication, transcription, and translation occur in the cytoplasm.


Key Terms

- **DNA**: a biopolymer of deoxyribonucleic acids (a type of nucleic acid) that has four different chemical groups, called bases: adenine, guanine, cytosine, and thymine
- **messenger RNA**: Messenger RNA (mRNA) is a molecule of RNA that encodes a chemical “blueprint” for a protein product.
- **protein**: any of numerous large, complex naturally-produced molecules composed of one or more long chains of amino acids, in which the amino acid groups are held together by peptide bonds

Genes and Proteins

Since the rediscovery of Mendel’s work in 1900, the definition of the gene has progressed from an abstract unit of heredity to a tangible molecular entity capable of replication, transcription, translation, and mutation. Genes are composed of DNA and are linearly arranged on chromosomes. Some genes encode structural and regulatory RNAs. There is increasing evidence from research that profiles the transcriptome of cells (the complete set all RNA transcripts present in a cell) that these may be the largest classes of RNAs produced by eukaryotic cells, far outnumbering the protein-encoding messenger RNAs (mRNAs), but the 20,000 protein-encoding genes typically found in animal cells, and the 30,000 protein-encoding genes typically found in plant cells, nonetheless have huge impacts on cellular functioning.

Protein-encoding genes specify the sequences of amino acids, which are the building blocks of proteins. In turn, proteins are responsible for orchestrating nearly every function of the cell. Both protein-encoding genes and the proteins that are their gene products are absolutely essential to life as we know it.



Genes Encode Proteins: Genes, which are carried on (a) chromosomes, are linearly-organized instructions for making the RNA and protein molecules that are necessary for all of processes of life. The (b) interleukin-2 protein and (c) alpha-2u-globulin protein are just two examples of the array of different molecular structures that are encoded by genes.

Replication, Transcription, and Translation are the three main processes used by all cells to maintain their genetic information and to convert the genetic information encoded in DNA into gene products, which are either RNAs or proteins, depending on the gene. In eukaryotic cells, or those cells that have a nucleus, replication and transcription take place within the nucleus while translation takes place outside of the nucleus in cytoplasm. In prokaryotic cells, or those cells that do not have a nucleus, all three processes occur in the cytoplasm.

Replication is the basis for biological inheritance. It copies a cell's DNA. The enzyme DNA polymerase copies a single parental double-stranded DNA molecule into two daughter double-stranded DNA molecules. Transcription makes RNA from DNA. The enzyme RNA polymerase creates an RNA molecule that is complementary to a gene-encoding stretch of DNA. Translation makes protein from mRNA. The ribosome generates a polypeptide chain of amino acids using mRNA as a template. The polypeptide chain folds up to become a protein.

This page titled 28.1B: The Relationship Between Genes and Proteins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.1C: Chromosomal Theory of Inheritance

The Chromosomal Theory of Inheritance identified chromosomes as the genetic material responsible for Mendelian inheritance.

Learning Objectives

- List the reasons that fruit flies are excellent model organisms for genetic research

Key Points


- Homologous chromosome pairs are independent of other chromosome pairs.
- Chromosomes from each homologous pair are sorted randomly into pre- gametes.
- Parents synthesize gametes that contain only half of their chromosomes; eggs and sperm have the same number of chromosomes.
- Gametic chromosomes combine during fertilization to produce offspring with the same chromosome number as their parents.
- Eye color in fruit flies was the first X-linked trait to be discovered; thus, Morgan's experiments with fruit flies solidified the Chromosomal Theory of Inheritance.

Key Terms

- **autosome**: any chromosome other than sex chromosomes
- **hemizygous**: having some single copies of genes in an otherwise diploid cell or organism
- **wild type**: the typical form of an organism, strain, gene or characteristic as it occurs in nature

Chromosomal Theory of Inheritance

The speculation that chromosomes might be the key to understanding heredity led several scientists to examine Mendel's publications and re-evaluate his model in terms of the behavior of chromosomes during mitosis and meiosis. In 1902, Theodor Boveri observed that proper embryonic development of sea urchins does not occur unless chromosomes are present. That same year, Walter Sutton observed the separation of chromosomes into daughter cells during meiosis. Together, these observations led to the development of the Chromosomal Theory of Inheritance, which identified chromosomes as the genetic material responsible for Mendelian inheritance.



(a) (b)

Sutton and Boveri: (a) Walter Sutton and (b) Theodor Boveri are credited with developing the Chromosomal Theory of Inheritance, which states that chromosomes carry the unit of heredity (genes).

The Chromosomal Theory of Inheritance was consistent with Mendel's laws and was supported by the following observations:

- During meiosis, homologous chromosome pairs migrate as discrete structures that are independent of other chromosome pairs.
- The sorting of chromosomes from each homologous pair into pre-gametes appears to be random.
- Each parent synthesizes gametes that contain only half of their chromosomal complement.
- Even though male and female gametes (sperm and egg) differ in size and morphology, they have the same number of chromosomes, suggesting equal genetic contributions from each parent.
- The gametic chromosomes combine during fertilization to produce offspring with the same chromosome number as their parents.

Despite compelling correlations between the behavior of chromosomes during meiosis and Mendel's abstract laws, the Chromosomal Theory of Inheritance was proposed long before there was any direct evidence that traits were carried on chromosomes. Critics pointed out that individuals had far more independently segregating traits than they had chromosomes. It was only after several years of carrying out crosses with the fruit fly, *Drosophila melanogaster*, that Thomas Hunt Morgan provided experimental evidence to support the Chromosomal Theory of Inheritance.

In 1910, Thomas Hunt Morgan started his work with *Drosophila melanogaster*, a fruit fly. He chose fruit flies because they can be cultured easily, are present in large numbers, have a short generation time, and have only four pair of chromosomes that can be easily identified under the microscope. They have three pair of autosomes and a pair of sex chromosomes. At that time, he already knew that X and Y have to do with gender. He used normal flies with red eyes and mutated flies with white eyes and cross bred them. In flies, the wild type eye color is red (XW) and is dominant to white eye color (Xw). He was able to conclude that the gene for eye color was on the X chromosome. This trait was thus determined to be X-linked and was the first X-linked trait to be identified. Males are said to be hemizygous, in that they have only one allele for any X-linked characteristic.



Eye Color in Fruit Flies: In Drosophila, the gene for eye color is located on the X chromosome. Red eye color is wild type and is dominant to white eye color.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44533/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- somatic. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/somatic. **License**: ***CC BY-SA: Attribution-ShareAlike***
- proteome. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/proteome. **License**: ***CC BY-SA: Attribution-ShareAlike***
- genome. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/genome. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44533/latest...e_16_00_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44518/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Principles of Biochemistry/Cell Metabolism I: DNA replication. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Princip...NA_replication. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cell Biology/Genes/Gene translation. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Cell_Bi...ne_translation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- translation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/translation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gene. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/gene%23English. **License**: ***CC BY-SA: Attribution-ShareAlike***
- transcription. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/transcription. **License**: ***CC BY-SA: Attribution-ShareAlike***
- protein. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/protein. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, How Genes Are Regulated. October 30, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m45480/latest/. **License**: ***CC BY: Attribution***
- Structural Biochemistry/Nucleic Acid/Translation. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Structu...id/

Translation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Principles of Biochemistry/Cell Metabolism II: RNA transcription. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Princip..._transcription**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- protein. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/protein**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- messenger RNA. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/messenger%20RNA**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- DNA. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/DNA**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44533/latest...e_16_00_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44518/latest...e_15_00_01.jpg**. **License**: ***CC BY: Attribution***
- hemizygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemizygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44481/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Robert Bear and David Rintoul, Extensions of the Laws of Inheritance. October 31, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m47304/latest/**. **License**: ***CC BY: Attribution***
- Structural Biochemistry/Chromosomes. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...ry/Chromosomes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autosome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/autosome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- wild type. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/wild_type**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44533/latest...e_16_00_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44518/latest...e_15_00_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Chromosomal Theory and Genetic Linkage October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44481/latest...e_13_01_01.jpg**. **License**: ***CC BY: Attribution***
- Robert Bear and David Rintoul, Extensions of the Laws of Inheritance. October 31, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m47304/latest/**. **License**: ***CC BY: Attribution***

This page titled 28.1C: Chromosomal Theory of Inheritance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.2: INHERITED VARIATION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.2A: Advantages and Disadvantages of Sexual Reproduction

This page titled 28.2: Inherited Variation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.2A: Advantages and Disadvantages of Sexual Reproduction

The genetic diversity of sexual reproduction, observed in most eukaryotes, is thought to give species better chances of survival.

Learning Objectives

- Describe the benefits of sexual reproduction

Key Points

- The variation that sexual reproduction creates among offspring is very important to the survival and reproduction of the population.
- In sexual reproduction, different mutations are continually reshuffled from one generation to the next when different parents combine their unique genomes; this results in an increase of genetic diversity.
- On average, a sexually-reproducing population will leave more offspring than an otherwise similar asexually-reproducing population.

Key Terms

- **sexual reproduction**: Sexual reproduction is the creation of a new organism by combining the genetic material of two organisms. There are two main processes during sexual reproduction: meiosis, involving the halving of the number of chromosomes, and fertilization, involving the fusion of two gametes and the restoration of the original number of chromosomes.
- **asexual reproduction**: any form of reproduction that involves neither meiosis nor fusion of gametes

An Introduction to Sexual Reproduction


Sexual reproduction was an early evolutionary innovation after the appearance of eukaryotic cells. During sexual reproduction, the genetic material of two individuals is combined to produce genetically-diverse offspring that differ from their parents. The fact that most eukaryotes reproduce sexually is evidence of its evolutionary success. In many animals, it is actually the only mode of reproduction. The genetic diversity of sexually-produced offspring is thought to give species a better chance of surviving in an unpredictable or changing environment.

Scientists recognize some real disadvantages to sexual reproduction. On the surface, creating offspring that are genetic clones of the parent appears to be a better system. If the parent organism is successfully occupying a habitat, offspring with the same traits would be similarly successful. Species that reproduce sexually must maintain two different types of individuals, males and females, which can limit the ability to colonize new habitats as both sexes must be present. Therefore, there is an obvious benefit to an organism that can produce offspring whenever circumstances are favorable by asexual budding, fragmentation, or asexual eggs. These methods of asexual reproduction do not require another organism of the opposite sex. Indeed, some organisms that lead a solitary lifestyle have retained the ability to reproduce asexually. In addition, in asexual populations, every individual is capable of reproduction. In sexual populations, the males are not producing the offspring themselves. In theory, an asexual population could grow twice as fast.

Nevertheless, multicellular organisms that exclusively depend on asexual reproduction are exceedingly rare. Why is sexuality (and meiosis) so common? This is one of the important unanswered questions in biology and has been the focus of much research beginning in the latter half of the twentieth century. There are several possible explanations, one of which is that the variation that sexual reproduction creates among offspring is very important to the survival and reproduction of the population. Thus, on average, a sexually-reproducing population will leave more descendants than an otherwise similar asexually-reproducing population. The only source of variation in asexual organisms is mutation. This is the ultimate source of variation in sexual organisms, but, in addition, those different mutations are continually reshuffled from one generation to the next when different

parents combine their unique genomes and the genes are mixed into different combinations by the process of meiosis. Meiosis is the division of the contents of the nucleus, dividing the chromosomes among gametes.

The process of meiosis produces unique reproductive cells called gametes, which have half the number of chromosomes as the parent cell. Fertilization, the fusion of haploid gametes from two individuals, restores the diploid condition. Thus, sexually-reproducing organisms alternate between haploid and diploid stages. However, the ways in which reproductive cells are produced and the timing between meiosis and fertilization vary greatly. There are three main categories of sexual life cycles: diploid-dominant, demonstrated by most animals; haploid-dominant, demonstrated by all fungi and some algae; and the alternation of generations, demonstrated by plants and some algae.



The Sexual Life Cycle: In animals, sexually-reproducing adults form haploid gametes from diploid germ cells. Fusion of the gametes gives rise to a fertilized egg cell, or zygote. The zygote will undergo multiple rounds of mitosis to produce a multicellular offspring.

The Red Queen Hypothesis

It is not in dispute that sexual reproduction provides evolutionary advantages to organisms that employ this mechanism to produce offspring. But why, even in the face of fairly stable conditions, does sexual reproduction persist when it is more difficult and costly for individual organisms? Variation is the outcome of sexual reproduction, but why are ongoing variations necessary? Possible answers to these questions are explained in the Red Queen hypothesis, first proposed by Leigh Van Valen in 1973.

All species co-evolve with other organisms; for example, predators evolve with their prey and parasites evolve with their hosts. Each tiny advantage gained by favorable variation gives a species an edge over close competitors, predators, parasites, or even prey. The only method that will allow a co-evolving species to maintain its own share of the resources is to also continually improve its fitness. As one species gains an advantage, this increases selection on the other species; they must also develop an advantage or they will be out-competed. No single species progresses too far ahead because genetic variation among the progeny of sexual reproduction provides all species with a mechanism to improve rapidly. Species that cannot keep up become extinct. The Red Queen's catchphrase was,"It takes all the running you can do to stay in the same place." This is an apt description of co-evolution between competing species.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- asexual reproduction. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/asexual_reproduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44470/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Reproduction Methods. October 26, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44836/latest/. **License**: *CC BY: Attribution*
- OpenStax College, Sexual Reproduction. October 26, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m45465/latest/. **License**: *CC BY: Attribution*
- sexual reproduction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/sexual%20reproduction. **License**: *CC BY-SA: Attribution-ShareAlike*
- Biological life cycle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Biological_life_cycle. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 28.2A: Advantages and Disadvantages of Sexual Reproduction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.3: CHROMOSOMAL BEHAVIOR IN MEIOSIS AND FERTILIZATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

28.3A: The Structure and Sequence of DNA

This page titled 28.3: Chromosomal Behavior in Meiosis and Fertilization is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.3A: The Structure and Sequence of DNA

DNA is a double helix of two anti-parallel, complementary strands having a phosphate-sugar backbone with nitrogenous bases stacked inside.

Learning Objectives

- Describe the structure of DNA and summarize the importance of DNA sequencing


Key Points

- The two DNA strands are anti-parallel in nature; that is, the 3′ end of one strand faces the 5′ end of the other strand.
- The nucleotides that comprise DNA contain a nitrogenous base, a deoxyribose sugar, and a phosphate group which covalently link with other nucleotides to form phosphodiester bonds.
- Nucleotide bases can be classified as purines (containing a double-ring structure) or pyrimidines (containing a single-ring structure).
- Adenine (purine) and thymine (pyrimidine) are complementary base pairs as are guanine (purine) and cytosine (pyrimidine).
- DNA sequencing is the process of determining the precise order of nucleotides within a DNA molecule.

Key Terms

- **deoxyribose**: a derivative of the pentose sugar ribose in which the 2′ hydroxyl (-OH) is reduced to a hydrogen (H); a constituent of the nucleotides that comprise deoxyribonucleic acid, or DNA
- **hydrogen bond**: A weak bond in which a hydrogen atom already covalently bonded to a oxygen or nitrogen atom in one molecule is attracted to an electronegative atom (usually nitrogen or oxygen) in the same or different molecule.
- **nucleotide**: the monomer comprising DNA or RNA molecules; consists of a nitrogenous heterocyclic base that can be a purine or pyrimidine, a five-carbon pentose sugar, and a phosphate group

The monomeric building blocks of DNA are deoxyribomononucleotides (usually referred to as just nucleotides), and DNA is formed from linear chains, or polymers, of these nucleotides. The components of the nucleotide used in DNA synthesis are a nitrogenous base, a deoxyribose, and a phosphate group. The nucleotide is named depending on which nitrogenous base is present. The nitrogenous base can be a purine such as adenine (A) and guanine (G), characterized by double-ring structures, or a pyrimidine such as cytosine (C) and thymine (T), characterized by single-ring structures. In polynucleotides (the linear polymers of nucleotides) the nucleotides are connected to each other by covalent bonds known as phosphodiester bonds or phosphodiester linkages.



Nucleotide Structure: Each nucleotide is made up of a sugar, a phosphate group, and a nitrogenous base. The sugar is deoxyribose in DNA and ribose in RNA. In their mononucleotide form, nucleotides can have one, two, or three phosphates attached to them. When linked together in polynucleotide chains, the nucleotides always have just one phosphate. A molecule with just a nitrogenous base and a sugar is known as a nucleoside. Once at least one phosphate is covalently attached, it is known as a nucleotide.

James Watson and Francis Crick, with some help from Rosalind Franklin and Maurice Wilkins, are credited with figuring out the structure of DNA. Watson and Crick proposed that DNA is made up of two polynucleotide strands that are twisted around each other to form a right-handed helix.

The two polynucleotide strands are anti-parallel in nature. That is, they run in opposite directions.

The sugars and phosphates of the nucleotides form the backbone of the structure, whereas the pairs of nitrogenous bases are pointed towards the interior of the molecule.

The twisting of the two strands around each other results in the formation of uniformly-spaced major and minor grooves bordered by the sugar-phosphate backbones of the two strands.

image

Three representations of DNA's double helical structure.: A is a spacefill model of DNA, where every atom is represented as a sphere. The two anti-parallel polynucleotide strands are colored differently to illustrate how they coil around each other. B is a cartoon model of DNA, where the sugar-phosphate backbones are represented as violet strands and the nitrogenous bases are represented as color-coded rings. C is another spacefill model, with the sugar-phosphate atoms colored violet and all nitrogenous base atoms colored green. The major and minor grooves, which wrap around the entire molecule, are apparent as the spaces between the sugar-phosphate backbones.


The diameter of the DNA double helix is 2 nm and is uniform throughout. Only the pairing between a purine and pyrimidine can explain the uniform diameter. That is to say, at each point along the DNA molecule, the two sugar phosphate backbones are always separated by three rings, two from a purine and one from a pyrimidine.

The two strands are held together by base pairing between nitrogenous bases of one strand and nitrogenous bases from the other strand. Base pairing takes place between a purine and pyrimidine stabilized by hydrogen bonds: A pairs with T via two hydrogen bonds and G pairs with C via three hydrogen bonds.

The interior basepairs rotate with respect to one another, but are also stacked on top of each other when the molecule is viewed looking up or down its long axis.

Each base pair is separated from the previous base pair by a height of 0.34 nm and each 360° turn of the helix travels 3.4 nm

along the long axis of the molecule. Therefore, ten base pairs are present per turn of the helix.



DNA Structure: DNA has (a) a double helix structure and (b) phosphodiester bonds. The (c) major and minor grooves are binding sites for DNA binding proteins during processes such as transcription (the copying of RNA from DNA) and replication.

DNA sequencing is the process of determining the precise order of nucleotides within a DNA molecule. Rapid DNA sequencing methods has greatly accelerated biological and medical research and discovery. Knowledge of DNA sequences has become indispensable for basic biological research, and in numerous applied fields such as diagnostics, biotechnology, forensic biology, and biological systematics. The rapid speed of sequencing attained with modern technology has been instrumental in obtaining complete DNA sequences, or genomes, of numerous types and species of life, including the human genome and those of other animal, plant, and microbial species.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44486/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- DNA sequencing. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/DNA_sequencing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- deoxyribose. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/deoxyribose**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleotide. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nucleotide**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- data-attribution-url=cnx.org/content/m44486/latest...e_14_02_01.jpg. **Provided by**: Connexions. **License**: ***CC BY: Attribution***
- OpenStax College, DNA Structure and Sequencing. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44486/latest...4_02_03abc.jpg**. **License**: ***CC BY: Attribution***

This page titled 28.3A: The Structure and Sequence of DNA is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.4: DNA

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.4A: DNA and RNA

28.4B: The DNA Double Helix

28.4C: DNA Packaging

28.4D: Types of RNA

This page titled 28.4: DNA is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.4A: DNA and RNA

DNA and RNA are nucleic acids that carry out cellular processes, especially the regulation and expression of genes.

Learning Objectives

- Describe the structure of nucleic acids and the types of molecules that contain them

Key Points

- The two main types of nucleic acids are DNA and RNA.
- Both DNA and RNA are made from nucleotides, each containing a five-carbon sugar backbone, a phosphate group, and a nitrogen base.
- DNA provides the code for the cell ‘s activities, while RNA converts that code into proteins to carry out cellular functions.
- The sequence of nitrogen bases (A, T, C, G) in DNA is what forms an organism’s traits.
- The nitrogen bases A and T (or U in RNA) always go together and C and G always go together, forming the 5′-3′ phosphodiester linkage found in the nucleic acid molecules.

Key Terms

- **nucleotide**: the monomer comprising DNA or RNA molecules; consists of a nitrogenous heterocyclic base that can be a purine or pyrimidine, a five-carbon pentose sugar, and a phosphate group
- **genome**: the cell’s complete genetic information packaged as a double-stranded DNA molecule
- **monomer**: A relatively small molecule which can be covalently bonded to other monomers to form a polymer.

Types of Nucleic Acids

The two main types of nucleic acids are deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). DNA is the genetic material found in all living organisms, ranging from single-celled bacteria to multicellular mammals. It is found in the nucleus of eukaryotes and in the chloroplasts and mitochondria. In prokaryotes, the DNA is not enclosed in a membranous envelope, but rather free-floating within the cytoplasm.

The entire genetic content of a cell is known as its genome and the study of genomes is genomics. In eukaryotic cells, but not in prokaryotes, DNA forms a complex with histone proteins to form chromatin, the substance of eukaryotic chromosomes. A chromosome may contain tens of thousands of genes. Many genes contain the information to make protein products; other genes code for RNA products. DNA controls all of the cellular activities by turning the genes “on” or “off.”


The other type of nucleic acid, RNA, is mostly involved in protein synthesis. In eukaryotes, the DNA molecules never leave the nucleus but instead use an intermediary to communicate with the rest of the cell. This intermediary is the messenger RNA (mRNA). Other types of RNA—like rRNA, tRNA, and microRNA—are involved in protein synthesis and its regulation.

Nucleotides

DNA and RNA are made up of monomers known as nucleotides. The nucleotides combine with each other to form a polynucleotide: DNA or RNA. Each nucleotide is made up of three components:

1. a nitrogenous base
2. a pentose (five-carbon) sugar
3. a phosphate group

Each nitrogenous base in a nucleotide is attached to a sugar molecule, which is attached to one or more phosphate groups.



DNA and RNA: A nucleotide is made up of three components: a nitrogenous base, a pentose sugar, and one or more phosphate groups. Carbon residues in the pentose are numbered 1′ through 5′ (the prime distinguishes these residues from those in the base, which are numbered without using a prime notation). The base is attached to the 1′ position of the ribose, and the phosphate is attached to the 5′ position. When a polynucleotide is formed, the 5′ phosphate of the incoming nucleotide attaches to the 3′ hydroxyl group at the end of the growing chain. Two types of pentose are found in nucleotides, deoxyribose (found in DNA) and ribose (found in RNA). Deoxyribose is similar in structure to ribose, but it has an H instead of an OH at the 2′ position. Bases can be divided into two categories: purines and pyrimidines. Purines have a double ring structure, and pyrimidines have a single ring.

Nitrogenous Base

The nitrogenous bases are organic molecules and are so named because they contain carbon and nitrogen. They are bases because they contain an amino group that has the potential of binding an extra hydrogen, and thus, decreasing the hydrogen ion concentration in its environment, making it more basic. Each nucleotide in DNA contains one of four possible nitrogenous bases: adenine (A), guanine (G) cytosine (C), and thymine (T).

Adenine and guanine are classified as purines. The primary structure of a purine consists of two carbon-nitrogen rings. Cytosine, thymine, and uracil are classified as pyrimidines which have a single carbon-nitrogen ring as their primary structure. Each of these basic carbon-nitrogen rings has different functional groups attached to it. In molecular biology shorthand, the nitrogenous bases are simply known by their symbols A, T, G, C, and U. DNA contains A, T, G, and C whereas RNA contains A, U, G, and C.

Five-Carbon Sugar

The pentose sugar in DNA is deoxyribose and in RNA it is ribose. The difference between the sugars is the presence of the hydroxyl group on the second carbon of the ribose and hydrogen on the second carbon of the deoxyribose. The carbon atoms of the sugar molecule are numbered as 1′, 2′, 3′, 4′, and 5′ (1′ is read as “one prime”).

Phosphate Group

The phosphate residue is attached to the hydroxyl group of the 5′ carbon of one sugar and the hydroxyl group of the 3′ carbon of the sugar of the next nucleotide, which forms a 5′3′ phosphodiester linkage. The phosphodiester linkage is not formed by simple dehydration reaction like the other linkages connecting monomers in macromolecules: its formation involves the removal of two phosphate groups. A polynucleotide may have thousands of such phosphodiester linkages.

This page titled 28.4A: DNA and RNA is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.4B: The DNA Double Helix

The DNA double helix looks like a twisted staircase, with the sugar and phosphate backbone surrounding complementary nitrogen bases.

Learning Objectives


- Describe the structure of DNA

Key Points

- The structure of DNA is called a double helix, which looks like a twisted staircase.
- The sugar and phosphate make up the backbone, while the nitrogen bases are found in the center and hold the two strands together.
- The nitrogen bases can only pair in a certain way: A pairing with T and C pairing with G. This is called *base pairing*.
- Due to the *base pairing*, the DNA strands are *complementary* to each other, run in opposite directions, and are called antiparallel strands.

Key Terms

- **mutation**: any error in base pairing during the replication of DNA
- **sugar-phosphate backbone**: The outer support of the ladder, forming strong covalent bonds between monomers of DNA.
- **base pairing**: The specific way in which bases of DNA line up and bond to one another; A always with T and G always with C.



DNA is a Double Helix: Native DNA is an antiparallel double helix. The phosphate backbone (indicated by the curvy lines) is on the outside, and the bases are on the inside. Each base from one strand interacts via hydrogen bonding with a base from the opposing strand.

A Double-Helix Structure


DNA has a double-helix structure, with sugar and phosphate on the outside of the helix, forming the sugar-phosphate backbone of the DNA. The nitrogenous bases are stacked in the interior in pairs, like the steps of a staircase; the pairs are bound to each other by hydrogen bonds. The two strands of the helix run in opposite directions. This *antiparallel* orientation is important to DNA

replication and in many nucleic acid interactions.

Base Pairs

Only certain types of base pairing are allowed. This means Adenine pairs with Thymine, and Guanine pairs with Cytosine. This is known as the base complementary rule because the DNA strands are complementary to each other.

If the sequence of one strand is AATTGGCC, the complementary strand would have the sequence TTAACCGG.



Antiparallel Strands: In a double stranded DNA molecule, the two strands run antiparallel to one another so one is upside down compared to the other. The phosphate backbone is located on the outside, and the bases are in the middle. Adenine forms hydrogen bonds (or base pairs) with thymine, and guanine base pairs with cytosine.

DNA Replication

During DNA replication, each strand is copied, resulting in a daughter DNA double helix containing one parental DNA strand and a newly synthesized strand. At this time it is possible a mutation may occur. A mutation is a change in the sequence of the nitrogen bases. For example, in the sequence AATTGGCC, a mutation may cause the second T to change to a G. Most of the time when this happens the DNA is able to fix itself and return the original base to the sequence. However, sometimes the repair is unsuccessful, resulting in different proteins being created.

This page titled 28.4B: The DNA Double Helix is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.4C: DNA Packaging

DNA packaging is an important process in living cells. Without it, a cell is not able to accommodate the large amount of DNA that is stored inside.

Learning Objectives

- Describe how DNA is packaged differently in prokaryotes and eukaryotes


Key Points

- In eukaryotic cells, DNA and RNA synthesis occur in a different location than protein synthesis; in prokaryotic cells, both these processes occur together.
- DNA is “supercoiled” in prokaryotic cells, meaning that the DNA is either under-wound or over-wound from its normal relaxed state.
- In eukaryotic cells, DNA is wrapped around proteins known as histones to form structures called nucleosomes.

Key Terms

- **nucleosomes**: The fundamental subunit of chromatin, composed of a little less than two turns of DNA wrapped around a set of eight proteins called histones.
- **histones**: The chief protein components of chromatin, which act as spools around which DNA winds.

A eukaryote contains a well-defined nucleus, whereas in prokaryotes the chromosome lies in the cytoplasm in an area called the nucleoid. In eukaryotic cells, DNA and RNA synthesis occur in a separate compartment from protein synthesis. In prokaryotic cells, both processes occur together. What advantages might there be to separating the processes? What advantages might there be to having them occur together?




Eukaryotic and prokaryotic cells: A eukaryote contains a well-defined nucleus, whereas in prokaryotes, the chromosome lies in the cytoplasm in an area called the nucleoid.

The size of the genome in one of the most well-studied prokaryotes, *E.coli,* is 4.6 million base pairs (approximately 1.1 mm, if cut and stretched out). So how does this fit inside a small bacterial cell? The DNA is twisted by what is known as supercoiling.

Supercoiling means that DNA is either under-wound (less than one turn of the helix per 10 base pairs) or over-wound (more than 1 turn per 10 base pairs) from its normal relaxed state. Some proteins are known to be involved in the supercoiling; other proteins and enzymes such as DNA gyrase help in maintaining the supercoiled structure.

Eukaryotes, whose chromosomes each consist of a linear DNA molecule, employ a different type of packing strategy to fit their DNA inside the nucleus. At the most basic level, DNA is wrapped around proteins known as histones to form structures called nucleosomes. The histones are evolutionarily conserved proteins that are rich in basic amino acids and form an octamer. The DNA (which is negatively charged because of the phosphate groups) is wrapped tightly around the histone core. This nucleosome is linked to the next one with the help of a linker DNA. This is also known as the “beads on a string” structure. This is further compacted into a 30 nm fiber, which is the diameter of the structure. At the metaphase stage the chromosomes are at their most compact, approximately 700 nm in width, and are found in association with scaffold proteins.



Eukaryotic chromosomes: These figures illustrate the compaction of the eukaryotic chromosome.

In interphase, eukaryotic chromosomes have two distinct regions that can be distinguished by staining. The tightly packaged region is known as heterochromatin, and the less dense region is known as euchromatin.

Heterochromatin usually contains genes that are not expressed, and is found in the regions of the centromere and telomeres. The euchromatin usually contains genes that are transcribed, with DNA packaged around nucleosomes but not further compacted.

This page titled 28.4C: DNA Packaging is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.4D: Types of RNA

RNA is the nucleic acid that makes proteins from the code provided by DNA through the processes of transcription and translation.

Learning Objectives

- Describe the structure and function of RNA

Key Points

- The nitrogen bases in RNA include adenine (A), guanine (G), cytosine (C), and uracil (U).
- Messenger RNA (mRNA) carries the code from the DNA to the ribosomes, while transfer RNA (tRNA) converts that code into a usable form.
- Ribosomes are the sites where tRNA and rRNA assemble proteins.
- RNA differs from DNA in that it is single stranded, has uracil instead of thymine, carries the code for making proteins instead of directing all of the cell ‘s functions, and has ribose as its five-carbon sugar instead of deoxyribose.


Key Terms

- **codon**: a sequence of three adjacent nucleotides, which encode for a specific amino acid during protein synthesis or translation
- **transcription**: the synthesis of RNA under the direction of DNA

RNA Structure and Function

The two main types of nucleic acids are deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). DNA is the genetic material found in all living organisms and is found in the nucleus of eukaryotes and in the chloroplasts and mitochondria. In prokaryotes, the DNA is not enclosed in a membranous envelope.

The other type of nucleic acid, RNA, is mostly involved in protein synthesis. Just like in DNA, RNA is made of monomers called nucleotides. Each nucleotide is made up of three components: a nitrogenous base, a pentose (five-carbon) sugar called ribose, and a phosphate group. Each nitrogenous base in a nucleotide is attached to a sugar molecule, which is attached to one or more phosphate groups.



RNA Structure: A nucleotide is made up of three components: a nitrogenous base, a pentose sugar, and one or more phosphate groups. Carbon residues in the pentose are numbered 1′ through 5′ (the prime distinguishes these residues from those in the base, which are numbered without using a prime notation). The base is attached to the 1′ position of the ribose, and the phosphate is attached to the 5′ position. When a polynucleotide is formed, the 5′ phosphate of the incoming nucleotide attaches to the 3′ hydroxyl group at the end of the growing chain. Two types of pentose are found in nucleotides, deoxyribose (found in DNA) and ribose (found in RNA). Deoxyribose is similar in structure to ribose, but it has an H instead of an OH at the 2′ position. Bases can be divided into two categories: purines and pyrimidines. Purines have a double ring structure, and pyrimidines have a single ring.

In RNA, the nitrogenous bases vary slightly from those of DNA. Adenine (A), guanine (G), and cytosine (C) are present, but instead of thymine (T), a pyrimidine called uracil (U) pairs with adenine. RNA is a single stranded molecule, compared to the double helix of DNA.

The DNA molecules never leave the nucleus but instead use an intermediary to communicate with the rest of the cell. This intermediary is the messenger RNA (mRNA). When proteins need to be made, the mRNA enters the nucleus and attaches itself to one of the DNA strands. Being complementary, the sequence of nitrogen bases of the RNA is opposite that of the DNA. This is called transcription. For example, if the DNA strand reads TCCAAGTC, then the mRNA strand would read AGGUUCAG. The mRNA then carries the code out of the nucleus to organelles called ribosomes for the assembly of proteins.

Once the mRNA has reached the ribosomes, they do not read the instructions directly. Instead, another type of RNA called transfer RNA (tRNA) needs to translate the information from the mRNA into a usable form. The tRNA attaches to the mRNA, but with the opposite base pairings. It then reads the sequence in sets of three bases called codons. Each possible three letter arrangement of A,C,U,G (e.g., AAA, AAU, GGC, etc) is a specific instruction, and the correspondence of these instructions and the amino acids is known as the “genetic code.” Though exceptions to or variations on the code exist, the standard genetic code holds true in most organisms.

The ribosome acts like a giant clamp, holding all of the players in position, and facilitating both the pairing of bases between the messenger and transfer RNAs, and the chemical bonding between the amino acids. The ribosome has special subunits known as ribosomal RNAs (rRNA) because they function in the ribosome. These subunits do not carry instructions for making a specific proteins (i.e., they are not messenger RNAs) but instead are an integral part of the ribosome machinery that is used to make proteins from mRNAs. The making of proteins by reading instructions in mRNA is generally known as “translation.”

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- genome. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/genome. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleotide. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/nucleotide. **License**: ***CC BY-SA: Attribution-ShareAlike***
- monomer. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/monomer. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- mutation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/mutation. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: www.boundless.com//biology/de...n/antiparallel. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_03.png. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_02.jpg. **License**: ***CC BY: Attribution***
- DNA structure. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/GFy_h8cu@9.8...and-Sequencing. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Histones. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Histone. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_03.png. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_02.jpg. **License**: ***CC BY: Attribution***
- figure7.jpg. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72...f21b5eabd@9.87. **License**: ***CC BY-SA: Attribution-ShareAlike***
- figure6.jpg. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72...f21b5eabd@9.87. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...ol11448/latest. **License**: ***CC BY: Attribution***
- Fact Sheet: DNA-RNA-Protein. **Provided by**: microBEnet. **Located at**: http://microbe.net/simple-guides/fac...a-rna-protein/. **License**: ***CC BY-SA: Attribution-ShareAlike***
- codon. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/codon. **License**: ***CC BY-SA: Attribution-ShareAlike***
- transcription. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/transcription. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_01.jpg. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_03.png. **License**: ***CC BY: Attribution***
- OpenStax College, Nucleic Acids. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44403/latest...e_03_05_02.jpg. **License**: ***CC BY: Attribution***
- figure7.jpg. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72...f21b5eabd@9.87. **License**: ***CC BY-SA: Attribution-ShareAlike***
- figure6.jpg. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72...f21b5eabd@9.87. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Nucleic Acids. May 20, 2015. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/cb178029-ce1.../Nucleic_Acids. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 28.4D: Types of RNA is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.5: CELL DIVISION

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.5A: The Role of the Cell Cycle

28.5B: Genomic DNA and Chromosomes

28.5C: Eukaryotic Chromosomal Structure and Compaction

This page titled 28.5: Cell Division is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.5A: The Role of the Cell Cycle

The cell cycle allows multiicellular organisms to grow and divide and single-celled organisms to reproduce.

Learning Objectives

- Explain the role of the cell cycle in carrying out the cell's essential functions

Key Points


- All multicellular organisms use cell division for growth and the maintenance and repair of cells and tissues.
- Single-celled organisms use cell division as their method of reproduction.
- Somatic cells divide regularly; all human cells (except for the cells that produce eggs and sperm) are somatic cells.
- Somatic cells contain two copies of each of their chromosomes (one copy from each parent).
- The cell cycle has two major phases: interphase and the mitotic phase.
- During interphase, the cell grows and DNA is replicated; during the mitotic phase, the replicated DNA and cytoplasmic contents are separated and the cell divides.

Key Terms

- **somatic cell**: any normal body cell of an organism that is not involved in reproduction; a cell that is not on the germline
- **interphase**: the stage in the life cycle of a cell where the cell grows and DNA is replicated
- **mitotic phase**: replicated DNA and the cytoplasmic material are divided into two identical cells

Introduction: Cell Division and Reproduction

A human, as well as every sexually-reproducing organism, begins life as a fertilized egg or zygote. Trillions of cell divisions subsequently occur in a controlled manner to produce a complex, multicellular human. In other words, that original single cell is the ancestor of every other cell in the body. Once a being is fully grown, cell reproduction is still necessary to repair or regenerate tissues. For example, new blood and skin cells are constantly being produced. All multicellular organisms use cell division for growth and the maintenance and repair of cells and tissues. Cell division is tightly regulated because the occasional failure of regulation can have life-threatening consequences. Single-celled organisms use cell division as their method of reproduction.




Cell Division and Growth: A sea urchin begins life as a single cell that (a) divides to form two cells, visible by scanning electron microscopy. After four rounds of cell division, (b) there are 16 cells, as seen in this SEM image. After many rounds of cell division, the individual develops into a complex, multicellular organism, as seen in this (c) mature sea urchin.

While there are a few cells in the body that do not undergo cell division, most somatic cells divide regularly. A somatic cell is a general term for a body cell: all human cells, except for the cells that produce eggs and sperm (which are referred to as germ

cells), are somatic cells. Somatic cells contain two copies of each of their chromosomes (one copy received from each parent). Cells in the body replace themselves over the lifetime of a person. For example, the cells lining the gastrointestinal tract must be frequently replaced when constantly “worn off” by the movement of food through the gut. But what triggers a cell to divide and how does it prepare for and complete cell division?

The cell cycle is an ordered series of events involving cell growth and cell division that produces two new daughter cells. Cells on the path to cell division proceed through a series of precisely timed and carefully regulated stages of growth, DNA replication, and division that produces two identical (clone) cells. The cell cycle has two major phases: interphase and the mitotic phase. During interphase, the cell grows and DNA is replicated. During the mitotic phase, the replicated DNA and cytoplasmic contents are separated and the cell divides.



The Cell Cycle: The cell cycle consists of interphase and the mitotic phase. During interphase, the cell grows and the nuclear DNA is duplicated. Interphase is followed by the mitotic phase. During the mitotic phase, the duplicated chromosomes are segregated and distributed into daughter nuclei. The cytoplasm is usually divided as well, resulting in two daughter cells

This page titled 28.5A: The Role of the Cell Cycle is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.5B: Genomic DNA and Chromosomes

The genome of an organism consists of its entire complement of DNA, which encodes the genes that control the organism's characteristics.

Learning Objectives

- Explain the importance of a genome to an organism

Key Points


- A cell 's DNA, packaged as a double-stranded DNA molecule, is called its genome.
- In prokaryotes, the genome is composed of a single, double-stranded DNA molecule in the form of a loop or circle; the region in the cell containing this genetic material is called a nucleoid.
- In eukaryotes, the genome consists of several double-stranded linear DNA molecules; each species of eukaryotes has a characteristic number of chromosomes in the nuclei of its cells.
- Matched pairs of chromosomes in a diploid organism are called homologous chromosomes, which are the same length and have specific nucleotide segments called genes in exactly the same location, or locus.
- Each copy of a homologous pair of chromosomes originates from a different parent, so the genes themselves are not identical.
- The difference between the DNA sequences in pairs of homologous chromosomes is less than one percent; the sex chromosomes, X and Y, are the single exception to this rule since their genes are different.

Key Terms

- **genome**: the cell's complete genetic information packaged as a double-stranded DNA molecule
- **nucleoid**: the irregularly-shaped region within a prokaryote cell where the genetic material is localized
- **gene**: a unit of heredity; the functional units of chromosomes that determine specific characteristics by coding for specific proteins
- **chromosome**: a structure in the cell nucleus that contains DNA, histone protein, and other structural proteins
- **locus**: a fixed position on a chromosome that may be occupied by one or more genes


Genomic DNA

Before discussing the steps a cell must undertake to replicate, a deeper understanding of the structure and function of a cell's genetic information is necessary. A cell's DNA, packaged as a double-stranded DNA molecule, is called its genome. In prokaryotes, the genome is composed of a single, double-stranded DNA molecule in the form of a loop or circle. The region in the cell containing this genetic material is called a nucleoid. Some prokaryotes also have smaller loops of DNA called plasmids that are not essential for normal growth. Bacteria can exchange these plasmids with other bacteria, sometimes receiving beneficial new genes that the recipient can add to their chromosomal DNA. Antibiotic resistance is one trait that often spreads through a bacterial colony through plasmid exchange.



Prokaryotic Genome: Prokaryotes, including bacteria and archaea, have a single, circular chromosome located in a central region called the nucleoid.

In eukaryotes, the genome consists of several double-stranded linear DNA molecules packaged into chromosomes. Each species of eukaryotes has a characteristic number of chromosomes in the nuclei of its cells. Human body cells have 46 chromosomes, while human gametes (sperm or eggs) have 23 chromosomes each. A typical body cell, or somatic cell, contains two matched sets of chromosomes, a configuration known as diploid. The letter n is used to represent a single set of chromosomes; therefore, a diploid organism is designated 2n. Human cells that contain one set of chromosomes are called gametes, or sex cells; these are eggs and sperm, and are designated 1n, or haploid.



Eukaryotic Genome: There are 23 pairs of homologous chromosomes in a female human somatic cell. The condensed chromosomes are viewed within the nucleus (top), removed from a cell in mitosis and spread out on a slide (right), and artificially arranged according to length (left); an arrangement like this is called a karyotype. In this image, the chromosomes were exposed to fluorescent stains for differentiation of the different chromosomes. A method of staining called “chromosome painting” employs fluorescent dyes that highlight chromosomes in different colors.

Matched pairs of chromosomes in a diploid organism are called homologous (“same knowledge”) chromosomes. Homologous chromosomes are the same length and have specific nucleotide segments called genes in exactly the same location, or locus. Genes, the functional units of chromosomes, determine specific characteristics, or traits, by coding for specific proteins. For

example, hair color is a trait that can be blonde, brown, or black.

Each copy of a homologous pair of chromosomes originates from a different parent; therefore, the genes themselves are not identical. The variation of individuals within a species is due to the specific combination of the genes inherited from both parents. Even a slightly altered sequence of nucleotides within a gene can result in an alternative trait. For example, there are three possible gene sequences on the human chromosome that code for blood type: sequence A, sequence B, and sequence O. Because all diploid human cells have two copies of the chromosome that determines blood type, the blood type (the trait) is determined by which two versions of the marker gene are inherited. It is possible to have two copies of the same gene sequence on both homologous chromosomes, with one on each (for example, AA, BB, or OO), or two different sequences, such as AB, AO, or BO.

Minor variations of traits, such as blood type, eye color, and handedness, contribute to the natural variation found within a species. However, if the entire DNA sequence from any pair of human homologous chromosomes is compared, the difference is less than one percent. The sex chromosomes, X and Y, are the single exception to the rule of homologous chromosome uniformity. Other than a small amount of homology that is necessary to accurately produce gametes, the genes found on the X and Y chromosomes are different.

This page titled 28.5B: Genomic DNA and Chromosomes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.5C: Eukaryotic Chromosomal Structure and Compaction

Chromosomes must coil to pack DNA into the cell during cell division, a process involving 3 levels of compaction.

Learning Objectives

- Describe the levels of chromsomal structure and compaction

Key Points

- During some stages of the cell cycle, the long strands of DNA are condensed into compact chromosomes to fit in the cell's nucleus.
- In the first level of compaction, short stretches of the DNA double helix wrap around a core of eight histone proteins at regular intervals along the entire length of the chromosome.
- The DNA surrouding the histone core is called a nucleosome; the DNA-histone complex is called chromatin.
- The second level of compaction occurs as the nucleosomes and the linker DNA between them are coiled into a 30-nm chromatin fiber, which shortens the chromosome so it's about 50 times shorter than the extended form.
- After replication, the chromosomes are composed of two linked sister chromatids; when fully compact, the pairs of identically-packed chromosomes are bound to each other by cohesin proteins.
- The connection between the sister chromatids is closest in a region called the centromere; this region is highly condensed.


Key Terms

- **nucleosome**: any of the subunits that repeat in chromatin; a coil of DNA surrounding a histone core
- **histone**: any of various simple water-soluble proteins that are rich in the basic amino acids lysine and arginine and are complexed with DNA in the nucleosomes of eukaryotic chromatin
- **chromatin**: a complex of DNA, RNA, and proteins within the cell nucleus out of which chromosomes condense during cell division

Eukaryotic Chromosomal Structure and Compaction

If the DNA from all 46 chromosomes in a human cell nucleus was laid out end to end, it would measure approximately two meters. However, the diameter would be only 2 nm. Considering that the size of a typical human cell is about 10 µm (100,000 cells lined up to equal one meter), DNA must be tightly packaged to fit in the cell's nucleus. At the same time, it must also be readily accessible for the genes to be expressed. During some stages of the cell cycle, the long strands of DNA are condensed into compact chromosomes. There are a number of ways that chromosomes are compacted to fit in the cell's nucleus and be accessible for gene expression.

In the first level of compaction, short stretches of the DNA double helix wrap around a core of eight histone proteins at regular intervals along the entire length of the chromosome. The DNA-histone complex is called chromatin. The beadlike, histone DNA complex is called a nucleosome. DNA connecting the nucleosomes is called linker DNA. A DNA molecule in this form is about seven times shorter than the double helix without the histones. The beads are about 10 nm in diameter, in contrast with the 2-nm diameter of a DNA double helix. The next level of compaction occurs as the nucleosomes and the linker DNA between them are coiled into a 30-nm chromatin fiber. This coiling further shortens the chromosome so that it is now about 50 times shorter than the extended form. In the third level of packing, a variety of fibrous proteins is used to pack the chromatin. These fibrous proteins also ensure that each chromosome in a non-dividing cell occupies a particular area of the nucleus that does not overlap with that of any other chromosome.



Levels of DNA Compaction: Double-stranded DNA wraps around histone proteins to form nucleosomes that have the appearance of "beads on a string." The nucleosomes are coiled into a 30-nm chromatin fiber. When a cell undergoes mitosis, the chromosomes condense even further.

DNA replicates in the S phase of interphase. After replication, the chromosomes are composed of two linked sister chromatids. When fully compact, the pairs of identically-packed chromosomes are bound to each other by cohesin proteins. The connection between the sister chromatids is closest in a region called the centromere. The conjoined sister chromatids, with a diameter of about 1 µm, are visible under a light microscope. The centromeric region is highly condensed and will appear as a constricted area.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44457/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44460/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Growth and Division. November 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m46034/latest/**. **License**: ***CC BY: Attribution***
- interphase. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/interphase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mitotic phase. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/mitotic%20phase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- somatic cell. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/somatic_cell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44460/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44457/latest...0_00_02abc.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- locus. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/locus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleoid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nucleoid**. **License**: ***CC BY-SA: Attribution-***

ShareAlike

- chromosome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chromosome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- genome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/genome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gene. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gene**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44460/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44457/latest...0_00_02abc.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Division October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...e_10_01_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Division October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...e_10_01_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- histone. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/histone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nucleosome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/nucleosome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chromatin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chromatin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. November 5, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44460/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Introduction. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44457/latest...0_00_02abc.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Division October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...e_10_01_01.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Division October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...e_10_01_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Cell Division October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44459/latest...e_10_01_03.jpg**. **License**: ***CC BY: Attribution***

This page titled 28.5C: Eukaryotic Chromosomal Structure and Compaction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.6: SOURCES OF GENETIC VARIATION

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

28.6A: Genetic Variation
28.6B: Mendel's Law of Segregation
28.6C: Mendel's Law of Independent Assortment

This page titled 28.6: Sources of Genetic Variation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.6A: Genetic Variation

Genetic variation is a measure of the variation that exists in the genetic makeup of individuals within population.

Learning Objectives

- Assess the ways in which genetic variance affects the evolution of populations

Key Points

- Genetic variation is an important force in evolution as it allows natural selection to increase or decrease frequency of alleles already in the population.
- Genetic variation can be caused by mutation (which can create entirely new alleles in a population), random mating, random fertilization, and recombination between homologous chromosomes during meiosis (which reshuffles alleles within an organism's offspring).
- Genetic variation is advantageous to a population because it enables some individuals to adapt to the environment while maintaining the survival of the population.

Key Terms

- **genetic diversity**: the level of biodiversity, refers to the total number of genetic characteristics in the genetic makeup of a species
- **crossing over**: the exchange of genetic material between homologous chromosomes that results in recombinant chromosomes
- **phenotypic variation**: variation (due to underlying heritable genetic variation); a fundamental prerequisite for evolution by natural selection
- **genetic variation**: variation in alleles of genes that occurs both within and among populations

Genetic Variation

Genetic variation is a measure of the genetic differences that exist within a population. The genetic variation of an entire species is often called genetic diversity. Genetic variations are the differences in DNA segments or genes between individuals and each variation of a gene is called an allele.For example, a population with many different alleles at a single chromosome locus has a high amount of genetic variation. Genetic variation is essential for natural selection because natural selection can only increase or decrease frequency of alleles that already exist in the population.

Genetic variation is caused by:


- mutation
- random mating between organisms
- random fertilization
- crossing over (or recombination) between chromatids of homologous chromosomes during meiosis

The last three of these factors reshuffle alleles within a population, giving offspring combinations which differ from their parents and from others.



Genetic variation in the shells of Donax variabilis: An enormous amount of phenotypic variation exists in the shells of Donax varabilis, otherwise known as the coquina mollusc. This phenotypic variation is due at least partly to genetic variation within the coquina population.

Evolution and Adaptation to the Environment



Low genetic diversity in the wild cheetah population: Populations of wild cheetahs have very low genetic variation. Because wild cheetahs are threatened, their species has a very low genetic diversity. This low genetic diversity means they are often susceptible to disease and often pass on lethal recessive mutations; only about 5% of cheetahs survive to adulthood.

Variation allows some individuals within a population to adapt to the changing environment. Because natural selection acts directly only on phenotypes, more genetic variation within a population usually enables more phenotypic variation. Some new alleles increase an organism's ability to survive and reproduce, which then ensures the survival of the allele in the population. Other new alleles may be immediately detrimental (such as a malformed oxygen-carrying protein) and organisms carrying these new mutations will die out. Neutral alleles are neither selected for nor against and usually remain in the population. Genetic variation is advantageous because it enables some individuals and, therefore, a population, to survive despite a changing environment.

Geographic Variation

Some species display geographic variation as well as variation within a population. Geographic variation, or the distinctions in the genetic makeup of different populations, often occurs when populations are geographically separated by environmental barriers or when they are under selection pressures from a different environment. One example of geographic variation are clines: graded changes in a character down a geographic axis.

Sources of Genetic Variation

Gene duplication, mutation, or other processes can produce new genes and alleles and increase genetic variation. New genetic variation can be created within generations in a population, so a population with rapid reproduction rates will probably have high genetic variation. However, existing genes can be arranged in new ways from chromosomal crossing over and recombination in sexual reproduction. Overall, the main sources of genetic variation are the formation of new alleles, the altering of gene number or position, rapid reproduction, and sexual reproduction.

This page titled 28.6A: Genetic Variation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.6B: Mendel's Law of Segregation

Mendel's Law of Segregation states that a diploid organism passes a randomly selected allele for a trait to its offspring, such that the offspring receives one allele from each parent.

Learning Objectives

- Apply the law of segregation to determine the chances of a particular genotype arising from a genetic cross

Key Points


- Each gamete acquires one of the two alleles as chromosomes separate into different gametes during meiosis.
- Heterozygotes, which posess one dominant and one recessive allele, can receive each allele from either parent and will look identical to homozygous dominant individuals; the Law of Segregation supports Mendel's observed 3:1 phenotypic ratio.
- Mendel proposed the Law of Segregation after observing that pea plants with two different traits produced offspring that all expressed the dominant trait, but the following generation expressed the dominant and recessive traits in a 3:1 ratio.

Key Terms

- **law of segregation**: a diploid individual possesses a pair of alleles for any particular trait and each parent passes one of these randomly to its offspring

Equal Segregation of Alleles

Observing that true-breeding pea plants with contrasting traits gave rise to F_1 generations that all expressed the dominant trait and F_2generations that expressed the dominant and recessive traits in a 3:1 ratio, Mendel proposed the law of segregation. The law of segregation states that each individual that is a diploid has a pair of alleles (copy) for a particular trait. Each parent passes an allele at random to their offspring resulting in a diploid organism. The allele that contains the dominant trait determines the phenotype of the offspring. In essence, the law states that copies of genes separate or segregate so that each gamete receives only one allele.



The Law of Segregation states that alleles segregate randomly into gametes: When gametes are formed, each allele of one parent segregates randomly into the gametes, such that half of the parent's gametes carry each allele.

For the F_2 generation of a monohybrid cross, the following three possible combinations of genotypes could result: homozygous dominant, heterozygous, or homozygous recessive. Because heterozygotes could arise from two different pathways (receiving one dominant and one recessive allele from either parent), and because heterozygotes and homozygous dominant individuals are phenotypically identical, the law supports Mendel's observed 3:1 phenotypic ratio. The equal segregation of alleles is the reason we can apply the Punnett square to accurately predict the offspring of parents with known genotypes.

The physical basis of Mendel's law of segregation is the first division of meiosis in which the homologous chromosomes with their different versions of each gene are segregated into daughter nuclei. The behavior of homologous chromosomes during meiosis can account for the segregation of the alleles at each genetic locus to different gametes. As chromosomes separate into different gametes during meiosis, the two different alleles for a particular gene also segregate so that each gamete acquires one of the two alleles. In Mendel's experiments, the segregation and the independent assortment during meiosis in the F1 generation give rise to the F2 phenotypic ratios observed by Mendel. The role of the meiotic segregation of chromosomes in sexual reproduction was not understood by the scientific community during Mendel's lifetime.

This page titled 28.6B: Mendel's Law of Segregation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.6C: Mendel's Law of Independent Assortment

Independent assortment allows the calculation of genotypic and phenotypic ratios based on the probability of individual gene combinations.

Learning Objectives

- Use the probability or forked line method to calculate the chance of any particular genotype arising from a genetic cross

Key Points

- Mendel's law of independent assortment states that genes do not influence each other with regard to the sorting of alleles into gametes; every possible combination of alleles for every gene is equally likely to occur.
- The calculation of any particular genotypic combination of more than one gene is, therefore, the probability of the desired genotype at the first locus multiplied by the probability of the desired genotype at the other loci.
- The forked line method can be used to calculate the chances of all possible genotypic combinations from a cross, while the probability method can be used to calculate the chance of any one particular genotype that might result from that cross.


Key Terms

- **independent assortment**: separate genes for separate traits are passed independently of one another from parents to offspring

Independent Assortment

Mendel's law of independent assortment states that genes do not influence each other with regard to the sorting of alleles into gametes: every possible combination of alleles for every gene is equally likely to occur. The independent assortment of genes can be illustrated by the dihybrid cross: a cross between two true-breeding parents that express different traits for two characteristics. Consider the characteristics of seed color and seed texture for two pea plants: one that has green, wrinkled seeds (yyrr) and another that has yellow, round seeds (YYRR). Because each parent is homozygous, the law of segregation indicates that the gametes for the green/wrinkled plant all are yr, while the gametes for the yellow/round plant are all YR. Therefore, the F_1 generation of offspring all are YyRr.

For the F2 generation, the law of segregation requires that each gamete receive either an R allele or an r allele along with either a Y allele or a y allele. The law of independent assortment states that a gamete into which an r allele sorted would be equally likely to contain either a Y allele or a y allele. Thus, there are four equally likely gametes that can be formed when the YyRr heterozygote is self-crossed as follows: YR, Yr, yR, and yr. Arranging these gametes along the top and left of a 4 × 4 Punnett square gives us 16 equally likely genotypic combinations. From these genotypes, we infer a phenotypic ratio of 9 round/yellow:3 round/green:3 wrinkled/yellow:1 wrinkled/green. These are the offspring ratios we would expect, assuming we performed the crosses with a large enough sample size.




Independent assortment of 2 genes: This dihybrid cross of pea plants involves the genes for seed color and texture.

Because of independent assortment and dominance, the 9:3:3:1 dihybrid phenotypic ratio can be collapsed into two 3:1 ratios, characteristic of any monohybrid cross that follows a dominant and recessive pattern. Ignoring seed color and considering only seed texture in the above dihybrid cross, we would expect that three-quarters of the F_2 generation offspring would be round and one-quarter would be wrinkled. Similarly, isolating only seed color, we would assume that three-quarters of the F_2 offspring would be yellow and one-quarter would be green. The sorting of alleles for texture and color are independent events, so we can apply the product rule. Therefore, the proportion of round and yellow F_2 offspring is expected to be (3/4) × (3/4) = 9/16, and the proportion of wrinkled and green offspring is expected to be (1/4) × (1/4) = 1/16. These proportions are identical to those obtained using a Punnett square. Round/green and wrinkled/yellow offspring can also be calculated using the product rule as each of these genotypes includes one dominant and one recessive phenotype. Therefore, the proportion of each is calculated as (3/4) × (1/4) = 3/16.

Forked-Line Method

When more than two genes are being considered, the Punnett-square method becomes unwieldy. For instance, examining a cross involving four genes would require a 16 × 16 grid containing 256 boxes. It would be extremely cumbersome to manually enter each genotype. For more complex crosses, the forked-line and probability methods are preferred.

To prepare a forked-line diagram for a cross between F_1 heterozygotes resulting from a cross between AABBCC and aabbcc parents, we first create rows equal to the number of genes being considered and then segregate the alleles in each row on forked lines according to the probabilities for individual monohybrid crosses. We then multiply the values along each forked path to obtain the F_2offspring probabilities. Note that this process is a diagrammatic version of the product rule. The values along each forked pathway can be multiplied because each gene assorts independently. For a trihybrid cross, the F_2 phenotypic ratio is 27:9:9:9:3:3:3:1.



Independent assortment of 3 genes: The forked-line method can be used to analyze a trihybrid cross. Here, the probability for color in the F2 generation occupies the top row (3 yellow:1 green). The probability for shape occupies the second row (3 round:1 wrinked), and the probability for height occupies the third row (3 tall:1 dwarf). The probability for each possible combination of traits is calculated by multiplying the probability for each individual trait. Thus, the probability of F2 offspring having yellow, round, and tall traits is 3 × 3 × 3, or 27.

Probability Method

While the forked-line method is a diagrammatic approach to keeping track of probabilities in a cross, the probability method gives the proportions of offspring expected to exhibit each phenotype (or genotype) without the added visual assistance.

To fully demonstrate the power of the probability method, however, we can consider specific genetic calculations. For instance, for a tetrahybrid cross between individuals that are heterozygotes for all four genes, and in which all four genes are sorting independently in a dominant and recessive pattern, what proportion of the offspring will be expected to be homozygous recessive for all four alleles? Rather than writing out every possible genotype, we can use the probability method. We know that for each gene the fraction of homozygous recessive offspring will be 1/4. Therefore, multiplying this fraction for each of the four genes, (1/4) × (1/4) × (1/4) × (1/4), we determine that 1/256 of the offspring will be quadruply homozygous recessive.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Structural Biochemistry/Organismic and Evolutionary Biology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...ionary_Biology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 22, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44584/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- A-level Biology/Central Concepts/Classification, selection and evolution. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/A-level..._and_evolution**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Structural Biochemistry/Evolution of Populations. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...etic_Variation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- crossing over. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/crossing%20over**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- genetic variation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/genetic%20variation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phenotypic variation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/phenotypic%20variation**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Genetic Diversity. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Genetic_diversity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coquina variation3. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Co...variation3.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cheetah genetic diversity. **Provided by**: Wikimedia. **Located at**: **en.Wikipedia.org/wiki/File:Ch..._diversity.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44479/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- Structural Biochemistry/Foundation of Genetics. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Structu...on_of_Genetics**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- law of segregation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/law%20of%20segregation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coquina variation3. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Co...variation3.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cheetah genetic diversity. **Provided by**: Wikimedia. **Located at**: **en.Wikipedia.org/wiki/File:Ch..._diversity.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autorecessive. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...orecessive.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44479/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- independent assortment. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/independent%20assortment**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Coquina variation3. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Co...variation3.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cheetah genetic diversity. **Provided by**: Wikimedia. **Located at**: **en.Wikipedia.org/wiki/File:Ch..._diversity.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Autorecessive. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...orecessive.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Laws of Inheritance. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44479/latest...e_12_03_03.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Laws of Inheritance. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **cnx.org/content/m44479/latest...e_12_03_02.jpg**. **License**: ***CC BY: Attribution***

This page titled 28.6C: Mendel's Law of Independent Assortment is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.7: COMPLEX INHERITANCE PATTERNS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

28.3C: Mendel's Model System
28.7A: The Punnett Square Approach for a Monohybrid Cross
28.7B: Phenotypes and Genotypes

This page titled 28.7: Complex Inheritance Patterns is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.3C: Mendel's Model System

The garden pea has several advantageous characteristics that allowed Mendel to develop the laws of modern genetics.

Learning Objectives

- Describe the scientific reasons for the success of Mendel's experimental work

Key Points

- Mendel used true-breeding plants in his experiments. These plants, when self-fertilized, always produce offspring with the same phenotype.
- Pea plants are easily manipulated, grow in one season, and can be grown in large quantities; these qualities allowed Mendel to conduct methodical, quantitative analyses using large sample sizes.
- Based on his experiments with the garden peas, Mendel found that one phenotype was always dominant over another recessive phenotype for the same trait.

Key Terms

- **phenotype**: the observable characteristics of an organism, often resulting from its genetic information or a combination of genetic information and environmental factors
- **genotype**: the specific genetic information of a cell or organism, usually a description of the allele or alleles relating to a specific gene.
- **true-breeding plant**: a plant that always produces offspring of the same phenotype when self-fertilized; one that is homozygous for the trait being followed.

Mendel's Model System

Mendel's seminal work was accomplished using the garden pea, *Pisum sativum*, to study inheritance. Pea plant reproduction is easily manipulated; large quantities of garden peas could be cultivated simultaneously, allowing Mendel to conclude that his results did not occur simply by chance. The garden pea also grows to maturity within one season; several generations could be evaluated over a relatively short time.

Pea plants have both male and female parts and can easily be grown in large numbers. For this reason, garden pea plants can either self-pollinate or cross-pollinate with other pea plants. In the absence of outside manipulation, this species naturally self-fertilizes: ova (the eggs) within individual flowers are fertilized by pollen (containing the sperm cell) from the same flower. The sperm and the eggs that produce the next generation of plants both come from the same parent. What's more, the flower petals remain sealed tightly until after pollination, preventing pollination from other plants. The result is highly inbred, or "true-breeding," pea plants. These are plants that always produce offspring that look like the parent. Today, we know that these "true-breeding" plants are homozygous for most traits.

A gardener or researcher, such as Mendel, can cross-pollinate these same plants by manually applying sperm from one plant to the pistil (containing the ova) of another plant. Now the sperm and eggs come from different parent plants. When Mendel cross-pollinated a true-breeding plant that only produced yellow peas with a true-breeding plant that only produced green peas, he found that the first generation of offspring is always all yellow peas. The green pea trait did not show up. However, if this first generation of yellow pea plants were allowed to self-pollinate, the following or second generation had a ratio of 3:1 yellow to green peas.

In this and all the other pea plant traits Mendel followed, one form of the trait was "dominant" over another so it masked the presence of the other "recessive" form in the first generation after cross-breeding two homozygous plants.. Even if the phenotype (visible form) is hidden, the genotype (allele controlling that form of the trait) can be passed on to next generation and produce the recessive form in the second generation. By experimenting with true-breeding pea plants, Mendel avoided the appearance of

unexpected (recombinant) traits in offspring that might occur if the plants were not true breeding.



Mendel's Experiments With Peas: Experimenting with thousands of garden peas, Mendel uncovered the fundamentals of genetics.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- homozygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/homozygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- heterozygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/heterozygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- monohybrid. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/monohybrid**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Punnett square. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Punnett_square**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **cnx.org/content/m44478/latest...e_12_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 3, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- genotype. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/genotype**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phenotype. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/phenotype**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 1, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44476/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **cnx.org/content/m44478/latest...e_12_02_02.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 3, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/**

latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. November 1, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44476/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44476/latest...ol11448/latest. **License**: *CC BY: Attribution*
- Human Physiology/Genetics and inheritance. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P...nd_inheritance. **License**: *CC BY-SA: Attribution-ShareAlike*
- phenotype. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/phenotype. **License**: *CC BY-SA: Attribution-ShareAlike*
- true-breeding. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/true-breeding. **License**: *CC BY-SA: Attribution-ShareAlike*
- genotype. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/genotype. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: cnx.org/content/m44478/latest...e_12_02_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Biology. November 3, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44478/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Biology. November 1, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44476/latest...ol11448/latest. **License**: *CC BY: Attribution*
- OpenStax College, Introduction. November 1, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44474/latest...2_00_01new.jpg. **License**: *CC BY: Attribution*

This page titled 28.3C: Mendel's Model System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.7A: The Punnett Square Approach for a Monohybrid Cross

A Punnett square applies the rules of probability to predict the possible outcomes of a monohybrid cross and their expected frequencies.

Learning Objectives

- Describe the Punnett square approach to a monohybrid cross

Key Points

- Fertilization between two true-breeding parents that differ in only one characteristic is called a monohybrid cross.
- For a monohybrid cross of two true-breeding parents, each parent contributes one type of allele resulting in all of the offspring with the same genotype.
- A test cross is a way to determine whether an organism that expressed a dominant trait was a heterozygote or a homozygote.


Key Terms

- **monohybrid**: a hybrid between two species that only have a difference of one gene
- **homozygous**: of an organism in which both copies of a given gene have the same allele
- **heterozygous**: of an organism which has two different alleles of a given gene
- **Punnett square**: a graphical representation used to determine the probability of an offspring expressing a particular genotype

Punnett Square Approach to a Monohybrid Cross

When fertilization occurs between two true-breeding parents that differ in only one characteristic, the process is called a monohybrid cross, and the resulting offspring are monohybrids. Mendel performed seven monohybrid crosses involving contrasting traits for each characteristic. On the basis of his results in F_1 and F_2 generations, Mendel postulated that each parent in the monohybrid cross contributed one of two paired unit factors to each offspring and that every possible combination of unit factors was equally likely.


To demonstrate a monohybrid cross, consider the case of true-breeding pea plants with yellow versus green pea seeds. The dominant seed color is yellow; therefore, the parental genotypes were YY (homozygous dominant) for the plants with yellow seeds and yy (homozygous recessive) for the plants with green seeds, respectively. A Punnett square, devised by the British geneticist Reginald Punnett, can be drawn that applies the rules of probability to predict the possible outcomes of a genetic cross or mating and their expected frequencies.To prepare a Punnett square, all possible combinations of the parental alleles are listed along the top (for one parent) and side (for the other parent) of a grid, representing their meiotic segregation into haploid gametes. Then the combinations of egg and sperm are made in the boxes in the table to show which alleles are combining. Each box then represents the diploid genotype of a zygote, or fertilized egg, that could result from this mating. Because each possibility is equally likely, genotypic ratios can be determined from a Punnett square. If the pattern of inheritance (dominant or recessive) is known, the phenotypic ratios can be inferred as well. For a monohybrid cross of two true-breeding parents, each parent contributes one type of allele. In this case, only one genotype is possible. All offspring are Yy and have yellow seeds.



Punnett square analysis of a monohytbrid cross: In the P generation, pea plants that are true-breeding for the dominant yellow phenotype are crossed with plants with the recessive green phenotype. This cross produces F1 heterozygotes with a yellow phenotype. Punnett square analysis can be used to predict the genotypes of the F2 generation.

A self-cross of one of the Yy heterozygous offspring can be represented in a 2 × 2 Punnett square because each parent can donate one of two different alleles. Therefore, the offspring can potentially have one of four allele combinations: YY, Yy, yY, or yy. Notice that there are two ways to obtain the Yy genotype: a Y from the egg and a y from the sperm, or a y from the egg and a Y from the sperm. Both of these possibilities must be counted. Recall that Mendel's pea-plant characteristics behaved in the same way in reciprocal crosses. Therefore, the two possible heterozygous combinations produce offspring that are genotypically and phenotypically identical despite their dominant and recessive alleles deriving from different parents. They are grouped together. Because fertilization is a random event, we expect each combination to be equally likely and for the offspring to exhibit a ratio of YY:Yy:yy genotypes of 1:2:1. Furthermore, because the YY and Yy offspring have yellow seeds and are phenotypically identical, applying the sum rule of probability, we expect the offspring to exhibit a phenotypic ratio of 3 yellow:1 green. Indeed, working with large sample sizes, Mendel observed approximately this ratio in every F_2 generation resulting from crosses for individual traits.

Beyond predicting the offspring of a cross between known homozygous or heterozygous parents, Mendel also developed a way to determine whether an organism that expressed a dominant trait was a heterozygote or a homozygote. Called the test cross, this technique is still used by plant and animal breeders. In a test cross, the dominant-expressing organism is crossed with an organism that is homozygous recessive for the same characteristic. If the dominant-expressing organism is a homozygote, then all F_1 offspring will be heterozygotes expressing the dominant trait. Alternatively, if the dominant expressing organism is a heterozygote, the F_1 offspring will exhibit a 1:1 ratio of heterozygotes and recessive homozygotes. The test cross further validates Mendel's postulate that pairs of unit factors segregate equally.



Example of a test cross: A test cross can be performed to determine whether an organism expressing a dominant trait is a homozygote or a heterozygote.

This page titled 28.7A: The Punnett Square Approach for a Monohybrid Cross is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.7B: Phenotypes and Genotypes

The observable traits expressed by an organism are referred to as its phenotype and its underlying genetic makeup is called its genotype.

Learning Objectives

- Distinguish between the phenotype and the genotype of an organism

Key Points

- Mendel used pea plants with seven distinct traits or phenotypes to determine the pattern of inheritance and the underlying genotypes.
- Mendel found that crossing two purebred pea plants which expressed different traits resulted in an F_1 generation where all the pea plants expressed the same trait or phenotype.
- When Mendel allowed the F_1 plants to self-fertilize, the F_2 generation showed two different phenotypes, indicating that the F_1 plants had different genotypes.


Key Terms

- **phenotype**: the appearance of an organism based on a multifactorial combination of genetic traits and environmental factors, especially used in pedigrees
- **genotype**: the combination of alleles, situated on corresponding chromosomes, that determines a specific trait of an individual, such as “Aa” or “aa”

Phenotypes and Genotypes

The observable traits expressed by an organism are referred to as its phenotype. An organism’s underlying genetic makeup, consisting of both physically visible and non-expressed alleles, is called its genotype. Johann Gregor Mendel’s (1822–1884) hybridization experiments demonstrate the difference between phenotype and genotype.

Mendel crossed or mated two true-breeding (self-pollinating) garden peas, *Pisum saivum*, by manually transferring pollen from the anther of a mature pea plant of one variety to the stigma of a separate mature pea plant of the second variety. Plants used in first-generation crosses were called P_0, or parental generation one, plants. Mendel collected the seeds belonging to the P_0 plants that resulted from each cross and grew them the following season. These offspring were called the F_1, or the first filial (filial = offspring, daughter or son), generation. Once Mendel examined the characteristics in the F_1 generation of plants, he allowed them to self-fertilize naturally. He then collected and grew the seeds from the F_1 plants to produce the F_2, or second filial, generation.



Mendelian crosses: In one of his experiments on inheritance patterns, Mendel crossed plants that were true-breeding for violet flower color with plants true-breeding for white flower color (the P generation). The resulting hybrids in the F_1 generation all had violet flowers. In the F_2 generation, approximately three-quarters of the plants had violet flowers, and one-quarter had white flowers.

When true-breeding plants in which one parent had white flowers and one had violet flowers were cross-fertilized, all of the F1 hybrid offspring had violet flowers. That is, the hybrid offspring were phenotypically identical to the true-breeding parent with violet flowers. However, we know that the allele donated by the parent with white flowers was not simply lost because it reappeared in some of the F2 offspring. Therefore, the F1 plants must have been genotypically different from the parent with violet flowers.

In his 1865 publication, Mendel reported the results of his crosses involving seven different phenotypes, each with two contrasting traits. A trait is defined as a variation in the physical appearance of a heritable characteristic. The characteristics included plant height, seed texture, seed color, flower color, pea pod size, pea pod color, and flower position. To fully examine each characteristic, Mendel generated large numbers of F_1 and F_2 plants, reporting results from 19,959 F_2 plants alone. His findings were consistent. First, Mendel confirmed that he had plants that bred true for white or violet flower color. Regardless of how many generations Mendel examined, all self-crossed offspring of parents with white flowers had white flowers, and all self-crossed offspring of parents with violet flowers had violet flowers. In addition, Mendel confirmed that, other than flower color, the pea plants were physically identical.

This page titled 28.7B: Phenotypes and Genotypes is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.8: PATTERNS OF INHERITANCE

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

28.8A: Lethal Inheritance Patterns

28.8B: Sex-Linked Traits

This page titled 28.8: Patterns of Inheritance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.8A: Lethal Inheritance Patterns

Inheriting two copies of mutated genes that are nonfunctional can have lethal consequences.

Learning Objectives

- Describe recessive and dominant lethal inheritance patterns

Key Points

- An inheritance pattern in which an allele is only lethal in the homozygous form and in which the heterozygote may be normal or have some altered non-lethal phenotype is referred to as recessive lethal.
- The dominant lethal inheritance pattern is one in which an allele is lethal both in the homozygote and the heterozygote; this allele can only be transmitted if the lethality phenotype occurs after reproductive age.
- Dominant lethal alleles are very rare because the allele only lasts one generation and is, therefore, not usually transmitted.
- In the case where dominant lethal alleles might not be expressed until adulthood, the allele may be unknowingly passed on, resulting in a delayed death in both generations.

Key Terms


- **mutation**: any heritable change of the base-pair sequence of genetic material
- **recessive lethal**: an inheritance pattern in which an allele is only lethal in the homozygous form and in which the heterozygote may be normal or have some altered non-lethal phenotype
- **dominant lethal**: an inheritance pattern is one in which an allele is lethal both in the homozygote and the heterozygote; this allele can only be transmitted if the lethality phenotype occurs after reproductive age

Lethal Inheritance Patterns

A large proportion of genes in an individual's genome are essential for survival. Occasionally, a nonfunctional allele for an essential gene can arise by mutation and be transmitted in a population as long as individuals with this allele also have a wild-type, functional copy. The wild-type allele functions at a capacity sufficient to sustain life and is, therefore, considered to be dominant over the nonfunctional allele. However, consider two heterozygous parents that have a genotype of wild-type/nonfunctional mutant for a hypothetical essential gene. In one quarter of their offspring, we would expect to observe individuals that are homozygous recessive for the nonfunctional allele. Because the gene is essential, these individuals might fail to develop past fertilization, die in utero, or die later in life, depending on what life stage requires this gene. An inheritance pattern in which an allele is only lethal in the homozygous form and in which the heterozygote may be normal or have some altered non-lethal phenotype is referred to as recessive lethal.

For crosses between heterozygous individuals with a recessive lethal allele that causes death before birth when homozygous, only wild-type homozygotes and heterozygotes would be observed. The genotypic ratio would therefore be 2:1. In other instances, the recessive lethal allele might also exhibit a dominant (but not lethal) phenotype in the heterozygote. For instance, the recessive lethal Curly allele in Drosophila affects wing shape in the heterozygote form, but is lethal in the homozygote.

Dominant Lethal Alleles



Effects of Huntington's disease on neurons: The neuron in the center of this micrograph (yellow) has nuclear inclusions characteristic of Huntington's disease (orange area in the center of the neuron). Huntington's disease occurs when an abnormal dominant allele for the Huntington gene is present.

A single copy of the wild-type allele is not always sufficient for normal functioning or even survival. The dominant lethal inheritance pattern is one in which an allele is lethal both in the homozygote and the heterozygote; this allele can only be transmitted if the lethality phenotype occurs after reproductive age. Individuals with mutations that result in dominant lethal alleles fail to survive even in the heterozygote form. Dominant lethal alleles are very rare because, as you might expect, the allele only lasts one generation and is not transmitted.

However, just as the recessive lethal allele might not immediately manifest the phenotype of death, dominant lethal alleles also might not be expressed until adulthood. Once the individual reaches reproductive age, the allele may be unknowingly passed on, resulting in a delayed death in both generations.

An example of this in humans is Huntington's disease in which the nervous system gradually wastes away. People who are heterozygous for the dominant Huntington allele (Hh) will inevitably develop the fatal disease. However, the onset of Huntington's disease may not occur until age 40, at which point the afflicted persons may have already passed the allele to 50 percent of their offspring.

This page titled 28.8A: Lethal Inheritance Patterns is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.8B: Sex-Linked Traits

A gene present on one of the sex chromosomes (X or Y in mammals) is a sex-linked trait because its expression depends on the sex of the individual.

Learning Objectives

- Distinguish between sex-linked traits and other forms of inheritance

Key Points

- In mammals, females have a homologous pair of X chromosomes, whereas males have an XY chromosome pair.
- The Y chromosome contains a small region of similarity to the X chromosome so that they can pair during meiosis, but the Y is much shorter and contains fewer genes.
- Males are said to be hemizygous because they have only one allele for any X-linked characteristic; males will exhibit the trait of any gene on the X-chromosome regardless of dominance and recessiveness.
- Most sex-linked traits are actually X-linked, such as eye color in *Drosophila* or color blindness in humans.


Key Terms

- **hemizygous**: Having some single copies of genes in an otherwise diploid cell or organism.
- **X-linked**: Associated with the X chromosome.
- **carrier**: A person or animal that transmits a disease to others without itself contracting the disease.
- **sex chromosomes**: A chromosome involved with determining the sex of an organism, typically one of two kinds.

Sex Determination

In humans, as well as in many other animals and some plants, the sex of the individual is determined by sex chromosomes. However, there are other sex determination systems in nature. For example, temperature-dependent sex determination is relatively common,
and there are many other types of environmental sex determination. Some species, such as some snails, practice sex change adults start out male, then become female. In tropical clown fish, the dominant individual in a group becomes female while the others are male.

The sex chromosomes are one pair of non-homologous chromosomes. Until now, we have only considered inheritance patterns among non-sex chromosomes, or autosomes. In addition to 22 homologous pairs of autosomes, human females have a homologous pair of X chromosomes, whereas human males have an XY chromosome pair. Although the Y chromosome contains a small region of similarity to the X chromosome so that they can pair during meiosis, the Y chromosome is much shorter and contains many fewer genes. When a gene being examined is present on the X chromosome, but not on the Y chromosome, it is said to be X-linked.



Human male karyotype: A human males possesses XY chromosomes, as seen in the bottom left of this karyotype. The Y chromosome is much shorter than the X chromosome, unlike all of the other homologous chromosome pairs.

X-Linked Traits



Eye color in Drosophila is an example of a X-linked trait: In *Drosophila*, the gene for eye color is located on the X chromosome. Clockwise from top left are brown, cinnabar, sepia, vermilion, white, and red. Red eye color is wild-type and is dominant to white eye color.

Insects also follow an XY sex-determination pattern and like humans, *Drosophila* males have an XY chromosome pair and females are XX. Eye color in *Drosophila* was one of the first X-linked traits to be identified, and Thomas Hunt Morgan mapped this trait to the X chromosome in 1910.


In fruit flies, the wild-type eye color is red (X^W) and is dominant to white eye color (X^w). Because this eye-color gene is located on the X chromosome only, reciprocal crosses do not produce the same offspring ratios. Males are said to be hemizygous, because they have only one allele for any X-linked characteristic. Hemizygosity makes the descriptions of dominance and recessiveness irrelevant for XY males because each male only has one copy of the gene. *Drosophila*males lack a second allele copy on the Y

chromosome; their genotype can only be X^WY or X^wY. In contrast, females have two allele copies of this gene and can be X^WX^W, X^WX^w, or X^wX^w.

X-Linked Crosses

In an X-linked cross, the genotypes of F_1 and F_2 offspring depend on whether the recessive trait was expressed by the male or the female in the P_1 generation. With regard to *Drosophila* eye color, when the P_1 male expresses the white-eye phenotype and the female is homozygous red-eyed, all members of the F_1generation exhibit red eyes. The F_1 females are heterozygous (X^WX^w), and the males are all X^WY, having received their X chromosome from the homozygous dominant P_1 female and their Y chromosome from the P_1 male.


A subsequent cross between the X^WX^w female and the X^WY male would produce only red-eyed females (with X^WX^W or X^WX^wgenotypes) and both red- and white-eyed males (with X^WY or X^wY genotypes). Now, consider a cross between a homozygous white-eyed female and a male with red eyes. The F_1 generation would exhibit only heterozygous red-eyed females (X^WX^w) and only white-eyed males (X^wY). Half of the F_2 females would be red-eyed (X^WX^w) and half would be white-eyed (X^wX^w). Similarly, half of the F_2 males would be red-eyed (X^WY) and half would be white-eyed (X^wY).



Punnett square analysis of *Drosophila* eye color: Punnett square analysis is used to determine the ratio of offspring from a cross between a red-eyed male fruit fly (X^WY) and a white-eyed female fruit fly (X^wX^w).

X-Linked Recessive Disorders in Humans


Sex-linkage studies provided the fundamentals for understanding X-linked recessive disorders in humans, which include red-green color blindness and Types A and B hemophilia. Because human males need to inherit only one recessive mutant X allele to be affected, X-linked disorders are disproportionately observed in males. Females must inherit recessive X-linked alleles from both of their parents in order to express the trait.



Color perception in different types of color blindness: In this chart you can see what people with different types of color blindness can see versus the normal color vision line at top.

Recessive Carriers

When they inherit one recessive X-linked mutant allele and one dominant X-linked wild-type allele, they are carriers of the trait and are typically unaffected. Carrier females can manifest mild forms of the trait due to the inactivation of the dominant allele located on one of the X chromosomes. However, female carriers can contribute the trait to their sons, resulting in the son exhibiting the trait, or they can contribute the recessive allele to their daughters, resulting in the daughters being carriers of the trait. Although some Y-linked recessive disorders exist, typically they are associated with infertility in males and are, therefore, not transmitted to subsequent generations.



Inheritance of a recessive X-linked disorder: The son of a woman who is a carrier of a recessive X-linked disorder will have a 50 percent chance of being affected. A daughter will not be affected, but she will have a 50 percent chance of being a carrier like her mother.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- mutation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mutation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...cessive-lethal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...ominant-lethal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...e_12_02_10.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- X-linked. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/X-linked**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carrier. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/carrier**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hemizygous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hemizygous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sex-determination systems. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sex-determination_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...e_12_02_10.jpg**. **License**: ***CC BY: Attribution***
- Human_male_karyotpe_high_resolution.jpg. **Provided by**: Wikipedia. **Located at**: **commons.wikimedia.org/wiki/F...resolution.jpg**. **License**: ***Public Domain: No Known Copyright***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...e_12_02_08.jpg**. **License**: ***CC BY: Attribution***
- OpenStax College, Biology. November 3, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- OpenStax College, Characteristics and Traits. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44478/latest...e_12_02_09.jpg**. **License**: ***CC BY: Attribution***
- upload.wikimedia.org/wikiped...ndness.svg.png. **Provided by**: Wikipedia. **Located at**: **upload.wikimedia.org/wikiped...ndness.svg.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 28.8B: Sex-Linked Traits is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.9: GENETIC DISEQUILIBRIUM

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

28.9A: Genetic Linkage and Distances

28.9B: Chromosomal Structural Rearrangements

This page titled 28.9: Genetic Disequilibrium is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.9A: Genetic Linkage and Distances

Linked genes can become unlinked during recombination; the probability of genes separating depends on their distance from each other.

Learning Objectives

- Discuss how linked genes can be inherited separately

Key Points

- Alleles positioned on the same chromosome are not always inherited together because during meiosis linked genes can became unlinked.
- Frans Janssen suggested chromosomes become unlinked during homologous recombination, a process where homologous chromosomes exchange segments of DNA.
- Alfred Sturtevant hypothesized that alleles that were closer together on a gene were more likely to be inherited together rather than alleles that were farther apart and used measurements of recombination between genes to create the first genetic map.
- When genes are perfectly linked, they have a recombination frequency of 0.
- When genes are unlinked, they have a recombination frequency of 0.5, which means 50 percent of offspring are recombinants and the other 50 percent are parental types.

Key Terms

- **homologous recombination**: a type of genetic recombination in which nucleotide sequences are exchanged between two similar or identical molecules of DNA
- **linkage**: the property of genes of being inherited together
- **synapsis**: the association of homologous maternal and paternal chromosomes during the initial part of meiosis

Genetic Linkage and Distances


Mendel's work suggested that traits are inherited independently of each other. Morgan identified a 1:1 ratio between a segregating trait and the X chromosome, suggesting that the random segregation of chromosomes was the physical basis of Mendel's model. This also demonstrated that linked genes disrupt Mendel's predicted outcomes. The fact that each chromosome can carry many linked genes explains how individuals can have many more traits than they have chromosomes. However, observations by researchers in Morgan's laboratory suggested that alleles positioned on the same chromosome were not always inherited together. During meiosis, linked genes somehow became unlinked.

Homologous Recombination


In 1909, Frans Janssen observed chiasmata (the point at which chromatids are in contact with each other and may exchange segments) prior to the first division of meiosis. He suggested that alleles become unlinked when chromosomes physically exchange segments. As chromosomes condensed and paired with their homologs, they appeared to interact at distinct points. Janssen suggested that these points corresponded to regions in which chromosome segments were exchanged. It is now known that the pairing and interaction between homologous chromosomes, known as synapsis, does more than simply organize the homologs for migration to separate daughter cells. When synapsed, homologous chromosomes undergo reciprocal physical exchanges of DNA at their arms in a process called homologous recombination, or more simply,"crossing over."

Genetic Maps

In 1913, Alfred Sturtevant, a student in Morgan's laboratory, created the first "chromosome map," a linear representation of gene order and relative distance on a chromosome.To construct a chromosome map, Sturtevant assumed that genes were ordered serially on threadlike chromosomes. He also assumed that the incidence of recombination between two homologous chromosomes could occur with equal likelihood anywhere along the length of the chromosome. Operating under these assumptions, Sturtevant hypothesized alleles that were far apart on a chromosome were more likely to dissociate during meiosis simply because there was a larger region over which recombination could occur. Conversely, alleles that were close to each other on the chromosome were likely to be inherited together. The average number of crossovers between two alleles, or their recombination frequency, correlated with their genetic distance from each other, relative to the locations of other genes on that chromosome. Sturtevant divided his genetic map into map units, or centimorgans (cM), in which a recombination frequency of 0.01 corresponds to 1 cM.



Inheritance Patterns of Unlinked and Linked Genes: In (a), two genes are located on different chromosomes so independent assortment occurs during meiosis. The offspring have an equal chance of being the parental type (inheriting the same combination of traits as the parents) or a nonparental type (inheriting a different combination of traits than the parents). In (b), two genes are very close together on the same chromosome so that no crossing over occurs between them. The genes are, therefore, always inherited together and all of the offspring are the parental type. In (c), two genes are far apart on the chromosome such that crossing over occurs during every meiotic event. The recombination frequency will be the same as if the genes were on separate chromosomes. (d) The actual recombination frequency of fruit fly wing length and body color that Thomas Morgan observed in 1912 was 17 percent. A crossover frequency between 0 percent and 50 percent indicates that the genes are on the same chromosome and crossover occurs some of the time.



Genetic Map Based on Recombination Frequencies in *Drosophila*			
MUTANT			WILD TYPE
Short aristae	0		Long aristae
Black body	48.5		Gray body
Cinnabar eyes	57.5		Red eyes
Vestigial wings	65.5		Normal wings
Brown eyes	104.5		Red eyes
Values in centimorgan (cM) map units; recombination frequency of 0.01 = 1 cM			

Genetic Maps: This genetic map orders Drosophila genes on the basis of recombination frequency.

By representing alleles in a linear map, Sturtevant suggested that genes can range from being perfectly linked (recombination frequency = 0) to being perfectly unlinked (recombination frequency = 0.5) when genes are on different chromosomes or genes are separated very far apart on the same chromosome. Perfectly unlinked genes correspond to the frequencies predicted by Mendel to assort independently in a dihybrid cross. A recombination frequency of 0.5 indicates that 50 percent of offspring are recombinants and the other 50 percent are parental types. That is, every type of allele combination is represented with equal frequency. This allowed Sturtevant to calculate distances between several genes on the same chromosome.

This page titled 28.9A: Genetic Linkage and Distances is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

28.9B: Chromosomal Structural Rearrangements

Structural rearrangements of chromosomes include both inversions and translocations, which may have detrimental effects on an organism.

Learning Objectives

- Describe the various types of structural rearrangements of chromosomes and how they can impact an organism

Key Points

- A chromosome inversion is the detachment, 180° rotation, and reinsertion of part of a chromosome; this may have no effect on the organism, but if the inversion occurs within a gene or moves a gene away from its regulatory elements it can have an adverse effect.
- Pericentric inversions include the centromere, while paracentric inversions occur outside of the centromere; a pericentric inversion can change the length of the chromosome arms above and below the centromere.
- A pericentric inversion on chromsome 18 appears to have been involved in the evolution of humans.
- A translocation occurs when a segment of a chromosome dissociates and reattaches to a different, nonhomologous chromosome and can be benign or detrimental; in reciprocal translocations, there is no gain or loss of genetic information, so these are usually benign.

Key Terms

- **inversion**: a segment of DNA in the context of a chromosome that is reversed in orientation relative to a reference karyotype or genome
- **translocation**: a transfer of a chromosomal segment to a new position, especially on a nonhomologous chromosome


Chromosomal Structural Rearrangements

Cytologists have characterized numerous structural rearrangements in chromosomes, but chromosome inversions and translocations are the most common. Both are identified during meiosis by the adaptive pairing of rearranged chromosomes with their former homologs to maintain appropriate gene alignment. If the genes carried on two homologs are not oriented correctly, a recombination event could result in the loss of genes from one chromosome and the gain of genes on the other. This would produce aneuploid gametes.

Chromosome Inversions


A chromosome inversion is the detachment, 180° rotation, and reinsertion of part of a chromosome. Inversions may occur in nature as a result of mechanical shear, or from the action of transposable elements (special DNA sequences capable of facilitating the rearrangement of chromosome segments with the help of enzymes that cut and paste DNA sequences). Unless they disrupt a gene sequence, inversions only change the orientation of genes and are likely to have milder effects than aneuploid errors. However, altered gene orientation can result in functional changes because regulators of gene expression could be moved out of position with respect to their targets, causing aberrant levels of gene products.

An inversion can be pericentric and include the centromere, or paracentric and occur outside of the centromere. A pericentric inversion that is asymmetric about the centromere can change the relative lengths of the chromosome arms, making these inversions easily identifiable.



Inversions can be pericentric or paracentric: Pericentric inversions include the centromere, and paracentric inversions do not. A pericentric inversion can change the relative lengths of the chromosome arms; a paracentric inversion cannot.

When one homologous chromosome undergoes an inversion, but the other does not, the individual is described as an inversion heterozygote. To maintain point-for-point synapsis during meiosis, one homolog must form a loop, and the other homolog must mold around it. Although this topology can ensure that the genes are correctly aligned, it also forces the homologs to stretch and can be associated with regions of imprecise synapsis.



Inversion heterozygotes: When one chromosome undergoes an inversion, but the other does not, one chromosome must form an inverted loop to retain point-for-point interaction during synapsis. This inversion pairing is essential to maintaining gene alignment during meiosis and to allow for recombination.

Not all structural rearrangements of chromosomes produce nonviable, impaired, or infertile individuals. In rare instances, such a change can result in the evolution of a new species. In fact, a pericentric inversion in chromosome 18 appears to have contributed to the evolution of humans. This inversion is not present in our closest genetic relatives, the chimpanzees. Humans and chimpanzees differ cytogenetically by pericentric inversions on several chromosomes and by the fusion of two separate chromosomes in chimpanzees that correspond to chromosome two in humans.


The pericentric chromosome 18 inversion is believed to have occurred in early humans following their divergence from a common

ancestor with chimpanzees approximately five million years ago. Researchers characterizing this inversion have suggested that approximately 19,000 nucleotide bases were duplicated on 18p, and the duplicated region inverted and reinserted on chromosome 18 of an ancestral human.

A comparison of human and chimpanzee genes in the region of this inversion indicates that two genes—ROCK1 and USP14—that are adjacent on chimpanzee chromosome 17 (which corresponds to human chromosome 18) are more distantly positioned on human chromosome 18. This suggests that one of the inversion breakpoints occurred between these two genes. Interestingly, humans and chimpanzees express USP14 at distinct levels in specific cell types, including cortical cells and fibroblasts. Perhaps the chromosome 18 inversion in an ancestral human repositioned specific genes and reset their expression levels in a useful way. Because both ROCK1 and USP14 encode cellular enzymes, a change in their expression could alter cellular function. It is not known how this inversion contributed to hominid evolution, but it appears to be a significant factor in the divergence of humans from other primates.

Translocations

A translocation occurs when a segment of a chromosome dissociates and reattaches to a different, nonhomologous chromosome. Translocations can be benign or have devastating effects depending on how the positions of genes are altered with respect to regulatory sequences. Notably, specific translocations have been associated with several cancers and with schizophrenia. Reciprocal translocations result from the exchange of chromosome segments between two nonhomologous chromosomes such that there is no gain or loss of genetic information.



Reciprocal translocations do not involve loss of genetic information: A reciprocal translocation occurs when a segment of DNA is transferred from one chromosome to another, nonhomologous chromosome.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: **http://cnx.org/content/m44481/latest...ol11448/latest**. **License**: ***CC BY: Attribution***
- homologous recombination. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/homolog...0recombination**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- synapsis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/synapsis**. **License**: ***CC BY-SA: Attribution-***

ShareAlike
- linkage. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/linkage. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Chromosomal Theory and Genetic Linkage October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44481/latest...e_13_01_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Chromosomal Theory and Genetic Linkage October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44481/latest...e_13_01_03.png. **License**: *CC BY: Attribution*
- translocation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/translocation. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Biology. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44483/latest...ol11448/latest. **License**: *CC BY: Attribution*
- inversion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/inversion. **License**: *CC BY-SA: Attribution-ShareAlike*
- OpenStax College, Chromosomal Theory and Genetic Linkage October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44481/latest...e_13_01_02.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Chromosomal Theory and Genetic Linkage October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44481/latest...e_13_01_03.png. **License**: *CC BY: Attribution*
- OpenStax College, Chromosomal Basis of Inherited Disorders. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44483/latest...e_13_03_09.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Chromosomal Basis of Inherited Disorders. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44483/latest...e_13_03_08.jpg. **License**: *CC BY: Attribution*
- OpenStax College, Chromosomal Basis of Inherited Disorders. October 16, 2013. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/content/m44483/latest...e_13_03_07.jpg. **License**: *CC BY: Attribution*

This page titled 28.9B: Chromosomal Structural Rearrangements is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30: APPENDIX B: DEVELOPMENT AND AGING OF THE ORGAN SYSTEMS

- Topic hierarchy

30.10: Development of the Heart

30.11: Development of Blood Vessels

30.12: Development of Lymphatic Tissues

30.13: Development of the Immune System

30.14: Development of the Respiratory System

30.15: Development of the Digestive System

30.16: Development of Metabolism

30.17: Development of Bodily Fluids

30.18: Development of the Reproductive System

30.1: Integumentary System Development

30.1A: Development of the Integumentary System

30.1B: Effects of Aging on the Integumentary System

30.2: Bone Development

30.2A: Stages of Bone Development

30.2B: Exercise and Bone Tissue

30.2C: Bone Tissue and the Effects of Aging

30.3: Development of the Skeleton

30.3A: Development of the Skeleton

30.3B: Aging and the Skeleton

30.4: Joint Development

30.4A: Development of Joints

30.4B: Joints and Aging

30.4C: Arthroplasty

30.5: Muscle Development

30.5A: Muscle Development

30.6: Development of the Nervous System

30.6A: Development of the Central Nervous System

30.6B: Development of the Peripheral Nervous System

30.6C: Regeneration of Nerve Fibers

30.6D: Aging and the Nervous System

30.7: Development of the Special Senses

30.47C: Development of Hearing and Balance

30.7A: Taste and Smell at Birth and in Old Age

30.7B: Development of Vision

30.8: Development of the Endocrine System

30.9: Development of Blood

This page titled 30: APPENDIX B: Development and Aging of the Organ Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.10: Development of the Heart

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.10: Development of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.11: Development of Blood Vessels

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.
Topic hierarchy

This page titled 30.11: Development of Blood Vessels is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.12: Development of Lymphatic Tissues

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.
Topic hierarchy

This page titled 30.12: Development of Lymphatic Tissues is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.13: Development of the Immune System

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.13: Development of the Immune System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.14: Development of the Respiratory System

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.14: Development of the Respiratory System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.15: Development of the Digestive System

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.
Topic hierarchy

This page titled 30.15: Development of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.16: Development of Metabolism

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.16: Development of Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.17: Development of Bodily Fluids

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.17: Development of Bodily Fluids is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.18: Development of the Reproductive System

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.18: Development of the Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.1: INTEGUMENTARY SYSTEM DEVELOPMENT

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

30.1A: Development of the Integumentary System

30.1B: Effects of Aging on the Integumentary System

This page titled 30.1: Integumentary System Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.1A: Development of the Integumentary System

The integumentary system develops from all embryonic layers (ectoderm, mesoderm, and neural crest cells).

Learning Objectives

- Outline the developmental process of the integumentary system

Key Points


- Ectoderm forms at 4 weeks.
- Between 4 and 12 weeks, the stratified epithelium forms and mesoderm forms the blood vessels and connective tissue.
- At 16 weeks, the basement membrane folds and melanoblasts start producing melanin.
- At 20 weeks, hair begins to grow from sebaceous glands and sweat glands form.

Key Terms

- **sebaceous gland**: A gland of the skin that secretes an oily substance, sebum, usually into a hair follicle near the surface of the skin.
- **neural crest cells**: A transient, multipotent, migratory cell population that gives rise to a diverse cell lineage including melanocytes, craniofacial cartilage and bone, smooth muscle, peripheral and enteric neurons, and glia.
- **fetal skin**: This forms from three layers: ectoderm, mesoderm, and neural crest cells.

The integumentary system is the largest organ system in the human body, responsible for protection from physical and environmental factors. The integumentary system is both a barrier and a sensory organ, and includes the skin (the largest bodily organ), as well as appendages, sweat and sebaceous glands, hair, nails and arrectores pillorum (tiny muscles at the root of each hair that cause goose bumps).

Fetal Skin Formation



Human Skin: This image details the parts of the integumentary system.

Fetal skin forms from three layers: ectoderm, mesoderm, and neural crest cells. Figure 1 shows a diagram of the skin structure.

At 4 weeks gestation, simple ectoderm epithelium forms. Between 4 and 12 weeks, the basal cells divide repeatedly to form the stratified epithelium while the mesoderm forms the blood vessels and connective tissues. Epidermal ridges (e.g. fingerprints) begin to develop around 10 weeks gestation and are completed by 17 weeks gestation. Sensory nerves also develop.

At 16 weeks gestation, the basement membrane folds. Melanoblasts that form melanocytes migrate with neural crests cells to the epithelium and begin producing melanin prior to birth. The connective tissue differentiates into the various layers of the dermis. Ectoderm thickens into fingernails and toenails. Other regions of the ectoderm form into epithelial columns called cords which become hair follicles and sebaceous and sweat glands.

At 20 weeks gestation, hair begins to grow from sebaceous glands, while sweat glands are formed from coiled cords. Other cords begin to form mammary glands.

This page titled 30.1A: Development of the Integumentary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.1B: Effects of Aging on the Integumentary System

The integumentary system undergoes intrinsic and extrinsic cutaneous aging, primarily involving the dermis.

Learning Objectives

- Contrast intrinsic aging with extrinsic aging of the integumentary system

Key Points

- Intrinsic aging is caused by internal factors alone, such as diminished collagen synthesis, and is sometimes known as chronological aging.
- Extrinsic aging is caused by external factors, such as exposure to ultraviolet radiation, which can result in photodamage and may lead to skin cancer formation.
- A validated comprehensive grading scale categorizes the clinical findings of skin aging as laxity, rhytids (wrinkles), and the various facets of photoaging, including erythema/telangiectasia, dyspigmentation, solar elastosis, keratoses, and poor texture.

Key Terms

- **extrinsic aging**: Aging caused by external factors such as exposure to radiation or toxic substances.
- **intrinsic aging**: Aging caused by internal factors related to degeneration of physiologic processes.

Intrinsic and extrinsic aging are terms used to describe cutaneous aging of the skin and other parts of the integumentary system. Aging primarily involves the dermis and has limited effects on the epidermis.

Intrinsic Aging



Wrinkled Skin: Wrinkles form due to intrinsic (loss of collagen) and extrinsic (photodamage) aging.

The effects of intrinsic aging are caused solely by internal factors. Sometimes called chronological aging, intrinsic aging is an inherent degenerative process due to declining physiologic functions and capacities. This may include qualitative and quantitative changes such as diminished or defective synthesis of collagen and elastin in the dermis. As skin ages, it becomes thinner and more easily damaged. Intensifying this effect is the decreasing ability of skin to heal itself with age. Skin aging is also noted by a decrease in volume and elasticity and the increased incidence of wrinkles. Aging skin receives less blood flow and lower glandular activity. Cortisol (associated with stress) causes degradation of collagen, accelerating the aging process.

Extrinsic Aging



Skin Cancer: Image of a basal cell carcinoma caused by extrinsic aging.

Extrinsic aging of skin is caused by external factors such as ultraviolet radiation, cigarette smoking, and air pollution. Of all extrinsic causes, radiation from sunlight has the most widespread documentation of negative effects on the skin. Because of this, extrinsic aging is often referred to as photoaging, defined as skin changes caused by chronic exposure to UV light. Photodamage implies changes beyond those associated with aging alone. It is defined as cutaneous damage caused by chronic exposure to solar radiation and is associated with emergence of neoplastic (cancer) lesions. Photoaging causes two main concerns: an increased risk for skin cancer, and the appearance of damaged skin. In younger skin, sun damage will heal faster since the cells in the epidermis have a faster turnover rate, while in older adults, thinner skin and slower healing may result in damage to the dermal layer.

A validated comprehensive grading scale has categorized the clinical findings of skin aging as laxity (sagging), rhytids (wrinkles), and the various facets of photoaging, including erythema/telangiectasia (redness), dyspigmentation (brown discoloration), solar elastosis (yellowing), keratoses (abnormal growths), and poor texture.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Embryological Development of the Integumentary System. **Provided by**: embryological-development-integumentary Wikispace. **Located at**: http://embryological-development-int...mentary+System. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Scarless wound healing. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Scarless_wound_healing. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neural crest cells. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/neural%20crest%20cells. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fetal skin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/fetal%20skin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sebaceous gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sebaceous_gland. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Integumentary System. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_P..._System%23Skin. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Intrinsic and extrinsic aging. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Intrins...xtrinsic_aging. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human skin. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Human_skin%23Aging. **License**: ***CC BY-SA:***

Attribution-ShareAlike
- extrinsic aging. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/extrinsic%20aging**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intrinsic aging. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intrinsic%20aging**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Human Physiology/Integumentary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P..._System%23Skin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi..._carcinoma.jpg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...y_cropped).jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 30.1B: Effects of Aging on the Integumentary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.2: BONE DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

30.2A: Stages of Bone Development
30.2B: Exercise and Bone Tissue
30.2C: Bone Tissue and the Effects of Aging

This page titled 30.2: Bone Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.2A: Stages of Bone Development

Although bone initially forms during fetal development, it undergoes secondary ossification after birth and is remodeled throughout life.

Learning Objectives

- Describe the process and purpose of bone remodeling

Key Points

- The formation of bone during the fetal stage of development occurs by two processes: intramembranous ossification and endochondral ossification.
- Secondary ossification occurs after birth and forms the epiphyses of long bones and the extremities of irregular and flat bones.
- After initial bone development, bones are remodeled throughout life to regulate calcium homeostasis and repair micro-damaged bones (from everyday stress), as well as to shape the skeleton during growth.

Key Terms

- **diaphysis**: The central shaft of any long bone.
- **epiphyses**: The rounded ends of a long bone at its joint with adjacent bone(s).

EXAMPLES

When a tooth is lost and not replaced, bone remodeling will fill in much of the socket. Although the remodeling will be obvious within a few weeks (especially when smiling), the process will continue for some months.

Bones are rigid organs that constitute part of the endoskeleton of vertebrates. They support and protect the various organs of the body, produce red and white blood cells, and store minerals. Bone tissue is a type of dense connective tissue that appears static, but is actually constantly remodeled throughout the life of the vertebrate organism. This occurs with the synchronized action of osteoclasts and osteoblasts, cells that reabsorb and deposit bone, respectively. Bone remodeling also occurs in response to trauma, such as following an accidental fracture or placement of dental implants.

Initial Bone Formation

The formation of bone during the fetal stage of development occurs by two processes: intramembranous ossification and endochondral ossification.

Intramembranous Ossification

Intramembranous ossification mainly occurs during the formation of the flat bones of the skull, as well as the mandible, maxilla, and clavicles. The bone is formed from connective tissue such as mesenchyme tissue rather than from cartilage. The steps in intramembranous ossification are:

- Development of ossification center
- Calcification
- Formation of trabeculae
- Development of periosteum

Endochondral Ossification

Endochondral ossification begins with points in the cartilage called “primary ossification centers.” They mostly appear during fetal development, though a few short bones begin their primary ossification after birth. These cartilage poitns are responsible for the formation of the diaphyses of long bones, short bones, and certain parts of irregular bones.


Secondary ossification occurs after birth and forms the epiphyses of long bones and the extremities of irregular and flat bones. The diaphysis and both epiphyses of a long bone are separated by a growing zone of cartilage (the epiphyseal plate). When the child reaches skeletal maturity (18 to 25 years of age), all cartilage is replaced by bone, fusing the diaphysis and both epiphyses together (epiphyseal closure).

Remodeling

Remodeling or bone turnover is the process of resorption followed by replacement of bone with little change in shape, and occurs throughout a person’s life, long beyond the initial development of bone. Osteoblasts and osteoclasts, coupled together via paracrine cell signalling, are referred to as a bone remodeling unit. Approximately 10% of the skeletal mass of an adult is remodeled each year.

The bone remodeling period consists of the duration of the resorption, the osteoclastic reversal (the phase marked by shifting of resorption processes into formative processes), and the formation periods of bone growth and development. The bone remodeling period refers to the average total duration of a single cycle of bone remodeling at any point on a bone surface.

The purpose of remodeling is to regulate calcium homeostasis and repair micro-damage from everyday stress, as well as to shape the skeleton during growth. Repeated stress, such as weight-bearing exercise or bone healing, results in the bone thickening at the points of maximum stress (Wolff’s law).



Osteoclasts and Osteoblasts: Bone tissue is removed by osteoclasts, and then new bone tissue is formed by osteoblasts. Both processes utilize cytokine (TGF-β, IGF) signalling.

This page titled 30.2A: Stages of Bone Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.2B: Exercise and Bone Tissue

Bones adapt to the muscle force loads placed on them, becoming thicker and stronger under stress and use and weaker and thinner when unused.

Learning Objectives

- Distinguish among the responses of bone to activity and hormones

Key Points

- Bone mass is lost if unused because it is metabolically costly to maintain it.
- Gender differences in sex hormones contribute to larger, stronger bones in men since testosterone stimulates muscle mass, which increases bone density.
- As a result of declines in estrogen, aging women suffer from decreased responsiveness to exercise and therefore have difficulty maintaining skeletal strength.
- To maintain skeletal strength, older women need to increase their exercise levels by walking more.

Key Terms

- **Wolff's law**: Bone in a healthy person or animal will adapt to the loads under which it is placed.
- **skeletal strength**: Determined by the relationship of trabecular bone to cortical bone.
- **muscle forces**: The result of increased muscle mass, producing increases in bone dimension and strength.

EXAMPLES

Although we often think of the elderly as feeble and weak, regular exercise can fight osteoporosis and maintain strength and flexibility. This is demonstrated by Johanna Quaas, an 86-year-old gymnast who can still perform an amazing routine on the parallel bars.




NASA Shuttle Astronaut: Astronauts who spend a long time in space will often return to earth with weaker bones, since gravity hasn't been exerting a load. Their bodies have reabsorbed much of the mineral that was previously in their bones.

According to Wolff's law, bone in a healthy person or animal will adapt to the load under which it is placed. If loading on a particular bone increases, the bone will remodel itself to provide the strength needed for resistance. The internal architecture of the trabeculae undergoes adaptive changes, followed by secondary changes to the external cortical portion of the bone, perhaps becoming thicker as a result. The opposite is true as well. If the load on a bone decreases, the bone will become weaker due to turnover. It is less metabolically costly to maintain and there is no stimulus for continued remodeling required to maintain bone mass.

Muscle force is a strong determinant of bone structure, particularly during growth and development. The gender divergence in the bone-muscle relationship becomes strongly evident during adolescence. In females, growth is characterized by increased estrogen levels and increased mass and strength of bone relative to that of muscle. In men, increases in testosterone fuel large increases in muscle, resulting in muscle force that coincides with substantial growth in bone dimensions and strength.

In adulthood, significant age-related losses are observed for both bone and muscle tissues. A large decrease in estrogen levels in women appears to diminish the skeleton's responsiveness to exercise more than in men. In contrast, the aging of the muscle-bone axis in men is a function of age-related declines in both hormones. In addition to the well-known age-related changes in the mechanical loading of bone by muscle, newer studies appear to provide evidence of age and gender-related variations in molecular signaling between bone and muscle that are independent of purely mechanical interactions. In summary, gender differences in acquisition and age-related loss in bone and muscle tissues may be important for developing gender-specific strategies for ways to reduce bone loss with exercise.



Tim Henman performs a backhand volley at the Wimbledon tournament in 2004.: The racquet-holding arm bones of tennis players become much stronger than those of the other arm. Their bodies have strengthened the bones in their racquet-holding arm since they are routinely placed under higher than normal stress.

Simple aerobic exercises like walking, jogging, and running could provide an important role in maintaining and/or increasing bone density in women. Walking is an inexpensive, practical exercise associated with low injury rates and high acceptability among the elderly. For these reasons, walking could be an appropriate approach to prevent osteoporosis and maintain bone mass.

This page titled 30.2B: Exercise and Bone Tissue is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.2C: Bone Tissue and the Effects of Aging

As individuals age, bone resorption can outpace bone replacement, which can lead to osteoporosis and fractures.

Learning Objectives

- Analyze osteoclastic activity

Key Points

- Bone resorption is the process by which osteoclasts break down bone and release its minerals, resulting in a transfer of calcium from bone to blood.
- Bone resorption is highly regulated. It can be stimulated or inhibited by signals from other parts of the body depending on the demand for calcium.
- As people age, the rate of bone resorption far exceeds the rate of bone formation. Thus, bones can weaken, leading to conditions like osteoporosis.
- Osteoporosis can be prevented with lifestyle changes such as proper nutrition and exercise.
- The risks of osteoporosis can also be reduced through the prevention of falls. Fall-prevention advice includes exercise to tone deambulatory muscles, proprioception-improvement exercises, and equilibrium therapies.

Key Terms

- **osteon**: Any of the central canals and surrounding bony layers found in compact bone.
- **periosteum**: A membrane surrounding a bone.

EXAMPLES


Joints undergo substantial wear and tear. The longer one lives, the more a joint is used. Gardeners and flooring installers put a lot of stress on their knees. Use of knee pads can relieve some of that stress.

Bone resorption is the process by which osteoclasts break down bone and release the minerals, resulting in a transfer of calcium from bone to blood.

The Role of Osteoclasts

The osteoclasts are multi-nucleated cells that contain numerous mitochondria and lysosomes. These cells are responsible for the resorption of bone and are generally present on the outer layer of bone, just beneath the periosteum. Attachment of the osteoclast to the osteon begins the process. The osteoclast then induces an infolding of its cell membrane and secretes collagenase and other enzymes important in the resorption process. High levels of calcium, magnesium, phosphate, and collagen products are released into the extracellular fluid as the osteoclasts tunnel into the mineralized bone. Osteoclasts are also prominent in the tissue destruction commonly found in psoriatic arthritis and other rheumatology-related disorders.

Regulation of Bone Tissue



Osteoclast: Osteoclast, displaying many nuclei within its “foamy” cytoplasm above a bone’s surface.

Bone resorption is highly constructible, stimulated or inhibited by signals from other parts of the body depending on the demand for calcium.

Calcium-sensing membrane receptors in the parathyroid gland monitor calcium levels in the extracellular fluid. Low levels of calcium stimulate the release of parathyroid hormone (PTH) from chief cells of the parathyroid gland. In addition to its effects on the kidney and the intestine, PTH also increases the number and activity of osteoclasts to release calcium from bone, thus stimulating bone resorption. High levels of calcium in the blood, on the other hand, lead to decreased PTH release from the parathyroid gland. This decreases the number and activity of osteoclasts, resulting in less bone resorption.

Aging

As people get older, the rate of resorption tends to exceed the rate of replacement, leading to conditions like osteoporosis. Bone resorption can also be the result of disuse and the lack of stimulus for bone maintenance. For instance, astronauts undergo a certain amount of bone resorption due to the lack of gravity providing the proper stimulus for bone maintenance. In addition, certain medical conditions such as hormone imbalances can cause bone resorption to increase, leading to increased susceptibility to fractures.

Osteoporosis risks can be reduced with lifestyle changes and sometimes medication. Lifestyle change includes diet, exercise, and fall-prevention measures. Medication includes calcium, vitamin D, and bisphosphonates. Fall-prevention advice includes exercise to tone deambulatory muscles, proprioception-improvement exercises, and equilibrium therapies. With its anabolic effect, exercise may simultaneously stop or reverse osteoporosis, a component of frailty syndrome.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Bone remodeling period. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone_remodeling_period**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- epiphyses. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/epiphyses**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diaphysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diaphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bonemetabolism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Bonemetabolism.svg**. **License**: ***CC BY-***

SA: Attribution-ShareAlike

- The Bone-Muscle Relationship in Men and Women. **Provided by**: Hindawi Publishing Company. **Located at**: http://www.hindawi.com/journals/josteo/2011/702735/. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Aerobic Exercise and Whole-Body Vibration in Offsetting Bone Loss in Older Adults. **Provided by**: Hindawi Publishing Company. **Located at**: http://www.hindawi.com/journals/jar/2011/379674/. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wolff's law. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wolff's_law. **License**: ***CC BY-SA: Attribution-ShareAlike***
- skeletal strength. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/skeletal%20strength. **License**: ***CC BY-SA: Attribution-ShareAlike***
- muscle forces. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/muscle%20forces. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Wolff's law. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Wolff's%20law. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bonemetabolism. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bonemetabolism.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ledon_2004.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...verage_can.jpg. **License**: ***Public Domain: No Known Copyright***
- Bone resorption. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone_resorption. **License**: ***CC BY-SA: Attribution-ShareAlike***
- WikiArc/ The Wiki Archaeological Information Resource - Joint Disease and Bone Growth. **Provided by**: Wikiarc. **Located at**: http://www.wikiarc.org/jointdisease. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Osteoporosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Osteoporosis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone resorption. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Bone_resorption. **License**: ***CC BY-SA: Attribution-ShareAlike***
- periosteum. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/periosteum. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteon. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/osteon. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bonemetabolism. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Bonemetabolism.svg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...ledon_2004.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...verage_can.jpg. **License**: ***Public Domain: No Known Copyright***
- Osteoclast. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Osteoclast.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 30.2C: Bone Tissue and the Effects of Aging is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.3: DEVELOPMENT OF THE SKELETON

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

30.3A: Development of the Skeleton

30.3B: Aging and the Skeleton

This page titled 30.3: Development of the Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.3A: Development of the Skeleton

Early in gestation, a fetus has a cartilaginous skeleton that becomes bone in the gradual process of endochondral ossification.

Learning Objectives

- Compare and contrast osteogenesis by intramembranous and endochondral ossification


Key Points

- At birth, a newborn baby has over 300 bones, while on average an adult human has 206 bones.
- Ossification or osteogenesis is the process of generating new bone material by precursor cells called osteoblasts.
- Two processes result in the formation of normal, healthy bone tissue: intramembranous ossification and endochondral ossification.
- Intramembranous ossification is the direct laying of bone into the primitive connective tissue (mesenchyme). The membrane that occupies the place of the future bone resembles connective tissue and ultimately forms the periosteum. It is composed of fibers and granular cells in a matrix.
- Endochondral ossification involves cartilage which is then mineralized.
- Bones are dynamic structures that respond to applied force over time. With chronic weight bearing and the forces from daily use and movement, bone mass can be maintained or increased.

Key Terms

- **intramembranous ossification**: Production of bone tissue during fetal development without a cartilage template. The membrane that occupies the place of the future bone resembles connective tissue and ultimately forms the periosteum, or outer bone layer.
- **trabecula**: A small, mineralized spicule that forms a network in spongy bone.
- **mesenchyme**: Undifferentiated cells of the early embryo that develop into the tissues of the lymphatic and circulatory systems, as well as connective tissues throughout the body such as bone and cartilage.
- **endochondral ossification**: The process by which bone tissue is created using a cartilage template during fetal development.

Early in gestation, a fetus has a cartilaginous skeleton from which the long bones and most other bones gradually form throughout development and for years after birth in a process called endochondral ossification. Ossification or osteogenesis is the process of laying new bone material by cells called osteoblasts. Two processes result in the formation of normal, healthy bone tissue. Intramembranous ossification is the direct laying of bone into the primitive connective tissue (mesenchyme). This is how the flat bones of the skull and the clavicles are formed. Endochondral ossification involves cartilage as a precursor. Ossification of the mandible occurs in the fibrous membrane covering the outer surfaces of Meckel's cartilages.



Endochondral Ossification: The development of the primary and secondary ossification centers

Intramembranous Ossification

In bones developed in membrane, no cartilaginous mold precedes the appearance of the bony tissue. The membrane that occupies the place of the future bone resembles connective tissue and ultimately forms the periosteum; it is composed of fibers and granular cells in a matrix. The peripheral portion is more fibrous, while in the interior the cells or osteoblasts predominate. The whole tissue is richly supplied with blood vessels. As the osteogenetic fibers grow out to the periphery, they continue to calcify and give rise to fresh bone spicules. Thus, a network of bone is formed, the meshes of which contain the blood vessels and a delicate connective tissue crowded with osteoblasts. Osteoblasts add layers of fresh bone to the trabeculae. These layers subsequently encroach upon the mesh, under the periosteum, and around the larger vascular channels that become the Haversian canals, thickening and developing the bones.

Endochondral Ossification

The perichondrium becomes the periosteum, which contains a layer of undifferentiated cells (osteoprogenitor cells) that later become osteoblasts. These osteoblasts secrete osteoid against the shaft of the cartilage model that serves as support for the new bone. Chondrocytes in the primary center of ossification begin to grow (hypertrophy). They stop secreting collagen and other proteoglycans and begin secreting alkaline phosphatase, an enzyme essential for mineral deposition. Then calcification of the matrix occurs and the hypertrophic chondrocytes begin to die, creating cavities within the bone. The hypertrophic chondrocytes (before apoptosis) secrete vascular endothelial cell growth factor, which induces the sprouting of blood vessels from the perichondrium. Blood vessels forming the periosteal bud invade the cavity left by the chondrocytes and branch in opposite directions along the length of the shaft. The blood vessels carry hemopoietic cells, osteoprogenitor cells, and other cells inside the cavity. The hemopoietic cells will later form the bone marrow. Osteoblasts, differentiated from the osteoprogenitor cells that entered the cavity via the periosteal bud, use the calcified matrix as a scaffold and begin to secrete osteoid, which forms the bone trabecula. Osteoclasts, formed from macrophages, break down spongy bone to form the medullary (bone marrow) cavity.

This page titled 30.3A: Development of the Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.3B: Aging and the Skeleton

In osteoporosis, bone mineral density (BMD) is reduced and the integrity of bone proteins is altered, increasing the risk of fracture.

Learning Objectives

- Construct a diagnosis and treatment plan for osteoporosis

Key Points

- The risk of osteoporosis can be reduced with lifestyle changes and sometimes medication, both aimed at restoring the amounts and types of necessary bone-associated proteins.
- Osteoporosis itself has no symptoms; its main consequence is the increased risk of bone fractures.
- Treatments for osteoporosis include changes in diet and exercise, increased consumption of calcium and vitamin D, and prescriptions for bisphosphonates.

Key Terms

- **bisphosphonates**: A class of drugs that prevents the loss of bone mass, used to treat osteoporosis and similar diseases. They are the most commonly prescribed drugs used to treat osteoporosis.
- **bone mineral density**: Abbreviated BMD, the amount of mineral matter per square centimeter of bone. BMD is used in medicine as an indirect indicator of osteoporosis and fracture risk.
- **osteoporosis**: A disease in which the bones become extremely porous and are subject to fracture most common in post-menopausal women.

Osteoporosis is a bone disease that leads to an increased risk of fracture. In osteoporosis, the bone mineral density (BMD) is reduced, bone microarchitecture deteriorates, and the amount and variety of proteins in bone is altered.

Types of Osteoporosis



(a) Osteoporotic Fractures and (b) Woman with Osteoporosis: Image shows compression fractures of the lumbar spine.

Hyperkyphosis, colloquially called a “dowager’s hump,” refers to an anteriorly concave curvature of the thoracic region of the spine.

The disease may be classified as primary type one, primary type two, or secondary. The form of osteoporosis most common in women after menopause is referred to as primary type one or postmenopausal osteoporosis.

Primary type two osteoporosis or senile osteoporosis occurs after age 75 and is seen in both females and males at a ratio of two to one. Finally, secondary osteoporosis may arise at any age and affect men and women equally. This form of osteoporosis results from chronic illness or prolonged use of medications such as glucocorticoids. Osteoporosis affects 55% of Americans aged 50 and above. Of these, approximately 80% are women.

Symptoms

Osteoporosis itself has no symptoms; its main consequence is the increased risk of fractures caused by weakened bones. Osteoporotic fractures, called fragility fractures, occur in situations where healthy people would not normally break a bone. These often affect the vertebral column, rib, hip, and wrist. The underlying mechanism in all cases of osteoporosis is an imbalance between bone resorption and bone formation.

Physical deformation can occur secondary to compromised bone densities. One example is hyperkyphosis, colloquially called a “dowager’s hump,” which refers to an anteriorly concave curvature of the thoracic region of the spine.

Treatment

Osteoporosis risks can be reduced with lifestyle changes and sometimes medication. Lifestyle changes include diet, exercise, and fall prevention. Exercises to tone deambulatory muscles and improve proprioception are recommended to decrease risk for falls. In addition, the anabolic effect of exercise can sometimes prevent progression of osteoporosis. Medication includes calcium, vitamin D, and bisphosphonates.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Embryogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skeleta...Embryogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- II. Osteology. **Provided by**: Saylor. **Located at**: **http://www.saylor.org/site/wp-conten...h2with6c6d.pdf**. **License**: ***Public Domain: No Known Copyright***
- Ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- endochondral ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/endocho...20ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- trabecula. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/trabecula**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mesenchyme. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mesenchyme**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- intramembranous ossification. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/intrame...20ossification**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone Growth. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...one_growth.png**. **License**: ***Public Domain: No Known Copyright***
- Osteoporosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Osteoporosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- bone mineral density. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bone%20mineral%20density**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- bisphosphonates. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/bisphosphonates**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteoporosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/osteoporosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone Growth. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...one_growth.png**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...l_fracture.jpg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...steoCutout.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 30.3B: Aging and the Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.4: JOINT DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

30.4A: Development of Joints
30.4B: Joints and Aging
30.4C: Arthroplasty

This page titled 30.4: Joint Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.4A: Development of Joints

Movable joints are formed at the intersection of bones and are covered by cartilage, which allows them to move freely.

Learning Objectives

- Explain the development of joints

Key Points


- Cartilage is a flexible material found between joints, initially formed by chondrification (also known as chondrogenesis).
- Following the initial chondrification, cartilage growth consists mostly of the development of immature cartilage to a more mature state.
- Remodeling of cartilage is predominantly affected by changes to the collagen matrix, which responds to tensile and compressive forces.
- Cartilage growth generally refers to collagen matrix deposition, but can include both growth and remodeling of the extracellular matrix.
- The pressure epiphyses, which take the weight of the long bones, include the head of the femur at the hip joint and the head of the humerus at the shoulder joint.

Key Terms

- **Chondrogenesis**: The process by which cartilage is developed.
- **cartilage**: A type of dense, non-vascular connective tissue, usually found at the end of joints, the rib cage, the ear, the nose, in the throat and between intervertebral disks.
- **Pressure epiphysis**: The region of the long bone that forms the joint.

EXAMPLES

Knowledge of joint growth and development is useful in forensic contexts. For instance, if a skeleton is found and the epiphyses are missing or unfused to the diaphyses, then the examiner knows that the skeleton is that of a juvenile or young adult: if female, probably under age 18; if male, probably under age 25. Other clues, such as dental eruption, can refine the age estimate.




Epiphyseal Plate: Image shows the location of the epiphyseal plates (or lines) and the articular surfaces of long bones.

Articular cartilage is a flexible material found between bones at movable joints. Chondrification (also known as chondrogenesis) is the process by which cartilage is formed from condensed mesenchyme tissue, which differentiates into chondroblasts and begins secreting the molecules that form the extracellular matrix. Following the initial chondrification that occurs during embryogenesis, cartilage growth consists mostly of the development of immature cartilage to a more mature state.

Cell division within cartilage occurs very slowly, so growth is not based on a mass increase of the cartilage itself. Articular cartilage function is dependent on the molecular composition of its extracellular matrix (ECM), which consists mainly of proteoglycans and collagens. Articular cartilage is maintained by embedded chondrocytes that comprise only 1% of the cartilage volume. Remodeling of cartilage is predominantly affected by changes to the collagen matrix, which responds to tensile and compressive forces. Cartilage growth generally refers to matrix deposition, but can include both growth and remodeling of the extracellular matrix.

The epiphysis is the rounded end of a long bone, at its joint with adjacent bone(s). Between the epiphysis and diaphysis (the long midsection of the long bone) lies the metaphysis, including the epiphyseal plate (growth plate). At the joint, the epiphysis is covered with articular cartilage; below that covering is a zone similar to the epiphyseal plate, known as subchondral bone. The region of the long bone that forms the joint is called pressure epiphysis. For example, the head of the femur (which is a part of the hip joint complex) is a pressure epiphyses. These epiphyses assist in transmitting the weight of the human body and are the regions of the bone that are under pressure during movement or locomotione. Another example of pressure epiphysis is the head of humorus, part of the shoulder complex.



Hyaline Cartilage: Histological image of the hyaline cartilage which coats the articular surfaces of joints.

This page titled 30.4A: Development of Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.4B: Joints and Aging

Aging is a common cause of chronic joint pain, as wear and tear from years of use results in inflammation and osteoarthritis.

Learning Objectives

- Describe the effects of aging on joints

Key Points

- As a person ages, the body's ability to regenerate and repair damage to the joints diminishes, resulting in chronic inflammation. This is one of the most common causes of joint pain.
- Osteoarthritis, the degenerative inflammation of the joints, is the leading cause of disability in people older than age 55. Osteoarthritis causes the joint's protective cartilage to erode over time, exposing the bone underneath.
- Joint injuries become more common with age. Contributing factors include decreased balance and motor control, which can lead to falls, and degeneration of the bone from osteoporosis, which can result in greater damage from small injuries.
- Heating pads, braces, and physical therapy can fully or partially ease symptoms. Prescription pain medications can also be used to alleviate joint pain. In cases of severe osteoarthritis, surgical joint replacement is often recommended.

Key Terms

- **osteoporosis**: A disease in which the bones become extremely porous and are subject to fracture, most common among postmenopausal women.
- **fibromyalgia**: Chronic pain, stiffness, and tenderness of the muscles, tendons, and joints.
- **osteoarthritis**: A form of arthritis affecting mainly older people, caused by chronic degeneration of the cartilage and synovial membrane of the joints leading to pain and stiffness.

EXAMPLES

Analysis of skeletal material from archaeological sites has shown that an individual's primary activities can be deduced from development of osteoarthritis. For instance, osteoarthritis of the toes, knees, and hips of female skeletons from the site of Abu Hureyra is clear evidence that they spent most of their days grinding flour on stone querns while kneeling on the ground.

Joint pain can be caused by a variety of health problems, from arthritis to fibromyalgia. Many temporary conditions, such as a virus or the flu, can also cause joint pain. Joint pain occurs in all age groups, but becomes more common with age as wear and tear on the joints from years of use results in inflammation and osteoarthritis. Older adults are also more prone to injuries that can cause joint pain.

Inflammation



Osteoarthritis: X-ray image of osteoarthritis of the big toe. The red box shows the normal joint.

One of the most common causes of joint pain, inflammation is the body's response to pain, infection, and other threats. As the joints become inflamed, their regeneration and repair process slows, making them weaker. As a person ages, the body's ability to regenerate and repair damage to the joints diminishes, resulting in chronic inflammation.

Osteoarthritis

Osteoarthritis, the degenerative inflammation of the joints, is the leading cause of disability in people older than age 55. Osteoarthritis is typically caused by mechanical stress on the joints, resulting in chronic inflammation leading to the loss of cartilage. With this condition, the protective cartilage of the joint erodes over time, exposing the bone underneath. The bones of the joint then rub together during movement, causing pain and stiffness. As a result of decreased movement secondary to pain, regional muscles may atrophy and ligaments become more lax.

Injuries

Sprains, bruising, broken bones, and many other temporary conditions can result in joint pain. Many of these injuries may not be to the joints themselves, but to their surrounding cartilage and ligaments. Joint injuries become more common with age as decreased balance and motor control lead to falls. In addition, bone degeneration caused by osteoporosis can result in greater damage from small injuries.

Treatment Options

Many remedies for joint pain are available. Heating pads, braces, and physical therapy can often minimize symptoms. Prescription pain medications can also be used to alleviate joint pain. In cases of severe osteoarthritis, surgical joint replacement is often recommended.

This page titled 30.4B: Joints and Aging is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.4C: Arthroplasty

Arthroplasty is a surgical procedure in which an arthritic or dysfunctional joint surface is replaced or repaired.

Learning Objectives

- Describe the purpose and process of arthroplasty

Key Points

- Arthroplasty can include removing or remodeling damaged surfaces and replacing them with silicone.
- Prosthetic replacements of hip and knee joints have become common arthroplastic surgeries.
- Wear of the bearing surfaces of components is an issue of joint replacement that can lead to damage to surrounding bone and contribute to eventual failure of the implant.

Key Terms

- **arthroplasty**: The surgical repair of a joint, or creation of an artificial joint, such as a hip replacement.

EXAMPLES

Hip fractures from falls are common among the elderly. Partial arthroplasty (replacing the hip socket and femur head and neck) may be done to relieve pain and improve mobility. However, as with all surgeries, complications can occur that for the elderly may prevent life-threatening challenges. Special care must be made when deciding to proceed with arthroplasty in the elderly patient.

Arthroplasty (literally “surgical repair of joint”) is an orthopedic surgical procedure in which the arthritic or dysfunctional joint surface is replaced and/or the joint is remodeled or realigned.



Planning Arthroplasty: CAD-assisted planning for a total hip arthroplasty.

Common types of arthroplasty include:

- Interpositional arthroplasty with interposition of skin, muscle, or tendon to keep inflammatory surfaces apart.
- Excisional arthroplasty in which the joint surface and bone are removed, leaving scar tissue to fill in the gap.

Other forms of arthroplasty include resectional arthroplasty, resurfacing arthroplasty, mold arthroplasty, cup arthroplasty, and silicone replacement arthroplasty, among others. Osteotomy to restore or modify joint congruity is also an arthroplasty.

For the last 45 years, the most successful and common form of arthroplasty has been the surgical replacement of degenerated joint surfaces with prostheses. For example, a hip joint that is affected by osteoarthritis may be replaced entirely (total hip arthroplasty) with a prosthetic hip. This would involve replacing both the acetabulum (hip socket) and the head and neck of the femur. The purpose of this procedure is to relieve pain, restore range of motion, and improve walking ability, thus leading to the improvement of muscle strength.

One of the main problems with joint replacements is wear of the bearing surfaces of components. This can lead to damage to surrounding bone and contribute to eventual failure of the implant. Use of alternative bearing surfaces has increased in recent years, particularly in younger patients, in an attempt to improve the wear characteristics of joint replacement components. These include ceramics and all-metal implants (as opposed to the original metal-on-plastic). The plastic (actually ultra high-molecular-weight polyethylene) can also be altered in ways that may improve durability.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Epiphysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epiphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cartilage. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cartilage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cartilage. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cartilage**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Pressure epiphysis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Pressure%20epiphysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Chondrogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chondrogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3f/Cartilage_polarised.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Long Bone. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: ***Public Domain: No Known Copyright***
- Joint. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Joint**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Osteoarthritis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Osteoarthritis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fibromyalgia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/fibromyalgia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteoarthritis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/osteoarthritis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- osteoporosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/osteoporosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3f/Cartilage_polarised.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Long Bone. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fc/Roe-h-rigidus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Orthopedic surgery. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Orthopedic_surgery**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Arthroplasty. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arthroplasty**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- arthroplasty. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/arthroplasty**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3f/Cartilage_polarised.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Long Bone. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/9/94/Illu_long_bone.jpg**. **License**: ***Public Domain: No Known Copyright***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fc/Roe-h-rigidus.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/2/22/Planification_computer.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 30.4C: Arthroplasty is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.5: MUSCLE DEVELOPMENT

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

30.5A: Muscle Development

This page titled 30.5: Muscle Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.5A: Muscle Development

Myogenesis is the formation of muscle tissue during embryonic development from stem cells in the mesoderm.

Learning Objectives

- Describe the process of myogenesis, the formation of muscular tissue

Key Points

- Human embryonic stem cells are pluripotent, meaning they differentiate into all cell types, including muscle cells.
- Muscle tissue is formed in the mesoderm layer of the embryo in response to signals from fibroblast growth factor, serum response factor, and calcium.
- In the presence of fibroblast growth factor, myoblasts fuse into multi-nucleated mytotubes, which form the basis of muscle tissue.
- Unused myoblasts dedifferentiate into myosatellite cells, which remain in the muscle fiber until needed to differentiate into new muscle cells when a muscle is damaged or stressed.
- Myocytes are tubular muscle cells or fibers that develop from myoblasts.
- Myocytes are specialized as cardiac, skeletal, or smooth muscle cells.

Key Terms


- **myogenesis**: The formation of muscle tissue during the development of an embryo.
- **mesoderm**: One of the three tissue layers in the embryo of a metazoan animal. Through embryonic development, it produces many internal organs of the adult, including the muscles, spine, and circulatory system.
- **myoblasts**: A type of embryonic stem cell that gives rise to muscle cells.

EXAMPLES

Body builders enhance the natural developmental processes of myoblast muscle fusion and dedifferentiation into myosatellites to dramatically increase myosatellite mass, and thus muscle size and weight.

Embryogenesis is the process by which the embryo is formed and develops, until it becomes a fetus.

Characteristics of Embryoblasts



Skeletal Satellite Muscle Cell: Satellite cells are located between the basement membrane and sarcolemma (cell membrane) of individual muscle fibers. They are able to differentiate and fuse to augment existing muscle fibers and to form new ones. These cells represent the oldest known adult stem cell niche, and are involved in the normal growth of muscle, as well as regeneration following injury or disease.

The cells of the inner cell mass (embryoblast), known as human embryonic stem cells (hESCs), differentiate to form four structures: the amnion, the yolk sac, the allantois, and the embryo itself. Human embryonic stem cells are pluripotent; that is, they can differentiate into any of the cell types present in the adult human, and into any of the intermediate progenitor cell types that eventually turn into the adult cell lines. hESCs are also immortal; they can divide and grow in number indefinitely, without undergoing either differentiation or cellular aging (cellular senescence).


The first differentiation of the hESCs that form the embryo proper, is in three cell types known as the germ layers: the *ectoderm*, the *mesoderm*, and the *endoderm*. The ectoderm eventually forms the skin (including hair and nails), mucous membranes, and nervous system. The mesoderm forms the skeleton and muscles, heart and circulatory system, urinary and reproductive systems, and connective tissues inside the body. The endoderm forms the gastrointestinal tract (stomach and intestines), the respiratory tract, and the endocrine system (liver and endocrine glands).

The Process of Myogenesis

Myogenesis is the formation of muscular tissue, particularly during embryonic development. Muscle fibers form from the fusion of myoblasts into multi-nucleated fibers called myotubes. In early embryonic development, these myoblasts proliferate if enough fibroblast growth factor (FGF) is present. When the FGF runs out, the myoblasts cease division and secrete fibronectin onto their extracellular matrix. The second stage involves the alignment of the myoblasts into the myotubes. The third stage is the actual cell fusion itself. In this stage, calcium ions are critical for development. Myocyte enhance factors (MEFs) promote myogenesis. Serum response factor (SRF) plays a central role during myogenesis, required for the expression of striated alpha-actin genes. Expression of skeletal alpha-actin is also regulated by the androgen receptor, which means steroids can regulate myogenesis.

Characteristics of Myoblasts

A myoblast is a type of embryonic progenitor cell that differentiates to form muscle cells. Skeletal muscle fibers are made when myoblasts fuse together, so muscle fibers have multiple nuclei. The fusion of myoblasts is specific to skeletal muscle (e.g., biceps brachii), not cardiac or smooth muscle.



Mesoderm: The embryonic layer from which muscle tissues develop, including cardiac muscle, skeletal muscles cells, tubule cell of the kidney, red blood cells, and smooth muscle in gut.

Myoblasts that do not form muscle fibers dedifferentiate back into satellite (myosatellite) cells. These cells remain adjacent to a muscle fiber, situated between the sarcolemma and the endomysium (the connective tissue that divides the muscle fascicles into individual fibers). Satellite cells are able to differentiate and fuse to augment existing muscle fibers and form new ones. In undamaged muscle, the majority of satellite cells are quiescent; they neither differentiate nor undergo cell division. In response to mechanical strain, satellite cells become activated and initially proliferate as skeletal myoblasts before undergoing myogenic differentiation.

A myocyte (also known as a muscle cell or muscle fiber) is the type of cell found in muscle tissue. These long, tubular cells arise developmentally from myoblasts to form muscles. There are various specialized forms of myocytes with various properties, including cardiac, skeletal, and smooth muscle cells. Cardiac myocytes generate the electrical impulses that control heart rate and other functions.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Myoblasts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myoblasts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Satellite cells. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Satellite_cells**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Embryogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myoblasts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myoblasts%23Myoblasts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Embryomics. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Embryom...3Embryogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myoblasts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/myoblasts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- mesoderm. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/mesoderm**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- myogenesis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/myogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- University of Michigan. **Provided by**: University of Michigan. **Located at**: **http://umich.edu/education/med/m1/ce...2009/materials**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Mesoderm. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Mesoderm.png**. **License**: ***Public***

Domain: No Known Copyright

This page titled 30.5A: Muscle Development is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.6: DEVELOPMENT OF THE NERVOUS SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

30.6A: Development of the Central Nervous System
30.6B: Development of the Peripheral Nervous System
30.6C: Regeneration of Nerve Fibers
30.6D: Aging and the Nervous System

This page titled 30.6: Development of the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.6A: Development of the Central Nervous System

The central nervous system (CNS) develops from a longitudinal groove on the neural plate that forms the rudimentary nervous system.

Learning Objectives

- Describe the development of the central nervous system

Key Points

- During early development of the vertebrate embryo, a longitudinal groove on the neural plate gradually deepens and the ridges on either side of it (the neural folds) ultimately meet, transforming it into a closed tube, the ectodermal wall of which forms the rudimentary nervous system.
- This tube initially differentiates into three vesicles (pockets): the prosencephalon at the front, the mesencephalon, the rhombencephalon
between the mesencephalon and spinal cord.
- The telencephalon differentiates into, among other things, the striatum, the hippocampus, and the neocortex, and its cavity becomes the first and second ventricles. Diencephalon elaborations include the subthalamus, hypothalamus, thalamus, and epithalamus, and its cavity forms the third ventricle.
- The survival of neurons is regulated by survival factors, called trophic factors.
- Agrin is not a central mediator of CNS synapse formation and there is active interest in identifying signals that mediate CNS synaptogenesis.

Key Terms


- **diencephalon**: The region of the human brain, specifically the forebrain, that includes the thalamus, hypothalamus, epithalamus, prethalamus or subthalamus, and pretectum.
- **metencephalon**: The portion of the embryonic rhombencephalon from which the pons and cerebellum arise.
- **telencephalon**: The anterior part of the forebrain; the endbrain.

The central nervous system (CNS) consists of the brain and the spinal cord. It is the part of the nervous system that integrates the information that it receives from, and coordinates the activity of, all parts of the bodies of bilaterian animals—that is, all multicellular animals except radially symmetric animals such as sponges and jellyfish.

Some classifications of the CNS also include the retina and the cranial nerves. Together with the peripheral nervous system, it has a fundamental role in behavior control. The CNS is contained within the dorsal cavity, with the brain in the cranial cavity and the spinal cord in the spinal cavity. In vertebrates, the brain is protected by the skull while the spinal cord is protected by the vertebrae, and both are enclosed in the meninges.

During early development of the vertebrate embryo, a longitudinal groove on the neural plate gradually deepens and the ridges on either side of it (the neural folds) become elevated and ultimately meet, transforming the groove into a closed tube, the ectodermal wall of which forms the rudiment of the nervous system. This tube initially differentiates into three vesicles (pockets): the prosencephalon at the front, the mesencephalon, and between the mesencephalon and the spinal cord, the rhombencephalon. At six weeks in human embryo development, the prosencephalon divides further into the telencephalon and diencephalon and the rhombencephalon divides into the metencephalon and myelencephalon.

As the vertebrate grows, these vesicles differentiate further still. The telencephalon differentiates into, among other things, the striatum, the hippocampus, and the neocortex, and its cavity becomes the first and second ventricles. Diencephalon elaborations include the subthalamus, hypothalamus, thalamus, and epithalamus, and its cavity forms the third ventricle. The tectum, pretectum, cerebral peduncle, and other structures develop out of the mesencephalon, and its cavity grows into the mesencephalic duct (cerebral aqueduct). The metencephalon becomes, among other things, the pons and the cerebellum; the myelencephalon forms the medulla oblongata; and their cavities develop into the fourth ventricle.



Morphogenisis and the Change in the Shape of Cell: Development of the neural tube in human embryos (Prentiss-Arey). A: An early embryo (Keibel) B: At 2 mm. (Graf Spee) C: At 2 mm. (Mall) D: At 2.7 mm (Kollmann).

This page titled 30.6A: Development of the Central Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.6B: Development of the Peripheral Nervous System

The peripheral nervous system develops from two strips of tissue called the neural crest, running lengthwise above the neural tube.

Learning Objectives

- Describe peripheral nervous system development


Key Points

- The first sign of the nervous system is the appearance of a thin strip of cells along the center of the back, called the neural plate. The inner portion of the neural plate is destined to become the central nervous system (CNS), while the outer portion will become the peripheral nervous system (PNS).
- Neurulation (neural development) progresses with the formation of the neural groove which closes to form the neural tube and neural crest.
- Neural crest cells from the roof plate of the neural tube migrate through the periphery where they differentiate into varied cell types, including pigment cells and the cells of the PNS.

Key Terms

- **neurulation**: The process by which the beginnings of the vertebrate nervous system is formed in embryos.
- **neural plate**: A thick, flat bundle of ectoderm formed in vertebrate embryos after induction by the notochord.
- **neural crest**: A strip of ectodermal material in the early vertebrate embryo inserted between the prospective neural plate and the epidermis.

In vertebrates, the first sign of the nervous system is the appearance of a thin strip of cells along the center of the back, called the neural plate. The inner portion of the neural plate (along the midline) is destined to become the central nervous system (CNS), the outer portion the peripheral nervous system (PNS). As development proceeds, a fold called the neural groove appears along the midline. This fold deepens and then closes up at the top. At this point the future CNS appears as a cylindrical structure called the neural tube, whereas the future PNS appears as two strips of tissue called the neural crest, running lengthwise above the neural tube. The sequence of stages from neural plate to neural tube and neural crest is known as neurulation.



Formation of the Fetal Nervous System: The neural tube will give rise to the central nervous system, while the neural crest will give rise to the peripheral nervous system. This diagram shows the epidermis, neural fold, and convergence.

Neural crest cells are a transient, multipotent, migratory cell population unique to vertebrates that gives rise to a diverse cell lineage including melanocytes, craniofacial cartilage and bone, smooth muscle, peripheral and enteric neurons and glia. After gastrulation, neural crest cells are specified at the border of the neural plate and the non-neural ectoderm. During neurulation, the borders of the neural plate, also known as the neural folds, converge at the dorsal midline to form the neural tube. Subsequently, neural crest cells from the roof plate of the neural tube undergo an epithelial to mesenchymal transition, delaminating from the neuroepithelium and migrating through the periphery where they differentiate into varied cell types, including pigment cells and the cells of the PNS. The emergence of the neural crest was important in vertebrate evolution because many of its structural derivatives are defining features of the vertebrate clade.

This page titled 30.6B: Development of the Peripheral Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.6C: Regeneration of Nerve Fibers

The peripheral nervous system (PNS) is able to repair and regenerate itself, but the central nervous system is incapable of doing so.

Learning Objectives

- Describe nerve fiber regeneration in the PNS


Key Points

- In the PNS, some self-regeneration after injury is possible if the cell body (soma) and portions of the membrane (neurilemma) are left intact.
- When an axon is damaged, the distal segment undergoes Wallerian degeneration, losing its myelin sheath. The proximal segment can either die by apoptosis or undergo the chromatolytic reaction, which is an attempt at repair.
- Injury to the PNS immediately elicits the migration of phagocytes, Schwann cells, and macrophages to the lesion site to clear away debris such as damaged tissue.
- In the case of a severed nerve, the proximal end begins to sprout axons and the presence of growth cones can be detected. Meanwhile, the distal end experiences Wallerian degeneration within hours of the injury; the axons and myelin degenerate, but the endoneurium remains.
- The environment within the CNS, especially following trauma, counteracts the repair of myelin and neurons. Glial scars rapidly form, and the glia actually produce factors that inhibit remyelination and axon repair.

Key Terms

- **neurilemma**: The outer membranous covering of a nerve fiber.
- **microglia**: The glial cells of the immune system; they can function as phagocytes.
- **neuroregeneration**: The regrowth or repair of damaged nervous tissue.
- **Schwann cell**: Glia of the peripheral nervous system, involved in many important aspects of peripheral nerve biology.

Nerve damage can be caused by physical injury or swelling (*e.g.* carpal tunnel syndrome), autoimmune disease (*e.g.* Guillain-Barré syndrome), infection (*e.g.* neuritis), diabetes, or failure of the blood vessels surrounding the nerve. Nerves can be easily damaged in a traumatic event, but can regenerate if the soma and a small portion of the neurilemma remain. Neuroregeneration refers to the regrowth or repair of nervous tissues, cells, or cell products. Such mechanisms may include generation of new neurons, glia, axons, myelin, or synapses. Neuroregeneration differs between the peripheral nervous system (PNS) and the central nervous system (CNS) by the functional mechanisms and especially by the extent and speed. When an axon is damaged, the distal segment undergoes Wallerian degeneration, losing its myelin sheath. The proximal segment can either die by apoptosis or undergo the chromatolytic reaction, an attempt at repair. In the CNS, synaptic stripping occurs as glia foot processes invade the dead synapse.



Nerve injury: Micrograph of a nerve with a decrease in myelinated nerve fibres (pink) and an abnormal increase in fibrous tissue (yellow), as may be seen in nerve injuries.

Nervous system injuries affect more than 90,000 people every year, 10,000 of which are spinal cord injuries. As a result, the field of nerve regeneration and repair, a subfield of neural tissue engineering dedicated to the discovery of new ways to recover nerve functionality after injury, is growing rapidly. The nervous system is divided into two parts: the CNS, which consists of the brain and spinal cord, and the PNS, which consists of cranial and spinal nerves along with their associated ganglia. While the PNS has an intrinsic ability for repair and regeneration, the CNS is for the most part incapable of self-repair and regeneration. There is no current treatment to recover human nerve function after injury to the central nervous system. In addition, multiple attempts at nerve re-growth across the PNS-CNS transition have not been successful. Although the PNS has the capability for regeneration, much research still needs to be done to optimize the environment for maximum regrowth potential. Nerve regeneration is part of the pathogenesis of many diseases, including multiple sclerosis.

PNS Injury

Neuroregeneration in the PNS occurs to a significant degree. Axonal sprouts form at the proximal stump and grow until they enter the distal stump. The growth of the sprouts are governed by chemotactic factors secreted from Schwann cells (neurolemmocytes). Injury to the PNS immediately elicits the migration of phagocytes, Schwann cells, and macrophages to the lesion site to clear away debris such as damaged tissue. When a nerve axon is severed, the end still attached to the cell body is labeled the proximal segment, while the other end is called the distal segment. After injury, the proximal end swells and experiences some retrograde degeneration, but once the debris is cleared, it begins to sprout axons and the presence of growth cones can be detected. The proximal axons are able to regrow as long as the cell body is intact and they have made contact with the Schwann cells in the endoneurial channel. Human axon growth rates can reach 2 mm/day in small nerves and 5 mm/day in large nerves. The distal segment, however, experiences Wallerian degeneration within hours of the injury; the axons and myelin degenerate, but the endoneurium remains. In the later stages of regeneration, the remaining endoneurial tube directs axon growth back to the correct targets. During Wallerian degeneration, Schwann cells grow in ordered columns along the endoneurial tube. This creates a band of Büngner (boB) that protects and preserves the endoneurial channel. Also, macrophages and Schwann cells release neurotrophic factors that enhance regrowth.

CNS Injury

Unlike PNS injury, injury to the CNS is not followed by extensive regeneration. It is limited by the inhibitory influences of the glial and extracellular environment. The hostile, non-permissible growth environment is in part created by the migration of myelin-associated inhibitors, astrocytes, oligodendrocytes, oligodendrocyte precursors, and microglia. The environment within the CNS, especially following trauma, counteracts the repair of myelin and neurons. Glial scars rapidly form and the glia actually produce factors that inhibit remyelination and axon repair. The axons themselves also lose the potential for growth with age. Slower degeneration of the distal segment than that which occurs in the peripheral nervous system also contributes to the inhibitory environment; inhibitory myelin and axonal debris are not cleared away as quickly. All these factors contribute to the formation of what is known as a glial scar, which axons cannot grow across.

This page titled 30.6C: Regeneration of Nerve Fibers is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.6D: Aging and the Nervous System

One of the effects of aging on the nervous system is the loss of neurons in the cerebral cortex.

Learning Objectives

- Describe the effects of aging on the nervous system

Key Points

- As the brain ages, neurons in the central nervous system (CNS), which do not typically regenerate, are lost. This reduces the brain's capacity to send and receive nerve impulses and slows information processing.
- Dementia is the age-related progressive decline in cognitive function. This may affect memory, attention, language, and problem-solving.
- Alzheimer's disease is a neurodegenerative disease characterized by progressive cognitive deterioration, declining activities of daily living, and neuropsychiatric symptoms or behavioral changes. It is the most common cause of dementia.


Key Terms

- **cerebral cortex**: The gray, folded, outermost layer of the cerebrum that is responsible for higher brain processes such as sensation, voluntary muscle movement, thought, reasoning, and memory.
- **dementia**: A progressive decline in cognitive function due to brain damage or disease. Areas particularly affected include memory, attention, judgement, language, and problem solving.

Neuron loss is an effect of aging on the nervous system. By the age of 30, the brain begins to lose thousands of neurons each day, causing a decreased capacity to send nerve impulses to and from the brain and slowing information processing. In addition, voluntary motor movements slow down, reflex time increases, and conduction velocity decreases.

Dementia (from Latin de- "apart, away" + mens (genitive mentis) "mind") is the progressive decline in cognitive function due to damage or disease in the brain beyond what is expected from normal aging. Dementia most commonly affects memory, attention, language, and problem solving, although those in the later stages in this condition may be disoriented in time, place, and person. Symptoms of dementia can be classified as either reversible or irreversible depending upon the etiology of the disease. Less than 10% of all dementia cases are reversible.

Alzheimer's disease (AD) is a neurodegenerative disease characterized by progressive cognitive deterioration, declining activities of daily living, and neuropsychiatric symptoms or behavioral changes. It is the most common cause of dementia. The most striking early symptom is short term memory loss (amnesia), which usually manifests as minor forgetfulness that becomes steadily more pronounced with illness progression. Older memories are often preserved. As the disorder progresses, cognitive (intellectual) impairment extends to the domains of language (aphasia), skilled movements (apraxia), recognition (agnosia), and functions (such as decision-making and planning) closely related to the frontal and temporal lobes of the brain as they become disconnected from the limbic system, reflecting extension of the underlying pathological process.



Effects of Alzheimer's Disease: Comparison of a normal aged brain (left) and the brain of a person with Alzheimer's (right). Differential characteristics are pointed out in the hippocampus, cerebral cortex, and ventricles.

The symptoms of Alzheimer's are triggered by neuron loss and an inflammatory response to the deposition of amyloid plaques and neurofibrillary tangles. Alzheimer's has a genetic component, and autosomal dominant mutations in three different genes have been identified that account for a small number of cases of familial, early-onset AD. For late onset AD (LOAD), only one susceptibility gene has so far been identified: the epsilon 4 allele of the apolipoprotein E gene. Age of onset itself has a heritability of around 50%.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Central nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Central_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Central nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Central_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neural_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Central nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Central_nervous_system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neural_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Neural development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Neural_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diencephalon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- metencephalon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/metencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- telencephalon. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/telencephalon**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Development of the neural tube. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Development_of_the_neural_tube.png**. **License**: ***Public Domain: No Known Copyright***
- Nervous system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nervous_system%23Development**.

License: *CC BY-SA: Attribution-ShareAlike*
- Neural crest. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neural_crest. **License**: *CC BY-SA: Attribution-ShareAlike*
- neurulation. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurulation. **License**: *CC BY-SA: Attribution-ShareAlike*
- neural crest. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neural_crest. **License**: *CC BY-SA: Attribution-ShareAlike*
- neural plate. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neural_plate. **License**: *CC BY-SA: Attribution-ShareAlike*
- Development of the neural tube. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Development_of_the_neural_tube.png. **License**: *Public Domain: No Known Copyright*
- Neural Crest. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Neural_Crest.png. **License**: *Public Domain: No Known Copyright*
- Neuroregeneration. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Neuroregeneration. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nerve fiber. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve_fiber%23Regeneration_of_peripheral_nerve_fibers. **License**: *CC BY-SA: Attribution-ShareAlike*
- Nerve. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Nerve. **License**: *CC BY-SA: Attribution-ShareAlike*
- neurilemma. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neurilemma. **License**: *CC BY-SA: Attribution-ShareAlike*
- neuroregeneration. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/neuroregeneration. **License**: *CC BY-SA: Attribution-ShareAlike*
- microglia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/microglia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Schwann cell. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/Schwann_cell. **License**: *CC BY-SA: Attribution-ShareAlike*
- Development of the neural tube. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Development_of_the_neural_tube.png. **License**: *Public Domain: No Known Copyright*
- Neural Crest. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Neural_Crest.png. **License**: *Public Domain: No Known Copyright*
- Endoneurial fibrosis - very high mag - cropped. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Endoneurial_fibrosis_-_very_high_mag_-_cropped.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Human Physiology/Development: birth through death. **Provided by**: Wikibooks. **Located at**: en.wikibooks.org/wiki/Human_Physiology/Development:_birth_through_death%23Nervous_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- dementia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/dementia. **License**: *CC BY-SA: Attribution-ShareAlike*
- cerebral cortex. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cerebral_cortex. **License**: *CC BY-SA: Attribution-ShareAlike*
- Development of the neural tube. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Development_of_the_neural_tube.png. **License**: *Public Domain: No Known Copyright*
- Neural Crest. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Neural_Crest.png. **License**: *Public Domain: No Known Copyright*
- Endoneurial fibrosis - very high mag - cropped. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Endoneurial_fibrosis_-_very_high_mag_-_cropped.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Alzheimer's disease brain comparison. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Alzheimer's_disease_brain_comparison.jpg. **License**: *Public Domain: No Known Copyright*

This page titled 30.6D: Aging and the Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.7: DEVELOPMENT OF THE SPECIAL SENSES

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

30.47C: Development of Hearing and Balance
30.7A: Taste and Smell at Birth and in Old Age
30.7B: Development of Vision

This page titled 30.7: Development of the Special Senses is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.47C: Development of Hearing and Balance

Critical periods have been identified for the development of the hearing and vestibular system.

Learning Objectives

- Describe the development of the inner ear for hearing and balance


Key Points

- The human inner ear develops during week four of embryonic development from the auditory placode, a thickening of the ectoderm that gives rise to the bipolar neurons of the cochlear and vestibular ganglions.
- The auditory vesicle gives rise to the utricular and saccular components of the membranous labyrinth.
- Beginning in the fifth week of development, the auditory vesicle also gives rise to the cochlear duct, which contains the spiral organ of Corti and the endolymph that accumulates in the membranous labyrinth.
- In our vestibular system, neurons are undeveloped at neuronal birth and mature during the critical period of the first two to three postnatal weeks.
- Many studies have supported a correlation between the type of auditory stimuli present in the early postnatal environment and the topographical and structural development of the auditory system.

Key Terms

- **inner ear**: The portion of the ear located within the temporal bone that includes the semicircular canals, vestibule, and cochlea. It is responsible for hearing and balance.
- **organ of Corti**: Found only in mammals, this inner ear organ contains auditory sensory cells, or “hair cells.”
- **vestibular system**: The sensory system that provides the leading contribution about movement and sense of balance in most mammals.

The human inner ear develops during week four of embryonic development from the auditory placode, a thickening of the ectoderm that gives rise to the bipolar neurons of the cochlear and vestibular ganglions. As the auditory placode invaginates towards the embryonic mesoderm, it forms the auditory vesicle or otocysts.



Formation of the Mammalian Ear: The cochlea and vestibule viewed from above.

The auditory vesicle gives rise to the utricular and saccular components of the membranous labyrinth. They contain the sensory hair cells and otoliths of the macula of utricle and of the saccule, respectively, which respond to linear acceleration and the force of gravity. The utricular division of the auditory vesicle also responds to angular acceleration, as do the endolymphatic sac and duct that connect the saccule and utricle.

Beginning in the fifth week of development, the auditory vesicle also gives rise to the cochlear duct, which contains the spiral organ of Corti and the endolymph that accumulates in the membranous labyrinth. The vestibular wall will separate the cochlear duct from the perilymphatic scala vestibuli, a cavity inside the cochlea. The basilar membrane separates the cochlear duct from the scala tympani, a cavity within the cochlear labyrinth. The lateral wall of the cochlear duct is formed by the spiral ligament and the stria vascularis, which produces the endolymph. The hair cells develop from the lateral and medial ridges of the cochlear duct, which together with the tectorial membrane make up the organ of Corti.

Critical periods have been identified for the development of hearing and the vestibular system. In our vestibular system, neurons are undeveloped at neuronal birth and mature during the critical period of the first two to three postnatal weeks. Disruption of maturation during this period can cause changes in normal balance and movement through space. Animals with abnormal vestibular development tend to have irregular motor skills. Studies have consistently shown that animals with genetic vestibular deficiencies during this critical period have altered vestibular phenotypes, most likely as a result of lack insufficient input from the semicircular canals and dopaminergic abnormalities.

Moreover, exposure to abnormal vestibular stimuli during the critical period is associated with irregular motor development. Many studies have supported a correlation between the type of auditory stimuli present in the early postnatal environment and the topographical and structural development of the auditory system. First reports on critical periods came from studies of deaf children and animals that received a cochlear implant to restore hearing. Approximately at the same time, other studies demonstrated that the adaptation to the cochlear implant is subject to an early developmental critical (sensitive) period. These corresponding data demonstrated both on children and in animals that this sensitive period has consequences for medical therapy of hearing loss.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Olfactory memory. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Olfactory_memory**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Taste bud. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Taste_bud**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory memory. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Olfactory_memory**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Developmental psychology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Developmental_psychology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory memory. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Olfactory_memory**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Loss of smell. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Loss_of_smell**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Loss of taste. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Loss_of_taste**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Developmental psychology. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Developmental_psychology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- taste bud. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/taste%20bud**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anosmia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anosmia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- olfactory bulb. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/olfactory%20bulb**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Taste bud. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Taste_bud.svg**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Olfactory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ol...ory_system.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Eye development. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Eye_development**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- optic sulcus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/optic%20sulcus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lens placode. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/lens%20placode**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- optic cup. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/optic%20cup**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Taste bud. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Taste_bud.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ol...ory_system.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray865. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray865.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray460. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray460.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- inner ear. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/inner_ear**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Critical period. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Critica...tibular_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Inner ear. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Inner_ear%23Embryology**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Critical period. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Critica...tibular_System**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- organ of Corti. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/organ%20of%20Corti**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//biology/de...tibular-system**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Taste bud. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Taste_bud.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Olfactory system. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ol...ory_system.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray865. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray865.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray460. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray460.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray923. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray923.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 30.47C: Development of Hearing and Balance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.7A: Taste and Smell at Birth and in Old Age

The senses of taste and smell develop in the intrauterine environment and can deteriorate with age.

Learning Objectives

- Describe the status of taste and smell at birth and in old age

Key Points

- Newborns are born with odor and taste preferences acquired in the womb from the smell and taste of amniotic fluid, influenced by the mother's diet.
- A significant change takes place in the regulation of olfaction just after birth so that odors related to the offspring are no longer aversive, allowing the female to positively respond to her babies.
- Anosmia is a lack of functioning olfaction: an inability to perceive odors.
- Ageusia is the loss of taste function, particularly the inability to detect sweetness, sourness, bitterness, saltiness, and umami (savory taste).

Key Terms

- **olfactory bulb**: The structure of the vertebrate forebrain involved in olfaction, the perception of odors.
- **anosmia**: The inability to perceive odors.
- **taste bud**: Sensory receptors located around the small structures on the upper surface of the tongue, soft palate, upper esophagus, and epiglottis.

EXAMPLES

Failure to detect odors (anosmia) is one reason that older individuals may not enjoy eating, since an inability to smell is related to an inability to taste (ageusia). Caretakers of such individuals must ensure they continue to eat healthy foods.

The senses of taste and smell first develop in neonates and can be diminished by the effects of aging.

Smell

At birth, infants can show expressions of disgust or pleasure when presented with pleasant (honey, milk) or unpleasant (rotten egg) odors and tastes. Newborns have inherent smell and taste preferences acquired in the womb from the smell and taste of amniotic fluid, which is influenced by the mother's diet.


Intrauterine Olfactory Learning

Intrauterine olfactory learning is demonstrated by behavioral evidence that newborns respond positively to the smell of their own amniotic fluid. Infants recognize and react favorably to scents emitted from their own mother's breasts, despite the fact that they also may be attracted to breast odors from unfamiliar nursing females in a different context. The unique scent of the mother (to the infant) is referred to as her olfactory signature. Infants are also able to recognize and respond with familiarity and preference to their mother's underarm scent.

In newborn mammals, the nipple area of the mother is the sole nutritional source, so the maternal olfactory scent becomes associated with food intake. Newborns who do not gain access to the mother's breasts would die shortly after birth, so this olfactory cue helps maintain and establish effective breast feeding. The mother's olfactory signature is experienced with reinforcing stimuli such as food, warmth, and tactile stimulation, enhancing further learning of that cue.

As demonstrated by animals in the wild (apes, for example), offspring are held by the mother immediately after birth without cleaning and continually exposed to the familiar odor of the amniotic fluid (making the transition from the intrauterine to extrauterine environment less overwhelming).

Studies demonstrate that the changes to the olfactory bulb and main olfactory system following birth are extremely important and influential for maternal behavior. Pregnancy and childbirth result in a high state of plasticity of the olfactory system that may facilitate olfactory learning within the mother. A significant change takes place in the regulation of olfaction just after birth so that odors related with the offspring are no longer aversive, allowing the female to positively respond to her babies.




Olfactory System: Human olfactory system. 1: Olfactory bulb 2: Mitral cells 3: Bone 4: Nasal epithelium 5: Glomerulus (olfaction) 6: Olfactory receptor cells

Anosmia and Aging

Older people experience a decline in the sense of smell. Anosmia, a lack of functioning olfaction (inability to perceive odors), may be temporary, but traumatic anosmia can be permanent. Anosmia is due to an inflammation of the nasal mucosa, blockage of nasal passages, or a destruction of one temporal lobe. A common cause of anosmia is old age.

Taste



Taste Receptors in Humans: Structure of the taste bud, including afferent nerve, connective tissue, basal cell, taste receptor cell, lingual epithelium, oral cavity, and taste pore.

Taste buds contain the receptors for taste and are located around the small structures (papillae) on the upper surface of the tongue, soft palate, upper esophagus, and epiglottis. These papillae are involved in detecting the five known elements of taste perception: salty, sour, bitter, sweet, and umami. Via small openings in the tongue epithelium (taste pores), parts of the food dissolved in saliva come into contact with taste receptors (taste buds). The taste receptor cells send information detected by clusters of various receptors and ion channels to the gustatory areas of the brain via the seventh, ninth, and tenth cranial nerves. On average, the human tongue has 2,000–8,000 taste buds.

The average life of a taste bud is 10 days. Ageusia is the loss of taste function, particularly the inability to detect sweetness, sourness, bitterness, saltiness, and umami. It is sometimes confused with anosmia, a loss of the sense of smell. Because the tongue can only indicate texture and differentiate between sweet, sour, bitter, salty, and umami, most of what is perceived as the sense of taste is actually derived from smell. True ageusia is relatively rare compared to hypogeusia (a partial loss of taste) and dysgeusia (a distortion or alteration of taste). Local damage and inflammation that interferes with the taste buds or local nervous system such as that stemming from radiation therapy, glossitis, tobacco use, and denture use also cause ageusia. Other known causes include loss of taste sensitivity from aging causing a difficulty detecting salty or bitter taste, anxiety disorder, cancer, renal failure and liver failure.

This page titled 30.7A: Taste and Smell at Birth and in Old Age is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.7B: Development of Vision

The eye forms from the neural tube, epidermis, and periocular mesenchyme, with sequential inductions of tissue during development.

Learning Objectives

- Describe the development of vision

Key Points


- Organogenesis of the eye is an example of a developmental cascade of inductions.
- Development of the optic vesicles starts in the three-week embryo from a progressively deepening groove in the neural plate called the optic sulcus.
- The optic vesicles come into contact with the epithelium and induce the epidermis to thicken to form the lens placode.
- The periocular mesenchyme migrates in during the formation of the optic cup and is critical for the induction of the retinal pigment epithelium and the optic nerve.

Key Terms

- **optic cup**: During embryonic development of the eye, the outer wall of the bulb of the optic vesicles becomes thickened and invaginated, converting it to a cup consisting of two strata of cells. These strata are continuous with each other at the cup margin, which ultimately overlaps the front of the lens and reaches as far forward as the future aperture of the pupil.
- **optic sulcus**: A progressively-deepening groove in the neural plate from which the optic vesicles develop.
- **lens placode**: A thickened portion of ectoderm which serves as the precursor to the lens.

The eye develops from the neural tube, the epidermis, and the periocular mesenchyme, which receives contributions from both the neural crest and mesoderm lineages. The organogenesis of the eye is an example of a developmental cascade of inductions, with three different tissues contributing to its differentiations.

Neural Tube




Chick embryo head with optic vesicle: The eyes make their appearance before the closure of the anterior end of the neural tube. After the closure of the tube they are known as the optic vesicles.

First, an outpocketing of the neural tube occurs, creating optic vesicles. Development of the optic vesicles starts in the three-week embryo from a progressively deepening groove in the neural plate called the optic sulcus. As this expands, the rostral neuropore (the exit of the brain cavity out of the embryo) closes and the optic sulcus and the neural plate becomes the optic vesicle.

Epidermis

The optic vesicles come into contact with the epithelium and induce the epidermis. The epithelium thickens to form the lens placode. The lens differentiates and invaginates until it detaches from the epithelium. The lens then acts as an inducer back to the optic vesicle to transform it into the optic cup and back to the epidermis to transform it into the cornea. The optic cup then delaminates into two layers: the neural retina and the retinal pigment epithelium.

Periocular Mesenchyme



The Optic Stalk and Optic Cup: During embryonic development of the eye, the outer wall of the bulb of the optic vesicles becomes thickened and invaginated, and the bulb is thus converted into a cup, the optic cup.

The periocular mesenchyme migrates inward during the formation of the optic cup and is critical for the induction of the retinal pigment epithelium and the optic nerve. The mesenchyme contributes to the cornea, iris, ciliary body, sclera, and blood vessels of the eye.

Chordamesoderm

Chordamesoderm induces the anterior portion of the neural tube to form the precursors of the synapomorphic tripartite brain of vertebrates, a bulge called the diencephalon. Further induction by the chordamesoderm forms a protrusion: the optic vesicle. This vesicle is subsequently invaginated by further inductions from the chordamesoderm, and induces the ectoderm that thickens (lens placode) and further invaginates to a point that detaches from the ectoderm and forms a neurogenic placode. The lens placode is triggered by the chordamesoderm to invaginate and form the optic cup, composed of an outer layer of neural retina and inner layer of pigmented retina that unite and form the optic stalk. The pigmented retina is formed by rods and cones and composed of small cilia typical of the ependymal epithelium of the neural tube. Some cells in the lens vesicle will form the cornea and the lens vesicle will develop completely to form the definitive lens. Iris is formed from the optic cup cells.

This page titled 30.7B: Development of Vision is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.8: Development of the Endocrine System

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.8: Development of the Endocrine System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

30.9: Development of Blood

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

This page titled 30.9: Development of Blood is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29: APPENDIX A: DISEASES, INJURIES, AND DISORDERS OF THE ORGAN SYSTEMS

29.10: Circulatory Diseases and Disorders

29.10A: Angiogenesis and Disease

29.10B: Varicose Veins

29.10C: Syncope

29.10D: Hypertension

29.11: Diseases and Disorders of the Immune System

29.11A: Immunodeficiencies

29.11B: Autoimmune Diseases

29.11C: Infectious Mononucleosis

29.11D: Psoriasis

29.12: Respiratory Diseases and Disorders

29.12A: Asthma

29.12B: Respiratory Distress Syndrome

29.12C: Cystic Fibrosis

29.12D: Hypoxia

29.13: Diseases and Disorders of the Digestive System

29.13A: Peritonitis

29.13B: Mumps

29.13C: Lactose Intolerance

29.13D: Peptic Ulcer Disease

29.14: Diseases and Disorders of the Metabolism

29.14A: Phenylketonuria (PKU)

29.14B: Galactosemia and Glycogen Storage Disease

29.15: Diseases and Disorders of the Urinary System

29.15A: Renal Disease and Failure

29.15B: Nephroptosis (Floating Kidney)

29.15C: Polycystic Kidney Disease

29.16: Body Fluid-Related Diseases and Disorders

29.16A: Acidosis

29.16B: Alkalosis

29.16C: Edema

29.17: Diseases and Disorders of the Reproductive System

29.17A: Sexually Transmitted Diseases (STDs)

29.17B: Erectile Dysfunction and the Blue Pill

29.17C: Cervical Cancer

29.18: Pregnancy-Related Disorders

29.18A: Ectopic Pregnancy

29.18B: Infertility

29.1: Diseases, Injuries, and Disorders of the Integumentary System

29.1A: Burns

29.1B: Allergies

29.1C: Acne

29.2: Diseases and Disorders of the Skeletal System

29.2A: Osteoporosis

29.2B: Paget's Disease

29.3: Injuries and Disorders of the Skeleton

29.3A: Spina Bifida

29.3B: Fractures

29.4: Diseases, Disorders, and Injuries of the Joints

29.4A: Sprain and Strain

29.4B: Rheumatism and Arthritis

29.4C: Common Joint Injuries

29.5: Diseases, Disorders, and Injuries of the Muscular System

29.5A: Muscular Dystrophy

29.5B: Exercise-Induced Muscle Damage

29.5C: Carpal Tunnel Syndrome

29.6: Injuries, Diseases, and Disorders of the Central Nervous System

29.6A: Traumatic Brain Injury

29.6B: Cerebrovascular Accident

29.6C: Transient Ischemic Attacks

29.6D: Alzheimer's Disease

29.6E: Brain Tumors

29.6F: Amyotrophic Lateral Sclerosis

29.7: Diseases and Disorders of the Endocrine System

29.7A: Pancreatic Islet Disorders: Diabetes and Hyperinsulinism

29.7B: Posttraumatic Stress Disorder (PTSD)

29.8: Blood Diseases and Disorders

29.8A: Hemostasis Disorders

29.8B: Anemia

29.8C: Sickle-Cell Disease

29.8D: Leukemia

29.9: Diseases and Disorders of the Heart

29.9A: Congenital Heart Defects

29.9B: Myocarditis and Endocarditis

29.9C: Myocardial Ischemia and Infarction

29.9D: Arrhythmia

This page titled 29: APPENDIX A: Diseases, Injuries, and Disorders of the Organ Systems is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.10: CIRCULATORY DISEASES AND DISORDERS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

29.10A: Angiogenesis and Disease

29.10B: Varicose Veins

29.10C: Syncope

29.10D: Hypertension

This page titled 29.10: Circulatory Diseases and Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.10A: Angiogenesis and Disease

Angiogenesis, the growth of new blood vessels from existing vessels, contributes to both normal tissue growth and tumorigenesis in cancer.

Learning Objectives

- Examine the relationship between angiogenesis and disease

Key Points

- Angiogenesis is a fundamental step in the transition of tumors from a dormant state to a malignant one, leading to the use of angiogenesis inhibitors as possible cancer therapeutics.
- Angiogenesis is also a target for combating diseases characterized by either poor vascularization or abnormal vasculature.
- Application of specific compounds that may inhibit or induce the creation of new blood vessels in the body may help combat such diseases as in the case of ischemic chronic wounds and age-related macular degeneration.

Key Terms


- **Malignant tumor**: In cancer, cells divide and grow uncontrollably, forming malignant tumors, and invade nearby parts of the body. The cancer may also spread to more distant parts of the body through the lymphatic system or bloodstream.
- **Angiogenesis inhibitors**: A substance that inhibits the growth of new blood vessels (angiogenesis). Some angiogenesis inhibitors are a normal part of the body's control, some are administered as drugs, and some come from diet.
- **VEGF**: Signal protein produced by cells that stimulates vasculogenesis and angiogenesis. It is part of the system that restores the oxygen supply to tissues when blood circulation is inadequate.

Angiogenesis is the physiological process involving the growth of new blood vessels from preexisting vessels. It is a normal and vital process in growth and development, as well as in wound healing and in the formation of granulation tissue. It is also a fundamental step in the transition of tumors from a dormant state to a malignant one, leading to the use of angiogenesis inhibitors. Tumors cannot grow beyond a certain size, generally 1–2 mm, due to a lack of oxygen and other essential nutrients.


Angiogenesis and Cancer

Cancer cells are cells that have lost their ability to divide in a controlled fashion. A malignant tumor consists of a population of rapidly-dividing and growing cancer cells. Mutations quickly accrue within the population. These mutations (variations) allow the cancer cells (or sub-populations of cancer cells within a tumor) to develop drug resistance and escape therapy.

Tumors induce blood vessel growth (angiogenesis) by secreting various growth factors (e.g., VEGF). Growth factors such as bFGF and VEGF can induce capillary growth into the tumor, which some researchers suspect supply the required nutrients that allow for tumor expansion. Bone marrow cell recruitment also participates in tumor angiogenesis.



Bone-marrow derived cells in tumor angiogenesis: Diverse chemoattractant factors promote the recruitment and infiltration of bone marrow cells to the tumor microenvironment where they suppress the antitumor immunity or promote tumor angiogenesis and vasculogenesis



Tumor Angiogenesis: The switch to the angiogenic phenotype occurs during multistage tumorigenesis. As malignancy develops, cells progress from a prevascular stage (normal to early hyperplasia) to a vascular stage (late hyperplasia to dysplasia to invasive carcinoma). Angiogenesis becomes clearly evident during dysplasia and is critical for further growth. Targeting tumor angiogenesis may be a novel strategy for preventing cancer.

In 2007, it was discovered that cancerous cells stop producing the anti-VEGF enzyme PKG. In normal cells (but not in cancerous ones), PKG apparently limits beta-catenin, which solicits angiogenesis. Other clinicians believe angiogenesis really serves as a waste pathway, taking away the biological end products secreted by rapidly-dividing cancer cells. In either case, angiogenesis is a necessary and required step for the transition from a small, harmless cluster of cells, often said to be about the size of the metal ball at the end of a ball-point pen, to a large tumor.

Angiogenesis is also required for the spread of a tumor, or metastasis. Single cancer cells can break away from an established solid tumor, enter the blood vessel, and be carried to a distant site, where they can implant and begin the growth of a secondary tumor.

Evidence now suggests the blood vessel in a given solid tumor may, in fact, be mosaic vessels, composed of endothelial cells and tumor cells. This mosaicity allows for substantial shedding of tumor cells into the vasculature, possibly contributing to the appearance of circulating tumor cells in the peripheral blood of patients with malignancies. The subsequent growth of such metastases will also require a supply of nutrients and oxygen and a waste disposal pathway.

Angiogenesis and Cardiovascular Disease

Angiogenesis also represents an excellent therapeutic target for the treatment of cardiovascular disease. It is a potent, physiological process that underlies the natural manner in which our bodies respond to a diminution of blood supply to vital

organs, namely the production of new collateral vessels to overcome the ischemic insult.

A large number of preclinical studies have been performed with protein-, gene- and cell-based therapies in animal models of cardiac ischemia, as well as models of peripheral artery disease. Reproducible and credible successes in these early animal studies led to high enthusiasm that this new therapeutic approach could be rapidly translated to a clinical benefit for millions of patients in the Western world suffering from these disorders.

A decade of clinical testing both gene- and protein-based therapies designed to stimulate angiogenesis in underperfused tissues and organs, however, has led from one disappointment to another. Although all of these preclinical readouts, which offered great promise for the transition of angiogenesis therapy from animals to humans were, in one fashion or another, incorporated into early stage clinical trials, the FDA has, to date (2007), insisted that the primary endpoint for approval of an angiogenic agent must be an improvement in exercise performance of treated patients.

Overexpression of VEGF causes increased permeability in blood vessels in addition to stimulating angiogenesis. In wet macular degeneration, VEGF causes proliferation of capillaries into the retina. Since the increase in angiogenesis also causes edema, blood, and other retinal fluids leak into the retina, causing loss of vision. A novel treatment of this disease is to use a VEGF-inhibiting siRNA to stop the main signaling cascade for angiogenesis.

This page titled 29.10A: Angiogenesis and Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.10B: Varicose Veins

Varicose veins, typically found in the legs, are those that have become enlarged and tortuous due to malfunctioning valves.

Learning Objectives

- Describe the cause of varicose veins

Key Points


- Veins have leaflet valves to prevent blood from flowing backwards; this backflow is called retrograde or reflux flow. The muscles of the leg pump the veins to return blood to the heart against the effects of gravity.
- When veins become varicose, the leaflets of the valves no longer meet properly, and the valves no longer work, a situation called valvular incompetence. This causes pooling of blood and enlargement of the veins.
- Varicose veins are more common in women than in men, and are linked with heredity. Other related factors are pregnancy, obesity, menopause, aging, prolonged standing, leg injury, abdominal straining, and crossing legs at the knees or ankles.
- Non-surgical treatments for varicose veins include elastic stockings, elevating the legs, and exercise. The traditional surgical treatment has been vein stripping, or removal of the affected veins.

Key Terms

- **superficial veins**: Those veins that are close to the surface of the body.
- **tortuous**: Twisted; having many turns; convoluted.

Varicose veins are those that have become enlarged and tortuous, commonly affecting the legs.

Causes of Varicose Veins



Varicose Veins: This image shows a normal vein with a working valve and normal blood flow. Figure B shows a varicose vein with a deformed valve, abnormal blood flow, and thin, stretched walls. The middle image shows where varicose veins might appear in a leg.

Veins have leaflet valves to prevent blood from flowing backwards; this backflow is called retrograde or reflux flow. The muscles of the leg pump the veins to return blood to the heart against the effects of gravity. When veins become varicose, the leaflets of the valves no longer meet properly, and the valves no longer work. This change is called valvular incompetence, and it allows blood to flow backwards. To accommodate the retrograde or reflux flow, the veins enlarge even more.



Leg with Varicose Veins: A man with many varicose veins in the lower leg.

Varicose veins are most often found in the superficial veins of the legs, which are subject to high pressure when standing. In addition to cosmetic concerns, varicose veins are often painful, especially when standing or walking. They can often itch, and scratching them can lead to skin ulcers.

Varicose veins are more common in women than in men, and are linked with heredity. Other related factors are pregnancy, obesity, menopause, aging, prolonged standing, leg injury, abdominal straining, and crossing legs at the knees or ankles. They are distinguished from “spider veins,” which are also due to valvular incompetence, by their size. Spider veins are less than one millimeter in diameter, whereas varicose veins are larger than this in diameter.

Treatments

Non-surgical treatments for varicose veins include elastic stockings, elevating the legs, and exercise. The traditional surgical treatment has been vein stripping, or removal of the affected veins. Because most of the blood in the legs is returned by the deep veins, the superficial veins, which return only about 10 per cent of the total blood of the legs, can usually be removed without serious harm.

This page titled 29.10B: Varicose Veins is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.10C: Syncope

Syncope, the medical term for fainting, is a transient loss of consciousness.

Learning Objectives

- Identify the main factor associated with syncope


Key Points

- This loss of consciousness can arise due to changes in postural tone.
- The two broad categories of syncope are cardiogenic and reflex; of the two, cardiogenic syncope is more likely to result in morbidity or mortality.
- Vasovagal syncope is situational syncope which may arise in situations where the patient feels scared, threatened, embarrassed, or uneasy.

Key Terms

- **cardiogenic syncope**: The form of syncope that is more likely to produce serious morbidity or mortality and require prompt treatment.
- **reflex syncope**: Involuntary fainting.
- **vasovagal syncope**: Situational syncope which may arise in situations where the patient feels scared, threatened, embarrassed, or uneasy.

Syncope, the medical term for fainting, is defined as a transient loss of consciousness and postural tone characterized by rapid onset, short duration, and spontaneous recovery due to global cerebral hypoperfusion that most often results from hypotension.



Vagus Nerve: The vagus nerve, labeled at the top left and colored in yellow, may lower heart rate, reducing blood flow to the brain and causing vasovagal syncope.

Many forms of syncope are preceded by a prodromal state that often includes dizziness and loss of vision (“blackout”) (temporary), loss of hearing (temporary), loss of pain and feeling (temporary), nausea and abdominal discomfort, weakness, sweating, a feeling of heat, palpitations, and other phenomena, which–if they do not progress to loss of consciousness and postural tone–are

often denoted “presyncope.” Abdominal discomfort prior to loss of consciousness may be indicative of seizure which should be considered different than syncope.

Types of Syncope

There are two broad categories of syncope, cardiogenic or reflex, which underlie most forms of syncope. Cardiogenic forms are more likely to produce serious morbidity or mortality and require prompt or even immediate treatment. Although cardiogenic syncope is much more common in older patients, an effort to rule out arrhythmic, obstructive, ischemic, or cardiomyopathic causes of syncope and circulatory inadequacy is mandatory in each patient.

Variants of reflex syncope often have characteristic histories, including precipitants and time course which are made evident by skilled history taking. Thus, the clinical history is the foremost tool used in the differential diagnosis of syncope. Physical examination, and electrocardiogram are part of the initial evaluation of syncope and other more specific tools such as loop recorders may be necessary in clinically uncertain cases.

Vasovagal Syncope

Vasovagal (situational) syncope, one of the most common types, may occur in scary, embarrassing or uneasy situations, or during blood drawing, coughing, urination, or defecation. In vasovagal syncope, the vagus nerve decreases blood flow to the brain. Other types include postural syncope (caused by a change in body posture), cardiac syncope (due to heart-related conditions), and neurological syncope (due to neurological conditions). There are many other causes of syncope, including low blood sugar levels and lung disease such as emphysema and a pulmonary embolus. The cause of the fainting can be determined by a doctor using a complete history, physical, and various diagnostic tests.

The vasovagal type can be considered in two forms:

- Isolated episodes of loss of consciousness, unheralded by any warning symptoms for more than a few moments. These tend to occur in the adolescent age group, and may be associated with fasting, exercise, abdominal straining, or circumstances promoting vasodilation (e.g., heat, alcohol). The subject is invariably upright. The tilt-table test, if performed, is generally negative.
- Recurrent syncope with complex associated symptoms called neurally mediated syncope. It is associated with any of the following: preceding or succeeding sleepiness, preceding visual disturbance (“spots before the eyes”), sweating, and light-headedness. The subject is usually but not always upright. The tilt-table test, if performed, is generally positive.

This page titled 29.10C: Syncope is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.10D: Hypertension

Hypertension is elevated blood pressure, clinically defined as at or greater than 140/90 (systolic/diastolic) mm/Hg.

Learning Objectives

- Evaluate the factors involved in hypertension

Key Points

- Primary hypertension has multiple or unknown causes, and secondary hypertension has known causes.
- In most cases of primary hypertension, increased peripheral resistance is the reason for the elevated pressure.
- Even moderate elevation of arterial blood pressure is associated with a shortened life expectancy.
- Chronically elevated blood pressure leads to widespread damage to the circulatory system, arterial disease, cardiac failure, and even neurological impairments.

Key Terms

- **total peripheral resistance**: Total peripheral resistance (TPR) is the sum of the resistance of all peripheral vasculature in the systemic circulation. Vasculature throughout the entire body can be thought of as two separate circuits—one is the systemic circulation, while the other is the pulmonary circulation.
- **hypertension**: The disease or disorder of abnormally high blood pressure.
- **blood pressure**: The pressure exerted by the blood against the walls of the arteries and veins; it varies during the heartbeat cycle, and according to a person's age, health, and physical condition.

Hypertension (HTN) or high blood pressure, sometimes called arterial hypertension, is a chronic medical condition in which the blood pressure in the arteries is elevated. This requires the heart to work harder than normal to circulate blood through the blood vessels. Blood pressure involves two measurements, systolic and diastolic, which depend on whether the heart muscle is contracting (systole) or relaxed between beats (diastole).


Normal blood pressure at rest is within the range of 100–140mmHg systolic (top reading) and 60–90mmHg diastolic (bottom reading). High blood pressure is said to be present if it is persistently at or above 140/90 mmHg. The repercussions of chronically elevated blood pressure included widespread damage to the circulatory system, arterial disease, cardiac failure and even neurological impairments

Primary and Secondary Hypertension

Hypertension is classified as either primary (essential) hypertension or secondary hypertension; about 90–95% of cases are categorized as "primary hypertension" which means high blood pressure with no obvious underlying medical cause. The remaining 5–10% of cases (secondary hypertension) are caused by other conditions that affect the kidneys, arteries, heart, or endocrine system. In most people with established essential (primary) hypertension, increased resistance to blood flow (total peripheral resistance) accounts for the high pressure while cardiac output remains normal. This increased peripheral resistance is mainly attributable to structural narrowing of small arteries and arterioles, although a reduction in the number or density of capillaries may also contribute. Secondary hypertension results from an identifiable cause. Renal disease is the most common secondary cause of hypertension. Hypertension can also be caused by endocrine conditions, such as Cushing's syndrome, hyperthyroidism, hypothyroidism, and acromegaly. Other causes of secondary hypertension include obesity, sleep apnea, pregnancy, and coarctation of the aorta.

There are many complications of chronic hypertension. Hypertension is a major risk factor for stroke, myocardial infarction (heart attacks), heart failure, aneurysms of the arteries (e.g., aortic aneurysm), peripheral arterial disease, and is a cause of chronic kidney disease. Even moderate elevation of arterial blood pressure is associated with a shortened life expectancy. Dietary and lifestyle changes can improve blood pressure control and decrease the risk of associated health complications, although drug

treatment is often necessary in people for whom lifestyle changes prove ineffective or insufficient.



High Blood Pressure Complications: Persistant high blood pressure can affect many areas of the body.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Angiogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Angiogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Angiogenesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Angiogenesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Malignant tumor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Malignant%20tumor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Angiogenesis inhibitors. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Angioge...s%20inhibitors**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vegf. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vegf**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myeloid Cells in the Tumor Microenvironment: Modulation of Tumor Angiogenesis and Tumor Inflammation. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2871549/**. **License**: ***Public Domain: No Known Copyright***
- PubMed Central, Figure 1: J Oncol.n2012; 2012: 879623.nPublished online 2011 September 29. doi:u00a010.1155/2012/879623. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/arti...8/figure/fig1/**. **License**: ***Public Domain: No Known Copyright***
- Varicose veins. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Varicose_veins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- tortuous. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/tortuous**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology...erficial-veins**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myeloid Cells in the Tumor Microenvironment: Modulation of Tumor Angiogenesis and Tumor Inflammation. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2871549/**. **License**: ***Public Domain: No Known Copyright***
- PubMed Central, Figure 1: J Oncol.n2012; 2012: 879623.nPublished online 2011 September 29. doi:u00a010.1155/2012/879623. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/arti...8/figure/fig1/**.

License: ***Public Domain: No Known Copyright***
- Leg Before 1. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Leg_Before_1.jpg**. **License**: ***Public Domain: No Known Copyright***
- Varicose veins. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...cose_veins.jpg**. **License**: ***Public Domain: No Known Copyright***
- Syncope (medicine). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Syncope...e)%23Vasovagal**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- vasovagal syncope. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/vasovagal%20syncope**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cardiogenic syncope. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cardiogenic%20syncope**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- reflex syncope. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/reflex%20syncope**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myeloid Cells in the Tumor Microenvironment: Modulation of Tumor Angiogenesis and Tumor Inflammation. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2871549/**. **License**: ***Public Domain: No Known Copyright***
- PubMed Central, Figure 1: J Oncol.n2012; 2012: 879623.nPublished online 2011 September 29. doi:u00a010.1155/2012/879623. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/arti...8/figure/fig1/**. **License**: ***Public Domain: No Known Copyright***
- Leg Before 1. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Leg_Before_1.jpg**. **License**: ***Public Domain: No Known Copyright***
- Varicose veins. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Varicose_veins.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gray793. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray793.png**. **License**: ***Public Domain: No Known Copyright***
- Hypertension. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypertension. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypertension. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- General Anatomy/Heart. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/General_Anatomy/Heart**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypertension. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- total peripheral resistance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/total%20peripheral%20resistance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypertension. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypertension**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- blood pressure. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/blood_pressure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myeloid Cells in the Tumor Microenvironment: Modulation of Tumor Angiogenesis and Tumor Inflammation. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2871549/**. **License**: ***Public Domain: No Known Copyright***
- PubMed Central, Figure 1: J Oncol.n2012; 2012: 879623.nPublished online 2011 September 29. doi:u00a010.1155/2012/879623. **Provided by**: National Insitute of Health. **Located at**: **http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3184418/figure/fig1/**. **License**: ***Public Domain: No Known Copyright***
- Leg Before 1. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Leg_Before_1.jpg**. **License**: ***Public Domain: No Known Copyright***
- Varicose veins. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Varicose_veins.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gray793. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray793.png**. **License**: ***Public Domain: No Known Copyright***
- Main complications of persistent high blood pressure. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Main_complications_of_persistent_high_blood_pressure.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 29.10D: Hypertension is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.11: DISEASES AND DISORDERS OF THE IMMUNE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.11A: Immunodeficiencies
29.11B: Autoimmune Diseases
29.11C: Infectious Mononucleosis
29.11D: Psoriasis

This page titled 29.11: Diseases and Disorders of the Immune System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.11A: Immunodeficiencies

Immunodeficiency is a state where the immune system's ability to fight infectious disease is impaired or absent.

Learning Objectives

- Describe immunodeficiencies of the immune system


Key Points

- A person with an immunodeficiency of any kind is said to be immunocompromised.
- Primary immunodeficiencies are also known as congenital immunodeficiencies as these disorders are inherited.
- Secondary immunodeficiencies are acquired from various environmental factors such as malnutrition, immunosuppressive drugs, and aging.

Key Terms

- **primary immunodeficiency**: Any of the many genetic disorders due to an aspect of the body's immune system being either absent or failing to function normally.

Immunodeficiency (or immune deficiency) is a state in which the immune system's ability to fight infectious disease is compromised or absent entirely. Immunodeficiency may also decrease cancer immunosurveillance. Most cases of immunodeficiency are acquired (secondary) but some people are born with a defective immune system, or primary immunodeficiency. As an anti-rejection protocol, transplant patients take medications to suppress their immune system, as do some patients suffering from an over-active immune system. A person with an immunodeficiency of any kind is said to be immunocompromised. An immunocompromised person may be particularly susceptible to opportunistic infections, as well as normal infections affecting the general population.



Genetic Immodeficiencies: Chart showing locations of malfunction leading to common errors of lymphocyte development.

Types of Immunodeficiencies

Humoral immune deficiency (with signs or symptoms depending on the cause) generally includes signs of hypogammaglobulinemia (decrease of one or more types of antibodies) with presentations including repeated mild respiratory infections, and/or agammaglobulinemia (lack of all or most antibody production) resulting in frequent severe infections and is often fatal.

T-cell deficiency is often caused secondary disorders such as acquired immune deficiency syndrome.

Granulocyte deficiency refers to decreased numbers of granulocytes (called granulocytopenia or, if absent, agranulocytosis) such as of neutrophil granulocytes (termed neutropenia). Granulocyte deficiencies also include decreased function of individual granulocytes, such as in chronic granulomatous disease. Asplenia is a granulocyte deficiency in which the spleen does not function.

Complement deficiency is where the function of the complement system is deficient.

Primary vs. Secondary

Immunodeficiency often affects multiple components, with notable examples including severe combined immunodeficiency (which is primary) and acquired immune deficiency syndrome (which is secondary). Distinctions between primary versus secondary immunodeficiencies are based, respectively, on whether the cause originates within the immune system itself or, in turn, is due to insufficiency of a supporting component of it, or an external decreasing factor of it.

A number of rare diseases feature a heightened susceptibility to infections from childhood onward. Primary Immunodeficiency is also known as congenital immunodeficiency. Many of these disorders are hereditary and are autosomal recessive or X-linked. There are over 80 recognized primary immunodeficiency syndromes—generally grouped by the part of the immune system that is malfunctioning, such as lymphocytes or granulocytes. The treatment of primary immunodeficiencies depends on the nature of the defect, and may involve antibody infusions, long-term antibiotics, and (in some cases) stem cell transplantation.

Secondary immunodeficiencies, also known as acquired immunodeficiencies, can result from various immunosuppressive agents, for example, malnutrition, aging, and particular medications (e.g., chemotherapy, disease-modifying antirheumatic drugs, immunosuppressive drugs after organ transplants, and glucocorticoids). For medications, the term immunosuppression generally refers to both beneficial and potential adverse effects of decreasing the function of the immune system. The term immunodeficiency generally refers solely to the adverse effect of increased risk for infection. Many specific diseases directly or indirectly cause immunosuppression. This includes many types of cancer, particularly those of the bone marrow and blood cells (those with leukemia, lymphoma, multiple myeloma, etc.), and certain chronic infections. Immunodeficiency is also the hallmark of acquired immunodeficiency syndrome (AIDS), caused by the human immunodeficiency virus (HIV). HIV directly infects a small number of T helper cells, and also impairs other immune system responses indirectly.

This page titled 29.11A: Immunodeficiencies is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.11B: Autoimmune Diseases

Autoimmune diseases are an inappropriate immune response against tissues in the body.

Learning Objectives

- Describe autoimmune diseases

Key Points

- Autoimmune diseases arise from an immune system that does not function properly.
- The common cause hypothesis of autoimmune diseases describes how autoimmune diseases are triggered by genes located in the same regions on the chromosome.
- Despite genetic factors being involved in the onset of autoimmune disease, much is not know about the other causes of autoimmunity.

Key Terms

- **autoimmune diseases**: Arise from an inappropriate immune response of the body against substances and tissues normally present in the body.
- **common cause hypothesis**: Predicts that common disease-causing alleles, or variants, will be found in all human populations which manifest a given disease.

Autoimmune diseases arise from an inappropriate immune response of the body against substances and tissues normally present in the body. In other words, the immune system mistakes some part of the body as a pathogen and attacks its own cells. Autoimmune diseases are commonly considered complex immune disorders. While many autoimmune diseases are rare, collectively these diseases afflict millions of patients. Approximately 5–8% of the U.S. population suffers from this group of chronic, debilitating diseases. Despite their clinical diversity, they have one similarity, namely the dysfunction of the immune system. It is suspected that genetic defects play a role in the etiology of these diseases.

Causes and Treatments

Modern high throughput technologies, like mRNA microarrays, have enabled researchers to investigate diseases at a genome-wide level. In contrast to classical inherited genetic diseases, like sickle cell anemia, autoimmune diseases are not caused by the defect of a single gene, but by the dysfunction of the complex interaction of a group of genes. Although no autoimmune disease has been completely analyzed, there has been tremendous success in recent years in identifying major players in the development of autoimmune diseases. There are over 50 publications that list gene variants that are associated with a certain autoimmune disease. Interestingly, a lot of these genes are located in the same regions on the chromosomes, the so called susceptibility regions. This has led to a “common cause hypothesis” of autoimmune disorders. Several organizations and institutes have established programs to investigate this common cause hypothesis. However, defects of one or more of these genes do not cause an autoimmune disease, but only predispose a person for an autoimmune disease.

The factors that trigger an autoimmune disease are still unknown. Studies with monogenetic twins have revealed that genetic influences only account for 25–40% of the disease risk making gene-environment interactions or environmental influences the predominant factors. The environmental influences are very diverse, rendering research in this area extremely difficult. These influences may be toxic substances like mercury in one case and ultraviolet light or even certain nutrients in another. Moreover, several bacteria, viruses, or hormones are among the suspected triggers of autoimmune disorders. The treatment of autoimmune diseases is typically with immunosuppression: medication which decreases the immune response.



Autoimmune Response: Erythema multiform is a rash that appears as an auto-immune response to a predisposing agent: a bacteria, an immunization, an antibiotic (in this case, amoxicillin).

This page titled 29.11B: Autoimmune Diseases is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.11C: Infectious Mononucleosis

Mononucleosis is an infectious disease caused by the Epstein-Barr virus (EBV) and results in flu-like symptoms.

Learning Objectives

- Describe infectious mononucleosis

Key Points

- Infectious mononucleosis is an infectious disease caused by the Epstein-Barr virus (EBV).
- Infectious mononucleosis (also called mono or kissing disease) is spread orally and is characterized by symptoms such as fever, sore throat, and fatigue.
- Once infected, the virus lives dormantly in B lymphocytes in a person.

Key Terms


- **Epstein-Barr virus (EBV)**: Virus responsible for causing infectious mononucelosis.
- **infectious mononucleosis**: An infectious, widespread viral disease caused by the Epstein–Barr virus (EBV), one type of herpes virus. Most people are exposed to the virus as children, when the disease produces no noticeable or only flu-like symptoms.

Infectious mononucleosis is an infectious, widespread viral disease caused by the Epstein–Barr virus (EBV), one type of herpes virus, to which more than 90% of adults have been exposed. Occasionally, the symptoms can recur at a later period. It is sometimes colloquially known as the “kissing disease” from its oral transmission. Most people are exposed to the virus as children, when the disease produces no noticeable or only flu-like symptoms. In developing countries, people are exposed to the virus in early childhood more often than in developed countries. As a result, the disease in its observable form is more common in developed countries. It is most common among adolescents and young adults.

Symptoms

The disease is characterized by fever, sore throat, and fatigue, along with several other possible signs and symptoms, especially in adolescents and young adults. The infection is spread via saliva and has an incubation period of four to seven weeks. Symptoms usually persist for two to three weeks, but fatigue is often more prolonged. Infectious mononucleosis is primarily diagnosed by observation of symptoms, but suspicion can be confirmed by several diagnostic tests. The most commonly used diagnostic criterion is the presence of 50% lymphocytes with at least 10% atypical lymphocytes (large, irregular nuclei), while the person also has fever, pharyngitis, and adenopathy. Infectious mononucleosis is generally self-limiting, so only symptomatic and/or supportive treatments are used. Rest is recommended during the acute phase of the infection.

Once the acute symptoms of an initial infection disappear, they often do not return. But once infected, the patient carries the virus for the rest of his or her life. The virus typically lives dormantly in B lymphocytes. Independent infections of mononucleosis may be contracted multiple times, regardless of whether the patient is already carrying the virus dormantly.



Main Symptoms of Infectious Mononucleosis: This graph illustrates the location of the symptoms.

This page titled 29.11C: Infectious Mononucleosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.11D: Psoriasis

Psoriasis is an autoimmune disease that affects the skin, caused by faulty immune signals that speed up the growth cycle of skin cells.

Learning Objectives

- Describe psoriasis

Key Points

- The plaque form is most common form of psoriasis and usually presents as red and white hues of scaly patches on the epidermis.
- Psoriasis is a chronic recurring condition that varies in severity from minor, localized patches to complete body coverage.
- The cause of psoriasis is not fully understood, but it is believed to have a genetic component.
- There is currently no cure, but various treatments can help to control the symptoms. Typically, topical agents are used for mild disease, phototherapy for moderate disease, and systemic immunosupressent agents for severe disease.

Key Terms

- **psoriasis**: A noncontagious disease whose main symptom is gray or silvery flaky patches on the skin which are red and inflamed underneath when scratched.
- **plaque**: A plaque is a raised area of inflamed skin covered with silvery white scaly skin and connective tissue.
- **autoimmune disease**: A disease where one's immune system attacks one's own tissues.

Psoriasis is an autoimmune disease that affects the skin. It occurs when the immune system mistakes the skin cells as a pathogen, and sends out faulty signals that speed up the growth cycle of skin cells. Psoriasis is not contagious. The most common form, plaque psoriasis, is commonly seen as red and white colored scaly patches appearing on the top first layer of the skin. Plaques frequently occur on the skin of the elbows and knees, but can affect any area, including the back, scalp, palms of hands and soles of feet, and genitals. In contrast to eczema, psoriasis is more likely to be found on the skin of outer side of the joint rather than in the crease.

Symptoms and Complications

Depending on the severity and location of outbreaks, individuals may experience significant physical discomfort and some disability. Itching and pain can interfere with basic functions such as self-care, walking, and sleep. Plaques on hands and feet can prevent individuals from working at certain occupations, playing some sports, and caring for family members or a home. Quality of life is an issue for psoriasis patients, as they may feel self-conscious about the appearance of their skin and have a poor self-image that stems from fear of public rejection and psychosexual concerns. Psychological distress can lead to significant depression and social isolation.

Treatment

Psoriasis is typically a lifelong condition. There is currently no cure, but various treatments can help to control the symptoms. Typically, topical agents are used for mild disease, phototherapy for moderate disease, and systemic immunosuppressant agents for severe disease.



Psoriasis: A person whose back and arms are affected by psoriasis.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Immunodeficiency. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Immunodeficiency**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- primary immunodeficiency. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/primary_immunodeficiency**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Primary immunodeficiency. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Primary_immunodeficiency**. **License**:
- Autoimmune disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Autoimmune_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- BMC Bioinformatics | Full text | The Autoimmune Disease Database: a dynamically compiled literature-derived database. **Provided by**: BioMed Central. **Located at**: **http://www.biomedcentral.com/1471-2105/7/325**. **License**: ***CC BY: Attribution***
- common cause hypothesis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/common%...e%20hypothesis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autoimmune diseases. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/autoimmune%20diseases**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Primary immunodeficiency. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Primary_immunodeficiency**. **License**:
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/a/a6/Erythema_multiforme.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Infectious mononucleosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Infectious_mononucleosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Epstein-Barr virus (EBV). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Epstein...0virus%20(EBV)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- infectious mononucleosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/infectious%20mononucleosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Primary immunodeficiency. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Primary_immunodeficiency**. **License**:
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/a/a6/Erythema_multiforme.jpg**.

License: *CC BY-SA: Attribution-ShareAlike*
- Infectious mononucleosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Infectious_mononucleosis. **License**:
- Psoriasis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Psoriasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- psoriasis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/psoriasis. **License**: *CC BY-SA: Attribution-ShareAlike*
- autoimmune disease. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/autoimmune+disease. **License**: *CC BY-SA: Attribution-ShareAlike*
- plaque. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/plaque. **License**: *CC BY-SA: Attribution-ShareAlike*
- Primary immunodeficiency. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Primary_immunodeficiency. **License**:
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/a/a6/Erythema_multiforme.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Infectious mononucleosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Infectious_mononucleosis. **License**:
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/f/f5/Psoriasis_on_back1.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 29.11D: Psoriasis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.12: RESPIRATORY DISEASES AND DISORDERS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.12A: Asthma
29.12B: Respiratory Distress Syndrome
29.12C: Cystic Fibrosis
29.12D: Hypoxia

This page titled 29.12: Respiratory Diseases and Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.12A: Asthma

Asthma is a common chronic inflammatory disease of the airways characterized by reversible airflow obstruction and bronchospasm.

Learning Objectives

- Identify the triggers that cause asthma (reversible blockage of bronchi) attacks


Key Points

- Symptoms of asthma include wheezing, coughing, chest tightness, and shortness of breath.
- Asthma may also be classified as atopic (extrinsic) or non-atopic (intrinsic) and is thought to be caused by a combination of genetic and environmental factors.
- Treatment of acute symptoms is usually with an inhaled short-acting beta-2 agonist (such as salbutamol).

Key Terms

- **asthma**: A long-term respiratory condition, in which the airways may unexpectedly and suddenly narrow, often in response to an allergen, cold air, exercise, or emotional stress. Symptoms include wheezing, shortness of breath, chest tightness, and coughing.
- **beta-2 agonist**: β2-adrenergic agonists, also known as β2-adrenergic receptor agonists, are a class of drugs used to treat asthma and other pulmonary disease states.
- **wheezing**: A wheeze (formally called “sibilant rhonchi” in medical terminology) is a continuous, coarse, whistling sound produced in the respiratory airways during breathing. For wheezes to occur, some part of the respiratory tree must be narrowed or obstructed, or airflow velocity within the respiratory tree must be heightened.

Asthma is a common chronic inflammatory disease of the airways characterized by variable and recurring symptoms, reversible airflow obstruction, and bronchospasm. Symptoms include wheezing, coughing, chest tightness, and shortness of breath. The prevalence of asthma has been rising steadily since the early 1980s. Asthma is clinically classified according to the frequency of symptoms, forced expiratory volume over one second (FEV1), and peak expiratory flow rate. The latter can be measured with a peak flow meter. Asthma may also be classified as atopic (extrinsic) or non-atopic (intrinsic). It is thought to be caused by a combination of genetic and environmental factors. Treatment of acute symptoms is usually with an inhaled short-acting beta-2 agonist (such as salbutamol). Symptoms can be prevented by avoiding triggers such as allergens and irritants and by inhaling corticosteroids. Leukotriene antagonists are less effective than corticosteroids and thus less preferred. Its diagnosis is usually made based on the pattern of symptoms and/or response to therapy over time.




Two peak flow meters.: Peak flow meters are used to measure peak expiratory flow rate.

Although asthma is a chronic obstructive condition, it is not considered as a part of chronic obstructive pulmonary disease as this term refers specifically to irreversible combinations of disease such as bronchiectasis, chronic bronchitis, and emphysema. Unlike these diseases, the airway obstruction in asthma is usually reversible; however, if left untreated, the chronic inflammation from asthma can lead the lungs to become irreversibly obstructed due to airway remodeling. In contrast to emphysema, asthma affects the bronchi, not the alveoli.

Asthma Attacks

An acute asthma exacerbation is commonly referred to as an asthma attack. The classic symptoms are shortness of breath, wheezing, and chest tightness. While these are the primary symptoms of asthma, in severe cases air motion may be significantly impaired such that no wheezing is heard. Signs that occur during an asthma attack include the use of accessory muscles of respiration (sternocleidomastoid and scalene muscles of the neck); there may be a paradoxical pulse (a pulse that is weaker during inhalation and stronger during exhalation) and over-inflation of the chest. A blue color of the skin and nails may occur from lack of oxygen.



Alveolar Effects of Asthma: During an asthma episode, inflamed airways react to environmental triggers such as smoke, dust,

or pollen. The airways narrow and produce excess mucus, making it difficult to breathe.

Causes

Asthma is caused by environmental and genetic factors. These factors influence how severe asthma is and how well it responds to medication. For example, increased exposure to indoor allergens in infancy and early childhood has been analyzed as a primary cause of the rise in asthma. Primary prevention studies aimed at the aggressive reduction of airborne allergens in a home with infants have shown mixed findings. Strict reduction of dust mite allergens, for example, reduces the risk of allergic sensitization to dust mites, and modestly reduces the risk of developing asthma up until the age of 8 years old. However, studies also showed that the effects of exposure to cat and dog allergens worked in the converse fashion; exposure during the first year of life was found to reduce the risk of allergic sensitization and of developing asthma later in life. Similarly, many environmental risk factors have been associated with asthma development and morbidity in children. For example, recent studies show a direct relationship between increased exposure to air pollutants and incidence of childhood asthma. Lastly, viral respiratory infections (e.g., rhinovirus, Chlamydia pneumoniae) are both a leading trigger of asthma exacerbation and may increase the risk of developing asthma, especially in young children.

Diagnosis

There is currently no precise physiologic, immunologic, or histologic test for diagnosing asthma. The diagnosis is usually made based on the pattern of symptoms (airways obstruction and hyperresponsiveness) and/or response to therapy (partial or complete reversibility) over time. The U.S. National Asthma Education and Prevention Program (NAEPP) uses a 'symptom patterns' approach. Their guidelines for the diagnosis and management of asthma state that a diagnosis of asthma begins by assessing for the presence of multiple key indicators such as wheezing, coughing, or viral infections which will increases the probability of a diagnosis of asthma. Spirometry is needed to establish a diagnosis of asthma.

Treatment

Upon diagnosis, bronchodilators are recommended for short-term relief of symptoms. In those with occasional attacks, no other medication is needed. If mild persistent disease is present (more than two attacks a week), low-dose inhaled glucocorticoids or alternatively, an oral leukotriene antagonist or a mast cell stabilizer is recommended. For those who suffer daily attacks, a higher dose of inhaled glucocorticoid is used. In a severe asthma exacerbation, oral glucocorticoids are added to these treatments. Avoidance of triggers is a key component of improving control and preventing attacks. The most common triggers include allergens, smoke (tobacco and other), air pollution, non selective beta-blockers, and sulfite-containing foods. Cigarette smoking and second-hand smoke (passive smoke) may reduce the effectiveness of management medications such as steroid/corticosteroid therapies.

This page titled 29.12A: Asthma is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.12B: Respiratory Distress Syndrome

Acute respiratory distress syndrome (ARDS) is a serious reaction to various forms of injuries or acute infection to the lung.

Learning Objectives

- Identify the factors involved in respiratory distress syndrome (RDS)

Key Points


- ARDS is characterized by inflammation of the lung parenchyma leading to impaired gas exchange with concomitant systemic release of inflammatory mediators causing inflammation, hypoxemia, and frequently resulting in multiple organ failure.
- This condition is often fatal, usually requiring mechanical ventilation and admission to an intensive care unit.
- People with ARDS usually present with shortness of breath, tachypnea leading to hypoxia, and providing less oxygen to the brain occasionally causing confusion.

Key Terms

- **acute respiratory distress syndrome**: Acute respiratory distress syndrome (ARDS) is a serious reaction to various forms of injuries or acute infection to the lung. ARDS is a severe lung syndrome (not a disease) caused by a variety of direct and indirect issues.

Acute respiratory distress syndrome (ARDS), also known as respiratory distress syndrome (RDS) or adult respiratory distress syndrome, is a serious reaction to various forms of injuries or acute infection to the lung. ARDS is a severe lung syndrome (not a disease) caused by a variety of direct and indirect issues. It is characterized by inflammation of the lung parenchyma leading to impaired gas exchange with concomitant systemic release of inflammatory mediators causing inflammation, hypoxemia, and frequently resulting in multiple organ fail. This condition is often fatal, usually requiring mechanical ventilation and admission to an intensive care unit.

Symptoms



X-ray of patient with ARDS.: Chest x-ray of patient with ARDS.

People with ARDS usually present with shortness of breath, tachypnea leading to hypoxia, and providing less oxygen to the brain occasionally causing confusion. ARDS can occur within 24 to 48 hours of an injury (trauma, burns, aspiration, massive blood transfusion, drug/alcohol abuse) or an acute illness (infectious pneumonia, sepsis, acute pancreatitis). It is characterized by bilateral infiltrates on chest radiograph sparing costophrenic angles and pulmonary artery wedge pressure < 18 mmHg (obtained by pulmonary artery catheterization).

Treatment

Mechanical ventilation is an essential part of the treatment of ARDS to relieve respiratory muscles of their work, and to protect the usually obtunded patient's airways. Ventilation is usually delivered through oro-tracheal intubation, or tracheostomy whenever prolonged ventilation (≥2 weeks) is deemed inevitable. However, mechanical ventilation may constitute a risk factor for the development, or the worsening, of ARDS.

Aside from the infectious complications arising from invasive ventilation with tracheal intubation, positive-pressure ventilation directly alters lung mechanics during ARDS. The result is higher mortality, i.e., through baro-trauma, when these techniques are used. If the underlying disease or injurious factor is not removed, the amount of inflammatory mediators released by the lungs in ARDS may result in a systemic inflammatory response syndrome (or sepsis if there is lung infection).

The evolution towards shock and/or multiple organ failure follows paths analogous to the pathophysiology of sepsis. This adds up to the impaired oxygenation which is the central problem of ARDS, as well as to respiratory acidosis, which is often caused by ventilation techniques such as permissive hypercapnia which attempt to limit ventilator-induced lung injury in ARDS. The result is a critical illness in which the "endothelial disease" of severe sepsis/SIRS is worsened by the pulmonary dysfunction, which further impairs oxygen delivery. Appropriate antibiotic therapy must be administered as soon as microbiological culture results are available. Empirical therapy may be appropriate if local microbiological surveillance is efficient.

This page titled 29.12B: Respiratory Distress Syndrome is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.12C: Cystic Fibrosis

Cystic fibrosis (CF) is an autosomal recessive disorder leading to respiratory congestion, multiple organ failure, and metabolic changes.

Learning Objectives

- Evaluate the factors involved in cystic fibrosis (CF)


Key Points

- Cystic fibrosis is characterized by abnormal transport of chloride and sodium across the epithelium, leading to thick, viscous secretions.
- CF is caused by a mutation in the gene coding for cystic fibrosis transmembrane conductance regulator (CFTR), which is required to regulate the components of sweat, digestive juices, and mucus.
- The hallmark symptoms of cystic fibrosis are salty tasting skin, poor growth, poor weight gain despite a normal food intake, accumulation of thick, sticky mucus, frequent chest infections, and coughing or shortness of breath.

Key Terms

- **CFTR gene**: The gene that encodes the CFTR protein is found on the human chromosome 7, on the long arm at position q31.2. from base pair 116,907,253 to base pair 117,095,955. CFTR orthologs have also been identified in all mammals for which complete genome data are available.
- **cystic fibrosis**: An inherited condition in which the exocrine glands produce abnormally viscous mucus, causing chronic respiratory and digestive problems.
- **autosomal recessive**: A mode of inheritance of genetic traits located on the allosomes (the sex determining chromosomes).

Cystic fibrosis (also known as CF or mucoviscidosis) is an autosomal recessive genetic disorder affecting most critically the lungs and also the pancreas, liver, and intestine. It is characterized by abnormal transport of chloride and sodium across an epithelium, leading to thick, viscous secretions. The name cystic fibrosis refers to the characteristic scarring (fibrosis) and cyst formation within the pancreas. Difficulty breathing is the most serious symptom and results from frequent lung infections that are treated with antibiotics and other medications. Other symptoms, including sinus infections, poor growth, and infertility affect other parts of the body. CF is caused by a mutation in the gene for the protein cystic fibrosis transmembrane conductance regulator (CFTR). The ΔF508 human genome mutation is characterized by the deletion of three base pairs in the CFTR nucleotide sequence, causing the loss of the amino acid phenylalanine located at position 508. This protein is required to regulate the components of sweat, digestive juices, and mucus. CFTR regulates the movement of chloride and sodium ions across epithelial membranes, such as the alveolar epithelia located in the lungs. Although most people without CF have two working copies of the CFTR gene, only one is needed to prevent cystic fibrosis due to the disorder's recessive nature. CF develops when neither gene works normally (as a result of mutation) and therefore has autosomal recessive inheritance.




Cystic Fibrosis Respiratory Infections by Age: A graph depicting common bacteria found in the respiratory tracts of cystic fibrosis patients as a function of age.

Diagnosis and Symptoms

CF is most common among caucasians and can be diagnosed before birth by genetic testing, or by a sweat test in early childhood. Ultimately, lung transplantation is often necessary as CF worsens. The hallmark symptoms of cystic fibrosis are salty tasting skin, poor growth, poor weight gain despite a normal food intake, accumulation of thick, sticky mucus, frequent chest infections, and coughing or shortness of breath. Symptoms often appear in infancy and childhood, such as bowel obstruction due to meconium ileus in newborn babies.

Complications


Lung disease develops in CF as a result of clogging of the airways due to mucus build-up, decreased mucociliary clearance, and resulting inflammation. Inflammation and infection cause injury and structural changes to the lungs, leading to a variety of symptoms. In the early stages, incessant coughing, copious phlegm production, and decreased ability to exercise are common. Many of these symptoms occur when bacteria that normally inhabit the thick mucus grow out of control and cause pneumonia. In later stages, changes in the architecture of the lung, such as pathology in the major airways (bronchiectasis), further exacerbate difficulties in breathing.



Clinical Manifestations of Cystic Fibrosis: A diagram showing clinical manifestations of cystic fibrosis.

Treatment

Many CF patients are on one or more antibiotics at all times, even when healthy, to prophylactically suppress infection. Antibiotics are absolutely necessary whenever pneumonia is suspected or there has been a noticeable decline in lung function, and are usually chosen based on the results of a sputum analysis and the patient's past response. Several mechanical techniques are used to dislodge sputum and encourage its expectoration. In the hospital setting, chest physiotherapy (CPT) is utilized. As lung disease worsens, mechanical breathing support may become necessary. Bi-lateral lung transplantation often becomes necessary for individuals with cystic fibrosis as lung function and exercise tolerance declines.



A Deletion in the CFTR Gene is Responsible for CF: The ΔF508 human genome mutation is characterized by the deletion of three base pairs in the CFTR nucleotide sequence, causing the loss of the amino acid phenylalanine located at position 508.

Gene therapy has been explored as a potential cure for cystic fibrosis. Ideally, gene therapy attempts to place a normal copy of the CFTR gene into affected cells. Transferring the normal CFTR gene into the affected epithelium cells would result in the production of functional CFTR in all target cells, without adverse reactions or an inflammation response. Studies have shown that to prevent the lung manifestations of cystic fibrosis, only 5–10% the normal amount of CFTR gene expression is needed.

Finally, a number of small molecules that aim at compensating various mutations of the CFTR gene are under development. About

10% of CF cases result from a premature stop codon in the DNA, leading to early termination of protein synthesis and truncated proteins. One approach to combating a faulty receptor is to develop drugs that get the ribosome to overcome this premature stop codon and synthesize a full-length CFTR protein.

This page titled 29.12C: Cystic Fibrosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.12D: Hypoxia

Hypoxia, or hypoxiation, is inadequate oxygen supply to the body as a whole (generalized hypoxia) or a region of the body (tissue hypoxia).

Learning Objectives

- Differentiate between hypoxia and hypoxemia

Key Points

- A mismatch between oxygen supply and its demand at the cellular level may result in a hypoxic condition.
- Hypoxia in which there is complete deprivation of oxygen supply is referred to as anoxia.
- Hypoxia occurs in healthy people when they ascend to high altitude, where it causes altitude sickness leading to high altitude pulmonary edema (HAPE) and high altitude cerebral edema (HACE).
- Hypoxia also occurs in healthy individuals when breathing mixtures of gases with a low oxygen content.
- Hypoxia is a serious consequence of preterm birth in the neonate.
- In humans, hypoxia is detected by chemoreceptors in the carotid body, which override the signals from central chemoreceptors in the hypothalamus, increasing pO_2 despite a falling pCO_2.

Key Terms

- **hyperventilation**: The state of breathing faster or deeper than necessary.
- **hypoxemia**: An abnormal deficiency in the concentration of oxygen in the blood, be it the partial pressure of oxygen (mm Hg), the content of oxygen (ml oxygen per dl of blood) or the percent saturation of the blood's hemoglobin, singly or in combination.
- **anoxia**: A condition in which tissues are severely or totally deprived of oxygen, severe hypoxia.

Hypoxia, or hypoxiation, is a pathological condition in which the body as a whole (generalized hypoxia) or a region of the body (tissue hypoxia) is deprived of adequate oxygen supply. Variations in arterial oxygen concentrations can be part of the normal physiology, for example, during strenuous physical exercise. A mismatch between oxygen supply and its demand at the cellular level may result in a hypoxic condition. Hypoxia in which there is complete deprivation of oxygen supply is referred to as anoxia.

Hypoxia vs. Hypoxemia

Hypoxia differs from hypoxemia in that in the latter the oxygen concentration within the arterial blood is abnormally low. It is possible to experience hypoxia and have a low oxygen content (e.g., due to anemia) but maintain high oxygen partial pressure (pO_2). Incorrect use of these terms can easily lead to confusion, especially as hypoxemia is among the causes of hypoxia (in hypoxemic hypoxia). Generalized hypoxia occurs in healthy people when they ascend to high altitude where it causes altitude sickness leading to potentially fatal complications: high altitude pulmonary edema (HAPE) and high altitude cerebral edema (HACE). Hypoxia also occurs in healthy individuals when breathing mixtures of gases with a low oxygen content, e.g., while diving underwater especially when using closed-circuit rebreather systems that control the amount of oxygen in the supplied air. A mild and non-damaging intermittent hypoxia is used intentionally during altitude trainings to develop an athletic performance adaptation at both the systemic and cellular level.

Hypoxia is also a serious consequence of preterm birth in the neonate. The main cause for this is that the lungs of the human fetus are among the last organs to develop during pregnancy. To assist the lungs to distribute oxygenated blood throughout the body, infants at risk of hypoxia are often placed inside an incubator capable of providing continuous positive airway pressure (also known as a humidicrib). In humans, hypoxia is detected by chemoreceptors in the carotid body. This response does not control ventilation rate at normal pO_2, but below normal the activity of neurons innervating these receptors increases dramatically, so much so to override the signals from central chemoreceptors in the hypothalamus, increasing pO_2 despite a falling pCO_2.

Hypoxemic hypoxia is a generalized hypoxia, with an inadequate supply of oxygen to the body as a whole. The term “hypoxemic hypoxia” specifies hypoxia caused by low partial pressure of oxygen in arterial blood. In the other causes of hypoxia that follow, the partial pressure of oxygen in arterial blood is normal. Hypoxemic hypoxia may be due to hypoventilation or inadequate pulmonary minute ventilation (e.g., respiratory arrest or by drugs such as opiates).

Symptoms of Hypoxia



Cyanosis during hypoxia: Cyanosis in the hand of patient suffering from hypoxia.

The symptoms of generalized hypoxia depend on its severity and acceleration of onset. In the case of altitude sickness, where hypoxia develops gradually, the symptoms include headaches, fatigue, shortness of breath, a feeling of euphoria, and nausea. In severe hypoxia, or hypoxia of very rapid onset, changes in levels of consciousness, seizures, coma, priapism, and death occur. Severe hypoxia induces a blue discoloration of the skin, called cyanosis. Because hemoglobin is a darker red when it is not bound to oxygen (deoxyhemoglobin), as opposed to the rich red color that it has when bound to oxygen (oxyhemoglobin), when seen through the skin it has an increased tendency to reflect blue light back to the eye. In cases where the oxygen is displaced by another molecule, such as carbon monoxide, the skin may appear “cherry red” instead of cyanotic.

Body’s Response to Hypoxia

In most tissues of the body, the response to hypoxia is vasodilation. By widening the blood vessels, the tissue allows greater perfusion. By contrast, in the lungs, the response to hypoxia is vasoconstriction. This is known as “Hypoxic pulmonary vasoconstriction” or “HPV.” To counter the effects of high-altitude diseases, the body must return arterial pO_2 toward normal. Acclimatization, the means by which the body adapts to higher altitudes, only partially restores pO_2 to standard levels. Hyperventilation, the body’s most common response to high-altitude conditions, increases alveolar pO_2 by raising the depth and rate of breathing.

However, while pO_2 does improve with hyperventilation, it does not return to normal. Studies of miners and astronomers working at 3,000 meters and above show improved alveolar pO_2 with full acclimatization, yet the pO_2 level remains equal to or even below the threshold for continuous oxygen therapy for patients with chronic obstructive pulmonary disease (COPD). In addition, there are complications involved with acclimatization. Polycythemia, in which the body increases the number of red blood cells in circulation, thickens the blood, raising the danger that the heart can’t pump it.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/beta-2-agonist**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- wheezing. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/wheezing**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- asthma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/asthma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma before-after-en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Asthma_before-after-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***Public Domain: No Known Copyright***
- Acute respiratory distress syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acute_respiratory_distress_syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/acute-respiratory-distress-syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma before-after-en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Asthma_before-after-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***Public Domain: No Known Copyright***
- Acute respiratory distress syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acute_respiratory_distress_syndrome**. **License**: ***Public Domain: No Known Copyright***
- Cystic Fibrosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cystic_Fibrosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autosomal recessive. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/autosomal%20recessive**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Boundless. **Provided by**: Boundless Learning. **Located at**: **www.boundless.com//physiology/definition/cftr-gene**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cystic fibrosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cystic_fibrosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma before-after-en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Asthma_before-after-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***Public Domain: No Known Copyright***
- Acute respiratory distress syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acute_respiratory_distress_syndrome**. **License**: ***Public Domain: No Known Copyright***
- Cystic Fibrosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cystic_Fibrosis**. **License**: ***Public Domain: No Known Copyright***
- CFTRdel4. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:CFTRdel4.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cystic Fibrosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cystic_Fibrosis**. **License**: ***Public Domain: No Known Copyright***
- Hypoxia (medical). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hypoxia_(medical)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoxemia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoxemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- anoxia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/anoxia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hyperventilation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hyperventilation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Asthma before-after-en. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Asthma_before-after-en.svg**. **License**: ***Public Domain: No Known Copyright***
- Asthma. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Asthma**. **License**: ***Public Domain: No Known Copyright***
- Acute respiratory distress syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acute_respiratory_distress_syndrome**. **License**: ***Public Domain: No Known Copyright***
- Cystic Fibrosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cystic_Fibrosis**. **License**: ***Public Domain: No***

Known Copyright
- CFTRdel4. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:CFTRdel4.gif. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cystic Fibrosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cystic_Fibrosis. **License**: *Public Domain: No Known Copyright*
- Hypoxia (medical). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoxia_(medical). **License**: *Public Domain: No Known Copyright*

This page titled 29.12D: Hypoxia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.13: DISEASES AND DISORDERS OF THE DIGESTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.13A: Peritonitis
29.13B: Mumps
29.13C: Lactose Intolerance
29.13D: Peptic Ulcer Disease

This page titled 29.13: Diseases and Disorders of the Digestive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.13A: Peritonitis

Peritonitis is an inflammation of the peritoneum, usually caused by an infectious organism that is introduced into the abdominal cavity.

Learning Objectives

- Describe the causes and prognosis of peritonitis


Key Points

- Perforation of part of the gastrointestinal tract is the most common cause of peritonitis.
- The main manifestations of peritonitis are acute abdominal pain, abdominal tenderness, and abdominal guarding.
- Depending on the severity of the patient's state, the management of peritonitis may include supportive measures, antibiotics, or surgery.
- If properly treated, typical cases of surgically correctable peritonitis (e.g., perforated peptic ulcer, appendicitis, and diverticulitis) have a mortality rate of less than 10%.

Key Terms

- **Spontaneous bacterial peritonitis (SBP)**: Spontaneous bacterial peritonitis (SBP) is the development of peritonitis (infection in the abdominal cavity) despite the absence of an obvious source for the infection.
- **peritoneal dialysis**: A treatment for patients with severe chronic kidney disease in which fluids are exchanged from the blood in the patient's peritoneum in the abdomen.
- **Blumberg sign**: Blumberg's sign is a sign that is elicited during physical examination in medicine and is indicative of peritonitis. A positive sign shows rebound tenderness, meaning that pressing a hand on the abdomen elicits less pain than releasing the hand abruptly, which will aggravate the pain, as the peritoneum snaps back into place.

Peritonitis is an inflammation of the peritoneum, the thin tissue that lines the inner wall of the abdomen and covers most of the abdominal organs. Peritonitis may be localized or generalized, and may result from infection (often due to rupture of a hollow organ as may occur in abdominal trauma or appendicitis) or from a non-infectious process.



Peritoneum: The peritoneum, colored in blue, is the serous membrane that forms the lining of the abdominal cavity. It covers

most of the intra-abdominal, supporting them and serving as a conduit for their blood and lymph vessels and nerves. Peritonitis is an inflammation of the peritoneum often causing abdominal pain and tenderness.

Symptoms and Complications

The main manifestations of peritonitis are acute abdominal pain, abdominal tenderness, and abdominal guarding, which are exacerbated by moving the peritoneum by coughing, flexing one's hips, or eliciting the Blumberg sign. Diffuse abdominal rigidity ("washboard abdomen") is often present, especially in generalized peritonitis. Other symptoms include fever, sinus tachycardia, and development of intestinal paralysis, which also causes nausea, vomiting, and bloating.

Complications of peritonitis include sequestration of fluid and electrolytes, as revealed by decreased central venous pressure, which may cause electrolyte disturbances, as well as significant hypovolemia, possibly leading to shock and acute renal failure. A peritoneal abscess may form and sepsis may develop, so blood cultures should be obtained.

Causes

Perforation of part of the gastrointestinal tract is the most common cause of peritonitis. Examples include perforation of the distal esophagus, of the stomach by a peptic ulcer or gastric carcinoma, of the duodenum, of the remaining intestine by appendicitis, diverticulitis, inflammatory bowel disease (IBD), intestinal infarction, intestinal strangulation, colorectal carcinoma, or of the gallbladder. Other possible reasons for perforation include abdominal trauma, ingestion of a sharp foreign body (such as a fish bone, toothpick or glass shard), perforation by an endoscope or catheter, and internal leakage of a colostomy site. In most cases of perforation of a hollow viscus, mixed bacteria are isolated from the infection; the most common agents include Gram-negative bacilli like E. coli and anaerobic bacteria. Fecal peritonitis results from the presence of feces in the peritoneal cavity. It can result from abdominal trauma and occurs if the large bowel is perforated during surgery.

Disruption of the peritoneum, even in the absence of perforation of a hollow viscus, may also cause infection simply by letting micro- organisms into the peritoneal cavity. Examples include trauma, surgical wound, continuous ambulatory peritoneal dialysis, and intra-peritoneal chemotherapy. Again, in most cases, mixed bacteria are isolated; the most common agents include cutaneous species such as Staphylococcus aureus and coagulase-negative staphylococci, but many others are possible, including fungi such as Candida.

Spontaneous bacterial peritonitis (SBP) is a peculiar form of peritonitis occurring in the absence of an obvious source of contamination. It occurs in patients with excess fluid in their abdomens, particularly in children. Intra-peritoneal dialysis predisposes a patient to peritoneal infection. Systemic infections such as tuberculosis may rarely have a peritoneal localization.

Sterile abdominal surgery, under normal circumstances, causes localized or minimal generalized peritonitis. However, peritonitis may also be caused by the rare case of a sterile foreign body inadvertently left in the abdomen after surgery (e.g., gauze, sponge). Much rarer non-infectious causes may include familial Mediterranean fever, porphyria, and systemic lupus erythematosus.

Treatment

Depending on the severity of the patient's state, the management of peritonitis may include supportive measures, antibiotics, or surgery. General supportive measures include vigorous intravenous rehydration and correction of electrolyte disturbances. Antibiotics are usually administered intravenously, but they may also be infused directly into the peritoneum. The empiric choice of broad-spectrum antibiotics often consist of multiple drugs, and should be targeted against the most likely agents, depending on the cause of peritonitis; once one or more agents are actually isolated, therapy will of course be targeted on them. Surgery is needed to perform a full exploration and lavage of the peritoneum, as well as to correct any gross anatomical damage that may have caused peritonitis. The exception is spontaneous bacterial peritonitis, which does not always benefit from surgery and may be treated with antibiotics in the first instance.

If properly treated, typical cases of surgically correctable peritonitis (e.g., perforated peptic ulcer, appendicitis, and diverticulitis) have a mortality rate of <10% in otherwise healthy patients, which rises to about 40% in the elderly, and/or in those with significant underlying illness, as well as in cases that present late (after 48 hours). If untreated, generalized peritonitis is almost always fatal.

This page titled 29.13A: Peritonitis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.13B: Mumps

Mumps was a common childhood viral disease, but widespread vaccination has now made it rare in developed countries.

Learning Objectives

- Analyze the cause, symptoms, and prevention of mumps

Key Points


- Mumps is a contagious disease that is spread from person to person through contact with respiratory secretions, such as saliva from an infected person. The common symptoms of mumps include inflammation of the salivary glands, pancreas, and testicles; fever; and headache.
- A physical examination confirms the presence of the swollen glands. Usually, the disease is diagnosed on clinical grounds and no confirmatory laboratory testing is needed.
- The most common preventative measure against mumps is a vaccination with a mumps vaccine. The vaccine may be given separately or as part of the routine MMR immunization vaccine which also protects against measles and rubella.
- Like many other viral illnesses, there is no specific treatment for mumps, other than supportive treatment. Death from mumps is very unusual. The disease is self-limiting, and general outcome is good. Known rare complications of mumps include infertility in men and profound hearing loss.

Key Terms

- **orchitis**: A painful inflammation of one or both testes.
- **salivary gland**: Any of several exocrine glands that produce saliva to break down carbohydrates in food enzymatically.
- **prodromal symptoms**: A prodrome is an early symptom (or set of symptoms) that might indicate the start of a disease before specific symptoms occur.
- **parotid gland**: Either of a pair of salivary glands located in front of, and below each ear in humans.

Mumps, also known as epidemic parotitis, was a common childhood viral disease caused by the mumps virus. Before the development of vaccination and the introduction of a vaccine in 1949, it was common worldwide, but now, outbreaks are largely confined to developed countries.


Symptoms



Child with mumps: This child with mumps displays the typical swelling of the salivary glands caused by the mumps virus.

The common symptoms of mumps include inflammation of the salivary glands, pancreas, and testicles; fever, and headache. Swelling of the salivary glands, specifically the parotid gland, is known as parotitis, and it occurs in 60–70% of infections and 95% of patients with symptoms. Parotitis causes swelling and local pain, particularly when chewing. It can occur on one side but is more common on both sides in about 90% of cases. Painful inflammation of the testicles in mumps in known as orchitis. Other symptoms of mumps can include dry mouth, sore face and/or ears and occasionally, in more serious cases, loss of voice. In addition, up to 20% of persons infected with the mumps virus do not show symptoms, so it is possible to be infected and spread the virus without knowing it. Fever and headache are prodromal symptoms of mumps, together with malaise and loss of appetite.

Causes



Mumps virus: This transmission electron micrograph (TEM) shows the ultrastructure of the mumps virus. It is a roughly spherical particle made up of layers of fatty lipids, large protein molecules, and nucleic acids. The virus is dotted with large protein “spikes” that enable it to gain entry to host cells. Inside lies a core of a single, long molecule of RNA wrapped up in protein that is released into the host cell.

Mumps is a contagious disease that is spread from person to person through contact with respiratory secretions, such as saliva from an infected person. When an infected person coughs or sneezes, the droplets aerosolize and can enter the eyes, nose, or mouth of another person. Mumps can also be spread by sharing food and drinks. The virus can survive on surfaces and then be spread after contact in a similar manner. A person infected with mumps is contagious from approximately six days before the onset of symptoms until about nine days after symptoms start. The incubation period can be anywhere from 14–25 days, but is more typically 16–18 days.

Diagnostics

A physical examination confirms the presence of the swollen glands. Usually, the disease is diagnosed on clinical grounds, and no confirmatory laboratory testing is needed. If there is uncertainty about the diagnosis, a test of saliva or blood may be carried out. An estimated 20–30% of cases are asymptomatic. As with any inflammation of the salivary glands, the level of amylase in the blood is often elevated.

Prevention

The most common preventative measure against mumps is a vaccination with a mumps vaccine. The vaccine may be given separately or as part of the routine MMR immunization vaccine which also protects against measles and rubella. The MMR vaccine is given at ages 12–15 months and then again at four to six years.

Treatment and Complications

Like many other viral illnesses, there is no specific treatment for mumps. Symptoms may be relieved by the application of intermittent ice or heat to the affected neck/testicular area and by the acetaminophen or ibuprofen for pain relief. Warm salt water gargles, soft foods, and extra fluids may also help relieve symptoms. Patients are advised to avoid acidic foods and beverages, since these stimulate the salivary glands, which can be painful.

Death from mumps is very unusual. The disease is self-limiting, and general outcome is good, even if other organs are involved. Known complications of mumps include:

- In teenage males and men, complications from orchitis such as infertility or sub-fertility are rare, but present.
- Spontaneous abortion in about 27% of cases during the first trimester of pregnancy.
- Mild forms of meningitis in up to 10% of cases.
- Profound hearing loss is very rare, but mumps was the leading cause of acquired deafness before the advent of the mumps vaccine.

After the illness, life-long immunity to mumps generally occurs; re-infection is possible but tends to be mild and atypical.

This page titled 29.13B: Mumps is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.13C: Lactose Intolerance

Lactose intolerance is the inability to digest lactose, a sugar found in milk, due to a lack of the enzyme lactase.

Learning Objectives

- Analyze lactose intolerance and tolerance

Key Points

- Symptoms of lactose intolerance include abdominal bloating and cramps, flatulence, diarrhea, nausea, borborygmi (rumbling stomach), and/or vomiting after consuming significant amounts of lactose. Individuals may be lactose intolerant to varying degrees, depending on the severity of these symptoms.
- Most mammals normally become lactose intolerant after weaning, but some human populations have developed lactase persistence, in which lactase production continues into adulthood. These populations (northern Europe, India, and a few groups in Africa) are dairying cultures.
- Lactase deficiency has a number of causes, and is therefore classified as one of three types: primary, secondary, or congenital. Primary lactase deficiency is genetic and normal as most adults worldwide do not maintain lactase production.
- Secondary/acquired/transient lactase deficiency is caused by an injury to the small intestine. Congenital lactase deficiency prevents lactase expression from birth, making nourishment from breast milk impossible.
- Lactose intolerance is not an allergy, because it is not an immune response, but rather a problem with digestion caused by lactase deficiency. Milk allergy is a separate condition, with distinct symptoms that occur when the presence of milk proteins trigger an immune reaction.
- Primary lactase deficiency is genetic, secondary/acquired/transient lactase deficiency is caused by an injury to the small intestine, and congenital lactase deficiency prevents lactase expression from birth.


Key Terms

- **lactose**: The disaccharide sugar of milk and dairy products, C12H22O11, (a product of glucose and galactose) used as a food and in medicinal compounds.
- **lactose intolerance**: The inability to fully metabolize lactose.
- **lactase**: A β-galactosidase enzyme that is involved in the hydrolysis of the disaccharide lactose into constituent galactose and glucose monomers.

EXAMPLES

Those who are lactose intolerant can be more tolerant of traditionally made yogurt than of milk because it contains lactase produced by the bacterial cultures used to make the yogurt. Frozen yogurt, if cultured similarly to its unfrozen counterpart, will contain similarly reduced lactose levels. However, many commercial brands contain milk solids, increasing the lactose content.

Lactose intolerance, also called lactase deficiency and hypolactasia, is the inability to digest lactose, a sugar found in milk and, to a lesser extent, in milk-derived dairy products. Lactose intolerant individuals have insufficient levels of lactase, the enzyme that metabolizes lactose into glucose and galactose, in their digestive system. In most cases, this causes symptoms such as abdominal bloating and cramps, flatulence, diarrhea, nausea, borborygmi (rumbling stomach), and/or vomiting after consuming significant amounts of lactose. Some studies in the U.S. and elsewhere suggest that milk consumption by lactose intolerant individuals may be a significant cause of irritable bowel syndrome.



Lactose: Lactose, a disaccharide of β-D-galactose and β-D-glucose, is normally broken down by the enzyme lactase into its component monosaccharides. Individuals who suffer from lactose intolerance have insufficient levels of lactase to break down the lactose in milk and dairy products.

Most mammals normally become lactose intolerant after weaning, but some human populations have developed lactase persistence, in which lactase production continues into adulthood. It is estimated that 75% of adults worldwide show some decrease in lactase activity during adulthood. The frequency of decreased lactase activity ranges from 5% in northern Europe through 71% for Sicily to more than 90% in some African and Asian countries. This distribution is now thought to have been caused by recent natural selection favoring lactase persistent individuals in cultures that rely on dairy products. While it was first thought that this would mean that populations in Europe, India, and Africa that had high frequencies of lactase persistence shared a single mutation, it has now been shown that lactase persistence is caused by several independently occurring mutations.

Causes

Primary lactase deficiency is genetic, only affects adults, and is caused by the absence of a lactase persistence allele. It is the most common cause of lactose intolerance as a majority of the world's population lacks these alleles. Secondary, acquired, or transient lactase deficiency is caused by an injury to the small intestine, usually during infancy, from acute gastroenteritis, diarrhea, chemotherapy, intestinal parasites, or other environmental causes. Congenital lactase deficiency is a very rare, autosomal recessive genetic disorder that prevents lactase expression from birth. It is particularly common in Finland. People with congenital lactase deficiency who are unable to digest lactose from birth are unable to digest breast milk.

Effects on Nutrition

While dairy products can be a significant source of nutrients in some societies, there is no evidence that lactose intolerance has any adverse impact on nutrition where consumption is the norm among adults. Congenital lactase deficiency (CLD), where the production of lactase is inhibited from birth, can be dangerous in any society because of infants' nutritional reliance on breast milk during their first months. Before the 20th century, babies born with CLD were not expected to survive, but these death rates can now be lowered using soybean-derived infant formulas and manufactured lactose-free dairy products. Beyond infancy, individuals affected by CLD usually have the same nutritional concerns as any lactose-intolerant adult.

Lactose cannot be directly absorbed through the wall of the small intestine into the bloodstream so, in the absence of lactase, it passes intact into the colon. Bacteria in the colon are able to metabolize lactose and the resulting fermentation produces copious amounts of gas (a mixture of hydrogen, carbon dioxide, and methane) that causes the various abdominal symptoms. The unabsorbed sugars and fermentation products also raise the osmotic pressure of the colon, resulting in an increased flow of water into the bowels (diarrhea).

Assessment and Treatment of Lactose Intolerance

To assess lactose intolerance, intestinal function is challenged by ingesting more dairy products than can be readily digested. Clinical symptoms typically appear within 30 minutes, but may take up to two hours, depending on other foods and activities. Substantial variability in response (symptoms of nausea, cramping, bloating, diarrhea, and flatulence) is to be expected, as the extent and severity of lactose intolerance varies among individuals.

It is important to distinguish lactose intolerance from milk allergy, an abnormal immune response (usually) to milk proteins. This may be done in diagnosis by giving lactose-free milk, producing no symptoms in the case of lactose intolerance, but the same reaction as to normal milk if it is a milk allergy. An intermediate result might suggest that the person has both conditions. Since lactose intolerance is the normal state for most adults worldwide, it is not considered a disease and a medical diagnosis is not normally required, although tests are available if confirmation is necessary.

Lactose intolerance is not considered a condition that requires treatment in societies where the diet contains relatively little dairy. However, those living among societies that are largely lactose-tolerant may find lactose intolerance troublesome. Although there are still no methodologies to reinstate lactase production, some individuals have reported that their intolerance varies over time, depending on health status and pregnancy. About 44% of lactose intolerant women regain the ability to digest lactose during pregnancy. This might be caused by slow intestinal transit and intestinal flora changes during pregnancy.

This page titled 29.13C: Lactose Intolerance is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.13D: Peptic Ulcer Disease

A peptic ulcer, also known as peptic ulcer disease, is an erosion in the wall of the stomach, duodenum, or esophagus.

Learning Objectives

- List the causes of and treatments for peptic ulcer disease

Key Points


- 70–90% of peptic ulcers are associated with Helicobacter pylori, a spiral-shaped bacterium that lives in the acidic environment of the stomach.
- Diagnosis is mainly established based on the characteristic symptoms. Stomach pain is usually the first signal of a peptic ulcer.
- Treatment of H. pylori usually leads to clearing of infection, relief of symptoms, and eventual healing of ulcers.
- Gas in the peritoneal cavity, shown on an erect chest x-ray or supine lateral abdominal x-ray, is an omen of perforated peptic ulcer disease, which requires emergency surgery.
- Gastrointestinal bleeding is the most common complication. Sudden large bleeding can be life-threatening. It occurs when the ulcer erodes one of the blood vessels, such as the gastroduodenal artery.
- Most bleeding ulcers require endoscopy urgently to stop bleeding with cautery, injection, or clipping.
- During the active phase, the base of the ulcer shows 4 zones: inflammatory exudate, fibrinoid necrosis, granulation tissue and fibrous tissue.
- A gastric peptic ulcer is a mucosal defect which penetrates the muscularis mucosae and muscularis propria, produced by acid-pepsin aggression.
- Ulcers are not purely an infectious disease and that psychological factors do play a significant role.
- Diagnosis is mainly established based on the characteristic symptoms. Stomach pain is usually the first signal of a peptic ulcer.
- Gastric ulcers are most often localized on the lesser curvature of the stomach.
- Gas in the peritoneal cavity, shown on an erect chest X-ray or supine lateral abdominal X-ray, is an omen of perforated peptic ulcer disease.
- Gastrointestinal bleeding is the most common complication. Sudden large bleeding can be life-threatening. It occurs when the ulcer erodes one of the blood vessels, such as the gastroduodenal artery.
- Burning or gnawing feeling in the stomach area lasting between 30 minutes and 3 hours commonly accompanies ulcers.
- Typical ulcers tend to heal and recur and as a result the pain may occur for few days and weeks and then wane or disappear.

Key Terms

- **prostaglandin**: Any of a group of naturally occurring lipids derived from the C20 acid prostanoic acid; they have a number of physiological functions and may be considered to be hormones.
- **NSAID**: Any drug of the non-steroidal anti-inflammatory class used as a pain reliever.
- **gastrin**: A hormone that stimulates the production of gastric acid in the stomach.
- **gastritis**: Inflammation of the lining of the stomach, characterized by nausea, loss of appetite, and upper abdominal discomfort or pain.

A peptic ulcer, also known as peptic ulcer disease, is an erosion in the wall of the stomach, duodenum, or esophagus. As many as 70–90% of such ulcers are associated with Helicobacter pylori, a spiral-shaped bacterium that lives in the acidic environment of the stomach. Ulcers can also be caused or worsened by drugs such as aspirin, ibuprofen, and other NSAIDs.

Symptoms



Deep gastric ulcer: This image, acquired via endoscope, shows a deep gastric ulcer.

Symptoms of a peptic ulcer include abdominal pain, classically near the stomach with severity relating to mealtimes, about three hours after eating a meal; bloating and abdominal fullness; nausea; copious vomiting; loss of appetite and weight loss; vomiting of blood; and melena, which are tarry, foul-smelling feces due to oxidized iron from hemoglobin. Rarely, an ulcer can lead to a gastric or duodenal perforation, which leads to acute peritonitis. This is extremely serious and requires immediate surgery.

Causes


A major causative factor of ulcers is chronic inflammation due to Helicobacter pylori that colonizes the mucosa. The immune system is unable to clear the infection, despite the appearance of antibodies. Thus, the bacterium can cause a chronic active gastritis, resulting in a defect in the regulation of gastrin production by that part of the stomach. Gastrin secretion can either be increased, or as in most cases, decreased, resulting in a too basic or too acidic stomach environment, respectively. A decrease in acid can promote H. pylori growth and an increase in acid can contribute to the erosion of the mucosa and therefore ulcer formation.

Another major cause is the use of NSAIDs. The gastric mucosa protects itself from gastric acid with a layer of mucus, the secretion of which is stimulated by certain prostaglandins. NSAIDs block the function of cyclooxygenase 1 (cox-1), which is essential for the production of these prostaglandins.

Researchers also continue to look at stress as a possible cause, or at least complication, in the development of ulcers. There is debate as to whether psychological stress can influence the development of peptic ulcers. Burns and head trauma, however, can lead to physiologic stress ulcers, which are reported in many patients who are on mechanical ventilation.

Diagnosis

The diagnosis is mainly established based on the characteristic symptoms. Stomach pain is usually the first signal of a peptic ulcer. In some cases, doctors may treat ulcers without diagnosing them with specific tests and observe whether the symptoms resolve, this indicating that their primary diagnosis was accurate.



Gastric ulcer: This endoscopic image shows a gastric ulcer, which upon biopsy was shown to be gastric cancer.

Confirmation of the diagnosis is made with the help of tests such as endoscopies or barium contrast x-rays. The tests are typically ordered if the symptoms do not resolve after a few weeks of treatment. Tests are also given when first appear in a person who is over age 45 or who has other symptoms such as weight loss, because stomach cancer can cause similar symptoms. Also, when severe ulcers resist treatment, particularly if a person has several ulcers or the ulcers are in unusual places, a doctor may suspect an underlying condition that causes the stomach to overproduce acid.

An esophagogastroduodenoscopy (EGD), a form of endoscopy, also known as a gastroscopy, is carried out on patients in whom a peptic ulcer is suspected. By direct visual identification, the location and severity of an ulcer can be described. Moreover, if no ulcer is present, EGD can often provide an alternative diagnosis.

If a peptic ulcer perforates, air will leak from the inside of the gastrointestinal tract (which always contains some air) to the peritoneal cavity (which normally never contains air). This leads to “free gas” within the peritoneal cavity. If the patient stands erect, as when having a chest x-ray, the gas will float to a position underneath the diaphragm. Therefore, gas in the peritoneal cavity, shown on an erect chest x-ray or supine lateral abdominal x-ray, is an omen of perforated peptic ulcer disease.

Treatment



Benign gastric ulcer: This gastric ulcer was found in tissue removed during a gastrectomy.

Younger patients with ulcer-like symptoms are often treated with antacids. The ability of antacids to neutralize acidity by increasing the pH or blocking the secretion of acid by gastric cells is critical in reducing acidity in the stomach. Patients who are taking NSAIDs may also be prescribed a prostaglandin analogue in order to help prevent peptic ulcers by replacing the prostaglandins whose formation is blocked by NSAID use.

When H. pylori infection is present, the most effective treatments are combinations of two antibiotics, such as Clarithromycin, Amoxicillin, Tetracycline, and Metronidazole; and one proton pump inhibitor, sometimes in combination with antacids. In complicated, treatment-resistant cases, three antibiotics may be used together with a proton pump inhibitor. Treatment of H. pylori usually leads to clearing of infection, relief of symptoms and eventual healing of ulcers. Recurrence of infection can occur and retreatment may be required, if necessary with other antibiotics.

Perforated peptic ulcer is a surgical emergency and requires surgical repair of the perforation. Most bleeding ulcers require endoscopy urgently to stop bleeding with cautery, injection, or clipping.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Peritonitis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peritonitis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Blumberg sign. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Blumberg%20sign**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spontaneous bacterial peritonitis (SBP). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spontan...onitis%20(SBP)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peritoneal dialysis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/peritoneal_dialysis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1040. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1040.png**. **License**: ***Public Domain: No Known Copyright***
- Mumps. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Mumps**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- prodromal symptoms. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/prodromal%20symptoms**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- orchitis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/orchitis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- parotid gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/parotid_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- salivary gland. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/salivary_gland**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1040. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1040.png**. **License**: ***Public Domain: No Known Copyright***
- Mumps virus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mumps_virus.jpg**. **License**: ***Public Domain: No Known Copyright***
- Mumps PHIL 130 lores. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mumps_PHIL_130_lores.jpg**. **License**: ***Public Domain: No Known Copyright***
- Lactose Intolerance. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Lactose_Intolerance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactase. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lactase**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactose. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lactose**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lactose intolerance. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lactose_intolerance**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1040. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1040.png**. **License**: ***Public Domain: No***

Known Copyright
- Mumps virus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mumps_virus.jpg**. **License**: ***Public Domain: No Known Copyright***
- Mumps PHIL 130 lores. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mumps_PHIL_130_lores.jpg**. **License**: ***Public Domain: No Known Copyright***
- Lactose Haworth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lactose_Haworth.svg**. **License**: ***Public Domain: No Known Copyright***
- Peptic ulcer disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Peptic_ulcer_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gastrin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gastrin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- prostaglandin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/prostaglandin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- NSAID. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/NSAID**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- gastritis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/gastritis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gray1040. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gray1040.png**. **License**: ***Public Domain: No Known Copyright***
- Mumps virus. **Provided by**: Wikipedia. **Located at**: **http://en.Wikipedia.org/wiki/File:Mumps_virus.jpg**. **License**: ***Public Domain: No Known Copyright***
- Mumps PHIL 130 lores. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Mumps_PHIL_130_lores.jpg**. **License**: ***Public Domain: No Known Copyright***
- Lactose Haworth. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Lactose_Haworth.svg**. **License**: ***Public Domain: No Known Copyright***
- Deep gastric ulcer. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Deep_gastric_ulcer.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Benign gastric ulcer 1. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Benign_gastric_ulcer_1.jpg**. **License**: ***Public Domain: No Known Copyright***
- Gastric ulcer 3. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Gastric_ulcer_3.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 29.13D: Peptic Ulcer Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.14: DISEASES AND DISORDERS OF THE METABOLISM

- Topic hierarchy

29.14A: Phenylketonuria (PKU)

29.14B: Galactosemia and Glycogen Storage Disease


This page titled 29.14: Diseases and Disorders of the Metabolism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.14A: Phenylketonuria (PKU)

Learning Objectives

- Describe the factors involved in phenylketonuria (PKU)

Phenylketonuria (PKU) is an autosomal recessive metabolic genetic disorder characterized by a mutation in the gene for the hepatic enzyme phenylalanine hydroxylase (PAH), rendering it nonfunctional. Other non-PAH mutations can also cause PKU. This is an example of non-allelic genetic heterogeneity. The PAH gene is located on chromosome 12 in the bands 12q22-q24.1. More than 400 disease-causing mutations have been found in the PAH gene. PAH deficiency causes a spectrum of disorders, including classic phenylketonuria and hyperphenylalaninemia (a less severe accumulation of phenylalanine). Because PKU is an autosomal recessive genetic disorder, both parents must have at least one mutated allele of the PAH gene. The child must inherit both mutated alleles, one from each parent. Therefore, it is possible for a parent with the disease to have a child without it if the other parent possesses one functional allele of the gene for PAH. Yet, a child from two parents with PKU will inherit two mutated alleles every time, and therefore will also inherit the disease.



Inheritance of PKU: Phenylketonuria is inherited in an autosomal recessive fashion.

PAH enzyme is necessary for the metabolism of the amino acid phenylalanine (Phe) to the amino acid tyrosine. When PAH activity is reduced, phenylalanine accumulates. Excessive phenylalanine can be metabolized into phenylketones through the minor route, a transaminase pathway with glutamate. Metabolites include phenylacetate, phenylpyruvate, and phenethylamine. Elevated levels of phenylalanine in the blood and detection of phenylketones in the urine is diagnostic.

Phenylalanine is a large, neutral amino acid (LNAA). LNAAs compete for transport across the blood–brain barrier (BBB) via the large neutral amino acid transporter (LNAAT). If phenylalanine is in excess in the blood, it will saturate the transporter. Excessive levels of phenylalanine tend to decrease the levels of other LNAAs in the brain. However, as these amino acids are necessary for protein and neurotransmitter synthesis, Phe buildup hinders the development of the brain, causing mental retardation.

Treatment

The mainstream treatment for classic PKU patients is a strict PHE-restricted diet supplemented by a medical formula containing amino acids and other nutrients. In the United States, the current recommendation is that the PKU diet should be maintained for life. Patients who are diagnosed early and maintain a strict diet can have a normal life span with normal mental development. However, recent research suggests that neurocognitive, psychosocial, quality of life, growth, nutrition, and bone pathology are slightly suboptimal.

Detection

PKU is commonly included in the newborn screening panel of most countries, with varied detection techniques. Most babies in developed countries are screened for PKU soon after birth. Screening for PKU is done with bacterial inhibition assay (Guthrie test), immunoassays using fluorometric or photometric detection, or amino acid measurement using tandem mass spectrometry (MS/MS). Measurements done using MS/MS determine the concentration of Phe and the ratio of Phe to tyrosine, both of which will be elevated in PKU.

Incidence

A rarer form of hyperphenylalaninemia occurs when PAH is normal, but there is a defect in the biosynthesis or recycling of the cofactor tetrahydrobiopterin (BH_4) by the patient. This cofactor is necessary for proper activity of the enzyme. The coenzyme (called biopterin) can be supplemented as treatment. Those who suffer from PKU must be supplemented with tyrosine to account for PAH deficiency in converting phenylalanine to tyrosine sufficiently. Dihydrobiopterin reductase activity is to replenish quinonoid-dihydrobiopterin back into its tetrahydrobiopterin form, which is an important cofactor in many metabolic reactions in amino acid metabolism. Those with this deficiency may produce sufficient levels of PAH, but since tetrahydrobiopterin is a cofactor for PAH activity, deficient dihydrobiopterin reductase renders any PAH enzyme non-functional. Tetrahydrobiopterin is also a cofactor in the production of L-DOPA from tyrosine and 5-hydroxy-l-tryptophan from tryptophan, which must also be supplemented as treatment in addition to the supplements for classical PKU. Levels of dopamine can be used to distinguish between these two types. Low levels of dopamine lead to high levels of prolactin. By contrast, in classic PKU, prolactin levels would be relatively normal. Tetrahydrobiopterin deficiency can be caused by defects in four different genes.

The mean incidence of PKU varies widely in different human populations. Caucasians are affected at a rate of 1 in 10,000. Turkey has the highest documented rate in the world, with 1 in 2,600 births, while countries such as Finland and Japan have extremely low rates with fewer than one case of PKU in 100,000 births.

Key Points

- PKU is an autosomal recessive genetic disorder that occurs due to a mutation in phenylalanine hydroxylase gene, which makes this enzyme non-functional.
- Phenylalanine hydroxylase is the enzyme that is necessary for the conversion of the amino acid phenylalanine (Phe) to the amino acid tyrosine.
- Phenylalanine is a large, neutral amino acid (LNAA), which can compete for transport across the blood – brain barrier (BBB) via the large neutral amino acid transporter (LNAAT).
- Untreated PKU can lead to mental retardation, seizures, and other serious medical problems. Therefore, PKU is commonly included in the newborn screening panel of most countries, with varied detection techniques.
- The mainstream treatment for classic PKU patients is a strict PHE-restricted diet supplemented by a medical formula containing amino acids and other nutrients.
- PKU is an autosomal recessive genetic disorder.

Key Terms

- **autosomal recessive**: Both copies of the gene in each cell have mutations. The parents of an individual with an **autosomal recessive** condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.
- **Phenylketonuria**: An autosomal recessive metabolic genetic disorder characterized by a mutation in the gene for the hepatic enzyme phenylalanine hydroxylase, rendering it nonfunctional.
- **phenylalanine**: An essential amino acid $C_9H_{11}NO_2$ found in most animal proteins; it is essential for growth; the inability to metabolize it leads to phenylketonuria; it is a constituent of aspartame.

This page titled 29.14A: Phenylketonuria (PKU) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.14B: Galactosemia and Glycogen Storage Disease

Learning Objectives

- Differentiate between galactosemia and glycogen storage disease

Galactosemia (British Galactosaemia) is a rare genetic metabolic disorder that affects an individual's ability to metabolize the sugar galactose properly. Although the sugar and lactose metabolizes to galactose, galactosemia is not related to and should not be confused with lactose intolerance. Galactosemia follows an autosomal recessive mode of inheritance that confers a deficiency in an enzyme responsible for adequate galactose degradation. Its incidence is about one per 60,000 births for Caucasians. In other populations the incidence rate differs.




Figure $29.14B.1$: Galactose: Galactosemia is caused by the inability to metabolize galactose, shown here.

Lactose in food (such as dairy products) is broken down by the enzyme lactase into glucose and galactose. In individuals with galactosemia, the enzymes needed for further metabolism of galactose are severely diminished or missing entirely, leading to toxic levels of galactose 1-phosphate in various tissues as in the case of classic galactosemia, resulting in hepatomegaly (an enlarged liver), cirrhosis, renal failure, cataracts, brain damage, and ovarian failure. Without treatment, mortality in infants with galactosemia is about 75%.

Diagnosis

Infants are routinely screened for galactosemia in the United States. Infants affected by galactosemia typically present with symptoms of lethargy, vomiting, diarrhea, failure to thrive, and jaundice. If the family of the baby has a history of galactosemia, doctors can test prior to birth by taking a sample of fluid from around the fetus or from the placenta.

Detection of the disorder through newborn screening (NBS) does not depend on protein or lactose ingestion, and, therefore, it should be identified on the first specimen unless the infant has been transfused. A specimen should be taken prior to transfusion. The enzyme is prone to damage if analysis of the sample is delayed or exposed to high temperatures.

Treatment

The only treatment for classic galactosemia is eliminating lactose and galactose from the diet. Even with an early diagnosis and a restricted diet, however, some individuals with galactosemia experience long-term complications such as speech difficulties, learning disabilities, neurological impairment (e.g., tremors, etc.), and ovarian failure in females. Infants with classic galactosemia cannot be breast-fed due to lactose in human breast milk and are usually fed a soy-based formula.

Galactosemia is sometimes confused with lactose intolerance, but galactosemia is a more serious condition. Lactose intolerant individuals have an acquired or inherited shortage of the enzyme lactase, and experience abdominal pains after ingesting dairy

products, but no long-term effects. In contrast, a galactosemic individual who consumes galactose can cause permanent damage to their bodies.

Glycogen Storage Disease

Glycogen storage disease (GSD, also glycogenosis and dextrinosis) is the result of defects in the processing of glycogen synthesis or breakdown within muscles, liver, and other cell types. GSD has two classes of cause: genetic and acquired. Genetic GSD is caused by any inborn error of metabolism (genetically defective enzymes) involved in these processes. In livestock, acquired GSD is caused by intoxication with the alkaloid castanospermine. Overall, according to a study in British Columbia, approximately 2.3 children per 100,000 births (one in 43,000) have some form of glycogen storage disease. In the United States, they are estimated to occur in one per 20,000–25,000 births. A Dutch study estimated it to be one in 40,000.

There are 11 distinct diseases that are commonly considered to be glycogen storage diseases (some previously thought to be distinct have been reclassified).

Key Points

- Galactosemia follows is a rare genetic metabolic disorder that affects an individual's ability to metabolize the sugar galactose properly.
- Infants affected by galactosemia typically present with symptoms of lethargy, vomiting, diarrhea, failure to thrive, and jaundice.
- A galactosemia test is a blood test (from the heel of the infant) or urine test that checks for three enzymes that are needed to change galactose sugar that is found in milk and milk products into glucose, a sugar that your body uses for energy.
- The only treatment for classic galactosemia is eliminating lactose and galactose from the diet.
- Glycogen storage disease (GSD, also glycogenosis and dextrinosis) is the result of defects in the processing of glycogen synthesis or breakdown within muscles, liver, and other cell types.
- The only treatment for classic galactosemia is eliminating lactose and galactose from the diet.
- There are eleven (11) distinct diseases that are commonly considered to be glycogen storage diseases.

Key Terms

- **enzyme**: A globular protein that catalyses a biological chemical reaction.
- **glycogen**: A polysaccharide that is the main form of carbohydrate storage in animals; converted to glucose as needed.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Phenylketonuria. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Phenylketonuria**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autosomal recessive. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/autosomal%20recessive**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Phenylketonuria. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Phenylketonuria**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- phenylalanine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/phenylalanine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Phenylketonuria. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Phenylketonuria**. **License**: ***Public Domain: No Known Copyright***
- enzyme. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/enzyme**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Galactosemia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Galactosemia**. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- Glycogen storage disease. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glycogen_storage_disease. **License**: *CC BY-SA: Attribution-ShareAlike*
- glycogen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/glycogen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Phenylketonuria. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Phenylketonuria. **License**: *Public Domain: No Known Copyright*
- Galactosemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Galactosemia. **License**: *Public Domain: No Known Copyright*

This page titled 29.14B: Galactosemia and Glycogen Storage Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.15: DISEASES AND DISORDERS OF THE URINARY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.15A: Renal Disease and Failure
29.15B: Nephroptosis (Floating Kidney)
29.15C: Polycystic Kidney Disease

This page titled 29.15: Diseases and Disorders of the Urinary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.15A: Renal Disease and Failure

Renal failure uremia is a syndrome of renal failure characterized by elevated levels of urea and creatinine in the blood.

Learning Objectives

- Distinguish between acute kidney injury and chronic kidney disease

Key Points

- The type of renal failure is determined by the trend in the serum creatinine.
- In renal failure, there may be problems with increased fluid in the body (leading to swelling), increased acid levels, raised levels of potassium, decreased levels of calcium, increased levels of phosphate, and in later stages, anemia.
- The two main forms are acute kidney injury, which is often reversible with adequate treatment, and chronic kidney disease, which is often not reversible. In both cases, there is usually an underlying cause.

Key Terms

- **renal calculus**: A solid concretion or crystal aggregation formed in the kidneys from dietary minerals in the urine; a kidney stone.
- **creatinine**: A heterocyclic amine 2-amino-1-methyl-5H-imidazol-4-one formed by the metabolism of creatine.
- **renal failure uremia**: Uremia is a term used to loosely describe the illness accompanying kidney failure (also called renal failure), in particular the nitrogenous waste products associated with the failure of this organ.

EXAMPLES

Before dialysis or kidney transplants were widely available treatments, patients with kidney failure had progressive uremic poisoning. These days, most patients have some early symptoms of uremia as their kidney function declines, but prompt dialysis treatments or transplantation avoids most symptoms.

Renal failure (also kidney failure or renal insufficiency) is a medical condition in which the kidneys fail to adequately filter waste products from the blood. The two main forms are acute kidney injury, which is often reversible with adequate treatment, and chronic kidney disease, which is often not reversible. In both cases, there is usually an underlying cause.

Diagnosis


Renal failure is mainly determined by a decrease in the glomerular filtration rate, which is the rate at which blood is filtered in the glomeruli of the kidney. This is detected by a decrease in or absence of urine production or determination of waste products (creatinine or urea) in the blood. Depending on the cause, hematuria (blood loss in the urine) and proteinuria (protein loss in the urine) may be noted.

In renal failure, there may be problems with increased fluid in the body (leading to swelling), increased acid levels, raised levels of potassium, decreased levels of calcium, increased levels of phosphate, and in later stages, anemia. Bone health may also be affected. Long-term kidney problems are associated with an increased risk of cardiovascular disease.

Categories of Renal Failure

Renal failure can be divided into two categories: acute kidney injury or chronic kidney disease. The type of renal failure is determined by the trend in the serum creatinine. Other factors that may help differentiate acute kidney injury from chronic kidney disease include anemia and the kidney size on ultrasound. Chronic kidney disease generally leads to anemia and small kidney

size.



Kidney Transplantation: Kidney transplantation requires a person to be at the end stage of renal failure.

Acute kidney injury (AKI), previously called acute renal failure (ARF), is a rapidly progressive loss of renal function, generally characterized by oliguria (decreased urine production, quantified as less than 400 mL per day in adults, less than 0.5 mL/kg/h in children or less than 1 mL/kg/h in infants); and fluid and electrolyte imbalance. AKI can result from a variety of causes, generally classified as prerenal, intrinsic, and postrenal. An underlying cause must be identified and treated to arrest the progress, and dialysis may be necessary to bridge the time gap required for treating these fundamental causes.

Chronic kidney disease (CKD) can also develop slowly and, initially, show few symptoms. CKD can be the long-term consequence of irreversible acute disease or part of a disease progression.

Acute kidney injuries can be present on top of chronic kidney disease, a condition called acute-on-chronic renal failure (AoCRF). The acute part of AoCRF may be reversible, and the goal of treatment, as with AKI, is to return the patient to baseline renal function, typically measured by serum creatinine. Like AKI, AoCRF can be difficult to distinguish from chronic kidney disease if the patient has not been monitored by a physician and no baseline (i.e., past) blood work is available for comparison.

Symptoms can vary from person to person. Someone in early stage kidney disease may not feel sick or notice symptoms as they occur. When kidneys fail to filter properly, waste accumulates in the blood and the body, a condition called azotemia. Very low levels of azotaemia may produce few, if any, symptoms. If the disease progresses, symptoms become noticeable (if the failure is of sufficient degree to cause symptoms). Renal failure accompanied by noticeable symptoms is termed uraemia.

Renal Failure Uremia

Renal failure uremia is a syndrome of renal failure that includes elevated blood urea and creatinine levels. Acute renal failure can be reversed if diagnosed early. Acute renal failure can be caused by severe hypotension or severe glomerular disease. Diagnostic tests include BUN and plasma creatinine level tests. It is considered to be chronic renal failure if the decline of renal function is to less than 25%.

This page titled 29.15A: Renal Disease and Failure is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.15B: Nephroptosis (Floating Kidney)

Nephroptosis is an abnormal condition in which the kidney drops down into the pelvis when the patient stands up.

Learning Objectives

- Describe nephroptosis or floating kidney

Key Points


- It is more common in women than men and it is asymptomatic in most patients.
- It can be characterized by violent attacks of colicky flank pain, nausea, chills, hypertension, hematuria and proteinuria.
- Nephropexy was performed in the past to stabilize the kidney, but presently surgery is not recommended in asymptomatic patients.

Key Terms

- **hematuria**: The presence of blood in the urine.
- **nephroptosis**: Nephroptosis (also called floating kidney or renal ptosis) is an abnormal condition in which the kidney drops down into the pelvis when the patient stands up and it is more common in women than in men.

Nephroptosis, also called floating kidney or renal ptosis, is an abnormal condition in which the kidney drops down into the pelvis when the patient stands up. It is more common in women than in men. It has been one of the most controversial conditions among doctors in both its diagnosis and its treatments.

It is believed to result from deficiency of supporting perirenal fasciae. The renal fascia is a layer of connective tissue encapsulating the kidneys and the suprarenal glands. The deeper layers below the renal fascia are, in order, the adipose capsule of the kidney (or perirenal fat), the renal capsule and finally the parenchyma of the renal cortex. The spaces about the kidney are typically divided into three compartments: the perinephric space and the anterior and posterior pararenal spaces.



Renal fascia: Transverse section, showing the relations of the capsule of the kidney.

- Anterior attachment: Passes anterior to the kidney, renal vessels, abdominal aorta, and inferior vena cava and fuses with the

anterior layer of the renal fascia of the opposite kidney.
- Posterior attachment: Fuses with the psoas fascia and side of the body of the vertebrae.
- Superior attachment: The anterior and posterior layers fuse at the upper pole of the kidney and then split to enclose the suprarenal gland. At the upper part of the suprarenal gland they again fuse to form the suspensory ligament of the suprarenal gland and fuse with the diaphragmatic fascia.
- Inferior attachment: The layers don't fuse. The posterior layer descends downwards and fuses with the iliac fascia. The anterior layer blends with the connective tissue of the iliac fossa.

The anterior fascia and posterior fascia fuse laterally to form the lateroconal fascia which fuses with the fascia transversalis.

Symptoms and Diagnosis

Nephroptosis is asymptomatic in most patients. However, nephroptosis can be characterized by violent attacks of colicky flank pain, nausea, chills, hypertension, hematuria, and proteinuria.

Patients with symptomatic nephroptosis often complain of sharp pains that radiate into the groin. Many patients also suggest a weighing feeling on the abdomen. Pain is typically relieved by lying down. The attack of colic pain is called *Dittel's crisis* or *renal paroxysm*.

Diagnosis is contemplated based upon patient symptoms. Diagnosis is confirmed during intravenous urography, by obtaining erect and supine films. Nephropexy was performed in the past to stabilize the kidney, but presently surgery is not recommended in asymptomatic patients. Laparoscopic nephropexy has recently become available for selected symptomatic patients.

This page titled 29.15B: Nephroptosis (Floating Kidney) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.15C: Polycystic Kidney Disease

Polycystic kidney disease (PKD) is a cystic genetic disorder of the kidneys.

Learning Objectives

- Distinguish between autosomal dominant and autosomal recessive polycystic kidney disease

Key Points

- There are two types of PKD: autosomal dominant and autosomal recessive.
- PKD can damage the liver and the pancreas, and, in some rare instances, the heart and brain.
- Studies show that the incidence of autosomal recessive polycystic kidney disease (ARPKD) is 1:20,000 live births, and is typically identified in the first few weeks after birth.
- Autosomal dominant polycystic kidney disease (ADPKD) is the most common of all the hereditary cystic kidney diseases with an incidence of 1:1,000 to 2:1,000 live births.

Key Terms

- **autosomal recessive polycystic kidney disease**: The recessive form of polycystic kidney, called ARPKD (autosomal recessive polycystic kidney disease), is less common than autosomal dominant polycystic kidney
- **polycystic kidney disease**: Polycystic kidney disease (PKD or PCKD, also known as polycystic kidney syndrome) is a cystic genetic disorder of the kidneys.
- **autosomal dominant polycystic kidney disease**: Autosomal dominant polycystic kidney disease ("ADPKD," "autosomal dominant PKD," or "Adult-onset PKD") is an inherited systemic disorder that predominantly affects the kidneys, but may affect other organs including the liver, pancreas, brain, and arterial blood vessels
- **kidney**: An organ in the body that filters the blood, producing urine.


EXAMPLES

PKD treatment involves controlling symptoms and preventing complications. High blood pressure may be hard to control, but controlling it is the most important part of treatment. Treatment may include: blood pressure medicines, diuretics, and a low-salt diet

Polycystic kidney disease (PKD or PCKD, also known as polycystic kidney syndrome) is a cystic genetic disorder of the kidneys. There are two types of PKD: autosomal dominant polycystic kidney disease (ADPKD), and the less-common autosomal recessive polycystic kidney disease (ARPKD). PKD is characterized by the presence of multiple cysts (hence,"polycystic"), typically in both kidneys. The cysts are numerous and are fluid-filled, resulting in massive enlargement of the kidneys. The disease can also damage the liver, pancreas, and, in some rare cases, the heart and brain. The two major forms of polycystic kidney disease are distinguished by their patterns of inheritance. Polycystic kidney disease is one of the most common life-threatening genetic diseases, affecting an estimated 12.5 million people worldwide.

Autosomal dominant polycystic kidney disease (ADPKD) is the most common of all the hereditary cystic kidney diseases, with an incidence of 1:1,000 to 2:1,000 live births. Studies show that 10% of end-stage renal disease (ESRD) patients treated with hemodialysis in Europe and the U.S. were initially diagnosed and treated for ADPKD. ADPKD does not appear to demonstrate a preference for any particular ethnicity.

Characterization



Polycystic Kidney Disease: Polycystic kidney disease, or PKD, is a cystic genetic disorder of the kidneys

ADPKD is characterized by progressive cyst development and bilaterally enlarged kidneys with multiple cysts. There are three genetic mutations in the PKD-1, PKD-2, and PKD3 gene with similar phenotypical presentations. Gene PKD-1 is located on chromosome 16, and codes for a protein involved in regulation of cell cycle and intracellular calcium transport in epithelial cells; it is responsible for 85% of the cases of ADPKD. PKD-2, on chromosome 4, codes for a group of voltage-linked calcium channels. PKD3 recently appeared in research papers as a postulated 3rd gene. At this time, PKD3 has not been proven. Fewer than 10% of cases of ADPKD appear in non-ADPKD families.

Cyst formation begins in utero from any point along the nephron, although fewer than 5% of nephrons are thought to be involved. As the cysts accumulate fluid, they enlarge, separate entirely from the nephron, compress the neighboring renal parenchyma, and progressively compromise renal function.

Under the function of gene defect, epithelial cells of renal tubule turn into epithelial cells of cyst wall after phenotype change and begin to have the function of secreting cyst fluid, which leads to continuous cysts enlargement.

Studies show that the incidence of autosomal recessive polycystic kidney disease (ARPKD) is 1:20,000 live births, and is typically identified in the first few weeks after birth. Unfortunately, resulting hypoplasia results in a 30% death rate in neonates with ARPKD. In ARPKD, kidneys retain their shape, but are larger than the normal anatomical range with dilated collecting ducts from the medulla to the cortex.

Complications

The major extrarenal complications of ADPKD include cerebral aneurysms, hepatic cysts, pancreatic cysts, cardiac valve disease (especially mitral valve prolapse), colonic diverticula, and aortic root dilatation.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/The Urinary System. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_P..._of_the_Kidney**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Renal failure. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Renal_failure**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- renal failure uremia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/renal%20failure%20uremia**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- Renal calculus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Renal_calculus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- creatinine. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/creatinine**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Kidtransplant. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Kidtransplant.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Floating kidney. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Floating_kidney**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Nephroptosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Nephroptosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hematuria. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hematuria**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- nephroptosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/nephroptosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Kidtransplant. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Kidtransplant.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2/Gray1126.png**. **License**: ***Public Domain: No Known Copyright***
- Polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Polycys...kidney_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Polycys...kidney_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autosomal dominant polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/autosom...dney%20disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/polycys...dney%20disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- autosomal recessive polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/autosom...dney%20disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- kidney. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/kidney**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Kidtransplant. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Kidtransplant.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi...2/Gray1126.png**. **License**: ***Public Domain: No Known Copyright***
- Polycystic kidney disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Polycys...kidney_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 29.15C: Polycystic Kidney Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.16: BODY FLUID-RELATED DISEASES AND DISORDERS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

29.16A: Acidosis

29.16B: Alkalosis

29.16C: Edema

This page titled 29.16: Body Fluid-Related Diseases and Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.16A: Acidosis

Learning Objectives

- Describe acidosis and its medical implications

Acidosis is an increased acidity in the blood and other body tissue (i.e., an increased hydrogen ion concentration). If not further qualified, it usually refers to acidity of the blood plasma. Acidosis is said to occur when arterial pH falls below 7.35, while its counterpart (alkalosis) occurs at a pH over 7.45. Arterial blood gas analysis and other tests are required to determine the main causes.

The term “acidemia” describes the state of low blood pH, while acidosis is used to describe the processes leading to these states. Nevertheless, the terms are sometimes used interchangeably. The distinction may be relevant where a patient has factors causing both acidosis and alkalosis, wherein the relative severity of both determines whether the result is a high or a low pH.

The rate of cellular metabolic activity affects and, at the same time, is affected by the pH of the body fluids. In humans, the normal pH of arterial blood lies between 7.35 and 7.45. Blood pH values compatible with life in all mammals are limited to a pH range between 6.8 and 7.8. Changes in the pH of arterial blood (and therefore the extracellular fluid) outside this range result in irreversible cell damage.

Types of Acidosis

Metabolic acidosis may result from increased production of metabolic acids or disturbances in the ability to excrete acid via the kidneys. Renal acidosis is associated with an accumulation of urea and creatinine as well as metabolic acid residues of protein catabolism.

A rise in lactate out of proportion to the level of pyruvate is termed “excess lactate,” and may be an indicator of fermentation due to anaerobic metabolism occurring in muscle cells, as seen during strenuous exercise. Once oxygenation is restored, the acidosis clears quickly. Another example of increased production of acids occurs in starvation and diabetic acidosis. It is due to the accumulation of ketoacids (ketosis) and reflects a severe shift from glycolysis to lipolysis for energy needs.

Metabolic Acidosis

Metabolic acidosis is compensated for in the lungs, as increased exhalation of carbon dioxide promptly shifts the buffering equation to reduce metabolic acid. This is a result of stimulation to chemoreceptors, which increases alveolar ventilation, leading to respiratory compensation, otherwise known as Kussmaul breathing (a specific type of hyperventilation). Should this situation persist, the patient is at risk for exhaustion leading to respiratory failure.

The Henderson-Hasselbalch approximation is useful for calculating blood pH, because blood is a buffer solution. The amount of metabolic acid accumulating can also be quantitated by using buffer base deviation, a derivative estimate of the metabolic as opposed to the respiratory component. In hypovolemic shock for example, approximately 50% of the metabolic acid accumulation is lactic acid, which disappears as blood flow and oxygen debt are corrected.


Treatment of uncompensated metabolic acidosis is focused upon correcting the underlying problem. When metabolic acidosis is severe and can no longer be compensated for adequately by the lungs, neutralizing the acidosis with infusions of bicarbonate may be required.

Respiratory Acidosis

Respiratory acidosis results from a build-up of CO_2 in the blood (hypercapnia) due to hypoventilation. It is most often caused by pulmonary problems, although head injuries, drugs (especially anesthetics and sedatives), and brain tumors can cause this acidemia. Pneumothorax, emphysema, chronic bronchitis, asthma, severe pneumonia, and aspiration are among the most frequent causes. It can also occur as a compensatory response to chronic metabolic alkalosis.

One key to distinguish between respiratory and metabolic acidosis is that in respiratory acidosis, the CO_2 is increased while the bicarbonate is either normal (uncompensated) or increased (compensated). Compensation occurs if respiratory acidosis is present, and a chronic phase is entered with partial buffering of the acidosis through renal bicarbonate retention.

However, in cases where chronic illnesses that compromise pulmonary function persist, such as late-stage emphysema and certain types of muscular dystrophy, compensatory mechanisms will be unable to reverse this acidotic condition. As metabolic bicarbonate production becomes exhausted, and extraneous bicarbonate infusion can no longer reverse the extreme buildup of carbon dioxide associated with uncompensated respiratory acidosis, mechanical ventilation will usually be applied.



***Symptoms of Acidosis**: General symptoms of acidosis are described in relation to their location on the body. These symptoms usually accompany symptoms of another primary defect (respiratory or metabolic).*

Key Points

- Acidemia is the state of low blood pH; acidosis is the process leading to the state of acidemia.
- The normal pH of arterial blood is between 7.35 and 7.45.
- Metabolic acidosis may occur as a result of increased metabolic acids, either from increased production of metabolic acids, or decreased excretion of acids by the kidneys. The lungs compensate for metabolic acidosis by increasing the exhalation of CO2.
- Blood pH is calculated using the Henderson-Hasselbach equation, appropriate here because blood is a buffer solution.
- Respiratory acidosis occurs from an increased concentration of carbon dioxide in the blood, usually due to hypoventilation resulting from pulmonary problems. In respiratory acidosis, carbon dioxide concentration rises, but bicarbonate is normal or increased.

Key Terms

- **Henderson-Hasselbalch equation**: In chemistry, the Henderson–Hasselbalch equation describes the derivation of pH as a measure of acidity (using pKa, the negative log of the acid dissociation constant) in biological and chemical systems. The equation is also useful for estimating the pH of a buffer solution and finding the equilibrium pH in acid-base reactions (it is widely used to calculate the isoelectric point of proteins).
- **respiratory acidosis**: Respiratory acidosis is a medical condition in which decreased ventilation (hypoventilation) causes increased blood carbon dioxide concentration and decreased pH (a condition generally called acidosis).
- **metabolic acidosis**: In medicine, metabolic acidosis is a condition that occurs when the body produces too much acid or when the kidneys are not removing enough acid from the body.

This page titled 29.16A: Acidosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.16B: Alkalosis

Alkalosis is the increased alkalinity of blood and other tissues, generally occurring when the blood pH is above 7.45.

Learning Objectives

- Describe the medical conditions caused by alkalosis

Key Points

- Alkalosis may be respiratory or metabolic.
- Respiratory alkalosis may be caused by hyperventilation, which causes a loss of carbon dioxide.
- Metabolic alkalosis may be caused by prolonged vomiting, which reduces the amount of hydrochloric acid in the stomach, diuretics, dehydration, endocrine disorders, or consumption of alkali.
- Metabolic alkalosis typically occurs with disturbed muscle function such as weakness, pain, cramps, and spasms, and may also cause hypocalcemia.
- Respiratory alkalosis may be acute or chronic. Acute respiratory alkalosis occurs rapidly, and the patient may lose consciousness and return to normal ventilation. Chronic respiratory alkalosis is maintained by metabolic compensation, where bicarbonate ion concentrations are reduced to compensate for decreased levels of carbon dioxide in the blood.
- Metabolic alkalosis may be chloride-responsive or chloride-resistant.

Key Terms

- **alkalosis**: An abnormally increased alkalinity in the blood.
- **metabolic alkalosis**: Metabolic alkalosis is a metabolic condition in which the pH of tissue is elevated beyond the normal range (7.35–7.45). This is the result of decreased hydrogen ion concentration, leading to increased bicarbonate, or alternatively a direct result of increased bicarbonate concentrations.
- **respiratory alkalosis**: Respiratory alkalosis is a medical condition in which increased respiration (hyperventilation) elevates the blood pH (a condition generally called alkalosis).

Alkalosis refers to a condition reducing hydrogen ion concentration of arterial blood plasma (alkalemia). Generally, alkalosis is said to occur when pH of the blood exceeds 7.45. The opposite condition is acidosis (when pH falls below 7.35).


Alkalosis can refer to respiratory alkalosis or metabolic alkalosis. The main cause of respiratory alkalosis is hyperventilation, resulting in a loss of carbon dioxide. Compensatory mechanisms for this would include increased dissociation of the carbonic acid buffering intermediate into hydrogen ions, and the related excretion of bicarbonate, both of which lower blood pH.

Metabolic Alkalosis

Metabolic alkalosis is a metabolic condition in which the pH of tissue is elevated beyond the normal range (7.35–7.45). This is the result of decreased hydrogen ion concentration, leading to increased bicarbonate or alternatively a direct result of increased bicarbonate concentrations. Metabolic alkalosis can be caused by prolonged vomiting, resulting in a loss of hydrochloric acid with the stomach content. Severe dehydration and the consumption of alkali are other causes. It can also be caused by administration of diuretics and endocrine disorders such as Cushing's syndrome. Compensatory mechanism for metabolic alkalosis involves slowed breathing by the lungs to increase serum carbon dioxide, a condition leaning toward respiratory acidosis. As respiratory acidosis often accompanies the compensation for metabolic alkalosis, and vice versa, a delicate balance is created between these two conditions.

Metabolic alkalosis is usually accompanied with hypokalemia, causing, e.g., muscular weakness, myalgia, and muscle cramps (owing to disturbed function of the skeletal muscles), and muscle spasms (from disturbed function of smooth muscles). It may also cause hypocalcemia. As the pH of blood increases, the protein in the blood becomes more ionized into anions. This causes the free calcium present in blood to bind strongly with protein. If severe, it may cause tetany (alkalotic tetany).

Figure 12



Davenport Diagram: A Davenport Diagram, as shown, is a graphical tool developed by Allan Jones Davenport that allows a clinician or investigator to describe blood bicarbonate concentrations and blood pH following a respiratory and/or metabolic acid-base disturbance.

This page titled 29.16B: Alkalosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.16C: Edema

Edema is an abnormal accumulation of fluid beneath the skin or in one or more cavities of the body that produces swelling.

Learning Objectives

- Evaluate factors involved in edema

Key Points

- During edema, there is either increased secretion of fluid into the interstitium or impaired removal of this fluid from the interstitium, resulting in fluid accumulation.
- Cutaneous edema is considered pitting edema when pressure applied to an area leaves an indentation that persists for some time after release.
- Edema may also occur as a result of cardiac failure due to the rise in hydrostatic pressure. A fall in osmotic pressure occurs in nephrotic syndrome and liver failure, and may cause edema.

Key Terms

- **edema**: An excessive accumulation of serum in tissue spaces or a body cavity
- **interstitium**: An interstitial space within a tissue or organ.
- **cutaneous edema**: Edema, formerly known as dropsy or hydropsy, is an abnormal accumulation of fluid beneath the skin or in one or more cavities of the body that produces swelling.

EXAMPLES

Causes of edema which are generalized to the whole body can cause edema in multiple organs and peripherally. For example, severe heart failure can cause pulmonary edema, pleural effusions, ascites, and peripheral edema.

Edema is an abnormal accumulation of fluid beneath the skin or in one or more cavities of the body that produces swelling. Edema is caused by increased secretion of fluid into the interstitium or impaired removal of this fluid.



Peripheral pitting edema: Peripheral pitting edema results from water retention. It can be caused by systemic diseases, pregnancy in some women, and as a result of heart failure or varicose veins.

Cutaneous edema is referred to as pitting when, after pressure is applied to a small area, the indentation persists for some time after the release of the pressure. Peripheral pitting edema, as shown in, is the more common type, resulting from water retention. It can be caused by systemic diseases, pregnancy in some women, either directly or as a result of heart failure, or local conditions such as varicose veins, thrombophlebitis, insect bites, and dermatitis. Non-pitting edema is observed when the indentation does not persist. It is associated with such conditions as lymphedema, lipoedema, and myxedema.

Edema may also occur as a result of cardiac failure due to the rise in hydrostatic pressure. A fall in osmotic pressure occurs in nephrotic syndrome and liver failure, and may cause edema.

Causes

Causes of edema which are generalized to the whole body can cause edema in multiple organs and peripherally. For example, severe heart failure can cause pulmonary edema, pleural effusions, ascites, and peripheral edema.

As well as the previously mentioned conditions, edemas often occur during the late stages of pregnancy in some women. Edema is more common with those of a history of pulmonary problems or poor circulation, and may be affected by the presence of arthritis. Edemas that occur during pregnancy are usually found in the lower part of the leg, usually from the calf down.

Edema will occur in specific organs as part of inflammations, tendinitis, or pancreatitis, for instance. Certain organs develop edema through tissue specific mechanisms. Examples of edema in specific organs include cerebral edema, pulmonary edema, periorbital edema (eye puffiness), and cutaneous edema due to mosquito bites, spider bites, bee stings, or contact dermatitis). Another cutaneous form of edema is myxedema, which is caused by increased deposition of connective tissue. In myxedema edema is caused by an increased tendency of the tissue to hold water within its extracellular space. In myxedema this is because of an increase in hydrophilic carbohydrate-rich molecules (perhaps mostly hyaluronan) deposited in the tissue matrix.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- metabolic acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/metabolic%20acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Henderson-Hasselbalch equation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Henders...lch%20equation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- respiratory acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/respiratory%20acidosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Symptoms of acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sy...f_acidosis.png**. **License**: ***Public Domain: No Known Copyright***
- Metabolic alkalosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Metabolic_alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Respiratory alkalosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Respiratory_alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Alkalosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- alkalosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- respiratory alkalosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/respiratory%20alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- metabolic alkalosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/metabolic%20alkalosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Symptoms of acidosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sy...f_acidosis.png**. **License**: ***Public Domain: No Known Copyright***

- Davenport Fig 12. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Da...ort_Fig_12.jpg. **License**: *Public Domain: No Known Copyright*
- Edema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Edema. **License**: *CC BY-SA: Attribution-ShareAlike*
- interstitium. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/interstitium. **License**: *CC BY-SA: Attribution-ShareAlike*
- cutaneous edema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/cutaneous%20edema. **License**: *CC BY-SA: Attribution-ShareAlike*
- edema. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/edema. **License**: *CC BY-SA: Attribution-ShareAlike*
- Symptoms of acidosis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sy...f_acidosis.png. **License**: *Public Domain: No Known Copyright*
- Davenport Fig 12. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Da...ort_Fig_12.jpg. **License**: *Public Domain: No Known Copyright*
- Edema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Edema. **License**: *CC BY: Attribution*

This page titled 29.16C: Edema is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.17: DISEASES AND DISORDERS OF THE REPRODUCTIVE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.17A: Sexually Transmitted Diseases (STDs)

29.17B: Erectile Dysfunction and the Blue Pill

29.17C: Cervical Cancer

This page titled 29.17: Diseases and Disorders of the Reproductive System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.17A: Sexually Transmitted Diseases (STDs)

A sexually transmitted infection is passed between people during unprotected sexual intercourse with an infected partner.

Learning Objectives

- Describe sexually transmitted infections

Key Points

- Many STIs are transmitted through the mucous membranes of the genitals, rectum, mouth, and throat.
- Common sexually transmitted infections include: gonorrhea, syphillis, chlamydia, genital warts, genital herpes, the human papillomavirus (HPV), and human immunodeficiency virus (HIV).
- Healthcare professionals suggest safer sex, such as the use of condoms, as the most reliable way of decreasing the risk of contracting STIs during sexual activity, but any activity that involves the exchange of body fluids (sharing needles, blood transfusions, etc.) can lead to an STI.
- Chlamydia is an STD that is characterized by symptoms such as urethritis, vaginal discharge, abdominal pain, and fever.
- Genital warts occur in clusters in the genital area and are caused by HPV, a virus which has been implicated in certain types of cancer.
- Genital herpes is caused by the Herpes simplex virus and has symptoms such as lesions on the genitalia, pain, and itching.
- HIV, the virus that causes AIDS, is an STD that is spread via blood, semen, or vaginal fluid and ultimately causes the immune system to fail.

Key Terms

- **sexually transmitted diseases (STD)**: Sexually transmitted infections (STI), also referred to as sexually transmitted diseases (STD) and venereal diseases (VD), are illnesses that have a significant probability of transmission between humans by means of human sexual behavior, including vaginal intercourse, oral sex, and anal sex.



Halting the spread of STDs: U.S. propaganda poster targeted at World War II soldiers and sailors appealed to their patriotism in urging them to protect themselves. The text at the bottom of the poster reads,"You can't beat the Axis if you get VD." Images of women were used to catch the eye on many VD posters.

Sexually transmitted infections (STI), also referred to as sexually transmitted diseases (STD) and venereal diseases (VD), are illnesses that have a significant probability of transmission between humans by means of human sexual behavior, including vaginal intercourse, oral sex, and anal sex. While in the past, these illnesses have mostly been referred to as STDs or VD, in recent years the term sexually transmitted infections (STIs) has been preferred, as it has a broader range of meaning; a person may be infected, and may potentially infect others, without having a disease. Some STIs can also be transmitted via the use of IV drug needles after its use by an infected person, as well as through childbirth or breastfeeding. STIs have been well known for hundreds of years.

Transmission

As may be noted from the name, sexually transmitted infections are transmitted from one person to another by certain sexual activities, rather than being actually caused by those sexual activities. Bacteria, fungi, protozoa or viruses are still the causative agents. It is not possible to catch any STI from a sexual activity with a person who is not carrying a disease; conversely, a person who has an STI contracted it from contact (sexual or otherwise) with someone who had it, or his/her bodily fluids. Some STIs such as HIV can be transmitted from mother to child either during pregnancy or breastfeeding.

Prevention

Healthcare professionals suggest safer sex, such as the use of condoms, as the most reliable way of decreasing the risk of contracting STIs during sexual activity, but safer sex should by no means be considered an absolute safeguard. The transfer of, and exposure to, bodily fluids, such as blood transfusions and other blood products, shared injection needles, needle-stick injuries (when medical staff are inadvertently jabbed or pricked with needles during medical procedures), shared tattoo needles, and childbirth are other avenues of transmission. These different means put certain groups, such as medical workers, haemophiliacs, and drug users particularly at risk.

Common sexually transmitted infections include: gonorrhea, syphillis, chlamydia, genital warts, genital herpes, the human papillomavirus (HPV), and human immunodeficiency virus (HIV).

This page titled 29.17A: Sexually Transmitted Diseases (STDs) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.17B: Erectile Dysfunction and the Blue Pill

Erectile dysfunction (ED) is the inability to develop or maintain an erection during sexual intercourse.

Learning Objectives

- Describe the causes of and treatments for erectile dysfunction

Key Points


- A penile erection is managed by a reflex erection (obtained by touching the penis) and/or a psychogenic erection (achieved by erotic/emotional stimuli).
- There are many causes of ED, including neurogenic disorders, Parkinson's disease, aging, drugs, and psychological factors.
- Treatment for ED includes drugs such as Viagra (Sildenafil), Cialis (Tadalifil), exercise, and vacuum therapy.
- Psychological impotence is where erection or penetration fails due to thoughts or feelings (psychological reasons) rather than physical impossibility; this is somewhat less frequent but can often be helped.
- As can be understood from the mechanisms of a normal erection, impotence may develop due to hormonal deficiency, disorders of the neural system, lack of adequate penile blood supply, or psychological problems.

Key Terms

- **psychogenic erection**: Erection achieved by erotic or emotional stimuli
- **chronic periodontitis**: Chronic periodontitis is a common disease of the oral cavity consisting of chronic inflammation of the periodontal tissues, often caused by accumulation of profuse amounts of dental plaque.
- **Erectile dysfunction**: Erectile dysfunction (ED) is sexual dysfunction characterized by the inability to develop or maintain an erection of the penis during sexual performance.

Erectile dysfunction (ED) is sexual dysfunction characterized by the inability to develop or maintain an erection of the penis during sexual performance. A penile erection is the hydraulic effect of blood entering and being retained in sponge-like bodies within the penis. The process is often initiated as a result of sexual arousal, when signals are transmitted from the brain to nerves in the penis. Erectile dysfunction is indicated when an erection is difficult to produce. There are various circulatory causes, including alteration of the voltage-gated potassium channel, as in arsenic poisoning from drinking water. The most significant organic causes are cardiovascular disease and diabetes, neurological problems (for example, trauma from prostatectomy surgery), hormonal insufficiencies (hypogonadism), and drug side effects. Psychological impotence is where erection or penetration fails due to thoughts or feelings (psychological reasons) rather than physical impossibility; this is somewhat less frequent but can often be helped. Notably, there is a strong response to placebo treatment in psychological impotence.

Erectile Function



Corpus Cavernosum Urethrae now known as Corpus Spongiosum: The corpus cavernosum penis is one of a pair of sponge-like regions of erectile tissue which contain most of the blood in the penis during penile erection.

Penile erection is managed by two different mechanisms. The first one is the reflex erection, which is achieved by directly touching the penile shaft. The second is the psychogenic erection, which is achieved by erotic or emotional stimuli. The former uses the peripheral nerves and the lower parts of the spinal cord, whereas the latter uses the limbic system of the brain. In both circumstances, an intact neural system is required for a successful and complete erection. Stimulation of penile shaft by the nervous system leads to the secretion of nitric oxide (NO), which causes the relaxation of smooth muscles of corpora cavernosa (the main erectile tissue of penis), and subsequently penile erection. Additionally, adequate levels of testosterone (produced by the testes) and an intact pituitary gland are required for the development of a healthy erectile system. As can be understood from the mechanisms of a normal erection, impotence may develop due to hormonal deficiency, disorders of the neural system, lack of adequate penile blood supply, or psychological problems. Restriction of blood flow can arise from impaired endothelial function due to the usual causes associated with coronary artery disease, but can also be caused by prolonged exposure to bright light.

There are several possible causes of ED, which include the following: drugs, neurogenic disorders (generalized anxiety disorder, personality disorders or traits), psychological problems, negative feelings, aging (making it four times higher in men in their 60s than in men in their 40s), kidney failure, diseases such as diabetes and multiple sclerosis (MS), and lifestyle.

Iatrogenic (Medical) Causes

Surgical intervention for a number of different conditions may remove anatomical structures necessary for erection, damage nerves, or impair blood supply. Complete removal of the prostate gland or external beam radiotherapy of the gland are common causes of impotence; both are treatments for prostate cancer. A study in 2002 found that ED can also be associated with bicycling. The number of hours on a bike and/or the pressure on the penis from the saddle of an upright bicycle is directly related to erectile dysfunction A recent study suggests an epidemiological association between chronic periodontitis (periodontal inflammation) and erectile dysfunction, similar to the association between periodontitis and coronary heart diseases, as well as cerebrovascular diseases. In all the three conditions (erectile dysfunction, coronary heart disease, and cerebrovascular diseases), despite the epidemiological association with periodontitis, no causative connection has yet been proven.

Treatments

Besides treating the major underlying causes such as potassium deficiency or arsenic contamination in drinking water, the primary treatment for erectile dysfunction consists of a trial of PDE5 inhibitor drugs (the first of which is sildenafil or Viagra). In some cases, treatment can involve prostaglandin tablets in the urethra, injections into the penis, a penile prosthesis, a

penis pump, or vascular reconstructive surgery. Additional treatments for ED are varied and include a purpose-designed external vacuum pump used to attain erection, with a separate compression ring fitted to the penis to maintain it. These pumps should be distinguished from other penis pumps (supplied without compression rings) which, rather than being used for temporary treatment of impotence, are claimed to increase penis length if used frequently, or vibrate as an aid to masturbation. More drastically, inflatable or rigid penile implants may be fitted surgically. Medications carry risk of priapism.

The cyclic nucleotide phosphodiesterases constitutes a group of enzymes that catalyze the hydrolysis of the cyclic nucleotides cyclic AMP and cyclic GMP. They exist in different molecular forms and are unevenly distributed throughout the body. One of the forms of phosphodiesterase is termed PDE5. The prescription PDE5 inhibitors sildenafil (Viagra), vardenafil (Levitra), and tadalafil (Cialis) are prescription drugs which are taken orally. They work by blocking the action of PDE5, which causes cGMP to degrade.


This page titled 29.17B: Erectile Dysfunction and the Blue Pill is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.17C: Cervical Cancer

Learning Objectives

- List causes and methods of prevention in cervical cancer

Cervical cancer is the term for a malignant neoplasm arising from cells originating in the cervix uteri. One of the most common symptoms of cervical cancer is abnormal vaginal bleeding, but in some cases there may be no obvious symptoms until the cancer has progressed to an advanced stage. Treatment usually consists of surgery (including local excision) in early stages, and chemotherapy and/or radiotherapy in more advanced stages of the disease.



Squamous carcinoma of the cervix: This large squamous carcinoma (bottom of picture) has obliterated the cervix and invaded the lower uterine segment. The uterus also has a round leiomyoma up higher.

The cervix is the narrow portion of the uterus where it joins with the top of the vagina. Most cervical cancers are squamous cell carcinomas, arising in the squamous (flattened) epithelial cells that line the cervix. Adenocarcinoma, arising in glandular epithelial cells is the second most common type. Very rarely, cancer can arise in other types of cells in the cervix.

Symptoms

The early stages of cervical cancer may be completely asymptomatic. Vaginal bleeding, contact bleeding, or (rarely) a vaginal mass may indicate the presence of malignancy. Also, moderate pain during sexual intercourse and vaginal discharge are symptoms of cervical cancer. In advanced disease, metastases may be present in the abdomen, lungs or elsewhere.

Symptoms of advanced cervical cancer may include: loss of appetite, weight loss, fatigue, pelvic pain, back pain, leg pain, swollen legs, heavy bleeding from the vagina, bone fractures, and/or (rarely) leakage of urine or feces from the vagina (rarely).

Diagnosis

Cancer screening using the Pap smear can identify precancerous and potentially precancerous changes in cervical cells and tissue. Treatment of high-grade changes can prevent the development of cancer in many victims. In developed countries, the widespread use of cervical screening programs has reduced the incidence of invasive cervical cancer by 50% or more. While the pap smear is an effective screening test, confirmation of the diagnosis of cervical cancer or pre-cancer requires a biopsy of the

cervix.

HPV

Human papillomavirus (HPV) infection appears to be a necessary factor in the development of almost all cases (90+%) of cervical cancer. HPV vaccines effective against the two strains of this large family of viruses that currently cause approximately 70% of cases of cervical cancer have been licensed in the U.S., Canada, Australia, and the EU. Since the vaccines only cover some of the cancer causing ("high-risk") types of HPV, women should seek regular Pap smear screening, even after vaccination.

Infection with some types of human papilloma virus (HPV) is the greatest risk factor for cervical cancer, followed by smoking. Other risk factors include human immunodeficiency virus. Not all of the causes of cervical cancer are known, however, and several other contributing factors have been implicated.

Key Points

- Human papillomavirus (HPV) is a major cause of cervical cancer.
- Pap smears are used to identify precancerous tissue in the cervix.
- Aside from HPV, smoking and STDs are other major risk factors for cervical cancer.
- The early stages of cervical cancer may be completely asymptomatic, but symptoms of advanced stages may include loss of appetite, weight loss, fatigue, pain, swollen legs, and bleeding from the vagina.

Key Terms

- **cervical cancer**: A type of cancer of the cervix of the uterus which can afflict some female mammals including humans. The term usually does not refer to cancer of the neck or other parts of the anatomy which use the term cervix.
- **Pap smear**: The Papanicolaou test (also called Pap smear, Pap test, cervical smear, or smear test) is a screening test used to detect potentially pre-cancerous and cancerous processes in the endocervical canal (transformation zone) of the female reproductive system.
- **human papillomavirus**: A virus that affects humans, sometimes causing cervical or other cancer; it is sometimes classified as a sexually transmitted disease.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Syphilis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Syphilis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Gonorrhea. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Gonorrhea**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Genital warts. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Genital_warts**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- HIV. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/HIV**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sexually transmitted. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Sexually_transmitted**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Chlamydia infection. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Chlamydia_infection**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Genital herpes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Genital_herpes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- HPV. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/HPV**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sexually transmitted diseases (STD). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/sexuall...seases%20(STD)**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- SheMayLookCleanBut. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sh...okCleanBut.jpg. **License**: *Public Domain: No Known Copyright*
- Erectile dysfunction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Erectile_dysfunction. **License**: *CC BY-SA: Attribution-ShareAlike*
- psychogenic erection. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/psychogenic%20erection. **License**: *CC BY-SA: Attribution-ShareAlike*
- Erectile dysfunction. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Erectile%20dysfunction. **License**: *CC BY-SA: Attribution-ShareAlike*
- chronic periodontitis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/chronic%20periodontitis. **License**: *CC BY-SA: Attribution-ShareAlike*
- SheMayLookCleanBut. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sh...okCleanBut.jpg. **License**: *Public Domain: No Known Copyright*
- Gray1154. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1154.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cervical cancer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cervical_cancer. **License**: *CC BY-SA: Attribution-ShareAlike*
- Pap smear. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Pap%20smear. **License**: *CC BY-SA: Attribution-ShareAlike*
- human papillomavirus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/human_papillomavirus. **License**: *CC BY-SA: Attribution-ShareAlike*
- cervical cancer. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/cervical_cancer. **License**: *CC BY-SA: Attribution-ShareAlike*
- SheMayLookCleanBut. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:SheMayLookCleanBut.jpg. **License**: *Public Domain: No Known Copyright*
- Gray1154. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Gray1154.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Cervical cancer. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Cervical_cancer. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 29.17C: Cervical Cancer is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.18: PREGNANCY-RELATED DISORDERS

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

29.18A: Ectopic Pregnancy

29.18B: Infertility

This page titled 29.18: Pregnancy-Related Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.18A: Ectopic Pregnancy

An ectopic pregnancy is implantation and development of the embryo outside the uterus, typically in the fallopian tubes.

Learning Objectives

- Identify the factors involved in ectopic pregnancies

Key Points

- An ectopic pregnancy, or eccysis, is a complication of pregnancy in which the embryo implants outside the uterine cavity. With rare exceptions, ectopic pregnancies are not viable. Furthermore, they are dangerous for the mother, since internal haemorrhage is a life threatening complication.
- Most ectopic pregnancies occur in the Fallopian tube (called tubal pregnancies), but implantation can also occur in the cervix, ovaries, and abdomen. An ectopic pregnancy is a potential medical emergency, and, if not treated properly, can lead to death.
- If left untreated, about half of ectopic pregnancies will resolve without treatment and are called tubal abortions.
- A heterotopic pregnancy occurs when there are two fertilized eggs, one outside the uterus and the other inside.
- The mortality rate from ectopic pregnancies in Western countries is very low (less than one percent), but in underdeveloped countries, especially in Africa, the maternal death rate is extremely high and are a major cause of death among women of childbearing age.

Key Terms

- **heterotopic pregnancy**: A heterotopic pregnancy is a rare complication of pregnancy in which both extra-uterine (ectopic pregnancy) and intrauterine gestation occur simultaneously.
- **ectopic pregnancy**: A pregnancy in which the fertilized ovum is implanted in any tissue other than the uterine wall.


An ectopic pregnancy, or eccysis, is a complication of pregnancy in which the embryo implants outside the uterine cavity. With rare exceptions, ectopic pregnancies are not viable. Furthermore, they are dangerous for the mother, since internal haemorrhage is a life threatening complication that may result. Most ectopic pregnancies occur in the Fallopian tube (so-called tubal pregnancies), but implantation can also occur in the cervix, ovaries, and abdomen. An ectopic pregnancy is a potential medical emergency, and, if not treated properly, can lead to death.

In a normal pregnancy, the fertilized egg enters the uterus and settles into the uterine lining where it has plenty of room to divide and grow. About 1% of pregnancies are in an ectopic location with implantation not occurring inside of the womb; of these pregnancies, 98% occur in the Fallopian tubes.

Detection

Detection of ectopic pregnancy in early gestation has been achieved mainly due to enhanced diagnostic capability. Despite all these notable successes in diagnostics and detection techniques ectopic pregnancy remains a source of serious maternal morbidity and mortality worldwide, especially in countries with poor prenatal care.

Treatment



Ectopic pregnancy: schematic view of ectopic pregnancy

In a typical ectopic pregnancy, the embryo adheres to the lining of the fallopian tube and burrows into the tubal lining. Most commonly, this invades vessels and will cause bleeding. This intratubal bleeding hematosalpinx expels the implantation out of the tubal end as a tubal abortion. Tubal abortion is a common type of miscarriage. There is no inflammation of the tube in ectopic pregnancy. The pain is caused by prostaglandins released at the implantation site, and by free blood in the peritoneal cavity, which is a local irritant. Sometimes the bleeding might be heavy enough to threaten the health or life of the woman. Usually, this degree of bleeding is due to delay in diagnosis, but sometimes, especially if the implantation is in the proximal tube (just before it enters the uterus), it may invade into the nearby Sampson artery, causing heavy bleeding earlier than usual.

If left untreated, about half of ectopic pregnancies will resolve without treatment. These are the tubal abortions. The advent of methotrexate treatment for ectopic pregnancy has reduced the need for surgery; however, surgical intervention is still required in cases where the Fallopian tube has ruptured or is in danger of doing so. This intervention may be laparoscopic or through a larger incision, known as a laparotomy.

While a fetus of ectopic pregnancy is typically not viable, very rarely, a live baby has been delivered from an abdominal pregnancy. In such a situation the placenta sits on the intraabdominal organs or the peritoneum and has found sufficient blood supply. This is generally bowel or mesentery, but other sites, such as the renal (kidney), liver or hepatic (liver) artery or even aorta have been described.

In rare cases of ectopic pregnancy, there may be two fertilized eggs, one outside the uterus and the other inside. This is called a heterotopic pregnancy. Often the intrauterine pregnancy is discovered later than the ectopic, mainly because of the painful emergency nature of ectopic pregnancies. Since ectopic pregnancies are normally discovered and removed very early in the pregnancy, an ultrasound may not find the additional pregnancy inside the uterus. When hCG levels continue to rise after the removal of the ectopic pregnancy, there is the chance that a pregnancy inside the uterus is still viable. This is normally discovered through an ultrasound.

Risk Factors

There are a number of risk factors for ectopic pregnancies. However, in as many as one third to one half of ectopic pregnancies, no risk factors can be identified. Risk factors include: pelvic inflammatory disease, infertility, use of an intrauterine device (IUD), previous exposure to DES, tubal surgery, intrauterine surgery (e.g., D&C), smoking, previous ectopic pregnancy, and tubal ligation.

Prognosis

The prognosis in Western countries is very good; maternal death is rare. For instance, in the UK, between 2003 and 2005 there were 32,100 ectopic pregnancies resulting in 10 maternal deaths (meaning that 1 in 3,210 women with an ectopic pregnancy died). In the developing world, however, especially in Africa, the death rate is very high, and ectopic pregnancies are a major cause of death among women of childbearing age.

Fertility following ectopic pregnancy depends upon several factors, the most important of which is a prior history of infertility. The treatment choice, whether surgical or nonsurgical, also plays a role. For example, the rate of intrauterine pregnancy may be higher following methotrexate compared to surgical treatment. Rate of fertility may be better following salpingostomy than salpingectomy.

This page titled 29.18A: Ectopic Pregnancy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.18B: Infertility

Infertility, in both males and females, refers to the inability to contribute to conception.

Learning Objectives

- Evaluate the factors involved in infertility

Key Points

- Several clinical and epidemiological guidelines exist to determine infertility, often based on regular sexual intercourse in the absence of contraception without conception.
- Infertility can result from genetic factors or more general factors such as endocrine disorders (diabetes, thyroid disorders, etc.) and environmental factors.
- In the U.S., up to 20% of infertile couples have unexplained infertility. In these cases abnormalities are likely to be present but not detected by current methods.
- Medical treatment of infertility generally involves the use of fertility medication, medical device, surgery, or intrauterine insemination. If these treatments fail to achieve a full term pregnancy, in vitro fertilization or other assisted reproductive techniques (ART) can be used.

Key Terms

- **contraception**: The use of a device or procedure to prevent conception as a result of sexual activity.
- **infertility**: The inability to conceive children.
- **conception**: The fertilization of an ovum by a sperm to form a zygote.

Infertility primarily refers to the biological inability of a person to contribute to conception. Infertility may also refer to the state of a woman who is unable to carry a pregnancy to full term. There are many biological causes of infertility, some which may be bypassed with medical intervention.

Definitions of infertility differ, with demographers tending to define infertility as childlessness in a population of women of reproductive age, while the epidemiological definition is based on “trying for” or “time to” a pregnancy, generally in a population of women exposed to a probability of conception. Existing definitions of infertility lack uniformity, rendering comparisons in prevalence between countries or over time problematic, and therefore data estimating the prevalence of infertility cited by various sources differs significantly. A couple that has tried unsuccessfully to have a child after a certain period of time (often a short period, but definitions vary) is sometimes said to be subfertile, meaning less fertile than a typical couple. Both infertility and subfertility are defined as the inability to conceive after a certain period of time (the length of which vary), so often the two terms overlap. Couples with primary infertility have never been able to conceive, while, secondary infertility is difficulty conceiving after already having conceived (and either carried the pregnancy to term or had a miscarriage).

Causes

Factors that can cause male as well as female infertility are:

- Genetic factors
- Chromosomal defects
- Diabetes mellitus, thyroid disorders, adrenal disease
- Hypothalamic-pituitary factors
- Hyperprolactinemia
- Hypopituitarism
- Environmental factors
- Toxins such as glues, volatile organic solvents or silicones, physical agents, chemical dusts, and pesticides. Tobacco smokers

are 60% more likely to be infertile than non-smokers.

For a woman to conceive, certain things have to happen: intercourse must take place around the time when an egg is released from her ovary; the systems that produce eggs and sperm have to be working at optimum levels; and her hormones must be balanced. Some women are infertile because their ovaries do not mature and release eggs. In this case synthetic FSH by injection or Clomid (Clomiphene citrate) via a pill can be given to stimulate follicles to mature in the ovaries. Other factors that can affect a woman's chances of conceiving include being over- or underweight, or her age as female fertility declines sharply after the age of 35. Sometimes it can be a combination of factors, and sometimes a clear cause is never established.

The main cause of male infertility is low semen quality. Another possible cause is a low sperm count. In some cases, both the man and woman may be infertile or sub-fertile, and the couple's infertility arises from the combination of these conditions. In other cases, the cause is suspected to be immunological or genetic; it may be that each partner is independently fertile but the couple cannot conceive together without assistance.


Unexplained Infertility

In the U.S., up to 20% of infertile couples have unexplained infertility. In these cases abnormalities are likely to be present but not detected by current methods. Possible problems could be that the egg is not released at the optimum time for fertilization, that it may not enter the fallopian tube, sperm may not be able to reach the egg, fertilization may fail to occur, transport of the zygote may be disturbed, or implantation fails. It is increasingly recognized that egg quality is of critical importance and women of advanced maternal age have eggs of reduced capacity for normal and successful fertilization. Also, polymorphisms in folate pathway genes could be one reason for fertility complications in some women with unexplained infertility.

Treatment

Medical treatment of infertility generally involves the use of fertility medication, medical device, surgery, or a combination of the following. If the sperm are of good quality and the mechanics of the woman's reproductive structures are good (patent fallopian tubes, no adhesions or scarring), physicians may start by prescribing a course of ovarian stimulating medication. The physician may also suggest intrauterine insemination (IUI), in which the doctor introduces sperm into the uterus during ovulation, via a catheter. In these methods, fertilization occurs inside the body. If conservative medical treatments fail to achieve a full term pregnancy, the physician may suggest the patient undergo in vitro fertilization (IVF). IVF and related techniques are called assisted reproductive technology (ART) techniques.

ART techniques generally start with stimulating the ovaries to increase egg production. After stimulation, the physician surgically extracts one or more eggs from the ovary, and unites them with sperm in a laboratory setting, with the intent of producing one or more embryos. Fertilization takes place outside the body, and the fertilized egg is reinserted into the woman's reproductive tract, in a procedure called embryo transfer.



Causes of Infertility: Numerous factors may contribute to infertility.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- ectopic pregnancy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ectopic_pregnancy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ectopic pregnancy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ectopic_pregnancy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- heterotopic pregnancy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/heterotopic%20pregnancy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ectopic. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ectopic.png**. **License**: ***Public Domain: No Known Copyright***
- Infertility causes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:In...ity_causes.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- infertility. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/infertility**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- conception. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/conception**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- contraception. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/contraception**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ectopic. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ectopic.png**. **License**: ***Public Domain: No Known Copyright***
- Infertility causes. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Infertility_causes.png**. **License**: ***Public Domain: No Known Copyright***

This page titled 29.18B: Infertility is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.1: DISEASES, INJURIES, AND DISORDERS OF THE INTEGUMENTARY SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.1A: Burns
29.1B: Allergies
29.1C: Acne

This page titled 29.1: Diseases, Injuries, and Disorders of the Integumentary System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.1A: Burns

A burn is a type of injury to flesh caused by heat, electricity, chemicals, light, radiation, or friction.

Learning Objectives

- Classify burns by depth of injury

Key Points

- Most burns affect only the first two layers of the skin, the epidermis and dermis.
- Burns are classified by severity, surface area involvement, and depth of injury.
- Burns are also classified as major or minor depending on a combination of factors including age, thickness of burns, surface area of the body involved, and the presence of inhalation injury.
- First degree burns are limited to the epidermis. Second degree burns extend into the superficial papillary dermis. Third degree burns extend through the entire dermis. Fourth degree burns involve all the skin layers as well as the underlying muscle and bone.

Key Terms

- **avascular**: Lacking blood vessels.
- **burn**: A type of injury to flesh caused by heat, electricity, chemicals, light, radiation, or friction.
- **eschar**: A dry, dark scab or scar, especially as a result of burning.

A burn is a type of injury to flesh caused by heat, electricity, chemicals, light, radiation, or friction. Most burns affect only the first two layers of the skin, the epidermis and dermis. Burns are classified by severity, surface area involvement, and depth of injury. In the latter system, burns are classified as first, second, third, or fourth degree burns based on the depth of injury to the dermis. Burns are also classified as major or minor depending on a combination of factors including age, thickness of burns, surface area of the body involved, and the presence of inhalation injury.

Types of Burns


First degree burns are limited to the epidermis. Second degree burns extend into the superficial papillary dermis. Third degree burns extend through the entire dermis. Fourth degree burns involve all the skin layers as well as the underlying muscle and bone.



First Degree Burn: Mild sunburn is an example of a first degree burn.



Second Degree Burn: Serious sunburn is an example of a second degree burn.



Third Degree Burn: Third degree burns to the lower leg with partial tissue charring.



Fourth Degree Burn: Eight day old fourth degree burn on arc of foot caused by motorcycle muffler.

Consequences of Burns

While large burns can be fatal, modern treatments developed in the last 60 years have significantly improved the prognosis of such burns, especially in children and young adults. In the United States, approximately four out of every 100 people to suffer burns will die from their injuries. The majority of these fatalities occur either at the scene or on the way to hospital.

Infection is a major complication of burns. Infection is likely caused by impaired resistance from disruption of the skin's mechanical integrity by the burn and generalized immune suppression. As a deep burn starts to heal, the skin barrier is replaced by eschar. This moist, protein-rich avascular environment encourages microbial growth. Migration of immune cells is hampered by destruction of the vasculature (blood vessels), and there is a release of inflammatory intermediaries that impede the immune response. Eschar also restricts distribution of systemically administered antibiotics because of its avascularity.

This page titled 29.1A: Burns is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.1B: Allergies

An allergy is a hypersensitivity disorder of the immune system.

Learning Objectives

- Evaluate the types and treatments of allergic reactions

Key Points

- Allergy is one of four forms of hypersensitivity and is formally called type I (or immediate) hypersensitivity.
- Mild allergies like hay fever are very common in the human population and cause symptoms such as red eyes, itchiness, runny nose, eczema, hives, or an asthma attack.
- Treatments for allergies include avoiding known allergens, use of medications such as anti-histamines that specifically prevent allergic reactions, steroids that modify the immune system in general, and medications such as decongestants that reduce the symptoms.

Key Terms

- **anaphylaxis**: A severe and rapid systemic allergic reaction to an allergen, causing a constriction of the trachea, preventing breathing; anaphylactic shock.
- **allergen**: A substance which causes an allergic reaction.
- **patch test**: A patch test is a method used to determine if a specific substance causes allergic inflammation of the skin.

An allergy is a hypersensitivity disorder of the immune system. Allergic reactions occur when a person's immune system reacts to normally harmless substances in the environment. A substance that causes a reaction is called an allergen. These reactions are acquired, predictable, and rapid. Allergy is one of four forms of hypersensitivity and is formally called type I (or immediate) hypersensitivity. Allergic reactions are distinctive because of excessive activation of certain white blood cells called mast cells and basophils by a type of antibody called Immunoglobulin E (IgE). This reaction results in an inflammatory response that can range from uncomfortable to dangerous.

Mild allergies like hay fever are very common in the human population and cause symptoms such as red eyes, itchiness, runny nose, eczema, and hives. Allergies can play a major role in conditions such as asthma. In some people, severe allergies to environmental or dietary allergens or to medication may result in life-threatening reactions called anaphylaxis. Food allergies and reactions to the venom of stinging insects such as wasps and bees are often associated with these severe reactions.

Risk factors for allergy can be placed in two general categories: namely, host and environmental factors. Host factors include heredity, gender, race, and age, with heredity being by far the most significant. However, there have been recent increases in the incidence of allergic disorders that cannot be explained by genetic factors alone. Four major environmental candidates are alterations in exposure to infectious diseases during early childhood, environmental pollution, allergen levels, and dietary changes.

Allergy Tests



Allergy Testing: A patient receiving a skin allergy test.

A variety of tests exist to diagnose allergic conditions. Blood tests can be done to look for an allergen-specific IgE. Skin allergy testing is another method for medical diagnosis of allergies through the attempt to provoke a small, controlled, allergic response. A microscopic amount of an allergen (for example, tree pollen) is introduced to a patient's skin by means of either a prick test, which employs a needle or pin to break the skin's surface, or a patch test, where a patch containing the allergen is applied to the skin.

If an immuno-response is seen, it can be concluded that the patient has a hypersensitivity (or allergy) to that allergen. A response can take the the form of a rash (urticaria, also known as hives), or, worse—anaphylaxis (a serious reaction that can result in death). Further testing can then be performed in order to identify the specific allergen.

Allergy Treatments

Treatments for allergies include avoiding known allergens, use of medications such as antihistamines that specifically prevent allergic reactions, steroids that modify the immune system in general, and medications such as decongestants that reduce the symptoms. Many of these medications are taken by mouth; however, epinephrine, which is used to treat anaphylactic reactions, is injected. Immunotherapy employs the injection of allergens in order to gradually desensitize the body's response.

This page titled 29.1B: Allergies is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.1C: Acne

Acne, clinically known as acne vulgaris, is a common human skin disease affecting the skin of the face, upper parts of the chest, and back.

Learning Objectives


- Describe the development of acne

Key Points

- Acne occurs most commonly during adolescence, and often continues into adulthood. For most people, acne diminishes over time and tends to disappear—or at the very least decreases—by age 25.
- Acne develops as a result of blockages in hair follicles. Enlargement of sebaceous glands and an increase in sebum production occur with increased androgen production during puberty and directly before menstruation.
- Sebaceous glands become clogged with sebum, a naturally occurring skin oil, and dead skin cells. Bacteria becomes trapped in these clogged follicles, producing pus and inflammation as the immune system attempts to destroy the bacteria. This buildup of pus results in a pimple.
- The most common bacteria that causes acne is Propionibacterium acnes, an anaerobic bacteria that is part of the natural bacterial flora of the face.

Key Terms

- **androgen**: A steroid hormone that stimulates or controls the development and maintenance of masculine characteristics in vertebrates.
- **dermabrasion**: A cosmetic medical procedure in which the surface of the epidermis of the skin is removed by abrasion, typically to remove scarring or sun-damaged skin.
- **sebaceous gland**: A gland of the skin that secretes an oily substance, sebum, usually into a hair follicle near the surface of the skin.



Hair Follicle: The sebaceous gland produces oily secretions that can block the follicle, resulting in a pimple.

Acne, clinically known as acne vulgaris, is a common human skin disease affecting skin with the densest population of sebaceous

follicles such as the face, upper parts of the chest, and back. Acne affects 40 to 50 million people of all racial and ethnic groups in the United States. Acne occurs most commonly during adolescence, and often continues into adulthood. For most people, acne diminishes over time and tends to disappear—or at the very least decreases—by age 25.

Development and Effects of Acne

Acne develops as a result of blockages in hair follicles. Enlargement of sebaceous glands and an increase in sebum production occur with increased androgen production during puberty and directly before menstruation. Sebaceous glands become clogged with sebum, a naturally occurring skin oil, and dead skin cells. Bacteria becomes trapped in these clogged follicles, producing pus and inflammation as the immune system attempts to destroy the bacteria. This buildup of pus results in a pimple. The most common bacteria that causes acne is Propionibacterium acnes, an anaerobic bacteria that is part of the natural bacterial flora of the face.

As acne heals, it can cause scarring. Aside from scarring, its main effects are psychological, such as reduced self-esteem and in very extreme cases, depression or suicide. Acne usually appears during adolescence, when people already tend to be most socially insecure.



Acne: Acne of a 14-year-old male during puberty.

Treatments for Acne

Many different treatments exist for acne including benzoyl peroxide, antibiotics, retinoids, antiseborrheic medications, anti-androgen medications, hormonal treatments, salicylic acid, alpha hydroxy acid, azelaic acid, nicotinamide, and keratolytic soaps. They are believed to work in at least four different ways, including: normalizing shedding into the pore to prevent clogging, killing Propionibacterium acnes, anti-inflammatory effects, and hormonal manipulation. Laser treatment and dermabrasion can be used to reduce the appearance of scarring caused by acne.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Skin burn. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Skin_burn**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- avascular. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/avascular**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- burn. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/burn**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- eschar. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/eschar. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sunburned chest. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...38/Sunburn.jpg. **License**: *Public Domain: No Known Copyright*
- Foot with fourth-degr eeburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...egree-burn.jpg. **License**: *Public Domain: No Known Copyright*
- Second degree sunburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...n_Shoulder.jpg. **License**: *Public Domain: No Known Copyright*
- Leg with third-degree burn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Burns_foot.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Allergen. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Allergen. **License**: *CC BY-SA: Attribution-ShareAlike*
- Skin allergy test. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Skin_allergy_test. **License**: *CC BY-SA: Attribution-ShareAlike*
- Anaphylaxis. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Anaphylaxis%23Skin. **License**: *CC BY-SA: Attribution-ShareAlike*
- Allergy. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Allergy. **License**: *CC BY-SA: Attribution-ShareAlike*
- allergen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/allergen. **License**: *CC BY-SA: Attribution-ShareAlike*
- patch test. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/patch%20test. **License**: *CC BY-SA: Attribution-ShareAlike*
- anaphylaxis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/anaphylaxis. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sunburned chest. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...38/Sunburn.jpg. **License**: *Public Domain: No Known Copyright*
- Foot with fourth-degr eeburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...egree-burn.jpg. **License**: *Public Domain: No Known Copyright*
- Second degree sunburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...n_Shoulder.jpg. **License**: *Public Domain: No Known Copyright*
- Leg with third-degree burn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...Burns_foot.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Allergy prick test. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...gy_testing.jpg. **License**: *Public Domain: No Known Copyright*
- Acne. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Acne. **License**: *CC BY-SA: Attribution-ShareAlike*
- dermabrasion. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/dermabrasion. **License**: *CC BY-SA: Attribution-ShareAlike*
- androgen. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/androgen. **License**: *CC BY-SA: Attribution-ShareAlike*
- sebaceous gland. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/sebaceous_gland. **License**: *CC BY-SA: Attribution-ShareAlike*
- Sunburned chest. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/3/38/Sunburn.jpg. **License**: *Public Domain: No Known Copyright*
- Foot with fourth-degr eeburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/78/8-day-old-3rd-degree-burn.jpg. **License**: *Public Domain: No Known Copyright*
- Second degree sunburn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/0/01/Second_Degree_Burn_on_Shoulder.jpg. **License**: *Public Domain: No Known Copyright*
- Leg with third-degree burn. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/Wikipedia/commons/7/78/Burns_foot.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Allergy prick test. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...gy_testing.jpg. **License**: *Public Domain: No Known Copyright*
- HairFollicle. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:HairFollicle.png. **License**: *Public Domain: No Known Copyright*
- Forehead with Acne. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...kne-jugend.jpg. **License**: *Public Domain: No Known Copyright*

This page titled 29.1C: Acne is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.2: DISEASES AND DISORDERS OF THE SKELETAL SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.2A: Osteoporosis
29.2B: Paget's Disease

This page titled 29.2: Diseases and Disorders of the Skeletal System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.2A: Osteoporosis

Learning Objectives

- Differentiate among primary type 1, primary type 2, and secondary osteoporosis

Women who are thin are at higher risk of developing osteoporosis than are heavier women. Weight-bearing exercise such as walking helps maintain bone strength. Heavier women require their skeletons to bear more weight, so their bones are stronger than is the case for thin women. Ballet dancers are at particular risk, as are women who diet to be as thin as these dancers. Gwyneth Paltrow is an example of such a woman. In 2010 at age 37, she found out that she had osteopenia, the early stage of osteoporosis.



***Woman with Osteoporosis**: Hyperkyphosis, colloquially called a "dowager's hump," refers to an anteriorly concave curvature of the thoracic region of the spine.*

Osteoporosis is a disease of the bones that leads to an increased risk of fracture. In osteoporosis, the bone mineral density (BMD) is reduced, bone microarchitecture deteriorates, and the amount and variety of proteins in bone is altered. The three main mechanisms by which osteoporosis develop are an inadequate peak bone mass (the skeleton develops insufficient mass and strength during growth), excessive bone resorption, and inadequate formation of new bone during remodeling. An interplay of these three mechanisms underlies the development of fragile bone tissue. The form of osteoporosis most common in women after menopause is referred to as primary type 1 or postmenopausal osteoporosis. Primary type 2 osteoporosis or senile osteoporosis occurs after age 75 and is seen in both females and males at a ratio of 2:1. Finally, secondary osteoporosis may arise at any age and affect men and women equally. This form of osteoporosis results from chronic predisposing medical problems or disease, or prolonged use of medications such as glucocorticoids.

Treatment of Osteoporosis

Osteoporosis risks can be reduced with lifestyle changes and sometimes medication, or treatment may involve both. Lifestyle change includes diet, exercise, and preventing falls. Medication includes calcium, vitamin D, bisphosphonates, and several others. Fall-prevention advice includes exercise to tone deambulatory muscles, proprioception-improvement exercises, and equilibrium therapies. Exercise, with its anabolic effect, may at the same time stop or reverse osteoporosis.

Risks of Osteoporosis

Osteoporosis itself has no symptoms. Its main consequence is the increased risk of bone fractures. Osteoporotic fractures are those that occur in situations where healthy people would not normally break a bone. They are, therefore, regarded as fragility fractures. Typical fragility fractures occur in the vertebral column, rib, hip, and wrist.

Key Points

- The three main mechanisms by which osteoporosis develops are an inadequate peak bone mass (the skeleton develops insufficient mass and strength during growth), excessive bone resorption, and inadequate formation of new bone during remodeling.
- Primary type 1 osteoporosis is the most common type in post-menopausal women. Primary type 2 or senile osteoporosis is common in both men and women over age 75.
- Secondary osteoporosis may arise at any age as the result of predisposing medical problems or prolonged use of of medications such as glucocorticoids.
- Primary type 2 osteoporosis is twice as prevelent in females as in males, but secondary osteoporosis affects men and women equally.
- Risk of osteoporosis can be reduced by exercises such as walking and by taking vitamin D supplements.
- Fragility fractures are most common in the vertebrae, hip and wrist joints, and ribs.

Key Terms

- **peak bone mass**: Peak bone mass is the average bone mass of healthy, young adults.
- **fragility fractures**: Fragility fracture is a type of pathologic fracture that occurs as result of normal activities, such as a fall from standing height or less, suggesting a weakness of the skeleton.

This page titled 29.2A: Osteoporosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.2B: Paget's Disease

Paget's disease is a chronic bone disorder that causes affected bones to become large and misshapen.

Learning Objectives

- Differentiate Paget's disease from osteoporosis

Key Points

- Men are more commonly affected by Paget's disease than women (ratio of 3:2) and people are not usually diagnosed before the age of 40.
- The process of bone remodeling is disrupted so that excessive bone resorption is followed by chaotic bone growth.
- Patients require vitamin D and calcium supplementation, along with moderate exercise (in the sun) to maintain joint mobility, but there is no cure.

Key Terms

- **Paget's disease**: Paget's disease of bone is a chronic disorder that can result in enlarged and misshapen bones.

EXAMPLES

Along with osteomalacia, rickets, and osteoporosis, Paget's disease is associated with vitamin D deprivation. If one chooses to avoid exposing unprotected skin to the UVB radiation of the sun, then one must compensate for this lack by taking vitamin D supplements or risk the development of one of these diseases.

Paget's disease of bone is a chronic disorder that can result in enlarged and misshapen bones. The excessive breakdown and formation of bone tissue causes affected bone to weaken, resulting in pain, misshapen bones, fractures, and arthritis in the joints near the affected bones. Paget's disease typically is localized, affecting just one or a few bones, as opposed to osteoporosis, for example, which affects all the bones in the body. Although there is no cure for Paget's disease, medications (bisphosphonates and calcitonin) can help control the disorder and lessen pain and other symptoms.




Paget's disease of the bone: Paget's disease of bone is shown in the left pelvis. Notice the seeds in the prostate after brachytherapy.

Diagnosis of Paget's Disease

Paget's disease is rarely diagnosed in people less than 40 years of age. Men are more commonly affected than women (ratio of 3:2). Prevalence of Paget's disease ranges from 1.5 to 8.0 percent, depending on age and country of residence. Because early diagnosis and treatment is important, after age 40, siblings and children of someone with Paget's disease may wish to have an alkaline phosphatase blood test every two or three years. If the alkaline phosphatase level is above normal, other tests such as a bone-specific alkaline phosphatase test, bone scan, or X-ray can be performed.

Initially, in Paget's disease there is a marked increase in the rate of bone resorption at localized areas caused by large and numerous osteoclasts. These localized areas of osteolysis are seen radiologically as an advancing lytic wedge in long bones or osteoporosis circumscripta in the skull. The osteolysis is followed by a compensatory increase in bone formation induced by osteoblasts recruited to the area. This is associated with accelerated deposition of lamellar bone in a disorganized fashion. This intense cellular activity produces a chaotic picture of trabecular bone ("mosaic" pattern), rather than the normal linear lamellar pattern. The resorbed bone is replaced and the marrow spaces are filled by an excess of fibrous connective tissue with a marked increase in blood vessels, causing the bone to become hypervascular. The bone hypercellularity may then diminish, leaving a dense "pagetic bone," also known as burned-out Paget's disease.



Paget's disease bone tissue: High magnification micrograph of Paget's disease of the bone. Notice the prominent jigsaw-puzzle like pattern.

Treatment

In general, patients with Paget's disease should receive 1000–1500 mg of calcium, adequate sunshine, and at least 400 units of vitamin D daily. Exercise is very important in maintaining skeletal health, avoiding weight gain, and maintaining joint mobility. Since undue stress on affected bones should be avoided, patients should discuss any exercise program with their physicians before beginning.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Osteoporosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Osteoporosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- peak bone mass. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/peak%20bone%20mass**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- fragility fractures. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/fragility%20fractures**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Elderly woman with osteoporosis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fe/OsteoCutout.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Paget's disease of bone. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Paget's_disease_of_bone**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Paget's disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Paget's%20disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Elderly woman with osteoporosis. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/f/fe/OsteoCutout.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Paget's disease. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/e/e0/Paget_disease_of_bone_-_high_mag.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Paget's disease. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/3/3e/Morbus_Paget_Becken2.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 29.2B: Paget's Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.3: INJURIES AND DISORDERS OF THE SKELETON

Close message

Learn more

Welcome to your new guide!

Click on "New" to build out your guide.

- Topic hierarchy

29.3A: Spina Bifida

29.3B: Fractures

This page titled 29.3: Injuries and Disorders of the Skeleton is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.3A: Spina Bifida

Spina bifida is a developmental congenital disorder caused by the incomplete closing of the embryonic neural tube.

Learning Objectives

- Describe spina bifida and why fewer babies are being born with it

Key Points

- In spina bifida, some vertebrae overlying the spinal cord are not fully formed and remain unfused and open.
- There is neither a single cause of spina bifida nor any known way to prevent it entirely; however, dietary supplementation with folic acid prior to pregnancy has been shown to be helpful in reducing the incidence of spina bifida.
- Neural tube defects can usually be detected during pregnancy by testing the mother's blood (AFP screening) or by a detailed fetal ultrasound.

Key Terms

- **neural tube**: A hollow longitudinal dorsal tube formed in the folding and subsequent fusion of the opposite ectodermal folds in the embryo that gives rise to the brain and spinal cord.
- **AFP**: Alpha-Fetal Protein (AFP); a major plasma protein produced by the yolk sac and the liver during fetal development that is thought to be the fetal form of serum albumin. AFP is measured in pregnant women through the analysis of maternal blood or amniotic fluid as a screening test for a subset of developmental abnormalities; it is principally increased in open neural tube defects and omphalocele and is decreased in Down syndrome.



Spina bifida: Location of spina bifida on an infant.


Spina bifida (Latin: "split spine") is a developmental congenital disorder caused by the incomplete closing of the embryonic neural tube. Some vertebrae overlying the spinal cord are not fully formed and remain unfused and open.

The protruded portion of the spinal cord and the nerves that originate at that level of the cord are damaged or not properly developed. As a result, there is usually some degree of paralysis and loss of sensation below the level of the spinal cord defect. Thus, the higher the level of the defect, the more severe the associated nerve dysfunction and resultant paralysis may be. People may have ambulatory problems; loss of sensation; deformities of the hips, knees, or feet; and loss of muscle tone.

Treatment for Spina Bifida

Spina bifida can be surgically closed after birth, but this does not restore normal function to the affected part of the spinal cord. Intrauterine surgery for spina bifida has also been performed; the safety and efficacy of this procedure is currently being investigated. The spinal cord lesion or the scarring due to surgery may result in a tethered spinal cord. In some individuals, this causes significant traction and stress on the spinal cord. This may lead to a worsening of associated paralysis, scoliosis, back pain, and worsening bowel and/or bladder function.

Causes of Spina Bifida



Spina Bifida: Spina bifida in the lumbar area. (1) External sac with cerebrospinal fluid. (2) Spinal cord wedged between the vertebrae.

There is neither a single cause of spina bifida nor any known way to prevent it entirely. However, dietary supplementation with folic acid prior to pregnancy has been shown to be helpful in reducing the incidence of spina bifida. Natural sources of folic acid include whole grains, fortified breakfast cereals, dried beans, leafy vegetables, and fruits. Neural tube defects can usually be detected during pregnancy by testing the mother's blood (AFP screening) or by a detailed fetal ultrasound. Ultrasound screening for spina bifida is partly responsible for the decline in new cases because many pregnancies are terminated out of fear that a newborn might have a poor future quality of life. With modern medical care, the quality of life of patients has greatly improved.

This page titled 29.3A: Spina Bifida is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.3B: Fractures

A bone fracture is a medical condition in which there is damage to the continuity of a bone.

Learning Objectives

- Describe the different types of bone fractures

Key Points

- A spinal fracture is a fracture of the vertebral column. A compression fracture is the collapse of vertebrae, while a cervical fracture is a fracture of any of the vertebral bones in the neck.
- A rib fracture is a break or fracture in one or more of the bones making up the rib cage.
- A hip fracture is a fracture that occurs in the proximal end of the femur, near the hip.
- A clavicle fracture is a bone fracture in the clavicle, or collarbone.

Key Terms

- **fracture**: Damage to the continuity of a bone.

A bone fracture is a medical condition in which there is damage to the continuity of a bone. A bone fracture can be the result of high force impact or stress, or a minimal trauma injury as a result of certain medical conditions that weaken the bones, such as osteoporosis or bone cancer. Although broken bone and bone break are common colloquialisms for a bone fracture,"break" is not a formal orthopedic term. Fractures can occur in bones throughout the body.

Fractures of the Vertebral Column

A spinal fracture (or vertebral fracture) is a fracture affecting the bones of the spinal column. They can affect the cervical vertebrae (a cervical fracture), the thoracic, or the lumbar regions.

Compression Fractures

A compression fracture is a collapse of vertebra. It may be due to trauma or a weakening of the vertebra. This weakening is seen in patients with osteoporosis or osteogenesis imperfecta, lytic lesions from metastatic or primary tumors, or infection. In healthy patients, it is most often seen in individuals suffering extreme vertical shocks, such as ejecting from an ejection seat. Seen in lateral views in plain x-ray films, compression fractures of the spine characteristically appear as wedge deformities, with greater loss of height anteriorly than posteriorly and intact pedicles in the anteroposterior view.

Cervical Fractures



Vertebral fracture: Lateral spine X-ray showing osteoporotic wedge fractures of the spine.

A cervical fracture is commonly called a broken neck. There are seven cervical vertebrae (neck bones) in the human neck, and the fracture of any can be catastrophic. The most common causes are traffic accidents or diving into shallow water. Abnormal movement of bones or pieces of bone can cause spinal cord injury resulting in loss of sensation, paralysis, or death. Considerable force is needed to cause a cervical fracture. Motor vehicle collisions and falls are common causes. A severe, sudden twist to the neck or a severe blow to the head or neck area can cause a cervical fracture. Sports involving violent physical contact carry a risk of cervical fracture.

Rib Fracture

A rib fracture is a break or fracture in one or more of the bones making up the rib cage. Fractures of the first and second ribs may be more likely to be associated with head and facial injuries than other rib fractures. The middle ribs are the ones most commonly fractured. Fractures usually occur from direct blows or from indirect crushing injuries. A rib fracture has the complication of potentially causing a pulmonary contusion. Rib fractures are usually quite painful because the ribs have to move to allow for breathing. When several ribs are broken in several places a flail chest results, and the detached bone sections will move separately from the rest of the chest.

Hip Fracture

A hip fracture is a serious femoral fracture that occurs in the proximal end of the femur (the long bone running through the thigh), near the hip. The term “hip fracture” is commonly used to refer to four different fracture patterns and is often due to osteoporosis; in the vast majority of cases, a hip fracture is a fragility fracture due to a fall or minor trauma in someone with weakened osteoporotic bone. Most hip fractures in people with normal bone are the result of high-energy trauma such as car accidents, falling from heights, or sports injuries.

Clavicle Fracture

A clavicle fracture is a bone fracture in the clavicle, or collarbone. It is often caused by a fall onto an outstretched upper extremity, a fall onto a shoulder, or a direct blow to the clavicle. Many research projects are underway regarding the medical healing process of clavicle fractures.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Spina Bifida. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spina_Bifida**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- AFP. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/AFP**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- neural tube. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/neural_tube**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spina bifida on an infant. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/83/Spina_bifida_drawing.gif**. **License**: ***Public Domain: No Known Copyright***
- Spina bifida. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/1/13/Meningomyelocele_diagram.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Clavicle fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Clavicle_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spinal fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Spinal_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cervical_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rib fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rib_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Compression fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Compression_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Bone fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Bone_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cervical vertebrae. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/cervical%20vertebrae**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hip fracture. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hip_fracture**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Spina bifida on an infant. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/8/83/Spina_bifida_drawing.gif**. **License**: ***Public Domain: No Known Copyright***
- Spina bifida. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/Wikipedia/commons/1/13/Meningomyelocele_diagram.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- 404px-L1_2_vertebral_fracture.jpg. **Provided by**: Wikimedia Commons. **Located at**: **commons.wikimedia.org/wiki/File:L1_2_vertebral_fracture.jpg**. **License**: ***Public Domain: No Known Copyright***

This page titled 29.3B: Fractures is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.4: DISEASES, DISORDERS, AND INJURIES OF THE JOINTS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.4A: Sprain and Strain
29.4B: Rheumatism and Arthritis
29.4C: Common Joint Injuries

This page titled 29.4: Diseases, Disorders, and Injuries of the Joints is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.4A: Sprain and Strain

A strain is a tear in tendon fibers resulting from overstretching, while a sprain is an equivalent injury to a ligament.

Learning Objectives

- Describe a strain and its treatment

Key Points

- Strains and sprains can happen to anyone, but athletes are at higher risk.
- P.R.I.C.E. is a treatment method for muscular strain: protection, rest, ice, compression, elevation.
- Cold compression therapy uses ice packs and wrapping to reduce swelling and inflammation to encourage the healing process.

Key Terms

- **strain**: An injury to a muscle or tendon in which the muscle fibers tear as a result of overstretching.
- **sprain**: To weaken a ligament by sudden and excessive exertion, as by wrenching; to overstrain, or stretch injuriously, but without luxation; as, to sprain one's ankle.

EXAMPLES

Gymnasts frequently suffer from mild ankle sprains. Cold compression therapy is the recommended treatment. As soon as possible, the gymnast needs to begin to take brief walks as part of rehabilitative therapy. Failure to rehabilitate the ankle can permanently weaken it.

A strain is an injury to a muscle or tendon in which the muscle fibers tear as a result of overstretching. A strain is also colloquially known as a pulled muscle. The equivalent injury to a ligament is a sprain. Typical symptoms of a strain include: localized pain, stiffness, discoloration, and bruising around the strained muscle. Strains can happen while doing everyday tasks and are not restricted to athletes. Nevertheless, people who play sports are more at risk of developing a strain due to increased muscle use.



Muscle Strain: Pulled hamstring after four days. Two images of the same leg. One of the pictures was taken in a mirror.

Treatment of Strains and Sprains

The first-line treatment for a muscular strain in the acute phase includes five steps commonly known as P.R.I.C.E.

- Protection: Apply soft padding to minimize impact with objects.
- Rest: Don't use the injured limb.
- Ice: Apply ice to reduce swelling by reducing blood flow to the injury site. Never ice for more than 20 minutes at a time.
- Compression: Wrap the strained area to reduce swelling with an ACE soft-wrapped bandage.
- Elevation: Keep the strained area as close to the level of the heart as is conveniently possible to keep blood from pooling in the injured area.

The ice and compression (cold compression therapy) will stop the pain and swelling while the injury starts to heal itself. Controlling the inflammation is critical to the healing process and the icing further restricts fluid leaking into the injured area as well as controlling pain.

This immediate treatment is usually accompanied by the use of non-steroidal anti-inflammatory drugs (e.g., ibuprofen), which both reduce the immediate inflammation and relieve pain. However, NSAIDs, including aspirin and ibuprofen, affect platelet function (this is why they are known as "blood thinners") and should not be taken during the period when tissue is bleeding because they will tend to increase blood flow and inhibit clotting. After the bleeding has stopped, NSAIDs can be used with some effectiveness to reduce inflammation and pain. It is recommended that the injured person should consult a medical provider if the injury is accompanied by severe pain, if the limb cannot be used, or if there is noticeable tenderness over an isolated spot. These can be signs of a broken or fractured bone, a sprain, or a complete muscle tear.

This page titled 29.4A: Sprain and Strain is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.4B: Rheumatism and Arthritis

Rheumatoid arthritis is a chronic, systemic inflammatory disorder that affects many tissues and organs, but mainly attacks flexible joints.

Learning Objectives

- Differentiate rheumatoid arthritis from osteoarthritis

Key Points

- Rheumatism is an archaic term for problems related to joints and connective tissues that has been replaced with the field of rheumatology.
- Although the cause of rheumatoid arthritis is unknown, autoimmunity plays a pivotal role in both its chronicity and progression; it is considered a systemic autoimmune disease.
- The discovery that vitamin D plays a role in immune regulation has led research showing improvement in patients with rheumatoid arthritis who had high plasma levels of 25(OH)D3.
- There are over 100 types of arthritis and osteoarthritis is the most common.
- Arthritis is the most common cause of disability in the U.S. as it makes physical activity very difficult for those suffering from it.

Key Terms

- **rheumatism**: Any disorder of the muscles, tendons, joints, bones, nerves, characterized by pain, discomfort, and disability.
- **rheumatoid arthritis**: A chronic and progressive disease in which the immune system attacks the joints. It is characterised by pain, inflammation and swelling of the joints, stiffness, weakness, loss of mobility, and deformity. Tissues throughout the body can be affected including the skin, blood vessels, heart, lungs, and muscles.

EXAMPLES

Kathleen Turner, the actress from the movie Body Heat found out when she was in her mid-40s that she suffered from rheumatoid arthritis. The crippling pain made her life almost intolerable. But with the help of medical treatment and exercise, she battled back to return to acting.

Rheumatism or rheumatic disorder is a non-specific term for medical problems affecting the joints and connective tissue. The study of, and therapeutic interventions in, such disorders is called rheumatology. The term “rheumatism” is still used in colloquial speech and historical contexts, but is no longer frequently used in medical or technical literature; there is no longer any recognized disorder simply called “rheumatism.” Some countries use the word rheumatism to describe fibromyalgia syndrome. The term “Rheumatic Diseases” is used to refer to connective tissue disorders.



Rheumatoid Arthritis: A hand affected by rheumatoid arthritis.

Rheumatoid arthritis (RA) is a chronic, systemic inflammatory disorder that may affect many tissues and organs, but principally attacks flexible (synovial) joints. The process involves an inflammatory response of the capsule around the joints (synovium) secondary to swelling (hyperplasia) of synovial cells, excess synovial fluid, and the development of fibrous tissue (pannus) in the synovium. The pathology of the disease process often leads to the destruction of articular cartilage and ankylosis (fusion) of the joints. Rheumatoid arthritis can also produce diffuse inflammation in the lungs, the membrane around the heart (pericardium), the membranes of the lung (pleura), and the white of the eye (sclera), and also nodular lesions, most common in subcutaneous tissue. Although the cause of rheumatoid arthritis is unknown, autoimmunity plays a pivotal role in both its chronicity and progression, and RA is considered a systemic autoimmune disease.

Treatment of Rheumatoid Arthritis

There is no known cure for rheumatoid arthritis, but many different types of treatment can alleviate symptoms and/or modify the disease process. The goals of treatment include minimizing clinical symptoms such as pain and swelling, as well as preventing bone deformity and radiographic damage (for example, bone erosions visible in x-rays), and maintaining the quality of life in terms of day-to-day activities. The discovery of the vitamin D receptor (VDR) in the cells of the immune system and the fact that activated dendritic cells produce the vitamin D hormone suggested that vitamin D could have immunoregulatory properties. 25(OH)D3 plasma levels have been found inversely correlated, with the RA disease activity showing a circannual rhythm (more severe in winter). Recently, greater intake of vitamin D was associated with a lower risk of RA. A significant clinical improvement was strongly correlated with the immunomodulating potential in vitamin D-treated RA patients.



Osteoarthritis: Severe osteoarthritis and osteopenia of the carpal joint and first carpometacarpal joint.

There are over 100 different forms of arthritis. The most common form, osteoarthritis (degenerative joint disease), is a result of trauma to the joint, infection of the joint, or age. Osteoarthritis is characterized by a lack of primary inflammation. Other forms of arthritis include rheumatoid arthritis, psoriatic arthritis, and related autoimmune diseases. Septic arthritis is caused by joint infection. The major complaint by individuals who have arthritis is joint pain. Pain is often a constant and may be localized to the joint affected. The pain from arthritis is due to inflammation that occurs around the joint, damage to the joint from disease, daily wear and tear of joint, muscle strains caused by forceful movements against stiff, painful joints, and fatigue.

Arthritis is the most common cause of disability in the United States. More than 20 million individuals with arthritis have severe limitations in function on a daily basis. Absenteeism and frequent visits to the physician are common in individuals who have arthritis. Arthritis makes it very difficult for individuals to be physically active and many become home bound. Treatment options vary depending on the type of arthritis and include physical therapy, lifestyle changes (including exercise and weight control), orthopedic bracing, and medications. Joint replacement surgery may be required in eroding forms of arthritis. Medications can help reduce inflammation in the joint which decreases pain. Moreover, by decreasing inflammation, the joint damage may be slowed.

This page titled 29.4B: Rheumatism and Arthritis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.4C: Common Joint Injuries

Common joint injuries include shoulder injuries, elbow injuries, and knee injuries.

Learning Objectives

- Describe common joint injuries

Key Points

- Common shoulder injuries include rotator cuff tears, dislocation, and separated shoulder.
- Common elbow injuries are primarily due to overuse. These include tennis elbow and golfer's elbow.
- Common knee injuries can be due to overuse or due to traumatic tears, such as the common ACL tear that occurs most often in sports.

Key Terms

- **ACL**: The most commonly injured ligament of the knee.

Shoulder Injuries

This instability increases the likelihood of joint injury, often leading to a degenerative process in which tissues break down and no longer function well.

Rotator Cuff Tear

The term "rotator cuff" refers to a group of four tendons that blend together as they attach to the upper end of the arm bone (humerus). Normally these tendons transmit the force of muscles originating on the shoulder Blase (scapula) to the arm providing motion and stability. Defects in the rotator cuff can come from an injury (cuff tear) or from degeneration (cuff wear). The degree to which a tendon is reparable depends on its quantity and quality. Degenerated tendons are often frail and retracted and may not be amenable to repair. Individuals that are elderly, smokers, or those having had cortisone injections often have weaker tendon tissue that fails without a significant injury. By contrast, those whose tendon was torn by a substantial fall often have good quality tendon that can be repaired if surgery is performed promptly after the injury.

Dislocated Shoulder

The shoulder joint is the most frequently dislocated major joint of the body. In a typical case of a dislocated shoulder, a strong force that pulls the shoulder outward (abduction) or extreme rotation of the joint pops the ball of the humerus out of the shoulder socket. Dislocation commonly occurs when there is a backward pull on the arm that either catches the muscles unprepared to resist or overwhelms the muscles. When a shoulder dislocates frequently, the condition is referred to as shoulder instability. A partial dislocation where the upper arm bone is partially in and partially out of the socket is called a subluxation. In the medical community, dislocation is commonly referred to as luxation.

Separated Shoulder

A shoulder separation occurs where the collarbone (clavicle) meets the shoulder blade (scapula). When ligaments that hold the AC (acromioclavicular) joint together are partially or completely torn, the outer end of the clavicle may slip out of place, preventing it from properly meeting the scapula. Most often the injury is caused by a blow to the shoulder or by falling on an outstretched hand. After injury it is hard to do a 180 degrees rotation.

Elbow Injuries

The types of disease most commonly seen at the elbow are due to injury.

Tendonitis

Two of the most common injuries at the elbow are overuse injuries: tennis elbow and golfer's elbow. Golfer's elbow involves the tendon of the common flexor origin which originates at the medial epicondyle of the humerus (the "inside" of the elbow). Tennis elbow is the equivalent injury, but at the common extensor origin (the lateral epicondyle of the humerus).

Fractures

There are three bones at the elbow joint, and any combination of these bones may be involved in a fracture of the elbow.

Knee Injuries

Knee pain can be caused by trauma, misalignment, and degeneration as well as by conditions like arthritis. The most common knee disorder is generally known as patellofemoral syndrome.The majority of minor cases of knee pain can be treated at home with rest and ice but more serious injuries do require surgical care.

Physical fitness is related integrally to the development of knee problems. The same activity such as climbing stairs may cause pain from patellofemoral compression for someone who is physically unfit, but not for someone else (or even for that person at a different time).

ACL Tear

The anterior cruciate ligament (ACL) is the most commonly injured ligament of the knee. The injury is common during sports. Twisting of the knee is a common cause of over-stretching or tearing the ACL. When the ACL is injured a popping sound may be heard, and the leg may suddenly give out. Besides swelling and pain, walking may be painful and the knee will feel unstable. Minor tears of the anterior cruciate ligament may heal over time, but a torn ACL requires surgery. After surgery, recovery is prolonged and low impact exercises are recommended to strengthen the joint.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Muscle strain. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscle_strain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- strain. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/strain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- sprain. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/sprain**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle strain. **Provided by**: Wikimedia. **Located at**: **upload.wikimedia.org/wikipedi..._hamstring.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rheumatism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rheumatism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Arthritis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arthritis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Rheumatoid arthritis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Rheumatoid_arthritis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- rheumatism. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/rheumatism**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- rheumatoid arthritis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/rheumatoid_arthritis. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle strain. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._hamstring.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...f_the_hand.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._Arthritis.JPG. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Knee. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Knee. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Shoulder problem. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Shoulder_problem. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Elbow. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Elbow. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Muscle strain. **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._hamstring.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi...f_the_hand.jpg. **License**: ***CC BY-SA: Attribution-ShareAlike***
- **Provided by**: Wikimedia. **Located at**: upload.wikimedia.org/wikipedi..._Arthritis.JPG. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 29.4C: Common Joint Injuries is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.5: DISEASES, DISORDERS, AND INJURIES OF THE MUSCULAR SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.5A: Muscular Dystrophy

29.5B: Exercise-Induced Muscle Damage

29.5C: Carpal Tunnel Syndrome

This page titled 29.5: Diseases, Disorders, and Injuries of the Muscular System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.5A: MUSCULAR DYSTROPHY

Muscular dystrophy (MD) is a group of muscle diseases characterized by non-functional muscle proteins that impair proper function.

Learning Objectives

- List the factors associated with muscular dystrophy (MD)


Key Points

- In muscular dystrophy (MD) the proteins dystrophin and the dystrophin protein complexes are altered.
- Progression of muscular dystrophies includes death of muscle tissue.
- Several types of MD exist including Duchenne MD, Becker, limb-girdle, congenital, facioscapulohumeral, myotonic, oculopharyngeal, distal, and Emery-Dreifuss MD.

Key Terms

- **dystrophin**: A cytoplasmic structural protein that is deficient in some forms of muscular dystrophy.
- **muscular dystrophy**: A group of genetic diseases that cause progressive skeletal muscle weakness, defects in muscle proteins, and the death of muscle cells and tissue.

Muscular dystrophy (MD) is a group of muscle diseases characterized by the creation of non-functional muscle proteins that weakens muscles and impairs proper function. Specifically the proteins dystrophin and the dystrophin protein complexes are altered and nonfunctional, unable to join with actin filaments. This leads to disruptions of mechanical stabilization and calcium concentration regulation within the sarcomeres, altering the ability of filaments to bind and cause contraction. Progression of the disease includes death of muscle tissue.



Muscular Dystrophy: In affected muscle the tissue becomes disorganized and the concentration of dystrophin (green) is greatly reduced.

Types of Muscular Dystrophy

Several types of MD exist, including Duchenne MD, the first description of a MD disease, noted by Guillame Duchenne, a French neurologist, after study of 13 young boys presenting symptoms. Other forms include Becker, limb-girdle, congenital, facioscapulohumeral, myotonic, oculopharyngeal, distal, and Emery-Dreifuss MD. Most of these diseases affect multiple organ systems including the cardiovascular, gastrointestinal, nervous, and endocrine systems. Although not typically affecting intellectual abilities, a subset of MD patients do exhibit cognitive impairment, behavioral, vision, and speech problems likely linked to MD, and this is seen particularly in DMD. MD-affected individuals with susceptible intellectual impairment are diagnosed through molecular characteristics but not through problems associated with disability.

Causes of Muscular Dystrophy

Muscular dystrophy is generally an inherited disease, with men affected at much higher rates than women. Women are often carriers of the disease gene variant, but not affected themselves, although other patterns of inheritance exist for the different types of MD. However, mutations of the dystrophin gene and nutritional defects (with no genetics history) at the prenatal stage are also possible in about 33% of people affected by DMD.

29.5B: Exercise-Induced Muscle Damage

This page titled 29.5A: Muscular Dystrophy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.5B: Exercise-Induced Muscle Damage

Exercise damages muscles due to eccentric and concentric muscle loading and often results in delayed onset muscle soreness (DOMS).

Learning Objectives

- Explain the process of exercise-induced muscle damage


Key Points

- Delayed onset muscle soreness is a result of damage in muscle sarcomeres during contractions.
- The acute but delayed nature of muscle soreness is indicative of an inflammatory immune response.
- This disruption of the muscle fibers triggers white blood cells to increase following the induced muscle soreness, leading to the inflammatory response observation from the induced muscle soreness.

Key Terms

- **eccentric contraction**: The elongation of a muscle while under tension due to an opposing force being greater than the force generated by the muscle.
- **concentric contraction**: A type of muscle contraction in which the muscles shorten while generating force.

Exercise damages muscles due to eccentric and concentric muscle loading. Resistance training, and particularly high loading during eccentric contractions, results in delayed onset muscle soreness (DOMS). Previously attributed to the accumulation of lactic acid during exercise, it is now understood that DOMS is due to structural damage in sarcomeres, particularly to the z-disks and contractile filaments. Z-disks provide the structural support for the contractile filaments of the sliding filament mechanism. Overloading of muscles damages these connections and the orientation of the contractile filaments.



The Sarcomere: Delayed onset muscle soreness is caused by structural damage to the Z disk and myosin and actin filaments.

Causes of Exercise-Induced Muscle Damage

Acute inflammation of the muscle cells, as understood in exercise physiology, can result after induced eccentric and concentric muscle training. Participation in eccentric training and conditioning, including resistance training and activities that emphasize eccentric lengthening of the muscle such as downhill running on a moderate to high incline, can result in considerable soreness within 24 to 48 hours, even though blood lactate levels, previously thought to cause muscle soreness, were much higher with level running. This has been noted especially in marathon runners whose muscle fibers revealed remarkable damage after both training and marathon competition. The onset and timing of this gradient damage to the muscle parallels the degree of muscle soreness experienced by the runners.

This disruption of the muscle fibers triggers white blood cells to increase following the induced muscle soreness, leading to the inflammatory response observation from the induced muscle soreness. Elevations in plasma enzymes, myoglobinemia, and abnormal muscle histology and ultrastructure are concluded to be associated with the inflammatory response. High tension in the contractile-elastic system of muscle results in structural damage to the muscle fiber and plasmalemma and its epimysium, perimysium, and endomysium. The mysium damage disrupts calcium homeostasis in the injured fiber and fiber bundles, resulting in necrosis that peaks about 48 hours after exercise. The products of the macrophage activity and intracellular contents (such as histamines, kinins, and K+) accumulate outside the cells. These substances then stimulate the free nerve endings in the muscle; a process that appears accentuated by eccentric exercise, in which large forces are distributed over a relatively small cross-sectional area of the muscle.

The acute, and delayed nature of muscle soreness is indicative of an inflammatory immune response. Damage to the sarcomeres causes aninflux of white blood cells, leading to inflammation, which is itself associated with increased plasma enzyme concentration, myoglobinemia, and abnormal muscle structure and histology. A further response to sarcomere damage is necrosis following damage to the mysium, which peaks about 48 hours following exercise.

The muscle adapts rapidly to the structural damage caused by exercise, and further soreness and damage in later exercise events is mitigated.

This page titled 29.5B: Exercise-Induced Muscle Damage is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.5C: Carpal Tunnel Syndrome

Carpal tunnel syndrome is an entrapment median neuropathy of the median nerve due to its compression at the wrist in the carpal tunnel.

Learning Objectives

- Describe the factors associated with carpal tunnel syndrome


Key Points

- Carpal tunnel syndrome is a neuropathy resulting from compression of the median nerve in the wrist as it travels through the carpal tunnel.
- The resulting symptoms include pain, numbness, and paresthesia (a tingling or burning sensation).
- The causes of carpal tunnel syndrome are not completely understood.
- Factors that exert pressure on the tunnel, such as benign tumors, inflammation, and repetitive tasks such as typing, are also strongly associated with CTS.
- CTS is diagnosed by clinical assessments that seek to elicit symptoms and include Phalen's maneuver, Tinel's sign, and Drukan test.
- Treatments of CTS include immobilization of the wrist, steroid injection, activity modification, physical therapy, massage, medication, or surgery to release the transverse carpal ligament
- Repetitive movements are believed to contribute to carpal tunnel syndrome. Symptoms of carpal tunnel syndrome arise from compression of the median nerve at the wrist.

Key Terms

- **carpal tunnel syndrome**: Carpal tunnel syndrome is an entrapment median neuropathy, causing paresthesia, pain, numbness, and other symptoms in the distribution of the median nerve due to its compression at the wrist in the carpal tunnel.

Carpal tunnel syndrome (CTS) is an entrapment median neuropathy, causing paresthesia, pain, numbness, and other symptoms in the distribution of the median nerve due to its compression at the wrist in the carpal tunnel. The pathophysiology is not completely understood but can be attributed to the compression of the median nerve traveling through the carpal tunnel.



Untreated Carpal Tunnel Syndrome: Patients with CTS experience numbness, tingling, or burning sensations in the thumb and fingers, in particular the index, middle fingers, and radial half of the ring fingers, which are innervated by the median nerve. Less-specific symptoms may include pain in the wrists or hands and loss of grip strength.

Symptoms of Carpal Tunnel Syndrome

Patients with CTS experience numbness, tingling, or burning sensations in the thumb and fingers, in particular the index, middle fingers, and radial half of the ring fingers, which are innervated by the median nerve. Less-specific symptoms may include pain in the wrists or hands and loss of grip strength. The numbness often occurs at night, with the hypothesis that the wrists are held flexed during sleep. It can be relieved by wearing a wrist splint that prevents flexion. Long-standing CTS leads to permanent nerve damage with constant numbness, atrophy of some of the muscles of the thenar eminence, and weakness of palmar abduction. Pain in carpal tunnel syndrome is primarily numbness that is so intense that it wakes one from sleep. Pain in electrophysiologically verified CTS is associated with misinterpretation of nociception and depression. Palliative treatments for CTS include use of night splints and corticosteroid injection. The only scientifically established disease modifying treatment is surgery to cut the transverse carpal ligament.

Causes of Carpal Tunnel Syndrome

Most cases of CTS are of unknown causes, or idiopathic. Carpal tunnel syndrome can be associated with any condition that causes pressure on the median nerve at the wrist. A variety of patient factors can lead to CTS, including heredity, size of the carpal tunnel, associated local and systematic diseases, and certain habits. Some common conditions that can lead to CTS include obesity, oral contraceptives, hypothyroidism, arthritis, diabetes, and trauma. Carpal tunnel syndrome is also associated with repetitive activities of the hand and wrist, in particular with a combination of forceful and repetitive activities. Non-traumatic causes generally happen over a period of time, and are not triggered by one certain event. Many of these factors are manifestations of physiologic aging.

Treatments of Carpal Tunnel Syndrome

Treatments of CTS include immobilization of the wrist, steroid injection, activity modification, physical therapy, massage, medication, or surgery to release the transverse carpal ligament.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Muscular dystrophy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Muscular_dystrophy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- dystrophin. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/dystrophin**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- muscular dystrophy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/muscular_dystrophy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- MuscularDystrophy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MuscularDystrophy.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Inflammation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Inflammation%23Exercise-induced_acute_inflammation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Delayed onset muscle soreness. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Delayed_onset_muscle_soreness**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Inflammation. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Inflammation%23Exercise_and_Inflammation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Concentric contraction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Concentric_contraction%23Concentric_contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- eccentric contraction. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/eccentric_contraction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- MuscularDystrophy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MuscularDystrophy.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Carpal Tunnel Syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Carpal_Tunnel_Syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- carpal tunnel syndrome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/carpal_tunnel_syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- MuscularDystrophy. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:MuscularDystrophy.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Sarcomere. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Sarcomere.svg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Untreated Carpal Tunnel Syndrome. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Untreated_Carpal_Tunnel_Syndrome.JPG**. **License**:

This page titled 29.5C: Carpal Tunnel Syndrome is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6: INJURIES, DISEASES, AND DISORDERS OF THE CENTRAL NERVOUS SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.6A: Traumatic Brain Injury
29.6B: Cerebrovascular Accident
29.6C: Transient Ischemic Attacks
29.6D: Alzheimer's Disease
29.6E: Brain Tumors
29.6F: Amyotrophic Lateral Sclerosis

This page titled 29.6: Injuries, Diseases, and Disorders of the Central Nervous System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6A: Traumatic Brain Injury

Traumatic brain injury (TBI) occurs when an external force injures the brain and can be caused by a direct impact or by acceleration alone.

Learning Objectives

- Recognize the severity of traumatic brain injuries

Key Points

- Traumatic brain injury (TBI) occurs when an external force injures the brain and can be caused by a direct impact or by acceleration alone.
- Following the initial injury, brain trauma causes secondary events, like changes in cerebral blood flow and intracranial pressure, that contribute to the damage caused by the trauma.
- Brain injuries can be classified into mild, moderate, and severe categories; the Glasgow Coma Scale (GCS) is the most commonly used system for classifying TBI severity.
- Symptoms of TBI depend on the severity of the injury, whether the injury is diffuse or focal, and the part of the brain that is affected.
- A famous case of personality change following TBI is that of Phineas Gage, who survived a construction accident where a large iron rod was driven through the front of his head.

Key Terms

- **Glasgow Coma Scale**: (GCS) A neurological scale that aims to give a reliable, objective way of recording the conscious state of a person for initial as well as subsequent assessment.

EXAMPLES

Examples of the causes of TBI include violence, transportation accidents, construction accidents, and sports. Examples of the symptoms of TBI include headache, nausea, dizziness, confusion, and trouble with memory, which can vary depending on the severity of the injury. Common long-term symptoms of moderate to severe TBI are changes in social behavior and cognitive changes, including problems with attention and planning.

Traumatic brain injury (TBI), also known as intracranial injury, occurs when an external force traumatically injures the brain. TBI can be classified based on severity, mechanism (closed or penetrating head injury), or other features, such as whether it occurs in a specific location or over a widespread area. Head injury usually refers to TBI, but is a broader category because it can involve damage to structures other than the brain, such as the scalp and skull.

TBI is a major cause of death and disability worldwide, especially in children and young adults. Causes include falls, vehicle accidents, and violence. Prevention measures include use of technology to protect those who suffer from automobile accidents, such as seat belts and sports or motorcycle helmets. Efforts to reduce the number of automobile accidents are also implemented, such as safety education programs and enforcement of traffic laws.


Causes of Traumatic Brain Injury

Brain trauma can be caused by a direct impact or by acceleration alone. In addition to the damage caused at the moment of injury, brain trauma causes secondary injury, a variety of events that take place in the minutes and days following the injury. These processes, which include alterations in cerebral blood flow and the pressure within the skull, contribute substantially to the damage from the initial injury.

Types of Brain Injuries

Brain injuries can be classified into mild, moderate, and severe categories. The Glasgow Coma Scale (GCS), the most commonly used system for classifying TBI severity, grades a person's level of consciousness on a scale of 3–15 based on verbal, motor, and eye-opening reactions to stimuli. It is generally agreed that a TBI with a GCS of 13 or above is mild, 9–12 is moderate, and 8 or below is severe.

Side Effects and Complications



Subdural hematoma is a possible result of traumatic head injury: In this image, the single arrow marks spread of the subdural haematoma and the double arrow marks the midline shift.

TBI can cause a host of physical, cognitive, social, emotional, and behavioral effects, and the outcome can range from complete recovery to permanent disability or death. The 20th century saw critical developments in diagnosis and treatment that decreased death rates and improved outcome. Some of the current imaging techniques used for diagnosis and treatment include CT scans (computed tomography) and MRIs (magnetic resonance imaging). Depending on the injury, treatment required may be minimal or may include interventions such as medications, emergency surgery or surgery years later. Physical therapy, speech therapy, recreation therapy, and occupational therapy may be employed for rehabilitation.

Complications are distinct medical problems that may arise as a result of the TBI. TBI can cause prolonged or permanent effects on consciousness, such as coma, brain death, persistent vegetative state (in which patients are unable to achieve a state of alertness to interact with their surroundings), and minimally conscious state. Lying still for long periods can cause complications including pressure sores, pneumonia or other infections, progressive multiple organ failure, and deep venous thrombosis, which can cause pulmonary embolism. Complications involving the blood vessels include vasospasm, in which vessels constrict and restrict blood flow, the formation of aneurysms, in which the side of a vessel weakens and balloons out, and stroke. Movement disorders that may develop after TBI include tremor, ataxia (uncoordinated muscle movements), myoclonus (shock-like contractions of muscles), and loss of movement range and control (in particular with a loss of movement repertoire). The risk of post-traumatic seizures increases with severity of trauma and is particularly elevated with certain types of brain trauma such as cerebral contusions or hematomas.

This page titled 29.6A: Traumatic Brain Injury is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6B: Cerebrovascular Accident

A cerebrovascular accident results from loss of oxygenated blood to a region of the brain and is typically accompanied by neuronal loss.

Learning Objectives

- Describe what is a cerebrovascular accident or stroke

Key Points

- A stroke, or cerebrovascular accident (CVA), is the rapid loss of brain function due to disturbance in the blood supply to the brain.
- Different types of strokes have different underlying causes.
- Stroke is currently the second leading cause of death in the western world.
- The most important modifiable risk factors for stroke are high blood pressure and atrial fibrillation, although other risk factors can also be modified to decrease risk.


Key Terms

- **embolus**: An obstruction causing an embolism: a blood clot, air bubble, or other matter carried by the blood stream and causing a blockage or occlusion of a blood vessel.
- **thrombosis**: The formation of thrombi in the blood vessels of a living organism, causing obstruction of the circulation.
- **embolism**: An obstruction or occlusion of an artery by an embolus, that is by a blood clot, air bubble, or other matter that has been transported by the blood stream.

EXAMPLES

- Examples of risk factors for stroke include high blood pressure, high cholesterol, diabetes, cigarette smoking, lack of physical activity, and obesity.
- The physical disabilities that can result from stroke are varied and depend on the severity of the stroke and the affected areas of the nervous system. Examples of possible disabilities caused by stroke include loss of motor control, muscle weakness, numbness, appetite loss, speech loss, vision loss, and pain.

A stroke, or cerebrovascular accident (CVA), is the rapid loss of brain function(s) due to disturbance in the blood supply to the brain. This can be due to ischemia (lack of blood flow) caused by blockage (thrombosis, arterial embolism) or a hemorrhage (leakage of blood). As a result, the affected area of the brain cannot function, which might result in an inability to move one or more limbs on one side of the body, inability to understand or formulate speech, or an inability to see one side of the visual field.



An Example of a Stroke: A intra-parenchymal bleed with surrounding edema.

A stroke is a medical emergency and can cause permanent neurological damage, complications, and death. It is the leading cause of adult disability in the United States and Europe and the second leading cause of death worldwide.

Risk Factors

Risk factors for stroke include old age, hypertension (high blood pressure), previous stroke or transient ischemic attack (TIA), diabetes, high cholesterol, cigarette smoking, and atrial fibrillation.

High blood pressure is the most important modifiable risk factor of stroke. An ischemic stroke is occasionally treated in a hospital with thrombolysis (also known as a “clot buster”), and some hemorrhagic strokes benefit from neurosurgery. Treatment to recover any lost function is termed “stroke rehabilitation,” ideally in a stroke unit and involving health professions such as speech and language therapy, physical therapy, and occupational therapy. Prevention of recurrence may involve the administration of antiplatelet drugs such as aspirin and dipyridamole, control and reduction of hypertension, and the use of statins. Selected patients may benefit from carotid endarterectomy and the use of anticoagulants.

Causes

Different types of strokes are caused by different mechanisms.

In thrombotic stroke a thrombus (blood clot) usually forms around atherosclerotic plaques. Since blockage of the artery is gradual, onset of symptomatic thrombotic strokes is slower. A thrombus itself (even if non-occluding) can lead to an embolic stroke if the thrombus breaks off, at which point it is called an ” embolus.” Two types of thrombosis can cause stroke: large vessel disease involves the common and internal carotids, vertebral, and the Circle of Willis.

An embolic stroke refers to the blockage of an artery by an arterial embolus, a travelling particle or debris in the arterial bloodstream originating from elsewhere. An embolus is most frequently a thrombus, but it can also be a number of other substances including fat (e.g., from bone marrow in a broken bone), air, cancer cells, or clumps of bacteria (usually from infectious endocarditis).

Systemic hypoperfusion is the reduction of blood flow to all parts of the body. It is most commonly due to cardiac pump failure from cardiac arrest or arrhythmias, or from reduced cardiac output as a result of myocardial infarction, pulmonary embolism, pericardial effusion, or bleeding. Hypoxemia (low blood oxygen content) may precipitate the hypoperfusion.

Cerebral venous sinus thrombosis leads to stroke due to locally increased venous pressure, which exceeds the pressure generated by the arteries. Infarcts are more likely to undergo hemorrhagic transformation (leaking of blood into the damaged area) than

other types of ischemic stroke.

A silent stroke is a stroke that does not have any outward symptoms, and the patients are typically unaware they have suffered a stroke. Despite not causing identifiable symptoms, a silent stroke still causes damage to the brain and places the patient at increased risk for both transient ischemic attack and major stroke in the future. Conversely, those who have suffered a major stroke are at risk of having silent strokes.

Prevention and Treatment

Given the disease burden of strokes, prevention is an important public health concern. Primary prevention is less effective than secondary prevention (as judged by the number needed to treat to prevent one stroke per year). Because a stroke may indicate underlying atherosclerosis, it is important to determine the patient's risk for other cardiovascular diseases such as coronary heart disease. Conversely, aspirin confers some protection against first stroke in patients who have suffered a myocardial infarction or patients with a high cardiovascular risk.

The most important modifiable risk factors for stroke are high blood pressure and atrial fibrillation (although magnitude of this effect is small). The evidence from the Medical Research Council trials is that 833 patients have to be treated for one year to prevent one stroke. Other modifiable risk factors include high blood cholesterol levels, diabetes, cigarette smoking, heavy alcohol consumption, and unhealthy diet.

This page titled 29.6B: Cerebrovascular Accident is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6C: Transient Ischemic Attacks

A transient ischemic attack is similar to a stroke; though without permanent damage, it can serve as an important risk factor for stroke.

Learning Objectives

- Describe the effects of transient ischemic attacks

Key Points

- A transient ischemic attack (TIA) is often referred to as mini-stroke and is a transient episode of brain dysfunction caused by ischemia, or loss of blood flow to the brain.
- TIAs and strokes cause similar symptoms such as paralysis, weakness, numbness, slurred speech, and mental confusion.
- Unlike a stroke, the symptoms of a TIA can resolve within 24 hours.
- Treatment is limited. A TIA may be prevented by changes in lifestyle.
- Having a TIA is a risk factor for future stroke.

Key Terms

- **ischemia**: Local disturbance in blood circulation due to mechanical obstruction of the blood supply, such as by a narrowed artery or clot.


EXAMPLES

Example drugs used to prevent future TIAs include anti-coagulant medications, like heparin and warfarin, and anti-platelet medications such as aspirin.

A transient ischemic attack (abbreviated as TIA, often referred to as mini stroke) is a transient episode of neurologic dysfunction caused by ischemia—either focal brain, spinal cord, or retinal—without acute infarction (tissue death).

TIAs share the same underlying etiology (cause) as strokes: a disruption of cerebral blood flow (CBF). TIAs and strokes cause the same symptoms. Unlike a stroke, the symptoms of a TIA can resolve within a few minutes or 24 hours. Brain injury may still occur in a TIA lasting only a few minutes. Having a TIA is a risk factor for eventually having a stroke or a silent stroke.

Symptoms



Heart-Great Vessels: Cardiac Myxoma: This image is the left atrial mass of a 71 year old man who had TIAs.

Symptoms vary widely from person to person, depending on the area of the brain involved. The most frequent symptoms include:

- temporary loss of vision (typically amaurosis fugax)
- difficulty speaking (aphasia)
- weakness on one side of the body (hemiparesis)
- numbness or tingling (paresthesia), usually on one side of the body.

There have also been cases of temporary and partial paralysis affecting the face and tongue of the afflicted. The symptoms of a TIA are short-lived and usually last a few seconds to a few minutes, and most symptoms disappear within 60 minutes. Some individuals may have a lingering feeling that something odd happened to the body. Dizziness, lack of coordination, or poor balance are also symptoms related to TIA. Symptoms vary in severity.

Causes

The most common cause of a TIA is an embolus that occludes an artery in the brain. This usually arises from a dislodged atherosclerotic plaque in one of the carotid arteries or from a thrombus (i.e., a blood clot) in the heart because of atrial fibrillation. In a TIA, the blockage period is very short-lived and hence there is no permanent damage.

Other reasons include excessive narrowing of large vessels resulting from an atherosclerotic plaque and increased blood viscosity caused by some blood diseases. TIA is related to other medical conditions such as hypertension, heart disease (especially atrial fibrillation), migraine, cigarette smoking, hypercholesterolemia, and diabetes mellitus.

Risk Factors

Risk factors include:

- Family history of stroke or TIA
- People 55 years or older
- Males have a slightly higher risk of TIA than females, but females are more likely to die from a stroke.
- African Americans generally tend to have a high risk of dying from a stroke, chiefly due to high blood pressure and uncontrolled diabetes.

Prevention and Treatment

A TIA may be prevented by changes in lifestyle. Although most of these recommendations have no solid empirical data, most medical professionals believe them to be so. These include:

- Avoiding smoking
- Cutting down on fats and cholesterol to help reduce plaque build up
- Eating a healthy diet including plenty of fruits and vegetables
- Limiting sodium in the diet, thereby reducing blood pressure
- Exercising regularly
- Moderating alcohol intake
- Maintaining a normal weight
- Controlling blood pressure and keeping blood sugars under control

The mainstay of treatment following acute recovery from a TIA should be to diagnose and treat the underlying cause. It is not always immediately possible to tell the difference between a CVA (stroke) and a TIA. Most patients who are diagnosed at a hospital's emergency department as having suffered from a TIA will be discharged home and advised to contact their primary physician to organize further investigations. A TIA can be considered as the last warning. The reason for the condition should be immediately examined by imaging of the brain.

The initial treatment is aspirin, second line is clopidogrel (PLAVIX), third line is ticlopidine. If TIA is recurrent after aspirin treatment, the combination of aspirin and dipyridamole is needed (Aggrenox).

This page titled 29.6C: Transient Ischemic Attacks is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6D: Alzheimer's Disease

Alzheimer's disease is an age-linked neurodegenerative disorder characterized by marked dementia.

Learning Objectives

- Describe Alzheimer's disease and its effect on the brain

Key Points

- Alzheimer's disease (AD) is the most common form of dementia and was first described by German psychiatrist Alois Alzheimer in 1906.
- There is no cure for AD, which worsens as it progresses and eventually leads to death. Current treatments only help with the symptoms of the disease.
- A small number of cases of AD are inherited and usually have an onset before age 65. This form of the disease is known as early-onset familial AD.
- AD is associated with deposits of amyloid-beta peptide, called plaques, and aggregates of the microtubule-associated protein tau, called tangles, in the brain, which contribute to neuron death.

Key Terms

- **amyloid-beta peptide**: A peptide of 36–43 amino acids that is processed from the Amyloid precursor protein, best known as a component of amyloid plaques in association with Alzheimer's disease.

EXAMPLES

Examples of common symptoms of AD include difficulty remembering recent events and, as the disease advances, confusion, irritability, aggression, mood swings, trouble with language, and long-term memory loss.

Alzheimer's disease (AD), also known in medical literature as Alzheimer disease, is the most common form of dementia. There is no cure for the disease, which worsens as it progresses, and eventually leads to death. It was first described by German psychiatrist and neuropathologist Alois Alzheimer in 1906 and was named after him.

Most often, AD is diagnosed in people over 65 years of age, although the less-prevalent early-onset Alzheimer's can occur much earlier. In 2006, there were 26.6 million sufferers worldwide. Alzheimer's is predicted to affect 1 in 85 people globally by 2050.

Symptoms and Prognosis


Although Alzheimer's disease develops differently for every individual, there are many common symptoms. Early symptoms are often mistakenly thought to be "age-related" concerns or manifestations of stress. In the early stages, the most common symptom is difficulty in remembering recent events. When AD is suspected, the diagnosis is usually confirmed with tests that evaluate behavior and thinking abilities, often followed by a brain scan if available.

As the disease advances, symptoms can include confusion, irritability, aggression, mood swings, trouble with language, and long-term memory loss. As the sufferer declines they often withdraw from family and society. Gradually, bodily functions are lost, ultimately leading to death. Since the disease is different for each individual, predicting how it will affect the person is difficult. AD develops for an unknown and variable amount of time before becoming fully apparent, and it can progress undiagnosed for years. On average, the life expectancy following diagnosis is approximately seven years. Fewer than three percent of individuals live more than fourteen years after diagnosis.

Causes


The cause for most Alzheimer's cases is still essentially unknown (except for 1% to 5% of cases where genetic differences have been identified). Several competing hypotheses exist trying to explain the cause of the disease. The oldest, on which most currently available drug therapies are based, is the cholinergic hypothesis, which proposes that AD is caused by reduced synthesis of the neurotransmitter acetylcholine. The cholinergic hypothesis has not maintained widespread support, largely because medications intended to treat acetylcholine deficiency have not been very effective. Other cholinergic effects have also been proposed, for example, initiation of large-scale aggregation of amyloid, leading to generalized neuroinflammation.

In 1991, the amyloid hypothesis postulated that amyloid beta (Aβ) deposits are the fundamental cause of the disease. Support for this postulate comes from the location of the gene for the amyloid beta precursor protein (APP) on chromosome 21, together with the fact that people with trisomy 21 (Down syndrome) who have an extra gene copy almost universally exhibit AD by 40 years of age. Also APOE4, the major genetic risk factor for AD, leads to excess amyloid buildup in the brain. Further evidence comes from the finding that transgenic mice that express a mutant form of the human APP gene develop fibrillar amyloid plaques and Alzheimer's-like brain pathology with spatial learning deficits.



Amyloid Plaque Formation: Enzymes act on the APP (Amyloid precursor protein) and cut it into fragments of protein, one of which is called beta-amyloid and its crucial in the formation of senile plaques in AD

In 2009, this theory was updated, suggesting that a close relative of the beta-amyloid protein, and not necessarily the beta-amyloid itself, may be a major culprit in the disease. The theory holds that an amyloid-related mechanism that prunes neuronal connections in the brain in the fast-growth phase of early life may be triggered by ageing-related processes in later life to cause the neuronal withering of Alzheimer's disease. N-APP, a fragment of APP from the peptide's N-terminus, is adjacent to beta-amyloid and is cleaved from APP by one of the same enzymes. N-APP triggers the self-destruct pathway by binding to a neuronal receptor called death receptor 6 (DR6, also known as TNFRSF21). DR6 is highly expressed in the human brain regions most affected by Alzheimer's, so it is possible that the N-APP/DR6 pathway might be hijacked in the ageing brain to cause damage. In this model, beta-amyloid plays a complementary role, by depressing synaptic function.



Brain Scans: This is a combination of two brain diagrams in one for comparison. In the left normal brain, in the right brain of a person with Alzheimer's disease.

This page titled 29.6D: Alzheimer's Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6E: Brain Tumors

A brain tumor is a pathological abnormal growth of cells in the brain.

Learning Objectives

- Describe the affects of brain tumors

Key Points

- A brain tumor is a neoplasm, or abnormal growth of cells, within the brain or the spinal canal.
- Detection of brain tumors usually occurs in advanced stages when the presence of the tumor has caused unexplained neurological symptoms.
- Neurological symptoms of a brain tumor depend on size and location, but may include impaired judgment, memory loss, lack of recognition, spatial orientation disorders, personality or emotional changes, visual impairment, and more.
- Treatment for a brain tumor depends on the tumor type and location and may include surgery, radiotherapy, and chemotherapy.
- Detection of brain tumors usually occurs in advanced stages when the presence of the tumor has caused unexplained neurological symptoms.
- Neurological symptoms of a brain tumor depend on size and location, but may include impaired judgment, memory loss, lack of recognition, spatial orientation disorders, personality or emotional changes, visual impairment, and more.
- The most common primary brain tumors are gliomas, meningiomas, pituitary adenomas, and nerve sheath tumors.

Key Terms

- **chemotherapy**: Chemical treatment to kill or halt the replication and/or spread of cancerous cells in a patient.
- **glioma**: A tumor that arises from glial cells in the brain or spinal cord.


EXAMPLES

An example of a highly treatable brain tumor subtype is medulloblastoma. An example of a brain tumor subtype with a poor prognosis is glioblastoma multiforme. Other examples of brain tumor subtypes include oligodendrogliomas and astrocytomas.

A brain tumor is an intracranial solid neoplasm—a tumor (defined as an abnormal growth of cells)—within the brain or the central spinal canal.

Brain tumors include all tumors inside the cranium or in the central spinal canal. They are created by an abnormal and uncontrolled cell division, usually in the brain itself, but also in lymphatic tissue, blood vessels, the cranial nerves, the brain envelopes (meninges), skull, pituitary gland, or pineal gland. Within the brain itself, the involved cells may be neurons or glial cells (which include astrocytes, oligodendrocytes, and ependymal cells). Brain tumors may also spread from cancers primarily located in other organs (metastatic tumors).

Signs and Symptoms



Metastasis in the Right Hemisphere: Brain metastasis in the right cerebral hemisphere from lung cancer shown on T1-weighted magnetic resonance imaging with intravenous contrast.

Visibility of signs and symptoms of brain tumors mainly depends on two factors: tumor size (volume) and tumor location. The moment that symptoms become apparent, either to the person or people around him (symptom onset), is an important milestone in the course of the diagnosis and treatment of the tumor. The symptom onset—in the timeline of the development of the neoplasm—depends in many cases on the nature of the tumor but in many cases is also related to the change of the neoplasm from “benign” (i.e., slow-growing/late symptom onset) to more malignant (fast growing/early symptom onset).

Symptoms of solid neoplasms of the brain (primary brain tumors and secondary tumors alike) can be divided into three main categories:

- Consequences of intracranial hypertension: Large tumors or tumors with extensive perifocal swelling (edema) inevitably lead to elevated intracranial pressure (intracranial hypertension), which translates clinically into headaches, vomiting (sometimes without nausea), altered state of consciousness (somnolence, coma), dilation of the pupil on the side of the lesion (anisocoria), papilledema (prominent optic disc at the funduscopic eye examination).
- Dysfunction: Depending on the tumor location and the damage it may have caused to surrounding brain structures, either through compression or infiltration, any type of focal neurologic symptoms may occur, such as cognitive and behavioral impairment, personality changes, ataxia, visual field impairment, impaired sense of smell, impaired hearing, double vision, dizziness; more severe symptoms might occur, such as including partial paralysis or impairment to swallow.
- Irritation: abnormal fatigue, weariness, absences, tremors, and epileptic seizures.

Treatment

Neurosurgeons take the time to observe the evolution of the neoplasm before proposing a management plan to the patient and his/her relatives. Various types of treatment are available depending on neoplasm type and location, and they may be combined to give the best chances of survival:

- Surgery: complete or partial resection of the tumor to remove as many tumor cells as possible
- Radiotherapy: the most commonly used treatment for brain tumors, the tumor is targeted with alpha and beta rays to shrink the tumor
- Chemotherapy: a treatment option for cancer, however it is seldom used to treat brain tumors as the blood and brain barrier prevents the drugs from reaching the cancerous cells

A variety of experimental therapies are available through clinical trials. Survival rates in primary brain tumors depend on the type

of tumor, age, functional status of the patient, the extent of surgical tumor removal, and other factors specific to each case.

This page titled 29.6E: Brain Tumors is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.6F: Amyotrophic Lateral Sclerosis

Amyotrophic lateral sclerosis (ALS) is caused by degeneration of upper and lower motor neurons, resulting in muscle weakness and atrophy.

Learning Objectives

- Describe amytrophic lateral sclerosis and its effect on the body

Key Points

- ALS is sometimes referred to as Lou Gehrig's disease after the New York Yankees baseball player who was diagnosed with the disease in 1939.
- ALS is characterized by rapidly progressive weakness, muscle atrophy, muscle spasticity, difficulty speaking, difficulty swallowing, and decline in breathing ability. Most patients with ALS die of respiratory failure or pneumonia.
- For the majority of patients without a family history of the disease, there is no known cause for ALS. Familial ALS is a heritable condition that can be traced to the dysfunction of specific genes.
- Most people with ALS die of respiratory failure or pneumonia. The median survival time from onset to death is around 39 months, and only 4% survive longer than 10 years.
- In "respiratory onset" ALS, the intercostal muscles that support breathing are affected first.
- For the majority of patients without a family history of the disease, there is no known cause for ALS. Familial ALS is a heritable condition that can be traced to the dysfunction of specific genes.

Key Terms

- **lateral**: Pertaining to the left or right of the body; further from the midline.
- **amyotrophic lateral sclerosis**: A chronic, progressive disease characterized by degeneration of the central nervous system and the loss of voluntary muscle control.

EXAMPLES

An example of a gene mutated in some cases of familial ALS is superoxide dismutase, or SOD1, which is an enzyme that acts as a powerful antioxidant.

Amyotrophic lateral sclerosis (ALS), also referred to as motor neuron disease in British English, is the most common form of the motor neuron diseases. The condition is sometimes called Lou Gehrig's disease in the U.S., after the New York Yankees baseball player who was diagnosed with the disease in 1939. The disorder is characterized by rapidly progressive weakness, muscle atrophy and fasciculations, muscle spasticity, difficulty speaking (dysarthria), difficulty swallowing (dysphagia), and decline in breathing ability.

Symptoms of ALS



Coronal MRI of an ALS patient: MRI (parasagittal FLAIR) demonstrates increased T2 signal within the posterior part of the internal capsule and can be tracked to the subcortical white matter of the motor cortex, outlining the corticospinal tract), consistent with the clinical diagnosis of ALS.

The disorder causes muscle weakness and atrophy throughout the body as a result of degeneration of the upper and lower motor neurons. Unable to function, the muscles weaken and atrophy. Affected individuals may ultimately lose the ability to initiate and control all voluntary movement, although bladder and bowel sphincters and the muscles responsible for eye movement are usually, but not always, spared. Cognitive function is generally spared for most patients although some (~5%) also have frontotemporal dementia. A higher proportion of patients (~30%–50%) also have more subtle cognitive changes that may go unnoticed but are revealed by detailed neuropsychological testing. Sensory nerves and the autonomic nervous system are generally unaffected meaning the majority of people with ALS will maintain sight, hearing, touch, smell, and taste. Bladder and bowel functions are also rarely affected by ALS.

The earliest symptoms of ALS are typically obvious weakness and/or muscle atrophy. Other presenting symptoms include muscle fasciculation (twitching), cramping, or stiffness of affected muscles; muscle weakness affecting an arm or a leg; and/or slurred and nasal speech. The parts of the body affected by early symptoms of ALS depend on which motor neurons in the body are damaged first. About 75% of people contracting the disease experience “limb onset” ALS (i.e., first symptoms in the arms or legs). Patients with the leg onset form may experience awkwardness when walking or running or notice that they are tripping or stumbling, often with a “dropped foot” which drags gently along the ground. Arm-onset patients may experience difficulty with tasks requiring manual dexterity such as buttoning a shirt, writing, or turning a key in a lock. Occasionally, the symptoms remain confined to one limb for a long period of time or for the whole length of the illness; this is known as monomelic amyotrophy.

Over time, patients experience increasing difficulty moving, swallowing (dysphagia), and speaking or forming words (dysarthria). Symptoms of upper motor neuron involvement include tight and stiff muscles (spasticity) and exaggerated reflexes (hyperreflexia) including an overactive gag reflex. Symptoms of lower motor neuron degeneration include muscle weakness and atrophy, muscle cramps, and fleeting twitches of muscles that can be seen under the skin (fasciculations). To be diagnosed with ALS, patients must have signs and symptoms of both upper and lower motor neuron damage that cannot be attributed to other causes.

Difficulty swallowing and chewing making eating normally very difficult and increase the risk of choking or aspirating food into the lungs. In later stages of the disease, aspiration pneumonia and maintaining a healthy weight can become a significant problem and may require insertion of a feeding tube. As the diaphragm and intercostal muscles (rib cage) that support breathing weaken, measures of lung function such as forced vital capacity and inspiratory pressure diminish. In respiratory onset ALS, this may occur before significant limb weakness is apparent. Most people with ALS die of respiratory failure or pneumonia. The median survival time from onset to death is around 39 months, and only 4% survive longer than 10 years. The best-known person with ALS, Stephen Hawking, has lived with the disease for more than 40 years, though his is an unusual case.

Causes and Treatments of ALS

For patients without a family history of the disease, which includes ~95% of cases, there is no known cause for ALS. Potential causes for which there is inconclusive evidence includes head trauma, military service, and participation in elite sports. Many other potential causes including chemical exposure, electromagnetic field exposure, occupation, physical trauma, and electric shock have been investigated but without consistent findings.

Riluzole (Rilutek) as of 2011 is the only treatment that has been found to improve survival, but only to a modest extent. It lengthens survival by several months, and may have a greater survival benefit for those with a bulbar onset. It also extends the time before a person needs ventilation support. Other treatments for ALS are designed to relieve symptoms and improve the quality of life for patients. This supportive care is best provided by multidisciplinary teams of health care professionals working with patients and caregivers to keep patients as mobile and comfortable as possible.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: *CC BY-SA: Attribution-ShareAlike*

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Traumatic brain injury. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Traumatic_brain_injury. **License**: *CC BY-SA: Attribution-ShareAlike*
- Glasgow Coma Scale. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Glasgow%20Coma%20Scale. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg. **License**: *Public Domain: No Known Copyright*
- Stroke. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Stroke. **License**: *CC BY-SA: Attribution-ShareAlike*
- thrombosis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/thrombosis. **License**: *CC BY-SA: Attribution-ShareAlike*
- embolus. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/embolus. **License**: *CC BY-SA: Attribution-ShareAlike*
- embolism. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/embolism. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg. **License**: *Public Domain: No Known Copyright*
- Parachemableedwithedema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parachemableedwithedema.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Transient ischemic attack. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Transient_ischemic_attack. **License**: *CC BY-SA: Attribution-ShareAlike*
- ischemia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/ischemia. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg. **License**: *Public Domain: No Known Copyright*
- Parachemableedwithedema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parachemableedwithedema.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Atrial myxoma. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Atrial_myxoma.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Alzheimer's disease. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Alzheimer's_disease. **License**: *CC BY-SA: Attribution-ShareAlike*
- amyloid-beta peptide. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/amyloid-beta%20peptide. **License**: *CC BY-SA: Attribution-ShareAlike*
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg. **License**: *Public Domain: No Known Copyright*
- Parachemableedwithedema. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Parachemableedwithedema.png. **License**: *CC BY-SA: Attribution-ShareAlike*
- Atrial myxoma. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Atrial_myxoma.jpg. **License**: *CC*

BY-SA: Attribution-ShareAlike
- Amyloid-plaque formation-big. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Amyloid-plaque_formation-big.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:Alzheimer's disease brain comparison.jpg - Wikimedia Foundation. **Provided by**: Wikimedia Foundation. **Located at**: **wikimediafoundation.org/wiki/File:COMPARISONSLICE_HIGH.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Brain tumor. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Brain_tumor**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- chemotherapy. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/chemotherapy**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- glioma. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/glioma**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg**. **License**: ***Public Domain: No Known Copyright***
- Parachemableedwithedema. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Parachemableedwithedema.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Atrial myxoma. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Atrial_myxoma.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amyloid-plaque formation-big. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Amyloid-plaque_formation-big.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:Alzheimer's disease brain comparison.jpg - Wikimedia Foundation. **Provided by**: Wikimedia Foundation. **Located at**: **wikimediafoundation.org/wiki/File:COMPARISONSLICE_HIGH.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hirnmetastase MRT-T1 KM. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Hirnmetastase_MRT-T1_KM.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amyotrophic lateral sclerosis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Amyotrophic_lateral_sclerosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- lateral. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/lateral**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- amyotrophic lateral sclerosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/amyotrophic_lateral_sclerosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Trauma subdural arrows. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Trauma_subdural_arrows.jpg**. **License**: ***Public Domain: No Known Copyright***
- Parachemableedwithedema. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Parachemableedwithedema.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Atrial myxoma. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/File:Atrial_myxoma.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Amyloid-plaque formation-big. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Amyloid-plaque_formation-big.jpg**. **License**: ***Public Domain: No Known Copyright***
- File:Alzheimer's disease brain comparison.jpg - Wikimedia Foundation. **Provided by**: Wikimedia Foundation. **Located at**: **wikimediafoundation.org/wiki/File:COMPARISONSLICE_HIGH.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hirnmetastase MRT-T1 KM. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Hirnmetastase_MRT-T1_KM.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ALS Coronal. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:ALS_Coronal.jpg**. **License**: ***CC BY-SA: Attribution-ShareAlike***

This page titled 29.6F: Amyotrophic Lateral Sclerosis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.7: DISEASES AND DISORDERS OF THE ENDOCRINE SYSTEM

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.7A: Pancreatic Islet Disorders: Diabetes and Hyperinsulinism

29.7B: Posttraumatic Stress Disorder (PTSD)

This page titled 29.7: Diseases and Disorders of the Endocrine System is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.7A: Pancreatic Islet Disorders: Diabetes and Hyperinsulinism

Hyperinsulinism refers to an above-normal level of insulin in the blood of a person or animal.

Learning Objectives

- Distinguish between the two types of hyperinsulinism: hyperglycemia and hypoglycemia

Key Points

- Hyperinsulinism can be associated with several types of medical problems, which can be roughly divided into two broad categories: hyperglycemia and hypoglycemia.
- Hyperglycemia or Hyperglycæmia, or high blood sugar, is a condition in which an excessive amount of glucose circulates in the blood plasma. This is generally a glucose level higher than (200 mg/dl).
- Hypoglycemia, or low blood sugar is an abnormally-diminished content of glucose in the blood. The principal problems arise from an inadequate supply of glucose to the brain, resulting in impairment of function (neuroglycopenia).

Key Terms

- **hyperinsulinism**: Hyperinsulinism refers to an above normal level of insulin in the blood of a person or animal.
- **diabetes**: A group of metabolic diseases whereby a person (or other animal) has high blood sugar due to an inability to produce, metabolize, or respond to the hormone insulin.
- **hypoglycemia**: A too low level of blood glucose.

Hyperinsulinemia

This refers to an above-normal level of insulin in the blood of a person or animal. Normal insulin secretion and blood levels are closely related to the level of glucose in the blood, so that a given level of insulin can be normal for one blood glucose level but low or high for another. Hyperinsulinism can be associated with several types of medical problems, which can be roughly divided into two broad and largely non-overlapping categories: those tending toward reduced sensitivity to insulin and high blood glucose levels (hyperglycemia), and those tending toward excessive insulin secretion and low glucose levels (hypoglycemia).

Hyperglycemia

Also known as hyperglycæmia, or “high blood sugar,” this is a condition in which an excessive amount of glucose circulates in the blood plasma. This is generally a glucose level higher than (200 mg/dl). Reference ranges for blood tests are 11.1 mmol/l, but symptoms may not start to become noticeable until even higher values such as 250–300 mg/dl or 15–20 mmol/l.

A subject with a consistent range above 126 mg/dl or 7 mmol/l is generally held to have hyperglycemia. Chronic levels exceeding 7 mmol/l (125 mg/dl) can produce organ damage.

Hypoglycemia

Hypoglycemia (not to be confused with hyperglycemia), or “low blood sugar,” is an abnormally-diminished content of glucose in the blood. The term literally means “low sugar blood.”

It can produce a variety of symptoms and effects but the principal problems arise from an inadequate supply of glucose to the brain, resulting in impairment of function (neuroglycopenia). Effects can range from mild dysphoria to more serious issues such

as seizures, unconsciousness, and (rarely) permanent brain damage or death.


The most common forms of hypoglycemia occur as a complication of treatment of diabetes mellitus with insulin or oral medications. Hypoglycemia is less common in non-diabetic persons, but can occur at any age. Among the causes are excessive insulin produced in the body (hyperinsulinemia), inborn error of metabolism, medications and poisons, alcohol, hormone deficiencies, prolonged starvation, alterations of metabolism associated with infection, and organ failure.

Hypoglycemia is treated by restoring the blood glucose level to normal by the ingestion or administration of dextrose or carbohydrate foods. In more severe circumstances it is treated by injection or infusion of glucagon.

Recurrent hypoglycemia may be prevented by reversing or removing the underlying cause, by increasing the frequency of meals, with medications like diazoxide, octreotide, or glucocorticoids, or by surgical removal of much of the pancreas.

The level of blood glucose low enough to define hypoglycemia may be different for different people, in different circumstances, and for different purposes, and occasionally has been a matter of controversy. Most healthy adults maintain fasting glucose levels above 4.0 mmol/L (72 mg/dl), and develop symptoms of hypoglycemia when the glucose falls below 4 mmol/L.

It can sometimes be difficult to determine whether a person's symptoms are due to hypoglycemia. Criteria referred to as Whipple's triad are used to determine a diagnosis of hypoglycemia.



Glucose and Insulin Levels: Insulin rises in healthy adults as glucose intake increases.

This page titled 29.7A: Pancreatic Islet Disorders: Diabetes and Hyperinsulinism is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.7B: Posttraumatic Stress Disorder (PTSD)

Posttraumatic stress disorder (PTSD) develops after exposure to an event that is so stressful for an individual that it becomes traumatic.

Learning Objectives

- Describe the role of the endocrine system in post-traumatic stress disorder (PTSD)

Key Points

- Three areas of the brain whose function may be altered in PTSD are the prefrontal cortex, the amygdala, and the hippocampus.
- SSRIs (selective serotonin reuptake inhibitors) are considered to be a first-line drug treatment. SSRIs include citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, and sertraline.
- Cognitive behavioral therapy (CBT) seeks to change the way a trauma victim feels and acts by changing the patterns of thinking or behavior, or both, responsible for negative emotions.
- PTSD symptoms may result when a traumatic event causes an over-reactive adrenaline response, which creates deep neurological patterns in the brain. These patterns can persist long after the event that triggered the fear, making an individual hyper-responsive to future fearful situations.
- Multiple studies show that parental PTSD and other posttraumatic disturbances in parental psychological functioning can, despite the best efforts of traumatized parents, interfere with their response to their child as well as their child's response to trauma.

Key Terms

- **Posttraumatic stress disorder**: Posttraumatic stress disorder (PTSD) develops after exposure to an event that is so stressful for an individual that it becomes traumatic.
- **cortisol**: A steroid hormone (also called hydrocortisone), produced by the adrenal cortex, that regulates the metabolism of carbohydrates and maintains blood pressure.
- **SSRI**: Selective serotonin re-uptake inhibitors are a class of compounds typically used as antidepressants in the treatment of depression, anxiety disorders, and some personality disorders.

Posttraumatic stress disorder (PTSD) develops after exposure to an event that is so stressful for an individual that it becomes traumatic. Historically (DSM-IV) categorized as an anxiety disorder, it is now categorized under "Trauma- and Stressor-Related Disorders" in the DSM 5.

The traumatic event may involve the threat of death to oneself or to someone else, or to one's own or someone else's physical, sexual, or psychological integrity, overwhelming the individual's ability to cope. As an effect of psychological trauma, PTSD is less frequent and more enduring than the more commonly seen acute stress response. Diagnostic symptoms for PTSD include re-experiencing the original trauma through flashbacks or nightmares, avoidance of stimuli associated with the trauma, and increased arousal, such as difficulty falling or staying asleep, anger, and hypervigilance. Additionally, PTSD and other posttraumatic disturbances in parental psychological functioning can, despite the best efforts of traumatized parents, interfere with their response to their child as well as their child's response to trauma.

PTSD is believed to be caused by experiencing any of a wide range of events which produces intense negative feelings of "fear, helplessness, or horror" in the observer or participant. Sources of such feelings may include (but are not limited to):

- Experiencing or witnessing childhood or adult physical, emotional, or sexual abuse
- Experiencing or witnessing physical assault, adult experiences of sexual assault, accidents, drug addiction, illnesses, and medical complications
- Employment in occupations exposed to war (such as soldiers) or disaster (such as emergency service workers)
- Getting a diagnosis of a life-threatening illness

Causes

Evolutionary psychology views different types of fears and reactions to fears as adaptations that may have been useful in the ancestral environment in order to avoid or cope with various threats. Mammals generally display several defensive behaviors roughly dependent on how close the threat is: avoidance, vigilant immobility, withdrawal, aggressive defense, appeasement, and finally complete, frozen immobility (the last possibly to confuse a predator's attack reflex or to simulate a dead and contaminated body). PTSD may correspond to and be caused by overactivation of such fear circuits. Thus, PTSD avoidance behaviors may correspond to mammal avoidance of and withdrawal from threats.

PTSD symptoms may result when a traumatic event causes an over-reactive adrenaline response, which creates deep neurological patterns in the brain. PTSD causes biochemical changes in the brain and body that differ from other psychiatric disorders such as major depression. Most people with PTSD also show a low secretion of cortisol and high secretion of catecholamines in urine, with a norepinephrine to cortisol ratio consequently higher than comparable non-diagnosed individuals. These findings suggest abnormality in the hypothalamic-pituitary-adrenal (HPA) axis.

Although most people (50–90%) encounter trauma over a lifetime, only about 8% develop full PTSD. Vulnerability to PTSD presumably stems from an interaction of biological diathesis, early childhood developmental experiences, and trauma severity. Predictor models have consistently found that childhood trauma, chronic adversity, and familial stressors increase risk for PTSD as well as biological markers of risk for PTSD after a traumatic event in adulthood. This effect of childhood trauma, which is not well-understood, may be a marker for both traumatic experiences and attachment problems.


Treatments

Modest benefits have been seen from early access to cognitive behavioral therapy, as well as from some medications such as propranolol. Critical incident stress management has been suggested as a means of preventing PTSD, but subsequent studies suggest the likelihood of its producing iatrogenic outcomes. A review of multiple studies, involving a number of different post-event psychological interventions structured to prevent PTSD ."..did not find any evidence to support the use of an intervention offered to everyone," and that ."..multiple session interventions may result in a worse outcome than no intervention for some individuals."


Cognitive behavioral therapy (CBT) seeks to change the way a trauma victim feels and acts by changing the patterns of thinking or behavior, or both, responsible for negative emotions. CBT have been proven to be an effective treatment for PTSD and is currently considered the standard of care for PTSD by the United States Department of Defense. In CBT, individuals learn to identify thoughts that make them feel afraid or upset and replace them with less distressing thoughts.

Eye movement desensitization and reprocessing (EMDR) is specifically targeted as a treatment for PTSD. Based on the evidence of controlled research, the American Psychiatric Association and the United States Department of Veterans Affairs and Department of Defense have placed EMDR in the highest category of effectiveness and research support in the treatment of trauma.

SSRIs (selective serotonin reuptake inhibitors) are considered to be a first-line drug treatment. SSRIs for which there are data to support use include: citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, and sertraline. Among the anti-depressants described in this section, bupropion and venlafaxine have the lowest patient drop-out rates. Sertraline, fluoxetine, and nefazodone have a modestly higher drop-out rate (~15%), and the heterocyclics and paroxetine have the highest rates (~20%+).



The hypothalmic-pituitary-adrenal axis.: The HPA axis is a major biological mechanism of stress and stress response



Post-traumatic stress disorder worldwide.: Incidences of PTSD by country (per 100,000 persons). The redder the country, the higher the incidence of PTSD.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Hyperglycemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hyperglycemia. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hyperinsulinism. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hyperin..._diabetes_risk. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hypoglycemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Hypoglycemia. **License**: ***CC BY-SA:***

Attribution-ShareAlike

- hyperinsulinism. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/hyperinsulinism**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- diabetes. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/diabetes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoglycemia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoglycemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Principles of Biochemistry/Glucose,Glycogen and Diabetes. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Princip...n_and_Diabetes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Post-traumatic stress disorder. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Post-tr...tress_disorder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cortisol. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cortisol**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Posttraumatic stress disorder. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/Posttr...tress+disorder**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- SSRI. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/SSRI**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Principles of Biochemistry/Glucose,Glycogen and Diabetes. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Princip...n_and_Diabetes**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Basic HPA Axis. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Basic_HPA_Axis.jpeg**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Post-traumatic stress disorder. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Post-tr...tress_disorder**. **License**:

This page titled 29.7B: Posttraumatic Stress Disorder (PTSD) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.8: BLOOD DISEASES AND DISORDERS

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.8A: Hemostasis Disorders

29.8B: Anemia

29.8C: Sickle-Cell Disease

29.8D: Leukemia

This page titled 29.8: Blood Diseases and Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.8A: HEMOSTASIS DISORDERS

The most common blood disorders involve defects in the amount or activity of factors involved in coagulation.

Learning Objectives

- Differentiate among the hemostasis disorders of Von Willebran disease, hemophilia, and factor V Leiden thrombophilia

Key Points

- Von Willebrand disease is the most common inherited bleeding disorder. People with type 1 and type 2 von Willebrand disease may have mild-to-moderate bleeding symptoms, while those with type 3 can have severe bleeding episodes.
- Hemophilia is a disease where there is a low level of, or are no, blood proteins important for clotting, causing an inability to produce blood clots. There are two types: Type A, which is a deficiency in factor VIII, and Type B, also known as Christmas disease, which is a deficiency in factor IX.
- The opposite of hemophilia is Factor V Leiden thrombophilia, a disorder of human factor V blood protein that causes a hypercoagulability disorder or overactive clotting resulting in dangerous blood clots.
- In sickle- cell disease, red blood cells undergo a change in shape upon deoxygenation because of polymerization of the abnormal sickle hemoglobin.
- Polycythemia is a condition in which there is a net increase in the total circulating erythrocyte (red blood cell).
- Leukemia is a cancer of the blood or bone marrow characterized by an abnormal proliferation of blood cells, usually white blood cells (leukocytes).

Key Terms

- **Hemophilia**: A group of hereditary genetic disorders that impair the body's ability to control blood clotting or coagulation, which is used to stop bleeding when a blood vessel is broken.
- **coagulation**: The process by which blood forms solid clots.
- **thrombophilia**: The tendency to develop thrombosis (blood clots) due to an abnormality in the system of coagulation.

The most common blood disorders, Von Willebrand disease, hemophilia, and Factor V Leiden thrombophilia, all involve defects in the amount or activity of factors involved in coagulation.

Von Willebrand Disease

Von Willebrand disease is the most common inherited bleeding disorder, and it affects both men and women equally. Von Willebrand disease is similar to hemophilia in that it involves a deficiency in the ability of blood to clot properly. Those affected by von Willebrand disease will have low levels of von Willebrand factor, a protein that helps the blood to clot, and/or their von Willebrand factor does not work properly.

There are three types of von Willebrand disease: type 1, type 2, and type 3. People with type 1 and type 2 von Willebrand disease may have the following mild-to-moderate bleeding symptoms: easy bruising, nosebleeds, bleeding from the gums after a dental procedure, heavy menstrual bleeding in women, blood in their stools or urine, and excessive bleeding after a cut or other accident or surgery.

Type 1 von Willebrand disease is the mildest and most common form of the disease. In this type of disease, there is a low level of von Willebrand factor. The level of factor VIII, another protein involved in clotting, may also be lower than normal. About 3 out of 4 people diagnosed with von Willebrand disease have type 1.


Type 2 von Willebrand disease has four subtypes and ranges from mild to moderate in severity. In this type of disease, there is a defect in von Willebrand factor resulting in improper activity.

Type 3 von Willebrand disease is very rare and is the most severe form. In this type of disease, individuals usually do not produce von Willebrand factor and have very low levels of factor VIII. Symptoms of type 3 von Willebrand disease might include any of the symptoms of types 1 and 2 and also include severe bleeding episodes for no reason, which can be life threatening if not treated immediately. Bleeding into soft tissues or joints causing severe pain and swelling is another symptom.

Many people with von Willebrand disease do not require treatment to manage the disease. However, if treatment is necessary, it may include a range of different interventions depending on the severity. These involve medicine to increase the level of von Willebrand factor in the blood, medicine to prevent the breakdown of clots, medicine to control heavy menstrual bleeding in women, or injection of clotting factor concentrates.

Hemophilia

Hemophilia is a disease where there is a low levels of, or none, of a blood protein important for clotting, causing an inability to produce blood clots. There are two types of hemophilia: Type A, which is a deficiency in factor VIII, and Type B, also known as Christmas disease, which is a deficiency in factor IX. Since people with hemophilia do not have the ability to make blood clots, even a small cut can result in severe bleeding, or the smallest bump or jar to the body could cause severe bruising that does not heal for months. Hemophilia is a recessive, sex-linked inheritable disorder, which is more common in men than women. Treatment typically involves replacement of the missing blood factors through blood transfusions.




Inheritance of X-linked Recessive Traits: Hemophilia is an x-linked recessive inheritable disorder.

Since the 1980s, the primary leading cause of death of people with severe hemophilia has shifted from hemorrhage to HIV/AIDS acquired through treatment with contaminated blood products. The second leading cause of death related to severe hemophilia complications is intracranial hemorrhage, which today accounts for one-third of all deaths of patients with hemophilia.

Factor V Leiden Thrombophilia

The opposite of hemophilia, is Factor V Leiden thrombophilia, a disorder of human factor V blood protein that causes a hypercoagulability disorder, or, overactive clotting, resulting in dangerous blood clots. In this disorder, the Leiden variant of factor V cannot be inactivated by activated protein C, as it would be in a person with normal factor V, resulting in excess clotting. Factor V Leiden is the most common hereditary hypercoagulability disorder among Eurasians. Those that have it are at a slightly higher risk of developing blood clots than those without. Those that test positive for factor V should avoid oral contraceptives, obesity, smoking, and high blood pressure to reduce their risk of developing blood clots.



Blood clot diagram.: Blood clot diagram.

29.8B: Anemia

This page titled 29.8A: Hemostasis Disorders is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.8B: Anemia

Anemia is a decrease in number of red blood cells or less than the normal quantity of hemoglobin in the blood resulting in tissue hypoxia.

Learning Objectives

- Distinguish among the types of anemia

Key Points

- The three main classes of anemia include anemia due to hemorrhage (excessive blood loss), hemolysis (blood cell destruction), or ineffective hematapoiesis (deficient red blood cell production).
- Symptoms of anemia include fatigue, malaise, heart enlargement, and impaired concentration.
- Anemia may be diagnosed using a complete blood count (CBC) and/or reticulocyte counts.
- A “kinetic” approach analyzes decreased RBC production vs. increased RBC destruction or loss.
- Iron deficiency anemia is the most common type of anemia and is particularly common in infants, toddlers, and pregnant women who have above-average needs for iron. It is treated with iron supplements.
- Severe anemia with blood loss may be treated with transfusions, but these come with their own adverse outcomes.

Key Terms

- **hematopoiesis**: The process by which blood cells are produced; hematogenesis.
- **reticulocyte**: An immature red blood cell, having a reticular network of RNA.
- **complete blood count (CBC)**: A test panel requested by a doctor or other medical professional that gives information about the cells in a patient’s blood.

EXAMPLES

At the beginning of the 20th century, hookworm infection was a major and severe problem in the southern United States. Hookworms mature in the small intestine where they bite into the villi of the small intestine and drink blood causing severe anemia. Individuals suffering from hookworm are lethargic, have trouble focusing, and do not appear to be very intelligent. Many negative stereotypes of the South are based on individuals suffering from hookworm. When the Rockefeller Foundation provided funding to purge the worms from these individuals and to build outhouses so that they would not be re-infected, these Southerners became bright, energetic, and able to focus on school and work.

Anemia is a decrease in number of red blood cells (RBCs), or less than the normal quantity of hemoglobin in the blood. It can include the decreased oxygen-binding ability of each hemoglobin molecule due to deformity or lack in numerical development as in some other types of hemoglobin deficiencies.

Because hemoglobin (found inside RBCs) normally carries oxygen from the lungs to the tissues, anemia leads to hypoxia (lack of oxygen) in organs. Since all human cells depend on oxygen for survival, varying degrees of anemia can have a wide range of clinical consequences.

Types of Anemia

Anemia is the most common disorder of the blood. There are several kinds, produced by a variety of underlying causes. It can be classified in a variety of ways, based on the morphology of RBCs, underlying etiologic mechanisms, and discernible clinical spectra, to mention a few.


The three main classes of anemia include excessive blood loss (acutely, such as a hemorrhage or chronically, through low-volume

loss), excessive blood cell destruction (hemolysis), or deficient red blood cell production (ineffective hematopoiesis).

Signs and Symptoms

Anemia goes undiagnosed in many people, as symptoms can be minor or vague. The signs and symptoms can be related to the anemia itself, or the underlying cause. Most commonly, people report non-specific symptoms of a feeling of weakness, or fatigue, general malaise, and, sometimes, poor concentration. They may also report dyspnea (shortness of breath) on exertion.

In very severe anemia, the body may compensate for the lack of oxygen-carrying capability of the blood by increasing cardiac output. This would result in symptoms such as palpitations, angina (if preexisting heart disease is present), intermittent claudication of the legs, and symptoms of heart failure.



Symptoms of Anemia: Image displays the common symptoms of anemia.

Diagnosis

Anemia is typically diagnosed on a complete blood count. Apart from reporting the number of red blood cells and the hemoglobin level, the automatic counters also measure the size of the red blood cells by flow cytometry, which is an important tool in distinguishing between the causes of anemia.

Examination of a stained blood smear using a microscope can also be helpful and is sometimes a necessity in regions of the world where automated analysis is less accessible. Reticulocyte counts, and the “kinetic” approach to anemia, have become more common than in the past in the large medical centers of the United States and some other wealthy nations; in part, because some automatic counters now have the capacity to include reticulocyte counts. This is a quantitative measure of the bone marrow ‘s production of new red blood cells. The reticulocyte production index is a calculation of the ratio between the level of anemia and the extent to which the reticulocyte count has risen in response.

If the degree of anemia is significant, even a “normal” reticulocyte count actually may reflect an inadequate response.

In the morphological approach, anemia is classified by the size of red blood cells. This either done automatically or on microscopic examination of a peripheral blood smear. The size is reflected in the mean corpuscular volume (MCV). This scheme quickly exposes some of the most common causes of anemia. For instance, a microcytic anemia (smaller than usual red blood cells) is often the result of iron deficiency.

In clinical workup, the MCV will be one of the first pieces of information available; so even among clinicians who consider the “kinetic” approach more useful philosophically, morphology will remain an important element of classification and diagnosis.

Iron Deficiency Anemia

This is the most common type of anemia overall and it has many causes. Iron is an essential part of hemoglobin, and low iron levels result in decreased incorporation of hemoglobin into red blood cells. It is due to insufficient dietary intake or absorption of iron to meet the body's needs. Infants, toddlers, and pregnant women have higher-than-average needs. Increased iron intake is also needed to offset blood losses due to digestive tract issues, frequent blood donations, or heavy menstrual periods.

In the United States, the most common cause of iron deficiency is bleeding or blood loss, usually from the gastrointestinal tract. Fecal occult blood testing, upper endoscopy, and lower endoscopy should be performed to identify bleeding lesions. In older men and women, the chances are higher that bleeding from the gastrointestinal tract could be due to colon polyp or colorectal cancer.

Worldwide, the most common cause of iron deficiency anemia is parasitic infestation (hookworm, amebiasis, schistosomiasis, and whipworm).

This page titled 29.8B: Anemia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.8C: Sickle-Cell Disease

Sickle-cell disease is an autosomal recessive genetic blood disorder in which red blood cells assume a rigid sickle shape.

Learning Objectives

- List the ways in which sickle-cell disease can present in affected individuals

Key Points

- Sickle-cell disease is common in tropical and sub-tropical regions because being a carrier, carrying a single sickle-cell mutation (sickle cell trait), affords some resistance to malaria.
- Sickle cell crisis, an acute condition occurring in patients with sickle-cell disease, include: vaso-occlusive crisis, splenic sequestration crisis, aplastic crisis, and hemolytic crisis.
- Complications from sickle-cell disease include stoke and silent stroke, decreased immunity due to hypersplenism, avascular necrosis of joints, chronic renal failure, and pulmonary hypertension.
- Sickle-cell disease is often diagnosed with hemoglobin levels from a complete blood count and detection of sickling of the red blood cells.


Key Terms

- **fitness**: A central idea in evolutionary biology; describes the ability to both survive and reproduce.
- **tachycardia**: A rapid resting heart rate, especially one above 100 beats per minute.

EXAMPLES

In the United States, sickle cell trait is most common among African Americans. However, anyone with ancestors from malarial zones should undergo genetic screening for sickle cell trait prior to pregnancy. This includes Southerners of European ancestry both because some of their ancestors may have come from circum-Mediterranean malarial zones, but also because recent DNA analysis has shown that a significant percentage have African ancestry. This is not surprising since those of African descent formed the largest population segment in the South for 200 or 300 years.

Sickle-cell disease (SCD), or sickle-cell anemia, is an autosomal recessive genetic blood disorder with overdominance characterized by red blood cells that assume an abnormal, rigid, and sickle shape. Sickling decreases the cells' flexibility and results in a risk of various complications, including shortened life expectancy. It is caused by a change in hemoglobin's primary structure.



Sickle cell anemia is caused by a change in hemoglobin's primary structure: This scanning electron micrograph shows red blood cells. Some exhibit the normal donut-shaped morphology while others exhibit the sickle shape characteristic of those with sickle-cell trait or sickle-cell disease.

Disease Presentation


Sickle-cell disease, usually presenting in childhood, occurs more commonly in people (or their descendants) from parts of tropical and sub-tropical regions where malaria is or was common. One-third of all indigenous inhabitants of Sub-Saharan Africa carry the gene because, in areas where malaria is common, there is a fitness benefit in carrying only a single sickle-cell gene (sickle cell trait). Those with only one of the two alleles of the sickle-cell disease, while not totally resistant, are more tolerant to malarial infection and show less severe symptoms when infected. The prevalence of the disease in the United States is approximately 1 in 5,000, mostly affecting Americans of Sub-Saharan African descent, according to the National Institutes of Health. In the United States, about 1 out of 500 African-American children born will have sickle-cell anemia.

The term “sickle cell crisis” is used to describe several independent acute conditions occurring in patients with sickle cell disease. Sickle cell disease results in anemia and several types of crises. Most episodes of sickle cell crises last between five and seven days. The vaso-occlusive crisis is caused by sickle-shaped red blood cells that obstruct capillaries and restrict blood flow to an organ, resulting in ischaemia, pain, necrosis, and often organ damage. The frequency, severity, and duration of these crises vary considerably. Because of its narrow vessels and function in clearing defective red blood cells, the spleen is frequently affected in splenic sequestration crisis. This autosplenectomy increases the risk of infection from encapsulated organisms. Preventative antibiotics and vaccinations are recommended for those with such asplenia. Aplastic crises are acute worsenings of the patient’s baseline anemia producing pallor, tachycardia, and fatigue. This crisis is triggered by parvovirus B19, which directly affects erythropoiesis (production of red blood cells) by invading the red cell precursors and multiplying in them and destroying them. Parvovirus infection nearly completely prevents red blood cell production for two to three days. In normal individuals, this is of little consequence, but the shortened red cell life of sickle-cell patients results in an abrupt, life-threatening situation. Hemolytic crises are acute accelerated drops in hemoglobin level. The red blood cells break down at a faster rate. This is particularly common in patients with co-existent G6PD deficiency. Management is supportive, sometimes with blood transfusions.

Diagnosis and Prevention

Sickle cell can be diagnosed by a blood test. In HbSS, the full blood count reveals hemoglobin levels in the range of 6–8 g/dL with a high reticulocyte count (as the bone marrow compensates for the destruction of sickle cells by producing more red blood cells). In other forms of sickle-cell disease, Hb levels tend to be higher. Abnormal hemoglobin forms can also be detected on hemoglobin electrophoresis, a form of gel electrophoresis on which the various types of hemoglobin move at varying speeds.

People who are known carriers of the disease often undergo genetic counseling before they have a child. A test to see if an unborn child has the disease takes either a blood sample from the fetus or a sample of amniotic fluid. Since taking a blood sample from a fetus has greater risks, the latter test is usually used. Children born with sickle-cell disease will undergo close observation by the pediatrician and will require management by a hematologist to assure they remain healthy. These patients will take a 1 mg dose of folic acid daily for life. From birth to five years of age, they will also have to take penicillin daily due to the immature immune system that makes them more prone to early childhood illnesses.



Sickle-cell disease: This image shows normal red blood cells flowing freely in a blood vessel. The inset image shows a cross-section of a normal red blood cell with normal hemoglobin. Figure B shows abnormal, sickled red blood cells blocking blood flow in a blood vessel. The inset image shows a cross-section of a sickle cell with abnormal (sickle) hemoglobin forming abnormal strands.

This page titled 29.8C: Sickle-Cell Disease is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.8D: Leukemia

Leukemia is a term covering a spectrum of cancers of the blood or bone marrow characterized by an increase in immature white blood cells.

Learning Objectives

- Describe the diagnosis and treatment of leukemia

Key Points

- Acute leukemia is a rapid increase in immature white blood cells, which requires immediate treatment and is the most common form of leukemia in children.
- Chronic leukemia is a much slower process with a buildup of more mature but abnormal white blood cells and occurs more often in older patients.
- Myeloid or myelogenous leukemias describe a cancerous change in bone marrow cells that typically mature into red blood cells, some types of white blood cells, and platelets.
- Leukemia results from DNA mutations that activate oncogenes or deactivate tumor suppressor genes.
- The symptoms of leukemia may all be explained by other disease; therefore, leukemia must always be diagnosed by a complete blood count or bone marrow examination.
- Leukemia is typically treated with multi-drug chemotherapy regimens, radiation, or in some cases bone marrow transplant.

Key Terms

- **leukemia**: Any specific form or type of cancer of the blood-forming tissues
- **bone marrow**: The fatty vascular tissue that fills the cavities of bones; the place where new blood cells are produced.
- **complete blood count**: A measure of the number, concentration, and size of white blood cells, platelets, and red blood cells in one's blood.

EXAMPLES

Children with acute leukemia may be candidates for bone marrow transplants. Generally, the best match is another family member. However, in one famous case, no one in the family was a match, so the parents conceived another baby, hoping the baby would provide a match. The baby was born in 1990, and did, indeed, turn out to be a match. She later provided bone marrow for a transplant for her older sister. The transplant was a success. The younger sister, now in her 20s, is happy that she was able to save the life of her older sister, who was dying in 1990, but has now had two decades and counting of a healthy life thanks to her "baby" sister.

Leukemia is a type of cancer of the blood or bone marrow characterized by an abnormal increase in the number of immature white blood cells called "blasts." Leukemia is a broad term covering a spectrum of diseases. It is part of a broader group of diseases affecting the blood, bone marrow, and lymphoid system, which are all known as hematological neoplasms. In 2000, approximately 256,000 children and adults around the world developed some form of leukemia, and 209,000 died from it. About 90% of all leukemias are diagnosed in adults.

Characterization

Clinically and pathologically, leukemia is subdivided into a variety of large groups. The first division is between its acute and chronic forms. Acute leukemia is characterized by a rapid increase in the number of immature blood cells. Crowding due to such cells makes the bone marrow unable to produce healthy blood cells. Immediate treatment is required in acute leukemia due to the rapid progression and accumulation of the malignant cells, which then spill over into the bloodstream and spread to other organs of the body. Acute forms of leukemia are the most common forms of leukemia in children. Chronic leukemia is characterized by

the excessive build up of relatively mature, but still abnormal, white blood cells. Typically taking months or years to mature, the cells are produced at a much higher rate than normal, resulting in many abnormal white blood cells. Whereas acute leukemia must be treated immediately, chronic forms are sometimes monitored for some time before treatment in order to ensure maximum effectiveness of therapy. Chronic leukemia is common in older people, but it can theoretically occur in any age group.


Additionally, the disease is subdivided according to which kind of blood cell is affected. Along these lines, it is divided into lymphoblastic or lymphocytic leukemia, and myeloid or myelogenous leukemia. In lymphoblastic or lymphocytic leukemia, the cancerous change takes place in a type of marrow cell that normally forms lymphocytes, which are infection -fighting immune system cells. In myeloid or myelogenous leukemia, the cancerous change takes place in a type of marrow cell that normally goes on to form red blood cells, some other types of white cells, and platelets.

Causes

Leukemia, like other cancers, results from mutations in the DNA. Certain mutations can trigger leukemia by activating oncogenes or deactivating tumor suppressor genes, thereby disrupting the regulation of cell death, differentiation or division. These mutations may occur spontaneously or as a result of exposure to radiation or carcinogenic substances. Some people have a genetic predisposition towards developing leukemia. In addition to these genetic issues, people with chromosomal abnormalities or certain other genetic conditions have a greater risk of leukemia. For example, people with Down syndrome have a significantly increased risk of developing forms of acute leukemia (especially acute myeloid leukemia), and Fanconi anemia is a risk factor for developing acute myeloid leukemia.

Diagnosis and Treatment

People with leukemia may easily become bruised, bleed excessively, or develop pinprick bleeds (petechiae). White blood cells, which are involved in fighting pathogens, may be suppressed or dysfunctional. This could cause the patient's immune system to be unable to fight off a simple infection or to start attacking other body cells. Because leukemia prevents the immune system from working normally, some patients experience frequent infection, ranging from infected tonsils, sores in the mouth, or diarrhea, to life-threatening pneumonia or opportunistic infections. Finally, the red blood cell deficiency leads to anemia, which may cause dyspnea and pallor. All the symptoms associated with leukemia can be attributed to other diseases. Consequently, leukemia is always diagnosed through medical tests.




Symptoms of Leukemia: Image illustrates the common symptoms of acute or chronic leukemia are shown. Because these can often be attributed to other conditions, a complete blood count and/or bone marrow aspirate are required to successfully diagnose leukemia.

Diagnosis is usually based on repeated complete blood counts and a bone marrow examination following observation of the symptoms. However, in rare cases, blood tests may not show if a patient has leukemia; usually, this may be because the leukemia

is in the early stages or has entered remission. A lymph node biopsy can also be performed to diagnose certain types of leukemia in certain situations. Following diagnosis, blood chemistry tests can be used to determine the degree of liver and kidney damage or the effects of chemotherapy on the patient. Many people have not been diagnosed because many of the symptoms remain vague, unspecific, and may refer to other diseases. As a result, the American Cancer Society predicts that at least one-fifth of the population living with leukemia have not yet been diagnosed.

Most forms of leukemia are treated with pharmaceutical medication, typically combined into a multi-drug chemotherapy regimen. Some are also treated with radiation therapy. In some cases, a bone marrow transplant is useful. Treatment using gene therapy is currently being pursued. One such approach turns T cells into cancer-targeting attackers. As of August 2011, a year after treatment, two of the three patients are cancer-free.



Bone Marrow Aspirate Smear: A Wright's stained bone marrow aspirate smear of patient with precursor B-cell acute lymphoblastic leukemia.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Human Physiology/Blood physiology. **Provided by**: Wikibooks. **Located at**: **en.wikibooks.org/wiki/Human_Physiology/Blood_physiology%23Diseases_of_the_Blood**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Hemophilia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Hemophilia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- thrombophilia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/thrombophilia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- coagulation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/coagulation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Thrombus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Thrombus**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- XlinkRecessive. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...kRecessive.jpg**. **License**: ***Public Domain: No Known Copyright***
- Anemia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Anemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- reticulocyte. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/reticulocyte**. **License**: ***CC BY-SA: Attribution-ShareAlike***

- hematopoiesis. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/hematopoiesis. **License**: *CC BY-SA: Attribution-ShareAlike*
- complete blood count (CBC). **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/complet...0count%20(CBC). **License**: *CC BY-SA: Attribution-ShareAlike*
- Thrombus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thrombus. **License**: *CC BY-SA: Attribution-ShareAlike*
- XlinkRecessive. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...kRecessive.jpg. **License**: *Public Domain: No Known Copyright*
- Symptoms of anemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sy..._of_anemia.png. **License**: *Public Domain: No Known Copyright*
- Sickle-cell disease. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Sickle-cell_disease. **License**: *CC BY-SA: Attribution-ShareAlike*
- tachycardia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/tachycardia. **License**: *CC BY-SA: Attribution-ShareAlike*
- fitness. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/fitness. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thrombus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thrombus. **License**: *CC BY-SA: Attribution-ShareAlike*
- XlinkRecessive. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:XlinkRecessive.jpg. **License**: *Public Domain: No Known Copyright*
- Symptoms of anemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sy..._of_anemia.png. **License**: *Public Domain: No Known Copyright*
- Sicklecells. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/Fi...icklecells.jpg. **License**: *Public Domain: No Known Copyright*
- Sickle cell 01. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sickle_cell_01.jpg. **License**: *Public Domain: No Known Copyright*
- bone marrow. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/bone_marrow. **License**: *CC BY-SA: Attribution-ShareAlike*
- Leukemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Leukemia. **License**: *CC BY-SA: Attribution-ShareAlike*
- leukemia. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/leukemia. **License**: *CC BY-SA: Attribution-ShareAlike*
- complete blood count. **Provided by**: Wiktionary. **Located at**: en.wiktionary.org/wiki/complete_blood_count. **License**: *CC BY-SA: Attribution-ShareAlike*
- Thrombus. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/Thrombus. **License**: *CC BY-SA: Attribution-ShareAlike*
- XlinkRecessive. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:XlinkRecessive.jpg. **License**: *Public Domain: No Known Copyright*
- Symptoms of anemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Symptoms_of_anemia.png. **License**: *Public Domain: No Known Copyright*
- Sicklecells. **Provided by**: Wikimedia. **Located at**: commons.wikimedia.org/wiki/File:Sicklecells.jpg. **License**: *Public Domain: No Known Copyright*
- Sickle cell 01. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Sickle_cell_01.jpg. **License**: *Public Domain: No Known Copyright*
- Acute leukemia-ALL. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Acute_leukemia-ALL.jpg. **License**: *Public Domain: No Known Copyright*
- Symptoms of leukemia. **Provided by**: Wikipedia. **Located at**: en.Wikipedia.org/wiki/File:Symptoms_of_leukemia.png. **License**: *Public Domain: No Known Copyright*

This page titled 29.8D: Leukemia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.9: DISEASES AND DISORDERS OF THE HEART

Close message
Learn more
Welcome to your new guide!
Click on "New" to build out your guide.

- Topic hierarchy

29.9A: Congenital Heart Defects
29.9B: Myocarditis and Endocarditis
29.9C: Myocardial Ischemia and Infarction
29.9D: Arrhythmia

This page titled 29.9: Diseases and Disorders of the Heart is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.9A: Congenital Heart Defects

A congenital heart defect is a defect in the structure of the heart and great vessels that is present at birth.

Learning Objectives

- Describe congenital heart defects and their causes

Key Points

- Many types of heart defects exist, most of which either obstruct blood flow in the heart or vessels near it, or cause blood to flow through the heart in an abnormal pattern. Other defects, such as long QT syndrome, affect the heart's rhythm.
- Heart defects are among the most common birth defects and are the leading cause of birth defect-related deaths.
- There is a complex sequence of events that result in a well formed heart at birth and disruption of any portion may result in a defect: the orderly timing of cell growth, cell migration, and programmed cell death (apoptosis).
- The cause of congenital heart disease may be either genetic or environmental, but is usually a combination of both.
- Ventricular septal defects are collectively the most common type of congenital defect, although approximately 30% of adults have a type of atrial septal defect called probe patent foramen ovale.
- Some children with congenital heart defects have no signs while others may exhibit shortness of breath, cyanosis, syncope, heart murmur, under-developing of limbs and muscles, poor feeding or growth, or respiratory infections.


Key Terms

- **hypoplasia**: Underdevelopment or incomplete development of a tissue or organ, especially when caused by an inadequate or below-normal number of cells.
- **cyanosis**: A blue discoloration of the skin due to the circulation of blood low in oxygen.
- **long QT syndrome**: A heart disease in which there is an abnormally long delay between the electrical excitation (or depolarization) and relaxation (repolarization) of the ventricles of the heart.

A congenital heart defect is a defect in the structure of the heart and great vessels that is present at birth. Many types of heart defects exist, most of which either obstruct blood flow in the heart or vessels near it or cause blood to flow through the heart in an abnormal pattern. Other defects, such as long QT syndrome, affect the heart's rhythm. Heart defects are among the most common birth defects and are the leading cause of birth defect-related deaths. Approximately nine people in 1,000 are born with a congenital heart defect. Many defects do not require treatment, but some complex congenital heart defects require medication or surgery.

Signs and Symptoms

Signs and symptoms are related to the type and severity of the heart defect. Some children have no signs while others may exhibit shortness of breath, cyanosis, syncope, heart murmur, under-developing of limbs and muscles, poor feeding or growth, or respiratory infections. Congenital heart defects cause abnormal heart structure resulting in production of certain sounds called heart murmur. These can sometimes be detected by auscultation; however, not all heart murmurs are caused by congenital heart defects.



Phonocardiograms from normal and abnormal heart sounds

Auscultogram of heart sounds, including murmurs: Changes in heart sounds can indicate specific congenital defects in heart valves and chambers.

Causes of Heart Defects

There is a complex sequence of events that result in a well formed heart at birth and disruption of any portion may result in a defect. The orderly timing of cell growth, cell migration, and programmed cell death (apoptosis) has been studied extensively and the genes that control the process are being elucidated. In the fetus, the lungs are unexpanded and cannot accommodate the full circulatory volume. Cells in part of the septum primum die creating a hole while muscle cells, the septum secundum, grow along the right atrial side the septum primum, except for one region, leaving a gap, the foramen ovale, through which blood can pass from the right artium to the left atrium, circumventing the pulmonary circuit. A small vessel, the ductus arteriosus allows blood from the pulmonary artery to pass to the aorta. Defects in any of these structures are associated with an increased incidence of other symptoms, together referred to as VACTERL association: V – Vertebral anomalies, A – Anal atresia, C – Cardiovascular anomalies, T – Tracheoesophageal fistula, E – Esophageal atresia, R – Renal (Kidney), and/or radial anomalies, L – Limb defects.

Genetics

Most of the known causes of congenital heart disease are sporadic genetic changes, either focal mutations or deletion or addition of segments of DNA. Small chromosomal abnormalities also frequently lead to congenital heart disease. The genes regulating the complex developmental sequence have only been partly elucidated. Some genes are associated with specific defects. Mutations of a heart muscle protein, α-myosin heavy chain (MYH6) are associated with atrial and septal defects. Several proteins that interact with MYH6 are also associated with cardiac defects. Another factor, the homeobox (developmental) gene, NKX2-5 also interacts with MYH6. Mutations of all these proteins are associated with both atrial and ventricular septal defects.

Fetal Environment


Known antenatal environmental factors that may lead to congenital heart defects include maternal infections (Rubella), drugs (alcohol, hydantoin, lithium, and thalidomide), and maternal illness (diabetes mellitus, phenylketonuria, and systemic lupus erythematosus). As noted in several studies following similar body mass index (BMI) ranges, prepregnant and gestating women, who were obese (BMI ≥ 30), carried a statistically significant risk of birthing children with congenital heart defects (CHD) compared to normal-weight women (BMI = 19-24.9). Although there are minor conflicting reports, there was significant support for the risk of fetal CHD development in overweight mothers (BMI = 25-29.9).

Hypoplasia

Hypoplasia can affect the heart, typically resulting in the underdevelopment of the right ventricle or the left ventricle. This results in only one side of the heart capable of pumping blood to the body and lungs effectively. Hypoplasia of the heart is rare, but is the most serious form of CHD. It is called hypoplastic left heart syndrome when it affects the left side of the heart and hypoplastic right heart syndrome when it affects the right side of the heart. In both conditions, emergency heart surgery will permit blood to circulate to the body (or lungs, depending on which side of the heart is defective). Hypoplasia of the heart is generally a cyanotic heart defect.

Treatment

Sometimes CHD improves without treatment. Other defects are so small that they do not require any treatment. Most of the time, CHD is serious and requires surgery and/or medications. Medications include diuretics, which aid the body in eliminating water, salts, and digoxin for strengthening the contraction of the heart. This slows the heartbeat and removes some fluid from tissues. Interventional cardiology now offers patients minimally invasive alternatives to surgery. Most patients require life-long specialized cardiac care, first with a pediatric cardiologist and later with and adult congenital cardiologist. There are more than 1.8 million adults living with congenital heart defects.



Congenital heart defects: This image shows the structure and blood flow in the interior of a normal heart, as well as shows two common locations for a ventricular septal defect. The defect allows oxygen-rich blood from the left ventricle to mix with oxygen-poor blood in the right ventricle.

This page titled 29.9A: Congenital Heart Defects is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.9B: Myocarditis and Endocarditis

Endocarditis and myocarditis are driven by inflammation of the heart.

Learning Objectives

- Distinguish between myocarditis and endocarditis and their effects on the heart

Key Points

- Inflammation of the inner layer of the heart wall, the endocardium, often including the heart valves, is called endocarditis.
- Endocarditis may be infective or non-infective. Diagnosis of endocarditis includes clinical identification of the vegetation, echocardiogram, and blood cultures to demonstrate the infective microorganisms involved in infective endocarditis.
- In subacute infective endocarditis, the vegetation may include granulomatous tissue which can become fibrotic or calcify.
- In non-infective endocarditis such as nonbacterial thrombic endocarditis (NBTE), also called marantic endocarditis, vegetations are typically small and sterile (do not contain microorganisms) and typically do not engender an inflammatory response.
- Inflammation of the myocardium, the heart muscle, is called myocarditis, typically characterized by infection of the heart, an inflammatory infiltrate, and heart muscle damage with or without necrosis (tissue death).
- Myocarditis most commonly has an infectious cause, such as the parovirus B19, lyme disease, or other viral and nonviral pathogens, but may also be an autoimmune reaction.

Key Terms

- **inflammation**: A condition of any part of the body, consisting in congestion of the blood vessels, with obstruction of the blood current, and growth of morbid tissue. It is manifested outwardly by redness and swelling, attended with heat and pain.

Endocarditis is an inflammation of the inner layer of the heart, the endocardium. It usually involves the heart valves (native or prosthetic valves). Other structures that may be involved include the interventricular septum, the chordae tendineae, the mural endocardium, or even on intracardiac devices.

Endocarditis is characterized by a prototypic lesion, the vegetation, which is a mass of platelets, fibrin, microcolonies of microorganisms, and scant inflammatory cells. In the subacute form of infective endocarditis, the vegetation may also include a center of granulomatous tissue, which may fibrose or calcify.

Classification and Diagnosis

There are multiple ways to classify endocarditis. The simplest classification is based on etiology: either infective or non-infective, depending on whether a microorganism is the source of the inflammation or not. Regardless, diagnosis of endocarditis is based on the clinical features, investigations such as echocardiogram, as well as any blood cultures demonstrating the presence of endocarditis-causing microorganisms.

Since the valves of the heart do not receive any dedicated blood supply, defensive immune mechanisms (such as white blood cells) cannot directly reach the valves via the bloodstream. If an organism (such as bacteria) attaches to a valve surface and forms a vegetation, the host immune response is blunted. The lack of blood supply to the valves also has implications on treatment, since drugs also have difficulty reaching the infected valve. Normally, blood flows smoothly through these valves. If they have been damaged (from rheumatic fever, for example) the risk of bacteria attachment is increased.

Types of Endocarditis

Nonbacterial thrombic endocarditis (NBTE) or marantic endocarditis is most commonly found on previously undamaged valves.

As opposed to infective endocarditis, the vegetations in NBTE are small, sterile, and tend to aggregate along the edges of the valve or the cusps. Also unlike infective endocarditis, NBTE does not cause an inflammation response from the body. NBTE usually occurs during a hypercoagulable state such as system wide bacterial infection, or pregnancy, though it is also sometimes seen in patients with venous catheters.

Another form of sterile endocarditis is termed Libman-Sacks endocarditis; this form occurs more often in patients with lupus erythematosus and is thought to be due to the deposition of immune complexes. Like NBTE, Libman-Sacks endocarditis involves small vegetations, while infective endocarditis is composed of large vegetations. These immune complexes precipitate an inflammation reaction, which helps to differentiate it from NBTE. Also unlike NBTE, Libman-Sacks endocarditis does not seem to have a preferred location of deposition and may form on the undersurfaces of the valves or even on the endocardium.

Myocarditis

Myocarditis or inflammatory cardiomyopathy is inflammation of heart muscle (myocardium). Myocarditis is most often due to infection by common viruses, such as parvovirus B19, less commonly nonviral pathogens such as Borrelia burgdorferi (Lyme disease) or Trypanosoma cruzi, or as a hypersensitivity response to drugs.

Myocarditis is often an autoimmune reaction. Streptococcal M protein and coxsackievirus B have regions (epitopes) that are immunologically similar to cardiac myosin. After the virus is gone, the immune system may attack cardiac myosin.

Symptoms and Outcomes

The consequences of myocarditis thus also vary widely. It can cause a mild disease without any symptoms that resolves itself, or it may cause chest pain, heart failure, or sudden death. An acute myocardial infarction-like syndrome with normal coronary arteries has a good prognosis. Heart failure, even with dilated left ventricle, may have a good prognosis. Ventricular arrhythmias and high-degree heart block have a poor prognosis. Loss of right ventricular function is a strong predictor of death.

Symptoms in infants and toddlers tend to be more nonspecific, with generalized malaise, poor appetite, abdominal pain, and/or chronic cough. Later stages of the illness will present with respiratory symptoms with increased work of breathing and is often mistaken for asthma.

Since myocarditis is often due to a viral illness, many patients give a history of symptoms consistent with a recent viral infection, including fever, rash, diarrhea, joint pains, and easy fatigueability.

A large number of causes of myocarditis have been identified, but often a cause cannot be found. In Europe and North America, viruses are common culprits. Worldwide, however, the most common cause is Chagas' disease, an illness endemic to Central and South America that is due to infection by the protozoan Trypanosoma cruzi.

This page titled 29.9B: Myocarditis and Endocarditis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.9C: Myocardial Ischemia and Infarction

Ischemic heart disease is caused by reduced blood flow to the heart muscle, usually due to atherosclerosis, and can lead to a heart attack.

Learning Objectives


- Distinguish between the effects of myocardial ischemia and infarction on the heart

Key Points

- Ischemic heart disease (IHD), or myocardial ischaemia, is a disease characterized by ischaemia, or, reduced blood supply, to the heart muscle, usually due to coronary artery disease and myocardial infarction.
- Coronary artery disease is the narrowing of the heart arteries due to atherosclerosis, the deposition of fat on the walls. If these deposits come loose, they can cause a clot that will block blood flow to the heart muscle, causing a myocardial infarction.
- Symptoms of stable ischemic heart disease include angina, which is characteristic chest pain on exertion, and decreased exercise tolerance. Symptoms of myocardial infarction include acute chest pain, shortness of breath, nausea, vomiting, palpitations, sweating, and/or anxiety.
- Treatment of IHD is dependent on the underlying disease. Stable IHD may be treated with anti-angina drugs. Treatments for myocardial infarction include coronary angioplasty, treatment with “clot buster” medications, and coronary bypass, where blocked heart blood vessels are replaced.
- The phrase “heart attack” is sometimes used incorrectly to describe sudden cardiac death, which may or may not be the result of acute myocardial infarction.

Key Terms

- **angioplasty**: The mechanical widening of a narrowed or totally obstructed blood vessel generally caused by atheroma.
- **ischemia**: Local disturbance in blood circulation due to mechanical obstruction of the blood supply, such as by a narrowed artery or clot.
- **angiography**: A medical imaging technique in which an x-ray image is taken to visualize the inside of blood vessels and organs of the body, with particular interest in the arteries, veins, and the heart chambers.



Acute myocardial infarction: The diagram depicts a myocardial infarction (2) of the tip of the anterior wall of the heart (an apical infarct) after occlusion (1) of a branch of the left coronary artery (LCA). Right coronary artery (RCA).


Ischemic heart disease (IHD), or myocardial ischaemia, is a disease characterized by ischaemia, or reduced blood supply to the heart muscle, usually due to coronary artery disease and myocardial infarction. Coronary artery disease is the narrowing of the heart arteries due to atherosclerosis, the deposition of fat on the walls. If these deposits come loose, they can cause a clot that will block blood flow to the heart muscle, causing a myocardial infarction. The risk increases with age, smoking, high cholesterol levels, diabetes, and high blood pressure, and is more common in men and those who have close relatives with ischaemic heart disease and a history of myocardial infarction. It is the most common cause of death in most western countries, and a major cause of hospital admissions.

Symptoms

Symptoms of stable ischemic heart disease include angina, which is characteristic chest pain on exertion, and decreased exercise tolerance.

Ischaemic heart disease may be present with any of the following problems:

- Angina—chest pain on exertion, in cold weather or emotional situations
- Acute chest pain—acute coronary syndrome, unstable angina or myocardial infarction
- Heart failure—difficulty in breathing or swelling of the extremities due to weakness of the heart muscle
- Heartburn




Atherosclerosis: This diagram depicts the progression of atherosclerosis in a blood vessel. In coronary artery disease, these atherosclerotic plaques build up in the coronary arteries and rupture may lead to ischemic heart disease and myocardial infarction.

Diagnosis

Ischemic heart disease is typically associated with an underlying specific heart disease.

In “stable” angina, chest pain with typical features occurs at predictable levels of exertion. Various forms of cardiac stress tests may be used to induce both symptoms and detect changes with electrocardiography (EKG), echocardiography (using ultrasound of the heart), or scintigraphy (using a radioactive marker to examine blood uptake by the heart muscle). If part of the heart seems to receive an insufficient blood supply, coronary angiography may be used to identify narrowing of the coronary arteries and suitability for angioplasty or bypass surgery.



Cardiac Stress Test: A patient undergoing a stress test. This test examines blood flow to the heart during exertion, and can be used as a diagnostic tool in ischemic heart disease, coronary artery disease, and heart failure.

Diagnosis of acute coronary syndrome and myocardial infarction generally take place in the emergency department when a patient presents with acute chest pain, combined with shortness of breath, nausea, vomiting, palpitations, sweating, and/or anxiety. An EKG is performed, looking for changes that are characteristic of a myocardial infarction. In the absence of these characteristic markers, heart damage is detected by cardiac markers in the blood that identify heart muscle damage. If there is no evidence of damage or infarction, a diagnosis of “unstable angina” is given. Depending on the risk assessment, stress testing or angiography may be used to identify coronary artery disease in these patients.

In patients with heart failure, stress testing or coronary angiography may be performed to identify and treat underlying coronary artery disease.

Causes

The disease process underlying most ischemic heart disease is atherosclerosis of the coronary arteries, which decreases blood flow to the heart muscle. Atherosclerosis is the thickening of the artery wall as a result of plaque formation, which is the accumulation of fatty materials such as cholesterol. Atherosclerosis reduces blood flow to the heart muscle, causing ischemia. When these plaques weaken, they can develops a tear, which forms blood clot that obstructs blood flow, causing a myocardial infarction.

Treatment

Treatment of ischemic heart disease is dependent on the underlying disease process. In stable IHD, antianginal drugs may be used to reduce the rate of occurrence and severity of angina attacks. Treatments for acute coronary syndrome and myocardial infarction include coronary angioplasty, treatment with thrombolysis “clot buster” medication, and coronary bypass, where narrowed blood vessels on the heart are replaced.

Various treatments are offered in people deemed to be at high risk of coronary artery disease. These include control of cholesterol levels in those with known high cholesterol, smoking cessation, and control of high blood pressure.

This page titled 29.9C: Myocardial Ischemia and Infarction is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

29.9D: Arrhythmia

Learning Objectives

- Describe what are cardiac arrhythmias

Cardiac arrhythmia (irregular heartbeat) is any of a large and heterogeneous group of conditions in which there is abnormal electrical activity in the heart. The heartbeat may be too fast or too slow and may be regular or irregular. Some arrhythmias are life-threatening medical emergencies that can result in cardiac arrest. Others cause symptoms such as an abnormal awareness of heartbeat (palpitations) and may be merely uncomfortable. These palpitations have also been known to be caused by atrial/ventricular fibrillation, wire faults, and other technical or mechanical issues in cardiac pacemakers/defibrillators. Still others may not be associated with any symptoms at all, but may predispose the patient to potentially life threatening stroke or embolism. SADS, or sudden arrhythmic death syndrome, is a term (as part of sudden unexpected death syndrome) used to describe sudden death due to cardiac arrest brought on by an arrhythmia in the absence of any structural heart disease on autopsy.

The Conduction System

Automaticity refers to a cardiac muscle cell firing off an impulse on its own. All of the cells in the heart have the ability to initiate an action potential; however, only some of these cells are designed to routinely trigger heart beats. These cells are found in the conduction system of the heart and include the SA node, AV node, Bundle of His, and Purkinje fibers. The sinoatrial node is a single specialized location in the atrium which has a higher automaticity (a faster pacemaker) than the rest of the heart and, therefore, is usually responsible for setting the heart rate and initiating each heart beat.



Figure 1: Cardiac conduction system: Electrical impulse arising in SA node, traversing atria to AV node, then entering ventricle. Rhythm originating at or above AV node constitutes supraventricular tachycardia (SVT).

In normal electrical activity, each heart beat originates as an electrical impulse from a small area of tissue in the right atrium of the heart called the sinus node or sino-atrial node or SA node. The impulse initially causes both atria to contract, then activates the atrioventricular (or AV) node which is normally the only electrical connection between the atria and the ventricles (main pumping chambers). The impulse then spreads through both ventricles causing a synchronized contraction of the heart muscle and, thus, the pulse. In adults the normal resting heart rate ranges from 60 to 80 beats per minute. The resting heart rate in children is much faster. A slow rhythm (less than 60 beats/min), is labeled bradycardia. This may be caused by a slowed signal from the sinus node (sinus bradycardia), a pause in the normal activity of the sinus node (sinus arrest), or by blocking of the electrical impulse on its way from the atria to the ventricles (AV block or heart block). Heart block comes in varying degrees and severity. Bradycardias may be present in the normally functioning heart of endurance athletes or other well-conditioned persons.

Arrhythmias

Arrhythmia may be classified by rate (normal, tachycardia, bradycardia), or mechanism (automaticity, reentry, fibrillation). It is also appropriate to classify by site of origin. *Arrhythmias with atrial origin include:* Premature Atrial Contractions (PACs), Wandering Atrial Pacemaker, Atrial flutter, and Atrial fibrillation (Afib). *Arrhythmias arising at the junction:* Supraventricular tachycardia (SVT), Junctional rhythm, and Junctional tachycardia. *Arrythmias arising from pathologies at the AV node (AV blocks):* First degree heart block, which manifests as PR prolongation, Second degree heart block, and Third degree heart block, also known as complete heart block.




Figure 2: Atrial flutter: Atrial flutter, shown here, is a cardiac arrhythmia that occurs in the atria of the heart and falls into the category of supra-ventricular tachycardias. While this rhythm occurs most often in individuals with cardiovascular disease (e.g., hypertension, coronary artery disease, and cardiomyopathy) and diabetes, it may occur spontaneously in people with otherwise normal hearts. It is typically not a stable rhythm, and frequently degenerates into atrial fibrillation (AF). However, it does rarely persist for months to years.

Fibrillation

When an entire chamber of the heart is involved in a multiple micro-reentry circuits and, therefore, quivering with chaotic electrical impulses, it is said to be in fibrillation. Fibrillation can affect the atrium (atrial fibrillation) or the ventricle (ventricular fibrillation); ventricular fibrillation is imminently life-threatening. Atrial fibrillation affects the upper chambers of the heart, known as the atria. Atrial fibrillation may be due to serious underlying medical conditions and should be evaluated by a physician. It is not typically a medical emergency. Ventricular fibrillation occurs in the ventricles (lower chambers) of the heart; it is always a medical emergency. If left untreated, ventricular fibrillation (VF, or V-fib) can lead to death within minutes. When a heart goes into V-fib, effective pumping of the blood stops. V-fib is considered a form of cardiac arrest.




Figure 3: Normal sinus rhythm: Schematic diagram of normal sinus rhythm for a human heart as seen on ECG. In atrial fibrillation, one form of cardiac arrhythmia, the P waves, which represent depolarization of the atria, are absent.

Treatment

Dysrhythmias may also be treated electrically, by applying a shock across the heart—either externally to the chest wall or internally to the heart via implanted electrodes. Cardioversion is either achieved pharmacologically or via the application of a shock synchronized to the underlying heartbeat. It is used for treatment of supraventricular tachycardias. Defibrillation differs in that the shock is not synchronized. It is needed for the chaotic rhythm of ventricular fibrillation and is also used for pulseless ventricular tachycardia. Often, more electricity is required for defibrillation than for cardioversion. In most defibrillation, the recipient has lost consciousness so there is no need for sedation.

Key Points

- Some arrhythmias are life -threatening medical emergencies that can result in cardiac arrest. Others cause symptoms such as an abnormal awareness of heart beat (palpitations), and may be merely uncomfortable.
- The term cardiac arrhythmia covers a very large number of very different conditions. The most common symptom of arrhythmia is an abnormal awareness of heartbeat, called palpitations.
- Automaticity refers to a cardiac muscle cell firing off an impulse on its own. All of the cells in the heart have the ability to initiate an action potential; however, only some are designed to routinely trigger heart beats. These cells are found in the conduction system of the heart.
- Ventricular fibrillation is imminently life-threatening. Atrial fibrillation is not typically a medical emergency.
- Arrhythmia may be classified by rate (normal, tachycardia, bradycardia) or mechanism (automaticity, reentry, fibrillation). It is also appropriate to classify by site of origin.
- Causes of SADS in young people include viral myocarditis, long QT syndrome, Brugada syndrome, Catecholaminergic polymorphic ventricular tachycardia, hypertrophic cardiomyopathy, and arrhythmogenic right ventricular dysplasia.

Key Terms

- **automaticity**: The ability of the heart muscles to contract independently of each other when separated or in unison when reattached in whole or in groups.
- **supraventricular tachycardia**: Any rapid heart rhythm originating at or above the atrioventricular node.
- **Cardiac arrhythmia (irregular heartbeat)**: A large and heterogeneous group of conditions in which there is abnormal electrical activity in the heart.

LICENSES AND ATTRIBUTIONS

CC LICENSED CONTENT, SHARED PREVIOUSLY

- Curation and Revision. **Authored by**: Boundless.com. **Provided by**: Boundless.com. **License**: ***CC BY-SA: Attribution-ShareAlike***

CC LICENSED CONTENT, SPECIFIC ATTRIBUTION

- Congenital heart defect. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Congenital_heart_defect**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- hypoplasia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/hypoplasia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- cyanosis. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/cyanosis**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- long QT syndrome. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/long_QT_syndrome**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- VSD image. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:VSD_image.jpg**. **License**: ***Public Domain: No Known Copyright***
- Phonocardiograms from normal and abnormal heart sounds. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ph...art_sounds.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Valvular heart disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Valvular_heart_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- inflammation. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/inflammation**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- VSD image. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:VSD_image.jpg**. **License**: ***Public Domain: No Known Copyright***
- Phonocardiograms from normal and abnormal heart sounds. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ph...art_sounds.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Myocardial infarction. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Myocardial_infarction**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Ischaemic heart disease. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Ischaemic_heart_disease**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- angioplasty. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/angioplasty**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- ischemia. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/ischemia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- angiography. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/angiography**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- VSD image. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:VSD_image.jpg**. **License**: ***Public Domain: No Known Copyright***
- Phonocardiograms from normal and abnormal heart sounds. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ph...art_sounds.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Stress test. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Stress_test.jpg**. **License**: ***Public Domain: No Known Copyright***
- Endo dysfunction Athero. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion_Athero.PNG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- AMI scheme. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:AMI_scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Arrhythmia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Arrhythmia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- automaticity. **Provided by**: Wiktionary. **Located at**: **en.wiktionary.org/wiki/automaticity**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- supraventricular tachycardia. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/suprave...%20tachycardia**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Cardiac arrhythmia (irregular heartbeat). **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/Cardiac...r%20heartbeat)**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- VSD image. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:VSD_image.jpg**. **License**: ***Public Domain: No***

Known Copyright

- Phonocardiograms from normal and abnormal heart sounds. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Ph...art_sounds.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Stress test. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Stress_test.jpg**. **License**: ***Public Domain: No Known Copyright***
- Endo dysfunction Athero. **Provided by**: Wikimedia. **Located at**: **commons.wikimedia.org/wiki/Fi...ion_Athero.PNG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- AMI scheme. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:AMI_scheme.png**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Atrial flutter34. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:At..._flutter34.JPG**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- Heart conduct sinus. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:He...duct_sinus.gif**. **License**: ***CC BY-SA: Attribution-ShareAlike***
- SinusRhythmLabels. **Provided by**: Wikipedia. **Located at**: **en.Wikipedia.org/wiki/File:Si...ythmLabels.svg**. **License**: ***Public Domain: No Known Copyright***

This page titled 29.9D: Arrhythmia is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

INDEX

A

acidosis
29.16A: Acidosis
afferent nerve transmission
12.4B: Classification of Nerves
arrhythmia
29.9D: Arrhythmia

B

Bell's palsy
12.5H: Facial (VII) Nerve
bone remodeling
6.4C: Bone Remodeling
bone repair
6.4D: Bone Repair
bronchi
21.3A: Bronchi and Subdivisions
bronchoconstriction
21.3A: Bronchi and Subdivisions

C

cervical cancer
29.17C: Cervical Cancer
coronary artery
17.2A: Heart Circulation
coronary circulation
17.2A: Heart Circulation
cranial nerves
12.5A: Brief Overview of Cranial Nerves
cytokines
20.4: Cytokines

D

dehydration synthesis
2.4C: Dehydration Synthesis
dendrites
4.4A: Characteristics of Nervous Tissue
diastole
17.2A: Heart Circulation

E

efferent nerve transmission
12.4B: Classification of Nerves
epidermis
5.1B: Structure of the Skin: Epidermis
erythrocyte
16.2A: RBC Anatomy

F

femoral artery
18.10F: Arteries of the Pelvis and Lower Limbs
Fetal Alcohol Syndrome
27.9B: Teratogens
fibular artery
18.10F: Arteries of the Pelvis and Lower Limbs
flat bones
6.3A: Gross Anatomy

G

Galactosemia
29.14B: Galactosemia and Glycogen Storage Disease
gastric phase
22.14B: Gastric Phase
glycogen
29.14B: Galactosemia and Glycogen Storage Disease

H

Haldane effect
21.9D: Carbon Dioxide Transport
Hering–Breuer inflation
21.10D: Proprioceptor Regulation of Breathing
homeostasis
23.9A: Homeostatic Process
hyoid bone
7.2A: The Hyoid Bone
Hyperkyphosis
29.2A: Osteoporosis

I

inflammation
20.1D: Inflammation
integumentary system
5: Integumentary System

J

Jendrassik maneuver
12.10C: Spinal Reflexes

L

long bones
6.3A: Gross Anatomy
lymphatic system
19: Lymphatic System

M

membrane potential
10.5D: Resting Membrane Potentials
metabolic alkalosis
6.3D: Chemical Composition of Bone
mitral valve
17.2B: Operation of Atrioventricular Valves
mouth
22.4A: Mouth
myelin sheath
4.4A: Characteristics of Nervous Tissue
myocardial infarction
17.2A: Heart Circulation

N

nerves
12.4B: Classification of Nerves
neuroglia
4.4A: Characteristics of Nervous Tissue
neurons
4.4A: Characteristics of Nervous Tissue

O

Orbits
7.1D: Orbits
osteoporosis
29.2A: Osteoporosis
oxygen transport
21.9C: Oxygen Transport

P

pancreas
22.9: The Pancreas
peritoneum
22.3: The Peritoneum
Peroxisomes
3.6E: Peroxisomes
phrenic nerves
21.10B: Neural Mechanisms (Cortex)
plantar artery
18.10F: Arteries of the Pelvis and Lower Limbs
popliteal artery
18.10F: Arteries of the Pelvis and Lower Limbs
primary motor cortex
21.10B: Neural Mechanisms (Cortex)

R

Resting Membrane Potentials
10.5D: Resting Membrane Potentials
ribs
7.4B: Thoracic Cage: Ribs

S

Schwann cell
4.4A: Characteristics of Nervous Tissue
secondary ossification
6.4B: Postnatal Bone Growth
skull
7.1: The Skull
sternum
7.4A: Thoracic Cage: Sternum
Subvalvular Apparatus
17.2B: Operation of Atrioventricular Valves
synovial joints
8.4: Synovial Joints
systole
17.2A: Heart Circulation

T

Teratogens
27.9B: Teratogens
thorax
7.4: The Thorax
tibial artery
18.10F: Arteries of the Pelvis and Lower Limbs
Transcytosis
18.8B: Transcytosis

tricuspid valve
17.2B: Operation of Atrioventricular Valves

V

vestibular system
13.3B: The Vestibular System
vulva
26.5F: Vulva

This page titled Index is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

INDEX

A

acidosis
29.16A: Acidosis
afferent nerve transmission
12.4B: Classification of Nerves
arrhythmia
29.9D: Arrhythmia

B

Bell's palsy
12.5H: Facial (VII) Nerve
bone remodeling
6.4C: Bone Remodeling
bone repair
6.4D: Bone Repair
bronchi
21.3A: Bronchi and Subdivisions
bronchoconstriction
21.3A: Bronchi and Subdivisions

C

cervical cancer
29.17C: Cervical Cancer
coronary artery
17.2A: Heart Circulation
coronary circulation
17.2A: Heart Circulation
cranial nerves
12.5A: Brief Overview of Cranial Nerves
cytokines
20.4: Cytokines

D

dehydration synthesis
2.4C: Dehydration Synthesis
dendrites
4.4A: Characteristics of Nervous Tissue
diastole
17.2A: Heart Circulation

E

efferent nerve transmission
12.4B: Classification of Nerves
epidermis
5.1B: Structure of the Skin: Epidermis
erythrocyte
16.2A: RBC Anatomy

F

femoral artery
18.10F: Arteries of the Pelvis and Lower Limbs
Fetal Alcohol Syndrome
27.9B: Teratogens
fibular artery
18.10F: Arteries of the Pelvis and Lower Limbs
flat bones
6.3A: Gross Anatomy

G

Galactosemia
29.14B: Galactosemia and Glycogen Storage Disease
gastric phase
22.14B: Gastric Phase
glycogen
29.14B: Galactosemia and Glycogen Storage Disease

H

Haldane effect
21.9D: Carbon Dioxide Transport
Hering–Breuer inflation
21.10D: Proprioceptor Regulation of Breathing
homeostasis
23.9A: Homeostatic Process
hyoid bone
7.2A: The Hyoid Bone
Hyperkyphosis
29.2A: Osteoporosis

I

inflammation
20.1D: Inflammation
integumentary system
5: Integumentary System

J

Jendrassik maneuver
12.10C: Spinal Reflexes

L

long bones
6.3A: Gross Anatomy
lymphatic system
19: Lymphatic System

M

membrane potential
10.5D: Resting Membrane Potentials
metabolic alkalosis
6.3D: Chemical Composition of Bone
mitral valve
17.2B: Operation of Atrioventricular Valves
mouth
22.4A: Mouth
myelin sheath
4.4A: Characteristics of Nervous Tissue
myocardial infarction
17.2A: Heart Circulation

N

nerves
12.4B: Classification of Nerves
neuroglia
4.4A: Characteristics of Nervous Tissue
neurons
4.4A: Characteristics of Nervous Tissue

O

Orbits
7.1D: Orbits
osteoporosis
29.2A: Osteoporosis
oxygen transport
21.9C: Oxygen Transport

P

pancreas
22.9: The Pancreas
peritoneum
22.3: The Peritoneum
Peroxisomes
3.6E: Peroxisomes
phrenic nerves
21.10B: Neural Mechanisms (Cortex)
plantar artery
18.10F: Arteries of the Pelvis and Lower Limbs
popliteal artery
18.10F: Arteries of the Pelvis and Lower Limbs
primary motor cortex
21.10B: Neural Mechanisms (Cortex)

R

Resting Membrane Potentials
10.5D: Resting Membrane Potentials
ribs
7.4B: Thoracic Cage: Ribs

S

Schwann cell
4.4A: Characteristics of Nervous Tissue
secondary ossification
6.4B: Postnatal Bone Growth
skull
7.1: The Skull
sternum
7.4A: Thoracic Cage: Sternum
Subvalvular Apparatus
17.2B: Operation of Atrioventricular Valves
synovial joints
8.4: Synovial Joints
systole
17.2A: Heart Circulation

T

Teratogens
27.9B: Teratogens
thorax
7.4: The Thorax
tibial artery
18.10F: Arteries of the Pelvis and Lower Limbs
Transcytosis
18.8B: Transcytosis

tricuspid valve
17.2B: Operation of Atrioventricular Valves

V

vestibular system
13.3B: The Vestibular System
vulva
26.5F: Vulva

This page titled Index is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

INDEX

A

acidosis
29.16A: Acidosis
afferent nerve transmission
12.4B: Classification of Nerves
arrhythmia
29.9D: Arrhythmia

B

Bell's palsy
12.5H: Facial (VII) Nerve
bone remodeling
6.4C: Bone Remodeling
bone repair
6.4D: Bone Repair
bronchi
21.3A: Bronchi and Subdivisions
bronchoconstriction
21.3A: Bronchi and Subdivisions

C

cervical cancer
29.17C: Cervical Cancer
coronary artery
17.2A: Heart Circulation
coronary circulation
17.2A: Heart Circulation
cranial nerves
12.5A: Brief Overview of Cranial Nerves
cytokines
20.4: Cytokines

D

dehydration synthesis
2.4C: Dehydration Synthesis
dendrites
4.4A: Characteristics of Nervous Tissue
diastole
17.2A: Heart Circulation

E

efferent nerve transmission
12.4B: Classification of Nerves
epidermis
5.1B: Structure of the Skin: Epidermis
erythrocyte
16.2A: RBC Anatomy

F

femoral artery
18.10F: Arteries of the Pelvis and Lower Limbs
Fetal Alcohol Syndrome
27.9B: Teratogens
fibular artery
18.10F: Arteries of the Pelvis and Lower Limbs
flat bones
6.3A: Gross Anatomy

G

Galactosemia
29.14B: Galactosemia and Glycogen Storage Disease
gastric phase
22.14B: Gastric Phase
glycogen
29.14B: Galactosemia and Glycogen Storage Disease

H

Haldane effect
21.9D: Carbon Dioxide Transport
Hering–Breuer inflation
21.10D: Proprioceptor Regulation of Breathing
homeostasis
23.9A: Homeostatic Process
hyoid bone
7.2A: The Hyoid Bone
Hyperkyphosis
29.2A: Osteoporosis

I

inflammation
20.1D: Inflammation
integumentary system
5: Integumentary System

J

Jendrassik maneuver
12.10C: Spinal Reflexes

L

long bones
6.3A: Gross Anatomy
lymphatic system
19: Lymphatic System

M

membrane potential
10.5D: Resting Membrane Potentials
metabolic alkalosis
6.3D: Chemical Composition of Bone
mitral valve
17.2B: Operation of Atrioventricular Valves
mouth
22.4A: Mouth
myelin sheath
4.4A: Characteristics of Nervous Tissue
myocardial infarction
17.2A: Heart Circulation

N

nerves
12.4B: Classification of Nerves
neuroglia
4.4A: Characteristics of Nervous Tissue
neurons
4.4A: Characteristics of Nervous Tissue

O

Orbits
7.1D: Orbits
osteoporosis
29.2A: Osteoporosis
oxygen transport
21.9C: Oxygen Transport

P

pancreas
22.9: The Pancreas
peritoneum
22.3: The Peritoneum
Peroxisomes
3.6E: Peroxisomes
phrenic nerves
21.10B: Neural Mechanisms (Cortex)
plantar artery
18.10F: Arteries of the Pelvis and Lower Limbs
popliteal artery
18.10F: Arteries of the Pelvis and Lower Limbs
primary motor cortex
21.10B: Neural Mechanisms (Cortex)

R

Resting Membrane Potentials
10.5D: Resting Membrane Potentials
ribs
7.4B: Thoracic Cage: Ribs

S

Schwann cell
4.4A: Characteristics of Nervous Tissue
secondary ossification
6.4B: Postnatal Bone Growth
skull
7.1: The Skull
sternum
7.4A: Thoracic Cage: Sternum
Subvalvular Apparatus
17.2B: Operation of Atrioventricular Valves
synovial joints
8.4: Synovial Joints
systole
17.2A: Heart Circulation

T

Teratogens
27.9B: Teratogens
thorax
7.4: The Thorax
tibial artery
18.10F: Arteries of the Pelvis and Lower Limbs
Transcytosis
18.8B: Transcytosis

tricuspid valve
17.2B: Operation of Atrioventricular Valves

V

vestibular system
13.3B: The Vestibular System
vulva
26.5F: Vulva

This page titled Index is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

INDEX

A

acidosis
29.16A: Acidosis
afferent nerve transmission
12.4B: Classification of Nerves
arrhythmia
29.9D: Arrhythmia

B

Bell's palsy
12.5H: Facial (VII) Nerve
bone remodeling
6.4C: Bone Remodeling
bone repair
6.4D: Bone Repair
bronchi
21.3A: Bronchi and Subdivisions
bronchoconstriction
21.3A: Bronchi and Subdivisions

C

cervical cancer
29.17C: Cervical Cancer
coronary artery
17.2A: Heart Circulation
coronary circulation
17.2A: Heart Circulation
cranial nerves
12.5A: Brief Overview of Cranial Nerves
cytokines
20.4: Cytokines

D

dehydration synthesis
2.4C: Dehydration Synthesis
dendrites
4.4A: Characteristics of Nervous Tissue
diastole
17.2A: Heart Circulation

E

efferent nerve transmission
12.4B: Classification of Nerves
epidermis
5.1B: Structure of the Skin: Epidermis
erythrocyte
16.2A: RBC Anatomy

F

femoral artery
18.10F: Arteries of the Pelvis and Lower Limbs
Fetal Alcohol Syndrome
27.9B: Teratogens
fibular artery
18.10F: Arteries of the Pelvis and Lower Limbs
flat bones
6.3A: Gross Anatomy

G

Galactosemia
29.14B: Galactosemia and Glycogen Storage Disease
gastric phase
22.14B: Gastric Phase
glycogen
29.14B: Galactosemia and Glycogen Storage Disease

H

Haldane effect
21.9D: Carbon Dioxide Transport
Hering–Breuer inflation
21.10D: Proprioceptor Regulation of Breathing
homeostasis
23.9A: Homeostatic Process
hyoid bone
7.2A: The Hyoid Bone
Hyperkyphosis
29.2A: Osteoporosis

I

inflammation
20.1D: Inflammation
integumentary system
5: Integumentary System

J

Jendrassik maneuver
12.10C: Spinal Reflexes

L

long bones
6.3A: Gross Anatomy
lymphatic system
19: Lymphatic System

M

membrane potential
10.5D: Resting Membrane Potentials
metabolic alkalosis
6.3D: Chemical Composition of Bone
mitral valve
17.2B: Operation of Atrioventricular Valves
mouth
22.4A: Mouth
myelin sheath
4.4A: Characteristics of Nervous Tissue
myocardial infarction
17.2A: Heart Circulation

N

nerves
12.4B: Classification of Nerves
neuroglia
4.4A: Characteristics of Nervous Tissue
neurons
4.4A: Characteristics of Nervous Tissue

O

Orbits
7.1D: Orbits
osteoporosis
29.2A: Osteoporosis
oxygen transport
21.9C: Oxygen Transport

P

pancreas
22.9: The Pancreas
peritoneum
22.3: The Peritoneum
Peroxisomes
3.6E: Peroxisomes
phrenic nerves
21.10B: Neural Mechanisms (Cortex)
plantar artery
18.10F: Arteries of the Pelvis and Lower Limbs
popliteal artery
18.10F: Arteries of the Pelvis and Lower Limbs
primary motor cortex
21.10B: Neural Mechanisms (Cortex)

R

Resting Membrane Potentials
10.5D: Resting Membrane Potentials
ribs
7.4B: Thoracic Cage: Ribs

S

Schwann cell
4.4A: Characteristics of Nervous Tissue
secondary ossification
6.4B: Postnatal Bone Growth
skull
7.1: The Skull
sternum
7.4A: Thoracic Cage: Sternum
Subvalvular Apparatus
17.2B: Operation of Atrioventricular Valves
synovial joints
8.4: Synovial Joints
systole
17.2A: Heart Circulation

T

Teratogens
27.9B: Teratogens
thorax
7.4: The Thorax
tibial artery
18.10F: Arteries of the Pelvis and Lower Limbs
Transcytosis
18.8B: Transcytosis

tricuspid valve
17.2B: Operation of Atrioventricular Valves

V

vestibular system
13.3B: The Vestibular System
vulva
26.5F: Vulva

This page titled Index is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

INDEX

A

acidosis
29.16A: Acidosis
afferent nerve transmission
12.4B: Classification of Nerves
arrhythmia
29.9D: Arrhythmia

B

Bell's palsy
12.5H: Facial (VII) Nerve
bone remodeling
6.4C: Bone Remodeling
bone repair
6.4D: Bone Repair
bronchi
21.3A: Bronchi and Subdivisions
bronchoconstriction
21.3A: Bronchi and Subdivisions

C

cervical cancer
29.17C: Cervical Cancer
coronary artery
17.2A: Heart Circulation
coronary circulation
17.2A: Heart Circulation
cranial nerves
12.5A: Brief Overview of Cranial Nerves
cytokines
20.4: Cytokines

D

dehydration synthesis
2.4C: Dehydration Synthesis
dendrites
4.4A: Characteristics of Nervous Tissue
diastole
17.2A: Heart Circulation

E

efferent nerve transmission
12.4B: Classification of Nerves
epidermis
5.1B: Structure of the Skin: Epidermis
erythrocyte
16.2A: RBC Anatomy

F

femoral artery
18.10F: Arteries of the Pelvis and Lower Limbs
Fetal Alcohol Syndrome
27.9B: Teratogens
fibular artery
18.10F: Arteries of the Pelvis and Lower Limbs
flat bones
6.3A: Gross Anatomy

G

Galactosemia
29.14B: Galactosemia and Glycogen Storage Disease
gastric phase
22.14B: Gastric Phase
glycogen
29.14B: Galactosemia and Glycogen Storage Disease

H

Haldane effect
21.9D: Carbon Dioxide Transport
Hering–Breuer inflation
21.10D: Proprioceptor Regulation of Breathing
homeostasis
23.9A: Homeostatic Process
hyoid bone
7.2A: The Hyoid Bone
Hyperkyphosis
29.2A: Osteoporosis

I

inflammation
20.1D: Inflammation
integumentary system
5: Integumentary System

J

Jendrassik maneuver
12.10C: Spinal Reflexes

L

long bones
6.3A: Gross Anatomy
lymphatic system
19: Lymphatic System

M

membrane potential
10.5D: Resting Membrane Potentials
metabolic alkalosis
6.3D: Chemical Composition of Bone
mitral valve
17.2B: Operation of Atrioventricular Valves
mouth
22.4A: Mouth
myelin sheath
4.4A: Characteristics of Nervous Tissue
myocardial infarction
17.2A: Heart Circulation

N

nerves
12.4B: Classification of Nerves
neuroglia
4.4A: Characteristics of Nervous Tissue
neurons
4.4A: Characteristics of Nervous Tissue

O

Orbits
7.1D: Orbits
osteoporosis
29.2A: Osteoporosis
oxygen transport
21.9C: Oxygen Transport

P

pancreas
22.9: The Pancreas
peritoneum
22.3: The Peritoneum
Peroxisomes
3.6E: Peroxisomes
phrenic nerves
21.10B: Neural Mechanisms (Cortex)
plantar artery
18.10F: Arteries of the Pelvis and Lower Limbs
popliteal artery
18.10F: Arteries of the Pelvis and Lower Limbs
primary motor cortex
21.10B: Neural Mechanisms (Cortex)

R

Resting Membrane Potentials
10.5D: Resting Membrane Potentials
ribs
7.4B: Thoracic Cage: Ribs

S

Schwann cell
4.4A: Characteristics of Nervous Tissue
secondary ossification
6.4B: Postnatal Bone Growth
skull
7.1: The Skull
sternum
7.4A: Thoracic Cage: Sternum
Subvalvular Apparatus
17.2B: Operation of Atrioventricular Valves
synovial joints
8.4: Synovial Joints
systole
17.2A: Heart Circulation

T

Teratogens
27.9B: Teratogens
thorax
7.4: The Thorax
tibial artery
18.10F: Arteries of the Pelvis and Lower Limbs
Transcytosis
18.8B: Transcytosis

tricuspid valve
17.2B: Operation of Atrioventricular Valves

V

vestibular system
13.3B: The Vestibular System
vulva
26.5F: Vulva

This page titled Index is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

GLOSSARY

S

Sample Word 1 | Sample Definition 1

This page titled Glossary is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning).

DETAILED LICENSING

Detailed licensing information for this resource is not available at this time.

This page titled Anatomy and Physiology (Boundless) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Boundless (now LumenLearning) via source content that was edited to the style and standards of the LibreTexts platform.