PHYSICAL ANTHROPOLOGY ANTHRO-1



Emma Blackthorne
Yuba College

LibreTexts™

Physical Anthropology Anthro-1

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 04/16/2026

TABLE OF CONTENTS

Licensing

1: Introduction to Biological/Physical Anthropology

- 1.1: Introduction to Physical Anthropology
  - 1.1.1: Science
  - 1.1.2: Anthropology as a Science
- 1.2: Introduction to Biology
  - 1.2.1: Introduction
  - 1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology
  - 1.2.3: Evolutionary Theory (Part 1)
  - 1.2.4: Evolutionary Theory (Part 2)
  - 1.2.5: Evolutionary Theory (Part 3)
  - 1.2.6: "Forces" of Evolution
  - 1.2.7: Genetics, Cellular Biology, and Variation (Part 1)
  - 1.2.8: Genetics, Cellular Biology, and Variation (Part 2)
  - 1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria
- 1.3: What is Biological Anthropology?

2: Mechanisms and Evolutionary Thought

- 2.1: Evolutionary Theory
  - 2.1.1: What is Biological Anthropology?
  - 2.1.2: Scientific Method
  - 2.1.3: History of Evolutionary Thought
  - 2.1.4: Darwinian Evolution
  - 2.1.5: Cell Biology and Basic Genetics
  - 2.1.6: Mendelian Genetics
  - 2.1.7: Modern Synthesis
  - 2.1.8: Modern Human Variation
- 2.2: Taxonomy
- 2.3: Mammals
- 2.4: Human Taxonomy
- 2.5: Osteology
- 2.6: Macroevolution
- 2.7: Species vs. Paleospecies
- 2.8: Interspecific vs. Intraspecific Variation
- 2.9: Adaptive Radiation
- 2.10: Analogy vs. Homology

3: Primates

- 3.1: Introduction to Primatology
- 3.2: Primate Classification
- 3.3: Primate Evolution
- 3.4: Primate Social Organization
- 3.5: Non-Human Primates

- 3.5.1: Classification - Ordering the Natural World
- 3.5.2: Primate Skeletal Anatomy
- 3.5.3: Modern Primates
- 3.5.4: Taxonomy of the Living Primates
- 3.5.5: Primate Ecology
- 3.5.6: Overviews of Living Primates (Strepsirhines)
- 3.5.7: Overviews of Living Primates (Haplorhines)
- 3.5.8: More Haplorhines (Old World Monkeys)
- 3.5.9: Even more Haplorhines (The Apes)
- 3.5.10: Primate Evolution
- 3.6: Ethology
- 3.7: Conservation
- 3.8: Lab 5 and 6- Primates

4: Paleontology and Primate Evolution

- 4.1: Introduction to Paleoanthropology
- 4.2: Introduction to Paleoanthropology
  - 4.2.1: Paleoanthropology
  - 4.2.2: What is a Hominim
  - 4.2.3: Miocene Epoch
    - 4.2.3.1: Introduction to the Miocene Epoch
    - 4.2.3.2: Sahelanthropus tchadensis
    - 4.2.3.3: Orrorin tugenensis
    - 4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba
    - 4.2.3.5: Early Hominid Cranium Comparison Checklist
- 4.3: Pliocene Epoch
  - 4.3.1: Introduction to the Pliocene Epoch
  - 4.3.2: Australopithecus anamensis
  - 4.3.3: Gracile Australopiths
  - 4.3.4: Australopithecus afarensis
  - 4.3.5: Australopithecus bahrelghazali
  - 4.3.6: Kenyanthropus platyops
  - 4.3.7: Australopithecus prometheus or africanus
  - 4.3.8: Australopithecus africanus
  - 4.3.9: Middle Hominid Cranium Comparison Checklist

5: Human Fossil Record

- 5.1: Pleistocene Epoch
  - 5.1.1: Introduction to the Pleistocene Epoch
  - 5.1.2: Paranthropines
  - 5.1.3: Australopithecus
    - 5.1.3.1: Australopithecus/Paranthropus aethiopicus
  - 5.1.4: Paranthropus boisei
  - 5.1.5: Paranthropus robustus
  - 5.1.6: Australopithecus garhi
  - 5.1.7: Australopithecus sediba
  - 5.1.8: Genus Homo
  - 5.1.9: Homo habilis

- 5.1.10: Homo rudolfensis
- 5.1.11: Homo species indeterminate
- 5.1.12: Homo naledi
- 5.1.13: The “erectus Grade”
- 5.1.14: Homo ergaster
- 5.1.15: Homo erectus
- 5.1.16: Homo georgicus
- 5.1.17: Homo antecessor
- 5.1.18: Homo floresiensis
- 5.1.19: Homo heidelbergensis
- 5.1.20: The Denisovans
- 5.1.21: Homo neanderthalensis
- 5.1.22: Homo sapiens
- 5.1.23: Recent Hominid Cranium Comparison Checklist

6: Race and Human Variation

- 6.1: The History of "Race" Concepts
- 6.2: Human Variation in Biological Anthropology Today
- 6.3: Talking About Human Biological Variation Going Forward
- 6.4: Key Terms
- 6.5: Human Variation- An Adaptive Significance Approach
  - 6.5.1: Stress and Homeostasis
  - 6.5.2: Adjustments and Adaptations
  - 6.5.3: Review Questions
  - 6.5.4: Key Terms

7: New Frontiers

- 7.1: Homo sapiens Futures - Doom, Gloom, and Hope?
  - 7.1.1: Introduction
  - 7.1.2: Doom and Gloom
  - 7.1.3: Hope
- 7.2: Bioarchaeology and Forensic Anthropology

Index

Glossary

Detailed Licensing

Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

CHAPTER OVERVIEW

1: Introduction to Biological/Physical Anthropology

1.1: Introduction to Physical Anthropology

1.1.1: Science

1.1.2: Anthropology as a Science

1.2: Introduction to Biology

1.2.1: Introduction

1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology

1.2.3: Evolutionary Theory (Part 1)

1.2.4: Evolutionary Theory (Part 2)

1.2.5: Evolutionary Theory (Part 3)

1.2.6: "Forces" of Evolution

1.2.7: Genetics, Cellular Biology, and Variation (Part 1)

1.2.8: Genetics, Cellular Biology, and Variation (Part 2)

1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria

1.3: What is Biological Anthropology?

1: Introduction to Biological/Physical Anthropology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

1.1: Introduction to Physical Anthropology

This chapter will introduce you to the field of anthropology, define basic terms and concepts and explain why it is important, and how it can change your perspective of the world around you.

1.1.1: Science

1.1.2: Anthropology as a Science

This page titled 1.1: Introduction to Physical Anthropology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

1.1.1: Science

Science is a specific way of looking at the universe.

science : empiricism :: religion : faith

Anthropology is mostly based on science. Anthropology is holistic. The four main subfields of anthropology are cultural anthropology, physical (biological) anthropology, archaeology, and linguistics. It is worthwhile to back up and introduce both science and anthropology. And before talking about science, we should back up even farther, and talk about *epistemology*, the study of how we know things.

The word science comes from the Latin for “knowledge”, but in modern English it means a very specific kind of knowledge, and implies a method of obtaining knowledge.

Scientific Method


Note

Another good summary of the scientific method

Here are a few terms to clarify: law vs. theory, quantitative vs. qualitative, inductive vs. deductive

A scientific "Law" is just an archaic term for an accepted theory; we could talk about Newton and his theory of gravity, or Darwin and the Law of Evolution, and what we mean is that neither hypothesis has been disproven yet.





Anthropology uses both quantitative (statistics) and qualitative (detailed description) methods, but leans towards qualitative research.

Other good terms to understand are induction and deduction. Induction is where you take what you can observe and make generalizations, like hypotheses and theories. Deduction is where you start with the general laws of the universe and you use them to predict how a specific event will play out. Both are important aspects of science: the ability to make generalizations, and the ability to predict future events. For example, Sherlock Holmes kept a notebook of all his previous cases, and from this he made inductive generalizations about human nature. When a client came to see him, he would deductively apply his criminal theories to solving the specific case.

Quantitative versus qualitative is another pair to distinguish, and have to do with what kind of data you use. Quantitative science is based on a large quantity of objects, qualitative science is based on intensive scrutiny of a small number of objects. For example, sociology tends to use quantitative methods – studying humans by asking many people a few questions; while anthropology tends to use qualitative methods – studying humans by asking a few people many questions.

All this terminology is relatively new in human history, but the foundation of science, empiricism, is ancient. I don't want to back up too far into philosophy, so let me just say that science is based on what you can experience. Scientists use what you can see for yourself with your own two eyes (or some extension of your eyes, like an electron scanning microscope). If a scientist makes an argument that a fossil belongs to an ancestor of *Homo sapiens*, they need to point out the same details that led them to that conclusion, and as scientists they are required to explain their ideas in a way that anyone else could see the same thing they are seeing and come to same conclusions. This is also an example of how science is "reproducible". Scientists don't get so excited about the **first** person to discover cold fusion, what makes it science is the **second** person to verify the results, or better said, fails to disprove the original hypothesis. Science is about disproving hypotheses. Science can say you´re wrong, but it can´t say you´re right. The goal of science is not about establishing Truth, and for that reason it is often not as satisfying as other branches of knowledge such as religion or art that can claim Truth with a capital "T". Try not to get too frustrated with statements that hedge their conclusions, or just admit that we just don´t know yet: this is a characteristic of good science.

Exercise 1.1.1.1

Read Deduction vs. Induction

How to Critically Review Scientific Articles

Scientific writing can be pretty dry. If you're chatting about a movie, probably the first thing you say is whether you liked it or not, and you'll build up an emotional story without any spoilers.

You need to shift gears for science. This may be disturbing to your self-esteem, but busy scientists usually don't care about how a scientific article makes you feel or whether you liked it or not. They want to know how much the data supports or disproves the hypotheses they are working with. They don't need spoiler alerts, they want as much spoiling as possible. There are more scientific articles than minutes in a human life, so scientists need the conclusion in the first paragraph, so they can decide whether to keep

reading or not. The process of doing science entails communicating several necessary components. With good scientific articles you should be able to easily find all of the following elements:

The Citation

A citation is how the article you are reviewing would look in a long list of references, works cited, or in a bibliography. It functions as a link between how you use the ideas in the article, and how the reader can get ahold of the article and read it themselves. It's important to stay anal-retentive about the format of a citation, so people can find the article. MLA and APA are popular formats, but much of anthropology uses the Chicago Style.

The citation functions as the title of your review and goes on top, like the format of an annotated bibliography.

This is the main difference between an essay and a critical review. An essay has a topic and a title, a critical review (or an annotated bibliography) just has a source, and then your thoughts about it below. So what goes below the citation? read on...

The Introduction to Your Review

Even though this is the order of elements in the final version, as you're working on your review, you want to skip ahead and come back here after the Conclusion.

Cover all the following sections and then summarize them into a single paragraph, like the annotation in a typical annotated bibliography, or an abstract that summarizes a longer work. If you organize your review with a paragraph for each section, and each paragraph begins with a topic sentence, then you can pretty much just copy the topic sentences word-for-word and you're done with the Introduction.

Because the introduction is basically a summary of your critical review, you need to do the critical review first, before the introduction.

Why all this jumping around between the article, introduction, and the rest of the critical review?

Your job as the writer is to make it easy for the reader to find the information they need as quickly as possible, and decide if they need to keep reading.

The Hypothesis

Don't waste time! Go straight to the core of the article. What is the author trying to prove? Hypotheses come from “Problems” and “Research Questions” but they are reworded as answers. The hypothesis shouldn't end in a question mark – it's the answer, stated as a concise declarative sentence that is either descriptive ("This is that.") or causal ("This causes that."). The question you and the scientists want to answer is how well was the hypothesis supported or disproved.

A Background

Now that you've stated the hypothesis, you can back-up and put it in context. Why is it important? How does the hypothesis connect to other research? For this class you want to refer to the other articles in the same section and explain how this research fits into those broader topics. Why did Arnie put this article in this section? What other sections might it fit into? What does this article have to do with physical anthropology?

This is sometimes called the “Problem” or “Research Question”. In a larger scientific write-up you might make this section into a “Literature Review”, which is a summary of everything that has been written about the subject before your contribution.

The Methods

Now that you've given the essential hypothesis, and given its background, you can add more details about the article itself. What did the scientist do? What techniques or technology were used? How did they look at something? What empirical senses were involved? This is very different from what the scientist thinks–which you find in the hypothesis or the conclusion.

Some Data

If methods are **how** the scientists looked at something, then data are **what** they saw. What did they see? feel? hear? touch? sense? What senses they used are methods, how their senses responded are data. Data are often reported as “Findings” or “Results”.

Scientists are obligated to make their data public so that other scientists can attempt to reproduce their conclusions. An article is already summarizing the data, and a review of an article should summarize it even more. In the humanities and social sciences, hypotheses are rarely completely falsified or overwhelmingly supported, how “true” a hypothesis is, depends on the quality and quantity of the data that supports it.

You want to give enough details to connect the hypothesis to the conclusion.

The Conclusion

The conclusion has two parts: yours and the author's. You want to present the conclusions that the article came to, and you want to wrap your review up, summarizing what you've done so far. In the conclusion, the scientists let us know what they think. How well did the data support the hypothesis? Was the hypothesis testable? Was it reliable (usually a review of the methods)? Was it valid (usually a critique of the background)? Was it verifiable; would it be possible for someone to repeat the same process? How did the sources cited or the “Literature Review” connect to the data? What further research do the scientists suggest?

In a *critical* ***review***, you want to present how the scientists answered these questions. As a ***critical*** *review,* you also want to answer these questions yourself, evaluating the strengths and weaknesses of the article. If this were a ***peer*** *review* you would be a peer of the author, a colleague, you would have the same background knowledge as the author, and be more likely to thoroughly understand the article and be able to evaluate it. But, you're taking an introductory class, you're not an expert yet, so try not to get cocky and feel like you are supposed to attack the author.

Science thrives on criticism. The goal of a peer review is to help the author fix their mistakes and get better.

Note

Here's a great video of Austin's Butterfly: Building Excellence in Student Work, that reflects the ideal way the scientific community comes to consensus.



Austin's Butterfly: Building Excellence in Student Work from EL Education on Vimeo.

Now that you're done with the body of your review, go back and write a short introduction to YOUR work in the form of an abstract or annotated bibliography.




Figure **1.1.1.2** - *http://xkcd.com/552/* (CC BY-NC 2.5)

- a good source for critically evaluating articles
- Twenty tips for interpreting scientific claims
- Science can't prove anything: how scientists define *theory*, *proof*, *uncertainty*
- The Research Methods Knowledge Base provides a good introduction to these concepts and terms

A Rough Guide to

SPOTTING BAD SCIENCE

Being able to evaluate the evidence behind a scientific claim is important. Being able to recognise bad science reporting, or faults in scientific studies, is equally important. These 12 points will help you separate the science from the pseudoscience.

1. SENSATIONALISED HEADLINES

Article headlines are commonly designed to entice viewers into clicking on and reading the article. At times, they can over-simplify the findings of scientific research. At worst, they sensationalise and misrepresent them.

2. MISINTERPRETED RESULTS

News articles can distort or misinterpret the findings of research for the sake of a good story, whether intentionally or otherwise. If possible, try to read the original research, rather than relying on the article based on it for information.

3. CONFLICTS OF INTEREST

Many companies will employ scientists to carry out and publish research - whilst this doesn't necessarily invalidate the research, it should be analysed with this in mind. Research can also be misrepresented for personal or financial gain.

4. CORRELATION & CAUSATION

Be wary of any confusion of correlation and causation. A correlation between variables doesn't always mean one causes the other. Global warming increased since the 1800s, and pirate numbers decreased, but lack of pirates doesn't cause global warming.

5. UNSUPPORTED CONCLUSIONS

Speculation can often help to drive science forward. However, studies should be clear on the facts their study proves, and which conclusions are as yet unsupported ones. A statement framed by speculative language may require further evidence to confirm.

6. PROBLEMS WITH SAMPLE SIZE

In trials, the smaller a sample size, the lower the confidence in the results from that sample. Conclusions drawn can still be valid, and in some cases small samples are unavoidable, but larger samples often give more representative results.

7. UNREPRESENTATIVE SAMPLES USED

In human trials, subjects are selected that are representative of a larger population. If the sample is different from the population as a whole, then the conclusions from the trial may be biased towards a particular outcome.

8. NO CONTROL GROUP USED

In clinical trials, results from test subjects should be compared to a 'control group' not given the substance being tested. Groups should also be allocated randomly. In general experiments, a control test should be used where all variables are controlled.

9. NO BLIND TESTING USED

To try and prevent bias, subjects should not know if they are in the test or the control group. In 'double blind' testing, even researchers don't know which group subjects are in until after testing. Note, blind testing isn't always feasible, or ethical.

10. SELECTIVE REPORTING OF DATA

Also known as 'cherry picking', this involves selecting data from results which supports the conclusion of the research, whilst ignoring those that do not. If a research paper draws conclusions from a selection of its results, not all, it may be guilty of this.

11. UNREPLICABLE RESULTS

Results should be replicable by independent research, and tested over a wide range of conditions (where possible) to ensure they are consistent. Extraordinary claims require extraordinary evidence - that is, much more than one independent study!

12. NON-PEER REVIEWED MATERIAL

Peer review is an important part of the scientific process. Other scientists appraise and critique studies, before publication in a journal. Research that has not gone through this process is not as reputable, and may be flawed.

© COMPOUND INTEREST 2015 - WWW.COMPOUNDCHEM.COM | @COMPOUNDCHEM

Shared under a

***Figure* 1.1.1.3** - *2014 CompoundChem.com* (CC BY-NC-ND 4.0)

Faith

I introduced this section by discussing epistemology, the theory of knowledge. Science is one kind of knowledge, faith is another kind of knowledge. What you know can come from what you experience with your own empirical senses, or you can believe something that someone told you. Faith is complex and varies from person to person, but a concise definition can be found on the popular bumper sticker that reads: “God said it, I believe it, that settles it.”




***Figure* 1.1.1.3** - *Circular Reasoning* (CC BY-NC 4.0)

Radical fundamentalist Christianity in the US makes what should be a parlour room discussion between science and religion into a political debate with real educational consequences for all us. Scientists struggle to understand the complex mechanisms of evolutionary theory, but for many, the struggle is made more difficult by ideological barriers set up by faith-based opposition to science.




***Figure* 1.1.1.4** - *Jesus Camp. good 2006 documentary on radical fundamentalism*




***Figure* 1.1.1.5** - *Big Think*

We need to stress evolutionary theory because it is a fundamental explanatory device in biology. There is a famous quote by geneticist Theodosius Dobzhansky where he states “nothing in biology makes sense, except in the light of evolution.” Trying to make sense of biological systems, including human beings, without evolution is like trying to understand physics while claiming that the force of gravity doesn't exist. There are very few belief systems in the world that deny evolution, and it is totally compatible with most religions around the world, even the Pope has come out supporting evolution (* 2014), so you have no conflict if you are a Catholic. But unfortunately, we happen to live in a culture that prefers faith over science. Our technology has changed rapidly in the last couple of millennia but our mindset has not quite kept up. Many fierce battles are fought today in school boards around the country over the separation of church and state, and whether faith-based ideas (Creationism/Intelligent Design) should be taught in public schools. The battles have spilled out onto the streets with people declaring their beliefs with little symbols on the backs of their cars.



***Figure* 1.1.1.6** - *Al Seckel, John Edwards 1983*

Personally, I wish I didn't have to dwell on the issue so much, I have friends who are Creationists and we get along fine. But, in my role as professor of physical anthropology, I cannot accept Creationism or Intelligent Design as anything more than dangerous fallacies that interfere with students' ability to learn the required curriculum. For me to teach "both sides" would be a form of repressive tolerance (Marcuse 1965). I'm not leading any crusades to banish Medieval thinking from society, but I get worked-up about the issue. It's like professional frustration – that I've failed at my job as an educator – when I see people who are proud of their ignorance. They act like being stupid is somehow cool or something. It reminds me of how Cornel West describes the problem of nihilism in society today.

It bothers me too how religion is used reinforce socioeconomic class. Many of you are smart enough to transfer to Harvard with a scholarship and get six-figure jobs in the budding genetics industry in major cities around the globe, but if during your job interview, you start spouting Intelligent Design ideology, your scientific credentials go down the drain, and you're back to flipping burgers at In-and-Out.

So, if you think Creationism/Intelligent Design is a load of crap, fine, so do 99.9% of the scientists in the world. But, if you are Christian, you don't have to abandon your faith for this class. I would get in trouble if half the class reasoned, "Well since my teacher has proven that a few poetic lines written thousands of years ago can't be interpreted literally, now I should do the exact opposite of all the moral precepts in the Bible, and become a Satanic mass-murdering tweeker." There are millions and millions of scientists who believe in Christ and evolution and find no contradiction between the two. If that doesn't console you, maybe it'll help to think of this class as an exercise in "know thyne enemy". You don't have to sign a "God is dead!" pledge, and you don't even have to actually believe in evolution to pass this class, you just have to understand it well enough to be able to regurgitate a few of the things I want to hear. But, be forewarned, I'm not a minion of the Devil; it's in my job description to test your faith and proselytize the wisdom of evolutionary theory.

There are very, very few people who actually take The Bible literally. Almost everyone interprets the meaning of the words in The Bible relative to their own language, historical milieu, and personal circumstances. Biblical scholars use the word *hermeneutics* to talk about the different ways that passages in The Bible can be interpreted. For example, there are several passages in the King James' version that says it's harder for a rich man to get to heaven than a camel to go through the eye of a needle. Some Biblical scholars think that the passage was poorly translated from the Aramaic to the Greek, and the mistake continued to Latin, and then English. They have found similar sayings from around that historical period that refer to the difficulty of threading a needle with a camel **hair**, or thread or rope made of camel hair, because camel hairs are so thick. Another interpretation is that Jerusalem had a small gate, called the Camel **Gate**, that was difficult to pass through if you were rich and carrying all your stuff. For me the metaphors of a camel hair through a needle or a rich guy with gear squeezing through a small door make a lot more sense than this image of an actual camel floating through some giant needle. Stuff gets lost in translation. Have you ever played that game, "telephone", where you get a big circle with your friends and one person makes up a complicated sentence and whispers in the ear of the person next to them, who whispers it in the ear of the person next to them, and as the message gets passed around the circle, people miss-hear things, or forget things, and the message changes. And, when you compare the original message to how it ended up, it often sounds silly.




***Figure* 1.1.1.7 -** *Flammarion engraving 1888, notice the pillars that hold up the heavens*

Some of the few people who take the Bible literally have been ridiculed into the closet. The Flat Earth Society took literally a passage from The Bible that said the heavens are held up by four pillars at the corners of the earth, and the image of the sun "rising" literally. They actually believed that if you walked far enough to some corner of the earth you would bump into a big pillar. Nowadays, it's hard to find an honest proponent of the Flat Earth theory, but they were around by the 1950s. Now, everyone just laughs at them, like you would probably laugh at someone one who claims the sun revolves around the earth. But on February 17, 1600, this was no laughing matter, when in the Campo di Fiore, Rome, Giordano Bruno was burned at the stake for advocating the heretical belief proposed by Copernicus known as *heliocentrism.*



Figure 1.1.1.8 - *Statue of Giordano Bruno, Campo di Fiore, Rome, Italy* (CC BY-NC 4.0)




Figure **1.1.1.9** - *http://i.imgur.com/AReqgfP.gif*

I'm glad that we're at a point in history where I don't have to worry about being burned at the stake for teaching the theory of evolution, and I'm glad that we're at a point where we can just poke fun at Creationists through Darwin fish on our cars, or spoofs such as The Church of the Flying Spaghetti Monster. But I don't think the battle is over. Obama undid many of attacks on science that characterized the Bush administration, but the Trump administration promises a radical attack on science, and an era of "alternate facts". We'll see...

Notes

- A TED talk video of how science can help cure cancer
- A primer on logic and arguments

Vocabulary

- literature review
- abstract
- research question
- epistemology
- anthropology
- archaeology

- biological anthropology
- Creationism
- deduction
- empiricism
- faith
- hermeneutics
- hypothesis
- induction
- Intelligent Design
- laws
- linguistics
- physical anthropology
- qualitative
- quantitative
- science
- taxonomy
- theory

Imagination Questions

- What would happen to science if radical Christian fundamentalists organized a military coup and took control of education in the US?
- What if artificial intelligence advances to the point where robots can do their own science?
- How strong is your belief in science or faith? If you are religious and someone convinced you your religion was wrong, would you give it up? If you believe in science, and you saw God, would you give up your scientific beliefs?

This page titled 1.1.1: Science is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **1.1: Science** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.1.2: Anthropology as a Science

Anthropological Imagination

The anthropological imagination (anthropological perspective) is how anthropologists see the world. Anthropology differs from other sciences because it emphasizes holism and genealogy. The emphasis on genealogy for cultural anthropology implies a focus on the family (domestic structure). The emphasis on genealogy for physical anthropology extends the metaphor of the family tree from an individual and their family, to a family tree writ-large that uses phylogenetic taxonomy to contextualize the human species. Anthropology's emphasis on holism implies a balance between different approaches and many subfields. The four main subfields of anthropology are cultural anthropology, physical (biological) anthropology, archaeology, and linguistics. Anthropologists balance objective and subjective epistemologies.

I think the best way to get a sense of how anthropology differs from other branches of science is to understand the anthropological imagination. I borrowed the concept of the anthropological imagination from one of my professors, Dr. Wade Pendleton, who in turn borrowed it from an introductory anthropology book (Dimen-Schein 1977), who based it on an important sociology book, *The Sociological Imagination* by C. Wright Mills (1959).

There isn't much published about the anthropological imagination, but it is basically the same thing as what most anthropologists call the "anthropological perspective" (Jurmain 2011:19-20; Field 2011). I like the connotations of "imagination" in the way it has been used by John Lennon and recent social movements to recognize the agency that people have to go beyond their cultural constraints. Franz Boas (1858-1942), one of the founders of anthropology, described this as people's need to break the "shackles of tradition" (* *Franz Boas: Shackles of Tradition*). It is especially related to cultural anthropology, where "the world is as you see it", the idea that if people believe in ghosts, then you as a scientist need to start with the hypothesis that those ghosts really exist. That might seem weird to many scientists, but anthropologists need to balance a detached, objective, way of seeing, with the subjective reality of the people they join to study.

Anthropologists balance several seemingly contradictory philosophies. I like to see the anthropological imagination as tendencies between two extreme poles, and though they may lean towards one side or the other, they can never really go to the extreme in any direction.




Figure 1.1.2.1 - *The anthropological imagination balances extremes* (CC BY-NC 4.0)

The principle method of fieldwork in cultural anthropology is called participant-observation, and participant observation exemplifies the balance of anthropology. Anthropologists must objectively study people as an outsider, but they also become part of that culture. They must be culturally relative, and not judge a foreign culture by the standards of the researcher's culture, but they also have their own ethical principles that come from the anthropologist's own culture, and there are limits to how dogmatic anthropologists can be about cultural relativism, and scientists (myself included) need to be a little bit ethnocentric to support things like the *United Nations Declaration of Human Rights and tell another culture that they're doing it wrong. Because our science is so tied to humans we can't avoid asking ethical questions, or as Sir Raymond Firth put it "Cui bonum?" *For Whose Good?(Schepher-Huhges1981)* What is the purpose of doing anthropology?

This is true to a lesser extent in our class, regarding physical anthropology. We are objectively discussing some aspects of a biological species that has been around hundreds of thousands of years, and has a few distinct characteristics from other animals, but at the same time, we are talking about ourselves, my relatives, the people who gave me the genes I have now, that enable me to think, and type, and wish that this font was easier to read on this crappy screen.

I think the two most distinctive characteristics of anthropology are that it is holistic and it emphasizes genealogy. Holism means that it tries to understand all facets of the human condition. This has many implications. Anthropology is multi-disciplinary, it involves many branches of knowledge. By the early 1900's Franz Boas solidified anthropology into four interrelated subfields: cultural anthropology, physical anthropology, archaeology, and linguistics. As anthropologists began solving real world problems, some advocated for a fifth subfield: applied anthropology. For each of these sub-fields you can combine practically any other branch of science to make sub-sub-fields, depending on your specialization. Don't get too hung up about the correct taxonomy for these branches of knowledge. It can be rewritten in many forms depending which branch you want to emphasize. But in anthropology, all specialists need to have a broad overview to fit their research into the larger questions of what it means to be

human, and this incorporation of specific issues into broad questions requires a holistic approach. A good example for physical anthropology is the concept of *biocultural evolution*, the idea that to understand human evolution we need to look at both biology and culture.

If you take a cultural anthropology class you will see the study of culture requires a holistic approach in its own right because culture is integrated and all-encompassing; you need to study all the elements of culture together and their interaction. Another consequence of holism, and the multi-disciplinary approach of anthropology, is that anthropologists tend to be skeptical of unicausal arguments. A unicausal argument is something like "people have wars because they have an aggressive nature." Anthropologists understand that human nature is supremely complex, and that culture can drastically change any human characteristic that people try to claim is biologically determined. Sure, people are aggressive, you can look at chimpanzees and hominid weapons, but humans are also peaceful, you can look at bonobos and the amazing art of the Upper Paleolithic. Try to keep this point in mind when you´re writing for this class: avoid unicausal arguments, give all sides of an issues, avoid oversimplification, explore the evidence that supports each position.

Exercise 1.1.2.1

Compare ETHNOGRAPHY to ETHNOLOGY

Another general emphasis in anthropology is on genealogy. In cultural anthropology, the structure of the family is usually a core element of a culture. In physical anthropology, I like to view our emphasis on classification and taxonomy as just an attempt to better understand the branches of our own family tree.

The anthropological imagination is also something that you as an individual will use to better understand yourself and your place in the world. Biological and cultural explanations can be useful in solving your own problems. Having an answer to “Why am I sweating right now?” means understanding the cooling mechanisms that our ancestors evolved over tens of millions of year, and the fight-or-flight response in response to stress that involves putting your own personal financial problems into a cultural context where education is touted as the method of class mobility, yet restricted by public policy that raises tuition, textbook prices, and limits financial aid.

Subfields of Anthropology

Exercise 1.1.2.1

Read Dennis O'Neil: The Subfields of Anthropology and Wikipedia's Explanation

Many anthropologists consider applied anthropology as a fifth subfield. I prefer to think of it as a research goal or purpose that cross-cuts all of the four subfields. For example, there are projects that can be considered applied linguistics [language renewal], or applied archaeology [Incan agronomy]. There are many applications for physical anthropology, especially in medicine. The word *forensic* means "legal", but most forensic anthropology tends to be a subfield of physical anthropology - we use what we know about human biology to help solve crimes.

Notes

- The Argentine Forensic Anthropology Team is a good example of forensic anthropology. They have been recently working to identify the 43 missing students from Ayotzinapa, Mexico.
- Check out careers in genetic counseling
- The Health and Medicine chapter from a cultural anthropology textbook
- Desmond Morris' The Language of the Body: a good video combining physical anthropology, cultural anthropology and linguistics

Conclusion: Anthropology and Science

In conclusion, anthropology is mostly a science, but has many aspects of humanism. For a more traditional introduction to anthropology and science read the Dennis O'Neil overview of anthropology.

Epistemology for Physical Anthropology

Epistemology means the study of how we know what we do. *Taxonomy* comes from the Greek word for “branches”. Here is a taxonomy of knowledge for this class:




***Figure* 1.1.2.2** - *An epistemology for physical anthropology* (CC BY-NC 4.0)

You add or subtract boxes, and draw the arrows differently depending on what you want to focus on, and because anthropology is holistic these charts don't make much difference; anthropologists include all relevant aspects of knowledge. We will focus on the bottom part of the flowchart, but always keep the bigger picture in the back of your mind.

Vocabulary

- anthropology
- archaeology
- biology
- culture
- epistemology
- ethnography
- ethnology
- linguistics
- participant-observation
- phylogenetic
- taxonomy

Imagination Question

How does the anthropological approach to understanding human beings differ from other classes you've taken?

This page titled 1.1.2: Anthropology as a Science is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **1.2: Anthropology as a Science** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

CHAPTER OVERVIEW

1.2: Introduction to Biology

Physical anthropology is also called biological anthropology. You need to understand the basics of biology before you apply it to humans. We are going to use these ideas as the basis for explaining phenomena in later sections, such as: Why primates scream at each other? How are those two fossils related? Why do people look different from each other? What are people going to look like in a million years? Now is a great time to review your notes from your High School biology class.

1.2.1: Introduction
1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology
1.2.3: Evolutionary Theory (Part 1)
1.2.4: Evolutionary Theory (Part 2)
1.2.5: Evolutionary Theory (Part 3)
1.2.6: "Forces" of Evolution
1.2.7: Genetics, Cellular Biology, and Variation (Part 1)
1.2.8: Genetics, Cellular Biology, and Variation (Part 2)
1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria

Thumbnail: Darwin's finches (also known as the Galápagos finches) are a group of about fifteen species of passerine birds.[1][2][3][4] They are well known for their remarkable diversity in beak form and function. From "Voyage of the Beagle." (Public Domain).

This page titled 1.2: Introduction to Biology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

1.2.1: Introduction

Physical anthropology is also called biological anthropology. You need to understand the basics of biology before you apply it to humans. We are going to use these ideas as the basis for explaining phenomena in later sections, such as: Why primates scream at each other? How are those two fossils related? Why do people look different from each other? What are people going to look like in a million years? Now is a great time to review your notes from your High School biology class.

Biology is a broad field that ranges in scale from the microscopic to the geographical. Evolutionary theory is crucial to biology. Charles Darwin's theory of natural selection came from a historical context where many other scientists were contributing to evolutionary theory. Natural selection means that individuals compete for resources and those with the variations that make them more fit to survive in a certain environment tend to survive and reproduce more. Sexual selection is the part of natural selection that focuses on competition for reproduction. Darwin understood the importance of variation.

Gregor Mendel used math to explain how variation is inherited. Population genetics used math to show how inheritance works in large populations. The modern synthesis combined Mendelian genetics and population genetics, to codify evolutionary theory into four forces: mutation, natural selection, migration, and genetic drift. Inheritance, mutation, and other sources of variation can now be understood through cellular biology and genetics. Humans are made up of cells. Cells have DNA. DNA is the code that directs protein synthesis. Proteins direct the functions of life.

Sickle cell anemia is an example of how the holistic approach of anthropology can use many subfields, mostly from biology, to understand the origins and consequences of an important human disease.

This page titled 1.2.1: Introduction is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.1: Introduction** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology

Anthropology is a broad field that incorporates many sub-fields and borrows from many other disciplines, so the space that is relevant to physical anthropology is also vast. It ranges from the atomic particle that causes genetic mutation—the primary cause of evolution—to the plasticity of the human body because of our need to adapt to seasonal changes caused by the elliptical orbit of the planet in the solar system. Anything bigger or smaller is beyond the scope of physical anthropology. Why atomic particles behave the way they do is a question of physics. Whether life exists on other planets is a question of astrobiology. Both are good themes for philosophy and science fiction, but are beyond the scope of this class.

Meters	Common Measurement	Abbreviation	Relevance
10^{-10}	1 ångström	1 Å	Size of an atom
10^{-9}	1 nanometer	1 nm	Diameter of DNA helix, size of a base
10^{-8}	10 nanometers	10 nm	Size of a codon (3 base pairs)
10^{-7}	100 nanometers	100 nm	Locus, length of a typical gene
10^{-6}	1 micron	1 µm	Cell nucleus, *Plasmodium falciparum*
10^{-5}	10 microns	10 µm	Typical primate cell
10^{-4}	100 microns (micrometer)	100 µm	Patch of melanin, width of hair, human egg
10^{-3}	1 millimeter	1 mm	The height of a cusp on a Y-5 molar
10^{-2}	1 centimeter	1 cm	Human ear bones,*Anopheles* mosquito
10^{-1}	10 centimeters	10 cm	Length of a human's opposable thumb
10^{0}	1 meter	1 m	Space that an individual primate occupies
10^{1}	10 meters	10 m	Typical sleeping area of a primate social group
10^{2}	hectare/ 2.5 acres, football field	100 m	Typical core area of a primate social group
10^{3}	10 hectares, 1 kilometer	1 km	Typical territory of a non-human primate social group
10^{4}	100 hectares	10 km	Typical home range of a non-human primate social group
10^{5}	100 kilometers	100 km	
10^{6}	1,000 kilometers, continent	1000 km	Typical range of a non-human primate species; range for most of human evolution
10^{7}	10,000 kilometers, the surface of the planet Earth	100000 km	The range of all biological evolution
10^{12}	1 billion kilometers, 1 terameter	1 Tm	The Earth's orbit in the solar system, includes meteors and solar radiation

Notice that most of the larger ranges are relevant for hominid evolution and primate behavior, the medium sizes are used in osteology and to compare human and hominid variation. The smaller sizes can describe human variation and genetics. Refer back to this chart as we cover those relevant sections.

Exercise 1.2.2.1

Use this chart and watch the movie POWERS OF TEN



> *"Recognize that the very molecules that make up your body, the atoms that construct the molecules, are traceable to the crucibles that were once the centers of high mass stars that exploded their chemically rich guts into the galaxy, enriching pristine gas clouds with the chemistry of life. So that we are all connected to each other biologically, to the earth chemically and to the rest of the universe atomically. That's kinda cool! That makes me smile and I actually feel quite large at the end of that. It's not that we are better than the universe, we are part of the universe. We are in the universe and the universe is in us."*
> *-Neil deGrasse Tyson*

Note

More on scale and measurements: human scale

Imagination Question

When you look in the mirror do you see yourself as a trillion cells, or a trillion, trillion atoms which came from stardust, or a collection of selfish genes, or an individual?

This page titled 1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.2: Scale, Human Space, Powers of Ten for Physical Anthropology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.3: Evolutionary Theory (Part 1)

Don't go to a dictionary and look up the definition of "evolution". Most of the definitions are just going to reinforce your confusion. For this class we are using the **biological** definitions of evolution: the splitting of a lineage into two new species, or the change in allele frequency in a breeding population from one generation to the next. For this class an individual can never evolve in their lifetime, a cell phone operating system can never evolve, only a population or a species can evolve. Darwin understood the problem with the word "evolution" and preferred the phrase "descent with modification", but that obviously never caught on.

Understanding evolutionary theory is crucial for all the chapters that follow. For example, in the paleoanthropology section we will make arguments about how natural selection gave our ancestors advantages in different environments, and we will argue how gene flow kept our recent ancestors from becoming separate species. If you don't learn about the forces of evolution now you won't understand the arguments we're going to make later.

Evolutionary theory is a difficult concept. In the last section we discussed the ideological barriers that extremely dogmatic religious people have that might keep them from understanding evolutionary theory. Another common misconception of evolution is that there is inherently something good about it, that to evolve means to progress. The simple definition of *evolution* is just “change”. It doesn't mean change for the better, or change for the worse, just that a species is different than what it used to be. It's important to separate how a word is used on the street from the very specific definition we are using here in our biological context. The main reason people have problems with a neutral definition of evolution is that we've had thousands of years of cultural baggage that clouds our empiricism. The philosopher Aristotle stressed the importance of finding the essence of a thing, and from this we get the concept of essentialism. People often think of humans as diverging from or progressing towards some ideal form, but these are cultural constructs, not what we empirically see in biology. We carry baggage around that makes simple concepts seem counterintuitive. Understanding the incorrect ideas helps us accept the correct ones.

Exercise 1.2.3.1

Read more about Darwin and design.

History of Evolutionary Theory, up to Darwin

It helps to understand evolutionary theory if you understand how it “evolved”. I put “evolved” in quotes because for this class I want to reserve the word *evolution* to mean “biological evolution” and not confuse it with historical and cultural changes. There are different ways to study history, and for this section we need to be careful not to fall into the trap of thinking of history as a series of Great Men – important people who change the world through their personal actions. This way of looking at history is very convenient for an introductory class, because you just need to make a handful of flashcards with the names on one side, and what they're famous for on the other. But, try to look beyond the individuals, and imagine the broader movements that were going on at the time these individuals lived.

It helps sometimes to put people and concepts into two columns: did they contribute to evolutionary theory or did they detract from the development of evolutionary theory? But, many historical figures can be put in both columns, depending on their effect on history, and what stage of their life you look at. For example, Linnaeus made amazing strides in biological taxonomy, but opposed evolutionary theory, at least until the end of his life; Lamarck created a great theory of evolution, but it was wrong; Cuvier contributed to our understanding of extinction, but argued against evolution and fixity of species, Lyell argued against biological evolution until after Darwin convinced him; Wallace's evolutionary theory was mixed with quirky spiritualism; Darwin's theory of Natural Selection was revolutionary, but his emphasis on blending made it more likely to ignore important contributions from scientists like Mendel, and gradualism made it harder to accept punctuated equilibrium. Just realize that for an introductory class, we only have time to give you a cardboard cut-out of these fascinating and complex individuals and their historical milieux.

Exercise 1.2.3.2

Read Dennis O'Neil's Intro

The Fixity of Species

In this modern world, we are so used to change that it's hard not be ethnocentric and imagine a time when things were pretty much the same as it was for your grandparents, and you expected things to be pretty much the same for your grandkids. Imagine growing up and liking the same music as your great-grandparents. Most religions around the world are conservative. God creates a world and some people to interact with it. And the reasoning often goes that if your deity is all-powerful, His creations would be perfect and He wouldn't need to keep tinkering with them. When people looked around at all the living creatures on the planet, they assumed that everything was stuck the way they saw it. This concept is called *the fixity of species*, and "fixity" in this case refers to being in a fixed position; unchanging.

> *Evolutionary theory is diametrically opposed to the fixity of species.*

The Great Chain of Being

The Great Chain of Being incorporates the concept of the fixity of species but ranks life into better or worse, noble animals like eagles and lions go towards the top, slimy animals like worms and eels go down to the bottom. Noah's Ark usually boards the same way, first-class passengers first. When applied to humans, the Great Chain of Being has always functioned to perpetuate political oppression such as racism, sexism, and classism. The dominance of the Great Chain of Being in Eurasian philosophy makes it hard at first for many students to accept the complete absence of any system of ranking or progress in evolutionary theory.




Figure **1.2.3.2** - *Wikipedia;1579 drawing of the Great Chain of Being from Didacus Valades, Rhetorica Christiana.*

John Ray

John Ray (1627-1705) is important to us because he invented the concept of the species (the word "species" is one of those Latin words that doesn't change between the singular and plural; "one species, two species"). Species are the most fundamental way of grouping life forms. You might ask how a book entitled "The Wisdom of God Manifested in the Works of Creation" could contribute to evolutionary theory but a common theme in all science is how scientists get some things right and some things wrong. In Ray's book *History of plants* he grapples with how to group the varieties of life, and he comes up with a concept that is still useful today – individuals that can reproduce are from the same species:

> *no surer criterion for determining species has occurred to me than the distinguishing features that perpetuate themselves in propagation from seed. Thus, no matter what variations occur in the individuals or the species, if they spring from the seed of one and*

> *the same plant, they are accidental variations and not such as to distinguish a species… Animals likewise that differ specifically preserve their distinct species permanently; one species never springs from the seed of another nor vice versa *[Ray 1686; Ray 1686]*

A species is something that exists over time and is capable of perpetuating itself, and variations within a single species occur. The Notice at the end of the quote how he insists on the fixity of species, and denies speciation, but we still use Ray's species concept today.

Carolus Linnaeus

Linnaeus is the father of taxonomy. Taxonomy was different from the Great Chain of Being because instead of grouping animals into biblical categories, he grouped them by biological characteristics: how they give birth, what they eat, aging, exterior movement, internal propulsion of fluids, diseases, death, glands, skin, and the shape of the inner ear (Foucault 1970). This list is very arbitrary compared to how today we compare the DNA of a species to classify it, but Linnaeus' taxonomy was radically different from the more poetic groupings of his time. Linnaeus was able to distinguish bats from birds, and snakes from eels and worms. His taxonomy changed over his lifetime and has been an important concern of biology to this day.

His approach to classifying humans codified scientific blunders regarding race that have continued over two centuries.

Exercise 1.2.3.3

Read this Introduction to Linnaeus




***Figure* 1.2.3.5 -** *1760 Hoppius based on Linnaeus, left to right: Troglodytes, Lucifer/Homo caudatus, Satyr, Pygmie*

Exercise 1.2.3.4

Briefly scan Linnaeus' system of nature 1740

Skim the pages on humans from this 1757 version

Buffon

George Louis Leclerc, Comte de Buffon wrote about how life changes according to the environment but didn't explain how.

Erasmus Darwin

Erasmus Darwin was the grandfather of Charles Darwin and his poetic writing about how life forms might change influenced his grandson, Charles.

* if you like prose, skim his chapter on generation

Jean-Baptiste Lamarck

Lamarck is known for his theory of Acquired Characteristics, also called the Use-Disuse theory. The theory goes that the physical traits you change during your lifetime get passed on to your kids. So if you believe in Lamarck's theory then you would expect Arnold Schwarzenegger's babies to be born with lots of muscles.

Of course, Lamarck was wrong. Physical characteristics acquired during an individual's lifetime are **not** transferred to its offspring (with a few epigenetic counterexamples). We'll show why he was wrong in the section on cellular biology. The reason Lamarck's

theory sounds so plausible is because **culture** is transmitted this way; what you **learn** during your lifetime can be taught to your children. But biological inheritance has very specific mechanical processes that we can now see in a microscope. You have to give Lamarck credit for coming up with an elegant theory of evolution. It has been disproven, but it was still a great theory for its time. Understanding why Lamarck was wrong will help you understand natural selection.

Exercise 1.2.3.5

Look at the classic example of Giraffes

Skim this Biography of Lamark

Georges Cuvier

Cuvier is important to evolutionary theory in promoting the concept of extinction. He saw fossils of animals no longer on the planet, such as giant elephants, and he correctly explained their disappearance as extinction, the big elephants just died out.




***Figure* 1.2.3.8 -** *evolution.berkeley.edu/evolib...cle/history_08*

His views on extinction contributed to evolutionary theory but his attempts to reconcile geology with The Bible did not. Cuvier is also known for his theory of Catastrophism: extinctions in the fossil record are only because of sudden changes in the environment, such as worldwide floods, like Noah's flood.

Geologists: James Hutton and Charles Lyell

Geology has always had a profound impact on evolutionary theory. Most of the scientists of the 18th and 19th Centuries called themselves "naturalists" and didn't distinguish between fields such as geology and biology. James Hutton and Charles Lyell were two of the most important scientists to use empirical observations to guide theories of how the Earth arose and became what we see today. Their two main contributions to evolutionary theory are the concept of geological time and the principle of uniformitarianism.

Geology is important to biological anthropology in many ways, such as dating and environmental context in paleoanthropology. In archaeology, Lyell's Law of Superposition is crucial – the deeper you dig, the older things are.

Geological Time

How long are you going to live? 50 years? 100 years? The length of time that you can perceive is limited by your own biological expiration date. The neat thing about geology is you start thinking like a rock. Imagine 1,000 years passing, 10,000 years, 100,000 years, 1,000,000 (a million) years, 1,000,000,000 (a billion) years… Lyell proved that the Earth is old, that rocks have been around for a long time. This was important to biologists and paleontologists because it showed that fossils are very old too. If life has been around for billions of years we can more easily accept that drastic changes could happen over that long a time span. Geological and astronomical time is sometimes called **Deep time,** and it is one of those very disturbing nonhuman concepts that science often forces us to think about.

Note

Bishop Ussher dates the world at 4004 BC.

The Principle of Uniformitarianism

What happened in the past continues to happen today. Most cosmologies around the world have some concept of a mythical past, where different laws of nature applied. Lyell demonstrated that huge geological formations can be explained by the same simple forces that we see today, such as erosion, wind, earth moved by earthquakes. We can explain something as awesome as the Grand Canyon with two simple processes that can be seen today: uplifting and erosion.

In biology, we can find simple processes that differentiate me from a banana slug. Both, the biological and the geological require huge quantities of time.

Exercise 1.2.3.6

Skim the 1830-3 Table of Contents of Lyell's Principle of Geology (10 pages)

Thomas Malthus

Malthus studied demography, how populations change. Overpopulation creates competition for limited resources. This idea of competition within a population is similar to the idea of competition within a species, and creates change. According to Malthus, humans can increase their population faster than they can increase their food supply. Having more people than food leads to starvation and competition for limited resources.

For example, imagine that Octamom has 8 daughters who each have 8 daughters who each have 8 daughters who each have 8 daughters, and in 5 generations you have a population of around 32,000. When Octamom works a plot of land she can produce one bushel of food per month, with her daughters helping she can produce 2 bushels, with her grandkids helping, 3 bushels, but by the time her great-great-grandkids have exhausted the nutrients in the soil, and run out of fertilizer made from non-renewable resources, her descendants are back to producing about 3 bushels a month, except it's like a Mad Max dystopia with thousands fighting over those few bushels, and most starving to death.

Darwin took Malthus' very specific observations of how humans compete for limited resources and generalized them to the broader field of biology.

Mary Anning

Mary Anning was the most famous fossilists of 19th century. A fossilist is someone who gathers fossils. A fossil is the imprint of a biological form, and we will come back to fossil in the sections on paleontology and paleoanthropology. Fossils represent biological data. If we go back to section 1.1 and consider the scientific method, it's important to remember that hypotheses are based on empirical observations, things we can see with our own senses. With a phenomenon like gravity you see the apple falling from the tree, but seeing evolution is more difficult. What many of the evolutionary thinkers of the 19th century got to see were the fossils collected by Mary Anning.

* Watch this puppet show of Mary Anning's life

Imagination Questions

- How big is life?
- How old is life?
- The line "same as it ever was", from the Talking Heads song "Once in a Lifetime" always reminds me of uniformitarianism and the song makes make think of deep time. Do you ever freak out about how short your lifespan is compared to how long the planet has been around?

This page titled 1.2.3: Evolutionary Theory (Part 1) is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.3: Evolutionary Theory (Part 1)** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.4: Evolutionary Theory (Part 2)

Charles Darwin

One of the most profound impacts Darwin had was to change how we ordered life, from a ladder (like the Great Chain of Being) to a tree.

Exercise 1.2.4.1

Read Dennis O'Neil's Darwin and Natural Selection




***Figure* 1.2.4.10** - *The page from Darwin's 1837 private notebook ("Notebook B on the transmutation of species," 1837–1838)*

Darwin's main Eureka! moment with evolutionary theory was that species (1) evolves into species A, B, C, and D, and everything in-between goes extinct:

> *"I think ... Case must be that one generation then should be as many living as now. To do this and to have many species in same genus (as is) requires extinction. Thus between A & B immense gap of relation. C & B the finest gradation, B & D rather greater distinction. Thus genera would be formed.—bearing relation."*

This is barely legible, confusing, and poorly written, so please don't use this as an example of anything you want to turn in for my class. Darwin's genius comes from translating this sketch into a 500 page bestseller by 1859, *On the Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*, which thoroughly explained Natural Selection and supported it with voluminous evidence. We remember Charles Darwin's not for the Aha! moment in 1837, but for the two decades of work he did afterwards to finish *On the Origin of Species* by 1859. The media tends to sensationalize scientific discovery (e.g. Isaac Newton gets hit by an apple and suddenly understands the mathematical equation that describes gravity), but for most scientists, it's more about hard work.




***Figure* 1.2.4.11** - *On the Origin of Species 1859*. Photo by Wellcome Collection (CC-BY-4.0)

Exercise 1.2.4.2

Open Charles Darwin's 1859 *ON THE ORIGIN OF SPECIES BY MEANS OF NATURAL SELECTION, OR THE PRESERVATION OF FAVOURED RACES IN THE STRUGGLE FOR LIFE.*; Read the description of the chapters carefully (page V to page X), then skim the rest for a few minutes

* A more readable version of On the Origin of Species, and more of Darwin's writtings,

Darwin had an unremarkable personal life. He wasn't a great student, he didn't have strong philosophical, political, or religious views.

* Good summary of Darwin's personal life

> *Darwin did not use evolution to promote atheism, or to maintain that no concept of God could ever be squared with the structure of nature. Rather, he argued that nature's factuality, as read withing the magisterium of science, cannot resolve, or even specify, the existence or character of God, the ultimate meaning of life the proper foundations of morality, or any other question with the different magisterium of religion. [Gould 1999]*

He wasn't oblivious to the social consequences of his findings, and was reluctant to publish.

Exercise 1.2.4.3

Watch *Creation*: a pretty good Hollywood movie about Darwin's personal and ethical problems; the story of how the movie was censored in the US says a lot about how this is still an important political issue.




***Figure* 1.2.4.12** - *Creation 2009*

Understanding Natural Selection




***Figure* 1.2.4.13** - "Evolution is no different than your family tree … except over a much longer period of time." by M.F. Bonnan/Florida Citizens for Science © 2010

A classic example of natural selection is the industrial melanism of the peppered moth. The same species of moth has black moths and white moths which can interbreed. When they land on white tree bark, the black moths tend to be eaten, and they become rare. Because of industrial pollution, the bark turned black, and now the white moths became rare. They cleaned up the pollution, the bark became lighter, the white moths survived more than the black moths. There are a few problems with the research but it is still a great example of how evolution works.

Note

- A video game that simulates Kettlewell's research
- Scientists may have found the gene that determines the color change of the moths.

Unfortunately, very few of us have grown up on a farm, so it's hard for us to understand where Darwin got the phrase "natural selection" from. We don't ever use the phrase "unnatural selection" but that's what human selection is. After every harvest the farmer notices the tastiest, biggest, or most fruitful plants and keeps their seeds until next season to plant, and animal breeders mate their best stock together. The farmers and breeders are selecting desirable characteristics and increasing the chance that they will be passed on to the next generation. From this process of selection we get all the food we eat today and an amazing range of domesticated pets.



***Figure* 1.2.4.13** "Big and Little Dog" By Ellen Levy Finch CC-BY-SA-3.0

We see In the archaeological record of Mesoamerica how teosinte was selected over thousands of years and became corn.



***Figure* 1.2.4.16 -** *http://teosinte.wisc.edu/index.html* photos by Hugh Iltis (left) and John Doebley (right), © The Doebley Lab, Department of Genetics, University of Wisconsin-Madison

Part of Darwin's genius was to recognize that the process farmers and animal breeders use to change a species, was also a natural phenomenon, that competition in an environment of limited resources would select those individuals who were more fit for that environment, and he coined the phrase "natural selection".

When we use the word "fit", don't think of 24-Hour Fitness, think of square peg fits into square hole, round peg fits into round hole. Some individuals fit into an ecosystem or an environment better than others.

Note

- Research on the origins of corn
- Article on genes and dog size

Sexual Selection

Darwin didn't stop with natural selection, he continued to expand evolutionary theory throughout his lifetime. Darwin avoided a few difficult ideas in *The Origin of Species* and left them for his 1871 work:* THE DESCENT OF MAN, AND SELECTION IN RELATION TO SEX**. This book tackled both a difficult scientific question – sexual selection, and a difficult social question – the origins of humans.** And within the book, Darwin left his most controversial chapters at the very end:* Part II Sexual selection of man (316-84) and General summary and conclusion (385-405).

Peacocks gave Darwin a headache. Natural selection says you are more likely to survive and reproduce if you are camouflaged, if you stay hidden and avoid predators. But, what gave Darwin, and the male chauvinist scientists of his time, the most trouble was the idea that the female was responsible for choosing the mate and driving evolution.



Figure 1.2.4.18 - *Female peahen and chick* (CC BY-NC 4.0)



Figure 1.2.4.19 - *Male peacock* (CC BY-NC 4.0)

Wallace was almost famous, but Darwin published before him.

Exercise 1.2.4.4

Read Dennis O'Neil's Darwin and Natural Selection

Watch the 'Animated Life of A.R. Wallace'

Skim Wallace's early, 1855, article on evolutionary theory: *On the Law Which Has Regulated the Introduction of New Species*, skip to the end and read the comments by Bernard Michaux who argued that Wallace probably believed in something close to natural selection because his article contains all the important themes of Darwinism: "gradualism, utility, adaptation to different environments, allopatric speciation, imperfection of the fossil record" (Michaux 2000).

Imagination Question

1) Natural Selection

In the novel *Cryptonomicon*, Neal Stephenson presents the origin of his protagonist as a series of survivors:

> *Like every other creature on the face of the earth, [he] was, by birthright, a stupendous badass, albeit in the somewhat narrow technical sense that he could trace his ancestry back up a long line of slightly less highly evolved stupendous badasses to that first self-replicating gizmo—which, given the number and variety of its descendants, might*

> *justifiably be described as the most stupendous badass of all time. Everyone and everything that wasn't a stupendous badass was dead.*

In the movie *Beast of the Southern Wild*, the teacher describes a mythical predator depicted in the Lascaux cave paintings as

> *a fierce, mean creature that walked the face of the earth back when we all lived in caves. They would gobble the cave-babies down right in front of their cave-parents. And the cavemen couldn't do nothing about it, because they were too poor, too stupid, too small.*
> *Who up in here think the caveman was sitting around crying like a bunch of pussies? Y'all gotta think about that.*
> *Any day now, the fabric of the universe is coming unraveled. Ice caps gonna melt, water's gonna rise, and everything south of the levee is going under. Y'all better learn how to survive now.*

Do you think of yourself as a badass? Do you give your ancestors credit for making you what you are? Do the hardships your ancestors overcame, inspire you to overcome the problems of the future?

2) Money talks

A) If Wallace's family had more money, evolutionary theory might have gone down a different path. Wallace was more spiritual than Darwin. Jonathan Marks defines "Atheistic Darwinism" as the use of Darwin to support atheism (2011:57-9). Maybe the backlash against evolutionary theory would have been less with Wallace at the helm?

B) The fictional protagonist of Elizabeth Gilbert's novel The Signature of All Things comes up with a "theory of competitive alteration" almost identical to Darwin and Wallace, through the same methods of empirical research and world travel, but she never publishes it, mostly because of her gender. How does power influence science?

3) Great Chain of Being

Below is a tongue-and-cheek reference to the Great Chain of Being from Margaret Atwood's novel *Life Before Man*:

> *Auntie Muriel is unambiguous about most things. Her few moments of hesitation have to do with the members of her own family. She isn't sure where they fit into the Great Chain of Being. She's quite certain of her own place, however. First comes God. Then comes Auntie Muriel and the Queen, with Auntie Muriel having a slight edge. Then come about five members of the Timothy Eaton Memorial Church, which Auntie Muriel attends. After this there is a large gap. Then white, non-Jewish Canadians, Englishmen, and white, non-Jewish Americans, in that order. Then there's another large gap, followed by all other human beings on a descending scale, graded according to skin color and religion. Then cockroaches, clothes moths, silverfish and germs, which are about the only forms of animal life with which Auntie Muriel has ever had any contact. Then all sexual organs, except those of flowers. [Atwood 1979]*

There has always been implicit racism in the Great Chain of Being; social inequalities were created by God. But there are broader questions. Why do humans feel the need to rank things? Why do we make top ten lists? Facebook started as a facial ranking network, Hot or Not. The grade you get in this class is basically part of a ranking system for employers and other schools. The difference in salaries between full-time and part-time faculty separates professors into two socioeconomic classes.

In the primate behavior section coming up, we'll see other primates similarly obsessed with dominance hierarchies. Perhaps ranking is human nature?

This page titled 1.2.4: Evolutionary Theory (Part 2) is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.4: Evolutionary Theory (Part 2)** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.5: Evolutionary Theory (Part 3)

Science doesn't stop with a founder. Creationists blame Darwin for evolutionary theory, but most biologists wouldn't call themselves “Darwinists”, any more than most physicists wouldn't call themselves “Newtonists.” Natural selection is just one factor of evolutionary theory. People know much more about evolution today than they did during Darwin's time. That doesn't mean that Darwin was wrong, just that science has progressed.


Darwin often gets blamed for *Social Darwinism*, the political ideology that extends "survival of the fittest" to justify exploiting the poor, but Darwin didn't come up with this phrase nor did he apply Natural Selection to human society. Social Darwinism was invented and promoted by others such as Herbert Spencer, Thomas Huxley, and Francis Galton. Charles Darwin objected to the application of his biological model to human social structure, and he definitely would have objected to the "Darwin Project" battle royale game, and "Darwin Awards" given out in his name.

So far, our overly broad unifying theories haven't been worth much more than interesting metaphors, so please try to separate nature from nurture: biology from culture.

Gregor Mendel

Even though neither Darwin nor Mendel knew about how heredity worked at the cellular level, it's almost impossible to talk about the consequences of their work without referring to what we know now. So we will introduce a few terms in this section that are anachronistic, and we'll wait to explain them in depth until the section on cellular biology.

The genius of Mendel is how he used mathematics to show how inheritance worked.



Courtesy of the Mendelianum, Moravian Museum, Brno.
Noncommercial, educational use only

***Figure* 1.2.5.22** - *Mendel counted peas*

Dominance and Recessiveness

Remember that Gregor Mendel (1822-1884) didn't know about DNA when he did his experiments, he didn't see meiosis in the microscope, he wasn't directly involved in the debates over evolution, but he found one of the sources of variation that Darwin's theory of natural selection relies on, and he discovered two important principles that are the foundation of genetics: The Principle of Segregation and The Principle of Independent Assortment. Darwin knew that variation was crucial to his theory, but he didn't know the source of variation.

The pea plant has variation. Some seeds are smooth, some wrinkled; some yellow, some green. Some pods are inflated, some constricted; some green, some yellow. Some flowers are purple, some white; some along the stem, some at the top. Some stems are tall, some are short. Mendel was careful to exclude other kinds of variation: how some plants are eaten by snails, some don't get enough water, some too much sun, some are cooked in soup, some peas are overcooked, some shot through straws. Mendel ignored all these things that happen to peas and only paid attention to this first set of variations, the either/or inherent characteristics that can be seen.

Mendel was rediscovered around 1900. Theories of inheritance at the time of Mendel focused on blending, for example, one parent with extremely dark skin and one parent with extremely light skin have a child who is neither very light, nor very dark, but a color that is in between the extremes. But when Mendel bred purple flowers with white flowers, he got only purple flowers, and then when he bred those purple flowers together, in the next generation he got mostly purple but some white ones. The white flower trait disappeared and then came back. The purple color dominated the white one, but the recessive white color was not gone for ever, it came back in a later generation. If you cross a purple flower with a white flower, Darwin would have expected a whitish-purple flower. What happened to the blending?

If you don't remember from your high-school biology class, here are some basics that we got from Mendel:

- Children resemble their parents.
- Genes come in pairs.
- Genes don't blend.
- Some genes are dominant.

The Principle of Segregation

Good science can come from unlikely sources. Mendel would momentarily escape from the duties of a monk in a cold monastery.

> *Mendel frequently took sanctuary in the little two-room building nestled into a corner of the monastery courtyard right up against the brewery next door. It gave him not only blessed warmth but also the space to engage in his scientific pursuits -- which would, he believed, prove important enough in time to earn him a place in the annals of horticulture. He had filled the glasshouse's long tables with pots of pea plants, each carefully labeled as to seed source and variety. His immediate goal was to breed these peas, thirty-four different seed types in all, after allowing them to self-fertilize for two full years. In the speeded-up growing seasons of the glasshouse, two years of growing meant perhaps six full generations -- enough to assure Mendel that the seeds were indeed what they appeared to be. [Marantz Henig 2000:14]*

Mendel isolated and bred different pea plants together and observed the characteristics of their offspring, and what really amazes me is that he counted them--and looking at the numbers he noticed patterns, and used simple math to work out the ratios of different traits.

First, he crossed true breeding plants (P) to get heterozygous plants, called hybrids (F_1), and then he crossed those hybrids with each other ($F_1 \times F_1 = F_2$) and counted:

> *Expt 1: Form of seed. — From 253 hybrids 7,324 seeds were obtained in the second trial year. Among them were 5,474 round or roundish ones and 1,850 angular wrinkled ones. Therefrom the ratio 2.96:1 is deduced.*
>
> *Expt 2: Color of albumen. — 258 plants yielded 8,023 seeds, 6,022 yellow, and 2,001 green; their ratio, therefore, is as 3.01:1.*
>
> *[...]*
>
> *Expt. 3: Color of the seed–coats. — Among 929 plants, 705 bore violet–red flowers and gray–brown seed–coats; 224 had white flowers and white seed–coats, giving the proportion 3.15:1.*
>
> *Expt. 4: Form of pods. — Of 1,181 plants, 882 had them simply inflated, and in 299 they were constricted. Resulting ratio, 2.95:1.*

> *[...]*
>
> *Expt. 5: Color of the unripe pods. — The number of trial plants was 580, of which 428 had green pods and 152 yellow ones. Consequently these stand in the ratio of 2.82:1.*
>
> *Expt. 6: Position of flowers. — Among 858 cases 651 had inflorescences axial and 207 terminal. Ratio, 3.14:1.*
>
> *Expt. 7: Length of stem. — Out of 1,064 plants, in 787 cases the stem was long, and in 277 short. Hence a mutual ratio of 2.84:1. In this experiment the dwarfed plants were carefully lifted and transferred to a special bed. This precaution was necessary, as otherwise they would have perished through being overgrown by their tall relatives. Even in their quite young state they can be easily picked out by their compact growth and thick dark–green foliage. [Mendel 1865]*

Notice that the ratios of the F_2 generation work out to about 3:1 which means that three plants have the dominant phenotype for every one plant that has the recessive phenotype. Mendel showed that each trait (seed color, seed shape, pod shape, pod color, flower color, flower position, stem length) is determined by a pair of characters, and they get them from their parents, one from the pollen cell and one from the egg cell, which come together to form the embryo. When the pollen and egg cells are made, these two characters are "segregated" so each egg and pollen cell has only one character. In genetics we now call these traits, *genes*, and the pair of characters is called a pair of *alleles*. From cellular biology, we now know that the segregation of alleles during the production of eggs and sperm is called *meiosis*. We'll come back to this in the sections on genetics and cellular biology.

The Principle of Independent Assortment

If you take true breeding plants with two different traits, like form of seed and color of seed-coat, cross them together, you first get all of the dominant trait. Then if you cross those new versions again, you get some interesting numbers 9:3:3:1 The numbers reveal that there's no connection between the traits; the traits are independently assorted. We can now explain this with cellular biology because the two traits are on different chromosomes.

Punnett squares

A Punnett square is a grid or matrix that represents the outcomes of different combinations. They are often presented as proofs of Mendel's Principle of Segregation and Principle of Independent Assortment, but Punnett squares came after Mendel, and I think it's important to understand the steps Mendel went through in his research: empirical observations of pea plant variations, breeding true-breeding plants, crossing specific traits, getting weird results, counting them, working out simple ratios, explaining the ratios as biological Principles as to how the peas (and all life, including humans) reproduce and transmit the information using traits from parent to offspring. Punnett squares are graphic representations of sexual reproduction: all the possible sperm are one axis, all the possible eggs on the other, and in the middle are all the possible combinations of fertilization – the individual zygotes (fertilized egg) who develop into fetuses, babies, and then adults. About a hundred years after Mendel's experiment we got to look in a microscope to confirm Mendel's mathematics and we continue to explore Mendelian traits in humans.

Exercise 1.2.5.1

Read: "MENDELIAN LAWS APPLY TO HUMAN BEINGS"

Here is an example using Tay-Sachs disease. The * HEXA gene on chromosome 15 makes part of an enzyme that is important for maintaining your central nervous system. If you have one or two normal alleles, you're OK, but if both your alleles have a Tay-Sachs mutation, then you'll have different neurological problems usually starting as an infant. If you are a genetic counselor and a couple comes to you planning to have kids, and they are both carriers (heterozygotes), you want to be able to tell them what is the

chance their baby will have Tay-Sachs. If we assign symbols to alleles, "t" = a Tay-Sachs mutation, and "T" = normal HEXA allele, then we can diagram the possible outcomes of fertilization.

	T	t
T	TT	Tt
t	Tt	tt

Statistically, 25% of their children will be normal (TT), 50% of their children will be carriers (Tt), and 25% of their children will be born with Tay-Sachs (tt). This principle works with most recessive diseases.




Figure **1.2.5.23**

Mendelian Traits Laboratory

Your phenotype results from the interaction of your genotype and the environment. Most traits are polygenic, meaning several genes contribute to how they are expressed. Even though your genes guide your development, the environment where you grow up influences how those genes are expressed. The combination of polygenic and environmental influence leads to an amazing variety of individuals. However, humans have a small number traits that are readily observable because they are: 1) Mendelian (determined by a single gene) so their expression is on/off, 2) tend not to be effected by the environment, 3) have high enough allele frequencies that someone in the class probably expresses them, and 4) are visible without genetic testing.

For each of the traits described below, your assignment is to: 1) record your phenotype, 2) assign letters to represent the alleles of the gene, and 3) list your possible genotypes. Your phenotype is best described in a complete sentence: "I can ____.", "I cannot ___."; "I have ____.", "I don't have ____." Include only the trait that you express. Mendelian genetics assigns letters or combinations of letters to represent alleles. For two-allele genes, often a single letter is used, the capital letter for the dominant allele and the lowercase letter for the recessive allele. For a two-allele gene, you can have three possible genotypes: heterozygous, homozygous dominant, and homozygous recessive. Depending on your phenotype, you may have more than one possible genotype that leads to the phenotype that you observed. [text adapted from Mendelian Traits Laboratory; missing source]

Example 1.2.5.1: PTC tasting

PTC, or phenylthiocarbamide, is a human-made chemical. While the majority of people find PTC to have a bitter taste, many find this substance tasteless. To discover your phenotype, chew a strip of filter paper that has been soaked in a concentrated solution of PTC. The ability to taste PTC is inherited as a dominant. There are some studies that compare tasting PTC to tasting broccoli. There are several genes and environmental influence involved but "PTC tasting is largely determined by a single gene, TAS2R8, with two common alleles, and the allele for tasting is mostly dominant over the allele for non-tasting" (McDonald 2012).

You would write:

PTC Tasting

My phenotype: Broccoli tastes bitter to me, and I can taste PTC.

Alleles: B= the allele that codes for tasting PTC; b= the allele that codes for not tasting PTC

My possible genotypes: BB or Bb

or depending on your phenotype you might write:

PTC Tasting

My phenotype: Broccoli tastes good to me, and I can't taste PTC.

Alleles: B= the allele that codes for tasting PTC; b= the allele that codes for not tasting PTC

My possible genotypes: bb only

Earwax

Earwax, or cerumen, occurs in two basic forms. The dry form is gray and brittle while the wet form is brown and sticky. The dry form is inherited as a recessive. The gene is located on *chromosome 16



Figure **1.2.5.24** - *"Mimikaki -- Japanese Ear Picks" © 2014 Dr. Timothy C. Hain*



Figure **1.2.5.25** - "Cotton swab" by Aney. (CC BY-SA 3.0)

Mid-Phalanx Hair

Look at the middle segment (phalanx) of your fingers. Note the presence or absence of hair. Complete absence of hair reflects a homozygous recessive genotype. Note that some types of work may wear the hair away.

Lactase Persistence

The ability to digest milk as an adult is a dominant trait. This is good example of biocultural evolution: a biological trait intertwined with cultural factors (how people produce food).

* more info at Wikipedia

Relative Finger Length

Lay your right hand on a piece of lined paper with the fingers perpendicular to the lines. Note the relative lengths of the second (index) and fourth (ring) fingers. There are three possible situations: the second finger is longer than the fourth, the second finger is shorter than the fourth, or the second finger is the same length as the fourth.

Individuals who are homozygous for the allele for short index finger have a shorter index finger. Individuals who are homozygous for the allele for long index fingers have longer index finger. Individuals whose index and ring fingers are equal length are heterozygous. However, this is not a true Mendelian trait because the expressions of the two alleles in the heterozygous individual show that this trait is influenced by at least one other gene on the 23rd chromosome. Heterozygous males express an index finger that is equal to or shorter than the ring finger. But heterozygous females will have an index finger that is equal or longer than the ring finger.

Exercise 1.2.5.2: IS YOUR RING FINGER LONGER THAN YOUR INDEX FINGER?



***Figure* 1.2.5.265** -"Index finger longer than ring finger" by Arnie Schoenberg, adapted from "Gaye Holud and Mehendi" by Russell eee from Wikimedia Commons (CC BY-SA 4.0)

If so, you're up to three times more likely to develop arthritis of the knee than women whose ring fingers are the same length as or shorter than their pointers. The British scientists who discovered the link speculate that it may have to do with hormones. This finger pattern is more common in women with low estrogen levels. Another clue: Men are more likely to have longer ring fingers, but for them, there's less of an arthritis link. [*Good Housekeeping* 2008:37]

* Article on cave art and relative finger length

ABO Blood type

Read Dennis O'Neil on Human Blood

Remember not to confuse the ABO blood type, a gene located on the ninth chromosome, with the Rhesus factor (Rh^+, Rh^-) a gene located on the first chromosome.




Figure* 1.2.5.27 *public domain




Figure 1.2.5.28 - *en.Wikipedia.org/wiki/ABO_blood_group_system*

Try to shift your thinking from Dominant means good/strong/prevalent, and Recessive means bad/weak/rare. Recessive just means it takes two alleles to expressive the trait. Dominant means you can have 2, or just 1 allele to express the trait. Don't conflate dominance with fitness. Dominance has to do with Mendel and inheritance, fitness is from Darwin's theory of natural selection.

Exercise 1.2.5.3

Review Dennis O'Neil's Mendelian Genetics

*Skim Mendel, Gregor. 1865. "Experiments in Plant Hybridization"

*Play around with the Online Mendelian Inheritance in Man database. Try typing in the name of a disease, or body part into the search engine, and follow the link.

Population Genetics

By the 1920s, a new definition of evolution became popular: "a change in allele frequency". We're looking at how many people have a trait in one generation and how it might change in the next generation.

To translate a Punnett square into math, just replace the sperm and eggs with variables that represent allele frequencies, and you get a pretty simple algebraic equation, that can be used to study how populations change over time. It was discovered independently in 1908 by two scientists: Hardy and Weinberg.

	p	q
p	pp	pq
q	pq	qq

A frequency is just another word for a percentage except we write it as a decimal e.g. $\bar{p}$ = .5 is the same thing as saying 50% are *p*, or that half the alleles are *p*. This implies that the other half are q, because with a two-allele gene, the percentages have to add up to 100% (the frequencies add up to 1.00), so

$$p + q = 1 \tag{1.2.5.1}$$

we can square both sides to get:

$$(p + q)^2 = p^2 + 2pq + q^2 = 1 \tag{1.2.5.2}$$

Now compare this to the Punnett square: *pp* is p^2, *qq* is q^2, and the two *pq* are 2*pq*. The p^2 and q^2represents the frequencies of the homozygotes and *2pq* represents the frequency of the heterozygote. So you can take a population, count the total alleles, and count the phenotypes of the individuals and compare them to see if anything looks weird. Mathematically, you expect the numbers to work out, and when they don't, you know some kind of evolution is occurring.




***Figure* 1.2.5.29** - *How the allele frequencies are supposed to work without evolution; the null hypothesis*

Exercise 1.2.5.4

Read the Dennis O'Neil Intro to the Hardy-Weinberg Equation.

*and here is a longer introduction to Population Genetics, if the math freaks you out, just skim to later sections.

The Modern Evolutionary Synthesis

By the 1920s, evolutionary theory had synthesized the macroevolution of Darwin, with the microevolution of Mendel and population genetics, and came up a short list of factors that cause evolution. They tested populations with the Hardy-Weinberg equation, and when they failed to get a null hypothesis, they started trying to figure out which of the four forces of evolution caused the change in allele frequency: mutation, natural selection, genetic drift, and migration. The theories have been refined in the last hundred years, the four forces of evolution is still a useful way to think of evolution.

Vocabulary

- antibodies
- antigens
- codominant
- dominant
- fixity of species
- Great Chain of Being
- Hardy-Weinberg Equation
- macroevolution
- microevolution
- population
- the Principle of Independent Assortment
- the Principle of Segregation
- Punnett Square
- recessive

This page titled 1.2.5: Evolutionary Theory (Part 3) is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.5: Evolutionary Theory (Part 3)** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.6: "Forces" of Evolution

One of the elegant things about evolutionary theory is it can describe phenomena on both the small scale and the large scale. We can use these same forces to explain microevolution – the change of an allele frequency of a population of the same species from one generation to the next; and we can use them to explain macroevolution – the change of one species into another species over long periods of time. This is similar to the way the theory of gravity can be used to describe the motion of molecular particles or large galaxies.

We use the word "force" to refer to a process that drives change, but thinking about evolution as a set of forces can be dangerous because it's easy to fall into the trap of thinking of evolution as a directional agent, pushing organisms towards an ultimate goal.

Mutation

Mutation is the prime mover, the creator of all new alleles. We'll learn more about how mutations happen in the section on cellular biology.

Exercise \(\PageIndex{1}\)

Review the last section on Darwin.

Read Dennis O'Neil's description of Natural Selection with an Emphasis on Population Genetics.

Note

- Article on carcinogenic traditional medicine, notice the kinds of mutations this plant causes



Migration

Migration (also called "gene flow") is where someone physically moves alleles from one population to another. When people move from one population to another they pack all 23 pairs of chromosomes inside the nuclei of their cells, and bring it all with them. If this changes the allele frequency of either population then it is by definition a kind of evolution. It can bring new alleles to a population that hadn't had that mutation recently, or just bring or remove a significant quantity of a certain allele to change the frequency of either population.

The individual doesn't actually have to migrate to the new population, they can just leave or pick up a few alleles. The stereotype that sailors have kids in every port, is probably better represented by today by the traveling businessman, soldier on leave, sexual tourist, or the sex trafficked. Migration is the geographic movement of alleles from one population to another. We'll come back to migration in future sections as an important point in understanding human origins and human variation. It is why there is only one species of hominid on the planet today.

Note

Article on coywolves and other hybrids which can be understood as a kind of gene flow and loosening of the species concept.

Genetic Drift

Random genetic drift, or genetic drift, is about statistics. The "drift" part has nothing to do with geographical movement (that would be migration/gene flow), what drifts is the allele frequency, like when you look at a graph of a complex system changing over time, and from a distance it looks like a straight line, but as you zoom in, the line becomes jagged, jumping up and down; the smaller your field of view, the more drastic the changes become.

A good way to understand genetic drift is to plan two trips to Viejas Casino, the first with $1,000,000,000 and the second with $100. Sit down at the cheapest table or slot machine you can find and start playing. For your first trip, your money will go up a little ($1,000,000,135) and down a little (down a little more because the House sets the odds $999,999,564) but after a few hours, you'll get bored and go home with around $1,000,000,000. Ok now go back with $100, your money will go up a little ($135) and down a little ($64), then up a little ($68), then down a little ($24) then up a little ($26) then down a little ($4) then up a little ($6) then down... whoops! no more money ($0), time to go home broke. The analogy here has to do with population size and alleles. Every generation alleles are shuffled and with a huge population statistically the allele frequency will stay pretty much the same, but with a small population, the random fluctuations are more drastic, and allele frequencies can drop to zero. If an allele frequency drops to zero, the game's over, and it's gone from the gene pool.

Here's a statistics exercise called the *Gambler's Fallacy* that also shows the difference between flipping a few coins and flipping a thousand coins.

A bottleneck is where the population shrinks to the point where lots of alleles drop out like this. The founder's effect is where a small group of people move to a new area and start a new population. The new population may grow quickly, but even though the number of people grows, if there is no other force of evolution, the allele frequencies of the new population is determined by the small number of founders who might happen to not represent the population they left. There's no way a small number of people can represent the diversity of a large population. In statistics this is known as sampling error. When comparing the old and new population, they have different allele frequencies, so by definition, evolution has occurred, and we attribute this kind of evolution to genetic drift.

Exercise \(\PageIndex{2}\)

- Read Dennis O'Neil's Small Population Size Effects
- Read Stanford's Encyclopedia of Philosophy: Random Drift

The "drift" in genetic drift comes from statistics, called *stochastic drift*. *Stochastic* just means "random", so stochastic drift is a fancy way to say that random stuff tends to happen with a small sample size.

Vocabulary For 2.3

- founder's effect
- gene flow
- genetic drift
- migration
- mutation
- stochastic drift

This page titled 1.2.6: "Forces" of Evolution is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.6: "Forces" of Evolution** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.7: Genetics, Cellular Biology, and Variation (Part 1)

By the middle of the 20th century, microscopes were good enough to actually see the "characters" that Mendel discovered, and by the end of the turn of this century, an outline of the human genome was completed, and now new alleles are discovered daily, and some even intentionally created.

We could easily do a whole class on this section, but you should focus on trying to understand the mechanisms of human variation, which we'll be dealing with for the rest of the class. What is the cause of human variation? How does it happen at the cellular level? Darwin knew that variation was essential for the functioning of natural selection, but he had no clue how it worked. Mendel's research in heredity suggested principles of simple variation, but failed to explain 90% of what you see in life. In order to understand where most of variation comes from you need a microscope, and understand basic cellular biology and genetics.

I find it helpful to get an overview of the scale of human variation. How big is the thing we're talking about? Go back and watch the 10 minute Powers of Ten movie, and focus on the cellular section.

Even the pictures can get confusing until you get a sense of how everything fits together. The same chromosomes can look different at different stages. During protein synthesis it's hanging loose, and during cell division it's all wound up like a dreadlock. Here's another graphic that goes from small to big and mentions a few stages:




***Figure* 1.2.7.2** - *Chromosomes are made of DNA.* Chromatin Structures By Richard Wheeler via Wikipedia (CC-BY-SA-3.0)

Cell size can vary drastically, but ova are big and sperm are small

A pair of sister chromatids and centromere and a pair of homologous chromosomes are about the same size; you get two chromatids because of DNA replication in preparation for cell division; you get two homologous chromosomes because of fertilization, one from Dad, and one from Mom.

All cells arise from pre-existing cells.

Cells

You are made up of about a trillion cells. Each cell has organelles inside them. A very important organelle is the nucleus. Inside the nucleus are your chromosomes. Your chromosomes direct protein synthesis, which determine how the cells interact with each other, and how you as an individual function.

Organelles

The ribosomes manufacture proteins. The mitochondria convert and store energy so the cell can use it. There are many others, but these are the most important to understanding human variation.

Nucleus

The nucleus contains the DNA, and is metaphorically like the brain of the cell.

Chromosomes

There are 23 pairs of chromosomes inside the nucleus of most human cells for most of the time. You have a pair, because as Mendel discovered, you get one from your dad and one from your mom. There are two kinds of chromosomes, autosomes and sex

chromosomes. The 22 pairs of autosomes are named for *auto* which means "self"; they're the chromosomes that stay with yourself. They are numbered from biggest to smallest. The last pair, the sex chromosomes, are named because they tend to determine the sex of an individual. The number of chromosomes doesn't make much difference. For an analogy, think of hard drive that can be formatted into different sectors.

A genome is an entire set of genes.

Different species have different number of chromosomes. Humans have 23 pairs, other apes have 24 pairs, hermit crabs have 127 pairs. Compare humans to a table of different species according to their number of chromosomes

Mitochondria

Mitochondria are the power plants of the cell, and they have their own separate DNA. The history of how mitochondria came to be is fascinating. We think they used to be independent living creatures swimming around, until 2 billion years ago, an ancestor of eukaryotic cells swallowed one, but instead of digesting it, that mitochondrion survived and began a symbiotic relationship with the host cell, reproducing inside the host's cytoplasm and being passed on to the next generation as the cell divided.

mtDNA

Mitochondrial DNA (mtDNA) can be used for genealogy and for dating the migrations of pre-historic populations.

Mitochondria are like cells within cells. Because, our cell's DNA is in the nucleus, and the mitochondrion in the cytoplasm, the mitochondrial DNA (mtDNA) was separated from the nuclear DNA of the host cell during reproduction. When sexual reproduction began, the eggs were bigger and they were almost always the source of mitochondria for the zygote. This means that you get your mtDNA from your mom, and it is inherited through matrilineal descent. Your mtDNA come:

from your great-great-great-great-great-great grandmother,

to your great-great-great-great-great grandmother,

to your great-great-great-great grandmother,

to your great-great-great grandmøther,

to your great-great grandmøther,

to your great grandmøther,

to your grandmøther,

to your møther,

to yøu.

And because of matrilineal descent, if your great-great-great grandmother had a mutation (o→ø) in the mtDNA of an egg, that mutation would be passed down to all of the descendants of that egg, and you would share the mutation with your mom, and your siblings, all those aunts and uncles, and fourth cousins on your great-great-great grandmother's side. The mutations in mtDNA accumulate and become markers to show ancestry, as well as demonstrate the evolutionary forces of migration and genetic drift. Because mitochondria are so simple, they have almost no functional variation – they either work or they don't – and without variation, natural selection doesn't happen. When you control for natural selection, the rate of neutral mutations of mtDNA becomes like a constant (one or two mutations every half-a-dozen millennia), and you can count how many different mutations two individuals have, and approximate how many generations ago they had a common ancestor. And by comparing large samples of indigenous populations, you can approximate where the mutation took place. We can correlate the genetic "when" and "where" with archaeological and historical data to test fascinating hypotheses of how humans moved across the globe.

Notes

National Geographic's Genographic Project

Skim Wikipedia: Mitochondria

Mitochondrial diseases

Cell Division

Explore Sex cells have one set of chromosomes; body cells have two.

Mitosis

Mitosis is the production of body cells for growth and healing. In mitosis, cells copy their chromosome and copy themselves, so that each daughter cell has the same number of chromosomes as the parent cell. Variations in the body cells can continue to be copied through mitosis (*e.g.* cancer), but the variations will not be passed down to the next generation.

Watch an 8 second Mitosis Movie (Try manually moving the cursor to see it slowly)

Meiosis

Meiosis is the production of gametes for sexual reproduction. In meiosis, cells copy themselves twice, but only copy their chromosomes once, so each of the viable daughter cells ends up with half the number of chromosomes as the parent cell. Individuals get the full number of chromosomes when two gametes combine during fertilization. Variations in gametes will be passed on the next generation. This is why when you get an x-ray, they put a lead blanket over your gametes – to block the radiation, and decrease birth defects.

Watch a quick Meiosis Movie

Oogenesis

Oogenesis makes ova or eggs.

Spermatogenesis

Spermatogenesis makes sperm.

Recombination

Meiosis is important because it increases variation by recombining your parents' genetic information.

Your genetic information comes in a small number of little packets, called chromosomes, and they were passed down from grandparent to parent to child. They come in pairs. One from one parent, one from the other. Meiosis splits the pairs, and shuffles them randomly so for example you might get one of your 3rd chromosomes from your paternal grandmother and one of your 4th chromosomes from your maternal grandfather.

My genome approximates my ancestry. If you know what to look for you can see which chromosomes came from my Mom and which from my Dad. My maternal grandparents were mostly descended from Britain and Ireland, and show up as light and dark blue on this chart. My paternal grandparents were mostly descended from Ashkenazim and show up as dark green. So for the first chromosome pair, the top one came from Dad and the bottom one from Mom. For chromosome pair 22, the top one is from Mom and the bottom one from Dad. For the sex chromosomes I got the Y from my Dad, and the single X from my Mom.

If I had my grandparents DNA, I could figure out whether the X chromosome that I got from my Mom, came from my maternal grandmother or my maternal grandfather. I definitely know that my Y chromosome came from my paternal grandfather. Each of my 46 chromosomes came from some great, great, great, ... grand-parent up in my family tree.

Crossing Over

Notice that that most of the chromosomes above aren't solid colors. The interspersed segments come from *crossing over*. During meiosis the homologous chromosomes are brought very close to each other. Because they are the same chromosome and have the same genes, pieces of one chromosome can "cross-over" to the one next to it.

Recombination includes the shuffling of chromosomes that you're getting from each parent, and a specific kind of recombination, called *crossing over*, where the chromosomes themselves can change, and genes can cross over from one grandparent's chromosome to another's. The discrete packages of chromosomes don't stay the same every generation, they open up and traits move from one to another.

Non-disjunction

While meiosis sorts and delivers the packets of genetic information we call chromosomes, one part of the process where variation can occur is that meiosis can deliver an **extra** packet, and we call this *non-disjunction*.

During meiosis the homologous chromosomes are brought together, and then pulled apart, but sometimes they aren't pulled apart hard enough and they stick to each other, and both chromosomes are pulled into one gamete, and the other gamete gets none. This is called non-disjunction; the junction between homologous chromosomes that is usually broken during meiosis is not. Having the wrong number of chromosomes is usually lethal, and the fertilized egg just doesn't reproduce, and you just don't get pregnant that month. Many people survive and do fine with more or fewer than 46 chromosomes. Chromosomes are numbered by size, so non-disjunction with higher numbered chromosomes tends to be less lethal. Down syndrome is also called Trisomy 21, having three of the somatic chromosome number 21.

Note

- Read the Down Syndrome Association of San Diego, Down Syndrome Facts
- New treatments for people with Down syndrome

This page titled 1.2.7: Genetics, Cellular Biology, and Variation (Part 1) is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.7: Genetics, Cellular Biology, and Variation (Part 1)** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.8: Genetics, Cellular Biology, and Variation (Part 2)

DNA

Deoxyribonucleic Acid is the chemical name for DNA we use to talk about its chemical structure. It comes in packages, called chromosomes which we can see in the microscope when it's all natted up as big dreadlocks inside the cell nucleus, when the cell is about to divide. Try not to confuse the doubling-thing in DNA and chromosomes because there are four very different kinds of doubling that happen at different times and scales, listed here from big to small:

1. First, the biggest scale of doubling is the pairing up of duplicated homologous chromosomes at the beginning of meiosis (metaphase I), but this doesn't last long. Karyotype pictures are often taken during this phase, so you see what looks like four chromosomes for each of the 23.
2. Second, for most of cell's life it has two of each of the 23 chromosomes, in what are called homologous pairs. The pair comes from fertilization, you get one from your mom and one from your dad; this is what Mendel was studying. Another pairing at this same scale is the duplicated sister chromatids after replication, and before the beginning of meiosis mentioned above.
3. Third, if you compare the upper and lower part of each of the 23 in a karyotype, you see the sister chromatids are joined in the center by a centromere, and the upper and lower parts of each side are called *arms.* The centromere is never exactly in the middle, so every chromosome has a shorter arm, p (for *petit*), and the longer arm, q. So, the X chromosome was named because after replication it has two sister chromatids joined in the middle by a centromere, and it looks like an "X", a dot with four arms.
4. Fourth, the smallest scale of doubling is at the structural level: the complementary strands of DNA wound together in a double helix. So if you have an A on one side, you have a T on the other; if you have a G on one side, you have a C on the other. These complementary strands are like the two sides of a zipper that come apart for protein synthesis and replication.

Replication

DNA likes to make copies of itself.

Protein Synthesis

There are two main stages in protein synthesis: transcription and translation. It starts in the nucleus with transcription, where enzymes take the message from the DNA and transcribes it into messenger RNA (mRNA). The mRNA takes the message out of the nucleus into the cytoplasm to the ribosome.

Translation occurs when the ribosome reads the message and puts the right amino acids in the right order. The ribosome needs help to gather and place the amino acids and uses transfer RNA (tRNA). The tRNA has an amazing property that a combination of three bases (codon) will stick to a particular amino acid, and as the ribosome reads the message from the mRNA it use the tRNA to transfer the correct amino acid in the correct order to make a protein.

After protein synthesis, the protein can leave the cell and do whatever it needs to do to keep the individual alive.

Check out this groovy example of hippy science from Stanford University in 1971: protein synthesis reenactment (the trip kicks-in at around 3:13)

A gene is a discrete sequence of DNA nucleotides.

Often one gene makes one protein, but not always. The expression of genes is influence by the enivorment. Some proteins require many gene. Some genes produce more than one protein.

Polygenic Traits

Polygenic traits are determined by a combination of many genes. For example, the hemoglobin protein takes 4 genes (6 for fetuses) and we'll look at a tiny change in one of the four genes that causes Sickle Cell anemia.

Pleiotropic Genes

Pleiotropy is where a single gene can effect multiple traits.

Locus > Gene > Allele

The words *locus*, *gene*, and *allele* can be very confusing, especially since people (including myself) get lazy and use them interchangeably, but technically they are distinct and you may as well get in the good habit of using them correctly. *Locus* has the same root as "location", and it refers to a place on a chromosome where a string of bases will fit. If that string of bases codes for a protein (a functional product), then that spot is also a *gene*. But in many cases, there can variations as to what bases fit in that same spot, and each of those possible variations is called a different *allele*, and often codes for different proteins and change the organism. In summary, if a *locus* does something it's a *gene*, all the different versions of the *gene* are *alleles*.

Introns and Exons

One of the ways to increase variation is have genes modify other genes, and there are many stages of protein synthesis where that can happen, one of them being during transcription where parts of the code are cut out.

Notes

- Read S-cool's description of introns and exons.
- The RNA message is sometimes edited.
- Some DNA does not encode proteins.
- Article "Exon Skipping: Borrowing from Nature to Treat Rare Genetic Diseases"

Codons

DNA words are three letters long.

Cells and the Source of Variation

The origin of all variation is mutation. Mutations can occur at many different scales. The smallest is the Single Nucleotide Polymorphism (SNP, called a "snip") that changes a single base.

If we think of meiosis as sorting and delivering genetic information to our kids in packets that we call chromosomes, there are many parts of the process where variation can occur, including: 1) meiosis shuffles the packets of informations in recombination as Mendel proved in his Principle of Independent Assortment, 2) meiosis moves traits from packet to packet in crossing over, and 3) meiosis can deliver an extra packet in non-disjunction.

Another possibility is that two different packages can stick together, like the chromosomal shifting that fused the greater ape chromosome 2a and 2b, into *Homo sapiens* chromosome 2. The same genes are there, just on one chromosome instead of two. These big changes are important for macroevolution (speciation), and makes it impossible for humans to reproduce with other apes (apologies to Jerry Springer and the *Weekly World News*).

The use of the word *recombination* can be confusing because most of the recombination that makes you different from your siblings is Mendel's principle of independent assortment, the shuffling of chromosomes, but there is also a type of recombination called *crossing over* that is a very different specific process on a smaller scale where genes jump between homologous chromosomes. Also, recombination is technically not a separate force of evolution, it's an aspect of all of them.

Because this is an introduction we have skipped many other processes that influence variation, such as genes that don't produce proteins directly, but just effect other genes that do. But, most genes make proteins, so lets get the basics down first.

for another review read Dennis O'Neil's description of mutation and recombination

Notes

- Single Nucleotide Polymorphisms (SNiPs)
- Some viruses store genetic information in RNA.
- Genetic disorders




Figure **1.2.8.5** - *La Plasmogenia (in Spanish) Alfonso Luis Herrera introduced Darwin to Mexico*




Figure **1.2.8.6** - *Leonardo DaVinci notebook*

Imagination Activities

- **Extract your own DNA**

Follow the instructions on this video, and take a selfie with your own DNA.
For each cell, how many membranes does the soap have to break down to release the DNA?

Imagination questions

- **Your tax dollars at work**

The Human Genome Project is federally funded. Do you think it's worth it?

- **Exploration**

Take a few minutes to explore the human genome with MapViewer at the National Center for Biotechnology Information. How is exploring our genome different from exploring unknown jungles, or the bottom of the sea, or outer space?

- **Life**

Is a virus alive? A DNA molecule?

What of humans is mechanistic? Where does free-will exist?

- **The whole is greater than the sum of its parts**

What is a person? A bunch of cells. Each individual is made up of about a trillion cells (1,000,000,000,000). Most of those cells have 46 chromosomes. Each chromosome has about a million and a half base pairs, and the human karyotype has about 3.2 billion base pairs total. The information in the 3,200,000,000 base pairs makes sure the 1,000,000,000,000 cells all work together. 3.2 billion seems like a lot of base pairs, but if you take computer memory as a metaphor it's not that much. If you think of a base pair like something between a bit and a byte, then all the genetic information fits on a 3.2 gigabyte (3.2 GB) thumbdrive. It's like the basic install of a video editing program. Not something you could attach to an email, but it wouldn't take that long to download. So what makes people so complicated, if their code is so small?

- **Multiple identities**

Here's an introduction to chimerism, the idea most of us are conjoined twins to a small extent. Does this change how you think of yourself?

Genetics and Ethics

Remember that the anthropological imagination avoids scientism: putting science on a pedestal protected from all criticism, and valuing science and scientific knowledge above people. Anthropology is the science **of** humans, so there is no way to avoid humanistic questions. The field of genetics has grown in a political context where world economic systems have loosened many ethical guidelines. It's like the wild, wild, west.

Identity and Ownership

If you think the government reading your emails is bad, think of having scientists steal your genetic code without your consent and owning you in your afterlife.

Exercise 1.2.8.1

- Read an update on Henrietta Lacks' genome
- New HBO movie

One of the great things about Obamacare is that it headed-off a growing problem of insurance companies using genetic information as a way to screen out members with a propensity for expensive medical conditions.

Our reliance on genetics in forensic anthropology (using anthropology for legal, usually police, situations) can lead to problems such as this case which seems straight out of the Jerry Springer show: * Man Fails Paternity Test Because Unborn Twin Is The Biological Father Of His Son

Stem Cells

Stem cells are undifferentiated, that means they divide and grow into many different kinds of cells. When you think of yourself as a trillion cells, once upon a time, you used to be one cell: a tiny, cute, little zygote (what happened!??!!). Stem cells are left over from that transition from one cell to billions of cells. They are great for medicine because if you are missing cells in your body, you tell the stem cells to become them.

The controversy was much worse a decade ago when the major source of stem cells was aborted fetuses, but today, they can be harvested from your own baby teeth, your blood, even from your left over liposuction.

Notes

- Article on stem cell therapy for Crohn's disease
- Article on stem cell therapy for diabetes, protein synthesis is how a cell produces insulin

Cloning

***Figure* 1.2.8.7** - *stuffistumbledupon.com*

To me, human cloning is no big deal. We're already dealing with the ethical issues that cloning raises.

Genetically identical humans? It's what we get with identical (monozygotic) twins, when the zygote divides and then separates into two embryos who become two individuals. They may be identical genetically, but variations in how they interact with their environment will make them physically distinct, and most importantly, their different cultural experiences will make them different people.

Exploiting people to harvest their organs? The demand for kidneys has led to transplant tourism and black markets in places like India and the Philippines, and a legal kidney market exists in Iran.The 1% are already playing God and cannibalizing the bodies of the 99%.

(Full disclosure: my tolerance for cloning may be biased because I am currently raising several clones of my own of various ages on a secret organic ranch in southern Utah which I plan to use to for parts as I get older. Here's a home video of my favorite "Little Arnie", AS0983-2342, getting in shape for his transplant surgery.)

GMOs

I think recombinant DNA, or genetically modified organisms (GMO) are a bit scarier... Will it kill you to eat them? No! Are they poisonous? No! Should you try avoid them? Yeah, it's probably a good idea to eat as much organic produce as as you can afford.

*Read this article that examines a historic period in agronomy known as the "Green Revolution"and how we might be able to use lessons learned from this technocratic disaster, to avoid potential disaster from the current GMO revolution.

We look at hackers today as some kind noble outlaw, or social bandit, but we need to keep asking ourselves, what are our limits, and how far we are willing to go to hack ourselves and the code that makes us who we are.

***Figure* 1.2.8.8** - *Frank Miller and Geof Darrow 1993. Hard Boiled*

The mad scientist trope in biology has a long history, from the Jewish golems, the alchemists of the Middle Ages, Mary Percy Shelly's *Frankenstein; or, the Modern Prometheus*, or some of my recent favorites, the Larry Fessenden 1991 movie, *The Telling*, and Margaret Atwood's 2003 novel, *Oryx and Crake*. But, another worry with the rapidly expanding technology of genetic modification is the chaos theory idea from *Jurassic Park*, or what I like to call the Homer Simpson effect – shit happens – genes can move around out of our control, and I think this next section is even scarier:

Lateral Gene Transfer

Lateral gene transfer (LGT), also called horizontal gene transfer (HGT), is distinguished from the kind of up-and-down verticalness of heredity that we represent with a family tree. It's important that you understand the classic Standard Evolutionary Theory of Darwin, Mendel, and the Modern Synthesis, because that's how things work 99.999999% percent of the time, but there are rare exceptions. Some evolutionary scientists have advocated including new research into an Extended Evolutionary Synthesis, others say that the old paradigm works fine.

Some consider LGT to be a kind of gene flow, but classic gene flow happens within the same species through migration. LGT is about genes that can be moved from one nucleus to another in ways other than meiosis. Humans have learned to do this intentionally as shown in the previous section, but it can also happen naturally.

I think the scariest thing about all the genetic engineering going on today is the combination of GMOs and LGT. Nature has dealt with LGT for billions of years and you can expect that we've evolved to deal with whatever genes are floating around in the environment. When you create a new gene, you can test how it will influence a certain organism in the laboratory, but when you release the gene into the environment, there is no way to test every possible combination of that gene inserted into every other species. So for example, you make a gene that allows leaf cells to produces their own pesticide and you put it in corn. Fine, the corn is great, no bugs, no need to spray pesticides. But, what happens if that gene moves to bacteria that live in your stomach. Whoops! Now you don't have to bother drinking pesticide, because you're producing it in your stomach already.

On the bright side, LGT does make a great back-story for horror movies.

Notes

- You are what you eat; a good introduction to LGT
- A recent editorial about "GMOs, Herbicides, and Public Health"
- An article on genomic infection.
- Scare tactics: Genetic Roulette movie trailer
- Good rebuttal to Genetic Roulette
- Article on Monsanto suing farmers for saving seeds
- Article on human embryo editing
- Website for the FDA's Cellular, Tissue, and Gene Therapies Advisory Committee
- An article on human babies with a two mothers and a father (a mitochondrial donor) or just two mothers, which the FDA is considering

Imagination Questions

- Designer Babies
  - 1) My niece was screened for chromosomal abnormalities *in utero*, and I've been avoiding asking my sister if she would have gotten an abortion if the fetus tested positive for Down syndrome. Would you abort your child if you found out it was going to be born with a disease? Would you like time to prepare? Would you even want to know?
  - 2) an article on a sperm donor clinic who filters out "the donor matches with an elevated risk of rare recessive paediatric conditions."
- Do you think births should be natural? Do you have limits on how far people should go to provide for the future of their children?

Imagination Actions

GMOs

Using what you've learned, find a recent campaign calling for the labeling of genetically modified foods (e.g. Just Label It), and write a letter to the appropriate policy maker, arguing for, or against, the legislation/action.

Vocabulary

- allele
- autosome
- cell
- centromere
- chromatid
- chromosome
- codon
- crossing-over
- double helix
- gamete
- gene
- individual
- karyotype
- locus
- meiosis
- mitochondria
- mitosis
- nucleotide
- nucleus
- ovum
- population
- somatic cell
- species
- sperm
- zygote

This page titled 1.2.8: Genetics, Cellular Biology, and Variation (Part 2) is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.8: Genetics, Cellular Biology, and Variation (Part 2)** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria

Exercise 1.2.9.1

Read the section called "Selection Against Both Homozygotes" from Dennis O'Neil Natural Selection

Anthropology is holistic because it covers many branches of knowledge. To understand sickle cell anemia we need look at the smallest change in a base pair, and at the global migration of alleles. We need to look two thousand years back in time to a transition from hunter-gatherers to horticulturalists, to the racial discrimination of the 20th Century. We apply the knowledge to the most deadly disease on the planet, and to mixology.

Mutation starts the process. In the sperm or egg on the 11th chromosome, at the 17th nucleotide of the gene for the beta chain of hemoglobin, there is a point mutation where an A is replaced by a T, which changes the codon GAG (for glutamic acid) to GTG (which encodes valine). Thus the 6th amino acid in the chain becomes valine instead of glutamic acid. The beautiful architecture of the hemoglobin molecule collapses, as if you took the capstone off of an arch, and the red blood cell takes on a sickle shape. The sickled cells get caught in blood vessels and don't carry oxygen as well.




***Figure* 1.2.9.1** National Institute of Health Sickle Cell Disease (public domain)

Exercise 1.2.9.2

Skim the medical literature on sickle cell anemia and click on the location to get a sense of what a gene is

This new allele is called the S allele, and as a Mendelian trait, you get one from each parent.

AA=normal hemoglobin
AS=sickle cell trait, sickle cell carrier
SS=sickle cell anemia

The allele frequency of any Single Nucleotide Polymorphism (SNP) is about 1 in 100,000, so you might expect the allele frequency of the S allele to stay at that rate: S=0.00001 But in some places the frequency of the S allele gets as high as 1 in 5, or S=0.2 When population geneticists see changes in allele frequencies they know that evolution is occurring, and the connection to malaria makes it clear that this is a case of natural selection.

The sickled cells are bad for blood flow and carrying oxygen, but they are good because they protect you from the parasites which cause malaria.

Malaria is an infectious disease caused by the parasite, *Plasmodium falciparum*, and is carried by mosquitoes, and easily spread. One little parasite gets into one of your red blood cells, reproduces 30,000 times, pops the cell, and then go on to infect thousands of other cells, which then go on to infect thousands more, until you are really sick.




Figure 1.2.9.3




Figure 1.2.9.4

But, when the parasite infects a sickle-shaped cell, there is less room to reproduce, and it doesn't pop the cell, so it's not spread. Sickle Cell Anemia is bad, but it gives you immunity to Malaria.

Natural selection acts on the mutation to change its allele frequency.

Fetuses produce a special kind of hemoglobin (HbF) that helps pull mom's oxygen across the placenta. Once born, the fetuses normally stop producing HbF, but some adults inherit a gene that tells their body to persist in producing fetal hemoglobin all their lives. The correlation between people with blood diseases like Sickle Cell anemia and hereditary persistence of fetal hemoglobin suggests that natural selection may have selected for the persistence of fetal blood to mitigate the effects of sickle cell anemia. Malaria makes it advantageous to have Sickle Cell, and then Sickle Cell makes it advantageous to have hereditary persistence of fetal hemoglobin.

Sickle Cell Anemia is an example of biocultural evolution because human cultural activity was the cause of people's genetic change. People in West Africa developed a new subsistence practice that produced more food by clearing land and planting crops. But it also created open spaces for mosquitoes to breed, and higher population densities that made it easier for malaria to spread. As malaria became endemic it became more advantageous to have the S allele.

Other cultural factors range from racism to mixology. Because of racism, and the misconception of Sickle-Cell Anemia as a racial disease, the US military initially prohibited African-Americans from flying planes fearing that all African-Americans would suffer sickling events at high altitudes. British Colonialists lacking malaria resistance turned to the bark of a tree from South America called quinine, and they preferred to drink this bitter tonic with gin for good measure. Unfortunately, because of natural selection, most malaria parasites are now resistant to quinine, and drinking gin & tonics in the tropics is more likely to cause dehydration than prevent malaria.

The holistic approach of anthropology allows us to understand sickle cell anemia through a wide range of disciplines including archaeological research on sites in West Africa, the genetics of humans, plasmodium parasites, and mosquitoes, racism in the US, and even mixology.




***Figure* 1.2.9.5** - *Bill Gates notes* © Copyright The Gates Notes, LLC

Notes

- History of the gin and tonic
- Exploring the use of fetal hemoglobin as a treatment for sickle cell anemia
- A 360º VR 3D video [you need VR goggles] on malaria, and how to prevent it:



Imagination Question

- Now that you understand natural selection, why are pesticides, malaria killing drugs, and a new insect repellent not as good in the long term as a new vaccine for malaria and a campaign to provide mosquito nets?
- What could go wrong with a genetically modified mosquito?



- People with Sickle Cell Trait may suffer during heavy exertion, and have been discriminated against because of their condition, and there is debate in sports and different branches of the US military as to whether to keep their condition confidential to protect them from discrimination, or publicize their condition to * protect them from the health risks.

Imagination Actions

- Read through the Nothing but Nets website, or Against Malaria, and take action to support providing mosquito nets to people from areas with endemic malaria.

Vocabulary

- malaria
- sickle cell anemia
- *Plasmodium falciparum*

This page titled 1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

1.3: What is Biological Anthropology?

Anthropology is the scientific study of humankind. The main purpose of anthropology is for us to understand human biological and cultural diversity, and the origins of humans. So, the humans that anthropologists study can be modern or ancient, and can span across the globe. Anthropologists are unique in that they study the entirety of human existence over time and space.



***Figure* 1.3.1** - *Pride parade, Seattle, WA, June 26, 2011*

Basic tenets of anthropology:

1. **Holism**: Holism means that a part of something can only truly be understood if examined within relation to the whole of it. For anthropologists, this means that they try to understand humankind through the interrelationships of all aspects of human existence -- for example, human biology has to be examined within the context of human cultures and vice versa. In addition, all of this must be examined within the context of the environment and historical processes. In an effort to be holistic, anthropology is often an interdisciplinary field that crosses over into other fields such as history, geology, and ecology.
2. **Relativism**: Relativism means that judgments, truths, or moral values have no absolutes, and can only be understood relative to the situation or individuals involved. For anthropologists, this means that they accept that all cultures are of equal value and must be studied from a neutral point of view. A good anthropologist must disregard their own beliefs, morals, and judgments when examining another culture. They must, instead, examine each culture within the context of its own beliefs.
3. **Universalism**: Universalism means that whatever the theoretical principle is, it's equally applicable to all. For anthropologists, universalism means that we believe all humans are equal -- in intelligence, complexity, etc.This is in contrast to *ethnocentrism*, which is the belief that some peoples are more important or culturally/biologically better than other peoples.
4. **Culture**: All humans have culture. Culture is the set of learned behaviors and knowledge that belong to a certain set of people. This is different from genetically hardwired behaviors (such as reflexes) in that they aren't biologically inherited. The most important thing to remember is that culture is learned.

Subfields in Anthropology

There are four subfields in anthropology: cultural anthropology, biological (or physical) anthropology, archaeology, and linguistic anthropology.

- **Sociocultural Anthropology**

Sociocultural anthropologists examine social patterns and practices across cultures, with a special interest in how people live in particular places and how they organize, govern, and create meaning. A hallmark of sociocultural anthropology is its concern with similarities and differences, both within and among societies, and its attention to race, sexuality, class, gender, and nationality. Research in sociocultural anthropology is distinguished by its emphasis on participant observation, which involves placing oneself in the research context for extended periods of time to gain a first-hand sense of how local knowledge is put to work in grappling with practical problems of everyday life and with basic philosophical problems of knowledge, truth, power, and justice. Topics of concern to sociocultural anthropologists include such areas as health, work, ecology and environment, education, agriculture and development, and social change.

- **Biological (or Physical) Anthropology**

Biological anthropologists seek to understand how humans adapt to diverse environments, how biological and cultural processes work together to shape growth, development and behavior, and what causes disease and early death. In addition, they are interested in human biological origins, evolution and variation. They give primary attention to investigating questions having to do with evolutionary theory, our place in nature, adaptation and human biological variation. To understand these processes, biological anthropologists study other primates (primatology), the fossil record (paleoanthropology), prehistoric people (bioarchaeology), and the biology (e.g., health, cognition, hormones, growth and development) and genetics of living populations.

- **Archaeology**

Archaeologists study past peoples and cultures, from the deepest prehistory to the recent past, through the analysis of material remains, ranging from artifacts and evidence of past environments to architecture and landscapes. Material evidence, such as pottery, stone tools, animal bone, and remains of structures, is examined within the context of theoretical paradigms, to address such topics as the formation of social groupings, ideologies, subsistence patterns, and interaction with the environment. Like other areas of anthropology, archaeology is a comparative discipline; it assumes basic human continuities over time and place, but also recognizes that every society is the product of its own particular history and that within every society there are commonalities as well as variation.

- **Linguistic Anthropology**

Linguistic anthropology is the comparative study of ways in which language reflects and influences social life. It explores the many ways in which language practices define patterns of communication, formulate categories of social identity and group membership, organize large-scale cultural beliefs and ideologies, and, in conjunction with other forms of meaning-making, equip people with common cultural representations of their natural and social worlds. Linguistic anthropology shares with anthropology in general a concern to understand power, inequality, and social change, particularly as these are constructed and represented through language and discourse.

(From: American Anthropological Association. 2015. What is Anthropology? <http://www.aaanet.org/about/WhatisAnthropology.cfm>. Accessed 2015 May 7.)

While the sheer scope of anthropological study requires people to choose a subfield and specialization within that subfield many anthropologists integrate multiple subfields in their work. It is common to find biological anthropologists working with cultural anthropologists to examine things such as disease patterns and archaeologists to look at migration patterns in the past.

What is Biological Anthropology?



***Figure* 1.3.2:** *Heather Hecht Edgar (University of New Mexico) conducting research at the Czech National Museum, 2005*

Biological anthropology is the study of the biology of humans and their nearest biological relatives. "Biological relatives" include both humankind's ancestors and our nearest living relatives, the non-human primates. Biological anthropologists fall under 3 major categories:

1. Human biology (human biological diversity, genetics, adaptations to environmental stressors, etc.)
2. Primatology (non-human primate biology, evolution, behavior, ecology, etc.)
3. Paleoanthropology (human origins and human evolution)

These are the topics we'll be studying in this course. Each unit follows each of the 3 major categories found in biological anthropology.

- Unit 1: In this unit, you'll learn the basic toolkit you'll need (evolutionary theory) in order to understand the rest of the course. This unit also covers human biology.
- Unit 2: This unit covers primatology.
- Unit 3: This unit covers paleoanthropology.

This page titled 1.3: What is Biological Anthropology? is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.1: What is Biological Anthropology?** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

CHAPTER OVERVIEW

2: Mechanisms and Evolutionary Thought

2.1: Evolutionary Theory

2.1.1: What is Biological Anthropology?

2.1.2: Scientific Method

2.1.3: History of Evolutionary Thought

2.1.4: Darwinian Evolution

2.1.5: Cell Biology and Basic Genetics

2.1.6: Mendelian Genetics

2.1.7: Modern Synthesis

2.1.8: Modern Human Variation

2.2: Taxonomy

2.3: Mammals

2.4: Human Taxonomy

2.5: Osteology

2.6: Macroevolution

2.7: Species vs. Paleospecies

2.8: Interspecific vs. Intraspecific Variation

2.9: Adaptive Radiation

2.10: Analogy vs. Homology

2: Mechanisms and Evolutionary Thought is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

2.1: Evolutionary Theory

Learning Objectives

At the end of this unit, students will be able to:

- answer the question, What is Biological Anthropology?
- explain the Scientific Method
- describe The History of Evolutionary Thought, including the roles of Charles Darwin & Darwinian Evolution
- relate Cell Biology & Genetics to molecular evolution
- apply Genetics & Evolution: Mendelian Genetics
- show how Genetics & Evolution: The Modern Synthesis works in population genetics
- discuss how population genetics applies to Human Variation & Race

In this unit students will learn about basic genetics, modern evolutionary theory, modern human variation, and the history of evolutionary thought.

2.1.1: What is Biological Anthropology?
2.1.2: Scientific Method
2.1.3: History of Evolutionary Thought
2.1.4: Darwinian Evolution
2.1.5: Cell Biology and Basic Genetics
2.1.6: Mendelian Genetics
2.1.7: Modern Synthesis
2.1.8: Modern Human Variation

This page titled 2.1: Evolutionary Theory is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

2.1.1: What is Biological Anthropology?

Anthropology is the scientific study of humankind. The main purpose of anthropology is for us to understand human biological and cultural diversity, and the origins of humans. So, the humans that anthropologists study can be modern or ancient, and can span across the globe. Anthropologists are unique in that they study the entirety of human existence over time and space.



***Figure* 2.1.1.1** - *Pride parade, Seattle, WA, June 26, 2011*

Basic tenets of anthropology:

1. **Holism**: Holism means that a part of something can only truly be understood if examined within relation to the whole of it. For anthropologists, this means that they try to understand humankind through the interrelationships of all aspects of human existence -- for example, human biology has to be examined within the context of human cultures and vice versa. In addition, all of this must be examined within the context of the environment and historical processes. In an effort to be holistic, anthropology is often an interdisciplinary field that crosses over into other fields such as history, geology, and ecology.
2. **Relativism**: Relativism means that judgments, truths, or moral values have no absolutes, and can only be understood relative to the situation or individuals involved. For anthropologists, this means that they accept that all cultures are of equal value and must be studied from a neutral point of view. A good anthropologist must disregard their own beliefs, morals, and judgments when examining another culture. They must, instead, examine each culture within the context of its own beliefs.
3. **Universalism**: Universalism means that whatever the theoretical principle is, it's equally applicable to all. For anthropologists, universalism means that we believe all humans are equal -- in intelligence, complexity, etc.This is in contrast to *ethnocentrism*, which is the belief that some peoples are more important or culturally/biologically better than other peoples.
4. **Culture**: All humans have culture. Culture is the set of learned behaviors and knowledge that belong to a certain set of people. This is different from genetically hardwired behaviors (such as reflexes) in that they aren't biologically inherited. The most important thing to remember is that culture is learned.

Subfields in Anthropology

There are four subfields in anthropology: cultural anthropology, biological (or physical) anthropology, archaeology, and linguistic anthropology.

- **Sociocultural Anthropology**

Sociocultural anthropologists examine social patterns and practices across cultures, with a special interest in how people live in particular places and how they organize, govern, and create meaning. A hallmark of sociocultural anthropology is its concern with similarities and differences, both within and among societies, and its attention to race, sexuality, class, gender, and nationality. Research in sociocultural anthropology is distinguished by its emphasis on participant observation, which involves placing oneself in the research context for extended periods of time to gain a first-hand sense of how local knowledge is put to work in grappling with practical problems of everyday life and with basic philosophical problems of knowledge, truth, power, and justice. Topics of concern to sociocultural anthropologists include such areas as health, work, ecology and environment, education, agriculture and development, and social change.

- **Biological (or Physical) Anthropology**

Biological anthropologists seek to understand how humans adapt to diverse environments, how biological and cultural processes work together to shape growth, development and behavior, and what causes disease and early death. In addition, they are interested in human biological origins, evolution and variation. They give primary attention to investigating questions having to do with evolutionary theory, our place in nature, adaptation and human biological variation. To understand these processes, biological anthropologists study other primates (primatology), the fossil record (paleoanthropology), prehistoric people (bioarchaeology), and the biology (e.g., health, cognition, hormones, growth and development) and genetics of living populations.

- **Archaeology**

Archaeologists study past peoples and cultures, from the deepest prehistory to the recent past, through the analysis of material remains, ranging from artifacts and evidence of past environments to architecture and landscapes. Material evidence, such as pottery, stone tools, animal bone, and remains of structures, is examined within the context of theoretical paradigms, to address such topics as the formation of social groupings, ideologies, subsistence patterns, and interaction with the environment. Like other areas of anthropology, archaeology is a comparative discipline; it assumes basic human continuities over time and place, but also recognizes that every society is the product of its own particular history and that within every society there are commonalities as well as variation.

- **Linguistic Anthropology**

Linguistic anthropology is the comparative study of ways in which language reflects and influences social life. It explores the many ways in which language practices define patterns of communication, formulate categories of social identity and group membership, organize large-scale cultural beliefs and ideologies, and, in conjunction with other forms of meaning-making, equip people with common cultural representations of their natural and social worlds. Linguistic anthropology shares with anthropology in general a concern to understand power, inequality, and social change, particularly as these are constructed and represented through language and discourse.

(From: American Anthropological Association. 2015. What is Anthropology? <http://www.aaanet.org/about/WhatisAnthropology.cfm>. Accessed 2015 May 7.)

While the sheer scope of anthropological study requires people to choose a subfield and specialization within that subfield many anthropologists integrate multiple subfields in their work. It is common to find biological anthropologists working with cultural anthropologists to examine things such as disease patterns and archaeologists to look at migration patterns in the past.

What is Biological Anthropology?



Figure* 2.1.1.2*: *Heather Hecht Edgar (University of New Mexico) conducting research at the Czech National Museum, 2005*

Biological anthropology is the study of the biology of humans and their nearest biological relatives. "Biological relatives" include both humankind's ancestors and our nearest living relatives, the non-human primates. Biological anthropologists fall under 3 major categories:

1. Human biology (human biological diversity, genetics, adaptations to environmental stressors, etc.)
2. Primatology (non-human primate biology, evolution, behavior, ecology, etc.)
3. Paleoanthropology (human origins and human evolution)

These are the topics we'll be studying in this course. Each unit follows each of the 3 major categories found in biological anthropology.

- Unit 1: In this unit, you'll learn the basic toolkit you'll need (evolutionary theory) in order to understand the rest of the course. This unit also covers human biology.
- Unit 2: This unit covers primatology.
- Unit 3: This unit covers paleoanthropology.

This page titled 2.1.1: What is Biological Anthropology? is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.1: What is Biological Anthropology?** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.2: Scientific Method

Science is a way to gain knowledge about natural phenomena using empirical observation and testing (Jurmain, et al 2013). While there are different protocols used in science, it is performed using a set of rules called **The Scientific Method** that guide scientific practice (Fancher 2000). The method stresses the need to develop a testable hypothesis, the use of objectivity and rationality, and the circularity of scientific research. This does not mean that science is infallible, but in using the scientific method, particularly with the necessity to have testable hypotheses and tests that are replicable by other researchers, rigorous conclusions are reached.

Why do students of biological anthropology need to understand the scientifc method? Because anthropology is a science.

Steps of the Scientific Method




***Figure* 2.1.2.1** - *Diagram of the scientific method*

1. Define the problem

    This is based on observation -- either something you've observed from nature or from something that's already been written. How many times have you seen something or read about something and thought up a question about it? If you have, then you've done the first step of the Scientific Method!

2. Develop Hypothesis

    Propose an explanation for the observed phenomenon. For a hypothesis to be a GOOD hypothesis, it must be testable. In other words, you must be ABLE to test it to see if it's supported or not supported. A hypothesis is still good, even if it's unsupported, as long as it's testable. There are lots of questions out there that are untestable, such as, "Somewhere on the planet, a pink elephant is dancing the can-can in a tutu." Because it's impossible to explore every inch of the world looking for our dancing elephant, this would not be feasibly testable. Along the same lines, a testable hypothesis would be, "Student group A, who have read the material on The Scientific Method, will have a higher average score on their quiz than will Student group B, who have not read the material on The Scientific Method."

3. Collect Data

    The experiment should be specifically designed to test the hypothesis. The experiment will provide data as to whether the hypothesis is supported or not. Experiments must be replicable by other researchers. For our example above with ever-so-fortunate quiz-takers, you'd have Group A read the material (Group B would not) and both groups would take the quiz. This

experimental design is replicable by other researchers because they too could find two groups of students, and follow the same protocol (one group reads the material, one not, and both take the quiz).

4. Organize/analyze data:

   Once the experiment is completed, you must organize the data and analyze the results. For our example, you would grade the quizzes and calculate the average score per group.

5. Conclusion

   You develop a statement that sums up what the data (collected during the experimental phase) says about the hypothesis. For our example, our concluding statement would be something like, "Group A, who read material on The Scientific Method, performed better on a Scientific Method quiz than did Group B, who did not read the material." If your hypothesis is supported, then you would move on to the next step in the process. If your hypothesis was not supported you would rework your hypothesis and start the process over.

6. Share the knowledge

   It is important to share the results of your work even (and maybe even *especially*) if your hypothesis was not supported. This hasn't been a step in the "old school version" of The Scientific Method, but remember -- information is only good if it's communicated to others!

Things to keep in mind:

- Science does not "prove" anything. Hypotheses are falsified/supported or not falsified/supported. For a hypothesis to be accepted as a theory, which is a generally accepted explanation of specific phenomena, it undergoes rigorous testing (Larsen 2008: 16). It can take decades for a hypothesis to become a theory.
- While we discuss various topics in biological anthropology that each piece of information was once a hypothesis that was tested and supported by the data. Contradictory supported-hypotheses are possible; some refer to this as **equifinality**. This just means that the data collected supports two (or more) hypotheses -- in most cases, there is not sufficient data available to support one more than the other (especially when we get to the information within human evolution). Hopefully, in these cases, one day we'll have enough evidence to overwhelmingly support one particular hypothesis, but at the present that's not possible. So, for the moment we have to open our minds to accept two potential conclusions.

The Scientific Method in Action!!! Watch the following clip and identify the steps of The Scientific Method. Was their hypothesis testable? Was their experiment replicable? What was the conclusion of their "scientific endeavor"? Clip: The Scientific Method in Action

References

1. Fancher LJ. 2000. The Great “SM.” < www.cod.edu/people/faculty/fancher/scimeth.htm>. Accessed May 7, 2015.
2. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
3. Kaziek CJ, Pearson D. 2014. Ask a biologist: using the scientific method to solve mysteries. < http://askabiologist.asu.edu/explore/scientific-method>. Accessed May 7, 2015.
4. Larsen CS. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.; 2008. 430 p.

This page titled 2.1.2: Scientific Method is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.2: Scientific Method** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.3: History of Evolutionary Thought

Contrary to popular opinion, evolutionary thought does not start with Charles Darwin. The history of evolutionary thought is a fascinating story spanning hundreds of years of both speculation and scientific discovery. Using the links below, explore the story from early studies of earth sciences, through 20th-century discoveries that enrich our understanding of life on earth. As you read each person's contribution, pay particular attention as to what exactly is the key contribution for evolutionary theory. Sometimes this will be clear, while other times it might be a bit more murky.

When navigating US Berkeley's *Understanding Evolution: The History of Evolutionary Thought* web-site, make sure you click on "next page" when you see it. All of the links below go to different pages on the *Understanding Evolution* web site. They are provided here so that you do not have to continually go the first page should you decide to read these pages in different sittings.

- **Pre-1800s**
  - Comparative Anatomy: Vesalius
  - Observation: Harvey & Palin
  - Fossils & Paleontology: Steno
  - Nested Hierarchies: Linneaus
  - Ancient Life: Leclerc & Buffon
  - Human Ecology: Malthus
- **1800-1900**
  - Extinction: Cuvier
  - Evolution Happens: Lamarck
  - Developmental Similarities: von Baer
  - Biostratigraphy: Smith
  - Uniformitarianism: Lyell
  - Discrete Genes: Mendel
  - Early Evolution and Development: Haeckel
  - Biogeography: Wallace & Wegener
  - Fossil Hominids: Huxley & Dubois
  - Chromosomes and Mutations: Morgan
- **1900-Present**
  - Random Mutation: Fisher, Haldane & Wright
  - The Modern Synthesis: Dobzhansky
  - Speciation: Mayr
  - DNA: Crick & Watson
  - Radiometric Dating: Clair Patterson
  - Endosymbiosis: Lynn Margulis
  - Evolution and Development for the 21st Century: Gould
  - Genetic Similarities: Wilson, Sarich, Sibley & Ahlquist

You may notice that Darwin is absent from the list above although you may have read about him if you followed the links on the *Understanding Evolution* website. That is because there is a special page devoted to Darwin. Even if you read the *Understanding Evolution* page you need to read the page included on this textmap.

This page titled 2.1.3: History of Evolutionary Thought is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.3: History of Evolutionary Thought** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.4: Darwinian Evolution

Who was Charles Darwin?

Ernst Mayr claims, and many would agree, that Charles Darwin has done more to influence the philosophy of science, evolutionary biology, and the modern zeitgeist than any other individual (Mayr 2000: 78). Darwin developed a comprehensive biological theory of evolution that provided a unity to the natural world never before seen. He demonstrated the importance of historicity in science as well as a methodology that involved observation, comparison and classification, not just experimentation. Darwin's ideas, which rocked the foundation of Western society, still create heated debate in the modern world (Mayr 2000). So, just who was Charles Darwin?




***Figure* 2.1.4.1** - *Charles Darwin in 1854*

For a quick introduction to Darwin, watch the PBS video "Who was Charles Darwin?"

Darwin: The Early Years

Darwin was born on February 12, 1809 in Shrewsbury, England. His father, Robert, and grandfather, Erasmus, were both physicians. His mother, Susannah, was a part of the Wedgwood family, famous even today for their fine china. Erasmus Darwin is remembered for his work *Zoonomia*, which attempts to explain the natural world using evolutionary principles (Mayr 2000). It is clear that Darwin was introduced to current intellectual thought about evolution when he was young (Darwin 1887).

Darwin grew up with a love of the natural world. Darwin himself tells us that early on he developed a love of travel and collecting things, including minerals and insects:

> *With respect to science, I continued collecting minerals with much zeal, but quite unscientifically—all that I cared about was a new-named mineral, and I hardly attempted to classify them. I must have observed insects with some little care, for when ten years old (1819) I went for three weeks to Plas Edwards on the sea-coast in Wales, I was very much interested and surprised at seeing a large black and scarlet Hemipterous insect, many moths (Zygæna), and a Cicindela which are not found in Shropshire. I almost made up my mind to begin collecting all the insects which I could find dead, for on consulting my sister I concluded that it was not right to kill insects for the sake of making a collection. From reading White's 'Selborne,' I took much pleasure in watching the habits of birds, and even made notes on the subject. In my simplicity I remember wondering why every gentleman did not become an ornithologist. -Darwin 1887: 34-35*

Darwin carried this love of the natural world to the University of Edinburgh, where he was introduced to the work of Cuvier, Lamarck, and Saint-Hilaire (Young 2007), and subsequently, the University of Cambridge. At Edinburgh, Darwin studied medicine, but when it became apparent that the young Darwin had no interest in medicine, his father sent him to study theology at Cambridge. It was at Cambridge that Darwin's love of natural history flourished. He studied and worked with noted botanist J. S. Henslow, who would be instrumental in getting Darwin on the H.M.S. Beagle, scientific generalist, William Whewell, and geologist, Adam Sedgwick, all of whom were influential in his intellectual development, although Henslow would be the most critical individual in Darwin's post-college life (Bowler 1992, Mayr 1991, Young 2007). While at Cambridge, Darwin became an avid collector of beetles. This activity helped him become familiar with comparative anatomy, a methodology that would be critical to the development of his evolutionary theory.

> *The pretty Panagaus crux-major was a treasure in those days, and here at Down I saw a beetle running across a walk, and on picking it up instantly perceived that it differed slightly from P. crux-major, and it turned out to be P. quadripunctatus, which is only a variety or closely allied species, differing from it very slightly in outline (Darwin 1887:51),*

After graduating from Cambridge, Henslow helped Darwin secure a position aboard the H.M.S. Beagle as the ship's naturalist. He almost did not make the trip as his father was against it, but his uncle, Josiah Wedgwood, convinced Darwin's father to let him go on the voyage (Eldredge 2005). The Beagle, under the captaincy of Robert Fitzroy, left England on December 27, 1831 with the goal of charting the waters of South America (Bowler 1992, Futuyma 1986). While on the Beagle, Darwin, who was still an orthodox member of the Church of England (Futuyma 1986), made numerous observations that would eventually lead him to develop his theory of evolution by natural selection.

The Beagle Voyage

During the five years Darwin was on board the Beagle, he read such scientific works as Charles Lyell's *Principles of Geology* and made observations that eventually led him to question his belief in the fixity of species, a commonly held notion that species were unchanging and perfect in their environment. He collected fossils, rocks, plants and animals, many of which were shipped back to England where eventually specialists such as the ornithologist John Gould examined the specimens. Darwin's discovery of a new species of the rhea living in the same area as the already known rhea raised questions about perfect adaptation...if it were true, why would there be two species of rhea in the same area [1]? However, it was the Beagle's stop in the Galápagos Islands that would be most important for the later development of his theory.



***Figure* 2.1.4.2 -** *HMS Beagle by Conrad Martens.*

In the Galapagos, Darwin studied birds, iguanas and tortoises. The natives of the Galápagos showed Darwin how to determine which island the tortoises came from based on the shape of their shells. After learning this, he returned to the birds he had collected while in the Galápagos and sorted them by island. This gave Darwin, "...an unrivalled [*sic*] opportunity to study the effects of geographical isolation in the production of new species" (Bowler 1992: 300). It would be the Galápagos birds, specifically the mockingbirds, that would lead Darwin to his ideas about natural selection (learn more about the mockingbirds).

> To read some of Darwin's journal entries for the voyage, check out The Beagle Voyage or for a complete collection, see Journal of Researches

> You can follow the Beagle voyage, read quotes and view sketches from Darwin's diary by visiting the Natural History Museum's Big Idea Big Exhibition.

After the Beagle

Upon his return to England in October 1836, Darwin plunged himself into the scientific community of London. He had worked hard while on the Beagle so that he would not have to become a clergyman.

> *As far as I can judge of myself, I worked to the utmost during the voyage from the mere pleasure of investigation, and from my strong desire to add a few facts to the great mass of facts in Natural Science. But I was also ambitious to take a fair place among scientific men--whether more ambitious or less so than most of my fellow-workers, I can form no opinion (Charles Darwin quoted in Eldredge 2005: 27).*

Henslow and Lyell had made Darwin well known in scientific circles while he was still at sea, so he readily became active in the Geological Society (Bowler 1992, Eldredge 2005) and the Athenaeum Club, a gentlemen's club that served as a meeting place for those of an intellectual bent (Eldredge 2005).

One of the first things Darwin did on his return was to organize and ship his collections to various specialists. Ornithologist John Gould quickly examined Darwin's collection of birds, informing him that the mockingbirds in the collection were in fact different species of mockingbirds, some from the various Galápagos Islands and some from the South American mainland. Darwin theorized that this could only be explained if the island birds' antecedents had found their way from the mainland to the islands. Once there, the descendents were gradually modified until they became new species (Young 2007: 108). Here we find Darwin's coherent thoughts on the role geographic isolation has in speciation, or the creation of new species. By March 1837, Darwin had rejected the idea of the fixity of species and creation by design and accepted the idea of evolution and the role of geographic isolation in speciation (Futuyma 1986, Mayr 1991).

In July 1837, Darwin opened his first notebook, 'Transmutation of Species,' which during this time in history was code for evolution. In this notebook, Darwin explored ideas about reproduction and variation and the nature of species. By the time he concluded this notebook, Darwin has laid out his ideas on common descent and branching evolution. What he still didn't know at this point was how it all occured (Young 2007).

A year later, Darwin began his studies of artificial selection. His second notebook outlines his thoughts on variation and inheritance and their relation to adaptation (Young 2007). Darwin's studies of artificial selection helped him understand how a natural mechanism might work (Bowler 1992). Breeders at the time talked about something akin to natural selection but only in the context of how it kept species and varieties "true to type" (Young 2007: 112). When Darwin read Thomas Malthus's work, An Essay on the Principle of Population, in 1838, the concepts Malthus outlined on exponential population growth and competition among members of a population for a limited food supply provided the crucial piece that was missing. Darwin realized that in light of competition among members of a population, variations of traits that allowed one member to have an advantage over other member would be preserved. He further concluded that disadvantageous variations would die out (Young 2007). Darwin now understood how it all occurred for all of nature, including humans.

> *As soon as I had become, in the year 1837 or 1838, convinced that species are mutable productions, I could not avoid the belief that man must come from under the same law (Darwin quoted in Eldredge 2005: 29.).*

Darwin married his cousin, Emma Wedgwood, in 1839. By 1842, the Darwins moved from London to Downe in Kent in order for their children to grow up in the country and also for Darwin's health. In his later years, Darwin was plagued with ill health. While we do not know exactly from what he suffered, the symptoms suggest a malfunctioning of the autonomous nervous system (Mayr 1991).



***Figure* 2.1.4.3** - *Emma Wedgwood Darwin.*

In 1842, Darwin outlined his ideas on evolution by natural selection, expanding further in an 1844 essay that was to published only upon the event of his premature death. Bowler (1992) suggests Darwin did not publish at this time for two reasons: 1) the social climate--there was quite a bit of debate over Lamarck's theory of evolution and Robert Chambers' *Vestiges* that discussed evolution as part of a divine plan, and 2) he could not explain why the fossil record showed "...that many families had been subject to a constant trend towards increasing levels of specialization" (Bowler 1992: 304). The answer to this question would come a decade later when Darwin figured out that specialization provided an advantage even if the environment was stable because it cut down on competition for resources (Bowler 1992). During his time at Downe, Darwin used the hypothetico-deductive method (more commonly known as the scientific method) to build his case for evolution by natural selection, particularly in his work with pigeons (Young 2007).

Darwin began to write his "Big Species" book in 1856. He first published his ideas in 1858 after receiving a manuscript from Alfred Russel Wallace that outlined his ideas on natural selection, which coincided with Darwin's. At the urging of close friends in the scientific community, Darwin quickly put together a paper based on his 1844 essay that was presented by Charles Lyell and Joseph Hooker at the Linnean Society on July 1, 1858 along with Wallace's manuscript. Darwin then set to work on putting together an abstract of his Big Species book, which was published in 1859. This abstract, *On the Origin of Species by Natural Selection*, sold out in one day (Futuyma 1986, Mayr 1991, Young 2007). It was "the book that shook the world" (Mayr 1991: 7).

After the publication of *Origin*, Darwin's work focused on expanding on ideas that were touched on in the abstract, but not completely described. His other contributions to biology include *The Variation of Animals and Plants under Domestication* (1868), *The Descent of Man and Selection in Relation to Sex* (1871), *The Expression of the Emotions in Man and Animals* (1872), *Insectivorous Plants* (1875), *The Effects of Cross- and Fertilization in the Vegetable Kingdom* (1876), *The Different Forms of Flowers on Plants of the Same Species* (1877), *The Power of Movement in Plants* (1880 with Sir Francis Darwin), and *The Formation of Vegetable Mold, through the Action of Worms, with Observations on Their Habits* (1881).

Darwin died in 1882, but his work continued to be reviewed and oftentimes vilified by philosophers, theologians, scientists. etc. Darwin had a number of friends that supported his work both before and after his death, and while most scientists of the time came to support Darwin's proposal about common descent, few adhered to his concept of natural selection (Mayr 1991). It would not be until the 1920s when a fundamental change occurred in the thinking about organisms that the true genius of Darwin's work would be recognized on a broader scale (Futuyma 1986).

Darwin's contribution is summed up nicely by Futuyma (1986:6):

> *Darwin was the first to marshall on so grand a scale the evidence of the first thesis [i.e., descent with modification from a common ancestor], the historical reality of evolution by drawing on all relevant sources of information: the fossil record, the geographic distribution of species, comparative anatomy and embryology, and the modification of domesticated organisms.*

Mayr (1992) states that Darwin was unique among scientists of his day because he was not only a great naturalist and biologist, but was a great theoretician and experimenter. It was these differences that allowed him to make links across fields and develop his

comprehensive, complex theor(ies) of evolution.

The Theory of Descent with Modification through Natural Selection

All information in this subsection is drawn from Mayr 1991.




***Figure* 2.1.4.4** - *Charles Darwin photographic portrait, 1881*

Darwin's theory is actually a series of five theories:

1. Evolution as such: species are not immutable; they change slowly and steadily over time. This idea challenged the commonly held belief that there was a perfect design to life on earth.
2. Common descent: groups of organisms come from a common ancestor and all organisms ultimately originate from a single origin of life on earth. This idea provided a unity to biological life that had never been seen before.
3. Multiplication of species: geographic isolation lead to the evolution of new species. This idea explained the natural biological diversity on earth.
4. Gradualism: change occurs over long periods of time. This idea was in opposition to the contention that new species arose suddenly (saltation).
5. Natural selection: those individuals with variations (traits) best suited to enable an individual to reproduce within an environment are more likely to pass on their traits to next generation. Each generation would produce more variation, when again, the traits best suited to enable survival would be passed on. The classic example of natural selection in action is the peppered moth. Learn more about this moth by watching a short video on YouTube or read this short article from the State University of New York.

But there were some things that Darwin didn't know, namely, how traits were inherited or what caused variation within a population. Darwin also could not account for how similar species got on different continents, e.g., the ostrich in Africa and the rhea in South America. It would take decades for geologists to develop the theory of plate tectonics. Geologists in Darwin's day did not estimate the earth over 100 million years old, not enough time for evolution to work, but none of these things take away from the brilliancy or resiliency of Darwin's work. Over 100 years of testing has refined Darwin's theory while at the same time demonstrating over and over that his basic ideas, particularly natural selection, hold true. For more on what Darwin didn't know when he published his theory, check out this article from the Smithsonian Magazine.

What was it about Darwin's theory that "shook the world"? Some of his ideas challenged religious dogma and others challenged secular philosophies. Religious dogma that was directly challenged was the belief in the constancy of the world. The predominant belief in Europe and the United States was that while there had been a few catastrophes, e.g., Noah's flood, the world remained unchanged from when it was created approximately 6000 years ago by a devine Creator. All organisms were created to perfectly adapted to their environment, therefore, they did not change. Man (and it was the common vernacular of the time), with his unique soul, was separate from the rest of nature. In the arena of secular philosophy, Darwin's ideas challenged essentialism, or the fixed

properties of an entity, the "...causal processes of nature as they had been elaborated by the physicists" (Mayr 1991: 39), e.g., Newton, and teleology, or a pre-determined end. While science has tested and retested Darwin's ideas and subsequently developed a synthetic theory of evolution (the modern synthesis), the debate continues on in many religious circles. To read more about the modern debate between religion and evolution visit the Talk Origins "God and Evolution" page.

References

1. Bowler PJ. 1992. The Environmental Sciences. New York (NY): W.W. Norton & Company, Inc.
2. Darwin F, editor. 1887. The Life and Letters of Charles Darwin. London: John Murrary, Ablemarle Street.
3. Eldredge N. 2005. Darwin: Discoering the Tree of Life. New York (NY): W. W. Norton & Company.
4. Futuyma DJ. 1986. Evolutionary Biology, 2nd edition. Sunderland (MA): Sinauer Associates, Inc.
5. Hayden T. 2009. What Charles Darwin Didn't Know. Smithsonian (February). http://www.smithsonianmag.com/science-nature/What-Darwin-Didnt-Know.html?c=y&page=1. Accessed. 2010 October 15.
6. Mayr E. 1991. One Long Argument: Charles Darwin and the Genesis of Modern Evolutionary Thought. Cambridge (MA): Harvard University Press.
7. Mayr E. 2000. Darwin's Influence on Modern Thought. Sci Am 283 (1): 78-83.
8. Young D. 2007. The Discovery of Evolution, 2nd edition. Cambridge: Cambridge University Press.

This page titled 2.1.4: Darwinian Evolution is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.4: Darwinian Evolution** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.5: Cell Biology and Basic Genetics

When Darwin developed his theory, there are two main things that he did not know: (1)where the variation comes from, and (2) how the traits are passed from one generation to the next. In the decades since *On the Origin of Species* was published, subsequent generations of biologists, natural scientists, geneticists, paleoanthropologists, etc., have conducted research that has refined our knowledge about evolution. We now can address the things that Darwin did not know. Cell biology and genetics explain where variation comes from and how those traits are inherited.

Cell biology

All living organisms are made of ***cells***. Some organisms are single-celled and spend their life made of just one cell and some organisms, like humans, are multi-cellular (have many cells). Here are some basic points that you always have to remember about cells:

- cells are the structural and functional units of all living things
- all cells come from pre-existing cells
- cells contain hereditary information (which is passed from "parent" cells to "daughter" cells during cell division).

Cell function

Cells are often called "the building blocks of life". Not only are all living things made of cells (the 'building block' part), they also are living functional units themselves. This is because cells do the following:

- obtain nutrients and other molecules through the cellular membrane (or cell wall) and convert that to biologically useful energy and useful molecules for building new cells (such as proteins and nucleic acids)
- make more cells (reproduction)

Types of cells

There are two types of cells: prokaryotic and eukaryotic. Prokaryotes include bacteria and blue-green algae. These are single-celled organisms in which the genetic material is not distinctly separated from the other components of the cell. Eukaryotes include everyone else. These can be single-celled, but are more often multi-celled organisms in which the genetic material is separated from all of the other cellular components (by something called the **nucleus**). Because our focus is on humans, the rest of our discussion will center on eukaryotes and eukaryotic cells.

Cell structure




Figure **2.1.5.1** - *A typical animal cell and its parts*

Each animal cell consists of a selectively permeable membrane (plants have cell walls) which contains the cytoplasm.

Organelles are like "little organs" within the cytoplasm, each performing a different vital cellular function. These are some of the more important (to us) organelles.

- Mitochondria: These are the "powerhouses" of the cell, responsible for production of the energy-rich molecule, ATP (adenine triphosphate), which powers the activities of the cell. Each cell has hundreds to thousands of mitochondria. Mitochondria have their own DNA, known as mitochondrial DNA (mtDNA), which is different from nuclear DNA (see below). Mitochondrial DNA has become an important tool in evolutionary research.
- Endoplasmic reticulum: This is a transport network for molecules that have specific destinations or require certain modifications. It comes in two types: smooth and rough. Rough endoplasmic reticulum has ribosomes (see below) on the surface.
- Golgi apparatus: Processes and packages large molecules such as proteins and lipids (fats) that are produced by the cell.
- Lysosomes: Breaks down non-usable organelles, food, viruses, and bacteria using enzymes.
- Ribosomes: Protein synthesis
- Nucleus: This is where you'll find the hereditary material, DNA (deoxyribonucleic acid) and RNA (ribonucleic acid).

Exercise 2.1.5.1

Explore the cell!!

In this interactive exercise, you can identify the parts of a typical animal cell and review the functions of the various organelles.

Typical animal cell

DNA

DNA (deoxyribonucleic acid) is the genetic blueprint of the cell. The structure of DNA is what's known as a "double helix", which looks much like a twisted ladder. The basic unit of DNA is a molecule called a nucleotide. Nucleotides are made of a sugar (deoxyribonucleic acid), a phosphate group, and a nitrogenous base. There are 4 different types of nucleotides, depending on what type of nitrogenous base they have.




***Figure* 2.1.5.2** - *DNA structure*

- adenine (A)
- thymine (T)
- cytosine (C)
- guanine (G)

These are separated into 2 groups:

- purines: adenine and guanine
- pyrimidines: cytosine and thymine

On the DNA "rungs", the bases must go in pairs: adenine (A) always bonds with thymine (T) and cytosine (C) with guanine (G). So, for base-pairs remember it's always A+T and C+G! (That rule is sometimes called "The base-pair rule".) If you want to explore further, check out what the Human Genome Project has learned about the human DNA sequence

RNA structure

RNA (ribonucleic acid) has a very similar structure to DNA, except that it's single-stranded, has a different sugar (ribonucleic acid, rather than deoxyribonucleic acid), and does not contain thymine (T). Instead, it contains uracil (U), which like thymine, bonds with adenine (A). So, in this case, the base-pair rule is A+U and C+G.

DNA function

For our purposes, we will concern ourselves with two of DNA's functions: replication and protein synthesis.

DNA replication

In this case, the goal is to replicate a DNA double helix to produce more DNA. DNA replication occurs in the nucleus of the cell.

The basic steps of DNA replication are:

1) The DNA is "unzipped" into two single strands.

2) Each unzipped strand acts as a template for reproduction of the complimentary strand.

3) The product is two copies of the original DNA, each containing one strand from the original DNA and one strand that has been added on.

Here's a slightly more thorough illustration of the process: DNA replication - simple.

And for those who are really curious about the details of the process: DNA replication - complex.

And for those who are looking for a little fun with their DNA replication: DNA - the double helix game

Important note: The important thing to remember about DNA replication is that when the complimentary strand is created, sometimes accidents or mistakes happen. For example, a G gets matched up with a T instead of the proper A. This is where some of the variation in hereditary information comes from.

Protein synthesis




***Figure* 2.1.5.3** - *Protein synthesis*

Proteins are the most common large molecules in cells. Bones, muscles, and red blood cells (among lots of other body parts) are made mostly of proteins. Therefore, the production of proteins is obviously a very important process in the body.

Proteins are made of smaller units called amino acids. There are 20 different amino acids, 9 of which are "essential amino acids", which means that they must be consumed through the diet, rather than being synthesized by the body. The sequence of amino acids

in a protein determines its structure and function.

The base pair sequence of the DNA molecule is known as **the genetic code**. The genetic code consists of 3-base sequences called **codons**.(Remember our friends A, T, G, and C -- the nitrogenous bases? That's what we're talking about here.) Each codon either codes for an amino acid, or signals that the protein chain is starting (an initiation codon) or stopping (a termination codon). The table on the right shows which codons code for which amino acids.

Each DNA molecule contains the information to make up many different proteins. A portion of a DNA molecule responsible for making up a single protein (or sometimes just part of a protein, called a polypeptide) is a **gene**. Therefore, each DNA molecule consists of many genes, that code for many proteins.

Protein synthesis has two basic steps: (1) transcription and (2) translation.

Definitions

Transcription

1) The gene (DNA) is copied onto RNA. The RNA copy of the gene is called the messenger RNA (or mRNA).

2) The mRNA leaves the nucleus and goes to the rough endoplasmic reticulum. (Remember this organelle from the cell structure section above?)

Translation

1) The mRNA goes into the ribosomes, where tRNA (translation RNA) reads the mRNA.

2) As the tRNA reads the mRNA, it attaches complementary amino acids to the newly synthesized amino acid chain (AKA the growing protein chain).

Note

Important note: The important thing to remember about protein synthesis is that when a protein is synthesized, sometimes accidents or mistakes happen. This is where some of the variation in protein synthesis comes from. Additionally, remember the possible mistakes in DNA replication? If the original DNA strand is changed so that a gene is altered, that can affect protein synthesis as well.

This video has fantastic digital animations illustrating DNA replication and protein synthesis.

Cell replication

To make more cells, we not only have to have replicated DNA, we have to replicate the rest of the cell as well in order to physically divide that replicated DNA. For body cells (AKA somatic cells), this process is called **mitosis**. For sex cells (AKA sperm and eggs), this process is called **meiosis**. But first, a little bit about DNA and chromosomes.




Figure **2.1.5.4** - *Chromosome*

Chromosomes

Most of the time, DNA in cells is in loosely arranged, dispersed strands. When DNA is like this, it's called **chromatin**. However, during cell replication DNA condenses into structures called **chromosomes**.

Each chromosome is made of (the numbers reference the illustration to the right): 1) two "sister" chromatids, 2) a centromere, which is a highly condensed area on the chromosome, 3) the "short arm" of the chromatid, 4) the "long arm" of the chromatid, and the 5) telomere, which is the end of a chromosome and is involved in both replication and chromosomal stability. In this diagram, the different colors represent genes on the chromosome.

Each somatic cell (sex cells are different) has the same number of chromosomes. This number is consistent within species, but often differs between species. The number of chromosomes in each body cell is called the **diploid number**. Sex cells have half as many chromosomes as do somatic cells. This number is called the **haploid number**.

Most species have two copies of each chromosome in each somatic cell. The member-chromosomes of each pair are called **homologous chromosomes**.




***Figure* 2.1.5.5** - *A typical human (male) karyotype*

On the left is a **karyotype**, which is an image of the full diploid complement of chromosomes from an individual. Above each number is each pair of homologous chromosomes. This is from a human, who typically have 23 pairs of chromosomes in each somatic cell, for a diploid total of 46 chromosomes. For humans, sex cells normally have 23 chromosomes.

Genes are found on each chromosome at a particular location called a **locus** (sing.) or **loci** (pl.). Different versions of the same gene are called **alleles**. If someone has the same version of a gene (the same alleles) on both of their homologous chromosomes, they have **homozygous alleles**. If, however, someone has two different versions of a gene (different alleles) on their homologous chromosomes, they have **heterozygous alleles**. For example, if on this person's chromosome 9 pair they have one allele for blood type A and one allele for blood type B, we would say that they are heterozygous for blood type (AB). If they have two alleles for blood type O, we would say that they are homozygous for blood type (OO).

Note

Explore human chromosomes

Mitosis

Somatic cells (AKA body cells) are replicated via the process of mitosis. This is where all new body cells come from. So, when you scratch yourself on something, mitosis is what makes new skin cells for healing that scratch.




Figure **2.1.5.6** - *Mitosis*

Mitosis has 4 phases (note that DNA has already replicated when mitosis starts):

1) Prophase:

- the nuclear envelope disappears
- chromatin condenses and forms chromosomes
- the chromosomes move toward the equator of the cell

2) Metaphase

- the chromosomes are lined up along the equator of the cell

3) Anaphase

- the sister chromatids split up and move toward the opposite ends of the cell

4) Telophase

- the cytoplasm divides between the two cells
- the chromosomes turn back into chromatin

The result is two identical "daughter" somatic cells, each of which contains one chromosome from the "parent" cell and one replicated chromosome.

Here's a video that illustrates the process: mitosis in action.

Meiosis

Sex cells (eggs and sperm) are replicated via the process of meiosis. For human females this process occurs and is completed before birth, but for males meiosis begins at puberty and continues throughout their lifespans. Meiosis is more complicated than mitosis in that it has two "cycles".




Figure **2.1.5.7** - *Meiosis*

1) Prophase 1

- chromatin condenses and forms chromosomes
- DNA is exchanged between homologous chromosomes. This is called homologus **recombination**. Recombination is an important source of genetic variation.
- the chromosomes start to line up at the equator of the cell

2) Metaphase 1

- the chromosomes are lined up along the equator of the cell

3) Anaphase 1

- the homologous chromosomes move toward the opposite ends of the cell

4) Telophase 1

- the cytoplasm divides between the two cells
- the chromosomes uncoil and turn back into chromatin

5) Prophase 2

- nuclear envelope disappears
- chromatin condenses into chromosomes
- the chromosomes start to line up at the equator of the cell

6) Metaphase 2

- the chromosomes are lined up along the equator of the cell

7) Anaphase 2

- the sister chromatids split apart at the centromeres and move toward the opposite ends of the cell

8) Telophase 2

- chromosomes turn back into chromatin
- the cytoplasm divides between the two cells

Here's a video that illustrates the process: meiosis in action

Errors during mitosis and meiosis




***Figure* 2.1.5.8** - *The karyotype of an individual with Down syndrome*

The chromosomes separate during meiosis 1 and the sister chromatids separate in meiosis 2 and mitosis. If this proceeds normally, it's called **disjunction**. If the separations occur abnormally, it's called **nondisjunction**. What results is an abnormal number of chromosomes (either too many, AKA trisomy OR too few, AKA monosomy).

Some of the more familiar nondisjunctions include:

Down Syndrome - trisomy of chromosome 21 (a karyotype of this is to the right)

Klinefelter Syndrome - extra X chromosomes in males (ex. XXY)

Turner Syndrome - lacking an X chromosome in females (XO)

This page titled 2.1.5: Cell Biology and Basic Genetics is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.5: Cell Biology and Basic Genetics** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.6: Mendelian Genetics

In the previous section, we explored the mechanisms of biological variation and evolution at the most fundamental level - cells and DNA. In this section, we are going to look at how genetics functions for individuals. This field of genetics is often called Classical or Mendelian Genetics. It focuses on how various traits are passed from one individual to the next.

Some working vocabulary

- **Locus** (sing.) and **Loci** (pl.): the specific position of a gene on a chromosome
- **Gene**: a discrete unit of a DNA sequence that codes for a protein (or polypeptide) or for a characteristic, process, or trait
- **Allele**: alternate forms of a gene at a given locus (ex. T and t)
- **Genotype**: the genetic makeup of an individual
- **Phenotype**: the outward expression of the genotype, as affected by the environment and culture of an individual
- **Dominant allele**: an allele that masks the effect of other alleles in the phenotype (ex. in the case of Tt, T masks the effect of t)
- **Recessive allele**: an allele that is masked in the phenotype unless present in a double dose (ex. in the case of tt, there is no T to mask the effect of t)
- **Codominant allele**: both alleles are expressed in the phenotype (ex. in the ABO blood types, there are three alleles (A,B,O); A and B are codominant alleles)
- **Homozygous**: two of the same alleles at a given locus on homologous chromosomes (ex. TT or tt)
- **Homozygous dominant**: both alleles are the dominant allele (ex. TT)
- **Homozygous recessive**: both alleles are the recessive allele (ex. tt)
- **Heterozygous**: two different alleles (ex. Tt)

Note: Capital letters are usually shorthand for the dominant allele and lower-case is usually shorthand for the recessive allele when discussing Mendelian genetics.

Traits

Traits come in two types: Simple (also called Mendelian or discrete traits) and complex (sometimes called *continuous* or quantitative traits).

Simple traits

These are either/or traits (they usually have 2 phenotypes):

- The trait is either present or absent (e.g., hairy pinna).
- or the trait is one state or the other (e.g., attached or unattached earlobes: see image).
- It's controlled by one gene, which has one effect.

They are biologically (genetically) determined:

- They cannot be changed.
- There is little to no influence from the environment.
- They have a finite number of genotypes, usually very small. For example, 1 trait with 2 options (T and t) = 3 genotypes (ex. TT, tt, Tt).

There are approximately 4000 traits identified as discrete or simple traits of inheritance. Some additional examples include:

- Soft vs. flaky earwax
- Widow's peak
- Blood type
- Dimples
- Chin cleft (Figure **2.1.6.1**)




***Figure* 2.1.6.1** - *Cleft chin*

Complex traits

- These traits have a continuous distibution (e.g., height: there are people ranging from short to tall, not just short and tall).
- The phenotype is the result of two factors; the number of loci involved and environmental influence (this includes culture).
- The trait is influenced by more than one gene; this trait is referred to as **polygenic**).
- Depending on the number of genes involved, there can be many genotypes. The more genes that are involved, the finer the genotype categories until they converge into a continuous distribution.
  - 1 gene = 3 genotypes
  - 2 genes = 9 genotypes
  - 3 genes = 27 genotypes
  - 4 genes = 81 genotypes
- It's often difficult to try to divide them into categories due to an infinite number of phenotypes

The opposite of this, where one gene has multiple effects, is called **pleiotropy**.

Due to the large role that the environment plays in complex traits, we have a statistical concept in genetics called heritability. **Heritability** is the proportion of the total variation observed in the population that can be attributed to genetics rather than to the environment.

Gregor Mendel and his Peas

For simple traits, it is easy to trace patterns of inheritance and use them to show how traits are passed on from one generation to the next. Charles Darwin thought that traits were passed down by "**blending**": each parent contributes equally to the offspring, and these contributions are mixed together and then halved at each successive generation (Figure **2.1.6.2**).




Figure* 2.1.6.2*: *Blending inheritance*

However, there was a contemporary of his that Darwin didn't know about, a monk named Gregor Mendel (pictured below), who figured out something important about simple traits and genes. Mendel is famous for a series of experiments that he completed with pea plants (between about 1856 and 1868) and the paper that he wrote about it (*Experiments with Plant Hybrids*). Unfortunately, despite the fact that Mendel published this work in 1865, it was relatively unknown until 1900. What he concluded was that genetic information is inherited in discrete units (he called them "factors", we call them "genes"). His work demonstrated that inheritance is non-blending or "**particulate**".



***Figure* 2.1.6.3** - *Gregor Mendel*

Mendel's pea experiments

Mendel first noticed that pea plants have seven simple traits:

1. Seed shape and color
2. Cotyledon color
3. Flower color
4. Pod shape
5. Pod color
6. Location of pods and flowers on the stem
7. Plant size

He also noted that when one selectively bred the plants with the various traits together, he could predict the ratios of offspring characteristics. For example, he noticed that when he bred pure-bred plants with violet flowers with those with white flowers, the first generation (F1) all had violet flowers. If blending inheritance was functioning, one would expect all of the flowers to be a pale violet, halfway between the original violet and white.




***Figure* 2.1.6.4** - *Mendel's seven traits*

Then when he took those F1 violet-flowered plants and bred them together, the second generation (F2) had a ratio of 3:1 violet to white flowers (there were still no pale-violet hybrids). This phenomenon was consistent across the pea plants' other six traits.

He determined that this is what was going on:

- The pure-bred violet plant was homozygous for its violet allele (BB).
- The pure-bred white plant was homozygous for its white allele (bb).
- So, when the two plants were crossed -- all of the offspring (F1) were heterozygous (Bb).
- And, when the heterozygous (Bb) offspring were crossed, this was the result: the genotype is 1 BB: 2 Bb: 1 bb and the phenotype is 3 violet: 1 white.




***Figure* 2.1.6.5** - *A Punnett square of pea flowers*

Punnett squares

This diagram in Figure **2.1.6.5** is called a Punnett square (after an early 20th c. English geneticist named Reginald Punnett). They're really pretty simple to construct. One just puts the alleles of each parent (one parent on the top and the other down the side) and then you fill in the boxes of what the offspring will be. In order to devise the ratios of the various genotypes and phenotypes, just count the number of boxes of each.

For further information on how to construct Punnett squares, go here or watch this video from Bozeman Science.

Mendel's very thorough experiments (some estimates say that he bred approximately 29,000 pea plants) lead him to write these four postulates.

Mendels' postulates

1. Genetic characteristics are controlled by genes that exist in pairs in individual organisms.
   - The pairs of genes that he's referring to are those on each pair of homologous chromosomes.
2. When an individual has two different genes for a particular trait, only one is expressed and is dominant to the other, which is recessive.
   - In other words, when an individual has a T allele and a t allele, the T allele will mask the expression of the t allele. Therefore, the phenotype expressed will be that of T.
3. During the formation of gametes (AKA sex cells, sperm and eggs), the genes separate, or segregate, randomly so that each sex cell receives one or the other with equal likelihood.
   - In other words, the offspring have an equal likelihood of getting a particular gene from one parent or the other.
   - This is also known as "Mendel's Law of Segregation".
4. During the formation of gametes, segregating pairs of genes assort independently of each other.
   - In other words, the segregation of any particular pair of genes does not affect the segregation of any other pair.
   - This is also known as "*Mendel's Law of Independent Assortment*"

Some examples of inheritance patterns in humans

Autosomal dominant traits




***Figure* 2.1.6.6** - *Autosomal dominant inheritance*

- An individual only has to have one copy of the allele in order to exhibit these traits.
- If an individual has the trait, then at least one of their parents has the trait.
- Both males and females are affected equally and should be capable of transmitting the trait.
- There is no skipping of generations.
- If one of the parents has it, then on average, 50% of the children will show the trait.

Examples of this include:

Huntington's disease

Neurofibromatosis-1

Autosomal recessive traits

- An individual needs both recessive alleles to express the trait
- Heterozygotes are considered to be "carriers"
- Both males and females should be affected equally and should be capable of transmitting the trait.




Figure **2.1.6.7** - *Autosomal recessive inheritance*

Examples of this include:

Cystic fibrosis, sickle cell anemia, and Tay Sachs disease

Sex-linked traits

- ***Remember*** that all females have two X sex chromosomes (XX) and males have one X and one Y sex chromosome (XY)
- These traits are inherited on the sex chromosomes (X and Y)
- There have different rates for males and females

There are three types:

1. X-linked dominant
2. X-linked recessive
3. Y-linked

X-linked dominant




***Figure* 2.1.6.8** - *X linked dominant inheritance*

- Both males and females will express the trait if they possess at least one copy of the allele.

An example of this is:

Incontinentia pigmenti (IP)

X-linked recessive

- For females, they need two copies of the allele to express the trait, but only one to be a "carrier".
- For males, they need only one copy of the allele to express the trait. They will pass the allele down to all of their daughters.




***Figure* 2.1.6.9** - *X linked recessive inheritance*

Examples of this include:

Red-green colorblindness and Hemophilia A

Y-linked

- No female will carry or express the allele (because they don't have a y chromosome)
- All sons will carry and express the allele (because they must inherit a y chromosome from their father)

An example of this is:

Hairy pinna

References

1. Klug, WS and Cummings, MR. 1991. Concepts of Genetics, 3rd. edition. Macmillan Publishing Company: New York.

This page titled 2.1.6: Mendelian Genetics is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.6: Mendelian Genetics** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.7: Modern Synthesis

The Modern Synthesis incorporates data from multiple scientific disciplines: biology, the natural sciences, genetics, paleontology, and paleoanthropology. While the basis for evolutionary theory was established by the beginning of the twentieth-century, debate continued as to whether natural selection or mutation was more important in evolution. In the 1930s, these two points of views came together to form the basis of the modern synthesis. Fisher, Haldane, and Wright developed mathematical models that became the foundation of population genetics, which studies the frequency of alleles, genotypes, and phenotypes in populations (Miller 2011: 321). However, it was Theodosius Dobzhansky that married theoretical modeling and empirical research in his seminal work, Genetics and the Origin of Species (1937) and signals the start of the Modern Synthesis (Artmann 2006). Biologist Ernst Mayr was another major contributor to the development of the modern synthesis. His work on the isolating mechanisms of speciation led to new insight into the evolution of new species. Mayr's Systematics and the Origin of Species (1942) is one of the key works of the Modern Synthesis.




***Figure* 2.1.7.1** - *Note by Francis Crick.*

A New Definition

Emerging from the work of scientists of the Modern Synthesis, we know that evolution is a two-step process (Miller 2011: 85):

1. The production and redistribution of variation, and
2. Natural selection acts on variation resulting in differential reproduction among individuals in a population.

A **population** is a group of individuals from which mates are usually found. Sometimes this is referred to as a **breeding population**. All members of a population share a **gene pool**, which is all of the genes found among the individuals of the population.




***Figure* 2.1.7.2** - *Graphic of microevolution. Modeled on University of California Museum of Paleontology's Understanding Evolution (evolution.berkeley.edu)*

Scientists can study variation in a gene pool and differential reproduction in populations to determine **allele frequencies**, which is the frequency of an allele relative to that of other alleles of the same gene. Changing allele frequencies within a population from one generation to the next is the definition of evolution; specifically, this is **microevolution. Macroevolution** is change that occurs after many generations (hundred or thousands); e.g., **speciation**, which is the evolution of a species into a new species. "Macroevolution encompasses the grandest trends and transformation in evolution, such as the origin of mammals and the radiation

of flowering plants (Evolution 101, 2015). Macroevolutionary events are reconstructed using data from living organisms, geology, and fossils. These are generally represented in a graphic form called a **phylogentic tree** (pictured below). Phylogenetic trees are a visual representation of proposed evolutionary relationships. They help researchers organize data. To learn more about phylogenetic trees check out Reading a Phylogenetic Tree.




***Figure* 2.1.7.3** - *Simplified phylogenetic tree*

Step 1: Production and Redistribution of Variation

As we know, production and redistribution of variation is the first step of evolution. There are several factors involved in the production and redistribution of variation: 1. Mutation 2. Genetic drift 3. Gene flow, and 4. Recombination. These factors, along with natural selection, which is the second step of evolution, are frequently referred to as the **forces or mechanisms of evolution**.

Mutation

Mutation is a random change in the genetic coding and is the only new source of genetic material. Mutations occur during the DNA replication process. In an evolutionary sense, it is only mutations that occur during meiosis (replication of sex cells) that are hereditary. Many mutations do not have detrimental effects to the individual, but some do. If the mutations are extreme, pregnancy often results in spontaneous abortion (miscarriage). However, mutations provide a necessary pool of genetic variation in a population.




***Figure* 2.1.7.4** - *Illustration of a point mutation*

There are different types of mutations. The first is a **point mutation**, which is a small change in the DNA sequence. Point mutations may be **synonymous point mutations or silent mutations**, which means that while the codon is altered the same amino acid is produced, or **nonsynonymous point mutations**, which result in the production of a different amino acid that can have a negative effect on the individual. For instance, sickle cell anemia results in a point mutation where "…a mutation on human chromosome 11 converts a GAG codon into a GTG codon. The GAG codon is encoded to produce the amino acid valine, whereas the GTG codon is encoded to produce glutamic acid. This substitution results in sickle cell anemia…" (Larsen 2008: 93).

Frameshift mutations occur when one or more nitrogenous bases are inserted into or deleted from the DNA sequence. The insertion or deletion causes the reading of the gene to be altered or stopped. It results in the production of a nonfunctioning protein.




***Figure* 2.1.7.5** - *Illustration of a deletion mutation*




***Figure* 2.1.7.6** - *Illustration of an insertion mutation*

Transposable elements are mutations that occur when pieces of DNA copy themselves into new areas of chromosomes. If this occurs in a non-coding segment of the DNA then there are little to no effects on the individual. If it transposes to a coding segment of DNA then serious consequences can occur. In extreme cases, entire chromosomes may be copied (a trisomy) or deleted (a monosomy).

Mutations may have no known cause. These are called **spontaneous mutations**. Mutations caused by environmental factors are called **induced mutations**.

Genetic Drift




***Figure* 2.1.7.7** - *Illustration of founder effect*

Genetic drift is the "…chance variation in allele frequency in small populations" (Gould 2006:1036). This can happen in a couple of ways: founder effect and bottleneck effect (or population bottleneck). **Founder effect** can occur if a small group separates from its parent group and settles elsewhere. Geographic isolation and endogamous marriage (human marriage pattern where mates are chosen from within a specific group, narrowing the gene pool) or mating practices restrict the gene pool, resulting in a lack of variation in comparison to the original parent population. If an allele is rare in the parent population, the small breeding population of the new group creates an environment whereby the rare allele can become common.

There are numerous examples of the founder effect among human populations. These include,




Figure \(\PageIndex{8}\) *- Map of blood group O*

- Huntington's Disease in Afrikaner populations in South Africa and among Venezuelans living near Lake Maraciabo,
- Prevalence of O blood type in Native Americans
- Ellis-van-Creveld syndrome (polydactyly) in Amish communities, and
- The Blue People of Kentucky (learn more about the Blue People www.indiana.edu/~oso/lessons/Blues/TheBlues.htm).

Bottleneck effect occurs when a population is drastically reduced in size due to some disaster, e.g., natural disaster, habitat destruction, or disease, leaving the surviving population with different allele frequencies as the original population. Four human population bottlenecks all related to climatic events have been identified. The most severe occurred about 70,000 years ago when a supervolcano erupted in Sumatra. It is estimated that only 5000 females survived in Africa (Ambrose 2003).




Figure 2.1.7.9 *- Illustration of the bottleneck effect*

Gene Flow




Figure 2.1.7.10 *- Illustration of gene flow.*

Gene flow, also called **admixture** or migration (in reference to people), is the exchange of genetic material between populations (Miller 2011). For humans, gene flow generally occurs when people move from one location to another; however, in the modern

world it is not out of the realm of possibility that someone's genetic material could move with the actual human being leaving their hometown. It should also be noted that migration does not always result in gene flow. For gene flow to occur interbreeding must happen. Human populations have exchanged genetic materials for millennia, helping to explain why there is only one human species.

Gene flow effects evolution in a couple of important ways (Evolution 101, 2015): • With the introduction and reintroduction of genes to a population its genetic variation is increased, and • The movement of genetic material can make distant populations genetically similar to one another, which reduces the chance of speciation.

Recombination

Recombination, or crossing over, can occur during meiosis. As the term implies, segments of chromosomes are exchanged between non-sister chromatids. This creates new combinations of genes that are not found in either parent (Speer 2003).

Step 2: Natural Selection




***Figure* 2.1.7.11** - *Peppered moths.*




***Figure* 2.1.7.12** - *Types of selection.*

Once variation is produced in some combination of the above processes, natural selection goes to work, causing directional change in allele frequency

related to environmental conditions that may result in evolutionary change, i.e., speciation. Shifts in allele frequency are called **adaptation**. In phrasing it this way, it sounds like natural selection works consciously, but it does not. Natural selection is a process

that is mindless and mechanistic. Perfection and progress are not goals. It can only act on those variants that are present in a population.




***Figure* 2.1.7.13** - Male peacock display.

Darwin concluded, "…individuals with advantageous characteristics…survive in higher numbers and produce more offspring than members of a population lacking advantageous characteristics" (Larsen 2008: 96). This is referred to as **differential reproduction** or **fitness**. Specifically, fitness is defined by genotypes—some genotypes confer an advantage to an individual allowing that individual to reproduce at a greater frequency than another individual, characteristics that are inherited by their offspring. It could be something like beak shape that allows a bird to exploit resources better than other birds (think Darwin's finches) or the color of wings that either hide or expose the peppered moth to predators (watch youtu.be/sVVldxxbWig for the full explanation) or a coloring, like the plumage of a male peacock, that attracts mates (this is referred to as **sexual selection**, which is a special case of natural selection which refers to an individuals ability to attract and copulate with a mate) that lead to differential reproduction.

There are three types of selection: disruptive, stabilizing, and directional selection. With disruptive selection, both homozygotes (the phenotypic extremes) are selected for while with directional selection only one of the homozygotes is selected for. Disrupting selection may result in a speciation event. Stabilizing selection refers to selection for the heterozygote, resulting in decreasing genetic diversity within the population.

If you would like a bit more on the types of selection, check out Bio.edu: Stabilizing selection (http://www.brooklyn.cuny.edu/bc/ahp/LAD/C21/C21_Stabilizing.html) Disruptive selection (http://www.brooklyn.cuny.edu/bc/ahp/LAD/C21/C21_Disruptive.html) Directional selection (http://www.brooklyn.cuny.edu/bc/ahp/LAD/C21/C21_Directional.html)

Species and Speciation

Mayr proposed the most commonly referenced definition of species (1942; quoted in Provine 2004), "Species are a group of actually or potentially interbreeding natural populations, which are reproductively isolated from other such groups." This definition is often referred to as the **biological species concept**. A key factor is that a species actually or potentially breeds in nature. However, this definition can be difficult to apply; e.g., organisms that reproduce asexually or organisms that form hybrids (carrion crows and hooded crows sometimes interbreed—are they one species or two species?). The biological species concept falls apart when applied to the fossil record, as it is impossible for scientists to observe the breeding behaviors of extinct organisms. Because

of this, fossil species are called **paleospecies** and can cover long periods of time. Paleontologists and paleoanthropologists use comparative analysis with contemporary species that are closely related to determine whether variation seen in fossil specimens is **intraspecific** (differences within a species due to age and sex) or **interspecific** (differences due to reproductive isolation). It is important to remember that the species or paleospecies name is a classificatory tool used in the sciences. It helps scientists organize information and communicate more easily with one another.

Speciation is "…the process by which a new species evolves from an earlier species (Jurmain et al 2013: 105). This idea is one of Darwin's five theories, specifically those of common descent and multiplicity of species, whereby new species arise from earlier species.

Simsonian Evolution

Just for fun, check out how The Simpsons explain evolution: www.youtube.com/watch?v=faRlFsYmkeY




***Figure* 2.1.7.14** - *Darwin's first tree.*

Explore (optional)

If you would like to read more about the concepts introduced on this page, check out the following web sources:

What is evolution? A short essay by Laurence A. Moran exploring the definition of evolution. (bioinfo.med.utoronto.ca/Evolu...Evolution.html)

Evolution is a Fact and a Theory. Another short essay by Moran explaining how evolution is both a fact and a theory. (bioinfo.med.utoronto.ca/Evolu...nd_Theory.html)

Why is evolution controversial anyway? A short PBS video discussing the controversy of evolutionary theory. You will need Quicktime or RealPlayer to watch this video. This page has several videos on it—look for the video entitled Why is evolution controversial anyway? (http://www.pbs.org/wgbh/evolution/educators/teachstuds/svideos.html)

References

1. Ambrose S. Population bottleneck. In: Robinson R, editor. Genetics, Vol. 3. New York (NY): Macmillan Reference USA; 2003. p. 167-171.

2. Artmann S. Neo-Darwinism, Origin of. In: Birx, HJ, editor. Encyclopedia of anthropology, Vol. 4. Thousand Oaks (CA): SAGE Reference; 2006. p. 1727-1732.
3. Dusheck J. Population genetics. In: Robinson R, editor. Genetics, Vol. 3. New York (NY): Macmillan Reference, USA, 2003. p. 171-174.
4. Ernst Mayr and the evolutionary synthesis [Internet]. Boston (MA): WGBH Educational Foundation and Clear Blue Sky Productions; c2001 [cited 2015 May 12]. Available from <http://www.pbs.org/wgbh/evolution/library/06/2/l_062_01.html>.
5. Evol3000 [Internet]. 2013 November 13 [cited 2015 May 12]; Available from: wallace.genetics.uga.edu/groups/evol3000/wiki/fb221/Bottlenecks_and_Founder_Effects.html.
6. Evolution 101 [Internet]. Berkeley (CA): University of California Museum of Paleontology; c2015 [cited 2015 May 12]. Available from evolution.berkeley.edu/evolibrary/article/evo_01.
7. Ferrell K. The modern synthesis of evolutionary theory. In: Schlage N, Lauer J, editors. Science and its times, Vol. 6: 1900 to 1949. Detroit (MI): Gale; 2000. p. 104-107.
8. Gould JL. Genetic drift. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 3. Thousand Oaks (CA): SAGE Reference; 2006. p. 1036.
9. Heddle J. Mutation. In: Robinson R, editor. Genetics, Vol. 3. New York (NY): Macmillan Reference USA; 2003. p. 93-98.
10. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
11. Larsen CS. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.; 2008. 430 p.
12. Mayr, E. Systematics and the Origin of Species. New York (NY): Columbia University Press. Quoted in Provine, WB. Ernst Mayr: genetics and speciation. Genetics. 2004; 167(3): 1041-1046.
13. Speer, MC. Crossing over. In: Robinson R, editor. Genetics, Vol. 3. New York (NY): Macmillan Reference USA; 2003. p. 194-197.
14. Sykora, P. Darwinism, modern. In: Birx, HJ, editor. Encyclopedia of anthropology, Vol. 2. Thousand Oaks (CA): SAGE Reference; 2006. p. 692-696.

This page titled 2.1.7: Modern Synthesis is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.7: Modern Synthesis** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.1.8: Modern Human Variation

One of the most striking things about humans is the huge range of biological variation we have as a species. In the previous section, we explored the mechanisms of change involved in the evolution of populations (mutation, migration/gene flow, genetic drift, and natural selection). In this section, we'll look at how those affect human populations to create human diversity.

Polymorphisms

A **polymorphism** is when a gene has at least two different alleles (and each allele is in more than 1% of the population).

ABO blood type is a perfect example of a polymorphism, which, you will recall, means there are two or more alleles at a locus. In the ABO blood group there are three alleles (A, B, and O) prevalent in most human populations. There are a few populations in the world which are not polymorphic. The O allele occurs in almost 100% of South American indigenous populations and is remarkably high in aboriginal groups in northern Australia. This is most likely due to genetic drift. When alleles vary geographically, they have a **clinal** distribution. As you can see in the maps below, B is more common in Europe and Asia, and A is found in high numbers in Asia and central Africa.




***Figure* 2.1.8.1** - *Map of blood group A*




***Figure* 2.1.8.2** - *Map of blood group B*




***Figure* 2.1.8.3** - *Map of blood group O*

You can learn more about the ABO blood group from the American Red Cross.

Using polymorphisms: clines and migration

If you know the clinal distribution of a polymorphism, you can use the movement of the alleles as a proxy for the movement of people.

For example, the map below shows the distribution of a group of genes called Haplogroup F (found on the Y chromosome) throughout the Middle East, Europe, and Asia. Presumably the distribution of the genes follow the movement of peoples as they dispersed from India outward.




Figure **2.1.8.4** - *Haplogroup F (Y-DNA)*

Natural selection and polymorphisms

While some clinal distributions are due to migration (gene flow), others have been shaped by natural selection.

Lactose tolerance




Figure **2.1.8.5** - *Worldwide prevalence of lactose intolerance in recent populations.*

The original human condition (found in most southern European, Asian, and African populations) is an inability for adults to digest lactose, the primary sugar found in dairy products. This is known as lactose intolerance and it has a genetic basis. Lactose intolerance can be a particularly unpleasant condition for those who consume dairy products and cannot digest them. Symptoms include diarrhea and intestinal gas.

However, some populations of people have developed an ability to digest lactose.

This is how that came about:

- An individual (or individuals) in the population was born with a genetic mutation that allowed them to digest lactose.
- During the same time span, the population was domesticating dairy-producing animals, collecting milk, and consuming it.
- Those who could digest the lactose in the milk (and therefore had a good food source without the unpleasant side effects) had more children than those who could not digest the lactose.
- Those children inherited the gene for lactose tolerance from their parents and passed it down to their children, etc.

This is **biocultural coevolution** between the domestication of dairy-producing animals in those populations and an increase in the prevalence of the lactose tolerance gene. In other words, the biology of the people was affected by their cultural behaviors and vice versa.

Sickle cell anemia

Sickle cell anemia is a very serious genetic disease that affects the hemoglobin in the blood.




***Figure* 2.1.8.6** - *Sickle cell distribution*

There are two forms of the Hb (hemoglobin) allele: 1) HbA, which is for normal functioning hemoglobin, that which allows red blood cells to carry oxygen efficiently, and 2) HbB (sometimes written as HbS), which is for sickle cell anemia (the hemoglobin and red blood cells do not function normally).

So, as we learned before about alleles and genotypes, when there are two alleles, there are three genotypes. In this case, the genotypes and phenotypes are:

- HbA/HbA = homozygous person has totally normal hemoglobin and their blood transports oxygen perfectly.
- HbA/HbB = heterozygous person has "the sickle cell trait" and therefore is a genetic carrier, but usually has few to no associated medical problems.
- HbB/HbB = homozygous person has sickle cell anemia.

Sickle cell anemia is a good example of the interaction of biology, environment, and culture. It has a fascinating biological interaction with the infectious disease malaria, which in turn is associated with agricultural practices. It has been found that those with sickle cell trait, i.e., people with the HbA/HbB genotype, are less likely to get malaria. Malaria is a disease caused by a parasite, *Plasmodium*. This life-threatening disease is transmitted through the bites of *Anopheles* mosquitos. Symptoms include fever, headache, chills, and vomiting. In severe cases, anemia, respiratory distress, or death may occur. Malaria in sub-Saharan Africa is associated with the spread of slash-and-burn agriculture and irrigation.

To recap:

- those with HbA/HbA easily get malaria,
- those with HbB/HbB get sickle cell anemia,
- those with HbA/HbB have protection from malaria and not have sickle cell anemia,

This is called the **heterozygote advantage**, because the heterozygotes are better off than either one of the homozygotes. It's this advantage that keeps the HbB allele present in certain populations, which is an example of a **balanced polymorphism**.

Important note: Remember that this HbB advantage only exists in malarial environments. It is **environmentally specific**. In an area without malaria, there is no advantage to having the HbB allele and therefore, the frequency of that allele is lower in populations in non-malarial environments.

A little fun with malaria: Malaria the video game

Adaptations and adaptability

Humans have **biological plasticity**, or an ability to adapt biologically to our environment. An **adaptation** is any variation that can increase one's biological fitness in a specific environment; more simply it is the successful interaction of a population with its environment. Adaptations may be biological or cultural in nature. Biological adaptations vary in their length of time, anywhere from a few seconds for a reflex to a lifetime for developmental acclimatization or genetics. The biological changes that occur

within an individual's lifetime are also referred to as **functional adaptations**. What type of adaptation is activated often depends on the severity and duration of **stressors** in the environment. A stressor is anything that disrupts homeostasis, which is a "condition of balance, or stability, within a biological system..." (Jurmain et al 2013: 322). Stressors can be abiotic, e.g., climate or high altitude, biotic, e.g., disease, or social, e.g., war and psychological stress. Cultural adaptations can occur at any time and may be as simple as putting on a coat when it is cold or as complicated as engineering, building, and installing a heating system in a building.

Types of Biological Adaptation

Acclimatization

This form of adaptation can take moments to weeks to occur and is reversible within an individual's lifetime no matter if it occurs when one is a child or an adult.

Short-term acclimatization can occur within seconds of exposure to a stressor. This type of response quickly reverses when the stressor is no longer present. Imagine stepping out of an air-conditioned building or car into a 90 degree day. Your body will quickly begin to perspire in an attempt to cool your body temperature and return to homeostasis. When the temperature declines, so will your perspiration. Tanning is another short-term response, in this case to increased UV-radiation exposure especially during summer months, which can occur within hours. Tans are generally lost during the winter when UV-radiation decreases.

Developmental Acclimatization

Developmental acclimatization occurs during an individual's growth and development. It's also called **ontological** acclimatization or developmental adjustment. Note that these cannot take place once the individual is fully grown. There is usually a "magic time window" of when the acclimatization can occur. This adaptation can take months to years to acquire.

A famous example of this is those who have grown up at high altitude vs. those who have moved to high altitude as adults. Those who were born at high altitude tend to develop larger lung capacities than do those who were not born at high altitude, but moved there later in life. However, developmental adjustment occurs in response to cultural stressors as well. Intentional body deformation has been documented throughout human history. The ancient Maya elite used cradle boards to reshape the skull. Foot binding in China, now an illegal practice, was considered an mark of beauty and enabled girls to find a wealthy spouse.

Genetics

Genetic adaptations can occur when a stressor is constant and lasts for many generations (O'Neil 1998-2013). The presence of the sickle cell allele in some human populations is one example. Keep in mind that genetic adaptations are **environmentally specific**. In other words, while a particular gene may be advantageous to have in one environment (AKA a genetic adaptation), it may be detrimental to have in another environment.

Human genetic adaptations and human variation

Skin color

Click on this link to watch a fantastic video explaining the interplay of skin color, UV, and vitamin D.



Figure 2.1.8.7 - *Inuit women*



***Figure* 2.1.8.8** - *Young Maasai men*

There are two ecological rules, known as Bergmann's rule and Allen's rule, that explain the variation in size and shape of bodies and extremities using latitude and temperature.

Bergmann's rule: Warm-blooded animals tend to have increasing body size with increasing latitude (toward the poles) and decreasing average temperatures.

Allen's rule: A corollary of Bergmann's rule that applies to appendages. Warm-blooded animals tend to have shorter limbs with increasing latitude and decreasing average temperatures

When organisms are more compact, they tend to conserve heat (due to a high mass:surface area ratio). When organisms are more linear, they tend to lose more heat (due to a low mass:surface area ratio).

This has been applied to humans. The idea is that populations toward the pole tend to be shorter and have shorter limbs than do people on the equator.

For example, the Inuit people of Canada (pictured above) tend to be shorter than the Maasai people of Kenya (pictured below):

Race

Technically, a race is a biologically classifiable **subspecies**. So, when we are asking, "Do human races exist?", what we're really asking is, "Are there biologically classifiable subspecies in humans?".

Here's the American Anthropological Association's statement on race

and the American Association of Physical Anthropologists statement on race

What are they saying?

Basically:

- race is an arbitrary categorization, races are not biologically distinct groups (in other words, race is a cultural construct, not a biological one)
- while groups of people who have lived together for a long time may have some alleles in common (for example, those that code for skin color or hair color), there is more genetic variation within races than there is between races
- the concept of race has historically been a tool that some people use to subjugate others

Further explore the concept of race, its history, and human variation.

References

1. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
2. Larsen CS. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.; 2008. 430 p.
3. O'Neil D. Human biological adaptability: an introduction to human responses to common environmental stresses [Internet]. c1998-2013; [cited 2015 June 1]. Available from <anthro.palomar.edu/adapt/default.htm>.

4. Yoshida-Levine B. Human variation and adaptation [Internet]. El Cajon (CA): Grossmont College; c2015 [cited 2015 June 1]. Available from <www.grossmont.edu/people/bonn...daptation.aspx>.

This page titled 2.1.8: Modern Human Variation is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **1.8: Modern Human Variation** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

2.2: Taxonomy

The two classic lines of evidence have been comparative anatomy and fossils. Recently, genetics has almost completely taken over comparative anatomy research, because it's like going to the primary source, the gene is what codes for the anatomy. It used to be that using fossils was the only way to get dates, but recently, the differences between the mutations in the mitochondrial DNA (mtDNA) of a species can be used to calculate the time when that species split off from another species with different mtDNA.

The fossil evidence gives us a picture that looks like this:



Figure* 2.2.1*: *"The Geologic Time Spiral: A Path to the Past" by Joseph Graham, William Newman, and John Stacy (USGS) 2008) (Public Domain)*

The genetic evidence looks like this:




***Figure* 2.2.2 -** *Genealogy of life by genetic distance , "The Tree of Life" by David M. Hillis, Derrick Zwickl, and Robin Gutell, University of Texas © 2003*

Both imply the differences between living forms and the time when they branched off from each other. The genetic evidence is more concrete about living forms, the fossil evidence is more concrete about extinct life forms and gives more accurate dates.

Taxonomy is another way to think of your family tree. Biologists use different methods to make taxonomies, such as cladistics (better science) and evolutionary systematics (includes dates of speciation events) but as an anthropologist, I tend to emphasize the

"genealogy" of classification – how a particular life form fits into my family tree. I try to figure out what part of my DNA I share with that organism, and probably inherited from a common ancestor. With mitochondrial DNA dating, we can get pretty close to calculating when the common ancestor between two life forms lived and when the split started.

For this class you can use most terminology interchangeably: taxon = clade = branch = phylum = group.

The whole family tree metaphor has recently come under attack by the evolutionary implications of processes like lateral gene transfer, and fuzzy hybrids, and we are learning more and more that evolution doesn't always follow a nice clean straight line. As we saw in the section on lateral gene transfer, there have been examples of two species interchanging DNA without having sex. Also, the concept of a species is not as rigid as we'd like to make it. Every once in awhile the normal sterile hybrid offspring of two different species can be fertile (Horse + Donkey = normally sterile mule).

These events are rare, so it's still worth learning classic evolutionary theory, but we can extrapolate from Charles Lyell's principle of Uniformitarianism that small changes can cause big effects over long periods of time. Call me stubborn, but I'm still not ready to give up on the family tree metaphor. LGT is good evidence that evolution doesn't follow the kind of clean branches that Darwin first sketched, but I think we can still go back to trees to save the metaphor and illustrate how life separates over time into distinct species, but can still occasionally come back together to share genetic material. Have you ever seen how when two separate roots or branches grow right next to each other for a long time that they sometimes fuse and become one root or one branch?

Exercise 2.2.1

SKIM THIS PAGE OF TREES OF LIFE

The theme of separation and reconnection plays an important part in the interpretation of our own recent evolutionary history.




***Figure* 2.2.4** - *Reed DL, Smith VS, Hammond SL, Rogers AR, et al. (2004) Genetic Analysis of Lice Supports Direct Contact between Modern and Archaic Humans. PLoS Biol 2(11) (CC BY 4.0)*




***Figure* 2.2.5** - *Human taxonomy, based on Stringer, C. (2012). "What makes a modern human". Nature 485 (7396): 33–35. (CC BY 4.0)*




Figure **2.2.6** - *Extension to 600 kya (Homo sapiens). (CC BY 4.0)*

Note

Another good article about the "tree of life" metaphor

Astrobiology

This class focuses on the planet earth where humans evolved and share ancestry with all known life. But, it's fascinating to think of what life on other planets may be like. There are some serious hypotheses that some evolution may have taken place on other planets and been brought to the earth through meteors

Note

Life in extreme environments.

This page titled 2.2: Taxonomy is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.8: Taxonomy** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.3: Mammals

There are three main branches of mammals alive today, classified mostly on their reproductive system:

- monotremes lay eggs,
- marsupials have pouches, and
- placental mammals keep the kids inside for longer.

Much of what it means to be human is shared with all mammals.

Note

Humans who can wiggle their ears share the same vestigial auricular muscles with other mamas.

Protomammals

Dinosaur-like mammals existed around hundreds of millions of years ago. They evolved into reptiles, mammals, dinosaurs, and birds.

Note

A new protomammal fossil.

Examples of living mammals

The split between the three groups of mammals living today happened while they were dodging huge dinosaurs in the Triassic period (250-200 mya). The three types of mammals:



Monotremes

The monotremes, or protheria, were once a large group of mammals that roamed the world, but today there are only two surviving species: the echidna and the duck-billed platypus. They lay eggs, have poison glands, and have other features we commonly consider reptilian.



***Figure* 2.3.3** - *Duck-billed platypus egg*

Marsupials

Marsupials, also called metatheria, are animals where newborns must crawl to their mother's pouch to nurse until developed.



***Figure* 2.3.5** - *Koala (Wikipedia)*



Figure* 2.3.6 *Possums

Placental Mammals

Placental mammals, also called eutheria, are known for carrying their fetuses inside a protective placenta until birth.

Imagination Questions

Does taxonomy make you feel more part of the animal kingdom? Try the exercises below to broaden your family tree.

1) How does the Linnaean taxonomy of humans compare to other life? For every branch in the Linnaean taxonomy of humans, find another species that fits into the categories above ours, but not those below it. For example, for "Order" you might pick cats, which have the same Kingdom, Phylum, Subphylum, Class, and Infraclass as us, but belong to a different Order.

Kingdom: Animalia

Phylum: Chordata

Subphylum: Vertebrata

Class: Mammalia

Subclass: Theria

Infraclass: Eutheria

Order: Carnivora

2) Take the first five to ten letters of your name and pick an animal that starts with each of those letters. Find the most recent common ancestor by looking up the origin of their common groups, and draw a taxonomy that connects all the animals and gives the dates when they split from each other. For example: Aardvark, Rhinoceros beetle, Newt, Ibis, Elephant. Insects first arose about 400 million years ago (mya), amphibians around 300 mya. Mammals split around 260 mya from the reptiles that became dinosaurs and then birds. The oldest fossil ancestors of Elephants and Aardvarks have been found only about 10 million years apart, about 60 mya and 50 mya, which is around when the first primate ancestors were found too, because of the adaptive radiation of mammals after 65mya. So Aardvarks and Elephants are my closest cousins, and Rhinoceros beetles probably don't get invited to many family functions.




***Figure* 2.3.9**

3) Draw your family tree back to the Big Bang. Follow your maternal line back, skipping generations in powers of ten. Look up what ancestor was around at times below in the column "years ago" and briefly describe your sample "grandmothers".
I calculated the "years ago" column by multiplying the number of generations by the average generation length. I estimated the average time between generations by taking the average age of fertility, from menarche (first menstruation) to menopause (last menstruation)/death.

Power of ten	Generations	Average age of menarche (First menstruation)	Average reproductive lifespan or age of menopause	Average generation length (years)	**Years ago**	**Who was my Great, Great […] Grandma?**	Arnie's example:
1	1	13	50	25	**25**		Mom
2	10	17	40	20	**200**		great, great, great, great, great, great, great, great, great, great, grandmother, probably an Irish peasant
3	100	?	25	18	**1800**		great, [x100], great grandmother, possibly a Celtic peasant
4	1,000	?	?	17	**17,000**		great, [x1000],great grandmother, an anatomically modern *Homo sapiens* hunter-gatherer possibly in the Iberian Peninsula with Maternal Haplogroup H3 (I happen to know this because I had my mitochondrial DNA tested) See image below.
5	10,000	?	?	16	**160,000**		
6	100,000	?	?	15	**1,500,000**		
7	1,000,000	9	?	12	**12,000,000**		
8	10,000,000	N/A		6	**60,000,000**		
9	100,000,000	N/A		4	**400,000,000**		
10	1,000,000,000	N/A		0.7	**700,000,000**		
11	10,000,000,00	N/A		0.1	**1,000,000,000**		

Power of ten	Generations	Average age of menarche (First menstruation)	Average reproductive lifespan or age of menopause	Average generation length (years)	**Years ago**	**Who was my Great, Great […] Grandma?**	Arnie's example:
	0						
12	100,000,000,000	N/A		0.017	**1,700,000,000**		
13	1,000,000,000,000	N/A		0.002	**2,000,000,000**		great, […],great grandmother, an algae-like eukaryote
14	10, 000,000,000,000	N/A		0.00036	**3,600,000,000**		great, […],great grandmother, a cyanobacterium in a stromatolite
15	100,000,000,000,000	N/A		0.000044	**4,400,000,000**		great, […],great grandmother, a strand of self-replicating molecule swirling in primordial ooze
16	etc.				**13,800,000,000**		stardust

Vocabulary

- allopatric
- ancestral
- convergent evolution
- derived
- divergent evolution
- homoplasy
- mammal
- marsupial
- monotremes
- parapatric
- peripatric
- placental
- primitive
- shared

This page titled 2.3: Mammals is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.10: Mammals** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.4: Human Taxonomy

The classic Linnaean taxonomy is basically part of your family tree without drawing in all the branches. Linnaeus was great for the 1700s, but can be very misleading today if you try to cram all the different branches of life into arbitrary horizontal categories like: Kingdom, Phylum, Class, Order, Family, etc. All of these categories are arbitrary and the only way to make them work is through lots of "sub-" and "super-". But, it's still useful to ask why the split between two taxa is being made: What do they have in common? How are they different? Especially as it relates to us. So if I want to compare and contrast myself to a frog, I go up the tree to the place where vertebrates split into amphibians and mammals. I will share all the characteristics of other vertebrates with frogs, and I will be different from frogs in all the ways mammals are. This helps you determine who is a closer relative, and this is useful in many different ways, e.g., if you want to predict how pollution might effect humans, it is usually better to look at mammals rather than amphibians, if you test a drug on pregnant rhesus monkeys, it might act differently on human mothers because we have a different placenta.

Kingdom: Animalia

Phylum: Chordata

Subphylum: Vertebrata

Class: Mammalia

Subclass: Theria

Infraclass: Eutheria

Order: Primates

Suborder: Anthropoidea

Superfamily: Hominoidea

Family: Hominidae

Genus: *Homo*

Species: *sapiens*

[from Dennis O'Neil 2012]

Note

Article on fish with legs, and the evolution of vertebrate limbs.

This page titled 2.4: Human Taxonomy is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.9: Human Taxonomy** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.5: Osteology

Osteology is the study of bones. Osteology is important to studying human variation, and primatology. Paleoanthropology relies on osteology because most fossils come from bones. Forensic anthropology uses osteology to solve crimes. Like most other physical traits, the bones we see are a consequence of genes and environment. There is nothing particularly profound about bones compared to other biological systems, but their durability makes them special for anthropology because they are the main source of data for paleoanthropologists, important to archaeology, and before DNA testing, they were important to the study of human variation.

We tend to think of bones as dead, dry, and brittle, and when you leave them out in the sun for a few years they do get old. Their hardness comes from a calcium-based crystal structure of hydroxyapatite ($Ca_{10}(PO_4)_6(OH)_2$), which interconnects like columns of Lego blocks.

In a biology class you tend to think of a bone as a living organ, like your heart or your lungs, but in anthropology we are used to looking at dead bones, outside of the body, when they are just shells of the functions they had when they supported living organisms.




***Figure* 2.5.2** - *Bone Growth rozwój kos´ci*




***Figure* 2.5.3** - *By OpenStax College - Anatomy & Physiology, Connexions Web site.*

Genetics determines most of what your bones look like, for example, your 23rd chromosomes determine several shapes that are commonly used to say whether someone looks male or female, and when forensic anthropologists use these differences to identify skeletons.




***Figure* 2.5.4** - *By CNX OpenStax [CC BY 4.0 (http://creativecommons.org/licenses/by/4.0)], via Wikimedia Commons*

However, like the rest of your body, the environment also effects your physical structures. The muscle attachments on your bones suggest your activities during your life, and stress, i.e. malnutrition, can be read in cross-sections of your teeth like tree rings. It's important that we have a basic shared vocabulary so that we can compare humans to other vertebrates, to evaluate fossils, and to understand several aspects of human variation. Learn the bones of the human skeleton below:




***Figure* 2.5.5 -** *Mariana Ruiz Villarreal, 2007*

Make sure to learn the scientific name!

Common Name	Scientific Name
skull	cranium
jawbone	mandible
collarbone	clavicle
shoulder blade	scapula
breast bone	sternum
funny bone	humerus
spine	vertebrae
hip(s)	pelvis
wrist	carpals
thigh bone	femur
kneecap	patella
shin bone	tibia
ankle	tarsals

Here's an online practice quiz with more detail then you need for this class, but good to know if you're going on in anything health related.

? Exercise 2.5.1

Skim animal skeletons

Skim Primate Skeletons and More Primate Skeletons

Vocabulary

- carpals
- cervical vertebrae

- clavicle
- coccyx
- cranium
- epiphyseal
- femur
- fibula
- frontal bone
- humerus
- hyoid
- lumbar vertebrae
- mandible
- manubrium
- metacarpals
- metatarsals
- occipital bone
- parietal bone
- patella
- pelvic girdle
- periosteum
- phalanges
- phalanges
- radius
- ribs
- sacrum
- scapula
- sternum
- tarsals
- temporal bone
- thoracic vertebrae
- tibia
- ulna

Imagination Questions

- Go to Balboa Park and visit the San Diego Natural History museum, it's free with a resident ID on the first Tuesday of the month. How are we constrained by our skeleton?
- Compare and contrast primate anatomy. Why is it different? Why is it the same?

Thumbnail: Catacombs of Paris. Image taken in October 2007. (Djtox).

This page titled 2.5: Osteology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **3: Osteology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.6: Macroevolution

Another word for macroevolution is speciation, the production of species, this is the level of evolution that Darwin studied, the kind that occurs over immense periods of time, where small changes accumulate to make life diverge into often drastically different forms. Now that we've introduced Mendel and cellular biology we can explain the two main pieces that Darwin was missing, heredity and variation.

Hox genes

Hox genes are a good example of one mechanism that enables drastic biological changes. Minor variations in the same small set of genes direct a zygote to grow into the shape of a worm, fly, mouse, or a human.

Exercise 2.6.1

- Read Hox gene intro
- Review Micro and Macro Evolution
- Living things share common genes.
- *Read about Human Accelerated Regions
- *Article on the genes for arms and legs coming from fish

Allopatric

“-patric” refers to geography. “allo” is Greek for *other*. Allopatric speciation happens when two populations are separated geographically; if you can't meet, you can't breed; in the absence of gene flow, variations accumulate through mutation and natural selection effects the two populations differently until they become separate species.

Punctuated Equilibrium

Darwin saw evolution as the gradual accumulation of changes, but looking closer at the fossil record, especially shellfish, later paleontologists began to question Darwin's ideas of phyletic gradualism, and proposed the theory of punctuated equilibrium, where the equilibrium of stasis or lack of change, is punctuated or broken by rapid evolutionary change. Remember that we are using the word "rapid" in a geological sense, usually it means hundreds of thousands of years.

Exercise 2.6.2

Skim this intro to punctuated equilibrium

This page titled 2.6: Macroevolution is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.2: Macroevolution** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.7: Species vs. Paleospecies

The definition of a species is something we can demonstrate very clearly: if two individuals can have grandkids, then they're part of the same species. With fossils it's tricky to test this. If you bang two fossils together, they don't make baby fossils. In the next section on paleoanthropology, we will lump fossils into groups and label them with an official-looking genus and species designation written in Latin and in italics, but it's really important for you to remember that this designation is not a concrete fact, it's just a hypothesis that will continue to be tested (and often contested). Understanding that paleospecies are hypotheses will help you to understand the context for many of the debates in paleontology and especially paleoanthropology.

Exercise 2.7.1

Skim species for evolutionary biology

READ EVOLUTIONARY SPECIES VS. CHRONOSPECIES

*Lygers, Zonkeys, Jaglions… skim examples of hybrids

This page titled 2.7: Species vs. Paleospecies is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.3: Species vs. Paleospecies** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.8: Interspecific vs. Intraspecific Variation

When you lump a bunch of fossils together and call them a paleospecies, you are saying that all the differences between the fossils are *intraspecific variation*, variations within a species. If you split all the different looking fossils into separate species then you are seeing *interspecific variation*, variation between species; *intra-* means "among" and *inter-* means "between"

This page titled 2.8: Interspecific vs. Intraspecific Variation is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.4: Interspecific vs. Intraspecific Variation** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.9: Adaptive Radiation

When given the opportunity, a group of species will move into available ecological niches. A classic example is when around 65 million years ago, mammals took on ecological roles formerly held by dinosaurs after the northwest top of the Yucatán Peninsula was hit by a giant asteroid which left the Chicxulub crater, a rain of fire, mega-tsunamis, and a cloud that cooled the Earth for tens of thousands of years.



***Figure* 2.9.1** - *The impact of meteors can change the weather*

When the dust cleared, the surviving mammal species radiated into the ecological niches formally occupied by dinosaurs. One species' misfortune is another's opportunity.

A good article on *dinosaur paleoecology. Morphospace is from *morphology* meaning the study of shape and space implies a landscape, or in this case an ecosystem where a particular shape is more or less adaptive.

This page titled 2.9: Adaptive Radiation is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.6: Adaptive Radiation** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

2.10: Analogy vs. Homology

When I'm swimming in the ocean and see a vertical fin above the water in the distance, I always freak out and think it's a shark even though it always turns out to be dolphins. Dolphins evolved from a four-legged mammal that might have looked something like a hippo, and the dolphin's fin and the shark's fin are coded for by totally different genes, and have totally different evolutionary pathways. Traits that look the same but evolved separately are called *analogies*.

Homologies are traits that share a common evolutionary pathway and the same genes that code for them. The range of morphology of vertebrate forelimbs is incredible, but they all have the same genetic source, and thus share most of the same bones.




***Figure* 2.10.1** - *Homologous vertebrate forelimbs*

When you see two species with similar traits, the first question is are they similar because they had the same ancestors (parallel) or they had different ancestors and evolved in similar environments.




***Figure* 2.10.2** - *Parallel vs. convergent evolution*




***Figure* 2.10.3** - *Convergent evolution for water, ichthyosaur vs dolphin by Sceptic view from Wikimedia Commons (CC BY-SA 4.0)*

https://socialsci.libretexts.org/@go/page/129992

Example 2.10.1:

Convergent evolution: echolocation in bats and dolphins

This page titled 2.10: Analogy vs. Homology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **4.7: Analogy vs. Homology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

CHAPTER OVERVIEW

3: Primates

3.1: Introduction to Primatology
3.2: Primate Classification
3.3: Primate Evolution
3.4: Primate Social Organization
3.5: Non-Human Primates
3.5.1: Classification - Ordering the Natural World
3.5.2: Primate Skeletal Anatomy
3.5.3: Modern Primates
3.5.4: Taxonomy of the Living Primates
3.5.5: Primate Ecology
3.5.6: Overviews of Living Primates (Strepsirhines)
3.5.7: Overviews of Living Primates (Haplorhines)
3.5.8: More Haplorhines (Old World Monkeys)
3.5.9: Even more Haplorhines (The Apes)
3.5.10: Primate Evolution
3.6: Ethology
3.7: Conservation
3.8: Lab 5 and 6- Primates

3: Primates is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

3.1: Introduction to Primatology

We are primates. One way to learn about humans is to study them as a kind of primate. This works especially well to explain how we got the physical structure that we have. It works a little bit to explain a few of our behaviors. It doesn't work at all to explain our culture. The Darwin tubercle is a projection on the helix of the ear resulting from a thickening of the cartilage. The actual size of the tubercle varies. It is one of many vestiges of our primate ancestry.



***Figure* 3.1.1 -** *http://en.Wikipedia.org/wiki/Darwin's_tubercle*

A broad research question in primatology is to compare and contrast primates to other mammals, and then compare and contrast primates to themselves.

Note

KATHRYN ORZECH'S INTRO: WHAT MAKES A PRIMATE A PRIMATE?

Exercise 3.1.1: One goal of primatology is to use it to help us understand ourselves.

REVIEW PRIMATE SKELETONS AND MORE PRIMATE SKELETONS

- Jane Goodall is famous for studying chimpanzees. Watch this JANE GOODALL: WHAT SEPARATES US FROM CHIMPANZEES?
- Skim the first chapter of Augustín Fuentes *Race, Monogamy, and Other Lies They Told You Busting Myths about Human Nature*


This page titled 3.1: Introduction to Primatology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **5.1: Introduction to Primatology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

3.2: Primate Classification

SYSTEMATICS: THE SCIENCE OF CLASSIFICATION

There are two means by which scientists classify organisms, **classic taxonomy** and **cladistics**. Paleoanthropologists are trained in evolutionary theory, and both biologists and paleontologists rely principally upon cladistics. There is definite utility in using a combination of both systems, that is, the binomial nomenclature (genus and species) of classic taxonomy combined with the cladistic arrangement of species in terms of shared characteristics. Classic taxonomy is based on the system begun by John Ray and elaborated by Carolus Linnaeus: kingdom, phylum, class, order, family, genus, species, etc. It classifies organisms based on descent from a common ancestor, using similarities in physical characteristics. We are *Homo sapiens*, as distinct from other members of our genus, such as *Homo neanderthalensis* (i.e., Neandertals). Note that since the "h" in *neanderthalensis* is silent, it is sometimes omitted from the common name. Cladistics refines classic taxonomy by linking organisms, based on the presence or absence of unique characteristics, into "grades." For example, if three species share a suite of characteristics but only two of them have a particular trait that is not present in more distantly related species, those two are more closely related and would be depicted as a separate grade. Figure 2.1 depicts five primate grade.



Cladogram of Primates

"Cladogram of Primates" by Petter Bøckman, CC BY-SA 3.0.


The following terms are used to delineate characteristics in cladistics:

- **Plesiomorphy**—a primitive trait that is present in the ancestor as well as descendent species, for example, **pentadactyly** (five digits) in primates is an ancient trait seen in amphibians and reptiles.
- **Apomorphy**—a derived trait that is not found in the ancestor but is present in descendent species, for example, nails in primates.
- **Autapomorphy**—a unique derived trait present in member species of a particular grade, for example, the lack of a tail in apes.
- **Synapomorphy**—a trait inherited by members of two or more grades from their common ancestor, wherein the trait was an apomorphy, for example, **bipedalism**in the various grades within our tribe, **Hominini**.
- **Homoplasy**—a trait in more than one grade that evolved independently, for example, **brachiation** (swinging by one's arms) in some New World monkeys and apes.

PRIMATE CLASSIFICATION

We are primates, that is, members of the order **Primates** (prī-mā'-tēz). The pie chart in Figure 2.2 shows the various orders of animals within the class **Mammalia**. We are most closely related to **tree shrews** (order: Scandentia) and **colugos** (order: Dermoptera, also known as flying lemurs). Primates are distinguished by a suite of characteristics known as **evolutionary trends** (see table below). However, we do not exhibit all of them to the same degree, and some are absent in certain species or lineages.

For example, **prosimians** retain a claw on the second digit of their feet, whereas **anthropoids** do not (more about the two primate groups later). These trends were first proposed by Napier and Napier (1967) and Le Gros Clark (1959), and more recently primatologists have refined and added to the list.




Orders within the class Mammalia. "Mammal species pie chart" by Aranae is in the public domain.


PRIMATE EVOLUTIONARY TRENDS

- Generalized, unspecialized skeleton:
  - No loss of limb bones from the ancestral condition.
  - Presence of a clavicle that allows greater mobility.
  - Capable of varied movement and locomotion.
- Large, complex brain (relative to body size), especially cerebral cortex.
- Decreased reliance on olfaction:
  - Reduction of snout and olfactory bulb in frontal cortex.
- Increased reliance on vision:
  - Enlarged visual cortex, greater visual acuity, and color vision.
  - Forward-oriented, overlapping fields (**binocular**) of vision, and excellent depth perception.
- **Prehensile** (grasping) hands and feet and **opposable** thumb and big toe.
  - Nails instead of claws.
- Long pre- and post-natal life periods with greater reliance on learning.
- Tendency toward diurnality.


Taxonomic charts of the living primates can be found below. The primates are divided into two major taxonomic groups: strepsirrhines, which retain **primitive**characteristics, such as the lemurs of Madagascar and the bushbabies of Africa, and the more derived haplorrhines, that is, the tarsier, monkeys, and apes. The older terms for the suborders that are still in popular use are **Prosimii** (see figure 2.3) and **Anthropoidea**. However, tarsiers (small, nocturnal prosimian from the islands of the Southeast Asian archipelago) have characteristics of both groups. The strepsirrhine primates have more typical mammalian noses or rhinaria (see Figure 2.4) that are moist and more complex. They have a larger olfactory bulb in the frontal cortex of their brains and scent glands in various locations on their bodies. They use those glands to communicate to other members of their species. We haplorhines have simpler, dry noses and do not smell as good!




Prosimian classification.




Prosimian noses (A through D) and the nose of a New World monkey (E). "Prosimian noses" by Reginald Innes Pocock is in the public domain.



New World anthropoid classification.



Old World monkey classification.




Ape classification.


As we learn more about biochemical and evolutionary relationships among the various groups of primates, primate taxonomy is changing. The **New World monkeys** (see Figure 2.5) have changed substantially in recent years, with the creation of multiple families that were formerly grouped into two or three.

The Old World anthropoids (monkeys and apes) and New World monkeys are also distinguished by our noses. Old World anthropoids have more ovoid, downward-facing nostrils, whereas New World monkeys' nostrils are round and forward-facing.

The Old World monkeys (see figure 2.6) are divided into the **cercopithecines**, with their cheek pouches and more generalized diet, and the leaf-eating **colobines**, with their complex guts. Think baboon (Africa) or snow monkey (Japan) for the former and black-and-white colobus (Africa) or Hanuman langur (primarily Indian subcontinent) for the latter. When asked, most people are more familiar with the cercopithecines, but if they see a picture of a black-and-white colobus (also known as a guereza) leaping through the air with its white mantle of fur and tail flying (or not…see Figure 2.8! I couldn't find an action shot!) or a Hanuman langur sitting on the steps of a temple (OK, a fort . . . see Figure 2.9!) in India, they usually recognize them.



Black-and-white colobus monkey. "Colobus guereza Mantelaffen" by Yoky is licensed under CC BY-SA 3.0.



Hanuman langur. "Langur-Amber Fort" by McKay Savage is licensed under CC BY 2.0.

The taxonomy of the apes (see Figure 2.7) has finally been updated. Until recently, humans were separated from the other great apes at the "family" level. All great apes are too closely related to be separated into different families. The **lesser apes**, i.e. the gibbons and siamangs of Southeast Asia, are still separated into their own family, the Hylobatidae. All of the **great apes** are now in the family **Hominidae**, formerly our exclusive domain. The orangutans come out at the subfamily level, leaving the African great apes in the subfamily **Homininae**. The gorillas have their own tribe, Gorillini (using the genus *Gorilla* to form the name) and if the chimps (genus *Pan*) are taken out of our tribe (**Hominini)**, they are assigned the tribe **Panini**! I did not make that up! Some experts suggest that chimps and humans should be included in the same genus.

APE CHARACTERISTICS

> *"Why are sleeves always too short for me?"*

We apes share a suite of characteristics (in varying degrees), and we humans have radically changed as we abandoned a more typical ape habitat and adapted to a more open terrestrial landscape. The table below lists great ape characteristics.

Great Ape Characteristics

- Relatively large brains.
- Y-5 molar—apes have a characteristic pattern of cusps and fissures on one or more mandibular molars.
- **Honing complex** consisting of large canines that are sharpened (honed) on the first lower premolar, termed a **sectorial premolar**.
- Upright trunk posture.
- Short, shallow, wide rib cage.
- High degree of mobility in joints of shoulders and wrists, termed the "**suspensory hanging adaptation**."
- Long arms and short legs.
- Long, curved hand and foot bones.
- Variable degree of sexual dimorphism (i.e., differences between male and female morphology) in body size.
  - Low in humans, moderate in chimps, high in gorillas and orangutans.
  - Pronounced male prognathism (jutting jaws or muzzle) and large canines, depending on species.
- Long life stages, especially the juvenile dependency period.
- Build nests.
- Capable of learning and using symbols.
- Tool use with some modification (e.g. chimps and orangutans preparing a stick for probing).

The ancestor of the great apes was an arboreal climber. At some point, apes deviated from the more quadrupedal monkey-like morphology, in favor of (1) a more upright, shorter, broader, shallower trunk (just think of our thorax versus a dog's); (2) elongated upper limbs; and (3) more mobile shoulder and wrist joints. While we cannot swing by our arms as well as the lesser apes (brachiation is the technical term), we great apes retain the **suspensory hanging adaptation** and can swing to varying degrees, as long as the tree will support us. Adult male gorillas do not swing because they are too massive!

As hominins, or bipedal apes, came to rely less on an arboreal environment, the bones of our ancestors' hands and some foot bones became shorter and straighter and our legs became longer and more efficient for covering long distances on African landscapes. Beginning with the emergence of our own genus: *Homo* (~2 mya), we became increasingly **encephalized** (i.e. an increase in brain size relative to body size), leaving our fellow great apes behind. While all great apes are **sexually dimorphic** in terms of body size (i.e., males are larger than females), humans are less so and the trend began even prior to our own genus. Depending on which fossil hominins we include in our lineage, male canines were either monomorphic (i.e., male and female canines were the same size) or became less dimorphic over time. This is significant because male canine dimorphism is associated with competition for females, i.e. males bite one another! If we accept, for example, that we are descended from the **ardipiths** (see Chapter 8) that lived over 4 mya, ancestral males did not have large canines. However, there is better evidence that we are descended from the **australopiths** and within that lineage, male canines were initially larger than females' and became smaller over time.

Apes live a relatively long time and consequently all of our life stages are prolonged as well, especially our juvenile dependency period. Of the nonhuman primates, orangutans win. Female orangutans have an interbirth interval (i.e. a period of time in between births) of eight years and juveniles do not even start wandering off until they are seven. That is not good news from a conservation perspective!


All great apes build nests and scientists speculate that our ancestors likely did as well, at least until they left the trees. We also use tools. Chimps are the champs when it comes to tool use, e.g. nut cracking, ant fishing, etc. Orangutans are very adept at tool use as well, but their thumbs are short and more distant from their fingers, so that their opposability is poor. Consequently, they use their mouths to trim sticks and manipulate them for the desired task. In captivity, they also use their mouths to draw and paint. While gorilla tool use had been known from observations of gorillas in captivity, the use of a tool in the wild was finally recorded in 2005 when a female used a stick to test water depth. Of course, we humans are on a whole different level and we can trace the development of hominin technology in the archaeological record to over 3 mya.

We have two language centers in our brain, **Broca's** and **Wernicke's areas** (see Figure 2.10). Broca's area is found in all Old World monkeys and apes; it plays a role in the motor control involved with speech production. As we well know from language studies, all nonhuman great apes are capable of learning and using symbols and have even made up a few. Washoe, the famous chimp that was taught American Sign Language, strung together the signs for "water" and "bird," when she saw a duck for the first

time. Where they fall short is in using syntax; they are poor at correctly stringing symbols together into meaningful sentences. However, Kanzi, the super bonobo at Sue Savage Rumbaugh's lab, does pretty well. Check him out in the video below.



It is pretty convincing that he understands some syntax with following her commands, such as putting the pickles inside the vacuum cleaner versus the vacuum cleaner inside the pickles! Okay, I made that one up but the things she has him do, in order to convince us that he understands, are amusing and amazing! We and our ancestors, beginning with the genus *Homo*, also have Wernicke's area. While a homologous area is present in monkeys and apes, ours has much broader interconnections and is uniquely involved with speech comprehension. At what point our ancestors began to speak is a highly contested topic but the archaeological record provides some clues (see Chapters 23 and 28).



Broca's and Wernicke's areas (left side of brain). "BrocasAreaSmall" by the National Institutes of Health is in the public domain.

Great ape social organization varies by species. The chimps and bonobos (genus: *Pan*) are most like the majority of human traditional societies in that they live in **male philopatric,** multi-male/multi-female communities. **Philopatry** refers to the sex that remains in their **natal group**, i.e. the group into which they were born. This type of social organization is seen in some ripe fruit specialists, such as the chimps and bonobos of Africa and the New World spider monkeys. Since fruit is an ephemeral resource, females cannot cooperatively defend it. Groups of related males cooperatively defend a home range against outsider males. Females emigrate from their natal community and join a different community when they reach sexual maturity. Group members come together intermittently into larger aggregations wherein they may interact. This grouping pattern is termed **fission-fusion**. Since males patrol and protect the area and females are relatively large and powerful and can climb, the danger of predation is relatively low and they therefore have less of a need to congregate. The **mating pattern** is termed **polygynandry**, meaning that males and females are promiscuous and may have multiple sexual partners. Males may attempt to monopolize and coerce females to mate but females are adept at getting around bullying males and may even mate with males outside of their group.

Orangutans are considered to be solitary foragers and their prominent mating pattern is **polygyny** (males have multiple mates), wherein females in a given area usually mate with the resident, large, dominant male. Smaller males who are sexually mature but lacking the pronounced secondary sexual characteristics, such as facial pads (termed flanges) and an enlarged throat sac for loud

calls, may try to forcibly copulate with females. Females are thought to monitor the dominant male's location so that they can stay within calling distance if they are being harassed by subordinate males.


Gorillas live in one-male groups, except for the mountain gorillas, where two males may reside, an older dominant and younger subordinate. Both sexes tend to leave their natal group. Females join a male who may or may not already have other female mates. Thus the mating system is also polygynous. Males defend their females and offspring from outsider males who may be infanticidal. In mountain gorillas, females are thought to prefer groups with two males as they provide better protection for her offspring.

It can thus be seen that there are characteristics of the human sexes in all three great ape **genera** (plural of genus): male defense and mate-guarding, female choice for dominant males and good genes, male philopatry with females maximizing resources for themselves and offspring in chimps and bonobos, etc.

This page titled 3.2: Primate Classification is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.


- **1.2: Primate Classification** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

3.3: Primate Evolution




Anthropoid Evolution by Keenan Taylor.

While we have no primate fossil material prior to the Eocene Epoch, the first primates are thought to have evolved prior to the **Paleocene Epoch** (66–56 mya), possibly as far back as 90 mya, during the Late **Cretaceous Period**. With the extinction of the dinosaurs at the end of the Cretaceous, many terrestrial niches became available and predation pressures were somewhat relaxed. In addition, temperatures were higher than in the recent past (see Figure 3.2) and the angiosperms (flowering plants) were undergoing an **adaptive radiation**, i.e. relatively rapid speciation, and spreading globally. The spread of flowering plants resulted in an adaptive radiation of insect pollinators and herbivores (plant-eaters), as well as insectivorous and herbivorous arboreal vertebrates.



Temperature change over time. “65 Myr Climate Change” by Robert A. Rohde is licensed under CC BY-SA 3.0. Notes: Pal = Paleocene, Eo = Eocene, Ol = Oligocene, Mio = Miocene, Pli = Pliocene, and Plt = Pleistocene



Primate phylogeny. “Primate phylogeny” from “Strepsirrhini” in Wikipedia is licensed CC-BY-SA

The earliest primates likely descended from a small, nocturnal, insectivorous mammal. The **tree shrews** and **colugos** (also known as flying lemurs) are the closest living relatives to primates. The tree shrew is used as a living model for what the earliest primates, or primate predecessors, might have been like. At some point, primates or their ancestors moved into the trees and adapted to an

arboreal environment. Two theories regarding the evolution of some primate characteristics, such as grasping or prehensile hands, forward-oriented eyes, and depth perception, are the Arboreal and Visual Predation Theories. The **Arboreal Theory** posits that primate characteristics, such as grasping hands and feet and the presence of nails instead of claws, are the result of moving into and adapting to an arboreal environment. (Imagine the casualties!) The **Visual Predation Theory** asserts that characteristics that were well-suited to scurrying around in trees and visual features in particular, such as convergent orbits, are adaptations to insect predation. Short of a butterfly net, grasping hands, visual acuity, and depth perception are essential for catching insects, but I guess they would be kind of *handy* for using a butterfly net as well!



Tree shrew. “Tupaia cf javanica 050917 manc” by W. Djatmiko is licensed under CC BY-SA 3.0.



Hands and feet of apes and monkeys. “Hands and Feet of Apes and Monkeys” by Richard Lydekker is in the public domain.

While primates are thought to have evolved in Asia, the majority of the early fossil material is found in North America and Europe, dating to the **Eocene Epoch** (~56–34 mya). The map in Figure 3.6 indicates both living and fossil strepsirrhine sites. They are divided into two superfamilies, **Adapoidea** and **Omomyoidea**. In general, the adapoids were diurnal, lemur-like animals that are

thought to be the ancestors of the strepsirrhine primates, i.e. the lemurs of Madagascar and the lorisids of Africa and Southeast Asia (i.e. bushbabies and pottos of Africa and lorises of Southeast Asia) (see Figure 3.7). The smaller, nocturnal omomyoids are good candidates for the ancestors of modern-day tarsiers. However, due to the early dates for ancestral tarsiers, it is possible that the omomyoids and tarsiers were sister lineages.



Range of living strepsirrhine primates (green) and Eocene-Miocene fossil sites (red). "Extant strepsirrhine range with fossil sites," a derivative work by Maky, is in the public domain.

During the Eocene Epoch, the early strepsirrhine-like primates experienced an adaptive radiation and expanded into numerous niches over a broad geographic area. The northern expansion of early primates into Europe and North America was possible because Eurasia and North America were joined as the large landmass known as **Laurasia** and, as mentioned, it was warm enough for tropical animals to move into northern latitudes. Due to subsequent global cooling, the early primates in North America and Europe eventually went extinct. Strepsirrhine primates spread into Africa after it docked with Laurasia. They are also hypothesized to have "rafted" on floating mats of vegetation to Madagascar, where they evolved into the great diversity of extinct and extant lemur species.



Strepsirrhini. Notes: Top left: ring-tailed lemurs (Madagascar); top right: diademed sifaka (Madagascar); top middle left: aye-aye (Madagascar); top middle right: ruffed lemur (Madagascar); bottom middle left: mouse lemur (Madagascar); bottom middle right: slow loris (Asia); bottom left: slender loris (Asia); bottom right: greater bushbaby (Africa).

By at least the late Eocene, the first anthropoid primates had evolved. There is debate over the origin of the anthropoids, i.e. the ancestor of the monkeys and apes. There are four different theories of our ancestry, each with its share of supporters: (1) adapoid, (2) omomyoid, (3) tarsier, or (4) independent origin as yet undiscovered. Remains of early anthropoids dating to the late Eocene are found in Africa and Asia. A possible **stem** or **basal anthropoid**, meaning the original ancestor of all monkeys and apes, comes from the Shanghuang deposits of China. Termed genus: *Eosimias* (see Figure 3.8), it was as small as the smallest living anthropoid, the pygmy marmoset monkey of South America. While ring-tailed lemurs have striped tails, I do not know of any other striped primates so am not sure why the artist gave them stripes … but it sure is an intriguing little creature! Other late Eocene fossils have been discovered in Myanmar (genus: *Pondaungia*), Thailand (genus: *Siamopithecus*), Libya (genus: *Biretia*), Algeria, and the **Fayum Beds** of Egypt.



Eosimias sinesis. Illustration by Keenan Taylor.

During the **Oligocene Epoch** (~34–23 mya), the anthropoid primates underwent a great adaptive radiation. The richest location for Oligocene anthropoid fossils is the Fayum Beds of Egypt. Oligocene anthropoids are divided into three families: Parapithecidae, Oligopithecidae, and Propliopithecidae, from most primitive to most derived over time. The New World monkeys are thought to have branched off from the parapithecids, with which they share some characteristics. Genus: *Apidium* is a prime contender for a possible ancestor. Once again, a rafting hypothesis is proposed for the migration of that ancestor from Africa to South America.

The ancestors of the Old World monkeys and apes diverged from the family: **Propliopithecidae**. The propliopithecid, *Aegyptopithecus zeuxis* (also known as *Propliopithecus zeuxis*) is thought to be a common ancestor of the ape and Old World monkey lineages (see Figure 3.9). While the earliest anthropoids were more monkey- than ape-like, the apes (or hominoids) were the first to successfully adapt to changing environmental conditions in Africa.



Aegyptopithecus or Propliopithecus zeuxis. "Aegyptopithecus NT" by Nobu Tamura is licensed under CC BY-SA 3.0.

For years, people have asked me, "Barbara, you don't really believe that we came from monkeys, do you?" and I always answered, "No, we came from apes!" However, our common anthropoid ancestor was more monkey- than ape-like…. So, "YEAH, I suppose I do!"

During the **Miocene Epoch** (~23–5.3 mya), the adaptive radiation of the apes or hominoids can be observed in the fossil record. The earliest fossils are from Kenya and Uganda. There were 20 or more genera of apes during the Miocene and they exhibited a wide range of body sizes and adaptive strategies. *Proconsul* is a possible stem ape, dating to ~18 mya (see Figure 3.10 and 3.11). The ancestry of the lesser apes is unclear but they are thought to have branched off 18–16 mya. The great apes diversified and spread from Africa to Asia and Europe. The ancestors of the orangutans, the **sivapithecines**, moved into western and subsequently eastern Asia. Remains in Turkey have been dated to 14 mya. The largest primate that ever lived, i.e. the now extinct genus:

Gigantopithecus (known only from isolated dental and mandibular fragments), also had a sivapithecine ancestry. **Dryopithecine** apes moved into Europe during the late Miocene. Generally referred to as "dental apes," due to the scanty remains of jaws and teeth, that evolutionary side branch eventually went extinct due to global cooling, as with the earlier strepsirrhines in the northern latitudes.




"Proconsul NT" by Nobu Tamura is licensed under CC BY-SA 3.0.

While there were Old World monkeys in the Miocene Epoch, such as genus:*Victoriapithecus* from Kenya, the adaptive radiation of the Old World monkeys lagged behind the hominoids. However, the same environmental conditions that drove most ape genera to extinction in Africa led to an explosion of monkey species. Monkeys could more quickly adapt due to their shorter life stages and greater number of offspring. A baboon can give birth every two years versus four or five years for gorillas and chimps, respectively. While the leaf-eating ancestor of the colobines stayed in the trees, the ancestor of the cercopithecine or cheek pouch monkeys, such as macaques and baboons, adapted to traveling on the ground as well as in the trees. The ability to exploit both arboreal and terrestrial resources expanded their niche and they survived and thrived in Africa and Asia. With only two extant genera, the African colobines did not diversify to the same extent, having been confined to forests. However, the Asian colobines did not experience the same forest loss as their African cousins did and are thus much more diverse. When African forests later expanded, the ancestors of some cercopithecine species, such as the colorful arboreal guenons, went back to the trees.

It has been difficult to trace the origin of the human/chimp/gorilla lineage during the mid-Miocene due to a paucity of fossils from that time and many conflicting viewpoints. Some of the contenders for the stem African great ape are *Nakalipithecus*(10 mya) and *Samburupithecus* (9.5 mya) from Kenya. Other possible ancestors or related species are *Afropithecus* (18–16 mya) and *Nacholapithecus* (15 mya) from Kenya and *Otavipithecus* (13 mya) from Namibia.

The chimp and human lineages are thought to have diverged by the late Miocene. Global cooling in the latter part of the Miocene led to the extinction of all ape genera in northern latitudes. Forest cover in Africa was vastly reduced over time due to climatic fluctuations and while most apes went extinct, the newly emerged hominins thrived. Hominins experienced an adaptive radiation during the Pliocene Epoch (~5.3–2.6 mya), and late in the Pleistocene Epoch (~2.6 mya–11.7 kya) our own species, *Homo sapiens*, evolved (≤200 kya).



Proconsul africanus by Keenan Taylor.

This page titled 3.3: Primate Evolution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **1.3: Primate Evolution** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

3.4: Primate Social Organization

Most primates live in groups. The best explanation for why animals form groups and endure the costs of feeding competition is to minimize the risk of predation. Grouping patterns are tied to diet and the defensibility of resources. Females are out to maximize resources for themselves and their offspring, so as to maximize their reproductive success. If a species eats grass or leaves, it does not make sense to defend those resources. However, there is safety in numbers and those species (especially arboreal species) will normally be found living or foraging in small groups. If a species specializes on ripe fruit, they cannot defend them because of the patchy nature of fruit in geographic space and time. In the case of the few primate ripe fruit specialists, such as chimps and spider monkeys, males defend a home range that contains resources that females need, and thus females are attracted to join them. While orangutans are also preferentially frugivorous, they are solitary due to their large size and strict arboreality, which limits resources to those that are accessible from supporting branches. Finally, if a species can eat a variety of things that come in variable-sized patches, they can band together and defend those resources as they come across them in their daily ranging. In that case, females stay together in their natal group (termed female philopatry) and cooperate in resource defense.



Stump-tailed macaques. "Macaca arctoides" by Frans de Waal is licensed under CC BY 2.5.

Social organization involves several aspects of group life, such as (1) the average numbers of individuals in terms of age and sex; (2) whether group members remain in their natal group at maturity or leave, and hence whether individuals have relatives in the group; (3) whether those animals that join a group in adulthood stay permanently or tend to leave after a period of time; (4) the pattern of interactions between individuals, e.g. whether there is a dominance hierarchy and if so, if an individual's position in the hierarchy is permanent or temporary; and (5) the number of potential mates to which an individual has access. While we tend to categorize species by their grouping pattern or social organization, it is increasingly apparent that there is variability within primate species. Some species share our pattern of living in **multi-male/multi-female** groups. Other categories of primate social organization are **solitary**, **male-female pairs**, and **one-male/multi-female** groups. Interestingly, all of the **mating systems** seen in primates, i.e. **monogamy**, **polygyny** (one male mates with multiple females), **polyandry** (one female mates with multiple males), and **polygynandry** (both males and females are promiscuous), are also seen in humans. Some men and women marry or mate for life; some men have multiple wives or partners, and the same goes for some women. I will discuss each type of social organization and mating pattern seen in the primates, along with example species.

SOLITARY AND DISPERSED POLYGYNY

Except for the orangutans, **solitary foragers** are small nocturnal prosimians that forage primarily for insects and fruit. Examples of solitary foragers are the bushbabies (see Figure 4.2) and pottos of Africa, most of the nocturnal lemurs of Madagascar, and the lorises of Asia. I am going out on a limb (too much?) to suggest that our earliest primate ancestors were as well, since we are thought to be descended from a small, insectivorous, nocturnal mammal. Prosimian solitary foragers either avoid predation by stealth (i.e. the **slow climbers**, such as pottos and slow lorises) or a form of locomotion termed **vertical clinging and leaping** (e.g., bushbabies) that allows for quick getaways. Females usually forage alone and either park their young nearby or leave them in a "nest," such as a tree hole. Sleeping groups may consist of female relatives and their young and/or females, young, and males, depending on the species and female-female tolerance. Male home ranges often overlap multiple female home ranges, and males monitor female sexual cycles by "making the rounds" and monitoring their scent, hence the use of the term "**dispersed polygyny**," i.e. one male and multiple dispersed females. One male may dominate other smaller or less dominant males in an area and may suppress them from breeding, via pheromonal activity.



Bushbaby of Africa. "Bushbabies" by Wegmann is licensed under CC BY-SA 3.0.

As mentioned, orangutans are the odd man out. They are large and arboreal so they do not need to group for protection. They need a lot of resources to support them and at some sites, they suffer periodic food shortages, so that grouping would hinder foraging. Females and their dependent offspring forage together. Females maintain proximity and mate with a dominant male with developed **secondary sexual characteristics**, i.e. large size, a throat sac for loud calls, and facial flanges. Until there is an opportunity for males to acquire females, such as when a large male dies, males stay small and mate opportunistically. Scientists are stymied at how they can delay maturation and then facultatively develop into the larger morph.

TERRITORIAL PAIRS AND MONOGAMY

While a few species of primates are commonly referred to as monogamous, **extrapair copulations** have been observed in every one of them. The last primate to have lost the title of true monogamist was the night monkey of Central and South America. Prior to that revelation, it was always fun to ask my students who the only true monogamous primate species is and see if they answered, "humans."



Gibbon of Southeast Asia. "Gibbon Hoolock de l'ouest" by Programme HURO is licensed under CC BY-SA 3.0.

Monogamy begs the question, "why?" While females may benefit from a monogamous relationship, if their mate supports them or their offspring in some way, it is difficult to understand why males would tie themselves to one mate when mating is not costly for them. There are several theories regarding the adaptive significance of pairing in primates. First is the idea that the female needs help defending a territory in order to obtain enough resources for herself and her offspring. Couples may actively and/or passively defend their territories (hence the more appropriate term "**territorial pair**") via threats, fighting, and/or **duetting**, i.e. calling together to indicate that the territory is occupied by a bonded pair. In the majority of species, males help by carrying offspring. The second theory suggests that monogamy is a way for males to protect their offspring from **infanticide**. In those species that form one-male groups (see next section), when a new male takes over, he may kill nursing infants. Once nursing is interrupted, a female undergoes hormonal changes and may return to **estrus** (fertile period). It is in the new male's best interest to impregnate females as soon as possible, in the "hope" that some of his offspring will make it to the juvenile stage before the next male comes in and wipes out the infants. Why would females mate with a homicidal maniac, you ask? It is not in their best interest to wait to reproduce either. That is the way natural selection works! Those traits that maximize **fitness**, i.e. reproductive success, are favored. In addition, a male offspring that grows up to be infanticidal will be in a better position to reproduce, if he has what it takes to take over a group.

There are territorial paired species within the prosimians (indris and wooly lemurs of Madagascar and the tarsiers of the Southeast Asian archipelago), New World monkeys (night, titi, saki, and some marmoset monkeys of Central and/or South America), and the

lesser apes (gibbons and siamangs of Southeast Asia—see Figure 4.3). We are learning that the lesser apes are much less monogamous than was previously thought. Females of some gibbon species tend toward polyandry and thus males are polygynous, making those species polygynandrous. We are the only great ape to have a *tendency* for monogamy, in that we tend to "fall in love" with one person at a time.

ONE-MALE GROUPS AND POLYGYNY

In some species, one male with one or a few females is the grouping pattern. However in other species (Hamadryas baboons, geladas, mandrills, drills, and some odd-nosed monkeys, such as snub-nosed monkeys), one-male units (OMUs) congregate into larger and larger groupings, in a multi-tiered or nested fashion, depending on their current activity. I will discuss this more complex grouping pattern after the discussion of one-male groups (OMGs). Except for the gorillas, all OMG species are Old World monkeys. The majority of the colobines form OMGs, e.g. African colobus monkeys of the genus: *Colobus* and Asian langurs and leaf monkeys. Cercopithecines of the genus: *Cercopithecus* (commonly known as guenons—see figure 4.4) and patas monkeys (*Erythrocebus*) are also OMG species.



De Brazza's monkey, a type of guenon of Africa. "Lightmatter guenon" by Aaron Logan is licensed under CC BY 1.0.

In the majority of OMG species, females are related but as groups get larger, they split along **matrilines**, meaning that a group of closely related females may splinter when competition increases. In addition, females may move between groups, especially in the colobines. Males fiercely compete for access to groups and infanticide occurs during takeovers. In those species that are seasonal breeders, it is difficult for the male to monitor and mate with all of the females and outsider males may sire some of the offspring. One guy can only do so much and females only have a small window of opportunity.


While the OMG makes sense for the colobines and their high leaf diet, it is not as clear why the more generalist guenons exhibit the same pattern. Like the colobine monkeys of Africa and Asia, it is possible that the ancestor of the extant arboreal guenons never left the trees and thus did not evolve the tendency for a larger grouping pattern in response to terrestrial predators. In addition, if they remained arboreal in relict forests, they may have enjoyed a more stable resource base. They are small- to medium-sized monkeys and thus can subsist on a variety of foods, primarily insects and fruit, both of which are indefensible food items, from a female perspective. Thus while a group is beneficial, it does not need to be large. It may be a bit of an oversimplification that female resources drive primate social organization, but it is a useful model with demonstrated heuristic value.



Hamadryas baboons of East Africa and the Arabian Peninsula. "Hamadryas harems" by Brian Jeffery Beggerly is licensed under CC BY 2.0.

For those species with a nested grouping pattern of OMUs, I will describe the system in Hamadryas baboons (see Figure 4.5) and contrast it with geladas (see Figure 4.7). Both species consist of OMUs that congregate into three larger group levels. For some

strange reason (as if there are not enough terms in a primate course), some primatologists use different terms for the levels in each of the species. The basic unit is the OMU. The next level is termed the clan; it consists of several OMUs, along with bachelor males, and the members tend to forage together (see Figure 4.6). The third level is the band, and that is the result of several clans congregating to forage over a large area. While Hamadryas bands are somewhat stable, gelada bands are not. Finally the troop (Hamadryas) or herd (gelada) is a combination of multiple bands that come together to sleep on cliffs in the mainly treeless regions where both species live, primarily in Ethiopia. Troops consist of hundreds of animals, over 700 in the Hamadryas and slightly fewer in the geladas. This odd grouping pattern is related to their harsh environment. Hamadryas live in subdesert conditions in Ethiopia and the Arabian Peninsula. They are generalists that eat whatever they can find. They fission and fuse (i.e. come together and separate again) into the various grouping levels as resources allow, but predators abound and shelter is scarce, so there is safety in numbers via vigilance. The geladas' situation is a bit different. They live in high-altitude conditions in Ethiopia and eat a lot of grass and grass products, such as seeds and corms. Again, there is safety in numbers but resources are ubiquitous so they spread out a bit and mooch and munch (new foraging category!), i.e. sit-eat-move, along the ground. The strangest aspect of the two species (other than their bizarre faces!) is that Hamadryas are male philopatric and geladas are female philopatric. While the gelada pattern makes sense, considering their relatedness to female-philopatric baboon species, Hamadryas are even more closely related to those baboons yet appear to deviate from the pattern. However, the females do not go far; they transfer at the clan or band level and thus are not far from kin. Thus the real question is why do male Hamadryas stay?



A depiction of a gelada or Hamadryas baboon clan. Note: OMUs =red circles and bachelors = blue Xs.



Grazing geladas. “Geladas” by Alastair Raeis licensed under CC BY-SA 2.0.

While there are regular takeovers in gelada OMUs, as would be expected, they are not as frequent in Hamadryas, primarily due to the facts that males have control over their small group of females and OMUs are surrounded by male relatives. Hamadryas females are usually coerced away from their mothers when they are young and then herded and punished by their new male leader until they learn to obey and not stray. Female geladas have a say in male takeovers; they either side with the resident male and help keep the new male out or they do not and the resident male is on his own. It is interesting that if a new male becomes established in the group, the former male may stay and help defend his offspring from becoming the victims of infanticide, but he can no longer breed.

ONE-FEMALE GROUPS AND POLYANDRY

This type of social organization is seen only in the callitrichids, i.e. the tamarins (see Figure 4.8) and marmosets of Central and South America. Within those groups, there is usually only one breeding female and one or two breeding males. Females gestate as many as five fetuses but on average, only two survive. Hence we talk about “**twinning**” in the callitrichids. Those groups with an extra male have better offspring survival. At birth, the offspring average one-fourth of the female’s weight and thus foraging to support them is a full-time job for the females. The females nurse the young and the males carry and nurture them.



Emperor tamarin. “Emperor Tamarin SF ZOO” by Brocken Inaglory is licensed under CC BY-SA 3.0.

Females pheromonally suppress cycling in their daughters and while sons become fertile, they have no mating options in the group. Mature daughters and sons also help with the care of their younger siblings. Helping behavior, while delaying their **direct fitness** (genes they pass on via reproduction), increases their **inclusive fitness** (genes they share with relatives). Full siblings share half of their genes (the same as between parents and offspring) and half siblings share one-fourth, on average. While the **proximate causation** (current stimulus or condition favoring the behavior, versus **ultimate causation**, i.e. the behavior was favored by natural selection due to its fitness benefits) for older siblings to stay is unclear, it is likely adaptive in some situations to delay reproduction. For example, it may be difficult for young animals to compete for territory or mates, and they are small and inexperienced and thus easy prey.

MULTI-MALE/FEMALE GROUPS AND POLYGYNANDRY

There are two types of multi-male/female groups (MMF). The first is the more common. They are medium to large groups of related females (female philopatric) with a sex ratio skewed in favor of females. Outsider males may congregate in all-male bands. Females and males are promiscuous, the mating pattern known as polygynandry. Many New World monkey species and most of the Old World cercopithecines (e.g. macaques—see Figures 4.1 and 4.9) exhibit this type of social organization. Females cooperate in

resource defense and males may have a more peripheral position within the group, except during the mating season in seasonal breeders. Many semi-terrestrial species exhibit this type of social organization, e.g. baboons (Hamadryas are the exception) and macaques. Terrestriality is associated with larger body and group size, likely for predation avoidance. With more females, come more males and with more males, females can benefit from seasonal breeding. There are enough males to go around and the glut of offspring that are then born reduces the probability that any one of them is eaten; i.e. the dilution effect. In addition, related females help keep watch over the young that then have playmates. Seasonal breeding is tied to environmental conditions, so that females benefit by timing events to coordinate with resource availability.



Long-tailed (also known as crab-eating) macaque. "Ngarai Sianok Sumatran monkey" by Sakurai Midori is licensed under CC BY-SA 3.0.

The second type of MMF is commonly called a **community** social organization. Species that exhibit this type of social organization are **male philopatric** ripe fruit specialists. As mentioned, females cannot defend fruit, so they do not band together into matrilines. Related males defend a territory that contains enough resources to attract females. Females and their offspring forage independently but group members come together periodically into larger aggregations, hence the other term for this type of social organization, **fission-fusion**. New World spider and muriqui monkeys and the chimps and bonobos of Africa (see figure 4.10) are all categorized as community species.



"Bonobo group hug." "Bonobo group hug" by Magnus Manske is licensed under CC BY-SA 2.0

This page titled 3.4: Primate Social Organization is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **1.4: Primate Social Organization** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

CHAPTER OVERVIEW

3.5: Non-Human Primates

Learning Objectives

At the end of this unit, students will be able to:

- understand the classification of living things
- define the parts of Skeletal Anatomy
- identify shared primate traits
- understand the taxonomy of the living primates
- summarize Primate Ecology&Behavior
- identify some of the living primates: Strepsirhines; New World Monkeys; Old World Monkeys; Apes
- describe Fossils & Primate Evolution

In the second unit, students will examine skeletal anatomy, taxonomy, modern non-human primates, primate socioecology, and primate evolution.

3.5.1: Classification - Ordering the Natural World
3.5.2: Primate Skeletal Anatomy
3.5.3: Modern Primates
3.5.4: Taxonomy of the Living Primates
3.5.5: Primate Ecology
3.5.6: Overviews of Living Primates (Strepsirhines)
3.5.7: Overviews of Living Primates (Haplorhines)
3.5.8: More Haplorhines (Old World Monkeys)
3.5.9: Even more Haplorhines (The Apes)
3.5.10: Primate Evolution

Thuymbnail: Lémurien Propithèque de Verreaux. (CC BY 2.0; Jean-Louis Vandevivère).

This page titled 3.5: Non-Human Primates is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

3.5.1: Classification - Ordering the Natural World

As we learned in Unit 1, Carolus Linnaeus developed what is known as the Linnaean system of classification. This basic system is still used today, although it is focused on illustrating evolutionary relationships between organisms. This may sound familiar and it should---think phylogentic tree! Now we take a moment to look in more depth at classification.

Why do we bother with classification systems? Classification systems are important to help scientists organize the massive diversity of the natural world. It's a method that helps scientists from different countries talk about the natural world without confusion. Modern classification is considered a natural system representing real relationships between organisms. One thing to keep in mind when looking at classification systems is that it is basically a hypothesis about evolutionary relationships between species, which is why you may see different schemes as researchers may interpret data differently.

The Linnaean System

This hierarchical system organizes organisms into ever increasingly similar categories. Linnaeus' system has been modified from when he originally published *Systema Natura* in 1735. He published ten editions of the book, adding new levels to the hierarchy.




***Figure* 3.5.1.1** - *Classification diagram*

According to the Linnaean system, human beings are classified as follows (Larsen 2008: 36):

Category Level Common Characteristics

Kingdom	Animalia	mobile multicellular organisms that consume other organisms for food and develop during an embryo stage
Subkingdom	Eumetazoa	all major animals except sponges that contain true tissue layers, organized as germ layers, which develop into organs in humans
Phylum	Vertebrata	group of vertebrate and invertebrate animals that have a notochord, which becomes the vertebral column in humans and other primates
Superclass	Tetrapoda	vertebrate animals with four feet or legs, including fish, amphibians, birds, dinosaurs, and mammals
Class	Mammalia	group of warm-blooded vertebrate animals that produce milk for their young in mammary glands; they have hair or fur and specialized teeth
Subclass	Theria	group of mammals that produce live young without a shelled egg
Order	Primates	group of mammals specialized for life in the trees, with large brains, stereoscopic vision, opposable thumbs, and grasping hands and feet
Suborder	Anthropoidea	group of primates, including monkeys, apes, and humans, but not prosimians; they have long life cycles and are relatively large-bodied
Family	Hominidae	group of anthropoids, including humans and great apes, and human ancestors; they have the largest bodies and brain sizes of all the primates
Genus	*Homo*	group of hominids including modern humans, their direct ancestors, and extinct relatives; they are bipedal and have large brains
Species	*sapiens*	modern and ancestral modern humans; they have culture, use language, and inhabit every continent except Antarctica
Subspecies	*sapiens*	modern humans alone

Rules of Classification

Taxonomy or systematics is the field that establishes the rules of classification. As established by Linnaeus, organisms are first classified by their physical similarities. Remember, today classification is interested in illustrating evolutionary relationships, so it is not enough simply to establish that there are physical similarities; it is important to try and understand how the similarities arose…was it because of a close common ancestor or some other reason.

To begin classification, researchers identify anatomical features that have the same function as that of similar structures in other species. Then they determine whether the similar features are due to a homology or analogy. A **homology** is a trait found in two or more different organisms that was inherited from a common ancestor. The more homologies shared by two organisms, the closer the genetic relationship. The structure of your arm, the wing of a bird, the fin of a whale, and the forelimb of lizard are examples of homologies. Confusing? Think of this as cousins who inherit the same trait from their grandfather (Understanding Evolution). These types of traits are called homologous structures or simply, homologies. Click here for another image.

However, not all similarities are homologies. An octopus has limbs, but the structure is not the same as the tetrapods referred to above. Octopus limbs evolved independently after they split from their shared ancestor with tetrapods. This type of similarity is called an **analogy** or analogous structure. How can analogous structures evolve in separate species? Analogies are a response to similar pressures. The Understanding Evolution website has a short animation that helps to explain this: evolution.berkeley.edu/evolibrary/article/0_0_0/similarity_hs_08.

In order to determine whether a trait is a homology or analogy, researchers have some criteria to help them make the determination:

- Do the bones have the same basic structure?
- Do the bones have the same relationship to other features?
- Is the development of the structure the same?

Researchers also try to distinguish whether or not structures are due to parallel evolution or convergent evolution. **Parallel evolution** refers to situations where organisms that are unrelated or distantly related have similar adaptations because of similar environmental pressures. When parallel evolution occurs “in distantly related [organisms] that are morphologically similar in overall appearance” (Armstrong 2002-2008) it is called **convergent evolution**. An example that will hopefully make this concept more clear comes from the plant world. *Cactaceae* (North American cactuses) and *Euphorbiaceae* (South American euphorbias are distant relatives that both have characteristics we attribute to succulents—thick stems, spines, and adapted to deserts, but they belong to different plant families).



***Figure* 3.5.1.2** - *Cactaceae:* Opuntia bigelovii



***Figure* 3.5.1.3** - *Euphorbiaceae:* Euphorbia aggregata ies

Some scientists use the terms interchangeably, making them even more confusing then they already are; this group of scientists proposes that what we are seeing are two different ends of a continuum. They suggest that we simplify terminology and call it convergence. To help make the distinction between 1) similarity due to a common ancestor and 2) similarities due to distantly related ancestors, some researchers use the term homology to refer to the former condition (#1 above) and **homoplasy** to refer to the latter (#2 above) (Armstrong 2002-2008).

Building Classifications

There are two basic schools of thought in constructing classifications or evolutionary relationships: Evolutionary systematics and cladistics. **Evolutionary systematics** is the traditional approach that relies on analysis of homologous characters (or traits), phylogeny, divergence, and adaptational level to establish evolutionary relationships. It arose out of the Modern Synthesis, in particular the work of Ernst Mayr, G. G. Simpson, and Thomas Cavalier-Smith, and has a time component to it. In more recent times, DNA sequences have been incorporated into the construction of evolutionary relationships.

Cladistics, or phylogenetic systematics, also reconstructs evolutionary relationships but only uses derived homologous characters. **Derived traits** are those characters that are different from the ancestor in form or function. An example of a derived trait for humans is the chin. **Ancestral traits**, which are sometimes referred to as primitive traits, are those characters inherited from the ancestor. For humans, the opposable thumb is an ancestral primate trait. Cladistic evolutionary relationships are presented in a cladogram and do not indicate time. Groups of organisms that share a common ancestor are part of **clade**.

Learn more about Reading Phylogenies and Cladograms from the Understanding Evolution web site.

References

1. Armstrong WP. c.2002-2008. Parallel and convergent evolution [Internet] [cited 2015 Jun 8]. Available from: waynesword.palomar.edu/convevol.htm
2. Carnegia J, Bruno LC, editors. 2001. Classification. Detroit (MI): UXL. p. 132-136. (UXL Complete Life Science Resource; vol. 1).
3. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
4. Larsen CS. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.; 2008. 430 p.
5. Lerner JK, Lerner KL. Cladistics. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, Vol. 2. Farmington Hills (MI): Gale; 2014. p. 964-965
6. Lerner KL, BW Lerner, editors. 2014. Evolution, Convergent. Farmington Hills (MI): Gale. p. 1675-1676. (The Gale Encyclopedia of Science; vol. 3).
7. Lerner KL, BW Lerner, editors. 2014. Evolution, Parallel. Farmington Hills (MI): Gale. p. 1679-1680. (The Gale Encyclopedia of Science; vol. 3).
8. O'Neill D. c.1998-2012. Classification of living things: An introduction to the principles of taxonomy with a focus on human classification categories [Internet] [cited 2015 Jun 8]. Available from: anthro.palomar.edu/animal/default.htm
9. Similarities and differences: understanding homology and analogy [Internet]. Berkeley (CA): University of California Museum of Paleontology; c2015 [cited 2015 Jun 8]. Available from: evolution.berkeley.edu/evolibrary/article/similarity_hs_01
10. Understanding Phylogenies [Internet]. Berkeley (CA): University of California Museum of Paleontology; c2015 [cited 2015 Jun 9]. Available from: evolution.berkeley.edu/evolibrary/article/0_0_0/evo_05

This page titled 3.5.1: Classification - Ordering the Natural World is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.1: Classification - Ordering the Natural World** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.2: Primate Skeletal Anatomy

The study of bones is called **osteology**. It is necessary to have a basic understanding of the skeletal anatomy since the data used in the study of primate and human evolution is based on fossilized bones. The information presented here focuses on the human skeleton. Human beings have an **endoskeleton**, which means that the skeletal structure is on the inside of the body. The endoskeleton grows along with the rest of the body. Endoskeletons are lighter than exoskeletons (skeletal system on the outside of the body), which allowed for the evolution of large vertebrate animals. The endoskeleton is comprised of bone. Bone has three structural parts:

1. Compact bone (periosteum): dense outer layer
2. Spongy bone: light, softer inner layer
3. Marrow: fills the inside core and makes red and white blood cells.

For the skeletal system to work properly it must have ligaments that connect the bone at the joints, muscles connected to the bones with tendons that allow for movement of the bone, and cartilage that fills the spaces between the bones, preventing them from scraping against one another.

Human infants have 270 bones, several of which fuse together during growth so that adult humans have 206 bones. The skeleton is divided into two categories: the axial skeleton and the appendicular skeleton. The **axial skeleton** is comprised of the vertebral column, ribs, sternum, and skull.




***Figure* 3.5.2.1** - *Axial Skeleton*

The human skull has a number of bones. These bones were seperate at birth and then fuse together as an individual ages.




***Figure* 3.5.2.2** - *Human skull bones (simplified)*




***Figure* 3.5.2.3** - *Human skull front bones (simplified)*

The **appendicular skeleton** is composed of the shoulder girdle, upper limbs, pelvic girdle, and the lower limbs.




Figure **3.5.2.4** - *Appendicular Skeleton*

Function of Bone

Bones function to:

1. give structural support to the body
2. protect vital organs
3. provide an environment for the production of red blood cells
4. store minerals like calcium
5. aid in movement of the body

Bone Types

There are four types of bones based on shape.

1. Long bones: long central shaft (diaphysis) and two knobby ends (epiphysis); e.g., femur
2. Short bones: about as long as wide; e.g., patella
3. Flat bones: characterized by being relatively thin and flat, e.g., sternum
4. Irregular bones: odd-shaped bones; e.g., vertebrae

Bones can also be classified based on their origin and texture, but for the purposes of this class you need only be familiar with the types based on shape.

You should spend some time on the website eSkeletons in order to familiarize yourself with the look of the various bones listed above in the different primates.

Types of Bone Cells

1. Osteoblast: forms new bone tissue
2. Osteoclast: absorbs and removes unwanted tissue
3. Osteocyte: maintains bone as living tissue
4. Hematopoietic: produces red blood cells

References

1. Carnagie J, Bruno LC, editors. 2001. Skeletal system. Detroit (MI): UXL. p. 538-542. (UXL Complete Life Science Resource; vol. 3).
2. Martin EL. Skeletal system. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, 5th edition, Vol. 7. Farmington Hills (MI): Gale; 2014. p. 3975-3981.
3. University of Chicago Medicine. c2015. Anatomy of the bone. The University of Chicago Medical Center [Internet] [cited 2015 Aug 2]. Available from: www.uchospitals.edu/online-library/content=P00109

Study Guide




***Figure* 3.5.2.5** - *Human skeleton diagram*

Print this blank image to make your own study guide of the human skeleton

---

This page titled 3.5.2: Primate Skeletal Anatomy is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.2: Primate Skeletal Anatomy** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.3: Modern Primates

Anthropology is the study of humans and the human condition. Humans are primates. We belong to the taxonomic order Primates (pronounced *pry may tees*). This order encompasses humans as well as what we call non-human primates. Non-human primates are our closest biological and evolutionary relatives. So, we study them in anthropology in order to learn more about ourselves.



Figure* 3.5.3.1: Golden Lion Tamarin. Woodland Park Zoo, Seattle, WA 9/30/2011*

General Mammalian Characteristics

The earliest evidence of mammals is from the Mesozoic era, however, there is limited fossil evidence and the fossils that have been found are mouse-like forms with quadrupedal locomotion. Primates evolved from an ancestral mammal during the Cenozoic era and share many characteristics with other mammals. The Cenozoic is the era of the adaptive radiation of mammals with thirty different mammalian orders evolving.

Some of the general characteristics of mammals includes:

- mammary glands: females produce milk to feed young during their immediate post-natal growth period
- hair: mammals have hair (sometimes called fur) that covers all or parts of their body
- jaw: the lower jaw is a single bone
- ear: the middle ear contains three bones: stapes (stirrup), incus (anvil), and malleus (hammer)
- heart: four-chambered heart
- aortic artery: main artery leaving the heart curves to the left to form the aortic arch
- diaphragm: mammals have one
- endothermy: mammals regulate their body temperature to maintain homeostais (a constant body temperature)
- diphyodonty: teeth are replaced only once during lifetime

General primate characteristics

While primates share traits with other mammals, such as mammary glands and endothermy, there are a number of derived traits that all primates share. Because humans share an evolutionary history with non-human primates (we share an ancestral primate), we share certain biological and behavioral traits with them.

Body

Primates have a **flexible** and **generalized** limb structure that's able to move readily in many directions. Compare the flexibility of the spider monkeys on the left with the horse on the right.




***Figure* 3.5.3.2**: *Spider monkeys*



***Figure* 3.5.3.3**: *Horse*

Hands

Primates have **prehensile** hands (and most of them have prehensile feet also). This means that they have the ability to grasp and manipulate objects. Primates have 5 digits (**pentadactyly**) on their hands and feet. Note that a few primates, like spider monkeys, have what are called **vestigial** thumbs. This means that they have either a very small or non-existent external thumb (but in that case, they will still have a small internal thumb bone).



***Figure* 3.5.3.4**: *Primate hands and feet*

Primates have flattened nails instead of claws. Primates also have very sensitive tactile pads on their digits with fingerprints (**dermatoglyphs**). Some monkeys, like woolly monkeys, have similar pads with fingerprints on the ends of their prehensile tails. Therefore, it functions just like another hand. Check out the similarities and differences of the various primate hands in Figure **3.5.3.4**.

Teeth

Primates have **heterodonty**, meaning that they have different teeth that perform different tasks when processing food via biting or chewing. With this, they have an ability to have a more **generalized** diet (compared to a specialist diet), meaning that they have more dietary flexibility. Why would this be a good thing to have?

Compare the teeth of the cayman (a relative of crocodiles and alligators) with the human teeth on the right. While the cayman's teeth do vary in size, they don't vary in structure. However, the human has incisors, canines, premolars, and molars -- all of which perform different food processing tasks.



***Figure* 3.5.3.5**: *Cayman*



***Figure* 3.5.3.6**: *Human jawbone*

Senses

Compared to most other mammals, primates have an increased reliance on **vision** and a decreased reliance on their senses of smell and hearing. Associated with this are their smaller, flattened noses, loss of whiskers, and relatively small, hairless ears. Also associated with this are their forward facing eyes with accompanying **binocular or stereoscopic vision**. This type of vision means that both eyes have nearly the same field of vision with a lot of overlap between them. It provides wonderful depth perception (but a loss of peripheral vision).

Note first how the eyes of the monkey on the left are more front-facing than the eyes of the cow on the right. Also note how flat the monkey's nose is and how small its ears are, when compared to those of the cow.



***Figure* 3.5.3.7**: *Monkey*



***Figure* 3.5.3.8**: *Cow*

Brain

In relation to other mammals, primates have a more expanded and elaborate brain. Compare the complexity of the human brain on the left to the cat brain on the right.




***Figure* 3.5.3.9**: *Human brain*




***Figure* 3.5.3.10**: *Cat brain*

Life history

Life history refers to the pattern that an organism takes from conception to death. When compared to other mammals, primates have:

- longer gestation (pregnancy) periods
- reduced number of offspring (usually one, but some species commonly have twins)
- delayed maturation, with a long infancy and juvenile learning period
- extension of the entire lifespan



Figure* 3.5.3.11: Chimpanzee mother and baby*

Behavior

The behavioral traits we share with other primates include:

- a greater dependence on flexible, learned behavior
- a tendency to live in social groups



Figure* 3.5.3.12: Baboons crossing the road*

References

1. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
2. Larsen CS. Our origins: discovering physical anthropology, 3rd edition. New York (NY): W.W Norton & Company, Inc.; 2014. 478 p.
3. Thies ML. Mammalian characteristics [Internet] [cited 2015 Aug 2]. Available from: http://www.shsu.edu/~bio_mlt/mammals.html

---

This page titled 3.5.3: Modern Primates is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.3: Modern Primates** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.4: Taxonomy of the Living Primates

There are over 200 species and 600+ subspecies of primates living today. They vary in size from the tiny mouse lemur, weighing in at a whopping two ounces to the male silverback gorilla who can weigh up to 400 pounds. Some are vegetarian while others are omnivores. Many live in tropical areas, but there is at least one species that lives in a temperate region where it snows. Some live exclusively in trees and others live exclusively on the ground. The diversity of primates is expressed in the classification system.

Two Systems of Classification

The ***traditional classification*** approach put the primates into two suborders: Prosimii and Anthropoidea. The Prosimii contained lemurs, lorises, and tarsiers while anthropoidea consisted of monkeys and apes. The traditional classification approach used levels of taxonomic complexity to determine which suborder a primate should belong.

Most primatologists today use an approach that uses ancestral-descendent relationships to determine the suborder. In this biochemical ***evolutionary classification*** approach, there are also two suborders, but they are called **Strepsirhini**, which includes lemurs, aye ayes, galagos, and lorises, and **Haplorhini**, which includes tarsiers, monkeys, apes, and humans.




Figure **3.5.4.2** - *Strepsirhini classification*

Strepsirhini, which means "twist-nose," characteristics include:

- snout-like nose
- wet rhinarium (naked, moist area of skin on the snout--think dog nose)
- heightened sense of smell
- prominent whiskers
- divided upper lip tied to the gums by a membrane
- wooly fur
- grooming claw
- dental combs (tooth comb)

Most Strepsirhines are nocturnal and arboreal and have specialized features, such as large eyes with a light reflecting disk, to help with night vision. They were the first suborder of primates to evolve (O'Neil 1998-2012) and are sometimes referred to as the "lower primates."




***Figure* 3.5.4.3 -** *Abbreviated Haplorhini classification*

Haplorhini, which means "simple nose," characteristics include:

- relatively flattened faces (when compared to Strepsirhini)
- forward facing eyes
- postorbital enclosure (bony plate encloses back of eye socket)
- dry noses
- decreased reliance on sense of smell
- larger brains and body size (when compared to Strepsirhini)
- diastema (space between upper lateral incisor and upper canine tooth) except in humans
- increased gestation, maturation, and parental care
- more mutual grooming

References

1. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p.
2. Larsen CS. Our origins: discovering physical anthropology, 3rd edition. New York (NY): W.W Norton & Company, Inc.; 2014. 478 p.
3. O'Neill, D. 2014. The Primates [Internet]. San Marcos (CA): Palomar College; [cited 2015 Jun 22]. Available from: anthro.palomar.edu/primate/Default.htm
4. Suborder Strepsirhini [Internet]. c. 2001-2007. Edinburgh (Scotland): The Natural History Collections of the University of Edinburgh: [cited 2015 Jun 22]. Available from: http://www.nhc.ed.ac.uk/index.php?page=493.504.513

This page titled 3.5.4: Taxonomy of the Living Primates is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.4: Taxonomy of the Living Primates** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.5: Primate Ecology

Primate ecology is the study of the relationship between primates and their environment. Their environment includes not only the physical environment (e.g., trees, water, weather) but also the other animals in the environment, including other non-human primates and even humans. Why should we care about primates and their ecology? Remember that evolution is always environmentally dependent, so the environment is the main thing that we have to consider when examining a primate, its biology, and behavior. Also, keep in mind that primates are not only affected by their environments, they affect their environments as well (by eating plants and insects, dispersing seeds, etc.).



***Figure* 3.5.5.1** - *Monkey feeding*

There are two primary environmental factors that primate ecologists focus on: Food and Predators.

1) **Food**

- **Primatologists** (scientists who study primates) examine the quality, quantity, and distribution of food in a primate's environment.
- Why? Because food = babies. This means that those individuals who acquire more high-quality food more efficiently are likely to have more offspring and therefore will be more evolutionarily successful.
- This pressure is greater for females than for males, because of the biological costs associated with reproduction (**reproductive asymmetry**). Remember that primates spend a very long time as dependent nursing infants, so primate mothers have a heavy burden when caring for them (the only exceptions are some New World monkeys: the father cares for the infants, but the mother still nurses them).

2) **Predators**

- Primatologists also examine the types and distribution of potential predators in a primate's ecosystem.
- Why? Because no one wants to get eaten, right? Those who can most efficiently and effectively avoid predation are likely to have more offspring and therefore will be more evolutionarily successful.




***Figure* 3.5.5.2** - *Jaguar*

Primatologists also look at other environmental factors, such as:

- weather
- the distribution of water
- the distribution of sleeping sites, and
- the primate's relationships with other individuals within their social group, other groups of their own species, and other species (including humans).



***Figure* 3.5.5.3** - *Baboon examining a leaf*

Most primates are **herbivores** (they eat plant foods) and are fairly generalist in their dietary habits. Some primates are **omnivores** and eat lots of things (plant and animal). However, some primates are more specialized.

- **Folivores**: eat mainly leaves.
- **Frugivores**: eat mainly fruit.
- **Insectivores**: eat mainly insects.
- **Gummivores**: eat mainly tree sap.



***Figure* 3.5.5.4** - *Lemur eating fruit*

One of the challenges that primates face in their day-to-day life is a type of **evolutionary arms race** they have with their food. Some food items, like fruit, "want" to be eaten. In other words, it's evolutionarily advantageous for a fruit to be eaten by a primate, who then carries the fruit's seeds far away from the parent plant in its stomach, to deposit them elsewhere. The fruit benefits from this **seed dispersal**, thereby limiting competition between the parent plant and its offspring. This is one of the hypothesized reasons that fruits have high sugar or fat contents, to make them attractive to seed dispersers, such as primates and birds.

However, some food items, like leaves, don't "want" to be eaten. It's not evolutionarily advantageous for a plant to be stripped of its leaves. A leafless tree would not be able to perform photosynthesis and would promptly die. This is why plants have developed certain chemicals, called **secondary compounds** or **antifeedants**, in their leaves. These chemicals make the leaves unpalatable or indigestible to primates. In these cases, scientists have determined that primates have developed behavioral methods to avoid consuming antifeedants. For example, most leaf-eating monkeys prefer to eat young leaves, which are visually identifiable (either lighter in color or a different color) and higher in nutrients and lower in indigestible fiber and antifeedants, than mature leaves.

Not only have primates developed behavioral strategies to meet these challenges, some primates have evolved different types of color vision in order to forage for food and other primates have developed specialized digestive systems to deal with low-quality (high-fiber, high-antifeedant) foods.



***Figure* 3.5.5.5** - *Black and white colobus monkey resting*

Activity budgets

How a primate spends its time is called its **activity budget**. These are comprised of the major categories of:

- foraging/feeding
- social behavior
- locomotion/traveling, and
- resting/sleeping

Why should we care about activity budgets?

Examining a primate's activity budget gives primatologists a good idea of how that particular primate (or primate group) "makes a living". For example, if squirrel monkey (pictured right) Group A spends 30% of its time foraging and squirrel monkey Group B spends 40% of its time foraging, then one can assume that there is something going on that is causing the difference. One can form possible hypotheses such as: Group B is larger than Group A and has more mouths to feed and therefore has to spend more time looking for food to feed them OR Group B has less food in its area than does Group A and so has to spend more time searching for it.

Remember also that if a primate has to spend more time eating or moving around in order to find something to eat, then they are having to spend less time either resting or socializing. Primates only have a finite amount of time in their day, so they have to maximize the use of it the best they can within their environmental parameters.

Activity budgets denote **when** a primate is "making a living":

- **Diurnal**: active during the day. Most primates are diurnal.
- **Nocturnal**: active during the night.
- **Crepuscular**: active during dawn and dusk. Ring-tailed lemurs (pictured below left) have this activity pattern.
- **Cathemeral**: active during irregular periods during day and night. Black lemurs (pictured below right) are one of the few primate species who have this activity pattern.



***Figure* 3.5.5.6** - *Ringtailed lemur*



***Figure* 3.5.5.7** - *Black lemurs*

Note that these activity patterns are not hard and fast rules. As with diet, there is a lot of variation in how flexible a particular species (or group within a species) is when it comes to their activity patterns. For example, although traditionally thought of as strictly nocturnal, the activity patterns of owl monkeys have been found to be highly sensitive to environmental factors such as temperature and amount of moonlight.

Food and Feeding Competition

One of the main factors that affects a primate "making a living" is **feeding competition** (competing with others for food).

There are two groups that a primate competes with;

- the other primates in its social group (AKA **within-group competition**)
- primates (of the same species) in other groups (AKA **between-group competition**)



Figure 3.5.5.8 - *Macaques fighting*

There are also two types of competition. The type of competition prevalent in a situation depends on the quality and quantity of food available in the environment.

- **Scramble competition**: happens when there is a lot of low quality food. This is more common for leaf-eating primates, because trees tend to have large quantities of edible leaves (although some types, such as young leaves, may be preferable to others). It's an indirect competition in which whoever finds food faster or eats faster gets more food than do other individuals.
- **Contest competition**: happens when there is a small amount of high quality food. This is more common for fruit-eating primates, because a fruiting tree has a limited amount of high-quality fruit. It's a direct competition, where certain individuals (stronger, higher social status) get more food than do other individuals via squabbling or fighting.

In total, one can have:

- within-group scramble competition
- within-group contest competition
- between-group scramble competition, and
- between-group contest competition.

Feeding competition can be severe and have serious effects on the health, well-being, and reproductive capacities of primates. In other words, it has a direct effect on their evolutionary fitness. As expected, primates demonstrate a lot of behavioral strategies to deal with the feeding competition they face. For example, a group with high scramble competition may spread out more when feeding so as to access more resources OR in a group with high contest competition, some individuals may form coalitions in order to "gang up on" other primates to take their food.

Spatial use: home ranges and territories

The area that a primate uses is its **home range**. The part of the home range that's used most often is the **core area**. Some species, such as snub-nosed monkeys, have very large home ranges (32 sq. km) and other species, such as the pygmy marmoset, have very small ones (0.003 sq. km). A primate's home range has to contain all of the resources it needs in order to survive (food, water, sleeping sites, etc.). The home ranges of different primate groups often overlap, sometimes only a little, but sometimes a lot. If the home range is physically defended, it's a **territory**. Chimpanzees are famous for their territorial conflicts, which are so organized and violent that scientists have begun calling it "warfare".

Primate Behavior

As noted in a previous section (The Modern Non-Human Primates), one of the fundamental traits that primates share is that we are **social** creatures. All primates share some form of social structure with others of their own species, whether that be in a large permanent social group or as an individual who has repeated short-term interactions with others. The social relationships that primates have with other members of their own species have a huge impact on the individuals involved and are incredibly

important to their health, well-being, and reproductive success. Therefore, the study of primate social behavior has been a focus of primatologists for the last 50 years.

Primate behavior studies come in three formats:



***Figure* 3.5.5.9** - *Chimpanzees in captivity*

- **Captive**: The animals are in captivity. Variables are easy to control in these situations (for example, the number of individuals in a group or food availability) and the primates are easier to observe at close-range. However, because of the artificiality of the situation, the primates may not be exhibiting normal behavior.
- **Semi-captive (AKA semi-free-ranging)**: The animals are captive, but in a very large area like an island or a fenced-in compound. Variables are still relatively easy to control and the primates are easier to observe (than in the wild). The primates tend to exhibit more natural behavior patterns in this type of setting than when compared to a completely captive situation.
- **Free-ranging**: The animals are living in the wild, in their natural environment. This is the most logistically difficult type of primate study. The animals are more difficult to observe (often requiring a **habituation** period to get them accustomed to the presence of observers). However, the primates are most likely to perform their normal array of behaviors in this type of study.

Social structure: the "whys" and the "hows"

The basic premise of why primates have certain social behaviors is due to **reproductive asymmetry**. Remember (from "Food", on this page) that females are under a lot more pressure than males to forage effectively because of the biological pressures they're under (due to the burden of reproduction). It is precisely this inequality that causes the sexes to have different evolutionary priorities:

- For **females**, their priority is **food**. Females can only reproduce at a certain rate, due to the length of time it takes to be pregnant, raise an infant to independence, and have their bodies recover to do it all over again. Access to lots of good quality food through little effort is the key to accomplishing this task.
- For **males**, their priority is finding **mates**. Males can reproduce at a much faster rate than can females. The only real constraint on their reproductive rate is how frequently they can get fertile females to mate with them.

So, **primate social structure** works like this:

- male distribution follows how the females distribute themselves, and
- the females distribute themselves depending on how the food is laid out in their environment.

Social groups have two **immigration/emigration** patterns:



***Figure* 3.5.5.10** - *Baboon social group*

Female philopatry:

- Females do not emigrate from their birth group at sexual maturity.
- Males usually emigrate from their birth group at sexual maturity.

- Females form the core of the group, are biologically related, and have tight social bonds (often exhibited through social cooperation and grooming: seen in these baboons, pictured right). Males do not have much in the way of positive social relationships .
- This is the most common form of social system in primates.

Male philopatry:

- Males do not emigrate from their birth group at sexual maturity.
- Females usually emigrate from their birth group at sexual maturity.
- Males form the core of the group, are related. and have tight social bonds. Females do not have much in the way of positive social relationships.
- This type of social system tends to happen when resources are widely dispersed, so females are widespread and difficult for the males to consistently access (as in the spider monkey, pictured right).

Social strategies



***Figure* 3.5.5.11**

In order to achieve their priorities, male and females must utilize certain social strategies to gain success.

Dominance

- As with chickens, primates have a "pecking order" or **dominance hierarchy** in their social groups.
- One's position in the dominance hierachy often gives them access to preferred resources, including not only food and mates, but other resources such as sleeping sites and water.
- Sometimes dominance hierarchies are determined through fighting, but more often they are sorted out through a series of aggressive/submissive non-contact interactions, such as approach/avoidance, facial expressions, and body postures. Primates try to avoid direct aggressive contact in an effort to avoid risk of bodily harm.



***Figure* 3.5.5.12** - *Macaques grooming*

Cooperation

- Dominance hierarchies aren't completely linear. One's rank in the hierarchy often depends on who they can get to cooperate with them during conflicts. For example, Monkey 2 may be submissive to Monkey 1 when alone, but when her buddy Monkey 3 is around, the two of them cooperate and chase Monkey 1 away from food together. Therefore, Monkey 2's position in the dominance hierarchy is situationally dependent.
- These cooperative relationships are usually between relatives. In male philopatric groups, they're usually brothers. In female philopatric groups, it's often mother-offspring or siblings.
- The relationships are cultivated through **affiliative** behaviors, such as play, grooming, and other forms of body contact (e.g., hugging). So, primates practice *You scratch my back, I'll scratch yours* both literally and figuratively.

VIDEO: How bonobos use play as a tool

Why be in a social group?



Figure **3.5.5.13**

So, why would a primate choose to be in a social group if it is just one big mess of food competition, mate competition, and political posturing? Well, because there are benefits, of course!

Living in a social group:

- provides access to **mates**. While there is, of course, competition for mates within the group, a social group ensures that there are at least mates available to fight over.
- provides more eyes looking for **food** and more brains remembering where food is in the ecosystem
- provides **anti-predator defense**

- There are more eyes on the lookout for predators.

- An additional anti-predator benefit is known as "The Selfish Herd" (AKA safety in numbers). It basically comes down to "you don't have to run faster than the predator, just faster than the other guy".

- Some social groups will "mob" predators and drive them away. This only works on smaller predators, especially those who rely on surprise attacks (such as the Harpy Eagle, pictured right).

Types of primate social groups

Solitary:



Figure **3.5.5.14** - *Galago*

- Males are alone most of the time, except when seeking out mates. Females live with their dependent offspring.
- One male will have territorial overlap with several females.
- Some prosimians have this social system (like the galago, pictured right).

Monogamy:



Figure **3.5.5.15** - *Titi monkey*

- A social group is comprised of a male, a female, and their dependent offspring.
- This is also called "pair bonded".
- This is common in New World monkeys (for example, titi monkeys, pictured right) and the "lesser apes" (siamangs and gibbons).

Single-male multifemale:



Figure **3.5.5.16** - *Diana monkey*

- A social group is comprised of one adult male, several (to many) adult females, and their dependent offspring.
- Males that don't belong to a social group may either live alone or in multimale (bachelor) groups that have no females.
- It's a difficult life for the resident male in these social groups, because not only did he have to fight his way into the group (either driving the previous resident male out or stealing females from another male), he has to continually fight to keep other males out of his group, while keeping the females in the group.
- When males take over groups, some may commit infanticide (killing the previous resident male's infants).
- This is a common social group in Old World monkeys (like the Diana monkey, pictured right)

Multimale-multifemale



Figure **3.5.5.17** - *Hanuman langur*

- A social group is comprised of more than one adult male, more than one adult female, and their dependent offspring
- In these groups, in order to have preferential access to females, males will form dominance hierarchies or develop biological characteristics that the females find attractive. For example, during the mating season, squirrel monkey males begin storing large amounts of water and fat on their bodies.
- Some species (such as Hanuman langurs - pictured right) have both single-male and multimale multifemale social groups in the same population.

Fission-fusion



Figure **3.5.5.18** - *Bonobo*

- In this situation, individuals belong to a large group called a community. Each community has a home range and consistent community membership. Within the community, individuals form temporary foraging groups called parties. Parties have an unpredictable membership. So, on any given day (or even part of a day), one cannot predict who will be foraging with whom.
- Females always travel with their dependent offspring.
- Sometimes males form partnerships and forage together.
- This type of social group is thought to have been an adaptation to environments with patchy, unpredictable fruit availability.
- It's found in chimpanzees, bonobos (pictured right), and spider monkeys.

Single-female multimale



Figure **3.5.5.19** - *Tamarins*

- A social group is comprised of one adult female, more than one adult male, and their dependent offspring.
- This is rare in primates.
- Found in marmosets and tamarins (pictured right), this type of social system is thought to be an adaptation to the twinning (giving birth to twins) that it common to these small primates. The males carry the dependent infants.

This page titled 3.5.5: Primate Ecology is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.5: Primate Ecology** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.6: Overviews of Living Primates (Strepsirhines)



***Figure* 3.5.6.1** - *White-headed Lemur, Masoala National Park, Madagascar*

The following pages provide a very brief overview of some of the primates living today. It is not in any way a complete list of the hundreds of primate species. It is intended to give you a basic idea of the behaviors of some of the Strepsirhines and Haplorhines. If you would like to learn more about the primates mentioned on the overview pages, check out the reference section, which has links to sources with more details.

The Strepsirhines

Lemuroidea




***Figure* 3.5.6.2** - *Distribution of lemurs (modified from original map from http://d-maps.com/carte.php?num_car=742&lang=en)*



***Figure* 3.5.6.3** - Varecia variegata *foot with toilet-claw*

Lemurs, which means “ghosts,” are small primitive primates ranging in size from about two ounces for the lesser mouse lemur to about 13 pounds for the Indri. They have a wet rhinarium, which indicates that the sense of smell is important. They have a dental comb and a “toilet claw” (grooming claw located on the second toe of the hind foot) both of which are used for grooming; the toilet claw is also used while eating. With the exception of the mouse lemur, lemurs are hindlimb-dominated quadrupedalists who are found equally in trees and on the ground. Many of the species, including sportive and ring-tailed lemurs, scent mark their territories.

There are two superfamilies of Strepsirhines: 1) Lemuroidea, which includes the families Cheirogaleidae, Daubentoniidae, Indriidae, Lemuridae, and Leipilemuridae, and 2) Lorisidea, which include the families Lorisidae and Galagidae.

Mouse and Dwarf Lemurs



Figure **3.5.6.4** - *Gray mouse lemur (Microcebus murinus)*

Cheirogaleidae, or mouse and dwarf lemurs, are nocturnal, arboreal insectivores that move around by hopping on their hind legs and leaping. They live in the forests of Madagascar. There are five genera: 1) *Cheirogaleus*, e.g., fat-tailed dwarf lemur and furry-eared dwarf lemur, 2) *Microcebus*, e.g., golden-brown mouse lemur and pygmy mouse lemur, 3) *Mirza*, i.e., Coquerel's giant mouse lemur and northern giant mouse lemur, 4) *Allocebus*, i.e., hairy-eared dwarf lemur, and 5) *Phaner*, e.g., pale fork-eared lemur and Amber Mountain fork-eared lemur, and twenty-one species. Cheirogaleus , Phaner , and Mirza live in monogamous family groups while Microcebus lives in multi-male, multi-female groups. Females bear two to three offspring at a time. Gestation periods vary from 60 days for the smallest species to 89 days for the larger species. Females have three nipples, which is a trait not shared by any of the other lemurs. Mouse lemurs' tails are at least as long as their bodies. There is some variety in diet although most dwarf and mouse lemurs are omnivorous, but some species specialize, such as Phaner that eats primarily gums.

True Lemurs



Figure **3.5.6.5** - *Eulemur mongoz*

Lemuridae, or true lemurs, are diurnal primates about the size of the typical domestic cat. They have long, heavily furred tails, tufts of fur at the tips of their ears, and a long snout. Their eyes are smaller in comparison to other Strepsirhines. Found on Madagascar and the Comoro Islands, true lemurs spend a lot of time of the ground, but general rest and sleep in trees. They are vertical climbers and leapers who eat leaves, fruits, and some insects. Several of the true lemur species live in large multi-male, multi-female groups where the females dominate the males. Females also tend to defend the group even though males are generally larger than females (in other words true lemurs exhibit sexual dimorphism). Other true lemurs are solitary. There are twenty-three species of true lemurs and five extant genera: 1) *Lemur*, i.e, ring-tailed lemur, 2) *Eulemur*, e.g., brown lemurs and mongoose lemurs, 3) *Varecia*, e.g., red ruffed lemur, 4) *Hapalemur*, e.g., golden bamboo lemur, and 5) *Prolemur*, i.e., greater bamboo lemur.

Sportive lemurs



Figure **3.5.6.6** - *Red-tailed Sportive Lemur, Kirindy, Madagascar.*

"The sportive lemur (*Lepilemur spp.*) received its name because, when threatened, it turns and raises its arms as if preparing to box" (Blashfield 2004: 2548). There are several species of sportive lemurs, including the red-tailed sportive lemur, the white-footed sportive lemur, and the northern sportive lemur. They live in the dry, evergreen forests of Madagascar. Sportive lemurs are also called weasel lemurs and weigh from 1.1 to 2.2 lbs. They have short, pointed head and large round ears. They also only have 32 teeth as compared to the true lemurs that have 36 teeth. Sportive lemurs are primarily folivores, but also eat bark, flowers, and fruit. Sportive lemurs are the only lemurs to exhibit cecotrophy [SEE cuh troh fee]. This means that they eat their own waste material that contains partially plant material in order to get as much nutrition as possible. Sportive lemurs are nocturnal vertical climbers and leapers. Males usually live alone and defend their territory from others using vocalizations, chasing, and, if necessary, fighting. Males can mate with multiple females during the mating season. Females bear one young after a gestation period of about four and a half months. Infants stay with the mother for about one year.

Leaping Lemurs



Figure **3.5.6.7** - *Indri female and infant.*

Indriids or leaping lemurs include sifakas (*Propithecus*), Indiri (*Indri*), and wooly lemurs (*Avahi*). They live in the rainforests, deciduous forests, and evergreen forests of Madagascar. The Indri is the largest of not only the Indriids, but the Strepsirhini in general. There is some variation in the appearance of the various Indriids. Sifakas do not have fur on their face and have sparse hair on their underside. Indris have silky fur and a stumpy tail and woolly lemurs have thick, woolly fur (as indicated by their name) and ears that are barely visible in their fur. One of the characteristics that unite the Indriids is the presence of 30 teeth instead of 36 like true lemurs. With the exception of the avahis, Indriids are diurnal. All of them are vertical climbers and leapers, leaping from tree to tree and leaping when on the ground (watch this short video to see some sifakas in (loco)motion: youtu.be/8RiZElNNpAY). In fact, Indris can leap up to 10 ft in a single bound. They live in small groups, which may be multi-male, multi-female or solitary/all male. Males, which are dominant, are responsible for defense of territories and indirectly contribute to infant care. Females bear one offspring at a time after a gestation period of 130-150 days. Indriids are vegetarian, focusing on leaves, buds, fruits, flowers, and bark.

Aye-aye



Figure **3.5.6.8** - *Aye-aye.*

Named after the sound it makes, the aye-aye has only 18 teeth; its incisors grow continually, and it lacks a toilet claw. It has large eyes, making them optimal for its nocturnal lifeway. Aye-ayes average about 3 ft in length. It has large, pointed ears and a pointed nose and chin. There is an extra-long middle finger with a hooked claw that is used when hunting for insects, to probe and scrape for insect larvae, and to clean its fur (watch this video from the Duke Lemur Center to see the aye-aye in action: youtu.be/HB_fQ1Rg_K4). While aye-ayes are classified as insectivores, they also eat parts of Ramy and palm nuts, fungi, and fruit. These solitary quadrupeds breed throughout the year. Gestation is about 170 days and females usually have one infant at a time, normally with over a year between births. Female ranges do not overlap, but male ranges may, which occasionally results in conflict between the two males.

Lorisoidea



Figure **3.5.6.9** - *Distribution of lorises, pottos, and galagos. Original blank map from http://d-maps.com/carte.php?num_car=3266&lang=en modified from world map to show range of lorises.*



Figure **3.5.6.10** - *Sri Lankan Slender Loris.*

Lorises and Pottos

There are five genera (*Arctocebus*, *Perodicticus*, *Pseudopotto*, *Loris*, and *Nycticebus*) and about fifteen species of Lorisidae (pottos and lorises) living in sub-Saharan Africa, southeast Asia, southwest China, southern India, and Sir Lanka. These small primates weigh in from nine ounces to two pounds. To survive in areas where monkeys live, these lorisids are nocturnal so that they do not have to compete directly with monkeys. Like other nocturnal Strepsirhines, they have a reflective layer at the back of their larger eyes to help them see in the dark. Also like many of the other Strepsirhines, lorisids have a wet rhinarium, albeit short, and dental comb. All of their digits have claws, but the one on the second toe is longer, thereby qualifying as a toilet claw. They have an opposable first toe that allows them to grasp branches securely; in fact, lorisids can be found feeding while hanging upside down

from tree branches! These arboreal primates prefer to forage for food on their own. While each species may have a food preference, in general lorisids are omnivores and eat insects, fruit, gum, small vertebrates, and birds' eggs. Many lorisids exhibit a form of quadrupedal locomotion called cryptic locomotion, which means that they move slowly and deliberately. This form of locomotion helps to keep them hidden from predators; they can stay still for hours at a time. In an unusual Strepsirhine predator-avoidance adaptation, Perodicticus potto has three or four skin-covered bony spines behind its neck. This potto will tuck its head and turn the spines toward attackers in an effort to ward off a predator. After one of the longest gestation periods, 193 days, females usually give birth to two offspring who are fully matured after ten to eighteen months.



***Figure* 3.5.6.11 -** *Galago granti*

Galagos

Galagos used to be classified as a subfamily of Lorisidae, but are now classified as a separate family, Galagidae. Generally referred to as bushbabies because their cry sounds like a mewling infant, galagos are small vertical climbers and leapers who can jump up to 15 feet, using their busy tails as balance. Their arms are longer than most primates in proportion to other primates. They have soft pads on the tips of their fingers that help them cling to branches. They can move quickly, snatching flying insects in midair, which they stir up using vocalizations. Galago ears "are so important that they can fold, like an accordion, when the little animal is moving quickly through prickly or otherwise dangerous leaves" (Blashfield 2014: 2608). After a night of hunting, galagos will sleep together in a communal sleeping hole in a tree. Like pottos and lorises, galago females usually give birth to two offspring who she carries around in her mouth for about two weeks after birth. A female galago and her offspring defend a territory from foreign females. Males also defend a territory that encompasses several female territories.

References

1. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Lorises and Pottos: Lorisidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2007. p.428-435
2. Blashfield JF. Lemurs. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, 5thedition, Vol. 5. Farmington Hills (MI): Gale; 2014. p. 2457-2550.
3. Blashfield JF. Lorises. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, 5thedition, Vol. 5. Farmington Hills (MI): Gale; 2014. p. 2606-2609.
4. Dewey T. 2008. Indriidae: indris, sifakas, and relatives. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 26]. Available from: animaldiversity.org/accounts/Indriidae/
5. Dewey T, Myers P. 2009. Cheirogaleidae: dwarf lemurs and mouse lemurs. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 26]. Available from: animaldiversity.org/accounts/Cheirogaleidae/
6. Duke Lemur Center [Internet]. Aye-aye. [cited 2015 Jun 26]. Available from: http://lemur.duke.edu/discover/meet-the-lemurs/aye-aye/
7. Duke Lemur Center [Internet]. Slow loris. [cited 2015 Jun 27]. Available from: Duke Lemur Center [Internet]. Aye-aye. [cited 2015 Jun 26]. Available from: http://lemur.duke.edu/discover/meet-the-lemurs/aye-aye/
8. Jurmain R, Kilgore L, Trevathan W. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning; 2013. 437 p. Larsen CS. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.; 2008. 430 p.
9. Myers P. 2000. Daubentoniidae: aya-aye. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 26]. Available from: animaldiversity.org/accounts/Daubentoniidae/

10. Myers P. 2000. Galagidae: bushbabies and galagos. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 27]. Available from: animaldiversity.org/accounts/Galagidae/
11. Myers P. 2000. Lemuridae: true lemurs. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 26]. Available from: animaldiversity.org/accounts/Lemuridae/
12. Allen CJ, Evan AV, McDade MC, Schlager N, Mertz LA, Harris MS, editors. 2007. Sportive lemurs: Lepilemuridae. Detroit (MI): UXL. p. 466-474. (Grizmek's Student Animal Life Resource; vol. 16).
13. Swartz JH. Lemurs. In: Birx HJ, editor. Encyclopedia of Anthropology, Vol. 4. Thousand Oaks (CA): SAGE Reference; 2006. p. 1459-1461.

This page titled 3.5.6: Overviews of Living Primates (Strepsirhines) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.6: Overviews of Living Primates (Strepsirhines)** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.7: Overviews of Living Primates (Haplorhines)

As you no doubt recall, Haplorhini means "simple nose." The shared characteristics of tarsiers, New World monkeys, Old World monkeys, and apes include:

- relatively flattened faces (when compared to Strepsirhini)
- forward facing eyes
- postorbital enclosure (bony plate encloses back of eye socket)
- dry noses
- decreased reliance on sense of smell
- larger brains and body size (when compared to Strepsirhini)
- diastema (space between upper lateral incisor and upper canine tooth) except in humans
- increased gestation, maturation, and parental care
- more mutual grooming




Figure* 3.5.7.1*: Hamadryas Baboon *(Papio hamadryas)* at the Leipzig Zoo.

Tarsiers

The three genera of Tarsiidae;

1. *Carlito*,
2. *Cephalopachus*, and
3. *Tarsius*,

Eighteen species living among the islands of southeast Asia, including Borneo, Sumatra, and some of the Philippines. Tarsiers are arboreal vertical climbers and leapers and can be (not so easily) found in dense rainforest. They are occasionally found in non-forested habitats only if there are enough vertical surfaces for leaping and clinging. They rarely spend time on the ground. Tarsiers are sometimes called “living fossils” because of their close resemblance to fossil primates dated to about 40 million years ago. They are small primates, weighing about 4 ounces and roughly 5 inches in length, although with their tail, which may have a tuft of fur on the end, they almost double in length. Like some Strepsirhines, tarsiers have toilet claws and 36 teeth. They are also nocturnal or crepuscular and have huge eyes (just one of their eyes weighs as much as its brain!). In fact, under the traditional classification scheme, tarsiers were classified as prosimians; however, in the new classification system, tarsiers are Haplorhines because they do not have a wet rhinarium. Tarsiers can turn their heads 180 degrees and have the longest hind limb to forelimb proportion of any mammal. Because tarsiers are difficult to locate in the wild, it appears that reproduction is seasonal. Females give birth to a single offspring who weigh from 25 to 30% of the mother’s body weight. When infants are born they can immediately cling and climb. The mother is the primary caretaker, although dominant males and subadult females may help. Most tarsiers spend their waking hours alone hunting (they are the only true carnivorous primate), but sleep in small groups during the day. Tarsiers are territorial with male and females ranges overlapping.




***Figure* 3.5.7.2** - *Distribution of tarsiers. Original map from http://d-maps.com/pays.php?num_pay=67&lang=en modified to show distribution of tarsiers.*



***Figure* 3.5.7.3** - Tarsier *(Carlito syrichta)*, Bohol, Philippines.

New World Monkeys

There are several families of New World Monkeys (NWMs):

1. Callitrichidae (i.e., marmosets and tamarins)
2. Cebidae (i.e., squirrel monkeys and capuchins)
3. Aotidae (i.e., night monkeys)
4. Pitheciidae (e.g., sakis and titis) and
5. Atelidae (i.e., howler monkeys and spider monkeys).

The 70+ species are found in Central and South America. All NWMs are arboreal although some will spend a little time on the ground. All are diurnal with the exception of the night monkey, which is nocturnal. All are quadrupeds, except for the spider monkey, which is a semi-brachiator. Some have prehensile tails, or “grasping” tails that serve as a fifth limb. Most, but not all, live in multi-male, multi-female groups. NWMs have flat noses with nostrils that are set far apart, wide open, and facing outwards.



***Figure* 3.5.7.4** - *Distribution of New World Monkeys. Original blank map from http://d-maps.com/pays.php?num_pay=299&lang=en modified to show range of New World Monkeys.*



***Figure* 3.5.7.5** - *White-eared Marmoset (notice the nostrils).*

They have strong big toes that are “strongly opposable but their thumbs are imperfectly opposable and their fingers cannot grip well” (Natural History Collections…2001-2007). NWMs have twelve premolars unlike their counterparts in the Old World who have eight premolars. The territories of different groups within various NWM species frequently overlap and sometimes “friendships” develop between the different species (Blashfield 2014).

New World Monkeys: Callitrichidae

Callitrichids are small primates, including the smallest monkey, the pygmy marmoset, which weighs about 3.5 ounces when fully mature and averages between 5-6 inches in length, who live in Central and South America. They generally have little fur on their faces, but have tufts of fur on their heads. Some of the most colorful primates belong to the Callitrichids, e.g., golden lion tamarins (pictured above). Their tails are non-prehensile and are used for balance when moving quadrupedally through the trees. Unlike most other monkeys, they have claws instead of nails except on the big toe (also called a hallux), which does have a nail, allowing them to dig into the bark of trees. Callitrichids do not have a third molar and are primarily insectivorous, but eat gums, fruit, buds, leaves, flowers, lizards, frogs, and snails as well. They live in monogamous (e.g., golden lion tamarin) or polyandrous (e.g., saddle-back tamarin) family groups. Females normally give birth to twins with the exception of Goeldi’s monkey that has single births. All individuals in the group contribute to infant care. Genera of callitrichids include *Callithrix*, e.g., buffy-headed marmoset and black-tufted marmoset, *Mico*, e.g., silvery marmoset and Emilia’s marmoset, *Cebuella*, i.e., pygmy marmoset, and *Saguinus*, e.g., Emperor tamarin and cottontop tamarin, *Callibella*, i.e., dwarf marmoset, *Callimico*, i.e., Goeldi’s marmoset, *Leontopithecus*, e.g., golden lion tamarin and black lion tamarin.



***Figure* 3.5.7.6** - *Golden lion tamarin family*



***Figure* 3.5.7.7** - Common marmoset *(Callithrix jacchus)* eating a cockroach

New World Monkeys: Cebidae

There are three genera of cebids: 1) *Cebus*, e.g., Kaapori capuchin and white-headed capuchin, 2) *Sapajus*, e.g., black capuchin and tufted capuchin, and 3) *Saimiri*, e.g., bare-eared squirrel monkey and Central American squirrel monkey. They are relatively small primates, ranging in weight from around 2.5 pounds (adult male squirrel monkey) to 8.5 pounds (adult male brown capuchin). Unlike callitrichids, cebids have a prehensile tail and the third molar. They are arboreal omnivores, focusing on fruits and insects. Females give birth to one or two offspring and are responsible for the care of their young. They live in relatively large multi-male, multi-female groups with a dominant male. Cebids perform urine washing; they urinate on their hands and then rub them on their fur and feet in order to scent mark their territory. These quadrupeds can be found in Central and South America as well as Trinidad and Tobago, Caribbean islands off the coast of Venezuela.



***Figure* 3.5.7.8** - *Bolivian-squirrel-monkey*



***Figure* 3.5.7.9** - White-faced capuchin *(Cebus capucinus)* on Rio Frio, Costa Rica

New World Monkeys: Aotidae

Night monkeys, also called owl monkeys, are arboreal primates living in Panama and parts of South America. They are small, weighing up to just under three pounds and exhibit no sexual dimorphism. Like other nocturnal primates, night monkeys have large eyes. Their faces are rounded and marked with three dark brown or black lines—one on either side of their eyes and one in the middle of the forehead. Sometimes it appears as if night monkeys have no ears, but actually their small, rounded ears can be hidden by thick fur. Night monkeys are monogamous; mated pairs live with two or three offspring. Females usually have a single offspring, with both parents participating in infant care. Primarily frugivores, night monkeys will also eat leaves, flowers, insects, eggs, birds, spiders, and bats. Like cebids, night monkeys practice urine washing.



***Figure* 3.5.7.10** - Night monkey *(Aotus zonalis)*

New World Monkeys: Pitheciidae

The long-coated pithecids live in the various rainforest habitats in South America. These arboreal quadrupeds live in small groups. Titi monkey groups are centered on a monogamous pair while the sakis and uakaris live in multi-male, multi-female groups ranging from 10 individuals up to 100. Pithecids are territorial but generally use vocalizations instead of physical confrontation to defend their territory. All pithecids are frugivores; in fact, they are specialized frugivores called granivores who focus primarily on fruit seeds. Most pithecid females give birth to a single offspring. In general, males do not participate in infant care, although male sakis may groom their young and male titi monkeys provide the majority of care for their offspring. The four genera of pithecids are 1) *Callcebus*, e.g., titi monkeys, 2) *Cacajao*, e.g., uakaris, 3) *Chiropotes*, e.g., bearded sakis, and 4) *Pithecia*, e.g., sakis.



Figure **3.5.7.11** - *White-faced saki monkey*



Figure **3.5.7.12** - *Titi Monkey*



Figure **3.5.7.13** - Uakari *(Cacajao calvus)*

New World Monkeys: Atelidae

The arboreal atelids are found in both Central and South America, although woolly monkeys are found only in the Amazon, muriquis are in Brazil's southeaster rainforest, and yellow-tailed woolly monkeys are only in the cloud forests in the Peruvian Andes. The atelids range is size from the smaller spider monkey, weighing about 13-20 pounds, to the howler monkey, weighing up to 25 pounds. Atelids have prehensile tails with a hairless, tactile pad on the underside of the tip of the tail. This sticky pad helps the atelids hold on to things, especially tree limbs as they hang from their tails to eat. They are primarily frugivores, but will eat leaves, flowers, plant gums, and insects depending on the season. Atelids live in multi-male, multi-female groups, anywhere from two individuals up to one hundred (male howler monkeys are known to have "harems" of females). Males are philopatric; females leave their birth group. Atelids have ranges, but do not defend territories. When ranges overlap, conflict can occur. Females usually give birth to one offspring in the dry season and solely responsible for care of the young. While all of the atelids use vocalizations to communicate, howler monkeys are known for their distinctive calls, or roars, that can be heard (by humans) over a mile away.



Figure **3.5.7.14** - Howler monkey (*Alouatta caraya*)



***Figure* 3.5.7.15** - *Geoffrey's spider monkey*



***Figure* 3.5.7.16** - Woolly monkey (*Lagothrix lagotricha*)

References

1. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Marmosets, tamarins, and Goeldi's monkey: Callitrichidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2005. p. 496-508.
2. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Howler monkeys and spider monkeys: Atelidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2005. p. 526-535.
3. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Night monkeys: Aotidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2005. p. 509-515.
4. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Sakis, Titis, and Uakaris: Pitheciidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2005. p. 516-525.
5. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Squirrel monkeys and capuchins: Cebidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2005. p. 487-495.
6. Blashfield JF. New World Monkeys. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, 5thedition, Vol. 6. Farmington Hills (MI): Gale; 2014. p. 3013-3018.
7. Blashfield JF. Tarsiers. In: Lerner KL , Lerner BW, editors. The Gale Encyclopedia of Science, 5thedition, Vol. 8. Farmington Hills (MI): Gale; 2014. p. 4309-4310.
8. Dewey T. 2007. Aotidae: night monkeys. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 28]. Available from: animaldiversity.org/accounts/Aotidae/
9. Dewey T. 2007. Atelidae: howler and prehensile tailed monkeys. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 28]. Available from: animaldiversity.org/accounts/Atelidae/
10. Dewey T. 2007. Pitheciidae: titi monkeys, sakis, and uakaris. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 28]. Available from: animaldiversity.org/accounts/Pitheciidae/
11. Dewey T, Myers P. Tarsiidae: tarsiers. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 28]. Available from: animaldiversity.org/accounts/Tarsiidae/
12. Jurmain R, Kilgore L, Trevathan W. 2013. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning.
13. Larsen CS. 2008. Our origins: discovering physical anthropology. New York (NY): W.W Norton & Company, Inc.

14. New World Monkeys [Internet]. c. 2001-2007. Edinburgh (Scotland): The Natural History Collections of the University of Edinburgh: [cited 2015 Jun 22]. Available from:http://www.nhc.ed.ac.uk/index.php?page=493.504.513
15. Shefferly N. Callitrichinae: marmosets and tamarins. Animal Diversity Web, University of Michigan Museum of Zoology [Internet] [cited 2015 Jun 28]. Available from: animaldiversity.org/accounts/Callitrichinae/

This page titled 3.5.7: Overviews of Living Primates (Haplorhines) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.7: Overviews of Living Primates (Haplorhines)** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.8: More Haplorhines (Old World Monkeys)

Old World monkeys (OWMs) are found in sub-Saharan Africa, southern Asia, and northern Japan. They live in a wide variety of habitats, from tropical jungle to seasonal snow and savannas. Some are arboreal and others terrestrial. There is variety of group structure, but most live in diurnal multi-male, multi-female groups. The traits that unite OWMs are:

- Lack of prehensile tail
- Forelimbs are shorter than hindlimbs
- Quadrupedal
- Narrow nostrils that are close together
- Well-developed opposable thumbs and big toes
- Presence of ischial callosities in many species (hard pads on the lower sides of their buttocks for sitting)
- More sexual dimorphism than Strepsirhines or NWMs
- Visible estrus in females
- Strong dominance hierarchies
- Bilophodont molars (molars form two parallel ridges)



***Figure* 3.5.8.1** - *Distribution of Old World Monkeys. Original blank map from http://d-maps.com/carte.php?num_car=3266&lang=en modified world map to show range of Old World Monkeys.*

All OWMs belong to the family Cercopithecidae. They are divided into two subfamilies: Cercopithecinae, monkeys with cheek pouches, and Colobinae, monkeys who are predominately leaf-eaters. There are eighteen genera and about ninety species of OWMs.




***Figure* 3.5.8.2** - *Macaca fascicularis* - crab-eating macaque (notice the nose)

Cercopithecinae



***Figure* 3.5.8.3** - Mandrill (*Mandrillus sphinx*)

The largest number of OWM species belong to the Cercopithecinae and include the following genera:

- *Allenopithecus*, i.e., Allen's swamp monkey

- *Miopithecus*, i.e., talapoin
- *Erythrocebus*, i.e., patas monkey
- *Chlorocebus*, e.g., vervet monkey and green monkey
- *Cercopithecus*, e.g., Diana monkey and Pluto monkey
- *Macaca*, e.g., Japanese macaque and booted macaque
- *Lophocebus*, e.g., grey-cheeked mangabey and Uganda mangabey
- *Rungwecebus* i.e., highland mangabey
- *Papio*, e.g., olive baboon and yellow baboon
- *Theropithecus*, i.e., gelada
- *Cercocebus*, white-eyelid mangabeys, e.g., sooty mangabey and collared mangabey
- *Mandrillus*, i.e., mandrill and drill



***Figure* 3.5.8.4** - Sexual dimorphism in Hamadryas baboons *(Papio hamadryas)*.

There are a number of semi-terrestrial cercopithecines, e.g., baboons, who spend their days foraging in the grasslands and sleep either on cliff-faces or in trees. These semi-terrestrial OWMs exhibit significant sexual dimorphism. Canine teeth are large and sharp; this is beneficial for eating but also for defense against rivals and predators. Ischial callosities are present on the ground-dwelling species. Their faces are bereft of fur except for perhaps a beard-like tuft on the chin. Some have interesting coloration of the face or genitals. Male mandrills (pictured above) have the most conspicuous coloring on their faces and rumps, in addition to their penises, which are bright red. Cercopithecines have elongated snouts with the OWM characteristic noses with nostrils that are narrow and set close together.



***Figure* 3.5.8.5** - Japanese macaques *(Macaca fuscata)* in hotspring at Nagano, Japan

Cercopithecines vary in size from the smallest at close to three pounds (*Miopithecus talapoin*) to the largest, the male mandrill (*Mandrillus sphinx*), at about forty-eight pounds. The smaller cercopithecines spend more time in trees than the large cercopithecines. They inhabit a wide-range of environments, from tropical rainforests, e.g., talapoins and mangabeys, to semi-arid deserts, e.g., chacma baboons (*Papio ursinus*), and even areas with seasonal snow, i.e., Japanese macaques (*Macaca fuscata*. In fact, Japanese macaques have an interesting cultural adaptation to living in a cold, snowy region…they hang out in hot springs. Most cheek-pouched monkeys are omnivorous, eating seeds, fruits, leaves, spiders, insects, and even small vertebrates. All cercopithecines live in large multi-male, multi-female groups or single-male, multi-female groups, with bachelor groups forming at times as they migrate from their natal groups. Most females have visible estrus, making it easy for males to tell when the female is fertile. The only species with no to little visible estrus are long-tailed macaques, forest-living guenons, and some mangabeys. Females typically give birth to one offspring at a time.



Figure **3.5.8.6** - Female vervet monkey *(Chlorocebus pygerythrus)* with infant

Colobinae



Figure **3.5.8.7** - Golden snub-nosed monkey *(Rhinopithecus roxellana)*

The Colobinae are quadrupedal folivores, including the colobus monkeys, langurs, and odd-nosed monkeys. There are ten different genera with long tails and various fur coloration:

- *Colobus*, e.g., black colobus and Angola colobus
- *Piliocolobus*, e.g., Zanzibar red colobus and Bouvier's red colobus
- *Procolobus*, i.e., olive colobus
- *Trachypithecus*, e.g., dusky leaf monkey and capped langur
- *Presbytis*, e.g., banded surili and Hose's langur
- *Semnopithecus*, e.g., purple-faced langur and Kashmir gray langur
- *Pygathrix*, e.g., red-shanked douc and gray-shanked douc
- *Rhinopithecus*, e.g., golden snub-nosed monkey and Myanmar snub-nosed monkey
- *Nasalis*, i.e., proboscis monkey
- *Simias*, i.e., pig-tailed langur



Figure **3.5.8.8** - Gray-shanked douc langur *(Pygathrix cinerea)*

Most are arboreal and are found in rainforests, mountain forests, savannas, and mangroves. Because of their leafy diet, colobines have multi-chambered, or sacculated, stomachs to break down plant cellulose into useable calories, and long intestines to increase the absorption of nutrients. They are the only ruminant primates. Colobines are smaller than many of the cercopithecines, weighing from about 10 to 34 pounds. They have short snouts in comparison to the cercopithecines and have teeth that are adapted for

shearing leaves. Their legs are somewhat longer than their arms. Colobus monkeys have a vestigial thumb, making fine manipulation of objects impossible. They have little to no hair on their faces other than tufts on the cheeks or chin. All except the snub-nosed monkey have long tails that are used to balance when moving. There is variety in group composition. Multi-male, multi-female groups or single-male, multi-female groups are common; monogamy is rare. Bachelor groups may form as males migrate from their natal group. Females usually give birth to one offspring at a time. During estrus, females have visible menstrual bleeding.



***Figure* 3.5.8.9** - King colobus *(Colobus polykomos);* note the thumb




***Figure* 3.5.8.10** - Proboscis Monkey (*Nasalis larvatus*)

References

1. Jurmain R, Kilgore L, Trevathan W. 2013. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning.
2. Kamilar JM, Roehrdanz NL. Monkeys, Old World. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 4. Thousand Oaks (CA): SAGE Reference; 2006. p. 1617-1618.
3. Martin RD. Old World monkeys I (Colobinae). In: Hutchins M, et al., editors. Grzimek's animal life encyclopedia, 2nd edition, Vol. 14. Detroit (MI): Gale; 2004. p. 171-186.
4. Martin RD. Old World monkeys II (Cercopithecinae). In: Hutchins M, et al., editors. Grzimek's animal life encyclopedia, 2nd edition, Vol. 14. Detroit (MI): Gale; 2004. p. 187-206.
5. Old World Monkeys [Internet]. c. 2001-2007. Edinburgh (Scotland): The Natural History Collections of the University of Edinburgh: [cited 2015 Jun 29]. Available from: http://www.nhc.ed.ac.uk/index.php?page=493.504.508.512
6. O'Neill D. 1998-2014. Old World Monkeys. Behavioral Sciences Department, Palomar College [Internet] [cited 2015 Jun 29]. Available from: anthro.palomar.edu/primate/prim_6.htm

This page titled 3.5.8: More Haplorhines (Old World Monkeys) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.8: More Haplorhines (Old World Monkeys)** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.9: Even more Haplorhines (The Apes)



Figure **3.5.9.1** *- Distribution of gibbons, siamangs, and orangutans. Original blank map from http://d-maps.com/pays.php?num_pay=67&lang=en modified to show distibution of the Southeast Asian great apes.*



Figure **3.5.9.2** *- Distribution of chimpanzees and gorillas. Original blank map from http://d-maps.com/carte.php?num_car=728&lang=en modified to show the distribution of the African great apes.*



Figure **3.5.9.3** - Chimpanzee *(Pan troglodytes)* female and offspring

All apes are members of the superfamily Hominoidea; sometimes they are called hominoids. Hominoids are divided into two families: Hylobatidae, the lesser apes, and Hominidae, the great apes. The Hominids are further divided into three subfamilies: Ponginae, Gorillinae, and Homininae, the latter of which is comprised of two tribes: Panini and Hominini. Which apes belong to each of the classifications will be discussed below.

Common Characteristics

Apes share some common characteristics; note these are in comparison to other primates (Jurmain et al. 2013: 142):

- Generally larger body size
- No tail
- Lower back is shorter and more stable
- Arms longer than legs (except for humans)
- Anatomical differences in shoulder joint that facilitate suspensory feeding and locomotion
- Generally more complex behavior
- More complex brain and enhanced cognitive abilities
- Increased period of infant development and dependency



***Figure* 3.5.9.4** - Siamang *(Symphalangus syndactylus)*




***Figure* 3.5.9.5** - Female and male white-browed gibbons *(Hoolock hoolock)*

There is one genera of hylobatids,*Hylobates* and four subgenera: 1) *Bunopithecus*, e.g., Western hoolock gibbon, 2) *Hylobates*, e.g., white-handed gibbon and Bornean white-bearded gibbon, 3) *Symphalangus*, i.e., siamang, and 4) *Nomascus*, e.g., black crested gibbon and yellow-cheeked gibbon, many of which are endangered or critically endangered. They are the smallest of the apes, which is why they are sometimes called the “lesser” apes. Hylobatids rarely spend time on the ground. Because they venture so infrequently to the ground they do not have to contend with predators. “Their lifestyle has enabled them to remain petite and agile” (Hamilton 2006[2]: 222). They are true brachiators. Their arms are much longer than their legs and the thumb is attached to the wrist instead of the palm. This enables them to use their hands as hooks as they swing through the trees. They can move quite fast as their arm span allows them to swing over nine feet in a single movement and they can leap up to almost 30 feet in a single bound (Hamilton 2006[2]). While feeding, hylobatids will use quadrumanous, or four-footed, climbing.

Like some Old World monkeys, hylobatids have ischial callosities, which are for sitting while sleeping (if they do not sit while sleeping, they curl up on their side). They are the only non-human ape that does not build a nest for sleeping. They are well-known for their vocalizations. Monogamous pairs will sing in the morning and evening. The calls can travel over half-a-mile and are used “for territorial defense, maintenance of the pair bond, and general communication” (Paciulli and Neary 2006: 1079). Gibbons may call out 80 to 90% of the day while siamangs only vocalize about 30% of the day. Check out the hylobatids sound gallery from the Gibbons Research Gallery to listen to various vocalizations: http://www.gibbons.de/main/sound.html.

The hylobatids exhibit some sexual dimorphism. They exhibit various colorations as well. For example, female white-browed gibbons (*H. hoolock*) are tan while the males are black; a condition referred to as sexual dichromatism, which means that males and females are different colors. Other species have various color fur, e.g., the silvery gibbon has a silver coat while the Kloss gibbon has a black coat and the Mueller’s Bornean gibbon is gray although there is no variation in color between males and females.

Hylobatids are frugivores, although they will eat leaves, flowers and small invertebrates as well. As mentioned previously, they are monogamous and live with their offspring, who may stay with their parents for up to ten years. At some point, offspring “bud off,” or take over part of their parents territory and start their own family groups. Families defend their territories, using vocalizations, chasing, and theatrics such as stick breaking and trashing vegetation; rarely are confrontations physical.

Hominidae



Figure **3.5.9.6** - Baby bonobo *(Pan paniscus)*

Orangutans, gorilla, chimpanzees, bonobos, and humans comprise the family Hominidae. Humans will not be covered in this section as Unit 3 is entirely about human evolution. The great apes are the largest primates, with the male eastern lowland gorilla weighing up to 450 pounds. There is extreme sexual dimorphism among orangutans and gorillas; less so in chimpanzees and bonobos. All are quadrupedal, although there are some differences in how quadrupedalism manifests itself. Orangutans exhibit quadrumanous scrambling when arboreal and fist-walking terrestrially while chimpanzees and gorillas are knuckle-walkers. Bonobos also exhibit frequent bipedalism.

Ponginae



Figure **3.5.9.7** - Bornean orangutan *(Pongo pygmaeus)*



Figure **3.5.9.8** - Sumatran orangutan *(Pongo abelii)*

Both species of pongids are critically endangered: *Pongo pygmaeus*, the Bornean orangutan, and *Pongo abelii*, the Sumatran orangutan with dark maroon hair and figure-eight-shaped, hairless faces. Bornean orangutans have ginger-colored hair with light hair on their elongated oval-shaped faces. Bornean orangutans are smaller and fatter than the Sumartran orangs. Second in size only to gorillas, orangutans exhibit sexual dimorphism with the adult males weighing up to 200 pounds with adult females about half that. Mature males have cheek flanges (the bigger the cheek flanges the better as females choose mates based on cheek flange size) and throat pouches with which they make booming calls that alert male rivals and females of their presence. They also have beards and mustaches (Sumatran orangs have longer facial hair than the Bornean orangs). Sexually mature (but not fully grown) males, and occasionally a flanged male, have been known to try to mate with unwilling females, a behavior human researchers have referred to as attempted rape. A study of Sumatran orangs suggests that this is a successful breeding strategy for the unflanged

males who sired as many offspring as flanged males without the stress of competing with a fully mature flanged male in an overlapping home range.



***Figure* 3.5.9.9** - Fully mature male Bornean orangutan *(Pongo pygmaeus)*

Orangutans are solitary primates with males and females coming together only to mate. Females may temporarily socialize with other females, but generally live only with one or two of their offspring. Females usually have one offspring at a time with birth spacing of at least six years. Of the non-human apes, female orangs have the longest period of parental investment, lasting up to ten years. Primatologists suggest that this long period of childcare is so that the young have sufficient time to learn "the complexities of life in the rain forest" (Groves 2006: 1787). Like the hylobatids, orangs spend the majority of their time in trees. Since they weigh too much to walk across tree branches to get to another tree, they return to the ground and walk to the next tree. Sumatran orangutans spend less time on the ground than their Bornean counterparts because they are subject to predators such as tigers and cloud leopards whereas the Bornean orangs do not have the same concern.

Orangutans are frugivores, but also eat bird eggs, termites, caterpillars, ants, bark, sap, and a variety of other foods depending on the season. During mast fruiting (a period occurring every two to ten years when many trees fruit at the same time), orangutans gorge themselves with fruit. This allows the orangs to increase their fat stores. Orangutans spend the majority of their day eating, resting, and moving between trees, upwards to 95% of their day. They spend two to three hours in the morning feeding before resting. In late afternoon they travel before preparing their nest for the night.

One of the major differences between the Sumatran and Bornean orangutans has to do with culture. The former manufacture and use tools. They use leaves for multiple purposes, including umbrellas and "toilet paper." They use branches as flyswatters and in display behavior. They have been observed modifying sticks so that they can be used to collect insects, open seedpods and large fruits, and dig holes. Stacks of leaves are used as padding when holding spiny fruits.

Gorillinae



***Figure* 3.5.9.10** - Western Gorilla *(Gorilla gorilla gorilla)*

The "gentle giant" of the great apes, there are two species of *Gorilla*: *G. gorilla* (Western lowland gorilla) and *G. beringei* (mountain gorilla) and four subspecies: *G. g. gorilla*, *G. g. diehli*, *G. b. beringei*, and *G. b. graueri*.




Figure 3.5.9.11 - Male silverback gorilla *(G. beringei)*

Both of the former are critically endangered while the latter is endangered. Gorillas are sexually dimorphic and typically live in single-male, multi-female groups with a dominant male, over twelve-years-old, who has a band of silver-gray hair on its back, thus is referred to as the silverback; however, there are some multi-male, multi-female groups. Both males and females migrate from their birth groups, although female mountain gorillas tend to migrate more than female lowland gorillas, which helps to avoid inbreeding. Also in the mountain gorillas, males may stay with their natal group. When this occurs, the male is subordinate to the silverback and is called a blackback male. If the silverback dies, one of the blackbacks has the opportunity to become the new silverback. For single-male, multi-female groups, when the silverback dies the females and their offspring may disperse to join other groups or they may stay together until a new silverback transfer. Without a silverback to protect the group, infanticide can occur. The silverback maintains bonds with his females through grooming and togetherness. Conflict between silverbacks involves vocalizations and threat displays, including chest beating; rarely is actual physical conflict involved.



Figure 3.5.9.12 - Mountain gorilla *(G. b. beringei)*

Group activity surrounds the silverback who makes all the decisions about when to travel, settling conflict, finding food resources, and defending the group. Blackbacks may assist the silverback with defense. These knuckle-walking primates, spend their days on the ground foraging for food. Females and juveniles sleep in trees, while the dominant male silverback sleeps on the ground. All gorillas are herbivores, although their food focus varies because of their habitats. Mountain gorillas (*G. b. beringei*) are folivores who favor bamboo. Eastern lowland gorillas (*G. b. graueri*) eat seasonally, focusing on fruit when available. They will also eat ants. Western lowland gorillas (*G. gorilla*) also focus on fruit, but also eat herbaceous vegetation, termites, and ants. Because of their diets, the home range of gorillas varies, with Western lowland gorillas having the largest range and mountain gorillas the smallest.

Females usually give birth to one offspring at a time. Infant mortality is high among mountain gorillas (up to 38%) so the mother's care is particularly important. Males are involved in socialization of infants but do not provide care beyond protection. Infants are weaned after three or four years, providing a decent interval between births and increasing the infant's chance of survival.



***Figure* 3.5.9.13** - Adult female and infant wild chimpanzees *(Pan troglodytes)* feeding on *Ficus sur (*figs*)*

There are two genera of extant hominins: *Pan* and *Homo*, with two species of *Pan*: 1) *Pan troglodytes* and 2) *Pan paniscus*, and one species of *Homo*: *Homo sapiens*.

The endangered *Pan troglodytes*, common chimpanzees, and *Pan paniscus*, bonobos are the apes most genetically similar to humans (98%+). Some researchers (Grove 2001 and Wildman et al. 2003) argue that chimpanzees are so closely related to humans that they should be genus *Homo*.



***Figure* 3.5.9.14** - *Pan paniscus (*note the part in the hair*)*

Chimpanzees live in West and Central Africa, primarily in Gabon and the Democratic Republic of Congo. They inhabit a number of habitats, including, savannas and montane, swamp, and evergreen rainforests. Bonobos are found only in the Democratic Republic of Congo in closed-canopy rainforest. Both chimps and bonobos exhibit some sexual dimorphism. Male bonobos weigh on average 86 pounds and females 63 pounds. Male chimps are slightly larger at an average of 88 to 132 pounds, and females between 70 and 104 pounds; they are sturdy and strong. As indicated by the average weights, chimpanzees are somewhat larger than bonobos. Chimpanzees have black hair and are born with pale faces that darken as they age. Males and females have beards and prominent ears. Bonobos have black fur and black faces with hair tufts on the sides of their faces and a part in the hair on their heads. While both species are quadrupedal knuckle-walkers, both can walk bipedally (although bonobos are better bipedal walkers) and bonobos also semi-brachiate.

Chimpanzees live in multi-male, multi-female groups, ranging anywhere from ten to over one hundred individuals. The core of the group is centered on an alpha male along with a small group of bonded males, although in West Africa, some groups appear to be oriented on females. The alpha male and his subordinates use physical force, display behavior, and aggression to control the community. They work together to defend the group's territory, patrolling and chasing out unfamiliar males. The size of the group's territory is dependent on the number of males in the group. Even though this small group of males works with the alpha, the dominant chimpanzee must constantly work to maintain his position of power.



Figure **3.5.9.15** - Mother chimpanzee *(Pan troglodytes)* and offspring

The strongest bond within chimpanzee social groups is between mothers and their offspring. Females generally have one baby at a time and give birth approximately every five to six years. Even though offspring are weaned by the time they are three-years old, juveniles may stay with their mother and siblings until age five through eight and help with childcare. Juvenile females leave the group at about ten-years-old, except for those groups that are centered on a core group of females. There is quite a bit of movement among group members who do not always spend the day together; they often break off into smaller groups of three to six individuals to forage.

Chimpanzees are omnivores, eating fruits, resin, bark, young leaves, tree seeds, and pith (Orcholl 2006). They incorporate over 300 species of plants in their diet. Some plants seem to be used only for medicinal purposes. Chimpanzees also incorporate animal protein in their diet, which comes from small mammals, including other primates like the colobus monkey, and insects such as termites. Male, and the occasional female, chimpanzees use cooperative techniques to hunt (watch this short video of chimps on the hunt: youtu.be/A1WBs74W4ik). At Mahale Mountain National Park, Tanzania, researchers have observed several occurrences of infanticide after failed hunts. Chimpanzees have been documented making and using spears to hunt and “fishing sticks” to acquire termites and honey (you can check out this short video about spear-wielding chimps: youtu.be/FyGxQq7jSA8). Chimpanzees have other cultural behaviors, including the hand-clasping grooming technique, using wadded up leaves as sponges, and hammers and anvils to break open nuts.



Figure **3.5.9.16** - Bonobos *(Pan paniscus)*

Bonobos are sometimes referred to as pygmy chimpanzees because they are slightly smaller and more slender than common chimpanzees, but the weight differences, as mentioned above, are not that great. There is less sexual dimorphism than chimps and about the same as in humans. Bonobos are efficient bipedal walkers and have been documented wading in streams while fishing, walking while carrying an armload of sugar cane, and with arms over each other’s shoulders (Orcholl 2006).



***Figure* 3.5.9.17** - Bonobo *(Pan paniscus)*

Bonobos live in fission-fusion social groups of over twenty individuals. Mother and son bonds are the strongest as sons use their mother's rank in the social hierarchy as a means to garner favor with other high-ranking females. It is usually the highest-ranking female's son who is alpha, but the alpha position is much different than that of the chimpanzee. The male and female alphas are almost equally dominant (Orcholl 2006). Unrelated females (juvenile females leave their natal group between ages seven to nine) work together to protect their offspring and secure resources at the expense of males. There is no core group of males that work together---after the alpha male, the hierarchy of males is murky. Also unlike chimpanzees, physical aggression is rarely used to dominate and intimidate. Intra-group conflict is settled using sexual interactions such as rump-rubbing and penis fencing among males and genital-to-genital rubbing (G-G rubbing) among females. Females wishing to join a group will use G-G rubbing with the alpha female and intercourse with the alpha male to gain admittance. While to humans, this might seem like an unusual ways to use sexual behavior, these behaviors, along with the fact that female bonobos are always receptive to sexual contact, have multiple partners, and copulate while pregnant, has eliminated infanticide from bonobo society. The fact that females leave their natal groups prevents incest. Females usually give birth to a single offspring; birth-spacing is from four to six years. Because males do not know which offspring belong to them they do not invest their time in infant care. Mothers are solely responsible for their offspring, which may help to explain why females band together.

Unlike chimpanzees, bonobos do not make and use tools, although captive bonobos have been observed using objects that are made available to them, such as tennis balls as water sponges. They do, however, exhibit emotions such as sympathy and empathy. They seem to respect other individuals' privacy, never entering another bonobo's sleeping nest. Like chimps, bonobos use at least one plant for medicinal purposes, specifically a leaf used when they have an intestinal parasite. Bonobos are omnivores, although about 57% of their diet is from fruit. They also eat leaves, flowers, bark, pith, roots, mushrooms, sprouts, and seeds. While bonobos do not hunt, they will opportunistically eat small mammals, insect larvae, earthworms, honey, eggs, and soil (Orcholl 2010[1]). Finding a food source can set off a flurry of excitement and sexual interaction.

References

1. Allen CJ, Evans AV, McDade MC, Schlager N, Mertz LA, Harris MS, et al., editors. Great Apes and Humans: Hominidae. In: Grzimek's student animal life resource, Vol. 16: Mammals: Vol. 3. Detroit (MI): UXL; 2007. p. 563-577.
2. Beckham EC, Cumberlidge N, Regenstein LG. Apes. In: Lerner KL, Lerner BW, editors. The Gale encyclopedia of science, 5th edition, Vol. 5. Farmington Hills (MI): Gale; 2014. p. 278-281.
3. Birx HJ. Gorillas. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 3. Thousand Oaks (CA): SAGE Reference; 2006. p. 1102-1103.
4. [1]Cawthon Lang KA. 2005. Primate Factsheets: Gorilla (Gorilla) Behavior. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/gorilla/behav
5. [2]Cawthon Lang KA. 2005. Primate Factsheets: Gorilla (Gorilla) Taxonomy, Morphology, & Ecology. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/gorilla
6. [3]Cawthon Lang KA. 2005. Primate Factsheets: Orangutan (Pongo) Behavior. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/orangutan/behav
7. [4]Cawthon Lang KA. 2005. Primate Factsheets: Orangutan (Pongo) Taxonomy, Morphology, & Ecology. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/orangutan
8. Cawthon Lang KA. 2006. Primate Factsheets: Chimpanzee (Pan troglodytes) Taxonomy, Morphology, & Ecology. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/chimpanzee

9. [1]Cawthon Lang KA. 2010. Primate Factsheets: Bonobo (Pan paniscus) Behavior. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/bonobo/behav
10. [2]Cawthon Lang KA. 2010. Primate Factsheets: Bonobo (Pan paniscus) Taxonomy, Morphology, & Ecology. Primate InfoNet [Internet] [cited 2015 Jul 02]. Available from: http://pin.primate.wisc.edu/factsheets/entry/bonobo/taxon
11. Cumberlidge N. Gorillas. In: Lerner KL, Lerner BW, editors. The Gale encyclopedia of science, 5th edition, Vol. 4. Farmington Hills (MI): Gale; 2014. p. 2023-2024.
12. Groves C. 2001. Primate taxonomy. Washington (DC): Smithsonian Institute Press.
13. Groves C. Orangutans. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 4. Thousand Oaks (CA): SAGE Reference; 2006. p. 1785-1787.
14. [1]Hamilton, GS. Apes, Lesser. In: Birx, HJ, editor. Encyclopedia of anthropology, Vol. 1. Thousand Oaks (CA): SAGE Reference; 2006. p. 220-221.
15. [2]Hamilton, GS. Apes, Lesser. In: Birx, HJ, editor. Encyclopedia of anthropology, Vol. 1. Thousand Oaks (CA): SAGE Reference; 2006. p. 221-222.
16. [ICUN] Internation Union for Conservation of Nature and Natural Resources [Internet]. 2015. The ICUN red list of threatened species [cited 2015 Jul 02]. Available from: www.icunredlist.org/
17. Jurmain R, Kilgore L, Trevathan W. 2013. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning.
18. O'Neill D. 1998-2014. Apes. Behavioral Sciences Department, Palomar College [Internet] [cited 2015 Jul 01]. Available from: anthro.palomar.edu/primate/prim_7.htm
19. [1]Orcholl JL. Bonobos. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 3. Thousand Oaks (CA): SAGE Reference; 2006. p. 389-392.
20. [2]Orcholl JL. Chimpanzees. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 3. Thousand Oaks (CA): SAGE Reference; 2006. p. 500-505.
21. Paciulli LM, Neary JL. Gibbons. In: Birx HJ, editor. Encyclopedia of anthropology, Vol. 3. Thousand Oaks (CA): SAGE Reference; 2006. p. 1078-1079.
22. Richman K. Chimpanzees. In: Lerner KL, Lerner BL, editors. The Gale encyclopedia of science, 5th edition, Vol. 2. Farmington Hills (MI): Gale; 2014. p. 905-910.
23. Wildman DE, Uddin M, Guozhen L, Grossman LI, Goodman M. 2003. Implications of natural selection in shaping 99.4% nonsynonymous DNA identity between humans and chimpanzees: enlarging genus Homo. Pro Natl Acad Sci U S A 100(12): 7181-7188.

This page titled 3.5.9: Even more Haplorhines (The Apes) is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.9: Even more Haplorhines (The Apes)** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.5.10: Primate Evolution



Figure **3.5.10.1** - Fossil ape*: Proconsul*

Now that you have an understanding of living primates' morphology and behavior, it is time to learn about the origins of primates. the study of primate evolution is multidisciplinary in nature and incorporates data and methods from paleontology, geology, anthropology, and archaeology to study the fossil record of primates.

Fossils



Figure **3.5.10.2** - *Permineralized wood*

Fossils are at the center of the study of ancestral primates. Animal fossils provide insight into morphology and behavior of ancient organisms while plant fossils help paleoanthropologists reconstruct ancient environments (paleoecology). One thing to keep in mind is that, "the fossil record is highly skewed in favor of organisms that died and were preserved in calm seas, estuaries, tidal flats, or the deep ocean floor (where there are few scavengers and little disruption of layers)" (Panell 2014: 1839). There are multiple ways that fossils form. Permineralization, or petrification, is the most common process. This occurs when organic remains decay and water seeps into the remains and deposits minerals that form crystals which harden. Generally, it is the bone that is fossilized as they are the hardest parts of the body and take the longest to decay, giving time for the minerals to crystalize and harden. Fossils can also form when an organic entity is encased in tar, resin, or even ice. When this occurs bodily tissues can be preserved, but it as extremely rare occurrence. Carbonization can lead to fossilization as well. When an organic item is buried in sediment, heat and pressure causes the release of hydrogen and oxygen, which leaves behind a carbon residue. To read more about other methods, check out the article, "How Do Fossils Form?" You can also learn more about the fossilization process by watching this short video, Becoming a Fossil.

When a fossil is found, it can be dated using a variety of methods. Some methods date the fossil itself; other methods use the surrounding matrices. There are relative dating techniques that provide an order of occurrence, but no absolute dates, and there are absolute, or chronometric, dating techniques that provide dates (usually a range of dates) for an object. Examples of relative dating techniques include stratigraphy, index fossil concept (biostratigraphy), and flourine dating. Stratigraphy involves mapping observable geologic strata and the artifacts (e.g., fossils) included in those layers. Building on stratigraphy, index fossils are used as time markers. Let's say that one type of fossil was found in a strata at Site A and the same type of fossil was found in a strata at Site B, then we can assume that the strata and fossils at both sites were deposited at the same time. Flourine dating is a process that uses the amount of flourine accumulation as a basis for comparison between objects, e.g., if two objects have similar amounts of

flourine then they were most likely deposited at the same time. Radiometric techniques are commonly used for dating fossils. Radiocarbon (^{14}C-dating) was the first radiometric dating technique developed; it revolutionized anthropological studies of the past. ^{14}C-dating can be used on organic materials and is accurate to about 70,000 years ago. It measures the rate of decay of carbon-14. Fission tracking involves measuring the track density of decaying uranium-238. As this method works best on igneous materials, it can be used to date the matrix surrounding fossils. Potassium-Argon dating (K/Ar-dating) is employed frequently in the study of primate origins. This method measures the rate of decay of potassium-40 into argon-40. Again, the rock surrounding fossils is dated and not the fossil itself. These are just a few of the dating methods used. You can learn more about these and other dating methods by reading the article, "Dating Rocks and Fossils Using Geologic Methods" (optional).

Taphonomy

The study of what happens to organisms after they die is called **taphonomy**. Taphonomic studies of fossils is important in order to understand the context of the fossil and the environment from which it came. Has the fossil been disturbed in any way? Has it been moved by water, scavenged by animals, etc?

Sometimes taphonomy is done with experimental methods.

Paleoecology

Paleoecologists study "...the physical structure and biological functions of organisms, their interactions with other, and their role in ancient ecosystems" (Lerner and Lerner 2014: 3217) to determine what the landscape may have looked like during the time that a particular organism, including primates, lived there (remember, evolution is environmentally dependent!). There are several specialized areas of study within paleoecology. **Paleobotany** is the study of fossil plants. For more on paleobotany, check out Dr. Robert Gastaldo's site: A Brief Introduction to Paleobotany. **Paleopalynology** is the study of fossil pollen while **phytoliths** are fossilized plant cells.




***Figure* 3.5.10.3** - *Geologic time scale*

Geologic time

From geology we have an understanding of the time involved in the development of earth. In biological anthropology, we are specifically focused on the late Mesozoic/early Cenozoic eras in relation to primate evolution. You should be familiar with the time ranges for the epochs of the Cenozoic. Below you will learn about primate developments associated with each epoch.

If you are feeling particularly investigatory today, you may want to check out this interactive geological time machine (optional)

Primate Evolution

The Mesozoic: the origin of mammals

The Mesozoic era is known as "the age of the dinosaurs" due to their ecological dominance at the time; however, it is during this era that the first mammals evolved, including a primate-like mammalian ancestor. Research using DNA analysis and fossils

suggests that by 75 mya (million years ago) all of the mammalian orders had diverged. The various mammalian families began to diversify in the Cenozoic era (Jurmain et al. 2013).

Several things happened at the end of this period that led to a rise in the mammals:

- a rise in the number and diversity of **angiosperms** (flowering plants)
- the K-T boundary extinction

The Cenozoic era

The Paleocene epoch

Proto-primates, primate-like mammals, evolved in the Paleocene epoch, about 65 mya. The proto-primates from this epoch are controversial; some argue that they are related to primates but are not actually primates (hence, "proto-primates"). The strongest evidence linking these proto-primates or Plesiadapiformes to primates comes in the form of *Carpolestes simpsoni*, a pleasiadapid found in Wyoming, USA. This fossil species has an opposable big toe, grasping fingers, and nails instead of claws (Bloch and Boyer 2002).



***Figure* 3.5.10.4 -** *Plesiadapiformes: Plesiadapis*

Plesiadapiformes Overview

- Flexible body, some had nails (not claws), bushy tail
- Small brain, **prognathic** face (projecting or snout-like). **diastema** (gap) between the incisors and the premolars
- Similar in lifestyle to an opossum, ate fruit and insects
- Found in North America and Europe (which were connected during this time period)
- Ranged from mouse-sized to small monkey-sized (11 lbs)

To learn more about the Plesiadapiformes, watch this optional short film: First Primates. Still want to learn more? Check out "Primate-like Mammals: A Stunning Diversity in the Treetops" (optional).

The Eocene epoch

The Eocene epoch (56-33 mya), with its warmer and wetter climate and adaptive radition of rainforests and angiosperms, sees the emergence of the first true primates, also called euprimates. Bloch and Boyer (2002) suggest that*Carpolestes* is ancestral to the euprimates. Over two hundred Eocene fossil primate species have been identified (Jurmain et al 2013). The euprimates are about the same size as modern squirrels, with prehensile hands and feet and more forward-facing eyes that gave them stereoscopic vision. These earliest primates are most like the strepsirhines and have been found in North America, Europe, North Africa, the Middle East, and Asia, which were all still connected during this time frame. Based on the fossils that have been discovered to date, we can suggest that Eocene primates were (Jurmain et al. 2013: 189):

- widely distributed
- mostly extinct by the end of the Eocene
- most are not ancestral to later primates

There are two main families of Eocene primates:

- Adapids: ancestor of strepsirhines
- Omomyids: ancestor of haplorhines

Adapids



***Figure* 3.5.10.5** - *Adapid*

- Found in North America, Europe, Asia, and Africa
- Resemble lemurs
- Forward-facing eyes, larger brain, less prognathic
- Nails (not claws), opposable thumb (and big toes)
- Flexible body
- Ate more leaves

Most recently, the discovery of *Darwinius* fossils in Messel, Germany will help to shed light on the evolution of the euprimates. *Darwinius*, dated to 47 mya, is the most complete fossil found from the Eocene. *Darwinius* appears to be a transitional adapid ancestor to monkeys and apes, which contradicts what most researchers claim that omomyids are the ancestors of monkeys and apes. As researchers other than the discoverers have opportunities to study the fossil, more will be made clear.

Omomyids



***Figure* 3.5.10.6** - *Omomyid*

- Found in North America, Europe, and Asia
- More closely resemble noctural prosimians, especially tarsiers
- Largest are around 1 kg
- Nails (not claws)
- Less prognathic, big eyes (close together), larger brain
- Ate more fruits and insects

The Oligocene epoch: the monkey/ape divergence (34 to 23 mya)

The global climate shifted again, cooling and drying, during the Oligocene epoch (33-23 mya). There is a reduction in the amount of rainforests, which retreated toward the equator, but an expansion of grasslands. So, there was an increase in **terrestrial** (ground) niches and a decrease in **arboreal** (tree) niches. While most of the fossils from this time are Old World monkeys, some are ancestors of New World monkeys; at this point the American continents had separated from Africa and Eurasia.

Parapithecids

- A good example of these was *Apidium*
- Size of squirrel, but more monkey-like
- **Quadruped** (walked on all fours)
- Dental formula 2-1-3-3 (so is possibly the **platyrrhine** or New World monkey ancestor)

Propliopithecids




***Figure* 3.5.10.7** - *Aegyptopithecus*

- A good example of these was *Aegyptopithecus* (pictured right)
- More ape-like
- Dental formula 2-1-2-3 (so is possibly the **catarrhine** or Old World monkey and ape ancestor)
- Aegyptopithecus in the news!

More on the origins of New World Monkeys: The Origins of Platyrrhine Monkeys in South America

The Miocene epoch: the Old World monkey/ape divergence (23 to 5 mya)

There were two climate shifts during this period, first there was a warm period with heavy forestation, followed by drier and cooler climates with decreasing forest and increased grasslands.

At this point, the Old World monkeys and apes split within the catarrhines. The fossils are from east Africa.

The ape ancestors had the following traits:



***Figure* 3.5.10.8** - *Sivapithecus*

- no tail
- arboreal quadrupeds (limb structure was still monkey-like)
- y-5 molars
- Some good examples of these are: *Proconsul*, *Sivapithecus* (pictured right), *and Gigantopithecus*

The Old World monkey ancestors at this point in time had the following traits:

- A tail
- Had macaque-like faces
- Were medium sized 7-11 lbs
- A good example of these is: *Victoriapithecus*

Where did primates come from?

There are three primary hypotheses about the origins of primates. The earliest hypothesis, the arboreal hypothesis, claims that the first primates evolved a suite of traits for living in trees, e.g., grasping hands and feet and stereoscopic vision. This hypothesis held sway from the early 1900s until the 1970s when the visual predation hypothesis was proposed. This hypothesis suggests that primates evolved because they provided an advantage for hunting small insects, e.g., orbital convergence that allowed for 3d-vision. Shortly thereafter, the angiosperm hypothesis was proposed, which claims that primate evolution is related to the adaptive radiation of angiosperms or flowering plants. Each of these hypotheses have been challenged over time, and it may be that each may play a part in explaining primate evolution. For more details on these hypotheses, you can check out this short article [optional]: The Origin of Primates.

In 2010, a controversial new hypothesis was proposed by Michael Heads. Heads suggests that volcanic eruptions and the breakup of the super continent, Pangea, lead to the emergence of the different lineages of primates. You can read more about Heads' hypothesis on LiveScience [optional].

References

1. Bloch JI, Boyer DM. 2002. Grasping primate origins. Science 298(5598): 1606-1610.
2. Castro J. 2013. How do fossils form? [Internet]. LiveScience [cited 2015 Jul 06]. Available from: http://www.livescience.com/37781-how-do-fossils-form-rocks.html
3. Jurmain R, Kilgore L, Trevathan W. 2013. Essentials of physical anthropology, 4th edition. Belmont (CA): Wadsworth, Cengage Learning.
4. Klaus HD. 2011. Non-human primate and human evolution. In: Hutchins M, editor. Grizmek's animal life encyclopedia: evolution. Detroit (MI): Gale; 2011. p. 299-309.
5. Lerner KL, Lerner BW, editors. Paleoecology. In: The Gale encyclopedia of science, 5th edition, Vol. 6. Farmington Hills (MI): Gale; 2014. p. 3217-3218.
6. O'Neill D. 1998-2014. Early primate evolution. Behavioral Sciences Department, Palomar College [Internet] [cited 2015 Jul 06]. Available from: anthro.palomar.edu/earlyprimates/Default.htm
7. Panell M. Fossils and fossilization. In: Lerner KL, Lerner BW, editors. The Gale encyclopedia of science, 5th edition, Vol. 3. Farmington Hills (MI): Gale; 2014. p. 1837-1842.
8. Peppe DJ, Deino AL. 2013. Dating rocks and fossils using geologic methods [Internet]. The Nature Education Knowledge Project [cited 2105 Jul 06]. Available from: http://www.nature.com/scitable/knowledge/library/dating-rocks-and-fossils-using-geologic-methods-107924044
9. Saneda TM. Dating techniques. In: Birx HJ, editor. 21st century anthropology: a reference handbook, Vol. 1. Thousand Oaks (CA): SAGE Reference; 2010. p. 352-360.

This page titled 3.5.10: Primate Evolution is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Tori Saneda & Michelle Field via source content that was edited to the style and standards of the LibreTexts platform.

- **2.10: Primate Evolution** by Tori Saneda & Michelle Field is licensed CC BY-SA 4.0. Original source: https://wikieducator.org/Biological_Anthropology.

3.6: Ethology

Ethology is the study of animal behavior. Don't confuse it with "eth**n**ology" the study of "ethnos", ethnicities, the comparative study of human cultures. I think videos are the best way to get a sense of both primate behavior and our place in the primate continuum. Captivity is a bad place to study behavior because the behavior has evolved in a certain environment, to solve problems in that environment, and you can't expect to see natural behavior outside of a natural setting, and I hate zoos because they justify the destruction of natural habit. But, some of these psychological experiments are useful to blur the line between human and non-human primate.

Note

Reintroducing Lemurs into Madagascar (7 part series with John Cleese)



2014 "Inside Ape Minds"

Behavioral Ecology

- Baboon adaptations to the savanna
- Safety in numbers, boa constrictor eats Purús red howler monkey
- The Monkey and the Snake: How the Primate Brain Reacts to Serpents

Primate Culture

Exercise 3.6.1

WATCH ORANGUTAN BEHAVIOR

- Orangutans plan ahead and share trip plans a day before leaving: Wild Apes Communicate Their Travel Plans for Tomorrow
- Chimp vs. Children
- Capuchin Monkey tool use:



Exercise 3.6.2

Watch Chimpanzee's from Bossou tool use

Article on social transmission of tool use, Sonso chimp leaf sponges

Ape Language?

Apes don't speak but they can learn sign language and symbolic keyboard languages. Book on Koko, a gorilla who was taught sign language

Ape Music?

Language is often defined as an exclusively human form of communication, but the line between human language and animal communication is not so sharp, as we see with Koko, Kanzi, and the dozens of other apes taught to use languages. Can the same be said about music? Gorillas make up little songs when they are eating. When we find regional variations of sounds produced by chimps, can we say they are like musical styles?

Exercise 3.6.3

Watch Chimpanzee's from Bossou clip leaves because they like the sound it makes, and they are bored, frustrated, or want to attract a mate (you need to turn the sound up on the video to hear it)



- Review of Chimpanzee drumming styles
- Article on Gorilla eating songs

Theory of Mind

Theory of mind refers to an individual's ability to think about what other individuals are thinking. The term has various definitions which can range from mirroring, copying another's actions, to mentalizing, predicting how another will react. Humans are definitely the best at this, but other animals demonstrate this behavior, including dogs, dolphins, elephants, some birds, and of course primates. Dogs are actually better at recognizing human pointing than chimpanzees.

* John Rubin "The Gap Between Humans and Apes"

Exercise 3.6.4

Watch a lecture by Tetsuro Matsuzawa, "Mind Reading" in Chimpanzees

Skim Charles Darwin's1872 "The Expressions of the Emotions in Man and Animals", especially look at the pictures:






***Figure* 3.6.2** - *"Chimpanzee disappointed and sulky. Drawn from life by Mr. Wood."*

Watch an amazing test of chimp short term memory:



* Apes and mirrors:



* Ape self-awareness:



Exercise 3.6.5

Read an article on contagious yawning in chimpanzees

Read an article on Chimp attitudes towards death

The three clips are arranged from agonistic (aggression) to affiliative (grooming and sex).

Agonistic Behavior

Agonistic means "aggressive", but it is usually more bluff and intimidation than physical violence. Natural selection is going to generally select for conflict resolution that avoids members of the same species injuring each other. Many primates are aggressive, but they don't kill each other very often. They learn hierarchies to avoid injury. But when push comes to shove, primates make bad pets.

- Watch a Violent Chimp Attack

Affiliative Behavior

Affiliative means "social". There is a push and pull of conflict and resolution in primate societies. Primates fight to see who's on top, and then make-up to keep the group together. Agonistic behavior helps to establish dominance hierarchies, and is usually followed by reconciliation, a kind of affiliative behavior. The most common primate affiliative behavior is grooming. We tend to think of grooming as keeping clean, but its main function for primates is social bonding.

- Read about Bonobos comfort each other
- James Fowler research on how humans chose their friends based on genetic closeness

K-selection vs. r-selection

If you say something is r-selected or K-selected, you are comparing a species or group of species to another, and comparing their strategies for growing their population. The terms come from variables in a math equation that describes how populations grow:

$$\frac{dN}{dt} = rN\left(\frac{K-N}{K}\right) \tag{3.6.1}$$

where

- N = the population density,
- r = the reproductive rate,
- K = the carrying capacity.

r-selected animals have plenty of habitat to grow into, so they crank out lots of kids and hope a few survive. K-selected animals have limitations on their resources, so they have few infants per birth, and longer birth spacing, and invest more parental care in making sure they survive. K-selection follows the human phylogenetic continuum closely. Vertebrates are more K-selected than invertebrates. Mammals are more K-selected than other vertebrates. Primates are more K-selected than other mammals. Anthropoids are more K-selected than prosimians. Hominoids are more K-selected than monkeys. Humans are one of the most K-selected species on the planet.

* Jane Goodall on chimpanzee motherhood:



* Verhulst equation of population growth or logistic growth equation

Vocabulary

- altruism
- language
- violence
- sexual dimorphism
- sexuality
- anthropomorphism


This page titled 3.6: Ethology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **5.4: Ethology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

3.7: Conservation

"Are primates going extinct?" This is a trick question. One answer is "no". The total world primate population has skyrocketed in the last 10,000 years and especially in the last 200 years. But, another answer is "yes". The total number of primate *species* has declined drastically and primate extinction is expected to continue. One primate is doing really well, at the expense of all the other primates. The primary cause is habitat loss, but a significant and very symbolic factor is that one primate is literally eating all the others.

* Jane Goodall video on conservation:



Research on Primates

Primates are valued as research subjects because their physiology is so similar to ours, but the same similarity makes captive breeding and research unethical. The Russians are good at sending primates into space, but haven't quite figured out how to get them back alive.

Habitat Loss

When we cut down forests we push primates towards extinction. This is happening to all primates around the world. The bulk of deforestation is to provide luxury food products (like hamburgers or palm oil) to meet consumer demand in the US.

Note

GRASP statement on palm oil plantations

Anthropozoonotic Diseases

Eco-tourism is generally good for primates because it adds monetary incentives for conservation, but because human and non-human primate biology is so similar (~98%), the diseases that infect us can often jump species and infect the primates we came to watch.

Note

Ecotourism can transfer disease from humans to primates

Bush Meat

I think the story of primate extinction is better told in pictures. What is important to remember is the root cause of why people resort to eating other primates: colonialism. The reason people are hungry is because of how resources are distributed. Even with overpopulation, there is enough food in the world to feed everyone. It is a question of distribution. There are the rich and the poor. Colonialism exacerbated existing class distinctions to better extract natural resources and send wealth to Europe. Colonialism led to post colonialism, where even after “independence”, patterns of corruption continue today.

It's easy for us to judge people and say we need to stop eating bush meat, but as anthropologists we strive for cultural relativity. Imagine your family was starving, and you had to choose between your children or another primate on the verge of extinction. Would you let YOUR family starve for the sake of another animal?

One of the most paralyzing factors in this issue is the history of Europe, the US, and Africa is so long and sordid, that our most well-intentioned actions often backfire. Look at the backlash to the recent Kony 2012 campaign. How can the US claim any moral superiority when they perpetuated slavery and racism? How can we claim that we have Africa's best interests at heart, after what what we did in the Congo in 1960--sending the CIA to assassinate the democratically elected leader? The sadness of these pictures goes way beyond the extinction of primates in our generation. It reflects a very difficult political situation, where everyone, especially the primates, loses.

Warning: Graphic Images

I was getting very depressed gathering these photos, but I stumbled on an innovative program that is trying to substitute beekeeping for the hunting of bushmeat, and it cheered me up for a moment. There is a crisis and we need to all do something about it, but things aren't hopeless, we just need to work quickly and do more.















https://socialsci.libretexts.org/@go/page/130010







https://socialsci.libretexts.org/@go/page/130010







https://socialsci.libretexts.org/@go/page/130010









https://socialsci.libretexts.org/@go/page/130010





- * Article on alternatives to bush meat
- * Mountain Gorilla census over 100
- * Article: Outlook is grim for mammals and birds as human population grows.
- * Legal rights for chimpanzees in the US
- * Vaccinating Lemurs:



* The Green Corridor Project includes research and reforestation to connect isolated chimpanzees to larger reserves, and help them coexist with humans:




Imagination Actions

- Participate in the latest Eyes on Apes action alerts
- Sign the Pledge not to torture slow lorises:



- Write letters to stop palm oil plantations to Herakles Farm, TIAA-CREF, another TIAA-CREF, Sara Lee, Pepsi on land used by thousands of orangutans, chimpanzees, and other primates:



- Sign this petition to ask Russia to stop sending primates into space.
- Sign this petition asking the Korea Central Zoo to stop giving cigarettes to chimps, and explain why chimps would be likely to get addicted to nicotine and suffer the same health problems as humans?
- Write a letter asking the NY Blood Center to not abandon chimps in Liberia, and sign this petition to ask IBM to pressure them.
- Comment on whether Naruto the selfie monkey should own the rights to the photos he took.



***Figure* 3.7.1** - *Self Portraits ©2008 Naruto the Celebes crested macaque, used without permission.*

This page titled 3.7: Conservation is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **5.5: Conservation** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

3.8: Lab 5 and 6- Primates

Name: _____

Section: _____

Student ID#:_____

Your group will be assigned one primate from below. You will research your primate and give a presentation next week to the class. In this presentation you must provide your classmates with the proper classification of your primate (from kingdom to species), its natural geography (no, not including zoos and private collections), its natural diet (not what it will eat if you feed it), natural group behaviors, specific diagnostic physical features, and any other traits you find interesting. The presentation is worth 10 points and should last around ten minutes. During the presentations, you will fill out the same information for the rest of the primates listed on this sheet. This sheet will be due the following week and will need all primates described as above (including those not presented). This group activity is worth 10 points, and ALL MEMBERS OF THE GROUP MUST PARTICIPATE. Should a group find that one or more of its members should be "voted extinct," please notify the instructor immediately.




Two Collared Lemurs. http://www.flickr.com/photos/joshbousel/2622755480/ Copyright: Josh Bousel



Ring Tail Lemur. http://www.flickr.com/photos/davidden/2269694846/

Copyright: David Dennis



Red Bellied Lemur http://www.flickr.com/photos/biggles621/2097775935/ Copyright: Steve Johnson



Crested Black Macaque http://www.flickr.com/photos/lipkee/616650100/ Copyright: Lip Kee



Bolivian Red Howler. http://www.flickr.com/photos/24932721@N04/2481039816/ Copyright : Nicky Jurd



Black Spider Monkeys. http://www.flickr.com/photos/wwarby/3209802292/ Copyright: William Warby



Baboon. (Primate Research Center, University of Wisconsin) Copyright: Anne Zeller




Common Marmosets. http://www.flickr.com/photos/mcwetboy/3607506156/ Copyright: Jonathan Crowe



Vervet. http://www.flickr.com/photos/mejymejy/2982075048/ Copyright: Mark Johnston



Orangutan. http://www.flickr.com/photos/jamiedfw/503804320/ Copyright: Jim Bowen



Gorilla. www.flickr.com/photos/chuqui/4737709648/ Copyright: Chuq Von Rospach



Chimpanzee. (Primate Research Center, University of Wisconsin) Copyright: Frans de Waal



Bonobo. www.flickr.com/photos/douglas...her/773973665/ Copyright: Douglas Fisher

Information Sources

- Lemur Center: Duke University: http://lemur.duke.edu/
- Primate Research Center, University of Wisconsin http://pin.primate.wisc.edu/:

3 Points of extra credit: Taxonomically completely classify the two primates below and defending your answer:



http://www.flickr.com/photos/acplinfo/4548285021/ Copyright: Allen County Public Library

This page titled 3.8: Lab 5 and 6- Primates is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Alex A. G. Taub via source content that was edited to the style and standards of the LibreTexts platform.

- **1.6: Lab 5 and 6- Primates** by Alex A. G. Taub is licensed CC BY-NC-SA 3.0. Original source: https://socialsci.libretexts.org/api/deki/files/94257/biological-anthropology-laboratory-activities.pdf.

CHAPTER OVERVIEW

4: Paleontology and Primate Evolution

4.1: Introduction to Paleoanthropology
4.2: Introduction to Paleoanthropology
4.2.1: Paleoanthropology
4.2.2: What is a Hominim
4.2.3: Miocene Epoch
4.2.3.1: Introduction to the Miocene Epoch
4.2.3.2: Sahelanthropus tchadensis
4.2.3.3: Orrorin tugenensis
4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba
4.2.3.5: Early Hominid Cranium Comparison Checklist
4.3: Pliocene Epoch
4.3.1: Introduction to the Pliocene Epoch
4.3.2: Australopithecus anamensis
4.3.3: Gracile Australopiths
4.3.4: Australopithecus afarensis
4.3.5: Australopithecus bahrelghazali
4.3.6: Kenyanthropus platyops
4.3.7: Australopithecus prometheus or africanus
4.3.8: Australopithecus africanus
4.3.9: Middle Hominid Cranium Comparison Checklist

4: Paleontology and Primate Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

4.1: Introduction to Paleoanthropology

Anthropology asks the broad question: “Who are we?” Paleoanthropology asks a more specific question: “How did we get to be what we are?"

In this part of the class we go back in time to follow the branch of the tree of life that leads to us. In the previous sections, we compared and contrasted ourselves to other vertebrates, then to other mammals, then to primates, and mostly to hominoids (apes). In paleoanthropology we start from the split that separates apes from humans and continue on that branch, and all its side branches, until we get to us.

Paleoanthropology deals with hominids (bipedal hominoids). In previous sections, we compared ourselves to living creatures that you can see running around in their natural habitat, but in the paleoanthropology section, most of our knowledge is based on data gathered through archaeology, and we focus on hominid fossils, and how to interpret them.

The next few sections are going to bombard you with specific dates, exotic places, hard to pronounce names, and plenty of Latin. Please try not to miss the forest for the trees. We need to sweat the details. The details are what paleoanthropologists use to support their very tenuous hypotheses. But make sure that when you hear a specific factoid, that you put it in context with a larger framework. So if you hear "Toros-Menalla" you should think "that´s kinda northwest of where most of those other hominid fossils were found." If you hear a species called *Homo ergaster*you think "well it's later than the australopithecines because it's of the genus *Homo*, but it's not quite us because the species name is different than *sapiens*." If you see a date for Lucy is 3.7-3.5mya you think: "Well she didn't use stone tools because those only appear around two and a half million years ago." If you read that *Sahelanthropus tchadensis* had a cranial capacity of 320-380 cc, you think "before australopithecines hominids had about the same size brain as chimps do now"

We have found hundreds of thousands of hominid fossils which represent tens of thousands of individuals. When someone claims that there is no "missing link" between apes and humans they are ignoring that huge body of evidence gathered by thousands of scientists. The real debates are about how to include those ten thousand individuals in our family tree, because these thousands of fossils are just a tiny fraction of the billions of hominid ancestors that have lived rich and important lives, but left no physical traces. A genealogical analogy is that you may know from your last name that you belong to certain clan, but you might not be able trace all your relatives back to that apical ancestor; in paleoanthropology there are many gaps.

Important factors for fossils are: dates, where found, morphology (shape), and associated tools.

hominid or hominins?

The confusion over taxonomic terminology gets worse. Recently, many scientists have redefined the term "hominid" to include great apes along with humans. This better reflects our genetic similarity to great apes. To distinguish humans and their bipedal ancestors from great apes they now call us "hominins". I don't really care which terms you use as long as you are consistent. It's always a good idea to define your terms anyway.

You don't want to get lost in the details, and always bring it back to the context of time, place, and relation to the major trends.

This page titled 4.1: Introduction to Paleoanthropology is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **6.1: Introduction to Paleoanthropology** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

CHAPTER OVERVIEW

4.2: Introduction to Paleoanthropology

4.2.1: Paleoanthropology

4.2.2: What is a Hominim

4.2.3: Miocene Epoch

4.2.3.1: Introduction to the Miocene Epoch

4.2.3.2: Sahelanthropus tchadensis

4.2.3.3: Orrorin tugenensis

4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba

4.2.3.5: Early Hominid Cranium Comparison Checklist


Thumbnail: Primate skulls provided courtesy of the Museum of Comparative Zoology, Harvard University. (CC BY 2.5; K. N'Diaye. Evolutionary History of a Gene Controlling Brain Size. PLoS Biol 2(5): e134. DOI: 10.1371/journal.pbio.0020134).

This page titled 4.2: Introduction to Paleoanthropology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

4.2.1: Paleoanthropology


WHAT IS PALEOANTHROPOLOGY?

Paleoanthropology, a subdiscipline of anthropology, is the study of extinct primates. While the majority of researchers doing this kind of work are anthropologists, paleontologists (within the discipline of geology) may also study fossil primates. The primary method used by paleoanthropologists is the analysis of fossil remains. However, they increasingly rely on other scientific disciplines to gain a better understanding of the environmental forces that played a role in our evolution, as well as the formation of the fossil record. For example, **geologists** identify processes of sedimentation and fossilization, and date fossils and their associated sediments using a variety of techniques (see DATING TECHNIQUES below). A variety of disciplines are involved in helping to reconstruct ancient environments and biological communities. **Paleontologists** identify ancient floral and faunal fossils. **Palynologists** analyze particles in ocean and lake cores, as well as pollen in terrestrial sediments (see Figure 1.2), to determine the predominant flora in a given area at a particular time. **Taphonomists** help determine how fossil assemblages were formed.



Human evolution. "Human Evolution Icon" by Magnetic Hyena is licensed under CC BY-SA 3.0.

In the 1920s, **Raymond Dart** proposed that early hominins (bipedal primates, like ourselves) found in South African caves had inhabited those caves. In addition, he interpreted puncture wounds found in some of the skulls as evidence that those hominins made and used weapons for hunting and male-male aggression. The taphonomist **C. K. Brain** argued in more recent times that either hominins fell through cracks into subterranean caves after having been cached in trees by leopards, or their bones were dragged in by rodents, such as porcupines, for gnawing. We now realize that while those early members of our tribe likely used simple tools, they were not big-game hunters or warmongers (see Chapter 15 for more information).



Pollen grains under scanning electron microscope. "Misc pollen colorized" by Dartmouth Electron Microscope Facility, Dartmouth College is in the public domain.


HISTORY OF THE DISCIPLINE

While paleoanthropology, as a formally recognized science, is fairly recent, questions and beliefs related to our origins extend back to the earliest members of our species and possibly even earlier. All modern humans living in traditional (e.g., hunter-gatherer

bands, tribes, or chiefdoms) or state-level societies have a set of beliefs associated with their origins. However, any ideas that fall outside the realm of science are part of a culture's religion and are termed creation myths.

The most influential fields to have contributed to the science of paleoanthropology are geology, biology, and archaeology. Geologists (even those who were not recognized as such, e.g., Charles Darwin) are primarily responsible for the realizations that (1) the earth is ancient, and it formed via natural processes; (2) the earth was originally covered with water, and life began in that "primordial sea"; (3) life on earth originated with simple forms, with some descendent species becoming more complex over time, as can be seen in the fossil record; (4) species change or go extinct in response to environmental change; (5) new species are the result of a portion of a population adapting to new or changed environmental conditions; (6) the same forces, such as volcanic eruptions, that operate today are those that shaped the earth and caused changes in the fossil record via extinctions and speciation events; and (7) layers and deposits are continually developing or eroding so that organisms are buried and fossils come to light, respectively. The idea that the same forces that operate today are those that shaped the earth and caused changes in the fossil record is termed **uniformitarianism**. Charles Lyell coined the term and is heralded as the father of modern geology. He greatly influenced Darwin and thus contributed to Darwin's synthetic view of the evolution of life on earth. Geologists use various methods to date fossils or fossil-containing sediments and have developed a **chronology** (i.e., a timeline) for the earth as a whole, as well as depositional layers in areas where fossils have been discovered.


Biologists and geneticists have refined the theory of **evolution by means of natural selection** by determining how traits are inherited. Scientists from a variety of disciplines have classified the known species of the world based on evolutionary relationships (also see Chapter 2).



Charles Darwin. "Charles Darwin 01" by J. Cameron is in the public domain.


Archaeology has played and continues to play a strong role in paleoanthropology via the study of the **archaeological record**, that is, the record of past human activity via cultural remains and **anthropogenic** (human-induced) changes to the environment. Thomas Jefferson has been referred to as the first archaeologist, in that his methods were more scientific than his fellow **antiquarians**. Antiquarians tended to be after the "goods," without regard for careful interpretation of the archaeological record. Most would be considered looters by today's standards. They took items of great cultural and historical significance for personal or museum collections. Some items have been returned to their countries of origin, but the damage is done when the archaeological record is disturbed or destroyed. Once an item has been removed from the area where it was found, scientists can no longer learn from its context, for example, from associated artifacts or the location of the artifact in geographic space and time.

Archaeologists and geologists played a key role in recognizing that "stones and bones" were evidence of earlier hominin activities. In addition, the fact that some of the bones were from extinct animals supported the idea that humans had been around for a long time. Archaeological methods of excavation and analyses, such as the **provenience** (i.e., the three-dimensional location within a site) and association of **artifacts** (i.e., portable human-made or altered objects), help archaeologists and paleoanthropologists reconstruct past behavior. Just as taphonomy plays a role in determining how fossil assemblages came to be, it is also useful for archaeological assemblages.



Eugène Dubois. “Eugene Dubois” is in the public domain.

According to *Merriam-Webster Online*, the first known use of the term “paleoanthropology” occurred in 1916. However, the earliest paleoanthropologists were not labeled as such and came from a variety of occupations, such as anatomists and physicians. The first hominin fossils discovered were the neandertals in the 1800s. However, paleoanthropologists disagreed about whether neandertals were ancestors of humans or were modern humans. **Eugène Dubois** was the first person to intentionally search for a fossil hominin. He went to Asia with the sole purpose of finding evidence that humans evolved there, as was the reigning belief in Western Europe. In 1891, he discovered a skull cap (known as a **calotte**) and femur on the **Solo River** in **Trinil**, Java. More discoveries in China and Java during the first half of the 20th century supported the Asian origin theory until Raymond Dart and his contemporary, **Robert Broom**, began discovering much more ancient material in South African quarries and caves. Further discoveries by **Louis** and **Mary Leakey** in East Africa cemented Africa as the birthplace of humanity, and the race to find human origins and ancestors was on.



Louis Leakey. “Louis Leakey” is in the public domain.

RECONSTRUCTING PALEOENVIRONMENTS

A variety of tools can be used to determine the type of environment past species occupied. As mentioned, paleontologists can use floral and faunal analyses and what they know about ancient species or their extant relatives to determine environment type, for example, the presence of aquatic-, grassland-, and/or forest-dwelling species. Palynologists examine particulates in aquatic and terrestrial strata (i.e., layers or sediments) to do the same, primarily focusing on floral analyses. A variety of isotopic tools can be used to categorize floral and/or faunal communities at a given site, such as hydrogen, oxygen, and carbon isotope fractionation and nitrogen isotope ratios. For example, calcium-rich remains such as eggshells, bones, and teeth can be analyzed isotopically to determine what types of vegetation those animals consumed and hence, the type of environment in which they lived. The strontium-to-calcium ratio in bones and teeth can be used to determine the amount of animal versus plant matter in the diet. Based

upon that technique, scientists now believe that paranthropines, a group of hominins in East and South Africa dating from the early- to mid-Pleistocene (see Chapter 16), ate some animal matter. However, whether they were consuming insects or larger prey is not known.

For more information on the aforementioned methods, consult Henke W, Tattersall I. 2006. *Handbook of paleoanthropology*. New York (NY): Springer.

DATING TECHNIQUES

Dating techniques fall into two categories, relative and absolute. **Relative dating techniques** (1) ordinally rank **strata** relative to one another through time (see Figure 1.6) or (2) use what is known about deposits in one area, such as volcanic ash or lava, to relatively date deposits in another area. Jefferson is credited with the **Law of Superposition**, which posits that as you go deeper into the earth, layers get older, as long as strata have not been disturbed due to human, animal, or geological activity. Thus artifacts or fossils found in one layer are either older or younger than those in a deeper or shallower layer, respectively. **Absolute dating techniques** use similarities in (1) floral and faunal assemblages or (2) sedimentary and/or chemical composition of deposits in order to match those of unknown age with those of known age and/or order the progression of environments, organisms, and climatic and geological activity within or between regions.




Diagram illustrating cross-cutting relations in geology. These relations can be used to give structures a relative age. Explanations: A – folded rock strata cut by a thrust fault; B – large intrusion (cutting through A); C – erosional angular unconformity (cutting off A & B) on which rock strata were deposited; D – volcanic dyke (cutting through A, B & C); E – even younger rock strata (overlying C & D); F – normal fault (cutting through A, B, C & E). Cross-cutting relations by Woudloper is licensed CC-BY-SA.

Absolute or chronometric dating techniques yield approximate dates in years **BP**(before the present) or **BCE** (before the Common Era). BCE and CE (Common Era) retain the BC/AD system of dating without the religious connotation. An abbreviated way to refer to a certain number of years ago, especially when considering the fossil record, is kya or mya (thousands or millions of years ago, respectively), thus eliminating all of those ungainly zeroes! While BP makes more sense in that you do not need to add 2,000+ years to the date, most people are accustomed to the BC/AD system, thus explaining the common use of BCE. The best-known absolute dating techniques are **radiometric dating methods**, for example, Carbon-14 (^{14}C). They are used to measure the half-life or replacement of radioactive elements in organic or fossil material or the layers in which they are found. Since those methods are time-limited and/or context-specific, the most appropriate technique(s) must be chosen based on a variety of parameters. The following techniques use radioactive decay for dating purposes:

[NOTE: For more information on the following methods, consult Henke and Tattersall (2006), *Handbook of Paleoanthropology*; and/or Davis (2009), "Other Dating Methods": www.geo.arizona.edu/palynology/geos462/11datingmeth.html.]

Carbon-14 dating (≤60 kya) measures the remaining ^{14}C in organic materials (i.e., carbon-containing). Since plants use carbon dioxide for photosynthesis, they contain all three isotopes of carbon (^{12}C, ^{13}C, and ^{14}C) in the approximate ratios present in the atmosphere. Animals eat plants and thus, at any particular time, they will all have approximately the same amount of ^{14}C. Once they die, they no longer accumulate carbon. The level of the more stable ^{12}C can then be compared to the remaining ^{14}C in organic remains to determine when they died. The half-life of ^{14}C is ~5,700 years, that is, half of the ^{14}C will have been lost in a specimen in that amount of time.

Uranium series dating (≤500 kya) examines the relative levels of two elements, Uranium-234 and Thorium-230, resulting from the former's decay into the latter. It is used to date calcium carbonate in coral and shells.

Potassium-Argon (K/Ar) and **Argon-Argon** (Ar/Ar) dating both measure the ratio of one isotope to another via the process of radioactive decay, Potassium-40 → Argon-40 and Argon-40 → Argon-39, respectively. They are often used to date volcanic layers but can also be used on other soil components, such as clay. While the age range for both methods may be reported to be unlimited, K/Ar dating is not useful for "young" materials because the half-life of potassium is so long—1.26 billion years.

Other methods that also rely on radioactivity are:

Electron spin resonance (ESR) (up to "a few" mya) examines the pattern of electrons that have "spun" out of their original location in mineral compounds (e.g., calcium compounds), leaving empty spaces behind, due to exposure to environmental radiation. Tooth enamel is the most useful application of ESR in paleoanthropology, but ESR can also be used to date quartz particles in sediments (Wagner 2006).

Fission track dating (20 mya—>10 kya) measures the number of "tracks" (pitting) in mineral compounds that result from the energy released when Uranium-238 spontaneously fissions over time. This method can be used to date a variety of minerals, such as mica, as well as products of volcanic (e.g., obsidian) and meteoric activities (Davis 2009; Wagner 2006).



Apatite crystals can be used in fission-track dating. "Apatite crystals" by OG59 is in the public domain.

Thermoluminescence (300–1 kya) measures radioactive decay particles in mineral compounds. It is useful for compounds that were exposed to intense heat (e.g., volcanic eruption) at some known point in time, when the "radioactive clock" was reset to zero and decay began anew. Thermoluminescence can be used to date artifacts (e.g., ceramics) and features (e.g., hearths), as well as products of sedimentation (e.g., speleothems, which are mineral deposits that form in caves) and volcanic activities (e.g., tephra, which are fragments from volcanic eruptions) (Davis 2009).

The following methods do not rely on radioactive activity but rather organic processes:

Dendrochronology uses tree rings in fossil or charred wood to date artifacts or fossils found in association with the wood. Each year, trees produce a new layer of peripheral tissue. When climatic conditions are favorable, more tissue is deposited and a thicker ring results, and vice versa. A cross-section of the tree tells the history of its growth (see Figure 1.8). However, in order to use dendrochronology as a dating method, a **chronology** (temporal record) needs to be constructed for a given region, in this case a map of the annual growth rate back through time. Living trees and dead wood can be used as long as there is overlap in ring patterns between them.



Dendrochronology: tree ring dating. "Dendrochronologie" by Stefan Kühn is licensed under CC BY-SA 3.0.

Amino acid racemization (2 mya–2 kya ± 15%) measures the ratio of two forms of an amino acid, one produced while an organism is alive and the accumulation of a second form after death. If the ambient temperature at the time of death can be approximated, the specimen can be dated and vice versa (Davis 2009).

Paleomagnetism (hundreds of thousands–millions of years, Fagan 2000) measures past changes in the earth's paleomagnetic fields that are preserved in some common minerals found in rocks and sediments. Since scientists have established a chronology of those changes, the materials can then be given approximate dates as to when they formed. When paleomagnetism is used to date archaeological materials, it is termed archaeomagnetic dating.

Obsidian hydration (100–1 mya) is used to date volcanic glass, that is, obsidian, by examining the amount of hydration that has occurred due to exposure to the elements. It is useful in dating obsidian artifacts as well as glacial and volcanic activities (Davis 2009).

Surface or Cosmogenic Nuclide Exposure Dating measures the amount of time that rocks have been exposed to the elements. It can be used to date glacial, lava, and rockslide movements and damage from extraterrestrial activities (e.g., solar flare-ups or meteorites) (Davis 2009; Wikipedia contributors 2015i).

ADDITIONAL STUDENT RESOURCES

1. Henke W, Tattersall I, editors. 2006. *Handbook of paleoanthropology*. New York (NY): Springer; [accessed 2015 Aug 15]. http://www.evolbiol.ru/large_files/handbook_paleoanthropology.pdf.

This page titled 4.2.1: Paleoanthropology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **1.1: Paleoanthropology** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.2.2: What is a Hominim

We are hominins, as are all those bipedal apes that came before us. Figure **4.2.2.1** shows most of the hominin species through time, some of which we are descended from and some that are side branches in our tribal tree!




Figure **4.2.2.1**: Hominin species through time. Model of the phylogeny of Hominini over the past 10 million years. (CC BY-SA 4.0; Dbachmann)

ANATOMICAL TERMS

To adequately understand a discussion of hominin evolution and appreciate changes over time, some basic anatomical information is necessary. It is also necessary to distinguish **primitive** or ape-like skeletal characteristics from those that are **derived**, i.e. those that arose later in time. Any discussion of anatomy assumes that one is speaking of a body in **anatomical position**, i.e. facing forward if standing or supine (face up) if lying down, with palms forward or up (see Figure 5.2—5.4). When referencing particular structures or regions of the body, we make use of the following terms:

- **Superior**—closer to the top of the head.
- **Inferior**—closer to the soles (or plantar surfaces) of the feet.
- **Medial**—closer to the midline of the body.
- **Lateral**—closer to the far right or left of the body, relative to the midline.

The following two terms are used in reference to the limbs:

- **Proximal**—closer to the base of a limb.
- **Distal**—closer to the end of a limb.

You may hear your professor use the above terms when describing aspects of human or fossil specimen anatomy.

Note: If you are thinking about a career in paleoanthropology, get as much training in anatomy as possible, even while an undergraduate. Anthropology and biology departments may offer courses in human anatomy, human osteology, forensic anthropology, and the like. Finds like “Lucy” (*Australopithecus afarensis*) come along once in a lifetime. Most discoveries consist of little pieces of fossilized bone or teeth and thus knowledge of human anatomy is crucial for recognizing possible landmarks on the remains.

HUMAN GROSS SKELETAL ANATOMY

The terms *gross anatomy* and *macroscopic anatomy* refer to the study of structures that are visible to the naked eye. In this section, we will consider whole bones versus the individual parts of bones and only those bones that are external, as those are most relevant in a discussion of bipedalism and human evolution. While the anatomy of the lower limb takes precedence in a discussion of

bipedalism, other parts of the body also changed over time. The anatomy of the skull is particularly important because the skulls and faces of hominin species changed over time and geographic space.




Figure **4.2.2.2**: Human axial skeleton (shaded). "Axial skeleton diagram" by Mariana Ruiz Villarreal is in the public domain. Human appendicular skeleton (shaded). "Appendicular skeleton diagram" by Mariana Ruiz Villarreal is in the public domain.

We will begin with regional anatomy. The **axial skeleton** consists of the head and torso. Regionally it is divided into the skull, thorax, and spinal column. The spinal column is divided into the seven cervical (neck), twelve thoracic (thorax/chest), five lumbar (lower back), five sacral (also known as sacrum), and four coccygeal (also known as coccyx or tailbone, they may also number three or five) vertebrae. The limbs and shoulder and hip regions make up the **appendicular skeleton**. The upper limb is also known

as the arm or forelimb. It is divided into the arm (upper arm), forearm, wrist, and hand and fingers. The lower limb is also known as the leg or hindlimb and is made up of the thigh, leg (equivalent of forearm), ankle, and foot and toes.

Individual bones of the axial and appendicular skeletons are labeled in Figure 5.4. They will not be discussed here.




Figure **4.2.2.3**: Human skeleton. "Human skeleton front en" by Mariana Ruiz Villarreal is in the public domain.

THE EVOLUTION OF BIPEDALISM

There are a variety of theories as to how bipedalism evolved and why it proved to be so successful for early hominins. One early idea suggested that by standing up, our ancestors would have been able to see above the grass and thus avoid predation. Baboons and patas monkeys provided living models for hypothesizing the environmental stresses early hominins might have faced on the open plains of Africa. While they likely traveled through open areas, we now know that the earliest hominins were exploiting forest

resources, as evidenced by their thinner molar enamel, relative to later hominins. There were also theories that involved the freeing up of the hands to make and use tools and for carrying resources to a safe place or home base. C. Owen Lovejoy believes that bipedalism allowed males to provision mates with resources (Lovejoy, 1981). Those males with the most advanced bipedal capabilities would have had an increased chance of mating and possibly offspring survival, and thus bipedalism would have spread throughout the population. While Lovejoy makes a good case for how a trait could be favored in a population, it is not clear why females would have needed to be provisioned unless their offspring had already lost their ability to hang on with their feet, and hence became a burden to foraging. We now know that ardipiths, while bipedal on the ground, had a divergent hallux, so that young animals could likely have hung on to their mothers in trees like modern primates. Although we cannot know for sure whether they were covered with hair, I can only speculate that when on the ground, young infants may have hung by clasping the fur on their mother's ventrum (anterior aspect of the trunk) while the mother supported their bottom. As the baby matured, it may have clasped its hands around the mother's neck or chest and hung on to her fur with its feet and later ridden "piggy-back" like modern quadrupedal monkeys and apes. Eventually, it would have walked beside her from place to place. However, if resources had become extremely scarce, bipedal males may have ventured out onto dangerous ground for resources with which to provision their mates. Another theory that sees males as being the impetus for bipedalism suggests that males may have been more terrestrial and females more arboreal, i.e. a case of niche partitioning, like gorillas and the mandrill and drill monkeys, where males forage on the ground and females and young spend more time in the trees. Other theories also suggest that bipedalism was a response to the changing nature of the resource base. For example, Meave Leakey and Kevin Hunt (a theory known as Hunt's Postural Feeding Hypothesis, Hunt 1996) believe that the ability to stand on two legs for long periods of time would have facilitated picking fruit from the terminal branches of low, scrubby trees in the increasingly open habitats of East Africa. While the aforementioned theories are not mutually exclusive and there was likely a synergistic effect that resulted from our ancestors' changing locomotor capabilities, a highly plausible model suggests that it was our ability to break out of the "ape habitat" that facilitated our evolutionary success.




Figure **4.2.2.4**: Running man. "Nude Man Running" by Eadweard Muybridge is in the public domain.

The following items will help us to better understand this great "step" in our ancestry:

- Apes are descended from an arboreal climber from the early Miocene of Africa.
- The ancestor of the African great apes possessed a suspensory hanging adaptation and thus had an upright trunk that was wide and shallow; mobile shoulder and wrist joints; long arms relative to leglength; long, curved hand and foot bones; and an opposable big toe (hallux). The next section explains how that thoracic morphology facilitated our ancestors' bipedalism.
- The common ancestor of chimps and hominins was likely a semi-terrestrial quadruped that was adapted to climbing, feeding, and sleeping in trees, as well as moving and foraging on the ground.
- The late Miocene of Africa was marked by climatic change that spurred floral and hence faunal changes. Equatorial Africa was cooler and increasingly drier than in earlier times. The Indian subcontinent continued to move north, resulting in the uplifting of the Himalayan mountain chain that produced a rain shadow, i.e. moisture-laden clouds that would have previously drifted down into Africa now lost their moisture on the mountains. Drying winds and cooler temperatures resulted in forest reduction and fragmentation in Africa, splitting and isolating resident faunal populations.
- The majority of African ape species went extinct due to loss of habitat.

- Bipedalism is an efficient means of locomotion for covering distance on fairly level ground.
- A more vertical posture reduces the surface area exposed to solar radiation in a more open environment. It also raises a large percentage of the body away from the hot ground, where it is exposed to cooling breezes.
- The ancestors of the hominins adapted to the changing environment by becoming bipedal on the ground. Over time they developed a more efficient heat exchange system for their bodies (sweating) and brains (large venous sinuses for rapid blood movement) and lost much of their body hair.
- Based upon early hominin teeth, they were generalists like chimps, likely getting the majority of their carbohydrates and fats from fruit, protein from young leaves, and possibly fat and protein from animal matter, e.g. social insects (chimps and gorillas eat a lot of ants and termites) and animals caught opportunistically. No early hominins exhibit the same degree of canine size or sharpness as chimp and gorilla males. Their canines stay sharp via a honing (sharpening) action with the first lower premolar, termed a **sectorial premolar** due to its unicuspid morphology. The combination of the action and morphology of the two teeth is termed a "**honing complex**". Males' canines are exceptionally large and, in combination with their high degree of sexual dimorphism, are used to compete with other males for access to females. Of course, they are also useful for predator defense. Thus, if the common ancestor of chimps, gorillas, and hominins possessed a premolar honing complex, as seems likely, the early hominin fossil evidence suggests that they were losing their fighting teeth. In addition, fossil hominins do not exhibit the same degree of sexual dimorphism as seen in chimps and especially gorillas. Later chapters contain more information on hominin socioecology.

Most apes went extinct as their habitats dwindled and they competed for limited resources. However, with an efficient means of locomotion to move between forest patches when resources became depleted, hominins could continue to exploit those resources to which they were adapted. They also likely evolved new capabilities for exploiting newly encountered food items as they moved through and between ecozones. Loss of habitat and resources often leads to local extinctions. By expanding their home ranges and dietary niches, hominins survived while the majority of their close relatives did not.

BIPEDAL ANATOMY

See Figure 5.4 for individual bones.

The majority of bipedal characteristics involve the hip (or pelvic girdle) and lower limb. However, as will be seen below, certain skull and trunk characteristics are also adaptations for bipedal locomotion. In addition, we have inherited many aspects of our upper bodies from our ape ancestors and those will all be discussed in the following sections.

SKULL

The skull consists of the bones of the braincase and face and the mandible (lower jaw). The **foramen magnum** is the hole in the occipital bone situated in the base of our skulls (see Figure 5.6). It is where our spinal cord exits the cranial vault. In hominins, the foramen magnum is positioned more anteriorly than in the other apes because our head sits on top of our vertebral column. Thus while the earliest hominins had very ape-like faces, the position of the foramen magnum shows that they were bipeds.




Figure **4.2.2.5**: Foramen magnum indicated from inside skull vault. "Crane4 Foramen magnum" by Didier Descouens is licensed under CC BY-SA 3.0. Bones of the skull. Plate 188 from Gray's Anatomy. "Side view of the skull" by Henry Vandyke Carter is in the public domain.

VERTEBRAE

Ape spines are not as flexible as monkeys' spines, giving us better upper body support since we are more upright than most other primates. Our vertebrae increase in size and robusticity from top to bottom so that our lumbar vertebrae are very large; they sit on the fused vertebrae of the sacrum, which is firmly attached to the hip bones. The sacrum is large and broad and curves inward (as does the coccyx) to help support the organs. Thus our spinal column is a strong supporting structure for the upper body. We hominins have two larger curves in our backs relative to the other apes, the **cervical curve** and the **lumbar curve**. The fact that our heads are more upright than nonhuman apes means that the cervical vertebrae must form a more concave curve, i.e. the superior aspect of our neck is arched back relative to theirs (Figure **4.2.2.6**). The more pronounced lumbar curve forms when we stand up and begin toddling about. The joints between the lumbar vertebrae are easily strained and it is thus important to maintain strong back and abdominal muscles throughout life, to aid in the stability of the region.




Figure **4.2.2.6**: Chimp (left) vs. human spine (right). Illustration by Keenan Taylor

THORAX

The thorax consists of the sternum, ribs, and thoracic vertebrae. The ape **thorax** is adapted to climbing and swinging in trees. It is wide (right to left) and shallow (front to back) relative to quadrupedal monkeys, which have a narrow and deep thorax like those of dogs. While the morphology of the thorax was originally adapted to arboreal climbing, the upright trunk also allowed for bipedal locomotion. In addition, the shallow nature of the thorax brings the center of gravity closer to the vertebral column for better weight concentration and transfer. Apes are better bipeds than most nonhuman primates. They can walk bipedally for short to moderate distances, depending on the species, but it is not efficient and they cannot maintain it for very long.

SHOULDER GIRDLE

The shoulder girdle consists of the clavicle, scapula, and humerus. The shoulder joint of extant nonhuman apes and early hominins is and was angled upward, demonstrating the arboreal ancestry of those hominins and, in combination with their long, curved fingers, it suggested that they could ascend and climb about in trees. Our **clavicles** stabilize our shoulder joints for swinging and hanging. The triangular shape of the **scapula** (shoulder blade) is more equilateral than that of a quadrupedal monkey's, which is more elongated. Our scapulae are more mobile than those of a typical quadruped and the articular surface of the scapula, i.e. where the round head of the humerus articulates (makes contact), is shallow and allows us to rotate our arms at the shoulder. This **suspensory hanging complex** of clavicle, scapula, and humerus morphology (also elbow and wrist adaptations—see below), along with the muscles and connective tissue involved, allows us to climb, hang, and swing by our arms.

UPPER LIMB

The upper limb consists of the humerus of the arm, the radius and ulna of the forearm, the eight carpal bones of the wrist, the five metacarpals of the body of the hand, and the phalanges of the digits (three per finger and two per thumb, or pollex). As mentioned, the head of the ape **humerus** is round, resulting in a very mobile shoulder joint. We can fully extend our arms at the **elbow** in order to hang or swing, whereas quadrupeds cannot. In addition, we can supinate and pronate our **forearm**, i.e. move our hand palm up or palm down. This movement is possible because the head of the radius is a concave disk that rotates on a ball-like structure termed the capitulum on the **distal** humerus (Figure 4.2.2.6). Our **wrist** joints are very flexible, allowing us to rotate and twist our hands in a variety of ways. Early hominins had short legs, long arms, and curved fingers. Over time, hominin legs lengthened so that their **intermembral index** (IMI) became reduced. The intermembral index is the ratio of armlength to leglength, calculated by the following equation:

$$IMI = \left(\frac{armlegnth}{leglength}\right) \times 100 \qquad (4.2.2.1)$$

If an animal has long legs and short arms (like us), they have a low IMI and vice versa; if fore- and hindlimbs are approximately equal in length, such as in terrestrial quadrupeds, the IMI will be close to 100.




Figure 4.2.2.7: *Human elbow. "Bones of the elbow" by Rob Swatski is licensed CC-BY-NC.*

HIP OR PELVIC GIRDLE

Our shoulders are somewhat analogous to our butts! This is because muscles originating from outside the limbs are crossing the joints to insert upon and move the limbs. Because we are bipeds, it is not as apparent as it is in quadrupeds, whose fore- and hindlimbs move similarly. However, if you compare the humerus and femur and the muscles that cross the respective joints (deltoids and gluteal muscles, respectively), you will definitely see similarities.

Our pelvis is very unique and interesting. It has changed significantly from an ape pelvis (see Figure 5.10). The pelvis is made up of three bones: the two lateral bones, termed innominates or os coxae, and the sacrum. Collectively, they form a basin-like structure that holds our internal organs while providing support for our upper bodies. Each **innominate** consists of three bones that fuse during development: the **ilium**, **ischium**, and **pubis** (Figure 4.2.2.6). They meet at the hip joint. Hominin innominates became shorter and broader, so that the ilium wraps around laterally from an earlier, more posterior position. This changed the orientation and action of our hip muscles, allowing for our striding gait and the ability to balance our weight on one fully extended leg while the other leg is in the swing phase. A portion of the *gluteus maximus* muscle inserts behind the hip joint in hominins (versus lateral in chimps), and thus instead of abducting the femur (moving it out laterally, as when doing jumping jacks), it changed to a powerful hip extensor (backward motion) for running.




Figure 4.2.2.8: *Left innominates of chimp (left), australopith (center), and human (right). Illustration by Keenan Taylor.*

Figures 5.11 and 5.12 illustrate the landmarks that are adaptations to bipedalism. The **iliac crest** is long and curved, as the bone wraps around laterally. The **iliac blade** is short but expanded horizontally. The iliac blades are buttressed or reinforced to handle the stress of our upper body weight. The thick section of bone, termed the **iliac pillar**, can be seen running from the iliac crest (at the iliac tubercle) down behind the hip joint. The articular area, termed the **acetabulum**, is large and deep, providing a stable socket for

the ball-like head of the femur. The joints of the pelvis are very strong and relatively immobile (compared to the shoulder joint). The **sacroiliac joint**(between the ilium and the sacrum) is large and more posterior and proximal to the hip joint than in quadrupedal apes. Together with the strong **pubic symphysis** (anterior joint where the pubis portions of the two innominates meet), these characteristics make for a very stable supporting structure.




Figure 4.2.2.9: *Human pelvic morphology: The three fused bones of the innominates (ilium, ischium, and pubis), sacrum, and coccyx. Plate 241 from Gray's Anatomy. "Male pelvis" by Henry Vandyke Carter is in the public domain.*




Figure 4.2.2.10: *Ossified joints of the bones of a left innominate (lateral view). Plate 237 from Gray's Anatomy. Note where the ilium, ischium, and pubis articulate at the acetabulum.*
"Plan of ossification of the hip bone" by Henry Vandyke Carter is in the public domain.

LOWER LIMB

The lower limb consists of the femur of the thigh, the tibia and fibula of the leg, seven tarsal bones of the ankle, five metatarsals of the body of the foot, and phalanges of the digits (three per toe and two per big toe or hallux). The head (proximal ball-like structure) of the hominin **femur** is large. The femur angles medially (inward) from hip to knee, so that our upper body weight is transferred down through our hip joints to our knees. This is termed the **carrying** or **bicondylar angle**. The knees of quadrupedal apes are directly below the hip joint, so there is more strain on the knee joints when they walk bipedally (Figure 4.2.2.11). We have two bulbous structures on the bottom of each femur, termed condyles. The innermost condyle, the **medial condyle**, has lengthened in hominins so that the femur sits on top of the flat tibial surface. If this were not the case, the medial condyle would not make contact with the **tibial plateau**, due to the bicondylar angle. Weight is transferred through our innominate, hip joint, and **lateral condyle** of the femur (Figure 4.2.2.11, "Human"). Unlike apes' knees that are chronically flexed, our knees are capable of full extension; each locks into place when the other leg is in swing phase, giving us a stable supporting leg. Each *gluteus medius* muscle alternately supports the opposite side of the torso and pelvis, so that it does not slump on the unsupported side.




Figure 4.2.2.11: *Pelvic girdle and weight loading on knee joint. Illustration by Keenan Taylor.*

Our feet (see Figure 5.14) have changed dramatically from a mobile, grasping structure to a rigid, supporting one. The tarsal bones of the human ankle are large and robust for support. The joint between the distal tibia and fibula is fairly immobile, so that the two bones are firmly lashed together. Together, they articulate with the **talus** (most superior tarsal bone) in a hinge joint. We have lost much of the mobility of an ape foot and thus have become less agile in climbing over time. The calcaneus or heel bone is very large and robust and, along with the ball of the foot (distal end of the first metatarsal) and the area below the baby toe (fifth metatarsophalangeal joint), forms a tripod structure. Our feet have three arches for support, shock absorption, and forward propulsion; they are the medial and lateral longitudinal arches and the transverse arch. Hominin toes became shorter and less curved over time. While shorter, the hallux became more robust and lost its degree of divergence and opposability, by coming into alignment with the lateral four toes, i.e. digits II through V.




Figure 4.2.2.12: *Human foot bones. Plates 290 and 291 from Gray's Anatomy. "Skeleton of foot. Medial aspect" and "Skeleton of foot. Lateral aspect" by Henry Vandyke Carter are in the public domain.*

HOMININ EVOLUTIONARY TRENDS

Just as there are evolutionary trends that characterize primates, there are also trends that characterize the hominins, i.e. traits that developed over time and occur to varying degrees in the various hominin species (Table 4.2.2.1). While all of the morphological

changes involved with bipedalism are hominin trends, there are also relevant characteristics in other parts of the body.

Table 4.2.2.1: **HOMININ EVOLUTIONARY TRENDS**

- Bipedalism and related morphological adaptations.
- Encephalization and corresponding prolonged juvenile dependency.
- Loss of craniofacial robusticity and muscle attachments.
- Reduced prognathism.
- Reduction in size of dentition.
- Molarization of premolars.
- Increased manual dexterity.
- Elongated legs and increased stature.
- Increased reliance on culture, technology, and language.
- Increased reliance on meat in the diet.

Most notable, our ancestors and their relatives became increasingly more intelligent. Our brains have increased in size more than four-fold, from a more chimp-sized brain (<400 cc) in the earliest hominins to a mean of ~1400 cc. This likely occurred in response to environmental stresses as well as competition with other hominins for resources. Skull size and shape changed in response to **encephalization**, i.e. increasing brain size.

Brains are very costly organs and researchers believe that in order for brain size to have increased, there would have had to have been a corresponding decrease in some other costly organ system. It is hypothesized that a higher quality diet allowed the hominin gut to shrink and, in turn, the brain to expand. Marked encephalization in the hominin lineage began with the first members of our own genus: *Homo*. While there is some evidence that earlier species (e.g. australopiths) manufactured tools, there is solid evidence that early *Homo* did, and the archaeological record suggests an increasing reliance on meat in their diet. While the dimensions of the thorax remained wide and shallow, the shape of the rib cage changed from a more conical, ape-like thorax that could accommodate a large gut, to our more barrel shape. As our brains became bigger, our ancestors gave birth to increasingly **altricial** (helpless) young that continued their development outside of the womb. Our offspring require protection and care for a much longer period than those of the other great apes, i.e. we have a prolonged juvenile dependency period. At least by the time of the earliest members of the *erectus* grade at 1.8 mya (see Chapter 27 and subsequent chapters), it is thought that infant requirements necessitated assistance from mates and/or relatives.

Many of the early hominins had pronounced, forward-oriented jaws, termed **prognathism** (*pro* = forward; *gnath* = jaw). Over time, hominins became more flat-faced, or **orthognathic**. While extant African apes retain primitive prognathism and the shearing/honing dental complex, hominins lost those pronounced canines, as well as the gaps in the corresponding tooth rows—termed **canine diastema** (singular) or diastemata (plural)—that allow apes to close their jaws. The hominin first lower/mandibular premolar developed into the characteristic **bicuspid** (two-cusped) morphology over time, as opposed to the apes' unicuspid sectorial premolar. The loss of the honing complex may have been in response to reduced competition for females as a result of male-female pair-bonding. Once the jaws no longer locked into place, our ancestors were free to grind their food, aiding in its digestion.

The size of jaw and neck muscle attachment sites on the skull became reduced in the hominin lineage over time, along with a reduction in the size of the teeth and craniofacial robusticity. The action of the powerful *temporalis* muscle (a muscle of **mastication**) changed from primarily acting on the front of the jaw, allowing apes to clamp their jaws powerfully shut during fighting, to acting on the molar region for grinding food (see Figure 5.15). The origin of the *temporalis* muscle moved over time from the midline of the top of the skull to a more inferior position on the lateral aspect of the frontal and parietal bones (see Figure 5.15), due to the reduction of the sagittal crest and decrease in *temporalis* power in hominins.




Figure **4.2.2.13**: *Temporalis muscle: Originates on frontal, parietal, and temporal bones and inserts on mandible. (Zygomatic is shown as having been cut to reveal underlying muscle. Plate 382 from Gray's Anatomy. "The temporalis" by Henry Vandyke Carter is in the public domain.*

Leglength increased as our hominin ancestors came to rely less on an arboreal environment and became primarily terrestrial. The pronounced increase in leglength and stature seen in the *erectus* grade (see Chapter 27), relative to past species, allowed them to cover vast distances efficiently and they moved from Africa to Asia in a relatively short period of time.

Hominin fingers became shorter and lost their curvature over time. By the time of the australopiths, hands had become more dexterous. There is evidence that *Australopithecus africanus* possessed a **"power" thumb**, giving them increased abilities for holding objects in one hand while manipulating or working them with the other hand. This was necessary for our ancestors to have made and efficiently used tools. The first metacarpal was adapted for such activity, and the **intrinsic thumb muscles** (within the hand versus extending from the forearm into the hand) were well-developed. There is some controversial evidence that by the time of the later robust australopiths, or paranthropines, expanded fingertips had appeared. Thus they may also have possessed the enlarged tactile pads and increased sensitivity and vascularization that allows for modern humans' fine degree of manual dexterity.

Finally there is the cultural side of things. We are like no other species that has ever lived. We are unique in our ability to practice, produce, and pass on culture. Our brains are predisposed to the acquisition of language and culture. Thus, the final evolutionary trend seen in the hominins is an increased reliance on culture and language over time.

HOMININ GROUPS

The hominins can be divided into five groups that will be covered in subsequent chapters, based on shared characteristics and/or phylogenetic affinity:

1. Earliest bipeds: *Orrorin* and possibly *Sahelanthropus*.
2. Forest-adapted bipeds with relatively thin molar enamel: ardipiths.
3. Bipeds that exploited a more open and drier niche with thick molar enamel: gracile australopiths, such as *Australopithecus anamensis* and *afarensis*.
4. Descendants of gracile australopiths with heavy masticatory apparatus, adapted to exploit tough or hard foods when necessary: paranthropines, also known as robust australopiths.
5. Hominins that retained the gracile masticatory apparatus of their australopith ancestors and exhibited a trend for encephalization and increasingly complex culture: *Homo* species.

This page titled 4.2.2: What is a Hominim is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **1.5: What is a Hominim** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

CHAPTER OVERVIEW

4.2.3: Miocene Epoch

The Miocene epoch extends from about 23.03 to 5.333 million years ago (Ma). The Miocene was named by Charles Lyell and means "less recent" because it has 18% fewer modern sea invertebrates than the Pliocene. The Miocene follows the Oligocene and is followed by the Pliocene epoch.

4.2.3.1: Introduction to the Miocene Epoch

4.2.3.2: Sahelanthropus tchadensis

4.2.3.3: Orrorin tugenensis

4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba

4.2.3.5: Early Hominid Cranium Comparison Checklist


Thumbnail: Miocene fauna of North America. Mural made for the US government-owned Smithsonian Museum (Public Domain).

This page titled 4.2.3: Miocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

4.2.3.1: Introduction to the Miocene Epoch

THE MIOCENE EPOCH (~23–5.3 MYA)

Relative to the Oligocene Epoch, the Miocene was initially warmer and by the mid-Miocene, primates had once again ventured into the northern latitudes. This time they were apes, versus the prosimians of the Eocene. However, like those earlier primates, the northern apes would eventually go extinct due to global cooling that began ~14 mya.



And, on this side of "the pond": Miocene North American fauna. "Miocene" by Jay Matternes is in the public domain.

Whenever there is climatic change, floral and faunal changes follow. Much of the dense equatorial forests became more open, and grasslands and deserts expanded so that the African landscape became more of a **mosaic environment**, versus deep, dark jungle. Initially the apes were very successful at diversifying and expanding into new niches; as mentioned in Chapter 3, this was the time of the **hominoid adaptive radiation**. Over time, however, as the landscape continued to change and forests became increasingly fragmented, the majority of ape species went extinct, leaving only a handful of species in Africa.

In recent years, three new genera of hominoids have been discovered in Africa. While there is debate as to whether the oldest, i.e. *Sahelanthropus*, is a true hominin, *Orrorin* and *Ardipithecus* are generally accepted as true hominins.

WARNING

Caution must be exercised in the interpretation of the fossil record. It is tempting to formulate a linear history for our species, e.g. species A gave rise to Species B which gave rise to Species C, etc. The fossil record is extremely incomplete and we may get only a glimpse of the species that lived at a particular time and place. Paleoanthropologists are continually revising their theories as new discoveries are made and old material is reinterpreted in light of new discoveries, methodologies, and working hypotheses.


This page titled 4.2.3.1: Introduction to the Miocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **2.1: Introduction to the Miocene Epoch** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.2.3.2: Sahelanthropus tchadensis

Sahelanthropus tchadensis **(<7mya)**

("HUMAN FROM THE SAHEL" / CHAD)



Sahelanthropus tchadensis by Keenan Taylor.

SITES

Toros-Menalla, Chad

PEOPLE

Michel Brunet

INTRODUCTION

The *Sahelanthropus tchadensis* specimen (see Figure 6.2) was discovered in 2001 at the site of **Toros-Menalla**, in the Djurab Desert of northern Chad, by **Michel Brunet** and associates. Brunet's incredible years-long quest for hominins in that area is documented in the NOVA series, *Becoming Human*(www.pbs.org/wgbh/nova/evolution/becoming-human.html). The species name translates to "human from the **sahel** of Chad." The sahel is the region of dry grasslands south of the Sahara desert. The skull has been nicknamed "**Toumai**" in the Dazaga language, meaning "hope of life."



Cast of the Sahelanthropus tchadensis holotype cranium. "Sahelanthropus tchadensis – TM 266-01-060-1" by Didier Descouens is licensed under CC BY-SA 4.0.


The location of the fossil material came as a surprise in that only one species of hominin had ever been discovered west of the Great Rift Valley of East Africa, i.e. *Australopithecus bahrelghazali* (see Chapter 12). However in 1998, Noel Boaz speculated that, contrary to the Rift theory for the origin of the hominins, a portion of the ancestral stock that gave rise to the chimp and human lineages became isolated in a riparian (i.e. riverine or gallery) forest zone in Chad that was surrounded by arid, open land. At a later point in time, a forest corridor allowed their movement into East Africa. Part of the problem at that point in paleoanthropology was that no species of hominins, prior to the australopiths, had been discovered in East Africa. They seemingly appeared *de novo* in the fossil record, beginning about 3.5 mya, with no intervening stages or "missing links" in evidence. We now have much older hominin species from Kenya and Ethiopia, i.e. *Orrorin tugenensis* and Ardipiths, respectively.

PHYLOGENY

While the phylogeny of *S. tchadensis* is unknown, some researchers believe that it may represent a **stem** or **basal hominin**, i.e. one of the earliest members of our tribal tree. (**Note**: Once a genus is used the first time in a document, it can subsequently be abbreviated.) Just as we do not know the ancestry of the species, we do not have any species that are good contenders for its descendants.

DISCOVERY AND GEOGRAPHIC RANGE

As mentioned, the holotype (the fossil(s) from a particular individual that are assigned to **and** used to define the characteristics of a species) was discovered at the desert site of Toros Menalla (see Figure 6.3). Unless fossils are discovered elsewhere, it is impossible to speculate about the extent of the geographic range of the species.



Toros Menalla site, Chad. "Sahelanthropus tchadensis – TM 266 location" by 120 is in the public domain.

PHYSICAL CHARACTERISTICS

The skull of *S. tchadensis* is very robust, with a chimp-sized brain and pronounced ape-like muscle attachments. While only fragmentary postcranial material has been discovered, some researchers claim that the foramen magnum is anteriorly oriented, suggesting an upright and bipedal hominin. Pronounced brow ridges are also concordant with early hominin status. The facial profile is surprisingly orthognathic and the jaws lack the honing complex, leading some researchers to speculate that *S. tchadensis* may lie near the base of our family tree, versus other phylogenetic scenarios. However, the pronounced posterior neck muscle attachments have led others to suggest that *S. tchadensis* may have been quadrupedal.



Sculpture of S. tchadensis by Élisabeth Daynès.

ENVIRONMENT AND WAY OF LIFE

Based upon fossilized faunal remains at the site, such as freshwater fish, rodents, and monkeys, it is likely that *S. tchadensis* inhabited a forest environment in close proximity to an ancient lake (Wayman 2012). Their way of life was likely that of a forest-dwelling ape. Like ardipiths (see Chapter 8), their molar enamel was thinner than that of the later australopiths and they thus likely had a chimp-like diet consisting of fruit, young leaves, and tender shoots.

This page titled 4.2.3.2: Sahelanthropus tchadensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **2.2: Sahelanthropus tchadensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.2.3.3: Orrorin tugenensis

Orrorin tugenensis (~6 mya)

("ORIGINAL" / TUGEN HILLS)

SITES


Aragai and Kapsomin in the Tugen Hills of Kenya

PEOPLE

Brigitte Senut and Martin Pickford

INTRODUCTION

In 2000, the team of **Brigitte Senut** and **Martin Pickford** discovered fossil material (see Figure 7.1) from the Lukeino Formation in the **Tugen Hills of Kenya**. Nicknamed "**Millenium Man**" due to its timely discovery, the fossils were dated to ~6 mya and given the taxonomic classification, *Orrorin tugenensis* ("original man from the Tugen hills"). Initially many paleoanthropologists were skeptical, especially since the fossils were not made available to the scientific community. While there is still debate, *O. tugenensis* is increasingly presented in published texts as a hominin.




Drawing of Orrorin tugenensis fossil material. "Orrorin tugenensis" by Lucius is licensed under CC BY-SA 3.0.

PHYLOGENY

The ancestry of *O. tugenensis* is unknown. Senut and Pickford believe that *Orrorin* is ancestral to humans. They suggest that the hominin tribe split prior to 6 mya with *Orrorin* and some species of australopiths (specifically *Australopithecus anamensis* and *Australopithecus afarensis*, which they place in the genus *Preanthropus*) in the human lineage and ardipiths and robust australopiths, or paranthropines (including *Australopithecus africanus*), on another branch that died out.

DISCOVERY AND GEOGRAPHIC RANGE

There are only two known sites for the species, Aragai and Kapsomin, both of which are in the Tugen Hills of Kenya (see Figure 7.2).



Ororrin tugenensis sites, Kenya. "Ororrin localities" by Chartep is licensed under CC BY-SA 3.0.

PHYSICAL CHARACTERISTICS

Few body parts have been recovered. The fossils consist of a partial humerus, femur, and mandible; a distal thumb bone (phalanx); and some teeth. Primitive characteristics include remnants of a honing complex, with large canines and a "**semi-sectorial**" premolar. The molars were covered with **thick enamel** like those of later hominins, and while they were small like our own, they were not as laterally expanded. Senut and Pickford assert that the femur of *O. tugenensis* is very human-like, with its large head and hominin-like carrying/bicondylar angle, and thus is part of the human lineage.

ENVIRONMENT AND WAY OF LIFE


It is generally accepted that *O. tugenensis* was bipedal, and that they likely practiced a similar way of life as the ardipiths and australopiths. Thus, they were likely semi-terrestrial, foraging both in trees and on the ground and using trees for sleep and safety. However, their more human-like femur and thick molar enamel allies them more with the australopiths and they thus likely spent more time on the ground than the ardipiths and exploited a greater percentage of terrestrial resources that cause more tooth wear. In addition, like *Australopithecus afarensis*, *O. tugenensis* exhibits vestiges of the honing complex and thus may be ancestral to australopiths.

This page titled 4.2.3.3: Ororrin tugenensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **2.3: Ororrin tugenensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.


4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba

GENUS: *Ardipithecus* (“ground ape”)



Ardi by Keenan Taylor.

Deposits within the **Afar triangle/depression** of Ethiopia (see Figure 8.2) have yielded multiple hominin species within the genera *Ardipithecus* and *Australopithecus*. This hotbed of hominin fossils is the northern limit of the **East African Rift Zone**, where the Arabian and African plates converge. The first species of ardipith to be discovered in the area was *Ar. ramidus* (4.4 mya), and the second and even older species was *Ar. kadabba* (5.8 mya). When *Ar. kadabba* was discovered by **Yohannes Haile-Selassie**, he believed that it was similar enough to *Ar. ramidus* that he included it in the same genus and species, thus warranting subspecies classification. Since *Ar. ramidus* was already named, its classification became *Ar. ramidus ramidus*, in the same manner that we became *Homo sapiens sapiens* when it was decided that Neandertals should be included in our species and they became *Homo sapiens neanderthalensis*. The original name is always preserved for the original fossil(s). It becomes known as the “**type specimen**” or “**holotype**” and is used to describe the characteristics that define the species. The older subspecies then became *Ar. ramidus kadabba*. Since that time, they have been split into two species within the genus: *Ardipithecus*. Each will be discussed below.




The East African Rift Zone. The Afar triangle is featured in darker pink at the most northerly limit (see ‘Erta ‘Ale site). “EAfrica” by the USGS is in the public domain.

Some paleoanthropologists have suggested that *Ardipithecus* may be a better candidate for our ancestry than one or more of the australopiths. It has also been suggested that australopiths are descended from *Ardipithecus* or that the ardipiths are a separate but related branch.

Ardipithecus ramidus **(4.4 mya)**

("GROUND APE" / "AT THE ROOT" IN AFAR LANGUAGE)



Map showing the fossil sites of the earliest hominids. "Map of the fossil sites of the earliest hominids (35.8-3.3M BP)" by Kameraad Pjotr and Sting is licensed under CC BY-SA 3.0.

SITES

Aramis, Ethiopia

PEOPLE

Tim White and colleagues

PHYLOGENY

The phylogeny of the ardipiths is unknown. We do not know if they left any descendants, but it is thought that Ardipithecus ramidus is likely descended from Ardipithecus kadabba.


DISCOVERY AND GEOGRAPHIC RANGE

During the early 1990s, fossils were unearthed at the site of **Aramis** in the **Middle Awash region** of the Afar Triangle of Ethiopia (see Figure 8.3). Since that time, material from more than 50 individuals has been recovered, in particular the famous "**Ardi**" skeleton (see Figures 8.4 and 8.5) that is ~50% complete. Prior to the discovery, all or most early African hominin fossils were considered to be australopiths. Tim White and his colleagues determined that the material was distinctive enough to warrant new genus classification.



Digital reconstruction of Ardipithecus ramidus specimen. "Ardi" by T. Michael Keesey is licensed under CC BY 2.0.

Fifteen years after its discovery, **Ardi** was presented to the world in a frenzy of media coverage. She shook the world, not just within paleoanthropology, but for anyone interested in our past. She was an upright ape! Yes, we are all apes, but Ardi looked like what we think of as an ape. She had an ape face, small brain, long and strong arms and fingers, and ape-like feet. However, she stood upright with straight rather than flexed legs. The film *Discovering Ardi* (2009, Discovery Communications) showcases her discovery, fossil processing and analyses, the artist's skeletal and full-body reconstructions, biomechanical computer-generated graphics, etc.—everything for the modern tech-appreciative student!



Artist's representation of Ardipithecus ramidus. "Ardipithecus Gesamt1" by Ori~ is licensed under CC BY-SA 3.0.

PHYSICAL CHARACTERISTICS

Ardipiths were likely arboreal climbers, like the ancient "basal" proconsulids. However, when they were on the ground, they walked bipedally, albeit with rather clumsy-looking feet. Characteristics that were initially used to designate hominin status were an anteriorly placed foramen magnum, aspects of the innominate and ulna, and especially the morphology of the humerus that suggested that it was not weight-bearing and therefore not involved in terrestrial locomotion.

Primitive characteristics of ardipiths can be seen in many body regions. Their brains were small. Depending on the source, their cranial base (the inferior portion of the occipital bone) was either flat like chimps and gorillas, or angled and tucked under the upper part of the cranium (termed **flexed cranial base** or **basicranial flexion**). The enamel on their molars was thin like that of chimps and other extinct forest-dwelling apes. They retained several primitive characteristics that are related to arboreal climbing:

- Elevated shoulder joints for reaching up for branches, etc.
- Long arms and long, curved fingers.
- Mobile wrists.
- Poorly differentiated thumb, meaning that they had poor opposability.

Orangutans are the only great ape with a similar thumb morphology and hence have poor manual dexterity. Ardipiths' **divergent hallux** (i.e. the big toe diverges away from the lateral four digits, like our thumbs) would also have been adaptive for climbing.

Derived postcranial characteristics of ardipiths are all in the hip and lower limb. While Ardi's species had short legs and long arms (i.e. high intermembral index), the morphology of their hip and leg bones reflect bipedality, according to Tim White, Owen Lovejoy, and other members of their team. The innominate and foot morphology combine adaptations for climbing and bipedalism. They were capable of full extension at the knee, unlike the flexed knee seen in extant apes. Their feet were stable and supported their body weight and the divergent hallux facilitated grasping and climbing.

Body mass is estimated from Ardi's skeleton and thus no estimate for males of the species is possible. Ardi is estimated to have been 3′11" (120 cm) tall and weighed about 110 lb (50 kg) (Gibbons 2009). The absence of both a honing complex and pronounced prognathism suggests to some researchers, such as C. Owen Lovejoy, that males were not competing for females and may have formed pair-bonds with them.

Review of Primitive Characteristics


- Small brain and flat cranial base.
- Thin molar enamel.
- Climbing complex of upper limb:
  - Elevated shoulder joint.
  - Long arms.
  - Long and curved fingers.
- Low manual dexterity.
- Divergent hallux.

Review of Derived Characteristics

- Anteriorly oriented foramen magnum.
- Bipedal hip, leg, and foot characteristics.

ENVIRONMENT AND WAY OF LIFE


The sites where ardipith fossils have been recovered were a mosaic environment consisting of wood- and grasslands during the late Miocene and early Pliocene. Those ancient apes likely subsisted on a combination of arboreal and terrestrial forest resources. Their thin molar enamel suggests that, like chimps and orangutans, their diet consisted of relatively soft food items, such as fruit, young leaves, and shoots and possibly ants and termites. They could locomote within and between trees when foraging, or climb down and travel between trees or forested areas. Since all great apes build nests, ardipiths may have made a new arboreal nest every night.



Ardipith by Keenan Taylor.

Some scientists see the lack of canine sexual dimorphism and honing complex as evidence for **pair-bonding**. There is good evidence from the animal literature to relate sexual **monomorphism** (no difference in size between males and females) and pair-bonding. As mentioned in Chapter 4, while there are several groups of primates that exhibit that grouping and mating strategy, they are all arboreal. The lesser apes are arboreal, medium-sized (<6 kg) primates that form territorial pairs. Their mode of locomotion is brachiation, i.e. swinging under branches. While that is a more efficient means by which to avoid predation relative to ardipiths climbing around in the trees, the latter were larger and heavier and thus were likely fairly safe. When they descended to the ground, however, pairs would have been fair game for predators.

The idea that male provisioning, in combination with pair-bonding, was the evolutionary stimulus for bipedalism is also problematic. As I discussed in Chapter 5, I do not understand why females would need to be provisioned unless their babies could no longer hang on. I just do not see males risking life and limb to bring food back to females. I know of no examples in nature of a mammalian male provisioning his mate. It is more likely that resources became fragmented to the point that upright locomotion was the most efficient way to move between trees and forest patches. Bipedalism was an adaptation to enable continued reliance on forest resources while minimizing competition. Ardipiths could move to another forest patch when resources became scarce. I think of them as being semi-terrestrial apes like chimps and gorillas but, unlike those extant species, they had less of a stable resource base. They thus evolved a more efficient locomotor strategy for expanding their home range.




Ardipith Parenting by Keenan Taylor.

Ardipithecus kadabba (5.8 mya)

(“GROUND APE” / “OLDEST ANCESTOR” IN AFAR LANGUAGE)

Yohannes Haile-Selassie discovered the second ardipith species in the Middle Awash region of the Afar Depression (see Figure 8.3). The species is thought to be ancestral to *Ar. ramidus*. Dental characteristics are more ape-like than those of *Ar. ramidus*. While the fossils are very fragmentary, a toe bone suggests bipedal foot movement involving the “toe-off” motion, i.e. when we push off with the toes of one foot as we plant our other foot following its “swing” phase.



Ardipith by Keenan Taylor.

This page titled 4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **2.4: Ardipithecus ramidus and Ardipithecus kadabba** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.2.3.5: Early Hominid Cranium Comparison Checklist

Name: _____

Section: _____

Student ID#:_____

EARLY HOMINID CRANIUM COMPARISON CHECKLIST

Martin Nickels

© 1999 ENSI (Evolution & the Nature of Science Institutes) www.indiana.edu/~ensiweb

This material may be copied only for noncommercial classroom teaching purposes, and only if this source is clearly cited.

PLEASE READ THESE 5 STATEMENTS BEFORE BEGINNING THIS ACTIVITY:

1. Work in groups of 3-4 students so that everyone can be involved in the activity.

2. BE SURE (!) TO TAKE TURNS doing different measurements and observations.

3. When taking a measurement, use the SLIDING CALIPERS (except for #11 & #12 which may require the HINGE calipers) and remember to...

4. ALWAYS MEASURE IN MILLIMETERS [mm] and round off to whole numbers.

5. PLEASE DO NOT ADD ANY PENCIL OR PEN MARK "TATTOOS" TO THESE CRANIA, OR STICK YOUR FINGERS IN THEIR EYE ORBITS OR NOSES!

I. BRAINCASE: (7 items #1-7)

1. Does the FOREHEAD (frontal bone) look more vertical OR flatter when the skull is held in normal anatomical position [NAP] (i.e., with the eyes oriented forward)?

2. Is a SUPRAORBITAL BROWRIDGE present?

3. If present, is the BROWRIDGE DIVIDED in the middle, or CONTINUOUS?

4. What is the SHAPE OF THE BRAINCASE (front to back) when viewed from above?

5. Is a SAGITTAL CREST present?

6. In NAP, is the FORAMEN MAGNUM oriented more downward OR more to the rear?

7. Is the MASTOID process relatively flat OR does it noticeably protrude (project)?

II. FACE: (5 items: #8-12)

8. Are the NASAL BONES raised (arched) OR flat?

9. Measure the MAXIMUM BREADTH (width) of the NASAL OPENING [mm].

10. Measure the MAXIMUM HEIGHT of the NASAL OPENING [mm].

11. Measure the LENGTH of the MAXILLA (the upper jaw) [mm]. (Measure down the middle of the palate from the front edge of the foramen magnum to either between or just in front of the two central incisors to determine how much the face projects forward.)

12. Measure the BIZYGOMATIC BREADTH using the hinge caliper if necessary [mm]. (This is the width or breadth of the face from the widest part of one zygomatic arch to the widest part of the other zygomatic arch.)

III. DENTITION: (6 items #13-18)

13. SHAPE OF THE DENTAL ARCADE: Do the tooth rows diverge towards the back OR are they more straight-sided and parallel to one another?

14. When viewed from the side, are the INCISORS angled out OR are they vertical?

15. Measure the COMBINED WIDTH or BREADTH of the 4 INCISORS together.

16. Does the CANINE tooth project above the chewing surfaces of the other teeth?

17. Is a CANINE DIASTEMA present?

18. Measure the COMBINED LENGTH of the LEFT 2 PREMOLARS and 3 MOLARS together by measuring from the back of the last molar to the front of the first premolar to determine the length of the chewing surface of the "cheek teeth". [mm]. (NOTE: Measure the right side if the left side is missing any of these 5 teeth.)

Source:

www.indiana.edu/~ensiweb/lessons/hom.cran.html

	Skull #1	Skull #2	Skull #3	Skull#4
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				

19) What trends do you notice? What are the major differences between these skulls?

This page titled 4.2.3.5: Early Hominid Cranium Comparison Checklist is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Alex A. G. Taub via source content that was edited to the style and standards of the LibreTexts platform.

- **1.8: Early Hominid Cranium Comparison Checklist** by Alex A. G. Taub is licensed CC BY-NC-SA 3.0. Original source: https://socialsci.libretexts.org/api/deki/files/94257/biological-anthropology-laboratory-activities.pdf.

CHAPTER OVERVIEW

4.3: Pliocene Epoch

The Pliocene Epoch is extends from 5.333 million to 2.58 million years BP and follows the Miocene Epoch and is followed by the Pleistocene Epoch.

4.3.1: Introduction to the Pliocene Epoch
4.3.2: Australopithecus anamensis
4.3.3: Gracile Australopiths
4.3.4: Australopithecus afarensis
4.3.5: Australopithecus bahrelghazali
4.3.6: Kenyanthropus platyops
4.3.7: Australopithecus prometheus or africanus
4.3.8: Australopithecus africanus
4.3.9: Middle Hominid Cranium Comparison Checklist


Thumbnail: impression of a Pliocene landscape. (Public Domain; Ridpath, John Clark).

This page titled 4.3: Pliocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

4.3.1: Introduction to the Pliocene Epoch

THE PLIOCENE EPOCH (5.3–2.6 mya)

The Pliocene Epoch (~5.3–2.6 mya) was characterized by global cooling and weather disruptions due to the formation of the Panama land bridge and resultant changes in ocean currents. The polar ice caps were expanding and sea levels had already begun to drop, as the Pleistocene Epoch (~2.6 mya–11.7 kya) approached. The geologic record shows us that Africa was subject to cooling and drying trends, with seasons becoming more pronounced. Grasslands were expanding and thus forest cover was shrinking. Intermittent wet and dry periods changed the African landscape. Lakes formed and dried and filled once again.



Pliocene fauna of North America. "Pliocene" by Jay Matternes is in the public domain.

All of the ardipiths and most australopiths went extinct during this epoch and by ~2.0 mya genus *Homo* appeared in the fossil record. The most tantalizing question with regard to the earliest members of our genus is what drove the encephalization process. By the time of the more derived paranthropines [*Paranthropus boisei* (2.5–1.4 mya) and *robustus* (1.9–1.3 mya)] and genus: *Homo* (≥2 mya), cranial capacity had increased ~100 cc and 200 cc, respectively, from the earlier hominins. It has generally been hypothesized that competition for resources was the driving force, and while that may be so, recent research has illuminated just how radical climatic changes were in East Africa. The film *Becoming Human* (2009, Discovery Communications) provides a nice overview of how scientists have pieced together the Pliocene environment of that region of the world from the local geology.

PLIOCENE HOMININS

The Pliocene Epoch can be considered the time of the great adaptive radiation of the hominins, when more than a dozen species evolved in the hominin corridor from Ethiopia to South Africa. While the ardipiths died out, concurrent with the reduction in forest cover during the late Miocene and early Pliocene, many australopiths came and went over the course of almost 3 mya of the Pliocene. The australopith lineage may pre-date the Pliocene; there is possible australopith material (possibly *Australopithecus anamensis* or *afarensis*) from the late Miocene from several Kenyan sites (e.g. Lothagam and Tabarin) for which no taxonomic designation has been established. While there is debate as to how much time australopiths spent outside of forested environments, they certainly filled distinctive niches, as evidenced by changing craniofaciodental morphology as well as more versatile hands that afforded them greater manipulative abilities. As we discover more about australopith limb morphology, it is apparent that some were more adapted to an arboreal environment whereas others were less so. Of even greater interest is that the East African australopiths (i.e. *Australopithecus anamensis*, *afarensis*, and *garhi*) may represent a different clade than those from South Africa (i.e. *Australopithecus africanus* and *sediba*). The East African species may have been more adapted to a terrestrial environment, while the South African forms may have retained or reverted to more primitive climbing adaptations. The various species may have walked differently as well.



Foraging paranthropines in South African grasslands. Artwork by Walter Voigt.

The question remains: from which group are we descended? Some researchers have suggested that *Au. sediba* from South Africa may be part of our lineage; this confounds matters for some researchers as to how the older East African species were seemingly less primitive in some ways than the more recent South African forms. We need to realize that we are only seeing part of the picture. Just as the ape population levels exploded on three continents in the Miocene, the same was apparently true for African hominins in the Plio-Pleistocene. Discoveries of new species of African apes and hominins are accelerating. We need to break out of our linear mindset and realize that there were numerous species adapting to localized conditions, so we should not expect to see uniform changes across time and geographic space. In addition, the same may hold true for grouping and mating practices. We should not jump to conclusions or get too caught up in how the various species were related and who begat whom! While everyone would like to be the person who found one of our long lost relatives, the history of paleoanthropology is fraught with those claims and debunks.



Modern landscape of Olduvai Gorge. "Panoramic view of Olduvai Gorge" by רנדום is licensed under CC BY-SA 3.0.

Upper limb morphology suggests that australopiths retained adaptations for climbing trees that were likely used for food and safety. However, when on the ground, they were habitual bipeds, albeit with some differences from ourselves.

This page titled 4.3.1: Introduction to the Pliocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.1: Introduction to the Pliocene Epoch** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.2: Australopithecus anamensis

(4.2 mya)

("SOUTHERN APE" / FROM THE "LAKE" IN THE TURKANA LANGUAGE)



Distal humerus of Australopithecus anamensis. "Australopithecus anamensis bone (University of Zurich)" by Nicolas Guérin is licensed under CC BY-SA 3.0.

SITES

- Allia Bay and Kanapoi, Kenya

PEOPLE

- Meave Leakey and Alan Walker

Australopithecus anamensis is the earliest known australopith. We do not know nearly as much about the species as about other australopiths due to a paucity of fossil material.

PHYLOGENY

Au. anamensis may be descended from the ardipith lineage or a heretofore undiscovered group. Lucy's species, *Au. afarensis*, may be descended from *Au. anamensis*.

DISCOVERY AND GEOGRAPHIC RANGE

Multiple paleoanthropologists (most notably Meave Leakey and Alan Walker) are credited with the discovery of *Au. anamensis* material. The species name refers to the Lake Turkana area of Kenya where the fossil sites of Kanapoi and Allia Bay are located (see Figure 10.2). There are also newer fossils from the Middle Awash area of the Afar Depression of Ethiopia (see Figure 10.2). The Ethiopian material is close in time and geographic space to an *Ardipithecus ramidus* site, lending some support to the possibility of their phylogenetic relatedness. There is some controversy over the lumping together of material from different levels and locations in Kenya that could have confounded the description of the species' characteristics.




Australopithecus anamensis sites. "Anamensis localities" by Chartep is licensed under CC BY-SA 4.0.

PHYSICAL CHARACTERISTICS

The species is thought to have been highly sexually dimorphic in body and canine size.

Much of the morphology is ape-like, and hence primitive. The jaws and teeth are the most primitive of any australopith, which is not surprising since it is the oldest. Unlike the parabolic tooth arrangement in the jaws of later hominins, *Au. anamensis*had an apelike, U-shaped dental arcade wherein the cheek teeth are nearly parallel (see Figure 10.3). Their jaws were also prognathic and their canines were larger than descendent species. However, the molars were expanded with thick enamel and low cusps like later hominins. In addition, aspects of the elbow, knee, and tibia were more derived, indicating its bipedal mode of locomotion.



Jaw of extinct ape illustrating U-shaped dental arcade. "Rangwapithecus gordoni jaw" by Ghedoghedo is licensed under CC BY-SA 3.0.

Characteristics

Review of Primitive Characteristics

- High degree of sexual dimorphism.
- Prognathic jaws with ape-like parallel cheek teeth.
- Larger canines than subsequent species.

Review of Derived Characteristics

- Expanded molars, low cusp relief, and thick enamel.
- Bipedal adaptations of elbow, knee, and tibia.

ENVIRONMENT AND WAY OF LIFE


Fossils have been found in a variety of paleoenvironmental settings, such as lakeside, woodland, and more open areas. The species likely slept in trees and foraged both in trees and on the ground, as they moved bipedally around their home range in search of resources and mates. The high degree of sexual dimorphism and the presence of the honing complex suggest a polygynous or polygynandrous mating system. The former may have involved a one-male/multi-female social organization and the latter, a multi-male/female pattern. A one-male system may have looked more like that of gorillas, where both males and females tend to leave their natal group. However, if we use chimps and bonobos as a model of our ancestral social organization, it was more likely a multi-male/female social organization, with males staying in their natal group (i.e. male-philopatric) and females leaving at sexual maturity to join another.

While it is easy to use our more closely related relatives to reconstruct our past behavior, we must remember that social organization is a function of both phylogeny and ecology. Taking both into consideration, in combination with their anatomy, it is more likely that *Au. anamensis* were more chimp-like. Their teeth do not have the higher cusps of the more folivorous gorilla, so their diet was likely more chimp-like and hence a combination of fruit, tender greens, and opportunistic animal matter. As mentioned in Chapter 5, that type of diet is harder to come by and females may have needed males to defend a territory for their nutritional needs and those of their offspring. Interestingly, isotopic analyses of *Paranthropus robustus* (robust australopith from South Africa) fossil material show that while males were from the area where the fossils were found, females were not. Thus we now have a fourth line of evidence favoring male philopatry.

This page titled 4.3.2: Australopithecus anamensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.


- **3.2: Australopithecus anamensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.3: Gracile Australopiths



Australopith fossil sites. "Map of the fossil sites of the early hominids (4.4-1M BP)" by Kameraad Pjotr and Sting is licensed under CC BY-SA 3.0.

Genus *Australopithecus* ("southern ape") was first used in 1924 by Raymond Dart for the "Taung Child," a juvenile *Au. africanus* specimen from the quarry site of Taung, in South Africa. *Au. anamensis*, *afarensis*, *africanus*, and *sediba* (depending on the evolutionary schema of individual paleoanthropologists) are popularly known as gracile australopiths, due to their more gracile masticatory apparatus relative to the robust paranthropines.



An australopith outlook. By Keenan Taylor, CC-BY-NC-SA.


This page titled 4.3.3: Gracile Australopiths is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.3: Gracile Australopiths** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.4: Australopithecus afarensis

(4.2 mya)

("SOUTHERN APE" / AFAR REGION OF ETHIOPIA)



Forensic facial reconstruction of Australopithecus afarensis. "Australopithecus afarensis" by Cicero Moraes is licensed under CC BY-SA 3.0.

SITES

- Ethiopia: Afar Depression (e.g., Hadar and Dikika)
- Tanzania: Laetoli
- Chad: Bahr el Ghazal

PEOPLE

- Donald Johanson, Mary Leakey, Zeresenay Alemseged

Australopithecus afarensis, or the "southern ape from Afar," is a well-known species due to the famous "Lucy" specimen. It has been extensively studied by numerous famous paleoanthropologists. As mentioned, it is categorized as a gracile form of australopith. The species survived for over a million years in the changing East African landscape, covering a broad geographic range. The famous Laetoli footprints are attributed to *Au. afarensis* (see Figures 11.5 and 11.6). They provided support for the then controversial idea of habitual bipedalism, as well as the species' presence in a more open environment.

PHYLOGENY


The most logical ancestor for *Au. afarensis* is *Au. anamensis*. The two species overlapped in time and geographic space. Some paleoanthropologists have always believed that genus: *Homo* is descended from *Au. afarensis*. Over time, others have changed their taxonomic scenarios from *Au. africanus* to *Au. afarensis* (which would formerly have been a sister lineage to *Au. africanus*) as our ancestor, and made *Au. africanus* a side branch of the robust forms. Part of the argument for classifying *Au. afarensis* outside of our lineage had to do with aspects of their anatomy being more derived than our own, e.g. the lateral flare of their ilia (the plural of ilium). Since the discovery of *Au. sediba* (Chapter 21), some scholars are back to favoring *Au. africanus*in our ancestry.



"Laetoli recreation." "Laetoli recreated" by Wapondaponda is licensed under CC BY-SA 3.0.

DISCOVERY AND GEOGRAPHIC RANGE


The geographic range of *Au. afarensis* extends over 1,600 km from the site of Hadar in the Afar Depression of Ethiopia to the Laetoli site in Tanzania (see Figure 11.3). The holotype comes from Laetoli. There is conjecture as to whether the Ethiopian and Tanzanian material should be attributed to the same species, since the sites are distant from one another and separated in time by 800 kya. In addition, if *Au. bahrelghazali* is included as a geographic variant of the species, their range expands 2,500 km westward into Chad (McHenry 2015). Thus this species was very successful at exploiting a variety of environments.



Map showing the major fossil sites where specimens of Australopithecus and Paranthropus have been found. From Clement and Hillson (2013), licensed under CC BY 3.0.

With the discovery of "Lucy" (3.2 mya) (see Figure 11.7) in 1974 by Donald Johanson's crew at the site of Hadar in the Afar Depression of Ethiopia, paleoanthropology gained momentum and the rush was on in East Africa to find more evidence of human origins. Certainly Louis and Mary Leakey recognized the importance of the Great Rift Valley, but Johanson "upped the ante" with his 3.2 mya find. In addition, since Lucy's skeleton was almost 40% complete (making it one of the six most complete fossilized hominin skeletons older than 100 kya), much could be said about her anatomy and locomotor capabilities.

Site AL 333 at Hadar yielded remains of 13 individuals, referred to as the "First Family." Some researchers speculated that they may have died together and thus possibly represent a social group. However, recent examination of the deposition pattern at the site suggests otherwise (see Behrensmeyer 2008).



Dikika Baby. "SelamAustralopithecus" by Jlorenz1 is licensed under CC BY-SA 3.0.

The more recently discovered "Dikika Baby" (3.3 mya) (see Figure 11.4), also known as "Selam" (meaning "peace" in the Afar language) has contributed greatly to our knowledge of development in early hominins. Dikika, meaning "nipple" in the Afar language, is the name of the nipple-shaped hill at the site of her discovery. Discovered by Zeresenay Alemseged in 2000, the three-year-old female has also been dubbed "Lucy's Baby" due to its proximity to Hadar where Lucy was discovered. Selam is now the oldest, most complete fossil hominin. It took five years to extract the fossils from the surrounding sandstone matrix in which they were embedded. Thus we can see that not only is there difficulty in locating fossils, along with their living conditions in the desert environments of East Africa, the fossils may take years to process before all of their secrets can be revealed.



Laetoli footprint cast. "Australopithecus afarensis footprint" by Tim Evanson is licensed under CC BY-SA 2.0.

Even more recent material from the Woranso Mille site in the Afar region has some scientists questioning whether the the *Au. afarensis* hypodygm (all fossil material attributed to a particular species) might represent at least two species. *Australopithecus deyiremeda* has been suggested for the newer material.

Mary Leakey discovered the first and oldest (4.2 mya) *Au. afarensis* material at Laetoli, Tanzania, and the holotype (type specimen) comes from that site. Her team recovered fossil material from 23 individuals, as well as the famous Laetoli footprints. The trail of footprints extends for almost 25 m. They were made by two individuals walking side by side, with a possible third, smaller individual hopping within tracks already made by one of the adults. The prints were formed when the hominins walked through wet ash that had erupted from a nearby volcano.

PHYSICAL CHARACTERISTICS

Note: Distinguishing primitive versus derived characteristics is difficult because we do not have all body parts to compare from one species to the next. In order to determine what changed, I am considering those aspects that are more *Homo*-like as derived characteristics, regardless of whether predecessors possessed them. In other words, it is not perfect science!

All body parts are represented in the hypodigm, i.e. all fossils assigned to a particular species. While some debate surrounds the gait and locomotor efficiency of the species, it is fairly well accepted that they were habitual bipeds that retained some arboreal characteristics in the form of upward-oriented shoulder joints, an ape-like scapula, a high intermembral index, and curved finger


bones. Their innominates and lower limbs were unquestionably those of a biped, and the big toe, while slightly divergent from the other four digits, was not nearly the grasping digit seen in apes. The buttressing of their ilium, in the form of the iliac pillar, shows that weight was being transferred through the bone in the same manner as our own. While there is evidence of the bicondylar or carrying angle of the femur, the femoral head was small and the neck (narrowing below the head) was longer in australopiths and paranthropines (i.e. robust australopiths—see Chapters 16–19), relative to *Homo*species. Lovejoy believes that the degree of lateral iliac flare and long femoral neck in australopiths were associated with increased leverage of the deep gluteal muscles so that they were more biomechanically efficient than modern humans (Lovejoy 1988). As we gave birth to larger-brained infants, our pelvic aperture had to expand laterally so that the femoral neck became shorter and the deep gluteal muscles became less biomechanically stable relative to australopiths. The bony elements of the *Homo*pelvis and hip had to become more robust to handle the increased force that the gluteal muscles generated on the bones (discussed in Lovejoy 1988 and Cartmill and Smith 2009). Based upon the Laetoli footprints, it appears that the feet of *Au. afarensis* were slightly inverted, which would have helped with climbing.



Replica of Laetoli footprints. "Laetoli footprints replica" by Momotarou2012 is licensed under CC BY-SA 3.0.

Au. afarensis had a prognathic, ape-like face (see Figure 11.8), a primitive skull morphology, and a small brain averaging 420 cc. They exhibited a slight sagittal crest for attachment of the temporalis muscle and a more pronounced nuchal crest, where their nuchal (posterior neck) muscles inserted on the posterior skull. The two crests were compound—a compound sagittal-nuchal crest—meaning that the sagittal crest converged at the center of the nuchal crest. Their teeth were large and their dental arcade was U-shaped, and thus more ape-like. The lower first premolar suggests a transitional phase, termed semisectorial, between the honing, sectorial (single-cusped) premolar of the apes and our more bicuspid premolars. The canines were monomorphic. Like *Au. anamensis*, their molars were expanded.

While their hands were capable of a precision grip, they did not have the same degree of mobility in their thumbs as later species of australopiths and paranthropines. The conical thorax is linked to climbing and a large gut, and possibly the degree of lateral flare of the iliac blades.



"Lucy." "Lucy Mexico" by danrha is in the public domain.

The Dikika Baby (see Figure 11.4) revealed inner ear adaptations that allowed them to distinguish their head from their torso; this is important for running. Her brain was of similar size to that of a same-aged chimp. This indicates that they had a more prolonged developmental period, since the adult brain was, for the most part, larger than those of chimps. While chimps' brains are ~380 cc, *Au. afarensis*' were on average 434 cc, and ranged from 342 to 540 cc. She had both deciduous and developing permanent teeth in her jaws. Finally, her ribs were in anatomical position, which confirmed the conical thorax.



Australopithecus afarensis reconstruction. "Australopithecusafarensis reconstruction" by Durova is licensed under CC BY-SA 4.0

Au. afarensis exhibited pronounced sexual dimorphism, with males and females averaging 4´11˝ and 3´5˝ tall, respectively. Weights ranged from 64 to 99 lb. While the differences in size and morphology between the sexes might suggest that they were promiscuous and that males competed for females, their canines were monomorphic, suggesting pair-bonding. This is also supported by the change in the first mandibular premolar to being semi-sectorial.

Characteristics

Review of Primitive Characteristics

- High degree of sexual dimorphism.
- Low cranial capacity.
- Compound sagittal-nuchal crests:
  - Slight sagittal crest.
  - Pronounced nuchal crest.
- Flat cranial base.
- Prognathic jaws with U-shaped dental arcade and large ape-like incisors.
- Arboreal characteristics:
  - Upward-oriented shoulder joints.

  - Ape-like scapula.
  - Long arms relative to legs.
  - Curved finger bones.
- Conical thorax.
- Some divergence of hallux.

Review of Derived Characteristics

- Inner ear adaptations allowed for more efficient running.
- Low cusp relief and thick enamel on molars.
- Loss of honing complex:
  - Semi-sectorial premolar.
  - Monomorphic canines.
- Precision grip adaptation in hands.
- Bipedal hips and lower limbs.
- Prolonged juvenile dependency and brain growth.

ENVIRONMENT AND WAY OF LIFE

This species inhabited a mixed woodland environment that is thought to have been more open than previous hominin habitats. They could thus have exploited arboreal resources and moved between trees and forested areas in a fairly efficient manner. They are considered to have been scavenger-foragers, collecting wild plant foods, opportunistically hunting animals, and scavenging large game from carnivore kills. There is evidence of stone tool use at the Dikika, Ethiopia, site. Since *Au. afarensis* are the only known hominin from that time and location, the tool use has been attributed to them. Researchers found cut marks on bones of two large animals that were dated to 3.4 mya. Even more exciting is the recent discovery of 3.3 mya tools in association with hominin fossil material at the West Lake Turkana, Kenya, site of Lomekwi 3. While it was commonly accepted that australopiths used tools, this is the first evidence that they made them. The tools have been designated as the Lomekwian industry and have displaced the Oldowan as the earliest tool industry, preceding it by 700,000 years (Harmund et al. 2015). The tools consist of anvils, cores (stones from which flakes for cutting are removed), and flakes (see *Homo habilis*: “Environment and Way of Life” for more information on stone tools and their production). Like extant great apes, they also would almost certainly have used biodegradable materials for tools, such as wooden, ivory, or antler digging sticks.

Au. afarensis exhibited premolar molarization and thick molar enamel for masticating a tougher, more dry-adapted diet, such as tubers (large edible roots, e.g. yams). However, they were not yet able to grind their food as well as later hominins whose jaws could move laterally due to the reduction in canine size.



Selam reconstruction at the National Museum of Addis Ababa, Ethiopia. “Selam” by Highrey is licensed under CC BY-SA 3.0.

The brain of Selam shows that the juvenile dependency period was prolonged relative to chimps and hence the chimp/hominin ancestor. In addition, once infants could not hang on with their feet, mothers would have had to put their babies down periodically. Dean Falk has suggested that this pattern of mother-infant care may have led to language, in the form of what she refers to as “motherese” (Falk 2009).

woman the gatherer

It is interesting that female chimps use tools more often than do males. In addition, “woman the gatherer” should share the limelight with “man the hunter,” as women in most traditional societies collected a larger share of their family’s food. Is it possible that women invented tools? How about language? For how long have we heard about the male provisioning model for the evolution of bipedalism, “man the toolmaker,” “man the hunter,” men romancing women with the first language? Let’s stir up that cooking pot!!!

As mentioned, we are unsure of their mating and thus grouping pattern. Regardless of whether the First Family died together and represented a social group, *Au. afarensis*likely lived in groups for protection and possibly cooperation. Males were much larger than females but had lost the large canines and honing complex of *Au. anamensis*. Thus while the degree of sexual dimorphism was much greater than in our own species, their monomorphic teeth suggest that they were transitioning toward pair-bonding while retaining polygynous tendencies. While females may have mated polyandrously, like a fair proportion of females in our own species, it may have been in their best interest to stick with their mate for help in raising their offspring, and not jeopardizing their safety with extra-pair copulations.



Lucy by Keenan Taylor.

This page titled 4.3.4: Australopithecus afarensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.4: Australopithecus afarensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.5: Australopithecus bahrelghazali

(<3.4 mya)

(“SOUTHERN APE” / BAHR EL GHAZAL [ARABIC: RIVER OF GAZELLES], CHAD)

Note: *Au. bahrelghazali* is out of order here, in terms of date, because of its affinity to *Au. afarensis*.



Bahr el Ghazal site, Chad. “Bahr el Ghazal, Chad; Australopithecus bahrelghazali 1995 discovery map” by En rouge is licensed under CC BY-SA 3.0.

SITES

- Bahr el Ghazal, Chad

PEOPLE

- Michel Brunet

While it is possible that *Au. bahrelghazali* was a distinct species, the majority view is that it was a geographic variant of *Au. afarensis*. If that is the case, one can see how far the species ranged, from the horn of Africa, south to Tanzania, and west to Chad (see Figure 12.1). The Chad population was close in time to the material from the Afar Depression in Ethiopia, e.g. Lucy at 3.2 mya and Selam at 3.3 mya, versus the older material from Laetoli, Tanzania, that is dated to as old as 4.2 mya. Remember also that some researchers believe that the Ethiopian and Tanzanian populations were too far removed in geographic space and time to be considered the same species and/or to be using the holotype from Laetoli for the other populations. The partial mandible that was discovered by Michel Brunet at Bahr el Ghazal differed from *Au. afarensis* only in having been less prognathic. A premolar was also found at the site. The ancient environment has been compared with that of Hadar (Lucy’s site), i.e. a mosaic environment consisting of wood- and grasslands.

This page titled 4.3.5: Australopithecus bahrelghazali is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.5: Australopithecus bahrelghazali** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.6: Kenyanthropus platyops

thropus platyops (3.5 mya)

("KENYAN HUMAN" / "FLAT-FACED")



Figure **4.3.6.1**: *Kenyanthropus platyops reconstruction. "Kenyanthropus platyops IMG 2946" by Rama is in the public domain.*

SITES

- Lake Turkana, Kenya

PEOPLE

- Meave Leakey

A surprisingly "flat-faced" hominin came to light with **Meave Leakey**'s discovery and naming of *Kenyanthropus platyops* ("flat-faced human from Kenya") in 1999. The degree of orthognathism was surprising for such an early hominin.

PHYLOGENY

Since the skull was crushed and reconstructed (Figure **4.3.6.1**), some paleoanthropologists discount the degree of orthognathism and would like to see the specimen assigned to genus: *Australopithecus*, and possibly to species: *afarensis*. Others feel that it is the base of our lineage, believing that a cladistic event occurred at *Au. afarensis*, leading on the one hand to the robust forms (including *Au. africanus*) and on the other hand toward humans. In the latter case, some would like to see *K. platyops* assigned to genus: *Homo*, believing it to be the ancestor of *Homo rudolfensis*, with which it shares a similar orthognathic profile. Of interest is that some have speculated that the species may have made the Laetoli footprints. However, if the *platyops* fossils really belong in *Au. afarensis*, it would not be news!



Figure **4.3.6.2**: *Lake Turkana, Kenya. "Kenya relief location map" by Uwe Dedering is licensed under CC BY-SA 3.0.*

DISCOVERY AND GEOGRAPHIC RANGE

K. platyops is known only from a site in the West Lake Turkana region of Kenya (see Figures 13.2 and 13.3), where Meave Leakey's team found the crushed skull in 1999. The **hypodigm** consists of a reconstructed skull, plus two partial maxillae and a temporal bone from other individuals.



Figure **4.3.6.3**: *Lake Turkana, Kenya. "LakeTurkanaSouthIsland" by Doron is licensed under CC BY-SA 3.0.*

PHYSICAL CHARACTERISTICS

While possessing primitive ape-like molars (elongated mesiodistally, i.e. front to back) and sharing similarities with *Au. anamensis* and *afarensis*, the lower face of *K. platyops*is surprisingly (and possibly mistakenly) orthognathic for its early date. The cranial capacity of *K. platyops* is also suspect due to the reconstruction but if accurate, it was fairly high relative to other species of the time, at 400–500 cc.

Characteristics

Review of Primitive Characteristics

- Elongated ape-like molars.

Review of Derived Characteristics

- Possibly more orthognathic.
- Possibly encephalized relative to previous and contemporary species.


This page titled 4.3.6: Kenyanthropus platyops is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.6: Kenyanthropus platyops** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.7: Australopithecus prometheus or africanus

"LITTLE FOOT" (~3.6–3.2 MYA)

("SOUTHERN APE" / PROMETHEUS / AFRICA)



Tarsals and metatarsal portion of "Little Foot." "Little Foot 01" by Tobias Fluegel is licensed under CC BY-SA 3.0.

SITES

- Sterkfontein and possibly Makapansgat, South Africa

PEOPLE

- Ronald Clarke

The controversial material that has come to be known as "Little Foot" is an almost complete skeleton from the site of Sterkfontein (see *Au. africanus* for more information on Sterkfontein). The story is remarkable in that the skeletal components were discovered at two different times. The earlier material was cataloged and stored as "cercopithecoid" (Old World monkey) remains. Fifteen years later, the rest of the skeleton was found at the same location at Sterkfontein (Silberburg Grotto) and matched to the previous material. The only other case that I know of where something like that happened was at the Swanscombe, England, site where three skull bones from a *Homo heidelbergensis* individual were found in three different years.

PHYLOGENY

Ronald Clarke believes that there were two species of australopith at Sterkfontein. He has assigned the species name *Australopithecus prometheus* to the Little Foot material, as well as two individuals that others assign to *Au. africanus*, one from Sterkfontein and the other from Makapansgat. While the phylogeny is unknown, Little Foot precedes and may be the ancestor of *Au. africanus*.

DISCOVERY AND GEOGRAPHIC RANGE

The "cercopithecoid" material was discovered by Ronald Clarke in a storeroom at the University of the Witwatersrand. Once he realized that the bones were hominin and had thus been miscatalogued, he sent his two assistants, Stephen Motsumi and Nkwane Molefe, out to the original site to see if they could find more … and they did! The species' geographic range is presently limited to the Sterkfontein area of South Africa (Duke 1998).

PHYSICAL CHARACTERISTICS

Little Foot is characterized as a climbing biped, as it had both arboreal and bipedal characteristics. Arboreal characteristics consisted of upward-oriented shoulder joints; ape-like arms that were, however, shorter than those of other southern australopiths; and curved hand and foot bones. Surprisingly, the few foot bones that were recovered are more complete than in any previous australopith specimen (see Figure 14.1). The morphology is transitional in that they retained some ape-like morphology. The hallux displayed the same degree of divergence as other australopiths. The upper limb combined both ape- and human-like characteristics. While the upper limb was shorter relative to australopiths, the hand bones remained curved. While the Little Foot specimen was ~4′ tall, males and females are estimated to have been 3′6″ and 4′6″ tall and 60–120 lb in weight, respectively.

ENVIRONMENT AND WAY OF LIFE

The environment of Sterkfontein is discussed in the section on *Au. africanus*.

Characteristics

Review of Primitive Characteristics

- Climbing adaptations:
  - Upward-oriented shoulder girdle.
  - Ape-like arms.
  - Curved hand and foot bones.

Review of Derived Characteristics

- Shorter arms relative to australopiths.

This page titled 4.3.7: Australopithecus prometheus or africanus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.7: Australopithecus prometheus or africanus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.8: Australopithecus africanus

(3–2 mya)

("SOUTHERN APE" / AFRICA)



Figure **4.3.8.1**: Cast of Taung Child fossils. "Australopithecus africanus – Cast of taung child" by Didier Descouens is licensed under CC BY-SA 4.0.

SITES

Taung, Makapansgat, Sterkfontein, Drimolen, and Gladysvale, South Africa

PEOPLE

Raymond Dart, Robert Broom, and C. K. Brain

Australopithecus africanus was the first fossil hominin discovered in Africa. In 1924, Raymond Dart (see his biographical sketch this chapter) identified the face, mandible, and endocast as being that of a juvenile bipedal ape (see Figure 15.1). Eugène Dubois's discovery of the Javanese *Homo erectus* fossils in 1891 refuted the reigning belief that "we got smart before we stood up." Once Dart's claims were accepted, the world realized the extent to which that idea was false. The small-brained *Au. africanus* showed that early hominins were bipedal apes as opposed to quadrupedal humans.

Unlike East African material that can be dated using a variety of techniques, primarily due to past volcanic activity, South African fossils and sites are much more difficult to date. Most dates can only be represented as ranges; hence the date for the earliest *Au. africanus* specimens falls between 3 and 2 mya.

PHYLOGENY

Au. africanus is considered to be a gracile australopith by some and a robust australopith by others. Traditionally, the species was favored as the immediate ancestor of the *Homo* lineage, specifically of *Homo habilis*. However, some researchers have always believed that *Au. afarensis* was the common ancestor of both *Au. africanus* and the *Homo* lineage, suggesting a cladistic event had occurred at *Au. afarensis*. This schema has gained popularity in recent years. However, with the new evidence being put forth for *Au. sediba*, it seems that the *Au. africanus* → *Homo*scenario was closer to the truth in that *Au. africanus* and *Au. sediba* are undoubtedly related and *Au. sediba* shares many characteristics with genus *Homo*. While there is some support for *Au. africanus* as ancestral to the more derived robust forms, that still leaves unresolved those characteristics shared between *Au. aethiopicus* and both *P. boisei* and *Au. afarensis* (see *Au. aethiopicus*, Chapter 17).

DISCOVERY AND GEOGRAPHIC RANGE

Au. africanus is known only from sites in South Africa (see map showing the major fossil sites in Chapter 11, Figure 11.3). Material from more than 200 individuals has been collected over more than 80 years. Most of the fossils came from caves, some of which were discovered during mining and blasting activities. Those caves formed via underground water activity. Fossilization was facilitated by water dripping on bones and calcifying, just as stalagmites form. Over millennia, many of the caves filled in with mineralized deposits and as the ground surfaces eroded, the underlying deposits were exposed and mined and then later excavated for fossils.

BIOGRAPHICAL SKETCH: RAYMOND DART (1893–1988) AND THE "TAUNG CHILD"

Raymond Dart was the first to recognize a fossil hominin in Africa. While his claim to have discovered a human ancestor was not initially accepted by the scientific community, he was vindicated when Robert Broom began finding similar material at other South African sites. The work of Dart and Broom played a pivotal role in directing attention to Africa as the birthplace of humanity, as Charles Darwin had earlier predicted.

Raymond Dart was a renowned Australian anatomist who was teaching at the University of Witwatersrand in South Africa at the time of the discovery. I have read several accounts over the years of the fateful day when the fossil material landed in Dart's hands. The Taung material was brought to him by either quarry workers or a colleague while he was dressing for the wedding of his daughter or friend, depending on the source you read. He is said to have attempted to use his wife's knitting needles to extract the fossils from the surrounding matrix. I am not sure why we need to know what he was doing when he got his hands on the goods. We certainly do not know what most paleoanthropologists were doing when they were presented with fossil finds. However, it conjures up amusing possibilities in my mind!

The fossilized remains came from the Taung Quarry in the process of blasting for lime. When presented with the material, Dart established that it was the face, mandible, and endocast (fossilized interior of the cranial vault) of a juvenile hominin. He based this on the anterior position of the foramen magnum and aspects of brain morphology reflected on the interior of the skull vault. Dart named his find *Australopithecus africanus*, meaning "southern ape of Africa," and the specimen became known as the "Taung Child."



Figure **4.3.8.2**: Raymond Dart. "Smithsonian Institution Archives – SIA-SIA2008-0845" from the Smithsonian Institution Archives is in the public domain.

The Taung child was two to three years old when it died. We now know that the caves of South Africa were formed by underground water activity and hominins that dropped or were dragged into those subterranean caves became fossilized during the process of speleothem formation. Over millennia, the caves filled with mineral deposits, and via erosion, the mineralized contents and underground cavities surfaced. Many fossils were likely destroyed during quarrying activities, but we are lucky that many were preserved.

Raymond Dart is also renowned for his "Killer Ape" theory and the osteodontokeratic (bone-tooth-antler) culture. Dart believed that animal long bones and carnivore mandibles that were found with australopith remains had been used as weapons to fight and kill one another. We now know that was not the case. They were most likely opportunistically hunting small prey and scavenging larger kills, and they were prey for larger animals.



Figure **4.3.8.3**: Sterkfontein Cave. "Sterkfontein Caves 16," photograph by Mike Peel (www.mikepeel.net), is licensed under CC BY-SA 4.0.

Raymond Dart is credited with the 1924 discovery and naming of *Au. africanus*. His now famous "Taung Child" came from the Taung quarry site. The two- to three-year-old juvenile is represented by its face, skull fragments, and mandible, and an endocast of its brain. Dart also worked at the site of Makapansgat. His contemporary, Robert Broom (see biographical sketch below), worked at the caves at Sterkfontein (Figure **4.3.8.3**), where he discovered a complete, female cranium known as "Mrs. Ples", along with other *Au. africanus* material. Broom also worked at the sites of Kromdraai and Swartkrans; the latter is where he discovered the first paranthropine, *Paranthropus robustus*. C. K. Brain, a famous taphonomist, also worked at Sterkfontein. He discounted Dart's views of the australopiths as "killer apes". He believed that bones either dropped into caves as part of large cats' prey or were dragged in by rodents for gnawing. Some individuals are thought to have accidentally been trapped in underground caverns. Some speculate that those individuals that show no evidence of having been preyed upon, due to their degree of completeness (e.g. Sts 573 from Sterkfontein and the *Au. sediba* party from the Malapa site), surely became trapped. The six *Au. sediba* individuals are thought to have possibly been attracted to the cave by water. Drimolen, a more recently discovered site, has yielded an almost complete cranium as well as material from approximately 80 individuals. A fifth site associated with the species is Gladysvale.

BIOGRAPHICAL SKETCH: ROBERT BROOM (1866–1951)

Robert Broom was a Scottish medical doctor and paleontologist who subsequently made a name for himself as a paleoanthropologist, even though the term did not exist at the time. He taught geology and zoology at a South African college until he was let go for his controversial beliefs in evolution that were contrary to the religious teachings at the college. Subsequent to his termination, his finances went downhill until Raymond Dart's influence secured him a position at the Transvaal Museum. While his specialty was mammalian-like reptiles, he became increasingly involved with fossil hominins. Together, Dart and Broom made paleoanthropological history, discovering the first and second species of African hominins, respectively. Broom worked at the *Au. africanus* sites of Sterkfontein and the *P. robustus* sites of Swartkrans and Kromdraai. All of those sites are now contained within the Cradle of Humankind World Heritage Site. He was the first to discover and name *Paranthropus robustus*, and his work with *Au. africanus* helped to support Dart's claims to have discovered a bipedal ape and human ancestor. Broom's most famous *Au. africanus* find was "Mrs. Ples" (possibly a male), which he originally named *Plesianthropus transvaalensis* or "primitive human" of the Transvaal.



Figure 4.3.8.4: Robert Broom. "Robert Broom00" from the American Museum of Natural History is in the public domain.

If accounts of Robert Broom are to be believed, he was a colorful character. He is said to have worn semi-formal attire while excavating and when media were present, he conveniently happened upon important discoveries. Supposedly, one of his team buried an artifact that was already labeled with a catalog number. Oh, to have been on the scene and witnessed what must have been an embarrassing situation! Not to mention that, according to numerous online sources, he thought nothing of stripping naked when it became "Africa Hot"! (I picked up that term from a couple of comedy movies but it is now in the Urban Dictionary.)

It is interesting that Broom did not believe in Darwinian evolution but rather, what we would now call Intelligent Design (Wikipedia contributors 2015h).

PHYSICAL CHARACTERISTICS

Au. africanus was more derived than *Au. afarensis*. This is not surprising considering that they lived at least one million years later, as well as the trend within the hominin lineage to become more encephalized and manually dexterous over time. *Au. africanus* were habitual bipeds with all of the corresponding lower limb adaptations. They also retained climbing characteristics, such as upward-oriented shoulder joints; long arms relative to legs; and long, curved hand and finger bones. However, in general their hands were more human-like than those of *Au. afarensis*.




Figure 4.3.8.5: Lobes of the brain. "BrainLobesLabelled" by Camazine is licensed under CC BY 3.0.

While the brain was small relative to later species, *Au. africanus* was not only more encephalized than past species, with a cranial capacity of 450 cc (range = 424–508 cc), but also possessed an enlarged cerebral cortex in the frontal and parietal regions (see Figure 15.5). Their encephalization quotient (EQ) was 2.7. The quotient is a method for comparing brain size among species. Anything greater than 1.0 means that the brain is larger than would be expected based on body size (FYI: our EQ is ~7.6). Broca's area is an area of the left lateral frontal cortex that is involved with the production of language (see Figure15.6). It is present in all Old World monkeys and apes, but it is enlarged in *Au. africanus* relative to previous species. These are all important developments, in that they herald the appearance of more complex thought processing and likely communication skills. There is debate over whether the lunate sulcus (see Figure 15.7), a fissure on both sides of the occipital lobe that is involved with vision, was more ape- or human-like. The sulcus is smaller in humans than in monkeys and apes.




Figure **4.3.8.6**: Broca's and Wernicke's areas (see areas surrounded by dotted lines). "Blausen 0102 Brain Motor&Sensory" by Blausen Medical Communications, Inc. is licensed under CC BY 3.0.Blausen.com staff. "Blausen gallery 2014." Wikiversity Journal of Medicine.
doi:10.15347/wjm/2014.010. ISSN 20018762.




Figure **4.3.8.7**: Lunate sulcus, also known as the simian or lateral occipital sulcus, outlined in red. "Gray726" vectorized by was_a_bee based on drawing by Henry Vandyke Carter is in the public domain.

The external skull reflected the cerebral expansion by becoming more rounded and exhibiting more of a forehead. In addition, the sagittal and pronounced nuchal muscle crests seen in *Au. afarensis* were not present. However, some males had convergent temporal lines that suggest that they may have had a slight sagittal crest. There is debate over whether the cranial base was flexed (see Figure 15.8), or whether that was a development seen only in more derived forms of *Homo*.




Figure **4.3.8.8**: Basicranial flexion in four hominin species. Illustration by Keenan Taylor.

The species' face was prognathic with a distinctive concavity in the midfacial region (see Figure 15.9). The dental arcade was more parabolic, and the teeth were smaller than those of *Au. afarensis*. The first premolar is considered to have been bicuspid, versus

semi-sectorial in *Au. afarensis*. However, their faces were more heavily buttressed for chewing a tougher diet, and some researchers thus consider them to be “robust” australopiths.



Figure **4.3.8.9**: Concave facial profile of Mrs. Ples (Au. africanus). “Australopithecus_africanus_STS_5-side_3204” by NCSSM is licensed CC-BY-NC-SA

They retained the primitive condition of long arms and their finger bones were somewhat curved. However, their hands were more human-like and they possessed our “power” thumb, giving them increased pinching and gripping capabilities. This was accomplished via better developed intrinsic thumb muscles, i.e. within the hand versus coming from the forearm to act on the thumb, and a specialized first metacarpal.

While they were considered to be habitual and very likely obligate bipeds, they still retained a divergent hallux. According to McHenry and Berger (1998) and Green et al. (2007) they may have been more arboreal than *Au. afarensis*.

Au. africanus were less sexually dimorphic than *Au. afarensis*, with males averaging 4′6″ (138 cm) tall and 90 lb (41 kg) and females at 3′9″ (115 cm) and 67 lb (29 kg).

Characteristics

Review of Primitive Characteristics

- Somewhat prognathic.
- Arboreal characteristics:
  - Upward-oriented shoulder joints.
  - Long arms relative to legs.
  - Long, curved metacarpals and phalanges.

Review of Derived Characteristics

- Skull changes:
  - More rounded skull vault.
  - Increased encephalization.
  - Frontal and parietal expansion.
  - More of a forehead.
  - Loss of sagittal crest and less pronounced nuchal crest.
- Facial buttressing.
- More of a parabolic dental arcade and reduced anterior dentition.
- Power thumb:
  - Well-developed intrinsic thumb muscles.
  - Specialized first metacarpal.
- Reduced sexual dimorphism.

ENVIRONMENT AND WAY OF LIFE

While there is some controversy as to how much time *Au. africanus* spent in open versus closed (forest) environments, their facial robusticity and molars reflect a greater reliance on tougher, more dry-adapted plant foods than earlier hominins. They retained

climbing adaptations and thus could have used trees for safety, sleeping, and food. When on the ground, it is thought that they were scavenger-foragers, foraging for plant foods and small prey and scavenging remains of carnivore kills. Crude stone tools have been found at the sites of Sterkfontein and Makapansgat, and while there is no evidence of tool manufacture, it appears that they were using stones for hammering and cutting. They are thought to have relied on tubers for part of their diet and they likely used digging sticks to unearth them, in much the same way as modern African peoples, such as the San-speaking groups of the Kalahari Desert.

As mentioned, it is thought that hominins may have fallen through the surface into subterranean caves and/or bones were dragged in by rodents for gnawing. While it is possible that they used open caves for shelter from sun and rain, there is no evidence that they lived in caves.

In nature, extractive foragers have relatively large brains. Examples within the primate world include the aye-aye (a prosimian of Madagascar) that hunts insects via a combination of audition (hearing) and extraction (see www.arkive.org for multiple videos); New World capuchin monkeys that steal young animals from tree holes and rip bark from trees to get at insects; open-country baboons that dig for tubers; and the brainy orangutans and chimps that use a variety of tools to extract a range of foods, from ants to honey (see Figures 15.10 and 15.11 for videos of chimps and orangutan). Chimps have even been observed using sticks to spear bushbabies (small nocturnal African prosimians) in tree holes (e.g. see "How Smart Are Planet's Apes?" on NationalGeographic.com). The same can be seen in some bird species, most notably tool-using crows and nut-burying nuthatches. Since our last common ancestor with chimps and bonobos was already relatively smart, I believe that while bipedalism was in response to diminishing habitat and resources, encephalization was in response to having to locate and learn to process new foods. Beginning with *Au. africanus* and the subsequent paranthropines, we see an upward trend for encephalization and expansion of those portions of the brain involved with association, i.e. complex thought processing, as well as increasing manual strength and dexterity for manipulating tools and objects. Thus we stepped out and then had to find and fix dinner!

We tend to think of "evolution by means of natural selection" as being a positive aspect of life, i.e. "survival of the fittest." However, when there is a "rapid" response to a changing environment, many individuals die out while those few survive.

Videos





Video* 4.3.8.1*: *Ant dipping 1 by TheFriendsAndAi, youtu.be/ScqG54B4KtE*






Figure* 4.3.8.2*: *Orangutan using tool by Alain Compost, youtu.be/pXOTAgq-eiY*

This page titled 4.3.8: Australopithecus africanus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **3.8: Australopithecus africanus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

4.3.9: Middle Hominid Cranium Comparison Checklist

Name: _____

Section: _____

Student ID#:_____

MIDDLE HOMINID CRANIUM COMPARISON CHECKLIST

Martin Nickels

© 1999 ENSI (Evolution & the Nature of Science Institutes) www.indiana.edu/~ensiweb

This material may be copied only for noncommercial classroom teaching purposes, and only if this source is clearly cited.

PLEASE READ THESE 5 STATEMENTS BEFORE BEGINNING THIS ACTIVITY:

1. Work in groups of 3-4 students so that everyone can be involved in the activity.

2. BE SURE (!) TO TAKE TURNS doing different measurements and observations.

3. When taking a measurement, use the SLIDING CALIPERS (except for #11 & #12 which may require the HINGE calipers) and remember to...

4. ALWAYS MEASURE IN MILLIMETERS [mm] and round off to whole numbers.

5. PLEASE DO NOT ADD ANY PENCIL OR PEN MARK "TATTOOS" TO THESE CRANIA, OR STICK YOUR FINGERS IN THEIR EYE ORBITS OR NOSES!

I. BRAINCASE: (7 items #1-7)

1. Does the FOREHEAD (frontal bone) look more vertical OR flatter when the skull is held in normal anatomical position [NAP] (i.e., with the eyes oriented forward)?

2. Is a SUPRAORBITAL BROWRIDGE present?

3. If present, is the BROWRIDGE DIVIDED in the middle, or CONTINUOUS?

4. What is the SHAPE OF THE BRAINCASE (front to back) when viewed from above?

5. Is a SAGITTAL CREST present?

6. In NAP, is the FORAMEN MAGNUM oriented more downward OR more to the rear?

7. Is the MASTOID process relatively flat OR does it noticeably protrude (project)?

II. FACE: (5 items: #8-12)

8. Are the NASAL BONES raised (arched) OR flat?

9. Measure the MAXIMUM BREADTH (width) of the NASAL OPENING [mm].

10. Measure the MAXIMUM HEIGHT of the NASAL OPENING [mm].

11. Measure the LENGTH of the MAXILLA (the upper jaw) [mm]. (Measure down the middle of the palate from the front edge of the foramen magnum to either between or just in front of the two central incisors to determine how much the face projects forward.)

12. Measure the BIZYGOMATIC BREADTH using the hinge caliper if necessary [mm]. (This is the width or breadth of the face from the widest part of one zygomatic arch to the widest part of the other zygomatic arch.)

III. DENTITION: (6 items #13-18)

13. SHAPE OF THE DENTAL ARCADE: Do the tooth rows diverge towards the back OR are they more straight-sided and parallel to one another?

14. When viewed from the side, are the INCISORS angled out OR are they vertical?

15. Measure the COMBINED WIDTH or BREADTH of the 4 INCISORS together.

16. Does the CANINE tooth project above the chewing surfaces of the other teeth?

17. Is a CANINE DIASTEMA present?

18. Measure the COMBINED LENGTH of the LEFT 2 PREMOLARS and 3 MOLARS together by measuring from the back of the last molar to the front of the first premolar to determine the length of the chewing surface of the "cheek teeth". [mm]. (NOTE: Measure the right side if the left side is missing any of these 5 teeth.)

Source:

www.indiana.edu/~ensiweb/lessons/hom.cran.html

	Skull #1	Skull #2	Skull #3	Skull#4
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				

19) What trends do you notice? What are the major differences between these skulls?

This page titled 4.3.9: Middle Hominid Cranium Comparison Checklist is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Alex A. G. Taub via source content that was edited to the style and standards of the LibreTexts platform.

- **1.9: Middle Hominid Cranium Comparison Checklist** by Alex A. G. Taub is licensed CC BY-NC-SA 3.0. Original source: https://socialsci.libretexts.org/api/deki/files/94257/biological-anthropology-laboratory-activities.pdf.

CHAPTER OVERVIEW

5: Human Fossil Record

5.1: Pleistocene Epoch

5.1.1: Introduction to the Pleistocene Epoch

5.1.2: Paranthropines

5.1.3: Australopithecus

5.1.3.1: Australopithecus/Paranthropus aethiopicus

5.1.4: Paranthropus boisei

5.1.5: Paranthropus robustus

5.1.6: Australopithecus garhi

5.1.7: Australopithecus sediba

5.1.8: Genus Homo

5.1.9: Homo habilis

5.1.10: Homo rudolfensis

5.1.11: Homo species indeterminate

5.1.12: Homo naledi

5.1.13: The “erectus Grade”

5.1.14: Homo ergaster

5.1.15: Homo erectus

5.1.16: Homo georgicus

5.1.17: Homo antecessor

5.1.18: Homo floresiensis

5.1.19: Homo heidelbergensis

5.1.20: The Denisovans

5.1.21: Homo neanderthalensis

5.1.22: Homo sapiens

5.1.23: Recent Hominid Cranium Comparison Checklist

5: Human Fossil Record is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

5.1: Pleistocene Epoch

The Pleistocene epoch is often colloquially referred to as the Ice Age and lasted from about 2,588,000 to 11,700 years ago, spanning the world's most recent period of repeated glaciations.


5.1.1: Introduction to the Pleistocene Epoch
5.1.2: Paranthropines
5.1.3: Australopithecus
5.1.3.1: Australopithecus/Paranthropus aethiopicus
5.1.4: Paranthropus boisei
5.1.5: Paranthropus robustus
5.1.6: Australopithecus garhi
5.1.7: Australopithecus sediba
5.1.8: Genus Homo
5.1.9: Homo habilis
5.1.10: Homo rudolfensis
5.1.11: Homo species indeterminate
5.1.12: Homo naledi
5.1.13: The “erectus Grade”
5.1.14: Homo ergaster
5.1.15: Homo erectus
5.1.16: Homo georgicus
5.1.17: Homo antecessor
5.1.18: Homo floresiensis
5.1.19: Homo heidelbergensis
5.1.20: The Denisovans
5.1.21: Homo neanderthalensis
5.1.22: Homo sapiens
5.1.23: Recent Hominid Cranium Comparison Checklist

Thumbnail: Pleistocene of South America showing Megatherium and two Glyptodon. CC BY-sa 3.0; DiBgd).

This page titled 5.1: Pleistocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.


5.1.1: Introduction to the Pleistocene Epoch

THE PLEISTOCENE EPOCH (~2.6 mya – 11.7 kya)




East African grassland and a local. Photo by the author.

The Pleistocene Epoch is commonly known as the Ice Age. The climate of Africa continued on the trajectory that began in the late Miocene and continued throughout the Pliocene (see Figures IV.2 and IV.3). While the Pleistocene was characterized as a period of global cooling, glacial advances, and dropping sea levels, the cold periods were interspersed with interglacial periods when the ice retreated and sea levels rose (see Figures IV.2, IV.3, and IV.4). Even within glacial periods, the climate varied. Animals in northern areas that were not adapted to arctic conditions went extinct or moved south when temperatures dropped and vice versa. They pushed in and out of Africa, in response to those climatic pulses.




Global temperature fluctuations from Pliocene (5–1.8 mya), through Pleistocene (1.8–0.1 mya), to present. "Five Myr Climate Change" by Robert A. Rohde is licensed under CC BY-SA 3.0.



Antarctic temperature changes during the last several glacial and interglacial cycles of the present ice age and a comparison to changes in global ice volume. "Ice Age Temperature" by Robert A. Rohde is licensed under CC BY-SA 3.0.

Elizabeth Vrba claims that there was a pulse-like turnover in faunal populations in East Africa at ~2.5 mya, in response to climatic change. Termed her "Turnover Pulse Hypothesis," she finds evidence of an increase in bovids (cow-like animals like wildebeests and water buffalo) and their predators and a decrease in forest-dwelling ungulates, corresponding to an increase in dry savanna environment (see Vrba 1985 and subsequent publications). Based on deep sea cores, we know that by 1.8 mya, there was a definite

increase in the sahel-type of environment. Thus evidence from past flora and fauna supports the more robust craniofaciodental anatomy of species that lived during the early Pleistocene, e.g., *Au. garhi*, *Au. aethhiopicus*, *P. boisei* and *P. robustus*. They would have needed stronger chewing abilities and thicker molar enamel to process more dry-adapted vegetation. Once the more gracile (in term of their masticatory apparatus) *Homo* species came on the scene, the East African hominins would come to represent an example of niche partitioning. Fossil sites suggest that *P. boisei*, *H. habilis*, and *H. egaster* could have been **sympatric** (i.e. overlapping geographic ranges). Thus while *Au. aethiopicus* and *P. boisei* survived by what might be termed masticatory brawn, the more encephalized *Homo* species used their brains and tools to consume a higher-quality diet. We can thus see that hominins were some of the colonizing species of the African savanna.



Temperature reductions during last glacial maximum 18 kya. “CLIMAP” Robert A. Rohde is licensed under CC BY-SA 3.0.

This page titled 5.1.1: Introduction to the Pleistocene Epoch is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.1: Introduction to the Pleistocene Epoch** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.2: Paranthropines

The paranthropines are three species of fossil hominins that exhibited hyper-robust masticatory apparatus, as evidenced by their heavy faces and mandibles, huge molars, and muscle insertions. They were included in the genus *Australopithecus* for many years, but the original genus invented by Robert Broom for the South African form, *Paranthropus robustus*, has been revived for at least two of the species. Chronologically, the earliest species was *aethiopicus*, and some researchers still assign them to *Australopithecus*. The second oldest is *Paranthropus boisei*, and both *Au. aethiopicus* and *P. boisei* are from East African sites. The South African and youngest species is *P. robustus*. The last of the paranthropines died out ~1 mya, 1.7 mya after *Au. aethiopicus*'s first appearance in the fossil record. They were a very interesting and apparently successful group of animals that adapted to the changing African landscape by expanding their dietary niche. The geographic range of the eastern species stretched down the "**hominin corridor**" from Ethiopia to Malawi, whereas the South African form is known only from South Africa. While it is thought that their preferred foods were similar to the more gracile forms, they could fall back on tougher and harder foods when resources became scarce.

This page titled 5.1.2: Paranthropines is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.2: Paranthropines** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.3: Australopithecus

This page was auto-generated because a user created a sub-page to this page.

5.1.3: Australopithecus is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

5.1.3.1: Australopithecus/Paranthropus aethiopicus

Australopithecus/Paranthropus aethiopicus (2.7–2.3 mya)

("SOUTHERN APE" / "BESIDE HUMAN" / ETHIOPIA)



Model of Paranthropus aethiopicus. "Paranthropus aethiopicus" by Nrkpan is licensed under CC BY-SA 3.0.

SITES

- Ethiopia: Sungura Deposits
- Kenya: Westh Lake Turkana


PEOPLE

- Yves Coppens, Camille Arambourg, and Alan Walker

Australopithecus aethiopicus is the most primitive of the robust species. I use genus *Australopithecus* because it is thought to be descended from *Au. afarensis*. In addition, *Paranthropus* was the genus name assigned to the South African robust form, *P. robustus*, and questions remain as to whether the two species are related.

PHYLOGENY

There are multiple lines of evidence to support *Au. aethiopicus* as a descendent species of *Au. afarensis*. While some believe that *Au. aethiopicus* gave rise to *P. boisei*, others link *P. boisei* with *P. robustus* in a different clade, with *Au. africanus* as their common ancestor. More recently discovered material within the geographic range of *Au. aethiopicus* supports the *Au. aethiopicus* → *P. boisei* evolutionary scenario. The dates of the new fossils fall between the two species, and they possess intermediate or transitional characteristics. Figure 17.2 shows one cladistic schema that illustrates how some researchers suggest these species were related. This particular scenario shows the authors' belief that *Au. africanus* is a robust form.



Cladistic analysis of early hominins. "Cladistic analysis of early hominins" by Charles T. G. Clarke is licensed under CC BY-SA 3.0.

DISCOVERY AND GEOGRAPHIC RANGE

In 1967, the earliest *Au. aethiopicus* fossils were discovered by Yves Coppens and Camille Arambourg in the Shungura deposits at the site of Omo in southern Ethiopia. They assigned them to a new genus and species, *Paraustralopithecus aethiopicus*. While it was debatable as to whether they actually had a new species, the discovery of the "Black Skull" (see Figure 17.1) in the West Lake

Turkana region of Kenya by Alan Walker in 1985 put any doubts to rest. At that time, the species was added to genus *Australopithecus* because it was thought to be descended from *Au. afarensis*. There were then three recognized species of robust australopiths in Africa, and efforts to determine their phylogenetic relationships began.

The Black Skull or KNM-WT (Kenya National Museum – West Turkana) 15000 was a magnificent find. The almost complete skull was stained from manganese, but it is always fun to sing scary movie music to my students when introducing … THE BLACK SKULL! (Figure 17.3 … for fun!)



Skull and crossbones. "Crossbones (PSF)" by Pearson Scott Foresman is in the public domain.


PHYSICAL CHARACTERISTICS

A unique characteristic that ties *Au. aethiopicus* to *P. boisei* is a heart-shaped foramen magnum, as opposed to the more ovoid form seen in *Au. africanus* and *P. robustus*. Primitive characteristics shared with *Au. afarensis* are the flat cranial base, small brain (~410 cc), long molars (mesiodistally, i.e. front to back versus side to side), and the degree of prognathism in the lower face. Because their faces were so broad and their brains so small, they exhibit a high degree of postorbital constriction (also known as waisting), i.e. the area between the face and braincase is narrow. Derived robust characteristics are buttressing of the skull, face, and mandible. Their muscles of mastication were incredibly strong, as evidenced by the sagittal crest running down the midline of their skull where the temporalis muscle originated. The sagittal crest was higher and more posteriorly placed than in the two more derived robust species.

The zygomatics were large and flared to allow for passage of the temporalis muscle from the sagittal crest to insert on the mandible and to expand the attachment site for the masseter muscle, the other large muscle of mastication (Figure 17.4). The zygomatics were more forwardly flared than in the other two robust species. They also had heavy nuchal (posterior neck) muscles to support the weight of their heavy face and skull, and the attachment sites of those muscles on the posterior skull was also an expanded crest that converged with the sagittal crest, i.e. a compound sagittal-nuchal crest. Large brow ridges in the robust species offset the stress generated by biting down on hard foods. However, *Au. aethiopicus*'s brows were smaller than the other two species. Their third maxillary molars were convergent, meaning they were positioned more medially than the first and second. While some researchers disagree, others find evidence for a more derived venous sinus system in the crania of the robust forms. The system consists of large collecting veins that ultimately empty into the jugular veins, allowing for rapid, gravity-fed blood drain from the brain, in order to keep fresh blood pumping in via several arterial systems. The largest and most superficial of those sinuses (see Figure 17.5) leave impressions on the inside of the skull vault. Dean Falk's "Radiator Theory" argues that our ancestors needed to keep their brains cool as they increased in size in a hot, open environment. The system became more complex after the time of the australopiths.



Muscles of mastication (Red=temporalis and peach=masseter) Illustration by Keenan Taylor



Cranial sinus system (SIN. = sinus). Based on Gray's Anatomy Plate 488. "Gray488 blue" by Wikid77 is in the public domain.

Characteristics

Review of Primitive Characteristics

- Flat cranial base.
- Small brain.
- Long molars.
- Prognathic.

Review of Derived Characteristics

- Heart-shaped foramen magnum.
- Robust skull, face, and mandible.
- Large, compound sagittal-nuchal crest.
- Large brow ridges.
- Pronounced postorbital constriction.
- Large, powerful muscles of mastication.
- Large forwardly flared zygomatics with a large zygomatic arch.
- Convergent maxillary third molars.
- Possible derived venous sinus system.

ENVIRONMENT AND WAY OF LIFE

As mentioned, there is evidence that *Au. afarensis* was more terrestrial than the southern australopith clade, suggesting that the classic ape environment had diminished. We know that grasslands were expanding and woodlands were shrinking. Since it is assumed that *Au. aethiopicus* is descended from *Au. afarensis*, the environment was favoring species with dietary adaptations that allowed them to survive.



“Australopithecus aethiopicus tool use” by Keenan Taylor.

This page titled 5.1.3.1: Australopithecus/Paranthropus aethiopicus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.3: Australopithecus/Paranthropus aethiopicus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.4: Paranthropus boisei

Paranthropus boisei (2.5 mya)

("BESIDE HUMAN" / BOISE: SURNAME OF ONE OF THE LEAKEY'S FINANCIERS)



Paranthropus boisei. "Paranthropus boisei side (University of Zurich)" by Nicolas Guérin is licensed under CC BY-SA 3.0.

SITES

- **Ethiopia:** Shungura Deposits and Konso
- **Kenya:** Chesowanja, West Lake Turkana, and Koobi Fora
- **Tanzania:** Olduvai and Peninj

PEOPLE

- Mary and Richard Leakey

In the first course that I took in physical anthropology, I was most fascinated by the *Paranthropus boisei* face from Olduvai Gorge (see Figures 18.1 and 18.5) and the Natron/Peninj mandible from the Peninj site near Lake Natron. I still remember the first time I saw them, and the species has always been for me one of the more interesting discoveries in paleoanthropology. Their faces, jaws, and cheek teeth were massive and truly unforgettable. While the Olduvai material is attributed to Mary Leakey, it was her husband Louis who announced to the world that he had his "man." He had been finding tools of what are now termed the Oldowan tradition and was hoping to discover who had made them when Mary discovered the first *P. boisei* fossils. They were assigned to a new genus and species: *Zinjanthropus boisei*, or "human from East Africa." "Zinj" is a derivation of the archaic term "Zanj" for an area of the East African coast, and Boise was the surname of one of the Leakeys' financial supporters (Wikipedia contributors 2015g).



Figure 5.1.4.2: Olduvai Gorge, by David Berkowitz, CC BY 2.0

Louis claimed that they had their toolmaker, but many did not agree with his conclusion. They thought he had found a big, dumb, herbivorous, bipedal ape that would not have had the cognitive abilities to produce stone tools. Hand bones have been attributed to the species by some researchers. If they are correct in their assignment, the species may have been capable of tool manufacture and efficient use. However, while their brains were somewhat larger than earlier species, we still do not know if they had the cognitive abilities and thus whether tools from East African sites can be attributed to them.

Fossils from more than 100 individuals have been recovered in the last 55 years. Over time, the genus has changed from *Zinjanthropus* to *Australopithecus* to *Paranthropus*, but some researchers are still using genus: *Australopithecus*.

Phylogeny

Support for *P. boisei* being descended from *Au. aethiopicus* has steadily increased. However, some still group *P. boisei* as a sister species of *P. robustus* and believe that they descended from *Au. africanus*. The two more derived forms share a molar trait with *Au. africanus*, in that the second molar (M2) is larger than the third (M3). Members of the latter group, such as Henry McHenry, contend that the more primitive *Au. aethiopicus* is an example of *homoplasy* or *convergent evolution*, making the robust traits homoplasies.



Figure **5.1.4.3**: Scientific reconstruction of Paranthropus boisei. "Paranthropus boisei" by Lillyundfreya is licensed under CC BY-SA 3.0.

Paleoanthropologists have tended to be conservative in their acceptance of homoplasies; common ancestry is more parsimonious.Except for the possible *Au. aethiopicus* → *P. boisei* scenario, the robust australopiths were evolutionary dead-ends as far as we know.

DISCOVERY AND GEOGRAPHIC RANGE

Mary Leakey discovered the first material in 1959 at Olduvai Gorge, Tanzania (see Figure 18.2). Nicknamed "Nutcracker Man," "Zinj," or "Dear Boy," the skull and face were dated to 1.7 mya. Fossils attributed to the species have since been found at other sites in Tanzania (Peninj), Kenya (Chesowanja, West Lake Turkana, and Koobi Fora in the East Lake Turkana area), and Ethiopia (the Shungura Deposits and Konso). Richard Leakey discovered the Koobi Fora fossils. Thus, like *Au. afarensis*, the species had a broad geographic range and survived for over a million years.

PHYSICAL CHARACTERISTICS

While the robust forms are somewhat larger than the gracile forms, they do not differ much postcranially. It is their skulls that set them apart; *P. boisei* had the most pronounced masticatory adaptations, so that relative to the other two species, they are termed "hyper-robust." Along with the other robust forms, they shared a buttressed skull, face, and mandible; large molars and premolars; a compound sagittal-nuchal crest (not compound in *P. robustus*); large muscles of mastication and nuchal muscles to support their heavy skulls; large, flared zygomatic arches; and a supraorbital torus to absorb the stress generated by chewing. According to some researchers, they shared the following with *P. robustus* and *Homo*: flexed skull base; more orthognathic face; lower face tucked under the braincase, thus increasing chewing force on the molars; a more parabolic dental arcade; and a longer thumb with broad, flat, distal phalanges that gave them better opposition and gripping power. The sagittal crest in *P. boisei* and *P. robustus* was more anteriorly positioned relative to *Au. aethiopicus*. In combination with less prognathism and the fact that their face was tucked under the braincase, the action and strength of the temporalis muscle was concentrated on the cheek teeth. The degree of robusticity and the size of craniofaciodental characteristics were unique to *P. boisei*. The zygomatics were large, heavy, and widely flared, making their faces very broad and the temporal fossa (the space between the zygomatic and the temporal bone) very deep. Again, this facilitated the passage of the temporalis muscle to insert on the mandible and expanded the attachment and anchoring of the

masseter muscle. The mandible was massive, with a very robust and deep body and tall rami (vertical side portions), and the temporomandibular (jaw) joint was exceptionally large. Their front teeth were so dwarfed by their enormous cheek teeth, they almost look juvenile. Their premolars were molarized, and some had a third root. The third molar exhibited a unique wrinkling pattern. Their **megadontia quotient** (MQ) was 2.7, meaning that their teeth were 2.7 times larger than would be expected. *P. boisei* had the largest supraorbital torus of the robust forms. They were somewhat more encephalized than past species, with a cranial capacity of 514 cc (range = 494–537 cc). Like all australopiths, the species was sexually dimorphic, with males at 4’6″ (137 cm) tall and 108 lb (49 kg) and with more pronounced sagittal-nuchal crests and females at 4’1″ (124 cm) and 70 lb (34 kg).



Figure **5.1.4.15**: Natron mandible. “Peninj mandible. Paranthropus boisei” by Matt Celeskey is licensed under CC BY-SA 2.0.

Review of Derived Characteristics

- Same robust characteristics as seen in *Au. aethiopicus* with the following differences:
  - More robust craniofaciodental characteristics.
  - Sagittal crest more anteriorly placed.
  - Large supraorbital torus.
  - Huge molars and premolars.
  - Large, heavier mandible.
- Encephalized.
- Flexed skull base.
- More orthognathic face.
- Molarized premolars.
- Lower face tucked under braincase, maximizing chewing force on molars and premolars.
- More parabolic dental arcade.
- Longer thumb with broad, flat distal phalanges.

ENVIRONMENT AND WAY OF LIFE

While they still had relatively long arms, one seldom hears anything about the arboreal habits of the robust forms. In East Africa, that is not too surprising since the forests had receded and the grasslands had expanded. By 1.8 mya, deep sea cores reveal an increase in the **sahel** type of environment, i.e. the area south of the Sahara that consists of dry grasslands and scrub forest. Based on **microwear** evidence from their molars, all or most of which are ground flat, *P. boisei* likely subsisted on open, dry-adapted plants. Silt on terrestrial foods would have contributed to the wear patterns. The availability of high-quality foods must have gotten even worse by the time of *P. boisei*, because their chewing and bite-force capabilities were increased relative to *Au. aethiopicus*. They ate a high proportion of C_4 plants, i.e. open and more dry-adapted grasses and shrubs. It is thought that they likely survived on hard, tough, and brittle fallback foods during periods of preferred resource scarcity. This is another possible example of niche partitioning, in that sympatric *Homo* species may not have been able to utilize or digest either food category. Isotopic analyses to calculate the strontium-to-calcium ratio in fossilized bones of *Au. africanus* and *P. robustus* have determined that they ate some animal matter. It is possible that *P. boisei*did as well. While we do not know if they were scavenging or hunting animals, it is very

possible that early hominins ate insects and insect larvae. All extant great apes consume social insects (ants and termites), and they derive a surprising degree of nutritional value from them. In traditional human societies, larvae are a favored resource.




Figure **5.1.4.6**: Paranthropus boisei: “King of the termite mound” by Keenan Taylor.

It seems likely that *P. boisei* lived in small groups … or did they? Living in a more open environment in equatorial Africa, there would seemingly have been safety in numbers. Since they were adapted to low-quality savanna resources and could fall back on food items that were unavailable to other sympatric species, they may have avoided strong within- and between-group feeding competition. We also know that pair-bonding may have been necessary, as females were not as self-sufficient as their quadrupedal ancestors. They had babies that could not hold on and were surrounded by males that could rape or commit infanticide. There is an argument that monogamy is a male adaptation that increases paternity assurance while reducing the risk of infanticide. We know that males compete for mates, and the high degree of sexual dimorphism in *P. boisei* supports that fact. With all of the preceding arguments in mind, is it possible for apes with a tendency toward monogamy to live together without all of the rules and laws seen in modern humans to keep male competition and jealousy in check? While we could argue that they also could have lived in pairs with their dependent offspring and never ventured far from trees, how did females climb with their infants? We still have many unanswered questions about hominin behavioral ecology!

This page titled 5.1.4: Paranthropus boisei is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.4: Paranthropus boisei** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.5: Paranthropus robustus

Paranthropus robustus (2.3 mya)

("BESIDE HUMAN" / ROBUST)



Paranthropus robustus. "Sterkfontein Caves 66," photograph by Mike Peel (www.mikepeel.net), is licensed under CC BY-SA 4.0.

SITES

- Swartkrans, Kromdraai, Drimolen, Gondolin, and Coopers Cave, South Africa

PEOPLE

- Robert Broom and Andre Keyser

In 1938, Robert Broom discovered the first *Paranthropus robustus* material at the site of Swartkrans, South Africa. He later found material at Kromdraai, and because the molar teeth were more primitive at that site, he changed the species name at Swartkrans to *P. crassidens* but used *P. robustus* for the Kromdraai material. Molar characteristics from the more recent material from the Drimolen site are thought to be intermediate between the Swartkrans and Kromdraai molars, and most researchers now consider the material from all three sites to be species: *robustus*.

PHYLOGENY

Au. africanus is the favored ancestor of *P. robustus*. However, others believe that *P. boisei* and *robustus* are descended from *Au. aethiopicus*. Of the former school, some believe that *P. boisei* is also descended from *Au. africanus* and thus a sister species to *P. robustus*. The two species would then have inherited their flexed skull base from *Au. africanus*. Like *P. boisei*, *P. robustus* appears to be an evolutionary dead end.

DISCOVERY AND GEOGRAPHIC RANGE

The species is known only from sites within the Cradle of Humankind World Heritage Site in South Africa. As mentioned in the introduction, Robert Broom discovered the first material at Swartkrans and subsequent specimens at the site of Kromdraai. Andre Keyser discovered the jaw and dental material at Drimolen in 1994.

PHYSICAL CHARACTERISTICS

Like *P. boisei*, *P. robustus* exhibited buttressing of the skull, face, and mandible; orthognathism in that the teeth were tucked under the cranial base; small anterior dentition; molarized premolars; large muscles of mastication; large zygomatic arches for passage of the temporalis muscle; sagittal crests in males and a nuchal crest that did not converge with the sagittal crest; tall mandibular rami

to increase the strength of the masseter and medial pterygoid (another muscle of mastication) muscles for their tough, fibrous diet and a large mandibular body; and a high degree of postorbital constriction due to their large face (see skull cast in Figure 19.1).

Characteristics shared with *Au. africanus* are a flexed skull base, megadontia (*P. robustus*'s molars were 17% larger, but their MQ of 2.2 was lower), molarized premolars, a larger second than third molar, and facial buttressing. While *P. robustus*appears to have been more orthognathic than *Au. africanus*, they were not. Their anteriorly oriented orbits and zygomatics reduced the distance between their midface and jaws, making it appear so. The nasal bones were recessed relative to the forward-oriented zygomatics, so that they too had a somewhat dish-shaped midface like *Au. africanus*. They had less of a forehead than *Au. africanus* because of the forward-oriented orbits. Finally, unlike *Au. africanus*, where maximum force was on the molars, the maximum force was on the premolars due to a combination of the forward placement of the zygomatics and an enlarged anterior portion of the temporalis muscle that was offset by the positioning of the dental arcade under the braincase (Cartmill and Smith 2009).

Average cranial capacity is estimated to have been 530 cc, giving them the highest EQ, 3.0, of any australopith (Cartmill and Smith 2009).

The postcranial morphology of *P. robustus* shares some similarities with *Homo*, such as broad distal finger phalanges that are thought to indicate enlarged tactile pads and increased vascularization, sensitivity, and motor control, as well as a large attachment for the *flexor pollicis longus* muscle that acts as a powerful thumb flexor. They thus are thought to have had a great degree of manual dexterity and were likely capable of making and certainly using tools. However, they retained some primitive australopith characteristics, including long arms; small vertebral bodies, especially inferiorly; small sacroiliac and hip joints; more posteriorly oriented ilia; and a long femoral neck. Males of the species are thought to have stood 4′ (1.2 m) tall and weighed 120 lb (54 kg) and females, 3′2″ (<1.0 m) and 90 lb (40 kg).

Characteristics

Review of Primitive Characteristics

- Retain prognathism, concave facial profile, long arms, small vertebral bodies, small sacroiliac joints, posteriorly-oriented ilia, and long femoral neck of Au. africanus.

Review of Derived Characteristics

- Same characteristics as P. boisei, except not as robust and large.
- Maximum bite force on premolars.
- Encephalized.

ENVIRONMENT AND WAY OF LIFE

Like *P. boisei*, *P. robustus* are thought to have been generalist herbivores that may have consumed some animal matter and could fall back on hard and brittle items, such as nuts, seeds, and hard-skinned fruits, when preferred items were not available. C. K. Brain found wooden digging sticks in association with *P. robustus* remains. In addition, polish on bone and horn cores (bone interior of animal horns) attributed to *P. robustus*is consistent with repeated digging, such as for tubers.

Of interest is recent research that indicates that they were more male- than female-philopatric, supporting the notion that we have inherited the chimp and bonobo pattern of females relocating to join and range with a group of males. Amazingly, isotopic analyses of strontium in their teeth indicates that females did not grow up where their fossils are found (Copeland et al. 2011).

This page titled 5.1.5: Paranthropus robustus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.5: Paranthropus robustus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.6: Australopithecus garhi

(2.5 mya)

(“SOUTHERN APE” / “SURPRISE” IN THE AFAR LANGUAGE)



Figure 5.1.6.1: Reconstructed Australopithecus garhi skull. “Musée national d’Ethiopie-Australopithecus garhi (2)” by Ji-Elle is licensed under CC BY-SA 3.0.

SITES

Bouri site in the Middle Awash area of the Afar Depression, Ethiopia

PEOPLE

Tim White and Berhane Asfaw

In 1996, researchers recovered portions of the frontal and parietal bones as well as a maxilla that contained teeth (Figure 5.1.6.1). These materials were attributed to *Australopithecus garhi*. While nearby limb bones could not be attributed to the species with absolute certainty, they have been used by some paleoanthropologists to describe the species’ characteristics. Thus there is very little useful material to “reconstruct” this species. Fossil-containing sediments also contained bones with cut marks and a few surface cores (shaped and modified rock) and flakes (sharp pieces of rock struck from a core) were found, suggesting that *Au. garhi* butchered animal remains and possibly made tools. However, tools in fossil-bearing layers would be better evidence. Some believe that the manufactured stone tools at the nearby Gona, Ethiopia, site may have been manufactured by *Au. garhi*.

PHYLOGENY

The species is likely descended from *Au. afarensis*, possibly directly so. Like *Au. afarensis*, *Au. garhi* is thought to have been more terrestrial than the South African australopiths. As is the case for the rest of the australopiths, some researchers believe them to be ancestral to genus *Homo*, and what little evidence there is for tool manufacture accords with our lineage.

DISCOVERY AND GEOGRAPHIC RANGE

The species was discovered in 1996 by Tim White’s crew, and later Berhane Asfaw became involved. The only fossil material comes from the Bouri site in the Middle Awash region of the Afar Depression in Ethiopia.

PHYSICAL CHARACTERISTICS

The maxilla and teeth are larger and more robust than those of *Au. afarensis*, and some researchers lump them with the paranthropines. Thus like *Au. aethiopicus* and *P. boisei*, *Au. garhi* were adapted to a broader dietary niche in response to environmental changes, particularly expanding grasslands. If the limb bones are rightfully attributed to the species, they had longer, more humanlike legs than other australopiths. Their arms were still apelike, based on the ratio of the arm to forearm length. The cranial capacity was 446 cc, falling midrange within that of *Au. afarensis*.

Characteristics

Review of Primitive Characteristics

- Ape-like arms.
- Small brain.

Review of Derived Characteristics

- Robust craniofaciodental characteristics.
- Possibly longer, more human-like legs.

This page titled 5.1.6: Australopithecus garhi is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.6: Australopithecus garhi** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.7: Australopithecus sediba

(“SOUTHERN APE / “FOUNTAIN OR WELLSPRING” IN THE SOTHO LANGUAGE)



Figure **5.1.7.1**: Australopithecus sediba holotype. “Australopithecus sediba” by Brett Eloff is licensed under CC BY-SA 4.0.

SITES

Malapa, South Africa

PEOPLE

Lee and Matthew Berger and subsequent colleagues

Six well-preserved individuals of a new species of *Australopithecus* were discovered, beginning in 2008, at the cave site of Malapa, South Africa. Lee Berger’s crew is credited with the discovery after Berger’s nine-year-old son Matthew (Figure **5.1.7.2**) happened upon the fossils of a juvenile male (MH1) that became the holotype for the species (Figure **5.1.7.1**). The other five individuals were an adult male, an adult female (MH2) and, remarkably, an infant. It is of great utility in terms of interpreting morphology that much of the fossil material was found *in situ* (i.e. where the body settled versus missing or scattered about), demonstrating that they became buried and began fossilizing fairly rapidly. It is thought that they fell into a cavern, possibly lured by the presence of water in that then dry environment.



Figure **5.1.7.2**: Nine-year-old Matthew Berger with fossil discovery. “Matthew Berger with Malapa Hominin 1” by Lee R. Berger is licensed under CC BY-SA 3.0.

PHYLOGENY

Since *Au. sediba* shares characteristics with both *Au. africanus* and *Homo*, it is thought to possibly be intermediate between the two species. That scenario, of course, disagrees with *Au. afarensis* as being our direct ancestor, with *Au. africanus* as a side branch. According to Berger et al. (2010), *Au. sediba* is more distinct from *Au. africanus*than the latter is from *Au. afarensis* in hand, pelvis, foot, and ankle morphology. However, the team working with the material believe it should stay within the genus *Australopithecus* because it lacks the *Homo* encephalization trend. Surprisingly, *Au. sediba* is more similar to *Homo erectus* in some respects than to Early *Homo* (*H. habilis*or *rudolfensis*). There has also been speculation that *Au. sediba* is a derived form of *Au. africanus*, but it is problematic in that the Malapa specimens are only 100 kya more recent than the youngest *Au. africanus* material. Furthermore, because the Malapa material is contemporary with Early *Homo* material (*H. habilis* and *rudolfensis*), some insist that they cannot

be ancestral to those species. However, Pickering et al. (2011) dispute that claim, since Early *Homo* material is still problematic, both taxonomically and spatiotemporally. Berger et al. (2010) have proposed four phylogenetic scenarios—that *Au. sediba* was the ancestor of *Homo habilis*, *Homo rudolfensis*, or *Homo erectus*, or was a sister species to the ancestor of the *Homo* lineage.



Figure **5.1.7.3**: *Malapa site. "Malapa fossil site, August 2011 site of discovery of Australopithecus sediba – view North" by Lee R. Berger is licensed under CC BY-SA 3.0.*

DISCOVERY AND GEOGRAPHIC RANGE

Because the "Malapa Party" (my term) discovery is such a great story, I used it to introduce the species and will not repeat it here. Malapa (Figure **5.1.7.3**) is 9.3 km northeast of Sterkfontein and 45 km north-northwest of Johannesburg in the Cradle of Humankind World Heritage Site (Figure **5.1.7.4**) (Wikipedia contributors 2015f). No fossil material has been found elsewhere.




Figure **5.1.7.4**: *Malapa area indicated on map of South African provinces. "Map of South Africa with provinces shaded and districts numbered (2011)" by Htonl is licensed under CC BY-SA 3.0.*

PHYSICAL CHARACTERISTICS

In general, the species' morphology is a mosaic of australopith- (especially *Au. africanus*) and *Homo*-like characteristics, but there are multiple lines of evidence to support its classification as a separate species (Berger et al. 2010). Since the Malapa material provides some of the best *in situ* evidence for many body parts, relative to other contemporary and previous species, I will spend more time discussing the physical and biomechanical significance.

The brain of *Au. sediba* was australopith-like in its size and "convolutional patterns" (Carlson et al. 2011). However, derived aspects of the frontal lobe show that *Homo*-like reorganization preceded pronounced encephalization (Carlson et al. 2011). This is an exciting discovery, in that we do not see so much of a gradual increase in size but rather that the hominin brain evolved in a mosaic fashion and that an increase in the association region was favored at the time. If cranial capacity alone was all that was available to infer cognition in *Au. sediba*, that important fact would be missed.




Figure **5.1.7.5**: Lee Berger with MH1. "Lee Berger and the Cranium of Australopithecus sediba MH1" by Brett Eloff is licensed under CC BY-SA 3.0.

Like most areas of the body, the cranial morphology reflects australopith- (especially *Au. africanus*) and *Homo*-like characteristics (Berger et al. 2010). Relative to *Au. africanus*, *Au. sediba* had smaller teeth and a less robust face (Figure **5.1.7.5**). Of interest is that they are described as having an "incipient" nose, i.e. the beginnings of a protruding nose (Berger et al. 2010). They share various mandibular and dental characteristics with *Au. africanus*, Early *Homo*, and *Homo erectus* (Irish et al. 2013; de Ruiter et al. 2013). Mandibular size and shape and tooth size most closely resemble Early *Homo*, and while some tooth measures and growth pattern are most similar to *H. erectus*, the overall growth trajectory is unique compared to other fossil hominins (de Ruiter et al. 2013). Analysis of tooth shape, molar cusp patterns, and root number of the various teeth shows that *Au. sediba* and *Au. africanus* share five synapomorphies (shared traits among related species), thus supporting a South African australopith clade. In addition, *Au. sediba* shares five synapomorphies with *H. habilis/rudolfensis/erectus*, suggesting a South African australopith ancestry for our lineage (Irish et al. 2013).

Au. sediba retained relatively long arms and elevated shoulder joints for climbing (see Figure 21.6) (Kivell et al. 2011; Schmid et al. 2013). Except for three wrist bones and the distal phalanges of the four fingers, the right distal forearm, wrist, and hand have been recovered intact for the adult female MH2. While the morphology of the hand is unique in some ways and they were still capable of strong flexion for climbing, some traits are associated with tool production in *Homo* species, especially the longer, more dexterous thumb and shorter fingers that facilitate our strong precision pinching. The only other hand that is well preserved from the late Pliocene is that of OH 7, attributed to *Homo habilis*. However, the Malapa hand was better suited to tool production and there is still speculation that OH 7 was actually *P. boisei*, a robust australopith not in our lineage. Thus, *Au. sediba* may have been responsible for the tools that have been recovered at South African sites (Kivell et al. 2011).



Figure **5.1.7.6**: *Australopithecus sediba compared with Lucy. From left to right: MH1, Lucy, MH2. "Australopithecus sediba and Lucy" by Peter Schmid is licensed under CC BY-SA 3.0.*

While determining the thoracic shape of fossil hominins has been a problem due to the lack of intact specimens, it is generally thought that australopiths had a conical thorax like extant great apes. A conical thorax is involved with arboreal locomotion, whereas our barrel-shaped thorax is (1) adapted for endurance walking and running, (2) related to shoulder breadth and arm-swinging to counter trunk rotation and facilitate our greater respiratory abilities, and (3) facilitated by a decrease in gut volume related to encephalization and a less herbaceous diet (first reviewed in Bramble and Lieberman 2004 and later in Schmid et al.

2013). The *Au. sediba*specimens support the notion of a conical shape to the upper thorax, but surprisingly, the lower thorax is not as flared as was previously thought, being more human-like. Schmid et al. (2013) assert that the “uncoupling” of the upper and lower trunk morphology in *Au. sediba* argues against a physiological response, in favor of a biomechanical one. The conical upper thorax would have been useful for climbing but would not have allowed for effective arm-swinging or heavy breathing and thus they were likely not capable of exertive walking and running. Their gut must have been smaller than a typical ape’s, suggesting a less herbaceous diet and setting the stage for the expensive tissue trade-off allowing for encephalization in our lineage (Schmid et al. 2013).

Au. sediba likely possessed the modern number of regional vertebrae. While the vertebrae of all hominoids increase in size from superior to inferior, earlier hominins lacked enlarged lumbar and sacral vertebral bodies and our pronounced sacral curve (see Williams et al. 2013 for references). *Au. sediba* is the earliest hominin to exhibit those characteristics, which are present in *Homo erectus* and all subsequent species. While the vertebrae in the five regions of our vertebral column have distinctive characteristics, they morphologically grade into one another so that there are transitional vertebrae, e.g. the seventh cervical vertebra resembles the first thoracic vertebra and so on. Unlike humans, the transitional thoracolumbar vertebra in *Au. sediba*, *H. erectus* (based on the Nariokotome specimen), and possibly *Au. africanus* is 11 versus 12. Thus while vertebra 12 is rib-bearing, it also likely functioned to elongate the lumbar region, increasing lower back flexibility (Williams et al. 2013).

The innominates were australopith-like in dimensions involved with childbirth but exhibited buttressing and other characteristics of *Homo*, such as iliac shape and vertical positioning, shortened ischium, and so forth (Kibii et al. 2011). The proximal femur was australopith-like with a small head and long neck, but the bicondylar/carrying angle and insertion of the gluteus maximus were humanlike. The knee exhibits both australopith- and human-like characteristics. Like us, they were capable of fully extending their knees during the swing phase of walking (DeSilva et al. 2013).

The partially articulated ankle and foot elements present a unique combination of primitive and derived traits. Humanlike characteristics include the morphology of the distal tibia, Achilles tendon insertion (calcaneal tuberosity), possible valgus (medially inverted) knee indicative of the bicondylar angle, probable arched foot, etc. Some aspects of the bones are more australopith-like, e.g. more gracile calcaneus; and some are more ape-like, e.g. robust medial malleolus (distal fibula, what we think of as our medial ankle “bone”) and some joints and muscle attachment sites (Zipfel et al. 2011).

Like in apes, initial ground contact was via the heel and lateral aspect of the foot, so that the foot was inverted. They then hyperpronated the foot by transferring weight medially so that the ankle was inverted and the plantar surface of the foot rolled medially and became everted. Modern humans with hyperpronation have problems with increased stress and wear on the joints, and the same is evident in *Au. sediba*remains (DeSilva et al. 2013). Hyperpronation, along with a robust medial malleolus, flexible midfoot, and upper limb adaptations, was apparently an adaptive response to bipedal terrestrial locomotion and retention of arboreal climbing abilities (J.M. DeSilva, personal communication with author, 2014). As in the ardipiths, the divergent hallux likely aided in climbing and would have given them more of a tripod foundation to stabilize their upright stance when standing and walking (Lovejoy et al. 2009).

Taken together, the mosaic of lower limb characteristics suggest that *Au. sediba*may have practiced a form of bipedalism unique from australopiths, or that they were descended from a more terrestrial ancestor and they then reverted to a more semi-arboreal lifestyle (DeSilva et al. 2013).

Characteristics

Review of Primitive Characteristics

- Brain size and convolutions.
- Australopith-like growth trajectory of teeth and jaws.
- Climbing adaptations: elevated shoulder joint, long arms, and strong hand flexion.
- Conical upper thorax.
- Australopith-like femoral head and neck.
- Some primitive aspects of ankle and foot.
- Retention of semi-divergent hallux.

Review of Derived Characteristics

- Reorganization of the frontal cortex.
- Small teeth and jaws and less robust face.
- *Homo*-like tooth shape and growth pattern.
- Humanlike lower thorax.
- Increased manual dexterity and precision grip.
- Longer, more dexterous thumb.
- Shorter fingers.
- Enlarged lumbar and sacral bodies and sacral curve.
- Buttressed and more vertical ilium.
- *Homo*-like carrying angle and *gluteus maximus* insertion.
- Full knee extension.
- Humanlike aspects of tibia, ankle, and foot.
- Possibly unique form of bipedalism, e.g. hyperpronation.

ENVIRONMENT AND WAY OF LIFE

The habitat of *Au. sediba* is thought to have been a mosaic environment of wood- and grasslands. Phytoliths from their tooth enamel indicate that they had access to forest products. Phytoliths are species-specific silica bodies in plants that can be used to identify what plant species were consumed. Fruit, leaves, wood, and bark were identified, with the latter being a first for fossil hominins (Henry et al. 2012).

Unlike other australopiths and paranthropines, the diet of *Au. sediba* most closely resembled savanna-dwelling chimps and secondarily *Ar. ramidus*, in that they specialized on C_3 plants, such as herbaceous plants, shrub and tree plant products, and grasses, and possibly animals that consumed those items. We know that C_4 plants were available at the time because remains of those more open, dry-adapted plants have been recovered in the same layers within which the Malapa material was found. In addition, C_4-consuming rodents, horses, and bovids were resident at the time. Their diet was thus less “carbonically” (I made that up for fun!) varied than contemporary hominin species inhabiting similar environments, such as *Au. africanus*and *Paranthropus robustus*, as well as *Homo* species that had an even broader diet. Microwear analyses show that the two Malapa individuals (MH1 and MH2) under study ate a greater percentage of hard food items relative to other australopiths, being more similar to *P. robustus* or *H. erectus* (Henry et al. 2012).

Being C_3 specialists in a C_4-dominated environment may have involved more time invested in foraging, or they may have favored a more closed environment for feeding. It is thus likely that there were enough woodland or riverine environments to have supported the population. Additionally, they could have consumed hard items when available or as fallback foods.

This page titled 5.1.7: Australopithecus sediba is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.7: Australopithecus sediba** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.8: Genus Homo

It is generally thought that by 2.5 mya, there were at least two species of *Homo* in East Africa, *Homo habilis* and *Homo rudolfensis*. The inclusion of those fossils in our genus is not accepted by all and is somewhat arbitrary. Some argue that *H. habilis* does not differ enough from australopiths to warrant different genus designation. Its inclusion in *Homo* was prompted by the fact that they are thought to have made and used tools and thus to have been cognitively advanced. *H. habilis* was more encephalized than the australopiths, and the skull vault is flexed as in *Au. africanus*, *P. boisei*, *P. robustus*, and later species of *Homo*.

There is debate as to whether *H. habilis* or *H. rudolfensis* gave rise to *Homo ergaster* (African form of the *erectus*-grade) and hence should be included in our lineage. Both species overlap the more derived *H. ergaster* in time and geographic space. The size and architecture of the brain of *H. habilis* make it a contender in the minds of some researchers. However, their limb proportions, i.e. retention of long arms and short legs, do not resemble *H. ergaster*. According to some researchers, *H. rudolfensis* possessed more modern femora (plural of femur), limb proportions, and a more orthognathic face, thus making them a better candidate for the ancestor of *H. ergaster*. However, it has not been determined that postcranial remains can be definitively assigned to the species, because postcrania are rarely found in association with cranial material (from which species designation derives). In addition, the claim to a more orthognathic face has been called into question based upon biometric impossibilities—the reconstruction of the face is impossible in terms of the relative positioning of key features (Bromage et al. 2008). While some claim that *H. rudolfensis* were more encephalized than *H. habilis*, others believe that their relative brain size (i.e. brain size taking into account body size) was lower and the temporal region, which is a conservative region of the skull, was more primitive. They also had more robust faces and teeth, unlike *H. ergaster*.

This page titled 5.1.8: Genus Homo is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.8: Genus Homo** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.9: Homo habilis

(2.3 mya)

("SAME" / "HANDY," "ABLE," ETC.)



Scientific reconstruction of Homo habilis. "Homo habilis" by Lillyundreya is licensed under CC BY-SA 3.0.

SITES

- Ethiopia: Hadar (and possibly Omo)
- Kenya: Koobi Fora
- Tanzania: Olduvai Gorge
- South Africa: Swartkrans and Sterkfontein

PEOPLE

- Mary and Louis Leakey, Donald Johanson, Tim White, and others

Of the two species of Early *Homo*, *Homo habilis* is the favored ancestor of *Homoergaster* and all subsequent hominin species.

PHYLOGENY

While the origin of *Homo habilis* has been in a state of flux in recent years, the discovery of *Au. sediba* has raised more questions about the origin of our genus. The discovery of Lucy in the early 1970s led some researchers to turn away from *Au. africanus* in favor of *Au. afarensis* as the ancestor of genus: *Homo*. In recent years, the idea that a cladistic event had occurred with *Au. afarensis*, leading to *Au. africanus* and the more derived robust forms on the one hand and genus *Homo* on the other, gained in popularity. *Au. sediba* now seems to have bridged the gap between the australopiths and genus *Homo*, sharing characteristics with *Au. africanus*, *H. habilis*, and *H. ergaster*. The similarities with the two *Homo* species may help resolve the problem as to which of the two species of "Early *Homo*" gave rise to *H. ergaster*. There are proponents in support of each of the evolutionary scenarios, with their share of pros and cons.



KNM-ER 1813, Koobi Fora, Kenya. "Homo habilis-KNM ER 1813" by Locutus Borg is in the public domain.

DISCOVERY AND GEOGRAPHIC RANGE

Louis and Mary Leakey discovered the first fossil material in 1960 at their site in Olduvai Gorge, Tanzania. Louis had been recovering stone tools from the site for years, but the manufacturer of those tools had previously eluded him. He named the species *Homo habilis* or "handy-man." Fossils attributed to *H. habilis* have also been found at Hadar (and possibly Omo), Ethiopia; Koobi Fora, Kenya (see Figure 23.2); and the South African sites of Swartkrans and Sterkfontein.

PHYSICAL CHARACTERISTICS

H. habilis exhibited a high degree of sexual dimorphism, with males and females weighing 114 and 70 lb and standing 5´2˝ and 4´1˝, respectively. Their skull, face, and dentition were more gracile than the australopiths. Their teeth and dental arcades were very human-like. The skull base was flexed, as seen in *Au. africanus* and the more derived robust australopiths and, relative to past species, the skull was rounder and higher, reflecting architectural changes in the brain. Cranial capacity ranged from 500 to 800 cc with a mean of 631 cc. This gave them an EQ of 3.1–3.5. At this point in hominin evolutionary history, we see increased asymmetry in the two hemispheres of the brain, termed lateralization or left hemispheric dominance. The left side of our brain is involved with language and analytical processes. Like all Old World monkeys and apes, *H. habilis* possessed Broca's area, which is involved with language production. However, it was larger than in past hominin species, and they also possessed Wernicke's area, which plays a role in language comprehension. They thus had the neural capacity for language. The left hemisphere is also related to right-handedness. They may have exhibited our tendency to hold objects with our left hand while working on them with our right. The frontal lobe, important in association processes, was expanded and resulted in more of a vertical forehead. The enlarged brain may have been facilitated by a decrease in gut volume, combined with a higher-quality diet that resulted from increased cognitive capabilities and an expanded technology base.

H. habilis had a smaller supraorbital torus and its face was more orthognathic than its supposed ancestor, *Au. africanus*, but they retained some prognathism in the lower face. They had fairly large ape-like incisors, but their canines, premolars, and molars were reduced in size. The mandible was more gracile, reflecting their reduced masticatory capabilities.

Like the majority of the australopiths, *H. habilis* possessed elongated arms, possibly suggesting continued reliance on an arboreal environment. While the digits were still curved, they had increased gripping capabilities for tool manufacture and use, as evidenced by the pronounced attachment site for the *flexor pollicis longus* muscle, which acts to flex the thumb.

The femoral head was enlarged and the neck shortened. Those changes are thought to have been the result of increased strain generated by an expanded pelvis for birthing larger-brained infants. However, no fossilized pelvic fossils have been found. Their foot was more modern, in that the hallux was no longer divergent but rather aligned with the lateral four digits, and the toes were shorter. They had less mobility in their feet, in that the foot had become more of a support structure like our own. The metatarsals were thick relative to modern feet, and the morphology of the third metatarsal suggests that they did not yet exhibit the degree of weight transfer and propulsive capabilities seen in modern humans.

Characteristics

Review of Primitive Characteristics

- Some prognathism.
- Large incisors.
- Curved phalanges.
- Long arms and short legs.
- Thick metatarsals.

Review of Derived Characteristics

- Gracile craniofaciodental characteristics:
  - Thin skull vault.
- More globular cranium.
- Expanded frontal lobe.
- Left hemispheric dominance.

- Enlarged Broca's and Wernicke's areas.
- Reduced supraorbital torus.
- Smaller mandible, canines, and cheek teeth.
- Parabolic dental arcade.
- Increased manual dexterity.
- Larger femoral head (and hence acetabulum) and shorter neck.
- More stable foot:
  - Loss of divergent hallux.
  - Shorter toes.

ENVIRONMENT AND WAY OF LIFE

Certainly one of the most interesting things about *H. habilis* is the appearance of a much more extensive archaeological record. The cultural period at that time, and extending through *Homo erectus*, is termed the Early Paleolithic, or the early portion of the Old Stone Age. While other species apparently preceded *H. habilis* in the manufacture of tools, it was thought for many years that they were the first to do so. The Oldowan or Olduwan tradition (industry and technology are also used synonymously with "tradition"), named after Olduvai Gorge, consisted of simple core tools and flakes. The technique involved the selection of a cobble (a workable-sized rock), followed by the use of a hammerstone to remove the outer rough surface (see Figure 23.3) or "cortex" and then to shape it into a core tool, by the removal of flakes. The flakes that are removed may be suitable for cutting and slicing. The process is called hard percussion, and the shaping is known as lithic reduction. "Lithic" refers to stone and is also used to denote a stone tool. Stone resources for the manufacture of tools were chosen for their suitability and transported across the landscape. Of course, this indicates a level of cognitive complexity, but we must remember that chimps and orangutans choose sticks and grass of particular widths and strengths, trim them to the appropriate length, and transport them in their mouths to their site of intended use.

Apes learn by trial and error, innovation and imitation, and cultural transmission, i.e. traits spread throughout a group by observation. Cultural transmission of innovations is even seen in monkeys, e.g. Japanese macaques washing sweet potatoes, skimming grain kernels floating on the surface to separate them from beach sand, and bathing in volcanic springs. While we do not know which species was the first to invent stone tools that were modified from their original form via lithic reduction and shaping, we can see the precursors of innovation and cultural transmission in our primate relatives. The real skill comes with having the manual dexterity to do so, making a tool that can accomplish a variety of uses, and the ability to teach others. I would argue that the earliest members of our genus had "theory of mind," i.e. the realization of another's thoughts. There is only one example of teaching in nonhuman primates and that was a mother chimp in the Tai Forest of the Ivory Coast that helped her daughter crack a nut, using their unique hammer and anvil technique. Our closest relatives, with all of their intelligence, symbolic capabilities as demonstrated in language studies, and similarities to our own behavior, do not know enough to teach their children. They are not capable of realizing that "I know something that you don't know" and vice versa. We go on and on about encephalization in the hominin lineage and technological advancements in the archaeological record over time, but what may have been the true dividing line between ourselves and the apes, whether bipeds or not, was the ability to teach our young, kin, and other group members and thus increase their chance of survival. The vehicle for developing a theory of mind is language. Human children develop a theory of mind at three or four years of age. Prior to that time, they do not realize that they or others may have incomplete information.

anecdote

Here is a fun anecdotal account that I always relay to my students:

My brother Michael was visiting my brother Jimmy. Jimmy was nowhere to be found when Michael realized that Jimmy's 18-month-old son had messed his diaper. Jimmy's older son must have been about three years old at the time. He helped Michael find everything that he needed to clean the baby. After Jimmy had reappeared and Michael had left for the day, the older boy remarked to his dad, "Uncle Mike is so dumb!" When asked what he meant by that, he replied, "He didn't know where the towels were; he didn't even know how to use the Diaper Genie® [a gizmo that turns dirty diapers into self-contained plastic coated links—truly magical!]."

This indicates that my nephew had not developed a theory of mind. He did not understand that Michael did not know things that he knew.




Hard hammer percussion. “Hard Hammer” by ZenTrowel is in the public domain.

Homo habilis was the first species to exhibit enlarged Broca’s and Wernicke’s areas. They thus may have had the motor control that allowed more lingual activity and the ability to comprehend the resulting sounds they could produce. Great apes can comprehend symbols, i.e. *this* stands for *that* even though *this* bears no resemblance to *that*. They have been taught American Sign Language, various computer languages, and spoken language. Where they fall short is in syntax—they cannot string together symbols into meaningful sentences. I firmly believe that the descendant species of Early *Homo*, i.e. *Homo ergaster*, had theory of mind, based on their stereotypical production of tools. There had to be teaching, learning, and training involved in order to produce an implement that is readily recognized as an Acheulian hand axe (see Figure 23.4). Thus, since we see an earlier stage of tool production in Early *Homo*, I would argue that they had rudimentary language and theory of mind.



Acheulian hand axe. “Bifaz en mano” by José-Manuel Benito Alvarez is licensed under CC BY-SA 2.5.

The Oldowan tradition lasted from approximately 2.5 to 1.5 mya but survived in some areas until 600 kya. Tools consisted of crude choppers (see Figure 23.6) and scrapers, as well as simple flake tools, some of which indicate that they were “retouched,” i.e. secondarily shaped and/or sharpened. In addition, there is evidence of possible wooden digging sticks or spears at the site of Koobi Fora, in the East Lake Turkana region of Kenya and possible bone tools at Olduvai Gorge.



Homo habilis Leopard Confrontation by Keenan Taylor.

Tools were likely used for acquiring and processing both animal (scavenging, butchering, disarticulation, skinning, cutting flesh, chopping bones open, etc.) and plant (digging tubers, cutting stalks, pounding to break down fiber, etc.) foods. Indications of hominins having butchered and scavenged animals comes from several lines of evidence. First, tools have been found with *H. habilis* remains. Second, there are concentrations of tools and fossilized animal bones that exhibit signs of cutting, disarticulation, and marrow extraction. Mary Leakey mapped one such area with a high accumulation of stone tools and bones, known as site DK. Third, the high frequency of particular bones at some sites is indicative of the hominins having "brought back the good stuff," i.e. skulls for brain and limb bones for meat and marrow. Fourth, microscopic analyses indicate that cut marks on some bones overlay predators' teeth marks, showing that the hominins arrived afterward. How they got meat away from scary scavengers is anyone's guess. Finally, experiments with modern-made stone tools in the Oldowan style reveal (1) that it is possible to butcher an elephant and (2) wear patterns that result from the butchering process match those found on ancient tools.

The following sites contain evidence of stone tools and their manufacture:

- Lomekwi 3, Lake Turkana region, Kenya (3.3 mya): cores, anvils, and flakes.
- Gona, Ethiopia (>2.5 mya): 3,000 stone artifacts.
- Hadar, Ethiopia (2.4 mya): tools were found with a *H. habilis* mandible.
- Olduvai Gorge, Tanzania (1.8 mya): numerous tools.
- Koobi Fora, Kenya: high concentration of flakes suggesting repeated use.



Oldowan choppers. "Pierre taillée Melka Kunture Éthiopie fond" by Didier Descouens is licensed under CC BY-SA 4.0.

H. habilis are thought to have been forager-scavengers that collected wild plant foods, hunted small animals opportunistically, and scavenged carcasses from large predators. While there is evidence of "repeated-use" sites, meaning that individuals returned to particular areas to meet, they are not thought to have settled in any one area but rather moved about the landscape in their quest for

food. They may have made use of those sites for a variety of communal or individual activities, such as grouping for “central place foraging” activities (from the animal literature, meaning to move out from and possibly return to a particular place), making new and/or using cached tools, butchering carcasses, sharing food, etc. Mary Leakey believed that her Site DK was indicative of a home base. While it is a romantic notion to look to modern hunter-gatherers with modern intelligence and advanced weaponry as being able to stay in one place until resources became scarce, it is not likely that those primitive hominins were camped out on a lake shore. It would have been a very dangerous place to be for long periods of time.

While they could have climbed trees and made sleeping nests in trees or on the ground, we do not know how much time they spent in the two microenvironments.

THE FIRST FAMILY OF PALEOANTHROPOLOGY: THE LEAKEYS

Louis Leakey (1903–1972) was born to British missionary parents residing in Kenya. He and his wife Mary made names for themselves with their pioneering work, searching for and discovering fossil hominins in East Africa. Louis is credited with the discovery of three hominin species, the first of which is considered to be a possible basal or stem ape, *Proconsul africanus* (“before Consul” [a famous chimp at the London Zoo]/“from Africa”). Louis was an early believer in an African human origin (Cartmill and Smith 2009). He became interested in the search for ancient hominins after his discovery of stone tools that he attributed to human ancestors. The Leakeys worked at Olduvai Gorge in Tanzania for many years. When Mary discovered the robust australopith that she named *Zinjanthropusboisei* (later to be changed to *Australopithecus boisei* and later to *Paranthropus boisei*), Louis proclaimed to the world that they had found his predicted “man the toolmaker.” According to legend, he was ridiculed by some because they felt that “Zinj” (also known as “Dear Boy” or “Nutcracker Man”), as the specimen came to be known, was an herbivorous ape that would not have had the mental capabilities to manufacture the tools that became known as the Oldowan technology. Louis was later rewarded with the discovery of fossils of a more derived hominin with a larger cranial capacity. He named the species *Homo habilis* (“Handy man”) as the first tool makers. There was and still is some controversy surrounding the classification of the species. He and his colleagues were accused of using cultural versus physical attributes to justify their inclusion of the fossil material in our genus *Homo*. Some still believe the species should be assigned to genus: *Australopithecus*. Regardless of the controversies, Louis made a name for himself and added to our knowledge of human ancestry. At Olduvai, he also discovered the cranium (missing its face) of a 1.2 mya *H. ergaster* individual. Another great accomplishment was sending the three “grand dames” of ape primatology into the field. He correctly believed that we can learn about ourselves from our closest relatives. He thus funded Jane Goodall to study the chimps of Gombe, Tanzania; Dian Fossey for her work with mountain gorillas in the Virunga Volcano region of Rwanda; and Biruté Galdikas to study the orangutans of Borneo.

Mary Leakey (née Mary Douglas Nikol, 1913–1996) is described on the Leakey website (www.leakey.com) as “one of the world’s most distinguished fossil hunters.” She is credited with the discovery of two species of early hominins, *Au. afarensis* at Laetoli and *P. boisei* at Olduvai, as well as the Laetoli footprints. (Laetoli is also in Tanzania.) Mary had an early interest in archaeology and, like Louis, excavated stone tools; in her case in France as a mere child. By age 17, she was auditing university courses in archaeology and geology. She met Louis in 1933 and accompanied him to Kenya to illustrate stone tools for a book he was writing. They married several years later and had three sons, Jonathan, Richard, and Philip. Jonathan hunted fossils along with his parents and discovered the first *H. habilis* specimen, a mandible known as “Jonny’s Child.” Richard moved into Kenya to work at sites around Lake Turkana, and his team discovered the oldest *H. ergaster* specimen (1.75 mya) in the West Lake Turkana region. In addition to his paleoanthropological work, he is a champion of wildlife conservation. His wife Meave is a renowned paleoanthropologist with several hominin species discoveries to her credit, and their daughter, Louise, is well on her way to making a name for herself (www.leakey.com).

This page titled 5.1.9: Homo habilis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.9: Homo habilis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.10: Homo rudolfensis

(2.4 mya)

("SAME" / FROM LAKE RUDOLF)

SITES

- Koobi Fora, Kenya; Chiwondo Beds, Malawi; possibly Olduvai Gorge, Tanzania

PEOPLE

- Richard and Meave Leakey, Bernard Ngeneo



Figure **5.1.10.1**: Homo rudolfensis holotype: KNM-ER 1470 from Koobi Fora, Kenya. "Homo rudolfensis" by Durova is licensed under CC BY-SA 4.0.

The second species of Early *Homo* to be discovered is now known as *Homo rudolfensis*, since it was discovered at the site of Koobi Fora on the east side of Lake Turkana, which was formerly known as Lake Rudolf.

PHYLOGENY

Homo rudolfensis may be a descendent of *Kenyanthropus platyops*. Those who favor that scenario would either assign *rudolfensis* to genus: *Kenyanthropus* or move *platyops* into genus: *Homo*. There are proponents on both sides of the argument as to whether *H. habilis* or *H. rudolfensis* gave rise to the earliest member of the "*erectus*grade," i.e. *Homo ergaster* in Africa, and its four descendent species: *erectus* in Asia, *antecessor* in Western Europe, *georgicus* in the Republic of Georgia, and *floresiensis* on the island of Flores.

DISCOVERY AND GEOGRAPHIC RANGE

The majority of *Homo rudolfensis* material comes from the Koobi Fora site where Richard Leakey discovered the type specimen, KNM-ER 1470 (see Figure 24.1) (Kenya National Museum – East Rudolf), as well as subsequent material. Material has also been recovered from the Chiwondo Beds of Malawi, and there is possible material at Olduvai Gorge. Thus, the species ranged from northern Kenya down through Tanzania and into Malawi, along the north-south hominin corridor.

Leakey originally called the material *Homo indet.* ("indeterminate") because he did not know in what, if any, group to include it. The use of species: *rudolfensis* began to gain in popularity as researchers came to realize they were dealing with two distinct species. For a while, some were content with the two species being geographic species, i.e. *H. habilis* at Olduvai and *H. rudolfensis* at Koobi Fora. However, *habilis*-like material was discovered at Koobi Fora (KNM-ER 1813) and some material at Olduvai resembled *H. rudolfensis*. The two species went through a tumultuous period when the species names were abandoned by some, in favor of "Early *Homo*." However, critics complained that it had become a "wastebasket" species. During that time, Richard Leakey visited SUNY Geneseo, and I had the opportunity to chat with him about the taxonomic problem. He was frustrated with the state of things and wished that paleoanthropologists would work out exactly what characterized the two species. Over time, paleoanthropologists have seemingly become more comfortable with the *habilis* designation for the majority of fossils from Olduvai Gorge, even if some of them prefer genus: *Australopithecus*. The same is true for the type specimen, KNM-ER 1470, as *H. rudolfensis*.

PHYSICAL CHARACTERISTICS

Relative to *H. habilis*, *H. rudolfensis* was larger and more robust, with a heavier face and jaws and larger dentition. Their faces were long, and the lower face was thought to be somewhat orthognathic based on the type specimen, KNM-ER 1470. New material discovered at Koobi Fora was thought to support the reconstructed facial morphology of KNM-ER 1470 and elucidate heretofore unknown aspects of the face and jaw. However, Bromage et al. (2008) have conducted architectural constraints analyses demonstrating that the degree of orthognathism would be biomechanically impossible. The molars were large with complex crowns and thick enamel like other contemporary and descendant species, but they were mesiodistally long and hence more primitive than *H. habilis*. The brain averaged 751 cc, but their encephalization quotient may have been only 3.0, versus the 3.1–3.5 range estimated for *H. habilis*. As mentioned, it is not known for certain that any of the postcranial material that some researchers assign to the species is justified. If the postcranial material belongs in the *H. rudolfensis* hypodigm, the following can be stated about the species: their innominate, femur (larger head, shorter neck, and robust shaft), and limb proportions were more modern, with longer legs and shorter arms, and the femoral robusticity corresponds to increased strain generated by a wider pelvic aperture.

Characteristics

Review of Primitive Characteristics

- Robust craniofaciodental characteristics.
- Primitive temporal region.
- Long molars (mesiodistally).

Review of Derived Characteristics

- Long orthognathic face?
- Large brain.
- Human-like limb proportions, innominate, and femora?

ENVIRONMENT AND WAY OF LIFE

While their environment did not differ from that of *H. habilis*, they may have been able to process tougher food items with their more robust masticatory apparatus. While tools are known from the time period and geographic region occupied by the species, tools have not been found in association with fossil material. It is possible that the more gracile masticatory apparatus of *H. habilis* could have resulted from tool use in food-processing activities and relaxed selection for robust features.

This page titled 5.1.10: Homo rudolfensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.10: Homo rudolfensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.11: Homo species indeterminate

Homo species indeterminate (2.8 mya)

A possible new species of Early *Homo* was recently discovered. Villmoare et al. (2015) reported on what they are calling *Homo* species indet. ("indeterminate"). The half mandible (LD 350-1) was discovered by Chalachew Seyoum in 2013 at the Ledi-Geraru site in the Afar region of Ethiopia. If indeed the jaw is part of the *Homo* lineage, it extends the origin of our genus to 2.8 mya, 400 kya older than the oldest *H. habilis*specimen, i.e. AL 666-1 from the Hadar site in Ethiopia where Lucy was discovered. Villmoare et al. (2015) believe that the specimen represents a transitional form between *Au. afarensis* and *Homo habilis*. Support for their supposition includes the combination of australopith and *Homo*-like characteristics seen in the mandible and teeth, as well as the fact that it is from the same area as *Au. afarensis* that is known to have survived there until 3 mya. They report on a fragmentary mandible from the Koobi Fora site in Kenya (KNM-ER 5431) that exhibits a similar combination of australopith and *Homo* morphology (see Villmoare et al. 2015 for references).

The anterior of the mandible exhibits the most primitive characteristics, along with some *Homo*-like characteristics. *Homo* characteristics include narrower molars with more derived dimensions and cusp morphology, and the canine/first premolar configuration is less ape- or australopith-like, as evidenced by wear patterns. They conclude that deviation in the teeth and jaws, from the australopith condition, occurred early in the *Homo* lineage.

DiMaggio et al. (2015) reported on the paleoenvironment of the site. The area was a mosaic environment consisting of open grass- and scrubland, gallery forest, and lakes and/or rivers. They see a faunal turnover 2.8–2.6 mya in the strata, in accordance with the area becoming more open and likely arid with an increase in grazers. They cite evidence for climatic and vegetation shifts resulting from "rifting processes and extensive volcanism [that] altered the architecture of sedimentary basins" (see DiMaggio et al., p. 1, for references). Their work fills an important gap in the paleoenvironmental history of the region.

The fact that *Au. garhi* is younger and their teeth are more robust leads the researchers to conclude parsimoniously that *Au. garhi* is not part of our ancestry, as earlier and later forms of *Homo* had more gracile teeth and jaws. The same would also hold true for *Au. sediba* due to their much later appearance in the fossil record at ~2 mya. However, Hawks et al. (2015) argue that the **osteometrics** (bone measurements) and anatomical features do not support firm taxonomic placement in genus: *Homo*. They believe that they have ample evidence showing that *Au. sediba* is closely related to early species of *Homo* (see Chapter 21).

This page titled 5.1.11: Homo species indeterminate is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.11: Homo species indeterminate** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.12: Homo naledi

(date unknown)

(“SAME” / “STAR” IN SOTHO LANGUAGE)




Figure 5.1.12.1: Skull and mandible of type specimen DH-1 by Lee Roger Berger research team is licensed under CC-BY-SA 4.0.

SITE

Rising Star cave system, South Africa

PEOPLE

Cavers Steven Tucker and Rick Hunter and investigated by Paul Dirks and Lee Berger and their associates

This newest member of our genus has once again confounded the evolutionary history of the *Homo* lineage. The most exciting aspect is the nature of the remains suggests that they were intentionally deposited in the deep cavern where they were discovered. *H. heidelbergensis* was heretofore the earliest species thought to have practiced intentional body disposal. Attempts at dating the remains have not been successful. However, Thackeray (2015) has estimated that the species may date to 2.0 ± 0.5 mya, based on comparisons of date and anatomical characteristics among *H. naledi*, *H. habilis*, *H. rudolfensis*, and *H. erectus* (Figure 5.1.12.1 and Chapter 27 for the *erectus* grade).




Figure 5.1.12.2: *Comparison among H. naledi, H. habilis, "African H. erectus", and H. floresiensis. By Chris Stringer, Natural History Museum, United Kingdom – Stringer, Chris (10september 2015). "The many mysteries of Homo naledi." eLife 4: e10627. DOI:10.7554/eLife.10627. PMC: 4559885. ISSN 2050-084X. Licensed under CC-BY 4.0.*

PHYLOGENY

It appears that the majority of researchers agree that the remains reflect a new hominin (see references this section, especially Randolph-Quinney 2015). Like most hominins, the phylogeny of the species is unknown but it likely descended from an australopith ancestry. What makes things even more difficult is that the species shares characteristics with possible extant or near extant species of *Homo* (*H. habilis*, *H. rudolfensis*, and *H. erectus*), more derived forms (e.g. neandertals and humans), as well as various australopiths. The mosaic of traits is interesting and further supports the bushy nature of the hominin tree.

DISCOVERY AND GEOGRAPHIC RANGE

The remains of a minimum of 15 individuals, totaling 1550 fossils (see Figure 26.4), were excavated in 2013 and 2014 from the Dinaledi Chamber, located within the Rising Star cave system in the Cradle of Humankind World Heritage Site, Gauteng Province, South Africa (Berger et al. 2015). The fossils are the largest collection of a hominin species in Africa (Dirks et al. 2015). The chamber is 30 m below ground and is only accessible via a 12 m narrow shaft (Randolph-Quinney 2015 and see Figure **5.1.12.3** – top right). Based on depositional data, the bodies were deposited over time (Dirks et al. 2015).




Figure 5.1.12.3: *Dinaledi Chamber by Paul H. G. M. Dirks, et al. is licensed under CC-BY 4.0.*

PHYSICAL CHARACTERISTICS

The remains are especially valuable as all body regions are represented, and some bones are articulated, so that anatomical positions and arrangements are preserved, e.g. an almost complete leg of a child and an adult hand (Dirks et al. 2015). The low cranial capacity, elevated shoulder joints, curved phalanges, and trunk and hip morphology are australopith-like. Crania, jaw and teeth morphology, and leg bones are, for the most part, *Homo*-like. The wrist is most similar to humans and neandertals. The foot is very human-like. (Berger et al. 2015, Harcourt-Smith et al. 2015, Kivell 2015, Thackeray 2015) Thus, we see an able terrestrial biped that could climb, forage, and take refuge in trees.



Figure **5.1.12.4**: *Homo naledi collection by Lee Roger Berger research team is licensed under CC-BY 4.0.*

Cranial capacity falls within the range for the australopiths, with males averaging 560 cc and females, 465 cc (Berger et al. 2015). The base of skull vault is flexed like members of the *erectus* grade and subsequent species of *Homo* (Figure **5.1.12.2** and 15.8). The vault bones are thin like those of *H. habilis*. *H. naledi* exhibits less postorbital constriction than the earliest australopiths, yet possesses a larger supraorbital torus than any gracile australopith (Berger et al. 2015). Taken together, it is an odd combination. A more gracile face would result in less postorbital constriction, yet the supraorbital torus is associated with chewing stress. Even more surprising, the teeth are smaller than those of *H. habilis*, *H. rudolfensis*, and the *erectus* grade, except for *H. floresiensis* (Berger et al. 2015). The skull vault is pentagonal in cross-section like Asian *H. erectus* (Figure **5.1.12.5**), due to the presence of a sagittal keel, i.e. a thickening along the midline of the skull, from front to back (Berger et al. 2015). Since the trait is poorly understood in *H. erectus*, its presence in another species may shed more light on the adaptive significance or ontological processes involved. However, unlike *H. erectus*, the keel begins in the parietal versus the frontal region (Figure **5.1.12.5**). While not well developed, *H. naledi* exhibits a canine fossa, i.e. a depression above the canine tooth, as seen in *H. antecessor*, *H. heidelbergensis*, and humans.



Figure **5.1.12.4**: *3-D scan of the fossilized skull of Homo naledi, an ancient human whose remains were discovered in a South African cave. The creature is one of more than 500 extinct species whose fossil scans are available for anyone to download at http://MorphoSource.org. Credit: Reconstruction by Peter Schmid and Ashley Kruger, University of the Witwatersrand*



Figure **5.1.12.5**: *A cast of Peking Man (H. erectus) to illustrate its sagittal keel – see ridge running across the top of the skull from front to back. By kevinzim. Licensed under CC-BY 2.0.*

While the hand of *H. naledi* (Figure **5.1.12.6**) shares characteristics with other hominins, the combination of characteristics is unique. They had long fingers and the two more proximal digit phalanges are curved even more than those of australopiths, suggestive of arboreal activities. Yet their wrist morphology is most similar to neandertals and modern humans and, along with their long, robust thumb, they were thus capable of strong manipulatory activities (Berger et al. 2015, Kivell et al. 2015).



Figure **5.1.12.6**: *Hand of H. naledi by Lee Roger Berger research team is licensed under CC-BY 4.0.*

The thorax and pelvis were flared like australopiths, but the vertebrae resemble those of the *erectus* grade and subsequent species of *Homo*.

While the combination of characteristics seen in the leg bones are distinctive, they are *Homo*-like, except that the femoral neck is long like that of australopiths. The foot (Figure **5.1.12.7**) is very human-like, with the primary differences being the curvature of their digits and less of a medial longitudinal arch (Harcourt-Smith 2015).




Figure **5.1.12.7**: *Foot of H. naledi by Lee Roger Berger research team is licensed under CC-BY 4.0.*

The seemingly advanced bipedal morphology of the two most recent hominin discoveries, i.e. *Au. sediba* and *H. naledi* suggest that strong selective forces favored the ability to move through the landscape, in search of food and other resources.

Berger et al. (2015) have calculated the weight of the sexes as follows: males at 55.8 kg and females at 39.7 kg. The only bone that they could use for calculating height was a tibia that yielded an estimate of 144.5 – 147.8 mm. They state that the species falls within the height range of modern small-statured human populations, as well as estimates for the *H. georgicus* hypodigm from Dmanisi (Lordkipanidze et al. 2007, cited in Berger et al. 2015).

ENVIRONMENT AND WAY OF LIFE

The environment of the region, at around 2 mya, has already been described in the "ENVIRONMENT AND WAY OF LIFE" sections of *Au. sediba* and *P. robustus*. Interestingly, the Dinaledi site is 800 m southwest of the *P. robustus* site of Swartkrans. It was a hominin neighborhood!


The evidence is compelling that the remains could not have been deposited via natural forces, but rather were carried at least part of the way, through a dark and narrow passage. We thus need to reassess our image of the cognitive capabilities and awareness of earlier members of our genus.

This page titled 5.1.12: Homo naledi is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.12: Homo naledi** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.13: The “erectus Grade”

The species that are collectively known as the *erectus* grade are believed to be descendants of the African “*erectus*” form, *Homo ergaster*, which in turn is thought to be descended from one of the two species of early *Homo*, *H. habilis* or *H. rudolfensis*. They lived in one form or another from 1.8 mya to as recently as 25–17 kya. *H. erectus* was first discovered by Eugène Dubois in 1891 (see Figure 27.1), at the Trinil site on the Solo River in Java. He had gone to Asia to find evidence of human ancestors, since the popular notion at the time was that we originated there.


PITHECANTHROPUS ERECTUS.

EINE

MENSCHENAEHNLICHE UEBERGANGSFORM

AUS

JAVA.

VON

EUG. DUBOIS,

MIT ZWEI TAFELN UND DREI IN DEN TEXT GEDRUCKTEN FIGUREN.

BATAVIA
LANDESDRUCKEREI
1894.

Title page from Dubois’s monograph. “Eugene Dubois, book” is in the public domain.

The first genus and species combination he assigned to the material was *Anthropopithecus erectus* (“upright man-ape”). He then changed it to *Pithecanthropus erectus* (“upright ape-man”). It was only later, after much more material had been unearthed, that Franz Weidenreich persuaded researchers that the species did not differ enough from our own to warrant different genus status. In more recent years, researchers have argued that the material from different continents should be assigned to different species. While we cannot know if the different continental assemblages represent true biological species, i.e. incapable of interbreeding, they are now minimally classified as geographic species. Since the holotype or “type specimen” came from Asia (Java), the species designation “*erectus*” was retained for Asian material. The African material has come to be known as *H. ergaster*, contemporary material from Dmanisi (in the Republic of Georgia) is known as *H. georgicus*, material from Spain as H. *antecessor*.

The strangely tiny and small-brained species from the island of Flores, *Homo floresiensis*, is thought to be a derived form of *H. erectus*. Some researchers place all of this material into the taxon *H. erectus*. Depending on which model people embrace for explaining the origin of our own species, one or more of those species would have evolved into archaic or premodern humans and, subsequently, **anatomically modern humans (AMH)**, i.e. *Homo sapiens sapiens*. The **Regional Continuity** or **Multiregional Model** supposes that whatever *erectus* forms were present in the various locations evolved through a premodern form, often termed Archaic *Homo sapiens* whether neandertal-like or otherwise, and then into AMH via gene flow between the populations. However with the more widely accepted “**Recent African Origin**” (**RAO**) theory, which holds that our ancestors arose in Africa ~200 kya and then moved out to populate the rest of the world, those “*erectus*” species that did not contribute to our lineage went extinct. The problem remains as to which of the later “*erectus*” forms gave rise to our premodern form, *Homo heidelbergensis*. At this point in time, the most plausible choices for the origin of *H. heidelbergensis*, the ancestor of both *Homo neanderthalensis* and AMH, are *Homo ergaster* or *Homo antecessor*.

This page titled 5.1.13: The “erectus Grade” is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.13: The “erectus Grade”** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.14: Homo ergaster

(1.8 mya)

("SAME" / "WORKING MAN")



Reconstruction of Homo erectus. "Homo erectus new" by Lillyundfreya is licensed under CC BY-SA 3.0.

SITES

- **Algeria:** Tighenif
- **Morocco:** Thomas Quarries and Sidi Abderrahman
- **Ethiopia:** Konso Gardula and Omo
- **Kenya:** Olorgesaillie and Nariokotome
- **Tanzania:** Olduvai Gorge
- **South Africa:** possibly Swartkrans

PEOPLE

- Richard Leakey, Kimoya Kimeu, Bernard Ngeneo

As mentioned, the African form of the "*erectus* grade" is termed *Homo ergaster*.

PHYLOGENY


H. ergaster is thought to have evolved from either *H. habilis* or *H. rudolfensis* in East Africa. However, it is possible that *H. habilis* may have been the first to leave Africa, after which it may have evolved into a pre-*ergaster*/*erectus* form that then moved into Africa and Asia. If *H. habilis* was in our ancestry, the latter scenario might explain how the more modernly proportioned *H. ergaster* appeared in the fossil record contemporary with *H. habilis* in East Africa. Of course, the hit and miss nature of the fossil record and fossil and species discoveries could also explain that phenomenon, i.e. the seeming lack of transitional forms in Africa.

DISCOVERY AND GEOGRAPHIC RANGE

The earliest *H. ergaster* material is from the East Lake Turkana site of Koobi Fora in Kenya. Richard Leakey is credited with this 1.8 mya discovery. Other sites outside of Africa are contemporary with African sites, e.g. the 1.8 mya Dmanisi site in the Republic of Georgia and the 1.8–1.6 mya site of Modjokerto in Java. (Note: There are problems with the Javanese dates because the fossil-containing layers are not conducive to more reliable dating methods.) The almost complete Nariokotome or Turkana Boy (see Figure 28.2) from the West Lake Turkana region of Kenya was discovered in 1984 by Kamoya Kimeu and dated to 1.6 mya. The

skeleton has been extremely important for reconstructing body morphology and limb proportions. The boy is thought to have been eight years old based upon tooth development patterns. He was formerly thought to be as old as 15, based on his height, stage of bone development, and hypothesized growth trajectories. However, dental calculations can accurately determine age due to the daily pattern of enamel deposition during tooth development. Scientists can count the microscopic, bead-like deposits that are laid down daily during the course of a tooth's development. Once it was determined that he was only eight years old yet 5´3˝ tall, it was apparent that *H. ergaster* developed at a much faster rate, more like a chimp than a human. Had Turkana Boy lived to adulthood, he would have been over 6´ tall. His morphology was adapted to the hot, dry conditions in equatorial East Africa, i.e. tall and long-limbed, similar to modern peoples of the region.

Other African sites include the North African sites of Tighenif (formerly Ternifine and sometimes assigned to *Homo mauritanicus*) in Algeria and Thomas Quarries and Sidi Abderrahman in Morocco; the East African sites of Konso Gardula and Omo in Ethiopia, Olorgesaillie in Kenya, and Olduvai Gorge in Tanzania; and possibly the South African site of Swartkrans, although it is not universally accepted that *H. ergaster* was there.




"Turkana Boy" by Mike Peel is licensed under CC BY-SA 4.0.

PHYSICAL CHARACTERISTICS

H. ergaster exhibited robust craniofacial characteristics relative to modern humans, but overall we see the continued reduction in dentition and masticatory apparatus. Their brains were large, with a maximum cranial capacity of 1200 cc and a range of 800–1200 cc, based on the material from the three continents. However, the relative brain size of early specimens may not have been much greater than that seen in early *Homo*. They had fairly robust supraorbital and nuchal regions. The skull vault was long and low, termed **platycephalic**, with a low maximum width. Our vaults are much higher, and we have high maximum width, due to subsequent cerebral expansion. The shape of the vault in cross-section has been described as that of a turtle shell. *H. ergaster*, like all hominins before and especially the robust australopiths, exhibits postorbital constriction between the orbits and the cranial vault. Their temporal lines were more pronounced than ours (see Figure 28.4), indicating that they had more powerful jaw muscles and chewing capabilities. They did not have much of a forehead due to several of the aforementioned craniofacial characteristics. Their nasal bones suggest a human-like, projecting nose. This would have been adaptive in arid as well as cold conditions. In addition to warming and humidifying inhaled air, the moisture in warm exhaled air condenses on the cooler nasal membranes, resulting in water conservation. The jaws were somewhat prognathic in the alveolar region, i.e. the bone that houses the teeth, and the mandibles were robust. They lacked a chin, a characteristic seen only in anatomically modern humans.

Relative to *H. habilis*, long thought to be the species from which *H. ergaster* was derived, *H. ergaster* possessed a longer skull vault, a larger brain (however, see previous paragraph), a smaller temporal fossa (the opening formed by the zygomatic arch where jaw muscles pass through and/or attach), a shorter face, a larger nose, reduced dentition and jaw robusticity, and a heavier nuchal region.

Relative to Asian *H. erectus*, *H. ergaster* possessed a higher cranial vault, and the bones of the vault were thinner. They also did not exhibit the **sagittal keel** (see *Homo erectus*, Chapter 29) typical of the Asian form. However, the Koobi Fora specimen exhibits slight keeling (see Figure 28.4).



Koobi Fora Homo ergaster. Postorbital constriction. Slight keeling Temporal line. "Homo ergaster skull replica, World Museum Liverpool" by Reptonix is licensed under CC BY 3.0.

Postcranially, *H. ergaster* was very human-like. While the thorax may still have been somewhat conical, they had more of a waist, demonstrating an uncoupling of the lower limb from the torso. The small birth canal relative to adult brain size suggests a long period of postnatal brain growth, as seen in subsequent hominins, especially our own species. They were tall, achieving heights of over 6´. They were also more robust than premodern humans that, in turn, were more robust than modern humans. Males were 20–30% larger than females. While the lifespan tended to be short, some individuals lived to be 50 to 60 years of age. There was high infant mortality, with 40% of fossil remains estimated to be less than 14 years of age.

Note

The ideal gas law is easy to remember and apply in solving problems, as long as you get the **proper values a**

Review of Primitive Characteristics

Retain:

- Initially brain possibly no larger than early *Homo*.
- Some robusticity in cranium and face.

- Supraorbital and nuchal tori.
- Postorbital constriction.
- Low forehead.
- Conical thorax.

Relative to later hominins:

- Platycephalic skull vault with low maximum breadth.
- Pronounced temporal lines.
- Alveolar prognathism.
- Large teeth but smaller than previous species.
- No chin.

Review of Derived Characteristics

- Protruding nose.
- Postcranially robust.
- Tall with long legs.
- Development of waist (decoupling between torso and lower limb).

ENVIRONMENT AND WAY OF LIFE

The environment of East Africa during the Pleistocene was hot and arid. Savannas had expanded and forests had become increasingly fragmented. There was increased volcanic activity in the Great Rift zone, and that is one hypothesis as to why at least a portion of the population moved out of the area. It is of great interest as to how and why *H. ergaster* (or a predecessor) first left East Africa. The "how" is not as difficult as the "why." While they may have crossed the Red Sea, they could also have come up through the Nile Valley and exited Africa across the Arabian land bridge that was exposed at the time. As to why they left, there are a variety of ideas that are not mutually exclusive. They may have ventured farther and farther due to (1) competition for resources with their own or other species of hominins or non-hominins; (2) local resource depletion; (3) similar environment (there is speculation that they were a grassland species that followed African grasslands into West Asian steppes, through the Indian subcontinent, and southeast to Indonesia); (4) following herd animals north and out of Africa; and/or (5) the aforementioned volcanic activity.

While there is debate as to whether the various "*erectus*" species hunted large game, it is not likely, since they lacked weaponry. It is generally accepted that they were mobile scavengers and foragers that hunted opportunistically. They certainly would have been capable of taking small prey and older and/or injured animals. Faunal remains at the Zhoukoudian, China, site consisted of deer, elephant, rhino, and beaver remains, but the bulk of the animal matter in their diet is thought to have come from smaller animals, such as rodents, hedgehogs, hares, and frogs, as well as ostrich eggs. However, Bunn and Gurtov (2014) have analyzed faunal remains at the Zinj FLK site in Olduvai Gorge, Tanzania (1.8 mya) and concluded that the high percentage (72%) of prime-aged large prey (especially bovids with weights of 250–750 lb) conforms solely to hominin hunting patterns, versus accidental deaths or carnivore kills. Thus at least some *H. ergaster* are thought to have been practicing ambush hunting and were therefore big-game hunters.



Acheulian hand axe. "Bifaz en mano" by José-Manuel Benito Álvarez is licensed under CC BY-SA 2.5.

While nomadic, they are thought to have stayed in an area for at least short periods of time, relative to past species. Early *H. ergaster* is associated with the Oldowan technology, and that is the technology that they took with them out of Africa. *H. ergaster* subsequently invented a tool tradition, termed **Acheulian**, that first appears in the archaeological record at 1.4 mya (newer data suggests possibly as early as 1.7 mya) and lasted to as late as 115 kya in some areas. The latter industry spread throughout Africa and as far east as the Indian subcontinent and west to Western Europe. It involved the use of better stone resources and tools that were more refined and standardized than in the Oldowan tradition. The most representative tool was a bifacially worked (shaped on both sides) hand axe in the shape of a teardrop (see Figure 28.5). The tool manifests what is called a "target design," in that the manufacturer knew what he or she was making and followed standardized steps in shaping the core into the resulting tool form. Once the core was prepared with the use of a stone hammer, the next step involved a "soft hammer" technique (see Figure 28.6). This required the use of a bone or antler hammer that could more carefully control the size and shape of the flakes to be removed on either side of the axe as it was shaped. Sharper and straighter edges resulted from the technique. The hominins had different tools for different purposes. The categories of tools included hand axes, smaller flake tools, choppers, picks, cleavers, and some bone tools. They would likely have been used to kill; butcher; scrape and cut hides; shape wood, bone, antler, and stone; and harvest and process plant foods.



Soft hammer percussion. "Soft Hammer" by ZenTrowel is in the public domain.

It would have been necessary for *H. erectus* to have had some sort of body covering prior to moving north into colder regions of Eurasia. There is some evidence to support *H. erectus*'s use of fire, but it is scant. However, even in later species wherein we know they had fire, such as *H. heidelbergensis* and *H. neanderthalensis*, there are fewer hearths apparent at archaeological sites than would be expected.

It was previously believed that the *erectus* grade lacked the technology to cross open water and therefore were limited to land travel. However, the discovery of a site dated to >8 kya on the island of Flores suggests that early *H. erectus* were capable of crossing fairly sizeable stretches of open sea. It is thus possible that *H. ergaster* may have rafted across the Red Sea to the Arabian

Peninsula or across the Mediterranean to Gibraltar or Italy (via Sicily) during periods of lower sea levels when much of the earth's water was tied up in ice.


While it is likely that *H. ergaster* and related species had some complex form of communication, due to their advanced cultural achievements, it is not known whether they had speech or language similar to our own. They possessed the arched palate, flexed cranial base, and cerebral centers that are necessary for speech production and comprehension. However, Turkana Boy had small intervertebral foramina (the openings in between the vertebrae where the spinal nerves exit the spinal cord) in the thoracic region. It is thus thought that they may not have had the necessary motor function for controlled breathing and speech. However as mentioned in the section on *Homo habilis* (Chapter 23), I believe the standardized nature of the Acheulian tool industry demonstrates active teaching and thus theory of mind that in turn assumes some sort of language.

This page titled 5.1.14: Homo ergaster is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.14: Homo ergaster** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.15: Homo erectus

(1.8 mya): (“SAME” / “UPRIGHT”)



Drawing of Trinil material. “Pithecanthropus-erectus” by 120 is in the public domain.

SITES

- Java: Trinil, Modjokerto, Sangiran, Ngandong
- China: Zhoukoudian, Taiwan, and sites in Yunxian, Hexian, and Lantian counties
- India: Narmada
- Turkey: Kocabas

PEOPLE

While there are many people associated with *Homo erectus*, I have listed a few of the historic names.

- Java: Eugène Dubois
- China: J. Gunnar Andersson, Davidson Black, Franz Weidenreich

Homo erectus is the genus and species combination that was retained for all mainland Asian, Taiwanese, and Javanese fossil material.

PHYLOGENY


The most popularly held notion is that *Homo erectus* is derived from *H. ergaster* or a pre-*ergaster* form that “quickly” moved out of Africa into Eastern Europe and Southeast Asia. However, *H. georgicus* is another possibility for the ancestor of *H. erectus*.

DISCOVERY AND GEOGRAPHIC RANGE

Eugène Dubois discovered the first *H. erectus* material at the Trinil site (see Figure 29.1) on the Solo River in Java in 1891. While there are problems with the dates, the oldest material from the Javanese site of Modjokerto may be “contemporary” with African and Georgian material at 1.8 mya. Other famous Javanese sites are Sangiran, Ngandong, and Trinil. Java is part of the Sunda shelf, and when initially colonized by *H. erectus*, it was connected to mainland Asia (see Figure 29.2). After reaching Java and possibly other areas of Southeast Asia, later groups of *H. erectus* moved north into China. The earliest Chinese fossils are dated to 1 mya. First assigned to the genus *Sinanthropus* (“Chinese man”), the material was later included in our own genus after Franz Weidenreich pointed to the similarities between the various assemblages of *erectus*-like fossils and other extinct and modern humans. The first fossils were discovered at the now famous site of Zhoukoudian (formerly Choukoudian), near Beijing (formerly Peking and hence the term, “Peking Man”). The local people called them “dragon bones” and were using them for medicinal purposes. Material from Zhoukoudian spans a time period of over 200,000 years, from 460 to 230 kya, with three distinct cultural periods thought to be in evidence.

One of the great mysteries of paleoanthropology surrounds the Zhoukoudian material. Weidenreich and his predecessors, Davidson Black and J. Gunnar Andersson, had amassed an unprecedented amount of fossil material from the site. Due to the imminent Japanese invasion, Weidenreich packed up the fossil material in 1941 with the intent of having it shipped to the United States. However, the material disappeared, and all that remains are Weidenreich's notes, drawings, and some casts of the original fossils.

Other Chinese sites are found in the counties of Lantian, Yunxian, and Hexian. A new discovery on the island of Taiwan has been linked to *H. erectus*, with the closest resemblance to the Hexian remains (Chang et al. 2015). Finally, the Narmada site in India has been a topic of debate for a long time but it has now been decided, at least by a portion of the paleoanthropological community, as being *Homo erectus*.



Sundaland (northwest of the Wallace Line). "Map of Sunda and Sahul" by Maximilian Dörrbecker is licensed under CC BY-SA 3.0.

PHYSICAL CHARACTERISTICS

While many of the physical characteristics of *H. erectus* are similar to *H. ergaster*, the Asian species is unique in a number of ways. Asian forms exhibit a thickening along the sagittal suture, termed a sagittal keel. The keel gives the skull a pentagonal shape in cross-section. It is unknown whether the keel served a function.



Homo erectus by Keenan Taylor.

Their incisors were shoveled, an adaptation that increases the stress resistance of teeth, especially when using them as tools. The molar enamel was characterized by a unique wrinkling pattern. Both of those dental characteristics are found in modern people of Asia and Asian ancestry and are interpreted by some scholars as evidence of regional continuity; in other words, there was a gradual evolution from *erectus*-like forms through archaic human populations and into modern populations in multiple areas via gene flow.



Homo erectus: Peking Man by Keenan Taylor.

Review of Derived Characteristics

- Sagittal keel.
- Shoveled incisors.
- Wrinkled molar enamel.

ENVIRONMENT AND WAY OF LIFE


Javanese sites in the early Pleistocene would have been conducive to tropical-adapted animals like *Homo erectus*. The area was part of the land bridge that was exposed beginning ~2.5 mya, making it accessible by land. Pleistocene Java was a mix of environments consisting of a variety of forest types, freshwater lakes and rivers, brackish marshes, and grasslands (Blain 2012).

At the time of *H. erectus* occupation, the site of Zhoukoudian, China, was in a transitional zone between temperate steppe and boreal forest. It would thus have been seasonally cold and would likely only have been habitable during the warmer months.

Culturally and technologically, Asian *H. erectus* are thought to have been somewhat similar to African *H. ergaster*. One of the key differences is the fact that the Acheulian industry never made it to Asia. The earliest inhabitants of Asia carried with them the Oldowan tool tradition, but the inventors of the Acheulian tradition apparently never followed. On maps, the Movius Line (see

Figure 29.5) demarcates the border between the two tool traditions during the Pleistocene. It has been suggested that bamboo would have been a suitable material for making tools, which could explain the paucity of stone tools found.

Populations of *H. erectus* survived in Asia for much of the Pleistocene Epoch. Recent redating of the Javanese site of Ngandong has yielded dates as recent as 53–27 kya. Even more surprising is the recent discovery of dwarfed hominins on the island of Flores, termed *H. floresiensis*, that have been dated to 18 kya. *H. floresiensis* is thought to be descended from a population of *H. erectus* that adapted to limited island resources by becoming dwarfed in size.



Movius Line: color change delineation extending from Eastern Europe to northeast India. "Biface Extension" by José-Manuel Benito Álvarez is in the public domain.

This page titled 5.1.15: Homo erectus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.15: Homo erectus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.16: Homo georgicus

(1.8 mya)

("SAME" / "GEORGIA")




Figure **5.1.16.1**: Reconstruction of Homo georgicus. "Homo georgicus" by 120 is licensed under CC BY-SA 3.0.

SITES

- Dmanisi, Georgia

PEOPLE

- Leo Gabunia, Yekva Abesalom, David Lordkipanidze

Located in the southern Caucasus region of the Republic of Georgia (Figure **5.1.16.2**), Dmanisi is the only known site for the geographic species *Homo georgicus*, of the *erectus* grade. Some treat it as a subspecies of *Homo erectus*, *H. erectus georgicus*, while others attribute it to *H. erectus*.



Figure **5.1.16.2**: Dmanisi site. "GeorgieKaart met Dmanisi" by Rasbak is in the public domain.

PHYLOGENY

H. georgicus is dated to the same time as the earliest African material at 1.8 mya. It is thus thought to be (1) closely related to or possibly a descendent species of *Homo ergaster* or (2) possibly the ancestor of *H. ergaster* and Asian *Homo erectus*.

DISCOVERY AND GEOGRAPHIC RANGE

Leo Gabunia and Vekua Abesalom discovered the site and first reported on the hominin fossil material. Beginning in 1991 and continuing to the present, David Lordkipanidze and his team have recovered teeth, multiple skulls, and numerous postcranial remains. A partial skeleton that was discovered in 2001 was thought to be more primitive than other *erectus*-grade material, and there was speculation that *H. habilis* may have preceded *H. ergaster* out of Africa, where it then evolved into an intermediate form. Since that time, researchers have come to believe it to be a more a primitive form of the *erectus* grade versus *H. habilis*.

The site has an interesting history (Figure **5.1.16.3**). The area has been settled since the Bronze Age. In the 6th century, an orthodox Christian cathedral was built on the site. By the 9th century, the region was under Arab rule. Because of its location on two

important trade routes, by medieval times it had become an important commercial center. It was conquered by the Seljuk Turks in the 11th century and later liberated by Georgian kings. The area was then attacked by the Turkomans in both the 14th and 15th centuries, from which it never fully recovered, in terms of its regional importance and economic development (Wikipedia contributors 2015b).



Figure **5.1.16.3**: *Dmanisi castle ruins with archeological site in background (under the white roof). "Ruins of Dmanisi Castle" by Larry V. Dumlao is licensed under CC BY-SA 4.0.*

PHYSICAL CHARACTERISTICS

H. georgicus exhibited a mix of primitive and *erectus*-like characteristics. They had small, robust skulls and habiline-like, prognathic faces (Figure **5.1.16.4**). Their upper limbs were australopith-like, while their spines and lower limbs were more modern. The cranial vault for one of the four adult skulls measured only 546 cc. Their molars were large, and tooth microwear suggests that they ate tough, fibrous plant foods and thus had a low-quality diet. Their canines were surprisingly long.



Figure **5.1.16.4**: *Cast of Homo georgicus. "Homo Georgicus IMG 2921" by Rama is in the public domain.*

Characteristics

Review of Primitive Characteristics

- Robust skull.
- Small brain.
- Prognathic face.
- Large molars and long canines.

Review of Derived Characteristics

- Human-like spine and lower limb.

ENVIRONMENT AND WAY OF LIFE

The environment at the time when *H. georgicus* lived in the area is thought to have been a mosaic of forest and steppes. A great many Oldowan tools and debitage (i.e. small chips of stone that are byproducts of tool manufacture) were found at the site, along with faunal remains. Due to their australopith-like upper limb morphology and the dangerous predators at the site, there is speculation that they may have slept and foraged in trees (de Lumley et al. 2005).

Of great interest is an older edentulous male. The tooth loss was antemortem because the alveolar bone had been resorbed, showing that he lived for quite a while after he began losing his teeth (de Lumley and Lordkipanidze 2006). He would likely have needed care during the intervening years, and his family or group-mates must have helped and supported him. This would be the earliest known evidence for kin selection and/or reciprocal altruism, respectively. Kin selection involves incurring a cost to oneself in order to benefit relatives, thus increasing their survival and propagating your genes because they share a portion of your genes in common. Reciprocal altruism is basically tit for tat. It explains why we will help nonrelatives. During our long hominin history prior to agriculture and sedentism, our ancestors lived in small bands. They were related to the members of their groups by blood or marriage. If Individual A incurred a cost to help a nonrelative, Individual B, there is a good chance that Individual B may have been around at a future point in time when Individual A needed help. Genes involved with those cooperative behaviors could thus have spread via natural selection. We are, and past hominins were, cooperative.



Figure **5.1.16.5**: *Homo georgicus by Keenan Taylor.*

This page titled 5.1.16: Homo georgicus is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.16: Homo georgicus** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.17: Homo antecessor

(1.2 mya)

("SAME" / "PREDECESSOR" OR "PIONEER")



Incomplete Homo antecessor skull from Gran Dolina, Spain. "Homo antecessor" by José-Manuel Benito is in the public domain.

SITES

- Gran Dolina and Sima del Elefante, Spain

PEOPLE

- Eudald Carbonell, Juan Luis Arsuaga, and J. M. Bermúdez de Castro

Until the recent discovery of hominin fossils dating to over 1.2 mya in the Atapuerca Mountains of Spain, the earliest known fossil material from the area was no more than 800 kya. The 800 kya material was assigned the taxonomic designation of *Homo antecessor*, or "Pioneer Man," as the first hominins to have ranged into Western Europe.

PHYLOGENY

Since its discovery, researchers have speculated that *Homo antecessor* was descended from a more derived form of *H. ergaster* that may have crossed the Mediterranean Sea from North Africa during a period of lowered sea levels. Now that the geological clock has been pushed back 300 kya, it is thought that they may have traveled through the Levant (i.e. the Eastern Mediterranean region). What is not known is whether the evolutionary event took place in Europe or North Africa. As far as descendant species, *Homo antecessor* is one contender for the ancestor of *Homo heidelbergensis* that in turn is thought to have given rise to neandertals and modern humans.



Homo antecessor by Keenan Taylor.

DISCOVERY AND GEOGRAPHIC RANGE

The species was discovered by Eudald Carbonell, Juan Luis Arsuaga, and J. M. Bermúdez de Castro. The first site to yield fossil material was the Gran Dolina site(see Figure 31.4) in the Atapuerca mountain range in Spain. Dated to 900 kya, the site has yielded both *H. antecessor* and *H. heidelbergensis* material. Eighty fossils from six individuals were recovered. It is of interest that much of the fossil material is immature and exhibits cut marks that may indicate that they were victims of cannibalism.



“Homo antecessor moves to Spain” by Keenan Taylor.


The earliest *H. antecessor* site is Sima del Elefante (“Pit of the Elephant”). Dated to 1.2 mya, only a handful of fossils have been recovered from the site. In addition to other sites in Spain, tools and footprints have been found in Happisburgh, England and Ceprano, Italy (also known as *Homo cepranensis*, some consider it more closely related to *H. heidelbergensis*).



Gran Dolina site. “Gran Dolina 2012 (Sierra de Atapuerca)” by Mario Modesto Mata is licensed under CC BY-SA 2.0.

PHYSICAL CHARACTERISTICS

The 800 kya *H. antecessor* material was more derived than earlier *erectus* forms, possessing a larger brain (1,000–1,150 cc) and a more modern skull and face. They had low foreheads and possessed an **occipital bun** (a bun or chignon is the hairstyle wherein one winds long hair into a doughnut shape on the top or back of the head), as did neandertals (see neandertal features in Figure 31.5). It has been suggested that the purpose of that posterior protrusion is to balance the weight of the anterior portion of the skull and face. Based upon cranial anatomy, they are thought to have been capable of detecting the same range of sounds as modern humans and were possibly right-handed. They were almost as tall as modern humans at 5.5–6′ (1.6–1.8 m), with males weighing ~200 lb (90 kg).



Neandertal cranial characteristics. "Neanderthal cranial anatomy" by Jason Potter is licensed under CC BY-SA 2.5.

Characteristics

Review of Derived Characteristics

- Increased encephalization.
- Occipital bun.
- Tall like Turkana Boy.

This page titled 5.1.17: Homo antecessor is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.17: Homo antecessor** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.18: Homo floresiensis

(74–12 kya)

("SAME" / FLORES)




Figure **5.1.18.1**: By Momotarou2012 commons.wikimedia.org/wiki/Fi...tion_model.jpg

SITES

- Liang Bua cave, Flores, Indonesia

PEOPLE

- Mike Morwood

The material assigned to the species *Homo floresiensis* comes only from the cave site of Liang Bua on the island of Flores in Indonesia Figure **5.1.18.2**). Because of its diminutive size, the new species took the world by storm when it was discovered in 2003 by Mike Morwood and his team. While tools attributed to the species have been dated to almost 100 kya, skeletal remains are dated to as young as 18 kya and as old as 95–74 kya (Brown et al. 2004).



Figure **5.1.18.2**: Liang Bua Cave. "Homo floresiensis cave" by Rosino is licensed under CC BY-SA 2.0.

Morwood et al. (1998) had previously discovered a much older site (840 kya) that they attributed to *H. erectus*, but no associated skeletal material was found. Thus the ancestor of *H. floresiensis* may have arrived much earlier.

PHYLOGENY

While there is controversy (see PHYSICAL CHARACTERISTICS below) surrounding this strange species, *H. floresiensis* is thought to have descended from a group of *H. erectus* that traveled across the sea from mainland Asia. Once there, they adapted to the island via a process known as insular or island dwarfism. Large mammalian species that become isolated on islands tend to decrease in size over time (as opposed to reptiles and small mammals that may increase in size), as smaller individuals require less food and thus have a better chance of survival and reproduction, when faced with limited space and resources and low risk of predation. (However, komodo dragons were present.) A dwarf stegodon (a relative of the Asian elephant) also inhabited the island and served as a prey species, as evidenced by juvenile bones with signs of butchering. In addition, modern "pygmy" human groups inhabit tropical rainforests where small body size is thought to possibly be adaptive for traveling through the tangled interior. Though short, they have low mass relative to surface area and relatively long limbs and extremities just like tall peoples of East Africa (e.g. the Maasai), to reduce metabolic heat and maximize cooling via sweating. The hominins may have survived until 12 kya when a volcanic eruption may have caused their extinction, as well as that of the dwarf stegodon. Since Flores was not inhabited when discovered by Portuguese traders in the 15th century, they may never have coexisted with modern humans (Wikipedia contributors 2015c, 2015d).



Figure **5.1.18.3**: Flores Island in red. "ID – Flores" by M. Minderhoud is in the public domain.

DISCOVERY AND GEOGRAPHIC RANGE

As mentioned, there is only one site on Flores where *H. floresiensis* material has been found, Liang Bua cave. Mike Morwood and his colleagues are credited with the discovery of this species that is so recent in time, the skeletal elements are not even fossilized (Brown et al. 2004)! Material from nine individuals has been recovered. The only complete skull, LB1, is from a 30-year-old adult female for which they have postcranial material as well. Nicknamed "The little lady of Flores" or "Flo," she serves as the type specimen for the species (Brown et al. 2004)



Figure **5.1.18.4**: Homo floresiensis. "Homo floresiensis" by Ryan Somma is licensed under CC BY-SA 2.0.

PHYSICAL CHARACTERISTICS

At only ~3.5′ (1.06 m) tall and 35–79 lb (16–36 kg), LB1 is very small relative to *H. erectus*, falling at the low end of *H. habilis*. Even more incredible is her brain size of 380 cc. Yet her encephalization quotient is estimated at 2.5–4.6. When compared with the brains of *H. erectus* and *H. ergaster* at 3.6–4.3 and *H. habilis* at 3.6–4.3, her brain is not as small as it first appears. Other aspects of *H. floresiensis*'s morphology are primitive relative to *Homo* species, such as its australopith-like hip and limb characteristics, especially in the shoulder and wrist. Craniofaciodental characteristics (Figure **5.1.18.4**) are consistent with *Homo* status, most closely aligning with *H. erectus*, including multiple mental foramina (i.e. small holes in the mandible through which multiple

branches of the trigeminal nerve exit the bone to innervate the face), whereas we have only one (Brown et al. 2004; Morwood et al. 2005). Postcranially, the limb bones were robust and the feet were flat and relatively large (Jungers et al. 2008).

While body size is expected to decrease via insular dwarfism, brain size does not usually keep pace and thus can be used to support a case for dwarfism. Some researchers have argued that LB1 is a modern human with microcephaly, an ontogenetic disorder that results in an exceptionally small brain and cranium and greatly limited cognitive capabilities. However, primitive skeletal characteristics, the complexity of the cultural remains, and the size of an important association area of the prefrontal cortex (Falk et al. 2005) do not support the microcephaly argument.

Characteristics

Review of Primitive Characteristics

- Australopith-like hip and upper limb morphology, especially shoulder and wrist components.

Review of Derived Characteristics

- Dwarfism.
- Brain reduction without apparent loss of cognition.

ENVIRONMENT AND WAY OF LIFE

Flores was primarily tropical forest during the time that *H. floresiensis* occupied the island. Tropical forests, while rich in biodiversity, are poor human habitats due to a low density of food species. In other words, on average you will find many different species of plants and animals per unit area but not very many of each. Compounded by the small area of the island, dwarfism and complex culture likely explain the survival of the species. They made and used tools, as evidenced by the presence of sharpened tools, prepared cores for the production of tools, debitage from their manufacture, anvils, etc., along with faunal remains from a variety of species, such as stegodon, komodo dragons, rats, and bats (Morwood et al. 2005). Their tools were small, compatible with their small body size. Burnt bones, fire-cracked rock, and a possible hearth consisting of a circle of fired rocks show that they made use of fire.

Large predators and scavengers like komodo dragons are known to “clean up” anything that dies, often without a trace. It is thus lucky that we have some remains of *H. floresiensis*.



Figure **5.1.18.5**: Homo floresiensis: Megalania encounter by Keenan Taylor.

This page titled 5.1.18: Homo floresiensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.18: Homo floresiensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.19: Homo heidelbergensis

(<600 kya)

("SAME" / HEIDELBERG, GERMANY)



Figure **5.1.19.1**: Model of Homo heidelbergensis head and shoulders. "Homo heidelbergensis adult male – head model – Smithsonian Museum of Natural History – 2012-05-17" by Tim Evanson is licensed under CC BY-SA 2.0<.

SITES

- **Europe:**
- **Spain:** Sima de los Huesos
- **France:** Arago
- **Germany:** Mauer, Steinheim, Ehringsdorf, Schöningen, Bilzingsleben
- **England:** Boxgrove, Swanscombe
- **Italy:** Ciampate del Diavolo (footprints)
- **Hungary:** Vértesszöllös
- **Greece:** Petralona, Apidima
- **Africa:**
- **Morocco:** Jebel Irhoud
- **Ethiopia:** Bodo, Omo (Kibish Beds)
- **Zambia:** Kabwe
- **Tanzania:** Lake Eyasi, Laetoli (Ngaloba Beds)
- **South Africa:** Florisbad, Elandsfontein
- **Asia:**
- **China:** Dali, Maba, Jinniushan

PEOPLE

- Otto Schoetensack and many more

For many years, fossil material from ~500–200 kya from Africa, Asia, and Europe that was more human- or *sapiens*-like was included in our own genus and species but was distinguished as "Early Archaic" *Homo sapiens* (EAHS). There was much debate as to when to draw the line between more *erectus*-like forms and more *sapiens*-like forms. The prevailing view was that material on all three continents was descended from *H. erectus*. (The various geographic species distinctions for *H. erectus* had not yet come into use.) Again, this is referred to as the **Regional Continuity Model** (**RCM**) for the origin of modern humans. In this scenario, *erectus*-like forms on each of the continents slowly evolved into modern humans via gene flow between populations. This is in contrast to the **Recent African Origin** (**RAO**) model whereby modern humans evolved in Africa and moved out to eventually replace archaic forms elsewhere (e.g., *H. erectus* in Asia).

The RAO model has gained in popularity due to a combination of the reevaluation of fossil material and especially DNA methods aimed at evaluating genetic distance between species, in terms of number of years since divergence from a common ancestor. The

material from Asia that had previously been assigned to EAHS has been relegated to *H. erectus*, with very little evidence for an intermediate form bridging the gap between *H. erectus* and AMH.

Fossil material from Europe and Africa that was formerly assigned to EAHS is now termed *Homo heidelbergensis*, with some researchers using *Homo rhodesiensis* for some of the African material. It is now well accepted that *H. heidelbergensis* was ancestral to both humans and neandertals (Figure **5.1.19.2**); note that in this scenario, *H. antecessor* is not considered to be ancestral to *H. heidelbergensis*).




Figure **5.1.19.2**: Human phylogeny. "Homo-Stammbaum, Version Stringer" by Chris Stringer is licensed under CC BY-SA 3.0 DE.

PHYLOGENY

H. heidelbergensis is thought to have evolved from a more derived form of *H. ergaster*. Some researchers believe that the material from Tighenif (formerly Ternifine), Algeria, known by some as *Homo mauritanicus*, is a possible transitional form between the two species. Mandibles from Tighenif are very similar to the type specimen, the Mauer mandible (see Figure 33.3) from the Heidelberg area of Germany, from which the species name is derived. In addition to the material from North Africa, the oldest material is from the Bodo site in Ethiopia, dated to 600 kya. Thus while an African origin is favored, some believe that *H. heidelbergensis* is descended from *H. antecessor*in Europe. Whether *H. heidelbergensis* evolved in Europe or Africa, they had to have migrated from one continent to the other. Thus if *H. heidelbergensis* evolved in Europe from *H. antecessor*, a portion of the population migrated to Africa and evolved toward AMH, leaving some regional population of the remaining European stock to evolve into *H. neanderthalensis*. The other more popular scenario has *H. heidelbergensis* evolving in Africa from some descendant form of *Homo ergaster*. A portion of the population then emigrated to Europe and evolved into neandertals while a portion of the remaining African stock evolved toward AMH.

In addition, a new species of hominin is also thought to be descended from *H. heidelbergensis*. The Denisovans, as they have come to be known due to their discovery in the Denisova Cave in the Altai Mountains of Russia, are thought to have branched off from the *H. heidelbergensis* lineage that led to neandertals. DNA analyses show that Denisovans interbred with neandertals, as well as the first wave of AMH that left Africa, possibly around 125 kya and subsequently settled Melanesia and Australia.



Figure 5.1.19.3: Type specimen: Mauer mandible. "Unterkiefer von Mauer (Replika)" by Gerbil is licensed under CC BY-SA 3.0.

Contrary new DNA evidence suggests that *H. heidelbergensis* branched off from some common ancestors prior to 800 kya and is thus not part of our recent lineage. *H. heidelbergensis* then divided into the neandertals and the Denisovan lineages, but interbreeding continued between them in areas of geographic overlap.

There is good evidence that *H. heidelbergensis* gave rise to neandertals. There are skulls from the sites of Erhingsdorf, Germany; Fontechavade, France; and Sima de los Huesos, Spain, that appear to be transitional in that they exhibit characteristics seen in neandertals, such as similar brow ridge morphology, enlarged facial sinuses, occipital bun, etc. It is rare in paleoanthropology to have such good support for continuity in physical characteristics between ancestor and descendent species. However, the fact that those species lived more recently in time and in developed and populated areas certainly helps, in terms of facilitating discoveries.



Figure 5.1.19.4: Homo heidelbergensis from Steinheim, Germany. "Homo steinheimensis, holotype" by Dr. Günter Bechly is licensed under CC BY-SA 3.0.

DISCOVERY AND GEOGRAPHIC RANGE

The earliest discoveries of *H. heidelbergensis* are from Germany. The type specimen was discovered in 1907 in Mauer, Germany. The oldest site is **Bodo**, Ethiopia (600 kya). There are numerous *H. heidelbergensis* sites in Europe (e.g. Steinheim, see Figure 33.4) that date from as early as 500 kya and range from Spain through Eastern Europe. The greatest number of individuals came from the **Sima de los Huesos** ("**Pit of Bones**") (see Figure 33.5) site in the Atapuerca Mountains of Spain. Both *H. antecessor* and *H. heidelbergensis* sites in that area were discovered when a railway was built. The *H. heidelbergensis* remains were found in a deep chamber inside of a cave, hence the name, "Pit of Bones." More than 32 individuals have been recovered, and most of them are juveniles. While there has been much speculation, the significance of this find in terms of cultural complexity (see ENVIRONMENT AND WAY OF LIFE below) remains elusive. There are a couple of sites in Asia that some researchers believe are also representative of *H. heidelbergensis*, including Dali, China. Sites in Africa (also referred to as *H. rhodesiensis*) include Jebel Irhoud in North Africa; Omo, Bodo, Ndutu, Eyasi, and Ngaloba in East Africa; Kabwe in Zambia; and Florisbad and Elandsfontein in South Africa.




Figure 5.1.19.5: Homo heidelbergensis from Sima de los Huesos, Spain. "Homo heidelbergensis-Cranium -5" by José-Manuel Benito Álvarez is licensed under CC BY-SA 2.5.

PHYSICAL CHARACTERISTICS

In general, *H. heidelbergensis* was less robust, both cranially and postcranially, than the *erectus* grade and more robust than AMH. There is intraspecific variability in craniofacial morphology, with some specimens being more robust and others more gracile. While this is not surprising, considering the temporal and geographic range for the species, it has led to disagreement as to how to define it.

H. heidelbergensis is primarily distinguished from *erectus*-like forms by its increased cranial capacity (1100–1400 cc—93% that of AMH) and more modern skull vault. Cerebral expansion, especially of the parietal lobes, led to increased cranial breadth in the superior aspect of the skull vault and thus a more vertically oriented skull. The occipital region is less angular due to reduced robusticity in the nuchal musculature. Some specimens have very pronounced brow ridges (see Figure 33.7) and some have speculated that those individuals represent males of the species. They share the presence of a canine fossa with AMH. The canine fossa is a depression above the canine on either side of the maxilla. The robusticity of their jaw morphology and tooth wear suggests that they were using their mouths and teeth as tools, for example, pulling meat off bones and working hides. Surprisingly, the middle ear bones were discovered for one specimen, and it could thus be determined that their hearing was comparable with our own (Martínez et al. 2004).



Figure 5.1.19.6: Homo heidelbergensis by Keenan Taylor.

Some *H. heidelbergensis* are considered to be transitional and exhibited characteristics seen in the subsequent neandertals. Their midfacial region was (1) prognathic with a corresponding large nose and (2) puffy due to large frontal and maxillary sinuses. Their zygomatics have a "swept back" appearance, and the posterior skull vault exhibited an occipital bun. The outer layer of their long bones (the cortical bone) was thick, and the interior medullary (marrow) cavity was narrow. The species was taller than neandertals, with males and females averaging 5′ 9″ (175 cm) and 5′ 2″ (157 cm), respectively. Average weights for males and females were 136 lb (62 kg) and 112 lb (51 kg), respectively (Smithsonian Institution 2015).




Figure 5.1.19.7: “Broken Hill Man” from Kabwe, Zambia (formerly “Rhodesian Man”). “Broken Hill Skull (Replica01)” by Gerbil is licensed under CC BY-SA 3.0.

Characteristics

Review of Derived Characteristics

- Increase in brain size & complexity of skull architecture, and corresponding vault changes.
- Loss of angulation in occipital region.
- Canine fossa.

Review of Characteristics Inherited by Neandertals

- Midfacial prognathism and large nose.
- Puffy appearance to face due to large frontal and maxillary sinuses.
- “Swept back” zygomatics.
- Occipital bun.
- Thick cortical bone and narrow medullary cavity.

ENVIRONMENT AND WAY OF LIFE

The Bodo site was of course, close to the equator and hence warm. *H. heidelbergensis*moved into Europe during the Holstein interglacial period (400–300 kya) (Wikipedia contributors 2015e).




Figure 5.1.19.8: Homo heidelbergensis hunting by Keenan Taylor.

Like those species that preceded them, *H. heidelbergensis* were mobile foragers. They left evidence for both seasonal and differential use camps. In addition to using rock shelters and caves for shelter, they are the first species for which we have evidence of building free-standing structures. At the site of **Terra Amata** in the south of France, the living floors of free-standing structures have been excavated. It is thought that a group returned to the site annually for fishing and other subsistence activities and

reconstructed their huts (up to 11 times) on the exact same site (see Figure 33.9). They are credited as having been the first to make the tools necessary for efficient fishing. Like past species, their big-game hunting capabilities are questionable. However, there is evidence that they may have ambushed large animals by forcing them off cliffs or cornering them in dead-end canyons. Support for ambush comes from faunal assemblages on the Channel Islands off the coast of France. The remains are from animals in prime condition, and the frequency of the various bones shows that the hominins were differentially removing the limbs and bringing them back to butcher at a home base. Javelin-type spears and thousands of faunal remains have been recovered at Schöningen, Germany. The spears were fire-hardened for strength and expertly weighted for flight, like modern javelins.



Figure **5.1.19.9**: Hut design at Terra Amata. "Terra-Amata-Hut" by Locutus Borg is in the public domain.

While some tools from *H. heidelbergensis* are Oldowan-like, most are of the Acheulian tradition. The species is credited with inventing a more conservative method, termed the **Levallois technique**, for controlling flake shape and maximizing their yield from a core. Flakes could then be worked into a variety of tools. They could also shape the core in such a way that a point could be struck off that was sharp on all sides (see Figure 33.10). *H. heidelbergensis* were the first to make compound tools, i.e. tools with more than one component, such as hammers and stone-tipped spears. There is a great website demonstrating and describing tool production at http://donsmaps.com/makingflinttools.html, as well as an animated visual for the Levallois technique at http://en.Wikipedia.org/wiki/Levallois_technique.




Figure **5.1.19.10**: Levallois point. "Levallois point" by José-Manuel Benito Álvarez is licensed under CC BY-SA 2.5.

H. heidelbergensis is the first species for which there is ample evidence of the controlled use of fire, in that hearths have been found at several sites. In addition to the aforementioned inventions, a couple of novel cultural practices have been suggested for the species. They may have made and used furniture, such as seaweed beds and stone blocks, and there is some evidence of art or written communication in the form of arcs and angles and the use of ocher (mineral pigments). A fine pink quartz hand axe, nicknamed "Excalibur," was found among the bodies in the "Pit of Bones." Some researchers believe it is the earliest evidence of ritual associated with burial, in that the artifact was seemingly unused and manufactured from exotic stone. The seclusion of the bodies may also represent an attempt at keeping them from being ravaged by scavengers.



Figure 5.1.19.11: Hand axe from Boxgrove, England. "Boxgrove handaxe" by Midnightblueowl is licensed under CC BY-SA 3.0.

All of these advancements and innovations are unequivocal support for the increase in cognition that resulted from the degree of encephalization and changes in brain architecture that are evident in the skull size and shape of *H. heidelbergensis*. Finally, if the new DNA evidence is correct and *H. heidelbergensis* branched off from our ancestry versus being our ancestor, the behavioral and cultural complexity apparent at *H. heidelbergensis* sites indicates that our common ancestor was cognitively advanced more than 800 kya!

This page titled 5.1.19: Homo heidelbergensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.19: Homo heidelbergensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.20: The Denisovans

THE DENISOVANS

A NEW SPECIES OF HOMININ

SITE

- Denisova and Okladnikov Caves in the Altai Mountains of Russia

PEOPLE

- Michael Shunkov and Anatoly Derevianko

In 2008, Russian scientists Michael Shunkov (paleontologist) and Anatoly Derevianko (archaeologist) discovered a terminal finger phalanx from a young girl, dubbed "X-woman," in the Denisova Cave in the Altai Mountains of Russia (see Figures 34.1 and 34.2). The Denisovans, as they have come to be called, inhabited the cave by 50 kya.



Denisova Cave (approximate location indicated by arrow). "Russia edcp relief location map" by Uwe Dedering is licensed under CC BY-SA 3.0.

PHYLOGENY

The phalanx was sequenced by Svante Pääbo's lab at the Max Planck Institute, where it was determined to be from a new form of extinct hominin. Its ancestor is thought to have split from our own lineage by >800 kya, subsequently splitting into the Denisovan and neandertal lineages ~640 mya (Callaway 2013). The two resulting lineages remained as genomically alike as two geographically distant modern human populations. Pääbo (2014) uses the example of Finns and the San people of South Africa. It is estimated that gene flow from neandertals to Denisovans was fairly low (≥0.5%) and seemingly occurred only locally in the Altai region (Prüfer et al. 2013). What is even more interesting from our perspective is that Denisovans seem to have interbred with the first wave of AMH as they passed through southern Asia after leaving Africa. These humans already carried neandertal genes from having interbred with them. Thus modern human populations that have descended from those early humans (i.e. indigenous Melanesians, Polynesians, Australians, and some Filipinos) carry 4.8% Denisovan genes, along with the mean of 2.5% neandertal genes that all Eurasians possess, meaning that a total of ~7% of their genes are derived from extinct hominins! Genes for dark skin, hair, and eyes were present in the Denisovan genome and are present in modern Melanesians (Marshall 2013). This is fascinating from two perspectives. First, it is interesting that those ancestral characteristics survived in a modern population. Second, we now know something about what the Denisovans likely looked like. Wow!

The Denisovan-like genes that the rest of Eurasians possess may have been inherited from neandertals, due to their close genetic relationship with the Denisovans. It is of great interest that the genetic variability in one of our important immunological systems, the human leucocyte antigen (HLA) system, is probably due to interbreeding with neandertals. Half of the HLA variant genes, termed alleles, seen in Eurasian populations are derived from those two extinct species (Abi-Rached et al. 2011). What is fascinating is that those alleles likely conferred a selective advantage on some of our ancestors, and they thus survived in their descendants. It is likely that neandertals and Denisovans inherited the allele(s) from *H. heidelbergensis*, lending further support for the latter species having diverged from our lineage prior to the mutation(s) that led to the forms present in the former two species.

Finally, a variant of the EPAS1 gene in Tibetans has also been traced to the Denisovans. The allele is an adaptation to the hypoxic (i.e. low oxygen) conditions of high altitude. The allele affords those individuals with better oxygen metabolism capabilities (Huerta-Sanchez et al. 2014).

Mitochondrial DNA (mtDNA) from a 400 kya femur from the Sima de los Huesos site in Spain has been found to most closely resemble Denisovans (Callaway 2013). The site contains *Homo heidelbergensis* material, and if the leg bone is from an individual of that species, it will further muddy the phylogenetic waters. However, it must be remembered that mtDNA gives only maternal lineage information, and it is thus possible that the individual was not more closely related to Denisovans but that the mtDNA alleles were conserved in his or her lineage.




Denisova Cave. "Turist den-peschera" by ЧуваевНиколай at ru.Wikipedia is licensed under CC BY-SA 3.0.

DISCOVERY AND GEOGRAPHIC RANGE

In addition to Denisova Cave, Shunkov and Derevianko also found bone fragments in the nearby Okladnikov Cave. They have since recovered two teeth and a toe bone.

The caves were also occupied by neandertals and humans at various times over the millennia. Prior to the discovery of neandertals in the area, it was thought that they ranged no farther east than Uzbekistan. This new discovery extends their geographic range by 2,000 km. The Denisovans also must have had a large geographic range, since they are thought to have bred with AMH as they passed through southern Asia (see Figure 34.3). It is amazing that for over 100 years, the only extinct hominins that we knew of in the northern hemisphere were the *erectus*-forms and neandertals and now, in just the last couple of years, we have two more, the Denisovans and the Red Deer Cave hominins. (Pääbo 2014, unless otherwise indicated.)



Possible routes taken by Late Pleistocene hominins. "Spread and evolution of Denisovans" by John D. Croft at English Wikipedia is licensed under CC BY-SA 3.0.

PHYSICAL CHARACTERISTICS

While there is not much to go on, the finger bone was broad and robust relative to modern humans and the molars were large with widely diverging roots, unlike the little or no space between neandertal molar roots. The robusticity of the phalanges suggests that the Denisovans may have been morphologically similar to neandertals (Reich et al. 2010).

ENVIRONMENT AND WAY OF LIFE

The environment of the area must have been at least seasonally cold. Mountains surrounding the area were covered with glacial ice during the Late Pleistocene (Lehmkuhl et al. 2011). However, it is possible that game may have been available year-round in the lower elevations. If large game migrated seasonally, then the hominins would have followed and thus would have only resided in the area when animals were on their northern grazing and birthing grounds.

Since *H. heidelbergensis* was cognitively and culturally advanced, the Denisovans were descended from them, and Denisovans interbred with neandertals in the Altai region at some point(s) in time, it is likely that they all had similar behavior and culture. While complex tools and an ivory bracelet were found, it is not known whether they are attributable to the Denisovans. They are more likely AMH artifacts, as they are usually only found at AMH sites.

This page titled 5.1.20: The Denisovans is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.20: The Denisovans** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.21: Homo neanderthalensis

(>300 kya)

("SAME" / NEANDER VALLEY, GERMANY)



Reconstruction of Neandertal at Neanderthal Museum. "Neandertaler reconst" by Stefan Scheer is licensed under CC BY-SA 3.0.

SITES

See "Neandertal sites" map, Figure 35.2

PEOPLE

Too many to mention (see text for some prominent researchers)

One of the best known and most enigmatic of the archaic hominins were the neandertals, *Homo neanderthalensis*. Although *Homo neanderthalensis* was originally included in our own genus and species but distinguished by subspecies status, i.e. *Homo sapiens neanderthalensis*, increasing evidence from DNA analysis suggests that the two lineages split sometime prior to 300 kya and, if new DNA evidence is correct, possibly prior to 800 kya. However, DNA evidence shows that they interbred, possibly as AMH migrated out of Africa one or more times or cohabited with neandertals in the Middle East. Eurasians and Australasians carry, on average, 2.5% neandertal genes. Thus while the RAO model for the origin of AMH is still favored and neandertals are considered a separate species, at least some populations were capable of interbreeding and thus were not true biological species at that point in time and geographic space.

Pronounciation

The material that became the holotype for the species was discovered in the Neander Valley near Dusseldorf, Germany. The German word for valley is "thal," and the "h" is silent. The "h" has been dropped for the common name in some sources. I can only imagine that someone got tired of people pronouncing the "th" and decided to head up a campaign to put a stop to it!

PHYLOGENY

There are two possible scenarios for the origin of neandertals and AMH. The first is based on fossil evidence and the second on DNA. The interpretation of the fossil record suggests that both species are derived from *H. heidelbergensis*, which in turn likely evolved from a derived form of *H. ergaster* (possibly *H. mauritanicus*) in Africa. At some point prior to 500 kya, *H. heidelbergensis* split into the AMH and neandertal lineages.

The latest genetic evidence supports a split between the AMH and neandertal lineages prior to 800 kya in Africa. The branch leading to neandertals and Denisovans is then considered *H. heidelbergensis*. The Denisovans and neandertals then split ~640 kya. However, localized groups continued to interbreed.

It is refreshing to learn that populations of hominins have been interbreeding and maintaining or forming genetic relationships since the beginning of "our" time. We modern humans are much more closely related to one another than were those ancient hominin

"species" and yet some of us do not see ourselves in others due to physical differences that mean no more than that we went different ways at different times and adapted to different environments.

Regardless of the neandertal/human/Denisovan phylogeny, a group of *H. heidelbergensis* moved into Western Europe, where a localized group then evolved into the neandertal lineage <300 kya. Transitional forms can be seen in several locales in Western Europe, especially Spain, France, and Germany.

As Pleistocene Europe became colder, neandertals adapted to the harsher conditions. The neandertals from Western Europe, with their stunted and cold-adapted bodies, are known as the "Classic" neandertals, as distinct from those to the east and southeast that retained a more gracile morphology. Dates for the Classic neandertals range from 75 to <30 kya. Figure 35.2 shows neandertal sites in Eurasia.

DISCOVERY AND GEOGRAPHIC RANGE

The earliest recognized discoveries were in Belgium and Gibraltar. The next discovery was the Neander Valley remains, which lent the name to the species. Fossil sites are ubiquitous in Western Europe, with the majority located in well-watered river valleys of France. More than 200 sites fall within a 20-mile radius of Les Ezies, France. There are also sites in Germany, Belgium, Spain, Portugal, and Italy. Some of the more famous sites are La Chapelle-aux-Saints, La Ferrassie, and St. Cesaire in France; the aforementioned Neander Valley in Germany; and Zafarraya Cave in Spain. The Chapelle-aux-Saints site has played a key role in the development of the myth of the neandertals as hulking, barbaric cavemen.



Figure **5.1.21.1**: Neandertal sites. "Carte Neandertaliens" by 120 is licensed under CC BY-SA 3.0.

The remains of an approximately 40-year-old male (see Figure 35.3) were excavated in 1908 and analyzed by Marcellin Boule, who characterized the individual as primitive, brutish, and hunched over. Researchers later realized that the adult was afflicted with arthritis, which accounted for his posture. While we cannot know how neandertals behaved relative to ourselves, they achieved a theretofore unprecedented level of cultural and technological complexity. The derogatory characterization stuck for many years until researchers realized just how much those ancient "peoples" had accomplished, such as intentional burial of their dead.



Figure 5.1.21.2: “Old Man” of La Chapelle-aux-Saints. “Homo sapiens neanderthalensis” by Luna04 is licensed under CC BY-SA 3.0.

From their supposed Western European origin, they spread east into the Middle East and as far east as Uzbekistan and northeast to Russia, in the area of the Denisovans. Some researchers do not accept that the nine-year-old boy at the site of Teshik Tash, Uzbekistan is neandertal, but rather they argue that he is AMH.

Non-classic neandertal sites are found in Croatia, the Czech Republic, Hungary, Syria, the Republic of Georgia, Russia, the Ukraine, Iraq, Uzbekistan, and Israel. Famous sites include Krapina and Vindija in Yugoslavia; the cave sites of Kebara, Amud, and Tabun in Israel; Shanidar in Iraq; and the aforementioned Teshik Tash in Uzbekistan.

The Israeli sites have been of interest for decades because they are seemingly contemporary with nearby AMH sites. There has been much speculation as to the nature of interactions between the two species. One theory is that when ice sheets blanketed much of Europe, the neandertals moved down into the Middle East along with other animals. The fact that AMH reached the Middle East by 120 kya but never entered Europe until after 40 kya suggested to some that the neandertals “held” Europe, preventing the encroachment of AMH. When moving down into the Middle East, neandertals may have pushed resident AMH out of the area. During subsequent warmer periods, AMH may have moved back into the area following the retreat of the neandertals to more northerly destinations. This idea of trading places has now been superseded by the idea of contemporaneity and interbreeding, at least by some groups at some point(s) in time.

Figure 35.2 illustrates the broad geographic range of the neandertals. It is likely that during glacial advances, populations moved south so that those in Western Europe were closer to the Mediterranean Sea and eastern neandertals may have pushed down into Israel and other warm areas, along with other animals. The fossil record indicates that animal herds moved up and down in latitude in accordance with climatic pulses, so it is very likely that hominin populations did as well. They were smart and may have inherited past cultural knowledge if they had language and theory of mind and, if for no other reason, they needed to eat and would have followed the game.

By the time that AMH moved into Western Europe ~35 kya, the neandertals had begun to die out. They likely succumbed to the increasingly harsh climate. They also went through an evolutionary bottleneck at some point and lost some of their genetic diversity, possibly leaving them more vulnerable to disease. As in the Middle East, there has been much speculation about what went on when AMH arrived in Western Europe. While they likely carried neandertal genes (unless those western AMH left no modern descendants), they themselves may not have mated with neandertals, and certainly western populations would have appeared somewhat different than Middle Eastern neandertals. However, most eastern neandertals were gone by the time AMH passed through their former eastern geographic range, en route to Western Europe. It has been suggested that AMH outcompeted them either directly, which is known as contest competition, or indirectly, which is known as scramble competition, or possibly even killed them as they encountered them. Contest competition involves one group preventing another group from accessing resources, whereas scramble competition involves one group being better at gaining access to resources than the other. I always think of a bully defending a buffet table from others versus kids scrambling at an Easter egg hunt where some are better than others at getting to and/or finding eggs. It has also been widely accepted that the neandertals were marginalized as AMH encroached on their territory. Except for a more recent date from the Croatian site of Vindija (28 kya), the most recent dates are from the Iberian Peninsula, where it is thought they retreated and died out. Regardless of what transpired between the two species, since it appears that neandertals were on their way out, it is likely a moot point. It is rather fitting that after all of the years of thinking that humans played a role in the demise of the neandertals, it appears they made love not war (at least as far as we know)!


PHYSICAL CHARACTERISTICS

See Figure 35.4 for a full skeletal view of a neandertal. As mentioned, western and eastern neandertals diverged morphologically over time. That clinal variation, i.e. a graded change in physical characteristics over geographic space, is thought to have been the result of time, varying adaptation and exposure to chronically cold Ice Age conditions, and possibly gene flow with AMH in the Middle East. Populations in Western Europe lived at higher latitudes, and the Classic neandertals exhibited cold adaptations that conform to Bergmann's and Allen's Rules. Bergmann's Rule states that as you move away from the equator, mass increases relative to surface area in order to conserve heat, as heat loss is a function of surface area. Allen's Rule pertains to limb or extremity length, so that organisms in colder environments exhibit shorter appendages. Thus in equatorial Africa, where people have adapted over the long term to hot and dry conditions, body morphology is long and gracile versus the short, stocky morphology of Arctic peoples. In addition to their stocky bodies, short appendages, and barrel chests, neandertals had facial adaptations to the cold. Like *H. heidelbergensis*, neandertals exhibited midfacial prognathism, large noses, and puffy faces due to enlarged sinuses.




Neandertal skeleton. "Neanderthalensis" by Claire Houck is licensed under CC BY-SA 2.0.

Due to their forward-oriented maxilla, the mandible also moved forward, leaving a space behind the third molar, termed a retromolar space. The internal nasal projections were large, thus increasing the internal surface area even further for warming and humidifying inspired air. In addition, unique projections extended from their internal nasal region, up into their orbits. Additional skull characteristics seen in both cold-adapted and eastern neandertals were large, smoothly rolled brow ridges above large, round, widely spaced orbits; "swept back" zygomatics; some unique inner ear characteristics; and, in the occipital region, an occipital bun and suprainiac fossae (two small depressions located above *inion*, or the external occipital protuberance; see Figure 35.5 for general area). While their skulls were longer and lower than those of AMH (see Figure 35.6), their absolute cranial capacity exceeded even that of modern humans. In accordance with Bergmann's Rule, a larger brain, while energetically costly in terms of calories, is more conservative from a heat generation and retention perspective. While the neandertal brain was larger, the frontal and parietal lobes (involved with higher thought processes) of AMH were expanded relative to those of neandertals. This may have given AMH an advantage in Ice Age Europe.



Neandertal cranial anatomy. "Neanderthal cranial anatomy" by Jason Potter is licensed under CC BY-SA 2.5.

Postcranially, neandertals have been described as a cross between a marathon runner (in terms of their endurance) and a wrestler. They were built for chasing down and killing prey. Their upper body was heavily muscled.



Homo sapiens and neandertal skulls compared. "Sapiens Neanderthal comparison" by hairymuseummatt is licensed under CC BY-SA 2.0.

ENVIRONMENT AND WAY OF LIFE

The neandertals have traditionally been portrayed as having endured harsh climatic conditions. However, there is debate as to how much of a tolerance they had for conditions in Ice Age Europe. Tattersall (2009) presents an overview of research that suggests that during the coldest periods neandertals lived in more southerly regions and only moved into higher latitudes when temperatures were warmer. While their morphology reflects chronic exposure to cold, like traditional arctic peoples, they did not have the modern technology of those modern humans, such as better shelter, tailored clothes, weapons, and millennia of advanced cultural traditions. They thus had to biologically adapt to the cold. However, when conditions deteriorated as the last glacial maximum approached, they went extinct along with other ill-suited species in the northern latitudes.



Homo neanderthalensis by Keenan Taylor.

It has been known for many years that there were two different types of neandertal settlements, and discussions were often biased as though some inhabited prime real estate and others eked out a living on the open plains. Cave sites in southern France were often described as well-watered river valleys with plentiful caves for shelter, game, water, stone resources for tools, and so forth, and they were thought to have been inhabited for millennia. Open-air sites were apparently inhabited by more nomadic groups that lived out in the open in free-standing structures and followed herd animals. Cave sites and open-air sites are now thought to represent the seasonal shift in subsistence strategies practiced by the same peoples.

Thus while neandertal populations may have been able to live year-round in more temperate regions, the rest were likely semi-nomadic like their ancestors. During the cold of winter in Western Europe, they likely sheltered in southern caves and, in summer, ventured north in pursuit of migrating herds. In addition to caves, they used rock shelters, whereby they built outward from a rock wall or overhang. They did the same inside of caves, building a shelter within a shelter. Post molds form when a wooden post inserted in the ground decays, so that a darker circle of humus is apparent. The pattern of post molds can be used to reconstruct the shape and size of houses and walls in the archaeological record. Based on post molds, neandertals are known to have covered cave entrances, probably during colder periods. Remains of constructed structures show that they used bones, posts, and rocks and likely covered them with skins and insulated with grass. At the Moldova site in the Ukraine, a 26 x 16´ oval ring of mammoth bones was excavated. The bones were likely covered with hides, forming a hut that contained numerous hearths.

During the Pleistocene, European winters are described as having been long and cold and summers were short and cool. Because of the seasonality, plant foods would primarily have been available during warmer months. European neandertals ate a high proportion of meat, with reindeer and mammoth making up the majority of the diet, based upon faunal assemblages and isotopic analyses, respectively. However, dietary composition varied by region. Horses, bovids, and goats inhabited plains whereas at higher elevations, mountain sheep and ibex dominated. At the site of Shanidar, Iraq, faunal remains included goat, sheep, bovid, pig, tortoise, bear, deer, fox, marten, and gerbil bones. At the same site, there is evidence of plant consumption and cooking. Henry (2011) found phytoliths and starch grains in calcium deposits (calculus) on neandertal teeth. Some of the starch from grasses showed damage that is characteristic of cooking. While we know that neandertals used fire, as evidenced by hearths at their sites, and likely ate plants when they were available, it is valuable to finally have supporting evidence. Since Shanidar is south of most of Europe and thus more temperate, it is likely that the neandertals had greater access to such resources.



Restoration of Le Moustier Neanderthals by Charles R. Knight. "Le Moustier" by 1920 is in the public domain.

While debate has raged for some time over whether neandertals practiced cannibalism, fossil material, especially from the French site of Moula-Guercy, provides convincing evidence that at least some groups did eat their own. Neandertal bones at the site exhibit the same signs of processing as animal bones. Bones were disarticulated and hammered open for marrow, and exhibit cut marks from muscle removal.

It is interesting how abhorrent cannibalism is to us. We identify with the neandertals and may feel disappointed that they practiced cannibalism. While it is difficult to say why they ate one another, there is evidence of dietary stress in the form of enamel hypoplasia at some sites, such as Krapina, Croatia. Thus some groups suffered periodic food shortages that resulted in faulty enamel deposition in developing children. If people are starving and there is a dead body available, historic accounts show that they will eat it. Thus it should not be surprising if neandertals consumed the dead, versus killing for consumption. There is no evidence that they practiced cannibalism in all times and places, and thus it could have been in response to extreme conditions.

Prior to the evidence from Moula Guercy, there was great reluctance anytime someone proposed cannibalism to explain damage to remains at particular sites. I think that the stimulus for many of us to feel an affinity for the neandertals is that they intentionally buried their dead. They seem so human. Now that we know that some of our ancestors interbred with them, we may feel an even greater connection to them and will have to accept the good along with possible acts of survival.




Discoid hand axe from Lyndford Quarry (near Mundford, Norfolk, UK), 60 kya. "Lyndford Hand axe-Discoid" by José-Manuel Benito Álvarez is licensed under CC BY-SA 4.0.

Neandertal culture falls within the period termed the Middle Paleolithic, i.e. the middle portion of the Old Stone Age. The neandertal tool tradition is termed the Mousterian Industry (see Figures 35.9 and 35.10 for examples of Mousterian tools), after the Le Moustier site in France (see Figure 35.8). While there are no known neandertal remains from North Africa, it is of interest that their tools have been found there (see Figure 35.11). The Mousterian method was an improvement on the Levallois technique that provided greater control over resulting flakes. Flakes were then modified into a variety of tools, such as scrapers and points, for various functions. Some of the tools were denticulate, meaning that they were saw-toothed. Like *H. heidelbergensis*, they made compound tools by hafting stone implements onto handles and shafts.




Mousterian flint artifacts. "Pointe levellois Beuzeville MHNT PRE.2009.0.203.2" by Didier Descouens is licensed under CC BY-SA 4.0.

A later tool tradition (35 kya) from the St. Cesaire site in France is categorized as an Upper Paleolithic industry, as the tools exhibit characteristics of AMH industries. Termed the Chatelperronian tradition (see Figure 35.12), it may be evidence of direct or indirect contact between neandertals and AMH, meaning they obtained the technology via contact with AMH or found one or more tools and used their own methods to replicate them. Some scholars claim the latter versus learning AMH modes of production. Several items from the neandertal site of Arcy-sur-Cure, France, have been interpreted as jewelry, another cultural achievement attributed solely to AMH.



Distribution of Mousterian lithic sites. "Distribution géographique des sites du Moustérien" by Manon Delamaison is licensed under CC BY-SA 3.0.



Chatelperronian tools. From Manuel d'archéologie préhistorique, celtique et gallo-romaine by Joseph Déchelette (1862–1914). "Pointes de chatelperron" by 120 is in the public domain.

While *H. naledi* and H. *heidelbergensis* deposited their dead in deep caves, the neandertals were the first species known to bury their dead in individual graves. Bodies are often found in a flexed position. There is very little evidence of ritual associated with neandertal burials. It appears that they dug a hole, folded the body into the hole, hence the flexed position, and possibly threw some other things in with it. Items are often interpreted as having some significance, but they are usually limited to animal bones and broken tools. However, at the site of Teshik Tash, Uzbekistan, a nine-year-old boy was buried with five sets of wild goat horns that may have adorned his body. While some have suggested that he was an AMH, if he was neandertal it appears to have been a ritualized burial.

The Shanidar site (see Figure 35.13) has always been the most romantic from my perspective. It is a cave site that experienced periodic cave-ins and has yielded the remains of several interesting individuals, some of which were intentionally buried. Shanidar 1 was an adult male. While ultimately the victim of a cave-in, he survived one or more earlier traumatic events in his life. He is thought to have been partially blind due to a head injury that involved one of his eyes. He was missing the end of one of his forearms and thus the hand as well. He suffered a leg injury that resulted in a permanent limp, and some of his teeth were completely worn down. The interesting question is, how did he survive? The oft-cited response is that his group mates helped him in life. He is thus heralded as another case of pre-human altruism or at least kin selection, if the care was provided by his relatives.

Shanidar 3 may be the earliest evidence for murder. Another adult male, he was possibly stabbed, as evidenced by a cut wound to one of his ribs. While the bone showed signs of healing, it is not known whether he died from the wound and was intentionally buried or died in a cave-in. Another interesting burial is Shanidar 4. He was also an adult male that was intentionally buried, and pollen from eight wildflower species was found in the grave. While highly controversial and possibly attributable to a seed-caching rodent species, many would like to think that neandertals not only buried their dead but also placed flowers on their remains.


As I was writing the previous section, it occurred to me that males may have been differentially buried relative to females. While I do not know the answer to that question, it is interesting and food for thought.



Shanidar Cave, Iraq (see the two men in the cave entrance for scale). "Erbil governorate shanidar cave" by JosephV at the English language Wikipedia is licensed under CC BY-SA 3.0.

The debate as to whether the neandertals could speak has raged for decades. For many years, experts thought that their larynx was situated too high in their throats to have allowed for speech. Our larynx drops during the course of development. Early on it is positioned high in the throat to allow simultaneous drinking and breathing. Babies cannot talk until the larynx drops and they then begin to babble. Thus while they are not mechanically able to speak early on, they are cognitively capable of learning language. Of interest is that some parents now teach their infants sign language so that they can communicate earlier.

While some researchers still doubt the neandertals' ability to speak, many have accepted that they likely had spoken language but would have been unable to produce the full range of sounds that characterize our own speech. The discovery of a neandertal hyoid bone at the Kebara site in Israel led many to accept their ability to talk, since its morphology was similar to our own. The hyoid is an important attachment site for the ligaments and cartilages of the larynx and for some extrinsic muscles of the tongue (i.e., geniohyoid, hyoglossus). The most telling evidence in support of neandertal speech, in addition to all of my previous arguments, is the presence of the FOX P2 gene in their genome. We also possess the gene, and it plays an important role in the acquisition of language.



Muscles involved with speech. Plate 386 from Gray's Anatomy. "Gray1019" by Henry Vandyke Carter is in the public domain.

This page titled 5.1.21: Homo neanderthalensis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.21: Homo neanderthalensis** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.22: Homo sapiens

ens (>200 kya)

("SAME" / "CAPABLE OF DISCERNING")



"Scene from the Upper Quaternary Paleolithic Period." Painting by José María Velasco (1840–1912) (Mexican)."José Maria Velasco – Scene from the Quaternary upper Paleolithic Period – Google Art Project" by Google Cultural Institute is in the public domain.

SITES

- **Africa:**
- **Tanzania:** Ngaloba
- **Kenya:** Guomde
- **Ethiopia:** Omo Kibish and Herto
- **Democratic Republic of Congo:** Katanda
- **South Africa:** Border Cave and Klasies River Mouth
- **Asia (the following are earliest regional sites):**
- **Israel:** Skhul and Jebel Qafzeh
- **Mongolia**: Ordos
- **Europe (the following are famous sites):**
- **France:** Cro Magnon (the most famous site in France)
- **Germany, Italy, Spain**
- **Australia (the following are earliest regional sites):**
- Mungo Lake and Kow Swamp
- **New World (the earliest regional site):**
- **Chile:** Mesa Verde

PEOPLE

Too numerous to mention, as one or more individuals work at each site and sites are numerous, especially in Eurasia

My discussion of early AMH will cover their existence and culture prior to the Mesolithic period (~12 kya) only and will focus primarily on European sites due to the paucity of information from earlier and non-European sites.

PHYLOGENY

The origin of our species is thought to have occurred in Africa sometime prior to 200 kya, based on fossil and genetic evidence. We may be descended from an African population of *Homo heidelbergensis* (sometimes referred to as *Homo rhodesiensis*) or an earlier common ancestor of *heidelbergensis*, based on recent genetic evidence. Fossils characteristic of a transitional form, termed *Homo helmei*, are found at the South African site of **Florisbad** and dated to 260 kya. Material from the **Herto** site in Ethiopia (Middle

Awash area of the Afar Depression) is dated to 160 kya and sometimes referred to as *Homo idaltu* or *Homo sapiens idaltu*. *Homo idaltu* may be descended from the earlier South African form(s) and the most recent ancestor of AMH.

Groups of AMH made one or more exoduses out of Africa during the Late Pleistocene. The ancestors of some Southeast Asians and the earliest Australians (as well as inhabitants of surrounding islands and those that were used as "stepping stones") may have left Africa ~125 kya. There are sites dating to ~120 kya in the Middle East. A later group left prior to 50 kya and populated Eurasia and the New World, and made their way to the South Pacific as well, where they must have come into contact and interbred with the previously existing humans there.


The AMH ancestors of Eurasians interbred with neandertals, so that living descendants have inherited an average of 2.5% of neandertal genes. Some Southeast Australasians inherited both neandertal and Denisovan genes, due to interbreeding, and they carry ~7% of genes from those two species.

DISCOVERY AND GEOGRAPHIC DISTRIBUTION

Most of the oldest AMH material has been discovered in more recent times, relative to early discoveries of Asian *Homo erectus* and neandertal remains. The earliest AMH material is dated to 200–150 kya in East Africa (e.g. Omo Kibish and Herto in Ethiopia, Guomde in Kenya, and Ngaloba in Tanzania), 180 kya in central Africa (Katanda in the Democratic Republic of Congo), and 160 kya in South Africa (e.g. Klasies River Mouth). Later sites dated to 115 kya are found in Israel (Skhul and Jebel Qafzeh caves) and South Africa (Border Cave).

Due to increasing aridification during the Late Pleistocene of sub-Saharan Africa, researchers speculate that there were few areas that would have been habitable for early AMH. South Africa and other coastal areas likely served as refugia during those times.

By 125 kya, it would have been possible for AMH to have left Africa during a short interglacial period (see Figure 36.2 for routes and dates of arrival throughout the world). Middle Eastern sites dated to <120 kya preserve evidence of AMH. The Arabian land bridge likely served as a corridor to Southeast Asia, just as it had for the ancestor of *Homo erectus*. By ~50 kya, that early wave of humans made it all the way to Australia. Archaeological sites dated to over 50 kya are found in Japan and Australia (with cave art possibly as old as 75 kya), and fossil material from Australia is dated to 40 kya. Some researchers see similarities between early Australians and early Africans, supporting the idea of at least two African exoduses and multiple human migrations into the South Pacific. The first wave left Africa first and arrived by 50 kya. Ancestors of AMH that left Africa during the "second" exodus followed at a later point in time. It is not known when they arrived in the various areas of Southeast Asia or whether they made it to the same destinations as the original colonizing group, prior to modern times.




Human arrival dates: 1 = Homo sapiens, 2 = neandertals, 3 = early hominins. "Spreading homo sapiens" by Magasjukur2 is licensed under CC BY-SA 2.5.

The earliest date for AMH in Eurasia is from the **Ordos** site in Mongolia at 50 kya. AMH reached Western Europe by 35 kya. Sites are found in Germany, France, Italy, and Spain, with the best-known site being the **Cro-Magnon** site in Les Ezies, France. The Cro-Magnon site gave the name to the earliest people of Western Europe and is the location where the "**Old Man of Cro-Magnon**" (see Figure 36.3) was found in 1868 by Louis Lartet.

By 15 kya, humans had spread throughout the world. They reached the New World, either by rafting along the shoreline from Asia during extremely low sea levels that characterized the last glacial maximum (~17 kya) or by crossing the Bering Land Bridge at a later point in time. Dates in South America (~14 kya) are older than those in North America and represent the former mode of travel. Sea levels dropped by as much as 120 m during that time. Crossing **Beringea** involved traveling between two ice shields and was likely a difficult undertaking.


PHYSICAL CHARACTERISTICS

AMH are distinguished from *Homo heidelbergensis* and *rhodesiensis* by a suite of cranial characteristics and continued loss of robusticity, both cranially and postcranially. AMH skulls were more vertically oriented with thinner bones. While the cranial capacity (mean = 1450 cc) was lower relative to neandertals, the brain was architecturally different, and corresponding behavior was more complex and indicative of greater lateralization. The parietal and frontal lobes were expanded, resulting in high maximum width and breadth and a more pronounced forehead. Those areas of the cerebrum are involved with higher thought-processing skills related to association, speech, and all of the other cognitive capabilities that make us unique relative to other species, past and present. The face was shorter, the orbits were more rectangular, and the brow ridges were less pronounced**.**




"The Old Man of Cro–Magnon." "Cro-Magnon" by 120 is licensed under CC BY-SA 3.0.

The brows are termed **bipartite** in that they are characterized by a slight bulge on either side of the nose, with a break in between. While the nose may have been pronounced, the nasal opening was narrower, and the midfacial prognathism seen in earlier forms was absent. Jaw and dental robusticity became further reduced. AMH are characterized by a **chin**, or **mental eminence** (an autapomorphic, or unique, trait in AMH). While the mandible was less robust, the morphology (described as an inverted T-shape) was strong and stress resistant.



Sternocleidomastoid muscle. Plate 1194 from Gray's Anatomy. "Gray1194" by Henry Vandyke Carter is in the public domain.



Mastoid process. From Gray's Anatomy. "Processusmastoideusossistemporalis" by Engusz is in the public domain.

Because they had more gracile and hence lighter faces, they needed stronger anterior neck muscles to flex the head from an extended position, i.e. from a back-tilted to a forward-tilted position. The **sternocleidomastoid** muscle (see Figure 36.4) accomplishes that action, along with some rotation. The muscle originates on the sternum and clavicle and inserts on the **mastoid process** (bony projections posterior to ears; see Figure 36.5). The mastoid process became more pronounced to handle the additional power that was necessary (Campbell 1998).

Postcranially, AMH exhibited narrow hips, long legs, and thinner long bones than *H. heidelbergensis* or *neanderthalensis*. While they were seemingly not as cold-adapted as neandertals, they moved into northern latitudes and survived through the last glacial maximum. It is strange that the seemingly more heat-adapted humans survived and the robust neandertals did not. However, their long legs and more gracile morphology were less energetically costly and afforded them greater endurance and a longer stride and hence greater speed. In addition, cultural adaptations to the climate must have occurred or they could not have survived. They are thought to have made better clothes, shelter, and weaponry and were skilled hunters.

ENVIRONMENT AND WAY OF LIFE

Early AMH arose and lived during the latter part of the Pleistocene Epoch, which was characterized by intermittent glacial and interglacial periods. They ventured into northern latitudes by ~50 kya and stayed and survived in extreme conditions during the period prior to, during, and after the last glacial maximum. Populations that stayed in Africa and other warm regions would certainly have continued with life as usual in the absence of climatic upheaval.




"Glyptodon old drawing" by Haplochromis is in the public domain. Paleoindians hunting a glyptodonin South America. Being large and slow-moving, they were hunted to extinction within two millennia. Painting by Heinrich Harder (1858–1935).

Until the advent of agriculture and the beginning of the modern geological epoch (i.e., the **Holocene**) approximately 10 kya, humans were mobile to semi-sedentary foragers. They exploited whatever flora and fauna were native and available in the various regions they inhabited and colonized, from mastodons in the far north to wallabies in the far south of the Old World. We really do not need to discuss much about how AMH made a living because we have living and historic examples in the ethnographic record to show us how people lived and adapted, even to environmental extremes, from the cold of the Arctic to the deserts of the world. We have a record of Stone Age populations whose way of life disappeared within my lifetime, such as Australian Aborigines and Amazonian Indians. Like modern foragers, population density would have been low, and life ranged from easy to hard, depending on the availability of resources, seasons, and climatic patterns and disasters. One estimate of mortality rates has 50% of people dying before 20 years of age, few females living beyond 30, and only 12% living beyond the age of 40.

Any discussion of early AMH culture is seemingly Eurocentric, because the majority of the known archaeological record falls between 30 and 20 kya in Western Europe. We know that ~30 kya, a warming trend occurred that lasted several thousand years. As glacial ice retreated, prime grazing land opened and spread from Spain to Siberia. As large game expanded their geographical range, so did humans and other predators. Human population numbers increased as groups spread throughout the habitable

landmass. However, as the last glacial maximum approached, ice reclaimed the land and humans were once again restricted in their range and movements.



Vegetation map for last glacial maximum (zoom in for a better look). "Last glacial vegetation map" by Jrockley is in the public domain.

The vegetation map in Figure 36.7 shows what the world was like beginning about 20 kya, during the last glacial maximum. By that time, AMH were living in Africa, Eurasia, island chains southeast of Asia, and Australia and the surrounding islands. A short time later, they ventured into the New World.

We certainly know that early humans had spoken language, and it would have facilitated their survival via group memory and tradition as well as problem-solving. They were qualitatively different than the neandertals, and their modes of communication were likely more advanced. Modern languages can be traced, showing their spread and evolution over time and geographic space. Some of the symbols that have survived from Paleolithic times, such as dots, dashes, and hand-prints, may have conveyed information. The same may be said for depictions of animals, humans, and hunting on cave walls.




"Ice age fauna of northern Spain" by Mauricio Antón. "Ice age fauna of northern Spain – Mauricio Antón" by FunkMonk is licensed under CC BY 2.5.

Early AMH culture falls within the period termed the **Upper Paleolithic** (40–12 kya). Relative to prior Middle Paleolithic sites, AMH cultural achievements are much more impressive. Over time, they made great technological advances, inventing a great variety of new and useful objects and modes of production. People left Africa armed with language, religion, and cultural identity, as they are cultural universals and there is some evidence in the archaeological record that suggests religious practices and initiation rituals (see below). Greater individual expression is apparent in the wonderful representational art that has survived in cave paintings and sculptures, and body adornment in the form of clothing, jewelry, and pigmentation. Complex aspects of culture,


such as rules regarding kinship and marriage, also may have preceded the African diaspora(s). They too are cultural universals, and while they likely changed over time in response to need and ecology, they certainly did not evolve independently in all places.

According to Stringer and Andrews (2005), cultural achievements in the various regions of the Old World were as follows:

- Europe ~40 kya:
  - All aspects of Upper Paleolithic culture.
  - Only early representational art.
- South Africa ~75 kya:
  - Blombos Cave.
    - Carved and decorated ocher crayons (see Figure 36.9).
- Australia ~30 kya:
  - Rafts.
  - Cremation.
  - Art.
  - Body adornment.
  - Bone artifacts.



Blombos Cave, South Africa: engraved ocher. "Blombos Cave engrave ochre" by Chris S. Henshilwood is licensed under CC BY-SA 3.0.




Reconstruction of tent from Magdalenian times at Pincevent, France. "Pincevent tent" by José-Manual Benito is in the public domain.

AMH settlements were larger and more permanent than those of neandertals and were often located on the tops of hills. Researchers have speculated that the view of the surrounding areas may have been beneficial for monitoring herd movements. They built better shelters, some of which had sunken and stone foundations. Materials that were utilized consisted of wooden posts and branches, stone, and hides, and where no wood was available, bones, tusks, and antlers. They also made tents (see Figure 36.10). The remains of a house in Russia show that it was approximately 50 by 120′ with multiple hearths along its length. That is four times the size of my house and architecturally impressive, considering their tool kit relative to ours. Cooking was done in stone-lined hearths and ovens. They dug storage pits to preserve and protect food stores from animals. Stone lamps consisted of a flat stone with a depression into which was placed animal fat that would have provided light for a couple of hours.

At the Sungir site near Moscow, clothes from 22 kya survived in permafrost. We thus know that Paleolithic peoples made tailored clothing like our own. Earlier hominins are thought to have worn ponchos and lashed skins onto their bodies. AMH tools of the trade that survive in the archaeological record are bone needles, pins, and awls for punching holes in the skins. In addition to animal hides, they used bark cloth and woven grass for insulation.

Upper Paleolithic tool industries were advanced relative to past traditions, with greater diversification and refinement. Tools consisted of knives, scrapers, chisels, borers, awls, needles, and a greater number of blade and compound tools. Stone, bone, ivory, and antler were used. Some of the more famous Upper Paleolithic industries are the **Aurignacian** (possibly <40 kya), **Gravettian** (~28–22 kya), **Solutrean**(~22–17 kya), and **Magdalenian** (~18–10 kya).



Map showing Aurignancian culture, 47–41 kya. "Aurignacian culture map-en" by Hughcharlesparker is licensed under CC BY-SA 3.0.


The Aurignacian industry (see Figure 36.11) may have originated in Asia, but it can be found throughout Europe (Stringer and Andrews 2005). The Cro-Magnon people that produced the tools, art, and other cultural achievements from that time are thus known as the Aurignacian or Cro-Magnon culture. Aurignacian and subsequent tool industries were predominantly blade-based. Like the Levallois technique and the Mousterian industry, AMH could produce a variety of tools from a single core, but they used a new mode of production known as the **punch technique**. It involved using a hammerstone, a hammer and chisel, or a long wooden spear (using upper body weight and strength) to "punch" blades from the core. For illustrations on this technique see Dennis O'Neil's Early Modern Human Culture at anthro.palomar.edu/homo2/.

The Gravettian industry had a more limited distribution in Eastern Europe and Western Asia (Stringer and Andrews 2005). Stereotypical tools of the industry were flat-end blades and burins (see Figure 36.12) that were used to work wood.



Brassempouy burins from the Gravettian culture. "Burin Brassempouy 213.3 Profil (3)" by Didier Descouens is licensed under CC BY 3.0.

The Solutrean industry (see Figure 36.13) is most famous for its beautiful and refined laurel leaf blades (see Figure 36.14). A method known as **pressure flaking** was used to finely and bifacially shape the blade. Pressure flaking involves the use of a pointed tool, such as antler or bone, to force tiny flakes from the surface and edges of the tool (see Figure 36.15).



Solutrean sites."Homo Sapiens in Europe – solutrean distribution map-fr" by Sémhur is licensed under CC BY-SA 4.0.



Solutrean leaf blade. "Biface feuille de laurier" by Calame is in the public domain.





FIG. 22.—Flaking Stone by outward pressure. (*After Holmes.*)

Flaking stone. From The origins of invention: a study of industry among primitive peoples by Otis Tufton Mason (1895). "19th century knowledge stone tools flaking stone by outward pressure" by Jonp154 is in the public domain.


Finally, the Magdalenian industry (see Figures 36.16, 36.17, and 36.18) is characterized by great advances in weaponry, such as the bow and arrow and the atlatl or spear thrower (see Figures 36.19 and 36.20), both of which allowed hunters to put distance between themselves and their prey.




Magdalenian blades and burins. “Burins and blades – Bernifal – Meyrals – MNP” by Sémhur is licensed under CC BY-SA 4.0.



Examples of antler weapons. From The Outline of History by H. G. Wells (1920). “Wells Reindeer Age articles” by SEWilco is in the public domain.



Geographic range of Magdalenian culture (in pink). "Homo Sapiens in Europe – magdalenian distribution map-fr" by Sémhur is licensed under CC BY-SA 4.0.



Atlatl being used to throw a spear. From Manuel d'archéologie préhistorique, celtique et gallo-romaine by Joseph Déchelette (1862–1914)."Propulseur-2" by 120 is in the public domain.




Atlatl. "Atlatl" by Maxim Razin is in the public domain.

Other weapons from the Upper Paleolithic are **stone missiles** or **bolas** (see Figure 36.21), **boomerangs**, spears, javelins, and clubs. AMH had refined fishing techniques, rafts, and canoes. **Harpoons** appear very early in the archaeological record, e.g. the **Katanda** site in the Democratic Republic of Congo is dated to 180–75 kya. They also constructed **traps**, **rope**, and **baskets**.



Bola from the site of Sidi Abderrahman, Morocco. "Bolo-Sidi-Abderrahman" by 120 is licensed under CC BY-SA 3.0.

Beginning ~25 kya, a cultural and symbolic explosion is evident in the archaeological record of Western Europe, possibly in response to the increasingly cold temperatures, such as if they spent more time inside caves or were performing rituals aimed at increasing their survival. The Cro-Magnon/Aurignacian people are known for their cave art and sculptures (see Figures 36.23 and 36.24). Figure 36.22 illustrates the incredible number of sites where early AMH art has been found. There are 150 sites in southwest France alone.



Paleolithic art sites. Aqua lines indicate “limits of the main glaciations.” “Upper Paleolithic Art in Europe” by Sugaar is in the public domain.

Common cave art themes are fauna; hunting; hands; dots and lines; some humans; and the occasional human costumed as an animal and sometimes dancing, such as “The Sorcerer” (see Figures 36.23 and 36.24). Men appear alone or in groups, but women are never pictured alone. Drawings of male and female genitalia are reported from multiple sites.



Half-bull/half-human etching. Cave site in Dordogne, France. “Gabillou Sorcier” by José-Manuel Benito is in the public domain.



Drawing by Breuil of the “Sorcerer” Cave painting. Trois-Frères, Ariège, France (15 kya). “Pintura Trois Freres” by Dcasawang1 is in the public domain.

Artists used charcoal and mineral pigments, such as ocher (see Figure 36.9 for a photo of an ocher crayon from Blombos Cave), and may have applied some of the shading by chewing the pigments and spitting them onto the walls, resulting in an air-brushed effect. There is a great demonstration of that practice in the third segment of the public television series *In Search of Human Origins* (NOVA 1994).

The art may have served a variety of purposes. First, it is difficult not to imagine their aesthetic and emotional value, e.g. accompanying stories and myths. They may have been used for more utilitarian purposes, such as hunting strategies or keeping records. Some might have served a social function, such as communication within and between groups or education. Finally, of

course, there is their proposed use in magic and ritual. Some art is found deep within caves in association with primitive stone lamps and children's footprints. Some of the caves were discovered by divers, and thus the archaeological record was not disturbed, i.e. they were not modern footprints. It is easy to see why this has been interpreted as possible evidence of initiation rituals.


While they drew and painted on outer walls as well as inside caves, the cave paintings survived better. The French site of **Lascaux Cave** (18 kya) (see Figures 36.25 and 36.26) is renowned for the color and beauty of its animal depictions, especially those seen in the "Great Hall of Bulls." At the cave site of **Grotte Chauvet**, France (~30 kya) (see Figure 36.27), more than 300 animals are depicted, many of which are not seen elsewhere, such as panthers and lions. Figures 36.29 and 36.30 show maps of the Spanish cave of La Pasiega (<20 kya), giving you some idea of the labyrinthine nature of some of the cave systems and richness of the galleries. Figures and forms in the cave number over 700 and consist of the following:

> *97 deer (69 females and 28 males), 80 horses, 32 ibex, 31 cattle (17 bison and 14 aurochs), two reindeer, a carnivorous animal, a chamois, a megaloceros, a bird and a fish; also there may be a mammoth and about 40 quadrupeds not clearly identified; also the ideomorphs, such as roof-shaped and other surprisingly varied symbols (more than 130), and very often including various anthropomorphs and hundreds of marks and partly erased traces. (Wikipedia contributors 2015a)*

Another famous Spanish cave site is **Altamira** (15 kya) (see Figure 36.28). The cave is about one kilometer long (Chivers, 2004) and contains polychromatic renderings of large mammals, especially bison, and human hand prints.



Grotte Chauvet cave, France. "Lions painting, Chauvet Cave (museum replica)" by HTO is in the public domain. Altamira cave, Spain. "AltamiraBison" by Rameessos is in the public domain.



Map of Gallery Zones within La Pasiega Cave, Spain. "La Pasiega-Plano (Cantabria)" by Locutus Borg is licensed under CC BY-SA 3.0.



Map of art of Gallery A, La Pasiega Cave, Spain (also see map above). "La Pasiega-Galería A-1er Santuario" by Locutus Borg is licensed under CC BY-SA 3.0.

It is of interest that some of the most magnificent animal art is located in the most acoustically resonant areas within caves. It conjures images of our ancestors having special ceremonies or gatherings. Some musical instruments have survived in the form of bone flutes, percussion instruments, and a possible lithiphone (a stone xylophone). Their music would have been amplified, and with the addition of flickering flames, the wall images would have seemingly come alive!

AMH invented pottery, with the earliest evidence being fired-clay animals from the Czech Republic. The most famous sculptures are the Venus figurines (i.e. fertility goddesses) that have been found from Western Europe to Siberia. They are usually clay or stone depictions (also wood, bone, and ivory) of obese women with pronounced breasts and buttocks. They were originally thought to have been produced by men for fertility purposes. A more recent interpretation is that they were self-sculptures by women. That may explain why they were usually faceless and why body parts closest to the eyes were large and disproportionate compared with their tiny feet. Another idea suggests their use as obstetrical aids. Other sculptures were created via bas-relief on walls and rocks, which involves carving some dimensionality into the façade (see the Venus of Laussel, Figure 36.31), and there were also engraved tools, jewelry, and so forth.

The idea of fecund women as fertility goddesses is an example of sympathetic magic, i.e. depicting what one would like to happen. Obese women would have been better able to carry and nurse their babies so that they had better chances of survival. The same could be said for hunting scenes and game animals. Men may have been hoping for safe and successful hunting or an increase in the local game.

Other interesting sculptures came from the following sites or areas:


- Le Tuc d'Audoubert, France:
  - Footprints (see Figure 36.32).
  - Two 1 m clay bisons found propped up in the middle of a small chamber inside the cave.
- Grotte de Montespan, France:
  - Children's footprints.
  - Clay figures of a bear, a lion, and other animals propped up against the cave's walls.
- Czech Republic:
  - Fired animal figurines (27 kya).

- “Oldest human portrait” (26 kya) carved from mammoth ivory and possibly attached to a staff or baton, showing great detail in the eyes, hair, mouth, and expression.

- El Juyo, Spain (14 kya):
  - Stone figure on which one side of the face is a smiling man and the other is a feline predator, discovered in a possible ceremonial sanctuary.



Venus sculptures from EuropeLeft: **Venus of Laussel.** “Venus-de-Laussel-vue-generale-noir” by 120 is licensed under CC BY 3.0. Top right: **Venus of Willendorf.** “A female Paleolithic figurine, Venus of Willendorf Wellcome M0000440” by Wellcome Images is licensed under CC BY 4.0. Bottom right: **Venus of Brassempouy.** “Venus of Brassempouy” by Jean-Gilles Berizzi is in the public domain.



Footprints in the Cave of Tuc d’Audoubert, France. “Upper Paleolithic period, view of gallery with heel prints, Wellcome M0015047” by Wellcome Images is licensed under CC BY 4.0.

In addition to murals and sculptures, early humans also decorated their tools and bodies. Ocher is found in burials and was likely used to color the body, just as seen in many modern indigenous groups. Depending on the mineral composition, ocher is found in a variety of colors: yellow, orange, rust, brown, etc. They also made and wore jewelry and decorated their clothes with beads. At the Sungir site near Moscow (see more in section on burial practices), two children and an old man were buried in garments that were covered with thousands of ivory beads, thought to have taken an hour each to produce.




"Lion Man" Aurignacian artifact. Oldest known zoo-/anthropo-morphic figurine (40 kya) from Hohlenstein-Stadel cave, Germany. "Lion man photo" by JDuckeck is in the public domain.

The earliest intentional burials that have been discovered for AMH are from the Middle East and dated to 120–80 kya. As mentioned, they not only buried their dead but also included grave goods and decorated the bodies in ritual fashion. Some Paleolithic cultures cremated remains. Mass burials have been found at some sites. Burial practices included placing the body in a flexed position, as the neandertals did, or supine and, in a few instances, covered with a slab of rock. Two interesting burials were found at Italian and Russian sites. At the site of **Grotte des Enfants**, Italy, two youngsters were decorated with hundreds of shells and pierced animal teeth. At the famous site of **Sungir** (see Figures 36.34 and 36.35) near Moscow, three interesting burials were found. In one grave, a nine-year-old and a twelve- or thirteen-year-old were buried together. They were flanked by two mammoth bone spears. The tusks would have had to have been boiled in a pit of water, using hot rocks, in order to straighten them. Ten thousand beads were sewn to their clothes, and the bodies were decorated with hundreds of perforated fox canines (remember that each fox has only four large canines!), carved ivory animals, pins, and pendants. They were placed on a bed of ocher. A 40-year-old man was also honored in death. His clothes were also decorated with thousands of beads, and he wore ivory bracelets. He too was placed on a bed of ocher.



Paleolithic burial at Sunghir site, Russia. "Sunghir-tumba paleolítica" by José-Manuel Benito Álvarez is in the public domain.



Map of Sungir site, Russia. "SungirMap" by ASekirin is licensed under CC BY-SA 3.0.

In conclusion, I found this great timeline of the Upper Paleolithic (Wikipedia contributors 2015j):

Timeline of the Upper Paleolithic

45,000–43,000

- Earliest evidence of modern humans found in Europe, in Southern Italy.

43,000–41,000

- At Ksar Akil in Lebanon, ornaments and skeletal remains of modern humans are dated to this period.
- Denisova hominin live in the Altai Mountains, Russia.

40,000–35,000 BC

- Early cultural center in the Swabian Alps.
- Earliest figurative art (Venus of Schelklingen, Germany).
- Beginning of the Aurignacian.
- The first flutes appear in Germany.
- Lion-Human created from Hohlenstein-Stadel.

39,000 BC

- Most of the giant vertebrates and megafauna in Australia became extinct, around the time of the arrival of humans.

38,000 BC

- Examples of cave art in Spain are dated to around 38,000 BC, making them the oldest examples of art yet discovered in Europe.

38,000 BC–29,000 BC

- Wall painting with horses, rhinoceroses, and aurochs, Chauvet Cave, Vallon-Pont-d'Arc, Ardéche gorge, France, is made.

35,000 BC

- Zar, Yataghyeri, Damjili and Taghlar caves in Azerbaijan
- First evidence of people inhabiting Japan.

32,000 BC

- Europeans understand how to harden clay figures by firing them in an oven at high temperatures.

30,000 BC

- First ground stone tools appear in Japan.
- Invention of the bow and arrow.

29,000–25,000 BC

- Venus of Dolní Věstonice, Czech Republic. It is the oldest known ceramic in the world.
- The "Red Lady of Paviland" lived around 29,000–26,000 years ago. Recent evidence has come to light that he was a tribal chief. (Note: it was a male skeleton dyed with red ocher.)

23,000 BC

- Venus of Petřkovice (*Petřkovická venuše* in Czech) from Petřkovice in Ostrava, Czech Republic, was made.

22,000 BC

- Last Glacial Maximum.
- Venus of Brassempouy, Grotte du Pape, Brassempouy, Landes, France, was made.
- Venus of Willendorf, Austria, was made.

20,000 BC

- Last Glacial Maximum. Mean sea levels are believed to be 110 to 120 meters (361 to 394 ft) *lower than present*, with the direct implication that many coastal and lower riverine valley archaeological sites of interest are today under water.

18,000 BC

- Spotted Horses, Pech Merle cave, Dordogne, France are painted.

18,000 BC–11,000 BC

- Ibex-headed spear thrower, from Le Mas d'Azil, Ariège, France.

18,000 BC–12,000 BC

- Mammoth-bone village in Mezhirich, Ukraine, is inhabited.

17,000 BC

- Spotted human hands, Pech Merle cave, Dordogne, France are painted.

17,000 BC–15,000 BC

- Hall of Bulls, Lascaux caves, France is painted.
- Bird-Headed man with bison and Rhinoceros, Lascaux caves, is painted.
- Lamp with ibex design, from La Mouthe cave, Dordogne, France, is made.

16,500 BC

- Paintings in Cosquer cave, where the cave mouth is now under water at Cap Margiou, France were made.

15,000 BC

- Bison, Le Tuc d'Audoubert, Ariège, France.

15,000 BC–12,000 BC

- Paleo-Indians move across North America, then southward through Central America.
- "Pregnant woman" from Laugerie-Basse, France was made.

14,000 BC

- Paleo-Indians searched for big game near what is now the Hovenweep National Monument in Colorado and Utah.
- Bison, on the ceiling of a cave at Altamira, Spain, is painted.
- Domestication of reindeer.

13,000 BC

- Beginning of the Holocene extinction.
- Earliest evidence of warfare (found in the Americas).

11,500 BC–10,000 BC

- Wooden buildings in South America (Chile), first pottery vessels (Japan).

11,000 BC

- First evidence of human settlement in Argentina.
- The "Arlington Springs Man" dies on the island of Santa Rosa, off the coast of California.
- Human remains deposited in caves which are now located off the coast of Yucatán.

10,500 BC

- Stone Age Creswellian culture settlement on Hengistbury Head, England, dates from around this year.

….And the rest, as they say, is "history"!

This page titled 5.1.22: Homo sapiens is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Barbara Welker via source content that was edited to the style and standards of the LibreTexts platform.

- **4.22: Homo sapiens** by Barbara Welker is licensed CC BY-NC-SA 4.0. Original source: https://milnepublishing.geneseo.edu/the-history-of-our-tribe-hominini.

5.1.23: Recent Hominid Cranium Comparison Checklist

Name: _____

Section: _____

Student ID#:_____

Martin Nickels

© 1999 ENSI (Evolution & the Nature of Science Institutes) www.indiana.edu/~ensiweb

This material may be copied only for noncommercial classroom teaching purposes, and only if this source is clearly cited.

PLEASE READ THESE 5 STATEMENTS BEFORE BEGINNING THIS ACTIVITY:

1. Work in groups of 3-4 students so that everyone can be involved in the activity.

2. BE SURE (!) TO TAKE TURNS doing different measurements and observations.

3. When taking a measurement, use the SLIDING CALIPERS (except for #11 & #12 which may require the HINGE calipers) and remember to...

4. ALWAYS MEASURE IN MILLIMETERS [mm] and round off to whole numbers.

5. PLEASE DO NOT ADD ANY PENCIL OR PEN MARK "TATTOOS" TO THESE CRANIA, OR STICK YOUR FINGERS IN THEIR EYE ORBITS OR NOSES!

I. BRAINCASE: (7 items #1-7)

1. Does the FOREHEAD (frontal bone) look more vertical OR flatter when the skull is held in normal anatomical position [NAP] (i.e., with the eyes oriented forward)?

2. Is a SUPRAORBITAL BROWRIDGE present?

3. If present, is the BROWRIDGE DIVIDED in the middle, or CONTINUOUS?

4. What is the SHAPE OF THE BRAINCASE (front to back) when viewed from above?

5. Is a SAGITTAL CREST present?

6. In NAP, is the FORAMEN MAGNUM oriented more downward OR more to the rear?

7. Is the MASTOID process relatively flat OR does it noticeably protrude (project)?----------

II. FACE: (5 items: #8-12)

8. Are the NASAL BONES raised (arched) OR flat?

9. Measure the MAXIMUM BREADTH (width) of the NASAL OPENING [mm].

10. Measure the MAXIMUM HEIGHT of the NASAL OPENING [mm].

11. Measure the LENGTH of the MAXILLA (the upper jaw) [mm]. (Measure down the middle of 33 the palate from the front edge of the foramen magnum to either between or just in front of the two central incisors to determine how much the face projects forward.)

12. Measure the BIZYGOMATIC BREADTH using the hinge caliper if necessary [mm]. (This is the width or breadth of the face from the widest part of one zygomatic arch to the widest part of the other zygomatic arch.)

III. DENTITION: (6 items #13-18)

13. SHAPE OF THE DENTAL ARCADE: Do the tooth rows diverge towards the back OR are they more straight-sided and parallel to one another?

14. When viewed from the side, are the INCISORS angled out OR are they vertical?

15. Measure the COMBINED WIDTH or BREADTH of the 4 INCISORS together.

16. Does the CANINE tooth project above the chewing surfaces of the other teeth?

17. Is a CANINE DIASTEMA present?

18. Measure the COMBINED LENGTH of the LEFT 2 PREMOLARS and 3 MOLARS together by measuring from the back of the last molar to the front of the first premolar to determine the length of the chewing surface of the "cheek teeth". [mm]. (NOTE: Measure the right side if the left side is missing any of these 5 teeth.)

Source: www.indiana.edu/~ensiweb/lessons/hom.cran.html

	Skull #1	Skull #2	Skull #3	Skull#4
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				

19) What trends do you notice? What are the major differences between these skulls?

20) How wide is the variety in Modern Skulls? Compare this to the historical differences you saw in weeks 8 and 9. Can bones be used to define membership in a past species? Why or why not?

This page titled 5.1.23: Recent Hominid Cranium Comparison Checklist is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Alex A. G. Taub via source content that was edited to the style and standards of the LibreTexts platform.

- **1.10: Recent Hominid Cranium Comparison Checklist** by Alex A. G. Taub is licensed CC BY-NC-SA 3.0. Original source: https://socialsci.libretexts.org/api/deki/files/94257/biological-anthropology-laboratory-activities.pdf.

6: Race and Human Variation

Michael B. C. Rivera, Ph.D., University of Cambridge

Learning Objectives

- Review the illustrious and (at times) troubling history of “race” concepts.
- Recognize human diversity and evolution as the thematic roots of our discipline.
- Critique earlier “race” concepts based on overall human diversity being lower compared to other species and human genetic variation being greater within a population than between populations.
- Explain how biological variation in humans is distributed clinally and in accordance with both isolation-by-distance and Out-of-Africa models.
- Identify phenotypic traits that reflect selective and neutral evolution.
- Relate a more nuanced view of human variation with today’s ongoing bioanthropological research, implications for biomedical studies, applications in forensic anthropology, and sociopolitical/economic concerns.

Humans exhibit biological diversity. Cognitively, humans also have a natural desire to categorize objects and other humans in order to make sense of the world around them. Since the birth of the discipline of **biological anthropology**, we have been interested in studying how humans vary biologically and what the sources of this variation are. Before we tackle these big problems, this first begs the question: Why *should* we study human diversity?

There are certainly academic reasons for studying **human diversity**. First, it is highly interesting and important to consider the evolution of our species and how our biological variation may be similar to (or different from) that of other species of animals (e.g., other primates and apes). Such investigation can give us clues as to how unique we are as a biological organism in relation to the rest of the animal kingdom. Second, anthropologists study modern human diversity to understand how different biological traits developed over evolutionary time. If we are able to grasp the evolutionary processes that produce and affect diversity, we can make more accurate inferences about evolution and adaptation among our hominin ancestors, complementing our study of fossil evidence and the archaeological record. Third, as will be discussed in more detail later on, it is important to consider that biological variation among humans has biomedical, forensic, and sociopolitical implications. For these reasons, the study of human variation and evolution has formed the basis of anthropological inquiry for centuries and continues to be a major source of intrigue and inspiration for scientific research conducted today.

An even more important role of the biological anthropologist is to improve *public* understanding of human evolution and diversity, outside of academic circles. Terms such as **race** and **ethnicity** are used in everyday conversations and in formal settings within and outside academia. The division of humankind into smaller, discrete categories is a regular occurrence in day-to-day life. This can be seen regularly when governments acquire census data with a heading like “geographic origin” or “ethnicity.” Furthermore, such checkboxes and drop-down lists are commonly seen as part of the identifying information required for surveys and job applications.

According to the Oxford English Dictionary (2018), *race* is a term that should be used to describe one or more of the following:

- a major division of the human species based on particular physical characteristics;
- the biological origin of a group of people, or ancestry;
- the fact or condition of belonging to a racial division or group, or the social qualities associated with this;
- a group of people sharing the same culture and language;
- any group of people or things with a common feature or features;
- a population within a species that is distinct in some way, especially a subspecies.

So many various definitions for one word already suggests that perhaps the concepts or meanings behind biological diversity are complicated. Even though the terms *race* and *ethnicity* are used often in commonplace settings, there is no consensus among biological anthropologists as to what races are, whether they even exist, and, if they do, how the term should be applied to the human species meaningfully. If biological anthropologists cannot reach a consensus on how to view human diversity, how can we possibly expect there to be a clear perspective on the nature and causes of biological variation outside of scientific academia? Ideas about ethnicity that people hold have huge social and political impacts, and notions of race have been part of the motivation behind

various forms of **racism** and **prejudice** today, as well as many wars and genocides throughout history. This is how the role of the biological anthropologist becomes crucial in the public sphere, as we may be able to debunk myths surrounding human diversity and shed light on how human variation is actually distributed worldwide for the non-anthropologists around us (Figure 13.1). Recent work in anthropological genetics has revealed the similarities amongst humans on a molecular level and the relatively few differences that exist between populations that one might be tempted to see as significantly distinctive.



Figure 13.1 Humans are biologically and culturally diverse. (Top left: Hadzabe members in Tanzania; top right: Inuit family in traditional seal and caribou clothing; bottom left: Andean man in traditional dress in Peru; bottom right: Dr. Jane Goodall.)

Science communication and education that centers upon race and our species' variation is interesting and important. Throughout this chapter, I will highlight how humans cannot actually be divided into discrete "races," because most traits instead vary on a continuous basis and human biology is, in fact, very **homogenous** compared to the greater genetic variation we observe in other closely related species. The reason we know this now is thanks to technological developments that have taken place over the last 50 or so years. Molecular anthropology, or anthropological genetics, revolutionized and continues to add new layers to our understanding of human biological diversity and the evolutionary processes that gave rise to the patterns of variation we observe in contemporary populations. The study of human variation has not always been unbiased, and thinkers and scientists have always worked in their particular sociohistorical context. For this reason, this chapter opens with a brief overview of race concepts throughout history, many of which relied on unethical and unscientific notions about different human groups.

About the Author

Michael B. C. Rivera

University of Cambridge, mbcr2@alumni.cam.ac.uk

The Arch and Anth Podcast, archandanthpodcast@gmail.com



Michael B. C. Rivera

Michael B. C. Rivera is a biological anthropologist and human bioarchaeologist, studying the transition into agriculture in coastal environments. His recently completed doctoral thesis brought together human skeletal biology, palaeopathology, and prehistoric archaeology to investigate the lives of ancient people on the northeastern European coastline. Being from Hong Kong and a student of human biological variation, Michael is also an advocate for greater inclusion, diversity, and equality in academia. Additionally,

as a believer in the value of science communication, and of the value of the discipline to greater society, he launched The Arch and Anth Podcast in May 2019, which disseminates scientific knowledge in a fun, educational, and informal interview-style audio format.

References

Antrosio, Jason. 2011. "'Race Reconciled': Race Isn't Skin Color, Biology, or Genetics." *Living Anthropologically Website*. https://www.livinganthropologically.com/biological-anthropology/race-reconciled-debunks-race/.

Beals, Kenneth L., Courtland L. Smith, Stephen M. Dodd, J. Lawrence Angel, Este Armstrong, Bennett Blumenberg, Fakhry G. Girgis, et al. 1984. "Brain Size, Cranial Morphology, Climate, and Time Machines [and Comments and Reply]." *Current Anthropology* 25 (3): 301–330.

Betti, Lia, François Balloux, William Amos, Tsunehiko Hanihara, and Andrea Manica. 2008. "Distance from Africa, Not Climate, Explains Within-Population Phenotypic Diversity in Humans." *Proceedings of the Royal Society B: Biological Sciences* 276 (Issue 165B): 809–814. doi:10.1098/rspb.2008.1563.

Betti, Lia, François Balloux, Tsunehiko Hanihara, and Andrea Manica. 2010. "The Relative Role of Drift and Selection in Shaping the Human Skull." *American Journal of Physical Anthropology* 141 (1): 76–82. doi:10.1002/ajpa.21115.

Betti, Lia, Noreen von Cramon-Taubadel, Andrea Manica, and Stephen J. Lycett. 2013. "Global Geometric Morphometric Analyses of the Human Pelvis Reveal Substantial Neutral Population History Effects, Even across Sexes." *PloS ONE* 8 (2): e55909. doi:10.1371/journal.pone.0055909.

———. 2014. "The Interaction of Neutral Evolutionary Processes with Climatically Driven Adaptive Changes in the 3D Shape of the Human Os Coxae." *Journal of Human Evolution* 73 (August): 64–74. doi:10.1016/j.jhevol.2014.02.021.

Boas, Franz. 1931. "Race and Progress." *Science* 74 (1905): 1–8.

Bowden, Rory, Tammie S. MacFie, Simon Myers, Garrett Hellenthal, Eric Nerrienet, Ronald E. Bontrop, Colin Freeman, Peter Donnelly, and Nicholas I. Mundy. 2012. "Genomic Tools for Evolution and Conservation in the Chimpanzee: *Pan troglodytes ellioti* Is a Genetically Distinct Population." *PLoS Genetics* 8 (3): e1002504. doi:10.1371/journal.pgen.1002504.

Gerbault, Pascale, Anke Liebert, Yuval Itan, Adam Powell, Mathias Currat, Joachim Burger, Dallas M. Swallow, and Mark G. Thomas. 2011. "Evolution of Lactase Persistence: An Example of Human Niche Construction." *Philosophical Transactions of the Royal Society B* 366 (1566): 863–877. doi:10.1098/rstb.2010.0268.

Hooton, Earnest A. 1936. "Plain Statements about Race." *Science* 83 (2161): 511–513.

Hrdlička, Aleš. 1918. "Physical Anthropology: Its Scope and Aims; Its History and Present Status in America. A: Physical Anthropology; Its Scopes and Aims." *American Journal of Physical Anthropology* 1 (1): 3–23.

Huxley, Julian. 1942. *Evolution: The Modern Synthesis*. London: Allen and Unwin.

Ingram, Catherine J. E., Charlotte A. Mulcare, Yuval Itan, Mark G. Thomas, and Dallas M. Swallow. 2009. "Lactose Digestion and the Evolutionary Genetics of Lactase Persistence." *Human Genetics* 124 (6): 579–591. doi:10.1007/s00439-008-0593-6.

Jablonski, Nina G. 2004. "The Evolution of Human Skin and Skin Color." *Annual Review of Anthropology* 33 (1): 585–623. doi:10.1146/annurev.anthro.33.070203.143955.

Jablonski, Nina G., and George Chaplin. 2000. "The Evolution of Human Skin Coloration." *Journal of Human Evolution* 39 (1): 57–106. doi:10.1006/jhev.2000.0403.

Kronenberg, Zev N., Ian T. Fiddes, David Gordon, Shwetha Murali, Stuart Cantsilieris, Olivia S. Meyerson, Jason G. Underwood, et al. 2018. "High-Resolution Comparative Analysis of Great Ape Genomes." *Science* 360 (6393): eaar6343. doi:10.1126/science.aar6343.

Lewontin, Richard. 1972. "The Apportionment of Human Diversity." *Evolutionary Biology* 6: 381–398. Eds. Dobzhansky, Theodosius, Hecht, Max K., Steere, William C. Springer, New York, NY

Linnaeus, Carl. 1758. *Systema Naturae*. http://www.cabdirect.org/abstracts/20057000018.html.

Liu, Hua, Franck Prugnolle, Andrea Manica, and François Balloux. 2006. “A Geographically Explicit Genetic Model of Worldwide Human-Settlement History.” *American Journal of Human Genetics* 79 (2): 230–237.

Livingstone, Frank B. 1962. “On the Nonexistence of Human Races.” *Current Anthropology* 3 (3): 279–281.

Long, Jeffery C., and Kittles, Rick A. 2003. “Human Genetic Diversity and the Nonexistence of Biological Races.” *Human Biology* 75 (4): 449–471.

Luzzatto, Lucio. 2012. “Sickle Cell Anaemia and Malaria.” *Mediterranean Journal of Hematology and Infectious Diseases* 4 (1). doi:10.4084/MJHID.2012.065.

Manica, Andrea, William Amos, François Balloux, and Tsunehiko Hanihara. 2007. “The Effect of Ancient Population Bottlenecks on Human Phenotypic Variation.” *Nature* 448 (7151): 346–348. doi:10.1038/nature05951.

Mourant, A. E., Ada C. Kopeć, and Kazimiera Domaniewska-Sobczak. 1976. *The Distribution of the Human Blood Groups and Other Polymorphisms*. 2nd edition. Oxford: Oxford University Press.

Morton, Samuel George. 1839.*Crania Americana, or, A comparative view of the skulls of various aboriginal nations of North and South America*. Philadelphia: J. Dobson.

National Research Council (U.S.) Committee on Human Genome Diversity. 1997. *Evaluating Human Genetic Diversity.* Washington, D.C.: National Academies Press

Ousley, Stephen D., Richard L. Jantz, and Donna Freid. 2009. “Understanding Race and Human Variation: Why Forensic Anthropologists Are Good at Identifying Race.” *American Journal of Physical Anthropology* 139 (1): 68–76. doi:10.1002/ajpa.21006.

Oxford English Dictionary. 2018. 3rd edition. Oxford: Oxford University Press.

Ponce de León, Marcia S., Toetik Koesbardiati, John David Weissmann, Marco Millela, Carlos S. Reyna-Blanco, Gen Suwa, Osamu Kondo, Anna-Sapfo Malaspinas, Tim D. White, and Christoph P. E. Zollikofer. 2018. “Human Bony Labyrinth Is an Indicator of Population History and Dispersal from Africa.” *Proceedings of the National Academy of Sciences* 115 (16): 4128–4133. doi:10.1073/pnas.1808125115.

Prado-Martinez, Javier, Peter H. Sudmant, Jeffrey M. Kidd, Heng Li, Joanna L Kelley, Belen Lorente-Galdos, Krishna R. Veeramah, et al. 2013. “Great Ape Genetic Diversity and Population History.” *Nature* 499 (7459): 471–475. doi:10.1038/nature12228.

Prugnolle, Franck, Andrea Manica, and François Balloux. 2005. “Geography Predicts Neutral Genetic Diversity of Human Populations.” *Current Biology* 15 (5): 159–160.

Rathmann, Hannes, Hugo Reyes-Centeno, Silvia Ghirotto, Nicole Creanza, Tsunehiko Hanihara, and Katerina Harvati. 2017. “Reconstructing Human Population History from Dental Phenotypes.” *Scientific Reports* 7: 12495. doi:10.1038/s41598-017-12621-y.

Relethford, John H. 2001. “Global Analysis of Regional Differences in Craniometric Diversity and Population Substructure.” *Human Biology* 73 (5): 629–636. doi:10.1353/hub.2001.0073.

———. 2002. “Apportionment of Global Human Genetic Diversity Based on Craniometrics and Skin Color.” *American Journal of Physical Anthropology* 118 (4): 393–398. doi:10.1002/ajpa.10079.

———. 2004. “Global Patterns of Isolation by Distance Based on Genetic and Morphological Data.” *Human Biology* 76 (4): 499–513. doi:10.1353/hub.2004.0060.

———. 2009. “Race and Global Patterns of Phenotypic Variation.” *American Journal of Physical Anthropology* 139 (1): 16–22. doi:10.1002/ajpa.20900.

Rosenberg, Noah A., Saurabh Mahajan, Sohini Ramachandran, Chengfeng Zhao, Jonathan K. Pritchard, and Marcus W. Feldman. 2005. “Clines, Clusters, and the Effect of Study Design on the Inference of Human Population Structure.” *PLoS Genetics* 1 (6): e70. doi:10.1371 /journal.pgen.0010070.

Rosenberg, Noah A., Jonathan K. Pritchard, James L. Weber, Howard M. Cann, Kenneth K. Kidd, Lev A. Zhivotovsky, and Marcus W. Feldman. 2002. “Genetic Structure of Human Populations.” *Science* 298 (5602): 2381–2385.

Sauer, Norman J. 1992. “Forensic Anthropology and the Concept of Race: If Races Don’t Exist, Why Are Forensic Anthropologists So Good at Identifying Them?” *Social Science and Medicine* 34 (2): 107–111. doi:10.1016/0277-9536(92)90086-6.

Staes, Nicky, Chet C. Sherwood, Katharine Wright, Marc de Manuel, Elaine E. Guevara, Tomas Marques-Bonet, Michael Krützen, et al. 2017. “FOXP2 Variation in Great Ape Populations Offers Insight into the Evolution of Communication Skills.” *Scientific Reports* 7 (1): 1–10. doi:10.1038/s41598-017-16844-x.

von Cramon-Taubadel, Noreen, and Stephen J. Lycett. 2008. “Brief Communication: Human Cranial Variation Fits Iterative Founder Effect Model with African Origin.” *American Journal of Physical Anthropology* 136 (1): 108–113. doi:10.1002/ajpa.20775.

Warren, Kerryn A. 2018. “AAPA Name Change.” *Bone and Evolution*, April 4. https://bonevolution.wordpress.com/2018/04/04/aapa-name-change/.

Weiss, Kenneth M., and Jeffrey C. Long. 2009. “Non-Darwinian Estimation: My Ancestors, My Genes’ Ancestors.” *Genome Research* 19: 703–710. doi:10.1101/gr.076539.108.19.

Wise, Tim. 2010. *Colorblind: The Rise of Post-Racial Politics and the Retreat from Racial Equity*. San Francisco: City Lights.

Yudell, Michael, Dorothy Roberts, Rob DeSalle, and Sarah Tishkoff. 2016. “Taking Race out of Human Genetics.” *Science* 351 (6273): 564–565. doi:10.1126/science.aac4951.

Figure Attributions

Figure 13.1a Tanzania – Hadzabe hunter (14533536392) by A_Peach from Berlin, Germany, is used under a CC BY 2.0 License.

Figure 13.1b Inuit-Kleidung 1 by Ansgar Walk is used under a CC BY-SA 3.0 License.

Figure 13.1c Andean Man by Cacophony is used under a CC BY-SA 4.0 License.

Figure 13.1d Jane Goodall GM byFloatjon is used under a CC BY-SA 3.0 License.

Figure 13.2 Egyptian races Drawing (1772-1846) by an unknown artist after a mural of the tomb of Seti I, Copy by Heinrich von Minutoli (1820), is in the public domain.

Figure 13.3 Naturalishistoria from the front page of Pliny the Elder’s Naturalis Historia is in the public domain.

Figure 13.4 Great Chain of Being 2 by Didacus Valades (Diego Valades) is in the public domain.

Figure 13.5 Carl von Linné by Alexander Roslin artist QS:P170,Q315102 is in the public domain.

Figure 13.6 Discovery of the Mississippi by William Henry Powell artist QS:P170,Q3568696 (photograph courtesy Architect of the Capitol) is in the public domain.

Figure 13.7 Blumenbach’s five races by Johann Friedrich Blumenbach is in the public domain.

Figure 13.8 (Ales Hrdlicka) SIA2009-4246 by Unknown photographer is in the public domain.

Figure 13.9 Eugenics congress logo scanned from Harry H. Laughlin, The Second International Exhibition of Eugenics held September 22 to October 22, 1921, is in the public domain.

Figure 13.10 Dobzhansky no Brasil em 1943 by Unknown photographer is in the public domain.

Figure 13.11 Julian Huxley 1-2 by Unknown photographer is in the public domain.

Figure 13.12 Skin color by S25454541 is used under a CC BY-SA 4.0 License.

Figure 13.13a Map of blood group a by Muntuwandi at en.Wikipedia is used under a CC BY-SA 3.0 License.

Figure 13.13b Map of blood group b by Muntuwandi at en.Wikipedia is used under a CC BY-SA 3.0 License.

Figure 13.13c Map of blood group o Based on diagrams from anthro.palomar.edu/vary/vary_3.htm reproduced from A. E. Mourant et.al., The Distribution of the Human Blood Groups and Other Polymorphisms, 2nd ed. (1976) is used under a CC BY-SA 3.0

License.

Figure 13.14 Sub-Saharan-Africa by Ezeu has been designated to the public domain (CC0).

Figure 13.15 Bottleneck effect by Tsaneda is used under a CC BY 3.0 License.

Figure 13.16 Chimpanzee IV (13968482163) by Chi King is used under a CC BY 2.0 License.

Figure 13.17 Human skulls by 22Kartika is used under a CC BY-SA 3.0 License.

Figure 13.18 Bony labyrinth by Selket (5 February 2007, UTC) has been designated to the public domain (CC0).

Figure 13.19 Forensic Anthropology Lab by Pp391 is used under a CC BY-SA 3.0 License.

Figure 13.20 Michael B. C. Rivera in Hong Kong original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

6: Race and Human Variation is shared under a CC BY-NC license and was authored, remixed, and/or curated by LibreTexts.

- **13: Race and Human Variation** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.1: The History of "Race" Concepts

“Race” in the Classical Era

The earliest classification systems used to understand human diversity are evidenced by ancient manuscripts, scrolls, and stone tablets recovered through archaeological, historical, and literary research. The Ancient Egyptians had the *Book of Gates*, dated to the New Kingdom between 1550 B.C.E. and 1077 B.C.E (Figure 13.2). In one part of this tome dedicated to depictions of the underworld, scribes used pictures and hieroglyphics to illustrate a division of Egyptian people into the four categories known to them at the time: the Aamu (Asiatics), the Nehesu (Nubians), the Reth (Egyptians), and the Themehu (Libyans). Though not rooted in any scientific basis like our current understandings of human variation today, the Ancient Egyptians believed that each of these groups were made of a distinctive category of people, distinguishable by their skin color, place of origin, and even behavioral traits.



Figure \(\PageIndex{1}\): (from left to right) Depicting a Berber (Libyan), a Nubian, an Asiatic (Levantine), and an Egyptian, copied from a mural of the tomb of Seti I.



Figure \(\PageIndex{2}\): Front page of Pliny the Elder’s Naturalis Historia.

The Roman philosopher Pliny the Elder (23–79 C.E.) also wrote about different groupings of people in his encyclopedia *Naturalis Historia* (Figure 13.3). In his opinion, all people fit under one of three categories: civilized peoples, barbarians, and monstrous individuals. Pliny the Elder’s work was deeply problematic. He believed that only Europeans were civilized and not monstrous-looking, while other groups of people lacked the ideal character and appearance. In both the cases of the *Book of Gates* and *Naturalis Historia*, the worldviews of those who wrote these volumes were also limited by how few and infrequent their encounters were with peoples elsewhere around the world—that is, those not residing in Europe, the Near East, or northern Africa. When faced with only the level of biological diversity they could see around them, distinguishing factors identified by these prominent thinkers relied simply on readily visible phenotypic traits, such as body size, skin color, and facial shape.

The most well-known of early documents is perhaps the Bible, where it is written that all humankind descends from one of three sons of Noah: Shem (the ancestor to all olive-skinned Asians), Japheth (the ancestor to pale-skinned Europeans), and Ham (the ancestor to darker-skinned Africans). Similar to the Ancient Egyptians, these distinctions were based on behavioral traits and skin color. More recent work in historiography and linguistics suggest that the branches of “Hamites,” “Japhethites,” and “Shemites” may also relate to the formation of three independent language groups some time between 1000 and 3000 B.C.E. With the continued proliferation of Christianity, this concept of approximately three racial groupings lasted until the Middle Ages and spread as far across Eurasia as crusaders and missionaries ventured at the time.


Figure \(\PageIndex{3}\): The Great Chain of Being from the Rhetorica Christiana by Fray Diego de Valades (1579).

Finally, there is also the "Great Chain of Being," conceived by ancient Greek philosophers like Plato (427–348 B.C.E.) and Aristotle (384–322 B.C.E.). They played a key role in laying the foundations of empirical science, whereby observations of everything from animals to humans were noted with the aim of creating taxonomic categories. Aristotle describes the Great Chain of Being as a ladder along which all objects, plants, animals, humans, and celestial bodies can be mapped in an overall hierarchy (in the order of existential importance, with humans placed near the top, just beneath divine beings) (Figure 13.4). Where he writes about humans, Aristotle expressed the belief that certain people are inherently (or genetically) more instinctive rulers, while others are more natural fits for the life of a worker or slave. Nowadays, based on research by biological anthropologists, we currently recognize that these early systems of classification and hierarchization are unhelpful in studying human biological diversity. Both behavioral traits and physical traits are coded for by multiple genes each, and how we exhibit those traits based on our genetics can vary significantly even between individuals of the same population.

"Race" during the Scientific Revolution

The 1500s and 1600s saw the beginnings of the "**Scientific Revolution**" in European societies, with thinkers like Copernicus, Galileo, and Da Vinci publishing some of their most important findings. While by no means the first or only scholars globally to use observation and experimentation to understand the world around them, early scientists living at the end of the medieval period in Europe increasingly employed more experimentation, quantification, and rational thought in their work. This is the main difference between the work of the ancient Egyptians, Romans, and Greeks, and that of workers like Isaac Newton and Carl Linnaeus in the 1600s and 1700s.




Figure \(\PageIndex{4}\): Carl Linnaeus.

Linnaeus is the author of *Systema Naturae* (1758), in which he classified all plants and animals he could observe under the first formalized naming system known as **binomial nomenclature** (i.e., how all organisms can be named by their genus and species, such as *Homo sapiens* or *Pan troglodytes*) (Figure 13.5). What was most anthropologically notable about Linnaeus's taxonomy was that he was one of the first to group humans with apes and monkeys, after noting the anatomical similarities between humans and

nonhuman primates. Linnaeus viewed the world in line with **essentialism**, a concept which dictates that there are a unique set of characteristics that organisms of a specific kind *must* have—organisms would fall outside taxonomic categorizations if they lacked any of the required criteria.

Despite these useful contributions to the biological sciences, Linnaeus still subdivided the human species into four varieties, with overtly racist categories based on skin color and "inherent" behaviors. According to him, Africans are all "black-skinned" and ruled by an erratic nature; Native Americans are "red" in skin tone and ruled by habit; Asians are "yellow-" or "brown-skinned" and ruled by belief; and Europeans are "white" and regulated by custom. These standards for categorization imply that Europeans are governed by carefully considered culture and custom, unlike the unthinking Asians and Indigenous Americans in his framework who normally act out of "habit" or "belief." Moreover, Linnaeus's traditional ranking also places sub-Saharan, dark-skinned Africans inferior to the other three. Wrongly so, European scientists during this period were not aware of their own biases skewing their interpretations of biological diversity. The conclusions and claims they came to, consciously or subconsciously, often fit such an age when the superiority of European cultures over others was a pervasive idea throughout these scientists' social and political lives.



Figure \(\PageIndex{5}\): Discovery of the Mississippi by Spanish colonialist explorer Hernando DeSoto in 1541 (painted in 1853 by William H. Powell).

Occurring alongside this Scientific Revolution was also the "**Age of Discovery**." Although much of Eurasia was linked by spice and silk trading routes, the European colonial period between the 1400s and 1700s was marked by many new and intentionally violent encounters overseas (Figure 13.6). When Europeans arrived by ship on the shores of continents that were already inhabited, it was their first meeting with the indigenous peoples of the Americas and Australasia, who looked, spoke, and behaved differently from peoples with whom they were familiar. Building on the idea of species and "subspecies," natural historians of this time invented the term *race*, from the French *rasse* meaning "local strain." The idea behind this terminology was rooted in the observation that geography plays a significant role in producing the biological traits we observe today. Naturalists like Comte de Buffon and Johann Blumenbach did believe that all people have a single origin, but they also believed that differences in environment could lead to biological changes between different groups of people (i.e., races). However, as they had no understanding of genetics, they were incorrect in assuming that factors such as skin color could change in a single lifetime depending on climate and diet and, essentially, behavior. Again, while drawing links between external physical characteristics and behavior is not scientific, differences in both were used to justify the **Othering** of "nonwhite" cultures. Establishing "otherness" and "inferiority" in other people's cultures was necessary at the time for colonialists to enforce European domination and the subordination of non-European people. Without genetic technologies, little was known at the time about the hereditary or evolutionary basis of skin color having little to do with innate differences between various "races."

Another such scientist at the time, Johann Friedrich Blumenbach (1752–1840), classified humans into five races based on his observations of cranial form variation as well as skin color. He thus dubbed the "original" form of the human cranium the "Caucasian" form, with the idea that the ideal climate conditions for early humans would have been in the Caucasus region near the Caspian Sea. The key insight Blumenbach presented was that human variation in any particular trait should be more accurately viewed as falling along a gradation (Figure 13.7). While some of his theories were correct according to what we observe today with

more knowledge in genetics, workers like him and Buffon believed erroneously that human "subspecies" were "degenerated" or "transformed" varieties of an ancestral Caucasian or European race. According to them, the Caucasian cranial dimensions were the least changed over human evolutionary time, while the other skull forms represented geographic variants of this "original." As will be discussed in greater detail later in this chapter, we have genetic and craniometric evidence for sub-Saharan Africa being the origin of the human species instead. Based on work that shows how most biological characteristics are coded for by nonassociated genes, it is not reasonable to draw links between individuals' personalities and their skull shapes.




Figure \(\PageIndex{6}\): Five skull drawings representing specimens for Blumenbach's "Mongolian," "American," "Caucasian," "Malayan," and "Aethiopian" races.

"Race" and the Dawn of Scientific Racism

Between the 1800s and mid-1900s, and contrary to what you might expect, an increased use of scientific methods to justify racial schemes developed in scholarship. Differing from Blumenbach and Buffon's views in earlier centuries, which saw all humans as environmentally deviated from one "original" humankind, classification systems after 1800 became more **polygenetic** (viewing all people as having separate origins) rather than **monogenetic** (viewing all people as having a single origin). Instead of moving closer to our modern-day understandings of human diversity, there was increased support for the notion that each race was created separately and with different attributes (intelligence, temperament, and appearance).

The 1800s were an important precursor to modern biological anthropology as we know it, given that the scientific measurement of human physical features (anthropometry) truly became popularized then. However, whether it was skin color, skull shape, or observations of behavior being analyzed as the data, empirical studies in the 1800s pushed the idea even further that Europeans were culturally and biologically superior. The leading figures in craniometry at this time, focusing on measurements of the skull, were also linked heavily with powerful individuals and wealthy sociopolitical institutions and financial bodies. Therefore, polygenetic ways of thinking were particularly influenced by sociohistorical and economic factors at the time. Theories in support of hierarchical racial schemes certainly helped continue the exploitative and unethical transatlantic slave trade between the 1500s and 1800s by justifying the transport and enslavement of African people on a "scientific" basis.

While considered one of the pioneers of American "physical" anthropology, Samuel George Morton (1799–1851) was a scholar who had a large role in 1800s scientific racism. By measuring cranial size and shape, he calculated that "Caucasians," on average, have greater cranial volumes than other groups, such as the Native Americans and "Negros." Today, we know that cranial size variation depends on such factors as Allen's and Bergmann's rules, which give the more likely explanation for the largest heads being found in people living among colder regions (i.e., Europeans) being climatic adaptation (Beals et al. 1984). In colder environments, it is advantageous for those living there to have larger and rounder heads because they conserve heat more effectively than slenderer heads (Beals et al. 1984).

Morton went on to write in his publication *Crania Americana* (1839) a number of views that fit with a concept called **biological determinism**. The idea behind biological determinism is that an association exists between people's physical characteristics and their behavior, intelligence, ability, values, and morals. If the idea is that some groups of people are *essentially* superior to others in cognitive ability and temperament, then it is easier to justify the unequal treatment of certain groups based on outward appearances. Based on his cranial measurements and observations of human nature, Morton claimed that Europeans were the most intelligent and "well-proportioned," while Asians were not fit for leadership and had short attention spans, Native Americans were slow in acquiring knowledge and fond of war, and Africans were superstitious, uninventive, and "barbarous."

Another such problematic thinker was Paul Broca (1824–1880), after which a region of the frontal lobe related to language use is named (Broca's area). Influenced by Morton, he likewise claimed that internal skull capacities could be linked with skin color and

cognitive ability. Considering his data taken from different parts of the globe, Broca thought that factors such as gender, education, and social status could have an influence on brain size for different groups, purporting that men had larger brains than women and that “eminent” men were superior to men of “mediocre talent.” He went on to justify the European colonization of other global territories by purporting it was noble for a biologically more “civilized” population to improve the “humanity” of more “barbaric” populations. Today, these theories of Morton, Broca, and others like them are known to have no scientific basis. If we could speak with them today, they would likely try to emphasize that their conclusions were based on empirical evidence and not *a priori* reasoning. However, we now can clearly see that their reasoning was biased and affected by prevailing societal views at the time.

“Race” and the Beginnings of Physical Anthropology

In the early 20th century, we saw a number of new figures coming into the science of human variation and shifting the theoretical focuses within. Most notably, these included Aleš Hrdlička and Franz Boas.




Figure \(\PageIndex{7}\): Aleš Hrdlička (1869–1943), a Czech anthropologist who founded the American Journal of Physical Anthropology.

Aleš Hrdlička (1869–1943) was a Czech anthropologist who moved to the United States. In 1903, he established the physical anthropology section of the National Museum of Natural History (Figure 13.8). In 1918, he founded the *American Journal of Physical Anthropology*, one of the foremost scientific journals disseminating bioanthropological research still today. As part of his work and the scope of the journal, he differentiated “**physical anthropology**” from other kinds of anthropology—he wrote that physical anthropology is “the study of racial anatomy, physiology, and pathology” and “the study of man’s variation” (Hrdlička 1918). In some ways, although the scope and technological capabilities of biological anthropologists have changed significantly, Hrdlička established an area of inquiry that has continued and prospered for over a hundred years.

Franz Boas (1858–1942) was a German American anthropologist who established the four-field anthropology system in the United States and founded the American Anthropological Association in 1902. He argued that the scientific method should be used in the study of human cultures and the comparative method for looking at human biology worldwide. Boas’s specialization was in the study of skull dimensions with respect to race. After a long-term research project, he demonstrated how cranial form was highly dependent on cultural and environmental factors and that human behaviors were influenced primarily not by genes but by social learning. He wrote in one essay for the journal *Science*: “While individuals differ, biological differences between races are small. There is no reason to believe that one race is by nature so much more intelligent, endowed with great willpower, or emotionally more stable than another, that the difference would materially influence its culture” (Boas 1931:6). This conclusion directly contrasted with the theories of the past that relied on biological determinism. Biological anthropologists today have found evidence that corroborates Boas’s explanations: societies do not exist on a hierarchy or gradation of “civilizedness” but instead are shaped by the world around them, their demographic histories, and the interactions they have with other groups.




Figure \(\PageIndex{8}\): Logo of the Second International Exhibition of Eugenics held in 1921.

The first half of the 1900s still involved some research that was essentialist and focused on proving racial determinism. Anthropologists like Francis Galton (1822–1911) and Earnest A. Hooton (1887–1954) created the field of **eugenics** as an attempt to formalize the social scientific study of "fitness" and "superiority" among members of 19th-century Europe. As a way of "dealing with" criminals, diseased individuals, and "uncivilized" people, eugenicists recommended prohibiting parts of the population from being married and sterilizing these members of society so they could no longer procreate (Figure 13.9). They instead encouraged "reproduction in individual families with sound physiques, good mental endowments, and demonstrable social and economic capability" (Hooton 1936). In the 1930s, Nazi Germany used this false idea of there being "pure races" to highly destructive effect. The need to be protected against admixture from "unfit" groups was their justification for their blatant racism and purging of citizens that fell under their subjective criteria.

Shortly after World War II and the Nazi Holocaust, the full extent of essentialist, eugenicist thinking became clear. Social constructions of race, and the notion that you could predict psychological or behavioral traits based on external appearance, had become unpopular both within and outside the discipline. It was up to those in the field of physical anthropology at the time to separate physical anthropology from race concepts that supported unscientific and socially damaging agendas. This does not mean that there are no physiological or behavioral differences between different members of the human species. However, going forward, a number of physical anthropologists saw human biological variation as more complicated than simple **typologies** could describe.

6.1: The History of "Race" Concepts is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **13.1: The History of "Race" Concepts** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.2: Human Variation in Biological Anthropology Today

"Populations" Instead of "Races"


Figure **6.2.1**: Theodosius Dobzhansky, an important scientist who formulated the 20th-century "modern synthesis" reconciling Charles Darwin's theory of evolution and Gregor Mendel's ideas on heredity.

After 1950, replacing the concept of "race" as a unit of diversity was the "**population**." This was outlined by those pioneering the "new physical anthropology," such as Sherwood Washburn, Theodosius Dobzhansky, and Julian Huxley, who borrowed this way of framing human groups from contemporary population geneticists (Figure 13.10). "Races" were then defined simply as populations that differ in the frequency of some gene or genes. And, on the other hand, a "population" is a group of individuals potentially capable of or actually interbreeding due to shared geographic proximity, language, ethnicity, culture, and/or values. Put another way, a population is a *local* interbreeding group with reduced gene flow between themselves and other groups of humans. Members of the same population may be expected to share many genetic traits (and, as a result, many phenotypic traits that may or may not be visible outwardly).


Figure **6.2.2**: Julian Huxley (1942).

Thinking of humans in terms of populations was part of Julian Huxley's (1942) "Modern Synthesis"—so named because it helped to reconcile two fundamental principles about evolution that had not been made sense of together before (Figure 13.11). As discussed in Chapter 3, Gregor Mendel (1822–1884) was able to show that inheritance was mediated by discrete particles (or genes) and not blended in the offspring. However, it was difficult for some 19th-century scientists to accept this model of genetic inheritance at the time because much of biological variation appeared to be continuous and not particulate (take skin color or height as examples). In the 1930s, it was demonstrated that traits could be polygenic and that multiple alleles could be responsible for any one phenotypic trait, thus producing the continuous variation in traits such as eye color that we see today. Thus, Huxley's "Modern Synthesis" outlines not only how human populations are capable of exchanging genes at the microevolutionary level but also how multiple alleles for one trait (polygenic exchanges) can cause gradual macroevolutionary changes.

Human Variation Is Clinal/Continuous (Not Discrete)

Human diversity cannot be broken into discrete "races," because most physical traits vary on a continuous or "clinal" basis. One obvious example of this is how human height does not only come in three values ("short," "medium," and "tall") but instead varies across a spectrum of vertical heights achievable by humans all over the world. (However, this is with the only difference being the huge divergence in how factors like body size and traits such as skin color have been viewed and used sociopolitically as a way of separating people throughout history.) The need to shift from typological "race" categories to a more nuanced understanding of continuously variable populations was realized by anthropologists working in the 1960s and 1970s who shifted their focus toward the study of individual traits rather than the study of groups (populations, races). Systematic evaluations of global biological

variation in humans only began then, when large numbers of genetic loci for large numbers of samples were sampled from human populations distributed worldwide. It was during the 1960s that "**clines**" in human genetic variation were first identified.

Frank B. Livingstone (1928–2005) wrote: "There are no races, only clines" (1962). A *cline* is a gradation in the frequency of an allele/trait between populations living in different geographic regions. In order to study human traits that are clinally distributed, it is often required to perform genetic testing to uncover the true frequencies of an allele or trait across a certain geographic space. One easily visible example of a clinal distribution seen worldwide is the patterning of human variation in skin color. Whether in southern Asia, sub-Saharan Africa, or Australia, dark brown skin is found. Paler skin tones are found in higher-latitude populations such as those who have lived in areas like Europe, Siberia, and Alaska for millennia. Skin color is easily observable as a phenotypic trait exhibiting **continuous variation**.

A clinal distribution still derives from genetic inheritance, but clines often correspond to some gradually changing environmental factor. Clinal patterns arise when selective pressures in one geographic area differ from those in another as well as when people procreate and pass on genes together with their most immediate neighbors. There are several mechanisms, selective and neutral, that can lead to the clinal distribution of an allele or a biological trait. **Natural selection** is the mechanism that produced a global cline of skin color, whereby darker skin color protects equatorial populations from high amounts of UV radiation; there is a transition of lessening pigmentation in individuals that reside further and further away from the tropics (Jablonski 2004; Jablonski and Chaplin 2000) (Figure 13.12). The ability and inability to digest lactose (milk sugar) among different world communities varies according to differential practices and histories of milk and dairy product consumption (Gerbault et al. 2011; Ingram et al. 2009). Where malaria seems to be most prevalent as a disease stressor on human populations, a clinal gradient of increasing sickle cell anemia experience toward these regions has been studied extensively by genetic anthropologists (Luzzatto 2012). Sometimes culturally defined mate selection based on some observable trait can lead to clinal variation between populations as well.



Figure **6.2.3**: Global map of indigenous skin colors.

Two neutral microevolutionary processes that may produce a cline in a human allele or trait are **gene flow** and **genetic drift**. The ways in which neutral processes can produce clinal distributions is seen clearly when looking at clinal maps for different blood groups in the human ABO blood group system (Figure 13.13). For instance, scientists have identified an east-to-west cline in the distribution of the blood type *B* allele across Eurasia. The frequency of *B* allele carriers decreases gradually westward when we compare the blood groups of East and Southeast Asian populations with those in Europe. This shows how populations residing nearer to one another are more likely to interbreed and share genetic material (i.e., undergo gene flow). We also see 90%–100% of native South American individuals, as well as between 70%–90% of Aboriginal Australian groups, carrying the *O* allele (Mourant et al. 1976). These high frequencies are likely due to random genetic drift and founder effects, in which population sizes were severely reduced by the earliest *O* allele-carrying individuals migrating into those areas. Over time, the *O* blood type has remained predominant.




Figure **6.2.4**: Global distribution of blood group A.




Figure **6.2.5**: Global distribution of blood type B.




Figure **6.2.6**: Global distribution of blood type O.

The Apportionment of Human Variation: Genetic Diversity Is Greater Within-Group Than Between-Groups

One problem with race-based classifications is they relied on an erroneous idea that people within a typological category were more similar to each other than they were to people in other groups. In other words, “race” concepts were predicated on the notion that individuals with particular characteristics would share more similar genes with each other within a particular “race” and share less with individuals of other “races” possessing different traits and genetic makeups. However, since around 50 years ago, scientific studies have shown that the majority of human genetic differences worldwide exist *within* groups (or “races”) individually rather than *between* groups.

Richard Lewontin (1929–) is a biologist and evolutionary geneticist who authored a paper evaluating where the total genetic variation in humans lies. This article, titled “The Apportionment of Human Diversity” (Lewontin 1972), addressed the following question: On average, how genetically similar are two randomly chosen people from the same group when compared to two randomly chosen people from different groups? Lewontin studied this problem by using genetic data. He obtained data for a large number of different human populations worldwide using 17 genetic markers (including alleles that code for various important enzymes and proteins, such as blood-group proteins). The statistical analysis he ran used a measure of human genetic differences in and among populations known as the fixation index (F_{ST}). Technically, F_{ST} can be defined as the proportion of total genetic variance within a *subpopulation* relative to the total genetic variance from an *entire population*. Therefore, F_{ST} values range from 0 to 1 (or, sometimes you will see this stated as a percentage between 0% and 100%). The closer the F_{ST} value of a population (e.g., the world’s population) approaches 1, the higher the degree of genetic differentiation among subpopulations relative to the overall

population. In his paper, Lewontin (1972) identified that most of human genetic differences (85.4%) were found within local subpopulations (e.g., the Germans or Easter Islanders), whereas 8.3% were found between populations within continental human groups, and 6.3% were attributable to traditional "race" groups (e.g., "Caucasian" or "Amerind"). These findings have been important for scientifically rejecting the existence of biological races (Long and Kittles 2008).

In 2002, another landmark article by Noah Rosenberg and colleagues (2002) explored worldwide human genetic variation using an even-greater genetic data set. They used 377 highly variable markers in the human genome and sampled from 1,056 individuals representative of 52 populations. The markers chosen for study were not ones that code for any expressed genes. Because these regions of the human genome were made of unexpressed genes, we may understand these markers as neutrally derived (as opposed to selectively derived) as they do not code for functional advantages or disadvantages. These neutral genetic markers likely reflect an intricate combination of regional **founder effects** and population histories. Analyses of these neutral markers allowed scientists to identify that a majority of global genetic variance (93%–95%) can be accounted for by within-population differences at the 377 genetic loci, while only a small proportion of genetic variance (3%–5%) can be attributed to differences among major groups (Rosenberg et al. 2002). Like Lewontin's (1972) findings, this lends support to the theory that distinct biological races do not exist, even though misguided concepts of race may still have real social and political consequences.

Biological Data Fit Isolation-By-Distance and Out-of-Africa Models

One further note is that while the world's population may be genetically divided into "groups," "subsets," "clumps," or "clusters" that reflect some degree of genetic similarity, it is more likely that these identifiable clusters reflect genetic or geographic distances —either with gene flow facilitated by proximity between populations or impeded by obstacles like oceans or environmentally challenging habitats (Rosenberg et al. 2005). Sometimes, inferred clusters using multiple genetic loci are interpreted by non-geneticists literally as "ancestral populations." However, it would be wrong to assume from these genetic results that highly differentiated and "pure" ancestral groups ever existed. These groupings reflect differences that have arisen over time due to clinal patterning, genetic drift, and/or restricted or unrestricted gene flow (Weiss and Long 2009). The clusters identified by scientists are arbitrary and the parameters used to split up the global population into groups is subjective and dependent on the particular questions or distinctions being brought into focus (Relethford 2009).



Figure **6.2.7**: Sub-Saharan Africa (shaded dark/green).

Additionally, research on worldwide genetic diversity has shown that human variation decreases with increasing distance from sub-Saharan Africa, where there is evidence for this vast region being the geographical origin of anatomically modern humans (Liu et al. 2006; Prugnolle et al. 2005) (Figure 13.14). Genetic differentiation decreases in human groups the further you sample data from relative to sub-Saharan Africa because of serial founder effects (Relethford 2004). Over the course of human colonization of the rest of the world outside Africa, populations broke away in expanding waves across continents into western Asia, then Europe and eastern Asia, followed by Oceania and the Americas. As a result, founder events occurred whereby genetic variation was lost, as the colonization of each new geographical region involved a smaller number of individuals moving from the original larger population to establish a new one (Relethford 2004). The most genetic variation is found across populations residing in different parts of sub-Saharan Africa, while other current populations in places like northern Europe and the southern tip of South America exhibit some of the least genetic differentiation relative to all global populations.

Besides fitting nicely into the Out-of-Africa model, worldwide human genetic variation conforms to an isolation-by-distance model, which predicts that genetic similarity between groups will decrease exponentially as the geographic distance between them increases. This is because of the greater and greater restrictions to gene flow presented by

geographic distance, as well as cultural and linguistic differences that occur as a result of certain degrees of isolation. Since genetic data conform to isolation-by-distance and Out-of-Africa models, these findings support the abolishment of “race” groupings. This research demonstrates that human variation is continuous and cannot be differentiated into geographically discrete categories. There are no “inherent” or “innate” differences between human groups; instead, variation derives from some degree of natural selection, as well as neutral processes like **population bottlenecking** (Figure 13.15), random **mutations** in the DNA, genetic drift, and gene flow through between-mate interbreeding.




Figure 6.2.8: The founder effect is a change in a small population’s gene pool due to a limited number of individuals breaking away from a parent population.

Humans Have Higher Homogeneity Compared to Many Other Species

An important fact to bear in mind is that humans are 99.9% identical to one another. This means that the apportionments of human diversity discussed above only concern that tiny 0.1% of difference that exists between all humans globally. Compared to other mammalian species, including the other great apes, human diversity is remarkably lower. This may be surprising given that the worldwide human population has already exceeded seven billion, and, at least on the surface level, we appear to be quite phenotypically diverse. Molecular approaches to human and primate genetics tells us that external differences are merely superficial. For a proper appreciation of human diversity, we have to look at our closest relatives in the primate order and mammalian class. Compared to chimpanzees, gibbons, and even gray wolves and giant pandas, humans have remarkably low average genome-wide **heterogeneity**.



Figure 6.2.9: Chimpanzee (Pan troglodytes).

When we look at chimpanzee genetic diversity, it is fascinating that western, central, eastern, and Cameroonian chimpanzee groups have substantially more genetic diversity between them than large global samples of human DNA (Bowden et al. 2012) (Figure 13.16). This is surprising given that all of these chimpanzee groups live relatively near one another in Africa, while measurements of human genetic diversity have been conducted using samples from entirely different continents. First, geneticists suppose that this could reflect differential experiences of the founder effect between humans and chimpanzees. Because all non-African human populations descended from a small number of anatomically modern humans who left Africa, it would be expected that all groups descended from that smaller ancestral group would be similar genetically. Second, our species is really young, given that we have

only existed on the planet for around 150,000 to 300,000 years. This gave humans little time for random genetic mutations to occur as genes get passed down through genetic interbreeding and meiosis. Chimpanzees, however, have inhabited different **ecological niches**, and less interbreeding has occurred between the four chimpanzee groups over the past six to eight million years compared to the amount of gene flow that occurred between worldwide human populations (Bowden et al. 2012).

Recent advances have now enabled the attainment of genetic samples from the larger family of great apes and the evaluation of genetic diversity among bonobos, orangutans, and gorillas alongside that of chimpanzees and humans (Prado-Martinez et al. 2013). Collecting such data and analyzing primate genetic diversity has been important not only to elucidate how different ecological, demographic, and climatic factors have shaped our evolution but also to inform upon conservation efforts and medical research. Genes that may code for genetic susceptibilities to tropical diseases that affect multiple primates can be studied through genome-wide methods. Species differences in the genomes associated with speech, behavior, or cognition could tell us more about how human individuals may be affected by genetically derived neurological or speech-related disorders and conditions (Prado-Martinez et al. 2013; Staes et al. 2017). In 2018, a great ape genomic study also reported genetic differences between chimpanzees and humans related to brain cell divisions (Kronenberg et al. 2018). From these results, it may be inferred that cognitive or behavioral variation between humans and the great apes might relate to an increased number of cortical neurons being formed during human brain development (Kronenberg et al. 2018). Comparative studies of human and nonhuman great ape genetic variation highlight the complex interactions of population histories, environmental changes, and natural selection between and within species. When viewed in the context of overall great ape diversity, we may reconsider how variable the human species is relatively and how unjustified previous “race” concepts really were.

Phenotypic Traits That Reflect Neutral Evolution

Most human traits are non-concordant. “Non-concordance” is a term used to describe how biological traits vary independent of each other—that is, they don’t get inherited in a correlative manner with other genetically controlled traits. For example, if you knew an individual had genes that coded for tall height, you would not be able to predict if they are lighter-skinned or have red hair. Depending on the trait being observed, different patterns of phenotypic variation may be found within and among groups worldwide. In this subsection, some phenotypic traits that reflect the aforementioned patterns of genetic variation will be discussed.



Figure **6.2.10**: Human skulls in Tana Toraja (Indonesia), common scenery in public graves.

Looking beyond genetic variation briefly, recent studies have revisited biological anthropology’s earlier themes of externally observable traits, such as skull shape (Figure 13.17). In the last 20 or so years, anthropologists have evaluated the level to which human cranial shape diversity reflects the results from genetic markers, such as those used previously to fit against Out-of-Africa models (Relethford 2004) or those used in the apportionment of human diversity between and within groups (Lewontin 1972; Rosenberg et al. 2002). Using larger sample sizes of cranial data collected from thousands of skulls worldwide and a long list of cranial measurements, studies demonstrate a similar decrease in diversity with distance from Africa and show that a majority of cranial variation occurs within populations rather than between populations (Betti et al. 2009; Betti et al. 2010; Manica et al. 2007; Relethford 2001; von Cramon-Taubadel and Lycett 2008). The greatest cranial diversity is found among skulls of sub-Saharan African origin, while the least variation is found among populations inhabiting places like Tierra del Fuego at the southern tip of

Argentina and Chile. While ancient and historical thinkers previously thought “race” categories could reasonably be determined based on skull dimensions, modern-day analyses using more informative sets of cranial traits simply show that migrations out of Africa and the relative distances between populations can explain a majority of worldwide cranial diversity (Betti et al. 2009).




Figure **6.2.11**: Diagram of the bony labyrinth in the inner ear.

This same patterning in phenotypic variation has even been found in studies examining shape variation of the pelvis (Betti et al. 2013; Betti et al. 2014), the teeth (Rathmann et al. 2017), and the human **bony labyrinth** of the ear (Ponce de León et al. 2018) (Figure 13.18). The skeletal morphology of these bones still varies worldwide, but a greater proportion of that variation can still be attributed to the ways in which human populations migrated across the world and exchanged genes with those closer to them rather than those further away. Human skeletal variation in these parts of the body is continuous and non-discrete. Given the important functions of the cranium and these other skeletal parts, we may infer that the genes that underpin their development have been relatively conserved by neutral evolutionary processes such as genetic drift and gene flow. It is also important to note that while some traits such as height, weight, cranial dimensions, and body composition are determined, in part, by genes, the underlying developmental processes behind these traits are underpinned by complex polygenic mechanisms that have led to the continuous spectrum of variation in such variables among modern-day human populations.

Phenotypic Traits That Reflect Natural Selection

Even though 99.9% of our DNA is the same between all humans worldwide, and many traits reflect neutral processes, there are parts of that remaining 0.1% of the human genome that code for individual and regional differences. Similarly to craniometric analyses that have been conducted in recent decades, human variation in skin color has also been reassessed using new methods and in light of greater knowledge of biological evolution.

New technologies allow scientists to use color photometry to sample and quantify the visible wavelength of skin color, in a way 19th- and 20th-century readers could not. In one report, it was found that 87.9% of global skin color variation can be attributed to genetic differences *between* groups, 3.2% to those among local populations within regions, and 8.9% *within* local populations (Relethford 2002). This apportionment differs significantly and is the reverse situation found in the distribution of genetic differences we see when we examine genetic markers such as blood type–related alleles. However, this pattern of human skin color worldwide is not surprising, given that we now understand that past selection has occurred for darker skin near the equator and lighter skin at higher latitudes (Jablonski 2004; Jablonski and Chaplin 2000). While most genetic diversity reflects neutral variation due to population migrations, geographic isolation, and restricted gene flow dynamics, some human genetic/phenotypic diversity is best explained as local adaptation to environmental conditions (i.e., selection). Given that skin color variation is atypical compared to other genetic markers and biological traits, this, in fact, goes against earlier “race” typologies. This is because recent studies ironically show how so much of genetic variation relates to neutral processes, while skin color does not. It follows that skin color *cannot* be viewed as useful in making inferences about other human traits.

On top of social implications, the quantification and interpretation of human variation has important medical and clinical applications (National Research Council Committee on Human Genome Diversity 1997). For instance, large-scale genomic studies sampling from human populations distributed worldwide have produced detailed knowledge on variation in disease resistance or susceptibility between and within populations. If we think about drug companies who develop medicines for African American patients particularly, the genetic diversity in predispositions to disease and/or good health is likely higher among people of African descent than these pharmaceutical businesses have taken into account. Through targeted sampling of various world groups, clinical geneticists may also identify genetic risk factors of certain common disorders such as chronic heart disease, asthma, diabetes, autoimmune diseases, and behavioral disorders. Having an understanding of population-specific biology is crucial in the development of therapies, medicines, and vaccinations, as not all treatments may be suitable for for every human, depending on their genotype. During diagnosis and treatment, it is important to have an evolutionary perspective on gene-environment

relationships in patients. Typological concepts of “race” are not useful, given that most racial groups (whether self-identified or not) popularly recognized lack homogeneity and are, in fact, variable. **Cystic fibrosis**, for instance, occurs in all world populations but can often be underdiagnosed in populations of African ancestry because it is thought of as a “white” disease (Yudell et al. 2016).



Figure **6.2.12**: The Forensic Anthropology Lab at the National Museum of Natural History, Smithsonian Institute, Washington, D.C.

Lastly, assignments of “race” to human remains is a common practice in forensic anthropology, especially in the United States and other worldwide contexts where bones are recovered and associated with criminal investigations (Figure 13.19). Forensic anthropologists have ascribed “race” or ancestry to sets of skeletons thanks to scientific research that has attempted to divide up different human groups into culturally constructed categories based on biologically “discrete” assortments (Sauer 1992). Rather than focusing on the neutral or selective *causes* of human biological variation, the concentration in forensic anthropology centers upon how *probabilistic* it may be to assign bones of certain dimensions to one of several identified racial categories. Forensic anthropologists do not agree with the typological “race” concepts of the past but, instead, root their racial categorization in methods of probability estimation (Sauer 1992). Based on many samples of skeletons from different world regions, statistical tests (such as discriminant function analysis) allow them to distinguish how *likely* certain skeletal dimensions may predict geographic **ancestry**.

It is important to remember that while it is possible to determine geographic origin (or ancestry) based on skull morphology, again, the amount of craniometric distinctiveness required to distinguish whether a cranium belongs to one group or another will make for arbitrary decisions (Relethford 2009). Individuals can vary in their skeletal dimensions by continental origin, country origin, regional origin, sex, age, environmental factors, and the time period in which they lived, making it difficult to assign individuals to particular categories in a completely meaningful way (Ousley et al. 2009). When forensic reports and scientific journal articles give an estimation of ancestry, it is crucial to keep in mind that responsible assignments of ancestry will be done through robust statistical testing and stated as a probability estimate. Today, we also live in a more globalized world where a skeletal individual may have been born originally to parents of two separate traditional racial categories. In contexts of great heterogeneity within populations, this definitely adds difficulty to the work of forensic scientists and anthropologists preparing results for the courtroom.

6.2: Human Variation in Biological Anthropology Today is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **13.2: Human Variation in Biological Anthropology Today** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.3: Talking About Human Biological Variation Going Forward

To conclude, utilizing *races* to describe human biological variation is not accurate or productive. Using a select few hundred genetic loci, or perhaps a number of phenotypic traits, it may be possible to assign individuals to a geographic ancestry. However, what constitutes a bounded genetic or geographical grouping is both arbitrary and potentially harmful owing to ethical and historical reasons (see Chapter 3 for more on the issues with commercial ancestry tests, for example). The discipline of biological anthropology has moved past typological frameworks that shoehorn continuously variable human populations into discrete and socially constructed subsets. Improvements in the number of markers, the genetic technologies used to study variation, and the number of worldwide populations sampled have led to more nuanced understandings of human diversity. It is of utmost importance that scientists and non-scientists, in theory, have each of the following clarified:

1. Today, we refer to different local human groups as "populations." What constitutes a population is carefully defined in scientific reports based on some geographical, linguistic, or cultural criteria and some degree of relativity to other closely or distantly related human groups.
2. Humans have significantly less genetic diversity than other primates and mammals, and all human beings on Earth share 99.9% of their overall DNA. Some of the remaining 0.1% of human variation varies on a clinal or continuous basis, such as can be seen when looking at ABO blood type **polymorphisms** worldwide.
3. Many biological characteristics in humans are actually determined non-concordantly and/or polygenically. Therefore, superiority or inferiority in human behavior or body form cannot justifiably be linked to fixed and innate differences between groups.
4. Genetic distances are correlated with geographic distances among the global human population. This is especially apparent when we consider that genetic diversity is highest in sub-Saharan Africa, and average genetic heterogeneity decreases in populations further away from the African continent in accordance with the migratory history of anatomically modern *Homo sapiens*.
5. The effects of gene flow, genetic drift, and population bottlenecking are reflected in some phenotypic traits, such as cranial shape.
6. Other traits, like skin color and lactase persistence, we recognize to be the product of many millennia of natural selective pressures influencing human biology from the external environment.

When taken altogether, genetic analyses of human diversity do not support 20th-century (or even earlier) concepts of race. In discussions about human diversity, each of these genomic results help clarify for all conversationalists how biological variation is distributed across the human population today. Taking care to think about and debate the nature of human variation is important, because although the effects and events that produced genetic differences among groups occurred in the ancient past, sociocultural concepts about race and ethnicity continue to have real social, economic, and political consequences in the modern era.

Beyond talking about diversity in the university setting, it is important that teachers, researchers, and students of anthropology recognize and assume the responsibility of influencing public perspectives of human diversity. Race-based classification systems were developed during the colonial era, transatlantic slave trade, and Scientific Revolution by some of the earliest scholars whom we may call the first "anthropologists" and students of humankind's variation. Unfortunately, some of their ideas put forward have persisted and evolved into present-day lived realities. Some of today's politicians and socioeconomic bodies have racially charged agendas that promote racism or certain kinds of economic or racial inequalities. As anthropologists, we must acknowledge that while human "races" are not a biological reality, their status as a (misguided) social construction does have real consequences for many people (Antrosio 2011). In other words, while "race" is a sociocultural invention in some people's minds, the treatment different individuals receive due to their perceived "race" can have significant financial, emotional, sociopolitical, and physiological costs. But assuming a "color-blind" position when it comes to the topics of "race" and ethnicity (especially in political discussions) is actually counterproductive since the negative social consequences of modern "race" ideas could be ignored, making it harder to examine and address instances of discrimination properly (Wise 2010). Rather than shy away from these topics, we can use our scientific findings to establish socially relevant and biologically accurate ideas concerning human diversity. Today, research into genetic and phenotypic differentiation among and within various human populations continues to expand in its scope, its technological capabilities, its sample sizes, and its ethical concerns. It is thanks to such scientific work done in the past few

decades that we now have a deeper understanding not only of how humans vary but also of how we are biologically a rather homogenous, intermixing world population.

SPECIAL TOPIC: MY EXPERIENCES AS A MINORITY ACADEMIC OF COLOR



Figure **6.3.1**: Michael B. C. Rivera in Hong Kong.

My name is Michael, and I am a researcher in biological anthropology (Figure 13.20). What strikes me as most interesting to investigate is human biological diversity today and the study of past human evolution. What I am really curious about is how we can use human skeletons to infer how people adapt to coastal environments. Relying on aquatic foods near rivers, lakes, and the sea is interesting because we have found evidence for positive effects of coastal living on dietary health and many unique adaptations in bones and teeth when living near rivers and beaches. I also really enjoy talking to students and non-scientists about our work, through teaching, science communication events, and writing book chapters like this one! I grew up in Hong Kong, a city in southern China. My father is from the Philippines and my mother is from Hong Kong, which makes me a mixed Filipino-Chinese academic. When I attended international schools in my youth, I saw that kids my age came in all shapes, sizes, and colors. It was not until I left Hong Kong that I realized people with my skin tone were somewhat rarer in British universities I attended.

Biological anthropology is not taught extensively back home in Hong Kong, but my initial motivation to enter this field was a great TV show called *Bones*. This TV series was about a brilliant anthropologist who examined human remains for the Smithsonian Institute in Washington, D.C., identified the individuals they belonged to through scientific analyses of bones and teeth, and told the stories of the deceased who could not tell their own. I went to the United Kingdom to earn my bachelor's, master's, and doctorate degrees. During my studies, I was taught about human genetics, apes and monkeys, forensics, human cultural and behavioral diversity, and the story of human evolution that began six million to eight million years ago. It was fascinating to me that we could answer important questions about human variation and history using scientific methods. While I was at university, I did not have many minority academic role models to look up to. Today, I look around and see other academics of color during conferences and perhaps one or two others around the places at which I work. I am inspired by all my colleagues who advocate for greater representation and diversity in universities (whether they are minority academics or not). I admire many of my fellow researchers who are underrepresented and do a great job of representing minority groups through their cutting-edge research and quality teaching at the undergraduate and graduate levels. The study of anthropology has really highlighted for me that we share a

common humanity and history. Being somebody who is “mixed race” and Asian likely played a key role in steering me toward a discipline that studies human diversity. As this chapter hopefully shows, there is a lot about race and ethnicity to discuss in terms of the discipline’s history and current understandings of human biological diversity. Some scientific and technological advancements today are unfortunately misused for reasons to do with money, politics, or the continuation of fairly antiquated ideas. It is my belief, alongside many of my friends and fellow anthropologists, that science should be more about empathy toward all members of our species and contributing to the intellectual and technological nourishment of society. After speaking to many members of the public, as well as my own undergraduate students, I have received lovely messages from other individuals of color expressing thanks and appreciation for my presence and understanding as a fellow minority and mentor figure. This is why anthropology needs more diversity and to make room for more personal routes into the discipline. All paths to anthropology are valuable and valid. I would encourage anyone to study anthropology as it really is a field for understanding and celebrating the intricacies of human diversity.

Review Questions

- How is the genetic variation of the human species distributed worldwide?
- What evolutionary processes are responsible for producing genotypic/phenotypic diversity within and between human populations?
- Should we continue to attribute any value in “race” concepts older than 1950, based on our current understandings of human biological diversity?
- How should we communicate scientific findings about human biological variation more accurately and responsibly to those outside the anthropological discipline?

6.3: Talking About Human Biological Variation Going Forward is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **13.3: Talking About Human Biological Variation Going Forward** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.4: Key Terms

Age of Discovery: A period between the late 1400s and late 1700s when European explorers and ships sailed extensively across the globe in pursuit of new trading routes and territorial conquest.

Ancestry: Biogeographical information about an individual, traced either through the study of an individual's genome, skeletal characteristics, or some other form of forensic/archaeological evidence. Anthropologists carry out probabilistic estimates of ancestry. They attribute sets of human remains to distinctive "ancestral" groups using careful statistical testing and should report ancestry estimations with statistical probability values.

Binomial nomenclature: A system of naming living things developed by Linnaeus in the 1700s using a scientific name made up of two Latin- or Greek-form words, with the first name capitalized and representative of an organism's genus and the second name indicating an organism's species (e.g., *Homo sapiens*, *Australopithecus afarensis*, *Pongo tapanuliensis*, etc.).

Biological anthropology: A subdiscipline of anthropology concerned with the biological origins, ecology, evolution, and diversity of humans and other primates. This term is increasingly preferred to *physical anthropology*, as many in the field now uncomfortably associate this original name (coined by Aleš Hrdlička) with the ways in which questions of human variation were studied in decades past and the sociohistorical context that made anthropology problematic before 1950 (see Warren 2018).

Biological determinism: The erroneous concept that an individual's behavioral characteristics are innate and determined by genes, brain size, or other physiological attributes, and with no influence of social learning or the environment around the individual during development.

Bony labyrinth: A system of interconnected canals within the auditory (ear- or hearing-related) apparatus, located in the inner ear and responsible for balance and the reception of sound waves.

Cline: A gradient of physiological or morphological change in a single character or allele frequency among a group of species across environmental or geographical lines (e.g., skin color varies clinally, as, over many generations, human groups living nearer the equator have adapted to have more skin pigmentation).

Continuous/clinal variation: Variation that exists between individuals and cannot be measured using distinct categories. Instead, differences between individuals within a population in relation to one particular trait are measurable along a smooth, continuous gradient.

Cystic fibrosis: A genetic disorder in which one defective gene causes overproduction and buildup of mucus in the lungs and other bodily organs, most common in northern Europeans (but also in other world populations more rarely).

Ecological niche: The position or status of an organism within its community and/or ecosystem, resulting from the organism's structural and functional adaptations (e.g., bipedalism, omnivory, lactose digestion, etc.).

Essentialism: A belief or view that an entity, organism, or human grouping has a specific set of characteristics that are fundamentally necessary to its being and classification into definitive categories.

Ethnicity: A complex term used commonly in an interchangeable way with the term *race* (see below).

Eugenics: A set of beliefs and practices that involves the controlled selective breeding of human populations with the hope of improving their heritable qualities, especially through surgical procedures like sterilization and legal rulings that affect marriage rights for interracial couples.

Founder effect: See population bottlenecking below.

Gene flow: A neutral (or nonselective) evolutionary process that occurs when genes get shared between populations.

Genetic drift: A neutral evolutionary process in which allele frequencies from generation to generation due to random chance.

Heterogeneity: The quality of being diverse genetically.

Homogenous: The quality of being uniform genetically.

Human diversity/differentiation/variation: Group differences involving variation in biology, physiology, body chemistry, behavior, and culture.

Isolation-by-distance model: A model that predicts a positive relationship between genetic distances and geographical distances between pairs of populations.

Monogenic: Characterized as being controlled by a single gene (or, in other words, one pair of alleles). Sickle cell anemia and cystic fibrosis are examples of disorders that are monogenically caused.

Monogenetic: Pertaining to the idea that the origin of a species is situated in one geographic region or time (as opposed to *polygenetic*).

Mutation: A gene alteration in the DNA sequence of an organism. As a random, neutral evolutionary process that occurs over the course of meiosis and early cell development, gene mutations are possible sources of diversity in any given human gene pool. Genetic mutations that occur in more than 1% of a population are termed *polymorphisms*.

Natural selection: An evolutionary process whereby certain traits are perpetuated through successive generations, likely owing to the advantages they give organisms in terms of chances of survival and/or reproduction.

Non-concordance: The fact of genes or traits not varying with one another and instead being inherited independently.

Othering: In postcolonial anthropology, we now understand "othering" to mean any action by someone or some group that establishes a division between "us" and "them" in relation to other individuals or populations. This could be based on linguistic or cultural differences, and it has largely been based on external characteristics throughout history.

Out-of-Africa model: A model that suggests that all humans originate from one single group of *Homo sapiens* in (sub-Saharan) Africa who lived between 100,000 and 315,000 years ago and who subsequently diverged and migrated to other regions across the globe.

Physical anthropology: See *biological anthropology* above.

Polygenetic: Having many different ancestries, as in older theories about human origins that involved multiple traditional groupings of humans evolving concurrently in different parts of the world before they merged into one species through interbreeding and/or intergroup warfare. These earlier suggestions have now been overwhelmed by insurmountable evidence for a single origin of the human species in Africa (see the "Out-of-Africa model").

Polymorphism: A genetic variant within a population (caused either by a single gene or multiple genes) that occurs at a rate of over 1% among the population. Polymorphisms are responsible for variation in phenotypic traits such as blood type and skin color.

Population: A group of humans living in a particular geographical area, with more local interbreeding within-group than interbreeding with other groups. A limited or restricted amount of gene flow between populations can occur due to geographical, cultural, linguistic, or environmental factors.

Population bottlenecking (or founder effect): An event in which genetic diversity is significantly reduced owing to a sharp reduction in population size. This can occur when environmental disaster strikes or as a result of human activities (e.g., genocides or group migrations). An important example of this loss in genetic variation occurred over the first human migrations out of Africa and into other continental regions.

Prejudice: An unjustified attitude toward an individual or group not based on reason, whether that is positive and showing preference for one group of people over another or negative and resulting in harm or injury to others.

Race: The identification of a group based on a perceived distinctiveness that makes that group more similar to each other than they are to others outside the group. This may be based on cultural differences, genetic parentage, physical characteristics, behavioral attributes, or something arbitrarily and socially constructed. As a social or demographic category, perceptions of "race" can produce effects that have real and serious consequences for different groups of people. This is despite the fact that biological anthropologists and geneticists have demonstrated that all humans are genetically homogenous and that more differences can be found within populations as opposed to between them in the overall apportionment of human biological variation.

Racism: Any action or belief that discriminates against someone based on perceived differences in race or ethnicity, and the characteristics, qualities, or abilities believed to be specific to a race that is inferior to another in some way.

Scientific Revolution: A period between the 1400s and 1600s when substantial shifts occurred in the social, technological, and philosophical sense, when a scientific method based on the collection of empirical evidence through experimentation was

emphasized and inductive reasoning used to test hypotheses and interpret their results.

Typology: An assortment system that relies on the interpretation of qualitative similarities or differences in the study of variation among objects or people. The categorization of cultures or human groups according to “race” was performed with a typological approach in the earliest practice of anthropology, but this practice has since been discredited and abandoned.

6.4: Key Terms is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **13.4: Key Terms** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.5: Human Variation- An Adaptive Significance Approach

Leslie E. Fitzpatrick, Ph.D., Mercyhurst University

Learning Objectives

- Describe how specific patterns of human adaptation are correlated to natural selection processes.
- Summarize the role of solar radiation in variations of human skin tone. In your explanation, include information as to why reduced pigmentation leading to lighter skin colors is advantageous for populations indigenous to northern latitudes.
- Compare and contrast the various genetic mutations present in Tibetan, Andean, and Ethiopian populations that allow them to survive at high altitudes.
- Define the relationship between specific genetic mutations in some human populations and certain infectious diseases, such as the sickle-cell trait mutation and malarial infection.

In the previous chapters of this text, we explored the role of evolutionary forces in human evolution as well as the basics of genetic variation. Within this framework, we now shift our focus toward examining the numerous challenges our species has faced throughout its evolutionary odyssey as well as how we have met those trials. Genetic variability within and between modern populations of humans has been influenced by years of evolutionary forces, most notably natural selection and genetic drift. As early humans left Africa and spread across the globe, they faced numerous challenges related to their new environments. Beyond genetically influenced changes in physiology as a result of evolution, humans have developed lifestyle strategies to cope with and even thrive in a wide range of habitats. The ways populations of humans met such challenges, coupled with their geographic separation throughout the majority of the last two centamillenia, have led to the many forms of adaptation in our species. This chapter focuses on the complexities of modern human variation through the lens of human evolutionary history.

About the Author

Leslie Fitzpatrick, Ph.D.

Mercyhurst University, Lfitzpatrick@mercyhurst.edu



Leslie Fitzpatrick

Leslie Fitzpatrick is an Assistant Professor in the Department of Biological Anthropology and Applied Forensic Sciences at Mercyhurst University in Erie, Pennsylvania. She earned a Ph.D. in Anthropology from the University of Wyoming (2017), an M.A. in Anthropology from Georgia State (2012), and a B.S. in Mechanical Engineering from Georgia Tech (2000). Her primary research focus is the stable-isotope analysis of human remains as a means of interpreting past mobility and diet profiles for both modern and archaeological populations. In addition to her work in the classroom and laboratory, she has worked as a bioarchaeologist at field sites in Germany, Spain, Croatia, Mexico, Peru, and throughout the United States.

References

American Academy of Pediatrics, Task Force on Infant Sleep Position and Sudden Infant Death Syndrome. 2000. “Changing concepts of sudden infant death syndrome: implications for infant sleeping environment and sleep position.” Pediatrics 105: 650–656, 116 (5): 1,245.

Beja-Pereira, Albano, David Caramelli, Carles Lalueza-Fox, Cristiano Vernesi, Nuno Ferrand, Antonella Casoli, Felix Goyache, Luis J. Royo, et. all., 2006. “The origin of European cattle: evidence from modern and ancient DNA.” PNAS 103 (21): 8,113–8,118.

Bigham, Abigail W. 2016. “Genetics of Human Origin and Evolution: High-Altitude Adaptations.” Current Opinion in Genetics & Development 41: 8–13.

Centers for Disease Control and Prevention. 2017. “Data & Statistics on Sickle Cell Disease.” Centers for Disease Control and Prevention website, August 9. https://www.cdc.gov/ncbddd/sicklecell/data.html.

Chaplin, George, and Nina G. Jablonski. 2009. “Vitamin D and the Evolution of Human Depigmentation.” American Journal of Physical Anthropology 139 (4): 451–461.

Crawford, Nicholas G., Derek E. Kelly, Matthew E. B. Hansen, Marcia H. Beltrame, Shaohua Fan, Shanna L. Bowman, Ethan Jewett, et.all., 2017. “Loci associated with skin pigmentation identified in African populations.” Science 358 (6,365): 1–49.

Cyrkalff, Marek, Cecilia P. Sanchez, Nicole Kilian, Curille Bisseye, Jacques Simpore, Friedrich Frischknecht, and Michael Lanzer. 2011. “Hemoglobins S and interfere with actin remodeling in Plasmodium falciparium-infected erythrocytes.” Science 334 (6,060): 1,283–1,286.

Dzialanski, Zbigniew, Michael Barany, Peter Engfeldt, Anders Magnuson Lovisa A. Olsson, and Torbjörn K. Nilsson. 2016. “Lactase persistence versus lactose intolerance: Is there an intermediate phenotype?” Clinical Biochemistry 49 (2,016): 248. 252.

Fernando, Sumadya D., Chaturaka Rodrigo, and Senaka Rajapakse. 2010. “The ‘hidden’ burden of malaria: cognitive impairment following infection.” Malaria Journal 9 (366): 1–11.

Gerbault, Pascale, Anke Liebert, Yuval Itan, Adam Powell, Mathias Currat, Joachim Burger, Dallas M. Swallow, and Mark G. Thomas. 2011. “Evolution of lactase persistence: an example of human niche construction.” Philosophical Transactions of the Royal Society B: Biological Sciences 366 (1,566): 863–877.

Hartman, T. K., S. J. Rogerson, and P. R. Fischer. 2013. “The impact of maternal malaria on newborns.” Annals of Tropical Paediatrics 30 (4): 271–282.

Longo, Dan L., Frédéric B. Piel, Martin H. Steinberg, and David C. Rees. 2017. “Sickle Cell Disease.” The New England Journal of Medicine 376 (16): 1,561–1,573.

Maddux, Scott D., Todd R. Yokley, Bohumil M. Svoma, and Robert G. Franciscus. 2016. “Absolute Humidity and the Human Nose: A Reanalysis of Climate Zones and Their Influence on Nasal Form and Function.” American Journal of Physical Anthropology 161 (2): 309–320.

Meyer, M. C., M. S. Alexander, Z. Wang, D. L. Hoffmann, J. A. Dahl, D. Degering, W. R. Haas and F. Schlütz. 2017. “Permanent human occupation of the central Tibetan Plateau in the Early Holocene.” Science 355 (6,320): 64–67.

Moore, Lorna G, Susan Niermeyer, and Stacy Zamudio. 1998. “Human Adaptation to High Altitude: Regional and Life-Cycle Perspectives.” Yearbook of Physical Anthropology 41: 25–64.

Noback, Marlijn L., Katerina Harvati, and Fred Spoor. 2011. “Climate-Related Variation of the Human Nasal Cavity.” American Journal of Physical Anthropology 145 (4): 599–614.

Peacock, A. J. 1998. “ABC of Oxygen: Oxygen at High Altitude.” BMJ 317 (7,165): 1,063–1,066.

Quillen, Ellen E., Heather L. Norton, Esteban J. Parra, Frida Loza-Durazo, Khai C. Ang, Florin Mircea Illiescu, Laurel N. Pearson, Mark D. Shriver, Tina Lasisi, Omer Gokcumen, Izzy Starr, Yen-Lung Lin, Alicia R. Martin, and Nina G. Jablonksi. 2018. “Shades of complexity: New perspectives on the evolution and genetic architecture of human skin.” American Journal of Physical Anthropology 1–23.

Ranciaro, Alessia, Michael C. Campbell, Jibril B. Hirbo, Wen-Ya Ko, Alain Froment, Paolo Anagnostou, Maritha J. Kotze, Muntaser Ibrahim, Thomas Nyambo, Sabah A. Omar, and Sarah A. Tishkoff. 2014. “Genetic Origins of Lactase Persistence and the Spread of Pastoralism in Africa.” American Journal of Human Genetics 94 (4): 496–510.

Roby, Brianne Barnett, Marsha Finkelstein, Robert J. Tibesar, and James D. Sidman. 2012. “Prevalence of Positional Plagiocephaly in Teens Born after the ‘Back to Sleep’ Campaign.” Otolaryngology—Head and Neck Surgery 146 (5): 823–828.

Shah, Mark B., Darren Braude, Cameron S. Crandall, Heemun Kwack, Lisa Rabinowitz, Thomas A. Cumbo, Buddha Basnyat, and Govin Bhasyall. 2006. “Changes in Metabolic and Hematologic Laboratory Values With Ascent to Altitude and the Development of

Acute Mountain Sickness in Nepalese Pilgrims.” Wilderness and Environmental Medicine 17 (3): 171–177.

Sherman, Paul W., and Jennifer Billing. 1999. “Darwinian Gastronomy: Why We Use Spices.” BioScience 49 (6): 453–463.

Snoddy, Anne Marie E., Hallie R. Buckley, and Siân E. Halcrow. 2016. “More than Metabolic: Considering the Broader Paleoepidemiological Impact of Vitamin D Deficiency in Bioarchaeology.” American Journal of Physical Anthropology 160 (2): 183–196.Sturm, Richard A., and David L. Duffy. 2012. “Human Pigmentation Genes Under Environmental Selection.” Genome Biology 13 (248): 1–15.

Tishkoff, Sarah A., Floyd A. Reed, Alessia Ranciaro, Benjamin F. Voight, Courtney C. Babbitt, Jesse S. Silverman, Kweli Powell, Holly M. Mortensen, Jibril B. Hirbo, Maha Osman, Muntaser Ibrahim, Sabah A. Omar, Godfrey Lema, Thomas B. Nyambo, Jilur Ghori, Suzannah Bumpstead, Jonathan K. Pritchard, Gregory A. Wray, and Panos Deloukas. 2007. “Convergent adaptation of human lactase persistence in Africa and Europe.” Nature Genetics 39 (1): 31–40.

Vukić, Maja, Antonio Neme, Sabine Seuter, Noora Saksa, Vanessa D. F. de Mello, Tarji Nurmi, Matti Uusitupa, Tomi-Pekka Tuomainen, Jyrki K. Virtanen, and Carsten Carlberg. 2015. “Relevance of Vitamin D Receptor Target Genes for Monitoring the Vitamin D Responsiveness of Primary Human Cells.” PLoS One, San Francisco 10 (4): 1–15.

World Health Organization. 2017. “Key points: World malaria report 2017.” World Health Organization website, November 29. http://www.who.int/malaria/media/world-malaria-report-2017/en/.

Figure Attributions

Figure 14.1 Deep water diver by Leslie E. Fitzpatrick is under a CC BY-NC 4.0 License.

Figure 14.2 Mechanisms of heat transfer original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.3 Body heat maintenance in cold and warm original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.4 Vasoconstriction and vasodilation original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.5 Atmospheric pressure original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.6 Premature infant by Leslie E. Fitzpatrick is under a CC BY-NC 4.0 License.

Figure 14.7 Quechua Woman in Peru by Alexander Fiebrandt (Alecconnell at de.Wikipedia) is under a CC BY-SA 2.0 DE License.

Figure 14.8 Плагиоцефалия (Plagiocephaly) by Medical advises at larece.ru/?p=27115 is under a CC BY-SA 3.0 License.

Figure 14.9 World Map of HVR adaptation in high altitude populations by Chkuu is used under a CC BY-SA 4.0 License.

Figure 14.10 Skin Pigmentation (Anatomy and Physiology, Figure 5.8) by OpenStax is used under a CC BY 4.0 License.

Figure 14.11 Penetration of skin layers by UVA and UVB rays a derivative work original to Explorations: An Open Invitation to Biological Anthropology by Katie Nelson is under a CC BY-NC 4.0 License. [Includes Skin Anatomy by NIH National Cancer Institute, public domain].

Figure 14.12 Evolutionary basis for human skin color variation original to Explorations: An Open Invitation to Biological Anthropology by Katie Nelson is under a CC BY-NC 4.0 License.

Figure 14.13 Rachitis, stages of development for children (slide numbers 7181 and 7182; photo number: M0003399) by Wellcome Collection is under a CC BY 4.0 license.

Figure 14.14 Eskimo Family NGM-v31-p564 by George R. King [original from National Geographic Magazine, Volume 31 (1917)] is in the public domain.

Figure 14.15a COLLECTIE TROPENMUSEUM Fulani vrouwen en kinderen putten water uit de waterput van Santaba TMnr 20010199 (Fulani women and children draw water from the Santaba water well) by Tropenmuseum, part of the National Museum of World Cultures is used under a CC BY-SA 3.0 License.

Figure 14.15b Greenland 1999 (01) by vadeve has been designated to the public domain (CC0).

Figure 14.16a Bergmann's Rule original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.16b Allen's Rule original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.17 Human nasal morphological variation original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 14.18 Malaria parasite life cycle-NIAID by NIH National Institute of Allergy and Infectious Diseases is in the public domain.

Figure 14.19 Sickle cell 01 by The National Heart, Lung, and Blood Institute (NHLBI) is in the public domain.

Figure 14.20 Red Blood Cell abnormalities by Armando Moreno Vranich has been designated to the public domain (CC0).

Figure 14.21 Sickle cell disease a derivative work original to Explorations: An Open Invitation to Biological Anthropology by Katie Nelson is under a CC BY-NC 4.0 License. [Includes two illustrations by Mary Nelson; Sickle cell anemia by Pkleong at English Wikibooks, public domain (CC0), modified (labels removed background erased).]

Figure 14.22 Lactose tolerance in the Old World by Joe Roe is used under a CC BY 4.0 License.

6.5: Human Variation- An Adaptive Significance Approach is shared under a CC BY-NC license and was authored, remixed, and/or curated by LibreTexts.

- **14: Human Variation- An Adaptive Significance Approach** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.5.1: Stress and Homeostasis

All organisms, including humans, must maintain a baseline of normal functions within their cells, tissues, and organs to survive. This constancy of internal functions is referred to as **homeostasis**; however, homeostatic regulation may be challenged by a variety of both external and internal factors. These stimuli are referred to as **stressors**, exposure to which leads to a period in which there is a potential for the disruption of homeostasis. Within limits, all organisms have evolved certain physiological mechanisms to respond to stressors in an effort to maintain homeostasis. For example, some organisms, such as dogs, will develop a thicker coat of fur during cooler periods and they will shed this additional fur during warmer periods. This cyclical fur development linked to seasonal weather changes is but one way that these organisms maintain their homeostasis. The range of changes in the physiology (function), morphology (form), and/or behavior of organisms in response to their environments and the potential stressors of those environments is regulated by its **phenotypic plasticity**. A special case of phenotypic plasticity, **polyphenism**, is broadly defined as the ability of a single genotype within the organism to produce multiple phenotypes when exposed to different environmental conditions or stressors. Changes to the organism's physiology, morphology, and/or behavior that are linked to its underlying phenotypic plasticity may result in changes that are merely temporary or those that are permanent. With respect to human phenotypic plasticity as well as evolutionary history, there are several primary mechanisms that have led to variations among individuals and between populations. These mechanisms, which are referred to as **adjustments** (behavioral, acclimatory, and developmental) and **adaptations**, are explained in detail in the following sections.

6.5.1: Stress and Homeostasis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **14.1: Stress and Homeostasis** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.5.2: Adjustments and Adaptations

Adjustments

The term “adjustment” refers to an organism’s non-genetic way of coping with the stressors of its environment. Although adjustments themselves are non-genetic in nature, the ability of an organism to experience or develop an adjustment is based in its phenotypic plasticity, which is linked to its evolutionarily guided genetic potential. Adjustments occur exclusively on the individual level. As such, different individuals within a population may experience a wide range of possible adjustments in response to a similar stressor. In general, the three main forms of adjustment are: behavioral, acclimatory, and developmental.

Behavioral Adjustments



Figure 6.5.2.1: Notice the lack of full spectrum color in this photo of a deep-water diver and the diver’s use of specialized equipment, such as a breathing apparatus to deliver gases for respiration, bodysuit to ensure thermal regulation, and flashlight to increase visibility in the low-light setting.

When you are cold, do you reach for a blanket? When you are warm, do you seek out shelter cooled by an air-conditioning system? If so, you have likely been influenced to do so by the culture in which you were raised. As noted earlier in the text, the term “culture” refers to a collection of shared, learned behaviors among individuals within a discrete population. **Behavioral adjustments** are regarded as cultural responses to environmental stressors. These adjustments are temporary in nature and, since they are nongenetic, must be constantly altered to meet novel situations posed by the environment. For example, through the use of a specialized mixture of gases for breathing, an apparatus for the delivery of the gases, protective clothing, and gear to increase visibility, divers are able to reach extraordinary depths (in excess of 300 meters below the surface) within the water. The deeper a diver descends, the more atmospheric pressure the diver experiences resulting in increased levels of potentially toxic byproducts of respiration within the body. In addition, with increased depth there is a decrease in the ambient temperature of the water as well as a decrease in the availability of light within the visible spectrum. Deep-water divers are well-versed in the environmental stressors of open waters and employ a variety of strategies based on behavioral adjustments to meet such demands. From wearing protective clothing to assist with maintaining the body’s core temperature to waiting at a specific depth for a prescribed period of time to facilitate the expulsion from the body of nitrogen gas that may have accumulated within the bloodstream, divers employ numerous behavioral adjustments to ensure their safety dive after dive (Figure 14.1). Without these culturally mediated behavioral adjustments, a deep-water diver’s first dive would be their last.

In many developing countries, the use of refrigeration for the storage of perishable food products is uncommon; therefore, individuals within these cultures have developed a variety of behavioral adjustment strategies related to food preparation to address the possibility of food spoilage. Through a cross-cultural analysis of spice use in recipes, Sherman and Billing (1999) determined that cultures closest to the equator, where temperatures are hotter, tend to use both a greater number and a wider variety of plant-based spices with bacteria-inhibiting phytochemical properties (e.g., garlic and onion) in their recipes than cultures located further from the equator. The antimicrobial properties of the spices permits the consumption of foods, particularly animal-based protein sources, for a period of time beyond that which would be considered safe. There are some acclimatory adjustment benefits to the use of some pungent spices as well, which are explored in the following section.

Acclimatory adjustments are temporary, reversible changes in an organism's physiology in response to environmental stressors. Although they are regarded as non-genetic in nature, the range of acclimatory adjustments an organism is capable of producing to accommodate a given stressor is linked to its underlying phenotypic plasticity and duration and severity of the stressor.




Figure **6.5.2.2**: Various thermodynamic mechanisms related to heat gain and loss in the human body.

Before we discuss how varying ambient temperatures affect the human body, we must detail the thermodynamic mechanisms through which heat may be gained or lost. There are four pathways for the loss of heat within the human body: conduction, convection, evaporation, and radiation (Figure 14.2).

Through **conduction** processes, heat will move from a warmer body to a cooler one through direct contact. An example of this is when you accidentally touch a hot cooktop with your hand and the heat is transferred from the cooktop to your skin.

With **convection**, when a warm body is surrounded by a cooler fluid (e.g. air or water) heat will be transferred from the warmer body to the cooler fluid. This is why we will often employ the behavioral adjustment of wearing multiple layers of clothing during the winter in an effort to prevent heat loss to the cooler atmosphere. Conversely, if your body temperature is cooler than that of the air surrounding you, your body will absorb heat.

Depending on your physical condition, most people will begin to sweat around 37.2℃ to 37.7℃ (98.9℉ – 99.9℉). Sweating is an example of **evaporation**, which occurs when a liquid, such as the water within our bodies, is converted to a gas. Phase conversions, such as those underlying the evaporative processes of transforming liquids to gases, require energy. In evaporation, this energy is in the form of heat, and the effect is to cool the body.

The final mechanism for heat loss within the human body is **radiation**, through which energy in the form of electromagnetic waves is produced at a wavelength that typically lies below that which is visible to the human eye. Although humans gain and lose heat from their bodies through radiation, this form of heat transfer is not visible. Humans are capable of losing and gaining heat through conduction, convection, and radiation; however, heat may not be gained through evaporation.

As the ambient temperature decreases, it becomes increasingly difficult for the human body to regulate its core temperature, which is central to the maintenance of homeostasis. The **hypothalamus** is a small portion of the human brain located near its base. It is responsible for numerous functions, including the regulation of body temperature. As measured orally, normal human body temperature averages 37 ℃ (98.6°F). When an individual's body temperature falls below 34.4 ℃ (93.9°F), the hypothalamus becomes impaired leading to issues with body temperature control. A total loss of the ability to regulate body temperature occurs around 29.4 ℃ (84.9°F), which may result in death. When the ambient temperature (e.g. air temperature) falls below the critical temperature of 31 ℃ (87.8°F), a nude human body that is at rest will respond with a series of physiological changes to preserve homeostasis (Figure 14.3).




Figure **6.5.2.3**: Example of overall body heat maintenance in cold and warm ambient environments.

In response to colder temperatures, the human body experiences two main types of physiological responses: those that increase the production of heat within the body and those that seek to retain the body's heat. The production of heat within the body is accomplished through short-term increases in the body's basal metabolic rate. This rate is a measure of the energy required to maintain necessary body processes while the body is in a resting state. As the body's basal metabolic rate increases, an individual must consume greater quantities of energy-providing nutrients to maintain the increase. Of course, such increases may not continue forever as they are energetically expensive. As with all acclimatory adjustments, an increase in the basal metabolic rate is merely temporary.

Another form of a temporary heat-generating acclimatory adjustment to cold stress is the physiological response of shivering. Shivering results when the hypothalamus stimulates increased muscular activity that leads to an elevation of the muscular metabolism. Much like the increased muscular metabolism that occurs during periods of strenuous exercise, the elevation of muscular metabolism rates during shivering leads to higher rates of body heat generation.

Other physiological mechanisms the body uses to assist with the maintenance of temperature related to homeostasis involve the preservation of heat already contained within the body. Of these mechanisms, the most notable is the constriction of peripheral capillaries in the skin through a process called **vasoconstriction**. The decreased surface area of the capillaries through vasoconstriction results in less heat reaching the surface of the skin where it would be dissipated into the atmosphere. In addition, vasoconstriction leads to the maintenance of heat near the core of the body where the vital organs are located. This is one of the reasons that an individual may experience cold-related injuries, such as frost-bite leading to tissue necrosis (tissue death) in regions of the body that are most distant from the core (e.g. fingers, toes, nose, ears, cheeks, chin, etc.).

Just as cold stress presents challenges to maintaining homeostasis within the body with respect to temperature, heat does as well. In hot climates, the body will begin to absorb extra heat from its surroundings (through conduction, convection, and radiation) resulting in potential heat-related disorders, such as heat exhaustion. When the human body is exposed to ambient temperatures above 35 ℃ (95°F), excess body heat will be lost primarily through evaporative processes, specifically through sweating. All humans, regardless of their environment, have approximately the same number of sweat glands within their bodies. Over time, individuals living in hot, arid environments will develop more sensitive forms of sweat glands resulting in the production of greater quantities of sweat. In an effort to prevent dehydration due to this form of acclimatory adjustment, there will be an additional reduction in the volume of urine produced by the individual.

As noted in the previous section, some cultural groups, particularly those in equatorial regions, add pungent spices to their foods to inhibit the colonization of bacteria (Sherman and Billing 1999). Although the addition of spices to foods to decrease spoilage rates is a behavioral adjustment, the application of some forms of peppers triggers an acclimatory adjustment process as well. Compounds referred to as capsaicinoids are the secondary byproducts of chili pepper plants' metabolism. Capsaicinoids are produced by the plants to deter their consumption by some forms of fungi and mammals. When mammals, such as humans, consume the capsaicinoids from chili peppers, a burning sensation may occur within their mouths and along their digestive tracts. This burning sensation is the result of the activation of capsaicin receptors along the body's nerve pathways. Although the peppers themselves may be at ambient temperature so their consumption is technically not causing any form of body temperature increase, the human body perceives the pepper as elevating its core temperature due to the activation of the capsaicin receptors. Even though the body's temperature has not actually been elevated due to the consumption of capsaicin, the hypothalamus will react as if it has, which leads to the initiation of sweating processes in an attempt to lower body temperature and maintain homeostasis. The increased piquancy (application of pungent spices to food) as a means of inhibiting food-borne bacterial colonization in warm climates, as well as spices' ability to trigger sweating processes as a method for cooling the body, is an example of the intersection between behavioral and acclimatory adjustments that may be utilized by individuals within certain populations.

In addition to increased sweat production in the body as a means of regulating internal body temperature to maintain homeostasis, **vasodilation** may occur (Figure 14.4). Vasodilation occurs when there is an expansion of the capillaries within the skin leading to a more effective transfer of heat from within the body to the exterior to allow conductive, convective, radiative, and evaporative (sweating) processes to occur.




Figure **6.5.2.4**: This image features an illustration of the vasoconstriction processes that occur within the peripheral vascular system when an individual is exposed to cold ambient temperatures and the vasodilation that occurs in warmer environments.

Physiologically-based acclimatory adjustments to hot, dry climates may be complemented by behavioral adjustments as well. For example, individuals in such climates may limit their physical activity during the times of day when the temperature is typically the hottest. Additionally, these individuals may wear loose fitting clothing that covers much of their skin. The looseness of the clothing allows for air to flow between the clothing and the skin to permit the effective evaporation of sweat. Although it may seem counterintuitive to cover one's body completely in a hot climate, the covering of the skin keeps the sun's rays from directly penetrating the skin and elevating the body's core temperature.

Acclimatory Adjustments: Altitudinal Stressors

The challenges posed by thermal conditions are but one form of environmental stressor humans must face. High-altitude environments, which are defined as altitudes in excess of 2400 meters above sea level (m.a.s.l.) or 7874 feet above sea level (f.a.s.l.), pose additional challenges to the maintenance of homeostasis in humans. Some of the main stressors encountered by those

living within high-altitude environments include: decreased oxygen availability, cold temperatures, low humidity, high wind speed, a reduced nutritional base, and increased solar radiation levels. Of these challenges, the most significant is the decreased availability of oxygen.




Figure **6.5.2.5**: As altitude increases, atmospheric pressure decreases, which allows for more space between air molecules.

To visualize how altitude affects the availability of oxygen, imagine two balloons that are each filled with the same quantity of oxygen molecules. One of these balloons is positioned at sea-level and the other is placed high upon a mountain peak. For the balloon at sea level, there is more atmospheric pressure pressing down on the molecules within this balloon. This leads to the oxygen molecules within the sea level balloon being forced into a more compact organization. In contrast, the mountain peak balloon has less atmospheric pressure pressing down on it. This leads to the oxygen molecules within that balloon spreading out from each other since they are not being forced together quite as strongly. This example highlights the availability of oxygen molecules in each breath than we take in low- versus high-altitude environments. At 5,500 m.a.s.l. (approximately 18,000 f.a.s.l.), the atmospheric pressure is approximately 50% of its value at sea level (Peacock 1998). At the peak of Mount Everest (8,900 m.a.s.l. or approximately 29,200 f.a.s.l.), the atmospheric pressure is equivalent to only about 30% of their sea level amounts (Peacock 1998) (Figure 14.5).

Due to the decreased availability of oxygen at higher altitudes, certain acclimatory adjustments are required to ensure the maintenance of homeostasis for individuals other than those who were gestated, born, and raised at high altitude. For these people, their rate of breathing will increase to permit greater quantities of air containing oxygen into the lungs when they ascend into higher altitude environments. An increased speed and depth of breathing, which is referred to as **hyperpnea**, is not sustainable indefinitely; thus, the rate of breathing begins to decrease as the person becomes acclimatized to the altitude. During the initial phases of high-altitude-related hyperpnea, the heart will begin to beat faster but the stroke volume (the amount of blood pushed through during each beat) will decrease slightly. In addition, the body will divert energy from non-critical bodily functions, such as digestive processes.

Once the atmospheric oxygen reaches the alveoli (small air sacs) in the lungs, it diffuses (spreads) across the alveolar membrane and enters **erythrocytes** (red blood cells). When the oxygen reaches the erythrocytes, it will loosely bind with hemoglobin, which is an iron-rich protein. It is within the alveoli that the oxygen combines with the hemoglobin. When the erythrocytes carrying the hemoglobin-binded oxygen molecules reach the capillaries where the partial pressure of oxygen is relatively low, the oxygen will be released by the hemoglobin so that it is free for diffusion into body cells. High-altitude-related hyperpnea leads to an increase in the pH of the blood, which makes the blood more alkaline. An increase in the alkalinity of blood is directly related to the partial pressures of oxygen and carbon dioxide in the bloodstream (Shah et al. 2006).

Similar to the acclimatory adjustments related to thermal conditions (e.g., shivering or sweating), those related to high altitude may not be sustained permanently due to their energetically expensive nature. Over a period of days or weeks, the human body will

begin to compensate for the increases in respiration as well as pH through the urinary excretion of bicarbonate (a metabolic byproduct). In addition, to assist with the transportation of oxygen in the absence of decreased respiration, there will be an increase in the following: **hematocrit** (percentage volume of erythrocytes in the blood), myoglobin (oxygen- and iron-binding protein in muscle tissues), red blood cell mass, pulmonary artery pressure, and quantity of capillaries in the skeletal muscular tissue.



Figure **6.5.2.6**: Premature infant born at 30 weeks, 4 days gestation to a mother with altitudinal-induced preeclampsia. Blue light assists the infant's liver with processing high levels of bilirubin.

Although the long-term acclimatory adjustments that an individual from low altitude experiences in a high-altitude environment may permit them to reside there successfully, reproduction within such settings is frequently complicated. With increased altitude comes an increased risk of miscarriage, lower birth weights, and higher infant mortality rates. As the mother's body seeks to preserve its own homeostasis when faced with challenges related to high-altitude living, there is often a decreased rate and volume of blood flow to the uterus as compared to a pregnant woman of similar physiological condition at a lower altitude (Moore et al. 1998). Of course, a decreased rate and volume of uterine blood flow results in a decrease in the amount of oxygen that will be passed through the uterus and placenta to the developing fetus. In addition, women who experience pregnancy at higher altitudes are more prone to developing preeclampsia (severe elevation of blood pressure), which is linked to increased rates of both fetal and maternal death (Moore et al. 1998) (Figure 14.6).

Developmental Adjustments

Although there are often significant challenges for a mother gestating a fetus at high altitude, individuals who are native to such environments experience a form of adjustment referred to as a developmental adjustment. Developmental adjustments occur only in individuals who spent their developmental period (i.e. childhood and adolescence) within a high-altitude environment and they do not apply to those who moved into these environments in the post-developmental (i.e., adult) phase. Furthermore, the degree of developmental adjustment within an individual is directly related to their underlying phenotypic plasticity (polyphenism) as well as the amount of time during the crucial growth and development period the individual resides within the challenging environment. Although humans have the remarkable capacity to develop and survive within environments that are not overly conducive to the successful maintenance of homeostasis, there are definitely physiological costs associated with this ability.



Figure **6.5.2.7**: Quechua woman from high-altitude region of the Peruvian Andes.

In general, high-altitude natives tend to grow more slowly and physically mature later than their low-altitude counterparts (Figure 14.7). Decreased growth and maturity rates are linked not only to the increased physiological demands placed on the body due to

the decreased partial pressure of oxygen but to decreases in the quality of the nutritional base at higher altitudes. Increased terrain complexity, elevated solar radiation levels, and higher wind speeds coupled with decreased temperatures and lower humidity levels at higher altitudes leads to difficulties with growing and maintaining crops and raising livestock. Overall, there is a decreased quality in the available nutritional base as altitude increases, which is correlated to a lack of the nutrients necessary to ensure proper physiological growth and development in humans. Thus, even though individuals may be able to develop and grow within high-altitude environments, they may not reach their full genetically mediated growth potential as they would in a lower-altitude environment.

The heart and lung capacity of individuals who are lifelong residents of high-altitude environments is larger than for individuals from lower-altitude regions. In addition, the high-altitude individuals are more efficient than those from low altitudes at diffusing oxygen from the bloodstream to the body's various tissues. Of course, as noted previously, the developmental adjustments that an individual may experience are defined by their underlying genetic composition (phenotypic plasticity) and researchers have begun to discover some of the genetic factors that appear to be unique to certain populations of humans who have resided in high-altitude environments for significant periods of time. The time frame necessary for natural selection processes to act upon the underlying genetic composition of a population is on the order of many millennia; thus, the physiological changes to indigenous high-altitude populations on a whole are best described in the following section on adaptations.

Not all developmental adjustments are linked to environmental pressures such as climate or altitude; rather, some of these adjustments are correlated to sociocultural or behavioral practices. As noted earlier, some behavioral adjustments permit individuals to maintain homeostasis in challenging environments (e.g., wearing heavy clothing layers to maintain body temperature in cold weather). Other behavioral adjustments, many of which are based on sociocultural principles, may affect the physiological appearance of an individual when they are practiced consistently during the development and growth phases.

Sudden infant death syndrome (SIDS) has no definitive cause; however, the American Academy of Pediatrics published a report in 1992 linking SIDS to infants (under the age of one) sleeping on their stomachs. The "Back to Sleep" campaign championed by the American Academy of Pediatrics helped educate members of the medical community as well as the public that the best sleep position for infants is on their backs. Per the American Academy of Pediatrics Task Force on Infant Sleep Position and Sudden Infant Death Syndrome (2000: 1,245), between 1992 and 2000 the frequency of infants being placed on their stomachs to sleep decreased from more than 70% to less than 20% and the SIDS rate decreased by more than 40%.



Figure **6.5.2.8**: Infant with positional plagiocephaly. Notice the irregular shape of the posterior (back) of the skull.

Placing infants on their backs to sleep has led to decreased infant mortality (death) rates due to SIDS; however, it has led to an unintended consequence: infant cranial deformation. The cranial deformations experienced by infants who sleep solely on their back tend to manifest in one of two forms: brachycephaly and plagiocephaly (Roby et al. 2012). With positional brachycephaly, the back of the infant's head appears rather uniformly flattened due to repetitive contact with a flat surface, such as a crib mattress or car seat back. In cases of positional plagiocephaly, the back of the infant's head appears asymmetrically flattened (Figure 14.8). This asymmetry is typically due to an uneven distribution of mechanical forces resulting from the manner in which the infant's head is in contact with a flat surface. The forms of cranial deformation resulting from sleep positioning do not affect the infant's brain development. For many individuals, the appearance of the deformation is minimized during later development. Still, some individuals will maintain the pattern of cranial deformation acquired during their infancy throughout their lives. The unintentional cranial deformation resulting from placing infants on their backs to sleep as a means of preventing SIDS-related deaths is a physiological indicator of a behavioral adjustment.

Adaptations

As we have just explored, survival and reproduction at high altitudes present numerous physiological challenges for most humans, but what if there were some humans who were specially adapted to life at high altitudes? The behavioral, acclimatory, and developmental adjustments discussed above are all related to the phenotypic plasticity of the individual; however, most adjustments are temporary in nature and they affect a single individual rather than all individuals within a population. But, what if the physiological changes were permanent? What if they affected all members of a population rather than just a single individual? The long-term, micro-evolutionary (i.e., genetic) changes that occur within a population in response to an environmental stressor are referred to as an **adaptation**. From an evolutionary standpoint, the term "adaptation" refers to a phenotypic trait (i.e., physiological/morphological feature or behavior) that has been acted upon by natural selection processes to increase a species' ability to survive and reproduce within a specific environment. Within the field of physiology, the term "adaptation" refers to traits that serve to restore homeostasis. The physiology-based interpretation of adaptations presumes that all traits serve a purpose and that all adaptations are beneficial in nature; however, this may be a fallacy, since some traits may be present without clear evidence as to their purpose. As such, during the following discussion of various forms of adaptations in human populations, we will focus our attention to phenotypic traits with an evidence-based purpose.

Adaptation: Altitudinal Adaptation

As mentioned in the previous section, there is genomic research supporting the evolutionary selection of certain phenotypes and their corresponding genotypes within indigenous high-altitude populations across the globe. The following discussion focuses on three high-altitude indigenous populations from Tibet, the Andes, and Ethiopia (Figure 14.9). Although these populations share many common genetic traits based on relatively similar evolutionary histories influenced by similar environmental stressors, there is support for local genetically based adaptation as well, based on different genes being acted upon by environmental stressors that may be unique to Tibet, the Andes, and Ethiopia (Bigham 2016).



Figure **6.5.2.10**: Highlighted regions feature (from left to right) the Andean, Simian (Ethiopian), and Tibetan Plateau high-altitude regions.

Tibetan populations have resided in the Tibetan Plateau and Himalayan Mountain regions at elevations exceeding 4,000 m.a.s.l. (13,100 f.a.s.l.) for at least the past 7,400 years (Meyer et al. 2017). A gene referred to as *EPAS1* is involved in the regulation of red blood cell (and hemoglobin) production as well as catecholamine homeostasis. Catecholamines are hormones (e.g., epinephrine) secreted as part of the sympathetic nervous system's acute stress ("fight-or-flight") response. An acute stress response typically includes an increase in: heart rate, blood pressure, and blood glucose levels. Long-term elevation of catecholamines in the body may lead to hypertension (elevated blood pressure), increased blood pH levels, the development of a form of cardiovascular disease leading to narrowing of the arteries (atherosclerosis), and blood clots. In the short term, an acclimatory adjustment leading to the increase of catecholamines and hemoglobin production by the hypoxia (low-oxygen level)-induced activation of the *EPAS1* gene may assist individuals from lower altitudes as they ascend to significantly higher altitudes; however, such increases may not be maintained for long before they cause damage to the body. For indigenous high-altitude populations of Tibet, a mutation in the *EPAS1* gene inhibits increased red blood cell production and assists with catecholamine regulation. The red blood cell count of high-altitude Tibetans with the *EPAS1* point mutation is about the same as for individuals residing at sea level.

Interestingly, individuals in populations from the high-altitude Andean Altiplano of Peru and Bolivia, such as the Quechua and Aymara, lack the *EPAS1* point mutation, so their red blood cell counts are relatively elevated compared with the Tibetan populations. In addition, populations from the Andean Altiplano also have an increased arterial oxygen saturation and low, hypoxia-

induced breathing pattern (minor hyperventilation) as compared to their Tibetan counterparts (Bigham 2016). Populations indigenous to the Semien Plateau of Ethiopia, such as the Oromo and Amhara, share a similar but not identical *EPAS1* point mutation with the Tibetan population (Bigham 2016). This suggests that the *EPAS1* mutations occurred independently from each other; however, their effects are still similar in that they permit the Tibetan and Ethiopian populations to survive at high altitudes. Not all adaptations are related to life in high-altitude environments, however. In the following sections, we will address two more general examples of adaptation in human populations: variations in skin color and differences in body build.

Adaptation: Skin Tone *Basics*

When you think about your own skin tone and compare it to members of your family, do you all possess exactly the same shade? Are some members of your family darker than others? What about your friends? Your classmates? Skin tone occurs along a continuum, which is a reflection of the complex evolutionary history of our species. The expression of skin tone is regulated primarily by both melanin and hemoglobin. **Melanin** is a dark brown-black pigment that is produced by the oxidation of certain amino acids (e.g., tyrosine, cysteine, phenylalanine) in the melanocytes. **Melanocytes** are specialized cells located in the base layer (stratum basale) of the skin's epidermis as well as several other areas within the body (Figure 14.10). Within the melanocytes, melanin is produced in the special organelle called a melanosome. Melanosomes serve as sites for the synthesis, storage, and transportation of melanin. Melanosomes transport the melanin particles through cellular projections to epidermal skin cells (keratinocytes) as well as to the base of the growing hair root (root sheath portion). In the eye, however, melanin particles produced by the melanosomes remain present within the iris (iridial melanocytes) and are not transported beyond their origin location. The two main forms of melanin related to skin, hair, and eye color are eumelanin and pheomelanin. All humans contain both eumelanin and pheomelanin within their bodies; however, the relative expression of these two forms of melanin determines an individual's overall coloring. Eumelanin is a brown-to-black colored melanin particle and pheomelanin is more pink-to-red. Individuals with darker skin or hair color have a greater expression of eumelanin than those with lighter-colored skin and blonde or red hair.




Figure **6.5.2.11**: Diagram featuring the relative numbers of melanocytes and melanosomes in light and dark shades of skin tone.

SPECIAL TOPIC: SKIN TONE GENETIC REGULATION

The melanocortin 1 receptor (*MC1R*) gene acts to control which types of melanin (eumelanin or pheomelanin) are produced by melanocytes. The *MC1R* receptor is located on the surface of the melanocyte cells (Quillen et al. 2018). Activation of the *MC1R* receptors may occur through exposure to specific environmental stimuli or due to underlying genetic processes. Inactive or blocked *MC1R* receptors results in melanocytes producing pheomelanin. If the *MC1R* gene receptors are activated, then the melanocytes will produce eumelanin. Thus, individuals with activated *MC1R* receptors tend to have darker pigmented skin and hair than individuals with inactive or blocked receptors.

The alleles of another gene, the major facilitator, superfamily domain-containing protein 12 (*MFSD12*) gene, affect the expression of melanocytes in a different way than the *MC1R* gene. Instead of affecting the activation of melanocyte receptors, the *MFSD12* alleles indirectly affect the membranes of melanocyte lysosomes (Quillen et al. 2018). The melanocyte's lysosomes are organelles containing digestive enzymes, which ultimately correlate to varying degrees of pigmentation in humans. Variations in the membranes of the melanocyte lysosomes ultimately correlate to differing degrees of pigmentation in humans.

Ancestral *MFSD12* allele variants are present in European and East Asian populations and are associated with lighter pigmentation of the skin (Crawford et al. 2017; Quillen et al. 2018). In addition, this ancestral variant is also associated with Tanzanian, San, and

Ethiopian populations of Afro-Asiatic ancestry (Crawford et al. 2017; Quillen et al. 2018). In contrast, the more-derived (i.e. more recent) allele variants that are linked to darker skin tones are more commonly present in East African populations, particularly those of Nilo-Saharan descent (Crawford et al. 2017; Quillen et al. 2018). The notion that ancestral alleles of *MFSD12* are associated with lighter skin pigmentation is in opposition to the commonly accepted idea that our pigmentation was likely darker throughout early human evolution (Crawford et al. 2017; Quillen et al. 2018). Due to the complexity of the human genome, *MFSD12* and *MC1R* are but two examples of alleles affecting human skin tone. Furthermore, there is genetic evidence suggesting that certain genomic variants associated with both darker and lighter skin color have been subject to directional selection processes for as long as 600,000 years, which far exceeds the evolutionary span of *Homo sapiens sapiens* (Crawford et al. 2017; Quillen et al. 2018). So, evolutionary processes may lead to skin becoming more darkly pigmented as well as more lightly pigmented.

Adaptation: *Melanogenesis*




Figure **6.5.2.13**: Penetration of skin layers by UVA and UVB rays.

Although all humans have approximately the same number of melanocytes within the epidermis, the production of melanin by these melanocytes varies. There are two forms of melanogenesis (the process through which melanocytes generate melanin): basal and activated. As discussed previously, the expression of eumelanin and pheomelanin by the melanocytes is genetically regulated through the expression of specific receptors (e.g., *MC1R*) or other melanocyte components (e.g., *MFSD12*). **Basal melanogenesis** is dependent upon an individual's inherent genetic composition and is not influenced by external factors. **Activated melanogenesis** occurs in response to ultraviolet radiation (UV) exposure, specifically UV-B (short UV wave) exposure. Increased melanogenesis in response to UV-B exposure serves to provide protection to the skin's innermost layer called the hypodermis, which lies below the epidermis and dermis (Figure 14.11). Melanin in the skin, specifically eumelanin, effectively absorbs UV-B radiation from light meaning that it will not reach the hypodermal layer. This effect is often more apparent during periods of the year when individuals tend to be outside more and the weather is warmer, which leads to those individuals donning fewer protective garments. The exposure of skin to sunlight is, of course, culturally mediated with some cultures encouraging the covering of skin at all times.

As previously noted, hemoglobin is an iron-rich protein that binds with oxygen in the bloodstream. For individuals with lighter-colored skin, blood vessels near the surface of the skin and the hemoglobin contained within those vessels is more apparent than in individuals with darker skin. The visible presence of hemoglobin coupled with the pink-to-red tone of the pheomelanin leads to lighter-skinned individuals having a pale pink skin tone. Individuals with lighter skin more readily absorb UV radiation as their basal melanin expression is directed more toward the production of pheomelanin than eumelanin. But, why are there so many variations in skin tone in humans? To answer this question, we now turn toward an exploration of an evolutionary-based adaptation of skin tone as a function of the environment.



Figure **6.5.2.14**: Evolutionary basis for human skin color variation.

Skin cancer is a significant concern for many individuals with light skin tone as the cumulative exposure of the epidermis and underlying skin tissues to UV radiation may lead to the development of abnormal cells within those tissues leading to malignancies. Although darker-skinned individuals are at risk for skin cancer as well, they are less likely to develop it due to increased levels of melanin, specifically eumelanin, in their skin. Even though skin cancer is a serious health concern for some individuals, most skin cancers occur in the post-reproductive years; therefore, it is improbable that evolutionary forces favoring varying melanin expression levels are related to a selective pressure to avoid such cancers. Furthermore, if avoiding skin cancer were the primary factor driving the evolution of various skin tones, then it reasons that everyone would have the most significant expression of eumelanin possible. So, why do we have different skin tones (Figure 14.12)? The term **cline** refers to the continuum or spectrum of gradations (i.e., levels or degrees) from one extreme to another. With respect to skin tone, the various tonal shades occur clinally

with darker skin being more prevalent near the equator and gradually decreasing in tone (i.e., decreased melanin production) in more distant latitudes. For individuals who are indigenous to equatorial regions, the increased levels of melanin within their skin provides them with a measure of protection against both sunburn and sunstroke as the melanin is more reflective of UV radiation than hemoglobin. In cases of severe sunburn, eccrine glands are affected, resulting in an individual's ability to sweat being

compromised. As sweat is the body's most effective means of reducing its core temperature to maintain homeostasis, damage to the eccrine glands may lead to numerous physiological issues related to heat that may ultimately result in death.

Even though avoiding severe sunburn and sunstroke is of great importance to individuals within equatorial regions, this is likely not the primary factor driving the evolutionary selection of darker skin within these regions. It has been proposed that the destruction of **folic acid**, which is a form of B-complex vitamin, by UV radiation may have led to the selection of darker skin in equatorial regions. For pregnant women, low levels of folic acid within the body during gestation may lead to defects in the formation of the brain and spinal cord of the fetus. This condition, which is referred to as spina bifida, often significantly reduces the infant's chances of survival without medical intervention. In men, low levels of folic acid within the body lead to an inhibition in the production of sperm. Thus, in geographic regions with high UV radiation levels (i.e., equatorial regions), there appears to be an evolutionarily driven correlation between darker skin and the maintenance of fertility.

If darker skin tone is potentially correlated to more successful reproduction, then why do lighter shades of skin exist? One hypothesis is that there is a relationship between lighter skin tone and vitamin D synthesis within the body. When skin is exposed to the UV-B radiation waves in sunlight, a series of chemical reactions occur within the epidermis leading to the production of vitamin D3. Before the body can use vitamin D3, it must travel to the liver and then to the kidneys where it is converted into different forms of bioactive molecules. Ultimately, it is converted into the bioactive molecule calcitriol (Vukić et al. 2015). Within the human body there are numerous cell types with binding receptors for calcitriol, so it is capable of adhering to the DNA of those cells (Snoddy et al. 2016). Calcitriol serves as a regulator in cellular-replication processes within the body, including those for pancreatic, breast, colon, and kidney cells (Snoddy et al. 2016). Insufficient calcitriol is associated with an increased risk of: some forms of cancer (colon, prostate, etc.), autoimmune diseases (multiple sclerosis, lupus, type I diabetes, etc.), cardiovascular diseases, and infections (e.g., tuberculosis, influenza) (Snoddy el al. 2016; Chaplin and Jablonski 2009). Deficiencies in calcitriol production and absorption within the human body may be linked to underlying genetic factors, such as a mutation in the vitamin D receptors present in some of the body's cells (Chaplin and Jablonski 2009). Alternatively, it may be linked to inadequate exposure to the UV-B rays necessary to stimulate calcitriol production or to a nutritional deficiency in vitamin D-rich foods. Regardless of the cause of the deficiency, individuals with a calcitriol (vitamin D3) deficiency may also be at risk for the development of certain skeletal abnormalities in addition to the previously mentioned health issues.




Figure 6.5.2.15: Children with rickets in various developmental stages.

Vitamin D is required for the absorption of certain nutrients, such as calcium and phosphorus, in the small intestine. These nutrients are among those that are critical for the proper growth and maintenance of bone tissue within the body. In the absence of adequate minerals, particularly calcium, bone structure and strength will be compromised leading to the development of rickets during the growth phase. Rickets is a disease affecting children during their growth phase and is characterized by inadequately calcified bones that are softer and more flexible than normal. Individuals with rickets will develop a true bowing of their femora, which may affect their mobility (Figure 14.13). In addition, deformation of pelvic bones in women may occur as a result of rickets leading to complications with reproduction. In adults, a deficiency in vitamin D will often result in osteomalacia, which is a general softening of the bones due to inadequate mineralization. This softening is the result of impaired bone metabolism that is primarily linked to

insufficient levels of bioavailable vitamin D, calcium, and phosphate. In addition, it may be linked to inadequate absorption of calcium in the bloodstream. As noted, a variety of maladies may occur due to the inadequate production or absorption of vitamin D, as well as the destruction of folate within the human body; so, from an evolutionary perspective, natural selection should favor a skin tone that is best suited to a given environment.

In general, the trend related to lighter skin pigmentation further from the equator follows a principle called Gloger's Rule. This rule states that within the same species of mammals the more heavily pigmented individuals tend to originate near the equator while lighter-pigmented members of the species will be found in regions further from the equator. Gloger's Rule applies latitudinally; however, it does not appear to hold for certain human populations near the poles. Specifically, the Inuit people (Figure 14.14), who are indigenous to regions near the North Pole and currently reside in portions of Canada, Greenland, Alaska, and Denmark. The Inuit have a darker skin tone that would not be anticipated under the provisions of Gloger's Rule. The high reflectivity of light off of snow and ice, which is common in polar regions, necessitates the darker skin tone of these individuals to prevent folic acid degradation just as it does for individuals within equatorial regions. The consumption of vitamin D–rich foods, such as raw fish, permits the Inuit to reside at high latitudes with darker skin tone while preventing rickets.


Figure **6.5.2.15**: Inuit family, 1917.

Genome studies have identified a number of genes (*TYR, OCA2/HERC2, TYRP1, SLC45A2, HPS6i*, etc.) related to the expression of melanin and pigmentation presentation in humans. Compared to the exceptionally large number of genes within the human genome, those regulating the expression of melanin are relatively few and appear on distinct loci. The genes at these loci are generally pleiotropic in nature, so there is a relatively predictable patterning in skin, hair, and eye color combinations (Sturm and Duffy 2012). For example, some populations that are indigenous to higher latitude regions tend to have lighter skin, hair, and eye color than their counterparts from equatorial regions. Still, since the genes affecting skin, hair, and eye color are actually independent, it is possible that variations may produce many phenotypic combinations. Turning again to our example of individuals indigenous to higher latitudes, it is theoretically possible to encounter an individual with dark hair, light-toned skin, and blue eyes within this region due to the variability of phenotypic combinations.

Adaptation: *Shape and Size Variations*

In addition to natural selection playing a role in the determination of melanin expression related to skin tone, which is correlated to the environment, it plays a significant role in the determination of the shape and size of the human body. As previously discussed, the most significant thermodynamic mechanism of heat loss from the body is radiation. At temperatures below 20℃ (68℉), the human body loses around 65% of its heat to radiative processes; however, the efficiency of radiation as a means of heat reduction is correlated to the overall body shape and size of the individual. There is a direct correlation between the ratio of an object's surface area to mass and the amount of heat that may be lost through radiation. For example, two metal objects of identical composition and mass are heated to the same temperature. One object is a cube and the other is a sphere. Which object will cool the fastest? Geometrically, a sphere has the smallest surface area per unit mass of any three-dimensional object, so the sphere will cool more slowly than the cube. In other words, the smaller the ratio of the surface area to mass an object has, the more it will retain heat. With respect to the cube in our example, mass increases by the cube, but surface area may increase only by the square, so size will affect the mass to surface area ratio. This, in general, holds true for humans, as well.

In regions where temperatures are consistently cold, the body shape and size of the individuals who are indigenous to the area tend to be more compact. These individuals have a relatively higher body mass to surface area (i.e., skin) than their counterparts from equatorial regions where the average temperatures are considerably warmer. Individuals from hot climates, such as the Fulani (Figure 14.15a) of West Africa, have limbs that are considerably longer than those of individuals from cold climates, such as the

Inuit of Greenland (Figure 14.15b). Evolutionarily, the longer limbs of individuals from equatorial regions (e.g., the Fulani) provide a greater surface area (i.e., lower body mass to surface area ratio) for the dissipation of heat through radiative processes. In contrast, the relatively short limbs of Arctic-dwelling people, such as the Inuit, allows for the retention of heat as there is a decreased surface area through which heat may radiate away from the body.



Figure **6.5.2.16**: The Fula people of Burkina Faso (pictured here in 1974) are from a tropical environment where the rapid dispersal of heat is necessary to maintain homeostasis.



Figure **6.5.2.17**: These Inuit people from Greenland live in an arctic environment where the conservation of heat in the body's core is of critical importance.




Figure **6.5.2.18**: These organisms are representative of Bergmann's rule. The animal on the left depicts an ungulate from a cooler environment with increased body weight and decreased surface area, compared to the slender ungulate on the right.

As described above, there are certain trends related to the general shape and size of human bodies in relation to the thermal conditions. To better describe these trends, we turn to a couple of general principles that are applicable to a variety of species beyond humans. Bergmann's Rule predicts that as average environmental temperature decreases, populations are expected to exhibit an increase in weight and a decrease in surface area (Figure 14.16a). Also, within the same species of homeothermic animals, the relative length of projecting body parts (e.g., nose, ears, and limbs) increases in relation to the average environmental temperature (Figure 14.16b). This principle, referred to as Allen's Rule, notes that longer, thinner limbs are advantageous for

the radiation of excess heat in hot environments and shorter, stockier limbs assist with the preservation of body heat in cold climates. A measure of the crural index (crural index=tibia length [divided by] femur length) of individuals from various human populations provides support for Allen's Rule since this value is lower in individuals from colder climates than it is for those from hot climates. The crural indices for human populations varies directly with temperature, so individuals with higher crural index values are generally from regions with a warmer average environmental temperature. Conversely, the crural indices are lower for individuals from regions where there are colder average temperatures.




Figure 6.5.2.19: These animals are representative of Allen's rule. Note the shorter limbs and ears of the rabbit on the left that you might find in cold temperatures. Note the length of the ears on the rabbit on the right that you might find in a warm climate. Rabbits do not sweat like humans, heat is dissipated primarily through their ears.

Nasal shape and size (Figure 14.17) is another physiological feature that is affected by way of an individual's ancestors' environments. The selective role of climate in determining human nasal variation is typically approached by dividing climates into four adaptive zones: hot-dry, hot-wet, cold-dry, and cold-wet (Maddux et al. 2016). One of the principal roles of the nasal cavity is to condition (i.e., warm and humidify) ambient air prior to its reaching the lungs. Given that function of the nasal cavity, it is anticipated that different nasal shapes and sizes will be related to varying environments. In cold-dry climates, an individual's nasal cavity must provide humidification and warmth to the dry air when breathing in through the nose (Noback et al. 2011). Also, in that type of climate, the nasal cavity must conserve moisture and minimize heat loss during when the individual exhales through the nose (Noback et al. 2011). From a physiological stress perspective, this is a stressful event.




Figure 6.5.2.20: Human nasal morphological variation as influenced by four major climate-based adaptive zones: hot-dry, hot-wet, cold-dry, and cold-wet. Note that images are presented left-to-right in relation to the climate-based adaptive zones, respectively.

Conversely, in hot-wet environments, there is no need for the nasal cavity to provide additional moisture to the inhaled air nor is there a need to warm the air or to preserve heat within the nasal cavity (Noback et al. 2011). So, in hot-wet climates, the body is under less physiological stress related to the inhalation of ambient air than in cold-dry climates. As with most human morphological elements, the shape and size of the nasal cavity occurs along a cline. Due to the environmental stressors of cold-dry environments requiring the humidification and warming of air through the nasal cavity, individuals indigenous to such environments tend to have taller (longer) noses with a reduced nasal entrance (nostril opening) size (Noback et al. 2011). This general shape is referred to as leptorrhine, and it allows for a larger surface area within the nasal cavity itself for the air to be warmed and humidified prior to entering the lungs (Maddux et al. 2016). In addition, the relatively small nasal entrance of leptorrhine noses serves as a means of conserving moisture and heat (Noback et al. 2011). Individuals indigenous to hot-wet climates tend to have platyrrhine nasal shapes, which are shorter with broader nasal entrances (Maddux et al. 2016). Since individuals in hot-wet climates do not need to humidify and warm the air entering the nose, their nasal tract is shorter and the nasal entrance wider to permit the effective cooling of the nasal cavity during respiratory processes.

Adaptation: Infectious Disease

Throughout our evolutionary journey, humans have been exposed to numerous infectious diseases. In the following section, we will explore some of the evolutionary-based adaptations that have occurred in certain populations in response to the stressors presented by select infectious diseases. One of the primary examples of natural selection processes acting on the human genome in response to the presence of an infectious disease is the case of the relationship between the sickle-cell anemia trait and malaria.

Malaria is a zoonotic disease (type of infectious disease naturally transmitted between animals and humans; covered in more detail in Chapter 16: Human Biology and Health) caused by the spread of the parasitic protozoa from the genus *Plasmodium* (Figure 14.18). These unicellular, eukaryotic protozoa are transmitted through the bite of a female *Anopheles* mosquito. During the bite process, the protozoan parasites that are present within an infected mosquito's saliva will enter the bloodstream of the individual where they will be transported to the liver. Within the liver, the parasites multiply and will eventually be released into the bloodstream where they will infect erythrocytes. Once inside the erythrocytes, the parasites will reproduce until they exceed the cell's storage capacity, causing it to burst and release the parasites into the bloodstream once again. This replication cycle will continue as long as there are viable erythrocytes within the host to infect.




Figure **6.5.2.21**: Life cycle of the malaria parasite.

General complications from malaria infections include: enlargement of the spleen (due to destruction of infected erythrocytes), lower number of thrombocytes (also called platelets, required for coagulation/clotting of blood), high levels of bilirubin (a byproduct of hemoglobin breakdown in the liver) in the blood, jaundice (yellowing of the skin and eyes due to increased blood bilirubin levels), fever, vomiting, retinal (eye) damage, and convulsions (seizures). According to the World Health Organization, in 2016 there were 445,000 deaths from malaria globally with the highest percentage of those deaths occurring in Africa (91%) and Southeast Asia (6%) (World Health Organization 2017). In sub-Saharan Africa, where incidents of malaria are the highest in the world, 125 million pregnancies are affected by malaria, resulting in 200,000 infant deaths (Hartman et al. 2013). Pregnant women who become infected during the gestational process are more likely to have low-birthweight infants due to prematurity or growth restriction inside the uterus (Hartman et al. 2013). After birth, infants born to malaria-infected mothers are more likely to develop infantile anemia (low red blood cell counts), a malaria infection that is not related to the maternal malarial infection, and they are more likely to die than infants born to non-malaria-infected mothers (Hartman et al. 2013).

For children and adolescents whose brains are still developing, there is a risk of cognitive (intellectual) impairment associated with some forms of malaria infections (Fernando et al. 2010). Given the relatively high rates of morbidity (disease) and mortality (number of deaths) associated with malaria, it leads to reason that this disease may have served as a selective pressure during human evolution. Support for natural selection related to malaria resistance is related to genetic mutations associated with sickle cell, thalassemia, glucose-6-phosphate dehydrogenase (G6PD) deficiency, and the absence of certain antigens (molecules capable of inducing an immune response from the host) on erythrocytes. For the purposes of this text, we will focus our discussion on the relationship between sickle cell disease and malaria.

Sickle cell disease is a group of genetically inherited blood disorders characterized by an abnormality in the shape of the hemoglobin within erythrocytes. It is important to note that there are multiple variants of hemoglobin, including, but not limited to: A, D, C, E, F, H, S, Barts, Portland, Hope, Pisa, and Hopkins. Each of these variants of hemoglobin may result in various conditions within the body; however, for the following explanation we will focus solely on variants A and S.




Figure **6.5.2.22**: Normal and sickled erythrocytes.

Individuals who inherit a mutated gene (hemoglobin with a sickled erythrocyte variety, HbS) on chromosome 11 from both parents will develop sickle cell anemia, which is the most severe form of the sickle cell disease family (Figure 14.19). The genotype of an individual with sickle cell anemia is HbSS; whereas, an individual without sickle cell alleles has a genotype of HbAA representing two normal adult hemoglobin type A variants. Manifestations of sickle cell anemia (HbSS) range from mild to severe with some of the more common symptoms being: anemia, blood clots, organ failure, chest pain, fever, and low blood oxygen levels. In high-income countries with advanced medical care, the median life expectancy of an HbSS individual is around 60 years; however, in low-income countries where advanced medical care is scarce, as many as 90% of children with sickle cell disease perish before the age of five (Longo et al. 2017).

Considering that advanced medical care was not available during much of human evolutionary history, it stands to reason that the majority of individuals with the HbSS genotype died before the age of reproduction. If that is the case though, why do we still have the HbS variant present in modern populations? As covered earlier in this textbook, the genotype of an individual is composed of genes from both biological parents. In the case of an individual with an HbSS genotype, the sickle cell allele (HbS) was inherited from each of the parents. For individuals with the heterozygous genotype of HbSA, they have inherited both a sickle cell allele (HbS) and a normal hemoglobin allele (HbA). Heterozygous (HbSA) individuals who reside in regions where malaria is endemic may have a selective advantage. They will experience a sickling of some, but not all, of their erythrocytes. Unlike an individual with the HbSS genotype, someone with HbSA may experience some of the symptoms listed above; however, they are generally less severe.

As noted earlier, the mechanism through which *Plasmodium* protozoan parasites replicate involves human erythrocyte cells. However, due to their sickled shape, as well as the presence of an abnormally shaped protein within the cell, the parasites are unable to replicate effectively in the erythrocyte cells coded for by the HbS allele (Cyrklaff et al. 2011). An individual who has an HbSA genotype and an active malaria infection will become ill with the disease to a lesser extent than someone with an HbAA genotype. Although normal erythrocytes (regulated by the HbA allele) allow for the replication of the parasite, the parasites will not be able to replicate in HbS erythrocytes of the heterozygote. So, individuals with the HbSA genotype are more likely to survive a malaria infection than an individual who is HbAA. Although individuals with the HbSA genotype may endure some physiological complications related to the sickling of some of their erythrocytes, their morbidity and mortality rates are lower than they are for HbSS members of the population. The majority of individuals who are heterozygous or homozygous for the HbS trait have ancestors who originated in sub-Saharan Africa, India, Saudi Arabia, regions in South and Central America, the Caribbean, and the Mediterranean (Turkey, Greece, and Italy) (Centers for Disease Control and Prevention 2017) (Figure 14.20).




Figure **6.5.2.23**: Distribution of sickle cell and associated erythrocytic abnormalities for Africa and Asia.

With respect to the history of these regions, during the early phases of settlement horticulture was the primary method of crop cultivation. Typically performed on a small scale, horticulture is based on manual labor and relatively simple hand tools rather than the use of draft animals or irrigation technologies. Common in horticulture is *swidden*, or the cutting and burning of plants in woodland and grassland regions. The swidden is the prepared field that results following a slash-and-burn episode. This practice fundamentally alters the soil chemistry, removes plants that provide shade, and increases the areas where water may pool. This anthropogenically altered landscape provides the perfect breeding ground for the *Anopheles* mosquito, as it prefers warm, stagnant pools of water (Figure 14.21).




Figure **6.5.2.24**: The effects of human horticultural activities on the balancing selection of populations in relation to sickle cell disease genotype variants.

Although swidden agriculture was historically practiced across the globe, it became most problematic in the regions where the *Anopheles* mosquito is endemic. These areas have the highest incidence rates of malaria infection. Over time, the presence of the *Anopheles* mosquito and the *Plasmodium* parasite that it transmitted acted as a selective pressure, particularly in regions where swidden agricultural practices were common, toward the selection of individuals with some modicum of resistance against the infection. In these regions, HbSS and HbSA individuals would have been more likely to survive and reproduce successfully. Although individuals and populations are far more mobile now than they have been throughout much of history, there are still regions where we can see higher rates of malaria infection as well as greater numbers of individuals with the HbS erythrocyte variant. The relationship between malaria and the selective pressure for the HbS variant is one of the most prominent examples of natural selection in the human species within recent evolutionary history.

Adaptation: Lactase Persistence

With the case of sickled erythrocytes and their resistance to infection by malaria parasites, there is strong support for a cause-and-effect-style relationship linked to natural selection. Although somewhat less apparent, there is a correlation between lactase persistence and environmental challenges. Lactase-phlorizin hydrolase (LPH) is an enzyme that is primarily produced in the small intestine and permits the proper digestion of lactose, a disaccharide (composed of two simple sugars: glucose and galactose) found in the milk of mammals. Most humans will experience a decrease in the expression of LPH following weaning, leading to an inability to properly digest lactose. Generally, LPH production decreases between the ages of two and five and is completely absent by the age of nine (Dzialanski et al. 2016). For these individuals, the ingestion of lactose may lead to a wide variety of gastrointestinal ailments including abdominal bloating, increased gas, and diarrhea. Although the bloating and gas are unpleasant,

the diarrhea caused by a failure to properly digest lactose can be life-threatening if severe enough due to the dehydration it can cause. Some humans, however, are able to produce LPH far beyond the weaning period.




Figure **6.5.2.25**: Interpolated map depicting the percentage of adults with the lactase persistence genotype in indigenous populations of the Old World. Circles denote sample locations.

Individuals who continue to produce LPH have what is referred to as the **lactase persistence** trait. The lactase persistence trait is encoded for a gene called *LCT*, which is located on human chromosome 2 (Ranciaro et al. 2014; see also Chapter 3). From an evolutionary and historical perspective, this trait is most commonly linked to cultures that have practiced cattle domestication (Figure 14.22). For individuals in those cultures, the continued expression of LPH may have provided a selective advantage. During periods of environmental stress, such as a drought, if an individual is capable of successfully digesting cow's milk, they have a higher chance of survival than someone who suffers from diarrhea-linked dehydration due to a lack of LPH. Per Tishkoff et al. , the "frequency of lactase persistence is high in northern European populations (more than 90% in Swedes and Danes), decreases in frequency across southern Europe and the Middle East (less than 50% in Spanish, French, and pastoralist Arab populations), and is low in non-pastoralist Asian and African populations (less than 1% in Chinese, less than 5% to 20% in West African agriculturalists)" (2007: 248). Although the frequency of the lactase persistence trait is relatively low among African agriculturalists, it is high among pastoralist populations that are traditionally associated with cattle domestication, such as the Tutsi and Fulani, who have frequencies of 90% and 50%, respectively (Tishkoff et al. 2007).

Cattle domestication began around 11,000 years ago in Europe (Beja-Pereira et al. 2006) and 7,500 to 9,000 years ago in the Middle East and North Africa (Tishkoff et al. 2007). Based on human genomic studies, it is estimated that the mutation for the lactase persistence trait occurred around 2,000 to 20,000 years ago for European populations (Tishkoff et al. 2007). For African populations, the lactase persistence trait emerged approximately 1,200 to 23,000 years ago (Gerbault et al. 2011). This begs the question: Is this mutation the same for both populations? It appears that the emergence of the lactase persistence mutation in non-European populations, specifically those in East Africa (e.g., Tutsi and Fulani), is a case of **convergent evolution**. With convergent evolution events, a similar mutation may occur in species of different lineages through independent evolutionary processes. Based on our current understanding of the genetic mutation pathways for the lactase persistence trait in European and African populations, these mutations are not representative of a shared lineage. In other words, just because a person of European origin and a person of African origin can each digest milk due to the presence of the lactase-persistence trait in their genotypes, it does not mean that these two individuals inherited it due to shared common ancestry.

Is it possible that the convergent evolution of similar lactase-persistence traits in disparate populations is merely a product of genetic drift? Or is there evidence for natural selection? Even though 23,000 years may seem like a long time, it is but a blink of the proverbial evolutionary eye. From the perspective of human evolutionary pathways, mutations related to the *LCT* gene have occurred relatively recently. Similar genetic changes in multiple populations through genetic drift processes, which are relatively slow and directionless, fail to accumulate as rapidly as have lactase-persistence traits (Gerbault et al. 2011). The widespread accumulation of these traits in a relatively short period of time supports the notion that an underlying selective pressure must be driving this form of human evolution. Although to date no definitive factors have been firmly identified, it is thought that

environmental pressures are likely to credit for the rapid accumulation of the lactase-persistence trait in multiple human populations through convergent evolutionary pathways.

Human Variation: Our Story Continues

From the time that the first of our species left Africa, we have had to adjust and adapt to numerous environmental challenges. The remarkable ability of human beings to maintain homeostasis through a combination of both nongenetic (adjustments) and genetic (adaptations) means has allowed us to occupy a remarkable variety of environments from high-altitude mountainous regions to the tropics near the equator. From adding piquant, pungent spices to our foods as a means of inhibiting food-borne illnesses due to bacterial growth to donning garments specially suited to local climates, behavioral adjustments have provided us with a nongenetic means of coping with obstacles to our health and well-being. Acclimatory adjustments, such as sweating when we are warm in an attempt to regulate our body temperature or experiencing increased breathing rates as a means of increasing blood oxygen levels in regions where the partial pressure of oxygen is low, have been instrumental in our survival with respect to thermal and altitudinal environmental challenges. For some individuals, developmental adjustments that were acquired during their development and growth phases (e.g., increased heart and lung capacities for individuals from high-altitude regions) provide them with a form of physiological advantage not possible for someone who ventures to such an environmentally challenging region as an adult. Genetically-mediated adaptations, such as variations in the pigmentation of our skin, have ensured our evolutionary fitness across all latitudes.

Will the human species continue to adjust and adapt to new environmental challenges in the future? If past performance is any measure of future expectations, then the human story will continue as long as we do not alter our environment to the point that the plasticity of our behavior, physiological, and morphological boundaries is exceeded. In the following chapters, you will explore additional information about our saga as a species. From the concept of race as a sociocultural construct to our epidemiological history, the nuances of evolutionary-based human variation are always present and provide the basis for understanding our history and our future as a species.

6.5.2: Adjustments and Adaptations is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **14.2: Adjustments and Adaptations** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.5.3: Review Questions

- Detail at least two examples of how natural selection has influenced human variation. Specifically, what was the selective pressure that may have led to a preference for a specific trait and how is that trait related to an increased level of fitness?
- Why is reduced pigmentation of the skin advantageous for individuals from northern latitudes? What role does darker skin pigmentation serve for individuals near the equator? What is the relationship between skin pigmentation and fitness?
- What are some of the risks associated with pregnancy at high altitude? Compare and contrast the various genetic mutations of the indigenous Tibetan, Andean, and Ethiopian high-altitude populations. In your answer, specifically address the issue of pregnancy at high altitudes.
- What is the relationship between the sickle cell mutation and the *Plasmodium* parasite? Would having the HbSA genotype still be advantageous in a region where such parasites are not common? Why or why not?

6.5.3: Review Questions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **14.3: Review Questions** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

6.5.4: Key Terms

Acclimatory adjustments: Processes by which an individual organism adjusts to maintain homeostasis in response to environmental challenges.

Activated melanogenesis: Increase in melanin production in response to ultraviolet radiation (UV) exposure.

Adaptation: alteration in population-level gene frequencies related to environmentally induced selective pressures; leads to a greater level of fitness for a population related to a specific environment.

Adjustment: Non-genetic-based ways in which organisms adjust to environmental stressors.

Allen's Rule: Due to thermal adaptation, homeothermic animals have body volume-to-surface ratios that vary inversely with the average temperature of their environment. In cold climates, the anticipated ratio is high and it is low in warm climates.

Basal melanogenesis: Genetically-mediated, non-environmentally influenced base melanin level.

Behavioral adjustments: An individual's culturally mediated responses to an environmental stressor in an effort to maintain homeostasis.

Bergmann's Rule: For a broadly distributed monophyletic group, species and populations of smaller size tend to be found in environments with warmer climates and those of larger size tend to be found in ones that are colder.

Cline: A continuum of gradations (i.e., degrees or levels) of a specific trait.

Conduction: Mechanism of heat transfer between objects through direct contact.

Convection: Movement of heat away from a warm object to the cooler surrounding fluid (i.e., gas or liquid).

Convergent evolution: Evolutionary process where organisms that are not closely related independently evolve similar traits as a product of adaptation to similar evolutionary parameters.

Erythrocyte: Red blood cell; most common form of blood cell—principle means of transporting oxygen throughout the circulatory system.

Evaporation: Mechanism of heat transfer where liquid is transformed into a gas utilizing energy (e.g., heat).

Folic acid: Form of B complex vitamin necessary for proper fetal development.

Gloger's Rule: For mammals of the same species, those with more darkly pigmented forms tend to be found closer to the equator and those with lighter forms are found in regions further from the equator.

Hematocrit: Volume percentage of red blood cells within the blood.

Homeostasis: Condition of optimal functioning for an organism.

Hyperpnea: Increased depth and rate of respiration.

Hypothalamus: Small portion of the human brain responsible for body temperature regulation.

Lactase persistence: Genetic mutation permitting the continued production of lactase-phlorizin hydrolase enzyme in the small intestine past the weaning period.

Melanin: Black-brown pigment produced by melanocytes; one of the primary pigments in skin.

Melanocytes: Specialized cells that produce melanin.

Phenotypic plasticity: Ability of one genotype to produce more than one phenotype dependent on environmental conditions.

Polyphenism: Multiple discrete phenotypes from an organism's genotype in response to the environment; a special form of phenotypic plasticity.

Radiation: Mechanism of heat transfer involving electromagnetic energy being emitted from an object.

Sickle cell disease: A group of genetically inherited blood disorders characterized by an abnormality in the shape of the hemoglobin within erythrocytes (red blood cells).

Stressor: Any stimulus resulting in an imbalance in an organism's homeostatic balance.

Vasoconstriction: Narrowing of the blood vessels due to contractions of the muscular vessel walls.

Vasodilation: Dilation of the blood vessels due to relaxation of the muscular vessel walls.

6.5.4: Key Terms is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **14.4: Key Terms** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

CHAPTER OVERVIEW

7: New Frontiers

7.1: Homo sapiens Futures - Doom, Gloom, and Hope?

7.1.1: Introduction

7.1.2: Doom and Gloom

7.1.3: Hope

7.2: Bioarchaeology and Forensic Anthropology

7: New Frontiers is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

7.1: Homo sapiens Futures - Doom, Gloom, and Hope?

7.1.1: Introduction

7.1.2: Doom and Gloom

7.1.3: Hope

Thumbnail: cjohnson7 from Rochester, Minnesota

This page titled 7.1: Homo sapiens Futures - Doom, Gloom, and Hope? is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

7.1.1: Introduction

I don't know a better way to say it, but as a species we've pretty much permanently screwed-up our planet. The real question is "What now?" Do we continue our nihilist craze to destroy the rest before going extinct, or do we try to salvage and restore as much we can? Most people in the Abrahamic Western tradition (Judeo-Christian-Islamic) have been enculturated to regard nature as something that needs to be dominated. Many Christians believe the hastened destruction of our world should be encouraged because it is a prerequisite for The Second Coming. Our current capitalist economy translates the domination of nature into the inalienable rights of the individual to exploit natural resources, even non-renewable ones, without taking into consideration the consequences for future generations. And to better use up those natural resources, capitalism creates a culture of consumerism, where our happiness is measured by conspicuous consumption.




Figure 7.1.1.1 - *2018 San Diego March for Science*

How could a sane hominid have such low self-esteem that they would take a credit card to a shopping mall to buy things they don't really need, from a factory in China that uses slave labor and bribes its way out of pollution controls, to the point where they're in debt and need to burn non-renewable fossil fuels sitting in rush hour traffic for hours every day to pay off the interest on their card, to the point where if you asked them to help solve some of the problems that effect their fellow hominids, they would respond that they don't have any free time, or they're afraid of getting fired if they speak up?

So where's the hope?

I hope you got a sense in this class of the difference between culture and biology, and in this case, the difference between biological change and cultural change, especially in terms of time, the rate of change, how much faster culture can change compared to biology. When we talk about drastic biological change we can talk about evolution and the adaptive radiation of large groups of species into ecological niches over millions of years. When we talk about drastic cultural change we are talking about revolution. In an instant of geological time, our species went from crude tools to stunningly beautiful cave art in what we studied as the Upper

Paleolithic revolution. I find hope in knowing how fast culture can change, that in the blink of an eye, the crude ideology of our dominant hominid could be replaced by something much more beautiful.

This page titled 7.1.1: Introduction is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.


- **8.1: Introduction** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

7.1.2: Doom and Gloom

> *we must acknowledge, as it seems to me, that man with all his noble qualities, with sympathy which feels for the most debased, with benevolence which extends not only to other men but to the humblest living creature, with his god-like intellect which has penetrated into the movements and constitution of the solar system—with all these exalted powers—Man still bears in his bodily frame the indelible stamp of his lowly origin. [Darwin 1871:405]*

Overpopulation

The word *overpopulation* is very subjective. There is no scientific way to decide how many people is too much. It is more of an aesthetic question: like would you prefer to live in a huge mansion all by yourself, or in a one-bedroom apartment with 10 roomates? Most people would choose something in between, probably leaning towards the mansion. The science of population change is called *demography*, literally a "map of people/districts", and the science has mapped a population explosion in the last eye-blink of human evolution. We are rapidly heading towards the one-bedroom apartment with 10 roomates, and the ensuing drama is taking place on a global scale.



Notes

World Population from Population Education on Vimeo.

Population growth

Where do you fit into the 7 billion?

Bioanthropological perspective: McKee, Jeffery 2012 “The Human Population Footprint on Global Biodiversity”

Video on transhumance

Loss of Biodiversity

Should we care about any other species besides our own? This question is like asking "Should I care about anyone else besides myself?" on a larger scale. Most people care about at least some of their family members. Through the genealogical emphasis of anthropology, many people can extend their altruism to include broader categories of people they're not related too, people of different "races", and hopefully other species.

To me biodiversity is an aesthetic question. Complex ecosystems are beautiful. Do you want kids to grow up in a world where the only other life they know is cockroaches, mosquitos, rats, and pigeons? Review the section on primate extinction

NoteS

Megafauna extinction due to humans

Humans as super-predators

Global Warming

If you compare a graph of population growth and average temperature you can see the correlation between human overpopulation and climate change. How much CO2 was produced by 6,000,000 (6 million) hunter-gatherers sitting around campfires? Almost nothing! That's about the equivalent of one campfire every two hundred square miles. How much CO_2 is produced by 6,000,000,000 (6 billion) people who are driving, consuming, deforesting? Our ecological footprint is massive compared to our hunter-gatherer ancestors, and there are thousand times more of us.

We should expect both massive extinction and some evolution as variations in populations are selected for in the new ecological niches caused by extreme weather, drought, flooding, acidification of the oceans. Global warming will cause more people to die of starvation and infectious disease. Many plants and animals will go extinct, humans will use cultural adaptations to survive, but it will be very painful and expensive, and the longer will put our heads in the sand and ignore the problem, the worse it will be.



Notes

Comparing human caused extinction to other mass extinctions

An economic approach, the environment as an “externality”

Pollution

Geological periods are usually bracketed by climate change and major extinction, and they are based on empirical evidence that a geologist with a rock hammer can go out and find. Human effects on the environment have begun to show up in the field of geology:

- "An anthropogenic marker horizon in the future rock record"

This page titled 7.1.2: Doom and Gloom is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.


- **8.2: Doom and Gloom** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

7.1.3: Hope

Is there hope for the human species? I think so. We have a millions of years before the sun engulfs us. One of the lessons of evolutionary psychology is that we do have free-will and we are not slaves to our instincts. The *Upper Paleolithic revolution* is an example of how quickly we can change. Skim the section on primate ethology. I believe understanding the primate origin of human behaviors helps us make cultural choices about how to suppress or embrace them.

We are starting to grant legal rights for non-human primates, Here's an example in Argentina. If you haven't already, read the the bee keepers for primate conservation article


* Primate have a sense of justice and reconciliation:



Exercise 7.1.3.1

WATCH THIS JANE GOODALL INTERVIEW

* "Get Whacked for Wildlife" 2016 campaign:



> *"Never doubt that a small group of thoughtful, committed citizens can change the world; indeed, it's the only thing that ever has." -Margaret Mead*

Imagination Actions

- Take this World Population Knowledge Quiz
- Volunteer with a non-profit group working to increase hope in the world

This page titled 7.1.3: Hope is shared under a CC BY-NC-ND 4.0 license and was authored, remixed, and/or curated by Arnie Daniel Schoenberg via source content that was edited to the style and standards of the LibreTexts platform.

- **8.3: Hope** by Arnie Daniel Schoenberg is licensed CC BY-NC-ND 4.0. Original source: https://www.arnieschoenberg.com/anth/bio/intro.

7.2: Bioarchaeology and Forensic Anthropology

Ashley Kendell, Ph.D., D-ABMDI, California State University, Chico

Alex Perrone, M.A., California State University, Chico

Colleen Milligan, Ph.D., California State University, Chico

Learning Objectives

- Define and differentiate bioarchaeology and forensic anthropology as subfields of biological anthropology.
- Describe the seven steps carried out during skeletal analysis.
- Outline the four major components of the biological profile.
- Contrast the four categories of trauma.
- Explain how to identify the different taphonomic agents that alter bone.
- Discuss ethical considerations for both bioarchaeology and forensic anthropology.

Bioarchaeology and forensic anthropology are both subfields of biological anthropology. While the goals of each subfield are different, each relies on skeletal analysis to gain information about humans, both past and present. This chapter will provide a general overview of the analysis of human skeletal remains, as it applies to both bioarchaeology and forensic anthropology.

About the Authors

Ashley Kendell

California State University, Chico, akendell@csuchico.edu



Ashley Kendell

Dr. Ashley Kendell is currently an assistant professor and forensic anthropologist at Chico State. Prior to beginning her position at Chico State, she was a visiting professor at the University of Montana and the forensic anthropologist for the state of Montana. Dr. Kendell obtained her doctorate from Michigan State University, and her research interests include skeletal trauma analysis and digitization and curation methods for digital osteological data. She is also a Registry Diplomate of the American Board of Medicolegal Death Investigators. Throughout her doctoral program, she worked as a medicolegal death investigator for the greater Lansing, Michigan, area and was involved in the investigation of over 200 forensic cases.

Alex Perrone

California State University, Chico, aperrone1@mail.csuchico.edu



Alex Perrone

Alex Perrone is a lecturer in anthropology and biological sciences at California State University, Chico, and a lecturer in anthropology at Butte Community College. She is also the Supervisor of the Human Identification Laboratory in the Department of Anthropology at California State University, Chico. Her research interests include bioarchaeology, paleopathology, forensic anthropology, skeletal biology, California prehistory, and public health. She has worked on bioarchaeological and archaeological projects in Antigua, California, Hawaii, Greece, and the UK, and was an archaeological technician for the USDA Forest Service.

She assists with training courses for local and federal law enforcement agencies and assists law enforcement agencies with the recovery and analysis of human remains.

Colleen Milligan

California State University, Chico, cfmilligan@csuchico.edu



Colleen Milligan

Dr. Colleen Milligan is a physical anthropologist with research interests in bioarchaeology, skeletal biology, and forensic anthropology. She has been a Fellow with the Department of Homeland Security and has assisted in forensic anthropology casework and recoveries in the State of Michigan and California. She has also assisted in community outreach programs in forensic anthropology and forensic science, as well as recovery training courses for local, state, and federal law enforcement officers. She is a certified instructor through Peace Officers Standards and Training (POST). Dr. Milligan serves as the current co-director of the Chico State Human Identification Laboratory.

References

Adams, Bradley J., and Lyle W. Konigsberg, eds. 2008. *Recovery, Analysis, and Identification of Commingled Remains*. Totowa, NJ: Humana Press.

Berg, Gregory E. 2017. “Sex Estimation of Unknown Human Skeletal Remains.” In *Forensic Anthropology: A Comprehensive Introduction, Second edition*, edited by Natalie R. Langley and MariaTeresa A. Tersigni-Tarrant, 143–159. Boca Raton, FL: CRC Press.

Brooks, S., and J. M. Suchey. 1990. “Skeletal Age Determination Based on the Os Pubis: A Comparison of the Acsádi-Nemeskéri and Suchey-Brooks Methods.” *Human Evolution* 5 (3): 227–238.

Cunningham, Craig, Louise Scheuer, and Sue Black. 2016. *Developmental Juvenile Osteology, Second edition*. London: Elsevier Academic Press.

Galloway, Alison. Broken Bones: Anthropological Analysis of Blunt Force Trauma. 1999. Springfield, IL: Charles C. THomas Publisher, LTD.

Hefner, Joseph T., and Kandus C. Linde. 2018. *Atlas of Human Cranial MacromorphoscopicTraits*. San Diego: Academic Press.

Hopkinson, Kimberly A., Sarah M. Yeats, and G. Richard Scott. 2008. “For Whom the Coin Tolls: Green Stained Teeth and Jaws in Medieval and Post-Medieval Spanish Burials.” *Dental Anthropology* 21 (1): 12–17.

Katz, Darryl, and Judy Myers Suchey. 1986. “Age Determination of the Male Os Pubis.” *American Journal of Physical Anthropology* 69 (4): 427–435.

Langley, Natalie R., Alice F. Gooding, and MariaTeresa Tersigni-Tarrant. 2017. “Age Estimation Methods.” In *Forensic Anthropology: A Comprehensive Introduction, Second edition*, edited by Natalie R. Langley and MariaTeresa A. Tersigni-Tarrant, 175–191. Boca Raton, FL: CRC Press.

Lovell, Nancy C. 1997. “Trauma Analysis in Paleopathology.” *Yearbook of Physical Anthropology* 104 (S25): 139–170.

İşcan, M. Y., S. R. Loth, and R. K. Wright. 1984. “Age Estimation from the Rib by Phase Analysis: White Males.” *Journal of Forensic Sciences* 29 (4): 1094–1104.

———. 1985. “Age Estimation from the Rib by Phase Analysis: White Females.” *Journal of Forensic Sciences* 30 (3): 853–863.

Native American Graves Protection and Repatriation Act (NAGPRA) 1990 (25 U.S. Code 3001 et seq.)

Nelson’s Caribbean Hell-hole: An Eighteenth-Century Navy Graveyard Uncovered. 2013. BBC Documentary. https://www.bbc.co.uk/programmes/p0187q6r.

NIH U.S. National Library of Medicine. 2019. “Genetics Home Reference: Achondroplasia.” Last modified February 5, 2019. https://ghr.nlm.nih.gov/condition/achondroplasia.

NIH. Genetic and Rare Diseases Information Center. Last modified October 13, 2017. https://rarediseases.info.nih.gov

NIST (National Institute of Standards and Technology). N.d. “The Organization of Scientific Area Committees for Forensic Science.” https://www.nist.gov/topics/organiza...rensic-science.

Ortner, Donald J., and Walter G. J. Putschar. 1981. *Identification of Pathological Conditions in Human Skeletal Remains*. Washington, D.C.: Smithsonian Institution Press.

Ousley, Stephen. 1995. “Should We Estimate Biological or Forensic Stature?” *Journal of Forensic Sciences*. 40, no. 5: 768–773.

Phenice, T. W. 1969. “A Newly Developed Visual Method of Sexing the Os Pubis.” *American Journal of Physical Anthropology* 30 (2): 297–302.

Rose, Jerome C., Thomas J. Green, and Victoria D. Green. 1996. “NAGPRA Is Forever: Osteology and the Repatriation of Skeletons.” *Annual Review of Anthropology* 25: 81–103.

Schaefer, Maureen, Sue Black, and Louise Scheuer. *Juvenile Osteology: A Laboratory and Field Manua*l. 2009. San Diego: Elsevier Academic Press.

Scientific Working Group for Forensic Anthropology (SWGANTH). 2010a. “Personal Identification.” Last modified June 3, 2010. www.nist.gov/sites/default/files/documents/2018/03/13/swganth_personal_identification.pdf.

Scientific Working Group for Forensic Anthropology (SWGANTH). 2010b. “Sex Assessment.” Last modified June 3, 2010. www.nist.gov/sites/default/files/documents/2018/03/13/swganth_sex_assessment.pdf.

Scientific Working Group for Forensic Anthropology (SWGANTH). 2011. “Trauma Analysis.” Last modified May 27, 2011. www.nist.gov/sites/default/files/documents/2018/03/13/swganth_trauma.pdf.

Scientific Working Group for Forensic Anthropology (SWGANTH). 2012. “Stature Estimation.” Last modified August 2, 2012. www.nist.gov/sites/default/files/documents/2018/03/13/swganth_stature_estimation.pdf.

Scientific Working Group for Forensic Anthropology (SWGANTH). 2013. “Age Estimation.” Last modified January 22, 2013. www.nist.gov/sites/default/files/documents/2018/03/13/swganth_age_estimation.pdf.

Tersigni-Tarrant, MariaTeresa A., and Natalie R. Langley. 2017. “Human Osteology.” In *Forensic Anthropology: A Comprehensive Introduction, Second edition*, edited by Natalie R. Langley and MariaTeresa A. Tersigni-Tarrant, 81–109. Boca Raton, FL: CRC Press.

Ubelaker, Douglas H. 2018. “A History of Forensic Anthropology.” Special issue: Centennial Anniversary Issue of AJPA, *American Journal of Physical Anthropology* 165 (4): 915–923.

White, Tim D., and Pieter A. Folkens. 2005. *The Human Bone Manual*. Burlington, MA: Elsevier Academic Press.

Figure Attributions

Figure 15.1 Example of burned sheetrock by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.2 Cross section of human long bone original to Explorations: An Open Invitation to Biological Anthropology by Mary Nelson is under a CC BY-NC 4.0 License.

Figure 15.3 Anatomy of a Flat Bone (Anatomy & Physiology, Figure 6.3.3 byOpenStax is used under aCC BY 4.0 License.

Figure 15.4 Bone (248 12) Bone cross section by Doc. RNDr. Josef Reischig, CSc. is used under a CC BY-SA 3.0 License.

Figure 15.5 Example of PVC pipe by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.6 Animal bone cross section by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.7 Tib fib growth plates by Gilo1969 at English Wikipedia is used under a CC BY 3.0 License.

Figure 15.8 Filling by Kauzio has been designated to the public domain (CC0).

Figure 15.9 Spanish Civil War – Mass grave – Estépar, Burgos by Mario Modesto Mata is used under a CC BY-SA 4.0 License.

Figure 15.10 Anterior and lateral view of a male and female cranium by Ashley Kendell a derivative work original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License. [Includes Human Female Asian Skull and Human Male Asian Skull by ©BoneClones used by permission.]

Figure 15.11 Features associated with the Phenice Method by Colleen Milligan original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.12 Skeletal traits commonly used in a morphological assessment of ancestry by Colleen Milligan original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.13 5-year-old Human Child Skull with Mixed Dentition Exposed by ©BoneClones is used by permission and available here is under a CC BY-NC 4.0 License.

Figure 15.14 Example of the progression of degenerative changes to the pubic symphysis a derived work by Ashley Kendell original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License. [Original photos by Dr. Julie Fleischman used by permission.]

Figure 15.15 Examples of degenerative changes to the sternal rib end by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.16 Measurement of the bicondylar length of the femur by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.17 Frontal bone sinuses by Alex Khimich is used under a CC BY-SA 4.0 License.

Figure 15.18 Shoulder replacement by Smithsonian [exhibit: Written in Bone, Today's Bones] has no known copyright restrictions.

Figure 15.19 Skull sword trauma by the National Institutes of Health, Health & Human Services [19th Century Collection, National Museum of Health and Medicine, Armed Forces Institute of Pathology, Washington, D.C. From exhibition "Visible Proofs: Forensic Views of the Body" U.S. National Library of Medicine] is in the public domain.

Figure 15.20 Skull hammer trauma by the National Institutes of Health, Health & Human Services [19th Century Collection, National Museum of Health and Medicine, Armed Forces Institute of Pathology, Washington, D.C. From exhibition "Visible Proofs: Forensic Views of the Body" U.S. National Library of Medicine] is in the public domain.

Figure 15.21 Trauma: Gunshot Wounds by Smithsonian [exhibit: Written in Bone, How Bone Biographies Get Written] has no known copyright restrictions.

Figure 15.22 Human Left Femur, Osteosarcoma by ©BoneClones is used by permission and available here is under a CC BY-NC 4.0 License.

Figure 15.23 Osteogenesis Imperfecta Type V by ShakataGaNai is used under a CC BY-SA 4.0 License.

Figure 15.24 Human Femur, Tibia and Fibula, Rickets by ©BoneClones is used by permission and available here is under a CC BY-NC 4.0 License.

Figure 15.25 Human Female Achondroplasia Dwarf Skeleton, Articulated by ©BoneClones is used by permission and available here is under a CC BY-NC 4.0 License.

Figure 15.26 Rodent gnawing by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.27 Carnivore damage by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.28 Burned bone by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.29 Root etching by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.30 Weathering by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

Figure 15.31 Cut marks by Alex Perrone original to Explorations: An Open Invitation to Biological Anthropology is under a CC BY-NC 4.0 License.

7.2: Bioarchaeology and Forensic Anthropology is shared under a CC BY-NC license and was authored, remixed, and/or curated by LibreTexts.

- **15: Bioarchaeology and Forensic Anthropology** by Beth Shook, Katie Nelson, Kelsie Aguilera, & Lara Braff, Eds. is licensed CC BY-NC 4.0. Original source: https://explorations.americananthro.org.

Index

A

acclimatory adjustments
6.5.2: Adjustments and Adaptations
activity budget
3.5.5: Primate Ecology
adaptations
6.5.1: Stress and Homeostasis
adaptive radiation
2.9: Adaptive Radiation
adjustments
6.5.1: Stress and Homeostasis
Age of Discovery
6.1: The History of "Race" Concepts
allopatric
2.6: Macroevolution
Anthropological Imagination
1.1.2: Anthropology as a Science
apes
3.5.9: Even more Haplorhines (The Apes)
Ardipithecus kadabba
4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba
Ardipithecus ramidus
4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba
Atelidae
3.5.7: Overviews of Living Primates (Haplorhines)
Australopithecus afarensis
4.3.4: Australopithecus afarensis
Australopithecus africanus
4.3.7: Australopithecus prometheus or africanus
4.3.8: Australopithecus africanus
Australopithecus anamensis
4.3.2: Australopithecus anamensis
Australopithecus bahrelghazali
4.3.5: Australopithecus bahrelghazali
Australopithecus garhi
5.1.6: Australopithecus garhi
Australopithecus prometheus
4.3.7: Australopithecus prometheus or africanus
Australopithecus sediba
5.1.7: Australopithecus sediba
axial skeleton
3.5.2: Primate Skeletal Anatomy

B

Beagle Voyage
2.1.4: Darwinian Evolution
behavioral adjustments
6.5.2: Adjustments and Adaptations
binomial nomenclature
6.1: The History of "Race" Concepts
biological anthropology
6: Race and Human Variation
biological determinism
6.1: The History of "Race" Concepts
bipedalism
4.2.2: What is a Hominim
bony labyrinth
6.2: Human Variation in Biological Anthropology Today
Bush Meat
3.7: Conservation

C

Cercopithecinae
3.5.8: More Haplorhines (Old World Monkeys)
Charles Darwin
2.1.4: Darwinian Evolution
cladistics
3.2: Primate Classification
clines
6.2: Human Variation in Biological Anthropology Today
Colobinae
3.5.8: More Haplorhines (Old World Monkeys)
conduction
6.5.2: Adjustments and Adaptations
conservation
3.7: Conservation
continuous variation
6.2: Human Variation in Biological Anthropology Today
convection
6.5.2: Adjustments and Adaptations
convergent evolution
3.5.1: Classification - Ordering the Natural World
Cro–Magnon man
5.1.22: Homo sapiens
cultural transmission
5.1.9: Homo habilis
cystic fibrosis
6.2: Human Variation in Biological Anthropology Today

D

Darwin tubercle
3.1: Introduction to Primatology
demography
7.1.2: Doom and Gloom
Denisovans
5.1.20: The Denisovans
Dikika Baby
4.3.4: Australopithecus afarensis

E

ecological niches
6.2: Human Variation in Biological Anthropology Today
enculturation
1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria
erectus grade
5.1.13: The “erectus Grade”
erythrocytes
6.5.2: Adjustments and Adaptations
essentialism
6.1: The History of "Race" Concepts
ethnicity
6: Race and Human Variation
ethology
3.6: Ethology
eugenics
6.1: The History of "Race" Concepts
evaporation
6.5.2: Adjustments and Adaptations
evolutionary classification
3.5.4: Taxonomy of the Living Primates

F

Familial Culture
1.2.7: Genetics, Cellular Biology, and Variation (Part 1)
founder effects
6.2: Human Variation in Biological Anthropology Today

G

gene flow
6.2: Human Variation in Biological Anthropology Today
genetic drift
1.2.6: "Forces" of Evolution
2.1.7: Modern Synthesis
6.2: Human Variation in Biological Anthropology Today
Gracile Australopiths
4.3.3: Gracile Australopiths

H

haplorhines
3.5.7: Overviews of Living Primates (Haplorhines)
Haplorhini
3.5.8: More Haplorhines (Old World Monkeys)
3.5.9: Even more Haplorhines (The Apes)
Haplorhini classification
3.5.4: Taxonomy of the Living Primates
hematocrit
6.5.2: Adjustments and Adaptations
heterogeneity
6.2: Human Variation in Biological Anthropology Today
holism
1.1.2: Anthropology as a Science
1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria
homeostasis
6.5.1: Stress and Homeostasis
Hominim
4.2.2: What is a Hominim
Homo antecessor
5.1.17: Homo antecessor
Homo erectus
5.1.15: Homo erectus
Homo ergaster
5.1.14: Homo ergaster
Homo floresiensis
5.1.18: Homo floresiensis

Homo georgicus
5.1.16: Homo georgicus
Homo habilis
5.1.9: Homo habilis
Homo heidelbergensis
5.1.19: Homo heidelbergensis
Homo naledi
5.1.12: Homo naledi
Homo neanderthalensis
5.1.21: Homo neanderthalensis
Homo rudolfensis
5.1.10: Homo rudolfensis
homogenous
6: Race and Human Variation
Hox genes
2.6: Macroevolution
human diversity
6: Race and Human Variation
Human Taxonomy
2.4: Human Taxonomy
hydroxyapatite
2.5: Osteology
hyperpnea
6.5.2: Adjustments and Adaptations
hypothalamus
6.5.2: Adjustments and Adaptations
hypothesis
2.1.2: Scientific Method

I
intermembral index (IMI)
4.2.2: What is a Hominim

K
Kenyanthropus platyops
4.3.6: Kenyanthropus platyops

L
Laetoli footprints
4.3.4: Australopithecus afarensis
Lamarckian Evolution
1.2.3: Evolutionary Theory (Part 1)
Lascaux Cave
5.1.22: Homo sapiens
Levels of culture
1.2.3: Evolutionary Theory (Part 1)
Linnaean System
3.5.1: Classification - Ordering the Natural World
Little Foot
4.3.7: Australopithecus prometheus or africanus

M
macroevolution
2.6: Macroevolution
marsupials
2.3: Mammals
megadontia quotient
5.1.4: Paranthropus boisei
melanin
6.5.2: Adjustments and Adaptations
melanocytes
6.5.2: Adjustments and Adaptations
Mendel's Law of Independent Assortment
2.1.6: Mendelian Genetics
Miocene Epoch
4.2.3: Miocene Epoch
Modern Synthesis
2.1.7: Modern Synthesis
monogenetic
6.1: The History of "Race" Concepts
monotremes
2.3: Mammals
mutation
1.2.6: "Forces" of Evolution
mutations
6.2: Human Variation in Biological Anthropology Today

N
natural selection
6.2: Human Variation in Biological Anthropology Today
neandertals
5.1.21: Homo neanderthalensis

O
occipital bun
5.1.17: Homo antecessor
Old World monkeys
3.5.8: More Haplorhines (Old World Monkeys)
Oldowan choppers
5.1.9: Homo habilis
Olduvai Gorge
5.1.4: Paranthropus boisei
Orrorin tugenensis
4.2.3.3: Orrorin tugenensis
osteology
2.5: Osteology
3.5.2: Primate Skeletal Anatomy
Othering
6.1: The History of "Race" Concepts
overpopulation
7.1.2: Doom and Gloom

P
paleoanthropology
4.1: Introduction to Paleoanthropology
4.2: Introduction to Paleoanthropology
Paleospecies
2.7: Species vs. Paleospecies
Parallel evolution
3.5.1: Classification - Ordering the Natural World
Paranthropines
5.1.2: Paranthropines
Paranthropus aethiopicus
5.1.3.1: Australopithecus/Paranthropus aethiopicus
Paranthropus boisei
5.1.4: Paranthropus boisei
Paranthropus robustus
5.1.5: Paranthropus robustus
phenotypic plasticity
6.5.1: Stress and Homeostasis
physical anthropology
6.1: The History of "Race" Concepts
placental mammals
2.3: Mammals
plate tectonics
2.9: Adaptive Radiation
Pliocene Epoch
4.3: Pliocene Epoch
polygenetic
6.1: The History of "Race" Concepts
polygyny
3.4: Primate Social Organization
polymorphism
2.1.8: Modern Human Variation
polymorphisms
6.3: Talking About Human Biological Variation Going Forward
polyphenism
6.5.1: Stress and Homeostasis
population
6.2: Human Variation in Biological Anthropology Today
population bottlnecking
6.2: Human Variation in Biological Anthropology Today
prejudice
6: Race and Human Variation
Primate Ecology
3.5.5: Primate Ecology
primatology
3.4: Primate Social Organization
3.5: Non-Human Primates
Protomammals
2.3: Mammals

R
race
6: Race and Human Variation
racism
6: Race and Human Variation
radiation
6.5.2: Adjustments and Adaptations
Recent African Origin (RAO) model
5.1.19: Homo heidelbergensis
Regional Continuity Model (RCM)
5.1.19: Homo heidelbergensis

S
Sahelanthropus tchadensis
4.2.3.2: Sahelanthropus tchadensis
scientific method
2.1.2: Scientific Method
Scientific Revolution
6.1: The History of "Race" Concepts
sickle cell anemia
1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria
social Darwinism
1.2.5: Evolutionary Theory (Part 3)

Social Organization (Primate)
3.4: Primate Social Organization
speciation
2.1.7: Modern Synthesis
2.6: Macroevolution
species
1.2.3: Evolutionary Theory (Part 1)
2.7: Species vs. Paleospecies
Strepsirhines
3.5.6: Overviews of Living Primates (Strepsirhines)
Strepsirhini classification
3.5.4: Taxonomy of the Living Primates
stressors
6.5.1: Stress and Homeostasis
subcultures
1.2.6: "Forces" of Evolution
T
tarsiers
3.5.7: Overviews of Living Primates (Haplorhines)
Taung Child
4.3.8: Australopithecus africanus
taxonomy
3.2: Primate Classification
traditional classification
3.5.4: Taxonomy of the Living Primates
Trisomy 21
1.2.7: Genetics, Cellular Biology, and Variation (Part 1)
typologies
6.1: The History of "Race" Concepts
V
vasoconstriction
6.5.2: Adjustments and Adaptations

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: Physical Anthropology Anthro-1

Webpages: 126

Applicable Restrictions: Noncommercial, No Derivatives

All licenses found:

- CC BY-NC-SA 4.0: 34.1% (43 pages)
- CC BY-NC-ND 4.0: 23.8% (30 pages)
- Undeclared: 19.8% (25 pages)
- CC BY-SA 4.0: 16.7% (21 pages)
- CC BY-NC-SA 3.0: 3.2% (4 pages)
- CC BY-NC 4.0: 2.4% (3 pages)

By Page

- Physical Anthropology Anthro-1 - *Undeclared*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Introduction to Biological/Physical Anthropology - *Undeclared*
    - 1.1: Introduction to Physical Anthropology - *CC BY-NC-ND 4.0*
      - 1.1.1: Science - *CC BY-NC-ND 4.0*
      - 1.1.2: Anthropology as a Science - *CC BY-NC-ND 4.0*
    - 1.2: Introduction to Biology - *CC BY-NC-ND 4.0*
      - 1.2.1: Introduction - *CC BY-NC-ND 4.0*
      - 1.2.2: Scale, Human Space, Powers of Ten for Physical Anthropology - *CC BY-NC-ND 4.0*
      - 1.2.3: Evolutionary Theory (Part 1) - *CC BY-NC-ND 4.0*
      - 1.2.4: Evolutionary Theory (Part 2) - *CC BY-NC-ND 4.0*
      - 1.2.5: Evolutionary Theory (Part 3) - *CC BY-NC-ND 4.0*
      - 1.2.6: "Forces" of Evolution - *CC BY-NC-ND 4.0*
      - 1.2.7: Genetics, Cellular Biology, and Variation (Part 1) - *CC BY-NC-ND 4.0*
      - 1.2.8: Genetics, Cellular Biology, and Variation (Part 2) - *CC BY-NC-ND 4.0*
      - 1.2.9: Summary Example- Holism in Anthropology, Sickle Cell Anemia and Malaria - *CC BY-NC-ND 4.0*
    - 1.3: What is Biological Anthropology? - *CC BY-SA 4.0*
  - 2: Mechanisms and Evolutionary Thought - *Undeclared*
    - 2.1: Evolutionary Theory - *CC BY-SA 4.0*
      - 2.1.1: What is Biological Anthropology? - *CC BY-SA 4.0*
      - 2.1.2: Scientific Method - *CC BY-SA 4.0*
      - 2.1.3: History of Evolutionary Thought - *CC BY-SA 4.0*
      - 2.1.4: Darwinian Evolution - *CC BY-SA 4.0*
      - 2.1.5: Cell Biology and Basic Genetics - *CC BY-SA 4.0*
      - 2.1.6: Mendelian Genetics - *CC BY-SA 4.0*
      - 2.1.7: Modern Synthesis - *CC BY-SA 4.0*
      - 2.1.8: Modern Human Variation - *CC BY-SA 4.0*
    - 2.2: Taxonomy - *CC BY-NC-ND 4.0*
    - 2.3: Mammals - *CC BY-NC-ND 4.0*
    - 2.4: Human Taxonomy - *CC BY-NC-ND 4.0*
    - 2.5: Osteology - *CC BY-NC-ND 4.0*
    - 2.6: Macroevolution - *CC BY-NC-ND 4.0*
    - 2.7: Species vs. Paleospecies - *CC BY-NC-ND 4.0*
    - 2.8: Interspecific vs. Intraspecific Variation - *CC BY-NC-ND 4.0*
    - 2.9: Adaptive Radiation - *CC BY-NC-ND 4.0*
    - 2.10: Analogy vs. Homology - *CC BY-NC-ND 4.0*
  - 3: Primates - *Undeclared*
    - 3.1: Introduction to Primatology - *CC BY-NC-ND 4.0*
    - 3.2: Primate Classification - *CC BY-NC-SA 4.0*
    - 3.3: Primate Evolution - *CC BY-NC-SA 4.0*
    - 3.4: Primate Social Organization - *CC BY-NC-SA 4.0*
    - 3.5: Non-Human Primates - *CC BY-SA 4.0*

- 3.5.1: Classification - Ordering the Natural World - *CC BY-SA 4.0*
- 3.5.2: Primate Skeletal Anatomy - *CC BY-SA 4.0*
- 3.5.3: Modern Primates - *CC BY-SA 4.0*
- 3.5.4: Taxonomy of the Living Primates - *CC BY-SA 4.0*
- 3.5.5: Primate Ecology - *CC BY-SA 4.0*
- 3.5.6: Overviews of Living Primates (Strepsirhines) - *CC BY-SA 4.0*
- 3.5.7: Overviews of Living Primates (Haplorhines) - *CC BY-SA 4.0*
- 3.5.8: More Haplorhines (Old World Monkeys) - *CC BY-SA 4.0*
- 3.5.9: Even more Haplorhines (The Apes) - *CC BY-SA 4.0*
- 3.5.10: Primate Evolution - *CC BY-SA 4.0*
- 3.6: Ethology - *CC BY-NC-ND 4.0*
- 3.7: Conservation - *CC BY-NC-ND 4.0*
- 3.8: Lab 5 and 6- Primates - *CC BY-NC-SA 3.0*

4: Paleontology and Primate Evolution - *Undeclared*

- 4.1: Introduction to Paleoanthropology - *CC BY-NC-ND 4.0*
- 4.2: Introduction to Paleoanthropology - *CC BY-NC-SA 4.0*
  - 4.2.1: Paleoanthropology - *CC BY-NC-SA 4.0*
  - 4.2.2: What is a Hominim - *CC BY-NC-SA 4.0*
  - 4.2.3: Miocene Epoch - *CC BY-NC-SA 4.0*
    - 4.2.3.1: Introduction to the Miocene Epoch - *CC BY-NC-SA 4.0*
    - 4.2.3.2: Sahelanthropus tchadensis - *CC BY-NC-SA 4.0*
    - 4.2.3.3: Orrorin tugenensis - *CC BY-NC-SA 4.0*
    - 4.2.3.4: Ardipithecus ramidus and Ardipithecus kadabba - *CC BY-NC-SA 4.0*
    - 4.2.3.5: Early Hominid Cranium Comparison Checklist - *CC BY-NC-SA 3.0*
- 4.3: Pliocene Epoch - *CC BY-NC-SA 4.0*
  - 4.3.1: Introduction to the Pliocene Epoch - *CC BY-NC-SA 4.0*
  - 4.3.2: Australopithecus anamensis - *CC BY-NC-SA 4.0*
  - 4.3.3: Gracile Australopiths - *CC BY-NC-SA 4.0*
  - 4.3.4: Australopithecus afarensis - *CC BY-NC-SA 4.0*
  - 4.3.5: Australopithecus bahrelghazali - *CC BY-NC-SA 4.0*
  - 4.3.6: Kenyanthropus platyops - *CC BY-NC-SA 4.0*
  - 4.3.7: Australopithecus prometheus or africanus - *CC BY-NC-SA 4.0*
  - 4.3.8: Australopithecus africanus - *CC BY-NC-SA 4.0*
  - 4.3.9: Middle Hominid Cranium Comparison Checklist - *CC BY-NC-SA 3.0*

5: Human Fossil Record - *Undeclared*

- 5.1: Pleistocene Epoch - *CC BY-NC-SA 4.0*
  - 5.1.1: Introduction to the Pleistocene Epoch - *CC BY-NC-SA 4.0*
  - 5.1.2: Paranthropines - *CC BY-NC-SA 4.0*
  - 5.1.3: Australopithecus - *Undeclared*
    - 5.1.3.1: Australopithecus/Paranthropus aethiopicus - *CC BY-NC-SA 4.0*
  - 5.1.4: Paranthropus boisei - *CC BY-NC-SA 4.0*
  - 5.1.5: Paranthropus robustus - *CC BY-NC-SA 4.0*
  - 5.1.6: Australopithecus garhi - *CC BY-NC-SA 4.0*
  - 5.1.7: Australopithecus sediba - *CC BY-NC-SA 4.0*
  - 5.1.8: Genus Homo - *CC BY-NC-SA 4.0*
  - 5.1.9: Homo habilis - *CC BY-NC-SA 4.0*
  - 5.1.10: Homo rudolfensis - *CC BY-NC-SA 4.0*
  - 5.1.11: Homo species indeterminate - *CC BY-NC-SA 4.0*
  - 5.1.12: Homo naledi - *CC BY-NC-SA 4.0*
  - 5.1.13: The “erectus Grade” - *CC BY-NC-SA 4.0*
  - 5.1.14: Homo ergaster - *CC BY-NC-SA 4.0*
  - 5.1.15: Homo erectus - *CC BY-NC-SA 4.0*
  - 5.1.16: Homo georgicus - *CC BY-NC-SA 4.0*
  - 5.1.17: Homo antecessor - *CC BY-NC-SA 4.0*
  - 5.1.18: Homo floresiensis - *CC BY-NC-SA 4.0*
  - 5.1.19: Homo heidelbergensis - *CC BY-NC-SA 4.0*
  - 5.1.20: The Denisovans - *CC BY-NC-SA 4.0*
  - 5.1.21: Homo neanderthalensis - *CC BY-NC-SA 4.0*
  - 5.1.22: Homo sapiens - *CC BY-NC-SA 4.0*
  - 5.1.23: Recent Hominid Cranium Comparison Checklist - *CC BY-NC-SA 3.0*

6: Race and Human Variation - *CC BY-NC 4.0*

- 6.1: The History of "Race" Concepts - *Undeclared*
- 6.2: Human Variation in Biological Anthropology Today - *Undeclared*
- 6.3: Talking About Human Biological Variation Going Forward - *Undeclared*
- 6.4: Key Terms - *Undeclared*
- 6.5: Human Variation- An Adaptive Significance Approach - *CC BY-NC 4.0*
  - 6.5.1: Stress and Homeostasis - *Undeclared*
  - 6.5.2: Adjustments and Adaptations - *Undeclared*

- 6.5.3: Review Questions - *Undeclared*
- 6.5.4: Key Terms - *Undeclared*

- 7: New Frontiers - *Undeclared*
  - 7.1: Homo sapiens Futures - Doom, Gloom, and Hope? - *CC BY-NC-ND 4.0*
    - 7.1.1: Introduction - *CC BY-NC-ND 4.0*
    - 7.1.2: Doom and Gloom - *CC BY-NC-ND 4.0*
    - 7.1.3: Hope - *CC BY-NC-ND 4.0*
  - 7.2: Bioarchaeology and Forensic Anthropology - *CC BY-NC 4.0*

- Back Matter - *Undeclared*
  - Index - *Undeclared*
  - Glossary - *Undeclared*
  - Detailed Licensing - *Undeclared*